

ON THE GLIVENKO-CANTELLI THEOREM FOR REAL-VALUED EMPIRICAL FUNCTIONS OF STATIONARY α -MIXING AND β -MIXING SEQUENCES

OUSMANE COULIBALY¹ AND HAROUNA SANGARÉ²

ABSTRACT. In this paper we extend the classical Glivenko-Cantelli theorem to real-valued empirical functions under dependence structures characterised by α -mixing and β -mixing conditions. We investigate sufficient conditions ensuring that families of real-valued functions exhibit the Glivenko-Cantelli (GC) property in these dependence settings. Our analysis focuses on function classes satisfying uniform entropy conditions, and we establish conditions on mixing coefficients to obtain GC theorems.

¹ Msc in Probabilities and Statistics
 Université des Sciences, des Techniques et des Technologies de Bamako (USTT-B), Mali
 Personal email : ocouwi@gmail.com

² DER MI, Faculté des Sciences et Techniques (FST), USTT-B, Mali
 LERSTAD, Gaston Berger University, Saint-Louis, Sénégal.
 Researcher at IMHO [<https://imhotepsciences.org>]
 Imhotep International Mathematical Center (imh-imc), <https://imhotepsciences.org>
 Institutional emails : harouna.sangare@usttb.edu.ml, harouna.sangare@mesrs.ml,
 sangare.harouna@ugb.edu.sn
 Personal email : harounasangareusttb@gmail.com

Keywords. α -mixing, β -mixing, Empirical process, Stationarity, Entropy number, Law of large numbers, Uniform convergence, Glivenko-Cantelli class

AMS 2020 Mathematics Subject Classification: 62G20; 62G30; 60F99; 60G10

1. INTRODUCTION

Empirical process theory plays a crucial role in probability and statistics, providing tools for analyzing the convergence properties of empirical measures and function classes. The classical Glivenko-Cantelli theorem asserts that the empirical distribution function converges uniformly to the true distribution function for independent and identically distributed (iid) samples. However, real-world data often exhibit dependence, necessitating an extension of this theory to dependent structures.

In this paper, we study the uniform convergence properties of real-valued empirical functions under α -mixing and β -mixing conditions. These mixing conditions describe dependence structures commonly encountered in time series analysis, stochastic processes, and machine learning. Our goal is to establish sufficient conditions for the Glivenko-Cantelli property in these settings and to derive implications for statistical applications.

Preliminaries and Definitions. Let $\{X, (X_n)_{n \geq 1}\}$ be a sequence of real-valued random variables (rrv's) defined on the same probability space $(\Omega, \mathcal{A}, \mathbb{P})$ and associated to the same cumulative distribution function (cdf) F . Let $\mathcal{L}_2(\mathbb{P}_X)$ be the class of all measurable functions $f : \mathbb{R} \rightarrow \mathbb{R}$ with $\mathbb{E}f(X)^2 < +\infty$. Let us define the functional empirical probability based on the n first observations, $n \geq 1$, by

$$(1.1) \quad \mathbb{P}_n(f) = \frac{1}{n} \sum_{i=1}^n f(X_i), \quad f \in \mathcal{L}_2,$$

which will be centered at the mathematical expectations

$$\mathbb{E}\mathbb{P}_n(f) = \mathbb{P}_X(f), \quad f \in \mathcal{L}_2.$$

In this paper, we are interested by functional Glivenko-Cantelli classes, that is classes $\mathcal{F} \subset \mathcal{L}_2$ for which we have

$$\sup_{f \in \mathcal{F}} |\mathbb{P}_n(f) - \mathbb{E}\mathbb{P}_n(f)| = \|\mathbb{P}_n - \mathbb{E}\mathbb{P}_n\|_{\mathcal{F}} \rightarrow 0, \quad \text{almost surely, as } n \rightarrow +\infty.$$

We might also consider a class \mathcal{C} of measurable sets in \mathbb{R} and define

$$(1.2) \quad \mathbb{P}_n(C) = \frac{1}{n} \sum_{i=1}^n \text{Card}\{X_i \in C\}, \quad C \in \mathcal{C},$$

with

$$\mathbb{E}\mathbb{P}_n(C) = \mathbb{P}_X(C) = \mathbb{P}(X \in C).$$

Taking $\mathcal{F}_c = \{\mathbb{I}_{]-\infty, x]}, x \in \mathbb{R}\}$ in Definition (1.1) or $\mathcal{C}_c = \{]-\infty, x], x \in \mathbb{R}\}$ in Definition (1.2), leads to the classical empirical function

$$F_n(x) = \frac{1}{n} \text{Card}\{i \in [1, n], X_i \leq x\}, \quad x \in \mathbb{R},$$

which in turn gives the Glivenko-Cantelli law for independent and identically distributed (*iid*) sequences $(X_n)_{n \geq 1}$ under the form :

$$(1.3) \quad \sup_{x \in \mathbb{R}} |F_n(x) - F(x)| \rightarrow 0, \quad \text{almost surely, as } n \rightarrow +\infty.$$

Such a result, also known as the fundamental theorem of statistics, is the frequentist paradigm (as opposed to the Bayesian paradigm) in statistics. In the form of (1.3), the Glivenko-Cantelli law has gone through a large number of studies for a variety of types of dependence. It has also been used extensively in statistical theory, both for finding asymptotically efficient estimators and for related statistical tests based on the Donsker

theorem, which we will not study here.

To give some examples, we cite the following results: Billingsley (1968) [Billingsley \(1968\)](#) showed the convergence in law on $D[0, 1]$ of the empirical process for ϕ -mixing rrv's under the condition $\sum_{k>0} k^2 \sqrt{\phi(k)} < +\infty$. Yoshihara (1975) [Yoshihara \(1975\)](#) obtained the same result for α mixing under the condition on the mixing coefficient $\alpha_n = O(n^{-a})$ with $a > 3$. This result was first improved by Shao (1995) [Shao \(1995\)](#) by assuming only $a > 2$, then by Shao and Yu (1996) [Shao and Yu \(1996\)](#) by assuming $a > 1 + \sqrt{2}$. Rio (2017) [Rio \(2017\)](#) obtained an optimal condition $a > 1$. Similar results are given by Shao and Yu (1996) [Shao and Yu \(1996\)](#) for ρ -mixing rrv's, Doukhan *et al.* (1995) [Doukhan et al. \(1995\)](#) for β -mixing rrv's. Next, Shao and Yu (1996) [Shao and Yu \(1996\)](#) weakened the covariance condition $a > \frac{3+\sqrt{33}}{2}$. Louhichi (2000) [Louhichi \(2000\)](#) gave another proof and an improvement with $a > 4$. Sangaré *et al.* (2020) [Sangaré et al. \(2020\)](#) gave a functional Glivenko-Cantelli theorem for ϕ mixing rrv's under the mixing coefficient condition $\phi_n = O(n^{-\frac{4}{1-\delta}})$ with $0 < \delta < 1$.

However, functional versions do not seem to have been developed for dependent data, although they are much more interesting than restricting to the particular case of \mathcal{F}_c . Moreover, the functional version has the advantage of being linear in the sense that

$$\forall (a, b) \in \mathbb{R}^2, \forall (f, g) \in (\mathcal{L}_1(\mathbb{P}_X))^2, \mathbb{P}_n(af + bg) = a\mathbb{P}_n(f) + b\mathbb{P}_n(g),$$

which allows the use of more mathematical latitudes.

In this paper, we use general Glivenko-Cantelli classes established by Sangaré *et al.* (2020) [Sangaré et al. \(2020\)](#), for real-valued empirical functions of dependent data and particularize them for α -mixing and β -mixing random variables to obtain GC classes.

The main results are Theorem [5](#), as a general GC theorem for arbitrary stationary sequences, Proposition [1](#), a Glivenko-Cantelli law for the real-valued empirical functions of stationary α -mixing sequences, and Proposition [2](#), a Glivenko-Cantelli law for the real-valued empirical functions of stationary β -mixing sequences.

This functional approach requires the use of concentration numbers, which we define in the next section. We will also need to make a number of calls to dependency types such as α -mixing and β -mixing, and other tools.

This leads us to the following organisation of the paper. In section [3](#), we recall the general strong law of large numbers of Sangaré and Lo (2015) [Sangaré and Lo \(2015\)](#), which is crucial for the proofs of Sangaré *et al.* (2020) [Sangaré et al. \(2020\)](#) general GC classes used in our results, and to give a brief introduction to the notions of α -mixing and β -mixing. But before that, section [2](#) is devoted to entropy numbers and Vapnik-Červonenkis classes. In section [4](#), we give our GC class regarding real-valued empirical functions for arbitrary stationary rv's sequences and particularize it for stationary α -mixing and

β -mixing rrv's sequences. Section 5 deals with conclusion and perspectives. The paper ends with [Appendix](#), where the detailed proof of the arbitrarily stationary Glivenko-Cantelli theorem is postponed.

2. ENTROPY NUMBERS AND VAPNIK-ČERVONENKIS

The following reminder about entropy numbers, which comes from combinatorial theory, might be perceived as making the paper heavier. However, we believe that it may help the reader who is not familiar with such techniques. Let $(E, \|\circ\|, \leq)$ be an ordered real normed space, which means that the order is compatible with the operations in the following sense

$$\forall(\lambda, z) \in \mathbb{R}_+ \vee \{0\} \times E, \forall(x, y) \in E^2, (x < y) \rightarrow ((x + z < y + z) \text{ and } (\lambda x < \lambda y)).$$

A bracket set of level $\varepsilon > 0$ in E is any set of the form

$$B(\ell_1, \ell_2, \varepsilon) = \{x \in E, \ell_1 \leq x \leq \ell_2\}, \ell_1 \leq \ell_2 \in E \text{ and } \|\ell_2 - \ell_1\| < \varepsilon.$$

For any subset F of E , the bracketing entropy number at level $\varepsilon > 0$, denoted $N_{(F, \|\circ\|, \varepsilon)}$ is the minimum of the numbers $p \geq 1$ for which we have p bracket sets (or simply p brackets) at level ε covering F , where the ℓ_i 's do not necessarily belong to F . The bracketing entropy number is closely related to the Vapnik-Červonenkis index (VC-index) of a Vapnik-Červonenkis set or class (VC-set or VC-class).

To define a VC-set, we need to recall some definitions. A subset D of $B \subset E$ is picked out by a subclass \mathcal{C} of the power set $\mathcal{P}(E)$ from B if and only if D is element of $\{B \cap C, C \in \mathcal{C}\}$. Next, B is shattered by \mathcal{C} if and only if all subsets of B are picked out by \mathcal{C} from B . Finally, the class \mathcal{C} is a VC-class if and only if there exists an integer $n \geq 1$ such that no set of cardinality n is shattered by \mathcal{C} . The minimum of those numbers n minus one is the index of that VC-class. The VC-class is the most quick way to bound bracketing entropy numbers $N_{(F, \|\circ\|, \varepsilon)}$, as stated in [van der Vaart and Wellner \(1996\)](#) : If \mathcal{C} is a VC-class of index $I(\mathcal{C})$, then

$$(2.1) \quad N_{(F, \|\circ\|, \varepsilon)} \leq KI(\mathcal{C})(4e)^{I(\mathcal{C})}(1/\varepsilon)^{rI(\mathcal{C}-1)},$$

where $K > 0$ and $r > 1$ are universal constants.

Finally, for a class \mathcal{F} of real-valued functions $f : E \rightarrow \mathbb{R}$, we may define the bracketing entropy number associated with \mathcal{F} to be the bracketing entropy number of the class $\mathcal{C}(\mathcal{F})$ of sub-graphs f of elements $f \in \mathcal{F}$

$$S_g(f) = \{(x, t) \in E \times \mathbb{R}, f(x) > t\},$$

in $E^* = E \times \mathbb{R}$ endowed with the product norm which is still a normed space. We denote by $N_{(F, \|\circ\|, \varepsilon)}$ to distinguish with the bracketing number using the norm of the functions $f \in \mathcal{F}$. Also, \mathcal{F} is a VC-supg-class if and only if $\mathcal{C}(\mathcal{F})$ is a VC-class and $I(\mathcal{F}) = IC(\mathcal{F})$. We will use such entropy numbers to formulate VC-classes.

In the next section, we describe the main tools that we used in our results and provide a brief reminder of the concepts of α mixing and β mixing.

3. THE SANGARÉ-LO SLNN, THE SANGARÉ-LO-TRAORÉ GENERAL FUNCTIONAL GLIVENKO-CANTELLI CLASSES, α -MIXING AND β -MIXING CONCEPT

In this section, we have a look at the tools at the heart of our findings.

3.1. General strong law of large numbers. We begin with this useful result proved by Sangaré and Lo (2015).

Theorem 1. (Sangaré and Lo (2015)). *Let X_1, X_2, \dots be an arbitrary sequence of rv's, and let $(f_{i,n})_{i \geq 1}$ be a sequence of measurable functions such that $\text{Var}[f_{i,n}(X_i)] < +\infty$, for $i \geq 1$ and $n \geq 1$. If for some $\delta, 0 < \delta < 3$*

$$(3.1) \quad C_1 = \sup_{n \geq 1} \sup_{q \geq 1} \text{Var} \left(\frac{1}{q^{(3-\delta)/4}} \sum_{i=1}^q f_{i,n}(X_i) \right) < +\infty$$

and for some $0 < \delta < 3$,

$$(3.2) \quad C_2 = \sup_{n \geq 0} \sup_{k \geq 0} \sup_{q: q^2+1 \leq k \leq (q+1)^2} \sup_{k \leq j \leq (q+1)^2} \text{Var} \left(\frac{1}{q^{(3-\delta)/2}} \sum_{i=1}^{j-q^2+1} f_{q^2+i,n}(X_{q^2+i}) \right) < +\infty$$

hold, then

$$\frac{1}{n} \sum_{i=1}^n (f_{i,n}(X_i) - \mathbb{E}(f_{i,n}(X_i))) \rightarrow 0, \text{ almost surely, as } n \rightarrow +\infty.$$

Remark 1. *We say that the sequence X_1, X_2, \dots, X_n satisfies the **(GCIP)** whenever Conditions (3.1) and (3.2) hold, and we denote \mathcal{C} , the class of measurable functions for which **(GCIP)** holds.*

In the next subsection, we recall general functional Glivenko-Cantelli classes due to Sangaré et al. (2020). These results will be use in Section 4 to find GC-classes for real-valued empirical function under α -mixing and β -mixing conditions.

3.2. General functional GC-classes. Let us begin by giving a slight different version of Theorem (2.4.1 in van der Vaart and Wellner (1996), page 122), established by Sangaré et al. (2020).

Theorem 2. (Sangaré et al. (2020)) *Let X, X_1, X_2, \dots be an arbitrary stationary sequence of rrv's with common cdf F .*

a) *Let \mathcal{C} be a class of measurable set such that*

(a1) for any $\varepsilon > 0$, $N_{[]}(\mathcal{C}, |\cdot|, \varepsilon) < +\infty$,

(a2) for any $C \in \mathcal{C}$, $\mathbb{P}_n(C) \rightarrow \mathbb{P}_X(C)$, as $n \rightarrow +\infty$.

Then \mathcal{C} is GC-class, that is

$$(3.3) \quad \lim_{n \rightarrow +\infty} \sup_{C \in \mathcal{C}} |\mathbb{P}_n(C) - \mathbb{P}_X(C)| = 0.$$

b) Let \mathcal{F} be a class of measurable set such that

(b1) for any $\varepsilon > 0$, $N_{[]}(\mathcal{F}, \|\cdot\|_{\mathcal{L}_2(\mathbb{P}_X)}, \varepsilon) < +\infty$,

(b2) for any $f \in \mathcal{F}$, $\mathbb{P}_n(f) \rightarrow \mathbb{P}_X(f)$, as $n \rightarrow +\infty$.

Then \mathcal{F} is GC-class, that is

$$(3.4) \quad \lim_{n \rightarrow +\infty} \sup_{f \in \mathcal{F}} |\mathbb{P}_n(f) - \mathbb{P}_X(f)| = 0.$$

Proof. The proof of the Theorem 2 is postponed in the Appendix (Section (6, page 15)) \square

Remark 2. If we apply Part (a) to \mathcal{C}_c , Condition (a1) can be dropped since it is VC-class of index 2. Indeed, for $A = \{x_1, x_2, x_3\}$ with $x < x_2 < x_3$. The subset $\{x_2\}$ cannot be picked out by \mathcal{C}_c from A since, for any $x \in \mathbb{R}$, $] -\infty, x] \cap A$ will be of on the five sets : \emptyset (for $x \leq x_1$), $\{x_1\}$ (for $x_1 < x \leq x_2$), $\{x_1, x_2\}$ (for $x_2 < x \leq x_3$) and A (for $x > x_3$).

Then finding GC-classes for \mathcal{C}_c reduces to establishing Condition (b1). For now, we are going to focus on the application of GC-classes of the functional empirical process for \mathcal{C}_c , that is, on results as in Formula (1.3). The focus will be on types of dependence, given it is known that \mathcal{C}_c is GC-class for *iid* data. Using the GCIP conditions in Theorem 1 leads to the applicable results as follows.

Theorem 3. (*Sangaré et al. (2020)*) Let X, X_1, X_2, \dots be an arbitrary stationary sequence of rrv's with common cdf F .

a) Let \mathcal{C} be a class of measurable set such that

(a1) for any $\varepsilon > 0$, $N_{[]}(\mathcal{C}, |\cdot|, \varepsilon) < +\infty$,

(a2) for any $C \in \mathcal{C}$, the following conditions hold : for some $0 < \delta < 3$,

$$(3.5) \quad \sup_{q \geq 1} \frac{1}{q^{(3-\delta)/2}} \sum_{i,j=1}^q [\mathbb{P}(X_i \in C, X_j \in C) - \mathbb{P}(X_i \in C)\mathbb{P}(X_j \in C)] < +\infty.$$

and

$$(3.6) \quad \sup_{k \geq 1} \sup_{q: q^2+1 \leq k \leq (q+1)^2} \sup_{k \leq j \leq (q+1)^2} \frac{1}{q^{(3-\delta)}} \sum_{i, \ell=1}^{j-q^2+1} \left[\begin{array}{l} \mathbb{P}(X_{q^2+i} \in C, X_{q^2+\ell} \in C) \\ -\mathbb{P}(X_{q^2+i} \in C) \mathbb{P}(X_{q^2+\ell} \in C) \end{array} \right] < +\infty.$$

Then \mathcal{C} is GC-class, that is

$$(3.7) \quad \lim_{n \rightarrow +\infty} \sup_{C \in \mathcal{C}} |\mathbb{P}_n(C) - \mathbb{P}_X(C)| = 0.$$

b) Let \mathcal{F} be a class of measurable set such that

(b1) for any $\varepsilon > 0$, $N_{[]}(\mathcal{F}, \|\circ\|_{\mathcal{L}_2(\mathbb{P}_X)}, \varepsilon) < +\infty$,

(b2) for any $f \in \mathcal{F}$, the following conditions hold : for some $0 < \delta < 3$,

$$(3.8) \quad \sup_{q \geq 1} \text{Var} \left(\frac{1}{q^{(3-\delta)/4}} \sum_{i=1}^q f(X_i) \right) < +\infty$$

and

$$(3.9) \quad \sup_{k \geq 0} \sup_{q: q^2+1 \leq k \leq (q+1)^2} \sup_{k \leq j \leq (q+1)^2} \text{Var} \left(\frac{1}{q^{(3-\delta)/2}} \sum_{i=1}^{j-q^2+1} f(X_{q^2+i}) \right) < +\infty.$$

Then \mathcal{F} is GC-class, that is

$$(3.10) \quad \lim_{n \rightarrow +\infty} \sup_{f \in \mathcal{F}} |\mathbb{P}_n(f) - \mathbb{P}_X(f)| = 0.$$

Proof. It follows the lines of that of Theorem 2 in the Appendix (Section (6, page 15)) □

Since we are also going to provide results for sequences verifying α -mixing and β -mixing conditions, we make a brief summary of these notions.

3.3. α -mixing. Let us have a brief recall on ϕ -mixing. Let $(\Omega, \mathcal{A}, \mathbb{P})$ be a probability space and $\mathcal{A}_1, \mathcal{A}_2$ two sub σ -algebras of \mathcal{A} . The α -mixing coefficient is given by

$$\alpha(\mathcal{A}_1, \mathcal{A}_2) = \sup \{ |\mathbb{P}(A)\mathbb{P}(B) - \mathbb{P}(A \cap B)|, A \in \mathcal{A}_1, B \in \mathcal{A}_2 \}$$

and

$$0 \leq \alpha(\mathcal{A}_1, \mathcal{A}_2) \leq 1/4.$$

We remind that we have (see [Doukhan \(1994\)](#))

Theorem 4. (Covariance inequality). *Let X and Y be measurable random variables with respect to \mathcal{A} and \mathcal{B} respectively, We have*

$$|\mathbb{C}ov(X, Y)| \leq 8\alpha^{\frac{1}{r}}(\mathcal{A}, \mathcal{B})\|X\|_p\|Y\|_q \text{ for } p, q, r \geq 1 \text{ and } \frac{1}{p} + \frac{1}{q} + \frac{1}{r} = 1.$$

To prove the above theorem, we need the following Lemma.

Lemma 1. *Under the same assumptions, we have the following result*

$$|\mathbb{C}ov(X, Y)| \leq 4\alpha(\mathcal{A}, \mathcal{B})\|X\|_\infty\|Y\|_\infty.$$

Proof. (of Lemma 1) Let $A = \text{sign}(\mathbb{E}(X | \mathcal{B}) - \mathbb{E}(X))$ and $B = \text{sign}(\mathbb{E}(Y | \mathcal{A}) - \mathbb{E}(Y))$. By the conditional expectation property, we have

$$\begin{aligned} |\mathbb{C}ov(X, Y)| &= |\mathbb{E}(XY) - \mathbb{E}(X)\mathbb{E}(Y)| \\ &= |\mathbb{E}(X\mathbb{E}(Y | \mathcal{A}) - \mathbb{E}(Y))| \\ &\leq \|X\|_\infty \mathbb{E}|\mathbb{E}(Y | \mathcal{A}) - \mathbb{E}(Y)| \\ &\leq \|X\|_\infty \mathbb{E}|B[(Y | \mathcal{A}) - \mathbb{E}(Y)]| \\ &\leq \|X\|_\infty |\mathbb{E}(BY) - \mathbb{E}(B)\mathbb{E}(Y)| \end{aligned}$$

and since

$$\begin{aligned} |\mathbb{E}(BY) - \mathbb{E}(B)\mathbb{E}(Y)| &= |\mathbb{E}[\mathbb{E}(YB | \mathcal{B}) - \mathbb{E}(B)\mathbb{E}(Y)]| \\ &= |\mathbb{E}[Y\mathbb{E}(B | \mathcal{B}) - \mathbb{E}(B)]| \\ &\leq \|Y\|_\infty \mathbb{E}|\mathbb{E}(B | \mathcal{B}) - \mathbb{E}(B)| \\ &\leq \|X\|_\infty \mathbb{E}|A[(B | \mathcal{B}) - \mathbb{E}(B)]| \\ &\leq \|Y\|_\infty |\mathbb{E}(AB) - \mathbb{E}(B)\mathbb{E}(A)|, \end{aligned}$$

therefore

$$|\mathbb{C}ov(X, Y)| \leq \|X\|_\infty\|Y\|_\infty|\mathbb{E}(AB) - \mathbb{E}(A)\mathbb{E}(B)|.$$

Let now $A^+ = \{a = 1\}$, $A^- = \{a = -1\}$, $B^+ = \{b = 1\}$, $B^- = \{b = -1\}$, then

$$\begin{aligned} |\mathbb{E}(AB) - \mathbb{E}(A)\mathbb{E}(B)| &= [|\mathbb{P}(A^+ \cap B^+) - \mathbb{P}(A^+)\mathbb{P}(B^+)| \\ &\quad + |\mathbb{P}(A^- \cap B^-) - \mathbb{P}(A^-)\mathbb{P}(B^-)| \\ &\quad + |\mathbb{P}(A^+ \cap B^-) - \mathbb{P}(A^+)\mathbb{P}(B^-)| \\ &\quad + |\mathbb{P}(A^- \cap B^+) - \mathbb{P}(A^-)\mathbb{P}(B^+)|]. \end{aligned}$$

Hence

$$|\mathbb{C}ov(X, Y)| \leq 4\alpha(\mathcal{A}, \mathcal{B})\|X\|_\infty\|Y\|_\infty.$$

□

We are now ready to prove the theorem 4.

Proof. (of Theorem 4)

$$\|X\|_p < \infty, \|Y\|_p < \infty, 1 < p < \infty.$$

Define $\overline{X} = X\mathbb{I}_{\{|X| \leq a\}}$, $\underline{X} = X\mathbb{I}_{\{|X| > a\}}$ and write $X = \overline{X} + \underline{X}$. Thus, by the Lemma 1,

$$\text{Cov}(X, Y) = |\text{Cov}(\overline{X}, Y) + \text{Cov}(\underline{X}, Y)| \leq 4\alpha(\mathcal{A}, \mathcal{B})a\|Y\|_\infty + 2\|Y\|_\infty\mathbb{E}|\underline{X}|.$$

Now Markov's inequality leads to $\mathbb{E}|\underline{X}| \leq 2a\frac{\mathbb{E}|X|^p}{a^p}$. Set $\frac{\mathbb{E}|X|^p}{a^p} = \alpha(\mathcal{A}, \mathcal{B})$ then

$$|\text{Cov}(X, Y)| \leq 6\alpha^{1-1/p}(\mathcal{A}, \mathcal{B})\|Y\|_\infty\|X\|_\infty.$$

Let $\|X\|_p < \infty$ and $\|Y\|_q < \infty$, with $\frac{1}{p} + \frac{1}{q} < 1$.

Let us define $\overline{Y} = Y\mathbb{I}_{\{|Y| \leq b\}}$, $\underline{Y} = Y\mathbb{I}_{\{|Y| > b\}}$ and write $Y = \overline{Y} + \underline{Y}$, hence analogously

$$|\text{Cov}(Y, X)| \leq 6\alpha^{1-1/p}(\mathcal{A}, \mathcal{B})b\|X\|_p + 2b\|\underline{Y}\|.$$

By setting $\frac{\mathbb{E}|Y|^q}{a^q} = \alpha^{1-1/p}(A, B)$, it yields

$$|\text{Cov}(X, Y)| \leq 8\alpha^{1/r}(\mathcal{A}, \mathcal{B})\|Y\|_q\|X\|_p.$$

□

We define the strong mixing coefficient $\alpha(n)$ by

$$\alpha(n) = \sup \{ \alpha(\mathcal{F}_1^k, \mathcal{F}_{n+k}^\infty), k \in \mathbb{N}^* \}$$

where \mathcal{F}_j^ℓ is the σ -algebra generated by the variables $(X_i, j \leq i \leq \ell)$. We say that $(X_n)_{n \geq 1}$ is α -mixing if $\alpha(n) \rightarrow 0$ as $n \rightarrow +\infty$. For some further clarification on this point, the reader may have a quick look at [Doukhan \(1994\)](#) and [Rio \(2017\)](#).

3.4. β -mixing. Let $\mathcal{A}_{1,n}$ be the σ -algebra generated by the finite subset X_1, \dots, X_n and $\mathcal{A}_{2,n}$ that generated by the infinite subset $X_{n+k+1}, X_{n+k+2}, \dots$. We define the mixing of type β by means of the quantity

$$\beta(n, k) = \beta(\mathcal{A}_{1,n}, \mathcal{A}_{2,n}) = \mathbb{E}[\text{ess - sup} \{ |\mathbb{P}(B|\mathcal{A}_{1,n}) - \mathbb{P}(B)|, B \in \mathcal{A}_{2,n} \}]$$

and

$$0 \leq \beta(\mathcal{A}_{1,n}, \mathcal{A}_{2,n}) \leq 1.$$

This coefficient is called the absolute regularity, or β -mixing coefficient, it may be rewritten as

$$\beta(n, k) = \sup_{A_i \in \mathcal{A}_{1,n}, B_i \in \mathcal{A}_{2,n}} \left\{ \frac{1}{2} \sum_{i=1}^I \sum_{j=1}^J |\mathbb{P}(A_i \cap B_j) - \mathbb{P}(A_i)\mathbb{P}(B_j)| \right\}.$$

For more details on this question, the reader may have a quick look at [Doukhan \(1994\)](#), [Rio \(2017\)](#) and [Bradley \(2005\)](#).

In the next section, we give our results for weakly dependent sequences, under assumptions of stationary α -mixing and β -mixing sequences for real-valued empirical functions.

4. RESULTS AND DISCUSSIONS

We begin with the main result.

4.1. GC-classes for real-valued empirical functions. We focus on \mathcal{F}_c and define the empirical function

$$F_n(x) = \frac{1}{n} \sum_{i=1}^n \mathbb{I}_{]-\infty, x]}(X_i), \quad x \in \mathbb{R},$$

and its expectation

$$F(x) = \mathbb{E}(\mathbb{I}_{]-\infty, x]}(X_i)), \quad x \in \mathbb{R}.$$

Now, by applying the Part (a) of Theorem (3, page 6) for $C_x =]-\infty, x]$, $x \in \mathbb{R}$, we get

Theorem 5. *Let X, X_1, X_2, \dots be an arbitrary stationary and square integrable sequence of rrv's, defined on a probability space $(\Omega, \mathcal{A}, \mathbb{P})$. We have*

$$\sup_{x \in \mathbb{R}} |F_n(x) - F(x)| \rightarrow 0, \quad \text{almost surely, as } n \rightarrow +\infty$$

whenever the following general conditions hold : for some $\delta \in]0, 3[$

$$(4.1) \quad \sup_{q \geq 1} \frac{1}{q^{(3-\delta)/2}} \left[\sum_{i=1}^q \text{Var}(\mathbb{I}_{\{X_i \leq x\}}) + \sum_{(i \neq j)=1}^q \text{Cov}(\mathbb{I}_{\{X_i \leq x\}}, \mathbb{I}_{\{X_j \leq x\}}) \right] < +\infty$$

and

$$(4.2) \quad \sup_{q \geq 1} \frac{1}{q^{(3-\delta)}} \left[\sum_{i=q^2+1}^{(q+1)^2} \text{Var}(\mathbb{I}_{\{X_i \leq x\}}) + \sum_{(i \neq j)=q^2+1}^{(q+1)^2} \text{Cov}(\mathbb{I}_{\{X_i \leq x\}}, \mathbb{I}_{\{X_j \leq x\}}) \right] < +\infty.$$

Remark 3. *Since in the expressions in (4.1) and (4.2), the covariances are zero whenever the variables are independent and identically distributed (iid), the iid case remains valid without any further condition, which is the classical Glivenko-Cantelli theorem.*

Proof. We apply the Part (a) of Theorem (3, page 6) for $C_x =]-\infty, x]$, $x \in \mathbb{R}$ and verify the conditions (3.1) and (3.2) in Theorem (1, page 5). The conditions become

$$\begin{aligned}
\sup_{q \geq 1} \mathbb{V}ar \left(\frac{1}{q^{(3-\delta)/4}} \sum_{i=1}^q \mathbb{I}_{\{X_i \leq x\}} \right) &= \sup_{q \geq 1} \frac{1}{q^{(3-\delta)/2}} \mathbb{V}ar \left(\sum_{i=1}^q \mathbb{I}_{\{X_i \leq x\}} \right) \\
&= \sup_{q \geq 1} \frac{1}{q^{(3-\delta)/2}} \sum_{i,j=1}^q \mathbb{C}ov \left(\mathbb{I}_{\{X_i \leq x\}}, \mathbb{I}_{\{X_j \leq x\}} \right) \\
&= \sup_{q \geq 1} \frac{1}{q^{(3-\delta)/2}} \left[\sum_{i=1}^q \mathbb{V}ar \left(\mathbb{I}_{\{X_i \leq x\}} \right) + \sum_{(i \neq j)=1}^q \mathbb{C}ov \left(\mathbb{I}_{\{X_i \leq x\}}, \mathbb{I}_{\{X_j \leq x\}} \right) \right] \\
&< +\infty
\end{aligned}$$

and

$$\begin{aligned}
&\sup_{k \geq 1} \sup_{q: q^2+1 \leq k \leq (q+1)^2} \sup_{k \leq j \leq (q+1)^2} \mathbb{V}ar \left(\frac{1}{q^{(3-\delta)/2}} \sum_{i=1}^{j-q^2+1} \mathbb{I}_{\{X_{q^2+i} \leq x\}} \right) \\
&= \sup_{k \geq 1} \sup_{q: q^2+1 \leq k \leq (q+1)^2} \sup_{k \leq j \leq (q+1)^2} \frac{1}{q^{(3-\delta)}} \mathbb{V}ar \left(\sum_{i=1}^{j-q^2+1} \mathbb{I}_{\{X_{q^2+i} \leq x\}} \right) \\
&= \sup_{q \geq 1} \frac{1}{q^{(3-\delta)}} \mathbb{V}ar \left(\sum_{i=q^2+1}^{(q+1)^2} \mathbb{I}_{\{X_i \leq x\}} \right) \\
&= \sup_{q \geq 1} \frac{1}{q^{(3-\delta)}} \sum_{i,j=q^2+1}^{(q+1)^2} \mathbb{C}ov \left(\mathbb{I}_{\{X_i \leq x\}}, \mathbb{I}_{\{X_j \leq x\}} \right) \\
&= \sup_{q \geq 1} \frac{1}{q^{(3-\delta)}} \left[\sum_{i=q^2+1}^{(q+1)^2} \mathbb{V}ar \left(\mathbb{I}_{\{X_i \leq x\}} \right) + \sum_{(i \neq j)=q^2+1}^{(q+1)^2} \mathbb{C}ov \left(\mathbb{I}_{\{X_i \leq x\}}, \mathbb{I}_{\{X_j \leq x\}} \right) \right] \\
&< +\infty.
\end{aligned}$$

□

Remark 4. Note that for sequences of stationary random variables, the condition (4.1) implies that of (4.2).

In the sequel, we use the Theorem (5 page 10) to give applications for stationary α -mixing and β -mixing sequences.

4.2. α -mixing case. We already made a brief recall on α -mixing in Subsection 3.3 in Section 3. We are now going to provide applications to it.

Proposition 1. Suppose that X'_j s form a α -mixing stationary and square integrable sequence of random variables with mixing condition α . Then

$$\sup_{x \in \mathbb{R}} |F_n(x) - F(x)| \rightarrow 0, \text{ almost surely, as } n \rightarrow +\infty$$

holds whenever we have, for some $\delta \in]0; 1[$,

$$\alpha(n) = O\left(n^{-\left(\frac{1+\delta}{1-\delta}\right)}\right).$$

Proof. We need to verify the condition (4.1) in Theorem (5, page 10)

$$\begin{aligned} \sup_{q \geq 1} \frac{1}{q^{(3-\delta)/2}} \mathbb{V}ar \left(\sum_{i=1}^q \mathbb{I}_{\{X_i \leq x\}} \right) &= \sup_{q \geq 1} \frac{1}{q^{(3-\delta)/2}} \left[\sum_{i=1}^q \mathbb{V}ar \left(\mathbb{I}_{\{X_i \leq x\}} \right) + \sum_{(i \neq j)=1}^q \mathbb{C}ov \left(\mathbb{I}_{\{X_i \leq x\}}, \mathbb{I}_{\{X_j \leq x\}} \right) \right] \\ &= \sup_{q \geq 1} \frac{1}{q^{(3-\delta)/2}} \left[\sum_{i=1}^q \mathbb{V}ar \left(\mathbb{I}_{\{X_i \leq x\}} \right) + 2 \sum_{1 \leq i < j \leq q} \mathbb{C}ov \left(\mathbb{I}_{\{X_i \leq x\}}, \mathbb{I}_{\{X_j \leq x\}} \right) \right] \\ &\leq \frac{q}{q^{(3-\delta)/2}} + \frac{16}{q^{(3-\delta)/2}} \sum_{1 \leq i \leq q-1, j=i+h, h \leq n} \alpha(|i-j|) \\ &\quad + \frac{16}{q^{(3-\delta)/2}} \sum_{1 \leq i \leq q-1, j=i+h, h > n} \alpha(|i-j|) \text{ by the Theorem 4} \\ &\leq \frac{q}{q^{(3-\delta)/2}} + \frac{8nq}{q^{(3-\delta)/2}} \times \frac{1}{4} + \frac{8q^2 - 8nq - q}{q^{(3-\delta)/2}} \alpha(n) \\ &\sim \frac{2nq}{q^{(3-\delta)/2}} + \frac{8q^2}{q^{(3-\delta)/2}} \alpha(n) < +\infty. \end{aligned}$$

Finally, the condition reduces to

$$n = \left[q^{(1-\delta)/2} \right] \text{ and } \alpha(n) = O\left(q^{\frac{1+\delta}{2}}\right) = O\left(n^{-\left(\frac{1+\delta}{1-\delta}\right)}\right).$$

□

Remark 5. In the case of α -mixing sequences, we do not need whole function α . Instead we may fix $x \in \mathbb{R}$ and consider the modulus $\alpha(x, n)$ related to the sequence $\mathbb{I}_{]-\infty, x]}(X_i)$'s. It is clear that

$$\forall n \geq 1, \forall x \in \mathbb{R}, \alpha(x, n) \leq \alpha(n).$$

We have the Glivenko-Cantelli theorem if and only if

$$\forall x \in \mathbb{R}, \alpha(n) = O\left(n^{-\left(\frac{1+\delta}{1-\delta}\right)}\right).$$

The same can be done for any particular class \mathcal{C} by using the sequences $\mathbb{I}_{\mathcal{C}}(X_i)$'s and the $\alpha(\mathcal{C}, n)$ mixing modulus, and get the condition

$$\forall x \in \mathcal{C}, \alpha(\mathcal{C}, n) = O\left(n^{-\left(\frac{1+\delta}{1-\delta}\right)}\right).$$

4.3. β -mixing case. We consider here a stationary process $(X_t)_{t \in \mathbb{Z}}$ satisfying a β -mixing condition. The β -mixing coefficient is defined as

$$\beta(n) = \sup_{A \in \mathcal{F}_t^{-\infty}, B \in \mathcal{F}_{t+n}^{\infty}} |\mathbb{P}(A \cap B) - \mathbb{P}(A)\mathbb{P}(B)|,$$

where $\mathcal{F}_t^{-\infty} = \sigma(X_\ell : \ell \leq t)$ and $\mathcal{F}_{t+n}^{\infty} = \sigma(X_\ell : \ell \geq t+n)$.

We begin with giving this useful result which formulates the covariance inequality for stationary β -mixing sequences.

Lemma 2. *For any stationary beta-mixing process $(X_t)_{t \in \mathbb{Z}}$ and for all $t, s \in \mathbb{Z}$, we have*

$$|\mathbb{Cov}(X_t, X_s)| \leq 2\beta(|t-s|)\|X_t\|_{\infty}\|X_s\|_{\infty},$$

where $\|X_t\|_{\infty} = \sup_{\omega \in \Omega} |X_t(\omega)|$ is the supremum norm of X_t .

Proof. The covariance of X_t and X_s is given by

$$\mathbb{Cov}(X_t, X_s) = \mathbb{E}[X_t X_s] - \mathbb{E}[X_t]\mathbb{E}[X_s].$$

We focus on the first term, $\mathbb{E}[X_t X_s]$. By introducing the σ -algebra $\mathcal{F}_t = \sigma(X_\ell : \ell \leq t)$, we can write

$$\mathbb{E}[X_t X_s] = \mathbb{E}[X_t \mathbb{E}[X_s | \mathcal{F}_t]].$$

Thus, the covariance becomes

$$\mathbb{Cov}(X_t, X_s) = \mathbb{E}[X_t(\mathbb{E}[X_s | \mathcal{F}_t] - \mathbb{E}[X_s])].$$

The definition of the coefficient $\beta(n)$ implies that for any random variable $X_s \in \mathcal{F}_{t+n}^{\infty}$, we have the bound

$$|\mathbb{E}[X_s | \mathcal{F}_t] - \mathbb{E}[X_s]| \leq 2\beta(n)\|X_s\|_{\infty},$$

where $n = |t-s|$.

By substituting into the previous expression, we obtain

$$|\mathbb{Cov}(X_t, X_s)| \leq \|X_t\|_{\infty} \cdot \mathbb{E}[|\mathbb{E}[X_s | \mathcal{F}_t] - \mathbb{E}[X_s]|].$$

Since $|\mathbb{E}[X_s | \mathcal{F}_t] - \mathbb{E}[X_s]| \leq 2\beta(|t-s|)\|X_s\|_{\infty}$, it follows that

$$|\mathbb{Cov}(X_t, X_s)| \leq 2\beta(|t-s|)\|X_t\|_{\infty}\|X_s\|_{\infty}.$$

Thus, the proof of Theorem 2 is complete. \square

The next result states the uniform convergence for empirical functions of β -mixing stationary sequences.

Proposition 2. *Suppose that X_j 's form a β -mixing stationary and square integrable sequence of random variables with mixing condition β . Then*

$$\sup_{x \in \mathbb{R}} |F_n(x) - F(x)| \rightarrow 0, \text{ almost surely, as } n \rightarrow +\infty$$

holds whenever we have, for some $\delta \in]0; 1[$,

$$\beta(n) = O\left(n^{-\left(\frac{1+\delta}{1-\delta}\right)}\right).$$

Proof. We need to verify the condition (4.1) in Theorem (5, page 10)

$$\begin{aligned} \sup_{q \geq 1} \frac{1}{q^{(3-\delta)/2}} \mathbb{V}ar \left(\sum_{i=1}^q \mathbb{I}_{\{X_i \leq x\}} \right) &= \sup_{q \geq 1} \frac{1}{q^{(3-\delta)/2}} \left[\sum_{i=1}^q \mathbb{V}ar \left(\mathbb{I}_{\{X_i \leq x\}} \right) + \sum_{(i \neq j)=1}^q \text{Cov} \left(\mathbb{I}_{\{X_i \leq x\}}, \mathbb{I}_{\{X_j \leq x\}} \right) \right] \\ &= \sup_{q \geq 1} \frac{1}{q^{(3-\delta)/2}} \left[\sum_{i=1}^q \mathbb{V}ar \left(\mathbb{I}_{\{X_i \leq x\}} \right) + 2 \sum_{1 \leq i < j \leq q} \text{Cov} \left(\mathbb{I}_{\{X_i \leq x\}}, \mathbb{I}_{\{X_j \leq x\}} \right) \right] \\ &\leq \frac{q}{q^{(3-\delta)/2}} + \frac{4}{q^{(3-\delta)/2}} \sum_{1 \leq i \leq q-1, j=i+h, h \leq n} \beta(|i-j|) \\ &\quad + \frac{4}{q^{(3-\delta)/2}} \sum_{1 \leq i \leq q-1, j=i+h, h > n} \beta(|i-j|) \text{ by the Lemma 2} \\ &\leq \frac{q}{q^{(3-\delta)/2}} + \frac{2nq}{q^{(3-\delta)/2}} + \frac{2q^2 - 2nq - q}{q^{(3-\delta)/2}} \beta(n) \\ &\sim \frac{2nq}{q^{(3-\delta)/2}} + \frac{2q^2}{q^{(3-\delta)/2}} \beta(n) < +\infty. \end{aligned}$$

Finally, the condition reduces to

$$n = \left[q^{(1-\delta)/2} \right] \text{ and } \beta(n) = O\left(q^{\frac{1+\delta}{2}}\right) = O\left(n^{-\left(\frac{1+\delta}{1-\delta}\right)}\right).$$

□

Remark 6. *In the case of β -mixing sequences too, we do not need whole function β . Instead we may fix $x \in \mathbb{R}$ and consider the modulus $\beta(x, n)$ related to the sequence $\mathbb{I}_{]-\infty, x]}(X_i)$'s. It is clear that*

$$\forall n \geq 1, \forall x \in \mathbb{R}, \beta(x, n) \leq \beta(n).$$

We have the Glivenko-Cantelli theorem if and only if

$$\forall x \in \mathbb{R}, \beta(n) = O\left(n^{-\left(\frac{1+\delta}{1-\delta}\right)}\right).$$

The same can be done for any particular class \mathcal{C} by using the sequences $\mathbb{I}_{\mathcal{C}}(X_i)$'s and the $\beta(\mathcal{C}, n)$ mixing modulus, and get the condition

$$\forall x \in \mathcal{C}, \beta(\mathcal{C}, n) = O\left(n^{-\left(\frac{1+\delta}{1-\delta}\right)}\right).$$

5. CONCLUSION

We have extended the classical Glivenko-Cantelli theorem to real-valued empirical function classes under α -mixing and β -mixing conditions. Our results provide sufficient conditions for uniform convergence and establish deviation bounds that account for dependence structures. Future research may explore tighter bounds and applications to high-dimensional settings.

6. APPENDIX

Proof of the Theorem (2 page 5). Consider a class \mathcal{F} of measurable real functions such that each $f \in \mathcal{F}$. Let $p = N_{[]}(\mathcal{F}, L_1, \epsilon)$ be finite, for every $\epsilon > 0$. Then there is a sequence of intervals $[f_i, g_i]$, $i = 1, \dots, p$ such that for every $f_i, g_i \in \mathcal{F}$, $1 \leq i \leq p$, and $\mathcal{F} \subseteq \bigcup [f_i, g_i]$, and such that $\mathbb{E}\mathbb{P}_n(g_i) - \mathbb{E}\mathbb{P}_n(f_i) \leq \epsilon$. For any $f \in \mathcal{F}$, there exist f_i and g_i such as $f_i \leq f \leq g_i$ and

$$(6.1) \quad \mathbb{P}_n(f) - \mathbb{E}\mathbb{P}_n(f) = \mathbb{P}_n(f) - \mathbb{E}\mathbb{P}_n(g_i) + \mathbb{E}\mathbb{P}_n(g_i) - \mathbb{E}\mathbb{P}_n(f).$$

On the other hand we have, by the monotonicity of the probability, that

$$\mathbb{P}_n(f_i) \leq \mathbb{P}_n(f) \leq \mathbb{P}_n(g_i).$$

Therefore

$$\begin{aligned} \mathbb{P}_n(f) - \mathbb{E}\mathbb{P}_n(f) &\leq \mathbb{P}_n(g_i) - \mathbb{E}\mathbb{P}_n(g_i) + \mathbb{E}\mathbb{P}_n(g_i) - \mathbb{E}\mathbb{P}_n(f) \\ &\leq (\mathbb{P}_n - \mathbb{E}\mathbb{P}_n)(g_i) + \epsilon. \end{aligned}$$

Next

$$\begin{aligned} \mathbb{E}\mathbb{P}_n(f) - \mathbb{P}_n(f) &\leq \mathbb{E}\mathbb{P}_n(f) - \mathbb{E}\mathbb{P}_n(f_i) + \mathbb{E}\mathbb{P}_n(f_i) - \mathbb{P}_n(f) \\ &\leq \epsilon + (\mathbb{E}\mathbb{P}_n - \mathbb{P}_n)(f_i). \end{aligned}$$

Thus

$$(6.2) \quad \begin{aligned} \|\mathbb{E}\mathbb{P}_n - \mathbb{P}_n\|_{\mathcal{F}} &= \sup_{f \in \mathcal{F}} |\mathbb{E}\mathbb{P}_n - \mathbb{P}_n|(f) \\ &= \sup_{f \in \mathcal{F}} \max((\mathbb{E}\mathbb{P}_n - \mathbb{P}_n)(f), (\mathbb{P}_n - \mathbb{E}\mathbb{P}_n)(f)) \\ &\leq \epsilon + \max_{1 \leq i \leq p} \max((\mathbb{E}\mathbb{P}_n - \mathbb{P}_n)(f_i), (\mathbb{P}_n - \mathbb{E}\mathbb{P}_n)(g_i)). \end{aligned}$$

Since for every i , $1 \leq i \leq p$, we have $\mathbb{P}_n(f_i) \rightarrow \mathbb{E}\mathbb{P}_n(f_i)$ by the Theorem (1, page 5), it follows that

$$\max_{1 \leq i \leq p} \max((\mathbb{E}\mathbb{P}_n - \mathbb{P}_n)(f_i), (\mathbb{P}_n - \mathbb{E}\mathbb{P}_n)(g_i)) \rightarrow 0.$$

By applying the upper limit on the left-hand side in (6.2), we get for every $\epsilon > 0$

$$\limsup_{n \rightarrow \infty} \|\mathbb{E}\mathbb{P}_n - \mathbb{P}_n\|_{\mathcal{F}} \leq \epsilon \text{ almost surely.}$$

□

REFERENCES

- Billingsley, P. (1968). *Convergence of probability measures*. Wiley, New York.
- Bradley, R. C. (2005). Basic Properties of Strong Mixing Conditions. A Survey and Some Open Questions. *Probability Surveys*. Vol. 2, 107–144. ISSN: 1549-5787, DOI: 10.1214/154957805100000104.
- Doukhan, P. (1994). *Mixing properties and exemples*. ISBN 0-387-94214-9. © Springer-Verlag New York, Inc.
- Doukhan, P., Massart, P. and Rio, E. (1995). Invariance principles for the empirical measure of a weakly dependent process. *Ann. Inst. H. Poincaré*. 31, 393-427.
- Louhichi, S. (2000). Weak convergence for empirical processes for associated sequences, *Ann. Inst. H. Poincaré, Proba. Stat.* 36, 5 547-567.
- Rio, E. (2017). *Asymptotic Theory of Weakly Dependent Random Processes*. Probability Theory and Stochastic Modelling 80. DOI 10.1007/978-3-662-54323-8, ©Springer-Verlag GmbH Germany.
- Sangaré, H. and Lo, G. S. (2015). A general strong law of large numbers and applications to associated sequences and to extreme value theory. *Annales Mathematicae et Informaticae*. 45, pp. 111-132.
- Sangaré, H., Lo, G. S. and Traoré, M. C. M. (2020). Arbitrary Functional Glivenko-Cantelli Classes and Applications to Different Types of Dependence. *Far East Journal of Theoretical Statistics*. 60, pp. 41-62. <http://dx.doi.org/10.17654/TS060020041>.
- Shao, Q. M. (1995). Weak convergence of multidimensional empirical processes for strong mixing sequences. *Chinese Ann. Math. Ser. A* 7, 547-552.
- Shao, Q. M. and Yu, H. (1996). Weak convergence for weighted empirical processes of dependent sequences. *The Annals of Probability*. Vol. 24, Number 4, 2097-2127.
- Van Der Vaart, A. W. and Wellner, J. A. (1996). *Weak convergence and empirical processes with application in statistics*. ISBN 0-387-94640-3, Springer-Verlag New York Berlin Heidelberg SPIN 10522999.
- Yoshihara, K. (1975). Billingsley's theorems on empirical processes of strong mixing sequences, *Yokohama Math. J.* 23, 1-7.