

Simple molecules and complex chemistry in a protoplanetary disk

A JWST investigation of the highly inclined disk d216-0939

A. Potapov^{1,*}, H. Linz^{1,2}, J. Bouwman², W. Rocha³, J. Martin⁴, S. Wolf⁴, Th. Henning², and H. Terada⁵

¹ Analytical Mineralogy Group, Institute of Geosciences, Friedrich Schiller University Jena, Jena, Germany
e-mail: alexey.potapov@uni-jena.de

² Max Planck Institute for Astronomy, Heidelberg, Germany
e-mail: [linz, bouwman]@mpia.de

³ Laboratory for Astrophysics, Leiden Observatory, Leiden University, Leiden, The Netherlands

⁴ Institute for Theoretical Physics and Astrophysics, Christian Albrecht University Kiel, Kiel, Germany

⁵ National Astronomical Observatory of Japan, Tokyo, Japan

Received December 10, 2024; accepted February 20, 2025

ABSTRACT

Context. While the number of detected molecules, particularly complex organic molecules, in the solid-state in astrophysical environments is still rather limited, laboratory experiments and astrochemical models predict many potential candidates. Detection of molecules in protoplanetary disks provides a bridge between the chemical evolution of the interstellar medium and the chemistry of planets and their atmospheres.

Aims. The excellent spectral sensitivity, broad wavelength coverage and high spatial resolution of the James Webb Space Telescope (JWST) allows for making progress in exploring chemical compositions of various astrophysical environments including planet-forming disks. They are a prerequisite for probing the disk content by means of sensitive absorption studies.

Methods. In this paper, we present initial results of the JWST Cycle 1 GO program 1741 on d216-0939, a highly inclined T Tauri disk located in the outskirts of the Orion Nebula Cluster. We utilise the NIRSpec and MIRI integral field unit spectrographs to cover its spectrum from 1.7 to 28 μm .

Results. In the d216-0939 disk, we give assignments of the composition of silicate grains. We unambiguously detect solid-state features of H_2O , CO_2 , $^{13}\text{CO}_2$, CO , OCN^- , and tentatively OCS ; species that had been detected recently also in other circumstellar disks. For the first time in disks, we provide unique detections of ices carrying NH_4^+ and the complex organic molecule ammonium carbamate ($\text{NH}_4^+\text{NH}_2\text{COO}^-$).

Conclusions. The latter detections speak for a very efficient NH_3 chemistry in the disk. We also show the very important role of scattering in the analysis of observational spectra of highly inclined disks.

Key words. Protoplanetary Disks – Solid state: volatile – Solid state: refractory – Telescopes: JWST

1. Introduction

In cold dense astrophysical environments, such as molecular clouds, protostellar envelopes and planet-forming disks beyond the snowline, the solid state is presented by dust grains mixed with molecular ices. The full composition of interstellar and circumstellar ices is still an open question. Water is the main constituent accounting for more than 60% of the ice in most lines of sight (e.g., Whittet 2003; van Dishoeck et al. 2021). In addition to water, several other molecules have been unambiguously detected in ices: CO , OCS , CO_2 , NH_3 , H_2CO , HCOOH , CH_4 , and CH_3OH (McGuire 2022) and recently OCN^- , NH_4^+ , and tentatively SO_2 (McClure et al. 2023). Basing on results of experimental and modelling studies simulating conditions and chemistry of astrophysical environments, we know that chemical reactions in/on such ices triggered by UV photons, heat, atoms, and cosmic rays may lead to the formation of many complex organic molecules (COMs) including direct precursors of prebiotic species and prebiotic species themselves, such as sugars, amino acids, and nucleobases. Here, we refer the reader to several review papers (Theulé et al. 2013; Linnartz et al. 2015;

Öberg 2016; Arumainayagam et al. 2019; Sandford et al. 2020; Potapov & McCoustra 2021; Fulvio et al. 2021).

However, detection of COMs in interstellar and circumstellar ices (in the solid state) is still a challenge because these molecules are less abundant, and their IR features are less distinct. In interstellar ice, methanol (CH_3OH), has been unambiguously detected (Pontoppidan et al. 2003). In the pre-JWST era, tentative detections of acetaldehyde (CH_3CHO Schutte et al. 1999), ethanol ($\text{C}_2\text{H}_5\text{OH}$ Terwisscha van Scheltinga et al. 2018), and urea (H_2NCONH_2 Raunier et al. 2004) have been proposed. New JWST data provide first tentative (e.g., Yang et al. 2022) and then statistically robust (Rocha et al. 2024) detections of $\text{C}_2\text{H}_5\text{OH}$, CH_3CHO and methyl formate (HCOOCH_3), as well as tentatively acetonitrile (CH_3CN , McClure et al. 2023) and ethyl cyanide ($\text{C}_2\text{H}_5\text{CN}$, Nazari et al. 2024) in interstellar ices.

Detection of molecules in planet-forming disks provides a bridge between the chemical evolution of the interstellar medium (ISM) and chemistries of planets and their atmospheres. Recent JWST/NIRSpec observations of the edge-on Class II protoplanetary disk HH 48 NE revealed spatially resolved absorption features of the major ice components H_2O , CO_2 , and CO , multiple weaker signatures from less abundant ices NH_3 , OCN^- , and

* Corresponding author

OCS and, for the first time in a protoplanetary disk, isotopologue $^{13}\text{CO}_2$ ice (Sturm et al. 2023a). JWST/MIRI observations of the same disk provided evidence for the presence of CH_4 and CH_3OH (Sturm et al. 2023b).

Despite these recent successes regarding smaller species, the general number of detected ice species in disks, particularly COMs, is still very limited. However, according to the laboratory experiments and astrochemical models there are many potential candidates. This manuscript presents results of the initial analysis of JWST observations of the protoplanetary disk d216-0939, a highly inclined disk in the outskirts of the Orion Nebula Cluster, and provides assignments of the ice and dust bands detected.

2. Observations, data reduction and analysis

2.1. Source selection

Ice bands have been observed in emission in planet-forming disks at far infrared wavelengths (e.g., Min et al. 2016). However, to observe the near- and mid-infrared ice bands in emission, one would typically need temperatures of several hundred Kelvin, which is far above the sublimation temperature. The best way to observe ice bands at near-infrared wavelengths is through absorption line studies of highly inclined disks (e.g., Terada & Tokunaga 2012). At high inclination angles the light of the central star is attenuated by the circumstellar disks allowing us to probe a wide region of the disk.

For our study, we selected the highly inclined disk d216-0939 (a.k.a. V* V2377 Ori or HOPS 65) in Orion, a high-mass star forming region likely to resemble the birthplace of the Solar System (cf. Young 2020). In the Hubble Space Telescope (HST) paper by Smith et al. (2005) this disk system stood out as featuring a large silhouette disk in HST optical imaging. A high-resolution ALMA study has derived an inclination of the outer disk of around $75\text{--}78^\circ$ (Sheehan et al. 2020). The selected target is among the few known suitably inclined disks that show indications of crystalline water ice (Terada & Tokunaga 2012; Terada et al. 2012) and strong reddening of the central star by the circumstellar disk material. As crystalline water ice forms at temperatures reached close to the snowline, this selection criterion means that we are likely probing a sightline in this system across the snowline. However, other scenarios may exist. Crystallization may also occur earlier in the history of the system when the outer disk was warmer, or material may have been transported outwards after being crystallized near the snowline. We note furthermore that the locations of snowlines are not constant over the evolution of the disk due to a variety of circumstances (e.g., Martin & Livio 2012; Owen 2020; Mori et al. 2021; Ros & Johansen 2024).

2.2. JWST observations

The highly inclined protoplanetary disk in Orion d216-0939 was observed with the integral field units (IFUs) of both the Near-Infrared Spectrograph (NIRSpec, Jakobsen et al. 2022; Böker et al. 2022) and the Mid-Infrared Instrument (MIRI, Wright et al. 2023) within our General Observation (GO) programme 1741 (PI: A. Potapov) at the end of the JWST Cycle 1. The NIRSpec observations used the medium resolution settings G235M/F170LP and G395M/F290LP to cover the 2 to 5-micron wavelength range, and especially to seamlessly cover the full extent of the water ice feature between 2.7 and 3.5 micron. As the target is expected to be slightly extended at the shortest wavelengths of NIRSpec we used a 4-point dither. Given the IFU

size we apply a target acquisition (TA) for the NIRSpec observations on 2 nearby stars to make sure our targets are properly centred on the image slicer. We used nearby stars with good-quality GAIA information for the TA as the target itself could show some deviation from point-source geometry at short wavelengths. As the IFU size of MIRI is large enough, no TA was needed. Observations with the MIRI Medium Resolution Spectrograph (MRS) were done for all 3 grating settings so the entire wavelength range between 5 and 28 microns was continuously covered. In the MIRI range we were especially interested in the silicate absorption features at 10 and 18 microns, as well as several faint absorption signals of complex organic molecule ices, mainly in the 5 - 10-micron range. For the MIRI observations we applied a 4-point dither, to get a proper spatial and wavelength sampling. At the longer wavelengths the source is expected to be compact, meaning the background can be estimated from the observations itself and no additional off-pointing was required.

We opted for medium resolution spectroscopy for both instruments, since the offered spectral resolution in the low-res mode ($R\sim 100$) is too coarse to allow a robust spectral decomposition of the H_2O ice features at the shortest wavelengths, and, in case of MIRI do not cover all required wavelengths (> 10 microns). This also facilitated a proper detection and separation of gas-phase lines (e.g., from water) from the ice features. We employed the IFUs of both instruments which provides a better control over the spatial distribution of the (continuum) emission/absorption and ice feature absorption signals.

2.3. Data reduction

All data reduction was performed with the JWST pipeline (see Bushouse et al. 2024) version 1.13.4, and was based on the pmap1210 calibration scheme. We processed the raw data with slightly modified pipeline scripts. Spectral de-fringing was applied to all data sets. We relied on the extract1D step to extract the 1D spectrum from the 2D spectral cubes. This extracts aperture photometry per cube slice. An aperture correction is applied afterwards. Both the aperture and the aperture correction are wavelength-dependent. For MIRI we experimented with aperture scaling between $1.5 \dots 2.0 \times \text{FWHM}$ and found that the smaller aperture does not change the continuum shape, but gives slightly higher signal-to-noise. For NIRSpec, the aperture size scaling within extract1D is fixed (to $2 \times \text{FWHM}$). For most of the data, we could use the autocenter function to enable a precise centroiding of the aperture on source before spectral extraction. We noticed that at the shortest wavelengths covered ($\lambda \lesssim 2.5\mu\text{m}$) the source more and more deviates from a point-source geometry, probably showing scattering at the more extended disk surface. We checked that the deviation from the point-source nature is negligible for the longer wavelengths where the important spectral features reside. Hence, an extraction approach relying on point source aperture corrections seems still warranted.

2.4. Approach for fitting the measurements

We do not have a well-constrained photospheric model for the central star, yet. Moreover, the IR continuum may be dominated by thermal emission from warm dust in the inner disk rather than from the stellar photosphere. At this point in the project we refrained from trying to develop such a detailed radiative transfer model for the star-disk system that could guide the determination of the continuum baseline. As a pragmatic approach, we therefore used a third-order polynomial fit for the




Fig. 1. NIRSpect (left) + MIRI (right) spectrum of the d216-0939 disk and assignments of the main spectral bands. Gas phase lines are not labelled.

NIRSpect range with continuum points of 2.5 - 2.55, 2.6 - 2.65, 2.69 - 2.80, 4.0 - 4.04, 4.06 - 4.07, 4.6 - 5.0, 5.15 - 5.2 μm and two linear fits for the MIRI ranges 5.72 - 7.88 and 8 - 15.9 μm . The uncertainties introduced by the continuum do not influence the band assignments presented in the paper.

We used the ENIGMA fitting tool (Rocha et al. 2021) to fit multiple features across the NIRSpect and MIRI spectral ranges by scaling laboratory spectra in optical depth scale of dust/ice and ice mixtures and pure ices to match the continuum-subtracted optical depths. This methodology is already described in different papers, such as Rocha et al. (2021, 2024, 2025). Briefly, these optical depth spectra (τ) are calculated as:

$$\tau_{\nu}^{\text{lab}} = 2.3 \text{Abs}_{\nu}, \quad (1)$$

where Abs_{ν} is the laboratory data (absorbance) as function of the wavenumber (ν) in units of cm^{-1} .

Some of the ices also play a role in the scattering, which is needed to correctly reproduce the observed spectral features. We applied the computational code OpTool (Dominik et al. 2021) to investigate the scattering and scattering opacities, we use the tool strictly for deriving the scattering opacities of different silicate grains covered with a variety of ices and ice mixtures. For the absorption opacities, we rely on the laboratory-measured ice data directly.

The relation between optical depth and opacity is given by:

$$\tau_{\nu} = a \kappa_{\nu}, \quad (2)$$

where a is a scaling factor.

In OpTool, we utilized the default approach implemented therein, i.e., Distribution of Hollow Spheres (DHS) with $f_{\text{max}} = 0.8$, with compact grains (porosity set to zero) and ice mantles of different compositions. We adopted an upper size limit of 1 μm and a power law of 3.5 for the grain size distribution. This seems to be a realistic approach; however, we note that Dartois et al. (2024) found that the grain sizes can significantly alter the shapes of the ice features. We experimented

with the grain size distributions, increasing the upper size limit to 2 and 3 μm . This did not, however, lead to a fit improvement. Furthermore, it worsened the overall SED fit we attempted.

We used two metrics to evaluate the goodness of the fit, the Root Mean Square Error (RMSE) and the Akaike Information Criterion (AIC). While the RMSE measures the average difference between model and observations, the AIC can quantify the quality of the fit based on the number of parameters used for the fit. The AIC is given by:

$$\text{AIC} = \chi^2 + 2p + \frac{2p(p+1)}{N-p-1}, \quad (3)$$

where p is the number of free parameters and N is the sample size. χ^2 is given by:

$$\chi^2 = \frac{1}{\text{dof}} \sum_{i=0}^{n-1} \left(\frac{\tau_{\nu,i}^{\text{obs}} - \sum_{j=0}^{m-1} w_j \tau_{\nu,j}^{\text{lab}}}{\gamma_{\nu,i}^{\text{obs}}} \right)^2, \quad (4)$$

where dof is the number of degrees of freedom (i.e. $N - p$), m is the number of laboratory spectra used in the fit, w_j the scaling factor applied to the individual spectra, γ is the error in the observational optical depth spectrum. Once AIC is calculated, different models can be compared using the ΔAIC , which is defined as $\text{AIC}_i - \min(\text{AIC})$, where AIC_i represents values for different models, and $\min(\text{AIC})$ corresponds to the minimum AIC value. ΔAIC below 2 means a substantial support of the model, while ΔAIC clearly above 2 indicates considerably less or no support of the model (Burnham & Anderson 2002). In this paper, the lowest RMSE and AIC is considered the best model, which is not necessarily the model containing more components.

3. Results

In Figure 1, we present the complete NIRSpect + MIRI spectrum up to 18 μm of the d216-0939 disk.

In the following, we split the spectrum into regions and discuss our assignments of the dust and ice bands done using the

Table 1. EMIIGMA fitting results for the 2.8 – 4.0 μm wavelength region.

Component and Fit	Combination #							
	1	2	3	4	5	6	7	8
MgSiO ₃ /H ₂ O 150 K (A)	x	x	x	x	x	x		
MgSiO ₃ /H ₂ O 150 K (S)	x	x	x	x	x	x	x	x
NH ₄ ⁺ /OCN ⁻ 80 K		x		x				
“Chemistry”			x	x	x	x	x	x
MgSiO ₃ /H ₂ O 10 K (A)					x		x	
MgSiO ₃ /H ₂ O 10 K (S)					x			
MgSiO ₃ /H ₂ O 100 K (A)						x		x
MgSiO ₃ /H ₂ O 100 K (S)						x		
AIC	19.1	9.8	7.4	9.5	11.3	11.4	7.4	7.2
RMSE	0.055	0.028	0.016	0.016	0.015	0.015	0.016	0.015

Notes. The Akaike Information Criterion (AIC) and Root-mean-square error (RMSE) for various input mixtures are listed. The best fit parameters are marked in boldface.

ENIIGMA fitting tool, from short to long wavelengths starting at the 3-micron region containing the stretching vibration band of H₂O and ending at the 8 – 16-micron region containing spectral signatures of silicates. We performed separate fits because determining a single global continuum for disks is rather challenging without an accurate radiative transfer model.

3.1. 2.8 – 4.0-micron region

The main spectral band in this region is the H₂O stretching mode. It may also contain signatures of NH₃, CH₃OH, NH₄⁺, and complex organic molecules. Particularly, ammonium salts including ammonium carbamate (NH₄⁺NH₂COO⁻) and ammonium formate (HCOO⁻NH₄⁺) may explain the red part of the band as was shown by the analysis of the Rosetta spectra of the comet 67P (Poch et al. 2020).

Based on our previous results demonstrating dust/ice mixing in cold astrophysical environments Potapov et al. (2021), we use spectral data obtained for physical mixtures of silicates and water ice. Crystalline water ice was previously detected in the d216-0939 disk (Terada & Tokunaga 2012; Potapov et al. 2021) and its presence was one of the main parameters while choosing the target for our JWST observations.

It was also clear from the analysis of the CO₂ stretching band at 4.27 μm that scattering plays a crucial role in the short-wavelength region (see Sect. 3.2). The scattering component of the H₂O band was obtained utilising OpTool, with the optical constants as inputs obtained for the MgSiO₃/H₂O mixtures with the mass ratio of 2.7 and presented in Potapov et al. (2018). The absorption spectra of these mixtures were also used as absorption components of the fits.

The best fit model is shown in Figure 2 and was obtained using the spectra of MgSiO₃/H₂O 150 K scattering (S) + MgSiO₃/H₂O 100 K absorption (A) + “chemistry”. “Chemistry” is the laboratory spectrum obtained after UV irradiation of a mixture of CO₂ + NH₃ at 75 K and its subsequent heating to 230 K (Potapov et al. 2022). The spectrum mainly contains spectral signatures of ammonium carbamate and carbamic acid (NH₂COOH). The high temperature (230 K) spectrum was chosen to reduce the contribution of NH₃, which has strong bands at 3 and 9 μm but is not unambiguously detected.




Fig. 2. Observational spectrum (black), the best fit model for the 2.8 – 4.0 μm region (red) and its components: MgSiO₃/H₂O 150 K (S) (blue), MgSiO₃/H₂O 100 K (A) (green), and “chemistry” (orange).

The Akaike Information Criterion (AIC) and Root-mean-square error (RMSE) were used to estimate the quality of the fits and are presented in Table 1. The criteria for the best fit were: (i) decrease of the RMSE and (ii) increase of the AIC by not more than 2 with respect to the lowest value. We started with the 150 K absorption (A) and scattering (S) spectra as ENIIGMA inputs. We do not see clear spectral signatures of NH₃ and CH₃OH in the 10-micron range, where the species have the strongest bands. NH₄⁺ is detected at 6.85 μm (see below), and addition of the NH₄⁺/OCN⁻ 80 K spectrum (originally presented in Novozamsky et al. 2001) from the LIDA database (Rocha et al. 2022) improved the fit considerably. However, addition of the “chemistry” spectrum to the fit provided a much better fit and was necessary to fit the red shoulder of the band. The peaks at 3.1 and 3.3 μm coincide with the peaks of several ammonium salts (ammonium formate, ammonium citrate, ammonium sulphate, ammonium carbamate), see Poch et al. (2020). In our case,




Fig. 3. Observations (black), the best fit model for the 4.0 – 4.5 μm region (red) and its components: blue – cold scattering on 13 K H_2O : CO_2 mantles, green – cold scattering on 13 K pure CO_2 mantles, orange – absorption in 35 K H_2O : CO_2 mantles.

ammonium carbamate was detected in the “chemistry” sample (Potapov et al. 2022). The 3.3 μm peak is clearly affected by PAH emission. Addition of the $\text{NH}_4^+/\text{OCN}^-$ spectrum to the fit after “chemistry” led to an increase of the AIC by more than 2 as a consequence of the ignorance of the $\text{NH}_4^+/\text{OCN}^-$ spectrum by the fit.

At the next step, we added $\text{MgSiO}_3/\text{H}_2\text{O}$ 10 and 100 K (A) and (S) inputs to consider the temperature gradient in the disk. Increase of the AIC in both cases indicated that not all spectra were considered by the fit. The ignored spectra were in both cases low-temperature scattering and high-temperature absorption. Finally, the best fit was obtained for the $\text{MgSiO}_3/\text{H}_2\text{O}$ 150 K (S) + $\text{MgSiO}_3/\text{H}_2\text{O}$ 100 K (A) + “chemistry” inputs. We note that the usage of $\text{MgSiO}_3/\text{H}_2\text{O}$ 150 K (A) or $\text{MgSiO}_3/\text{H}_2\text{O}$ 10 K (A) instead of $\text{MgSiO}_3/\text{H}_2\text{O}$ 100 K (A) also provides a reasonable fit. Thus, 150 K scattering provides a defining contribution to the spectrum.

3.2. 4.0 – 4.5-micron region

This spectral region contains the strong absorption feature of $^{12}\text{CO}_2$ ice at 4.27 μm and the weaker ice feature of the $^{13}\text{CO}_2$ variety at 4.39 μm . Note that we also detected the combination mode of $^{12}\text{CO}_2$ at 2.68 μm . Very interesting is the positive bump blueward of the $^{12}\text{CO}_2$ absorption feature at 4.27 μm . This bump is relatively broad, though it does not coincide in wavelength with known solid-state feature of ice, dust or PAHs. We assume that here we see the imprint of scattering on ice mantles covering dust grains. This effect has been described and modelled by Dartois et al. (2022).

The best fit model is shown in Figure 3 and was obtained using the spectra of scattering on 13 K H_2O : CO_2 mantles, cold scattering on 13 K pure CO_2 mantles, and absorption in 35 K H_2O : CO_2 mantles.

For modelling the main CO_2 feature, we started with pure absorption data. A detailed overview of the utilised absorbance and optical constants is given in Table 2. We first consider moderate-temperature ice mixtures where CO_2 is mixed with H_2O . Absorbance data exist for temperatures of 13 K (cold), 35 K (lukewarm), and 72 K (warm). It quickly turns out that such pure absorbance reference data cannot replicate the full shape of the

measured CO_2 feature well. They are still too narrow, they cannot capture the wide red wing, and obviously do not contain a positive (“negative absorption”) feature that could emulate the bump close to 4.2 μm . We therefore decided to include extinction caused by scattering also for the CO_2 fitting into the selection of input data for ENIIGMA. We used OpTool to compute the scattering properties with the model parameters as described in Section 2.4. To be consistent, we used optical constants for the grain core and mantle at identical or at least similar temperatures, respectively. For the grain core, we used the same $\text{MgSiO}_3/\text{H}_2\text{O}$ data as above. For the mantle material, we used the optical constants from several databases (cf. Table 2). We excluded the wavelength range containing the Brackett α line at around 4.05 μm and start at 4.065 μm .

We continued with two-component fits of absorbance + scattering at similar temperatures, respectively. The red absorption wing is much better fitted by the inclusion of scattering. Additionally, the scattering produces some positive feature blueward of the main absorption feature, although the exact shape does not agree. The fit continuously improves when going from warm absorption + warm scattering to cold absorption + cold scattering. Another clear step of improvement occurs when including scattering of a pure cold CO_2 mantle. Such data offer an additional positive component that helps to fit better the truly blue part of the positive bump. Furthermore, the faint $^{13}\text{CO}_2$ absorption feature¹ at 4.39 μm is better captured when the pure CO_2 data are included. (The CO_2 samples used for the laboratory reference data contain the ^{13}C and ^{12}C isotopologues in a ratio of 1:100.) It must be noted that the pure CO_2 component was also required to fit the 10-micron range (see Sect. 3.5).

When more than three components are combined, several acceptable solutions with roughly similar fit quality and RMSE are possible, however, AIC is increased by more than 2 in all the cases. One notes that in many cases, the warm and lukewarm scattering components are ignored by ENIIGMA. Inclusion of one variety of cold scattering is, however, mandatory to reach a good fit. Our modelling is not fully conclusive regarding the necessity to have (luke)warm absorption and (luke)warm scattering. If given a multitude of input choices, ENIIGMA will include a 72 K absorption component in the final fit, possibly since those warm spectra have less of a wiggle at around 4.3 μm (probably an ^{18}OCO feature) than cold absorption models from the used spectral libraries. The AIC and RMSE values are presented in Table 3.

3.3. 4.5 – 4.8-micron region

This spectral region contains the spectral signatures of OCN^- at 4.62 μm and CO at 4.67 μm . We fitted these two features together in ENIIGMA but removed the strong Pfund β line at 4.65 μm from the optical depth input data before fitting, since it turned out to be confusing for the fitting routine. The OCN^- feature is in the $\text{NH}_4^+/\text{OCN}^-$ spectrum from LIDA database. The best fit model is presented in Figure 4. The AIC and RMSE values are presented in Table 4.

There are data for $\text{NH}_4^+/\text{OCN}^-$ at 12 K and 80 K. We tested both data sets. The cold 12 K data come, however, with a quite pronounced secondary absorption feature at around 4.65 μm which is not observed in our data. The 80 K file does not show this and is smoother and better fitting to the JWST measure-

¹ We note in passing that also the $^{13}\text{CO}_2$ feature at around 4.39 μm seems to show a positive (scattering) bump blueward of its absorption maximum, similar to the one discussed for the $^{12}\text{CO}_2$ feature.

Table 2. Input data sets for fitting the CO₂ region around 4.27 μm.

Detailed Description	Rocha Model	Trivial name for Table 3
H ₂ O: CO ₂ :NH ₃ :CH ₄ 10:1:1:1 72 K absorbance	D13a	Warm absorption
H ₂ O: CO ₂ :NH ₃ :CH ₄ 10:1:1:1 35 K absorbance	D12a	Lukewarm absorption
H ₂ O: CO ₂ 10:1 13 K absorbance	D2a	Cold absorption
CO ₂ 13 K absorbance	G2	Cold absorption pure CO ₂
MgSiO ₃ /H ₂ O 100K core + D13a 72 K mantle scatter	D13a	Warm scattering
MgSiO ₃ /H ₂ O 10K core + D12a 35 K mantle scatter	D12a	Lukewarm scattering
MgSiO ₃ /H ₂ O 10K core + D2a 13 K mantle scatter	D2a	Cold scattering
MgSiO ₃ /H ₂ O 10K core + G2 13 K mantle scatter	G2	Cold scattering pure CO ₂

Notes. The paper references are Rocha & Pilling (2014) for model G2, and Rocha et al. (2017) for models D13a, D12a, and D2a.

Table 3. EMIIGMA fitting results for the 4.0 – 4.5 μm wavelength region.

Component and Fit	Combination #											
	1	2	3	4	5	6	7	8	9	10	11	12
Warm absorption	x				x		x		x	x		
Lukewarm absorption		x				x		x				
Cold absorption			x	x					x	x	x	x
Cold absorption pure CO ₂										x	x	(x)
Warm scattering	x						x		x	x		
Lukewarm scattering		x						x				
Cold scattering			x	x	x	x	x	x	x	x	x	x
Cold scattering pure CO ₂				x	x	x	x	x	x	x	x	x
AIC	8.09	9.60	6.55	7.05	7.10	7.02	9.02	9.13	11.06	14.22	9.08	9.17
RMSE	0.029	0.033	0.026	0.014	0.015	0.014	0.014	0.015	0.014	0.020	0.014	0.015

Notes. The Akaike Information Criterion (AIC) and Root-mean-square error (RMSE) for various input mixtures. The best fit parameters are marked in boldface.

Table 4. EMIIGMA fitting results for the 4.5 – 4.8 μm wavelength region.

Component and Fit	Combination #										
	1	2	3	4	5	6	7	8	9	10	
NH ₄ ⁺ /OCN ⁻ 12 K		x									
pure CO 15 K		x	x				x	x		(x)	
NH ₄ ⁺ /OCN ⁻ 80 K			x	x	x	x	x	x	x	x	
H ₂ O:CO ₂ :CO: NH ₃ :CH ₃ OH 15K			x			x	x				
C/H ₂ O				x	x	x					
Rocha D8a H ₂ O:NH ₃ :CO 13 K							x	x	x	x	
cold Rocha D8a-mantle scattering									x	x	
AIC		7.44	7.39	6.89	7.55	8.91	10.81	6.79	8.81	7.20	9.18
RMSE		0.027	0.026	0.024	0.027	0.024	0.024	0.024	0.024	0.0151	0.0148

Notes. The Akaike Information Criterion (AIC) and Root-mean-square error (RMSE) for various input mixtures are listed. The best fit parameters are marked in boldface.

ments. Also, a preference for the 80 K spectrum is a clear result of the 5.6 – 8 μm region fitting (see subsection 3.4).

Pure CO absorption data from LIDA give a rather sharp narrow absorption feature at around 4.67 μm, while our measured CO absorption profile is wider. We therefore also use CO in a H₂O mixture. As one example, we use the absorbance of a H₂O:CO₂:CO:NH₃:CH₃OH mixture at 15 K taken from LIDA. As a further choice, we utilize the model “D8a” lab data from

Rocha et al. (2017), which represents an H₂O:NH₃:CO mixture with 10:6:4 mass ratio, at 13 K. With such a dataset in the mix, the core of the absorption is filled out better. The general shape, especially the absorption wing at longer wavelengths, however, is not fully replicated. Hence, we included scattering on ice mantles also for the CO range. We used the aforementioned model D8a which has optical constants available to compute the scattering properties, taking the MgSiO₃/H₂O material properties




Fig. 4. Observations (black), the best fit model for the 4.5 – 4.8 μm region (red) and its components: dark blue – pure CO absorption (15 K), light blue – scattering on $\text{H}_2\text{O}:\text{CO}$ ice mantle (13 K), green – $\text{NH}_4^+/\text{OCN}^-$ 80 K, yellow – absorption on $\text{H}_2\text{O}:\text{CO}$ ice mantle (13 K).




Fig. 5. Observations (black), the best fit model for the 5.6 – 8 μm region (red) and its components: $\text{MgSiO}_3/\text{H}_2\text{O}$ 150 K (green) + $\text{NH}_4^+/\text{OCN}^-$ 80 K (blue) + “chemistry” (orange).

at 10 K for the grain core. The inclusion of scattering widens the profile and fills the red wing satisfactorily. We note that the pure CO absorption is quite strongly down-weighted in the four-component mix that gives a good fitting result. The inclusion of pure CO seems to improve the RMSE to a small degree. Still, we cannot exclude the possibility that the narrow pure CO absorption just contributes to better fitting some absorption in the deepest absorption trough of the global feature that may arise from superimposed roto-vibrational absorption lines of CO seen in the spectrum. We also tentatively detected OCS at 4.9 μm .

3.4. 5.6 – 8-micron region

This spectral region contains the H_2O bending vibration mode at 6.2 μm , the NH_4^+ band at 6.9 μm , and the OCN^- band at 7.6 μm . The best fit model is presented in Figure 5 and was obtained




Fig. 6. Intermediate model for the 5.6 – 8 μm region including $\text{MgSiO}_3/\text{H}_2\text{O}$ 150 K (green) and $\text{NH}_4^+/\text{OCN}^-$ 80 K (blue).

using the input mixture of the spectra of $\text{MgSiO}_3/\text{H}_2\text{O}$ (A) 150 K + $\text{NH}_4^+/\text{OCN}^-$ 80 K + “chemistry”.

We started with the $\text{MgSiO}_3/\text{H}_2\text{O}$ 100 and 150 K and $\text{NH}_4^+/\text{OCN}^-$ 12 and 80 K inputs. There is a red shift of the NH_4^+ band with temperature. Our fits clearly show that the 12 K band does not match the observational band, and that the 80 K spectrum must be used. This was the reason for trying the $\text{NH}_4^+/\text{OCN}^-$ 80 K spectrum also in the 3-micron range. The usage of the $\text{MgSiO}_3/\text{H}_2\text{O}$ 150 K spectrum provided a better fit and was used for further fits. This intermediate result is presented in Fig. 6 where the observational spectrum is fitted by two laboratory spectra, $\text{MgSiO}_3/\text{H}_2\text{O}$ 150 K and $\text{NH}_4^+/\text{OCN}^-$ 80 K. The NH_4^+ and OCN^- bands (6.85 and 7.6 μm correspondingly) are fitted well but it is obvious that there is missing signal between 6.2 and 6.7 and between 7.0 and 7.6 μm .

Exactly in these regions (6.2 – 6.7 and 7.0 – 7.3 μm), ammonium carbamate ($\text{NH}_4^+\text{NH}_2\text{COO}^-$), the product of $\text{CO}_2 + \text{NH}_3$ chemistry, has spectral signatures (Bossa et al. 2008; Potapov et al. 2019). The next input to the model was the “chemistry” spectrum, which contains the aforementioned bands of ammonium carbamate. This led to the best fit shown in Fig. 5. An excess signal around 7.5 μm in the observational spectrum points to possible additional components. As potential candidates we considered refractory hydrocarbons obtained by UV irradiation of CH_3OH ice (the spectrum was presented in Potapov et al. 2021), as well as CH_4 from LIDA database (presented in Rachid et al. 2020). Additions of these components did not improve the fit. The AIC and RMSE values are presented in Table 5.

3.5. 8 – 16-micron region

This spectral region contains the silicate stretching band around 10 μm , the water librational band around 12 μm , and the CO_2 bending band around 15 μm . The best fit model is presented in Fig. 7 and was obtained using the input mixture of the spectra of $\text{MgSiO}_3/\text{H}_2\text{O}$ 150 K, $\text{Mg}_2\text{SiO}_4/\text{H}_2\text{O}$ 150 K (both spectra presented in Potapov et al. 2021), CO_2 , $\text{H}_2\text{O}/\text{CO}_2$, and $\text{H}_2\text{O}/\text{NH}_3$ (the spectra were presented in Rocha & Pilling 2014; Rocha et al. 2017; McClure et al. 2023), correspondingly.

Table 5. EMIIGMA fitting results for the 5.6 – 8.0 μm wavelength region.

Component and Fit	Combination #						
	1	2	3	4	5	6	7
MgSiO ₃ /H ₂ O 100 K	x		x				
NH ₄ ⁺ 12 K	x	x					
MgSiO ₃ /H ₂ O 150 K		x		x	x	x	x
NH ₄ ⁺ 80 K			x	x	x	x	x
“Chemistry”					x	x	x
Hydrocarbons						x	
CH ₄							x
AIC	16.2	14.5	12.7	11.4	8.1	10.2	10.1
RMSE	0.049	0.046	0.042	0.038	0.020	0.020	0.020

Notes. The Akaike Information Criterion (AIC) and Root-mean-square error (RMSE) for various input mixtures are listed. The best fit parameters are marked in boldface.




Fig. 7. Observations (black), the best fit model for the 8 – 16 μm region (red) and its components: MgSiO₃/H₂O 150 K (green), Mg₂SiO₄/H₂O (blue), H₂O/NH₃ (magenta), H₂O/CO₂ (orange), CO₂ (brown).

We started with three-component models including CO₂ and H₂O/CO₂ spectra (basing on the CO₂ results presented above) and 10, 100 or 150 K MgSiO₃/H₂O spectra. Inclusion of the second CO₂ component was necessary to fit the CO₂ band at 15 microns. We clearly observe the splitting of the band caused by the segregation and/or distillation causing CO₂ to be present in pure form as well (cf. Isokoski et al. 2013). The MgSiO₃/H₂O 150 K spectrum provided the best fit from the three choices, and we used it in the following. At the next step, we added spectra of other silicates mixed with water ice, namely the Mg₂SiO₄/H₂O 150 K and Mg₂SiO₄/H₂O 150 K spectra. Addition of the Mg₂SiO₄/H₂O spectrum led to an improvement of the fit. At the next step, we tried to add low temperature components, such as MgSiO₃/H₂O 10 and 100 K, H₂O/CH₃OH, H₂O/NH₃ and H₂O/CO/CO₂/NH₃/CH₃OH at 10 K, and high-temperature components, such as H₂O/CO₂/NH₃ at 150 K and H₂O/CO₂/NH₃/CH₃OH at 100 K. Only the H₂O/NH₃ addition

improved the fit. Thus, a small amount of cold ammonia may be present in the disk ices. The AIC and RMSE values are presented in Table 6.

4. Discussion

4.1. NH₃ chemistry and complex organic molecules

Ammonium carbamate is a product of the NH₃ + CO₂ chemistry (reaction CO₂ + 2NH₃ → NH₄⁺NH₂COO⁻), which, as it was shown by laboratory experiments, can be triggered by heat (above 60 – 65 K, Bossa et al. 2008; Potapov et al. 2019, 2020) and much less efficiently by energetic processing (Bossa et al. 2008). The detection of ammonium carbamate is consistent with the detection of NH₄⁺ and OCN⁻, also the products of NH₃ chemistry, where NH₃ reacts with HNC leading to the production of both species via a direct pure thermal acid-base reaction or with CO leading, if energetically triggered (UV photons, ions), to the production of OCN⁻ (Novozamsky et al. 2001; Martinez et al. 2014). NH₄⁺ can be also produced by ultraviolet photolysis of NH₃-H₂O ice mixtures (Moon et al. 2010) and, in principle, can be a product of dissociation of ammonium carbamate.

Pure thermal reactions definitely take place in planet-forming disks, thus, the formation of all three species (NH₄⁺NH₂COO⁻, NH₄⁺, OCN⁻) can be directly related to the disk chemistry. The detection of these molecules is consistent with a questionable detection of NH₃ and non-detection of warm CO₂, clearly showing their efficient involvement into the disk chemistry and their evolution tracks. Energetic processing may take place in both, the disk and its parent molecular cloud environments, however, midplanes and warmer layers of planet-forming disks – the locations of molecular ices, are shielded from external UV fields. An open question is the survival of the molecules, if formed in the parent molecular cloud, during the chemical evolution of the environment (transformation of a cloud into a disk). Dissociation, desorption and further reactions of the species may lead to their disappearance. Moreover, ammonium carbamate has not been detected in molecular clouds. Thus, the synthesis of the detected species can be referred to both the disk studied and the molecular cloud from which it was formed with a clear preference to the disk.

One of the fundamental questions regarding small Solar System bodies (comets, asteroids and their meteoritic remains) is the

Table 6. EMIIGMA fitting results for the 8 – 16 μm wavelength region.

Component and Fit	Combination #											
	1	2	3	4	5	6	7	8	9	10	11	12
MgSiO ₃ /H ₂ O 10 K	x					x						
H ₂ O/CO ₂	x	x	x	x	x	x	x	x	x	x	x	x
pure CO ₂	x	x	x	x	x	x	x	x	x	x	x	x
MgSiO ₃ /H ₂ O 100 K		x					x					
MgSiO ₃ /H ₂ O 150 K			x	x	x	x	x	x	x	x	x	x
Mg ₂ SiO ₄ /H ₂ O 150 K				x		x	x	x	x	x	x	x
MgFeSiO ₄ /H ₂ O 150 K					x							
H ₂ O/CH ₃ OH ^a								x				
H ₂ O/NH ₃ ^a									x			
H ₂ O/CO/CO ₂ /NH ₃ /CH ₃ OH ^a										x		
H ₂ O/CO ₂ /NH ₃ ^b											x	
H ₂ O/CO ₂ /NH ₃ /CH ₃ OH ^c												x
AIC	18.6	16.2	14.2	15.5	16.2	17.5	17.4	17.4	15.6	17.9	17.9	17.5
RMSE	0.050	0.045	0.040	0.038	0.040	0.038	0.038	0.038	0.033	0.039	0.039	0.038

Notes. The Akaike Information Criterion (AIC) and Root-mean-square error (RMSE) for various input mixtures are listed. The best fit parameters are marked in boldface.

^(a) ice mixture at 10 K ^(b) ice mixture at 150 K ^(c) ice mixture at 100 K

origin of detected complex organics, particularly building blocks of biological macromolecules, such as sugars, amino acids, and nucleobases. Whether these species originate from the parent to the Solar System protoplanetary disk or even the interstellar cloud is a long-standing discussion in the astrochemical community. Many complex organic molecules have been detected in comets. Our particular attention attracted the detection of ammonium salts, such as ammonium carbamate ($\text{NH}_4^+\text{NH}_2\text{COO}^-$), ammonium formate ($\text{HCOO}^-\text{NH}_4^+$), and ammonium sulphate ($(\text{NH}_4^+)_2\text{SO}_4^{2-}$) (Poch et al. 2020). Ammonium salts may dominate the reservoir of nitrogen in comets and asteroids and, considering previous exogenous delivery hypotheses (e.g., Oró 1961; Brack 1991), on the early Earth. Being delivered to the early Earth and mixed in liquid water, ammonium salts could participate in prebiotic reactions leading to the synthesis of amino acids, nucleobases and sugars (Lerner & Cooper 2005; Dziejcz et al. 2009; Callahan et al. 2011). For instance, ammonium carbamate can be a precursor of urea $\text{CO}(\text{NH}_2)_2$ and urea, in turn, is a possible precursor of pyrimidine $\text{C}_4\text{H}_4\text{N}_2$ required for the synthesis of nucleobases in RNA molecules (Robertson & Miller 1995). Thus, the detection of ammonium carbamate in the protoplanetary disk links parent protoplanetary and daughter planetary environments and brings us one more step closer to the understanding of the processes and conditions leading to the formation of prebiotic worlds. Moreover, as the Solar System is believed to be formed in a massive star-forming region, the Orion disks most likely represent analogues of the protoplanetary disk phase of the Solar System (e.g., Young 2020; Dauphas & Chaussidon 2011, and references therein). The detection of ammonium salts in the Orion disks can then be directly linked to their detection in Solar System objects like the comet 67P.

4.2. Warm vs Cold

Highly inclined disks, which are not completely edge on (inclination angles of 70 – 85 degrees) present a particular interest. As shown for the HH 48 NE disk (see Sturm et al. 2024), the location of the ices detected in the NIRSpc range is mainly warm layers significantly above the midplane. This is consistent with our detections of warm water ice (100/150 K) at 3 microns, warm NH_4^+ (80 K) at 7 microns, and ammonium carbamate (a result of “warm” chemistry). Detection of cold CO speaks for its presence only in the midplane of the disk due to its very low desorption temperature (around 30 K) and Sturm et al. (2024) had the same view. Detection of only cold CO₂ (having the desorption temperature of 85 K) points to its presence only near the midplane of the disk and its depletion in the warmer layer of the disk due to the efficient $\text{NH}_3 + \text{CO}_2$ chemistry discussed above. In the 10-micron range, the absorption is predominantly from regions near the disk midplane (cf. Sturm et al. 2024), and here we detect a mixture cold and warm materials.

5. Conclusions

The highly inclined disk d216-0939 has been observed by JWST. The analysis of the absorption bands shows the presence of simple and complex molecular species. For the first time, a complex organic molecule ammonium carbamate has been detected in astrophysical environments pointing to the complex chemistry taking place in protoplanetary disks. Together with the detection of ammonium salts in the Solar System Comet 67P, this result links protoplanetary disks and minor bodies of the Solar System.

Acknowledgements. We thank the referee for a constructive report. We are grateful to the JWST team for the wonderful observations. This work is based on observations made with the NASA/ESA/CSA James Webb Space Telescope. The data were obtained from the Mikulski Archive for Space Telescopes at the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS 5-03127 for JWST.

These observations are associated with program #1741. The study was supported by the Federal Ministry for Economic Affairs and Climate Action on the basis of a decision by the German Bundestag (the German Aerospace Center project 50OR2215). A. P. acknowledges support from the Deutsche Forschungsgemeinschaft (Heisenberg grant PO 1542/7-1).

References

- Arumainayagam, C. R., Garrod, R. T., Boyer, M. C., et al. 2019, *Chemical Society Reviews*, 48, 2293
- Böker, T., Arribas, S., Lützgendorf, N., et al. 2022, *A&A*, 661, A82
- Bossa, J. B., Theulé, P., Duvernay, F., Borget, F., & Chiavassa, T. 2008, *A&A*, 492, 719
- Brack, A. 1991, *Grana*, 30, 505
- Burnham, K. & Anderson, D. 2002, *Model selection and multimodel inference: a practical information-theoretic approach* (Springer Verlag)
- Bushouse, H., Eisenhamer, J., Dencheva, N., et al. 2024, *JWST Calibration Pipeline*, 10.5281/zenodo.6984365
- Callahan, M. P., Smith, K. E., Cleaves, H. J., et al. 2011, *Proceedings of the National Academy of Science*, 108, 13995
- Dartois, E., Noble, J. A., Caselli, P., et al. 2024, *Nature Astronomy*, 8, 359
- Dartois, E., Noble, J. A., Ysard, N., Demyk, K., & Chabot, M. 2022, *A&A*, 666, A153
- Dauphas, N. & Chaussidon, M. 2011, *Annual Review of Earth and Planetary Sciences*, 39, 351
- Dominik, C., Min, M., & Tazaki, R. 2021, *OpTool: Command-line driven tool for creating complex dust opacities*, *Astrophysics Source Code Library*, record ascl:2104.010
- Dziedzic, P., Bartoszewicz, A., & Córdoba, A. 2009, *Tetrahedron Letters*, 50, 7242
- Fulvio, D., Potapov, A., He, J., & Henning, T. 2021, *Life*, 11, 568
- Isokoski, K., Poteet, C. A., & Linnartz, H. 2013, *A&A*, 555, A85
- Jakobsen, P., Ferruit, P., Alves de Oliveira, C., et al. 2022, *A&A*, 661, A80
- Lerner, N. R. & Cooper, G. W. 2005, *Geochim. Cosmochim. Acta*, 69, 2901
- Linnartz, H., Ioppolo, S., & Fedoseev, G. 2015, *International Reviews in Physical Chemistry*, 34, 205
- Martin, R. G. & Livio, M. 2012, *MNRAS*, 425, L6
- Martínez, R., Bordalo, V., da Silveira, E. F., & Boechat-Roberty, H. M. 2014, *MNRAS*, 444, 3317
- McClure, M. K., Rocha, W. R. M., Pontoppidan, K. M., et al. 2023, *Nature Astronomy*, 7, 431
- McGuire, B. A. 2022, *ApJS*, 259, 30
- Min, M., Bouwman, J., Dominik, C., et al. 2016, *A&A*, 593, A11
- Moon, E.-S., Kang, H., Oba, Y., Watanabe, N., & Kouchi, A. 2010, *ApJ*, 713, 906
- Mori, S., Okuzumi, S., Kunitomo, M., & Bai, X.-N. 2021, *ApJ*, 916, 72
- Nazari, P., Rocha, W. R. M., Rubinstein, A. E., et al. 2024, *A&A*, 686, A71
- Novozamsky, J. H., Schutte, W. A., & Keane, J. V. 2001, *A&A*, 379, 588
- Öberg, K. I. 2016, *Chemical Reviews*, 116, 9631–9663
- Oró, J. 1961, *Nature*, 190, 389
- Owen, J. E. 2020, *MNRAS*, 495, 3160
- Poch, O., Istiqomah, I., Quirico, E., et al. 2020, *Science*, 367, aaw7462
- Pontoppidan, K. M., Dartois, E., van Dishoeck, E. F., Thi, W. F., & d’Hendecourt, L. 2003, *A&A*, 404, L17
- Potapov, A., Bouwman, J., Jäger, C., & Henning, T. 2021, *Nature Astronomy*, 5, 78
- Potapov, A., Fulvio, D., Krasnokutski, S., Jäger, C., & Henning, T. 2022, *Journal of Physical Chemistry A*, 126, 1627
- Potapov, A., Jäger, C., & Henning, T. 2018, *ApJ*, 865, 58
- Potapov, A., Jäger, C., & Henning, T. 2020, *ApJ*, 894, 110
- Potapov, A. & McCoustra, M. 2021, *Int. Rev. Phys. Chem.*, 40, 299
- Potapov, A., Theulé, P., Jäger, C., & Henning, T. 2019, *ApJ*, 878, L20
- Rachid, M. G., Terwisscha van Scheltinga, J., Koletzki, D., & Linnartz, H. 2020, *A&A*, 639, A4
- Raunier, S., Chiavassa, T., Duvernay, F., et al. 2004, *A&A*, 416, 165
- Robertson, M. P. & Miller, S. L. 1995, *Nature*, 375, 772
- Rocha, W. R. M., McClure, M. K., Sturm, J. A., et al. 2025, *A&A*, 693, A288
- Rocha, W. R. M., Perotti, G., Kristensen, L. E., & Jørgensen, J. K. 2021, *A&A*, 654, A158
- Rocha, W. R. M. & Pilling, S. 2014, *Spectrochimica acta. Part A, Molecular and biomolecular spectroscopy*, 123, 436
- Rocha, W. R. M., Pilling, S., de Barros, A. L. F., et al. 2017, *MNRAS*, 464, 754
- Rocha, W. R. M., Rachid, M. G., Olsthoorn, B., et al. 2022, *A&A*, 668, A63
- Rocha, W. R. M., van Dishoeck, E. F., Ressler, M. E., et al. 2024, *A&A*, 683, A124
- Ros, K. & Johansen, A. 2024, *A&A*, 686, A237
- Sandford, S. A., Nuevo, M., Bera, P. P., & Lee, T. J. 2020, *Chemical Reviews*, 120, 4616
- Schutte, W. A., Boogert, A. C. A., Tielens, A. G. G. M., et al. 1999, *A&A*, 343, 966
- Sheehan, P. D., Tobin, J. J., Federman, S., Megeath, S. T., & Looney, L. W. 2020, *ApJ*, 902, 141
- Smith, N., Bally, J., Licht, D., & Walawender, J. 2005, *AJ*, 129, 382
- Sturm, J. A., McClure, M. K., Beck, T. L., et al. 2023a, *A&A*, 679, A138
- Sturm, J. A., McClure, M. K., Bergner, J. B., et al. 2023b, *A&A*, 677, A18
- Sturm, J. A., McClure, M. K., Harsono, D., et al. 2024, *A&A*, 689, A92
- Terada, H. & Tokunaga, A. T. 2012, *ApJ*, 753, 19
- Terada, H., Tokunaga, A. T., Pyo, T.-S., et al. 2012, *AJ*, 144, 175
- Terwisscha van Scheltinga, J., Ligterink, N. F. W., Boogert, A. C. A., van Dishoeck, E. F., & Linnartz, H. 2018, *A&A*, 611, A35
- Theulé, P., Duvernay, F., Danger, G., et al. 2013, *Advances in Space Research*, 52, 1567
- van Dishoeck, E. F., Kristensen, L. E., Mottram, J. C., et al. 2021, *A&A*, 648, A24
- Whittet, D. C. B., ed. 2003, *Dust in the galactic environment* (Institute of Physics (IOP) Publishing), 2nd Edition
- Wright, G. S., Rieke, G. H., Glasse, A., et al. 2023, *PASP*, 135, 048003
- Yang, Y.-L., Green, J. D., Pontoppidan, K. M., et al. 2022, *ApJ*, 941, L13
- Young, E. D. 2020, in *IAU Symposium, Vol. 345, Origins: From the Protosun to the First Steps of Life*, ed. B. G. Elmegreen, L. V. Tóth, & M. Güdel, 70–77