

SSL4EO-S12 v1.1: A Multimodal, Multiseasonal Dataset for Pretraining, Updated

Benedikt Blumenstiel
IBM Research Europe
benedikt.blumenstiel@ibm.com

Nassim Ait Ali Braham
German Aerospace Center
nassim.aitalibraham@dlr.de

Conrad M Albrecht
German Aerospace Center
conrad.albrecht@dlr.de

Stefano Maurogiovanni
Juelich Supercomputing Centre
s.maurogiovanni@fz-juelich.de

Paolo Fraccaro
IBM Research Europe
paolo.fraccaro@ibm.com

Abstract

This technical report presents SSL4EO-S12 v1.1, a multimodal, multitemporal Earth Observation dataset designed for pretraining large-scale foundation models. Building on the success of SSL4EO-S12, this extension addresses the previous challenges of data misalignment and a limited data structure for low-barrier, analysis-ready EO processing. SSL4EO-S12 v1.1 covers the world’s 10,000 largest cities and its surroundings within a 50km radius across four seasons, resulting in a diverse collection of nearly one million patches. SSL4EO-S12 v1.1 packages the data in Zarr file format for cloud-efficient loading and representation of meta-information such as including cloud masks and geolocation. Released under the CC-BY-4.0 license, SSL4EO-S12 v1.1 facilitates open research and provides a robust foundation for future advancements in self-supervised learning and geospatial analysis. The dataset is available online through <https://datapub.fz-juelich.de/ssl4eo-s12>, and we provided additional resources at <https://github.com/DLR-MF-DAS/SSL4EO-S12-v1.1>.

1. Introduction

Large-scale Earth Observation (EO) datasets have played a pivotal role in the development of powerful Foundation Models (FM) [4, 8, 9, 11]. By training on diverse and extensive collections of remote sensing imagery, these models achieve better accuracy and converge faster [8, 11]. Among such datasets, SSL4EO-S12 [10] has been widely adopted and enabled various new EO FMs.

Co-funded by the European Union (Horizon Europe, Embed2Scale, 101131841). We also thank Yi Wang, main architect of the SSL4EO-S12 dataset for valuable discussions.

Despite its success, the original SSL4EO-S12 dataset presented two main challenges: (1) misalignments between Sentinel-1 (S-1) and Sentinel-2 (S-2) modalities and (2) limited availability of analytics-ready data (ARD). To address these issues, we present SSL4EO-S12 v1.1. We ensure proper alignment by downloading larger areas and reprojecting S-1 GRD data to match S-2 L2A coordinates. Furthermore, we introduce stricter filtering and provide the dataset in an ARD format suitable for direct pretraining of EO FMs. Figure 1 illustrates S-2 L2A and S-2 GRD time series samples from SSL4EO-S12 v1.1.

This technical report is organized as follows: We first detail the data processing steps necessary to construct SSL4EO-S12 v1.1, including alignment, reprojection, and filtering. We then present an overview of the final dataset and discuss its applicability for EO FM training pipelines.

2. Related Work

The development of large-scale, multimodal Earth Observation (EO) datasets has been critical for advancing self-supervised learning in remote sensing. Existing datasets such as BigEarthNet-MM [7] and SEN12MS [6] have provided valuable baselines, yet they often focus on single timestamps, limiting their utility for pre-training foundation models that require diverse spatiotemporal inputs [8].

SSL4EO-S12 addressed many of these limitations by including multiple modalities (e.g., S-1 and S-2) and covering multiple seasons, thereby supporting a broader range of downstream tasks [10]. Several state-of-the-art models have leveraged SSL4EO-S12 to achieve significant improvements in accuracy and generalizability [4, 9, 10]. For instance, contrastive learning approaches and transformer-based architectures have demonstrated notable performance gains when pre-trained on SSL4EO-S12.




Figure 1. Example patches from SSL4EO-S12 v1.1 showing four columns with timestamps of Sentinel-2 (S-2) L2A (left) and Sentinel-1 (S-1) GRD (right) products. S-1 GRD is visualized using a VH-VV-VH pseudo coloring. VV is scaled from $-30 - 5$ db while VH is scaled from $-40 - 0$ db to account for their value ranges.

3. Data Processing

All data for SSL4EO-S12 v1.1 were acquired from Google Earth Engine [5]. In SSL4EO-S12, all samples were projected onto EPSG:4326, which can distort distances, especially near the poles. We retain the native UTM zones in v1.1 to preserve spatial accuracy and minimize unnecessary reprojections. For each location in v1, four timestamps from different seasons were selected from 2019 to 2021 with cloud coverage below 10% [10]. We selected the timestamp closest to the S-2 timestamps in v1. To ensure correct spatial alignment, we first downloaded a larger area of interest for both S-2 and S-1 to buffer spatial mismatch on reprojection of these modalities when paired with other datasets. Next, we checked that all S-2 timestamps for a location shared the same UTM zone. If timestamps spanned multiple zones—common at UTM boundaries—we picked the majority CRS for that location, cropped a reference timestamp to a common size of 264×264 pixels, and reprojected the remaining timestamps accordingly. Subsequently, we projected S-1 patches onto the curated 264×264 -pixel S-2 patches. Following v1 [10], we fixed the patch sizes to 264 pixels as it is divisible by six, accommodating S-2 pixel resolutions of 10 m, 20 m, and 60 m. All bands were upsampled to 10 m using nearest interpolation.

Next, we dropped any location with more than 1% of missing values (NaN) for any timestamp or channel. The remaining NaNs were filled in using nearest neighbor inter-

polation. We split the dataset into training and validation sets, following a 99%-to-1% partition defined by the *TerraMesh* dataset to avoid spatial overlap that may leak information while training models. We add a numerical offset of +1000 to the data to facilitate the latest processing standard defined by ESA [1].

The cloud mask provided by S-2 products is prone to errors. Correspondingly, it served for data sampling only. Following [3], we used the SEnSeI v2 model [2] to provide accurate cloud masks for SSL4EO-S12 v1.1.

Furthermore, we provide S-2 RGB data based on L2A products. For reasons of data compression, reflectance value ranges of 0–2000 get mapped to 0–255 (single-byte integers). However, this approach may lead to saturation in bright regions (e.g., clouds, snow, and sand). To address this issue, we first remove the offset from the data. Next, we determine the 2% and 98% quantiles over the RGB channels within each image and apply a factor of 0.5 to values outside these bounds to reduce extreme pixel intensities. We then map these adjusted values to an 8-bit (0–255) range, using the 0.2% and 99.8% quantiles as the lower and upper limits, respectively. If the computed upper limit falls below 2000, we clamp it to 2000. Finally, if the median of the RGB values in an image is below 1000, we set the lower limit to 0 in order to avoid further darkening already dim images. The exact transformation is provided in Listing 1.

¹TerraMesh is a larger pretraining dataset to be released in Q2 2025.

Listing 1. Pseudocode for the S-2-to-RGB transformation.

```

import numpy as np

def transform_image(img, offset=1000):
    img = img - offset
    Q2, Q98 = np.quantile(img, [0.02, 0.98])
    img = np.where(img >= Q2, img,
                   Q2 + (img - Q2) * 0.5)
    img = np.where(img <= Q98, img,
                   Q98 + (img - Q98) * 0.5)
    Q02, Q50, Q998 = np.quantile(img,
                                   [0.002, 0.5, 0.998])
    U = max(2000, Q998)
    L = 0 if Q50 < 1000 else Q02
    img = (img - L) / (U - L) * 255

    img = np.clip(img, 0, 255)
    return img.astype(np.uint8)

```

This pipeline ensures a more visually balanced and artifact-free RGB representation of the S-2 data, as visible in Figure 1.

For efficient data access and storage, we shuffled and stacked the samples into Zarr files (`ZipStore`, Zarr version 2) in chunks of 64 samples and four timestamps each. This approach reduces the total file count, which is often important for storing data on an HPC cluster or Hugging-Face. Using Zarr Zip files, the data remains compatible with the common tool `xarray` and enables lazy data loading via `dask`. The data is chunked by timestamp to enable efficient loading of individual time steps if no time series is needed. The S-2 bands are stored as 16-bit (2 bytes) integers, and the S-1 data is stored as 16-bit floats. Combined with Zarr’s built-in compression, these measures significantly reduce storage overhead.

4. Dataset

The final SSL4EO-S12 v1.1 dataset (ARD version) contains 246,144 distinct locations, each with four timestamps, resulting in a total of 984,576 samples. The entire dataset is split into 3,846 Zarr Zip files for efficient handling. Figure 2 visualizes the global distribution of these samples.

Like v1, the dataset covers all four seasons, including snowy imagery. Sampling within a 50 km radius of the world’s 10,000 most populated cities ensures near-global coverage while focusing on urban areas but also includes ocean patches and rural areas. This diversity makes EO FMs pre-trained on SSL4EO-S12 v1.1 suitable for numerous downstream tasks, such as land use and change detection, segmentation of floods, or object detection for urban planning. The dataset also supports the training of multi-

Table 1. Summary of modalities included in SSL4EO-S12 v1.1. The range includes the typical value range, including the offset, but values below and above the range are present as well.

Modality	DType	Range	Units	# Bands
S-1 GRD	float16	-50 – +1	dB	2
S-2 L1C	int16	1k – 11k	DN	13
S-2 L2A	int16	1k – 11k	DN	12
S-2 RGB	uint8	0 – 255	–	3

temporal models as well as multimodal fusion techniques with S-1 and S-2 spatially aligned. Table 1 includes further details about the modalities. Providing S-2 L1C and S-2 L2A ensures that pre-trained models can be used for many downstream tasks regardless of top-of-atmosphere (L1C) or bottom-of-atmosphere information (L2A).

We provide overviews of typical Zarr Zip files in the git repository. Each file contains 64 samples and four timestamps, making data loading more efficient during pre-training. Data loading speed is an important concern for small models—loading individual samples can be slow. Our chunk size allows an entire mini-batch of 64 patches to be read at once to improve GPU utilization and to reduce the overall I/O overhead. For small models, a batch size of 64 likely fits into memory of current GPUs. However, larger models may require smaller effective batch sizes (as of 2024). This is easily handled by subsampling chunks. For larger models, a longer processing time allows to load data in parallel. The main trade-off with fixed-chunk sizes is reduced shuffling within individual batches from epoch to epoch. However, multi-GPU training typically provides sufficient shuffling at a global scale.

Relevant metadata is stored with the image data in the Zarr files as additional data variables. The metadata includes center coordinates in latitude and longitude (EPSG:4326) and pixel coordinates in the corresponding UTM zones, the file identifiers, as well as timestamps as defined by ESA. The SEnSel v2 cloud mask includes classes: *land*, *water*, *snow*, *thin cloud*, *thick cloud*, *cloud shadow*, and *no data*. The shared storage in one Zarr file simplifies data management and allows users to efficiently reference image location and timestamps.

SSL4EO-S12 v1.1 is released under the CC-BY-4.0 license, which permits broad adoption and reuse in both academic and industrial environments. SSL4EO-S12 v1.1 adds to the family of SSL4EO-*, open-source datasets to serve the Earth observation data science community.

Zarr documentation: <https://zarr-specs.readthedocs.io/en/latest/v2/v2.0.html>




Figure 2. Global distribution of SSL4EO-S12 v1.1 training (green) and validation (magenta) center points (not to scale).

References

- [1] ESA. Sentinel-2 products specification document. 2021. <https://sentinel.esa.int/documents/247904/685211/S2-PDGS-TAS-DI-PSD-V14.9.pdf>.
- [2] Alistair Francis. Sensor independent cloud and shadow masking with partial labels and multimodal inputs. *IEEE Transactions on Geoscience and Remote Sensing*, 2024.
- [3] Alistair Francis and Mikolaj Czerkawski. Major tom: Expandable datasets for earth observation. *arXiv preprint arXiv:2402.12095*, 2024.
- [4] Anthony Fuller, Koreen Millard, and James Green. Croma: Remote sensing representations with contrastive radar-optical masked autoencoders. *Advances in Neural Information Processing Systems*, 36, 2024.
- [5] Noel Gorelick, Matt Hancher, Mike Dixon, Simon Ilyushchenko, David Thau, and Rebecca Moore. Google earth engine: Planetary-scale geospatial analysis for everyone. *Remote Sensing of Environment*, 2017.
- [6] Michael Schmitt, Lloyd Haydn Hughes, Chunping Qiu, and Xiao Xiang Zhu. Sen12ms—a curated dataset of georeferenced multi-spectral sentinel-1/2 imagery for deep learning and data fusion. *arXiv preprint arXiv:1906.07789*, 2019.
- [7] Gencer Sumbul, Arne De Wall, Tristan Kreuziger, Filipe Marcelino, Hugo Costa, Pedro Benevides, Mario Caetano, Begüm Demir, and Volker Markl. Bigearthnet-mm: A large-scale, multimodal, multilabel benchmark archive for remote sensing image classification and retrieval [software and data sets]. *IEEE Geoscience and Remote Sensing Magazine*, 9(3):174–180, 2021.
- [8] Daniela Szwarcman, Sujit Roy, Paolo Fraccaro, Porsteinn Elfi Gíslason, Benedikt Blumenstiel, Rinki Ghosal, Pedro Henrique de Oliveira, Joao Lucas de Sousa Almeida, Rocco Sedona, Yanghui Kang, et al. Prithvi-eo-2.0: A versatile multi-temporal foundation model for earth observation applications. *arXiv preprint arXiv:2412.02732*, 2024.
- [9] Yi Wang, Conrad M Albrecht, Nassim Ait Ali Braham, Chenying Liu, Zhitong Xiong, and Xiao Xiang Zhu. Decur: decoupling common & unique representations for multimodal self-supervision. *arXiv preprint arXiv:2309.05300*, 2023.
- [10] Yi Wang, Nassim Ait Ali Braham, Zhitong Xiong, Chenying Liu, Conrad M Albrecht, and Xiao Xiang Zhu. Ssl4eo-s12: A large-scale multimodal, multitemporal dataset for self-supervised learning in earth observation [software and data sets]. *IEEE Geoscience and Remote Sensing Magazine*, 11(3):98–106, 2023.
- [11] Zhitong Xiong, Yi Wang, Fahong Zhang, Adam J Stewart, Joëlle Hanna, Damian Borth, Ioannis Papoutsis, Bertrand Le Saux, Gustau Camps-Valls, and Xiao Xiang Zhu. Neural plasticity-inspired foundation model for observing the earth crossing modalities. *arXiv e-prints*, pages arXiv–2403, 2024.