Periodic Response Solutions to Multi-Dimensional Nonlinear Schrödinger equation with unbounded perturbation

Zuhong You $^{\ast a}$ and Xiaoping Yuan $^{\ast b}$

 a School of Mathematical Sciences, Fudan University, Shanghai 200433, P. R. China

^b School of Mathematical Sciences, Fudan University, Shanghai 200433, P. R. China

Abstract

By applying the Craig-Wayne-Bourgain (CWB) method, we establish the existence of periodic response solutions to multi-dimensional nonlinear Schrödinger equations (NLS) with unbounded perturbation.

1 Introduction

In the last 40 years, significant progress has been made in the KAM (Kolmogorov-Arnold-Moser) theory for nonlinear Hamiltonian partial differential equations (PDEs). The first existence results were established by Kuksin[Kuk87] and Wayne[Way90], who studied nonlinear wave (NLW) and Schrödinger equations (NLS) in one spatial dimension under Dirichlet boundary conditions. There are two main approaches: the classical KAM technique (e.g., [Kuk87], [Way90], [Kuk93], [P96], [CY00], [EK10], [Yua21]) and the Craig-Wayne-Bourgain (CWB) method (e.g., [CW93], [Bou94],[Bou95], [Bou98],[Bou05], [BP11], [BB13], [Wan16], [BB20]). There are too many works to list here in this field.

In the context of unbounded perturbations, where the nonlinearity involves derivatives, KAM theory encounters greater challenges compared to the bounded case. The initial KAM theorem for unbounded perturbations was pioneered by Kuksin[Kuk98][Kuk00], who studied small denominator equations with large variable coefficients and developed applicable KAM theorems to investigate the persistence of finite-gap solutions for the KdV equation. The estimate concerning the small-denominator equation with large variable coefficients is now called the Kuksin lemma[Kuk87]. Later, Liu and Yuan[LY10][LY11] extended the Kuksin lemma to the limiting case and established the KAM theorem for quantum Duffing oscillators, derivative nonlinear Schrödinger equations (DNLS) and and Benjamin-Ono equations. The Italian school (e.g., [BBP13], [BBM14], [BBM16], [BBHM18]) has developed a novel approach for addressing unbounded perturbations.

All existent approaches dealing with unbounded perturbations rely on classical KAM techniques. As for the CWB method, it was only briefly mentioned by Bourgain in [Bou94], where he suggested in the final remark that his analysis could be extended to unbounded perturbations, such as the Hamiltonian "derivative" wave equation. However, no details were provided, and there have been no further developments applying the CWB method for unbounded perturbations until today. Moreover, all existing results for unbounded perturbations are restricted to the case where the spatial dimension is equal to 1. To the best of our knowledge, there are currently no established results for cases where the perturbations are unbounded and the spatial dimension is greater than one. In this paper, we take a step towards

^{*}The work was supported by National Natural Science Foundation of China (Grant NO. 12371189).

addressing this gap by employing the CWB method to study periodic response solutions of multi-dimensional nonlinear Schrödinger equations with fractional derivative perturbations (unbounded ones).

Consider the multi-dimensional nonlinear Schrödinger equations (NLS) with fractional derivatives:

$$iu_t - \Delta u + u + D^{\alpha}(|u|^2 u) = \varepsilon P(x, \omega t)$$
(1)

under periodic boundary conditions (i.e., $x \in \mathbb{T}^d$). Here, $P(x, \theta) : \mathbb{T}^d \times \mathbb{T} \to \mathbb{R}$ is Gevrey smooth. Furthermore, the fractional derivative D^{α} is defined as follows:

Let $f \in L^2(\mathbb{T}^d)$ be a smooth periodic function on the torus \mathbb{T}^d . We define

$$D^{\alpha}f(x) = \sum_{n \in \mathbb{Z}^d} \langle n \rangle^{\alpha} \hat{f}(n) e^{in \cdot x},$$
(2)

where $\hat{f}(n)$ is the *n*-th Fourier coefficient of f and $\langle n \rangle = \sqrt{\sum_{1 \le j \le d} |n_j|^2 + 1}$.

Furthermore, we define the Gevrey norm of P as follows:

$$||P||_{c} = \sum_{n,k} |\hat{P}(n,k)| e^{2(|n|+|k|)^{c}},$$
(3)

where $\hat{P}(n,k)$ refers to the Fourier coefficients of P and c is referred to the Gevrey index.

We assume $\omega \in [1, 2]$ is the parameter. Our aim is to prove that there is a $\frac{2\pi}{\omega}$ -time periodic solution to (1) for "most" $\omega \in [1, 2]$. More precisely, we have the following theorem:

Theorem 1.1. Assume $\alpha < \frac{1}{30(d(2d+2)^{d+2}+2)}$, $P(x,\theta) : \mathbb{T}^d \times \mathbb{T} \to \mathbb{R}$ is Gevrey smooth, c is small enough and $\|P\|_c < 1$. Then, for $\varepsilon > 0$ small enough, there exists a set $I_{\varepsilon} \subset [1,2]$ with

mes
$$I_{\varepsilon} \to 0$$
, as $\varepsilon \to 0$, (4)

such that for any $\omega \in [1,2] \setminus I_{\varepsilon}$, there exists a Gevrey smooth function

$$u: \mathbb{T}^d \times \mathbb{R} \to \mathbb{R},\tag{5}$$

which solves (1) and is $\frac{2\pi}{\omega}$ periodic in time. Moreover, we have

$$\sum_{n,k} |\hat{u}(n,k)| e^{\frac{1}{2}(|n|+|k|)^c} < C\varepsilon^{1/4},\tag{6}$$

where $\hat{u}(n,k)$ is the Fourier coefficient of u and C is a constant depending on d, α and c.

Remark 1. In this paper, the Gevrey index c is taken to be small enough for technical simplicity. To extend the proof to arbitrary values of $c \in (0,1)$, a more refined coupling lemma and more precise estimates of the Green's function are required.

Remark 2. The nonlinearity $|u|^2 u$ can be replaced by any $\frac{\partial H}{\partial \bar{u}}$ where $H(u, \bar{u}) = \sum a_{j,l} u^j \bar{u}^l$, $a_{jl} = a_{lj} \in \mathbb{R}$, is real analytic. When H is a polynomial, the proof follows the same line as described in this paper. When H is real analytic instead of polynomial, combining the scheme in this paper and (1) of Section 14 in [Bou98] gives the proof.

Remark 3. The idea to handle the unbounded perturbation is to define non-singular sites to be (n,k) which makes $\min_{\pm} |D_{\pm,n,k}| = \min_{\pm} |\pm k\omega + |n|^2 + 1| > N^{\delta}$ for some small δ , which is different from the usual ones and allows us to handle the unbounded perturbation for $(\alpha < \delta)$. Only when δ is small, can the existing separation lemma for $|n|^2$ handle it; thus, α in this paper is small.

We introduce some useful notations.

For any positive numbers a, b the notation $a \leq b$ means $Ca \leq b$ for some constant C > 0. By $a \ll b$ we mean that the constant C is very large. The various constants can be defined by the context in which they arise. Finally, N^{a-} means $N^{a-\epsilon}$ with some small $\epsilon > 0$ (the precise meaning of "small" can again be derived from the context).

2 Initial approximate solution

Choose and fix the various constants c, M, C_1, C_2, C_3 such that

$$M > 100, \quad 0 < c < \frac{\log \frac{17}{16}}{\log M}, \quad C_1 = 20d, \quad C_2 > 2, \quad C_3 > C_2 + 2.$$
 (7)

Replacing u by $\varepsilon^{\frac{1}{3}}u$, equation (1) becomes

$$iu_t - \Delta u + u + \varepsilon^{\frac{2}{3}} D^{\alpha}(|u|^2 u) = \varepsilon^{\frac{2}{3}} P(x, \omega t).$$
(8)

To solve a periodic solution with frequency ω of (8) is equivalent to solve

$$i\partial_{\omega}u - \Delta u + u + \varepsilon^{\frac{2}{3}}D^{\alpha}(|u|^{2}u) = \varepsilon^{\frac{2}{3}}P(x,\theta),$$
(9)

where $\partial_{\omega} u = \omega \partial_{\theta}$. Let

$$F(u) = i\partial_{\omega}u - \Delta u + u + \varepsilon^{\frac{2}{3}}D^{\alpha}(|u|^{2}u) - \varepsilon^{\frac{2}{3}}P(x,\theta).$$

Definition 2.1. For $f \in L^2(\mathbb{T}^{d+1})$, we denote

$$\Gamma_N f(x,\theta) = \sum_{|n| < N, |k| < N} \hat{f}(n,k) e^{in \cdot x + ik\theta}.$$
(10)

Definition 2.2. We say $u(x,\theta) \in L^2(\mathbb{T}^{d+1})$ is an $O(\delta)$ -approximate solution to (9) if

$$|F(u)|| \le \delta,\tag{11}$$

where $\|\cdot\|$ denote the norm of $L^2(\mathbb{T}^{d+1})$.

Lemma 2.3. Let $j_0 = \lfloor \frac{\log \log \frac{1}{\varepsilon}}{2 \log M} \rfloor$. There exists $u_{j_0}(x, \theta; \omega) \in L^2(\mathbb{T}^{d+1})$ smoothly defined for $\omega \in [1, 2]$ such that

- (i) supp $\hat{u}_{j_0} \subset B(0, M^{j_0})$.
- (ii) $||u_{j_0}|| < \varepsilon^{\frac{1}{5}}$.
- (iii) $|\hat{u}_{j_0}(\xi)| < e^{-|\xi|^c}$ and $|\partial \hat{u}_{j_0}(\xi)| < 2e^{-|\xi|^c}$, where $\xi = (n,k) \in \mathbb{Z}^{d+1}$ and ∂ refers to the derivative with respect to ω .

Furthermore, there exists a subset $\Lambda_{j_0} \subset [1,2]$ which is a union of disjoint intervals of size $\exp(-j_0^{C_3})$ such that

mes
$$[1,2] \setminus \Lambda_{j_0} \lesssim (\log \frac{1}{\varepsilon})^{-1} + e^{-j_0 - 9}$$

For $\omega \in \Lambda_{j_0}$, u_{j_0} is an $O(e^{-2(M^{j_0})^c})$ -approximate solution to (9), i.e., $||F(u_{j_0})|| < e^{-2(M^{j_0})^c}$. Moreover, for $\omega \in \Lambda_{j_0}$, we also have

$$\|\partial F(u_{j_0})\| < e^{-2(M^{j_0})^c}$$

Proof. Partition [1,2] into intervals $\{I_{\zeta}\}$ of size $\frac{1}{1-e^{-j_0-9}}\exp(-j_0^{C_3})$. Let

$$\begin{split} &\Lambda_{j_0} \\ = \bigcup \{ (1 - e^{-j_0 - 9}) I_{\zeta} : \exists \ \omega' \in I_{\zeta} \text{ such that } |-k\omega' + |n|^2 + 1 | \ge (\log \frac{1}{\varepsilon})^{-1} (1 + |k|)^{-\tau}, \\ &|n| < M^{j_0}, |k| < M^{j_0} \}. \end{split}$$

Here, $\tau = d + 1$ and $(1 - e^{-j_0 - 9})I_{\zeta}$ refers to $(1 - e^{-j_0 - 9})$ -dilation of I_{ζ} . For $\omega \in \Lambda_{j_0}$, we have

$$|-k\omega + |n|^{2} + 1| > (\log \frac{1}{\varepsilon})^{-1} (1 + |k|)^{-\tau} - \exp(-\frac{1}{2}j_{0}^{C_{3}})$$
$$> \frac{1}{2} (\log \frac{1}{\varepsilon})^{-1} (1 + |k|)^{-\tau}.$$

Thus, we have

$$[1,2] \setminus \Lambda_{j_0}$$

$$\subset \bigcup_{\substack{|n| < M^{j_0} \\ |k| < M^{j_0}}} \{\omega \in [0,1] : |-k\omega + |n|^2 + 1| < (\log \frac{1}{\varepsilon})^{-1} (1+|k|)^{-\tau} \} \cup (\bigcup_{\zeta} I_{\zeta} \setminus (1-e^{-j_0-9}) I_{\zeta}).$$

$$(12)$$

Denote

$$R_{n,k} = \{\omega \in [1,2] : |-k\omega + |n|^2 + 1| < (\log\frac{1}{\varepsilon})^{-1}(1+|k|)^{-\tau}\}.$$
(13)

Note that $R_{n,0} = \emptyset$ and $R_{n,k} = \emptyset$ for |n| > 2|k|. Thus, we have

$$\max [1,2] \setminus \Lambda_{j_0} < \sum_{\substack{|k| < M^{j_0} \\ k \neq 0}} \sum_{\substack{n \le 2|k| \\ n \le 2|k|}} \max R_{n,k} + e^{-j_0 - 9} \\ \lesssim \sum_{\substack{|k| < M^{j_0} \\ k \neq 0}} (\log \frac{1}{\varepsilon})^{-1} (1+|k|)^{-d-1} \cdot |k|^{d-1} + e^{-j_0 - 9} \\ \lesssim (\log \frac{1}{\varepsilon})^{-1} + e^{-j_0 - 9}.$$

For $\omega \in \Lambda_{j_0}$, let

$$\hat{u}_{j_0}(n,k) = \begin{cases} \frac{\varepsilon^{\frac{2}{3}}\hat{P}(n,k)}{-k\omega+|n|^2+1}, & |k| < M^{j_0}, |n| < M^{j_0}, \\ 0, & \text{otherwise.} \end{cases}$$

We have

$$\begin{aligned} |\hat{u}_{j_0}(n,k)| &< \varepsilon^{\frac{2}{3}-} e^{-\frac{3}{2}(|n|+|k|)^c}, \\ |\partial \hat{u}_{j_0}(n,k)| &< \varepsilon^{\frac{2}{3}-} e^{-\frac{3}{2}(|n|+|k|)^c}. \end{aligned}$$

By the definition of \hat{u}_{j_0} and Plancherel's Theorem, we have

$$\begin{aligned} \|F(u_{j_{0}})\| \\ = \|i\partial_{\omega}u_{j_{0}} - \Delta u_{j_{0}} + u_{j_{0}} + \varepsilon^{\frac{2}{3}}D^{\alpha}(|u_{j_{0}}|^{2}u_{j_{0}}) - \varepsilon^{\frac{2}{3}}P(x,\theta)\| \\ = \|\varepsilon^{\frac{2}{3}}D^{\alpha}(|u_{j_{0}}|^{2}u_{j_{0}}) - \varepsilon^{\frac{2}{3}}(1 - \Gamma_{M^{j_{0}}})P\| \\ \leq \|\varepsilon^{\frac{2}{3}}D^{\alpha}(|u_{j_{0}}|^{2}u_{j_{0}})\| + \|\varepsilon^{\frac{2}{3}}(1 - \Gamma_{M^{j_{0}}})P\| \\ \leq \varepsilon + \varepsilon^{\frac{2}{3}}\exp(-(2-)(M^{j_{0}})^{c}) \\ \leq \exp(-2(M^{j_{0}})^{c}). \end{aligned}$$
(14)

Moreover, we have

$$\begin{aligned} \|\partial F(u_{j_0})\| \\ = \|\partial \varepsilon^{\frac{2}{3}} D^{\alpha}(|u_{j_0}|^2 u_{j_0}) - \partial \varepsilon^{\frac{2}{3}} (1 - \Gamma_{M^{j_0}}) P\| \\ = \|\partial \varepsilon^{\frac{2}{3}} D^{\alpha}(|u_{j_0}|^2 u_{j_0})\| \\ \le \varepsilon < \exp(-2(M^{j_0})^c). \end{aligned}$$
(15)

By Whitney extension theorem, we can extend $\hat{u}_{j_0}(n,k)$ to the whole parameter space [1,2] and we have

$$\|\hat{u}_{j_0}(n,k)\|_{C^1([1,2])} < \varepsilon^{\frac{2}{3}-} e^{-(|n|+|k|)^c}.$$
(16)

This completes the proof.

3 The Newton iteration scheme

Assume that $u_j(x,\theta;\omega) \in L^2(\mathbb{T}^{d+1})$ $(j \geq j_0)$ is defined smoothly for $\omega \in [1,2]$. Moreover, assume that

(j-i) supp $\hat{u}_j \subset B(0, M^j)$.

(j-ii)
$$||u_j|| < (1 + \sum_{j'=j_0}^j \frac{1}{j'^2})\varepsilon^{1/5}.$$

(j-iii) $|\hat{u}_j(\xi)| < (1 + \sum_{j'=j_0}^j \frac{1}{j'^2})e^{-|\xi|^c}$ and $|\partial \hat{u}_j(\xi)| < 2(1 + \sum_{j'=j_0}^j \frac{1}{j'^2})e^{-|\xi|^c}$, where $\xi = (n,k) \in \mathbb{Z}^{d+1}$, and ∂ refers to the derivative with respect to ω .

(j-iv) $||F(u_j)|| < e^{-2(M^j)^c}$ and $||\partial F(u_j)|| < e^{-2(M^j)^c}$ are valid for ω restricted to a subset Λ_j of ω -parameter set which, in particular, we assume to be a union of disjoint intervals of size $\exp(-j^{C_3})$.

Let $u = u_j + v$, then we have

$$F(u) = i\partial_{\omega}u_j + i\partial_{\omega}v - \Delta u_j - \Delta v + u_j + v + \varepsilon^{\frac{2}{3}}D^{\alpha}\left((u_j + v)^2(\bar{u}_j + \bar{v})\right) - \varepsilon^{\frac{2}{3}}P$$
$$= F(u_j) + i\partial_{\omega}v - \Delta v + v + \varepsilon^{\frac{2}{3}}D^{\alpha}(2u_j\bar{u}_jv + u_j^2\bar{v}) + \varepsilon^{\frac{2}{3}}D^{\alpha}(2u_jv\bar{v} + \bar{u}_jv^2 + v^2\bar{v}).$$
(17)

Thus, we have the linearized equation

$$i\partial_{\omega}v - \Delta v + v + \varepsilon^{\frac{2}{3}}D^{\alpha}(2u_j\bar{u}_jv + u_j^2\bar{v}) = -F(u_j).$$
⁽¹⁸⁾

The equation conjugate to (18) is

$$-i\partial_{\omega}\bar{v} - \Delta\bar{v} + \bar{v} + \varepsilon^{\frac{2}{3}}D^{\alpha}(\bar{u}_{j}^{2}v + 2u_{j}\bar{u}_{j}\bar{v}) = -\overline{F}(u_{j}).$$
(19)

Let

$$w = \begin{pmatrix} v \\ \overline{v} \end{pmatrix}, \quad Q_j = \begin{pmatrix} -F(u_j) \\ -\overline{F}(u_j) \end{pmatrix}.$$

Then, the homological equation is

$$\begin{pmatrix} i & 0\\ 0 & -i \end{pmatrix} \partial_{\omega} w - \Delta w + w + \varepsilon^{\frac{2}{3}} D^{\alpha} \begin{pmatrix} 2u_j \bar{u}_j & u_j^2\\ \bar{u}_j^2 & 2u_j \bar{u}_j \end{pmatrix} w = Q_j.$$
(20)

Passing to Fourier coefficients, we get a lattice system of equations:

$$(D + \varepsilon^{\frac{2}{3}} \Lambda S_j) \widehat{w} = \widehat{Q}_j, \tag{21}$$

where

$$D = \begin{pmatrix} -k\omega + |n|^2 + 1 & 0\\ 0 & k\omega + |n|^2 + 1 \end{pmatrix},$$
(22)

$$\Lambda = \begin{pmatrix} k \cdot 0 + \langle n \rangle^{\alpha} & 0\\ 0 & k \cdot 0 + \langle n \rangle^{\alpha} \end{pmatrix},$$
(23)

and

$$S_{j} = \begin{pmatrix} S_{2u_{j}\bar{u}_{j}} & S_{u_{j}^{2}} \\ S_{\bar{u}_{j}^{2}} & S_{2u_{j}\bar{u}_{j}} \end{pmatrix},$$
(24)

where S_{ϕ} represents the Toeplitz operator corresponding to ϕ (i.e., $S_{\phi}((n,k), (n,k)) = \hat{\phi}(n - n', k - k')$). The index of the matrix $T_j = D + \varepsilon^{\frac{2}{3}} \Lambda S_j$ is $(\pm, n, k) \in \mathbb{Z}_2 \times \mathbb{Z}^{d+1}$. Denote $T_{j,N} = T_j|_{|n|,|k| < N}$.

Lemma 3.1. Let $N = M^{j+1}$. Under assumptions (j-i), (j-ii), (j-ii), (j-iv). Assume that we have confirmed

$$||T_{j,N}^{-1}|| < 2\exp(\log N)^{C_2} := B(N),$$
(25)

and

$$|T_{j,N}^{-1}(\xi,\xi')| < 2e^{-\frac{1}{2}|\xi-\xi'|^c} \text{ for } |\xi-\xi'| > N^{\frac{1}{2}},$$
(26)

for ω restricted to a subset $\Lambda'_j \subset \Lambda_j$, which we assume to be a union of disjoint intervals of size $\frac{1}{1-e^{-j-10}} \exp(-(j+1)^{C_3})$. Then there exist a subset $\Lambda_{j+1} \subset \Lambda'_j$ and a function $u_{j+1}(x,\theta;\omega) \in L^2(\mathbb{T}^{d+1})$ defined smoothly for $\omega \in [1,2]$ such that

 $((j+1)-i) \text{ supp } \hat{u}_{j+1} \subset B(0, M^{j+1}).$

$$((j+1)-ii) ||u_{j+1}|| < (1 + \sum_{j'=j_0}^{j+1} \frac{1}{j'^2})\varepsilon^{1/5}$$

 $\begin{array}{l} ((\mathbf{j+1})\text{-}\mathbf{i}\mathbf{i}\mathbf{i}) \ |\hat{u}_{j+1}(\xi)| < (1 + \sum_{j'=j_0}^{j+1} \frac{1}{j'^2})e^{-|\xi|^c} \ and \ |\partial \hat{u}_{j+1}(\xi)| < 2(1 + \sum_{j'=j_0}^{j+1} \frac{1}{j'^2})e^{-|\xi|^c}, \ where \ \xi = (n,k) \in \mathbb{Z}^{d+1}, \ and \ \partial \ refers \ to \ the \ derivative \ with \ respect \ to \ \omega. \end{array}$

 $\begin{array}{l} ((j+1)\text{-iv}) \ \|F(u_{j+1})\| < e^{-2(M^{j+1})^c} \ and \ \|\partial F(u_{j+1})\| < e^{-2(M^{j+1})^c} \ are \ valid \ for \ \omega \ restricted \ to \ \Lambda_{j+1} \\ which \ is \ a \ union \ of \ disjoint \ intervals \ of \ size \ \exp(-(j+1)^{C_3}). \end{array}$

$$((j+1)-v) \|\partial^{\beta}(u_{j+1}-u_j)\| < e^{-\frac{11}{8}(M^{j+1})^c}, \text{ where } \beta = 0, 1.$$

 $((j+1)-vi) \mod (\Lambda'_j \setminus \Lambda_{j+1}) \lesssim e^{-j-10}.$

Proof. For simplicity, we omit the subscript j of S_j and T_j . By the definition of S and assumptions, we have

$$|S((\pm, n, k), (\pm, n', k'))| < e^{-(1-)(|n-n'|+|k-k'|)^c},$$
(27)

and

$$|\partial S((\pm, n, k), (\pm, n', k'))| < e^{-(1-)(|n-n'|+|k-k'|)^c}.$$
(28)

We omit the index \pm . Thus, we have

$$|T(\xi,\xi')| < \varepsilon^{2/3} \langle n \rangle^{\alpha} e^{-(1-)|\xi-\xi'|^c}, \text{ for } \xi \neq \xi', \xi = (n,k),$$
(29)

and

$$\partial T(\xi,\xi')| < \varepsilon^{2/3} \langle n \rangle^{\alpha} e^{-(1-)|\xi-\xi'|^c}, \text{ for } \xi \neq \xi', \xi = (n,k).$$
(30)

Let $\omega \in \Lambda'_j$. Since one has

$$\partial T_N^{-1} = -T_N^{-1} (\partial T_N) T_N^{-1},$$
(31)

it follows that

$$\|\partial T_N^{-1}\| \le \|T_N^{-1}\|^2 \|\partial T_N\| < B(N)^2 N < e^{3(\log N)^{C_2}}.$$
(32)

When $|\xi - \xi'| > N^{3/4}$, we have

$$\begin{aligned} |(\partial T_N^{-1})(\xi,\xi')| &\leq \sum_{|\xi_1|,|\xi_2| < N} \left| T_N^{-1}(\xi,\xi_1) \right| \left| \partial T_N(\xi_1,\xi_2) \right| \left| T_N^{-1}(\xi_2,\xi') \right| \\ &\leq N^{2(d+1)+3} e^{5(\log N)^{C_2}} e^{-\frac{1}{2}(|\xi-\xi'| - 3N^{1/2})^c} \\ &\leq e^{-(\frac{1}{2}-)|\xi-\xi'|^c}. \end{aligned}$$
(33)

Since \hat{Q}_j is expressed as a polynomial in \hat{u}_j and $\hat{\bar{u}}_j$, and given that supp $\hat{u}_j \subset B(0, M^j)$, we get

 $u_{j+1} = u_j + v_j.$

$$\operatorname{supp} \hat{Q}_j \subset B(0, CM^j) \subset B(0, \frac{1}{4}M^{j+1}).$$
(34)

Let

$$\hat{w}_j = \begin{pmatrix} \hat{v}_j \\ \hat{\bar{v}}_j \end{pmatrix} = T_N^{-1} \hat{Q}_j, \tag{35}$$

and

It is straightforward that

$$\operatorname{supp} \hat{v}_j \subset B(0, M^{j+1}).$$
(36)

By (35), we have

$$\begin{aligned} \|v_{j}\| &\leq \|T_{N}^{-1}\| \|Q_{j}\| \\ &\leq \exp(\log N)^{C_{2}} e^{-2(M^{j})^{c}} \\ &< e^{-(2-)(M^{j})^{c}} < e^{-\frac{3}{2}(M^{j+1})^{c}}, \end{aligned}$$
(37)

provided that $c < \frac{\log \frac{4}{3}}{\log M}$. Moreover, we have

$$\begin{aligned} \|\partial v_{j}\| &\leq \|\partial T_{N}^{-1}\| \|Q_{j}\| + \|T_{N}^{-1}\| \|\partial Q_{j}\| \\ &\leq e^{3(\log N)^{C_{2}}} e^{-2(M^{j})^{c}} + e^{(\log N)^{C_{2}}} e^{-2(M^{j})^{c}} \\ &< e^{-(2-)(M^{j})^{c}} < e^{-\frac{3}{2}(M^{j+1})^{c}}. \end{aligned}$$
(38)

Thus, (36), (37) and (38) permit us to confirm ((j+1)-i), ((j+1)-ii), ((j+1)-iii), ((j+1)-v) for $u_{j+1} = u_j + v_j$ provided the ω -parameters are restricted to Λ'_j .

By (17), we have

$$\begin{pmatrix} F(u_{j+1}) \\ \overline{F}(u_{j+1}) \end{pmatrix} = \left((T - T_N) \hat{w}_j \right)^{\vee} + \varepsilon^{2/3} \begin{pmatrix} D^{\alpha} (2u_j v_j \bar{v}_j + \bar{u}_j v_j^2 + v_j^2 \bar{v}_j) \\ D^{\alpha} (2\bar{u}_j v_j \bar{v}_j + u_j \bar{v}_j^2 + v_j \bar{v}_j^2) \end{pmatrix},$$
(39)

where $\hat{w}_j = \begin{pmatrix} \hat{v}_j \\ \hat{v}_j \end{pmatrix}$. For the second term, we have

$$\|D^{\alpha}(2u_jv_j\bar{v}_j + \bar{u}_jv_j^2 + v_j^2\bar{v}_j)\| < e^{-(3-)(M^{j+1})^c}$$
(40)

and

$$\|\partial D^{\alpha}(2u_{j}v_{j}\bar{v}_{j} + \bar{u}_{j}v_{j}^{2} + v_{j}^{2}\bar{v}_{j})\| = \|D^{\alpha}\partial(2u_{j}v_{j}\bar{v}_{j} + \bar{u}_{j}v_{j}^{2} + v_{j}^{2}\bar{v}_{j})\| < e^{-(3-)(M^{j+1})^{c}}.$$
 (41)

Now we consider the first term. Denote P_K the projection on B(0, K). We have

$$(T - T_N)\hat{w}_j = (I - P_N)TP_{\frac{N}{2}}\hat{w}_j + (T - T_N)(\hat{w}_j - P_{\frac{N}{2}}\hat{w}_j)$$

= $(I - P_N)TP_{\frac{N}{2}}\hat{w}_j + (T - T_N)(I - P_{\frac{N}{2}})T_N^{-1}\hat{Q}_j$
= $(I - P_N)TP_{\frac{N}{2}}\hat{w}_j + (T - T_N)(I - P_{\frac{N}{2}})T_N^{-1}P_{\frac{N}{4}}\hat{Q}_j.$ (42)

By (29), (30), (37) and (38), we obtain

$$\|\partial^{\beta}(I - P_{N})TP_{\frac{N}{2}}\hat{w}_{j}\| < N^{\alpha}e^{-(1-)(\frac{1}{2}N)^{c}}e^{-\frac{3}{2}(M^{j+1})^{c}} < \frac{1}{3}e^{-\frac{1}{2}N^{c}}e^{-\frac{3}{2}(M^{j+1})^{c}} = \frac{1}{3}e^{-2(M^{j+1})^{c}},$$
(43)

for $\beta = 0, 1$. By (j-iv), (26) and (33), we get

$$\begin{aligned} \|\partial^{\beta}(T-T_{N})(I-P_{\frac{N}{2}})T_{N}^{-1}P_{\frac{N}{4}}\hat{Q}_{j}\| &< e^{-(\frac{1}{2}-)(\frac{1}{4}N)^{c}}e^{-2(M^{j})^{c}} \\ &\leq \frac{1}{3}e^{-\frac{17}{8}(M^{j})^{c}} \\ &\leq \frac{1}{3}e^{-2(M^{j+1})^{c}}, \end{aligned}$$
(44)

for $\beta = 0, 1$, provided that $c < \frac{\log \frac{17}{16}}{\log M}$. Hence, we obtain

$$\|\partial^{\beta}(T-T_N)\hat{w}_j\| < \frac{2}{3}e^{-2(M^{j+1})^c},\tag{45}$$

for $\beta = 0, 1$ and $\omega \in \Lambda'_j$. Thus, we have

$$\|\partial^{\beta} F(u_{j+1})\| < e^{-2(M^{j+1})^{c}},\tag{46}$$

for $\beta = 0, 1$ and $\omega \in \Lambda'_j$. Note that in the above v_j is defined on Λ'_j . We need to extend its definition to the entire ω -parameter set [1,2]. Assume

$$\Lambda'_j = \underset{\zeta}{\cup} I_{\zeta},\tag{47}$$

where $\{I_{\zeta}\}$ are disjoint intervals of size $\frac{1}{1-e^{-j-10}} \exp(-(j+1)^{C_3})$. For each ζ , denote $I'_{\zeta} \subset I_{\zeta}$ the $(1-e^{-j-10})$ -dilation of I_{ζ} with the same center. Let $0 \leq \psi_{\zeta} \leq 1$ be a smooth function satisfying

$$\psi_{\zeta} = 0$$
 outside $I_{\zeta}, \ \psi_{\zeta} = 1$ on I'_{ζ} , (48)

$$|\partial \psi_{\zeta}| < \exp(2(j+1)^{C_3}).$$
 (49)

Let

$$\tilde{v}_j = \sum_{\zeta} \psi_{\zeta} v_j, \tag{50}$$

then we have

$$\operatorname{supp} \hat{\tilde{v}}_j \subset B(0, M^{j+1}), \tag{51}$$

$$\|\partial^{\beta} \tilde{v}_{j}\| < e^{-(\frac{3}{2}-)(M^{j+1})^{c}}, \text{ for } \beta = 0, 1.$$
 (52)

Let $u_{j+1} = u_j + \tilde{v}_j$. Thus, (51) and (52) permit us to confirm ((j+1)-i), ((j+1)-ii), ((j+1)-iii), ((j+1)-iii), ((j+1)-iv). Let $\Lambda_{j+1} = \bigcup_{\zeta} I'_{\zeta}$, then ((j+1)-iv) is confirmed. Moreover, we have

$$\operatorname{mes}\left(\Lambda_{j}^{\prime}\setminus\Lambda_{j+1}\right)\lesssim e^{-j-10}.$$
(53)

This completes the proof.

Thus, the key is to obtain (25) and (26). We refer to T_N^{-1} as Green function.

4 Estimate of Green function

In this section, we estimate $T_{j,N}^{-1}$ and find Λ_j' step by step. Recall that

$$T_j = D + \varepsilon^{2/3} \Lambda S_j. \tag{54}$$

Denote

$$\tilde{T}_j = \Lambda^{-1} D + \varepsilon^{2/3} S_j.$$
(55)

We have $T_j = \Lambda \tilde{T}_j$. Thus, we have $T_{j,N}^{-1} = \tilde{T}_{j,N}^{-1} \Lambda_N^{-1}$. To obtain (25) and (26), it suffices to ensure that

$$\|\tilde{T}_{j,N}^{-1}\| < 2\exp(\log N)^{C_2},\tag{56}$$

$$|\tilde{T}_{j,N}^{-1}(\xi,\xi')| < 2e^{-\frac{1}{2}|\xi-\xi'|^c} \text{ for } |\xi-\xi'| > N^{\frac{1}{2}}.$$
(57)

The reason we introduce \tilde{T} is that it is self-adjoint. Denote $\tilde{D} = \Lambda^{-1}D$, $\tilde{T}_j = \tilde{D} + \varepsilon^{2/3}S_j$ and $N_j = M^j$. Furthermore, we denote

$$\tilde{D}_{\pm,n,k} = \langle n \rangle^{-\alpha} (\pm k\omega + |n|^2 + 1).$$
(58)

For $N_{j+1} \leq \varepsilon^{-\frac{1}{30d}}$, we construct Λ_j and Λ'_j $(j \geq j_0)$ directly. Note that Λ_{j_0} has been constructed in Section 2. Now we construct Λ'_{j_0} . Partition [1, 2] into intervals $\{I_{j_{0+1},\zeta}\}$ of size $\frac{1}{1-e^{-j_0-10}} \exp(-(j_0+1)^{C_3})$. Let

$$\Lambda'_{j_0} = \bigcup \{ I_{j_0+1,\zeta} : \exists \ \omega' \in I_{j_0+1,\zeta} \text{ such that } |-k\omega'+|n|^2+1 | \ge (\log \frac{1}{\varepsilon})^{-1}(1+|k|)^{-\tau}, \\ |n| < N_{j_0+1}, |k| < N_{j_0+1} \}.$$

Note that $\frac{2}{1-e^{-j_0-10}} \exp(-(j_0+1)^{C_3}) < \exp(-j_0^{C_3})$. It is easy to check that $\Lambda'_{j_0} \subset \Lambda_{j_0}$. For $\omega \in \Lambda'_{j_0}$, we have

$$\begin{aligned} |-k\omega + |n|^2 + 1| &\geq (\log \frac{1}{\varepsilon})^{-1} (1 + |k|)^{-\tau} - 2N_{j_0+1} \cdot \frac{1}{1 - e^{-j_0 - 10}} \exp(-(j_0 + 1)^{C_3}) \\ &> \frac{1}{2} (\log \frac{1}{\varepsilon})^{-1} (1 + |k|)^{-\tau}, \end{aligned}$$

for $|n| < N_{j_0+1}, |k| < N_{j_0+1}$. Thus, we have

$$|\tilde{D}_{\pm,n,k}| > \frac{1}{2} (\log \frac{1}{\varepsilon})^{-1} N_{j_0+1}^{-\tau-\alpha},$$
(59)

for $|\boldsymbol{n}| < N_{j_0+1}, |\boldsymbol{k}| < N_{j_0+1}.$ We obtain

$$\|\tilde{T}_{j_0,N_{j_0+1}}^{-1}\| = \|(I + \varepsilon^{2/3}\tilde{D}_{N_{j_0+1}}^{-1}S_{j_0,N_{j_0+1}})^{-1}\tilde{D}_{N_{j_0+1}}^{-1}\| \\ \leq 4(\log\frac{1}{\varepsilon})N_{j_0+1}^{\tau+\alpha} < \exp(\log N_{j_0+1})^{C_2}.$$
(60)

Furthermore, we have

$$\tilde{T}_{j_0,N_{j_0+1}}^{-1} = \tilde{D}_{N_{j_0+1}}^{-1} + \sum_{l=1}^{\infty} (-1)^l \varepsilon^{\frac{2l}{3}} (\tilde{D}_{N_{j_0+1}}^{-1} S_{j_0,N_{j_0+1}})^l \tilde{D}_{N_{j_0+1}}^{-1}.$$
(61)

Note that we have

$$\left| \left((\tilde{D}_{N_{j_0+1}}^{-1} S_{j_0,N_{j_0+1}})^l \tilde{D}_{N_{j_0+1}}^{-1} \right) (\xi,\xi') \right| \le \sum_{|\xi_1|,\dots,|\xi_{l-1}| \le N_{j_0+1}} N_{j_0+1}^{10dl} e^{-(1-)|\xi-\xi'|^c} \le N_{j_0+1}^{12dl} e^{-(1-)|\xi-\xi'|^c}.$$

$$(62)$$

Thus, we obtain

$$\begin{split} |\tilde{T}_{j_0,N_{j_0+1}}^{-1}(\xi,\xi')| &\leq \sum_{l=1}^{\infty} (-\varepsilon^{\frac{2}{3}} N_{j_0+1}^{12d})^l e^{-(1-)|\xi-\xi'|^c} \\ &\leq e^{-\frac{1}{2}|\xi-\xi'|^c}, \text{ for } \xi \neq \xi'. \end{split}$$
(63)

By Lemma 3.1, we obtain Λ_{j_0+1} . Proceed this process until $N_{j+1} > \varepsilon^{-\frac{1}{30d}}$. We obtain

$$\Lambda_j \subset \Lambda'_{j-1} \subset \Lambda_{j-1} \subset \cdots \subset \Lambda'_{j_0} \subset \Lambda_{j_0}.$$

Lemma 3.1 and a standard measure estimate as in Section 2 tell us

mes
$$[1,2] \setminus \Lambda_j \lesssim (\log \frac{1}{\varepsilon})^{-1} + e^{-j_0 - 10}.$$
 (64)

For $N_{j+1} > \varepsilon^{-\frac{1}{30d}}$, we have the following lemma.

Lemma 4.1. Let $N_{j+1} > \varepsilon^{-\frac{1}{30d}}$. Assume there exist $\Lambda_j \subset \Lambda_{j-1} \subset \cdots \subset \Lambda_{j_0} \subset [1,2]$ and functions $u_{j'}(x,\theta;\omega) \in L^2(\mathbb{T}^{d+1})(j_0 \leq j' \leq j)$ defined smoothly for $\omega \in [1,2]$ such that (j'-i), (j'-ii), (j'-iv) $(j_0 \leq j' \leq j)$ and (j'-v) $(j_0 + 1 \leq j' \leq j)$ hold. Moreover, assume we have

$$\|\tilde{T}_{j',N_{j'+1}}^{-1}\| < 2\exp(\log N_{j'+1})^{C_2},\tag{65}$$

$$|\tilde{T}_{j',N_{j'+1}}^{-1}(\xi,\xi')| < 2e^{-\frac{1}{2}|\xi-\xi'|^c} \text{ for } |\xi-\xi'| > N_{j'+1}^{\frac{1}{2}}.$$
(66)

for $\omega \in \Lambda_{j'+1}$ for $j_0 \leq j' \leq j-1$. Then, there exists a subset $\Lambda'_j \subset \Lambda_j$ which is a union of disjoint intervals of size $\frac{1}{1-e^{-j-10}} \exp(-(j+1)^{C_3})$ such that

$$\|\tilde{T}_{j,N_{j+1}}^{-1}\| < 2\exp(\log N_{j+1})^{C_2}, \\ |\tilde{T}_{j,N_{j+1}}^{-1}(\xi,\xi')| < 2e^{-\frac{1}{2}|\xi-\xi'|^c}, \text{ for } |\xi-\xi'| > N_{j+1}^{\frac{1}{2}}.$$
(67)

Moreover, we have

$$\max \Lambda_j \setminus \Lambda'_j \le N_{j+1}^{-10d}.$$
(68)

Proof. To control $\tilde{T}_{j,N_{j+1}}^{-1}$, we cover $Q = [-N_{j+1}, N_{j+1}]^{d+1}$ by intervals $Q_0 = [-N_{\lfloor \frac{j+1}{3} \rfloor}, N_{\lfloor \frac{j+1}{3} \rfloor}]$ and intervals Q_r in \mathbb{Z}^{d+1} of size $N_{j+1}^{1/3}$ such that dist $(0,Q_r) > N_{j+1}^{1/4}$. By the inductive hypothesis, we have

$$\tilde{T}_{\lfloor (j+1)/3 \rfloor - 1, N_{\lfloor (j+1)/3 \rfloor}}^{-1} \| < 2 \exp(\log N_{\lfloor (j+1)/3 \rfloor})^{C_2},$$
(69)

$$|\tilde{T}_{\lfloor (j+1)/3 \rfloor - 1, N_{\lfloor (j+1)/3 \rfloor}}^{-1}(\xi, \xi')| < 2e^{-\frac{1}{2}|\xi - \xi'|^c} \text{ for } |\xi - \xi'| > N_{\lfloor (j+1)/3 \rfloor}^{\frac{1}{2}}.$$
(70)

Since $||v_{j'}|| < e^{-\frac{11}{8}(M^{j'+1})^c}$, we have

$$\|(u_j - u_{\lfloor (j+1)/3 \rfloor - 1})(\xi)\| < e^{-(\frac{3}{8} -)(M^{\lfloor (j+1)/3 \rfloor})^c} e^{-|\xi|^c}.$$
(71)

Thus, we have

$$(\tilde{T}_j - \tilde{T}_{\lfloor (j+1)/3 \rfloor - 1})(\xi, \xi')| < e^{-\frac{1}{4}(M^{\lfloor (j+1)/3 \rfloor})^c} e^{-|\xi - \xi'|^c}.$$
(72)

Form (69), (70), (72) and $e^{-\frac{1}{4}(M^{\lfloor (j+1)/3 \rfloor})^c} N^C_{\lfloor (j+1)/3 \rfloor} \exp(\log N_{\lfloor (j+1)/3 \rfloor})^{C_2} \ll 1$, we obtain

$$\|\tilde{T}_{j,N_{\lfloor (j+1)/3 \rfloor}}^{-1}\| < 4 \exp(\log N_{\lfloor (j+1)/3 \rfloor})^{C_2},$$
(73)

$$|\tilde{T}_{j,N_{\lfloor (j+1)/3 \rfloor}}^{-1}(\xi,\xi')| < 4e^{-\frac{1}{2}|\xi-\xi'|^c} \text{ for } |\xi-\xi'| > N_{\lfloor (j+1)/3 \rfloor}^{\frac{1}{2}}.$$
(74)

To estimate \tilde{T}_{j,Q_r}^{-1} , we need the following lemma:

Lemma 4.2. Fix any large number B. There is a partition $\{\pi_{\zeta}\}$ of \mathbb{Z}^d satisfying the properties

diam $\pi_{\zeta} < B^{\tilde{C}_d}$

and

$$n - n'| + ||n|^2 - |n'|^2| > B$$
, if $n \in \pi_{\zeta}, n' \in \pi_{\zeta'}, \zeta \neq \zeta'$.

Here, $\tilde{C}_d = (2d+2)^{d+2}$.

In the sequel, we omit the subscript j of T_j and the subscript j+1 of N_{j+1} , i.e., we denote $T_j = T$ and $N_{j+1} = N$. Fix ω . Denote

$$\Omega_r = \{ (n,k) \in Q_r : \min_{\pm} |\tilde{D}_{\pm,n,k}| = \min_{\pm} |\langle n \rangle^{-\alpha} (\pm k\omega + |n|^2 + 1)| < 1 \}.$$
(75)

Choose a constant δ such that $\delta > \alpha$ and $(d\tilde{C}_d + 2)\delta < \frac{1}{30}$, which can be chosen since we assume $\alpha < \frac{1}{30(d(2d+2)^{d+2}+2)}$. Let $B = N^{\delta}$ in Lemma 4.2, and denote by $\{\pi_{\zeta}\}$ the partition of \mathbb{Z}^d . By Lemma 4.2, we have

$$\operatorname{diam} \pi_{\zeta} < N^{\delta C_d},\tag{76}$$

and

$$|n - n'| + ||n|^2 - |n'|^2| > N^{\delta}, \text{ if } n \in \pi_{\zeta}, n' \in \pi_{\zeta'}, \zeta \neq \zeta'.$$
(77)

Now we consider the structure of Ω_r . If $(n,k), (n',k') \in \Omega_r$, we have

$$||n|^2 - |k\omega|| < N^{\alpha} + 1 \text{ and } ||n'|^2 - |k'\omega|| < N^{\alpha} + 1.$$
 (78)

Thus, we have

$$||n|^{2} - |n'|^{2}| < 2|k - k'| + 2N^{\alpha} + 2.$$
(79)

If $n \in \pi_{\zeta}$, $n' \in \pi_{\zeta'}$, $\zeta \neq \zeta'$, then

$$2|k - k'| + |n - n'| + 2N^{\alpha} + 2 > |n - n'| + ||n|^2 - |n'|^2| > N^{\delta},$$
(80)

which implies

$$|k - k'| + |n - n'| \gtrsim N^{\delta}.$$
(81)

Fix |n| < N, the number of k such that $\min_{\pm} |\langle n \rangle^{-\alpha} (\pm k\omega + |n|^2 + 1)| < 1$ is at most $4N^{\alpha}$. Hence, we have

$$\#\{(n,k)\in\Omega_r:n\in\pi_{\zeta}\}\lesssim N^{\delta\tilde{C}_dd}\cdot N^{\alpha}.$$
(82)

The above argument allows us to obtain a partition of Ω_r :

$$\Omega_r = \bigcup_{\kappa} \Omega_{r,\kappa},\tag{83}$$

such that

diam
$$\Omega_{r,\kappa} \lesssim N^{\delta \tilde{C}_d d + \alpha + \delta} < N^{(\tilde{C}_d d + 2)\delta},$$
(84)

dist
$$(\Omega_{r,\kappa}, \Omega_{r,\kappa'}) \gtrsim N^{\delta}$$
. (85)

Let ω' in the $O(N^{-2})$ -neighborhood of ω . We have

$$\min_{\pm} |\langle n \rangle^{-\alpha} (\pm k\omega' + |n|^2 + 1)| > \frac{1}{2}, \text{ for } (n,k) \notin \Omega_r.$$
(86)

Hence, $\{\Omega_{r,\kappa}\}$ remains fixed for ω' in the $O(N^{-2})$ -neighborhood of ω . Denote this neighborhood by $I_{r,\kappa,s}$ ($s < N^2$, and the subscript s corresponds to the neighborhood of ω). Let $\tilde{\Omega}_{r,\kappa,s}$ be the $N^{\frac{\delta}{2}}$ -neighborhood of $\Omega_{r,\kappa,s}$. Note that $(d\tilde{C}_d + 2)\delta < \frac{1}{30}$. As long as we ensure that

$$\|\tilde{T}_{\tilde{\Omega}_{r,\kappa,s}}^{-1}\| < N^{C_1} \tag{87}$$

for all κ , applying Lemma A.2 gives us

$$\|\tilde{T}_{Q_r}^{-1}\| \lesssim N^{C_1+1},$$
(88)

$$|\tilde{T}_{Q_r}^{-1}(\xi,\xi')| < e^{-\frac{1}{10}|\xi-\xi'|^c}, \text{for } |\xi-\xi'| > N^{\frac{1}{5}}.$$
(89)

Let $\omega'_1 = \omega'^{-1}$ and write

$$\omega_1' \tilde{T}_{\tilde{\Omega}_{r,\kappa,s}} = \begin{pmatrix} \langle n \rangle^{-\alpha} (-k + \omega_1'(|n|^2 + 1)) & 0\\ 0 & \langle n \rangle^{-\alpha} (k + \omega_1'(|n|^2 + 1)) \end{pmatrix} + \varepsilon^{2/3} \omega_1' S_{\tilde{\Omega}_{r,\kappa}}.$$
(90)

Then, we have

$$\partial_{\omega_1'}(\omega_1'\tilde{T}_{\tilde{\Omega}_{r,\kappa,s}}) = \begin{pmatrix} (|n|^2+1)\langle n \rangle^{-\alpha} & 0\\ 0 & (|n|^2+1)\langle n \rangle^{-\alpha} \end{pmatrix} + O(\varepsilon^{2/3}).$$
(91)

Denote $E_{\tilde{\Omega}_{r,\kappa,s}}(\omega'_1)$ as an eigenvalue of $\omega'_1 \tilde{T}_{\tilde{\Omega}_{r,\kappa,s}}$. The first-order eigenvalue variation implies that

$$\left|\partial_{\omega_1'} E_{\tilde{\Omega}_{r,\kappa,s}}\right| \gtrsim 1. \tag{92}$$

Thus, there exists a subset $\tilde{I}_{r,\kappa,s}$ of $I_{r,\kappa,s}$ satisfying

- mes $\tilde{I}_{r,\kappa,s} \lesssim N^{1/2} N^{-C_1}$.
- For $\omega' \in I_{r,\kappa,s} \setminus \tilde{I}_{r,\kappa,s}$, we have

$$\|\tilde{T}_{\tilde{\Omega}_{r,\kappa,s}}^{-1}\| < N^{C_1}$$

Thus, for $\omega \in \Lambda_{j-1} \setminus \bigcup_{r,\kappa,s} \tilde{I}_{r,\kappa,s}$, we have

$$\|\tilde{T}_{\tilde{\Omega}_{r,\kappa}}^{-1}\| < N^{C_1}, \text{ for all } r, \kappa.$$
(93)

By applying Lemma A.2, we have (88) and (89) for all r. Furthermore, by applying Lemma A.1, we have

$$\|\tilde{T}_{j,N_{j+1}}^{-1}\| < \exp(\log N_{j+1})^{C_2},\tag{94}$$

$$|\tilde{T}_{j,N_{j+1}}^{-1}(\xi,\xi')| < e^{-\frac{1}{2}|\xi-\xi'|^c} \text{ for } |\xi-\xi'| > N_{j+1}^{\frac{1}{2}},$$
(95)

for $\omega \in \Lambda_{j-1} \setminus \bigcup_{r,\kappa,s} \tilde{I}_{r,\kappa,s}$. Now we construct Λ'_j . Partition Λ_j into intervals $\{I_{j,\zeta}\}$ of size $\frac{1}{1-e^{-j-10}} \exp(-(j+1)^{C_3})$. Let

$$\Lambda'_{j} = \bigcup \{ I_{j,\zeta} : \exists \ \omega \in I_{j,\zeta} \text{ such that } (94), (95) \}.$$
(96)

By a perturbation argument, we have

$$\|\tilde{T}_{j,N_{j+1}}^{-1}\| < 2\exp(\log N_{j+1})^{C_2},\tag{97}$$

$$|\tilde{T}_{j,N_{j+1}}^{-1}(\xi,\xi')| < 2e^{-\frac{1}{2}|\xi-\xi'|^c} \text{ for } |\xi-\xi'| > N_{j+1}^{\frac{1}{2}},$$
(98)

for $\omega \in \Lambda'_i$. Moreover, we have

$$\max \Lambda_j \setminus \Lambda'_j < \max \bigcup_{\substack{r,\kappa,s \\ r,\kappa,s}} \tilde{I}_{r,\kappa,s} < N_{j+1}^{2d} N_{j+1}^2 N_{j+1}^{-C_1 + \frac{1}{2}} < N_{j+1}^{-C_1 + 2d + 3} < N_{j+1}^{-10d},$$
(99)

where the second inequality follows from the fact that the number of both r and κ is less than N_j^d .

Theorem 1.1 follows from Lemma 2.3, Lemma 3.1 and Lemma 4.1.

A Coupling Lemma

In this appendix, we state two coupling lemmas whose proof can be found in [Bou98, Lemma 5.3, Lemma 7].

Lemma A.1. Assume T satisfies

$$|T(\xi,\xi')| < e^{-|\xi-\xi'|^c} \text{ for } \xi \neq \xi'.$$
 (100)

Let $\Omega \subset \mathbb{Z}^d$ be an interval and assume $\Omega = \bigcup_{\zeta} \Omega_{\zeta}$ a covering of Ω with intervals Ω_{ζ} satisfying

- $|T_{\Omega_{\zeta}}^{-1}(\xi,\xi')| < B.$
- $|T_{\Omega_{\zeta}}^{-1}(\xi,\xi')| < K^{-C} \text{ for } |\xi \xi'| > \frac{K}{100}.$
- For each $\xi \in \Omega$, there is ζ such that

$$B_K(\xi) \cap \Omega = \{\xi' \in \Omega : |\xi' - \xi| \le K\} \subset \Omega_{\zeta}.$$
(101)

• diam $\Omega_{\zeta} < C'K$ for each ζ .

Here, C > C(d) and B, K are numbers satisfying the relation

$$\log B < \frac{1}{100} K^c \text{ and } K > K_0(c, C', r).$$
 (102)

Then

$$T_{\Omega}^{-1}(\xi,\xi')| < 2B,$$
 (103)

$$|T_{\Omega}^{-1}(\xi,\xi')| < e^{-\frac{1}{2}|x-y|^c} \text{ for } |\xi-\xi'| > (100C'K)^{\frac{1}{1-c}}.$$
(104)

Lemma A.2. Fix some constants $\frac{1}{10} > \varepsilon_1 > \varepsilon_2 > \varepsilon_3 > 0$ and let Ω be a subset of the *M*-ball in \mathbb{Z}^{d+1} $(M \to \infty)$. Assume $\{\Omega_{\kappa}\}$ a collection of subsets of Ω satisfying

$$\operatorname{diam}\,\Omega_{\kappa} < M^{\varepsilon_1},\tag{105}$$

dist
$$(\Omega_{\kappa}, \Omega_{\kappa'}) > M^{\varepsilon_2}$$
 for $\kappa \neq \kappa'$. (106)

Write T = D + S (D is a diagonal matrix) where

$$||S|| < \varepsilon, \quad |S(\xi,\xi')| < \varepsilon e^{-|\xi-\xi'|^c}, \tag{107}$$

where c is sufficiently small and

$$|D(\xi)| > \rho \gg \varepsilon \quad \text{if} \quad \xi \in \Omega \setminus \bigcup \Omega_{\kappa}, \tag{108}$$

$$\|(T|_{\tilde{\Omega}_{\kappa}})^{-1}\| < M^C \text{ for all } \kappa, \tag{109}$$

where $\tilde{\Omega}_{\kappa}$ is an M^{ε_3} -neighborhood of Ω_{κ} . Then

$$||(T|\Omega)^{-1}|| < \rho^{-1} M^{C+1}, \tag{110}$$

and

$$(T|\Omega)^{-1}(\xi,\xi')| < e^{-\frac{1}{10}|\xi-\xi'|^c} \text{ if } |\xi-\xi'| > M^{2\varepsilon_1}.$$
(111)

References

- [BB13] Massimiliano Berti and Philippe Bolle. Quasi-periodic solutions with Sobolev regularity of NLS on \mathbb{T}^d with a multiplicative potential. J. Eur. Math. Soc. (JEMS), 15(1):229–286, 2013.
- [BB20] Massimiliano Berti and Philippe Bolle. Quasi-periodic solutions of nonlinear wave equations on the d-dimensional torus. EMS Monographs in Mathematics. EMS Publishing House, Berlin, [2020] ©2020.
- [BBHM18] Pietro Baldi, Massimiliano Berti, Emanuele Haus, and Riccardo Montalto. Time quasi-periodic gravity water waves in finite depth. *Invent. Math.*, 214(2):739–911, 2018.
- [BBM14] Pietro Baldi, Massimiliano Berti, and Riccardo Montalto. KAM for quasi-linear and fully nonlinear forced perturbations of Airy equation. *Math. Ann.*, 359(1-2):471–536, 2014.

- [BBM16] Pietro Baldi, Massimiliano Berti, and Riccardo Montalto. KAM for autonomous quasi-linear perturbations of KdV. Ann. Inst. H. Poincaré C Anal. Non Linéaire, 33(6):1589–1638, 2016.
- [BBP13] Massimiliano Berti, Luca Biasco, and Michela Procesi. KAM theory for the Hamiltonian derivative wave equation. Ann. Sci. Éc. Norm. Supér. (4), 46(2):301–373, 2013.
- [Bou94] Jean Bourgain. Construction of quasi-periodic solutions for Hamiltonian perturbations of linear equations and applications to nonlinear PDE. Internat. Math. Res. Notices, (11), 1994.
- [Bou95] J. Bourgain. Construction of periodic solutions of nonlinear wave equations in higher dimension. *Geom. Funct. Anal.*, 5(4):629–639, 1995.
- [Bou98] J. Bourgain. Quasi-periodic solutions of Hamiltonian perturbations of 2D linear Schrödinger equations. Ann. of Math. (2), 148(2):363–439, 1998.
- [Bou05] Jean Bourgain. Green's function estimates for lattice Schrödinger operators and applications. Princeton University Press, Princeton, NJ, 2005.
- [BP11] Massimiliano Berti and Michela Procesi. Nonlinear wave and Schrödinger equations on compact Lie groups and homogeneous spaces. *Duke Math. J.*, 159(3):479– 538, 2011.
- [CW93] Walter Craig and C. Eugene Wayne. Newton's method and periodic solutions of nonlinear wave equations. *Comm. Pure Appl. Math.*, 46(11):1409–1498, 1993.
- [CY00] Luigi Chierchia and Jiangong You. KAM tori for 1D nonlinear wave equations with periodic boundary conditions. *Comm. Math. Phys.*, 211(2):497–525, 2000.
- [EK10] L. Hakan Eliasson and Sergei B. Kuksin. KAM for the nonlinear Schrödinger equation. Ann. of Math. (2), 172(1):371–435, 2010.
- [Kuk87] S. B. Kuksin. Hamiltonian perturbations of infinite-dimensional linear systems with imaginary spectrum. *Funktsional. Anal. i Prilozhen.*, 21(3):22–37, 95, 1987.
- [Kuk93] Sergej B. Kuksin. Nearly integrable infinite-dimensional Hamiltonian systems, volume 1556 of Lecture Notes in Mathematics. Springer-Verlag, Berlin, 1993.
- [Kuk98] Sergei B. Kuksin. A KAM-theorem for equations of the Korteweg-de Vries type. *Rev. Math. Math. Phys.*, 10(3):ii+64, 1998.
- [Kuk00] Sergei B. Kuksin. Analysis of Hamiltonian PDEs, volume 19 of Oxford Lecture Series in Mathematics and its Applications. Oxford University Press, Oxford, 2000.
- [LY10] Jianjun Liu and Xiaoping Yuan. Spectrum for quantum Duffing oscillator and small-divisor equation with large-variable coefficient. Comm. Pure Appl. Math., 63(9):1145–1172, 2010.
- [LY11] Jianjun Liu and Xiaoping Yuan. A KAM theorem for Hamiltonian partial differential equations with unbounded perturbations. *Comm. Math. Phys.*, 307(3):629– 673, 2011.

- [P96] Jürgen Pöschel. A KAM-theorem for some nonlinear partial differential equations. Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4), 23(1):119–148, 1996.
- [Wan16] W.-M. Wang. Energy supercritical nonlinear Schrödinger equations: quasiperiodic solutions. *Duke Math. J.*, 165(6):1129–1192, 2016.
- [Way90] C. Eugene Wayne. Periodic and quasi-periodic solutions of nonlinear wave equations via KAM theory. Comm. Math. Phys., 127(3):479–528, 1990.
- [Yua21] Xiaoping Yuan. KAM theorem with normal frequencies of finite limit points for some shallow water equations. *Comm. Pure Appl. Math.*, 74(6):1193–1281, 2021.