

# Objective Mackey and Tambara functors via parametrized categories

Ross Street

March 5, 2025

*2020 Mathematics Subject Classification:* 18M10; 18N10; 20J99; 20C05; 18M05; 18N25

*Key words and phrases:* finite group; Mackey functor; categorification; parametrized category; variable category; cocompletion; Tambara functor; pointwise extension; spans; monoidale

## Contents

<b>1</b>	<b>Parametrized categories and protocolibrations</b>	<b>3</b>
<b>2</b>	<b><math>\mathcal{R}</math>-cocompletion</b>	<b>7</b>
<b>3</b>	<b><math>\mathcal{R}</math>-Spans and <math>\mathcal{R}</math>-cocompleteness</b>	<b>11</b>
<b>4</b>	<b>A distributive law from compatibility</b>	<b>14</b>
<b>5</b>	<b>Polynomials as spans of spans</b>	<b>17</b>
<b>6</b>	<b>Finite coproducts for spans and polynomials</b>	<b>19</b>
<b>7</b>	<b>Objective Mackey functors</b>	<b>23</b>
<b>8</b>	<b>A monoidal structure on <math>\text{OMky}_{\mathcal{R}}\mathcal{E}</math></b>	<b>25</b>
<b>9</b>	<b>An objective Mackey functor of Mackey functors</b>	<b>27</b>
<b>10</b>	<b>Objective Tambara functors</b>	<b>28</b>

## Abstract

The first word in the title is intended in a sense suggested by Lawvere and Schanuel whereby finite sets are objective natural numbers. At the objective level, the axioms defining abstract Mackey and Tambara functors are categorically familiar. The first step was taken by Harald Lindner in 1976 when he recognized that Mackey functors, defined as pairs of functors, were single functors with domain a category of spans. We define objective Mackey and objective Tambara functors as parametrized categories which have local finite products and satisfy some parametrized completeness and cocompleteness restriction. However, we can replace the original parametrizing base for objective Mackey functors by a bicategory of spans while the replacement for objective Tambara functors is a bicategory obtained by iterating the span construction; these iterated spans are polynomials. There is an objective Mackey functor of ordinary Mackey functors. We show that there is a distributive law relating objective Mackey functors to objective Tambara functors analogous to the distributive law relating abelian groups to commutative rings. We remark on hom enrichment matters involving the 2-category  $\text{Cat}_+$  of categories admitting finite coproducts and functors preserving them, both as a closed base and as a skew-closed base.

## Introduction

The sense in which we use the terms “abstract” and “objective” is that of Lawvere and Schanuel (for example, see [30]). Coproduct and product of finite sets have the virtue, over abstract addition and multiplication of natural numbers, of universal properties which can help in understanding of combinatorial equations as bijections.

The process of moving from objective to abstract is very old. Like quantization in Physics, the reverse process is also rather old, yet it is not well defined even up to any reasonable notion of equivalence. The term “categorification” was coined by Crane–Frenkel in low-dimensional topology (see [11]) and then adopted by representation theorists.

The Introduction of David Chan’s paper [9] provides good background to the roles of Mackey and Tambara functors in equivariant stable homotopy theory. In particular, the third paragraph begins with the following sentence: *While Mackey functors should be thought of as the correct analogue of an abelian group in the equivariant setting, the role of commutative rings is played by objects known as Tambara functors.*

Motivated by terminology and constructions in [19, 5, 6, 9], my goal was to investigate in what way Tambara functors on a lextensive category  $\mathcal{E}$  are Mackey functors with extra structure. I soon moved to the objective level and found what I consider is a satisfactory confirmation of the analogy between Mackey functors and abelian groups, and commutative rings and Tambara functors.

The main idea is that the basic construction is a variant of Bénabou’s bicategory  $\text{Spn}\mathcal{E}$  of spans  $U \xleftarrow{u} S \xrightarrow{v} V$  in a category  $\mathcal{E}$  with pullbacks (see [2]). We allow  $\mathcal{E}$  to be replaced by a bicategory  $\mathcal{C}$  equipped with a chosen wide (= all object inclusive), locally full subbicategory  $\mathcal{R}$  with a few conditions making it what we here call a *protocalibration*. For our bicategory  $\text{Spn}_{\mathcal{R}}\mathcal{C}$ , we restrict our 1-morphisms to spans for which the left leg  $u$  is in  $\mathcal{R}$  yet we allow our 2-morphisms to be isomorphism classes of span morphisms which are pseudo on the left and lax on the right. Its universal property is that categories parametrized by  $\text{Spn}_{\mathcal{R}}\mathcal{C}$  are essentially categories parametrized by  $\mathcal{C}$  which are  $\mathcal{R}$ -cocomplete. This last term refers to the existence of pointwise left extensions along  $\mathcal{C}$ -parametrized functors of the form  $\mathcal{C}(-, U) \xrightarrow{\mathcal{C}(-, r)} \mathcal{C}(-, V)$  with  $r \in \mathcal{R}$ .

Other constructions are obtained by duality and iteration. If  $\mathcal{L}$  is a protocalibration of  $\mathcal{C}^{\text{co}}$  then categories parametrized by  ${}_{\mathcal{L}}\text{Spn}\mathcal{C} = (\text{Spn}_{\mathcal{L}}\mathcal{C}^{\text{co}})^{\text{co}}$  are essentially categories parametrized by  $\mathcal{C}$  which are  $\mathcal{L}$ -complete. Moreover, with the specific iteration  $\text{Spn}_{\mathcal{E}_*}({}_{\mathcal{L}}\text{Spn}\mathcal{E})$  we obtain the opposite of the bicategory  $\text{Ply}\mathcal{E}$  of polynomials in a category  $\mathcal{E}$  with pullbacks where  $\mathcal{P}$  consists of the powerful morphisms in  $\mathcal{E}$  and  $\mathcal{E}_*$  consists of spans with the left leg  $u$  invertible. The universal property (in the case where  $\mathcal{E}$  is locally cartesian closed) obtained by iteration agrees with that proved by Charles Walker [44].

## 1 Parametrized categories and protocalibrations

Categories varying over [36] (fibred over [4], indexed by [24], or parametrized by [31]) a category or bicategory  $\mathcal{C}$  can be regarded as pseudofunctors  $\mathbb{X} : \mathcal{C}^{\text{op}} \rightarrow \text{Cat}$ . We think of an object of  $\mathbb{X}U$  as a *generalised object* of the  $\mathcal{C}$ -variable category  $\mathbb{X}$ . We write

$$\text{CAT}\mathcal{C} = \text{Ps}(\mathcal{C}^{\text{op}}, \text{Cat})$$

for the (strict) bicategory of pseudofunctors (called “homomorphisms” in [2]), pseudonatural transformations, and modifications.

A  $\mathcal{C}$ -variable category  $\mathbb{X} \in \text{CAT}\mathcal{C}$  is *cocomplete with respect to a morphism*  $j : \mathbb{A} \rightarrow \mathbb{B}$  in  $\text{CAT}\mathcal{C}$  (see Section 7 of [36]) when every morphism

$f : \mathbb{A} \rightarrow \mathbb{X}$  has a pointwise (see Section 4 of [33]) left extension along  $j$ . Generally, pointwiseness translates to a lax form of the Chevalley-Beck (CB) condition. However, the pseudo form is relevant when dealing with groupoid fibrations as we will now remind the reader.

A morphism  $\kappa : z \rightarrow x$  in a category  $E$  is *cartesian* for a functor  $p : E \rightarrow B$  when the following square (1.1) is a pullback in  $\text{Set}$  for all objects  $k$  of  $E$ .

$$\begin{array}{ccc} E(k, z) & \xrightarrow{E(k, \kappa)} & E(k, x) \\ p \downarrow & & \downarrow p \\ B(pk, pz) & \xrightarrow{B(pk, p\kappa)} & B(pk, px) \end{array} \quad (1.1)$$

A functor  $p : E \rightarrow B$  is a *groupoid fibration (over  $B$ )* when every morphism of  $E$  is cartesian for  $p$  and, for all  $x \in E$  and  $\phi : b \rightarrow px$  in  $B$ , there exists a morphism  $\kappa : z \rightarrow x$  in  $E$  and an isomorphism  $b \cong pz$  whose composite with  $p\kappa$  is  $\phi$ . A functor  $p : E \rightarrow B$  is a *groupoid opfibration (over  $B$ )* when  $p : E^{\text{op}} \rightarrow B^{\text{op}}$  is a groupoid fibration.

A morphism  $p : E \rightarrow B$  in a bicategory  $\mathcal{C}$  is a *groupoid fibration* when, for all objects  $U \in \mathcal{C}$ , the functor  $\mathcal{C}(U, p) : \mathcal{C}(U, E) \rightarrow \mathcal{C}(U, B)$  is a groupoid fibration. The proof of Proposition 1.1 is essentially as for Proposition 23 of [33]; alternatively, check it in  $\text{Cat}$  and use Yoneda since all concepts are representable.

**Proposition 1.1.** *In any finitely complete bicategory, suppose  $p : E \rightarrow B$  is a groupoid opfibration. In diagram (1.2), the 2-morphism  $\sigma$  exhibits  $k$  as a pointwise left extension of  $f$  along  $p$  if and only if, for all morphisms  $b$  with the square a bicategorical pullback (= briefly, bipullback), the pasted composite at  $p$  exhibits  $kb$  as a left extension of  $f\tilde{b}$  along  $\tilde{p}$ .*

$$\begin{array}{ccc} E & \xrightarrow{p} & B \\ & \searrow f & \swarrow k \\ & X & \end{array} \quad \begin{array}{ccc} P & \xrightarrow{\tilde{p}} & V \\ \tilde{b} \downarrow \lrcorner & \xrightarrow{\cong} & \downarrow b \\ E & \xrightarrow{p} & B \end{array} \quad (1.2)$$

If  $\mathcal{R}$  is a set of morphisms of a bicategory  $\mathcal{C}$ , we say  $\mathbb{X}$  is  $\mathcal{R}$ -cocomplete when it is cocomplete with respect to all morphisms  $\mathcal{C}(-, j) : \mathcal{C}(-, U) \rightarrow \mathcal{C}(-, V)$  for  $j : U \rightarrow V$  in  $\mathcal{R}$ . A pseudonatural transformation  $\theta : \mathbb{X} \rightarrow \mathbb{Y}$  is  $\mathcal{R}$ -cocontinuous when it preserves left extensions along the  $\mathcal{C}(-, j)$  for  $j \in \mathcal{R}$ .

Let  $\text{CAT}_{\mathcal{R}}\mathcal{C}$  denote the subcategory of  $\text{CAT}\mathcal{C}$  consisting of the  $\mathcal{R}$ -cocomplete objects and  $\mathcal{R}$ -cocontinuous morphisms.

We are interested in those  $\mathcal{R}$  in possession of most of the properties in common with the “calibrations” of [40], and the calibrations of Bénabou [3] when  $\mathcal{C}$  is a category. In the case where  $\mathcal{C}$  is a category of  $G$ -sets, these conditions plus closure under coproducts (which we will also require in Section 6) are those called “indexing categories” in equivariant stable homotopy theory; see Chan [9].

**Definition 1.2.** *A protocalibration of a bicategory  $\mathcal{C}$  is a set  $\mathcal{R}$  of morphisms when it satisfies the four conditions:*

- I. all equivalences are in  $\mathcal{R}$  and any morphism isomorphic to one in  $\mathcal{R}$  is in  $\mathcal{R}$ ;*
- C. composites of morphisms in  $\mathcal{R}$  are in  $\mathcal{R}$ ;*
- P. for all  $U \xrightarrow{f} W$  in  $\mathcal{R}$  and all  $V \xrightarrow{g} W$  in  $\mathcal{C}$ , the bicategorical pullback  $P \xrightarrow{\bar{f}} V$  of  $f$  along  $g$  exists and is in  $\mathcal{R}$ ;*
- G. every morphism in  $\mathcal{R}$  is a groupoid opfibration.*

*Each protocalibration  $\mathcal{R}$  will be identified with the subcategory of  $\mathcal{C}$  containing all objects, only the morphisms in  $\mathcal{R}$ , and all 2-morphisms between them. Notice that a protocalibration of  $\mathcal{C}^{\text{co}}$  is the same as one for  $\mathcal{C}$  except for condition (G) where “groupoid opfibration” must be replaced by “groupoid fibration”.*

**Example 1.3.** *1. Suppose  $\mathcal{C}$  is just a category (locally discrete as a bicategory). Then every morphism is both a groupoid fibration and a groupoid opfibration, and bicategorical pullbacks are pullbacks. If  $\mathcal{C}$  has pullbacks, the set of all morphisms is a protocalibration.*

*2. Again suppose  $\mathcal{C}$  is a category. If pullbacks of powerful (= exponentiable) morphisms exist then the powerful morphisms form a protocalibration (see Corollary 2.6 of [41]). This applies to the category  $\mathcal{C} = \text{Cat}$  of categories and functors where the powerful morphisms are the Giraud-Conduché functors [10].*

*3. Let  $\mathcal{E}$  be a category with pullbacks and every morphism powerful. The groupoid fibrations (= left-adjoint morphisms) in the bicategory  $\text{Spn}\mathcal{E}$  of spans in  $\mathcal{E}$  is a protocalibration of  $(\text{Spn}\mathcal{E})^{\text{co}}$  (see Propositions 4.1 and 8.3 of [40]).*

**Proposition 1.4** (Compare [34] Proposition 9.12, [36] Proposition 7.12). *Let  $\mathcal{R}$  be a protocalibration of the bicategory  $\mathcal{C}$ . A  $\mathcal{C}$ -variable category  $\mathbb{X}$  is  $\mathcal{R}$ -cocomplete if and only if, for all  $f \in \mathcal{R}$ , the functor  $\mathbb{X}f$  has a left adjoint  $\mathbb{X}_*f$  and, for all bicategorical pullback squares*

$$\begin{array}{ccc} P & \xrightarrow{f_g} & V \\ g_f \downarrow \lrcorner & \xrightarrow{\cong} & \downarrow g \\ U & \xrightarrow{f} & W \end{array} \quad (1.3)$$

in  $\mathcal{C}$ , the mate

$$\mathbb{X}_*f_g \circ \mathbb{X}g_f \Rightarrow \mathbb{X}g \circ \mathbb{X}_*f$$

of the isomorphism

$$\mathbb{X}g_f \circ \mathbb{X}f \cong \mathbb{X}f_g \circ \mathbb{X}g$$

is invertible. A pseudonatural transformation  $\theta : \mathbb{X} \rightarrow \mathbb{Y}$  is  $\mathcal{R}$ -cocontinuous if and only if, for all  $U \xrightarrow{f} W$  in  $\mathcal{R}$ , the mate  $\theta_f : \mathbb{Y}_*f \circ \theta_U \rightarrow \theta_W \circ \mathbb{X}_*f$  of  $\theta_f : \theta_U \circ \mathbb{X}f \rightarrow \mathbb{Y}f \circ \theta_W$  is invertible.

**Remark 1.5.** Henceforth we occasionally allow ourselves to omit the invertible 2-morphisms in bipullbacks such as (1.3).

**Lemma 1.6.** *Suppose  $\mathcal{C}$  has binary bicategorical products<sup>1</sup> and  $\mathcal{R}$  is a protocalibration of  $\mathcal{C}$ . If  $X \xrightarrow{f} Y$  and  $X' \xrightarrow{f'} Y'$  are in  $\mathcal{R}$  then so too is  $X \times X' \xrightarrow{f \times f'} Y \times Y'$ . So, if  $\mathcal{C}$  has finite bicategorical products then  $\mathcal{R}$  is a monoidal subcategory of the cartesian monoidal bicategory  $\mathcal{C}$ .*

*Proof.* The morphism  $X \times X' \xrightarrow{f \times f'} Y \times Y'$  is the composite of  $f \times 1_{X'}$  and  $1_Y \times f'$  which are respectively bipullbacks of  $f$  and  $f'$  along projections.  $\square$

**Corollary 1.7.** *Suppose  $\mathcal{C}$  has binary biproducts and  $\mathcal{R}$  is a protocalibration of  $\mathcal{C}$ . If  $\mathbb{Z}$  is  $\mathcal{R}$ -cocomplete then so is  $\mathbb{Z}(- \times W)$  for all  $W \in \mathcal{C}$ .*

*Proof.* Cocompleteness of  $\mathbb{Z}(- \times W)$  with respect to any  $f$  amounts, by Yoneda, to cocompleteness of  $\mathbb{Z}$  with respect to  $f \times 1_W$ .  $\square$

**Proposition 1.8.** *Suppose  $\mathcal{C}$  has binary biproducts and  $\mathcal{R}$  is a protocalibration of  $\mathcal{C}$ . For  $\mathbb{Y}$  and  $\mathbb{Z}$  in  $\text{CAT}_{\mathcal{R}}\mathcal{C}$ , the object  $[\mathbb{Y}, \mathbb{Z}]_{\mathcal{R}} \in \text{CAT}\mathcal{C}$ , defined on objects by*

$$[\mathbb{Y}, \mathbb{Z}]_{\mathcal{R}}W = \text{CAT}_{\mathcal{R}}\mathcal{C}(\mathbb{Y}, \mathbb{Z}(- \times W))$$

and on morphisms  $W' \xrightarrow{h} W$  by composition with  $\mathbb{Z}(- \times h)$ , is in  $\text{CAT}_{\mathcal{R}}\mathcal{C}$ .

<sup>1</sup>We will abbreviate this to “biproduct” and use “direct sum” when product and co-product coincide.

*Proof.* By the last sentence of Proposition 1.4, we need to see that, if  $\theta : \mathbb{Y} \rightarrow \mathbb{Z}(- \times W)$  is  $\mathcal{R}$ -cocontinuous, so too is  $\mathbb{Z}(- \times h) \circ \theta$ . Take  $V \xrightarrow{r} U$  in  $\mathcal{R}$ . The right square in the diagram

$$\begin{array}{ccccc}
\mathbb{Y}U & \xrightarrow{\theta_U} & \mathbb{Z}(U \times W) & \xrightarrow{\mathbb{Z}(1_U \times h)} & \mathbb{Z}(U \times W') \\
\mathbb{Y}r \downarrow & \xrightarrow[\cong]{\theta_r} & \downarrow \mathbb{Z}(r \times 1_W) & \xrightarrow[\cong]{} & \downarrow \mathbb{Z}(r \times 1_{W'}) \\
\mathbb{Y}V & \xrightarrow{\theta_V} & \mathbb{Z}(V \times W) & \xrightarrow{\mathbb{Z}(1_V \times h)} & \mathbb{Z}(V \times W')
\end{array}$$

is the value of  $\mathbb{Z}$  on a bipullback in  $\mathcal{C}$ . The invertibility of the mate of the pasted 2-morphism follows from  $\mathcal{R}$ -cocontinuity of  $\theta$  and  $\mathcal{R}$ -cocompleteness of  $\mathbb{Z}$ .  $\square$

## 2 $\mathcal{R}$ -cocompletion

Suppose  $\mathcal{R}$  is a protocalibration of the bicategory  $\mathcal{C}$ . The  $\mathcal{R}$ -cocompletion  $\overrightarrow{\mathbb{X}}$  of  $\mathbb{X} \in \text{CAT}\mathcal{C}$  is described as follows (see [32, 4, 36]). An object of the category  $\overrightarrow{\mathbb{X}}U$  is a pair  $(U \xleftarrow{u} S, x \in \mathbb{X}S)$  with  $u \in \mathcal{R}$ ; we might think of such an object as a  $U$ -indexed family of generalised objects of  $\mathbb{X}$ . A morphism

$$(U \xleftarrow{u} S, x \in \mathbb{X}S) \rightarrow (U \xleftarrow{u'} S', x' \in \mathbb{X}S')$$

of  $\overrightarrow{\mathbb{X}}U$  is an isomorphism class of triples  $(S \xrightarrow{w} S', u'w \cong u, x \xrightarrow{\xi} (\mathbb{X}w)x')$  where  $\xi$  is a morphism of  $\mathbb{X}S$ ; if we identify representables with their representing objects, the picture of such a morphism is a lax morphism (2.4) of spans in  $\text{CAT}\mathcal{C}$ .

$$\begin{array}{ccccc}
& & S & & \\
& u \in \mathcal{R} \nearrow & \downarrow w & \xleftarrow{\xi} & x \\
U & \xleftarrow{u' \in \mathcal{R}} & S' & \xrightarrow{x'} & \mathbb{X}
\end{array} \tag{2.4}$$

For  $r : V \rightarrow U$  in  $\mathcal{C}$ , define the functor  $\overrightarrow{\mathbb{X}}r : \overrightarrow{\mathbb{X}}U \rightarrow \overrightarrow{\mathbb{X}}V$  to take  $(u, x)$  to  $(u_r, (\mathbb{X}r_u)x)$  where

$$\begin{array}{ccc}
P \xrightarrow{u_r} V & & P' \xrightarrow{u'_r} V \\
r_u \downarrow \lrcorner & & r_{u'} \downarrow \lrcorner \\
S \xrightarrow{u} U & & S' \xrightarrow{u'} U
\end{array} \tag{2.5}$$

are bipullbacks, and  $(w, \xi)$  to  $(\bar{w}, \bar{\xi})$  where  $\bar{w}$  is defined by compatible  $u'_r \bar{w} \cong u_r, r_u \bar{w} \cong w r_u$ , and  $\bar{\xi}$  by

$$\bar{\xi} := \left( (\mathbb{X}r_u)x \xrightarrow{(\mathbb{X}r_u)\xi} (\mathbb{X}r_u)(\mathbb{X}w)x' \cong (\mathbb{X}\bar{w})(\mathbb{X}r_u)x' \right) .$$

For a 2-morphism  $\alpha : r \Rightarrow s : V \rightarrow U$ , let  $S \xleftarrow{s_u} Q \xrightarrow{u_s} V$  be the bipullback of the cospan  $S \xrightarrow{u} U \xleftarrow{s} V$ . Since  $u$  is a groupoid opfibration, there exists a cocartesian 2-morphism  $\kappa : r_u \Rightarrow k$  and isomorphism  $su_r \cong uk$  such that  $u\kappa$  is the composite  $ur_u \cong ru_r \xrightarrow{\alpha u_r} su_r \cong uk$ . If  $r \in \mathcal{R}$  then  $\overrightarrow{\mathbb{X}}r$  has a left adjoint  $\overrightarrow{\mathbb{X}}_*r : \overrightarrow{\mathbb{X}}V \rightarrow \overrightarrow{\mathbb{X}}U$  taking  $(V \xleftarrow{v} T, y \in \mathbb{X}T)$  to  $(U \xleftarrow{rv} T, y \in \mathbb{X}T)$  (noting that  $rv \in \mathcal{R}$  by condition (C)). To verify the pseudo-Chevalley-Beck condition, we refer to the bipullbacks (2.6) with  $f, u \in \mathcal{R}$ .

$$\begin{array}{ccccc} Q & \xrightarrow{u_{gf}} & P & \xrightarrow{f_g} & V \\ g_{fu} \downarrow \lrcorner & & g_f \downarrow \lrcorner & & \downarrow g \\ S & \xrightarrow{u \in \mathcal{R}} & U & \xrightarrow{f \in \mathcal{R}} & W \end{array} \quad (2.6)$$

Then we see that

$$\begin{aligned} (\overrightarrow{\mathbb{X}}g \circ \overrightarrow{\mathbb{X}}_*f)(u, x) &= (\overrightarrow{\mathbb{X}}g)(fu, x) \cong (f_g u_{g_f}, (\mathbb{X}g_{fu})x) \\ &= (\overrightarrow{\mathbb{X}}_*f_g)(u_{g_f}, (\mathbb{X}g_{fu})x) \\ &\cong (\overrightarrow{\mathbb{X}}_*f_g \circ \mathbb{X}g_f)(u, x) \end{aligned}$$

naturally in  $(u, x) \in \overrightarrow{\mathbb{X}}U$ . So  $\overrightarrow{\mathbb{X}} \in \text{CAT}^{\mathcal{E}}$  is  $\mathcal{R}$ -cocomplete by Proposition 1.4.

**Remark 2.1.** Suppose  $\mathcal{C}$  is a category with pullbacks so that we can take  $\mathcal{R}$  to contain all morphisms. The terminal object  $\mathbf{1}$  of  $\text{CAT}^{\mathcal{C}}$  takes all objects of  $\mathcal{C}$  to the terminal category. Then  $\mathbb{C} := \overrightarrow{\mathbf{1}}$  takes  $U \in \mathcal{C}$  to the slice category  $\mathcal{C}_{/U}$  and is defined on morphisms using pullback. Notice that an ordinary category has coproducts if and only if all pointwise left Kan extensions into it exist along functors between discrete categories. Also  $\text{Set}$  is the coproduct cocompletion of the terminal ordinary category. Now we see that  $\mathbb{C}$  is the coproduct cocompletion of the terminal  $\mathcal{C}$ -variable category. This is one justification for viewing  $\mathbb{C}$  as playing the role in  $\mathcal{C}$ -variable category theory of  $\text{Set}$  in ordinary category theory.

**Definition 2.2.** For a protocolibration  $\mathcal{R}$  of a bicategory  $\mathcal{C}$ , we write  $\mathbb{C}_{\mathcal{R}}$  for the  $\mathcal{R}$ -cocompletion  $\overrightarrow{\mathbf{1}}$  of  $\mathbf{1}$ .



Directly from the explicit descriptions, we can verify the following.

**Proposition 2.3.** *For a protocalibration  $\mathcal{R}$  of a bicategory  $\mathcal{C}$  and a pseudofunctor  $\mathbb{X} \in \text{CAT}\mathcal{C}$ , the unique  $! : \mathbb{X} \rightarrow \mathbf{1}$  induces a pseudonatural transformation  $\mathbb{X}! : \overrightarrow{\mathbb{X}} \rightarrow \mathbb{C}_{\mathcal{R}}$  which is a componentwise fibration. If  $\mathcal{C}$  is a finitely complete category and each  $! : S \rightarrow \mathbf{1}$  is in  $\mathcal{R}$  then  $\mathbb{X}!_1 : \overrightarrow{\mathbb{X}}\mathbf{1} \rightarrow \mathbb{C}_{\mathcal{R}}\mathbf{1}$  is the fibration over  $\mathcal{C}$  constructed by Grothendieck from the pseudofunctor  $\mathbb{X} : \mathcal{C}^{\text{op}} \rightarrow \text{Cat}$ .*

There is a pseudonatural transformation  $\eta_{\mathbb{X}} : \mathbb{X} \rightarrow \overrightarrow{\mathbb{X}}$  whose component  $\eta_{\mathbb{X}U} : \mathbb{X}U \rightarrow \overrightarrow{\mathbb{X}}U$  at  $U \in \mathcal{C}$  is defined by  $\eta_{\mathbb{X}U}x = (U \xleftarrow{1_U} U, x \in \mathbb{X}U)$ .

**Proposition 2.4.** *Suppose  $\mathcal{R}$  is a protocalibration of the bicategory  $\mathcal{C}$ . Then  $\mathbb{X} \in \text{CAT}\mathcal{C}$  is  $\mathcal{R}$ -cocomplete if and only if  $\eta_{\mathbb{X}}$  has a left adjoint in  $\text{CAT}\mathcal{C}$ .*

*Proof.* A left adjoint to  $\eta_{\mathbb{X}U} : \mathbb{X}U \rightarrow \overrightarrow{\mathbb{X}}U$  precisely gives, for each  $(u, x) \in \overrightarrow{\mathbb{X}}U$ , an object  $u_*(x) \in \mathbb{X}U$  and an isomorphism  $\mathbb{X}U(u_*(x), y) \cong \overrightarrow{\mathbb{X}}U((u, x), (1_U, y))$  natural in  $y \in \mathbb{X}U$ . However,  $\overrightarrow{\mathbb{X}}U((u, x), (1_U, y)) \cong \mathbb{X}S(x, (\mathbb{X}u)y)$ . So this amounts precisely to having a left adjoint  $\mathbb{X}_*u$  to  $\mathbb{X}u$  for all  $u \in \mathcal{R}$ . Pseudonaturality of  $\eta_{\mathbb{X}}$  amounts to the  $\mathcal{R}$ -Chevalley-Beck condition for  $\mathbb{X}$ . The result follows now from Proposition 1.4.  $\square$

We think of the left adjoint to  $\eta_{\mathbb{X}}$  as assigning a coproduct to the indexed families of generalised objects.

**Corollary 2.5.** *The pseudonatural transformation  $\eta_{\overrightarrow{\mathbb{X}}} : \overrightarrow{\mathbb{X}} \rightarrow \overrightarrow{\overrightarrow{\mathbb{X}}}$  has a left adjoint  $\mu_{\mathbb{X}} : \overrightarrow{\overrightarrow{\mathbb{X}}} \rightarrow \overrightarrow{\mathbb{X}}$  in  $\text{CAT}\mathcal{C}$ .*

Much as in Section 2 of [33] and Section 3 of [35], for each protocalibration  $\mathcal{R}$  of  $\mathcal{C}$ , we see we have a Kock-Zöberlein [25, 26, 46] pseudomonad

$$\overrightarrow{(-)}_{\mathcal{R}} = ((\overrightarrow{-}), \eta, \mu)$$

on  $\text{CAT}\mathcal{C}$ .

**Corollary 2.6.** *The inclusion of the subcategory  $\text{CAT}_{\mathcal{R}}\mathcal{C}$  of  $\text{CAT}\mathcal{C}$ , consisting of the  $\mathcal{R}$ -cocomplete objects and  $\mathcal{R}$ -cocontinuous morphisms, is pseudomonadic. That is, the inclusion is biequivalent over  $\text{CAT}\mathcal{C}$  to the forgetful pseudofunctor from the bicategory of Eilenberg-Moore pseudo-algebras for the pseudomonad  $\overrightarrow{(-)}_{\mathcal{R}}$ . In particular,  $\text{CAT}_{\mathcal{R}}\mathcal{C}$  is a complete bicategory and bicategorical limits are preserved by the inclusion.*

The *opposite*  $\mathbb{X}^{\text{op}}$  of  $\mathbb{X} \in \text{CAT}\mathcal{C}$  is obtained by composition with the 2-isomorphism  $(-)^{\text{op}} : \text{Cat} \rightarrow \text{Cat}^{\text{co}}$ . This is the value on objects for a 2-isomorphism

$$(-)^{\text{op}} : \text{CAT}\mathcal{C} \rightarrow (\text{CAT}\mathcal{C}^{\text{co}})^{\text{co}} . \quad (2.7)$$

Suppose  $\mathcal{L}$  is a protocalibration of  $\mathcal{C}^{\text{co}}$ . We define  $\mathbb{X} \in \text{CAT}\mathcal{C}$  to be  $\mathcal{L}$ -complete when  $\mathbb{X}^{\text{op}}$  is  $\mathcal{L}$ -cocomplete. It follows that the  $\mathcal{L}$ -completion  $\overleftarrow{\mathbb{X}}$  of  $\mathbb{X} \in \text{CAT}\mathcal{C}$  is  $(\overrightarrow{\mathbb{X}^{\text{op}}})^{\text{op}}$ . If  $\mathbb{X}$  is  $\mathcal{L}$ -complete and  $V \xrightarrow{f} U$  is in  $\mathcal{L}$ , we write  $\mathbb{X}_! f$  for the right adjoint of  $\mathbb{X}f : \mathbb{X}U \rightarrow \mathbb{X}V$ .

We write  ${}_{\mathcal{L}}\text{CAT}\mathcal{C}$  for the locally full subbcategory of  $\text{CAT}\mathcal{C}$  consisting of the  $\mathcal{L}$ -complete objects and  $\mathcal{L}$ -continuous morphisms. The 2-isomorphism (2.7) induces an isomorphism

$${}_{\mathcal{L}}\text{CAT}\mathcal{C} \cong (\text{CAT}{}_{\mathcal{L}}\mathcal{C}^{\text{co}})^{\text{co}} . \quad (2.8)$$

We have a dual Kock-Zöberlein pseudomonad on  $\text{CAT}\mathcal{C}$  for which the Eilenberg-Moore pseudo-algebras are the  $\mathcal{L}$ -complete objects where the algebra action on the object is right adjoint to the unit.

An object of  $\overleftarrow{\mathbb{X}}U$  is a pair  $(U \xleftarrow{u} S, x \in \mathbb{X}S)$  with  $u \in \mathcal{L}$ , as for  $\overrightarrow{\mathbb{X}}U$ . A morphism

$$(U \xleftarrow{u} S, x \in \mathbb{X}S) \rightarrow (U \xleftarrow{u'} S', x' \in \mathbb{X}S')$$

of  $\overleftarrow{\mathbb{X}}U$  is an isomorphism class of triples  $(S' \xrightarrow{w} S, u' \cong uw, (\mathbb{X}w)x \xrightarrow{\zeta} x')$  where  $\zeta$  is a morphism of  $\mathbb{X}S'$ .

$$\begin{array}{ccccc} U & \xleftarrow{u \in \mathcal{L}} & S & \xrightarrow{x} & \mathbb{X} \\ & \searrow^{u' \in \mathcal{L}} & \uparrow \cong & \swarrow \zeta & \\ & & S' & \xrightarrow{x'} & \mathbb{X} \end{array} \quad (2.9)$$

For  $r : V \rightarrow U$  in  $\mathcal{E}$ , the functor  $\overleftarrow{\mathbb{X}}r : \overleftarrow{\mathbb{X}}U \rightarrow \overleftarrow{\mathbb{X}}V$  takes the object  $(u, x)$  to  $(u_r, (\mathbb{X}r_u)x)$  with notation as in (2.5) which is used also in defining  $\overleftarrow{\mathbb{X}}r$  on morphisms. The unit  $\rho_{\mathbb{X}} : \mathbb{X} \rightarrow \overleftarrow{\mathbb{X}}$  for the pseudomonad has components at  $U$  the functor  $\rho_U$  taking  $x \in \mathbb{X}U$  to  $(1_U, x) \in \overleftarrow{\mathbb{X}}U$  and  $\zeta : x \rightarrow x'$  to  $(1_U, \zeta)$ . For emphasis we state the dual form of Proposition 2.4:

**Corollary 2.7.** *Suppose  $\mathcal{L}$  is a protocalibration of the bicategory  $\mathcal{C}^{\text{co}}$ . Then  $\mathbb{X} \in \text{CAT}\mathcal{C}$  is  $\mathcal{L}$ -complete if and only if  $\rho_{\mathbb{X}}$  has a right adjoint in  $\text{CAT}\mathcal{C}$ .*

We think of the right adjoint to  $\rho_{\mathbb{X}}$  as assigning a product to the indexed families of generalised objects of  $\mathbb{X}$ .



where  $\rho$  is the composite  $gv_r \xrightarrow{\kappa} k \cong v'_r h$ . Now we define the desired composite of the two span morphisms  $[f, \sigma]$  and  $[g, \tau]$  appearing in diagram (3.11) to be  $[h, s'\rho]$ .

We have a pseudofunctor

$$(-)_* : \mathcal{C} \rightarrow \mathrm{Spn}_{\mathcal{R}}\mathcal{C} \quad (3.13)$$

taking the morphism  $U \xrightarrow{r} V$  to the  $\mathcal{R}$ -span  $r_* = (U \xleftarrow{1_U} U \xrightarrow{r} V)$  and the 2-morphism  $\alpha : r \Rightarrow s$  to the 2-morphism

$$\begin{array}{ccc} & U & \\ 1_U \in \mathcal{R} \swarrow & \downarrow 1_U & \searrow r \\ U & \xrightarrow{1_U \in \mathcal{R}} U & \xrightarrow{s} V \\ & \cong & \alpha \end{array}$$

If  $r \in \mathcal{R}$  then  $r_*$  has a right adjoint  $r^* = (V \xleftarrow{r} U \xrightarrow{1_U} U)$  in  $\mathrm{Spn}_{\mathcal{R}}\mathcal{C}$ . Every morphism  $U \xleftarrow{u} S \xrightarrow{v} V$  of  $\mathrm{Spn}_{\mathcal{R}}\mathcal{C}$  decomposes as  $v_* \circ u^*$ .

Restriction along the opposite of (3.13) defines a pseudofunctor

$$\mathrm{CATSpn}_{\mathcal{R}}\mathcal{C} \rightarrow \mathrm{CAT}\mathcal{C} \quad (3.14)$$

which is pseudomonadic since  $(-)_*$  is the identity on objects and the restriction has both left and right biadjoints. Re-identification of the pseudoalgebras as follows is an objective variant of Lindner's result [27].

**Proposition 3.1.** *Restriction along the opposite of (3.13) defines a bicategorical equivalence*

$$\mathrm{CATSpn}_{\mathcal{R}}\mathcal{C} \sim \mathrm{CAT}_{\mathcal{R}}\mathcal{C} .$$

*Proof.* We shall describe the inverse biequivalence. Each  $\mathbb{X} \in \mathrm{CAT}_{\mathcal{R}}\mathcal{C}$  extends along the opposite of  $(-)_*$  to a pseudofunctor  $\mathbb{X}_{\mathrm{sp}} : (\mathrm{Spn}_{\mathcal{R}}\mathcal{C})^{\mathrm{op}} \rightarrow \mathrm{Cat}$  which agrees with  $\mathbb{X}$  on objects and is defined on morphisms by  $\mathbb{X}_{\mathrm{sp}}(v_* \circ u^*) = \mathbb{X}_* u \circ \mathbb{X} v$ . For 2-morphisms, we define  $\mathbb{X}_{\mathrm{sp}}[f, \sigma] : \mathbb{X}_* u \circ \mathbb{X} v \rightarrow \mathbb{X}_* u' \circ \mathbb{X} v'$  to be the mate under the adjunction  $\mathbb{X}_* u \dashv \mathbb{X} u$  of the composite

$$\mathbb{X} v \xrightarrow{\mathbb{X} \sigma} \mathbb{X} f \circ \mathbb{X} v' \xrightarrow{\mathbb{X} f \circ \eta_{u'} \circ \mathbb{X} v'} \mathbb{X} f \circ \mathbb{X} u' \circ \mathbb{X}_* u' \circ \mathbb{X} v' \cong \mathbb{X} u \circ \mathbb{X}_* u' \circ \mathbb{X} v'$$

where  $\eta_{u'}$  is the unit for  $\mathbb{X}_* u' \dashv \mathbb{X} u'$ . The composition-preservation constraints come from the pointwiseness in  $\mathcal{R}$ -cocompleteness: referring to the

diagram (3.11), we have

$$\begin{aligned}
\mathbb{X}_{\text{sp}}(v_* \circ u^*) \circ \mathbb{X}_{\text{sp}}(s_* \circ r^*) &\cong \mathbb{X}_*u \circ \mathbb{X}v \circ \mathbb{X}_*r \circ \mathbb{X}s \\
&\cong \mathbb{X}_*u \circ \mathbb{X}_*r_v \circ \mathbb{X}v_r \circ \mathbb{X}s \\
&\cong \mathbb{X}_*(u \circ r_v) \circ \mathbb{X}(s \circ v_r) \\
&\cong \mathbb{X}_{\text{sp}}((s \circ v_r)_* \circ (u \circ r_v)^*) \\
&\cong \mathbb{X}_{\text{sp}}(s_* \circ v_{r_*} \circ r_v^* \circ u^*) \\
&\cong \mathbb{X}_{\text{sp}}(s_* \circ r^* \circ v_* \circ u^*) .
\end{aligned}$$

Clearly  $\mathbb{X}_{\text{sp}} \circ (-)_* \simeq \mathbb{X}$ .

For any  $\mathcal{R}$ -cocontinuous pseudonatural transformation  $\theta : \mathbb{X} \rightarrow \mathbb{Y}$ , we have the diagram

$$\begin{array}{ccccc}
\mathbb{X}V & \xrightarrow{\mathbb{X}v} & \mathbb{X}S & \xrightarrow{\mathbb{X}_*u} & \mathbb{X}U \\
\theta_V \downarrow & \xleftarrow[\cong]{\theta_v} & \downarrow \theta_S & \xleftarrow[\cong]{\theta_u^{-1}} & \downarrow \theta_U \\
\mathbb{Y}V & \xrightarrow{\mathbb{Y}v} & \mathbb{Y}S & \xrightarrow{\mathbb{Y}_*u} & \mathbb{Y}U
\end{array}$$

for all  $S \xrightarrow{u} U$  in  $\mathcal{R}$  and  $S \xrightarrow{v} V$  in  $\mathcal{C}$  (using terminology from Proposition 1.4 for the mate of  $\theta_u$ ). The pasted composite diagram exhibits the extension of  $\theta$  to a pseudonatural transformation  $\theta_{\text{sp}} : \mathbb{X}_{\text{sp}} \rightarrow \mathbb{Y}_{\text{sp}}$ . For a modification  $\theta \Rightarrow \phi : \mathbb{X} \rightarrow \mathbb{Y}$ , we can keep the same components on objects to define the extended modification  $\theta_{\text{sp}} \Rightarrow \phi_{\text{sp}} : \mathbb{X}_{\text{sp}} \rightarrow \mathbb{Y}_{\text{sp}}$ .  $\square$

Let  $\mathcal{L}$  be a protocalibration of the bicategory  $\mathcal{C}^{\text{co}}$ . So we can define

$$\mathcal{L}\text{Spn}\mathcal{C} = (\text{Spn}_{\mathcal{L}}\mathcal{C}^{\text{co}})^{\text{co}} .$$

The 2-morphisms are represented by diagrams of the form (3.15).

$$\begin{array}{ccccc}
& & S & & \\
& u \in \mathcal{L} & \uparrow & v & \\
& \cong & \uparrow & \cong & \\
U & \xleftarrow{u' \in \mathcal{L}} & S' & \xrightarrow{v'} & V
\end{array} \tag{3.15}$$

For completeness we state the dual form of Proposition 3.1.

**Proposition 3.2.** *Restriction along the opposite of  $(-)_* : \mathcal{C} \rightarrow \mathcal{L}\text{Spn}\mathcal{C}$  defines a bicategorical equivalence*

$$\text{CAT}(\mathcal{L}\text{Spn}\mathcal{C}) \sim \mathcal{L}\text{CAT}\mathcal{C} .$$

## 4 A distributive law from compatibility

Let  $\mathcal{E}$  be a category with pullbacks. The functor defined by pulling back along  $u : S \rightarrow U$  is denoted by  $\Delta_u : \mathcal{E}/U \rightarrow \mathcal{E}/S$ . There is always a left adjoint to  $\Delta_u$  denoted by  $\Sigma_u$  and obtained by composition with  $u$ . Recall that a morphism  $u : S \rightarrow U$  is called *powerful* (or “exponentiable”) when  $\Delta_u$  has a right adjoint  $\Pi_u$ . Then, for each  $a : A \rightarrow U$ , the counit of the adjunction  $\Delta_u \dashv \Pi_u$  gives a morphism  $e$  as in diagram (4.16).

$$\begin{array}{ccccc}
 A & \xleftarrow{e} & P & \xrightarrow{\bar{u}} & B \\
 & \searrow a & \downarrow \lrcorner & & \downarrow \Pi_u a \\
 & & S & \xrightarrow{u} & U
 \end{array}
 \quad (4.16)$$

Let  $\mathcal{P}$  denote the set of powerful morphisms in  $\mathcal{E}$ ; by Corollary 2.6 of [41],  $\mathcal{P}$  is a protocalibration in the sense of Definition 1.2.

For this section fix two protocalibrations  $\mathcal{L}$  and  $\mathcal{R}$  of the category  $\mathcal{E}$ . We are particularly interested in the  $\mathcal{R}$ -cocompletion and the  $\mathcal{L}$ -completion. So, in this section,  $\overrightarrow{\mathbb{X}}$  will mean the  $\mathcal{R}$ -cocompletion of  $\mathbb{X} \in \text{CAT}\mathcal{E}$  and  $\overleftarrow{\mathbb{X}}$  will mean the  $\mathcal{L}$ -completion of  $\mathbb{X}$ .

Our Definition 4.1 is motivated by concepts in Blumberg-Hill [5, 6] and Chan [9].

**Definition 4.1.** *Suppose  $\mathcal{L}$  and  $\mathcal{R}$  are protocalibrations of the category  $\mathcal{E}$ . We say the pair  $(\mathcal{L}, \mathcal{R})$  is compatible when  $\mathcal{L} \subseteq \mathcal{P} \cap \mathcal{R}$  and  $\Pi_r v \in \mathcal{R}$  whenever  $r \in \mathcal{L}$  and  $v \in \mathcal{R}$ .*

**Example 4.2.** *In a category  $\mathcal{E}$  with pullbacks, the maximum example of a compatible pair is  $(\mathcal{P}, \mathcal{E})$ .*

**Lemma 4.3.** *Suppose the pair  $(\mathcal{L}, \mathcal{R})$  of protocalibrations of  $\mathcal{E}$  is compatible. Spans in  $\mathcal{E}$  with left leg invertible and right leg in  $\mathcal{R}$  are morphisms in  $\mathcal{L}\text{Spn}\mathcal{E}$  and as such form a protocalibration of  $\mathcal{L}\text{Spn}\mathcal{E}$  denoted by  $\mathcal{R}_*$ .*

*Proof.*  $\mathcal{R}_*$  consists of the spans isomorphic to  $w_* = (V \xleftarrow{1_V} V \xrightarrow{w} V')$  for some  $V \xrightarrow{w} V'$  in  $\mathcal{R}$ . The only condition for a protocalibration which is not totally clear is (G); but this follows from the fact that each square (4.17) is a bipullback of a groupoid fibration.  $\square$

$$\begin{array}{ccc}
 \mathcal{L}\text{Spn}\mathcal{E}(U, V)^{\text{op}} & \xrightarrow{\text{incl.}} & \text{Spn}\mathcal{E}(U, V) \\
 w_* \circ - \downarrow \lrcorner & & \downarrow w_* \circ - \\
 \mathcal{L}\text{Spn}\mathcal{E}(U, V')^{\text{op}} & \xrightarrow{\text{incl.}} & \text{Spn}\mathcal{E}(U, V')
 \end{array}
 \quad (4.17)$$

Suppose  $(\mathcal{L}, \mathcal{R})$  is compatible. There is a bicategorical distributive law

$$\lambda : \overleftarrow{(-)} \circ \overrightarrow{(-)} \longrightarrow \overrightarrow{(-)} \circ \overleftarrow{(-)}$$

where the component functor

$$\lambda_U : \overleftarrow{\mathbb{X}}U \longrightarrow \overrightarrow{\mathbb{X}}U$$

takes  $(U \xleftarrow{u} S, S \xleftarrow{a} A, x \in \mathbb{X}A)$ , where  $u \in \mathcal{L}$  and  $a \in \mathcal{R}$ , to

$$(U \xleftarrow{\Pi_u a} B, B \xleftarrow{\bar{u}} P, (\mathbb{X}e)x \in \mathbb{X}P)$$

in the notation of diagram (4.16).

**Proposition 4.4.** *If  $\mathbb{X} \in \text{CAT}^{\mathcal{E}}$  is  $\mathcal{L}$ -complete then so is the  $\mathcal{R}$ -cocompletion  $\overrightarrow{\mathbb{X}}$ . Moreover, the unit  $\eta_{\mathbb{X}} : \mathbb{X} \rightarrow \overrightarrow{\mathbb{X}}$  is in  ${}_{\varphi}\text{CAT}^{\mathcal{E}}$ .*

*Proof.* Suppose  $V \xrightarrow{r} U$  is in  $\mathcal{L}$ . We will use the right adjoints  $\Pi_r$  and  $\mathbb{X}_!r$  available because of powerfulness and of  $\mathcal{L}$ -completeness of  $\mathbb{X}$ .

We must show that  $\overrightarrow{\mathbb{X}}r$  has a right adjoint. For each  $(V \xleftarrow{v} T, y \in \mathbb{X}T) \in \overrightarrow{\mathbb{X}}V$ , we must produce an object  $(U \xleftarrow{v'} R, y' \in \mathbb{X}R) \in \overrightarrow{\mathbb{X}}U$  and a bijection

$$\overrightarrow{\mathbb{X}}V((\Delta_r u, (\mathbb{X}\bar{r})x), (v, y)) \cong \overrightarrow{\mathbb{X}}U((u, x), (v', y')) \quad (4.18)$$

natural in  $(u, x) \in \overrightarrow{\mathbb{X}}U$ , where the following square is a pullback.

$$\begin{array}{ccc} P & \xrightarrow{\bar{r}} & S \\ \Delta_r u \downarrow \lrcorner & & \downarrow u \\ V & \xrightarrow{r} & U \end{array}$$

Take an element  $(w, \xi)$  as shown in (4.19) of the left-hand side of (4.18).

$$\begin{array}{ccccc} & & P & & \\ & \Delta_r u \swarrow & \downarrow w & \xleftarrow{\xi} & \searrow (\mathbb{X}\bar{r})x \\ V & \xleftarrow{v} & T & \xrightarrow{y} & \mathbb{X} \end{array} \quad (4.19)$$

Since  $r$  is powerful, morphisms  $w$  making the left triangle of (4.19) commute are in natural bijection with morphisms  $w' : S \rightarrow R$  such that  $\Pi_r v \circ w' = u$  where  $R$  is the domain of  $\Pi_r v$ . Since  $\bar{r}$  is in  $\mathcal{L}$  and  $\mathbb{X}$  is  $\mathcal{L}$ -complete,

morphisms  $\xi : (\mathbb{X}\bar{r})x \rightarrow (\mathbb{X}w)y$  are in natural bijection with morphisms  $\bar{\xi} : x \rightarrow (\mathbb{X}_1\bar{r})(\mathbb{X}w)y$ . The diagram below shows how  $w$  and  $w'$  are related.

$$\begin{array}{ccccc}
& & P & \xrightarrow{\bar{r}} & S \\
& & \downarrow \lrcorner & & \downarrow w' \\
& & \Delta_{\bar{r}}w' & & \\
& w \nearrow & Q & \xrightarrow{\bar{r}} & R \\
& e \leftarrow & \downarrow \lrcorner & & \downarrow \Pi_r v \\
& & \Delta_r \Pi_r v & & \\
& v \searrow & V & \xrightarrow{r} & U
\end{array}$$

Also, we have the Chevalley-Beck isomorphism  $(\mathbb{X}w')(\mathbb{X}_1\bar{r}) \cong (\mathbb{X}_1\bar{r})(\mathbb{X}\Delta_{\bar{r}}w')$ . Consequently, we have  $(\mathbb{X}_1\bar{r})(\mathbb{X}w) \cong (\mathbb{X}_1\bar{r})(\mathbb{X}\Delta_{\bar{r}}w')(\mathbb{X}e) \cong (\mathbb{X}w')(\mathbb{X}_1\bar{r})(\mathbb{X}e)$ . So  $\bar{\xi} : x \rightarrow (\mathbb{X}_1\bar{r})(\mathbb{X}w)y$  are in bijection with  $\bar{\xi} : x \rightarrow (\mathbb{X}w')(\mathbb{X}_1\bar{r})(\mathbb{X}e)y$ . Therefore we have the desired bijection (4.18) with  $v' = \Pi_r v$  and  $y' = (\mathbb{X}_1\bar{r})(\mathbb{X}e)y$ .

Since  $\overrightarrow{\mathbb{X}}$  is  $\mathcal{R}$ -cocomplete, it already satisfies the CB-condition for left adjoints  $\overrightarrow{\mathbb{X}}_* r$ . By replacing all functors in that invertibility condition by their right adjoints when  $r \in \mathcal{L}$ , we induce an invertible mate yielding the CB-condition for the right adjoints  $\overrightarrow{\mathbb{X}}_1 r$ . So  $\overrightarrow{\mathbb{X}}$  is  $\mathcal{L}$ -complete.

The second sentence of the proposition follows on checking that the canonical natural transformation

$$\begin{array}{ccc}
\mathbb{X}V & \xrightarrow{\eta_{\mathbb{X}}V} & \overrightarrow{\mathbb{X}}V \\
\mathbb{X}_1 r \downarrow & \Longrightarrow & \downarrow \overrightarrow{\mathbb{X}}_1 r \\
\mathbb{X}U & \xrightarrow{\eta_{\mathbb{X}}U} & \overrightarrow{\mathbb{X}}U
\end{array}$$

is invertible. □

The pseudomonad  $\overrightarrow{(-)}$  on  $\text{CAT}^{\mathcal{E}}$  therefore lifts to  ${}_{\mathcal{L}}\text{CAT}^{\mathcal{E}}$ .

$$\begin{array}{ccc}
{}_{\mathcal{L}}\text{CAT}^{\mathcal{E}} & \xrightarrow{{}_{\mathcal{L}}\overrightarrow{(-)}} & {}_{\mathcal{L}}\text{CAT}^{\mathcal{E}} \\
\text{incl} \downarrow & \cong & \downarrow \text{incl} \\
\text{CAT}^{\mathcal{E}} & \xrightarrow{\overrightarrow{(-)}} & \text{CAT}^{\mathcal{E}}
\end{array} \tag{4.20}$$

The pseudo-algebras for the lifted pseudomonad are those of the composite pseudomonad  $\mathbb{X} \mapsto \overrightarrow{\mathbb{X}}$  on  $\text{CAT}^{\mathcal{E}}$  form the 2-category we denote by  ${}_{\mathcal{L}}\text{CAT}_{\mathcal{R}}^{\mathcal{E}}$  at the top of the diamond (4.21) of inclusion 2-functors. Using Lemma 4.3 and Proposition 3.1, we have biequivalences

$${}_{\mathcal{L}}\text{CAT}_{\mathcal{R}}^{\mathcal{E}} \sim \text{CAT}_{\mathcal{R}_*}({}_{\mathcal{L}}\text{Spn}^{\mathcal{E}}) \sim \text{CATSpn}_{\mathcal{R}_*}({}_{\mathcal{L}}\text{Spn}^{\mathcal{E}}) .$$



While the composite pseudomonad is not Kock-Zöberlein or dual Kock-Zöberlein, the pseudo-algebra structures are unique up to isomorphism: it is a property of an  $\mathcal{E}$ -variable category to be a pseudo-algebra for both  $\overleftarrow{(-)}$  and  $\overrightarrow{(-)}$ , and to satisfy the distributive law.

$$\begin{array}{ccc}
 & \mathcal{L}\text{CAT}_{\mathcal{R}\mathcal{E}} & \\
 \swarrow & & \searrow \\
 \mathcal{L}\text{CAT}_{\mathcal{E}} & & \text{CAT}_{\mathcal{R}\mathcal{E}} \\
 \searrow & & \swarrow \\
 & \text{CAT}_{\mathcal{E}} &
 \end{array} \tag{4.21}$$

## 5 Polynomials as spans of spans

Let  $\mathcal{E}$  be a category with pullbacks.

There is a bicategory  $\text{Ply}_{\mathcal{E}}$  of polynomials in  $\mathcal{E}$ ; in the notation of Walker [44] it is  $\text{Poly}(\mathcal{E})$  and in the notation of Proposition 8.4 of [40] it is the bicategory  $\text{PolySpn}_{\mathcal{E}}$ . The objects are those of  $\mathcal{E}$ . The morphisms from  $X$  to  $Y$  are *polynomials*  $p = (X \xleftarrow{r} A \xrightarrow{n} B \xrightarrow{t} Y)$  in  $\mathcal{E}$  with  $n$  powerful. The morphisms between polynomials are as in Subsection 2.11 of [15] and Definition 8.2 of [40]: they are isomorphism classes of commutative diagrams

$$\begin{array}{ccccc}
 & & A & \xrightarrow{n} & B \\
 & & \uparrow \ell & & \uparrow 1_B \\
 X & \xleftarrow{r} & & & & \xrightarrow{t} & Y \\
 & & P & \xrightarrow{n'_g} & B \\
 & & \downarrow g_{n'} \lrcorner & & \downarrow g \\
 & & A' & \xrightarrow{n'} & B'
 \end{array} \tag{5.22}$$

wherein the square as indicated is a pullback. Composition uses bipullback of spans of spans (see [15, 45, 40]). As a consequence of Theorem 5.1 and Proposition 6.6 below, finite coproducts exist in each  $\text{Ply}_{\mathcal{E}}(X, Y)$  and are preserved by precomposition with morphisms, yet generally not by postcomposition making the bicategory  $\text{Ply}_{\mathcal{E}}$  enriched in a certain skew monoidal bicategory  $\text{Cat}_{+\text{sk}}$  (extending a bit the terminology of [16]).

Recall the philosophy of [40] that it is useful to regard  $p$  as a span

$$p = (X \xleftarrow{(n, A, r)} B \xrightarrow{(1_B, B, t)} Y) \tag{5.23}$$

in  $\text{Spn}^{\mathcal{E}}$ . Then the morphisms interpret as diagrams

$$\begin{array}{ccc}
 & B & \\
 (n,A,r) \swarrow & \downarrow (1,B,g) \xrightarrow{\lambda} & \searrow (1,B,t) \\
 X & B' & Y \\
 (n',A',r') \swarrow & & \searrow (1,B',t')
 \end{array} . \tag{5.24}$$

We modify this a bit by taking a compatible pair  $(\mathcal{L}, \mathcal{R})$  of protocalibrations on  $\mathcal{E}$  and restricting the polynomials (5.23) to those with  $n \in \mathcal{L}$  and  $t \in \mathcal{R}$ . Restricting the morphisms of  $\text{Ply}^{\mathcal{E}}$  in this way, we obtain a sub-bicategory which we denote by  $\mathcal{L}\text{Ply}_{\mathcal{R}}^{\mathcal{E}}$ .

**Theorem 5.1.** *If  $(\mathcal{L}, \mathcal{R})$  is a compatible pair of protocalibrations on the category  $\mathcal{E}$  then*

$$(\mathcal{L}\text{Ply}_{\mathcal{R}}^{\mathcal{E}})^{\text{op}} = \text{Spn}_{\mathcal{R}^*}(\mathcal{L}\text{Spn}^{\mathcal{E}}) .$$

*Proof.* The 2-morphisms of  $\text{Spn}_{\mathcal{R}}(\mathcal{L}\text{Spn}^{\mathcal{E}})$  are isomorphism classes of morphisms as pictured in (5.25). Unpacking this shows that the data involved is exactly as in the diagram (5.22) in  $\mathcal{E}$ .

$$\begin{array}{ccc}
 & B & \\
 (1,B,t) \swarrow & \downarrow (1,B,g) \xrightarrow{\lambda} & \searrow (n,A,r) \\
 Y & B' & X \\
 (1,B',t') \swarrow & & \searrow (n',A',r')
 \end{array} . \tag{5.25}$$

While unpacking, just note that the 2-morphism  $\lambda$  in  $\mathcal{L}\text{Spn}^{\mathcal{E}}$  is in the reverse direction as a morphism of spans.  $\square$

**Corollary 5.2.**  $\text{Ps}(\mathcal{L}\text{Ply}_{\mathcal{R}}^{\mathcal{E}}, \text{Cat}) \sim \mathcal{L}\text{CAT}_{\mathcal{R}}^{\mathcal{E}}$

*Proof.* Using Theorem 5.1 and Propositions 3.1 and 3.2, we have

$$\begin{aligned}
 \text{Ps}(\mathcal{L}\text{Ply}_{\mathcal{R}}^{\mathcal{E}}, \text{Cat}) &\sim \text{CAT}(\text{Spn}_{\mathcal{R}^*}(\mathcal{L}\text{Spn}^{\mathcal{E}})) \\
 &\sim \text{CAT}_{\mathcal{R}^*}(\mathcal{L}\text{Spn}^{\mathcal{E}}) \\
 &\sim \mathcal{L}\text{CAT}_{\mathcal{R}}^{\mathcal{E}} . \quad \square
 \end{aligned}$$

$$\begin{array}{ccc}
 & \mathcal{E} & \\
 (-)_* \swarrow & & \searrow (-)_* \\
 \mathcal{L}\text{Spn}^{\mathcal{E}} & & \text{Spn}_{\mathcal{R}}^{\mathcal{E}} \\
 & \searrow & \swarrow \\
 & (\mathcal{L}\text{Ply}_{\mathcal{R}}^{\mathcal{E}})^{\text{op}} &
 \end{array} \tag{5.26}$$

## 6 Finite coproducts for spans and polynomials

Examples of the concepts of this section are provided by lextensive categories. Recall that a category  $\mathcal{E}$  is *lextensive* (in the sense used by Schanuel [30]) when it has finite limits, finite coproducts, and, equivalences

$$\mathbf{1} \simeq \mathcal{E}/_0 \quad \text{and} \quad \mathcal{E}/_U \times \mathcal{E}/_V \simeq \mathcal{E}/_{U+V} .$$

The second equivalence takes an object  $(A \xrightarrow{h} U, B \xrightarrow{k} V) \in \mathcal{E}/_U \times \mathcal{E}/_V$  to the object  $(A + B \xrightarrow{h+k} U + V) \in \mathcal{E}/_{U+V}$ ; the fact that this inverse equivalence is fully faithful means that a commutative triangle

$$\begin{array}{ccc} A + B & \xrightarrow{f} & A' + B' \\ & \searrow^{h+k} & \swarrow_{h'+k'} \\ & U + V & \end{array} \quad (6.27)$$

implies there are unique  $r : A \rightarrow A'$  and  $s : B \rightarrow B'$  such that  $f = r + s$ ,  $h = h'r$  and  $h' = k's$ .

It follows that the initial object  $0$  is strictly initial and, for every coproduct diagram  $U \xrightarrow{i} U + V \xleftarrow{j} V$  in  $\mathcal{E}$  and every  $R \xrightarrow{f} U + V$ , there is a diagram

$$\begin{array}{ccccc} S & \xrightarrow{\bar{i}} & R & \xleftarrow{\bar{j}} & T \\ h \downarrow & & f \downarrow & & \downarrow k \\ U & \xrightarrow{i} & U + V & \xleftarrow{j} & V \end{array} \quad (6.28)$$

in which the squares are pullbacks and the rows are coproduct diagrams.

This should motivate the following bicategorical concepts.

Suppose the bicategory  $\mathcal{C}$  admits the binary bicoproduct  $U \xrightarrow{i} U + V \xleftarrow{j} V$  and  $\mathcal{R}$  is a protocalibration of  $\mathcal{C}$ . For the pseudofunctor  $\mathbb{C}_{\mathcal{R}} : \mathcal{C}^{\text{op}} \rightarrow \text{Cat}$  of Definition 2.2, we have a canonical comparison functor

$$\mathbb{C}_{\mathcal{R}}(U + V) \rightarrow \mathbb{C}_{\mathcal{R}}U \times \mathbb{C}_{\mathcal{R}}V$$

taking  $Z \xrightarrow{w} U + V$  to the pair  $(X \xrightarrow{w_i} U, Y \xrightarrow{w_j} V)$  defined by bipullbacks

$$\begin{array}{ccc} X & \xrightarrow{i_w} & Z \\ w_i \downarrow \lrcorner & \cong & \downarrow w \\ U & \xrightarrow{i} & U + V \end{array} \quad \begin{array}{ccc} Y & \xrightarrow{j_w} & Z \\ w_j \downarrow \lrcorner & \cong & \downarrow w \\ V & \xrightarrow{j} & U + V . \end{array}$$

**Lemma 6.1.** *Suppose the bicategory  $\mathcal{C}$  admits, and the protocalibration  $\mathcal{R}$  is closed under, finite bicomponents. Then the canonical comparison functors*

$$\mathbb{C}_{\mathcal{R}}0 \rightarrow \mathbf{1} \quad \text{and} \quad \mathbb{C}_{\mathcal{R}}(U + V) \rightarrow \mathbb{C}_{\mathcal{R}}U \times \mathbb{C}_{\mathcal{R}}V$$

*have left adjoints respectively defined on objects by*

$$(0 \in \mathbf{1}) \mapsto (0 \xrightarrow{1_0} 0) \quad \text{and} \quad (X \xrightarrow{u \in \mathcal{R}} U, Y \xrightarrow{v \in \mathcal{R}} V) \mapsto (X + Y \xrightarrow{u+v \in \mathcal{R}} U + V).$$

*Proof.* This is an exercise in using the universal properties of bicomponent and bipullback.  $\square$

**Definition 6.2.** *Suppose  $\mathcal{R}$  is a protocalibration of the bicategory  $\mathcal{C}$ . We say a bicategory  $\mathcal{C}$  is  $\mathcal{R}$ -extensive when it admits finite bicomponents and these are taken into finite biproducts by the pseudofunctor  $\mathbb{C}_{\mathcal{R}} : \mathcal{C}^{\text{op}} \rightarrow \text{Cat}$  of Definition 2.2.*

$$\mathbb{C}_{\mathcal{R}}0 \simeq \mathbf{1} \quad \text{and} \quad \mathbb{C}_{\mathcal{R}}(U + V) \simeq \mathbb{C}_{\mathcal{R}}U \times \mathbb{C}_{\mathcal{R}}V$$

*When  $\mathcal{R}$  is closed under finite bicomponents, the inverse equivalences are the left adjoints in Definition 6.1.*

**Example 6.3.** *Suppose  $\mathcal{E}$  is a lextensive category. Then the maximum protocalibration  $\mathcal{E}$  is closed under finite coproducts and  $\mathcal{E}$  is  $\mathcal{E}$ -extensive.*

**Example 6.4.** *Again suppose  $\mathcal{E}$  is a lextensive category. The subcategory  $\mathcal{P}$  consisting of the powerful morphisms is closed under finite coproducts. To see this, first realize that injections  $X \xrightarrow{i} X + Y$  are powerful since  $\mathcal{E}_{/(X+Y)} \xrightarrow{\Delta_i} \mathcal{E}_{/X}$  transports across equivalence to the projection  $\mathcal{E}_{/X} \times \mathcal{E}_{/Y} \rightarrow \mathcal{E}_{/X}$  which has a right adjoint since  $\mathcal{E}_{/Y}$  has a terminal object. Then notice that pullback along  $0 \rightarrow A$  is equivalent to the functor  $\mathcal{E}_{/A} \rightarrow \mathbf{1}$  whose right adjoint picks out the terminal object and, if  $A \xrightarrow{u} X$  and  $B \xrightarrow{v} X$  are powerful, then pullback along  $A + B \xrightarrow{[u,v]} X$  is equivalent to  $\mathcal{E}_{/X} \xrightarrow{(\Delta_u, \Delta_v)} \mathcal{E}_{/A} \times \mathcal{E}_{/B}$  which has a right adjoint taking  $(U \xrightarrow{a} A, V \xrightarrow{b} B)$  to the product  $\Pi_u(a) \times \Pi_v(b)$  in  $\mathcal{E}_{/X}$ . Indeed,  $\mathcal{E}$  is  $\mathcal{P}$ -extensive because the equivalence  $\mathcal{E}_{/(X+Y)} \simeq \mathcal{E}_{/X} \times \mathcal{E}_{/Y}$  restricts to the powerful objects.*

**Proposition 6.5.** *Suppose the bicategory  $\mathcal{C}$  is  $\mathcal{R}$ -extensive and that the protocalibration  $\mathcal{R}$  is closed under the finite bicomponents. If  $\underline{\mathbb{X}} \in \text{CAT}\mathcal{C}$  preserves finite bicategorical products then the  $\mathcal{R}$ -cocompletion  $\overline{\mathbb{X}}$  also preserves finite bicategorical products.*

*Proof.* The objects of  $\overrightarrow{\mathbb{X}}(U + V)$  are pairs  $(U + V \xleftarrow{w} R, z \in \mathbb{X}R)$  with  $w \in \mathcal{R}$ . Since  $\mathcal{C}$  is  $\mathcal{R}$ -extensive,  $R \xrightarrow{w} U + V$  is equivalent over  $U + V$  to some  $S + T \xrightarrow{u+v} U + V$  and the morphisms  $u$  and  $v$  are in  $\mathcal{R}$ . Since  $\mathbb{X}$  preserves finite biproducts,  $z \in \mathbb{X}R$  gives  $(x, y) \in \mathbb{X}S \times \mathbb{X}T$ . The assignment  $(R, z) \mapsto ((S, x), (T, y))$  is the required equivalence  $\overrightarrow{\mathbb{X}}(U + V) \simeq \overrightarrow{\mathbb{X}}U \times \overrightarrow{\mathbb{X}}V$ . It is clear that  $\overrightarrow{\mathbb{X}}0 \simeq \mathbf{1}$  since all  $0 \xleftarrow{u \in \mathcal{R}} S$  are isomorphic equivalences and  $\mathbb{X}0 \simeq \mathbf{1}$ .  $\square$

**Proposition 6.6.** *Suppose the bicategory  $\mathcal{C}$  is  $\mathcal{R}$ -extensive and that the protocalibration  $\mathcal{R}$  is closed under the finite bicoproducts. Then*

- i. *the bicategory  $\text{Spn}_{\mathcal{R}}\mathcal{C}$  admits finite bicoproducts, and*
- ii. *each homcategory  $\text{Spn}_{\mathcal{R}}\mathcal{C}(X, Y)$  admits finite coproducts preserved by composition on the diagrammatic right by any morphism  $Y \xrightarrow{(u, S, v)} Z$  where  $u \in \mathcal{R}$ .*

*Proof.* The bi-initial object of  $\text{Spn}_{\mathcal{R}}\mathcal{C}$  is that of  $\mathcal{C}$  denoted 0. For the 1-morphism part of the universal property: if  $0 \xleftarrow{u} S \xrightarrow{v} Y$  is a left  $\mathcal{R}$ -span then  $u$  is an equivalence (using  $\mathbb{C}_{\mathcal{R}}0 \sim \mathbf{1}$ ); so  $S$  is bi-initial and we conclude that all such spans are isomorphic in  $\text{Spn}_{\mathcal{R}}\mathcal{C}$  while the 2-morphism property of 0 follows from that of  $0 \in \mathcal{C}$ .

We will show that a bicoproduct  $U \xrightarrow{i} U + V \xleftarrow{j} V$  in  $\mathcal{C}$  gives one  $U \xrightarrow{i_*} U + V \xleftarrow{j_*} V$  in  $\text{Spn}_{\mathcal{R}}\mathcal{C}$ . Note that coprojections  $i, j$  and codiagonals are in  $\mathcal{R}$ . Take spans  $U \xleftarrow{u} S \xrightarrow{v} W$  and  $V \xleftarrow{r} T \xrightarrow{v} W$  with  $u, r \in \mathcal{R}$  (and hence  $u + r \in \mathcal{R}$ ). We obtain  $U + V \xleftarrow{u+r \in \mathcal{R}} S + T \xrightarrow{[v, s]} W$  whose composites with  $i_*$  and  $j_*$  are isomorphic to the given two spans because our assumptions give the bipullbacks

$$\begin{array}{ccc} S & \xrightarrow{\bar{i}} & S + T \\ \downarrow \lrcorner & \cong & \downarrow u+r \\ U & \xrightarrow{i} & U + V \end{array} \quad \begin{array}{ccc} T & \xrightarrow{\bar{j}} & S + T \\ \downarrow \lrcorner & \cong & \downarrow u+r \\ V & \xrightarrow{j} & U + V . \end{array}$$

The 2-morphism part of the universal property uses  $\mathcal{R}$ -extensivity and the properties of  $\mathcal{R}$ :

$$\begin{array}{ccccc} & & S + T & & \\ & \swarrow^{u+r \in \mathcal{R}} & \downarrow f+g & \searrow^{[v, s]} & \\ U + V & \xleftarrow{u'+r' \in \mathcal{R}} & S' + T' & \xrightarrow{[v', s']} & W . \end{array}$$

The coproduct of  $X \xleftarrow{u \in \mathcal{R}} S \xrightarrow{v} Y$  and  $X \xleftarrow{u' \in \mathcal{R}} S' \xrightarrow{v'} Y$  in  $\mathrm{Spn}_{\mathcal{R}} \mathcal{C}(X, Y)$  is  $X \xleftarrow{[u, u'] \in \mathcal{R}} S + S' \xrightarrow{[v, v']} Y$ . Contemplation of the diagram

$$\begin{array}{ccccc}
 & & R + R' & & \\
 & \swarrow^{a+b \in \mathcal{R}} & & \searrow_{[c, d]} & \\
 & S + S' & \cong & T & \\
 \swarrow_{[u, u'] \in \mathcal{R}} & & & & \searrow_s \\
 X & & Y & \xleftarrow{r \in \mathcal{R}} & Z
 \end{array}$$

(where the diamond is a bipullback) reveals the claimed right preservation property.  $\square$

For any bicategory  $\mathcal{D}$  admitting finite bicoproducts, we write  $\mathrm{fpCAT} \mathcal{D}$  for the full sub-bicategory of  $\mathrm{CAT} \mathcal{D}$  consisting of the finite biproduct preserving pseudofunctors  $\mathcal{D}^{\mathrm{op}} \rightarrow \mathrm{Cat}$ . If  $\mathcal{T}$  is a protocalibration of  $\mathcal{D}$ , we write  $\mathrm{fpCAT}_{\mathcal{T}} \mathcal{D}$  for the intersection of  $\mathrm{fpCAT} \mathcal{D}$  and  $\mathrm{CAT}_{\mathcal{T}} \mathcal{D}$ .

**Proposition 6.7.** *Under the hypotheses of Proposition 6.6, the biequivalence of Proposition 3.1 restricts to a biequivalence*

$$\mathrm{fpCAT} \mathrm{Spn}_{\mathcal{R}} \mathcal{C} \sim \mathrm{fpCAT}_{\mathcal{R}} \mathcal{C} .$$

*Proof.* The pseudofunctor  $(-)_* : \mathcal{C} \rightarrow \mathrm{Spn}_{\mathcal{R}} \mathcal{C}$  preserves finite bicategorical coproduct so restriction along its opposite restricts to a pseudofunctor from left to right which is an equivalence on homcategories. Also, if  $\mathbb{X} \in \mathrm{CAT}_{\mathcal{R}} \mathcal{C}$  preserves finite bicategorical products then so does its extension  $\mathbb{X}_{\mathrm{sp}}$  to  $(\mathrm{Spn}_{\mathcal{R}} \mathcal{C})^{\mathrm{op}}$  since  $\mathcal{R}$  is closed under finite bicoproduct.  $\square$

**Example 6.8.** *Continuing with Example 6.3, Proposition 6.6 tells us that  $\mathrm{Spn} \mathcal{E} = \mathrm{Spn}_{\mathcal{E}} \mathcal{E}$  has bicoproducts and local coproducts preserved by right composition. However,  $\mathrm{Spn} \mathcal{E} = (\mathrm{Spn} \mathcal{E})^{\mathrm{op}}$ , so local coproducts are preserved by composition on the left as well. Furthermore, if  $\mathcal{R}$  is any protocalibration of the left extensive category  $\mathcal{E}$  closed under finite coproducts then  $\mathcal{E}$  is  $\mathcal{R}$ -extensive. The bicoproducts and local coproducts in  $\mathrm{Spn}_{\mathcal{R}} \mathcal{E}$  are preserved by the inclusion pseudofunctor into  $\mathrm{Spn} \mathcal{E}$ . It follows that the local coproducts in  $\mathrm{Spn}_{\mathcal{R}} \mathcal{E}$  are preserved by composition on both sides. Consider the symmetric closed monoidal bicategory  $\mathrm{Cat}_+$  of categories with finite coproducts and finite-coproduct-preserving functors; the internal hom  $[\mathcal{A}, \mathcal{B}]_+$  of  $\mathrm{Cat}_+$  is  $\mathrm{Cat}_+(\mathcal{A}, \mathcal{B})$  with pointwise coproduct and the unit object is the category set of finite sets. Cartesian product  $A \times B$  of categories is a direct sum in  $\mathrm{Cat}_+$  and the terminal category  $\mathbf{1}$  is also initial. What we have is that  $\mathrm{Spn} \mathcal{E}$  is  $\mathrm{Cat}_+$ -enriched and to say a pseudofunctor  $(\mathrm{Spn} \mathcal{E})^{\mathrm{op}} \rightarrow \mathrm{Cat}_+$  is biproduct preserving is to say it is  $\mathrm{Cat}_+$ -enriched.*

**Lemma 6.9.** *Suppose  $(\mathcal{L}, \mathcal{R})$  is a compatible pair of protocalibrations on an  $\mathcal{R}$ -extensive category  $\mathcal{E}$ . Suppose both  $\mathcal{L}$  and  $\mathcal{R}$  are closed under finite coproducts in  $\mathcal{E}$ . The protocalibration  $\mathcal{R}_*$  of the bicategory  $\mathcal{L}\text{Spn}\mathcal{E}$  (see Lemma 4.3) is closed under bicoproducts and  $\mathcal{L}\text{Spn}\mathcal{E}$  is  $\mathcal{R}_*$ -extensive.*

*Proof.* For this proof, put  $\mathcal{C} = \mathcal{L}\text{Spn}\mathcal{E}$ . The injections into a bicoproduct and the codiagonals are in  $\mathcal{R}_*$ . The bicoproduct of  $f_*$  and  $f'_*$  is isomorphic to  $(f + f')_*$  and so in  $\mathcal{R}_*$ . It remains to prove  $\mathbb{C}_{\mathcal{R}_*}$  preserves biproducts. Up to equivalence, we can take the objects of  $\mathbb{C}_{\mathcal{R}_*}U$  to be morphisms  $X \xrightarrow{a} U$  in  $\mathcal{R}$  since every morphism  $X \rightarrow U$  in  $\mathcal{R}_*$  is isomorphic to some  $X \xrightarrow{a_*} U$  with  $a \in \mathcal{R}$ . Moreover, every morphism

$$\begin{array}{ccc} X & \xrightarrow{[u, S, v]} & Y \\ & \searrow a_* & \swarrow b_* \\ & U & \end{array} \quad \begin{array}{c} \cong \\ \Rightarrow \end{array}$$

in  $\mathbb{C}_{\mathcal{R}_*}U$  has a morphism of the form  $f_*$  in the isomorphism class  $[u, S, v]$ . Thus we have an equivalence of categories  $\mathbb{C}_{\mathcal{R}_*}U \simeq \mathbb{E}_{\mathcal{R}}U$ .<sup>3</sup> Also, the canonical functor  $\mathbb{C}_{\mathcal{R}_*}(U + V) \rightarrow \mathbb{C}_{\mathcal{R}_*}U \times \mathbb{C}_{\mathcal{R}_*}V$  transports across the equivalences to the lextensivity equivalence  $\mathbb{E}_{\mathcal{R}}(U + V) \simeq \mathbb{E}_{\mathcal{R}}U \times \mathbb{E}_{\mathcal{R}}V$ .  $\square$

As a corollary of Theorem 5.1, Proposition 6.6 and Lemma 6.9, we obtain:

**Theorem 6.10.** *Under the hypotheses of Lemma 6.9, the bicategory  $\mathcal{L}\text{Ply}_{\mathcal{R}}\mathcal{E}$  admits finite biproducts and its homs admit finite coproducts preserved by composition on the diagrammatic left.*

## 7 Objective Mackey functors

*Suppose the bicategory  $\mathcal{C}$  is finitely bicomplete and  $\mathcal{R}$ -extensive for a protocalibration  $\mathcal{R}$  which, when regarded as a wide subcategory of  $\mathcal{C}$ , is closed under finite bicoproducts.*

**Definition 7.1.** *An objective  $\mathcal{R}$ -Mackey functor on  $\mathcal{C}$  is an  $\mathcal{R}$ -cocomplete pseudofunctor  $\mathbb{M} : \mathcal{C}^{\text{op}} \rightarrow \text{Cat}$  which preserves finite biproducts (in the bicategorical sense). We define the bicategory of objective  $\mathcal{R}$ -Mackey functors by  $\text{OMky}_{\mathcal{R}}\mathcal{C} = \text{fpCAT}_{\mathcal{R}}\mathcal{C}$ . By Proposition 6.7,*

$$\text{fpCATSpn}_{\mathcal{R}}\mathcal{C} \sim \text{OMky}_{\mathcal{R}}\mathcal{C} . \quad (7.29)$$

<sup>3</sup>recalling Definition 2.2 although now we have  $\mathbb{E}$  instead of  $\mathbb{C}$  since we have  $\mathcal{E}$  instead of  $\mathcal{C}$ .

When  $\mathcal{R} = \mathcal{C}$ , we drop the subscripts  $\mathcal{C}$  and refer to objective Mackey functors.

**Example 7.2.** In the terminology of Definition 2.2, the object  $\mathbb{C}_{\mathcal{R}} \in \text{CAT}_{\mathcal{R}}\mathcal{C}$  is an objective  $\mathcal{R}$ -Mackey functor and, as suggested by the case  $\mathcal{C} = \text{set}^G$  of finite sets acted on by a finite group  $G$ , is called the objective  $\mathcal{R}$ -Burnside functor.

**Example 7.3.** In the terminology of Proposition 1.8, if  $\mathbb{Z}$  is an objective  $\mathcal{R}$ -Mackey functor, so is  $[\mathbb{Y}, \mathbb{Z}]_{\mathcal{R}}$ . For, we have

$$\begin{aligned} [\mathbb{Y}, \mathbb{Z}]_{\mathcal{R}}(W + W') &\simeq \text{CAT}_{\mathcal{R}}\mathcal{E}(\mathbb{Y}, \mathbb{Z}(- \times (W + W'))) \\ &\simeq \text{CAT}_{\mathcal{R}}\mathcal{E}(\mathbb{Y}, \mathbb{Z}(- \times W) \times \mathbb{Z}(- \times W')) \\ &\simeq \text{CAT}_{\mathcal{R}}\mathcal{E}(\mathbb{Y}, \mathbb{Z}(- \times W)) \times \text{CAT}_{\mathcal{R}}\mathcal{E}(\mathbb{Y}, \mathbb{Z}(- \times W')) \\ &\simeq [\mathbb{Y}, \mathbb{Z}]_{\mathcal{R}}W \times [\mathbb{Y}, \mathbb{Z}]_{\mathcal{R}}W' . \end{aligned}$$

(We are using the last sentence of Corollary 2.6 in the third step.)

As in Example 6.8, we consider the symmetric closed monoidal bicategory  $\text{Cat}_+$  of categories with finite coproducts and finite-coproduct-preserving functors; the internal hom  $[\mathcal{A}, \mathcal{B}]_+$  of  $\text{Cat}_+$  is  $\text{Cat}_+(\mathcal{A}, \mathcal{B})$  with pointwise coproduct. Cartesian product  $A \times B$  of categories is a direct sum in  $\text{Cat}_+$  and the terminal category  $\mathbf{1}$  is also initial.

**Proposition 7.4.** Every objective  $\mathcal{R}$ -Mackey functor  $\mathbb{M} : \mathcal{C}^{\text{op}} \rightarrow \text{Cat}$  factors through the forgetful 2-functor  $\text{Cat}_+ \rightarrow \text{Cat}$  by a pseudofunctor also denoted by  $\mathbb{M}$ .

*Proof.* By assumption, the codiagonal  $\nabla : U + U \rightarrow U$  and unique  $! : 0 \rightarrow U$  are in  $\mathcal{R}$ . The diagonal of  $\mathbb{M}U$  is isomorphic to the composite

$$\mathbb{M}U \xrightarrow{\mathbb{M}\nabla} \mathbb{M}(U + U) \simeq \mathbb{M}U \times \mathbb{M}U$$

and so has a left adjoint by Proposition 1.4. Thus  $\mathbb{M}U$  has binary coproducts. Similarly,  $\mathbb{M}U \xrightarrow{\mathbb{M}!} \mathbb{M}0 \simeq \mathbf{1}$  has a left adjoint so  $\mathbb{M}U$  has an initial object. For  $f : U \rightarrow V$ , the functor  $\mathbb{M}f$  preserves these finite coproducts as can be seen by applying the CB property to the pullbacks

$$\begin{array}{ccc} U + U & \xrightarrow{\nabla} & U \\ f+f \downarrow & & \downarrow f \\ V + V & \xrightarrow{\nabla} & V \end{array} , \quad \begin{array}{ccc} 0 & \xrightarrow{!} & U \\ 1_0 \downarrow & & \downarrow f \\ 0 & \xrightarrow{!} & V \end{array}$$

in our  $\mathcal{R}$ -extensive bicategory  $\mathcal{C}$ . □



**Corollary 7.5.** *If  $\mathbb{M}$  is an objective  $\mathcal{R}$ -Mackey functor and  $U \xrightarrow{i} U+V \xleftarrow{j} V$  is a coproduct in  $\mathcal{C}$  then*

$$\begin{array}{ccccc} \mathbb{M}U & \xrightarrow{\mathbb{M}_*i} & \mathbb{M}(U+V) & \xrightarrow{\mathbb{M}j} & \mathbb{M}V \\ & \perp & & \top & \\ & \xleftarrow{\mathbb{M}i} & & \xleftarrow{\mathbb{M}_*j} & \end{array}$$

*exhibits a (bicategorical) direct sum in  $\text{Cat}_+$ .*

**Corollary 7.6.** *For all  $f : X' \rightarrow X$  in  $\mathcal{C}$ , the functor  $\mathbb{C}_{\mathcal{R}}f : \mathbb{C}_{\mathcal{R}}X \rightarrow \mathbb{C}_{\mathcal{R}}X'$  preserves finite coproducts and the pseudofunctor*

$$\mathcal{C}^{\text{op}} \times \mathcal{C} \xrightarrow{(-)_* \times (-)^*} (\text{Spn}_{\mathcal{R}})^{\text{op}} \times \text{Spn}_{\mathcal{R}} \xrightarrow{\text{Spn}_{\mathcal{R}}(-,-)} \text{Cat}$$

*factors through the inclusion  $\text{Cat}_+ \hookrightarrow \text{Cat}$  of the sub-2-category of categories with finite coproducts and functors which preserve them.*

**Remark 7.7.** *Example 6.8 told of a hom enrichment of  $\text{Spn}_{\mathcal{R}}\mathcal{C}$  in  $\text{Cat}_+$  in the ordinary sense. For  $\text{Spn}_{\mathcal{R}}\mathcal{C}$ , we have an example of enrichment in a skew base which is an objective version of a base observed in 2012 by Cockett and Lack in connection with left additive (= left linear over the ring of integers) categories; see page 1101 of [16]. What we notice is that the sub-2-category  $\text{Cat}_+$  of  $\text{Cat}$  has a skew closed structure [39] with internal hom of  $A, B \in \text{Cat}_+$  taken as the ordinary functor category  $[\mathcal{A}, \mathcal{B}]$  where the finite coproducts are pointwise; write  $\text{Cat}_{+\text{sk}}$  for  $\text{Cat}_+$  equipped with this skew-monoidal structure. Then Proposition 6.6 and Corollary 7.6 tell us that  $\text{Spn}_{\mathcal{R}}\mathcal{C}$  is a skew  $\text{Cat}_{+\text{sk}}$ -enrichment of  $\mathcal{C}$  in the sense of Campbell [7] where the enriched hom is  $\text{Spn}_{\mathcal{R}}\mathcal{C}(X, Y)$ .*

*In particular, from Theorem 6.10, we have that  $\mathcal{L}\text{Ply}_{\mathcal{R}}\mathcal{C}$  is a skew enrichment of  $\mathcal{L}\text{Spn}_{\mathcal{R}}\mathcal{C}$  in  $\text{Cat}_{+\text{sk}}$ .*

We end this section with a consequence of Proposition 6.5.

**Proposition 7.8.** *The  $\mathcal{R}$ -cocompletion pseudomonad  $\overrightarrow{(-)}$  on  $\text{CAT}\mathcal{C}$  restricts to a pseudomonad on the bicategory  $\text{fpCAT}\mathcal{C}$  of finite biproduct preserving pseudofunctors  $\mathcal{C}^{\text{op}} \rightarrow \text{Cat}$ . The Eilenberg-Moore pseudo-algebras are the objective  $\mathcal{R}$ -Mackey functors.*

## 8 A monoidal structure on $\text{OMky}_{\mathcal{R}}\mathcal{C}$

The construction  $[\mathbb{Y}, \mathbb{Z}]_{\mathcal{R}}$  defined and discussed in Proposition 1.8 and Example 7.3 provides a closed structure (in the sense of [14]) on the bicategory

$\text{OMky}_{\mathcal{R}}\mathcal{C}$ . The closed structure is symmetric monoidal: there is a symmetric tensor product  $\mathbb{M} \star \mathbb{N}$  defined by a pseudonatural equivalence

$$\text{OMky}_{\mathcal{R}}\mathcal{C}(\mathbb{M} \star \mathbb{N}, \mathbb{L}) \simeq \text{OMky}_{\mathcal{R}}\mathcal{C}(\mathbb{M}, [\mathbb{N}, \mathbb{L}]_{\mathcal{R}}) .$$

There is a bicategorically universal pseudonatural transformation

$$\omega_{\mathbb{M}, \mathbb{N}} : \mathbb{M} \times \mathbb{N} \rightarrow \mathbb{M} \star \mathbb{N}$$

which is  $\mathcal{R}$ -cocontinuous in each variable separately. The unit for this tensor product is the objective  $\mathcal{R}$ -Burnside functor  $\mathbb{C}_{\mathcal{R}}$  (see Example (7.2)); for

$$\begin{aligned} [\mathbb{C}_{\mathcal{R}}, \mathbb{L}]_{\mathcal{R}} &= \text{CAT}_{\mathcal{R}}\mathcal{C}(\vec{\mathbf{1}}, \mathbb{L}(- \times W)) \\ &\simeq \text{CAT}\mathcal{C}(\mathbf{1}, \mathbb{L}(- \times W)) \\ &\simeq \lim \mathbb{L}(- \times W) \\ &\simeq \mathbb{L}(\mathbf{1} \times W) \\ &\simeq \mathbb{L}W \end{aligned}$$

where the (bicategorical) limit is over the category  $\mathcal{C}^{\text{op}}$  which has the (bi-categorical) initial object  $\mathbf{1}$ .

The monoidales (pseudomonoids) for the convolution monoidal structure  $\mathbb{A} \star \mathbb{B}$  we call *objective  $\mathcal{R}$ -Green functors on  $\mathcal{C}$*  and write  $\text{OGrn}_{\mathcal{R}}\mathcal{C}$  for the bicategory they, with the strong monoidal morphisms, form. An objective  $\mathcal{R}$ -Green functor is *symmetric* when it is symmetric as a monoidale.

The tensor unit for a symmetric monoidal structure is always a symmetric monoidale. So the objective  $\mathcal{R}$ -Burnside functor  $\mathbb{C}_{\mathcal{R}}$  is an example of an objective  $\mathcal{R}$ -Green functor.

Let us now describe the symmetric monoidal structure obtained by transporting the one on  $\text{OMky}_{\mathcal{R}}\mathcal{C}$  across the biequivalence (7.29).

By Lemma 1.6, we see that finite product in  $\mathcal{C}$  supplies the bicategory  $\text{Spn}_{\mathcal{R}}\mathcal{C}$  with a symmetric monoidal structure. Using Proposition 7.4, we see that  $\text{Ps}_{\times}(\text{Spn}_{\mathcal{R}}\mathcal{C}, \text{Cat})$  is the  $\text{Cat}_+$ -enriched bicategory  $\text{Ps}(\text{Spn}_{\mathcal{R}}\mathcal{C}, \text{Cat}_+)_+$  of  $\text{Cat}_+$ -enriched pseudofunctors. The desired transported symmetric closed monoidal structure  $\mathbb{A} \star \mathbb{B}$  on  $\text{Ps}(\text{Spn}_{\mathcal{R}}\mathcal{C}, \text{Cat}_+)_+$  is the Day convolution structure characterized by the existence of pseudonatural equivalences

$$\begin{aligned} &\text{Ps}(\text{Spn}_{\mathcal{R}}\mathcal{C}, \text{Cat}_+)_+(\mathbb{A} \star \mathbb{B}, \mathbb{C}) \\ &\simeq \text{Ps}(\text{Spn}_{\mathcal{R}}\mathcal{C} \times \text{Spn}_{\mathcal{R}}\mathcal{C}, \text{Cat}_+)_+(\mathbb{A} - \otimes \mathbb{B} \sim, \mathbb{C}(- \times \sim)) \\ &\simeq \text{Ps}(\text{Spn}_{\mathcal{R}}\mathcal{C} \times \text{Spn}_{\mathcal{R}}\mathcal{C}, \text{Cat}_+)_+(\mathbb{A} -, [\mathbb{B} \sim, \mathbb{C}(- \times \sim)]_+) . \end{aligned}$$

## 9 An objective Mackey functor of Mackey functors

Take a lextensive category  $\mathcal{E}$  equipped with a protocalibration  $\mathcal{R}$  closed under finite coproducts. We will consider the bicategory  $\text{Spn}_{\mathcal{R}}\mathcal{E}$  in the notation of Section 3 and will write  $\text{clSpn}_{\mathcal{R}}\mathcal{E}$  for the classifying category of  $\text{Spn}_{\mathcal{R}}\mathcal{E}$ ; the morphisms are isomorphism classes of spans (see [2]). We write  $\text{Spn}\mathcal{E}$  and  $\text{clSpn}\mathcal{E}$  for  $\text{clSpn}_{\mathcal{R}}\mathcal{E}$  and  $\text{clSpn}_{\mathcal{R}}\mathcal{E}$  in the case where  $\mathcal{R} = \mathcal{E}$ .

As we have pointed out, the bicategory  $\text{Spn}_{\mathcal{R}}\mathcal{E}$  is enriched in the closed symmetric monoidal bicategory  $\text{Cat}_+$  of categories with finite coproducts and finite-coproduct-preserving functors. The symmetric monoidal structure on the bicategory  $\text{Spn}_{\mathcal{R}}\mathcal{E}$  is  $\text{Cat}_+$ -enriched. The more abstract category  $\text{clSpn}_{\mathcal{R}}\mathcal{E}$ , along with its symmetric monoidal structure, is enriched in the usual symmetric closed monoidal category  $\text{CMon}$  of commutative monoids.

An  $\mathcal{R}$ -Mackey functor on  $\mathcal{E}$  is a  $\text{CMon}$ -enriched (or equally, a finite-product-preserving) functor

$$M : (\text{clSpn}_{\mathcal{R}}\mathcal{E})^{\text{op}} \rightarrow \text{CMon} .$$

This is consistent with Lindner [27] who looked at the case  $\mathcal{R} = \mathcal{E}$ .

**Example 9.1.** *i. Let  $G$  be a finite group, let  $\mathcal{E} = [G, \text{set}]$  be the category of finite  $G$ -sets, and let  $R : G \rightarrow \text{vect}$  be a finite-dimensional representation of  $G$ . Define  $M : \text{clSpn}\mathcal{E} \rightarrow \text{CMon}$  by  $MX = \text{Set}^G(X, R)$  and, for  $\tau : X \rightarrow R$  and  $y \in Y$ ,*

$$M(X \xrightarrow{[u, S, v]} Y)(X \xrightarrow{\tau} R)(y) = \sum_{v(s)=y} \tau(u(s)) .$$

*ii. There is the Burnside functor  $E_{\mathcal{R}} : (\text{clSpn}_{\mathcal{R}}\mathcal{E})^{\text{op}} \rightarrow \text{CMon}$  where*

$$E(X) = (\text{clSpn}_{\mathcal{R}}\mathcal{E})(X, 1) = \mathbb{E}_{\mathcal{R}}(X) \cong$$

*is the set of isomorphism classes of the slice category  $\mathcal{E}/_X$  made into a commutative monoid with addition induced by coproduct.*

Put

$$\text{Mky}_{\mathcal{R}}\mathcal{E} = [(\text{clSpn}_{\mathcal{R}}\mathcal{E})^{\text{op}}, \text{CMon}]_+ ,$$

the  $\text{CMon}$ -enriched functor category. It is a monoidal  $\text{CMon}$ -enriched category by Day convolution [12]. The monoids for this convolution structure are  $\mathcal{R}$ -Green functors on  $\mathcal{E}$ ; see [29] for more details in the case  $\mathcal{R} = \mathcal{E}$ .

We write  $\text{Grn}_{\mathcal{R}}\mathcal{E}$  for the  $\text{CMon}$ -enriched category of Green functors. The Burnside functor  $E_{\mathcal{R}}$  is an example.

Let  $\text{LLxt}$  denote the 2-category whose objects are pairs  $(\mathcal{A}, \mathcal{O})$  consisting of a lextensive category  $\mathcal{A}$  and a protocalibration  $\mathcal{O}$  closed under finite coproducts, and, whose morphisms  $F : (\mathcal{A}, \mathcal{O}) \rightarrow (\mathcal{A}', \mathcal{O}')$  are functors  $F : \mathcal{A} \rightarrow \mathcal{A}'$  preserving pullback and finite coproducts, and taking  $\mathcal{O}$  to  $\mathcal{O}'$ . Then we have a pseudofunctor

$$\text{Mky}_{\mathcal{R}} : \text{RLxt} \rightarrow \text{Cat}_+$$

taking  $(\mathcal{A}, \mathcal{O})$  to  $\text{Mky}_{\mathcal{O}}\mathcal{A}$  regarded as a category with finite coproducts.

It is useful to notice that the objective Burnside functor  $\mathbb{E}_{\mathcal{R}} : \mathcal{E}^{\text{op}} \rightarrow \text{Cat}_+$  lands in the category  $\text{RLxt}$  over  $\text{Cat}_+$  by equipping each lextensive category  $\mathbb{E}_{\mathcal{R}}(X) (\subseteq \mathcal{E}_{/X})$  with the protocalibration  $\mathcal{R}_{/X}$ .

**Proposition 9.2.** *The composite pseudofunctor*

$$\text{Mky}_{\mathcal{R}\mathcal{R}} := \mathcal{E}^{\text{op}} \xrightarrow{\mathbb{E}_{\mathcal{R}}} \text{RLxt} \xrightarrow{\text{Mky}_{\mathcal{R}}} \text{Cat}_+$$

*becomes a symmetric objective  $\mathcal{R}$ -Green functor when equipped with the multiplication induced by the finite-coproduct-preserving-in-each-variable functors*

$$\text{Mky}_{\mathcal{R}}\mathbb{E}_{\mathcal{R}}X \times \text{Mky}_{\mathcal{R}}\mathbb{E}_{\mathcal{R}}Y \rightarrow \text{Mky}_{\mathcal{R}}\mathbb{E}_{\mathcal{R}}(X \times Y)$$

*taking  $(M, N)$  to  $M \boxtimes N$  where*

$$(M \boxtimes N)(S \xrightarrow{(a,b)} X \times Y) = M(S \xrightarrow{a} X) \otimes N(X \xrightarrow{b} Y).$$

**Remark 9.3.** *In much of what we have just said the values of an objective Mackey functor could land in any bicategory  $\mathcal{K}$  with homs enriched in  $\text{Cat}_+$  rather than in  $\text{Cat}_+$  itself. That is, we can define a  $\mathcal{K}$ -valued objective Mackey functor on  $\mathcal{E}$  to be an  $\mathcal{R}$ -cocomplete  $\text{Cat}_+$ -enriched pseudofunctor  $\mathbb{M} : \mathcal{E}^{\text{op}} \rightarrow \mathcal{K}$ . We write  $\text{OMky}_{\mathcal{R}}(\mathcal{E}, \mathcal{K})$  for the bicategory of these. For the convolution tensor product to exist on this bicategory, we require  $\mathcal{K}$  to be appropriately cocomplete. We could do a similar thing in the abstract case by replacing  $\text{CMon}$  by a general  $\text{CMon}$ -enriched category with finite coproducts.*

## 10 Objective Tambara functors

Suppose  $\mathcal{E}$  is finitely complete and the pair  $(\mathcal{L}, \mathcal{R})$  satisfies the hypotheses of Lemma 6.9.

**Definition 10.1.** An objective  $(\mathcal{L}, \mathcal{R})$ -Tambara functor on  $\mathcal{E}$  is a finite biproduct preserving pseudofunctor  $\mathbb{T} : \mathcal{E}^{\text{op}} \rightarrow \text{Cat}$  which is in  ${}_{\mathcal{L}}\text{CAT}_{\mathcal{R}}\mathcal{E}$ . We define the bicategory of objective  $(\mathcal{L}, \mathcal{R})$ -Tambara functors by

$${}_{\mathcal{L}}\text{Tmb}_{\mathcal{R}}\mathcal{E} := \text{fp } {}_{\mathcal{L}}\text{CAT}_{\mathcal{R}}\mathcal{E} .$$

By Proposition 6.7 and Theorem 5.1,

$$\text{fpCAT}(({}_{\mathcal{L}}\text{Ply}_{\mathcal{R}}\mathcal{E})^{\text{op}}) \sim {}_{\mathcal{L}}\text{Tmb}_{\mathcal{R}}\mathcal{E} . \quad (10.30)$$

We write  $\mathbb{T}_{\text{pl}} : \text{Ply}_{\mathcal{E}} \rightarrow \text{Cat}$  for the inverse image of  $\mathbb{T}$  under this biequivalence.

**Lemma 10.2.**  $\mathcal{E}$  is  $\mathcal{L}$ -extensive.

*Proof.* The inclusion  $\mathbb{E}_{\mathcal{L}} \rightarrow \mathbb{E}_{\mathcal{R}}$  is componentwise fully faithful and the  $\mathcal{R}$ -extensivity equivalences restrict to  $\mathbb{E}_{\mathcal{L}}$ .  $\square$

Therefore Proposition 6.5 applies to both  $\mathcal{R}$  and  $\mathcal{L}$  yielding:

**Corollary 10.3.** If  $\mathbb{X} \in \text{CAT}$  preserves biproducts then so do  $\overleftarrow{\mathbb{X}}$  and  $\overrightarrow{\mathbb{X}}$ .

**Proposition 10.4.** The composite pseudomonad  $\overrightarrow{(-)}$  on  $\text{CAT}\mathcal{E}$  (see Section 4) restricts to a pseudomonad on the bicategory  $\text{fpCAT}\mathcal{E}$  of finite product preserving pseudofunctors  $\mathcal{E}^{\text{op}} \rightarrow \text{Cat}$ . The pseudo-algebras are the objective  $(\mathcal{L}, \mathcal{R})$ -Tambara functors.

An  $(\mathcal{L}, \mathcal{R})$ -Tambara functor on  $\mathcal{E}$  is a finite-product-preserving functor

$$T : \text{cl } {}_{\mathcal{L}}\text{Ply}_{\mathcal{R}}\mathcal{E} \rightarrow \text{CMon} .$$

This concept, for the case  $\mathcal{L} = \mathcal{R} = \mathcal{E} = G\text{-set}$ , arose in the paper [42] by Tambara; also see [28, 22, 9] for more general  $\mathcal{L}$  and  $\mathcal{R}$ .

---

## References

- [1] Paul Balmer and Ivo Dell’Ambrogio, *Mackey 2-Functors and Mackey 2-Motives*, EMS Monographs in Math. (2020).
- [2] Jean Bénabou, *Introduction to bicategories*, Lecture Notes in Mathematics **47** (Springer-Verlag, 1967) 1–77.

- [3] Jean Bénabou, *Théories relatives à un corpus*, Comptes Rendus de l'Académie des Sciences, Paris **281** (Série A, 17 novembre 1975) 831–834.
- [4] Jean Bénabou, *Fibrations petites et localement petites*, Comptes Rendus de l'Académie des Sciences, Paris **281** (Série A, 24 novembre 1975) 897–900.
- [5] A. J. Blumberg and M. A. Hill, *Incomplete Tambara functors*, Algebraic and Geometric Topology **18(2)** (2018) 723–766.
- [6] A. J. Blumberg and M. A. Hill, *Bi-incomplete Tambara functors*, London Mathematical Society Lecture Note Series **474** (CUP, 2022) 276–313.
- [7] Alexander Campbell, *Skew-enriched categories*, Applied Categorical Structures **26** (2018) 597–615.
- [8] A. Carboni, G.M. Kelly, R.F.C. Walters and R.J. Wood, *Cartesian bicategories II*, Theory and Applications of Categories **19** (2007) 93–124.
- [9] David Chan, *Bi-incomplete Tambara functors as  $\mathcal{O}$ -commutative monoids*, Tunisian Journal of Mathematics **6(1)** (2024) 1–47.
- [10] François Conduché, *Au sujet de l'existence d'adjoints à droite aux foncteurs 'image réciproque' dans la catégorie des catégories*, Comptes Rendus de l'Académie des Sciences, Paris **275** Série A (1972) 891–894.
- [11] Louis Crane and Igor B. Frenkel, *Four-dimensional topological quantum field theory, Hopf categories, and the canonical bases*, Journal of Mathematical Physics **35(10)** (1994) 5136–5154.
- [12] Brian J. Day, *On closed categories of functors*, Lecture Notes in Mathematics **137** (Springer, Berlin, 1970) 1–38.
- [13] Brian J. Day, *Limit spaces and closed span categories*, Lecture Notes in Mathematics **420** (Springer, Berlin, 1974) 65–74.
- [14] Brian Day and Ross Street, *Monoidal bicategories and Hopf algebroids*, Advances in Mathematics **129** (1997) 99–157.
- [15] N. Gambino and J. Kock, *Polynomial functors and polynomial monads*, Mathematical Proceedings Cambridge Philosophical Society **154** (2013) 153–192.

- [16] Richard Garner and Jean-Simon Lemay, *Cartesian differential categories as skew enriched categories*, Applied Categorical Structures **29** (2021) 1099–1150; <https://doi.org/10.1007/s10485-021-09649-7>.
- [17] Richard Garner and Daniel Schäppi, *When coproducts are biproducts*, Mathematical Proceedings Cambridge Philosophical Society **161** (2016) 47–51.
- [18] J. P. C. Greenlees and J. P. May, *Localization and completion theorems for MU-module spectra*, Annals of Mathematics (2) **146(3)** (1997) 509–544.
- [19] M. A. Hill and M. J. Hopkins, *Equivariant symmetric monoidal structures*, Preprint (2016); arXiv:1610.03114 [math.AT]
- [20] M. A. Hill, M. J. Hopkins, and D. C. Ravenel, *On the nonexistence of elements of Kervaire invariant one*, Annals of Mathematics (2) **184(1)** (2016) 1–262.
- [21] R. Houston, *Finite products are biproducts in a compact closed category*, Preprint (2006); arXiv:math.CT/0604542v1
- [22] Rolf Hoyer, *Two topics in stable homotopy theory*, (Ph.D. thesis, University of Chicago 2014) <https://www.proquest.com/docview/1559962111>.
- [23] Stefan Jackowski and James McClure, *Homotopy decomposition of classifying spaces via elementary abelian subgroups*, Topology **31(1)** (1992) 113–132.
- [24] P. T. Johnstone, R. Paré, R. D. Rosebrugh, D. Schumacher, R. J. Wood and G. C. Wraith, *Indexed Categories and Their Applications*, Lecture Notes in Mathematics **661** (1978).
- [25] Anders Kock, *Monads for which structures are adjoint to units*, Århus University Preprint Series **35** (1972-73) 1–15.
- [26] Anders Kock, *Monads for which structures are adjoint to units*, Journal of Pure and Applied Algebra **104** (1995) 41–59.
- [27] Harald Lindner, *A remark on Mackey functors*, Manuscripta Math. **18** (1976), 273–278.

- [28] Kristen Luise Mazur, *On the Structure of Mackey Functors and Tambara Functors*, (Ph.D. thesis, University of Virginia 2013) <https://www.proquest.com/docview/1445384784>.
- [29] Elango Panchadcharam and Ross Street, *Mackey functors on compact closed categories*, Journal of Homotopy and Related Structures **2(2)** (2007) 261–293.
- [30] Stephen H. Schanuel, *Objective number theory and the retract chain condition*, Journal of Pure and Applied Algebra **154** (2000) 295–298.
- [31] Dietmar Schumacher and Ross Street, *Some parametrized categorical concepts*, Communications in Algebra **16** (1988) 2313–2347.
- [32] Ross Street, *Two constructions on lax functors*, Cahiers de topologie et géométrie différentielle **13** (1972) 217–264.
- [33] Ross Street, *Fibrations and Yoneda’s lemma in a 2-category*, Lecture Notes in Mathematics **420** (1974) 104–133.
- [34] Ross Street, *Cosmoi of internal categories*, Transactions American Math. Soc. **258** (1980) 271–318.
- [35] Ross Street, *Fibrations in bicategories*, Cahiers de topologie et géométrie différentielle **21** (1980) 111–160.
- [36] Ross Street, *Conspectus of variable categories*, Journal of Pure and Applied Algebra **21** (1981) 307–338.
- [37] Ross Street, *Cauchy characterization of enriched categories*, Reprints in Theory and Applications of Categories **4** (2004) 1–16.
- [38] Ross Street, *Monoidal categories in, and linking, geometry and algebra*, Bulletin of the Belgian Mathematical Society – Simon Stevin **19(5)** (2012) 769–821.
- [39] Ross Street, *Skew-closed categories*, Journal of Pure and Applied Algebra **217(6)** (2013) 973–988.
- [40] Ross Street, *Polynomials as spans*, Cahiers de topologie et géométrie différentielle catégoriques **61(2)** (2020) 113–153.
- [41] Ross Street and Dominic Verity, *The comprehensive factorization and torsors*, Theory and Applications of Categories **23(3)** (2010) 42–75.



- [42] Daisuke Tambara, *On multiplicative transfer*, Communications in Algebra **21(4)** (1993) 1393–1420.
- [43] John Ullman, *Tambara functors and commutative ring spectra*, (preprint 2013) <http://www.arxiv.org/pdf/1304.4912v1>.
- [44] Charles Walker, *Universal properties of bicategories of polynomials*, Journal of Pure and Applied Algebra **223** (2019) 3722–3777.
- [45] Mark Weber, *Polynomials in categories with pullbacks*, Theory and Applications of Categories **30(16)** (2015) 533–598.
- [46] V. Zöberlein, *Doktrinen auf 2-Kategorien*, Manuscript (Math. Inst. Univ. Zürich, 1973).