

Microphysics of Circumgalactic Turbulence Probed by Fast Radio Bursts and Quasars

STELLA KOCH OCKER ^{1,2} MANDY C. CHEN ^{1,2} S. PENG OH ³ AND PRATEEK SHARMA ⁴

¹*Cahill Center for Astronomy and Astrophysics, California Institute of Technology, Pasadena, CA 91125, USA*

²*Observatories of the Carnegie Institution for Science, Pasadena, CA 91101, USA*

³*Department of Physics, University of California, Santa Barbara, CA 93106, USA*

⁴*Department of Physics, Indian Institute of Science, Bengaluru 560012, India*

ABSTRACT

The circumgalactic medium (CGM) is poorly constrained at the sub-parsec scales relevant to turbulent energy dissipation and regulation of multi-phase structure. Fast radio bursts (FRBs) are sensitive to small-scale plasma density fluctuations, which can induce multipath propagation (scattering). The amount of scattering depends on the density fluctuation spectrum, including its amplitude C_n^2 , spectral index β , and dissipation scale l_i . We use quasar observations of CGM turbulence at \gtrsim pc scales to infer C_n^2 , finding it to be $10^{-16} \lesssim C_n^2 \lesssim 10^{-9} \text{ m}^{-20/3}$ for hot ($T > 10^6$ K) gas and $10^{-9} \lesssim C_n^2 \lesssim 10^{-5} \text{ m}^{-20/3}$ for cool ($10^4 \lesssim T \lesssim 10^5$ K) gas, depending on the gas sound speed and density. These values of C_n^2 are much smaller than those inferred in the interstellar medium at similar physical scales. For most FRB sightline geometries, the scattering delays from the CGM are negligible ($\ll 10 \mu\text{s}$ at 1 GHz), but are more detectable from the cool gas as radio scintillation. Joint quasar-FRB observations of individual galaxies can yield lower limits on l_i , even if the CGM is not a significant scattering site. An initial comparison between quasar and FRB observations (albeit for different systems) suggests $l_i \gtrsim 200$ km in $\sim 10^4$ K gas in order for the quasar and FRB constraints to be consistent. If a foreground CGM is completely ruled out as a source of scattering along an FRB sightline then l_i may be comparable to the smallest cloud sizes (\lesssim pc) inferred from photoionization modeling of quasar absorption lines.

Keywords: Radio bursts – Circumgalactic medium – Turbulence – Scattering

1. INTRODUCTION

The flow of gas between intergalactic and interstellar media through the circumgalactic medium (CGM) fuels galaxy formation and subsequent evolution. Turbulence is fundamental to a range of ongoing processes in the CGM: it moderates heating, provides pressure support, generates magnetic dynamos, yields density fluctuations that mediate cooling, and mixes dust and metals, leading to CGM enrichment (for a review, see [Faucher-Giguère & Oh 2023](#)). While there is abundant evidence that the CGM is multi-phase, based primarily on high areal covering fractions of cool/warm ($\sim 10^4 - 10^5$ K) gas detected in large samples of quasar absorption lines spanning a range of halo sizes and redshifts ([Tumlinson et al. 2017](#); [Rudie et al. 2019](#)), the mass budget, origin, and regulation of this cooler gas by the surrounding hot ($\gtrsim 10^6$ K) medium, which is itself poorly constrained, remain uncertain. Turbulence likely plays a key role in cou-

pling and structuring these different phases ([Ji et al. 2019](#); [Fielding et al. 2020](#); [Tan et al. 2021](#); [Gronke et al. 2022](#); [Mohapatra et al. 2022a, 2023](#)).

Observational constraints from integral field spectroscopy (IFS) of emission around bright quasars ([Chen et al. 2023b, 2024, 2025](#)) and nonthermal broadening of quasar absorption lines by foreground gas ([Rudie et al. 2019](#); [Qu et al. 2022](#); [Chen et al. 2023a](#)) indicate that velocity fluctuations in the CGM are broadly consistent with subsonic Kolmogorov turbulence on roughly pc to 10 kpc scales in $10^4 - 10^5$ K gas ([Chen et al. 2023a](#)) and on $\approx 1 - 100$ kpc scales in $\gtrsim 10^6$ K gas (indirectly inferred through cool nebular tracers; [Chen et al. 2023b, 2024](#)), with evidence for injection on scales $\sim 10 - 100$ kpc depending on the system and its local environment ([Chen et al. 2025](#)). However, CGM turbulence remains poorly constrained at the small (\ll pc) spatial scales relevant to energy dissipation and the regulation of multi-phase structure (e.g. [Hummels et al. 2019](#); [Nelson et al. 2020](#); [Butsky et al. 2024](#)).

Fast radio bursts (FRBs) are one of the few astrophysical probes with the potential to constrain microscale physics in the CGM ([Vedantham & Phinney 2019](#); [Jow et al. 2024](#)).

Like other compact radio sources (e.g. pulsars), FRBs undergo multipath propagation, or scattering, through density fluctuations along the line-of-sight (LOS), and the characteristics of the scattering, including its frequency dependence and impact on the burst intensity profile, are related to the density fluctuation power spectrum (e.g. Cordes et al. 1986; Rickett 1990). FRB intersections through foreground halos are common; in addition to all FRBs probing the Galactic halo and a significant fraction probing halos in the Local Volume (Connor & Ravi 2022), intermediate-redshift halo intersections grow with increasing FRB source redshift (Macquart & Koay 2013), such that most FRBs at $z \gtrsim 0.5$ are expected to pass within twice the virial radius of multiple Milky Way-mass galaxies (Ocker et al. 2022a). As we will discuss, FRB scattering generically probes density fluctuations at \lesssim au scales in the CGM.

Placing meaningful constraints on the CGM from FRB scattering requires understanding the relative scattering contributions (or “scattering budget”) of all media along the LOS, including host galaxies and the Galactic interstellar medium (ISM). For FRBs with precise scattering budgets, scattering appears to predominantly arise from the ISMs of host galaxies and the Milky Way (Ocker et al. 2022b; Cordes et al. 2022; Sammons et al. 2022), with observational limits on CGM scattering at the $\lesssim 10$ s of μ s level at 1 GHz (Prochaska et al. 2019; Connor et al. 2020; Ocker et al. 2021a). These scattering upper limits suggest that the scattering strength (and hence the strength of turbulence) is over an order of magnitude smaller in the CGM than in the Galactic ISM (Ocker et al. 2021a). However, scattering from the CGM may increase for FRBs with multiple foreground halos (e.g. Faber et al. 2024), and if the strength of CGM turbulence evolves as a function of galaxy properties and/or cosmic time (Ocker et al. 2022a).

Theoretical predictions of CGM scattering are essential to the interpretation of FRB scattering budgets and their subsequent use as foreground probes. Previous theoretical examination of FRB scattering in the CGM has focused on simple analytic prescriptions for the McCourt et al. (2018) “misty” CGM model (Vedantham & Phinney 2019; Jow et al. 2024), and predicted more scattering from the CGM than is actually observed (Ocker et al. 2021a; Jow et al. 2024). In this paper, we leverage the growing number of observations of CGM turbulence in both emission and absorption to empirically predict how much scattering may occur in the CGM. While line emission and absorption probe disparate spatial scales from FRB radio wave scattering, it is possible that CGM turbulence extends to sub-parsec scales.

Observations of plasma turbulence across a wide range of conditions, including in situ measurements of the solar wind (Alexandrova et al. 2008, 2012) and very local ISM (Lee & Lee 2019) and remote measurements of the diffuse

ISM (Spangler & Gwinn 1990; Rickett et al. 2009; Geiger et al. 2024), indicate that dissipation occurs near or below the plasma kinetic scales, as expected for magnetized (and anisotropic) plasma turbulence (e.g. Lithwick & Goldreich 2001). X-ray observations of the intracluster medium similarly indicate that density fluctuations persist down to scales well below the collisional mean free path (Zhuravleva et al. 2019). Chromatic scattering of pulsars and masers is consistent with a dissipation scale $\sim 100 - 1000$ km in the ionized ISM (Spangler & Gwinn 1990; Rickett et al. 2009; Geiger et al. 2024), comparable to the ion inertial scale $\lambda_i = V_A/\Omega_i \approx 230 \text{ km}(n_e/\text{cm}^{-3})^{-1/2}$, where V_A is the Alfvén speed, Ω_i is the ion cyclotron frequency, and the latter equality is for an ionized hydrogen plasma with electron density n_e . In the CGM, $n_e \sim 10^{-4} - 10^{-2} \text{ cm}^{-3}$ implies ion inertial scales $\lambda_i \sim 10^3 - 10^4$ km. Even for a conservatively small density $n_e \sim 10^{-5} \text{ cm}^{-3}$, $\lambda_i \sim 10^5$ km.

Characteristic values of the plasma kinetic scales for the hot and cool CGM are shown in Table 1, along with length scales relevant to FRB scattering. These kinetic scales relevant to dissipation are orders of magnitude smaller than scales considered in CGM simulations, which typically focus on hydrodynamic (non-magnetized) turbulence but without explicit dissipation. Moreover, for typical CGM temperatures and densities the dissipation scale of hydrodynamic turbulence is larger than the collisional mean free path (which is $\approx 5000 \text{ au} - 1 \text{ pc}$; Spitzer 1956; Sarazin 1986). If magnetic turbulence is present in the CGM at sub-parsec scales, then FRBs may offer the only means of constraining it.

This paper makes empirical predictions of FRB scattering in the CGM, based on observational data at larger spatial scales. In doing so, we provide a methodology for contextualizing FRB scattering measurements and their physical interpretation in the landscape of other CGM probes, and we demonstrate how quasar and FRB observations may be used in tandem to constrain CGM microphysics. Section 2 describes the theoretical formalisms relating velocity and density fluctuation spectra, which are used to translate between observational constraints from quasars and FRBs summarized in Section 3. The results of converting quasar-inferred velocity fluctuations into estimates of density fluctuations and subsequent FRB scattering are given in Section 4. Implications are discussed in Section 5.

2. THEORY RELATING DENSITY AND VELOCITY FLUCTUATION SPECTRA

In the inertial range of a turbulent cascade, both velocity and density fluctuations follow power-law wavenumber spec-

Table 1. Length Scales Relevant to FRB Scattering in the CGM

| | Hot CGM | Cool CGM | |
|-------------------------------|--------------------------------------|---------------------|---------|
| Injection (Outer) Scale l_o | 10 – 100 kpc | $\gtrsim 10$ pc (?) | |
| Multipath Scale r_{mp} | $10^{-3} - 10^{-4}$ pc | 0.01 – 1 pc | |
| Fresnel Scale r_F | au | au | |
| Diffraction Scale r_{diff} | 0.2 au | 10^5 km | |
| Dissipation (l_i) | Ion Inertial Length λ_i | 2×10^4 km | 2300 km |
| | Ion Gyroradius r_i | 9000 km | 900 km |
| | Electron Inertial Length λ_e | 500 km | 50 km |
| | Electron Gyroradius r_e | 200 km | 20 km |
| | Debye Length λ_D | 7 km | 0.07 km |

NOTE—Order-of-magnitude estimates of length scales relevant to FRB scattering in the hot CGM assume $n_e \sim 10^{-4} \text{ cm}^{-3}$, $T \sim 10^6$ K, and $B \sim \mu\text{G}$, and in the cool CGM $n_e \sim 10^{-2} \text{ cm}^{-3}$, $T \sim 10^4$ K, and $B \sim \mu\text{G}$. Length scales are defined as follows: The multipath scale is the diameter of the scattering disk at the scattering screen (Eq. 11); the Fresnel scale approximately divides refractive from diffractive optics (Eq. 9); and the diffractive scale is the scale at which the rms phase perturbations are unity (Eq. 10). Microscales relevant to turbulent dissipation are evaluated for an ionized hydrogen plasma based on the following definitions: $\lambda_i = V_A/\Omega_i$, $r_i = v_{th,i}/\Omega_i$, $\lambda_e = V_A/\Omega_e$, $r_e = v_{th,e}/\Omega_e$, and $\lambda_D = \sqrt{\epsilon_0 k_B T/n_e}$, where V_A is the Alfvén speed, $\Omega_{i/e}$ is the ion/electron cyclotron frequency, $v_{th,i/e}$ is the ion/electron thermal speed, ϵ_0 is the permittivity of free space, and k_B is the Boltzmann constant.

tra of the form:

$$\begin{aligned}
 P_{\delta v}(q) &= C_v^2 q^{-\beta}, \\
 P_{\delta n_e}(q) &= C_n^2 q^{-\beta}, \\
 (q_o < q < q_i)
 \end{aligned}$$

where C_v^2 and C_n^2 are the spectral amplitudes, the wavenumber $q = 2\pi/l$ for a length scale l , and the spectra extend over an inertial range of wavenumbers between an outer scale $l_o = 2\pi/q_o$ and an inner scale $l_i = 2\pi/q_i$. For an inner scale $l_i \ll l$, a length scale of interest, the density fluctuation variance at the scale l can be found by integrating $P_{\delta n_e}$, which gives

$$\begin{aligned}
 \langle \delta n_e^2 \rangle &= \frac{2(2\pi)^{4-\beta} C_n^2 l^{\beta-3}}{\beta-3} \quad (\beta > 3, l \gg l_i) \\
 &\approx 3(2\pi)^{1/3} C_n^2 l^{2/3} \quad (\beta = 11/3, l \gg l_i)
 \end{aligned} \quad (1)$$

where the second approximation is for a Kolmogorov spectral index. The velocity fluctuation variance has a similar relation to C_v^2 . For $\beta = 11/3$, $\langle \delta n_e^2 \rangle$ in cm^{-6} , and l in au,

$$C_n^2 \approx 6.41 \times 10^3 \text{ m}^{-20/3} \frac{\langle \delta n_e^2 \rangle}{\text{cm}^{-6}} \left(\frac{l}{\text{au}} \right)^{-2/3}. \quad (2)$$

The motivation for $l \sim \text{au}$ comes from the very small Fresnel scales below which diffractive scattering occurs, as we discuss in §3.2. We relate the density and velocity fluctuation

spectra assuming that the density fluctuations are a passive scalar, and we focus on the subsonic regime relevant to the CGM (Mach numbers $\mathcal{M} \lesssim 1$). The hot ($T \gtrsim 10^6$ K) gas is treated separately from the cool ($10^4 \lesssim T \lesssim 10^5$ K) gas, as follows.

2.1. Hot CGM

For the hot CGM, adiabatic simulations of isotropic, homogeneous, subsonic turbulence indicate that the root-mean-square (rms) density fluctuations are related to the Mach number squared (Mohapatra & Sharma 2019; Mohapatra et al. 2022a),

$$\delta n_e / \langle n_e \rangle = b \mathcal{M}^2, \quad (3)$$

where $\langle n_e \rangle$ is the average, volume-weighted hot gas density and $b \approx 0.3 - 1$ (hereafter we assume $b \approx 1$; Federrath et al. 2010; Konstantin et al. 2012). The Mach number can be constrained from measured velocity structure functions (VSFs) via comparison of VSF slopes across different VSF orders, given an assumed sound speed (Chen et al. 2023b, 2024; for analogous simulation analysis, see Mohapatra et al. 2022b). For a Kolmogorov spectrum, the rms density fluctuations have the same power-law index in length scale l as the rms velocity fluctuations, giving

$$\frac{\delta n_e}{\langle n_e \rangle} \approx 3 \times 10^{-5} \left(\frac{\mathcal{M}}{0.3} \right)^2 \left(\frac{l}{\text{au}} \right)^{1/3} \left(\frac{100 \text{ kpc}}{l_o} \right)^{1/3}, \quad (4)$$

where we have chosen values for \mathcal{M} and l_o representative of VSFs measured in the CGM (Chen et al. 2024). The density fluctuation amplitude C_n^2 can thus be expressed in terms of \mathcal{M} as

$$C_n^2 \approx 10^{-12} \text{ m}^{-20/3} \times \left(\frac{\langle n_e \rangle}{10^{-4} \text{ cm}^{-3}} \right)^2 \left(\frac{\mathcal{M}}{0.3} \right)^4 \left(\frac{100 \text{ kpc}}{l_o} \right)^{2/3}, \quad (5)$$

where we have combined Equations 2 and 4 and scaled to densities typical of the hot CGM. This $C_n^2 \sim 10^{-12} \text{ m}^{-20/3}$ is many orders of magnitude smaller than that typical in the Galactic ionized ISM, for which $\langle n_e \rangle \sim 0.1 \text{ cm}^{-3}$, $\mathcal{M} \sim 1$, and $l_o \sim 100$ pc yields $C_n^2 \sim 10^{-3} \text{ m}^{-20/3}$ (directly comparable to values inferred from observations of pulsar scattering and DM variations; Armstrong et al. 1995; Krishnakumar et al. 2015; Ocker et al. 2021b). Hence, the contribution of the hot CGM to scattering is expected to be negligible. Note, that for $\mathcal{M} > 1$, Equation 3 should be modified to scale linearly with \mathcal{M} (Mohapatra & Sharma 2019). A linear scaling between density fluctuations and Mach number also holds if the hot CGM is highly stratified with strong buoyant restoring forces (Zhuravleva et al. 2014; Mohapatra et al. 2021). Even with these modifications, the hot CGM will still be orders of magnitude away from producing a detectable signal.

2.2. Cool CGM

For the cool CGM, density fluctuations are expected to behave similarly to those in an isothermal turbulence cascade in which δn_e scales linearly with \mathcal{M} (Padoan et al. 1997; Konstandin et al. 2012; Federrath et al. 2021). The amplitudes of the velocity and density fluctuation spectra, C_v^2 and C_n^2 , can be related to each other by expressing \mathcal{M} in terms of the rms velocity fluctuations, which are related to C_v^2 in a form analogous to Equation 2, yielding

$$C_v^2 = \left(\frac{c_s}{b \langle n_e \rangle} \right)^2 C_n^2 \quad (6)$$

where b is the same constant as in Equation 3, $\langle n_e \rangle$ is the average cool gas electron density, and c_s is the sound speed (Simard & Ravi 2021). For c_s , $\langle n_e \rangle$, and C_v^2 in their typical units,

$$C_n^2 \approx 10^{-6} \text{ m}^{-20/3} \left(\frac{C_v^2}{10^{-9} \text{ km}^{4/3} \text{ s}^{-2}} \right) \times \left(\frac{\langle n_e \rangle}{10^{-2} \text{ cm}^{-3}} \right)^2 \left(\frac{c_s}{30 \text{ km s}^{-1}} \right)^{-2}. \quad (7)$$

The velocity spectral amplitude C_v^2 is constrained by measurements of nonthermal line widths, which can be used to infer $\langle \delta v^2 \rangle$. For nonthermal broadening dominated by turbulence, the nonthermal line width b_{NT} is related to the rms velocity fluctuations δv as $\delta v = \sqrt{3/2} b_{\text{NT}}$, and $\delta v \propto l^{1/3}$ for $\beta = 11/3$ (Chen et al. 2023a). Given δv at a specific length scale l , C_v^2 can be derived analogously to Equation 1:

$$\begin{aligned} \langle \delta v^2 \rangle &\approx 3(2\pi)^{1/3} C_v^2 l^{2/3} \\ &\approx 600 \text{ km}^2 \text{ s}^{-2} \left(\frac{C_v^2}{10^{-9} \text{ km}^{4/3} \text{ s}^{-2}} \right) \left(\frac{l}{\text{kpc}} \right)^{2/3}. \end{aligned} \quad (8)$$

For $\delta v \approx 20 \text{ km/s}$ at $l = 1 \text{ kpc}$, $C_v^2 \approx 10^{-9} \text{ km}^{4/3} \text{ s}^{-2}$, the fiducial value adopted in Equation 7. Note that inferring C_n^2 from velocity fluctuations always requires an assumption of the sound speed, which is implicitly used to constrain \mathcal{M} from measured VSFs. Moving forward, we assume that the density-Mach relations used above also apply to the volume-averaged density within a single phase.

3. CGM TURBULENCE OBSERVABLES

Observations of CGM turbulence are primarily at $\gg \text{pc}$ scales, from IFS of emission around bright quasars (Chen et al. 2023b, 2024, 2025) and nonthermal broadening of quasar absorption lines (Rudie et al. 2019; Qu et al. 2022; Chen et al. 2023a). Here we summarize key observational results from emission and absorption studies, followed by the FRB scattering observables that will be estimated and interpreted.

3.1. Emission and Absorption

IFS yields spatially resolved maps of emission line radial velocities, which constrain velocity fluctuations over spatial scales of $\approx 1 - 100 \text{ kpc}$ (Chen et al. 2023b, 2024, 2025). Nonthermal broadening of quasar absorption lines yields the sightline-integrated velocity dispersion, but associating the resulting velocity fluctuation amplitude with a specific spatial scale requires photoionization modeling of sometimes degenerate line components in order to identify discrete clouds and estimate their sizes (Chen et al. 2023a). These two techniques have been applied to galaxy samples spanning different redshifts, masses, and active galactic nuclei (AGN) characteristics.

We make use of three key results from the quasar studies described above:

1. CGM turbulence is broadly consistent with a Kolmogorov spectral index¹ (Chen et al. 2023b,a, 2024). We thus assume $\beta = 11/3$ for both hot and cool gas in subsequent analysis, and discuss the effect of modifying β on our results.
2. In emission, the velocity fluctuations are consistent with Mach numbers $\approx 0.2 - 1.8$. These velocities are inferred from emission by cool $T \sim 10^4 \text{ K}$ nebular gas, and are thought to reflect the motions of cool clumps entrained in hot $T > 10^6 \text{ K}$ gas (Chen et al. 2023b, 2024). The Mach number is inferred by comparing the velocity dispersion in the plane of the sky, σ_{pos} , and the hot gas sound speed, and is corroborated by the VSF slopes. Given that the majority of systems appear to be subsonic, we restrict our analysis to Mach numbers ≤ 1 , and we focus on densities typical of the hot CGM in Milky Way-mass galaxies, $10^{-5} \leq n_e \leq 10^{-3} \text{ cm}^{-3}$ (e.g. Sharma et al. 2012; Miller & Bregman 2015; Singh et al. 2018; Voit 2019). While the hot CGM may have even higher densities in the core of the halo, we are primarily interested in FRB LOSs at $\sim 10 - 200 \text{ kpc}$ from the halo center, where observations are not contaminated by the ISM.
3. In absorption, nonthermal line broadening indicates a roughly Kolmogorov scaling relation between cloud size and nonthermal line width for $10^4 - 10^5 \text{ K}$ gas, where the cloud sizes probed are $\sim \text{pc} - 10 \text{ kpc}$, and the corresponding densities are $\sim 10^{-4} - 0.1 \text{ cm}^{-3}$. The fitted scaling relation gives the rms velocity fluctuations at 1 kpc , $\delta v(1 \text{ kpc}) \approx 22 \text{ km/s}$ (Chen et al. 2023a). The uncertainty in this anchor value of δv has

¹ While deviations from Kolmogorov turbulence are seen in some systems, it is unclear whether these deviations are bona fide and physical (Chen et al. 2024).

a miniscule effect on resulting estimates of C_v^2 when accounting for a range of possible densities and sound speeds, and so we leave δv at 1 kpc fixed to 22 km s^{-1} in subsequent analysis of the cool gas. Nonthermal broadening also yields estimates of the Mach number, which appears to be $\mathcal{M} < 1$ for a majority of absorbers (Rudie et al. 2019; Qu et al. 2022).

Based on the above results, we use the velocity fluctuations constrained in emission to estimate density fluctuations in the hot CGM, and fluctuations constrained in absorption for the cool CGM. Applying these results to our scaling relations between C_v^2 and C_n^2 (Section 2) assumes a one-to-one correspondence for density and velocity fluctuations across all scales, which may not necessarily be the case. We also note that these emission and absorption measurements trace different galaxy systems that are heterogeneous in mass, redshift, and AGN properties. There is initial evidence that turbulence may be more dominant in the CGM of quiescent galaxies than in star-forming galaxies and at $z \leq 1$, although the physical interpretation of this trend is unclear and larger samples are needed for a robust, uniform comparison (Qu et al. 2022). The inferred energy transfer rates per unit mass also appear to be $> 10\times$ larger in the CGM around luminous quasars than in typical L_* galaxies (Chen et al. 2023b, 2025). This difference may be related to both the differing gas phases probed and a more systematic difference in the galaxy properties like AGN activity. In future, direct comparisons between emission and/or absorption and FRB observations will preferably be performed within the same galaxy system.

3.2. FRB Scattering

A continuous spectrum of electron density fluctuations will produce a range of refractive and diffractive effects that are observable on different (and sometimes overlapping) time and radio frequency scales (Cordes et al. 1986). Refraction can include angle of arrival variations or image wandering, long-term (days to months) intensity modulations, and under certain conditions multiple imaging. Diffraction includes short-term (minutes to hours) intensity modulations, pulse broadening, and angular broadening. We use the term ‘‘scintillation’’ to refer to the generic phenomenon of chromatic intensity modulations caused by constructive and destructive interference of scattered radiation, which can be both refractive and diffractive in nature (Rickett 1990). The relevance of refractive and diffractive scattering is a function of spatial scale that depends on the shape, amplitude, and cutoff of the density fluctuation spectrum (Goodman & Narayan 1985; Cordes et al. 1986; Coles et al. 1987).

In general, diffractive effects dominate for Kolmogorov media when the inner scale is much smaller than the Fresnel scale, whereas if density fluctuations decline more strongly

at small scales (i.e., the power spectrum has a steeper power law scaling), refraction can dominate across a wide range of spatial scales. Modifications to the density fluctuation spectrum (e.g. local enhancements or shocks and energy injection at multiple scales) also affect the resulting radio scattering observables. For simplicity, we focus subsequent analysis on a statistically homogeneous medium with a single injection scale, and wherever a fiducial assumption of the spectral index β is needed we adopt $\beta = 11/3$. However, in principle the scattering observables treated here can be derived for a medium of arbitrary spectral index (see Appendix and for an in-depth treatment, Cordes et al. 1986).

3.2.1. Length Scales of Diffractive Scattering

Given that the majority of FRBs are only observed once, we focus on diffractive scattering effects that are routinely constrained within individual bursts, namely pulse broadening and scintillation. In the regime of diffractive scattering, radio waves undergo instantaneous multipath propagation, and the underlying plasma density fluctuations must extend to length scales significantly smaller than the Fresnel scale, r_F , which roughly separates the regimes of diffractive and refractive optics (see, however, Jow et al. 2023, for an in-depth treatment that accounts for the effect of lens strength in the refractive regime).

For a scattering medium (or screen/lens) at a cosmological redshift z_ℓ , the Fresnel scale is

$$r_F = \left(\frac{d_{l_o} d_{s_l} \lambda_{\text{obs}}}{2\pi d_{s_o} (1 + z_\ell)} \right)^{1/2} \quad (9)$$

where d_{l_o} , d_{s_l} , and d_{s_o} are angular diameter distances between the lens and observer, source and lens, and source and observer, respectively, and the observing wavelength λ_{obs} is a factor $(1 + z_\ell)$ smaller in the lens frame (Macquart & Koay 2013). Figure 1 shows r_F vs. z_ℓ for a range of source redshifts. For typical FRB redshifts ($z_s \lesssim 1$), r_F is at most 2 to 5 au, and even for larger z_s it does not exceed 10 au. The length scales relevant to diffractive scattering are thus orders of magnitude smaller than typical resolutions in numerical CGM simulations, and well below the spatial scales accessible to both absorption and emission line observations of the CGM.

The characteristic scattering angle, θ_d , is typically defined as $\theta_d = (d_{s_l}/d_{s_o})\lambda_{\text{obs}}/2\pi r_{\text{diff}}$, where the diffractive scale r_{diff} is the transverse separation in the observing plane across which there is an rms phase difference of 1 radian (equivalently, r_{diff} is the length scale for which the phase structure function is unity; see Rickett 1990 for a review). For plane waves (as appropriate for FRB sources at cosmological distances) and the regime where $r_{\text{diff}} > l_i$,

$$r_{\text{diff}} = \left[\frac{2^{2-\beta} \pi^2 r_e^2 \lambda_{\text{obs}}^2 \beta \text{SM} \Gamma(-\beta/2)}{(1 + z_\ell)^2 \Gamma(\beta/2)} \right]^{1/(2-\beta)}, \quad (10)$$




Figure 1. Fresnel scale vs. lens redshift (z_ℓ) for a range of FRB source redshifts (z_s). The Fresnel scale is evaluated using angular diameter distances (Equation 9). The diffractive scattering effects considered here arise from density fluctuations at length scales significantly smaller than the Fresnel scale.

where r_e is the classical electron radius, λ_{obs} is the observing wavelength ($\lambda_{\text{obs}} \approx 0.3$ m for $\nu = 1$ GHz), and $\text{SM} \approx C_n^2 L$ for a path length L through a screen of constant C_n^2 (Rickett et al. 1984; Cordes et al. 1986; Coles et al. 1987)². The diameter of the scattering cone at the scattering screen, also known as the multipath scale, is then

$$r_{\text{mp}} = 2d_{\text{lo}}\theta_d \quad (11)$$

(Cordes et al. 1986). The multipath scale defines the maximum length scale of the density fluctuations driving multipath propagation. All of our scattering expressions apply in the so-called strong scattering regime, for which $r_{\text{mp}} > r_F$. This regime applies for most of the conditions we are interested in; specific conditions under which this regime does not apply are discussed further in Section 4.

We will refer to the time delay between an undeflected ray and a ray scattered through angle θ_d as the diffractive scattering time, τ_d , which is distinct from the mean scattering time, τ , that will be used throughout the rest of the paper.

² Note that Equation 10 is appropriate for wavenumber defined as $q = 2\pi/l$ and applies to $2 < \beta < 4$. Cordes et al. (1986) also give expressions for r_{diff} (denoted Δx_d in that paper) for $4 < \beta < 6$. As in Cordes et al. (1986), we define r_{diff} as the $1/e$ half-width of the visibility function. As in Macquart & Koay (2013), we include a factor of $(1 + z_\ell)^{-2}$ in r_{diff} , but we note that their expression for r_{diff} differs from ours by a factor of $\pi^{1/(2-\beta)}$.

The diffractive scattering time is $\tau_d \propto (d_{\text{so}}d_{\text{lo}}/d_{\text{sl}})(\theta_d^2/2c)$. For a Gaussian scattered image, θ_d is the $1/e$ half-width of the image, and the corresponding time delay is the $1/e$ time of a scattered pulse that temporally decays as a one-sided exponential. This special case of a Gaussian scattered image is conventionally assumed when inferring scattering delays from observed pulse shapes.

However, in a turbulent medium with $\beta < 4$ and an inner scale $l_i \ll r_{\text{diff}}$, the scattered image is not Gaussian and instead has an extended halo that decays more slowly than a Gaussian, because the density fluctuations at scales $l \ll r_{\text{diff}}$ lead to scattering angles $\theta \sim \lambda/l \gg \lambda/r_{\text{diff}}$ (Lambert & Rickett 1999). In the time domain, the scattered pulse decays more slowly than an exponential, and the pulse has a significant amount of radiation at time delays $\gg \tau_d$. The mean scattering time of this pulse, τ , will be greater than the diffractive scattering time τ_d . Conversely, when $r_{\text{diff}} \ll l_i$ (or $\beta \geq 4$), the scattered image becomes Gaussian, the extended halo is absent, and $\tau = \tau_d$ (Rickett et al. 2009; Geiger et al. 2024).

In the CGM, inferred diffractive scattering times $\lesssim 10$ s of μs imply $r_{\text{diff}} \gtrsim 10^6$ km (Prochaska et al. 2019). If l_i is comparable to plasma kinetic scales, then we are in the regime where $l_i \ll r_{\text{diff}}$. In this regime, the scattered pulse will deviate from a one-sided exponential and include contributions from scattering angles $> \theta_d$. The relevant time delays $\tau > \tau_d$ hence yield information about density fluctuations extending down to the inner scale l_i .

3.2.2. Generic Ionized Cloudlet Model

We proceed by relating the mean scattering time to underlying density fluctuations using a generic ionized cloudlet model, in which the CGM is a statistically homogeneous medium consisting of ionized cloudlets with a volume filling factor f , internal cloudlet density n_{ec} , and internal density fluctuations following a Kolmogorov spectrum down to scales l_i smaller than the diffractive scale. The scattering strength of the medium is then recast in terms of a composite density fluctuation parameter (see Appendix)

$$\begin{aligned} \tilde{F} &= (\zeta \epsilon^2 / f) l_o^{3-\beta} l_i^{\beta-4} \\ &= (\zeta \epsilon^2 / f) (l_o^2 l_i)^{-1/3} \text{ for } \beta = 11/3, \end{aligned} \quad (12)$$

where $\zeta = \langle n_{ec}^2 \rangle / \langle n_{ec} \rangle^2$ is the density variation between cloudlets and $\epsilon^2 = \langle (\delta n_{ec})^2 \rangle / n_{ec}^2$ is the fractional density variance within a cloudlet (Cordes et al. 2016; Ocker et al. 2021a; Cordes et al. 2022).

This formulation has the advantage of yielding a simple relation between the mean scattering time τ_{obs} (observer frame) and the dispersion measure (DM) contributions of a given medium along the LOS,

$$\tau_{\text{obs}} \approx 48.03 \text{ ns } \tilde{F} G_{\text{scatt}} \text{DM}_\ell^2 \nu^{-4} (1 + z_\ell)^{-3} \quad (13)$$

where ν is the observing frequency, z_ℓ is the screen redshift, DM_ℓ is the screen DM (due to the gas phase under consideration) in its rest frame, and G_{scatt} is a geometric leverage factor that depends on the relative locations of the source, screen, and observer. For cosmological screens far from the observer and source,

$$G_{\text{scatt}} = 2d_{\text{sl}}d_{\text{lo}}/Ld_{\text{so}}, \quad (14)$$

for angular diameter distances defined as in Equation 9 and a path length through the screen L . The equivalent scintillation bandwidth, defined as the frequency width of the intensity autocorrelation function, is $\Delta\nu_{\text{d}} = C/(2\pi\tau)$, where C is a constant of order unity depending on the density fluctuation spectrum; for a uniform Kolmogorov medium filling the entire LOS, $C = 1.16$, whereas for a Kolmogorov thin screen $C = 0.96$ (Cordes & Rickett 1998). We adopt $C = 1$ in subsequent analysis, which has a negligible effect on our results.

Given that the majority of scattering times reported in the literature are $1/e$ times that assume a Gaussian scattered image, we require a means of converting the $1/e$ delay to the mean scattering time given by Equation 13. This conversion is accomplished through the dimensionless factor $A_\tau \equiv \tau_{\text{d}}/\tau$, which directly depends on the ratio l_i/r_{diff} (and therefore implicitly on the observing frequency, which changes the strength of scattering and hence r_{diff}). For a Kolmogorov spectrum and $\nu = 1$ GHz, $A_\tau \approx 1/6$ when $l_i/r_{\text{diff}} \ll 0.1$ (with little variation below this value), and $A_\tau \approx 0.7$ for $l_i/r_{\text{diff}} = 1$ (Cordes et al. 2022). In the CGM, $l_i \sim 10^3 - 10^4$ km and $r_{\text{diff}} \gtrsim 10^6$ km implying $l_i/r_{\text{diff}} \lesssim 10^{-3} - 10^{-2}$. We therefore adopt $A_\tau = 1/6$ when converting the $1/e$ delay τ_{d} to the mean scattering time, and all $1/e$ delays are scaled to 1 GHz assuming $\tau_{\text{d}} \propto \nu^{-4.4}$, as appropriate for $\beta = 11/3$. The mean scattering delay scales exactly as ν^{-4} , but A_τ depends on r_{diff} , and hence ν , such that $\tau_{\text{d}} \propto A_\tau \nu^{-4} \propto \nu^{-4.4}$, as expected.

For a volume-averaged density $\bar{n}_e = f\langle n_{ec} \rangle$, \tilde{F} is related to C_{n}^2 as

$$\tilde{F} \approx 0.5 (\text{pc}^2 \text{ km})^{-1/3} \left(\frac{C_{\text{n}}^2}{\text{m}^{-20/3}} \right) \left(\frac{\bar{n}_e}{\text{cm}^{-3}} \right)^{-2} \left(\frac{l_i}{\text{km}} \right)^{-1/3} \quad (15)$$

(see Appendix). Formally the dependence on \bar{n}_e cancels out when evaluating C_{n}^2 , which is proportional to \bar{n}_e^2 (Equation 5).

For typical Galactic pulsar sightlines probing the diffuse ISM, $C_{\text{n}}^2 \approx 10^{-3.5} \text{ m}^{-20/3}$, $l_i \approx 1000$ km, and $\bar{n}_e \approx 0.02 \text{ cm}^{-3}$, which yields $\tilde{F} \approx 0.04 (\text{pc}^2 \text{ km})^{-1/3}$. This estimate is entirely consistent with values of \tilde{F} inferred directly from pulse broadening and DM measurements in the disk (Ocker et al. 2021a). In the hot CGM, densities $\sim 10^{-4} \text{ cm}^{-3}$ yield $C_{\text{n}}^2 \sim 10^{-12} \text{ m}^{-20/3}$ (Equation 5), equivalent to $\tilde{F} \sim 10^{-5.3}$

$(\text{pc}^2 \text{ km})^{-1/3}$ for $l_i \sim 1000$ km. For the cool CGM, a more typical value of $C_{\text{n}}^2 \sim 10^{-6} \text{ m}^{-20/3}$ would yield $\tilde{F} \sim 10^{-3} (\text{pc}^2 \text{ km})^{-1/3}$ for the same inner scale. These estimates do not involve any assumptions about the underlying density fluctuations described by ζ , ϵ^2 , or f in this analysis; \tilde{F} is estimated purely based on C_{n}^2 and its relation to the velocity fluctuation spectrum.

4. ANALYSIS & RESULTS

The formalisms laid out in Sections 2-3 are used to convert velocity fluctuations into the density fluctuation amplitude C_{n}^2 and subsequent strength of FRB scattering. Figure 2 illustrates our approach. For the hot CGM, Mach numbers inferred from VSFs and projected velocity dispersion σ_{pos} (Chen et al. 2023b, 2024) are converted to C_{n}^2 for a range of possible n_e , and assuming fiducial values for the outer scale noted further below (Equation 5). For the cool CGM, nonthermal line widths indicate a typical velocity dispersion $\delta v \approx 22$ km/s at 1 kpc (Chen et al. 2023a), which is used to estimate C_v^2 (Equation 8), and subsequently C_{n}^2 for a range of sound speeds and densities (Equation 7). We then estimate \tilde{F} from C_{n}^2 for a range of possible dissipation scales l_i (Equation 15). The resulting constraints on C_{n}^2 and \tilde{F} are presented in Section 4.1, and are used to predict FRB scattering observables in Section 4.2. These predictions are compared to observed FRBs in Section 4.3. Prospects for constraining the dissipation scale of CGM turbulence are demonstrated in Sections 4.4-4.5.

4.1. Density Fluctuations in the Hot and Cool CGM

Figure 3 shows the density fluctuation spectral amplitude C_{n}^2 vs. electron density for Mach numbers inferred in the hot CGM and an outer scale $l_o = 100$ kpc. For a range of typical densities, C_{n}^2 in the hot CGM is five to twelve orders of magnitude smaller than values typical of the ionized ISM probed by Galactic pulsars. Figure 3 also shows the equivalent range of \tilde{F} vs. l_i . Depending on the outer scale and Mach number, \tilde{F} is between $\sim 10^{-8} (\text{pc}^2 \text{ km})^{-1/3}$ to $\sim 10^{-4} (\text{pc}^2 \text{ km})^{-1/3}$.

Constraints on C_{n}^2 and \tilde{F} for the cool gas probed in absorption are also shown in Figure 3. Here, C_{n}^2 is estimated from C_v^2 for sound speeds $15 \lesssim c_s \lesssim 50$ km/s (these are the sound speeds for $10^4 \lesssim T \lesssim 10^5$ K, the temperatures inferred from photoionization modeling; Chen et al. 2023a), and for densities between 10^{-3} and 0.1 cm^{-3} . While quasar absorption lines can constrain cloud densities $< 10^{-3} \text{ cm}^{-3}$, these smaller densities correspond to a minority of absorbers and to the largest cloud sizes (> 10 kpc), for which modeling uncertainties are exacerbated. Due to the higher densities expected in the cool gas, C_{n}^2 and \tilde{F} reach values comparable to those routinely inferred in the ionized ISM from pulsar scattering.

4.2. Scattering Predictions




Figure 2. Schematic view of the methods used to derive density fluctuations from velocity fluctuation measurements in the CGM and subsequently predict levels of FRB scattering. All equations referenced are in Sections 2-3. See Section 4 for further discussion.




Figure 3. Density fluctuation spectral amplitude C_n^2 vs. electron density (left) and density fluctuation parameter \tilde{F} vs. inner scale (right). For the hot CGM, C_n^2 is evaluated using Equation 5 for a range of Mach numbers inferred from IFS observations of the hot CGM in emission. The outer (injection) scale for the hot CGM is taken to be $l_o = 100$ kpc, and may be rescaled to different outer scales given $C_n^2 \propto l_o^{-2/3}$. For the cool CGM, C_n^2 is evaluated using Equation 6 for sound speeds corresponding to temperatures $10^4 \lesssim T \lesssim 10^5$ K. The fluctuation parameter \tilde{F} is evaluated from C_n^2 for a range of inner scales equivalent to the ion inertial scale at the densities chosen for each phase (Equation 15). The blue shaded regions show values typically inferred in the ionized ISM from Galactic pulsars; in the ISM, C_n^2 and \tilde{F} can be even larger than the ranges shown.




Figure 4. Mean pulse broadening time in the lens frame $\tau(1+z_\ell)^3$ (top) and scintillation bandwidth in the lens frame $\Delta\nu_d(1+z_\ell)^{-3}$ (bottom) vs. $DM_\ell^2 G_{\text{scatt}}$, a range of possible DMs and geometric configurations for the hot CGM of a halo intervening a LOS. Scattering parameters are estimated for Mach numbers typical of the hot CGM based on IFS velocity fluctuation measurements.

We use the \tilde{F} derived from quasar constraints to estimate the mean pulse broadening time τ and scintillation bandwidth $\Delta\nu_d$ induced by the CGM of a single halo for a range of possible FRB LOS configurations. Figures 4 and 5 show τ and $\Delta\nu_d$ for the hot and cool CGM, respectively, in the rest frame of the scattering medium and for a wide range of $DM_\ell^2 G_{\text{scatt}}$. Typical values of DM_ℓ are $\sim 1 - 10 \text{ pc cm}^{-3}$ in the cool CGM and $\sim 100 \text{ pc cm}^{-3}$ in the hot CGM (Prochaska & Zheng 2019), while $G_{\text{scatt}} \sim 10^2 - 10^4$ for typical LOS geometries and path lengths $\sim 0.1 \text{ Mpc}$ through the CGM (Cordes et al. 2022). We find that $\tau(1+z_\ell)^3$ is about 10 times larger in the cool CGM than in the hot CGM, and $\tau(1+z_\ell)^3$ is less than $10 \mu\text{s}$ at 1 GHz in both gas phases for a majority of plausible DMs and LOS configurations. Our inference from the quasar measurements suggests that CGM turbulence induces miniscule levels of pulse broadening, and that CGM scattering will only be detectable in the time domain when both DM_ℓ and G_{scatt} are large. The geometric leverage alone will only be sufficiently large to render detectable pulse broadening when an FRB is at high redshifts ($z > 1$) and/or the path length (L) through the scattering




Figure 5. Same as Figure 4 but for the cool CGM. Scattering parameters are estimated for sound speeds c_s typical of the cool CGM using density fluctuations inferred from nonthermal broadening measurements of quasar absorption lines. Since the DM contribution of the cool CGM may be significantly smaller than that of the hot CGM, smaller $DM_\ell^2 G_{\text{scatt}}$ are shown than in the previous figure.

medium is small ($L < 100 \text{ kpc}$) for a fixed DM_ℓ , provided that the expected amount of pulse broadening from the Milky Way in the same direction is subdominant.

These same results imply that scattering from the cool CGM may be more detectable in the frequency domain as diffractive scintillation, as we find $\Delta\nu_d(1+z_\ell)^{-3} \sim 0.01 - 10 \text{ MHz}$ at 1 GHz for the majority of plausible $DM_\ell^2 G_{\text{scatt}}$. These scintillation bandwidths are readily measurable for typical observing configurations, with the main barrier to detection being the Galactic foreground, which can induce scintillation at comparable levels depending on the LOS, and the host galaxy ISM and/or immediate FRB environment, which may scatter the FRB sufficiently to quench CGM scintillation. Scintillation from the hot CGM could be detectable ($\Delta\nu_d(1+z_\ell)^{-3} \sim 0.1 - 10 \text{ MHz}$ at 1 GHz) for large $DM_\ell^2 G_{\text{scatt}}$, but low Mach numbers and/or low $DM_\ell^2 G_{\text{scatt}}$ would correspond to $\Delta\nu_d(1+z_\ell)^{-3} \gg 10 \text{ MHz}$ at 1 GHz, which may require the observing bandwidth of an ultra-wideband receiver.

For typical source and lens separations, the τ and $\Delta\nu_d$ estimated for the cool CGM above are equivalent to angular

broadening in the lens frame $\theta_s \sim (d_{\text{so}}/d_{\text{sl}})\theta_d \sim 0.1 - 1 \mu\text{as}$ at 1 GHz. The hot CGM will produce even smaller θ_s . In the observer frame, these levels of angular broadening will likely be undetectable for most LOSs, due to both the Galactic foreground and typical radio imaging resolutions.

Scattering media near the source or observer have $G_{\text{scatt}} \rightarrow 1$ (Cordes et al. 2022). If the turbulence strengths inferred from quasar measurements are comparable to turbulence strengths in the Galactic CGM, then the expected amount of pulse broadening from the Galactic CGM is extremely small – for expected DM contributions $\sim 10 - 100 \text{ pc cm}^{-3}$ (Cook et al. 2023; Dutta et al. 2024), τ is at the nanosecond level or smaller, well below previously published upper limits (Ocker et al. 2021a). Similar levels of scattering would also be expected for the host galaxy CGM.

4.3. Comparison to Observed Scattering

The most stringent upper limits on scattering in the CGM are from FRBs with minimal scattering (based on observations with high time resolutions and/or large frequency bandwidths) and positively identified halos at known distances along their LOSs. About 15 precisely localized FRBs in the literature have confirmed foreground halos with known redshifts (Huang et al. 2024; van Leeuwen et al. 2023; Faber et al. 2024). Only a few of these have measured scattering times or scintillation bandwidths comparable to the values estimated for the CGM above, with the rest either having scattering significantly larger than expected for the CGM or lacking a scattering constraint altogether.

Two of the most stringent upper limits on CGM scattering are from FRB20191108A (aka FRB191108) and FRB20181112A (aka FRB181112). FRB191108 has a scintillation bandwidth of about 40 MHz at 1.37 GHz (equivalent to 16 ns at 1 GHz for $\tau \propto \nu^{-4.4}$) and intersects M33 at an impact parameter of 18 kpc (Connor et al. 2020). FRB181112 has a scattering time of $21 \pm 1 \mu\text{s}$ at 1.3 GHz (equivalent to $67 \mu\text{s}$ at 1 GHz; Cho et al. 2020) and intersects a foreground halo at $z_\ell = 0.36$ with an impact parameter of 29 kpc (Prochaska et al. 2019). We subtract the Milky Way scattering contribution for these two FRBs by taking the smallest measured scattering time at 1 GHz of pulsars at similar absolute Galactic latitudes $|b|$, based on the pulsar scattering database in Cordes et al. (2022). For FRB181112 ($b = -48^\circ$), $\tau_{\text{min}}^{\text{MW}} \approx 1.9 \text{ ns}$ at 1 GHz for pulsars near the same latitude, whereas for FRB191108 ($b = -30^\circ$), $\tau_{\text{min}}^{\text{MW}} \approx 2.6 \text{ ns}$ at 1 GHz. The Galactic scattering contribution to these FRB LOSs will be three times larger because when viewed from cosmological distances they behave like plane waves and are scattered through larger angles than Galactic sources, which are spherical waves (Cordes et al. 2016). The difference between the observed scattering and Milky Way scattering is then taken to be an upper limit on the total scattering from

the CGM along the FRB LOS; assuming a small Milky Way contribution thus yields a larger and more conservative upper limit on the scattering contribution of the intervening CGM.

Ocker et al. (2021a) used these scattering measurements to derive upper limits on \tilde{F} in the CGMs of the foreground galaxies, in addition to deriving \tilde{F} in the Milky Way disk and CGM using Galactic pulsars and two repeating FRBs with precise scattering budgets. Figure 6 shows \tilde{F} inferred for the Galactic CGM and disk based on these FRB and pulsar measurements, along with the \tilde{F} we infer for the hot and cool CGM based on quasars.

Ocker et al. (2021a) estimated \tilde{F} in the CGM by assuming the entire column density of hot halo gas contributed to scattering, and used a modified Navarro-Frenk-White (mNFW) profile (Prochaska & Zheng 2019) with a cutoff at twice the virial radius to estimate the corresponding DM. Here, we instead separate the CGM constraints from Ocker et al. (2021a) into the \tilde{F} that would be inferred if scattering were attributed to either the hot CGM or the cool CGM. For the hot CGM, we use the same mNFW profile and assumptions as in Ocker et al. (2021a), which yield DM contributions of about 90 pc cm^{-3} for FRB191108 and 135 pc cm^{-3} for FRB181112.

For the cool gas, column densities in neutral hydrogen are well constrained observationally from quasar absorption lines, and yield typical hydrogen column densities $N_{\text{H}} \sim 10^{18} - 10^{20} \text{ cm}^{-2}$ for neutral fractions $\sim 10^{-2} - 10^{-3}$ (Werk et al. 2014). We use the mean N_{H} of the COS-Haloes sample, $10^{19.6} \text{ cm}^{-2}$ (Werk et al. 2014), equivalent to a DM of 13 pc cm^{-3} . For the Galactic cool CGM, we use a mean DM of 20 pc cm^{-3} inferred from HI measurements of high-velocity clouds (Prochaska & Zheng 2019). While there is considerable uncertainty in the DM contribution of the CGM, these fiducial values sufficiently illustrate how the range of \tilde{F} compares between the different tracers shown in Figure 6.

The upper limits on \tilde{F} in the Milky Way CGM and the foreground CGM of FRB181112 are well above the range of values inferred from quasar measurements for both the hot and cool CGM, suggesting that these upper limits are too large to be constraining, and that the FRB scattering used to derive these limits is likely not dominated by the CGM. The upper limit on \tilde{F} for FRB191108 probing M33 comes closest to the ranges inferred for the cool and hot CGM from quasars; regardless of whether scattering is attributed to the cool or hot CGM, the resulting upper limit on \tilde{F} is comparable to the largest values inferred from quasars. This result affirms that the scattering of FRB191108 may have a significant contribution from the CGM.

4.4. The Dissipation Scale of CGM Turbulence

Consistency between the \tilde{F} inferred from FRB scattering and quasar-based velocity fluctuation measurements does not necessarily translate into positive proof that CGM density




Figure 6. The density fluctuation parameter \tilde{F} inferred for the CGM and Milky Way disk from pulsar (green), FRB (blue), and quasar (orange) observations. The range of values shown for the Milky Way disk is based on pulsar scattering measurements, while the upper limit on \tilde{F} in the Milky Way halo is based on two repeating FRBs for which a scattering budget can be constructed (Ocker et al. 2021a). Two other upper limits on \tilde{F} from FRBs are shown, the foreground galaxy CGM intersected by FRB181112 and the M33 CGM intersected by FRB191108. The FRB constraints are separated into the \tilde{F} that would be obtained if all of the scattering is attributed to cool gas (thick lines) or to hot gas (thin lines), based on the different expected DMs, and after subtracting the minimum scattering contribution of the Milky Way (see Section 4.3 and Equation 13). FRB and pulsar constraints account for the conversion between the $1/e$ delay measured from pulse shapes and the mean scattering time; for FRB scattering in the CGM, the assumed conversion factor is $A_\tau \approx 1/6$, whereas for pulsars we assume $A_\tau \approx 1$ because their larger scattering leads to smaller diffractive scales and hence larger A_τ . The range of \tilde{F} shown for the cool and hot CGM traced by quasars is based on Figure 3 above, and assumes an inner scale equal to the ion inertial scale.

fluctuations extend to \lesssim au scales (after all, we have assumed a single uniform cascade of density fluctuations across sub-au to $>$ kpc scales, when in fact there may be multiple injection and dissipation scales, which can also alter the shape of the density fluctuation spectrum and introduce deviations from Kolmogorov turbulence). However, since \tilde{F} depends on the inner scale l_i , in principle one can combine FRB-based constraints on \tilde{F} (which are based on observables and make no assumption about l_i) with quasar-inferred values of \tilde{F} (which necessarily cover a range of possible l_i), to yield either a lower limit or, with complete information, a constraining inference of the inner scale.

Figure 7 shows \tilde{F} vs. a wide range of possible values for the inner scale, based on the C_n^2 inferred for the hot and cool CGM in Figure 3. These quasar-based constraints on \tilde{F} are compared to FRB191108, which gives an upper limit on \tilde{F} ir-

respective of the inner scale. Taking this comparison at face value, the inner scale of the coldest ($\sim 10^4$ K) gas would need to be $\gtrsim 200$ km in order for the quasar and FRB191108 constraints to be consistent; this constraint falls between the electron inertial length and ion gyroradius typical of the cool CGM (Table 1). For the hot gas, the FRB upper limit has more constraining power if the outer scale is small ($l_o = 10$ kpc), and is consistent with an inner scale $\gtrsim 10^4$ km for Mach numbers $\lesssim 0.5$. If the outer scale is large ($l_o \sim 100$ kpc) and $\mathcal{M} \approx 1$, then $l_i \gtrsim 10^6$ km, much greater than the ion inertial length.

These lower limits are purely demonstrative and do not account for potentially large uncertainties in the DM contribution of the CGM, which dominates the total uncertainty in the constraint on l_i since $l_i \propto \text{DM}^{-6}$. A full error accounting should also include formal uncertainty errors on the scattering measurement, and uncertainty in the Milky Way scattering subtraction. For the purposes of this initial comparison, we note that reducing the Milky Way scattering contribution to an even smaller value, e.g. 1 ns, would yield a lower limit on l_i in the cool gas that is more comparable to the electron inertial length, but still compatible with dissipation at plasma microscales.

In principle, this comparison can be highly informative when the sound speed or Mach number is either known or more tightly bound, and when the FRB and quasar constraints are placed on the same system. Hydrogen column densities inferred from quasar absorption lines could also be used to place tighter constraints on the DM contribution of the CGM – albeit along a different line of sight – mitigating uncertainty in the FRB DM budget. If the CGM in question has detectable emission, then IFS measurements could also be used to infer the turbulence spectral index, and the subsequent analysis can be modified accordingly (rather than assuming a Kolmogorov spectral index as we have done here), though of course this still involves extrapolation over many orders of magnitude, along which the spectral index could change.

4.5. Conditions for Multipath Propagation

The results above all assume that the conditions for multipath propagation are satisfied in the CGM, i.e., that the CGM does produce some level of radio pulse broadening or scintillation. If, however, such scattering from the CGM can be firmly ruled out, then the inner scale of the density fluctuation spectrum must be greater than the multipath scale (Equation 11). CGM scattering could be ruled out by localization of scattering screens to the Milky Way and/or host galaxy ISMs, which requires some combination of pulse broadening, scintillation bandwidth, and/or angular broadening measurements. Detection of scintillation over months-long baselines (e.g., for a repeating FRB), or combining an angular




Figure 7. Density fluctuation parameter \tilde{F} vs. the dissipation scale, for the cool (left) and hot (middle, right) CGM as inferred from quasar observations. Upper limits on \tilde{F} from FRB191108 are shown by the black horizontal lines, separated into constraints for the cool and hot phases based on their different DM contributions, and after subtracting the minimum Milky Way scattering contribution (see Section 4.3).

broadening measurement with a scintillation bandwidth, can also constrain the scattering screen location, in some cases revealing it to be within the Milky Way ISM (Ocker et al. 2021a; Wu et al. 2024). When pulse broadening or scintillation is detected on distinct frequency scales, such that the observed scattering cannot be explained by a single screen, the scattering screens can typically be localized to within the Milky Way and host galaxy (Masui et al. 2015; Ocker et al. 2022b; Sammons et al. 2022; Nimmo et al. 2025). Such an analysis on an FRB with an intervening CGM would be the basis for determining whether the scattering is consistent or inconsistent with arising in that CGM.

Figure 8 shows r_{mp} vs. $\text{SM}_{100} \equiv C_n^2 \times 100 \text{ kpc}$, the scattering measure for the range of C_n^2 extrapolated from quasar measurements and a fiducial path length of 100 kpc. We assume the same path length for both gas phases, noting that SM_{100} can be easily rescaled; e.g., for a path length of 10 kpc SM_{100} would be 10 times smaller than the values shown in Figure 8. Only multipath scales greater than the Fresnel scale are relevant, because when $r_{\text{mp}} < r_F$ the strong scattering regime (and the scattering formulas used in this work) no longer apply. The multipath scale is significantly affected by the choice of β : For a given SM_{100} , r_{mp} differs by seven orders of magnitude between $\beta = 3.1$ and $\beta = 3.9$. By contrast, modifying λ and z_ℓ within their typical ranges ($0.1 \lesssim \lambda \lesssim 0.8 \text{ m}$, $z_\ell < 1$) changes r_{mp} by less than an order of magnitude from the values shown in Figure 8. For a fixed path length of 100 kpc, r_{mp} in the hot CGM ranges from a few au to $\sim 0.1 \text{ pc}$, depending on β . If β is small ($\lesssim 3.4$), r_{mp} is comparable to the Fresnel scale and no strong diffractive scattering will occur regardless of the inner scale – in this scenario, a scattering non-detection would not yield any information on the inner scale. For $\beta = 3.67 \approx 11/3$ (Kol-




Figure 8. The multipath scale r_{mp} (equivalent to the scattering cone diameter at the screen) vs. $\text{SM}_{100} \equiv C_n^2 \times 100 \text{ kpc}$, for the range of C_n^2 found in the CGM (see Figure 3). Here r_{mp} is evaluated as $r_{\text{mp}} = 2d_{10}\theta_d = (d_{10}d_{\text{sl}}/d_{\text{so}})\lambda/\pi r_{\text{diff}}$, where r_{diff} is defined as in Equation 10, for fiducial values $d_{\text{sl}}/d_{\text{so}} = 0.5$, $\lambda = 0.3 \text{ m}$, $z_\ell = 0.2$, and a range of β . The approximate division between SM_{100} characteristic of hot and cool gas is indicated by the grey dashed line. When $r_{\text{mp}} < r_F \sim 2 \times 10^{-5} \text{ pc}$ (black dash-dotted line), the strong scattering regime no longer applies.

mogorov), r_{mp} can be larger than the Fresnel scale, and a robust non-detection of scattering in this regime would suggest an inner scale much larger than plasma microscales, which could imply the presence of a damping process, e.g., ion-neutral damping, that terminates the cascade above kinetic scales (Lithwick & Goldreich 2001). For $\text{SM}_{100} \gtrsim 10^{-7} \text{ kpc m}^{-20/3}$, roughly characteristic of the cool CGM, $\beta \approx 11/3$ yields $r_{\text{mp}} \sim 0.01 - 10 \text{ pc}$, comparable to the smallest spa-

tial scales inferred for cool CGM clouds from quasar absorption lines (although these cloud sizes may be overestimated due to the limited velocity resolution of spectrographs; Rudie et al. 2019; Chen et al. 2023a). This result suggests that if the cool CGM produces scattering, then cool CGM cloudlets must sustain internal turbulence at scales much smaller than their sizes.

5. DISCUSSION

Quasar observations of CGM turbulence suggest that density fluctuations are extremely weak in the hot CGM ($10^{-16} \lesssim C_n^2 \lesssim 10^{-9} \text{ m}^{-20/3}$, equivalent to $10^{-8} \lesssim \tilde{F} \lesssim 10^{-4} (\text{pc}^2 \text{ km})^{-1/3}$) and moderately weak in the cool CGM ($10^{-9} \lesssim C_n^2 \lesssim 10^{-5} \text{ m}^{-20/3}$, equivalent to $10^{-4} \lesssim \tilde{F} \lesssim 10^{-3} (\text{pc}^2 \text{ km})^{-1/3}$). The expected level of FRB scattering from the CGM is thus miniscule: $\tau(1 + z_\ell)^3 \ll 10 \mu\text{s}$ at 1 GHz for most sightline geometries, with τ from the Milky Way and host galaxy CGMs predicted to be $\lesssim \text{ns}$. Our analysis indicates that even if CGM scattering is undetected, stringent upper limits on the scattering delay can be combined with quasar observations to yield lower limits on the dissipation scale. An initial comparison between quasar-inferred values of \tilde{F} and FRB191108 suggests that the FRB’s scattering is likely dominated by the foreground CGM, and that it is compatible with dissipation near plasma microscales. The dissipation scale can be more tightly constrained if a firm measurement of CGM scattering is made, but this scenario likely requires either negligible scattering from the ISMs of the FRB host galaxy and Milky Way, or highly robust determinations of their scattering contributions. The methodology pursued here is also applicable to FRB host galaxies, for which ISM turbulence may readily be studied in emission.

5.1. Modifying Our Assumptions

While we have assumed uniform, isotropic Kolmogorov turbulence for most of our analysis, there is evidence that CGM turbulence can deviate from a Kolmogorov spectrum, and that the turbulence spectrum may include local enhancements due to shocks and/or local energy sources (e.g., inflows and outflows). Turbulence spectra significantly shallower or steeper than Kolmogorov would change our scattering predictions. Keeping δn_e fixed at the outer scale (to, e.g., a value set by the Mach number), $3 < \beta < 11/3$ will yield C_n^2 larger than the values we estimate for $\beta = 11/3$, thus leading to larger scattering delays for a given halo and LOS geometry (provided that the multipath scale remains larger than the Fresnel and diffractive scales). Conversely, $11/3 < \beta < 4$ will yield C_n^2 and τ smaller than what we find for $\beta = 11/3$. For $\beta > 4$ refractive effects will dominate over a larger range of spatial scales and the diffractive scattering considered here may become irrelevant because density fluctuations would become too weak at the length scales relevant to diffraction

(Goodman & Narayan 1985; Cordes et al. 1986). If β is well determined from velocity fluctuation measurements, then its empirical value should be used to estimate C_n^2 , although this still involves extrapolation over many orders of magnitude.

In addition to assuming $\beta = 11/3$, we have adopted a range of sound speeds and Mach numbers based on the range of CGM systems with observed turbulence, none of which overlap with known FRBs. Targeted surveys of FRB-quasar pairs intersecting foreground galaxies, or FRBs intersecting foreground CGMs detected in emission, may directly yield estimates of the Mach number, β , and even l_o in those individual systems, enabling a more direct comparison between FRB and quasar observations. Such a comparison may already be achievable in M31, which covers a large enough area of the sky that dozens of quasars and FRBs intersect it (Lehner et al. 2020; Connor & Ravi 2022).

5.2. Dissecting the Cloudlet Model: Volume Filling Fraction and Outer Scale of the Cold CGM

Our formalism for relating scattering observables to density fluctuations is based on an ionized cloudlet model in which the CGM is composed of cloudlets that sustain internal turbulence, while allowing for fluctuations in the mean density of different cloudlets. Our analysis suggests that cooler ($10^4 - 10^5 \text{ K}$) cloudlets are the most likely to produce observable FRB scattering, as suggested in previous studies. One may then ask what additional properties of the cool CGM may be constrained by our model, other than the strength of density fluctuations and their dissipation scale. The size of individual cloudlets is not explicitly factored into the model, but it is implicitly related to the model parameters ζ , f , and l_o . If the CGM is composed of a single large cloud with uniform mean density, then $\zeta = \langle n_{ec}^2 \rangle / \langle n_{ec} \rangle^2$ and the volume filling factor f will be unity, with the path length through the scattering screen L comparable to the cloud size. If, on the other hand, the CGM is significantly “clumpy,” with a range of mean cloudlet densities along the LOS, then $\zeta > 1$ (this parameter is identical to a clumping factor; e.g. see Eq. 18 in Dutta et al. 2024). If cool gas is additionally concentrated into numerous tiny cloudlets spread uniformly in the CGM with a small volume fraction $f \ll 1$, L is comparable to the virial radius. A small volume filling fraction with a large area covering fraction of cool gas is implied by quasar absorption studies (see Hummels et al. 2024; Dutta et al. 2024; Singh Bisht et al. 2024 for a range of possible models). Regardless of the exact scenario, the cloudlet size may also be related to the outer scale of the density fluctuations within cloudlets. These parameters are all lumped together in the composite $\tilde{F} = \zeta \epsilon^2 / f (l_o^2 l_i)^{1/3}$, which we infer both from quasar observations of turbulence and upper limits on FRB scattering in the CGM.

While dissecting \tilde{F} is difficult, involving five degenerate parameters, it is not altogether impossible. In principle, ϵ^2 and l_o can both be inferred by leveraging quasar-based constraints on column densities and cloud sizes, and l_i can be constrained through the methods laid out in this paper. Given some prior on ϵ^2 , l_o , and l_i , one can then constrain the likely range of ζ/f , which will be close to unity for a highly uniform, volume-filling medium or much greater than unity for a highly clumpy/non-uniform medium. As an illustration, if we assume $\bar{n}_e \sim 10^{-2} \text{ cm}^{-3}$ and $l_o \sim 10 \text{ pc}$, we have $C_n^2 \sim 10^{-6} \text{ m}^{-20/3}$ (Figure 3) and $\epsilon^2 \sim 10^{-4}$ (Equation 1). For $l_i \sim 2000 \text{ km}$ (the ion inertial scale for $n_e \sim 10^{-2} \text{ cm}^{-3}$), and a total constraint on $\tilde{F} \sim 5 \times 10^{-4} (\text{pc}^2 \text{ km})^{-1/3}$ (Figure 3), we would then infer $\zeta/f \sim 300$. Given that $\zeta \geq 1$ by definition, this value of ζ/f would imply a volume filling factor of cloudlets $f \gtrsim 0.003$. Conversely, assuming a perhaps unphysically large outer scale $l_o \sim 100 \text{ kpc}$ would yield $\zeta/f \sim 100$, implying an unrealistically large filling fraction of cool cloudlets for any plausible range of cold gas overdensities and mass fractions. While these are purely demonstrative calculations, they illustrate that our empirically motivated constraints on \tilde{F} would imply insensibly large filling fractions of high-density gas if an outer scale comparable to the halo radius is assumed. The line of reasoning followed above offers one possible path to interpreting quasar and/or FRB-based constraints on \tilde{F} .

5.3. Comparison with VP19 and the Misty CGM

Applying this ionized cloudlet model to quasar-based inferences of turbulence yields scattering predictions that are at least 100 times smaller than the fiducial values predicted by Vedantham & Phinney (2019) (hereafter VP19), who used a simple analytic prescription for the McCourt et al. (2018) misty CGM, in which the the CGM is a volume of cool cloudlets with a volume filling factor $f \sim 10^{-4}$, internal density $n_e \sim 10^{-3} \text{ cm}^{-3}$, and cloudlet radius $r_c \sim 1 \text{ pc}$. In VP19, scattering arises from the random distribution of cloudlets along a given LOS, with the superposition of intersected cloudlets yielding phase perturbations that mimic a Kolmogorov spectrum with an outer scale comparable to the cloudlet size. The smallest scale cutoff of the phase perturbations (i.e., the inner scale) is not included in their model, because they focus on predicting the diffractive scattering time, which depends on r_{diff} rather than l_i . While their model sources density fluctuations not from turbulence, but from the number and mean density of intersected cloudlets, making a direct comparison between our results difficult (and perhaps unphysical), we can nonetheless investigate what modifications to their model would yield scattering predictions more similar to our own.

In the VP19 model, $\tau_d \propto r_c^{-2} \nu^{-4.4} \text{DM}^{2.4} f_a^{1.2} (d_{\text{sl}} d_{\text{lo}} / d_{\text{so}})$, where r_c is the cloud size and $f_a \sim fb/r_c$ is the number of

intersected clouds at an impact parameter b , for a volume-filling factor f . Their fiducial values $r_c \sim 1 \text{ pc}$, $\nu \sim 1 \text{ GHz}$, $\text{DM} \sim 0.03 \text{ pc cm}^{-3}$, $f_a \sim 10$, and $(d_{\text{sl}} d_{\text{lo}} / d_{\text{so}}) \sim 1 \text{ Gpc}$ yields $\tau_d \sim 0.4 \text{ ms}$. Since decreasing τ_d requires increasing r_c and/or decreasing f_a , the most natural way to produce a smaller scattering prediction is to thus increase r_c . A modest increase from $r_c \sim 1 \text{ pc}$ to $r_c \sim 10 \text{ pc}$ is sufficient to make VP19's predictions broadly consistent with our own (see their Equation 19), albeit for a different assumed DM. Increasing the DM in VP19 would require an even larger increase in r_c to produce scattering levels comparable to our predictions.

An important limitation of VP19 is the assumption of a universal cloud size. A key motivation for considering a misty CGM is the large area covering factor $f_A \sim \mathcal{O}(1)$ of cold gas, despite its small volume filling fraction $f \sim 10^{-3}$ (McCourt et al. 2018). Since $f_A \sim (R/r_c)f$, for an $R \sim 100 \text{ kpc}$ halo to have order unity f_A we require $r_c < 100 \text{ pc}$. McCourt et al. (2018) argued that the cooling length $l_s \sim \min(c_s t_{\text{cool}})$, the scale on which cooling clouds establish sonic contact with their surroundings, sets a characteristic cloud length scale. Since $l_s \propto n_e^{-1}$, this corresponds to a fixed column $N_e \sim 10^{17} \text{ cm}^{-2}$, or $l_s \sim 30 \text{ pc} (n_e/10^{-3})^{-1} \text{ cm}^{-3}$. This characteristic scale is much larger than $r_c \sim 1 \text{ pc}$ clouds frequently invoked in the literature, which is actually the value of r_c for l_s in higher redshift, denser halos³. Later work on multi-phase gas in a turbulent medium has found a broad power-law spectrum of cloud sizes, with $dN/dM \propto M^{-2}$, or equal mass per logarithmic interval (Gronke et al. 2022; Das & Gronke 2023; Singh Bisht et al. 2024). At fixed density, this is equivalent to equal volume per logarithmic interval. Since $f_A \propto f/r_c$, the smallest clouds will dominate the area covering factor, so a key parameter is the small-scale cutoff to this power-law spectrum. In Gronke et al. (2022), this small-scale cutoff is – modulo uncertainty due to limited numerical resolution – of order the survival radius (the minimal scale where cooling can overcome the mixing due to shear-induced Kelvin Helmholtz instabilities), which is larger than the scale predicted by McCourt et al. (2018). There are many complications (e.g. non-thermal pressure support) that can affect these estimates, but it is fair to say that the lower limit on cool cloud sizes for $z \sim 0$ halos is very likely to be significantly larger than $\sim 1 \text{ pc}$.

Increasing r_c also alleviates the problem posed by Jow et al. (2024), who argue that refractive multiple imaging of FRBs by CGM clouds would suppress Galactic scintillation (and hence that detection of Galactic scintillation may disfa-

³ McCourt et al. (2018) focused on such systems, as they found that – by comparing to observational data – the shattering hypothesis worked well at high redshift, but less well at low redshift.

vor some misty CGM models). Increasing r_c from their nominal value of $r_c \sim 0.1$ pc to $r_c \sim 10$ pc decreases the number of refractive images by a factor of 10^{-10} , to $N_{\text{image}} \approx 1$ (see Equation 27 in Jow et al. 2024). We thus find that VP19’s CGM model can produce scattering levels comparable to our results if the model’s cloud size is increased, and that under these same conditions refractive multiple imaging (and significant quenching of Galactic scintillation) is unlikely.

The single cloud-scale model adopted by VP19 is qualitatively reminiscent of early monoscale descriptions of scintillation in both the interplanetary medium (IPM) and the ISM, in which density irregularities were modeled as a Gaussian power spectrum with a single characteristic length scale, as opposed to a turbulent power-law spectrum covering a wide range of scales (Lovell 1970). Under certain conditions (e.g., $\beta \geq 4$, or $\beta < 4$ and $l_i \gtrsim r_{\text{diff}}$), a power-law medium can produce a phase structure function with a similar dependence on spatial separation as a Gaussian density power spectrum (see Rickett 1977 for a review). Gaussian density spectra were largely ruled out based on observational evidence for density fluctuations spanning a much wider range of spatial scales than allowed in the monoscale model (Armstrong et al. 1981, 1995; Chepurnov & Lazarian 2010). Distinguishing between monoscale and turbulent media in the CGM will similarly require assessment of observations spanning a wide range of spatial scales; measuring FRB scattering delays in a single radio frequency band will likely not suffice. However, the potential compatibility of non-thermal line broadening measurements with FRB scattering upper limits, as illustrated in this paper, already hints at fluctuations spanning a much wider range of scales than a monoscale model (such as VP19) would allow.

Importantly, we have shown that observed FRB scattering is not inconsistent with the misty CGM picture, as quasar observations of cool gas turbulence indicate tiny levels of scattering would occur even if that turbulence extends down to sub-au scales. We thus find that cool gas can be ubiquitous in the CGM without producing extreme amounts of FRB scattering incompatible with observations. Whether that cool gas is misty or concentrated in larger discrete clouds (more precisely, kinematically coherent structures) is difficult to assess. It is relatively straightforward to rule out scattering from a single discrete cloud because the small path length through the cloud would increase G_{scatt} by several orders of magnitude, yielding significantly larger scattering than we predict (Faber et al. 2024). Ultimately, targeted surveys of quasars and FRBs probing the same galaxy systems, with the requisite dedication of observing resources, offer the best prospects for constraining CGM structure across the full range of spatial scales necessary to test misty CGM models.

ACKNOWLEDGMENTS

The authors thank Cameron Hummels, Hsiao-Wen Chen, Gwen Rudie, Irina Zhuravleva, Jim Cordes, and Drummond Fielding for insightful conversations about this work. PS thanks Pawan Kumar for his introduction to the basic physics of radio scintillation. Part of this work was performed at the Aspen Center for Physics, which is supported by National Science Foundation grant PHY-2210452. SKO and MC are supported by the Brinson Foundation through the Brinson Prize Fellowship Program. SKO is a member of the NANOGrav Physics Frontiers Center (NSF award PHY-2020265). SPO acknowledges NSF grant AST240752 for support. Caltech and Carnegie Observatories are located on the traditional and unceded lands of the Tongva people.

APPENDIX

A. RELATING C_n^2 AND \tilde{F}

Here we lay out the formalism relating C_n^2 , the amplitude of the density fluctuation power spectrum, and \tilde{F} , the composite density fluctuation parameter in the ionized cloudlet model (Section 3.2), expanding on the approach laid out in Cordes et al. (2016). We define the mean scattering time for a source at distance d_{so}

$$\tau = (1/2c) \int_0^{d_{\text{so}}} ds \eta(s) s (1 - s/d_{\text{so}}) \quad (\text{A1})$$

where $\eta(s)$ is the mean-square scattering angle per unit length due to turbulent plasma at a distance s :

$$\eta(s) = \lambda^4 r_e^2 \int dq q^3 P_{\delta n_e}(q, s). \quad (\text{A2})$$

Here we have assumed an isotropic 3D density fluctuation spectrum $P_{\delta n_e}$, and for a spectral index $\beta < 4$ and an inner scale $l_i = 2\pi/q_i$ much smaller than the outer scale l_o , the mean-square scattering angle is (Cordes & Rickett 1998):

$$\begin{aligned}\eta(s) &\approx \frac{\lambda^4 r_e^2 \Gamma(3 - \beta/2)}{4 - \beta} q_i^{4-\beta} C_n^2(s) \\ &\approx \frac{1}{3} \lambda^4 r_e^2 \Gamma(7/6) q_i^{1/3} C_n^2(s) \text{ for } (\beta = 11/3).\end{aligned}\tag{A3}$$

In the cloudlet model, individual cloudlets sustain internal turbulence with rms density fluctuations $\langle \delta n_{ec}^2 \rangle$ related to $C_{n,c}^2$, where $C_{n,c}^2$ refers to the turbulence amplitude within individual cloudlets:

$$\begin{aligned}C_{n,c}^2 &= \frac{\beta - 3}{2(2\pi)^{4-\beta}} \langle \delta n_{ec}^2 \rangle l_o^{3-\beta} \\ &= [3(2\pi)^{1/3}]^{-1} \langle \delta n_{ec}^2 \rangle l_o^{-2/3} \text{ for } \beta = 11/3.\end{aligned}\tag{A4}$$

Letting $\epsilon^2 = \langle \delta n_{ec}^2 \rangle / n_{ec}^2$ and $C_{SM} = (\beta - 3)/2(2\pi)^{4-\beta}$, we thus have

$$C_{n,c}^2 = C_{SM} \epsilon^2 n_{ec}^2 l_o^{3-\beta},\tag{A5}$$

emphasizing that n_{ec} is the local mean density of a cloudlet. Ultimately, we are interested in evaluating $\eta(s)$ for the volume-averaged $C_n^2 = \overline{C_{n,c}^2}$ along the LOS:

$$\begin{aligned}\overline{C_{n,c}^2} &= f C_{SM} \langle \epsilon^2 n_{ec}^2 \rangle l_o^{3-\beta} \\ &= f C_{SM} \langle \epsilon^2 n_{ec}^2 \rangle l_o^{3-\beta} \times \frac{\langle f n_{ec} \rangle^2}{\langle f n_{ec} \rangle^2} \\ &= C_{SM} \frac{\zeta \epsilon^2}{f l_o^{\beta-3}} \bar{n}_e^2,\end{aligned}\tag{A6}$$

where $\zeta = \langle n_{ec}^2 \rangle / \langle n_{ec} \rangle^2$ describes the fractional variations in density between cloudlets, $\bar{n}_e = f \langle n_{ec} \rangle$ is the volume-averaged electron density along the LOS, and we have assumed the outer scale and ϵ^2 to be the same for all cloudlets.

Given the dependence of $\eta(s)$ on q_i , we lump all quantities describing the density fluctuations together into the parameter $\tilde{F} = \zeta \epsilon^2 / f l_o^{\beta-3} l_i^{4-\beta}$. We thus have

$$\begin{aligned}\overline{C_{n,i}^2} &= C_{SM} \tilde{F} \bar{n}_e^2 l_i^{4-\beta} \\ &= C_{SM} \tilde{F} \bar{n}_e^2 l_i^{1/3} \text{ for } \beta = 11/3.\end{aligned}\tag{A7}$$

The subsequent mean scattering time can be evaluated using Equation A1, as laid out in Cordes et al. (2016, 2022). For $\beta = 11/3$, this procedure ultimately yields the form given in Equation 13, where we have used the screen dispersion measure due to a given phase $DM_\ell = \bar{n}_e L$. The frequency dependence of $\eta(s)$, and hence the mean pulse broadening delay, is exactly ν^{-4} , whereas the diffractive scattering delay has a frequency dependence that depends on β . Modifying β would only change the constant prefactor in Equation 13 and the exponents of the inner and outer scales in \tilde{F} ; all other quantities would remain the same.

REFERENCES

- Alexandrova, O., Carbone, V., Veltri, P., & Sorriso-Valvo, L. 2008, ApJ, 674, 1153, doi: [10.1086/524056](https://doi.org/10.1086/524056)
- Alexandrova, O., Lacombe, C., Mangeney, A., Grappin, R., & Maksimovic, M. 2012, ApJ, 760, 121, doi: [10.1088/0004-637X/760/2/121](https://doi.org/10.1088/0004-637X/760/2/121)
- Armstrong, J. W., Cordes, J. M., & Rickett, B. J. 1981, Nature, 291, 561, doi: [10.1038/291561a0](https://doi.org/10.1038/291561a0)
- Armstrong, J. W., Rickett, B. J., & Spangler, S. R. 1995, ApJ, 443, 209, doi: [10.1086/175515](https://doi.org/10.1086/175515)
- Butsky, I. S., Hummels, C. B., Hopkins, P. F., Quinn, T. R., & Werk, J. K. 2024, MNRAS, 535, 1672, doi: [10.1093/mnras/stae2459](https://doi.org/10.1093/mnras/stae2459)
- Chen, H.-W., Qu, Z., Rauch, M., et al. 2023a, ApJL, 955, L25, doi: [10.3847/2041-8213/acf85b](https://doi.org/10.3847/2041-8213/acf85b)

- Chen, M. C., Chen, H.-W., Rauch, M., et al. 2023b, *MNRAS*, 518, 2354, doi: [10.1093/mnras/stac3193](https://doi.org/10.1093/mnras/stac3193)
- , 2024, *ApJ*, 962, 98, doi: [10.3847/1538-4357/ad1406](https://doi.org/10.3847/1538-4357/ad1406)
- , 2025, *ApJL*, 978, L18, doi: [10.3847/2041-8213/ad9bac](https://doi.org/10.3847/2041-8213/ad9bac)
- Chepurnov, A., & Lazarian, A. 2010, *ApJ*, 710, 853, doi: [10.1088/0004-637X/710/1/853](https://doi.org/10.1088/0004-637X/710/1/853)
- Cho, H., Macquart, J.-P., Shannon, R. M., et al. 2020, *ApJL*, 891, L38, doi: [10.3847/2041-8213/ab7824](https://doi.org/10.3847/2041-8213/ab7824)
- Coles, W. A., Frehlich, R. G., Rickett, B. J., & Codona, J. L. 1987, *ApJ*, 315, 666, doi: [10.1086/165168](https://doi.org/10.1086/165168)
- Connor, L., & Ravi, V. 2022, *Nature Astronomy*, 6, 1035, doi: [10.1038/s41550-022-01719-7](https://doi.org/10.1038/s41550-022-01719-7)
- Connor, L., van Leeuwen, J., Oostrum, L. C., et al. 2020, *MNRAS*, 499, 4716, doi: [10.1093/mnras/staa3009](https://doi.org/10.1093/mnras/staa3009)
- Cook, A. M., Bhardwaj, M., Gaensler, B. M., et al. 2023, *ApJ*, 946, 58, doi: [10.3847/1538-4357/acbbd0](https://doi.org/10.3847/1538-4357/acbbd0)
- Cordes, J. M., Ocker, S. K., & Chatterjee, S. 2022, *ApJ*, 931, 88, doi: [10.3847/1538-4357/ac6873](https://doi.org/10.3847/1538-4357/ac6873)
- Cordes, J. M., Pidwerbetsky, A., & Lovelace, R. V. E. 1986, *ApJ*, 310, 737, doi: [10.1086/164728](https://doi.org/10.1086/164728)
- Cordes, J. M., & Rickett, B. J. 1998, *ApJ*, 507, 846, doi: [10.1086/306358](https://doi.org/10.1086/306358)
- Cordes, J. M., Wharton, R. S., Spitler, L. G., Chatterjee, S., & Wasserman, I. 2016, arXiv e-prints, arXiv:1605.05890, <https://arxiv.org/abs/1605.05890>
- Das, H. K., & Gronke, M. 2023, *MNRAS*, doi: [10.1093/mnras/stad3125](https://doi.org/10.1093/mnras/stad3125)
- Dutta, A., Bisht, M. S., Sharma, P., et al. 2024, *MNRAS*, 531, 5117, doi: [10.1093/mnras/stae977](https://doi.org/10.1093/mnras/stae977)
- Faber, J. T., Ravi, V., Ocker, S. K., et al. 2024, arXiv e-prints, arXiv:2405.14182, doi: [10.48550/arXiv.2405.14182](https://doi.org/10.48550/arXiv.2405.14182)
- Faucher-Giguère, C.-A., & Oh, S. P. 2023, *ARAA*, 61, 131, doi: [10.1146/annurev-astro-052920-125203](https://doi.org/10.1146/annurev-astro-052920-125203)
- Federrath, C., Klessen, R. S., Iapichino, L., & Beattie, J. R. 2021, *Nature Astronomy*, 5, 365, doi: [10.1038/s41550-020-01282-z](https://doi.org/10.1038/s41550-020-01282-z)
- Federrath, C., Roman-Duval, J., Klessen, R. S., Schmidt, W., & Mac Low, M. M. 2010, *A&A*, 512, A81, doi: [10.1051/0004-6361/200912437](https://doi.org/10.1051/0004-6361/200912437)
- Fielding, D. B., Ostriker, E. C., Bryan, G. L., & Jermyn, A. S. 2020, *ApJL*, 894, L24, doi: [10.3847/2041-8213/ab8d2c](https://doi.org/10.3847/2041-8213/ab8d2c)
- Geiger, A., Cordes, J. M., Lam, M. T., et al. 2024, arXiv e-prints, arXiv:2411.08191, doi: [10.48550/arXiv.2411.08191](https://doi.org/10.48550/arXiv.2411.08191)
- Goodman, J., & Narayan, R. 1985, *MNRAS*, 214, 519, doi: [10.1093/mnras/214.4.519](https://doi.org/10.1093/mnras/214.4.519)
- Gronke, M., Oh, S. P., Ji, S., & Norman, C. 2022, *MNRAS*, 511, 859, doi: [10.1093/mnras/stab3351](https://doi.org/10.1093/mnras/stab3351)
- Huang, Y., Simha, S., Khrykin, I., et al. 2024, arXiv e-prints, arXiv:2408.12864, doi: [10.48550/arXiv.2408.12864](https://doi.org/10.48550/arXiv.2408.12864)
- Hummels, C. B., Rubin, K. H. R., Schneider, E. E., & Fielding, D. B. 2024, *ApJ*, 972, 148, doi: [10.3847/1538-4357/ad5965](https://doi.org/10.3847/1538-4357/ad5965)
- Hummels, C. B., Smith, B. D., Hopkins, P. F., et al. 2019, *ApJ*, 882, 156, doi: [10.3847/1538-4357/ab378f](https://doi.org/10.3847/1538-4357/ab378f)
- Ji, S., Oh, S. P., & Masterson, P. 2019, *MNRAS*, 487, 737, doi: [10.1093/mnras/stz1248](https://doi.org/10.1093/mnras/stz1248)
- Jow, D. L., Pen, U.-L., & Feldbrugge, J. 2023, *MNRAS*, 525, 2107, doi: [10.1093/mnras/stad2332](https://doi.org/10.1093/mnras/stad2332)
- Jow, D. L., Wu, X., & Pen, U.-L. 2024, *Proceedings of the National Academy of Science*, 121, e2406783121, doi: [10.1073/pnas.2406783121](https://doi.org/10.1073/pnas.2406783121)
- Konstandin, L., Girichidis, P., Federrath, C., & Klessen, R. S. 2012, *ApJ*, 761, 149, doi: [10.1088/0004-637X/761/2/149](https://doi.org/10.1088/0004-637X/761/2/149)
- Krishnakumar, M. A., Mitra, D., Naidu, A., Joshi, B. C., & Manoharan, P. K. 2015, *ApJ*, 804, 23, doi: [10.1088/0004-637X/804/1/23](https://doi.org/10.1088/0004-637X/804/1/23)
- Lambert, H. C., & Rickett, B. J. 1999, *ApJ*, 517, 299, doi: [10.1086/307181](https://doi.org/10.1086/307181)
- Lee, K. H., & Lee, L. C. 2019, *Nature Astronomy*, 3, 154, doi: [10.1038/s41550-018-0650-6](https://doi.org/10.1038/s41550-018-0650-6)
- Lehner, N., Berek, S. C., Howk, J. C., et al. 2020, *ApJ*, 900, 9, doi: [10.3847/1538-4357/aba49c](https://doi.org/10.3847/1538-4357/aba49c)
- Lithwick, Y., & Goldreich, P. 2001, *ApJ*, 562, 279, doi: [10.1086/323470](https://doi.org/10.1086/323470)
- Lovelace, R. V. E. 1970, PhD thesis, -
- Macquart, J.-P., & Koay, J. Y. 2013, *ApJ*, 776, 125, doi: [10.1088/0004-637X/776/2/125](https://doi.org/10.1088/0004-637X/776/2/125)
- Masui, K., Lin, H.-H., Sievers, J., et al. 2015, *Nature*, 528, 523, doi: [10.1038/nature15769](https://doi.org/10.1038/nature15769)
- McCourt, M., Oh, S. P., O’Leary, R., & Madigan, A.-M. 2018, *MNRAS*, 473, 5407, doi: [10.1093/mnras/stx2687](https://doi.org/10.1093/mnras/stx2687)
- Miller, M. J., & Bregman, J. N. 2015, *ApJ*, 800, 14, doi: [10.1088/0004-637X/800/1/14](https://doi.org/10.1088/0004-637X/800/1/14)
- Mohapatra, R., Federrath, C., & Sharma, P. 2021, *MNRAS*, 500, 5072, doi: [10.1093/mnras/staa3564](https://doi.org/10.1093/mnras/staa3564)
- , 2022a, *MNRAS*, 514, 3139, doi: [10.1093/mnras/stac1610](https://doi.org/10.1093/mnras/stac1610)
- Mohapatra, R., Jetti, M., Sharma, P., & Federrath, C. 2022b, *MNRAS*, 510, 2327, doi: [10.1093/mnras/stab3429](https://doi.org/10.1093/mnras/stab3429)
- Mohapatra, R., & Sharma, P. 2019, *MNRAS*, 484, 4881, doi: [10.1093/mnras/stz328](https://doi.org/10.1093/mnras/stz328)
- Mohapatra, R., Sharma, P., Federrath, C., & Quataert, E. 2023, *MNRAS*, 525, 3831, doi: [10.1093/mnras/stad2574](https://doi.org/10.1093/mnras/stad2574)
- Nelson, D., Sharma, P., Pillepich, A., et al. 2020, *MNRAS*, 498, 2391, doi: [10.1093/mnras/staa2419](https://doi.org/10.1093/mnras/staa2419)
- Nimmo, K., Pleunis, Z., Beniamini, P., et al. 2025, *Nature*, 637, 48, doi: [10.1038/s41586-024-08297-w](https://doi.org/10.1038/s41586-024-08297-w)
- Ocker, S. K., Cordes, J. M., & Chatterjee, S. 2021a, *ApJ*, 911, 102, doi: [10.3847/1538-4357/abeb6e](https://doi.org/10.3847/1538-4357/abeb6e)
- Ocker, S. K., Cordes, J. M., Chatterjee, S., & Dolch, T. 2021b, *ApJ*, 922, 233, doi: [10.3847/1538-4357/ac2b28](https://doi.org/10.3847/1538-4357/ac2b28)

- Ocker, S. K., Cordes, J. M., Chatterjee, S., & Gorsuch, M. R. 2022a, arXiv e-prints, arXiv:2203.16716. <https://arxiv.org/abs/2203.16716>
- Ocker, S. K., Cordes, J. M., Chatterjee, S., et al. 2022b, *ApJ*, 931, 87, doi: [10.3847/1538-4357/ac6504](https://doi.org/10.3847/1538-4357/ac6504)
- Padoan, P., Jones, B. J. T., & Nordlund, Å. P. 1997, *ApJ*, 474, 730, doi: [10.1086/303482](https://doi.org/10.1086/303482)
- Prochaska, J. X., & Zheng, Y. 2019, *MNRAS*, 485, 648, doi: [10.1093/mnras/stz261](https://doi.org/10.1093/mnras/stz261)
- Prochaska, J. X., Macquart, J.-P., McQuinn, M., et al. 2019, *Science*, 366, 231, doi: [10.1126/science.aay0073](https://doi.org/10.1126/science.aay0073)
- Qu, Z., Chen, H.-W., Rudie, G. C., et al. 2022, *MNRAS*, 516, 4882, doi: [10.1093/mnras/stac2528](https://doi.org/10.1093/mnras/stac2528)
- Rickett, B., Johnston, S., Tomlinson, T., & Reynolds, J. 2009, *MNRAS*, 395, 1391, doi: [10.1111/j.1365-2966.2009.14471.x](https://doi.org/10.1111/j.1365-2966.2009.14471.x)
- Rickett, B. J. 1977, *ARA&A*, 15, 479, doi: [10.1146/annurev.aa.15.090177.002403](https://doi.org/10.1146/annurev.aa.15.090177.002403)
- . 1990, *ARA&A*, 28, 561, doi: [10.1146/annurev.aa.28.090190.003021](https://doi.org/10.1146/annurev.aa.28.090190.003021)
- Rickett, B. J., Coles, W. A., & Bourgois, G. 1984, *A&A*, 134, 390
- Rudie, G. C., Steidel, C. C., Pettini, M., et al. 2019, *ApJ*, 885, 61, doi: [10.3847/1538-4357/ab4255](https://doi.org/10.3847/1538-4357/ab4255)
- Sammons, M. W., James, C. W., Trott, C. M., & Walker, M. 2022, *MNRAS*, 517, 5216, doi: [10.1093/mnras/stac3013](https://doi.org/10.1093/mnras/stac3013)
- Sarazin, C. L. 1986, *Reviews of Modern Physics*, 58, 1, doi: [10.1103/RevModPhys.58.1](https://doi.org/10.1103/RevModPhys.58.1)
- Sharma, P., McCourt, M., Parrish, I. J., & Quataert, E. 2012, *MNRAS*, 427, 1219, doi: [10.1111/j.1365-2966.2012.22050.x](https://doi.org/10.1111/j.1365-2966.2012.22050.x)
- Simard, D., & Ravi, V. 2021, arXiv e-prints, arXiv:2107.11334. <https://arxiv.org/abs/2107.11334>
- Singh, P., Majumdar, S., Nath, B. B., & Silk, J. 2018, *MNRAS*, 478, 2909, doi: [10.1093/mnras/sty1276](https://doi.org/10.1093/mnras/sty1276)
- Singh Bisht, M., Sharma, P., Dutta, A., & Nath, B. B. 2024, arXiv e-prints, arXiv:2411.17173, doi: [10.48550/arXiv.2411.17173](https://doi.org/10.48550/arXiv.2411.17173)
- Spangler, S. R., & Gwinn, C. R. 1990, *ApJL*, 353, L29, doi: [10.1086/185700](https://doi.org/10.1086/185700)
- Spitzer, L. 1956, *Physics of Fully Ionized Gases*
- Tan, B., Oh, S. P., & Gronke, M. 2021, *MNRAS*, 502, 3179, doi: [10.1093/mnras/stab053](https://doi.org/10.1093/mnras/stab053)
- Tumlinson, J., Peebles, M. S., & Werk, J. K. 2017, *ARA&A*, 55, 389, doi: [10.1146/annurev-astro-091916-055240](https://doi.org/10.1146/annurev-astro-091916-055240)
- van Leeuwen, J., Kooistra, E., Oostrum, L., et al. 2023, *A&A*, 672, A117, doi: [10.1051/0004-6361/202244107](https://doi.org/10.1051/0004-6361/202244107)
- Vedantham, H. K., & Phinney, E. S. 2019, *MNRAS*, 483, 971, doi: [10.1093/mnras/sty2948](https://doi.org/10.1093/mnras/sty2948)
- Voit, G. M. 2019, *ApJ*, 880, 139, doi: [10.3847/1538-4357/ab2bfd](https://doi.org/10.3847/1538-4357/ab2bfd)
- Werk, J. K., Prochaska, J. X., Tumlinson, J., et al. 2014, *ApJ*, 792, 8, doi: [10.1088/0004-637X/792/1/8](https://doi.org/10.1088/0004-637X/792/1/8)
- Wu, Z.-W., Main, R. A., Zhu, W.-W., et al. 2024, *Science China Physics, Mechanics, and Astronomy*, 67, 219512, doi: [10.1007/s11433-023-2186-2](https://doi.org/10.1007/s11433-023-2186-2)
- Zhuravleva, I., Churazov, E., Schekochihin, A. A., et al. 2019, *Nature Astronomy*, 3, 832, doi: [10.1038/s41550-019-0794-z](https://doi.org/10.1038/s41550-019-0794-z)
- Zhuravleva, I., Churazov, E. M., Schekochihin, A. A., et al. 2014, *ApJL*, 788, L13, doi: [10.1088/2041-8205/788/1/L13](https://doi.org/10.1088/2041-8205/788/1/L13)