

# ON TERWILLIGER $\mathbb{F}$ -ALGEBRAS OF DIRECT PRODUCTS OF GROUP DIVISIBLE ASSOCIATION SCHEMES

YU JIANG

**ABSTRACT.** The Terwilliger algebras of association schemes over an arbitrary field  $\mathbb{F}$  were briefly called the Terwilliger  $\mathbb{F}$ -algebras of association schemes in [9]. In this paper, the Terwilliger  $\mathbb{F}$ -algebras of direct products of group divisible association schemes are studied. The centers, the semisimplicity, the Jacobson radicals and their nilpotent indices, the Wedderburn-Artin decompositions of the Terwilliger  $\mathbb{F}$ -algebras of direct products of group divisible association schemes are obtained.

**Keywords.** Terwilliger algebra; Direct product; Group divisible association scheme  
**Mathematics Subject Classification 2020.** 05E30 (primary), 05E16 (secondary)

## 1. INTRODUCTION

Association schemes, briefly called schemes, are intensively studied as important objects in algebraic combinatorics. In particular, they are known to have connections with many different mathematical objects, such as groups, graphs, codes, designs, and so on. Conversely, many different tools have already been used to study schemes.

The subconstituent algebras of commutative schemes, introduced by Terwilliger in [17], are algebraic tools of studying schemes. They are finite-dimensional semisimple associative  $\mathbb{C}$ -algebras and are also known as the Terwilliger algebras of commutative schemes. In [6], Hanaki defined the Terwilliger algebras for an arbitrary scheme and an arbitrary commutative unital ring. In [9], we briefly called the Terwilliger algebras of schemes over an arbitrary field  $\mathbb{F}$  the Terwilliger  $\mathbb{F}$ -algebras of schemes. So the Terwilliger algebras of commutative schemes are exactly their Terwilliger  $\mathbb{C}$ -algebras.

The Terwilliger  $\mathbb{C}$ -algebras of many commutative schemes have been investigated (for example, see [2, 4, 13, 14, 15, 16, 17, 18, 19, 20]). However, the investigation of the Terwilliger  $\mathbb{F}$ -algebras of schemes is almost completely open (see [7]). In [3, 9], the authors presented some ring-theoretical properties of the Terwilliger  $\mathbb{F}$ -algebras of quasi-thin schemes. In this paper, we study the Terwilliger  $\mathbb{F}$ -algebras of direct products of group divisible schemes from the viewpoint of ring theory. As the main results of this paper, we obtain the centers, the semisimplicity, the Jacobson radicals and their nilpotent indices, the Wedderburn-Artin decompositions of these algebras (see Theorems 4.15, 5.21, 6.8, 8.13, respectively). The main results of this paper contribute to understanding the Terwilliger  $\mathbb{F}$ -algebras of direct products of schemes.

The outline of this paper is as follows: In Section 2, we list the basic notation and preliminaries. In Section 3, we give two  $\mathbb{F}$ -bases for a Terwilliger  $\mathbb{F}$ -algebra of a direct product of group divisible schemes. Theorems 4.15, 5.21, 6.8 are proved in Sections 4, 5, 6, respectively. Sections 7 and 8 contain the proof of Theorem 8.13.

## 2. BASIC NOTATION AND PRELIMINARIES

For a general background on association schemes, the reader may refer to [1, 22].

**2.1. Conventions.** Let  $\mathbb{N}$  be the set of all natural numbers. Define  $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$ . If  $g, h \in \mathbb{N}_0$ , set  $[g, h] = \{a : a \in \mathbb{N}_0, g \leq a \leq h\}$ . Fix a nonempty finite set  $\mathbb{X}$ . Each association scheme on  $\mathbb{X}$  is briefly called a scheme. Fix a field  $\mathbb{F}$  of characteristic  $p$ . Let  $\mathbb{F}^\times$  be the set of all nonzero elements in  $\mathbb{F}$ . The addition, the multiplication, the scalar multiplication of  $\mathbb{F}$ -matrices in this paper are usual matrix operations. For any subset  $\mathbb{U}$  of an  $\mathbb{F}$ -linear space  $\mathbb{V}$ , let  $\langle \mathbb{U} \rangle$  be the  $\mathbb{F}$ -linear subspace of  $\mathbb{V}$  spanned by  $\mathbb{U}$ . All  $\mathbb{F}$ -algebras in this paper are finite-dimensional associative unital algebras. All modules in this paper mean finite-dimensional left modules of some  $\mathbb{F}$ -algebras.

**2.2. Schemes.** Let  $\mathbb{S} = \{R_0, R_1, \dots, R_d\}$  denote a partition of the cartesian product  $\mathbb{X} \times \mathbb{X}$ . Then  $\mathbb{S}$  is called a  $d$ -class scheme if the following conditions hold together:

- (S1)  $R_0 = \{(a, a) : a \in \mathbb{X}\}$ ;
- (S2) If  $g \in [0, d]$ , then there is  $g^* \in [0, d]$  such that  $R_{g^*} = \{(a, b) : (b, a) \in R_g\} \in \mathbb{S}$ ;
- (S3) If  $g, h, i \in [0, d]$  and  $(x, y), (\tilde{x}, \tilde{y}) \in R_i$ , then there is a constant  $p_{gh}^i$  such that

$$p_{gh}^i = |\{a : (x, a) \in R_g, (a, y) \in R_h\}| = |\{a : (\tilde{x}, a) \in R_g, (a, \tilde{y}) \in R_h\}|.$$

From now on,  $\mathbb{S} = \{R_0, R_1, \dots, R_d\}$  is a fixed  $d$ -class scheme. Call  $\mathbb{S}$  a symmetric scheme if  $g = g^*$  for any  $g \in [0, d]$ . Call  $\mathbb{S}$  a commutative scheme if  $p_{gh}^i = p_{hg}^i$  for any  $g, h, i \in [0, d]$ . Notice that the symmetric schemes are commutative schemes. Set  $xR_g = \{(x, a) \in R_g\}$  and  $k_g = p_{gg}^0$  for any  $x \in \mathbb{X}$  and  $g \in [0, d]$ . If  $x, y \in \mathbb{X}$  and  $g \in [0, d]$ , (S3) and the inequality  $R_g \neq \emptyset$  imply that  $k_g = |xR_g| = |yR_g| > 0$ . Call  $\mathbb{S}$  a  $p'$ -valenced scheme if  $p \nmid k_g$  for any  $g \in [0, d]$ . Hence each scheme is a  $0'$ -valenced scheme. If  $x, y \in \mathbb{X}$ ,  $g, h \in [0, d]$ ,  $y \in xR_i$ , then  $p_{gh}^i = |xR_g \cap yR_{h^*}|$  by (S2) and (S3). Call  $\mathbb{S}$  a triply regular scheme if, for any  $x, y, z \in \mathbb{X}$ ,  $g, h, i, j, k, \ell \in [0, d]$ ,  $y \in xR_j$ , and  $z \in xR_k \cap yR_\ell$ ,  $|xR_g \cap yR_h \cap zR_i|$  only depends on the subscripts  $g, h, i, j, k, \ell$  and is independent of the choices of  $x, y, z$  satisfying  $y \in xR_j$  and  $z \in xR_k \cap yR_\ell$ .

For any nonempty subsets  $\mathbb{U}, \mathbb{V}$  of  $\mathbb{S}$ , set  $\mathbb{U}\mathbb{V} = \{R_a : \exists R_b \in \mathbb{U}, \exists R_c \in \mathbb{V}, p_{bc}^a \neq 0\}$ . Hence  $\mathbb{S}$  is a commutative scheme if and only if  $\mathbb{U}\mathbb{V} = \mathbb{V}\mathbb{U}$  for any nonempty subsets  $\mathbb{U}, \mathbb{V}$  of  $\mathbb{S}$ . For convenience, utilize  $R_g R_h$  to denote  $\{R_g\}\{R_h\}$  for any  $g, h \in [0, d]$ .

Assume that  $g, h \in \mathbb{N}$  and  $\mathbb{U}$  denotes a set with cardinality  $gh$ . Call a partition of  $\mathbb{U}$  a group divisible  $(g, h)$ -partition of  $\mathbb{U}$  if this partition has  $g$  members and each member in this partition has cardinality  $h$ . Call each member in a group divisible  $(g, h)$ -partition of  $\mathbb{U}$  a  $\mathbb{U}$ -group. Notice that a given  $\mathbb{U}$ -group is not necessary to have a finite group structure. Assume that  $x, y \in \mathbb{U}$  and  $\mathbb{G}$  denotes a group divisible  $(g, h)$ -partition of  $\mathbb{U}$ . Write  $x \sim_{\mathbb{G}} y$  if  $x \neq y$  and  $x, y$  belong to a single  $\mathbb{U}$ -group in  $\mathbb{G}$ . For completeness, write  $x \not\sim_{\mathbb{G}} y$  if  $x, y$  belong to two distinct  $\mathbb{U}$ -groups in  $\mathbb{G}$ .

Assume that  $g, h \in \mathbb{N}$ ,  $g, h \geq 2$ , and  $|\mathbb{X}| = gh$ . Assume that  $\mathbb{G}$  is a group divisible  $(g, h)$ -partition of  $\mathbb{X}$ . Call  $\mathbb{S}$  a group divisible scheme of parameter  $(g, h)$  if  $d = 2$ ,  $R_1 = \{(a, b) : a, b \in \mathbb{X}, a \sim_{\mathbb{G}} b\}$ ,  $R_2 = \{(a, b) : a, b \in \mathbb{X}, a \not\sim_{\mathbb{G}} b\}$  (see [1, 12]). If  $\mathbb{S}$  is a group divisible scheme of parameter  $(g, h)$ , recall that  $\mathbb{S}$  is a triply regular scheme (see [16, Lemma 5]). See [1, 12] for many applications of a group divisible scheme.

Fix  $n \in \mathbb{N}$ . If  $g \in [0, 3^n - 1]$ , let  $\sum_{h=1}^n g_{[h]} 3^{h-1}$  denote the 3-adic decomposition of  $g$  with  $n$  bits. Define  $\mathfrak{g}_0 = \{a : a \in [1, n], g_{[a]} = 0\}$ ,  $\mathfrak{g}_1 = \{a : a \in [1, n], g_{[a]} = 1\}$ , and  $\mathfrak{g}_2 = \{a : a \in [1, n], g_{[a]} = 2\}$  for any  $g \in [0, 3^n - 1]$ . Let  $\ell_1, \ell_2, \dots, \ell_n, m_1, m_2, \dots, m_n$  be a fixed sequence in  $\mathbb{N} \setminus \{1\}$ . Let  $n_1, n_2, n_3, n_4$  denote  $|\{a : a \in [1, n], \ell_a > m_a = 2\}|$ ,  $|\{a : a \in [1, n], \ell_a = m_a = 2\}|$ ,  $|\{a : a \in [1, n], \ell_a, m_a > 2\}|$ ,  $|\{a : a \in [1, n], m_a > \ell_a = 2\}|$ , respectively. For any sets  $\mathbb{U}, \mathbb{U}^1, \mathbb{V}, \mathbb{V}^1, \mathbb{W}, \mathbb{W}^1$ , let  $\mathbb{U} \Delta \mathbb{V} = (\mathbb{U} \setminus \mathbb{V}) \cup (\mathbb{V} \setminus \mathbb{U})$  and put  $(\mathbb{U}, \mathbb{V}, \mathbb{W}) \preceq (\mathbb{U}^1, \mathbb{V}^1, \mathbb{W}^1)$  if  $\mathbb{U} \subseteq \mathbb{U}^1, \mathbb{V} \subseteq \mathbb{V}^1, \mathbb{W} \subseteq \mathbb{W}^1$ . So  $\preceq$  is a partial order among all set triples. If  $\mathbb{U} \subseteq [1, n]$ , set  $\mathbb{U}^\bullet = \{a : a \in \mathbb{U}, \ell_a > 2\}$  and  $\mathbb{U}^\circ = \{a : a \in \mathbb{U}, m_a > 2\}$ .

Fix a set sequence  $\mathbb{U}_1, \mathbb{U}_2, \dots, \mathbb{U}_n$  that satisfies  $|\mathbb{U}_g| = \ell_g m_g$  for any  $g \in [1, n]$ . For any  $g \in [1, n]$ , fix a group divisible  $(\ell_g, m_g)$ -partition  $\mathbb{G}_g$  of  $\mathbb{U}_g$ . The cartesian product of  $\mathbb{U}_1, \mathbb{U}_2, \dots, \mathbb{U}_n$  is denoted by  $\mathbb{U}_1 \times \mathbb{U}_2 \times \dots \times \mathbb{U}_n$ . If  $\mathbf{u} \in \mathbb{U}_1 \times \mathbb{U}_2 \times \dots \times \mathbb{U}_n$ , let  $\mathbf{u}_g$  be the  $\mathbb{U}_g$ -component of  $\mathbf{u}$ . For any  $\mathbf{u}, \mathbf{v} \in \mathbb{U}_1 \times \mathbb{U}_2 \times \dots \times \mathbb{U}_n$  and  $g \in [0, 3^n - 1]$ , write  $\mathbf{u} =_g \mathbf{v}$  to indicate that  $\mathbf{u}_h \sim_{\mathbb{G}_h} \mathbf{v}_h$  and  $\mathbf{u}_i \not\sim_{\mathbb{G}_i} \mathbf{v}_i$  if and only if  $h \in \mathfrak{g}_1$  and  $i \in \mathfrak{g}_2$ . Following [1], call  $\mathbb{S}$  a direct product of group divisible schemes of parameters  $(\ell_1, m_1), (\ell_2, m_2), \dots, (\ell_n, m_n)$  if  $\mathbb{X} = \mathbb{U}_1 \times \mathbb{U}_2 \times \dots \times \mathbb{U}_n, d = 3^n - 1$ , and  $R_g$  is defined to be  $\{(\mathbf{a}, \mathbf{b}) : \mathbf{a}, \mathbf{b} \in \mathbb{X}, \mathbf{a} =_g \mathbf{b}\}$  for any  $g \in [1, d]$ . If  $\mathbb{S}$  is a direct product of group divisible schemes of parameters  $(\ell_1, m_1), (\ell_2, m_2), \dots, (\ell_n, m_n)$ , then  $\mathbb{S}$  is a symmetric scheme. Moreover, recall that  $\mathbb{S}$  is a triply regular scheme (see [21, Theorem 10]).

**2.3. Algebras.** Let  $\mathbb{Z}$  be the integer ring. Let  $\mathbb{F}_p$  be the prime subfield of  $\mathbb{F}$ . Given  $g \in \mathbb{Z}$ , let  $\bar{g}$  be the image of  $g$  under the unital ring homomorphism from  $\mathbb{Z}$  to  $\mathbb{F}_p$ .

Let  $\mathbb{A}$  be an  $\mathbb{F}$ -algebra with the identity element  $1_{\mathbb{A}}$  and the zero element  $0_{\mathbb{A}}$ . The center of  $\mathbb{A}$ , denoted by  $Z(\mathbb{A})$ , is defined to be  $\{a : a \in \mathbb{A}, ab = ba \forall b \in \mathbb{A}\}$ . So  $Z(\mathbb{A})$  is an  $\mathbb{F}$ -subalgebra of  $\mathbb{A}$  with the identity element  $1_{\mathbb{A}}$ . Let  $\mathbb{I}$  denote a two-sided ideal of  $\mathbb{A}$ . Let  $\mathbb{A}/\mathbb{I}$  be the quotient  $\mathbb{F}$ -algebra of  $\mathbb{A}$  with respect to  $\mathbb{I}$ . Call  $\mathbb{I}$  a nilpotent two-sided ideal of  $\mathbb{A}$  if there is  $h \in \mathbb{N}$  such that the product of any  $h$  elements in  $\mathbb{I}$  equals  $0_{\mathbb{A}}$ . If  $\mathbb{I}$  is a nilpotent two-sided ideal of  $\mathbb{A}$ , then the nilpotent index of  $\mathbb{I}$  is defined to be the smallest choice of  $h$ . The Jacobson radical  $\text{Rad}(\mathbb{A})$  of  $\mathbb{A}$  is the sum of all nilpotent two-sided ideals of  $\mathbb{A}$ . Recall that  $\text{Rad}(\mathbb{A})$  is also a nilpotent two-sided ideal of  $\mathbb{A}$ . Call  $\mathbb{A}$  a semisimple  $\mathbb{F}$ -algebra if  $\mathbb{A}$  equals a direct sum of its minimal two-sided ideals. It is already known that  $\mathbb{A}$  is a semisimple  $\mathbb{F}$ -algebra if and only if  $\text{Rad}(\mathbb{A}) = \{0_{\mathbb{A}}\}$ . Let  $e$  be an idempotent of  $\mathbb{A}$ . Set  $e\mathbb{I}e = \{eae : a \in \mathbb{I}\}$ . Notice that  $e\mathbb{I}e$  is an  $\mathbb{F}$ -subalgebra of  $\mathbb{A}$  with the identity element  $e$ .

Let  $\mathbb{V}$  be an  $\mathbb{A}$ -module. Call  $\mathbb{V}$  an irreducible  $\mathbb{A}$ -module if  $\mathbb{V}$  has no nonzero proper  $\mathbb{A}$ -submodule. Let  $\mathbb{K}$  be an extension field of  $\mathbb{F}$ . Let  $\mathbb{A}_{\mathbb{K}}$  be the tensor product  $\mathbb{K} \otimes_{\mathbb{F}} \mathbb{A}$  of  $\mathbb{F}$ -algebras. Let  $\mathbb{V}_{\mathbb{K}}$  be the tensor product  $\mathbb{K} \otimes_{\mathbb{F}} \mathbb{V}$  of  $\mathbb{F}$ -linear spaces. So  $\mathbb{V}_{\mathbb{K}}$  is an  $\mathbb{A}_{\mathbb{K}}$ -module by the diagonal action. Call  $\mathbb{V}$  an absolutely irreducible  $\mathbb{A}$ -module if  $\mathbb{V}_{\mathbb{L}}$  is an irreducible  $\mathbb{A}_{\mathbb{L}}$ -module for any extension field  $\mathbb{L}$  of  $\mathbb{F}$ . Two lemmas are needed.

**Lemma 2.1.** [5, Proposition 3.2.4] *Assume that  $e$  denotes an idempotent of  $\mathbb{A}$ . Then  $\text{Rad}(e\mathbb{A}e) = e\text{Rad}(\mathbb{A})e$ . In particular,  $\text{Rad}(e\mathbb{A}e) \subseteq \text{Rad}(\mathbb{A})$ .*

**Lemma 2.2.** [5, 11, Corollaries 3.1.7, 1.10.4] *Assume that  $g \in \mathbb{N}$  and  $\mathbb{A}/\text{Rad}(\mathbb{A})$  is isomorphic to a direct sum of  $g$  full matrix  $\mathbb{F}$ -algebras of square  $\mathbb{F}$ -matrices. Then the number of all pairwise nonisomorphic irreducible  $\mathbb{A}$ -modules equals  $g$ . Moreover, all irreducible  $\mathbb{A}$ -modules are precisely all absolutely irreducible  $\mathbb{A}$ -modules.*

**2.4. Terwilliger  $\mathbb{F}$ -algebras of schemes.** If  $g \in \mathbb{N}$ , let  $M_g(\mathbb{F})$  be the full matrix  $\mathbb{F}$ -algebra of  $(g \times g)$ -matrices whose entries are in  $\mathbb{F}$ . If  $g, h \in \mathbb{N}$ , let  $gM_h(\mathbb{F})$  denote the direct sum of  $g$  copies of  $M_h(\mathbb{F})$ . If  $g, h, i, j \in \mathbb{N}$ , notice that  $gM_h(\mathbb{F}) \cong iM_j(\mathbb{F})$  as  $\mathbb{F}$ -algebras if and only if  $g = i$  and  $h = j$ . Let  $\mathbb{U}$  be a nonempty finite set. Let  $M_{\mathbb{U}}(\mathbb{F})$  denote the full matrix  $\mathbb{F}$ -algebra of square  $\mathbb{F}$ -matrices whose rows and columns are labeled by the elements from  $\mathbb{U}$ . Then  $M_{\mathbb{U}}(\mathbb{F}) \cong M_{|\mathbb{U}|}(\mathbb{F})$  as  $\mathbb{F}$ -algebras. Let  $I, J, O$  be the identity matrix, the all-one matrix, the all-zero matrix in  $M_{\mathbb{X}}(\mathbb{F})$ , respectively. If  $x, y \in \mathbb{X}$ , let  $E_{x,y}$  be the  $\{\bar{0}, \bar{1}\}$ -matrix in  $M_{\mathbb{X}}(\mathbb{F})$  whose unique nonzero entry is the  $(x, y)$ -entry. The transpose of a matrix  $M$  in  $M_{\mathbb{X}}(\mathbb{F})$  is denoted by  $M^T$ .

If  $g, h \in \mathbb{N}_0$ , let  $\delta_{g,h}$  be the Kronecker delta of  $g$  and  $h$  whose values are from  $\mathbb{F}$ . If  $g \in [0, d]$ , the adjacency  $\mathbb{F}$ -matrix  $A_g$  with respect to  $R_g$  is defined to be  $\sum_{(x,y) \in R_g} E_{x,y}$ . If  $x \in \mathbb{X}$  and  $g \in [0, d]$ , the dual  $\mathbb{F}$ -idempotent  $E_g^*(x)$  with respect to  $x$  and  $R_g$  is defined to be  $\sum_{y \in xR_g} E_{y,y}$ . If  $x \in \mathbb{X}$  and  $g, h \in [0, d]$ , the following equations hold:

$$(2.1) \quad A_g^T = A_{g^*} \text{ and } E_g^*(x)^T = E_g^*(x) \text{ (} A_g^T = A_g \text{ if } \mathbb{S} \text{ is a symmetric scheme),}$$

$$(2.2) \quad E_g^*(x)E_h^*(x) = \delta_{g,h}E_g^*(x),$$

$$(2.3) \quad A_0 = I = \sum_{i=0}^d E_i^*(x).$$

Pick  $x \in \mathbb{X}$ . The Terwilliger  $\mathbb{F}$ -algebra of  $\mathbb{S}$  with respect to  $x$ , denoted by  $\mathbb{T}(x)$ , is the  $\mathbb{F}$ -subalgebra of  $M_{\mathbb{X}}(\mathbb{F})$  generated by  $A_0, A_1, \dots, A_d, E_0^*(x), E_1^*(x), \dots, E_d^*(x)$ . Therefore Equation (2.3) implies that the identity element of  $\mathbb{T}(x)$  equals  $I$ . By the definition of  $\mathbb{T}(x)$  and Equation (2.1), notice that  $M \in \mathbb{T}(x)$  if and only if  $M^T \in \mathbb{T}(x)$ .

If  $g, h, i \in [0, d]$ , then  $E_g^*(x)A_hE_i^*(x) \neq O$  if and only if  $p_{g^*i}^h \neq 0$  (see [6, Lemma 3.2]). So  $\mathbb{T}(x)$  has an  $\mathbb{F}$ -linearly independent subset  $\{E_a^*(x)A_bE_c^*(x) : a, b, c \in [0, d], p_{a^*c}^b \neq 0\}$  by the definition of  $\mathbb{T}(x)$  and Equation (2.2). In general, the algebraic structures of  $\mathbb{T}(x)$  and  $\text{Rad}(\mathbb{T}(x))$  may depend on the choices of the fixed field  $\mathbb{F}$  and the given  $x$  (see [6, 5.1]). See [3, 6, 8, 9, 10] for more progress on the algebraic structures of  $\mathbb{T}(x)$  and  $\text{Rad}(\mathbb{T}(x))$ . The following three lemmas are required in the following sections.

**Lemma 2.3.** [6, Theorem 3.4] *Assume that  $x \in \mathbb{X}$ .  $\mathbb{T}(x)$  is a semisimple  $\mathbb{F}$ -algebra only if  $\mathbb{S}$  is a  $p'$ -valenced scheme.*

**Lemma 2.4.** [16, Lemma 4] *Assume that  $x \in \mathbb{X}$  and  $\mathbb{S}$  is a triply regular scheme.  $\mathbb{T}(x)$  has an  $\mathbb{F}$ -basis  $\{E_a^*(x)A_bE_c^*(x) : a, b, c \in [0, d], p_{a^*c}^b \neq 0\}$ .*

**Lemma 2.5.** [10, Lemma 2.10] *Assume that  $x \in \mathbb{X}$  and  $\mathbb{S}$  is a symmetric scheme. If  $\mathbb{S}$  is also a triply regular scheme, then the  $\mathbb{F}$ -subalgebra  $E_g^*(x)\mathbb{T}(x)E_g^*(x)$  of  $\mathbb{T}(x)$  is a commutative  $\mathbb{F}$ -algebra for any  $g \in [0, d]$ .*

We end this section by simplifying the notation. From now on, assume that  $\mathbb{S}$  is a direct product of group divisible schemes of parameters  $(\ell_1, m_1), (\ell_2, m_2), \dots, (\ell_n, m_n)$ . We shall quote the fact that  $\mathbb{S}$  is a symmetric scheme without reference. Moreover, we shall also quote the fact that  $\mathbb{S}$  is a triply regular scheme without reference. Fix  $x \in \mathbb{X}$ . For convenience, we abbreviate  $\mathbb{T} = \mathbb{T}(x)$  and  $E_g^* = E_g^*(x)$  for any  $g \in [0, d]$ .

3. ALGEBRAIC STRUCTURE OF  $\mathbb{T}$ :  $\mathbb{F}$ -BASIS

In this section, we present two  $\mathbb{F}$ -bases for  $\mathbb{T}$ . We start with a sequence of lemmas.

**Lemma 3.1.** *Assume that  $g, h \in [0, d]$ . Then  $g = h$  if and only if  $\mathfrak{g}_1 = \mathfrak{h}_1$  and  $\mathfrak{g}_2 = \mathfrak{h}_2$ . Moreover,  $\mathfrak{g}_0, \mathfrak{g}_1, \mathfrak{g}_2$  are pairwise disjoint and  $[1, n] = \mathfrak{g}_0 \cup \mathfrak{g}_1 \cup \mathfrak{g}_2$ .*

*Proof.* The desired lemma follows from the 3-adic decompositions of integers.  $\square$

**Lemma 3.2.** *Assume that  $g, h, i \in [0, d]$ . Then  $p_{gh}^i \neq 0$  only if*

$$(3.1) \quad (\mathfrak{g}_0 \cap \mathfrak{h}_1) \cup (\mathfrak{g}_1 \cap \mathfrak{h}_0) \subseteq \mathfrak{i}_1 \subseteq (\mathfrak{g}_0 \cap \mathfrak{h}_1) \cup (\mathfrak{g}_1 \cap \mathfrak{h}_0) \cup (\mathfrak{g}_1 \cap \mathfrak{h}_1)^\circ \cup (\mathfrak{g}_2 \cap \mathfrak{h}_2).$$

*Proof.* As  $p_{gh}^i \neq 0$ , there are  $\mathbf{y}, \mathbf{z}, \mathbf{w} \in \mathbb{X}$  such that  $\mathbf{w} \in \mathbf{y}R_g \cap \mathbf{z}R_h$  and  $\mathbf{z} \in \mathbf{y}R_i$ . If  $j \in \mathfrak{g}_0 \cap \mathfrak{h}_1$ , notice that  $\mathbf{y}_j = \mathbf{w}_j \sim_{\mathbb{G}_j} \mathbf{z}_j$  and  $j \in \mathfrak{i}_1$ . Similarly, notice that  $j \in \mathfrak{i}_1$  if  $j \in \mathfrak{g}_1 \cap \mathfrak{h}_0$ . If  $j \in \mathfrak{g}_0 \cap \mathfrak{i}_1$ , then  $\mathbf{w}_j = \mathbf{y}_j \sim_{\mathbb{G}_j} \mathbf{z}_j$  and  $j \in \mathfrak{g}_0 \cap \mathfrak{h}_1$ . If  $j \in \mathfrak{g}_1 \cap \mathfrak{i}_1$ , then  $\mathbf{w}_j \sim_{\mathbb{G}_j} \mathbf{y}_j \sim_{\mathbb{G}_j} \mathbf{z}_j$ , which yields that  $j \in \mathfrak{g}_1 \cap \mathfrak{h}_0$  or  $j \in \mathfrak{g}_1 \cap \mathfrak{h}_1$  and  $m_j > 2$ . Similarly,  $j \in \mathfrak{g}_2 \cap \mathfrak{h}_2$  if  $j \in \mathfrak{g}_2 \cap \mathfrak{i}_1$ . The desired lemma thus follows from Lemma 3.1.  $\square$

**Lemma 3.3.** *Assume that  $g, h, i \in [0, d]$ . Then  $p_{gh}^i \neq 0$  only if*

$$(3.2) \quad \mathfrak{g}_2 \Delta \mathfrak{h}_2 \subseteq \mathfrak{i}_2 \subseteq (\mathfrak{g}_2 \Delta \mathfrak{h}_2) \cup (\mathfrak{g}_2 \cap \mathfrak{h}_2)^\bullet.$$

*Proof.* As  $p_{gh}^i \neq 0$ , there are  $\mathbf{y}, \mathbf{z}, \mathbf{w} \in \mathbb{X}$  such that  $\mathbf{w} \in \mathbf{y}R_g \cap \mathbf{z}R_h$  and  $\mathbf{z} \in \mathbf{y}R_i$ . If  $j \in \mathfrak{g}_2 \setminus \mathfrak{h}_2$ , then  $\mathbf{y}_j \not\sim_{\mathbb{G}_j} \mathbf{w}_j = \mathbf{z}_j$  or  $\mathbf{y}_j \not\sim_{\mathbb{G}_j} \mathbf{w}_j \sim_{\mathbb{G}_j} \mathbf{z}_j$ , which suggests that  $j \in \mathfrak{i}_2$ . Similarly,  $j \in \mathfrak{i}_2$  if  $j \in \mathfrak{h}_2 \setminus \mathfrak{g}_2$ . If  $j \in (\mathfrak{g}_0 \cap \mathfrak{i}_2) \cup (\mathfrak{g}_1 \cap \mathfrak{i}_2)$ , then  $\mathbf{w}_j = \mathbf{y}_j \not\sim_{\mathbb{G}_j} \mathbf{z}_j$  or  $\mathbf{w}_j \sim_{\mathbb{G}_j} \mathbf{y}_j \not\sim_{\mathbb{G}_j} \mathbf{z}_j$ , which yields that  $j \in \mathfrak{h}_2 \setminus \mathfrak{g}_2$ . If  $j \in \mathfrak{g}_2 \cap \mathfrak{h}_2 \cap \mathfrak{i}_2$ , notice that  $\mathbf{w}_j \not\sim_{\mathbb{G}_j} \mathbf{y}_j \not\sim_{\mathbb{G}_j} \mathbf{z}_j$  and  $\ell_j > 2$ . The desired lemma thus follows from Lemma 3.1.  $\square$

Lemmas 3.2 and 3.3 motivate us to give the following notation and four lemmas.

**Notation 3.4.** *Use  $\mathbb{P}$  to denote the set of all triples  $(g, h, i)$  in the cartesian product  $[1, d] \times [1, d] \times [1, d]$  that satisfy Equations (3.1) and (3.2). Let  $g, h, i \in [0, d]$ . Define  $\mathbb{P}_{g,h} = \{a : (g, h, a) \in \mathbb{P}\}$ . So Lemmas 3.2 and 3.3 imply that  $p_{gh}^i \neq 0$  only if  $i \in \mathbb{P}_{g,h}$ .*

**Lemma 3.5.** *Assume that  $g, h, i \in [0, d]$ . Then  $p_{gh}^i \neq 0$  if and only if  $(g, h, i) \in \mathbb{P}$ .*

*Proof.* Assume that  $(g, h, i) \in \mathbb{P}$ . Then there exist  $\mathfrak{j} \subseteq (\mathfrak{g}_1 \cap \mathfrak{h}_1)^\circ$ ,  $\mathfrak{k} \subseteq (\mathfrak{g}_2 \cap \mathfrak{h}_2)^\bullet$ , and  $\mathfrak{l} \subseteq (\mathfrak{g}_2 \cap \mathfrak{h}_2) \setminus \mathfrak{k}$  such that  $\mathfrak{i}_1 = (\mathfrak{g}_0 \cap \mathfrak{h}_1) \cup (\mathfrak{g}_1 \cap \mathfrak{h}_0) \cup \mathfrak{j} \cup \mathfrak{l}$  and  $\mathfrak{i}_2 = (\mathfrak{g}_2 \Delta \mathfrak{h}_2) \cup \mathfrak{k}$ . Assume that  $\mathbf{y}, \mathbf{z} \in \mathbb{X}$  and  $\mathbf{z} \in \mathbf{y}R_i$ . So  $\mathbf{y}_m = \mathbf{z}_m$ ,  $\mathbf{y}_o \sim_{\mathbb{G}_o} \mathbf{z}_o$ ,  $\mathbf{y}_q \not\sim_{\mathbb{G}_q} \mathbf{z}_q$  for any  $m \in \mathfrak{i}_0$ ,  $o \in \mathfrak{i}_1$ , and  $q \in \mathfrak{i}_2$ . Then there is  $\mathbf{w} \in \mathbb{X}$  such that  $\mathbf{w}_m = \mathbf{y}_m = \mathbf{z}_m$ ,  $\mathbf{w}_o \sim_{\mathbb{G}_o} \mathbf{y}_o = \mathbf{z}_o$ ,  $\mathbf{w}_q \not\sim_{\mathbb{G}_q} \mathbf{y}_q = \mathbf{z}_q$ ,  $\mathbf{w}_r = \mathbf{y}_r \sim_{\mathbb{G}_r} \mathbf{z}_r$ ,  $\mathbf{w}_s = \mathbf{z}_s \sim_{\mathbb{G}_s} \mathbf{y}_s$ ,  $\mathbf{w}_t \not\sim_{\mathbb{G}_t} \mathbf{y}_t \sim_{\mathbb{G}_t} \mathbf{z}_t$ ,  $\mathbf{w}_u = \mathbf{z}_u \not\sim_{\mathbb{G}_u} \mathbf{y}_u$ ,  $\mathbf{w}_v \sim_{\mathbb{G}_v} \mathbf{z}_v \not\sim_{\mathbb{G}_v} \mathbf{y}_v$ ,  $\mathbf{w}_w = \mathbf{y}_w \not\sim_{\mathbb{G}_w} \mathbf{z}_w$ ,  $\mathbf{w}_x \sim_{\mathbb{G}_x} \mathbf{y}_x \not\sim_{\mathbb{G}_x} \mathbf{z}_x$ ,  $\mathbf{w}_y \sim_{\mathbb{G}_y} \mathbf{y}_y \sim_{\mathbb{G}_y} \mathbf{z}_y$  and  $\mathbf{w}_y \sim_{\mathbb{G}_y} \mathbf{z}_y$ ,  $\mathbf{w}_z \not\sim_{\mathbb{G}_z} \mathbf{y}_z \not\sim_{\mathbb{G}_z} \mathbf{z}_z$  and  $\mathbf{w}_z \not\sim_{\mathbb{G}_z} \mathbf{z}_z$  for any  $m \in \mathfrak{g}_0 \cap \mathfrak{h}_0 \cap \mathfrak{i}_0$ ,  $o \in \mathfrak{g}_1 \cap \mathfrak{h}_1 \cap \mathfrak{i}_0$ ,  $q \in \mathfrak{g}_2 \cap \mathfrak{h}_2 \cap \mathfrak{i}_0$ ,  $r \in \mathfrak{g}_0 \cap \mathfrak{h}_1$ ,  $s \in \mathfrak{g}_1 \cap \mathfrak{h}_0$ ,  $t \in \mathfrak{l}$ ,  $u \in \mathfrak{g}_2 \cap \mathfrak{h}_0$ ,  $v \in \mathfrak{g}_2 \cap \mathfrak{h}_1$ ,  $w \in \mathfrak{g}_0 \cap \mathfrak{h}_2$ ,  $x \in \mathfrak{g}_1 \cap \mathfrak{h}_2$ ,  $y \in \mathfrak{j}$ ,  $z \in \mathfrak{k}$ . Hence  $\mathbf{w} \in \mathbf{y}R_g \cap \mathbf{z}R_h$  and  $p_{gh}^i \neq 0$  by Lemma 3.1. The other direction follows from Lemmas 3.2 and 3.3. The desired lemma follows.  $\square$

**Lemma 3.6.**  *$\mathbb{T}$  has an  $\mathbb{F}$ -basis  $\{E_a^* A_b E_c^* : (a, c, b) \in \mathbb{P}\}$  whose cardinality equals  $|\mathbb{P}|$ .*

*Proof.* The desired lemma follows from an application of Lemmas 2.4 and 3.5.  $\square$

**Lemma 3.7.** *Assume that  $g, h, i, j, k \in [0, d]$ . Then*

$$E_g^* A_h E_i^* A_j E_k^* = \sum_{(g,k,\ell) \in \mathbb{P}} c_{g,h,i,j,k,\ell} E_g^* A_\ell E_k^*,$$

where  $c_{g,h,i,j,k,\ell} = \overline{|\mathbf{y}R_h \cap \mathbf{x}R_i \cap \mathbf{z}R_j|} \in \mathbb{F}$  for any  $\mathbf{y} \in \mathbf{x}R_g$  and  $\mathbf{z} \in \mathbf{x}R_k \cap \mathbf{y}R_\ell$ . Moreover, the coefficient  $c_{g,h,i,j,k,\ell}$  only depends on the subscripts  $g, h, i, j, k, \ell$  and is independent of the choices of  $\mathbf{y}, \mathbf{z}$ . If  $c_{g,h,i,j,k,\ell} \in \mathbb{F}^\times$  for some  $\ell \in [0, d]$ , then  $R_\ell \in R_g R_k \cap R_h R_j$ .

*Proof.* The desired lemma follows from Lemma 3.6 and a direct computation.  $\square$

**Lemma 3.8.** *The  $\mathbb{F}$ -dimension of  $\mathbb{T}$  equals  $2^{n_2+2n_3} 3^{n_3} 5^{n_2} 11^{n_1+n_4}$ .*

*Proof.* Recall that  $n_1 = |\{a : a \in [1, n], \ell_a > m_a = 2\}|$ ,  $n_2 = |\{a : a \in [1, n], \ell_a = m_a = 2\}|$ ,  $n_3 = |\{a : a \in [1, n], \ell_a, m_a > 2\}|$ , and  $n_4 = |\{a : a \in [1, n], m_a > \ell_a = 2\}|$ . The desired lemma thus follows from combining Lemmas 3.6, 3.1, and a direct computation.  $\square$

For another  $\mathbb{F}$ -basis of  $\mathbb{T}$ , the following notations and five lemmas are necessary.

**Notation 3.9.** *Assume that  $g, h \in [0, d]$ . Then  $\mathbb{U}_{g,h}$  is defined to be the triple set  $\{(\mathfrak{a}(1), \mathfrak{a}(2), \mathfrak{a}(3)) : \mathfrak{a}(1) \subseteq (\mathfrak{g}_1 \cap \mathfrak{h}_1)^\circ, \mathfrak{a}(2) \subseteq (\mathfrak{g}_2 \cap \mathfrak{h}_2)^\bullet, \mathfrak{a}(3) \subseteq \mathfrak{g}_2 \cap \mathfrak{h}_2\}$ . Let  $\mathbb{U}_{g,h,i}$  be  $\{R_a : a \in \mathbb{P}_{g,h}, \mathfrak{a}_1 \cap (\mathfrak{g}_1 \cap \mathfrak{h}_1)^\circ \subseteq \mathfrak{i}(1), \mathfrak{a}_2 \cap (\mathfrak{g}_2 \cap \mathfrak{h}_2)^\bullet \subseteq \mathfrak{i}(2), \mathfrak{a}_1 \cap (\mathfrak{g}_2 \cap \mathfrak{h}_2) \subseteq \mathfrak{i}(3)\}$  for any  $\mathfrak{i} = (\mathfrak{i}(1), \mathfrak{i}(2), \mathfrak{i}(3)) \in \mathbb{U}_{g,h}$ . Set  $\mathfrak{o} = (\emptyset, \emptyset, \emptyset)$ . Notice that  $\mathfrak{o} \in \mathbb{U}_{g,h}$ . As there is  $j \in [0, d]$  such that  $\mathfrak{j}_1 = (\mathfrak{g}_0 \cap \mathfrak{h}_1) \cup (\mathfrak{g}_1 \cap \mathfrak{h}_0)$  and  $\mathfrak{j}_2 = \mathfrak{g}_2 \Delta \mathfrak{h}_2$ , notice that  $R_j \in \mathbb{U}_{g,h,i}$  for any  $\mathfrak{i} \in \mathbb{U}_{g,h}$ . Moreover,  $\mathbb{U}_{g,h} = \mathbb{U}_{h,g} = \mathbb{U}_{g,g} \cup \mathbb{U}_{h,h}$  and  $\mathbb{U}_{g,h,i} = \mathbb{U}_{h,g,i}$  for any  $\mathfrak{i} \in \mathbb{U}_{g,h}$ . As an example, assume that  $n = \ell_1 = 2, \ell_2 = m_1 = 3$ , and  $m_2 = 4$ . Hence  $d = 8$ . Notice that  $\mathbb{U}_{6,7} = \{(\emptyset, \emptyset, \emptyset), (\emptyset, \emptyset, \{2\}), (\emptyset, \{2\}, \{2\})\}$  and  $\mathbb{U}_{6,7,(\emptyset, \emptyset, \{2\})} = \{R_1, R_4\}$ .*

**Notation 3.10.** *Assume that  $g, h \in [0, d]$  and  $\mathfrak{i} \in \mathbb{U}_{g,h}$ . Set  $B_{g,h,i} = \sum_{R_j \in \mathbb{U}_{g,h,i}} E_g^* A_j E_h^*$ . As  $\mathbb{U}_{g,h,i} \neq \emptyset$  by Notation 3.9, notice that  $B_{g,h,i}$  is a defined nonzero matrix in  $\mathbb{T}$ .*

**Lemma 3.11.** *Assume that  $g, h \in [0, d]$  and  $\mathfrak{i} = (\mathfrak{i}(1), \mathfrak{i}(2), \mathfrak{i}(3)), \mathfrak{j} \in \mathbb{U}_{g,h}$ . Assume that  $k \in [0, d]$ ,  $\mathfrak{k}_1 = (\mathfrak{g}_0 \cap \mathfrak{h}_1) \cup (\mathfrak{g}_1 \cap \mathfrak{h}_0) \cup \mathfrak{i}(1) \cup (\mathfrak{i}(3) \setminus \mathfrak{i}(2))$ , and  $\mathfrak{k}_2 = (\mathfrak{g}_2 \Delta \mathfrak{h}_2) \cup \mathfrak{i}(2)$ . Then  $R_k \in \mathbb{U}_{g,h,i}$ . Moreover, if  $R_k \in \mathbb{U}_{g,h,i}$ , then  $\mathfrak{i} \preceq \mathfrak{j}$ .*

*Proof.* The first statement is obvious. Assume that  $\mathfrak{j} = (\mathfrak{j}(1), \mathfrak{j}(2), \mathfrak{j}(3))$ . If  $R_k \in \mathbb{U}_{g,h,i}$ , then  $\mathfrak{i}(1) \subseteq \mathfrak{j}(1), \mathfrak{i}(2) \subseteq \mathfrak{j}(2), \mathfrak{i}(3) = (\mathfrak{i}(3) \setminus \mathfrak{i}(2)) \cup \mathfrak{i}(2) \subseteq \mathfrak{j}(3)$ . The desired lemma follows.  $\square$

**Lemma 3.12.** *Assume that  $g \in [0, d]$ . Then  $B_{g,g,\mathfrak{o}} = E_g^*$ .*

*Proof.* Notation 3.4 implies that  $h \in \mathbb{P}_{g,g}$  if and only if  $\mathfrak{h}_1 \subseteq \mathfrak{g}_1^\circ \cup \mathfrak{g}_2$  and  $\mathfrak{h}_2 \subseteq \mathfrak{g}_2^\bullet$ . So  $\mathbb{U}_{g,g,\mathfrak{o}} = \{R_0\}$  by Lemma 3.1. The desired lemma follows from Equation (2.2).  $\square$

**Lemma 3.13.** *Assume that  $g, h \in [0, d]$  and  $\mathfrak{i} \in \mathbb{U}_{g,h}$ . Then  $B_{g,h,i}^T = B_{h,g,i}$ .*

*Proof.* As  $\mathbb{U}_{g,h,i} = \mathbb{U}_{h,g,i}$ , the desired lemma thus follows from Equation (2.1).  $\square$

**Lemma 3.14.** *Assume that  $g, h, i, j \in [0, d]$ ,  $\mathfrak{k} \in \mathbb{U}_{g,h}, \mathfrak{l} \in \mathbb{U}_{i,j}$ . Then  $B_{g,h,\mathfrak{k}} = B_{i,j,\mathfrak{l}}$  if and only if  $g = i, h = j$ , and  $\mathfrak{k} = \mathfrak{l}$ .*

*Proof.* One direction is obvious. For the other direction, assume that  $B_{g,h,\mathfrak{k}} = B_{i,j,\mathfrak{l}}$ . Notice that  $g = i$  and  $h = j$  by Equation (2.2). As  $B_{g,h,\mathfrak{k}} = B_{g,h,\mathfrak{l}}$ , Lemma 3.6 implies that  $\mathbb{U}_{g,h,\mathfrak{k}} = \mathbb{U}_{g,h,\mathfrak{l}}$ . The desired lemma thus follows from Lemma 3.11.  $\square$

**Lemma 3.15.**  $\mathbb{T}$  has an  $\mathbb{F}$ -linearly independent subset  $\{B_{a,b,c} : a, b \in [0, d], \mathbf{c} \in \mathbb{U}_{a,b}\}$ .

*Proof.* Let  $\mathbb{U} = \{B_{a,b,c} : a, b \in [0, d], \mathbf{c} \in \mathbb{U}_{a,b}\}$ . Let  $L$  be a nonzero  $\mathbb{F}$ -linear combination of the matrices in  $\mathbb{U}$ . Assume that  $L = O$ . If  $M \in \mathbb{U}$ , let  $c_M$  be the coefficient of  $M$  in  $L$ . Then there exists  $N \in \mathbb{U}$  such that  $c_N \in \mathbb{F}^\times$ . Equation (2.2) thus implies that  $N = E_g^* N E_h^*$  for some  $g, h \in [0, d]$ . Hence  $\mathbb{V} = \{A : A \in \mathbb{U}, c_A \in \mathbb{F}^\times, A = E_g^* A E_h^*\} \neq \emptyset$ . According to Lemma 3.14, there exist  $i \in \mathbb{N}$  and pairwise distinct  $j_1, j_2, \dots, j_i \in \mathbb{U}_{g,h}$  such that  $\mathbb{V} = \{B_{g,h,j_1}, B_{g,h,j_2}, \dots, B_{g,h,j_i}\}$ . Let us consider the cases  $i = 1$  and  $i > 1$ .

If  $i = 1$ , there is  $c \in \mathbb{F}^\times$  such that  $c B_{g,h,j_1} = E_g^* L E_h^* = O$ . This is a contradiction since  $B_{g,h,j_1} \neq O$ . Assume further that  $i > 1$ . Then there is no loss to assume that  $j_1$  is maximal in  $\{j_1, j_2, \dots, j_i\}$  with respect to the partial order  $\preceq$ . As  $E_g^* L E_h^* = O$ ,  $B_{g,h,j_1}$  is also an  $\mathbb{F}$ -linear combination of the matrices in  $\{B_{g,h,j_2}, B_{g,h,j_3}, \dots, B_{g,h,j_i}\}$ . The combination of Lemmas 3.1, 3.6, 3.11 thus implies that  $j_1 \preceq j \in \{j_2, j_3, \dots, j_i\}$ . Hence  $j_1 \in \{j_2, j_3, \dots, j_i\}$  by the choice of  $j_1$ . This is also a contradiction. The above contradictions thus imply that  $L \neq O$ . The desired lemma thus follows.  $\square$

We are now ready to present another  $\mathbb{F}$ -basis of  $\mathbb{T}$  and an additional notation.

**Theorem 3.16.** Assume that  $\mathbb{U} = \{(a, b, \mathbf{c}) : a, b \in [0, d], \mathbf{c} \in \mathbb{U}_{a,b}\}$ .  $\mathbb{T}$  has an  $\mathbb{F}$ -basis  $\{B_{a,b,c} : a, b \in [0, d], \mathbf{c} \in \mathbb{U}_{a,b}\}$  with cardinality  $|\mathbb{U}|$  and  $|\mathbb{U}| = 2^{n_2+2n_3} 3^{n_3} 5^{n_2} 11^{n_1+n_4}$ .

*Proof.* Assume that  $g, h \in [0, d]$ ,  $i = |(\mathfrak{g}_1 \cap \mathfrak{h}_1)^\circ|$ ,  $j = |(\mathfrak{g}_2 \cap \mathfrak{h}_2)^\bullet|$ , and  $k = |\mathfrak{g}_2 \cap \mathfrak{h}_2|$ . Notice that  $|\mathbb{P}_{g,h}| = |\mathbb{U}_{g,h}| = 2^{i-j+k} 3^j$  by Lemma 3.1 and a direct computation. As  $|\mathbb{P}_{g,h}| = |\mathbb{U}_{g,h}|$  and  $g, h$  are chosen from  $[0, d]$  arbitrarily, the following equality holds:

$$|\mathbb{P}| = \sum_{\ell=0}^d \sum_{m=0}^d |\mathbb{P}_{\ell,m}| = \sum_{\ell=0}^d \sum_{m=0}^d |\mathbb{U}_{\ell,m}| = |\mathbb{U}|.$$

Notice that  $|\{B_{a,b,c} : a, b \in [0, d], \mathbf{c} \in \mathbb{U}_{a,b}\}| = |\mathbb{U}|$  by Lemma 3.14. Then the desired theorem follows from combining Lemmas 3.6, 3.8, and 3.15.  $\square$

**Notation 3.17.** Assume that  $\mathbb{T}$  has an  $\mathbb{F}$ -basis  $\mathbb{A}$ . Let  $M \in \mathbb{T}$ . Then  $M$  is uniquely written as an  $\mathbb{F}$ -linear combination of the matrices in  $\mathbb{A}$ . If  $B \in \mathbb{A}$ , use  $c_{\mathbb{A}}(M, B)$  to denote the coefficient of  $B$  in this  $\mathbb{F}$ -linear combination that represents  $M$ . Define  $\text{Supp}_{\mathbb{A}}(M) = \{A : A \in \mathbb{A}, c_{\mathbb{A}}(M, A) \in \mathbb{F}^\times\}$ . So  $M = O$  if and only if  $\text{Supp}_{\mathbb{A}}(M) = \emptyset$ . Define  $\mathbb{B}_1 = \{E_a^* A_b E_c^* : (a, c, b) \in \mathbb{P}\}$  and  $\mathbb{B}_2 = \{B_{a,b,c} : a, b \in [0, d], \mathbf{c} \in \mathbb{U}_{a,b}\}$ . Hence both  $\text{Supp}_{\mathbb{B}_1}(M)$  and  $\text{Supp}_{\mathbb{B}_2}(M)$  can be defined by Lemma 3.6 and Theorem 3.16. Assume further that  $g, h \in [0, d]$  and  $i \in \mathbb{U}_{g,h}$ . Use  $c_{g,h,i}(M)$  to denote  $c_{\mathbb{B}_2}(M, B_{g,h,i})$ .

We next investigate the structure constants of  $\mathbb{B}_2$  in  $\mathbb{T}$ . We begin with a notation.

**Notation 3.18.** Assume that  $g, h, i \in [0, d]$ . Assume that  $\mathbf{j} = (j_{(1)}, j_{(2)}, j_{(3)}) \in \mathbb{U}_{g,h}$  and  $\mathfrak{k} = (k_{(1)}, k_{(2)}, k_{(3)}) \in \mathbb{U}_{h,i}$ . Let  $\mathbb{l}_{(1)} = ((\mathfrak{g}_1 \cap \mathfrak{i}_1)^\circ \setminus \mathfrak{h}_1) \cup (\mathfrak{g}_1 \cap \mathfrak{i}_1 \cap (j_{(1)} \cup k_{(1)}))$ ,  $\mathbb{l}_{(2)} = ((\mathfrak{g}_2 \cap \mathfrak{i}_2)^\bullet \setminus \mathfrak{h}_2) \cup (\mathfrak{g}_2 \cap \mathfrak{i}_2 \cap (j_{(2)} \cup k_{(2)}))$ ,  $\mathbb{l}_{(3)} = ((\mathfrak{g}_2 \cap \mathfrak{i}_2) \setminus \mathfrak{h}_2) \cup (\mathfrak{g}_2 \cap \mathfrak{i}_2 \cap (j_{(3)} \cup k_{(3)}))$ . Set  $(g, h, i, \mathbf{j}, \mathfrak{k}) = (\mathbb{l}_{(1)}, \mathbb{l}_{(2)}, \mathbb{l}_{(3)})$ . Notice that  $(g, h, i, \mathbf{j}, \mathfrak{k}) \in \mathbb{U}_{g,i}$  and  $B_{g,i,(g,h,i,\mathbf{j},\mathfrak{k})}$  is defined. If  $g = h = i$ , notice that  $(g, g, g, \mathbf{j}, \mathfrak{k}) = (j_{(1)} \cup k_{(1)}, j_{(2)} \cup k_{(2)}, j_{(3)} \cup k_{(3)})$  and denote  $(g, g, g, \mathbf{j}, \mathfrak{k})$  by  $\mathbf{j} \cup \mathfrak{k}$ . For example, assume that  $n = \ell_1 = 2$ ,  $\ell_2 = m_1 = 3$ , and  $m_2 = 4$ . Notice that  $d = 8$  and  $(6, 7, 6, (\emptyset, \emptyset, \{2\}), (\emptyset, \emptyset, \{2\})) = (\emptyset, \emptyset, \{2\}) \in \mathbb{U}_{6,6}$ .

**Lemma 3.19.** *Assume that  $g, h, i \in [0, d]$ ,  $j \in \mathbb{U}_{g,h}$ ,  $\mathfrak{k} \in \mathbb{U}_{h,i}$ ,  $R_\ell \in R_g R_i \cap \mathbb{U}_{g,h,j} \mathbb{U}_{h,i,\mathfrak{k}}$ . Then  $R_\ell \in \mathbb{U}_{g,i,(g,h,i,j,\mathfrak{k})}$ .*

*Proof.* As  $R_\ell \in R_g R_i$ , notice that  $\ell \in \mathbb{P}_{g,i}$  by Lemma 3.5. There are  $R_m \in \mathbb{U}_{g,h,j}$  and  $R_q \in \mathbb{U}_{h,i,\mathfrak{k}}$  such that  $R_\ell \in R_m R_q$ . Let  $j = (j_{(1)}, j_{(2)}, j_{(3)})$  and  $\mathfrak{k} = (k_{(1)}, k_{(2)}, k_{(3)})$ . As  $R_m \in \mathbb{U}_{g,h,j}$  and  $R_q \in \mathbb{U}_{h,i,\mathfrak{k}}$ , notice that  $(g_1 \cap h_1)^\circ \cap m_1 \subseteq j_{(1)}$ ,  $(g_2 \cap h_2)^\bullet \cap m_2 \subseteq j_{(2)}$ ,  $(g_2 \cap h_2) \cap m_1 \subseteq j_{(3)}$ ,  $(h_1 \cap i_1)^\circ \cap q_1 \subseteq k_{(1)}$ ,  $(h_2 \cap i_2)^\bullet \cap q_2 \subseteq k_{(2)}$ ,  $(h_2 \cap i_2) \cap q_1 \subseteq k_{(3)}$ ,  $l_1 \subseteq m_1 \cup q_1 \cup (m_2 \cap q_2)$ ,  $l_2 \subseteq m_2 \cup q_2$ ,  $m_2 \subseteq g_2 \cup h_2$ ,  $q_2 \subseteq h_2 \cup i_2$  by Notation 3.4.

Notice that  $(g_1 \cap i_1)^\circ \cap l_1 \subseteq ((g_1 \cap i_1)^\circ \setminus h_1) \cup ((g_1 \cap h_1 \cap i_1)^\circ \cap (m_1 \cup q_1 \cup (m_2 \cap q_2)))$ . Notice that  $(g_1 \cap i_1)^\circ \cap l_1 \subseteq ((g_1 \cap i_1)^\circ \setminus h_1) \cup (g_1 \cap i_1 \cap (j_{(1)} \cup k_{(1)}))$  by Lemma 3.1. Similarly, notice that  $(g_2 \cap i_2) \cap l_2 \subseteq ((g_2 \cap i_2) \setminus h_2) \cup (g_2 \cap i_2 \cap (j_{(3)} \cup k_{(3)}))$ . Notice that  $(g_2 \cap i_2)^\bullet \cap l_2 \subseteq ((g_2 \cap i_2)^\bullet \setminus h_2) \cup ((g_2 \cap h_2 \cap i_2)^\bullet \cap (m_2 \cup q_2))$ . Therefore  $(g_2 \cap i_2)^\bullet \cap l_2 \subseteq ((g_2 \cap i_2)^\bullet \setminus h_2) \cup (g_2 \cap i_2 \cap (j_{(2)} \cup k_{(2)}))$ . The desired lemma follows.  $\square$

**Lemma 3.20.** *Assume that  $g, h, i \in [0, d]$ ,  $j \in \mathbb{U}_{g,h}$ , and  $\mathfrak{k} \in \mathbb{U}_{h,i}$ . Then*

$$\text{Supp}_{\mathbb{B}_1}(B_{g,h,j} B_{h,i,\mathfrak{k}}) \subseteq \{E_g^* A_a E_i^* : R_a \in \mathbb{U}_{g,i,(g,h,i,j,\mathfrak{k})}\}.$$

*Proof.* The desired lemma is from combining Equation (2.2), Lemmas 3.7, 3.19.  $\square$

**Notation 3.21.** *Assume that  $g, h, i \in [0, d]$  and  $\mathbb{U}, \mathbb{V}, \mathbb{W} \subseteq [1, n]$ . Let  $k_{(\mathbb{U},\mathbb{V},\mathbb{W})}$  denote*

$$\prod_{j \in \mathbb{U}} (m_j - 1) \prod_{k \in \mathbb{V}} (\ell_k - 1) m_k \prod_{\ell \in \mathbb{W} \setminus \mathbb{V}} m_\ell,$$

where a product over the empty set equals one. Hence  $k_j$  is defined for any  $j \in \mathbb{U}_{g,h}$ . Assume that  $j = (j_{(1)}, j_{(2)}, j_{(3)}) \in \mathbb{U}_{g,h}$  and  $\mathfrak{k} = (k_{(1)}, k_{(2)}, k_{(3)}) \in \mathbb{U}_{h,i}$ . Let  $j \cap \mathfrak{k}$ ,  $j \setminus i$ ,  $i \cap j$  be  $(j_{(1)} \cap k_{(1)}, j_{(2)} \cap k_{(2)}, j_{(3)} \cap k_{(3)})$ ,  $(j_{(1)} \setminus i_1, j_{(2)} \setminus i_2, j_{(3)} \setminus i_3)$ ,  $(i_1 \cap j_{(1)}, i_2 \cap j_{(2)}, i_3 \cap j_{(3)})$ , respectively. So  $j \cap \mathfrak{k} \in \mathbb{U}_{g,h} \cap \mathbb{U}_{h,i}$ ,  $j \setminus i \in \mathbb{U}_{g,h}$ ,  $i \cap j \in \mathbb{U}_{i,i}$ ,  $k_{j \cap \mathfrak{k}}$ ,  $k_{j \setminus i}$ ,  $k_{i \cap j}$  are defined.

Set  $k_{\mathbb{U},\mathbb{V}} = k_{(\mathbb{U},\mathbb{V},\mathbb{V})}$ . If  $\mathbb{U} = (g_1 \cap i_1) \setminus h_1$  and  $\mathbb{V} = (g_2 \cap i_2) \setminus h_2$ , set  $k_{(g,h,i)} = k_{\mathbb{U},\mathbb{V}}$ . If  $\mathbb{U} = h_1 \setminus (g_1 \cup i_1)$  and  $\mathbb{V} = h_2 \setminus (g_2 \cup i_2)$ , set  $k_{[g,h,i]} = k_{\mathbb{U},\mathbb{V}}$ . For example, assume that  $n = \ell_1 = 2$ ,  $\ell_2 = m_1 = 3$ , and  $m_2 = 4$ . Then  $d = 8$ ,  $k_{(4,5,6)} = 1$ , and  $k_{[4,5,6]} = 3$ .

**Lemma 3.22.** *Assume that  $\mathbb{U}, \mathbb{V}, \mathbb{W} \subseteq [1, n]$ . Then  $k_{(\mathbb{U},\mathbb{V},\mathbb{W})} = k_{(\mathbb{U}^\circ, \mathbb{V}, \mathbb{W})}$ . Moreover, if  $\mathbb{V} \subseteq \mathbb{W}$ , then  $k_{(\mathbb{U},\mathbb{V},\mathbb{W})} = k_{(\mathbb{U}, \mathbb{V}^\bullet, \mathbb{W})}$ .*

*Proof.* The desired lemma follows from the hypotheses and a direct computation.  $\square$

**Lemma 3.23.** *Assume that  $\mathbb{U}, \mathbb{U}^1, \mathbb{V}, \mathbb{V}^1, \mathbb{W}, \mathbb{W}^1 \subseteq [1, n]$ . If  $\mathbb{U} \cap \mathbb{U}^1 = \emptyset$ ,  $\mathbb{V} \cap \mathbb{V}^1 = \emptyset$ ,  $\mathbb{V} \cap \mathbb{W}^1 = \emptyset$ ,  $\mathbb{V}^1 \cap \mathbb{W} = \emptyset$ ,  $\mathbb{W} \cap \mathbb{W}^1 = \emptyset$ , then  $k_{(\mathbb{U},\mathbb{V},\mathbb{W})} k_{(\mathbb{U}^1,\mathbb{V}^1,\mathbb{W}^1)} = k_{(\mathbb{U} \cup \mathbb{U}^1, \mathbb{V} \cup \mathbb{V}^1, \mathbb{W} \cup \mathbb{W}^1)}$ .*

*Proof.* The desired lemma follows from the hypotheses and a direct computation.  $\square$

**Lemma 3.24.** *Assume that  $g \in [0, d]$ . Then  $k_g = \prod_{h \in \mathfrak{g}_1} (m_h - 1) \prod_{i \in \mathfrak{g}_2} (\ell_i - 1) m_i$ .*

*Proof.* As  $k_g = |\mathbf{x}R_g| = k_{\mathfrak{g}_1, \mathfrak{g}_2}$ , the desired lemma is from a direct computation.  $\square$

**Lemma 3.25.** *Assume that  $\mathbb{U}, \mathbb{V}, \mathbb{W}$  are pairwise disjoint subsets of  $[1, n]$ . Assume that  $\mathbb{W}^1 = \{a : a \in [0, d], \mathbb{U} \subseteq \mathfrak{a}_1 \subseteq \mathbb{U} \cup \mathbb{W}, \mathfrak{a}_2 = \mathbb{V}\}$ . Then  $\sum_{g \in \mathbb{W}^1} k_g = k_{(\mathbb{U}, \mathbb{V}, \mathbb{W})}$ .*

*Proof.* The desired lemma is from combining Lemmas 3.1, 3.24, a computation.  $\square$

**Lemma 3.26.** *Assume that  $g, h, i \in [0, d]$ . Assume that  $\mathbf{j} = (\mathbf{j}_{(1)}, \mathbf{j}_{(2)}, \mathbf{j}_{(3)}) \in \mathbb{U}_{g,h}$  and  $\mathbf{k} = (\mathbf{k}_{(1)}, \mathbf{k}_{(2)}, \mathbf{k}_{(3)}) \in \mathbb{U}_{h,i}$ . Then  $[1, n]$  is a union of the pairwise disjoint subsets  $\mathfrak{h}_0$ ,  $\mathfrak{h}_1 \setminus (\mathfrak{g}_1 \cup \mathfrak{i}_1)$ ,  $\mathfrak{h}_2 \setminus (\mathfrak{g}_2 \cup \mathfrak{i}_2)$ ,  $\mathbf{j}_{(1)} \setminus \mathfrak{i}_1$ ,  $\mathbf{j}_{(2)} \setminus \mathfrak{i}_2$ ,  $\mathbf{k}_{(1)} \setminus \mathfrak{g}_1$ ,  $\mathbf{k}_{(2)} \setminus \mathfrak{g}_2$ ,  $\mathbf{j}_{(1)} \cap \mathbf{k}_{(1)}$ ,  $\mathbf{j}_{(2)} \cap \mathbf{k}_{(2)}$ ,  $(\mathfrak{g}_1 \cap \mathbf{k}_{(1)}) \setminus \mathbf{j}_{(1)}$ ,  $(\mathfrak{g}_1 \cap \mathfrak{h}_1) \setminus (\mathfrak{i}_1 \cup \mathbf{j}_{(1)})$ ,  $(\mathfrak{g}_2 \cap \mathbf{k}_{(3)}) \setminus \mathbf{j}_{(3)}$ ,  $(\mathfrak{g}_2 \cap \mathfrak{h}_2) \setminus (\mathfrak{i}_2 \cup \mathbf{j}_{(3)})$ ,  $\mathbf{j}_{(3)} \setminus (\mathfrak{i}_2 \cup \mathbf{j}_{(2)})$ ,  $(\mathbf{j}_{(3)} \cap \mathbf{k}_{(2)}) \setminus \mathbf{j}_{(2)}$ ,  $(\mathfrak{h}_1 \cap \mathfrak{i}_1) \setminus \mathbf{k}_{(1)}$ ,  $(\mathfrak{h}_2 \cap \mathfrak{i}_2) \setminus \mathbf{k}_{(3)}$ ,  $\mathbf{k}_{(3)} \setminus (\mathfrak{g}_2 \cup \mathbf{k}_{(2)})$ ,  $(\mathbf{j}_{(3)} \cap \mathbf{k}_{(3)}) \setminus \mathbf{k}_{(2)}$ .*

*Proof.* The desired lemma follows from Lemma 3.1 and a direct computation.  $\square$

**Notation 3.27.** *Assume that  $g, h, i \in [0, d]$ . Assume that  $\mathbf{j} = (\mathbf{j}_{(1)}, \mathbf{j}_{(2)}, \mathbf{j}_{(3)}) \in \mathbb{U}_{g,h}$  and  $\mathbf{k} = (\mathbf{k}_{(1)}, \mathbf{k}_{(2)}, \mathbf{k}_{(3)}) \in \mathbb{U}_{h,i}$ . Let  $\mathbb{U}_{g,h,i,j,\mathbf{k},1}$  be the union of  $\mathfrak{h}_0$ ,  $\mathfrak{h}_1 \setminus (\mathfrak{g}_1 \cup \mathfrak{i}_1)$ ,  $\mathfrak{h}_2 \setminus (\mathfrak{g}_2 \cup \mathfrak{i}_2)$ ,  $\mathbf{j}_{(1)} \setminus \mathfrak{i}_1$ ,  $\mathbf{j}_{(2)} \setminus \mathfrak{i}_2$ ,  $\mathbf{k}_{(1)} \setminus \mathfrak{g}_1$ ,  $\mathbf{k}_{(2)} \setminus \mathfrak{g}_2$ ,  $\mathbf{j}_{(1)} \cap \mathbf{k}_{(1)}$ ,  $\mathbf{j}_{(2)} \cap \mathbf{k}_{(2)}$ . Let  $\mathbb{U}_{g,h,i,j,\mathbf{k},2}$  be the union of  $(\mathfrak{g}_1 \cap \mathbf{k}_{(1)}) \setminus \mathbf{j}_{(1)}$ ,  $(\mathfrak{g}_1 \cap \mathfrak{h}_1) \setminus (\mathfrak{i}_1 \cup \mathbf{j}_{(1)})$ ,  $(\mathfrak{g}_2 \cap \mathbf{k}_{(3)}) \setminus \mathbf{j}_{(3)}$ ,  $(\mathfrak{g}_2 \cap \mathfrak{h}_2) \setminus (\mathfrak{i}_2 \cup \mathbf{j}_{(3)})$ ,  $\mathbf{j}_{(3)} \setminus (\mathfrak{i}_2 \cup \mathbf{j}_{(2)})$ ,  $(\mathbf{j}_{(3)} \cap \mathbf{k}_{(2)}) \setminus \mathbf{j}_{(2)}$ . Set  $\mathbb{U}_{g,h,i,j,\mathbf{k},3} = [1, n] \setminus (\mathbb{U}_{g,h,i,j,\mathbf{k},1} \cup \mathbb{U}_{g,h,i,j,\mathbf{k},2})$ . Let  $\mathbf{y}, \mathbf{z} \in \mathbb{X}$ . Then Lemmas 3.26 and 3.1 give a unique  $\mathbf{w} \in \mathbb{X}$ , where  $\mathbf{w}_\ell = \mathbf{x}_\ell$ ,  $\mathbf{w}_m = \mathbf{y}_m$ ,  $\mathbf{w}_q = \mathbf{z}_q$  for any  $\ell \in \mathbb{U}_{g,h,i,j,\mathbf{k},1}$ ,  $m \in \mathbb{U}_{g,h,i,j,\mathbf{k},2}$ , and  $q \in \mathbb{U}_{g,h,i,j,\mathbf{k},3}$ . Define  $(\mathbf{y}, \mathbf{z})_{g,h,i,j,\mathbf{k}} = \mathbf{w}$ .*

**Lemma 3.28.** *Assume that  $\mathbf{y}, \mathbf{z} \in \mathbb{X}$ ,  $g, h, i \in [0, d]$ ,  $\mathbf{j} = (\mathbf{j}_{(1)}, \mathbf{j}_{(2)}, \mathbf{j}_{(3)}) \in \mathbb{U}_{g,h}$ , and  $\mathbf{k} = (\mathbf{k}_{(1)}, \mathbf{k}_{(2)}, \mathbf{k}_{(3)}) \in \mathbb{U}_{h,i}$ . If  $R_\ell \in \mathbb{U}_{g,h,j}$ ,  $R_m \in \mathbb{U}_{h,i,\mathbf{k}}$ ,  $\mathbf{w} \in \mathbf{y}R_\ell \cap \mathbf{x}R_h \cap \mathbf{z}R_m$ , then there are  $q \in [0, d]$  and  $\mathbb{U} \subseteq (\mathbf{j}_{(3)} \setminus (\mathfrak{i}_2 \cup \mathbf{j}_{(2)})) \cup (\mathbf{k}_{(3)} \setminus (\mathfrak{g}_2 \cup \mathbf{k}_{(2)})) \cup ((\mathbf{j}_{(3)} \cap \mathbf{k}_{(3)}) \setminus (\mathbf{j}_{(2)} \cap \mathbf{k}_{(2)}))$  such that  $\mathbf{w} \in (\mathbf{y}, \mathbf{z})_{g,h,i,j,\mathbf{k}}R_q$ ,  $\mathfrak{q}_1 = (\mathfrak{h}_1 \setminus (\mathfrak{g}_1 \cup \mathfrak{i}_1)) \cup (\mathbf{j}_{(1)} \setminus \mathfrak{i}_1) \cup (\mathbf{k}_{(1)} \setminus \mathfrak{g}_1) \cup (\mathbf{j}_{(1)} \cap \mathbf{k}_{(1)}) \cup \mathbb{U}$ , and  $\mathfrak{q}_2 = (\mathfrak{h}_2 \setminus (\mathfrak{g}_2 \cup \mathfrak{i}_2)) \cup (\mathbf{j}_{(2)} \setminus \mathfrak{i}_2) \cup (\mathbf{k}_{(2)} \setminus \mathfrak{g}_2) \cup (\mathbf{j}_{(2)} \cap \mathbf{k}_{(2)})$ .*

*Proof.* The union of  $(\mathbf{j}_{(3)} \cap \mathfrak{l}_1) \setminus (\mathfrak{i}_2 \cup \mathbf{j}_{(2)})$ ,  $(\mathbf{j}_{(3)} \cap \mathbf{k}_{(2)} \cap \mathfrak{l}_1) \setminus \mathbf{j}_{(2)}$ ,  $(\mathbf{k}_{(3)} \cap \mathfrak{m}_1) \setminus (\mathfrak{g}_2 \cup \mathbf{k}_{(2)})$ , and  $(\mathbf{j}_{(3)} \cap \mathbf{k}_{(3)} \cap \mathfrak{m}_1) \setminus \mathbf{k}_{(2)}$  is denoted by  $\mathbb{U}$ . By Lemmas 3.1 and 3.26, there exists  $q \in [0, d]$  such that  $\mathfrak{q}_1 = (\mathfrak{h}_1 \setminus (\mathfrak{g}_1 \cup \mathfrak{i}_1)) \cup (\mathbf{j}_{(1)} \setminus \mathfrak{i}_1) \cup (\mathbf{k}_{(1)} \setminus \mathfrak{g}_1) \cup (\mathbf{j}_{(1)} \cap \mathbf{k}_{(1)}) \cup \mathbb{U}$  and  $\mathfrak{q}_2 = (\mathfrak{h}_2 \setminus (\mathfrak{g}_2 \cup \mathfrak{i}_2)) \cup (\mathbf{j}_{(2)} \setminus \mathfrak{i}_2) \cup (\mathbf{k}_{(2)} \setminus \mathfrak{g}_2) \cup (\mathbf{j}_{(2)} \cap \mathbf{k}_{(2)})$ . Write  $\mathbf{v} = (\mathbf{y}, \mathbf{z})_{g,h,i,j,\mathbf{k}}$ .

Notice that  $\mathbf{w}_r \sim_{\mathbb{G}_r} \mathbf{x}_r = \mathbf{v}_r$  and  $\mathbf{w}_s \not\sim_{\mathbb{G}_s} \mathbf{x}_s = \mathbf{v}_s$  if  $r \in \mathfrak{q}_1 \setminus \mathbb{U}$  and  $s \in \mathfrak{q}_2$ . Notice that  $\mathbf{w}_r \sim_{\mathbb{G}_r} \mathbf{y}_r = \mathbf{v}_r$  if  $r \in ((\mathbf{j}_{(3)} \cap \mathfrak{l}_1) \setminus (\mathfrak{i}_2 \cup \mathbf{j}_{(2)})) \cup ((\mathbf{j}_{(3)} \cap \mathbf{k}_{(2)} \cap \mathfrak{l}_1) \setminus \mathbf{j}_{(2)})$ . Notice that  $\mathbf{w}_r \sim_{\mathbb{G}_r} \mathbf{z}_r = \mathbf{v}_r$  if  $r \in ((\mathbf{k}_{(3)} \cap \mathfrak{m}_1) \setminus (\mathfrak{g}_2 \cup \mathbf{k}_{(2)})) \cup ((\mathbf{j}_{(3)} \cap \mathbf{k}_{(3)} \cap \mathfrak{m}_1) \setminus \mathbf{k}_{(2)})$ . If  $r \in \mathfrak{h}_0$ , then  $\mathbf{w}_r = \mathbf{x}_r = \mathbf{v}_r$ . If  $r$  is in the union of  $(\mathfrak{g}_1 \cap \mathbf{k}_{(1)}) \setminus \mathbf{j}_{(1)}$ ,  $(\mathfrak{g}_1 \cap \mathfrak{h}_1) \setminus (\mathfrak{i}_1 \cup \mathbf{j}_{(1)})$ ,  $(\mathfrak{g}_2 \cap \mathbf{k}_{(3)}) \setminus \mathbf{j}_{(3)}$ ,  $(\mathfrak{g}_2 \cap \mathfrak{h}_2) \setminus (\mathfrak{i}_2 \cup \mathbf{j}_{(3)})$ ,  $(\mathbf{j}_{(3)} \cap \mathfrak{l}_0) \setminus (\mathfrak{i}_2 \cup \mathbf{j}_{(2)})$ , and  $(\mathbf{j}_{(3)} \cap \mathbf{k}_{(2)} \cap \mathfrak{l}_0) \setminus \mathbf{j}_{(2)}$ , then  $\mathbf{w}_r = \mathbf{y}_r = \mathbf{v}_r$  as  $R_\ell \in \mathbb{U}_{g,h,j}$ . If  $r$  is in the union of  $(\mathfrak{h}_1 \cap \mathfrak{i}_1) \setminus \mathbf{k}_{(1)}$ ,  $(\mathfrak{h}_2 \cap \mathfrak{i}_2) \setminus \mathbf{k}_{(3)}$ ,  $(\mathbf{k}_{(3)} \cap \mathfrak{m}_0) \setminus (\mathfrak{g}_2 \cup \mathbf{k}_{(2)})$ , and  $(\mathbf{j}_{(3)} \cap \mathbf{k}_{(3)} \cap \mathfrak{m}_0) \setminus \mathbf{k}_{(2)}$ , then  $\mathbf{w}_r = \mathbf{z}_r = \mathbf{v}_r$  as  $R_m \in \mathbb{U}_{h,i,\mathbf{k}}$ . The desired lemma thus follows as  $R_\ell \in \mathbb{U}_{g,h,j}$ ,  $R_m \in \mathbb{U}_{h,i,\mathbf{k}}$ , and Lemma 3.26 holds.  $\square$

**Lemma 3.29.** *Assume that  $\mathbf{y}, \mathbf{z} \in \mathbb{X}$ ,  $g, h, i \in [0, d]$ ,  $\mathbf{j} = (\mathbf{j}_{(1)}, \mathbf{j}_{(2)}, \mathbf{j}_{(3)}) \in \mathbb{U}_{g,h}$ , and  $\mathbf{k} = (\mathbf{k}_{(1)}, \mathbf{k}_{(2)}, \mathbf{k}_{(3)}) \in \mathbb{U}_{h,i}$ . Assume that  $\mathbf{y} \in \mathbf{x}R_g$  and  $\mathbf{z} \in \mathbf{x}R_i$ . If there are  $\ell \in [0, d]$  and  $\mathbb{U} \subseteq (\mathbf{j}_{(3)} \setminus (\mathfrak{i}_2 \cup \mathbf{j}_{(2)})) \cup (\mathbf{k}_{(3)} \setminus (\mathfrak{g}_2 \cup \mathbf{k}_{(2)})) \cup ((\mathbf{j}_{(3)} \cap \mathbf{k}_{(3)}) \setminus (\mathbf{j}_{(2)} \cap \mathbf{k}_{(2)}))$  such that  $\mathbf{w} \in (\mathbf{y}, \mathbf{z})_{g,h,i,j,\mathbf{k}}R_\ell$ ,  $\mathfrak{l}_1 = (\mathfrak{h}_1 \setminus (\mathfrak{g}_1 \cup \mathfrak{i}_1)) \cup (\mathbf{j}_{(1)} \setminus \mathfrak{i}_1) \cup (\mathbf{k}_{(1)} \setminus \mathfrak{g}_1) \cup (\mathbf{j}_{(1)} \cap \mathbf{k}_{(1)}) \cup \mathbb{U}$ , and  $\mathfrak{l}_2 = (\mathfrak{h}_2 \setminus (\mathfrak{g}_2 \cup \mathfrak{i}_2)) \cup (\mathbf{j}_{(2)} \setminus \mathfrak{i}_2) \cup (\mathbf{k}_{(2)} \setminus \mathfrak{g}_2) \cup (\mathbf{j}_{(2)} \cap \mathbf{k}_{(2)})$ , then  $\mathbf{w} \in \mathbf{x}R_h$ .*

*Proof.* Set  $\mathbf{v} = (\mathbf{y}, \mathbf{z})_{g,h,i,j,\mathbf{k}}$ . Then  $\mathbf{w}_m \sim_{\mathbb{G}_m} \mathbf{v}_m = \mathbf{x}_m$  and  $\mathbf{w}_q \not\sim_{\mathbb{G}_q} \mathbf{v}_q = \mathbf{x}_q$  if  $m \in \mathfrak{l}_1 \setminus \mathbb{U}$  and  $q \in \mathfrak{l}_2$ . If  $m \in (\mathbf{j}_{(3)} \setminus (\mathfrak{i}_2 \cup \mathbf{j}_{(2)})) \cup ((\mathbf{j}_{(3)} \cap \mathbf{k}_{(2)}) \setminus \mathbf{j}_{(2)})$ , then  $\mathbf{w}_m = \mathbf{v}_m = \mathbf{y}_m \not\sim_{\mathbb{G}_m} \mathbf{x}_m$  or  $\mathbf{w}_m \sim_{\mathbb{G}_m} \mathbf{v}_m = \mathbf{y}_m \not\sim_{\mathbb{G}_m} \mathbf{x}_m$ . Then  $\mathbf{w}_m = \mathbf{v}_m = \mathbf{z}_m \not\sim_{\mathbb{G}_m} \mathbf{x}_m$  or  $\mathbf{w}_m \sim_{\mathbb{G}_m} \mathbf{v}_m = \mathbf{z}_m \not\sim_{\mathbb{G}_m} \mathbf{x}_m$  if  $m \in (\mathbf{k}_{(3)} \setminus (\mathfrak{g}_2 \cup \mathbf{k}_{(2)})) \cup ((\mathbf{j}_{(3)} \cap \mathbf{k}_{(3)}) \setminus \mathbf{k}_{(2)})$ . If  $m \in \mathfrak{h}_0$ , notice that  $\mathbf{w}_m = \mathbf{v}_m = \mathbf{x}_m$ . If  $m \in ((\mathfrak{g}_1 \cap \mathbf{k}_{(1)}) \setminus \mathbf{j}_{(1)}) \cup ((\mathfrak{g}_1 \cap \mathfrak{h}_1) \setminus (\mathfrak{i}_1 \cup \mathbf{j}_{(1)})) \cup ((\mathfrak{h}_1 \cap \mathfrak{i}_1) \setminus \mathbf{k}_{(1)})$ , then  $\mathbf{w}_m = \mathbf{v}_m = \mathbf{y}_m \sim_{\mathbb{G}_m} \mathbf{x}_m$  or  $\mathbf{w}_m = \mathbf{v}_m = \mathbf{z}_m \sim_{\mathbb{G}_m} \mathbf{x}_m$ . If  $m \in ((\mathfrak{g}_2 \cap \mathbf{k}_{(3)}) \setminus \mathbf{j}_{(3)}) \cup ((\mathfrak{g}_2 \cap \mathfrak{h}_2) \setminus (\mathfrak{i}_2 \cup \mathbf{j}_{(3)})) \cup ((\mathfrak{h}_2 \cap \mathfrak{i}_2) \setminus \mathbf{k}_{(3)})$ ,  $\mathbf{w}_m = \mathbf{v}_m = \mathbf{y}_m \not\sim_{\mathbb{G}_m} \mathbf{x}_m$  or  $\mathbf{w}_m = \mathbf{v}_m = \mathbf{z}_m \not\sim_{\mathbb{G}_m} \mathbf{x}_m$ . The desired lemma is from Lemma 3.26.  $\square$

**Lemma 3.30.** *Assume that  $\mathbf{y}, \mathbf{z} \in \mathbb{X}$ ,  $g, h, i \in [0, d]$ ,  $\mathbf{j} = (\mathbf{j}_{(1)}, \mathbf{j}_{(2)}, \mathbf{j}_{(3)}) \in \mathbb{U}_{g,h}$ , and  $\mathbf{k} = (\mathbf{k}_{(1)}, \mathbf{k}_{(2)}, \mathbf{k}_{(3)}) \in \mathbb{U}_{h,i}$ . Assume that  $R_\ell \in \mathbb{U}_{g,i,(g,h,i,j,\mathbf{k})}$  and  $\mathbf{y} \in \mathbf{x}R_g \cap \mathbf{z}R_\ell$ . If there are  $m \in [0, d]$  and  $\mathbb{U} \subseteq (\mathbf{j}_{(3)} \setminus (\mathbf{i}_2 \cup \mathbf{j}_{(2)})) \cup (\mathbf{k}_{(3)} \setminus (\mathbf{g}_2 \cup \mathbf{k}_{(2)})) \cup ((\mathbf{j}_{(3)} \cap \mathbf{k}_{(3)}) \setminus (\mathbf{j}_{(2)} \cap \mathbf{k}_{(2)}))$  such that  $\mathbf{w} \in (\mathbf{y}, \mathbf{z})_{g,h,i,j,\mathbf{k}} R_m$ ,  $\mathbf{m}_1 = (\mathbf{h}_1 \setminus (\mathbf{g}_1 \cup \mathbf{i}_1)) \cup (\mathbf{j}_{(1)} \setminus \mathbf{i}_1) \cup (\mathbf{k}_{(1)} \setminus \mathbf{g}_1) \cup (\mathbf{j}_{(1)} \cap \mathbf{k}_{(1)}) \cup \mathbb{U}$ ,  $\mathbf{m}_2 = (\mathbf{h}_2 \setminus (\mathbf{g}_2 \cup \mathbf{i}_2)) \cup (\mathbf{j}_{(2)} \setminus \mathbf{i}_2) \cup (\mathbf{k}_{(2)} \setminus \mathbf{g}_2) \cup (\mathbf{j}_{(2)} \cap \mathbf{k}_{(2)})$ , then  $\mathbf{w} \in \mathbf{y}R_q$  for some  $R_q \in \mathbb{U}_{g,h,i}$ .*

*Proof.* Set  $\mathbf{r} = \{a : a \in [1, n], \mathbf{w}_a \sim_{\mathbb{G}_a} \mathbf{y}_a\}$  and  $\mathbf{s} = \{a : a \in [1, n], \mathbf{w}_a \not\sim_{\mathbb{G}_a} \mathbf{y}_a\}$ . By Lemma 3.1, there is  $q \in [0, d]$  such that  $\mathbf{q}_1 = (\mathbf{g}_0 \cap \mathbf{h}_1) \cup (\mathbf{g}_1 \cap \mathbf{h}_0) \cup (\mathbf{j}_{(1)} \cap \mathbf{r}) \cup (\mathbf{j}_{(3)} \cap \mathbf{r})$  and  $\mathbf{q}_2 = (\mathbf{g}_2 \triangle \mathbf{h}_2) \cup (\mathbf{j}_{(2)} \cap \mathbf{s})$ . Notice that  $R_q \in \mathbb{U}_{g,h,j}$  by Lemma 3.5. Set  $\mathbf{v} = (\mathbf{y}, \mathbf{z})_{g,h,i,j,\mathbf{k}}$ .

If  $t \in \mathbf{g}_0 \cap \mathbf{h}_0$ , then  $\mathbf{w}_t = \mathbf{v}_t = \mathbf{x}_t = \mathbf{y}_t$ . If  $t \in \mathbf{h}_0 \setminus \mathbf{g}_0$ , then  $\mathbf{w}_t = \mathbf{v}_t = \mathbf{x}_t \sim_{\mathbb{G}_t} \mathbf{y}_t$  or  $\mathbf{w}_t = \mathbf{v}_t = \mathbf{x}_t \not\sim_{\mathbb{G}_t} \mathbf{y}_t$ . If  $t \in (\mathbf{h}_1 \setminus (\mathbf{g}_1 \cup \mathbf{i}_1)) \cup (\mathbf{k}_{(1)} \setminus \mathbf{g}_1)$ , then  $\mathbf{w}_t \sim_{\mathbb{G}_t} \mathbf{v}_t = \mathbf{x}_t = \mathbf{y}_t$  or  $\mathbf{w}_t \sim_{\mathbb{G}_t} \mathbf{v}_t = \mathbf{x}_t \not\sim_{\mathbb{G}_t} \mathbf{y}_t$ . If  $t \in (\mathbf{h}_2 \setminus (\mathbf{g}_2 \cup \mathbf{i}_2)) \cup (\mathbf{k}_{(2)} \setminus \mathbf{g}_2)$ , then  $\mathbf{w}_t \not\sim_{\mathbb{G}_t} \mathbf{v}_t = \mathbf{x}_t = \mathbf{y}_t$  or  $\mathbf{w}_t \not\sim_{\mathbb{G}_t} \mathbf{v}_t = \mathbf{x}_t \sim_{\mathbb{G}_t} \mathbf{y}_t$ . If  $t \in (\mathbf{j}_{(1)} \setminus \mathbf{i}_1) \cup (\mathbf{j}_{(1)} \cap \mathbf{k}_{(1)}) \cup (\mathbf{j}_{(3)} \setminus (\mathbf{i}_2 \cup \mathbf{j}_{(2)})) \cup ((\mathbf{j}_{(3)} \cap \mathbf{k}_{(3)}) \setminus \mathbf{j}_{(2)})$ , then  $\mathbf{w}_t = \mathbf{y}_t$  or  $\mathbf{w}_t \sim_{\mathbb{G}_t} \mathbf{y}_t$  as  $R_\ell \in \mathbb{U}_{g,i,(g,h,i,j,\mathbf{k})}$ . If  $t \in \mathbf{k}_{(3)} \setminus (\mathbf{g}_2 \cup \mathbf{k}_{(2)})$ , then  $R_\ell \in \mathbb{U}_{g,i,(g,h,i,j,\mathbf{k})}$  gives  $\mathbf{w}_t = \mathbf{v}_t = \mathbf{z}_t \not\sim_{\mathbb{G}_t} \mathbf{y}_t$  or  $\mathbf{w}_t \sim_{\mathbb{G}_t} \mathbf{v}_t = \mathbf{z}_t \not\sim_{\mathbb{G}_t} \mathbf{y}_t$ . If  $t$  is in the union of  $(\mathbf{g}_1 \cap \mathbf{k}_{(1)}) \setminus \mathbf{j}_{(1)}$ ,  $(\mathbf{g}_1 \cap \mathbf{h}_1) \setminus (\mathbf{i}_1 \cup \mathbf{j}_{(1)})$ ,  $(\mathbf{g}_2 \cap \mathbf{k}_{(3)}) \setminus \mathbf{j}_{(3)}$ , and  $(\mathbf{g}_2 \cap \mathbf{h}_2) \setminus (\mathbf{i}_2 \cup \mathbf{j}_{(3)})$ , then  $\mathbf{w}_t = \mathbf{v}_t = \mathbf{y}_t$ . If  $t \in ((\mathbf{g}_2 \cap \mathbf{h}_1 \cap \mathbf{i}_1) \setminus \mathbf{k}_{(1)}) \cup ((\mathbf{h}_2 \cap \mathbf{i}_2) \setminus (\mathbf{j}_{(3)} \cup \mathbf{k}_{(3)}))$ , then  $R_\ell \in \mathbb{U}_{g,i,(g,h,i,j,\mathbf{k})}$  gives  $\mathbf{w}_t = \mathbf{v}_t = \mathbf{z}_t = \mathbf{y}_t$  or  $\mathbf{w}_t = \mathbf{v}_t = \mathbf{z}_t \not\sim_{\mathbb{G}_t} \mathbf{y}_t$ . If  $t \in ((\mathbf{h}_1 \cap \mathbf{i}_1) \setminus (\mathbf{g}_2 \cup \mathbf{k}_{(1)})) \cup ((\mathbf{i}_2 \cap \mathbf{j}_{(3)}) \setminus (\mathbf{j}_{(2)} \cup \mathbf{k}_{(3)}))$ ,  $R_\ell \in \mathbb{U}_{g,i,(g,h,i,j,\mathbf{k})}$  gives  $\mathbf{w}_t = \mathbf{v}_t = \mathbf{z}_t = \mathbf{y}_t$  or  $\mathbf{w}_t = \mathbf{v}_t = \mathbf{z}_t \sim_{\mathbb{G}_t} \mathbf{y}_t$ . The desired lemma is from Lemma 3.26.  $\square$

**Lemma 3.31.** *Assume that  $\mathbf{y}, \mathbf{z} \in \mathbb{X}$ ,  $g, h, i \in [0, d]$ ,  $\mathbf{j} = (\mathbf{j}_{(1)}, \mathbf{j}_{(2)}, \mathbf{j}_{(3)}) \in \mathbb{U}_{g,h}$ , and  $\mathbf{k} = (\mathbf{k}_{(1)}, \mathbf{k}_{(2)}, \mathbf{k}_{(3)}) \in \mathbb{U}_{h,i}$ . Assume that  $R_\ell \in \mathbb{U}_{g,i,(g,h,i,j,\mathbf{k})}$  and  $\mathbf{z} \in \mathbf{x}R_i \cap \mathbf{y}R_\ell$ . If there are  $m \in [0, d]$  and  $\mathbb{U} \subseteq (\mathbf{j}_{(3)} \setminus (\mathbf{i}_2 \cup \mathbf{j}_{(2)})) \cup (\mathbf{k}_{(3)} \setminus (\mathbf{g}_2 \cup \mathbf{k}_{(2)})) \cup ((\mathbf{j}_{(3)} \cap \mathbf{k}_{(3)}) \setminus (\mathbf{j}_{(2)} \cap \mathbf{k}_{(2)}))$  such that  $\mathbf{w} \in (\mathbf{y}, \mathbf{z})_{g,h,i,j,\mathbf{k}} R_m$ ,  $\mathbf{m}_1 = (\mathbf{h}_1 \setminus (\mathbf{g}_1 \cup \mathbf{i}_1)) \cup (\mathbf{j}_{(1)} \setminus \mathbf{i}_1) \cup (\mathbf{k}_{(1)} \setminus \mathbf{g}_1) \cup (\mathbf{j}_{(1)} \cap \mathbf{k}_{(1)}) \cup \mathbb{U}$ ,  $\mathbf{m}_2 = (\mathbf{h}_2 \setminus (\mathbf{g}_2 \cup \mathbf{i}_2)) \cup (\mathbf{j}_{(2)} \setminus \mathbf{i}_2) \cup (\mathbf{k}_{(2)} \setminus \mathbf{g}_2) \cup (\mathbf{j}_{(2)} \cap \mathbf{k}_{(2)})$ , then  $\mathbf{w} \in \mathbf{z}R_q$  for some  $R_q \in \mathbb{U}_{h,i,\mathbf{k}}$ .*

*Proof.* Set  $\mathbf{r} = \{a : a \in [1, n], \mathbf{w}_a \sim_{\mathbb{G}_a} \mathbf{z}_a\}$  and  $\mathbf{s} = \{a : a \in [1, n], \mathbf{w}_a \not\sim_{\mathbb{G}_a} \mathbf{z}_a\}$ . By Lemma 3.1, there is  $q \in [0, d]$  such that  $\mathbf{q}_1 = (\mathbf{h}_0 \cap \mathbf{i}_1) \cup (\mathbf{h}_1 \cap \mathbf{i}_0) \cup (\mathbf{k}_{(1)} \cap \mathbf{r}) \cup (\mathbf{k}_{(3)} \cap \mathbf{r})$  and  $\mathbf{q}_2 = (\mathbf{h}_2 \triangle \mathbf{i}_2) \cup (\mathbf{k}_{(2)} \cap \mathbf{s})$ . Notice that  $R_q \in \mathbb{U}_{h,i,\mathbf{k}}$  by Lemma 3.5. Set  $\mathbf{v} = (\mathbf{y}, \mathbf{z})_{g,h,i,j,\mathbf{k}}$ .

If  $t \in \mathbf{h}_0 \cap \mathbf{i}_0$ , then  $\mathbf{w}_t = \mathbf{v}_t = \mathbf{x}_t = \mathbf{z}_t$ . If  $t \in \mathbf{h}_0 \setminus \mathbf{i}_0$ , then  $\mathbf{w}_t = \mathbf{v}_t = \mathbf{x}_t \sim_{\mathbb{G}_t} \mathbf{z}_t$  or  $\mathbf{w}_t = \mathbf{v}_t = \mathbf{x}_t \not\sim_{\mathbb{G}_t} \mathbf{z}_t$ . If  $t \in (\mathbf{h}_1 \setminus (\mathbf{g}_1 \cup \mathbf{i}_1)) \cup (\mathbf{j}_{(1)} \setminus \mathbf{i}_1)$ , then  $\mathbf{w}_t \sim_{\mathbb{G}_t} \mathbf{v}_t = \mathbf{x}_t = \mathbf{z}_t$  or  $\mathbf{w}_t \sim_{\mathbb{G}_t} \mathbf{v}_t = \mathbf{x}_t \not\sim_{\mathbb{G}_t} \mathbf{z}_t$ . If  $t \in (\mathbf{h}_2 \setminus (\mathbf{g}_2 \cup \mathbf{i}_2)) \cup (\mathbf{j}_{(2)} \setminus \mathbf{i}_2)$ , then  $\mathbf{w}_t \not\sim_{\mathbb{G}_t} \mathbf{v}_t = \mathbf{x}_t = \mathbf{z}_t$  or  $\mathbf{w}_t \not\sim_{\mathbb{G}_t} \mathbf{v}_t = \mathbf{x}_t \sim_{\mathbb{G}_t} \mathbf{z}_t$ . If  $t \in (\mathbf{k}_{(1)} \setminus \mathbf{g}_1) \cup (\mathbf{j}_{(1)} \cap \mathbf{k}_{(1)}) \cup (\mathbf{k}_{(3)} \setminus (\mathbf{g}_2 \cup \mathbf{k}_{(2)})) \cup ((\mathbf{j}_{(3)} \cap \mathbf{k}_{(3)}) \setminus \mathbf{k}_{(2)})$ , then  $\mathbf{w}_t = \mathbf{z}_t$  or  $\mathbf{w}_t \sim_{\mathbb{G}_t} \mathbf{z}_t$ . If  $t \in \mathbf{j}_{(3)} \setminus (\mathbf{i}_2 \cup \mathbf{j}_{(2)})$ ,  $R_\ell \in \mathbb{U}_{g,i,(g,h,i,j,\mathbf{k})}$  gives  $\mathbf{w}_t = \mathbf{v}_t = \mathbf{y}_t \not\sim_{\mathbb{G}_t} \mathbf{z}_t$  or  $\mathbf{w}_t \sim_{\mathbb{G}_t} \mathbf{v}_t = \mathbf{y}_t \not\sim_{\mathbb{G}_t} \mathbf{z}_t$ . If  $t$  is in the union of  $(\mathbf{g}_1 \cap \mathbf{k}_{(1)}) \setminus \mathbf{j}_{(1)}$ ,  $(\mathbf{g}_2 \cap \mathbf{k}_{(3)}) \setminus (\mathbf{j}_{(3)} \cup \mathbf{k}_{(2)})$ ,  $(\mathbf{h}_1 \cap \mathbf{i}_1) \setminus \mathbf{k}_{(1)}$ , and  $(\mathbf{h}_2 \cap \mathbf{i}_2) \setminus \mathbf{k}_{(3)}$ , then  $R_\ell \in \mathbb{U}_{g,i,(g,h,i,j,\mathbf{k})}$  gives  $\mathbf{w}_t = \mathbf{z}_t$  or  $\mathbf{w}_t \sim_{\mathbb{G}_t} \mathbf{z}_t$ . If  $t \in ((\mathbf{g}_1 \cap \mathbf{h}_1) \setminus (\mathbf{i}_1 \cup \mathbf{j}_{(1)})) \cup ((\mathbf{g}_2 \cap \mathbf{h}_2) \setminus (\mathbf{i}_2 \cup \mathbf{j}_{(3)}))$ , then  $\mathbf{w}_t = \mathbf{v}_t = \mathbf{y}_t \sim_{\mathbb{G}_t} \mathbf{z}_t$  or  $\mathbf{w}_t = \mathbf{v}_t = \mathbf{y}_t \not\sim_{\mathbb{G}_t} \mathbf{z}_t$  as  $R_\ell \in \mathbb{U}_{g,i,(g,h,i,j,\mathbf{k})}$ . The desired lemma is from Lemma 3.26.  $\square$

We are now ready to conclude this section by the structure constants of  $\mathbb{B}_2$  in  $\mathbb{T}$ .

**Theorem 3.32.** *Assume that  $g, h, i \in [0, d]$ ,  $\mathbf{j} \in \mathbb{U}_{g,h}$ ,  $\mathbf{k} \in \mathbb{U}_{\ell,i}$  for some  $\ell \in [0, d]$ . Then  $B_{g,h,j} B_{\ell,i,\mathbf{k}} = \delta_{h,\ell} \overline{k_{[g,h,i]} k_{\mathbf{j}} k_{\mathbf{k}} k_{\mathbf{j}} \cap \mathbf{k}} B_{g,i,(g,h,i,j,\mathbf{k})}$ .*

*Proof.* Assume that  $h = \ell$  by Equation (2.2). If  $\mathbf{y}, \mathbf{z} \in \mathbb{X}$ ,  $R_m \in \mathbb{U}_{g,i,(g,h,i,j,\mathbf{k})}$ ,  $\mathbf{z} \in \mathbf{y}R_m$ , notice that  $\sum_{R_q \in \mathbb{U}_{g,h,j}} \sum_{R_r \in \mathbb{U}_{h,i,\mathbf{k}}} |\mathbf{y}R_q \cap \mathbf{x}R_h \cap \mathbf{z}R_r| = k_{[g,h,i]} k_{\mathbf{j}} k_{\mathbf{k}} k_{\mathbf{j}} \cap \mathbf{k}$  by combining Lemmas 3.28, 3.29, 3.30, 3.31, 3.26, 3.25, 3.23. As  $R_m$  can be chosen from  $\mathbb{U}_{g,i,(g,h,i,j,\mathbf{k})}$  arbitrarily, the desired theorem thus follows from Lemmas 3.20 and 3.7.  $\square$

4. ALGEBRAIC STRUCTURE OF  $\mathbb{T}$ : CENTER

In this section, we present an  $\mathbb{F}$ -basis of  $Z(\mathbb{T})$  and list a property of this  $\mathbb{F}$ -basis. We first recall Notations 3.9, 3.10, 3.17, 3.18, 3.21 and give an additional notation.

**Notation 4.1.** Assume that  $g \in [0, d]$  and  $\mathfrak{h} \in \mathbb{U}_{g,g}$ . Define  $C_{\mathfrak{h}} = \sum_{i=0}^d \overline{k_{\mathfrak{h} \setminus i}} B_{i, i \cap \mathfrak{h}}$ . As  $k_{\mathfrak{h} \setminus g} = 1$ , Theorem 3.16 thus implies that  $C_{\mathfrak{h}}$  is a defined nonzero matrix in  $\mathbb{T}$ . As  $k_{\mathfrak{o} \setminus i} = 1$  for any  $i \in [0, d]$ , notice that  $C_{\mathfrak{o}} = I$  by Lemma 3.12 and Equation (2.3).

The following three lemmas give some properties of the objects in Notation 4.1.

**Lemma 4.2.** Assume that  $g, h \in [0, d]$ ,  $\mathfrak{i} \in \mathbb{U}_{g,g}$ , and  $\mathfrak{j} \in \mathbb{U}_{h,h}$ . Then  $c_{g,g,\mathfrak{i}}(C_{\mathfrak{i}}) = \bar{1}$ . In particular,  $C_{\mathfrak{i}} = C_{\mathfrak{j}}$  if and only if  $\mathfrak{i} = \mathfrak{j}$ .

*Proof.* As  $k_{\mathfrak{i} \setminus g} = 1$ , the first statement follows from Theorem 3.16. One direction in the second statement is obvious. For the other direction, assume that  $C_{\mathfrak{i}} = C_{\mathfrak{j}}$ . The first statement and Theorem 3.16 imply that  $B_{g,g,\mathfrak{i}} \in \text{Supp}_{\mathbb{B}_2}(C_{\mathfrak{i}}) = \text{Supp}_{\mathbb{B}_2}(C_{\mathfrak{j}})$ . So Theorem 3.16 gives  $\mathfrak{i} = g \cap \mathfrak{j} \preceq \mathfrak{j}$ . Similarly,  $\mathfrak{j} \preceq \mathfrak{i}$ . The desired lemma follows.  $\square$

**Lemma 4.3.**  $\mathbb{T}$  has an  $\mathbb{F}$ -linearly independent subset  $\{C_{\mathfrak{a}} : \exists b \in [0, d], \mathfrak{a} \in \mathbb{U}_{b,b}\}$ .

*Proof.* Let  $\mathbb{U} = \{C_{\mathfrak{a}} : \exists b \in [0, d], \mathfrak{a} \in \mathbb{U}_{b,b}\}$ . Let  $L$  be a nonzero  $\mathbb{F}$ -linear combination of the matrices in  $\mathbb{U}$ . Assume that  $L = O$ . If  $C_{\mathfrak{g}} \in \mathbb{U}$ , let  $c_{\mathfrak{g}}$  be the coefficient of  $C_{\mathfrak{g}}$  in  $L$ . There is  $C_{\mathfrak{h}} \in \mathbb{U}$  such that  $c_{\mathfrak{h}} \in \mathbb{F}^{\times}$ . Therefore  $\mathbb{V} = \{C_{\mathfrak{a}} : C_{\mathfrak{a}} \in \mathbb{U}, c_{\mathfrak{a}} \in \mathbb{F}^{\times}\} \neq \emptyset$ . According to Lemma 4.2, there must exist  $g \in \mathbb{N}$  and pairwise distinct  $\mathfrak{h}_1, \mathfrak{h}_2, \dots, \mathfrak{h}_g$  such that  $\mathbb{V} = \{C_{\mathfrak{h}_1}, C_{\mathfrak{h}_2}, \dots, C_{\mathfrak{h}_g}\}$ . Let us distinguish the cases  $g = 1$  and  $g > 1$ .

If  $g = 1$ , then  $O = L = c_{\mathfrak{h}_1} C_{\mathfrak{h}_1} \neq O$ , which is a contradiction. Assume that  $g > 1$ . There is no loss to assume further that  $\mathfrak{h}_1$  is maximal in  $\{\mathfrak{h}_1, \mathfrak{h}_2, \dots, \mathfrak{h}_g\}$  with respect to the partial order  $\preceq$ . As  $L = O$ ,  $C_{\mathfrak{h}_1}$  is an  $\mathbb{F}$ -linear combination of the matrices in  $\{C_{\mathfrak{h}_2}, C_{\mathfrak{h}_3}, \dots, C_{\mathfrak{h}_g}\}$ . The combination of Lemma 4.2, Theorem 3.16, and Notation 4.1 thus implies that  $\mathfrak{h}_1 = i \cap \mathfrak{h}_j \preceq \mathfrak{h}_j$  for some  $i \in [0, d]$  and  $j \in [2, g]$ . Then  $\mathfrak{h}_1 = \mathfrak{h}_j$  by the choice of  $\mathfrak{h}_1$ . This is absurd. Hence  $L \neq O$ . The desired lemma follows.  $\square$

**Lemma 4.4.** The  $\mathbb{F}$ -dimension of  $\langle \{C_{\mathfrak{a}} : \exists b \in [0, d], \mathfrak{a} \in \mathbb{U}_{b,b}\} \rangle$  equals  $2^{n_2+2n_3} 3^{n_1+n_4}$ .

*Proof.* Recall that  $n_1 = |\{a : a \in [1, n], \ell_a > m_a = 2\}|$ ,  $n_2 = |\{a : a \in [1, n], \ell_a = m_a = 2\}|$ ,  $n_3 = |\{a : a \in [1, n], \ell_a, m_a > 2\}|$ , and  $n_4 = |\{a : a \in [1, n], m_a > \ell_a = 2\}|$ . Notice that  $|\{C_{\mathfrak{a}} : \exists b \in [0, d], \mathfrak{a} \in \mathbb{U}_{b,b}\}| = |\{\mathfrak{a} : \exists b \in [0, d], \mathfrak{a} \in \mathbb{U}_{b,b}\}| = 2^{n_2+2n_3} 3^{n_1+n_4}$  by combining Lemmas 4.2, 3.1, a direct computation. The desired lemma is from Lemma 4.3.  $\square$

Our main goal is to find an  $\mathbb{F}$ -basis for  $Z(\mathbb{T})$ . We begin with a sequence of lemmas.

**Lemma 4.5.** Assume that  $g, h, i \in [0, d]$  and  $\mathfrak{j} \in \mathbb{U}_{g,h}$ . Then  $k_{\mathfrak{j}} = k_{\mathfrak{j} \setminus i} k_{i \cap \mathfrak{j}}$ .

*Proof.* The desired lemma follows from Lemma 3.23 and a direct computation.  $\square$

**Lemma 4.6.** Assume that  $g, h \in [0, d]$  and  $\mathfrak{i} \in \mathbb{U}_{g,h}$ . Then  $B_{g,h,\mathfrak{o}} B_{h,h,\mathfrak{i}} = B_{g,h,\mathfrak{i}}$ .

*Proof.* As  $k_{[g,h,h]} = k_{\mathfrak{o} \setminus h} = k_{\mathfrak{i} \setminus g} = k_{\mathfrak{o} \cap \mathfrak{i}} = 1$ , the desired lemma is from Theorem 3.32.  $\square$

**Lemma 4.7.** Assume that  $g, h \in [0, d]$  and  $\mathfrak{i} \in \mathbb{U}_{g,h}$ . Then  $B_{g,g,\mathfrak{i}} B_{g,h,\mathfrak{o}} = B_{g,h,\mathfrak{i}}$ .

*Proof.* The desired lemma is a direct application of Lemmas 4.6 and 3.13.  $\square$

**Lemma 4.8.** *Assume that  $g \in [0, d]$  and  $\mathfrak{h}, \mathfrak{i} \in \mathbb{U}_{g,g}$ . Then  $B_{g,g,\mathfrak{h}}B_{g,g,\mathfrak{i}} = \overline{k_{\mathfrak{h} \cap \mathfrak{i}}}B_{g,g,\mathfrak{h} \cup \mathfrak{i}}$  and the commutative  $\mathbb{F}$ -subalgebra  $E_g^* \mathbb{T} E_g^*$  of  $\mathbb{T}$  has an  $\mathbb{F}$ -basis  $\{B_{g,g,\mathfrak{a}} : \mathfrak{a} \in \mathbb{U}_{g,g}\}$ .*

*Proof.* As  $k_{[g,g,g]} = k_{\mathfrak{h} \setminus g} = k_{\mathfrak{i} \setminus g} = 1$ , the first statement is thus from Theorem 3.32. The desired lemma is from combining Lemma 2.5, Theorem 3.16, and Equation (2.2).  $\square$

**Lemma 4.9.** *Assume that  $g, h, i \in [0, d]$  and  $\mathfrak{j} \in \mathbb{U}_{i,i}$ . Then  $k_{\mathfrak{j} \setminus g} k_{(g \cap \mathfrak{j}) \setminus h} = k_{\mathfrak{j} \setminus h} k_{(h \cap \mathfrak{j}) \setminus g}$  and  $(g, g, h, g \cap \mathfrak{j}, \mathfrak{o}) = (g, h, h, \mathfrak{o}, h \cap \mathfrak{j})$ .*

*Proof.* Notice that  $k_{\mathfrak{j} \setminus g} k_{(g \cap \mathfrak{j}) \setminus h} k_{h \cap (g \cap \mathfrak{j})} = k_{\mathfrak{j}} = k_{\mathfrak{j} \setminus h} k_{(h \cap \mathfrak{j}) \setminus g} k_{g \cap (h \cap \mathfrak{j})}$  by Lemma 4.5. The first statement follows as  $k_{h \cap (g \cap \mathfrak{j})} = k_{g \cap (h \cap \mathfrak{j})}$  by a direct computation. If  $\mathfrak{j} = (\mathfrak{j}_{(1)}, \mathfrak{j}_{(2)}, \mathfrak{j}_{(3)})$ , then  $(g, g, h, g \cap \mathfrak{j}, \mathfrak{o}) = (g, h, h, \mathfrak{o}, h \cap \mathfrak{j}) = (g_1 \cap h_1 \cap \mathfrak{j}_{(1)}, g_2 \cap h_2 \cap \mathfrak{j}_{(2)}, g_3 \cap h_3 \cap \mathfrak{j}_{(3)})$  by a direct computation. The desired lemma follows from the above discussion.  $\square$

**Lemma 4.10.**  *$Z(\mathbb{T})$  has an  $\mathbb{F}$ -linearly independent subset  $\{C_{\mathfrak{a}} : \exists b \in [0, d], \mathfrak{a} \in \mathbb{U}_{b,b}\}$ .*

*Proof.* Pick  $g, h, i \in [0, d]$  and  $\mathfrak{j} \in \mathbb{U}_{i,i}$ . So  $B_{g,h,\mathfrak{o}}C_{\mathfrak{j}} = C_{\mathfrak{j}}B_{g,h,\mathfrak{o}}$  by combining Equation (2.2), Lemma 4.9, and Theorem 3.32. Assume that  $\mathfrak{k} \in \mathbb{U}_{g,h}$ . Then  $B_{g,h,\mathfrak{k}}C_{\mathfrak{j}} = C_{\mathfrak{j}}B_{g,h,\mathfrak{k}}$  by combining Equation (2.2), Lemmas 4.6, 4.7, and 4.8. Since  $g, h, i, \mathfrak{j}, \mathfrak{k}$  are chosen arbitrarily, the desired lemma thus follows from Theorem 3.16 and Lemma 4.3.  $\square$

**Lemma 4.11.** *Assume that  $M \in Z(\mathbb{T})$ . Then  $\text{Supp}_{\mathbb{B}_2}(M) \subseteq \{B_{a,a,b} : a \in [0, d], \mathfrak{b} \in \mathbb{U}_{a,a}\}$ .*

*Proof.* As  $M \in Z(\mathbb{T})$ , Theorem 3.16 implies that  $M$  is an  $\mathbb{F}$ -linear combination of the matrices in  $\mathbb{B}_2$ . Assume that  $\text{Supp}_{\mathbb{B}_2}(M) \not\subseteq \{B_{a,a,b} : a \in [0, d], \mathfrak{b} \in \mathbb{U}_{a,a}\}$ . Following Equations (2.3) and (2.2), there exist distinct  $g, h \in [0, d]$  such that  $E_g^* M E_h^* \neq O$ . As  $M \in Z(\mathbb{T})$  and  $E_g^*, E_h^* \in \mathbb{T}$ , notice that  $O \neq E_g^* M E_h^* = M E_g^* E_h^* = O$  by Equation (2.2). This is a contradiction. The desired lemma is thus from this contradiction.  $\square$

**Lemma 4.12.** *Assume that  $g, h \in [0, d]$  and  $M \in Z(\mathbb{T})$ . Assume that  $\mathfrak{i} \in \mathbb{U}_{g,g} \cap \mathbb{U}_{h,h}$  and  $\mathfrak{i}$  is maximal in  $\{\mathfrak{a} : \exists b \in [0, d], \mathfrak{a} \in \mathbb{U}_{b,b}, B_{b,b,\mathfrak{a}} \in \text{Supp}_{\mathbb{B}_2}(M)\}$  with respect to the partial order  $\preceq$ . Then  $c_{g,g,\mathfrak{i}}(M) = c_{h,h,\mathfrak{i}}(M)$ .*

*Proof.* As  $M \in Z(\mathbb{T})$ , Lemma 4.11 implies that  $M$  is an  $\mathbb{F}$ -linear combination of the matrices in  $\{B_{a,a,b} : a \in [0, d], \mathfrak{b} \in \mathbb{U}_{a,a}\}$ . According to Equation (2.2), observe that

$$(4.1) \quad \begin{aligned} M E_g^* &= c_{g,g,\mathfrak{i}}(M) B_{g,g,\mathfrak{i}} + \sum_{\mathfrak{j} \in \mathbb{U}_{g,g} \setminus \{\mathfrak{i}\}} c_{g,g,\mathfrak{j}}(M) B_{g,g,\mathfrak{j}} \text{ and} \\ E_h^* M &= c_{h,h,\mathfrak{i}}(M) B_{h,h,\mathfrak{i}} + \sum_{\mathfrak{j} \in \mathbb{U}_{h,h} \setminus \{\mathfrak{i}\}} c_{h,h,\mathfrak{j}}(M) B_{h,h,\mathfrak{j}}. \end{aligned}$$

By combining Equations (2.2), (4.1), Lemmas 4.6, 4.7, Theorem 3.32, notice that

$$(4.2) \quad \begin{aligned} M B_{g,h,\mathfrak{o}} &= M E_g^* B_{g,h,\mathfrak{o}} = c_{g,g,\mathfrak{i}}(M) B_{g,h,\mathfrak{i}} + \sum_{\mathfrak{j} \in \mathbb{U}_{g,g} \setminus \{\mathfrak{i}\}} c_{g,g,\mathfrak{j}}(M) \overline{k_{\mathfrak{j} \setminus h}} B_{g,h,h \cap \mathfrak{j}} \text{ and} \\ B_{g,h,\mathfrak{o}} M &= B_{g,h,\mathfrak{o}} E_h^* M = c_{h,h,\mathfrak{i}}(M) B_{g,h,\mathfrak{i}} + \sum_{\mathfrak{j} \in \mathbb{U}_{h,h} \setminus \{\mathfrak{i}\}} c_{h,h,\mathfrak{j}}(M) \overline{k_{\mathfrak{j} \setminus g}} B_{g,h,g \cap \mathfrak{j}}. \end{aligned}$$

If  $c_{g,g,\mathfrak{j}}(M) \in \mathbb{F}^\times$  and  $\mathfrak{i} = h \cap \mathfrak{j} \preceq \mathfrak{j}$  for some  $\mathfrak{j} \in \mathbb{U}_{g,g} \setminus \{\mathfrak{i}\}$ , then  $\mathfrak{i} = \mathfrak{j}$  by the choice of  $\mathfrak{i}$ . This is absurd. So  $c_{g,h,\mathfrak{i}}(M B_{g,h,\mathfrak{o}}) = c_{g,g,\mathfrak{i}}(M)$  by Equation (4.2) and Theorem 3.16. Similarly,  $c_{g,h,\mathfrak{i}}(B_{g,h,\mathfrak{o}} M) = c_{h,h,\mathfrak{i}}(M)$ . The desired lemma follows as  $M \in Z(\mathbb{T})$ .  $\square$

For further discussion, the following notation and an additional lemma are needed.

**Notation 4.13.** Assume that  $\mathbb{U}, \mathbb{V}, \mathbb{W}$  are sets. Define  $|(\mathbb{U}, \mathbb{V}, \mathbb{W})| = |\mathbb{U}| + |\mathbb{V}| + |\mathbb{W}|$ .

**Lemma 4.14.** Assume that  $M \in Z(\mathbb{T})$ . Then  $M \in \langle \{C_{\mathbf{a}} : \exists b \in [0, d], \mathbf{a} \in \mathbb{U}_{b,b}\} \rangle$ .

*Proof.* If  $L \in Z(\mathbb{T})$ , then  $\text{Supp}_{\mathbb{B}_2}(L) \subseteq \{B_{a,a,b} : a \in [0, d], \mathbf{b} \in \mathbb{U}_{a,a}\}$  by Lemma 4.11. The maximum number in  $\{|\mathbf{a}| : \exists b \in [0, d], \mathbf{a} \in \mathbb{U}_{b,b}, B_{b,b,\mathbf{a}} \in \text{Supp}_{\mathbb{B}_2}(L)\}$  is denoted by  $\max(L)$  if  $L \in Z(\mathbb{T}) \setminus \{O\}$ . Assume that  $M \neq O$ . Work by induction on  $\max(M)$ .

If  $\max(M) = 0$ , notice that  $M \in \langle \{I\} \rangle$  by combining the fact  $M \in Z(\mathbb{T})$ , Lemma 3.12, and Equation (2.3). Hence  $M \in \langle \{C_o\} \rangle$  as  $C_o = I$ . The base case is checked.

Assume that  $\max(M) > 0$  and  $L \in \langle \{C_{\mathbf{a}} : \exists b \in [0, d], \mathbf{a} \in \mathbb{U}_{b,b}\} \rangle$  for any  $L \in Z(\mathbb{T})$  and  $\max(L) < \max(M)$ . Assume that  $g \in \mathbb{N}$  and  $\mathfrak{h}_1, \mathfrak{h}_2, \dots, \mathfrak{h}_g$  are exactly all pairwise distinct elements in  $\{\mathbf{a} : \exists b \in [0, d], \mathbf{a} \in \mathbb{U}_{b,b}, |\mathbf{a}| = \max(M), B_{b,b,\mathbf{a}} \in \text{Supp}_{\mathbb{B}_2}(M)\}$ . So  $\mathfrak{h}_1, \mathfrak{h}_2, \dots, \mathfrak{h}_g$  are maximal in  $\{\mathbf{a} : \exists b \in [0, d], \mathbf{a} \in \mathbb{U}_{b,b}, B_{b,b,\mathbf{a}} \in \text{Supp}_{\mathbb{B}_2}(M)\}$  with respect to the partial order  $\preceq$ . Assume that  $B_{i_1, i_1, \mathfrak{h}_1}, B_{i_2, i_2, \mathfrak{h}_2}, \dots, B_{i_g, i_g, \mathfrak{h}_g} \in \text{Supp}_{\mathbb{B}_2}(M)$ . Set

$$(4.3) \quad N = M - \sum_{j=1}^g c_{i_j, i_j, \mathfrak{h}_j}(M) C_{\mathfrak{h}_j}.$$

There is no loss to assume that  $N \neq O$ . As  $M \in Z(\mathbb{T})$ , notice that  $N \in Z(\mathbb{T})$  by Lemma 4.10 and Equation (4.3). Notice that  $\max(N) \leq \max(M)$  by Notation 4.1 and Equation (4.3). Assume that  $\max(N) = \max(M)$ . There is  $B_{\ell, \ell, \mathfrak{k}} \in \text{Supp}_{\mathbb{B}_2}(N)$  such that  $|\mathfrak{k}| = \max(M)$ . Then the combination of Lemmas 4.2, 4.12, Equation (4.3), Theorem 3.16 implies that  $\mathfrak{k} \notin \{\mathfrak{h}_1, \mathfrak{h}_2, \dots, \mathfrak{h}_g\}$ . By the choice of  $\mathfrak{k}$  and Theorem 3.16, notice that  $B_{\ell, \ell, \mathfrak{k}} \in \text{Supp}_{\mathbb{B}_2}(M)$ . It contradicts the choice of  $g$ . So  $\max(N) < \max(M)$ . The desired lemma is thus from Equation (4.3) and the inductive hypothesis.  $\square$

We are now ready to present an  $\mathbb{F}$ -basis of  $Z(\mathbb{T})$  and a property of this  $\mathbb{F}$ -basis.

**Theorem 4.15.**  $Z(\mathbb{T})$  has an  $\mathbb{F}$ -basis  $\{C_{\mathbf{a}} : \exists b \in [0, d], \mathbf{a} \in \mathbb{U}_{b,b}\}$  whose cardinality is  $2^{n_2+2n_3}3^{n_1+n_4}$ . In particular, the  $\mathbb{F}$ -dimension of  $Z(\mathbb{T})$  is independent of the choices of the fixed field  $\mathbb{F}$  and  $\mathbf{x}$ .

*Proof.* The desired theorem follows from combining Lemmas 4.10, 4.14, and 4.4.  $\square$

**Lemma 4.16.** Assume that  $g \in [0, d]$  and  $\mathfrak{h} \in \mathbb{U}_{g,g}$ . Then  $C_{\mathfrak{h}}C_{\mathfrak{h}} = \overline{k_{\mathfrak{h}}}C_{\mathfrak{h}}$ . In particular, the  $\mathbb{F}$ -subalgebra  $Z(\mathbb{T})$  of  $\mathbb{T}$  is a semisimple  $\mathbb{F}$ -algebra only if  $p \nmid k_i$  for any  $i \in [0, d]$  and  $\mathfrak{j} \in \mathbb{U}_{i,i}$ .

*Proof.* For any  $k \in [0, d]$ , notice that  $k_{\mathfrak{h} \setminus k} k_{k \cap \mathfrak{h}} = k_{\mathfrak{h}}$  by Lemma 4.5. The desired lemma is thus from combining Equation (2.2), Lemma 4.8, and the above computation.  $\square$

We end this section by presenting an example of Theorem 4.15 and Lemma 4.16.

**Example 4.17.** Assume that  $n = \ell_1 = 2$ ,  $\ell_2 = m_1 = 3$ ,  $m_2 = 4$ . Then  $n_1 = n_2 = 0$  and  $n_3 = n_4 = 1$ . The  $\mathbb{F}$ -dimension of  $Z(\mathbb{T})$  is twelve by Theorem 4.15. By Theorem 4.15 and a direct computation,  $Z(\mathbb{T})$  has an  $\mathbb{F}$ -basis containing precisely  $I, C_{(\emptyset, \emptyset, \{1\})}, C_{(\emptyset, \emptyset, \{2\})}, C_{(\emptyset, \emptyset, \{1,2\})}, C_{(\emptyset, \{2\}, \{2\})}, C_{(\emptyset, \{2\}, \{1,2\})}, C_{(\{1\}, \emptyset, \emptyset)}, C_{(\{1\}, \emptyset, \{2\})}, C_{(\{1\}, \{2\}, \{2\})}, C_{(\{2\}, \emptyset, \emptyset)}, C_{(\{2\}, \emptyset, \{1\})}, C_{(\{1,2\}, \emptyset, \emptyset)}$ . By Lemma 4.16,  $C_{(\emptyset, \emptyset, \{1\})}C_{(\emptyset, \emptyset, \{1\})} = \overline{3}C_{(\emptyset, \emptyset, \{1\})}$ .

5. ALGEBRAIC STRUCTURE OF  $\mathbb{T}$ : SEMISIMPLICITY

In this section, we determine  $\text{Rad}(E_g^* \mathbb{T} E_g^*)$  for any  $g \in [0, d]$ . As an application, we also determine the semisimplicity of  $\mathbb{T}$ . For our purpose, we recall Notations 3.9, 3.10, 3.17, 3.18, 3.21, 4.13. We begin our discussion with two preliminary lemmas.

**Lemma 5.1.** *Assume that  $\mathbb{U}, \mathbb{V}, \mathbb{W} \subseteq [1, n]$ . Then  $p \mid k_{(\mathbb{U}, \mathbb{V}, \mathbb{W})}$  if and only if one of the following statements holds: There exists  $g \in \mathbb{U}$  such that  $p \mid m_g - 1$ ; There exists  $g \in \mathbb{V}$  such that  $p \mid (\ell_g - 1)m_g$ ; There exists  $g \in \mathbb{W} \setminus \mathbb{V}$  such that  $p \mid m_g$ .*

*Proof.* The desired lemma follows from the hypotheses and a direct computation.  $\square$

**Lemma 5.2.** *Assume that  $g \in [0, d]$ . The  $\mathbb{F}$ -subalgebra  $E_g^* \mathbb{T} E_g^*$  of  $\mathbb{T}$  has a two-sided ideal  $\langle \{B_{g,g,a} : a \in \mathbb{U}_{g,g}, p \mid k_a\} \rangle$ .*

*Proof.* Assume that  $B_{g,g,h}, B_{g,g,i} \in E_g^* \mathbb{T} E_g^*$  and  $p \mid k_i$ . Assume that  $\mathfrak{h} = (\mathfrak{h}_{(1)}, \mathfrak{h}_{(2)}, \mathfrak{h}_{(3)})$  and  $\mathfrak{i} = (\mathfrak{i}_{(1)}, \mathfrak{i}_{(2)}, \mathfrak{i}_{(3)})$ . If  $j \in \mathfrak{i}_{(1)}$  and  $p \mid m_j - 1$ , then  $j \in \mathfrak{h}_{(1)} \cup \mathfrak{i}_{(1)}$  and  $p \mid m_j - 1$ . If  $j \in \mathfrak{i}_{(2)}$  and  $p \mid (\ell_j - 1)m_j$ , then  $j \in \mathfrak{h}_{(2)} \cup \mathfrak{i}_{(2)}$  and  $p \mid (\ell_j - 1)m_j$ . If  $j \in (\mathfrak{h}_{(3)} \cap \mathfrak{i}_{(3)}) \setminus \mathfrak{i}_{(2)}$  and  $p \mid m_j$ , then  $j \in (\mathfrak{h}_{(3)} \cap \mathfrak{i}_{(3)}) \setminus (\mathfrak{h}_{(2)} \cap \mathfrak{i}_{(2)})$  and  $p \mid m_j$ . If  $j \in \mathfrak{i}_{(3)} \setminus (\mathfrak{h}_{(3)} \cup \mathfrak{i}_{(2)})$  and  $p \mid m_j$ , then  $j \in (\mathfrak{h}_{(3)} \cup \mathfrak{i}_{(3)}) \setminus (\mathfrak{h}_{(2)} \cup \mathfrak{i}_{(2)})$  and  $p \mid m_j$ . So  $p \mid k_{\mathfrak{h} \cap \mathfrak{i} k_{\mathfrak{h} \cup \mathfrak{i}}}$  by Lemma 5.1. As  $B_{g,g,h}, B_{g,g,i}$  are chosen arbitrarily, the desired lemma is from Lemma 4.8.  $\square$

Lemmas 5.1 and 5.2 motivate us to give the following notation and two lemmas.

**Notation 5.3.** *Assume that  $g, h \in [0, d]$  and  $\mathfrak{i} = (\mathfrak{i}_{(1)}, \mathfrak{i}_{(2)}, \mathfrak{i}_{(3)}) \in \mathbb{U}_{g,h}$ . Write  $\mathbb{U}_i$  for  $\{a : a \in \mathfrak{i}_{(1)}, p \mid m_a - 1\} \cup \{a : a \in \mathfrak{i}_{(2)}, p \mid (\ell_a - 1)m_a\} \cup \{a : a \in \mathfrak{i}_{(3)} \setminus \mathfrak{i}_{(2)}, p \mid m_a\}$ . So Lemma 5.1 shows that  $p \mid k_i$  if and only if  $\mathbb{U}_i \neq \emptyset$ . Set  $\mathbb{I}_g = \langle \{B_{g,g,a} : a \in \mathbb{U}_{g,g}, p \mid k_a\} \rangle$ . Lemma 5.2 thus implies that  $\mathbb{I}_g$  is a two-sided ideal of the  $\mathbb{F}$ -subalgebra  $E_g^* \mathbb{T} E_g^*$  of  $\mathbb{T}$ .*

**Lemma 5.4.** *Assume that  $g \in [0, d]$  and  $\mathbb{I}_g \neq \{O\}$ . Then there is a nonzero product of  $|\{a : a \in \mathfrak{g}_1, p \mid m_a - 1\}| + |\{a : a \in \mathfrak{g}_2, p \mid (\ell_a - 1)m_a\}|$  matrices in  $\mathbb{I}_g$ .*

*Proof.* Set  $\mathbb{U} = \{a : a \in \mathfrak{g}_1, p \mid m_a - 1\} \cup \{a : a \in \mathfrak{g}_2, p \mid (\ell_a - 1)m_a\}$ . As  $\mathbb{I}_g \neq \{O\}$ ,  $\mathbb{U} \neq \emptyset$ . Let  $h \in \mathbb{N}$  and  $\mathbb{U} = \{i_1, i_2, \dots, i_h\}$ . If  $j \in \mathfrak{g}_1$  and  $p \mid m_j - 1$ , set  $\mathfrak{k}_j = (\{j\}, \emptyset, \emptyset)$ . If  $j \in \mathfrak{g}_2^\bullet$  and  $p \mid (\ell_j - 1)m_j$ , set  $\mathfrak{k}_j = (\emptyset, \{j\}, \{j\})$ . If  $j \in \mathfrak{g}_2 \setminus \mathfrak{g}_2^\bullet$  and  $p \mid (\ell_j - 1)m_j$ , set  $\mathfrak{k}_j = (\emptyset, \emptyset, \{j\})$ . Hence  $\mathfrak{k}_{i_1}, \mathfrak{k}_{i_2}, \dots, \mathfrak{k}_{i_h} \in \mathbb{U}_{g,g}$  and  $B_{g,g,\mathfrak{k}_{i_1}}, B_{g,g,\mathfrak{k}_{i_2}}, \dots, B_{g,g,\mathfrak{k}_{i_h}} \in \mathbb{I}_g$ . Then  $B_{g,g,\mathfrak{k}_{i_1}} B_{g,g,\mathfrak{k}_{i_2}} \cdots B_{g,g,\mathfrak{k}_{i_h}} \neq O$  by Lemma 4.8. The desired lemma follows.  $\square$

**Lemma 5.5.** *Assume that  $g \in [0, d]$ . Then  $\mathbb{I}_g$  is a nilpotent two-sided ideal of the  $\mathbb{F}$ -subalgebra  $E_g^* \mathbb{T} E_g^*$  of  $\mathbb{T}$ . Furthermore, the nilpotent index of  $\mathbb{I}_g$  equals*

$$|\{a : a \in \mathfrak{g}_1, p \mid m_a - 1\}| + |\{a : a \in \mathfrak{g}_2, p \mid (\ell_a - 1)m_a\}| + 1.$$

*Proof.* Set  $\mathbb{U} = \{a : a \in \mathfrak{g}_1, p \mid m_a - 1\} \cup \{a : a \in \mathfrak{g}_2, p \mid (\ell_a - 1)m_a\}$ . So  $\mathbb{I}_g \neq \{O\}$  if and only if  $\mathbb{U} \neq \emptyset$ . So it is enough to check the case  $\mathbb{I}_g \neq \{O\}$ . Set  $h = |\mathbb{U}| > 0$ . Let  $B_{g,g,i_1}, B_{g,g,i_2}, \dots, B_{g,g,i_{h+1}} \in \mathbb{I}_g$ . So  $\mathbb{U}_{i_1}, \mathbb{U}_{i_2}, \dots, \mathbb{U}_{i_{h+1}}$  are nonempty subsets of  $\mathbb{U}$ . By the Pigeonhole Principle, there must exist  $j, \mathfrak{k} \in \{i_1, i_2, \dots, i_{h+1}\}$  such that  $\mathbb{U}_j \cap \mathbb{U}_{\mathfrak{k}} \neq \emptyset$ .

Assume that  $j = (\mathfrak{j}_{(1)}, \mathfrak{j}_{(2)}, \mathfrak{j}_{(3)})$  and  $\mathfrak{k} = (\mathfrak{k}_{(1)}, \mathfrak{k}_{(2)}, \mathfrak{k}_{(3)})$ . Assume that  $\ell \in \mathbb{U}_j \cap \mathbb{U}_{\mathfrak{k}}$ . Then  $\ell \in (\mathfrak{j}_{(1)} \cap \mathfrak{k}_{(1)}) \cup ((\mathfrak{j}_{(2)} \cap \mathfrak{k}_{(2)}) \cup ((\mathfrak{j}_{(3)} \cap \mathfrak{k}_{(3)}) \setminus (\mathfrak{j}_{(2)} \cap \mathfrak{k}_{(2)}))$ . Hence  $\ell \in \mathbb{U}_{j \cap \mathfrak{k}}$  and  $p \mid k_{j \cap \mathfrak{k}}$ . Hence the product of any  $h + 1$  matrices in  $\mathbb{I}_g$  is the zero matrix by Lemma 4.8. The desired lemma thus follows from the above discussion and Lemma 5.4.  $\square$

For further discussion, the next notations and a sequence of lemmas are required.

**Notation 5.6.** Assume that  $g, h \in [0, d]$  and  $\mathbf{i} = (\mathbf{i}_{(1)}, \mathbf{i}_{(2)}, \mathbf{i}_{(3)}) \in \mathbb{U}_{g,h}$ . Notice that  $((\mathfrak{g}_1 \cap \mathfrak{h}_1)^\circ, \mathbf{i}_{(3)}^\bullet, \mathfrak{g}_2 \cap \mathfrak{h}_2) \in \mathbb{U}_{g,h}$ . Use  $(g, h; \mathbf{i})$  to denote  $((\mathfrak{g}_1 \cap \mathfrak{h}_1)^\circ, \mathbf{i}_{(3)}^\bullet, \mathfrak{g}_2 \cap \mathfrak{h}_2)$ . Set  $n_{g,h,\mathbf{i}} = |(g, h; \mathbf{i})| - |\mathbf{i}|$ . Hence  $(g, h; \mathbf{i}) = (h, g; \mathbf{i})$  and  $n_{g,h,\mathbf{i}} = n_{h,g,\mathbf{i}}$ . Let  $j, k \in [0, n_{g,h,\mathbf{i}}]$ . Set  $\mathbb{U}_{g,h,\mathbf{i},j} = \{\mathbf{a} : \mathbf{i} \preceq \mathbf{a} \preceq (g, h; \mathbf{i}), |\mathbf{a}| - |\mathbf{i}| = j, p \nmid k_{\mathbf{a}}\}$ . Hence  $\mathbb{U}_{g,h,\mathbf{i},j} = \mathbb{U}_{h,g,\mathbf{i},j} \subseteq \mathbb{U}_{g,h}$ . If  $p \nmid k_{\mathbf{i}}$ , then  $\mathbb{U}_{g,h,\mathbf{i},0} = \{\mathbf{i}\}$ . If  $j \neq k$ , then  $\mathbb{U}_{g,h,\mathbf{i},j} \cap \mathbb{U}_{g,h,\mathbf{i},k} = \emptyset$ . For example, assume that  $p = n = \ell_1 = 2$ ,  $\ell_2 = m_1 = 3$ , and  $m_2 = 4$ . Then  $d = 8$ . Moreover, notice that  $(6, 7; (\emptyset, \emptyset, \{2\})) = (\emptyset, \{2\}, \{2\})$ ,  $n_{6,7,(\emptyset, \emptyset, \{2\})} = 1$ ,  $\mathbb{U}_{6,7,(\emptyset, \emptyset, \{2\}),0} = \mathbb{U}_{6,7,(\emptyset, \emptyset, \{2\}),1} = \emptyset$ .

**Notation 5.7.** Assume that  $g, h \in [0, d]$ ,  $\mathbf{i} \in \mathbb{U}_{g,h}$ , and  $p \nmid k_{[g,h,g]}$ . Let  $D_{g,h,\mathbf{i}}$  denote

$$\sum_{j=0}^{n_{g,h,\mathbf{i}}} \sum_{\mathfrak{k} \in \mathbb{U}_{g,h,\mathbf{i},j}} (-1)^j \overline{k_{[g,h,g]}}^{-1} \overline{k_{\mathfrak{k}}}^{-1} B_{g,h,\mathfrak{k}},$$

where the sum over the empty set equals the zero matrix. If  $p \nmid k_{\mathbf{i}}$ , Notation 5.6 and Theorem 3.16 imply that  $\mathbb{U}_{g,h,\mathbf{i},0} \neq \emptyset$  and  $D_{g,h,\mathbf{i}}$  is a defined nonzero matrix in  $\mathbb{T}$ . If  $g = h$ , notice that  $k_{[g,g,g]} = 1$  and  $D_{g,g,\mathbf{i}} = E_g^* D_{g,g,\mathbf{i}} E_g^* \in E_g^* \mathbb{T} E_g^*$  by Equation (2.2).

**Lemma 5.8.** Assume that  $g, h, i \in [0, d]$  and  $\mathbb{U} \subseteq \mathfrak{g}_1$ . Assume that  $\mathbb{U}^1 \subseteq \mathbb{V} \subseteq \mathbf{i}_1$  and  $\mathbb{U}^1 \subseteq \mathbb{V}^1 \subseteq (\mathfrak{h}_1 \cap \mathbf{i}_1)^\circ$ . Then  $\mathbb{U}^1, (\mathfrak{g}_1 \cap \mathbb{V}) \setminus (\mathbb{U} \cup \mathbb{U}^1), (\mathfrak{h}_1 \cap \mathbf{i}_1)^\circ \setminus (\mathfrak{g}_1 \cup \mathbb{U}^1), (\mathbf{i}_1 \cap \mathbb{U}) \setminus \mathbb{U}^1$  are pairwise disjoint. Moreover,  $\mathfrak{g}_1 \cap \mathbf{i}_1 \cap (\mathbb{U} \cup \mathbb{V}) = \mathfrak{g}_1 \cap \mathbf{i}_1 \cap (\mathbb{U} \cup \mathbb{V}^1)$  if and only if  $\mathbb{V}^1 = \mathbb{U}^1 \cup ((\mathfrak{g}_1 \cap \mathbb{V}) \setminus (\mathbb{U} \cup \mathbb{U}^1)) \cup \mathbb{W}$  and  $\mathbb{W} \subseteq ((\mathfrak{h}_1 \cap \mathbf{i}_1)^\circ \setminus (\mathfrak{g}_1 \cup \mathbb{U}^1)) \cup ((\mathbf{i}_1 \cap \mathbb{U}) \setminus \mathbb{U}^1)$ .

*Proof.* The first statement is clear. One direction of the second statement is obvious since  $\mathfrak{g}_1 \cap \mathbf{i}_1 \cap (\mathbb{U} \cup \mathbb{V}) = \mathfrak{g}_1 \cap \mathbf{i}_1 \cap (\mathbb{U} \cup \mathbb{U}^1 \cup ((\mathfrak{g}_1 \cap \mathbb{V}) \setminus (\mathbb{U} \cup \mathbb{U}^1)))$ . Assume further that  $\mathfrak{g}_1 \cap \mathbf{i}_1 \cap (\mathbb{U} \cup \mathbb{V}) = \mathfrak{g}_1 \cap \mathbf{i}_1 \cap (\mathbb{U} \cup \mathbb{V}^1)$ . Hence  $(\mathfrak{g}_1 \cap \mathbb{V}) \setminus (\mathbb{U} \cup \mathbb{U}^1) = (\mathfrak{g}_1 \cap \mathbb{V}^1) \setminus (\mathbb{U} \cup \mathbb{U}^1)$ . If  $j \in \mathbb{V}^1 \setminus (\mathfrak{g}_1 \cup \mathbb{U}^1)$ , then  $j \in (\mathfrak{h}_1 \cap \mathbf{i}_1)^\circ \setminus (\mathfrak{g}_1 \cup \mathbb{U}^1)$ . The desired lemma follows.  $\square$

**Lemma 5.9.** Assume that  $g, h, i \in [0, d]$  and  $\mathbb{U} \subseteq \mathfrak{g}_2$ . Assume that  $\mathbb{U}^1 \subseteq \mathbb{V} \subseteq \mathbf{i}_2$  and  $\mathbb{U}^1 \subseteq \mathbb{V}^1 \subseteq \mathbb{W} \subseteq (\mathfrak{h}_2 \cap \mathbf{i}_2)^\bullet$ . Then  $\mathbb{U}^1, (\mathfrak{g}_2 \cap \mathbb{V}) \setminus (\mathbb{U} \cup \mathbb{U}^1), \mathbb{W} \setminus (\mathfrak{g}_2 \cup \mathbb{U}^1), (\mathbb{U} \cap \mathbb{W}) \setminus \mathbb{U}^1$  are pairwise disjoint. Moreover,  $\mathfrak{g}_2 \cap \mathbf{i}_2 \cap (\mathbb{U} \cup \mathbb{V}) = \mathfrak{g}_2 \cap \mathbf{i}_2 \cap (\mathbb{U} \cup \mathbb{V}^1)$  if and only if  $\mathbb{V}^1 = \mathbb{U}^1 \cup ((\mathfrak{g}_2 \cap \mathbb{V}) \setminus (\mathbb{U} \cup \mathbb{U}^1)) \cup \mathbb{W}^1$  and  $\mathbb{W}^1 \subseteq (\mathbb{W} \setminus (\mathfrak{g}_2 \cup \mathbb{U}^1)) \cup ((\mathbb{U} \cap \mathbb{W}) \setminus \mathbb{U}^1)$ .

*Proof.* The first statement is clear. One direction of the second statement is obvious since  $\mathfrak{g}_2 \cap \mathbf{i}_2 \cap (\mathbb{U} \cup \mathbb{V}) = \mathfrak{g}_2 \cap \mathbf{i}_2 \cap (\mathbb{U} \cup \mathbb{U}^1 \cup ((\mathfrak{g}_2 \cap \mathbb{V}) \setminus (\mathbb{U} \cup \mathbb{U}^1)))$ . Assume further that  $\mathfrak{g}_2 \cap \mathbf{i}_2 \cap (\mathbb{U} \cup \mathbb{V}) = \mathfrak{g}_2 \cap \mathbf{i}_2 \cap (\mathbb{U} \cup \mathbb{V}^1)$ . Hence  $(\mathfrak{g}_2 \cap \mathbb{V}) \setminus (\mathbb{U} \cup \mathbb{U}^1) = (\mathfrak{g}_2 \cap \mathbb{V}^1) \setminus (\mathbb{U} \cup \mathbb{U}^1)$ . If  $j \in \mathbb{V}^1 \setminus (\mathfrak{g}_2 \cup \mathbb{U}^1)$ , notice that  $j \in \mathbb{W} \setminus (\mathfrak{g}_2 \cup \mathbb{U}^1)$ . The desired lemma follows.  $\square$

**Lemma 5.10.** Assume that  $g, h, i \in [0, d]$  and  $\mathbb{U} \subseteq \mathfrak{g}_2$ . Assume that  $\mathbb{U}^1 \subseteq \mathbb{V} \subseteq \mathbf{i}_2$  and  $\mathbb{U}^1 \subseteq \mathbb{V}^1 \subseteq \mathfrak{h}_2 \cap \mathbf{i}_2$ . Then  $\mathbb{U}^1, (\mathfrak{g}_2 \cap \mathbb{V}) \setminus (\mathbb{U} \cup \mathbb{U}^1), (\mathfrak{h}_2 \cap \mathbf{i}_2) \setminus (\mathfrak{g}_2 \cup \mathbb{U}^1), (\mathbf{i}_2 \cap \mathbb{U}) \setminus \mathbb{U}^1$  are pairwise disjoint. Moreover,  $\mathfrak{g}_2 \cap \mathbf{i}_2 \cap (\mathbb{U} \cup \mathbb{V}) = \mathfrak{g}_2 \cap \mathbf{i}_2 \cap (\mathbb{U} \cup \mathbb{V}^1)$  if and only if  $\mathbb{V}^1 = \mathbb{U}^1 \cup ((\mathfrak{g}_2 \cap \mathbb{V}) \setminus (\mathbb{U} \cup \mathbb{U}^1)) \cup \mathbb{W}$  and  $\mathbb{W} \subseteq ((\mathfrak{h}_2 \cap \mathbf{i}_2) \setminus (\mathfrak{g}_2 \cup \mathbb{U}^1)) \cup ((\mathbf{i}_2 \cap \mathbb{U}) \setminus \mathbb{U}^1)$ .

*Proof.* The first statement is clear. One direction of the second statement is obvious since  $\mathfrak{g}_2 \cap \mathbf{i}_2 \cap (\mathbb{U} \cup \mathbb{V}) = \mathfrak{g}_2 \cap \mathbf{i}_2 \cap (\mathbb{U} \cup \mathbb{U}^1 \cup ((\mathfrak{g}_2 \cap \mathbb{V}) \setminus (\mathbb{U} \cup \mathbb{U}^1)))$ . Assume further that  $\mathfrak{g}_2 \cap \mathbf{i}_2 \cap (\mathbb{U} \cup \mathbb{V}) = \mathfrak{g}_2 \cap \mathbf{i}_2 \cap (\mathbb{U} \cup \mathbb{V}^1)$ . Hence  $(\mathfrak{g}_2 \cap \mathbb{V}) \setminus (\mathbb{U} \cup \mathbb{U}^1) = (\mathfrak{g}_2 \cap \mathbb{V}^1) \setminus (\mathbb{U} \cup \mathbb{U}^1)$ . If  $j \in \mathbb{V}^1 \setminus (\mathfrak{g}_2 \cup \mathbb{U}^1)$ , then  $j \in (\mathfrak{h}_2 \cap \mathbf{i}_2) \setminus (\mathfrak{g}_2 \cup \mathbb{U}^1)$ . The desired lemma follows.  $\square$

**Lemma 5.11.** *Assume that  $g, h, i \in [0, d]$  and  $\mathbf{j} = (\mathbb{j}_{(1)}, \mathbb{j}_{(2)}, \mathbb{j}_{(3)}) \in \mathbb{U}_{g,h}$ . Assume that  $\mathfrak{k} = (\mathbb{k}_{(1)}, \mathbb{k}_{(2)}, \mathbb{k}_{(3)})$ ,  $\mathfrak{l} = (\mathbb{l}_{(1)}, \mathbb{l}_{(2)}, \mathbb{l}_{(3)})$ ,  $\mathfrak{m} = (\mathbb{m}_{(1)}, \mathbb{m}_{(2)}, \mathbb{m}_{(3)}) \in \mathbb{U}_{h,i}$ . Assume that  $\mathfrak{k} \preceq \mathfrak{l} \preceq (h, i; \mathfrak{k})$  and  $\mathfrak{k} \preceq \mathfrak{m} \preceq (h, i; \mathfrak{k})$ . Then  $(g, h, i, \mathbf{j}, \mathfrak{l}) = (g, h, i, \mathbf{j}, \mathfrak{m})$  if and only if  $\mathbb{m}_{(1)} = \mathbb{k}_{(1)} \cup ((\mathbb{g}_1 \cap \mathbb{l}_{(1)}) \setminus (\mathbb{j}_{(1)} \cup \mathbb{k}_{(1)})) \cup \mathbb{U}$ ,  $\mathbb{U} \subseteq ((\mathbb{h}_1 \cap \mathbb{i}_1)^\circ \setminus (\mathbb{g}_1 \cup \mathbb{k}_{(1)})) \cup ((\mathbb{i}_1 \cap \mathbb{j}_{(1)}) \setminus \mathbb{k}_{(1)})$ ,  $\mathbb{m}_{(2)} = \mathbb{k}_{(2)} \cup ((\mathbb{g}_2 \cap \mathbb{l}_{(2)}) \setminus (\mathbb{j}_{(2)} \cup \mathbb{k}_{(2)})) \cup \mathbb{V}$ ,  $\mathbb{V} \subseteq (\mathbb{k}_{(3)} \bullet \setminus (\mathbb{g}_2 \cup \mathbb{k}_{(2)})) \cup ((\mathbb{j}_{(2)} \cap \mathbb{k}_{(3)}) \setminus \mathbb{k}_{(2)})$ ,  $\mathbb{m}_{(3)} = \mathbb{k}_{(3)} \cup ((\mathbb{g}_2 \cap \mathbb{l}_{(3)}) \setminus (\mathbb{j}_{(3)} \cup \mathbb{k}_{(3)})) \cup \mathbb{W}$ ,  $\mathbb{W} \subseteq ((\mathbb{h}_2 \cap \mathbb{i}_2) \setminus (\mathbb{g}_2 \cup \mathbb{k}_{(3)})) \cup ((\mathbb{i}_2 \cap \mathbb{j}_{(3)}) \setminus \mathbb{k}_{(3)})$ .*

*Proof.*  $(g, h, i, \mathbf{j}, \mathfrak{l}) = (g, h, i, \mathbf{j}, \mathfrak{m})$  if and only if  $\mathbb{g}_1 \cap \mathbb{i}_1 \cap (\mathbb{j}_{(1)} \cup \mathbb{l}_{(1)}) = \mathbb{g}_1 \cap \mathbb{i}_1 \cap (\mathbb{j}_{(1)} \cup \mathbb{m}_{(1)})$ ,  $\mathbb{g}_2 \cap \mathbb{i}_2 \cap (\mathbb{j}_{(2)} \cup \mathbb{l}_{(2)}) = \mathbb{g}_2 \cap \mathbb{i}_2 \cap (\mathbb{j}_{(2)} \cup \mathbb{m}_{(2)})$ ,  $\mathbb{g}_2 \cap \mathbb{i}_2 \cap (\mathbb{j}_{(3)} \cup \mathbb{l}_{(3)}) = \mathbb{g}_2 \cap \mathbb{i}_2 \cap (\mathbb{j}_{(3)} \cup \mathbb{m}_{(3)})$ . The desired lemma thus follows from combining Lemmas 5.8, 5.9, and 5.10.  $\square$

**Lemma 5.12.** *Assume that  $g, h, i \in [0, d]$ ,  $\mathbf{j} \in \mathbb{U}_{g,h}$ ,  $\mathfrak{k}, \mathfrak{l}, \mathfrak{m} \in \mathbb{U}_{h,i}$ ,  $\mathfrak{k} \preceq \mathfrak{l} \preceq (h, i; \mathfrak{k})$ ,  $\mathfrak{k} \preceq \mathfrak{m} \preceq (h, i; \mathfrak{k})$ ,  $(g, h, i, \mathbf{j}, \mathfrak{l}) = (g, h, i, \mathbf{j}, \mathfrak{m})$ , and  $p \nmid k_{[g,h,g]} k_j k_\mathfrak{k} k_\mathfrak{l}$ . Then  $p \nmid k_\mathfrak{m}$ .*

*Proof.* Assume that  $\mathbf{j} = (\mathbb{j}_{(1)}, \mathbb{j}_{(2)}, \mathbb{j}_{(3)})$ ,  $\mathfrak{k} = (\mathbb{k}_{(1)}, \mathbb{k}_{(2)}, \mathbb{k}_{(3)})$ ,  $\mathfrak{l} = (\mathbb{l}_{(1)}, \mathbb{l}_{(2)}, \mathbb{l}_{(3)})$ , and  $\mathfrak{m} = (\mathbb{m}_{(1)}, \mathbb{m}_{(2)}, \mathbb{m}_{(3)})$ . As  $p \nmid k_j k_\mathfrak{k} k_\mathfrak{l}$ , notice that  $\mathbb{U}_\mathfrak{j} = \mathbb{U}_\mathfrak{k} = \mathbb{U}_\mathfrak{l} = \emptyset$ . As  $p \nmid k_{[g,h,g]}$ , notice that  $\{a : a \in \mathbb{h}_1 \setminus \mathbb{g}_1, p \mid m_a - 1\} \cup \{a : a \in \mathbb{h}_2 \setminus \mathbb{g}_2, p \mid (\ell_a - 1)m_a\} = \emptyset$ . By Lemma 5.11, notice that  $\mathbb{m}_{(1)} \subseteq \mathbb{j}_{(1)} \cup \mathbb{k}_{(1)} \cup \mathbb{l}_{(1)} \cup (\mathbb{h}_1 \setminus \mathbb{g}_1)$ ,  $\mathbb{m}_{(2)} \subseteq \mathbb{j}_{(2)} \cup \mathbb{k}_{(2)} \cup \mathbb{l}_{(2)} \cup (\mathbb{h}_2 \setminus \mathbb{g}_2)$ , and  $\mathbb{m}_{(3)} \setminus \mathbb{m}_{(2)} \subseteq \mathbb{j}_{(3)} \cup (\mathbb{k}_{(3)} \setminus \mathbb{k}_{(2)}) \cup (\mathbb{l}_{(3)} \setminus \mathbb{l}_{(2)}) \cup (\mathbb{h}_2 \setminus \mathbb{g}_2)$ . Hence  $\mathbb{U}_\mathfrak{m} = \emptyset$  by the above discussion and Notation 5.3. The desired lemma is from Notation 5.3.  $\square$

**Lemma 5.13.** *Assume that  $g, h, i \in [0, d]$ ,  $\mathbf{j} \in \mathbb{U}_{g,h}$ , and  $\mathfrak{k}, \mathfrak{l}, \mathfrak{m} \in \mathbb{U}_{h,i}$ . Assume that  $\mathfrak{k} \preceq \mathfrak{l} \preceq (h, i; \mathfrak{k})$ ,  $\mathfrak{k} \preceq \mathfrak{m} \preceq (h, i; \mathfrak{k})$ , and  $(g, h, i, \mathbf{j}, \mathfrak{l}) = (g, h, i, \mathbf{j}, \mathfrak{m})$ . Then  $k_{g \cap \mathfrak{m}} k_{\mathbf{j} \cap \mathfrak{l}} = k_{g \cap \mathfrak{l}} k_{\mathbf{j} \cap \mathfrak{m}}$ .*

*Proof.* Assume that  $\mathbf{j} = (\mathbb{j}_{(1)}, \mathbb{j}_{(2)}, \mathbb{j}_{(3)})$ ,  $\mathfrak{k} = (\mathbb{k}_{(1)}, \mathbb{k}_{(2)}, \mathbb{k}_{(3)})$ ,  $\mathfrak{l} = (\mathbb{l}_{(1)}, \mathbb{l}_{(2)}, \mathbb{l}_{(3)})$ , and  $\mathfrak{m} = (\mathbb{m}_{(1)}, \mathbb{m}_{(2)}, \mathbb{m}_{(3)})$ . By Lemma 5.11, there are  $\mathbb{U} \subseteq (\mathbb{i}_1 \cap \mathbb{j}_{(1)}) \setminus \mathbb{k}_{(1)}$ ,  $\mathbb{V} \subseteq (\mathbb{j}_{(2)} \cap \mathbb{k}_{(3)}) \setminus \mathbb{k}_{(2)}$ , and  $\mathbb{W} \subseteq (\mathbb{i}_2 \cap \mathbb{j}_{(3)}) \setminus \mathbb{k}_{(3)}$  such that  $\mathbb{g}_1 \cap \mathbb{m}_{(1)} = (\mathbb{j}_{(1)} \cap \mathbb{k}_{(1)}) \cup ((\mathbb{g}_1 \cap \mathbb{l}_{(1)}) \setminus \mathbb{j}_{(1)}) \cup \mathbb{U}$ ,  $\mathbb{g}_2 \cap \mathbb{m}_{(2)} = (\mathbb{j}_{(2)} \cap \mathbb{k}_{(2)}) \cup ((\mathbb{g}_2 \cap \mathbb{l}_{(2)}) \setminus \mathbb{j}_{(2)}) \cup \mathbb{V}$ ,  $\mathbb{g}_2 \cap \mathbb{m}_{(3)} = (\mathbb{j}_{(3)} \cap \mathbb{k}_{(3)}) \cup ((\mathbb{g}_2 \cap \mathbb{l}_{(3)}) \setminus \mathbb{j}_{(3)}) \cup \mathbb{W}$ ,  $\mathbb{j}_{(1)} \cap \mathbb{m}_{(1)} = (\mathbb{j}_{(1)} \cap \mathbb{k}_{(1)}) \cup \mathbb{U}$ ,  $\mathbb{j}_{(2)} \cap \mathbb{m}_{(2)} = (\mathbb{j}_{(2)} \cap \mathbb{k}_{(2)}) \cup \mathbb{V}$ , and  $\mathbb{j}_{(3)} \cap \mathbb{m}_{(3)} = (\mathbb{j}_{(3)} \cap \mathbb{k}_{(3)}) \cup \mathbb{W}$ . The desired lemma thus follows from Lemma 3.23 and a direct computation.  $\square$

**Lemma 5.14.** *Assume that  $g, h, i \in [0, d]$ . Assume that  $\mathbf{j} = (\mathbb{j}_{(1)}, \mathbb{j}_{(2)}, \mathbb{j}_{(3)}) \in \mathbb{U}_{g,h}$  and  $\mathfrak{k} = (\mathbb{k}_{(1)}, \mathbb{k}_{(2)}, \mathbb{k}_{(3)}) \in \mathbb{U}_{h,i}$ . Assume that  $p \nmid k_{[g,h,g]} k_{[h,i,h]} k_j k_\mathfrak{k}$ . Then  $B_{g,h,j} D_{h,i,\mathfrak{k}} \neq O$  only if the containments  $((\mathbb{h}_1 \cap \mathbb{i}_1)^\circ \setminus \mathbb{g}_1) \cup (\mathbb{i}_1 \cap \mathbb{j}_{(1)}) \subseteq \mathbb{k}_{(1)}$ ,  $(\mathbb{k}_{(3)} \bullet \setminus \mathbb{g}_2) \cup (\mathbb{j}_{(2)} \cap \mathbb{k}_{(3)}) \subseteq \mathbb{k}_{(2)}$ , and  $((\mathbb{h}_2 \cap \mathbb{i}_2) \setminus \mathbb{g}_2) \cup (\mathbb{i}_2 \cap \mathbb{j}_{(3)}) \subseteq \mathbb{k}_{(3)}$  hold.*

*Proof.* Assume that the three containments do not hold together and  $B_{g,h,j} D_{h,i,\mathfrak{k}} \neq O$ . Assume that  $B_{h,i,\mathfrak{l}} \in \text{Supp}_{\mathbb{B}_2}(D_{h,i,\mathfrak{k}})$  and  $B_{g,i,(g,h,i,j,\mathfrak{l})} \in \text{Supp}_{\mathbb{B}_2}(B_{g,h,j} D_{h,i,\mathfrak{k}})$  by Theorem 3.32. Therefore  $p \nmid k_{[g,h,g]} k_j k_\mathfrak{k} k_\mathfrak{l}$  by Notations 5.6 and 5.7. Set  $\mathfrak{l} = (\mathbb{l}_{(1)}, \mathbb{l}_{(2)}, \mathbb{l}_{(3)})$ . Let  $m = |(\mathbb{g}_1 \cap \mathbb{l}_{(1)}) \setminus (\mathbb{j}_{(1)} \cup \mathbb{k}_{(1)})| + |(\mathbb{g}_2 \cap \mathbb{l}_{(2)}) \setminus (\mathbb{j}_{(2)} \cup \mathbb{k}_{(2)})| + |(\mathbb{g}_2 \cap \mathbb{l}_{(3)}) \setminus (\mathbb{j}_{(3)} \cup \mathbb{k}_{(3)})|$ . Let  $q$  be the sum of  $|(\mathbb{h}_1 \cap \mathbb{i}_1)^\circ \setminus (\mathbb{g}_1 \cup \mathbb{k}_{(1)})|$ ,  $|(\mathbb{i}_1 \cap \mathbb{j}_{(1)}) \setminus \mathbb{k}_{(1)}|$ ,  $|\mathbb{k}_{(3)} \bullet \setminus (\mathbb{g}_2 \cup \mathbb{k}_{(2)})|$ ,  $|(\mathbb{j}_{(2)} \cap \mathbb{k}_{(3)}) \setminus \mathbb{k}_{(2)}|$ ,  $|(\mathbb{h}_2 \cap \mathbb{i}_2) \setminus (\mathbb{g}_2 \cup \mathbb{k}_{(3)})|$ , and  $|(\mathbb{i}_2 \cap \mathbb{j}_{(3)}) \setminus \mathbb{k}_{(3)}|$ . Therefore  $q > 0$ . By combining Lemmas 5.8, 5.9, 5.10, 5.11, 5.12, 5.13, 4.5, Theorems 3.32, and 3.16,

$$c_{g,i,(g,h,i,j,\mathfrak{l})}(B_{g,h,j} D_{h,i,\mathfrak{k}}) = \sum_{r=0}^q (-1)^{m+r} k_{[h,i,h]}^{-1} k_{g \cap \mathfrak{l}}^{-1} k_{[g,h,i]} k_{\mathbf{j} \cap \mathfrak{l}} k_{\mathbf{j} \cap \mathfrak{i}} \binom{q}{r} = \bar{0},$$

which is absurd as  $B_{g,i,(g,h,i,j,\mathfrak{l})} \in \text{Supp}_{\mathbb{B}_2}(B_{g,h,j} D_{h,i,\mathfrak{k}})$ . The desired lemma follows.  $\square$

**Lemma 5.15.** *Assume that  $g \in [0, d]$  and  $\mathfrak{h}, \mathfrak{i} \in \mathbb{U}_{g,g}$ . Assume that  $p \nmid k_{\mathfrak{h}}k_{\mathfrak{i}}$ . Then*

$$B_{g,g,\mathfrak{h}}D_{g,g,\mathfrak{i}} = D_{g,g,\mathfrak{i}}B_{g,g,\mathfrak{h}} = \begin{cases} \overline{k_{\mathfrak{h}}}D_{g,g,\mathfrak{i}}, & \text{if } \mathfrak{h} \preceq \mathfrak{i}, \\ O, & \text{otherwise.} \end{cases}$$

*Proof.* If  $\mathfrak{h} \preceq \mathfrak{i}$ , notice that  $B_{g,g,\mathfrak{h}}D_{g,g,\mathfrak{i}} = D_{g,g,\mathfrak{i}}B_{g,g,\mathfrak{h}} = \overline{k_{\mathfrak{h}}}D_{g,g,\mathfrak{i}}$  by Notation 5.7 and Lemma 4.8. By combining Lemmas 4.8, 5.14, and a direct computation, notice that  $B_{g,g,\mathfrak{h}}D_{g,g,\mathfrak{i}} = D_{g,g,\mathfrak{i}}B_{g,g,\mathfrak{h}} \neq O$  only if  $\mathfrak{h} \preceq \mathfrak{i}$ . The desired lemma follows.  $\square$

**Lemma 5.16.** *Assume that  $g \in [0, d]$  and  $\mathfrak{h}, \mathfrak{i} \in \mathbb{U}_{g,g}$ . Assume that  $p \nmid k_{\mathfrak{h}}k_{\mathfrak{i}}$ . Then*

$$D_{g,g,\mathfrak{h}}D_{g,g,\mathfrak{i}} = D_{g,g,\mathfrak{i}}D_{g,g,\mathfrak{h}} = \begin{cases} D_{g,g,\mathfrak{h}}, & \text{if } \mathfrak{h} = \mathfrak{i}, \\ O, & \text{otherwise.} \end{cases}$$

*Proof.* By combining Notation 5.7, Lemmas 4.8, 5.15,  $D_{g,g,\mathfrak{h}}D_{g,g,\mathfrak{i}} = D_{g,g,\mathfrak{i}}D_{g,g,\mathfrak{h}} \neq O$  only if  $\mathfrak{h} \preceq \mathfrak{i} \preceq \mathfrak{h}$ . Hence  $D_{g,g,\mathfrak{h}}D_{g,g,\mathfrak{i}} = D_{g,g,\mathfrak{i}}D_{g,g,\mathfrak{h}} \neq O$  only if  $\mathfrak{h} = \mathfrak{i}$ . By Notation 5.7 and Lemma 5.15, notice that  $D_{g,g,\mathfrak{h}}D_{g,g,\mathfrak{h}} = D_{g,g,\mathfrak{h}}$ . The desired lemma follows.  $\square$

We are now ready to give the first main result of this section and two corollaries.

**Theorem 5.17.** *Assume that  $g \in [0, d]$ . Then  $|\{\mathfrak{a} : \mathfrak{a} \in \mathbb{U}_{g,g}, p \nmid k_{\mathfrak{a}}\}| = 2^{n_{g,1}+n_{g,3}}3^{n_{g,2}}$ , where  $n_{g,1} = |\{a : a \in \mathfrak{g}_1^\circ, p \nmid m_a - 1\}|$ ,  $n_{g,2} = |\{a : a \in \mathfrak{g}_2^\bullet, p \nmid (\ell_a - 1)m_a\}|$ , and  $n_{g,3} = |\{a : a \in \mathfrak{g}_2 \setminus \mathfrak{g}_2^\bullet, p \nmid m_a\}|$ . Furthermore,  $E_g^* \mathbb{T} E_g^* / \mathbb{I}_g \cong 2^{n_{g,1}+n_{g,3}}3^{n_{g,2}}M_1(\mathbb{F})$  as  $\mathbb{F}$ -algebras. In particular,  $\text{Rad}(E_g^* \mathbb{T} E_g^*) = \mathbb{I}_g$  and the nilpotent index of  $\text{Rad}(E_g^* \mathbb{T} E_g^*)$  equals  $|\{a : a \in \mathfrak{g}_1, p \mid m_a - 1\}| + |\{a : a \in \mathfrak{g}_2, p \mid (\ell_a - 1)m_a\}| + 1$ .*

*Proof.* Set  $\mathbb{U} = \{D_{g,g,\mathfrak{a}} : \mathfrak{a} \in \mathbb{U}_{g,g}, p \nmid k_{\mathfrak{a}}\} \cup \{B_{g,g,\mathfrak{a}} : \mathfrak{a} \in \mathbb{U}_{g,g}, p \mid k_{\mathfrak{a}}\}$ . The  $\mathbb{F}$ -subalgebra  $E_g^* \mathbb{T} E_g^*$  of  $\mathbb{T}$  has an  $\mathbb{F}$ -basis  $\mathbb{U}$  by combining Notation 5.7, Theorem 3.16, Lemmas 5.16, and 4.8. Hence  $\{D_{g,g,\mathfrak{a}} + \mathbb{I}_g : \mathfrak{a} \in \mathbb{U}_{g,g}, p \nmid k_{\mathfrak{a}}\}$  is an  $\mathbb{F}$ -basis of  $E_g^* \mathbb{T} E_g^* / \mathbb{I}_g$ . Hence  $|\{D_{g,g,\mathfrak{a}} + \mathbb{I}_g : \mathfrak{a} \in \mathbb{U}_{g,g}, p \nmid k_{\mathfrak{a}}\}| = |\{\mathfrak{a} : \mathfrak{a} \in \mathbb{U}_{g,g}, p \nmid k_{\mathfrak{a}}\}| = 2^{n_{g,1}+n_{g,3}}3^{n_{g,2}}$  by a direct computation. The first two statements are from the above discussion and Lemma 5.16. The desired theorem is thus from the second statement and Lemma 5.5.  $\square$

**Corollary 5.18.** *Assume that  $g \in [0, d]$ ,  $\mathfrak{h} \in \mathbb{U}_{g,g}$ ,  $p \nmid k_{\mathfrak{h}}$ ,  $\text{Irr}(g, \mathfrak{h}) = \langle \{D_{g,g,\mathfrak{h}} + \mathbb{I}_g\} \rangle$ .  $\{\text{Irr}(g, \mathfrak{a}) : \mathfrak{a} \in \mathbb{U}_{g,g}, p \nmid k_{\mathfrak{a}}\}$  contains exactly the representatives of all isomorphic classes of irreducible  $E_g^* \mathbb{T} E_g^*$ -modules. Moreover, all irreducible  $E_g^* \mathbb{T} E_g^*$ -modules are precisely all absolutely irreducible  $E_g^* \mathbb{T} E_g^*$ -modules.*

*Proof.* Notice that  $\text{Irr}(g, \mathfrak{h})$  is an irreducible  $E_g^* \mathbb{T} E_g^*$ -module by combining Lemmas 4.8, 5.2, and 5.15. Assume that  $\mathfrak{i}, \mathfrak{j} \in \mathbb{U}_{g,g}$  and  $p \nmid k_{\mathfrak{i}}k_{\mathfrak{j}}$ . Lemma 5.16 thus can imply that  $\mathfrak{i} = \mathfrak{j}$  if and only if  $\text{Irr}(g, \mathfrak{i}) \cong \text{Irr}(g, \mathfrak{j})$  as irreducible  $E_g^* \mathbb{T} E_g^*$ -modules. Hence the desired corollary follows from Theorem 5.17 and Lemma 2.2.  $\square$

**Corollary 5.19.** *Assume that  $g \in [0, d]$ . Then the  $\mathbb{F}$ -subalgebra  $E_g^* \mathbb{T} E_g^*$  of  $\mathbb{T}$  is a semisimple  $\mathbb{F}$ -algebra if and only if  $p \nmid k_g$ . In particular, the  $\mathbb{F}$ -subalgebra  $E_g^* \mathbb{T} E_g^*$  of  $\mathbb{T}$  is a semisimple  $\mathbb{F}$ -algebra if and only if  $p \nmid \prod_{h \in \mathfrak{g}_1} (m_h - 1) \prod_{i \in \mathfrak{g}_2} (\ell_i - 1)m_i$ .*

*Proof.* The desired corollary follows from Lemma 3.24 and Theorem 5.17.  $\square$

We next introduce another lemma and the second main result of this section.

**Lemma 5.20.**  $\mathbb{S}$  is a  $p'$ -valenced scheme if and only if  $p \nmid k_d k_{d/2}$ . In particular,  $\mathbb{S}$  is a  $p'$ -valenced scheme if and only if  $p \nmid \prod_{g=1}^n (\ell_g - 1)(m_g - 1)m_g$ .

*Proof.* The desired lemma follows from Lemma 3.24 and a direct computation.  $\square$

**Theorem 5.21.**  $\mathbb{T}$  is a semisimple  $\mathbb{F}$ -algebra if and only if  $\mathbb{S}$  is a  $p'$ -valenced scheme. In particular,  $\mathbb{T}$  is a semisimple  $\mathbb{F}$ -algebra if and only if  $p \nmid \prod_{g=1}^n (\ell_g - 1)(m_g - 1)m_g$ .

*Proof.* Assume that  $\mathbb{S}$  is a  $p'$ -valenced scheme. Notice that  $p \nmid k_g$  for any  $g \in [0, d]$ . Corollary 5.19 implies that  $\text{Rad}(E_g^* \mathbb{T} E_g^*) = \{O\}$  for any  $g \in [0, d]$ . Pick  $M \in \text{Rad}(\mathbb{T})$ . Assume further that  $M \neq O$ . Lemma 2.1 implies that  $E_g^* M E_g^* = O$  for any  $g \in [0, d]$ . As  $M \neq O$  and Equation (2.3) holds,  $E_h^* M E_i^* \neq O$  for some distinct  $h, i \in [0, d]$ . By Lemma 3.1, there is  $j \in [0, d]$  such that  $j_1 = i_1 \setminus h_1$  and  $j_2 = i_2 \setminus h_2$ . So  $k_j = k_{[h, i, h]}$ .

By Equation (2.2), there must exist  $k \in \mathbb{N}$  and pairwise distinct  $l_1, l_2, \dots, l_k \in \mathbb{U}_{h, i}$  such that  $\text{Supp}_{\mathbb{B}_2}(E_h^* M E_i^*) = \{B_{h, i, l_1}, B_{h, i, l_2}, \dots, B_{h, i, l_k}\}$ . By a direct computation, notice that  $(h, i, h, l_1, \mathfrak{o}), (h, i, h, l_2, \mathfrak{o}), \dots, (h, i, h, l_k, \mathfrak{o})$  are pairwise distinct in  $\mathbb{U}_{h, h}$ . By Theorem 3.16,  $B_{h, h, (h, i, h, l_1, \mathfrak{o})}, B_{h, h, (h, i, h, l_2, \mathfrak{o})}, \dots, B_{h, h, (h, i, h, l_k, \mathfrak{o})}$  are pairwise distinct. As  $k_{l_1 \setminus h} = k_{\mathfrak{o} \setminus h} = k_{l_1 \cap \mathfrak{o}} = 1$ ,  $c_{h, h, (h, i, h, l_1, \mathfrak{o})}(E_h^* M E_i^* B_{i, h, \mathfrak{o}}) = c_{h, i, l_1}(E_h^* M E_i^*) \overline{k_j} \in \mathbb{F}^\times$  by combining Equation (2.2), Theorems 3.16, 3.32. In particular,  $E_h^* M E_i^* B_{i, h, \mathfrak{o}} \neq O$  by Theorem 3.16. As  $M \in \text{Rad}(\mathbb{T})$ , notice that  $E_h^* M E_i^* B_{i, h, \mathfrak{o}} \in \text{Rad}(E_h^* \mathbb{T} E_h^*) \setminus \{O\}$  by Equation (2.2) and Lemma 2.1. This contradicts the equality  $\text{Rad}(E_h^* \mathbb{T} E_h^*) = \{O\}$ . So  $\text{Rad}(\mathbb{T}) = \{O\}$  and  $\mathbb{T}$  is a semisimple  $\mathbb{F}$ -algebra. The first statement is from Lemma 2.3. The desired theorem is thus from the first statement and Lemma 5.20.  $\square$

We conclude this section by two corollaries of Theorem 5.21 and an example.

**Corollary 5.22.**  $\mathbb{T}$  is a semisimple  $\mathbb{F}$ -algebra if and only if the  $\mathbb{F}$ -subalgebra  $E_g^* \mathbb{T} E_g^*$  of  $\mathbb{T}$  is a semisimple  $\mathbb{F}$ -algebra for any  $g \in [0, d]$ .  $\mathbb{T}$  is a semisimple  $\mathbb{F}$ -algebra if and only if the  $\mathbb{F}$ -subalgebras  $E_d^* \mathbb{T} E_d^*$ ,  $E_{d/2}^* \mathbb{T} E_{d/2}^*$  of  $\mathbb{T}$  are semisimple  $\mathbb{F}$ -algebras.

*Proof.* The first statement is from Corollary 5.19 and Theorem 5.21. The desired corollary is from combining the first statement, Lemma 5.20, and Corollary 5.19.  $\square$

**Corollary 5.23.**  $\mathbb{T}$  is a semisimple  $\mathbb{F}$ -algebra if and only if the  $\mathbb{F}$ -subalgebra  $Z(\mathbb{T})$  of  $\mathbb{T}$  is a semisimple  $\mathbb{F}$ -algebra.

*Proof.* For any  $g \in [0, d]$ , Lemmas 3.24 and 3.22 imply that  $(g_1^\circ, g_2^\bullet, g_2) \in \mathbb{U}_{g, g}$  and  $k_g = k_{(g_1^\circ, g_2^\bullet, g_2)}$ . For any  $g \in [0, d]$  and  $\mathfrak{h} \in \mathbb{U}_{g, g}$ , notice that  $\mathfrak{h} \preceq (g_1^\circ, g_2^\bullet, g_2)$  and  $k_{\mathfrak{h}} \mid k_{(g_1^\circ, g_2^\bullet, g_2)}$  by a direct computation. The above discussion thus shows that  $\mathbb{S}$  is a  $p'$ -valenced scheme if and only if  $p \nmid k_{\mathfrak{h}}$  for any  $g \in [0, d]$  and  $\mathfrak{h} \in \mathbb{U}_{g, g}$ . The desired corollary thus follows from Lemma 4.16 and Theorem 5.21.  $\square$

**Example 5.24.** Assume that  $n = \ell_1 = 2$ ,  $\ell_2 = m_1 = 3$ ,  $m_2 = 4$ . Notice that  $d = 8$ . Corollary 5.19 implies that the  $\mathbb{F}$ -subalgebra  $E_7^* \mathbb{T} E_7^*$  of  $\mathbb{T}$  is a semisimple  $\mathbb{F}$ -algebra if and only if  $p \neq 2$ . If  $p \neq 2$ , Theorem 5.17 thus implies that  $E_7^* \mathbb{T} E_7^* \cong 6M_1(\mathbb{F})$  as  $\mathbb{F}$ -algebras. If  $p = 2$ , Theorem 5.17 thus implies that  $E_7^* \mathbb{T} E_7^* / \text{Rad}(E_7^* \mathbb{T} E_7^*) \cong M_1(\mathbb{F})$  as  $\mathbb{F}$ -algebras. By Theorem 5.21,  $\mathbb{T}$  is a semisimple  $\mathbb{F}$ -algebra if and only if  $p \notin [2, 3]$ .

6. ALGEBRAIC STRUCTURE OF  $\mathbb{T}$ : JACOBSON RADICAL

In this section, we determine  $\text{Rad}(\mathbb{T})$  and its nilpotent index. For this purpose, we recall Notations 3.9, 3.10, 3.17, 3.18, 3.21, 4.1, 5.3 and start with another notation.

**Notation 6.1.** Assume that  $g, h \in [0, d]$  and  $\mathbf{i} \in \mathbb{U}_{g,h}$ . Then  $\mathbb{V}_{g,h,\mathbf{i}}$  is defined to be  $\{a : a \in \mathfrak{g}_1 \Delta \mathfrak{h}_1, p \mid m_a - 1\} \cup \{a : a \in \mathfrak{g}_2 \Delta \mathfrak{h}_2, p \mid (\ell_a - 1)m_a\} \cup \mathbb{U}_i$ . As  $\mathbb{U}_{g,h} = \mathbb{U}_{h,g}$ , notice that  $\mathbb{V}_{g,h,\mathbf{i}} = \mathbb{V}_{h,g,\mathbf{i}}$ . Moreover, set  $\mathbb{I} = \langle \{B_{a,b,\mathbf{c}} : a, b \in [0, d], \mathbf{c} \in \mathbb{U}_{a,b}, \mathbb{V}_{a,b,\mathbf{c}} \neq \emptyset\} \rangle$ . By Lemma 5.1, it is clear that  $\mathbb{I} = \langle \{B_{a,b,\mathbf{c}} : a, b \in [0, d], \mathbf{c} \in \mathbb{U}_{a,b}, p \mid k_{[a,b,a]} k_{[b,a,b]} k_{\mathbf{c}}\} \rangle$ .

**Lemma 6.2.** Assume that  $g, h, i \in [0, d]$ ,  $\mathbf{j} \in \mathbb{U}_{g,h}$ , and  $\mathbf{k} \in \mathbb{U}_{h,i}$ . If  $B_{g,h,\mathbf{j}} B_{h,i,\mathbf{k}} \neq O$ , then  $\mathbb{V}_{g,h,\mathbf{j}} \cup \mathbb{V}_{h,i,\mathbf{k}} \subseteq \mathbb{V}_{g,i,(g,h,i,\mathbf{j},\mathbf{k})}$ . In particular,  $\mathbb{I}$  is a two-sided ideal of  $\mathbb{T}$ .

*Proof.* Assume that  $\mathbb{V}_{g,h,\mathbf{j}} \neq \emptyset$  and  $\mathbb{V}_{h,i,\mathbf{k}} \neq \emptyset$ . Pick  $q \in \mathbb{V}_{g,h,\mathbf{j}}$ . If  $q \in \mathfrak{g}_1 \setminus (\mathfrak{h}_1 \cup \mathfrak{i}_1)$ , then  $p \mid m_q - 1$  and  $q \in \mathbb{V}_{g,i,(g,h,i,\mathbf{j},\mathbf{k})}$ . If  $q \in (\mathfrak{g}_1 \cap \mathfrak{i}_1) \setminus \mathfrak{h}_1$ , then  $q \in (\mathfrak{g}_1 \cap \mathfrak{i}_1)^\circ \setminus \mathfrak{h}_1$ ,  $p \mid m_q - 1$ ,  $q \in \mathbb{V}_{g,i,(g,h,i,\mathbf{j},\mathbf{k})}$ . If  $q \in \mathfrak{h}_1 \setminus (\mathfrak{g}_1 \cup \mathfrak{i}_1)$ , then  $p \mid m_q - 1$ ,  $p \mid k_{[g,h,i]}$ ,  $B_{g,h,\mathbf{j}} B_{h,i,\mathbf{k}} = O$  by Theorem 3.32. If  $q \in (\mathfrak{h}_1 \cap \mathfrak{i}_1) \setminus \mathfrak{g}_1$ , then  $p \mid m_q - 1$  and  $q \in \mathbb{V}_{g,i,(g,h,i,\mathbf{j},\mathbf{k})}$ . If  $q \in \mathfrak{g}_2 \setminus (\mathfrak{h}_2 \cup \mathfrak{i}_2)$ , then  $p \mid (\ell_q - 1)m_q$  and  $q \in \mathbb{V}_{g,i,(g,h,i,\mathbf{j},\mathbf{k})}$ . If  $q \in (\mathfrak{g}_2 \cap \mathfrak{i}_2)^\bullet \setminus \mathfrak{h}_2$ , then  $p \mid (\ell_q - 1)m_q$  and  $q \in \mathbb{V}_{g,i,(g,h,i,\mathbf{j},\mathbf{k})}$ . If  $q \in (\mathfrak{g}_2 \cap \mathfrak{i}_2) \setminus ((\mathfrak{g}_2 \cap \mathfrak{i}_2)^\bullet \cup \mathfrak{h}_2)$ , then  $p \mid m_q$  and  $q \in \mathbb{V}_{g,i,(g,h,i,\mathbf{j},\mathbf{k})}$ . If  $q \in \mathfrak{h}_2 \setminus (\mathfrak{g}_2 \cup \mathfrak{i}_2)$ , then  $p \mid (\ell_q - 1)m_q$ ,  $p \mid k_{[g,h,i]}$ ,  $B_{g,h,\mathbf{j}} B_{h,i,\mathbf{k}} = O$  by Theorem 3.32. If  $q \in (\mathfrak{h}_2 \cap \mathfrak{i}_2) \setminus \mathfrak{g}_2$ , then  $p \mid (\ell_q - 1)m_q$  and  $q \in \mathbb{V}_{g,i,(g,h,i,\mathbf{j},\mathbf{k})}$ .

Assume that  $\mathbf{j} = (\mathbf{j}_{(1)}, \mathbf{j}_{(2)}, \mathbf{j}_{(3)})$  and  $\mathbf{k} = (\mathbf{k}_{(1)}, \mathbf{k}_{(2)}, \mathbf{k}_{(3)})$ . If  $q \in \mathbf{j}_{(1)} \setminus \mathfrak{i}_1$ , then  $p \mid m_q - 1$ ,  $p \mid k_{\mathbf{j} \setminus \mathbf{i}}$ ,  $B_{g,h,\mathbf{j}} B_{h,i,\mathbf{k}} = O$  by Theorem 3.32. If  $q \in \mathfrak{i}_1 \cap \mathbf{j}_{(1)}$ , then  $q \in \mathfrak{g}_1 \cap \mathfrak{i}_1 \cap (\mathbf{j}_{(1)} \cup \mathbf{k}_{(1)})$ ,  $p \mid m_q - 1$ ,  $q \in \mathbb{V}_{g,i,(g,h,i,\mathbf{j},\mathbf{k})}$ . If  $q \in \mathbf{j}_{(2)} \setminus \mathfrak{i}_2$ , then  $p \mid (\ell_q - 1)m_q$ ,  $p \mid k_{\mathbf{j} \setminus \mathbf{i}}$ ,  $B_{g,h,\mathbf{j}} B_{h,i,\mathbf{k}} = O$  by Theorem 3.32. If  $q \in \mathfrak{i}_2 \cap \mathbf{j}_{(2)}$ , then  $q \in \mathfrak{g}_2 \cap \mathfrak{i}_2 \cap (\mathbf{j}_{(2)} \cup \mathbf{k}_{(2)})$ ,  $p \mid (\ell_q - 1)m_q$ ,  $q \in \mathbb{V}_{g,i,(g,h,i,\mathbf{j},\mathbf{k})}$ . If  $q \in \mathbf{j}_{(3)} \setminus (\mathfrak{i}_2 \cup \mathbf{j}_{(2)})$ , then  $p \mid m_q$ ,  $p \mid k_{\mathbf{j} \setminus \mathbf{i}}$ ,  $B_{g,h,\mathbf{j}} B_{h,i,\mathbf{k}} = O$  by Theorem 3.32. If  $q \in (\mathbf{j}_{(3)} \cap \mathbf{k}_{(3)}) \setminus \mathbf{j}_{(2)}$ , then  $p \mid m_q$ ,  $p \mid k_{\mathbf{j} \cap \mathbf{k}}$ ,  $B_{g,h,\mathbf{j}} B_{h,i,\mathbf{k}} = O$  by Theorem 3.32. If  $q \in (\mathfrak{i}_2 \cap \mathbf{j}_{(3)}) \setminus (\mathbf{j}_{(2)} \cup \mathbf{k}_{(3)})$ , then  $q \in \mathfrak{g}_2 \cap \mathfrak{i}_2 \cap ((\mathbf{j}_{(3)} \cup \mathbf{k}_{(3)}) \setminus (\mathbf{j}_{(2)} \cup \mathbf{k}_{(2)}))$ ,  $p \mid m_q$ ,  $q \in \mathbb{V}_{g,i,(g,h,i,\mathbf{j},\mathbf{k})}$ . In conclusion, notice that  $\mathbb{V}_{g,h,\mathbf{j}} \subseteq \mathbb{V}_{g,i,(g,h,i,\mathbf{j},\mathbf{k})}$  as  $B_{g,h,\mathbf{j}} B_{h,i,\mathbf{k}} \neq O$ .

As  $\mathbb{V}_{g,h,\mathbf{j}} = \mathbb{V}_{h,g,\mathbf{j}}$ ,  $\mathbb{V}_{h,i,\mathbf{k}} = \mathbb{V}_{i,h,\mathbf{k}}$ ,  $(g, h, i, \mathbf{j}, \mathbf{k}) = (i, h, g, \mathbf{k}, \mathbf{j})$  by a direct computation, the first statement is thus from the containment  $\mathbb{V}_{g,h,\mathbf{j}} \subseteq \mathbb{V}_{g,i,(g,h,i,\mathbf{j},\mathbf{k})}$  and Lemma 3.13. The desired lemma thus follows from Theorem 3.16 and the first statement.  $\square$

**Lemma 6.3.** Assume that  $\mathbb{I} \neq \{O\}$ . There are  $2 \mid \{a : a \in [1, n], p \mid (\ell_a - 1)(m_a - 1)m_a\}$  matrices in  $\mathbb{I}$  such that a product of all these matrices is not the zero matrix.

*Proof.* Set  $\mathbb{U} = \{a : a \in [1, n], p \mid (\ell_a - 1)(m_a - 1)m_a\}$ . Since  $\mathbb{I} \neq \{O\}$ , notice that  $\mathbb{U} \neq \emptyset$ . By Lemma 3.1, there is  $g \in [1, d]$  such that  $\mathfrak{g}_1 = \{a : a \in [1, n], p \mid m_a - 1\}$  and  $\mathfrak{g}_2 = \{a : a \in [1, n], p \nmid m_a - 1, p \mid (\ell_a - 1)m_a\}$ . Let  $h \in \mathbb{N}$  and  $\mathbb{U} = \{i_1, i_2, \dots, i_h\}$ . If  $j \in \mathfrak{g}_1$ , define  $\mathbf{k}_j = (\{j\}, \emptyset, \emptyset)$ . If  $j \in \mathfrak{g}_2^\bullet$ , define  $\mathbf{k}_j = (\emptyset, \{j\}, \{j\})$ . If  $j \in \mathfrak{g}_2 \setminus \mathfrak{g}_2^\bullet$ , define  $\mathbf{k}_j = (\emptyset, \emptyset, \{j\})$ . So  $\mathbf{k}_\ell \in \mathbb{U}_{g,g}$  and  $\mathbb{V}_{g,g,\mathbf{k}_\ell} \neq \emptyset$  for any  $\ell \in \{i_1, i_2, \dots, i_h\}$ . Notice that  $B_{g,g,\mathbf{k}_{i_1}}, B_{g,g,\mathbf{k}_{i_2}}, \dots, B_{g,g,\mathbf{k}_{i_h}} \in \mathbb{I}$  and  $B_{g,g,\mathbf{k}_{i_1}} B_{g,g,\mathbf{k}_{i_2}} \cdots B_{g,g,\mathbf{k}_{i_h}} \neq O$  by Lemma 4.8.

If  $m \in \mathfrak{g}_1$ , then Lemma 3.1 gives  $q \in [0, d]$ ,  $\mathfrak{q}_1 = \mathfrak{g}_1 \setminus \{m\}$ , and  $\mathfrak{q}_2 = \mathfrak{g}_2$ . Define  $r_m = q$ . If  $m \in \mathfrak{g}_2$ , then Lemma 3.1 gives  $s \in [0, d]$ ,  $\mathfrak{s}_1 = \mathfrak{g}_1$ , and  $\mathfrak{s}_2 = \mathfrak{g}_2 \setminus \{m\}$ . Define  $r_m = s$ . For any  $t \in \{i_1, i_2, \dots, i_h\}$ ,  $\mathbb{V}_{g,r_t,\mathbf{o}} = \mathbb{V}_{r_t,g,\mathbf{o}} \neq \emptyset$  and  $B_{g,r_t,\mathbf{o}}, B_{r_t,g,\mathbf{o}} \in \mathbb{I}$ . For any  $t \in \{i_1, i_2, \dots, i_h\}$ , notice that  $k_{[g,r_t,g]} = k_{\mathbf{o} \setminus g} = k_{\mathbf{o}} = 1$  and  $B_{g,r_t,\mathbf{o}} B_{r_t,g,\mathbf{o}} = B_{g,g,\mathbf{k}_t}$  by Theorem 3.32. So  $B_{g,r_{i_1},\mathbf{o}} B_{r_{i_1},g,\mathbf{o}} B_{g,r_{i_2},\mathbf{o}} B_{r_{i_2},g,\mathbf{o}} \cdots B_{g,r_{i_h},\mathbf{o}} B_{r_{i_h},g,\mathbf{o}} \neq O$ . The desired lemma follows as the choices of matrices in  $\mathbb{I}$  are clear by the above discussion.  $\square$

The following lemmas allow us to show that  $\mathbb{I}$  is a nilpotent two-sided ideal of  $\mathbb{T}$ .

**Lemma 6.4.** *Assume that  $g, h, i, j \in [0, d]$ ,  $\mathfrak{k} \in \mathbb{U}_{g,h}$ ,  $\mathfrak{l} \in \mathbb{U}_{h,i}$ , and  $\mathfrak{m} \in \mathbb{U}_{i,j}$ . Assume that  $\mathbb{V}_{g,h,\mathfrak{k}} \cap \mathbb{V}_{h,i,\mathfrak{l}} \cap \mathbb{V}_{i,j,\mathfrak{m}} \neq \emptyset$ . Then  $B_{g,h,\mathfrak{k}} B_{h,i,\mathfrak{l}} B_{i,j,\mathfrak{m}} = O$ .*

*Proof.* Assume that  $\mathfrak{k} = (\mathbb{k}_{(1)}, \mathbb{k}_{(2)}, \mathbb{k}_{(3)})$ ,  $\mathfrak{l} = (\mathbb{l}_{(1)}, \mathbb{l}_{(2)}, \mathbb{l}_{(3)})$ ,  $\mathfrak{m} = (\mathbb{m}_{(1)}, \mathbb{m}_{(2)}, \mathbb{m}_{(3)})$ . Pick  $q \in \mathbb{V}_{g,h,\mathfrak{k}} \cap \mathbb{V}_{h,i,\mathfrak{l}} \cap \mathbb{V}_{i,j,\mathfrak{m}}$ . If  $q \in \mathbb{h}_1 \setminus (\mathbb{g}_1 \cup \mathbb{i}_1)$ , then  $p \mid m_q - 1$  and  $p \mid k_{[g,h,i]}$ . If  $q \in \mathbb{k}_{(1)} \setminus \mathbb{i}_1$ , then  $p \mid m_q - 1$  and  $p \mid k_{\mathfrak{k} \setminus i}$ . If  $q \in \mathbb{i}_1 \setminus (\mathbb{h}_1 \cup \mathbb{j}_1)$ , then  $p \mid m_q - 1$  and  $p \mid k_{[h,i,j]}$ . If  $q \in \mathbb{m}_{(1)} \setminus \mathbb{h}_1$ , then  $p \mid m_q - 1$  and  $p \mid k_{\mathfrak{m} \setminus h}$ . If  $q \in \mathbb{h}_2 \setminus (\mathbb{g}_2 \cup \mathbb{i}_2)$ , then  $p \mid (\ell_q - 1)m_q$  and  $p \mid k_{[g,h,i]}$ . If  $q \in \mathbb{k}_{(2)} \setminus \mathbb{i}_2$ , then  $p \mid (\ell_q - 1)m_q$  and  $p \mid k_{\mathfrak{k} \setminus i}$ . If  $q \in \mathbb{k}_{(3)} \setminus (\mathbb{i}_2 \cup \mathbb{k}_{(2)})$ , then  $p \mid m_q$  and  $p \mid k_{\mathfrak{k} \setminus i}$ . If  $q \in \mathbb{i}_2 \setminus (\mathbb{h}_2 \cup \mathbb{j}_2)$ , then  $p \mid (\ell_q - 1)m_q$  and  $p \mid k_{[h,i,j]}$ . If  $q \in \mathbb{m}_{(2)} \setminus \mathbb{h}_2$ , then  $p \mid (\ell_q - 1)m_q$  and  $p \mid k_{\mathfrak{m} \setminus h}$ . If  $q \in \mathbb{m}_{(3)} \setminus (\mathbb{h}_2 \cup \mathbb{m}_{(2)})$ , then  $p \mid m_q$  and  $p \mid k_{\mathfrak{m} \setminus h}$ . If  $q \in \mathbb{l}_{(1)} \setminus \mathbb{g}_1$ , then  $p \mid m_q - 1$  and  $p \mid k_{\mathfrak{l} \setminus g}$ . If  $q \in \mathbb{k}_{(1)} \cap \mathbb{l}_{(1)}$ , then  $p \mid m_q - 1$  and  $p \mid k_{\mathfrak{k} \cap \mathfrak{l}}$ . If  $q \in \mathbb{l}_{(2)} \setminus \mathbb{g}_2$ , then  $p \mid (\ell_q - 1)m_q$  and  $p \mid k_{\mathfrak{l} \setminus g}$ . If  $q \in \mathbb{l}_{(3)} \setminus (\mathbb{g}_2 \cup \mathbb{l}_{(2)})$ , then  $p \mid m_q$  and  $p \mid k_{\mathfrak{l} \setminus g}$ . If  $q \in (\mathbb{k}_{(3)} \cap \mathbb{l}_{(3)}) \setminus (\mathbb{k}_{(2)} \cap \mathbb{l}_{(2)})$ , then  $p \mid m_q$  and  $p \mid k_{\mathfrak{k} \cap \mathfrak{l}}$ . If  $q \in \mathbb{k}_{(2)} \cap \mathbb{l}_{(2)}$ , then  $p \mid (\ell_q - 1)m_q$  and  $p \mid k_{\mathfrak{k} \cap \mathfrak{l}}$ . Hence  $B_{g,h,\mathfrak{k}} B_{h,i,\mathfrak{l}} B_{i,j,\mathfrak{m}} = O$  by the above discussion and Theorem 3.32. The desired lemma thus follows.  $\square$

**Lemma 6.5.** *Assume that  $g \in \mathbb{N} \setminus [1, 2]$  and  $h_1, i_1, h_2, i_2, \dots, h_g, i_g \in [0, d]$ . Assume that  $\mathbb{j}_k \in \mathbb{U}_{h_k, i_k}$  for any  $k \in [1, g]$ . If there are pairwise distinct  $\ell, m, q \in [1, g]$  such that  $\mathbb{V}_{h_\ell, i_\ell, \mathbb{j}_\ell} \cap \mathbb{V}_{h_m, i_m, \mathbb{j}_m} \cap \mathbb{V}_{h_q, i_q, \mathbb{j}_q} \neq \emptyset$ , then  $B_{h_1, i_1, \mathbb{j}_1} B_{h_2, i_2, \mathbb{j}_2} \cdots B_{h_g, i_g, \mathbb{j}_g} = O$ .*

*Proof.* Assume that  $B_{h_1, i_1, \mathbb{j}_1} B_{h_2, i_2, \mathbb{j}_2} \cdots B_{h_g, i_g, \mathbb{j}_g} \neq O$ . As  $\ell, m, q$  are pairwise distinct, there is no loss to assume that  $\ell < m < q$ . According to Equation (2.2) and Theorem 3.32, there exist  $r, s \in \mathbb{F}^\times$  such that  $B_{h_1, i_1, \mathbb{j}_1} B_{h_2, i_2, \mathbb{j}_2} \cdots B_{h_{m-1}, i_{m-1}, \mathbb{j}_{m-1}} = r B_{h_1, i_{m-1}, \mathfrak{t}}$  and  $B_{h_{m+1}, i_{m+1}, \mathbb{j}_{m+1}} B_{h_{m+2}, i_{m+2}, \mathbb{j}_{m+2}} \cdots B_{h_g, i_g, \mathbb{j}_g} = s B_{h_{m+1}, i_g, \mathfrak{u}}$  for some  $\mathfrak{t} \in \mathbb{U}_{h_1, i_{m-1}}$  and  $\mathfrak{u} \in \mathbb{U}_{h_{m+1}, i_g}$ . Notice that  $\mathbb{V}_{h_\ell, i_\ell, \mathbb{j}_\ell} \subseteq \mathbb{V}_{h_1, i_{m-1}, \mathfrak{t}}$  and  $\mathbb{V}_{h_q, i_q, \mathbb{j}_q} \subseteq \mathbb{V}_{h_{m+1}, i_g, \mathfrak{u}}$  by Lemma 6.2. So  $\mathbb{V}_{h_1, i_{m-1}, \mathfrak{t}} \cap \mathbb{V}_{h_m, i_m, \mathbb{j}_m} \cap \mathbb{V}_{h_{m+1}, i_g, \mathfrak{u}} \neq \emptyset$  as  $\mathbb{V}_{h_\ell, i_\ell, \mathbb{j}_\ell} \cap \mathbb{V}_{h_m, i_m, \mathbb{j}_m} \cap \mathbb{V}_{h_q, i_q, \mathbb{j}_q} \neq \emptyset$ . Then

$$O \neq B_{h_1, i_1, \mathbb{j}_1} B_{h_2, i_2, \mathbb{j}_2} \cdots B_{h_g, i_g, \mathbb{j}_g} = r s B_{h_1, i_{m-1}, \mathfrak{t}} B_{h_m, i_m, \mathbb{j}_m} B_{h_{m+1}, i_g, \mathfrak{u}} = O$$

by Equation (2.2) and Lemma 6.4. The above inequality is an obvious contradiction. Therefore  $B_{h_1, i_1, \mathbb{j}_1} B_{h_2, i_2, \mathbb{j}_2} \cdots B_{h_g, i_g, \mathbb{j}_g} = O$ . The desired lemma thus follows.  $\square$

The following lemma is a direct application of the Pigeonhole Principle.

**Lemma 6.6.** *Assume that  $g \in \mathbb{N}$  and  $\mathbb{U}$  is a set with cardinality  $g$ . Assume that  $\mathbb{V}_1, \mathbb{V}_2, \dots, \mathbb{V}_{2g+1}$  are nonempty subsets of  $\mathbb{U}$ . Then there exist pairwise distinct  $h, i, j \in [1, 2g+1]$  such that  $\mathbb{V}_h \cap \mathbb{V}_i \cap \mathbb{V}_j \neq \emptyset$ .*

*Proof.* Work by induction on  $g$ . If  $g = 1$ , then  $\mathbb{V}_1 \cap \mathbb{V}_2 \cap \mathbb{V}_3 = \mathbb{U} \neq \emptyset$ . The base case is thus checked. Assume that  $g > 1$  and any sequence of  $2g - 1$  nonempty subsets of a set with cardinality  $g - 1$  has a nonempty intersection of three members.

By the Pigeonhole Principle, notice that there exists  $k \in \mathbb{U}$  such that  $k \in \mathbb{V}_\ell \cap \mathbb{V}_m$  for some distinct  $\ell, m \in [1, 2g+1]$ . There is no loss to require that  $\ell < m$ . If there exists  $q \in [1, 2g+1] \setminus \{\ell, m\}$  such that  $k \in \mathbb{V}_q$ , notice that  $\mathbb{V}_\ell \cap \mathbb{V}_m \cap \mathbb{V}_q \neq \emptyset$ .

Assume further that  $k \notin \mathbb{V}_r$  for any  $r \in [1, 2g+1] \setminus \{\ell, m\}$ . Therefore  $\mathbb{U} \setminus \{k\}$  has  $2g - 1$  nonempty subsets  $\mathbb{V}_1, \mathbb{V}_2, \dots, \mathbb{V}_{\ell-1}, \mathbb{V}_{\ell+1}, \dots, \mathbb{V}_{m-1}, \mathbb{V}_{m+1}, \dots, \mathbb{V}_{2g+1}$ . By the inductive hypothesis, there are pairwise distinct  $s, t, u \in [1, 2g+1] \setminus \{\ell, m\}$  such that  $\mathbb{V}_s \cap \mathbb{V}_t \cap \mathbb{V}_u \neq \emptyset$ . The desired lemma thus follows from the above discussion.  $\square$

**Lemma 6.7.** *The product of any  $2|\{a : a \in [1, n], p \mid (\ell_a - 1)(m_a - 1)m_a\}| + 1$  matrices in  $\mathbb{I}$  is the zero matrix. In particular,  $\mathbb{I}$  is a nilpotent two-sided ideal of  $\mathbb{T}$  whose nilpotent index is  $2|\{a : a \in [1, n], p \mid (\ell_a - 1)(m_a - 1)m_a\}| + 1$  and  $\mathbb{I} \subseteq \text{Rad}(\mathbb{T})$ .*

*Proof.* Set  $\mathbb{U} = \{a : a \in [1, n], p \mid (\ell_a - 1)(m_a - 1)m_a\}$  and  $g = |\mathbb{U}|$ . If  $g = 0$ , notice that  $\text{Rad}(\mathbb{T}) = \mathbb{I} = \{O\}$  by Theorem 5.21. Assume further that  $g > 0$ . Assume that  $B_{h_k, i_k, j_k} \in \mathbb{B}_2$  and  $\mathbb{V}_{h_k, i_k, j_k} \neq \emptyset$  for any  $k \in [1, 2g + 1]$ . Notice that  $\mathbb{V}_{h_k, i_k, j_k} \subseteq \mathbb{U}$  for any  $k \in [1, 2g + 1]$ . By Lemma 6.6, there are pairwise distinct  $\ell, m, q \in [1, 2g + 1]$  such that  $\mathbb{V}_{h_\ell, i_\ell, j_\ell} \cap \mathbb{V}_{h_m, i_m, j_m} \cap \mathbb{V}_{h_q, i_q, j_q} \neq \emptyset$ . So  $B_{h_1, i_1, j_1} B_{h_2, i_2, j_2} \cdots B_{h_{2g+1}, i_{2g+1}, j_{2g+1}} = O$  by Lemma 6.5. The first statement thus follows from the definition of  $\mathbb{I}$ . The desired lemma thus follows from Lemma 6.3 and the first statement.  $\square$

We are now ready to list the main result of this section and an additional corollary.

**Theorem 6.8.** *Assume that  $M \in \text{Rad}(\mathbb{T})$ . Then  $M \in \mathbb{I}$ . In particular,  $\text{Rad}(\mathbb{T}) = \mathbb{I}$  and the nilpotent index of  $\text{Rad}(\mathbb{T})$  is  $2|\{a : a \in [1, n], p \mid (\ell_a - 1)(m_a - 1)m_a\}| + 1$ .*

*Proof.* Assume that  $M \in \text{Rad}(\mathbb{T}) \setminus \mathbb{I}$ . By Equation (2.3), there are  $g, h \in [0, d]$  such that  $E_g^* M E_h^* \in \text{Rad}(\mathbb{T}) \setminus \mathbb{I}$ . As  $\mathbb{I} \subseteq \text{Rad}(\mathbb{T})$  and Theorem 3.16 holds, there is no loss to assume that  $\text{Supp}_{\mathbb{B}_2}(E_g^* M E_h^*) \cap \mathbb{I} = \emptyset$ . By combining Theorem 3.16, Equation (2.2), and Notation 6.1, there exist  $i \in \mathbb{N}$  and pairwise distinct  $j_1, j_2, \dots, j_i \in \mathbb{U}_{g, h}$  such that  $\text{Supp}_{\mathbb{B}_2}(E_g^* M E_h^*) = \{B_{g, h, j_1}, B_{g, h, j_2}, \dots, B_{g, h, j_i}\}$  and  $p \nmid k_{[g, h, g]} k_{[h, g, h]} k_{j_k}$  for any  $k \in [1, i]$ . Hence  $B_{g, h, j_1}, B_{g, h, j_2}, \dots, B_{g, h, j_i}$  are pairwise distinct by Theorem 3.16.

The combination of Equation (2.2), Theorem 5.17, and Lemma 2.1 thus implies that  $E_g^* M E_h^* B_{h, g, \mathfrak{o}} \in \text{Rad}(\mathbb{T}) \cap E_g^* \mathbb{T} E_g^* \subseteq \text{Rad}(E_g^* \mathbb{T} E_g^*)$ . As  $j_1, j_2, \dots, j_i$  are pairwise distinct, notice that  $(g, h, g, j_1, \mathfrak{o}), (g, h, g, j_2, \mathfrak{o}), \dots, (g, h, g, j_i, \mathfrak{o})$  are pairwise distinct by a direct computation. So  $B_{g, g, (g, h, g, j_1, \mathfrak{o})}, B_{g, g, (g, h, g, j_2, \mathfrak{o})}, \dots, B_{g, g, (g, h, g, j_i, \mathfrak{o})}$  are pairwise distinct and  $\text{Supp}_{\mathbb{B}_2}(E_g^* M E_h^* B_{h, g, \mathfrak{o}}) \subseteq \{B_{g, g, (g, h, g, j_1, \mathfrak{o})}, B_{g, g, (g, h, g, j_2, \mathfrak{o})}, \dots, B_{g, g, (g, h, g, j_i, \mathfrak{o})}\}$  by Theorems 3.16 and 3.32. Lemmas 3.23 and 3.22 imply that  $k_{(g, h, g, j_k, \mathfrak{o})} = k_{[h, g, h]} k_{j_k}$  for any  $k \in [1, i]$ . In particular,  $p \nmid k_{(g, h, g, j_k, \mathfrak{o})}$  for any  $k \in [1, i]$ . This thus forces that  $E_g^* M E_h^* B_{h, g, \mathfrak{o}} = O$  by Theorems 3.16 and 5.17. According to Theorem 3.32, notice that  $c_{g, g, (g, h, g, j_1, \mathfrak{o})}(E_g^* M E_h^* B_{h, g, \mathfrak{o}}) = c_{g, h, j_1}(E_g^* M E_h^*) \overline{k_{[g, h, g]}} \in \mathbb{F}^\times$ . So  $E_g^* M E_h^* B_{h, g, \mathfrak{o}} \neq O$  by Theorem 3.16. This contradiction  $O = E_g^* M E_h^* B_{h, g, \mathfrak{o}} \neq O$  thus can imply that  $M \in \mathbb{I}$ . As  $M$  is chosen from  $\text{Rad}(\mathbb{T})$  arbitrarily and Lemma 6.7 holds, notice that  $\text{Rad}(\mathbb{T}) = \mathbb{I}$ . The desired theorem thus follows from Lemma 6.7.  $\square$

**Corollary 6.9.**  *$\text{Rad}(Z(\mathbb{T}))$  has an  $\mathbb{F}$ -basis  $\{C_{\mathfrak{a}} : \exists b \in [0, d], \mathfrak{a} \in \mathbb{U}_{b, b}, p \mid k_{\mathfrak{a}}\}$ .*

*Proof.* Notice that  $\{C_{\mathfrak{a}} : \exists b \in [0, d], \mathfrak{a} \in \mathbb{U}_{b, b}, p \mid k_{\mathfrak{a}}\} \subseteq \text{Rad}(Z(\mathbb{T}))$  by Lemma 4.16. For any  $g \in [0, d]$ ,  $\mathfrak{h} \in \mathbb{U}_{g, g}$ , and  $p \nmid k_{\mathfrak{h}}$ ,  $\text{Supp}_{\mathbb{B}_2}(C_{\mathfrak{h}}) \subseteq \{B_{a, a, \mathfrak{b}} : a \in [0, d], \mathfrak{b} \in \mathbb{U}_{a, a}, p \nmid k_{\mathfrak{b}}\}$  by Lemma 4.5. As  $\text{Rad}(Z(\mathbb{T})) = Z(\mathbb{T}) \cap \text{Rad}(\mathbb{T})$ , Theorems 6.8 and 3.16 yield that any nonzero  $\mathbb{F}$ -linear combination of the matrices in  $\{C_{\mathfrak{a}} : \exists b \in [0, d], \mathfrak{a} \in \mathbb{U}_{b, b}, p \nmid k_{\mathfrak{a}}\}$  is not contained in  $\text{Rad}(Z(\mathbb{T}))$ . The desired corollary is thus from Theorem 4.15.  $\square$

We are now ready to end this section by listing an example of Theorem 6.8.

**Example 6.10.** *Assume that  $n = \ell_1 = 2$ ,  $\ell_2 = m_1 = 3$ , and  $m_2 = 4$ . If  $p \in [2, 3]$ , Theorem 6.8 thus implies that the nilpotent index of  $\text{Rad}(\mathbb{T})$  equals five. If  $p \notin [2, 3]$ , Example 5.24 and Theorem 6.8 implies that the nilpotent index of  $\text{Rad}(\mathbb{T})$  equals one.*

7. ALGEBRAIC STRUCTURE OF  $\mathbb{T}$ : QUOTIENT  $\mathbb{F}$ -ALGEBRA

In this section, we present an  $\mathbb{F}$ -basis for  $\mathbb{T}/\text{Rad}(\mathbb{T})$  and determine the structure constants of this  $\mathbb{F}$ -basis in  $\mathbb{T}/\text{Rad}(\mathbb{T})$ . For this purpose, we recall Notations 3.9, 3.10, 3.17, 3.18, 3.21, 4.13, 5.6, 5.7. By Notation 6.1 and Theorem 6.8, we recall that  $\text{Rad}(\mathbb{T}) = \langle \{B_{a,b,c} : a, b \in [0, d], \mathbf{c} \in \mathbb{U}_{a,b}, p \nmid k_{[a,b,a]}k_{[b,a,b]}k_{\mathbf{c}}\} \rangle$ . We list three lemmas.

**Lemma 7.1.** *Assume that  $g, h \in [0, d]$  and  $\mathbf{i} \in \mathbb{U}_{g,h}$ . Assume that  $p \nmid k_{[g,h,g]}k_{[h,g,h]}$ . Then  $\overline{k_{[g,h,g]}}D_{g,h,\mathbf{i}}^T = \overline{k_{[h,g,h]}}D_{h,g,\mathbf{i}}$ .*

*Proof.* As  $(g, h; \mathbf{i}) = (h, g; \mathbf{i})$ ,  $n_{g,h,\mathbf{i}} = n_{h,g,\mathbf{i}}$ , and  $\mathbb{U}_{g,h,\mathbf{i},j} = \mathbb{U}_{h,g,\mathbf{i},j}$  for any  $j \in [0, n_{g,h,\mathbf{i}}]$ , the desired lemma thus follows from Lemma 3.13 and a direct computation.  $\square$

**Lemma 7.2.** *Assume that  $g, h, i, j \in [0, d]$ ,  $\mathbf{k} \in \mathbb{U}_{g,h}$ ,  $\mathbf{l} \in \mathbb{U}_{i,j}$ . If  $p \nmid k_{[g,h,g]}k_{[i,j,i]}k_{\mathbf{k}}k_{\mathbf{l}}$ , then  $D_{g,h,\mathbf{k}} = D_{i,j,\mathbf{l}}$  if and only if  $g = i$ ,  $h = j$ , and  $\mathbf{k} = \mathbf{l}$ .*

*Proof.* One direction is obvious. Assume that  $D_{g,h,\mathbf{k}} = D_{i,j,\mathbf{l}}$ . Then  $O \notin \{D_{g,h,\mathbf{k}}, D_{i,j,\mathbf{l}}\}$  as  $p \nmid k_{\mathbf{k}}k_{\mathbf{l}}$ . Notice that  $g = i$  and  $h = j$  by Equation (2.2). Notice that  $\mathbb{U}_{g,h,\mathbf{k},0} = \{\mathbf{k}\}$ ,  $\mathbb{U}_{g,h,\mathbf{l},0} = \{\mathbf{l}\}$ ,  $B_{g,h,\mathbf{k}} \in \text{Supp}_{\mathbb{B}_2}(D_{g,h,\mathbf{l}})$ ,  $B_{g,h,\mathbf{l}} \in \text{Supp}_{\mathbb{B}_2}(D_{g,h,\mathbf{k}})$ , and  $\mathbf{k} \preceq \mathbf{l} \preceq \mathbf{k}$  by the definitions of  $D_{g,h,\mathbf{k}}$ ,  $D_{i,j,\mathbf{l}}$  and Theorem 3.16. The desired lemma thus follows.  $\square$

**Lemma 7.3.**  *$\mathbb{T}$  has an  $\mathbb{F}$ -linearly independent subset*

$$\{D_{a,b,\mathbf{c}} : a, b \in [0, d], \mathbf{c} \in \mathbb{U}_{a,b}, p \nmid k_{[a,b,a]}k_{[b,a,b]}k_{\mathbf{c}}\}.$$

*Proof.* Let  $\mathbb{U}$  be the above displayed set. Notice that  $D_{0,0,0} \in \mathbb{U}$ . Let  $L$  be a nonzero  $\mathbb{F}$ -linear combination of the matrices in  $\mathbb{U}$ . Assume that  $L = O$ . If  $M \in \mathbb{U}$ , let  $c_M$  be the coefficient of  $M$  in  $L$ . So there is  $N \in \mathbb{U}$  such that  $c_N \in \mathbb{F}^\times$ . By Equation (2.2),  $N = E_g^* N E_h^*$  for some  $g, h \in [0, d]$ . So  $\mathbb{V} = \{A : A \in \mathbb{U}, c_A \in \mathbb{F}^\times, A = E_g^* A E_h^*\} \neq \emptyset$ . By Lemma 7.2, there exist  $i \in \mathbb{N}$  and pairwise distinct  $j_1, j_2, \dots, j_i \in \mathbb{U}_{g,h}$  such that  $\mathbb{V} = \{D_{g,h,j_1}, D_{g,h,j_2}, \dots, D_{g,h,j_i}\}$ . We thus can distinguish the cases  $i = 1$  and  $i > 1$ .

If  $i = 1$ , there is  $c \in \mathbb{F}^\times$  such that  $E_g^* L E_h^* = c D_{g,h,j_1} = O$ . This is a contradiction as  $p \nmid k_{j_1}$  and  $D_{g,h,j_1} \neq O$ . Assume further that  $i > 1$ . There is no loss to assume that  $j_1$  is minimal in  $\{j_1, j_2, \dots, j_i\}$  with respect to the partial order  $\preceq$ . As  $E_g^* L E_h^* = O$ ,  $D_{g,h,j_1}$  is also an  $\mathbb{F}$ -linear combination of the matrices in  $\{D_{g,h,j_2}, D_{g,h,j_3}, \dots, D_{g,h,j_i}\}$ . As  $\mathbb{U}_{g,h,j_1,0} = \{j_1\}$  and Theorem 3.16 holds, this  $\mathbb{F}$ -linear combination of  $D_{g,h,j_1}$  shows that  $j \preceq j_1$  for some  $j \in \{j_2, j_3, \dots, j_i\}$ . Hence  $j_1 \in \{j_2, j_3, \dots, j_i\}$  by the choice of  $j_1$ . It is also a contradiction. The desired lemma is thus from the inequality  $L \neq O$ .  $\square$

We are now ready to introduce the first main result of this section.

**Theorem 7.4.**  *$\mathbb{T}/\text{Rad}(\mathbb{T})$  has an  $\mathbb{F}$ -basis*

$$\{D_{a,b,\mathbf{c}} + \text{Rad}(\mathbb{T}) : a, b \in [0, d], \mathbf{c} \in \mathbb{U}_{a,b}, p \nmid k_{[a,b,a]}k_{[b,a,b]}k_{\mathbf{c}}\}.$$

*Proof.* Let  $\mathbb{U}$  be the union of  $\{D_{a,b,\mathbf{c}} : a, b \in [0, d], \mathbf{c} \in \mathbb{U}_{a,b}, p \nmid k_{[a,b,a]}k_{[b,a,b]}k_{\mathbf{c}}\}$  and  $\{B_{a,b,\mathbf{c}} : a, b \in [0, d], \mathbf{c} \in \mathbb{U}_{a,b}, p \mid k_{[a,b,a]}k_{[b,a,b]}k_{\mathbf{c}}\}$ . According to Theorem 3.16 and Lemma 7.2, notice that  $|\mathbb{U}| = |\{(a, b, \mathbf{c}) : a, b \in [0, d], \mathbf{c} \in \mathbb{U}_{a,b}\}|$ . Therefore  $\mathbb{T}$  has an  $\mathbb{F}$ -basis  $\mathbb{U}$  by Lemma 7.3 and Theorem 3.16. The desired theorem thus follows.  $\square$

For the remaining main results of this section, the following lemmas are necessary.

**Lemma 7.5.** *Assume that  $g, h, i \in [0, d]$ . Assume that  $\mathbf{j} = (\mathbf{j}_{(1)}, \mathbf{j}_{(2)}, \mathbf{j}_{(3)}) \in \mathbb{U}_{g,h}$  and  $\mathbf{k} = (\mathbf{k}_{(1)}, \mathbf{k}_{(2)}, \mathbf{k}_{(3)}) \in \mathbb{U}_{h,i}$ . If the containments  $((\mathbf{h}_1 \cap \mathbf{i}_1)^\circ \setminus \mathbf{g}_1) \cup (\mathbf{i}_1 \cap \mathbf{j}_{(1)}) \subseteq \mathbf{k}_{(1)}$ ,  $(\mathbf{k}_{(3)}^\bullet \setminus \mathbf{g}_2) \cup (\mathbf{j}_{(2)} \cap \mathbf{k}_{(3)}) \subseteq \mathbf{k}_{(2)}$ ,  $((\mathbf{h}_2 \cap \mathbf{i}_2) \setminus \mathbf{g}_2) \cup (\mathbf{i}_2 \cap \mathbf{j}_{(3)}) \subseteq \mathbf{k}_{(3)}$  hold, then  $n_{g,i,(g,h,i,j),\mathbf{k}} = n_{h,i,\mathbf{k}}$ .*

*Proof.* As  $((\mathbf{h}_1 \cap \mathbf{i}_1)^\circ \setminus \mathbf{g}_1) \cup (\mathbf{i}_1 \cap \mathbf{j}_{(1)}) \subseteq \mathbf{k}_{(1)}$ ,  $(\mathbf{h}_1 \cap \mathbf{i}_1)^\circ \setminus \mathbf{k}_{(1)} = ((\mathbf{g}_1 \cap \mathbf{i}_1)^\circ \cap \mathbf{h}_1) \setminus \mathbf{k}_{(1)}$  and  $(\mathbf{g}_1 \cap \mathbf{i}_1)^\circ \setminus (((\mathbf{g}_1 \cap \mathbf{i}_1)^\circ \setminus \mathbf{h}_1) \cup (\mathbf{g}_1 \cap \mathbf{i}_1 \cap (\mathbf{j}_{(1)} \cup \mathbf{k}_{(1)}))) = ((\mathbf{g}_1 \cap \mathbf{i}_1)^\circ \cap \mathbf{h}_1) \setminus \mathbf{k}_{(1)}$ .

As  $(\mathbf{k}_{(3)}^\bullet \setminus \mathbf{g}_2) \cup (\mathbf{j}_{(2)} \cap \mathbf{k}_{(3)}) \subseteq \mathbf{k}_{(2)}$  and  $\mathbf{i}_2 \cap \mathbf{j}_{(3)} \subseteq \mathbf{k}_{(3)}$ ,  $\mathbf{k}_{(3)}^\bullet \setminus \mathbf{k}_{(2)} = (\mathbf{g}_2 \cap \mathbf{k}_{(3)}^\bullet) \setminus \mathbf{k}_{(2)}$ ,  $((\mathbf{g}_2 \cap \mathbf{i}_2)^\bullet \setminus \mathbf{h}_2) \cup (\mathbf{g}_2 \cap \mathbf{i}_2 \cap (\mathbf{j}_{(3)}^\bullet \cup \mathbf{k}_{(3)}^\bullet)) = ((\mathbf{g}_2 \cap \mathbf{i}_2)^\bullet \setminus \mathbf{h}_2) \cup (\mathbf{g}_2 \cap \mathbf{k}_{(3)}^\bullet)$ , and  $((\mathbf{g}_2 \cap \mathbf{i}_2)^\bullet \setminus \mathbf{h}_2) \cup (\mathbf{g}_2 \cap \mathbf{i}_2 \cap (\mathbf{j}_{(2)} \cup \mathbf{k}_{(2)})) = ((\mathbf{g}_2 \cap \mathbf{i}_2)^\bullet \setminus \mathbf{h}_2) \cup (\mathbf{g}_2 \cap \mathbf{k}_{(2)})$ .

As  $((\mathbf{h}_2 \cap \mathbf{i}_2) \setminus \mathbf{g}_2) \cup (\mathbf{i}_2 \cap \mathbf{j}_{(3)}) \subseteq \mathbf{k}_{(3)}$ ,  $(\mathbf{h}_2 \cap \mathbf{i}_2) \setminus \mathbf{k}_{(3)} = (\mathbf{g}_2 \cap \mathbf{h}_2 \cap \mathbf{i}_2) \setminus \mathbf{k}_{(3)}$  and  $(\mathbf{g}_2 \cap \mathbf{i}_2) \setminus (((\mathbf{g}_2 \cap \mathbf{i}_2) \setminus \mathbf{h}_2) \cup (\mathbf{g}_2 \cap \mathbf{i}_2 \cap (\mathbf{j}_{(3)} \cup \mathbf{k}_{(3)}))) = (\mathbf{g}_2 \cap \mathbf{h}_2 \cap \mathbf{i}_2) \setminus \mathbf{k}_{(3)}$ . The desired lemma thus follows from the definitions of  $n_{g,i,(g,h,i,j),\mathbf{k}}$  and  $n_{h,i,\mathbf{k}}$ .  $\square$

**Lemma 7.6.** *Assume that  $g, h, i \in [0, d]$ ,  $\mathbf{j} \in \mathbb{U}_{g,h}$ ,  $\mathbf{k} \in \mathbb{U}_{h,i}$ ,  $p \nmid k_{[h,i,h]}k_jk_\mathbf{k}$ . Then  $p \nmid k_{(g,h,i,j),\mathbf{k}}$ .*

*Proof.* Set  $\mathbf{l} = (\mathbf{l}_{(1)}, \mathbf{l}_{(2)}, \mathbf{l}_{(3)})$ ,  $\mathbf{l}_{(1)} = \mathbf{g}_1 \cap \mathbf{i}_1 \cap (\mathbf{j}_{(1)} \cup \mathbf{k}_{(1)})$ ,  $\mathbf{l}_{(2)} = \mathbf{g}_2 \cap \mathbf{i}_2 \cap (\mathbf{j}_{(2)} \cup \mathbf{k}_{(2)})$ , and  $\mathbf{l}_{(3)} = \mathbf{g}_2 \cap \mathbf{i}_2 \cap (\mathbf{j}_{(3)} \cup \mathbf{k}_{(3)})$ . As  $p \nmid k_jk_\mathbf{k}$ , notice that  $p \nmid (m_q - 1)(\ell_r - 1)m_r m_s$  for any  $q \in \mathbf{l}_{(1)}$ ,  $r \in \mathbf{l}_{(2)}$ , and  $s \in \mathbf{l}_{(3)} \setminus \mathbf{l}_{(2)}$ . Therefore  $p \nmid k_\mathbf{l}$ . As  $p \nmid k_{[h,i,h]}$ , notice that  $p \nmid (m_q - 1)(\ell_r - 1)m_r$  for any  $q \in (\mathbf{g}_1 \cap \mathbf{i}_1) \setminus \mathbf{h}_1$  and  $r \in (\mathbf{g}_2 \cap \mathbf{i}_2) \setminus \mathbf{h}_2$ . So  $p \nmid k_{(g,h,i)}$ . Since  $k_{(g,h,i,j),\mathbf{k}} = k_{(g,h,i)}k_\mathbf{l}$  by Lemmas 3.23 and 3.22, the desired lemma follows.  $\square$

**Lemma 7.7.** *Assume that  $g, h, i \in [0, d]$ . Assume that  $\mathbf{j} = (\mathbf{j}_{(1)}, \mathbf{j}_{(2)}, \mathbf{j}_{(3)}) \in \mathbb{U}_{g,h}$  and  $\mathbf{k} = (\mathbf{k}_{(1)}, \mathbf{k}_{(2)}, \mathbf{k}_{(3)}) \in \mathbb{U}_{h,i}$ . Assume that  $\ell \in [0, n_{h,i,\mathbf{k}}]$ ,  $\mathbf{m} \in \mathbb{U}_{h,i,\mathbf{k},\ell}$ , and  $p \nmid k_{[h,i,h]}k_j$ . Assume that  $((\mathbf{h}_1 \cap \mathbf{i}_1)^\circ \setminus \mathbf{g}_1) \cup (\mathbf{i}_1 \cap \mathbf{j}_{(1)}) \subseteq \mathbf{k}_{(1)}$ ,  $(\mathbf{k}_{(3)}^\bullet \setminus \mathbf{g}_2) \cup (\mathbf{j}_{(2)} \cap \mathbf{k}_{(3)}) \subseteq \mathbf{k}_{(2)}$ , and  $((\mathbf{h}_2 \cap \mathbf{i}_2) \setminus \mathbf{g}_2) \cup (\mathbf{i}_2 \cap \mathbf{j}_{(3)}) \subseteq \mathbf{k}_{(3)}$ . Then  $(g, h, i, \mathbf{j}, \mathbf{m}) \in \mathbb{U}_{g,i,(g,h,i,j),\mathbf{k},\ell}$ .*

*Proof.* As  $\mathbf{m} \in \mathbb{U}_{h,i,\mathbf{k},\ell}$ ,  $\mathbf{k} \preceq \mathbf{m} \preceq (h, i; \mathbf{k})$ ,  $|\mathbf{m}| - |\mathbf{k}| = \ell$ ,  $p \nmid k_\mathbf{m}$ . By a direct computation,  $(g, h, i, \mathbf{j}, \mathbf{k}) \preceq (g, h, i, \mathbf{j}, \mathbf{m}) \preceq (g, i; (g, h, i, \mathbf{j}, \mathbf{k}))$ . By Lemma 7.5,  $\mathbb{U}_{g,i,(g,h,i,j),\mathbf{k},\ell}$  is defined. Then  $|(g, h, i, \mathbf{j}, \mathbf{m})| - |(g, h, i, \mathbf{j}, \mathbf{k})| = |\mathbf{m}| - |\mathbf{k}| = \ell$  by a direct computation and the containments  $((\mathbf{h}_1 \cap \mathbf{i}_1)^\circ \setminus \mathbf{g}_1) \cup (\mathbf{i}_1 \cap \mathbf{j}_{(1)}) \subseteq \mathbf{k}_{(1)}$ ,  $(\mathbf{k}_{(3)}^\bullet \setminus \mathbf{g}_2) \cup (\mathbf{j}_{(2)} \cap \mathbf{k}_{(3)}) \subseteq \mathbf{k}_{(2)}$ , and  $((\mathbf{h}_2 \cap \mathbf{i}_2) \setminus \mathbf{g}_2) \cup (\mathbf{i}_2 \cap \mathbf{j}_{(3)}) \subseteq \mathbf{k}_{(3)}$ . As  $p \nmid k_{[h,i,h]}k_j$ , notice that  $p \nmid k_{[h,i,h]}k_jk_\mathbf{m}$  and  $p \nmid k_{(g,h,i,j),\mathbf{m}}$  by Lemma 7.6. The desired lemma is thus from the above discussion.  $\square$

**Lemma 7.8.** *Assume that  $g, h, i \in [0, d]$ . Assume that  $\mathbf{j} = (\mathbf{j}_{(1)}, \mathbf{j}_{(2)}, \mathbf{j}_{(3)}) \in \mathbb{U}_{g,h}$  and  $\mathbf{k} = (\mathbf{k}_{(1)}, \mathbf{k}_{(2)}, \mathbf{k}_{(3)}) \in \mathbb{U}_{h,i}$ . Assume that  $\ell \in [0, n_{h,i,\mathbf{k}}]$ ,  $\mathbf{m}, \mathbf{q} \in \mathbb{U}_{h,i,\mathbf{k},\ell}$ , and  $p \nmid k_{[h,i,h]}k_j$ . Assume that  $((\mathbf{h}_1 \cap \mathbf{i}_1)^\circ \setminus \mathbf{g}_1) \cup (\mathbf{i}_1 \cap \mathbf{j}_{(1)}) \subseteq \mathbf{k}_{(1)}$ ,  $(\mathbf{k}_{(3)}^\bullet \setminus \mathbf{g}_2) \cup (\mathbf{j}_{(2)} \cap \mathbf{k}_{(3)}) \subseteq \mathbf{k}_{(2)}$ , and  $((\mathbf{h}_2 \cap \mathbf{i}_2) \setminus \mathbf{g}_2) \cup (\mathbf{i}_2 \cap \mathbf{j}_{(3)}) \subseteq \mathbf{k}_{(3)}$ . Then  $\mathbf{m} = \mathbf{q}$  if and only if  $(g, h, i, \mathbf{j}, \mathbf{m}) = (g, h, i, \mathbf{j}, \mathbf{q})$ . Moreover, the map that sends  $\mathbf{r}$  to  $(g, h, i, \mathbf{j}, \mathbf{r})$  is injective from  $\mathbb{U}_{h,i,\mathbf{k},\ell}$  to  $\mathbb{U}_{g,i,(g,h,i,j),\mathbf{k},\ell}$ .*

*Proof.* Assume that  $\mathbf{m} = (\mathbf{m}_{(1)}, \mathbf{m}_{(2)}, \mathbf{m}_{(3)})$  and  $\mathbf{q} = (\mathbf{q}_{(1)}, \mathbf{q}_{(2)}, \mathbf{q}_{(3)})$ . Assume that  $(g, h, i, \mathbf{j}, \mathbf{m}) = (g, h, i, \mathbf{j}, \mathbf{q})$ . Notice that  $\mathbf{g}_1 \cap \mathbf{i}_1 \cap (\mathbf{j}_{(1)} \cup \mathbf{m}_{(1)}) = \mathbf{g}_1 \cap \mathbf{i}_1 \cap (\mathbf{j}_{(1)} \cup \mathbf{q}_{(1)})$ ,  $\mathbf{g}_2 \cap \mathbf{i}_2 \cap (\mathbf{j}_{(2)} \cup \mathbf{m}_{(2)}) = \mathbf{g}_2 \cap \mathbf{i}_2 \cap (\mathbf{j}_{(2)} \cup \mathbf{q}_{(2)})$ , and  $\mathbf{g}_2 \cap \mathbf{i}_2 \cap (\mathbf{j}_{(3)} \cup \mathbf{m}_{(3)}) = \mathbf{g}_2 \cap \mathbf{i}_2 \cap (\mathbf{j}_{(3)} \cup \mathbf{q}_{(3)})$ .

As  $\mathbf{i}_1 \cap \mathbf{j}_{(1)} \subseteq \mathbf{k}_{(1)}$ ,  $\mathbf{j}_{(2)} \cap \mathbf{k}_{(3)} \subseteq \mathbf{k}_{(2)}$ ,  $\mathbf{i}_2 \cap \mathbf{j}_{(3)} \subseteq \mathbf{k}_{(3)}$ ,  $\mathbf{k} \preceq \mathbf{m}$ , and  $\mathbf{k} \preceq \mathbf{q}$ , notice that  $\mathbf{g}_1 \cap \mathbf{m}_{(1)} = \mathbf{g}_1 \cap \mathbf{q}_{(1)}$ ,  $\mathbf{g}_2 \cap \mathbf{m}_{(2)} = \mathbf{g}_2 \cap \mathbf{q}_{(2)}$ , and  $\mathbf{g}_2 \cap \mathbf{m}_{(3)} = \mathbf{g}_2 \cap \mathbf{q}_{(3)}$ . Then  $\mathbf{m}_{(1)} \setminus \mathbf{g}_1 = \mathbf{k}_{(1)} \setminus \mathbf{g}_1 = \mathbf{q}_{(1)} \setminus \mathbf{g}_1$ ,  $\mathbf{m}_{(2)} \setminus \mathbf{g}_2 = \mathbf{k}_{(2)} \setminus \mathbf{g}_2 = \mathbf{q}_{(2)} \setminus \mathbf{g}_2$ ,  $\mathbf{m}_{(3)} \setminus \mathbf{g}_2 = \mathbf{k}_{(3)} \setminus \mathbf{g}_2 = \mathbf{q}_{(3)} \setminus \mathbf{g}_2$  as  $(\mathbf{h}_1 \cap \mathbf{i}_1)^\circ \setminus \mathbf{g}_1 \subseteq \mathbf{k}_{(1)}$ ,  $\mathbf{k}_{(3)}^\bullet \setminus \mathbf{g}_2 \subseteq \mathbf{k}_{(2)}$ , and  $(\mathbf{h}_2 \cap \mathbf{i}_2) \setminus \mathbf{g}_2 \subseteq \mathbf{k}_{(3)}$ . Hence  $\mathbf{m} = \mathbf{q}$ . The first statement thus follows. The desired lemma thus follows from Lemma 7.7.  $\square$

**Lemma 7.9.** *Assume that  $g, h, i \in [0, d]$ . Assume that  $\mathbf{j} = (\mathbb{j}_{(1)}, \mathbb{j}_{(2)}, \mathbb{j}_{(3)}) \in \mathbb{U}_{g,h}$  and  $\mathfrak{k} = (\mathbb{k}_{(1)}, \mathbb{k}_{(2)}, \mathbb{k}_{(3)}) \in \mathbb{U}_{h,i}$ . Assume that  $\ell \in [0, n_{h,i,\mathfrak{k}}]$ ,  $p \nmid k_{[h,i,h]}k_jk_{\mathfrak{k}}$ ,  $\mathbf{m} \in \mathbb{U}_{g,i,(g,h,i,j,\mathfrak{k}),\ell}$ . Assume that  $((\mathbb{h}_1 \cap \mathbb{i}_1)^\circ \setminus \mathfrak{g}_1) \cup (\mathbb{i}_1 \cap \mathbb{j}_{(1)}) \subseteq \mathbb{k}_{(1)}$ ,  $(\mathbb{k}_{(3)}^\bullet \setminus \mathfrak{g}_2) \cup (\mathbb{j}_{(2)} \cap \mathbb{k}_{(3)}) \subseteq \mathbb{k}_{(2)}$ , and  $((\mathbb{h}_2 \cap \mathbb{i}_2) \setminus \mathfrak{g}_2) \cup (\mathbb{i}_2 \cap \mathbb{j}_{(3)}) \subseteq \mathbb{k}_{(3)}$ . Then there is  $\mathbf{q} \in \mathbb{U}_{h,i,\mathfrak{k},\ell}$  such that  $\mathbf{m} = (g, h, i, \mathbf{j}, \mathbf{q})$ . Moreover, the map that sends  $\mathbf{r}$  to  $(g, h, i, \mathbf{j}, \mathbf{r})$  is bijective from  $\mathbb{U}_{h,i,\mathfrak{k},\ell}$  to  $\mathbb{U}_{g,i,(g,h,i,j,\mathfrak{k}),\ell}$ .*

*Proof.* Assume that  $\mathbf{m} = (\mathfrak{m}_{(1)}, \mathfrak{m}_{(2)}, \mathfrak{m}_{(3)})$  and  $\mathbf{q} = (\mathfrak{q}_{(1)}, \mathfrak{q}_{(2)}, \mathfrak{q}_{(3)})$ . Assume that  $\mathfrak{q}_{(1)} = (\mathbb{k}_{(1)} \setminus \mathfrak{g}_1) \cup (\mathbb{h}_1 \cap \mathfrak{m}_{(1)})$ ,  $\mathfrak{q}_{(2)} = (\mathbb{k}_{(2)} \setminus \mathfrak{g}_2) \cup (\mathbb{h}_2 \cap \mathfrak{m}_{(2)})$ ,  $\mathfrak{q}_{(3)} = (\mathbb{k}_{(3)} \setminus \mathfrak{g}_2) \cup (\mathbb{h}_2 \cap \mathfrak{m}_{(3)})$ .

As  $\mathbf{m} \in \mathbb{U}_{g,i,(g,h,i,j,\mathfrak{k}),\ell}$  and  $\mathfrak{k} \in \mathbb{U}_{h,i}$ , notice that  $(g, h, i, \mathbf{j}, \mathfrak{k}) \preceq \mathbf{m} \preceq (g, i; (g, h, i, \mathbf{j}, \mathfrak{k}))$ ,  $|\mathbf{m}| - |(g, h, i, \mathbf{j}, \mathfrak{k})| = \ell$ ,  $p \nmid k_{\mathbf{m}}$ ,  $\mathbf{q} \in \mathbb{U}_{h,i}$ , and  $\mathbf{m} = (g, h, i, \mathbf{j}, \mathbf{q})$  by a direct computation. As  $\mathbb{i}_1 \cap \mathbb{j}_{(1)} \subseteq \mathbb{k}_{(1)}$ ,  $\mathbb{j}_{(2)} \cap \mathbb{k}_{(3)} \subseteq \mathbb{k}_{(2)}$ , and  $\mathbb{i}_2 \cap \mathbb{j}_{(3)} \subseteq \mathbb{k}_{(3)}$ , notice that  $\mathfrak{k} \preceq \mathbf{q} \preceq (h, i; \mathfrak{k})$ ,  $\mathfrak{q}_{(1)} \setminus \mathbb{k}_{(1)} = (\mathbb{h}_1 \cap \mathfrak{m}_{(1)}) \setminus \mathbb{k}_{(1)} = \mathfrak{m}_{(1)} \setminus (((\mathbb{g}_1 \cap \mathbb{i}_1)^\circ \setminus \mathbb{h}_1) \cup (\mathbb{g}_1 \cap \mathbb{i}_1 \cap (\mathbb{j}_{(1)} \cup \mathbb{k}_{(1)})))$ ,  $\mathfrak{q}_{(2)} \setminus \mathbb{k}_{(2)} = (\mathbb{h}_2 \cap \mathfrak{m}_{(2)}) \setminus \mathbb{k}_{(2)} = \mathfrak{m}_{(2)} \setminus (((\mathbb{g}_2 \cap \mathbb{i}_2)^\bullet \setminus \mathbb{h}_2) \cup (\mathbb{g}_2 \cap \mathbb{i}_2 \cap (\mathbb{j}_{(2)} \cup \mathbb{k}_{(2)})))$ , and  $\mathfrak{q}_{(3)} \setminus \mathbb{k}_{(3)} = (\mathbb{h}_2 \cap \mathfrak{m}_{(3)}) \setminus \mathbb{k}_{(3)} = \mathfrak{m}_{(3)} \setminus (((\mathbb{g}_2 \cap \mathbb{i}_2) \setminus \mathbb{h}_2) \cup (\mathbb{g}_2 \cap \mathbb{i}_2 \cap (\mathbb{j}_{(3)} \cup \mathbb{k}_{(3)})))$ . Hence  $|\mathbf{q}| - |\mathfrak{k}| = |\mathbf{m}| - |(g, h, i, \mathbf{j}, \mathfrak{k})| = \ell$ . As  $p \nmid k_{\mathfrak{k}}k_{\mathbf{m}}$ , notice that  $p \nmid (m_s - 1)(\ell_t - 1)m_t m_u$  for any  $s \in \mathfrak{q}_1$ ,  $t \in \mathfrak{q}_2$ , and  $u \in \mathfrak{q}_3 \setminus \mathfrak{q}_2$ . So  $p \nmid k_{\mathbf{q}}$ . The first statement thus follows. The desired lemma thus follows from the first statement and Lemma 7.8.  $\square$

**Lemma 7.10.** *Assume that  $g, h, i \in [0, d]$ . Assume that  $\mathbf{j} = (\mathbb{j}_{(1)}, \mathbb{j}_{(2)}, \mathbb{j}_{(3)}) \in \mathbb{U}_{g,h}$  and  $\mathfrak{k} = (\mathbb{k}_{(1)}, \mathbb{k}_{(2)}, \mathbb{k}_{(3)}) \in \mathbb{U}_{h,i}$ . Assume that  $\mathfrak{l} \preceq \mathfrak{k}$ ,  $\mathbb{i}_1 \cap \mathbb{j}_{(1)} \subseteq \mathbb{k}_{(1)}$ ,  $\mathbb{j}_{(2)} \cap \mathbb{k}_{(3)} \subseteq \mathbb{k}_{(2)}$ , and  $\mathbb{i}_2 \cap \mathbb{j}_{(3)} \subseteq \mathbb{k}_{(3)}$ . Then  $k_j = k_{\mathbb{j} \setminus \mathbb{i}_1} k_{\mathbb{j} \cap \mathbb{i}_1}$  and  $k_{(g,h,i)} k_{\mathfrak{l}} = k_{\mathbb{i} \setminus \mathbb{g}} k_{(g,h,i,j,\mathfrak{l})}$ .*

*Proof.* Assume that  $\mathfrak{l} = (\mathbb{l}_{(1)}, \mathbb{l}_{(2)}, \mathbb{l}_{(3)})$ ,  $\mathbf{m} = (\mathfrak{m}_{(1)}, \mathfrak{m}_{(2)}, \mathfrak{m}_{(3)})$ ,  $\mathfrak{m}_{(1)} = \mathbb{g}_1 \cap \mathbb{i}_1 \cap (\mathbb{j}_{(1)} \cup \mathbb{l}_{(1)})$ ,  $\mathfrak{m}_{(2)} = \mathbb{g}_2 \cap \mathbb{i}_2 \cap (\mathbb{j}_{(2)} \cup \mathbb{l}_{(2)})$ , and  $\mathfrak{m}_{(3)} = \mathbb{g}_2 \cap \mathbb{i}_2 \cap (\mathbb{j}_{(3)} \cup \mathbb{l}_{(3)})$ . As  $\mathfrak{k} \preceq \mathfrak{l}$ ,  $\mathbb{i}_1 \cap \mathbb{j}_{(1)} \subseteq \mathbb{k}_{(1)}$ ,  $\mathbb{j}_{(2)} \cap \mathbb{k}_{(3)} \subseteq \mathbb{k}_{(2)}$ ,  $\mathbb{i}_2 \cap \mathbb{j}_{(3)} \subseteq \mathbb{k}_{(3)}$ , notice that  $\mathbb{j}_{(1)} \cap \mathbb{l}_{(1)} = \mathbb{i}_1 \cap \mathbb{j}_{(1)}$ ,  $\mathbb{j}_{(2)} \cap \mathbb{l}_{(2)} = \mathbb{i}_2 \cap \mathbb{j}_{(2)}$ ,  $\mathbb{j}_{(3)} \cap \mathbb{l}_{(3)} = \mathbb{i}_2 \cap \mathbb{j}_{(3)}$ ,  $\mathbb{g}_1 \cap \mathbb{i}_1 \cap (\mathbb{j}_{(1)} \cup \mathbb{l}_{(1)}) = \mathbb{g}_1 \cap \mathbb{l}_{(1)}$ ,  $\mathbb{g}_2 \cap \mathbb{i}_2 \cap (\mathbb{j}_{(2)} \cup \mathbb{l}_{(2)}) = \mathbb{g}_2 \cap \mathbb{l}_{(2)}$ , and  $\mathbb{g}_2 \cap \mathbb{i}_2 \cap (\mathbb{j}_{(3)} \cup \mathbb{l}_{(3)}) = \mathbb{g}_2 \cap \mathbb{l}_{(3)}$ . As  $k_j = k_{\mathbb{j} \setminus \mathbb{i}_1} k_{\mathbb{j} \cap \mathbb{i}_1}$ ,  $k_{\mathfrak{l}} = k_{\mathbb{l} \setminus \mathbb{g}} k_{\mathbb{l} \cap \mathbb{g}}$ ,  $k_{(g,h,i,j,\mathfrak{l})} = k_{(g,h,i)} k_{\mathbf{m}}$  by combining Lemmas 4.5, 3.23, and 3.22, the desired lemma thus follows.  $\square$

**Lemma 7.11.** *Assume that  $g, h, i \in [0, d]$ . Then  $k_{[g,h,g]} = k_{(h,g,i)} k_{[g,h,i]}$ . In particular,  $k_{[g,h,g]} k_{[h,i,h]} = k_{(g,h,i)} k_{[g,h,i]} k_{[g,i,g]}$ .*

*Proof.* As  $\mathbb{h}_1 \setminus \mathbb{g}_1 = (\mathbb{h}_1 \setminus (\mathbb{g}_1 \setminus \mathbb{i}_1)) \cup ((\mathbb{h}_1 \cap \mathbb{i}_1) \setminus \mathbb{g}_1)$  and  $\mathbb{h}_2 \setminus \mathbb{g}_2 = (\mathbb{h}_2 \setminus (\mathbb{g}_2 \setminus \mathbb{i}_2)) \cup ((\mathbb{h}_2 \cap \mathbb{i}_2) \setminus \mathbb{g}_2)$ , the first statement follows from 3.23. The first statement gives  $k_{[h,i,h]} = k_{(g,h,i)} k_{[g,i,h]}$  and  $k_{[g,i,g]} = k_{(h,g,i)} k_{[g,i,h]}$ . The desired lemma is thus from a direct computation.  $\square$

**Lemma 7.12.** *Assume that  $g, h, i \in [0, d]$ . Assume that  $\mathbf{j} = (\mathbb{j}_{(1)}, \mathbb{j}_{(2)}, \mathbb{j}_{(3)}) \in \mathbb{U}_{g,h}$ ,  $\mathfrak{k} = (\mathbb{k}_{(1)}, \mathbb{k}_{(2)}, \mathbb{k}_{(3)}) \in \mathbb{U}_{h,i}$ ,  $p \nmid k_{[g,i,g]} k_{[h,i,h]} k_j k_{\mathfrak{k}}$ ,  $((\mathbb{h}_1 \cap \mathbb{i}_1)^\circ \setminus \mathfrak{g}_1) \cup (\mathbb{i}_1 \cap \mathbb{j}_{(1)}) \subseteq \mathbb{k}_{(1)}$ ,  $(\mathbb{k}_{(3)}^\bullet \setminus \mathfrak{g}_2) \cup (\mathbb{j}_{(2)} \cap \mathbb{k}_{(3)}) \subseteq \mathbb{k}_{(2)}$ ,  $((\mathbb{h}_2 \cap \mathbb{i}_2) \setminus \mathfrak{g}_2) \cup (\mathbb{i}_2 \cap \mathbb{j}_{(3)}) \subseteq \mathbb{k}_{(3)}$ . Then  $p \nmid k_{(g,h,i,j,\mathfrak{k})}$  and  $B_{g,h,j} D_{h,i,\mathfrak{k}} = \overline{k_{[g,h,g]}} \overline{k_j} D_{g,i,(g,h,i,j,\mathfrak{k})}$ .*

*Proof.* As  $p \nmid k_{[g,i,g]} k_{[h,i,h]} k_j k_{\mathfrak{k}}$ , notice that  $D_{h,i,\mathfrak{k}} \neq O$  and  $p \nmid k_{(g,h,i,j,\mathfrak{k})}$  by Lemma 7.6. By combining Theorem 3.32, Lemmas 7.5, 7.9, 7.10, 7.11, and a direct computation,

$$B_{g,h,j} D_{h,i,\mathfrak{k}} = \overline{k_{[g,h,g]}} \overline{k_j} \sum_{\ell=0}^{n_{g,i,(g,h,i,j,\mathfrak{k})}} \sum_{(g,h,i,j,\mathfrak{m}) \in \mathbb{U}_{g,i,(g,h,i,j,\mathfrak{k}),\ell}} (-1)^\ell \overline{k_{[g,i,g]}}^{-1} \overline{k_{(g,h,i,j,\mathfrak{m})}}^{-1} B_{g,i,(g,h,i,j,\mathfrak{m})},$$

which implies that  $B_{g,h,j} D_{h,i,\mathfrak{k}} = \overline{k_{[g,h,g]}} \overline{k_j} D_{g,i,(g,h,i,j,\mathfrak{k})}$ . The desired lemma follows.  $\square$

**Lemma 7.13.** *Assume that  $g, h, i \in [0, d]$ . Assume that  $\mathbf{j} = (\mathbf{j}_{(1)}, \mathbf{j}_{(2)}, \mathbf{j}_{(3)}) \in \mathbb{U}_{g,h}$  and  $\mathbf{k} = (\mathbf{k}_{(1)}, \mathbf{k}_{(2)}, \mathbf{k}_{(3)}) \in \mathbb{U}_{h,i}$ . Then the containments  $((\mathfrak{g}_1 \cap \mathfrak{h}_1)^\circ \setminus \mathfrak{i}_1) \cup (\mathfrak{g}_1 \cap \mathbf{k}_{(1)}) \subseteq \mathbf{j}_{(1)}$ ,  $(\mathbf{j}_{(3)} \bullet \mathfrak{i}_2) \cup (\mathbf{j}_{(3)} \cap \mathbf{k}_{(2)}) \subseteq \mathbf{j}_{(2)}$ ,  $((\mathfrak{g}_2 \cap \mathfrak{h}_2) \setminus \mathfrak{i}_2) \cup (\mathfrak{g}_2 \cap \mathbf{k}_{(3)}) \subseteq \mathbf{j}_{(3)}$ ,  $((\mathfrak{h}_1 \cap \mathfrak{i}_1)^\circ \setminus \mathfrak{g}_1) \cup (\mathfrak{i}_1 \cap \mathbf{j}_{(1)}) \subseteq \mathbf{k}_{(1)}$ ,  $(\mathbf{k}_{(3)} \bullet \mathfrak{g}_2) \cup (\mathbf{j}_{(2)} \cap \mathbf{k}_{(3)}) \subseteq \mathbf{k}_{(2)}$ ,  $((\mathfrak{h}_2 \cap \mathfrak{i}_2) \setminus \mathfrak{g}_2) \cup (\mathfrak{i}_2 \cap \mathbf{j}_{(3)}) \subseteq \mathbf{k}_{(3)}$  hold if and only if  $(\mathfrak{g}_1 \cap \mathfrak{h}_1)^\circ \setminus \mathbf{j}_{(1)} = (\mathfrak{h}_1 \cap \mathfrak{i}_1)^\circ \setminus \mathbf{k}_{(1)}$ ,  $(\mathfrak{g}_2 \cap \mathfrak{h}_2) \bullet \setminus \mathbf{j}_{(2)} = (\mathfrak{h}_2 \cap \mathfrak{i}_2) \bullet \setminus \mathbf{k}_{(2)}$ ,  $(\mathfrak{g}_2 \cap \mathfrak{h}_2) \setminus \mathbf{j}_{(3)} = (\mathfrak{h}_2 \cap \mathfrak{i}_2) \setminus \mathbf{k}_{(3)}$ .*

*Proof.* Assume that the displayed containments hold together. According to a direct computation, notice that  $(\mathfrak{g}_1 \cap \mathfrak{h}_1)^\circ \setminus \mathbf{j}_{(1)} = (\mathfrak{g}_1 \cap \mathfrak{h}_1 \cap \mathfrak{i}_1)^\circ \setminus ((\mathbf{j}_{(1)} \cup \mathbf{k}_{(1)}) \setminus \mathfrak{i}_1) = (\mathfrak{h}_1 \cap \mathfrak{i}_1)^\circ \setminus \mathbf{k}_{(1)}$ ,  $(\mathfrak{g}_2 \cap \mathfrak{h}_2) \bullet \setminus \mathbf{j}_{(2)} = ((\mathfrak{g}_2 \cap \mathfrak{h}_2 \cap \mathfrak{i}_2) \bullet \setminus ((\mathbf{j}_{(3)} \cup \mathbf{k}_{(3)}) \setminus \mathfrak{i}_2)) \cup (((\mathfrak{g}_2 \cap \mathfrak{h}_2 \cap \mathfrak{i}_2) \bullet \cap \mathbf{j}_{(3)} \cap \mathbf{k}_{(3)}) \setminus ((\mathbf{j}_{(2)} \cup \mathbf{k}_{(2)})) \setminus \mathfrak{i}_2)$ ,  $(\mathfrak{h}_2 \cap \mathfrak{i}_2) \bullet \setminus \mathbf{k}_{(2)} = ((\mathfrak{g}_2 \cap \mathfrak{h}_2 \cap \mathfrak{i}_2) \bullet \setminus ((\mathbf{j}_{(3)} \cup \mathbf{k}_{(3)}) \setminus \mathfrak{i}_2)) \cup (((\mathfrak{g}_2 \cap \mathfrak{h}_2 \cap \mathfrak{i}_2) \bullet \cap \mathbf{j}_{(3)} \cap \mathbf{k}_{(3)}) \setminus ((\mathbf{j}_{(2)} \cup \mathbf{k}_{(2)})) \setminus \mathfrak{i}_2)$ , and  $(\mathfrak{g}_2 \cap \mathfrak{h}_2) \setminus \mathbf{j}_{(3)} = (\mathfrak{g}_2 \cap \mathfrak{h}_2 \cap \mathfrak{i}_2) \setminus ((\mathbf{j}_{(3)} \cup \mathbf{k}_{(3)}) \setminus \mathfrak{i}_2) = (\mathfrak{h}_2 \cap \mathfrak{i}_2) \setminus \mathbf{k}_{(3)}$ . One direction is checked.

For the other direction, the equality  $(\mathfrak{g}_1 \cap \mathfrak{h}_1)^\circ \setminus \mathbf{j}_{(1)} = (\mathfrak{h}_1 \cap \mathfrak{i}_1)^\circ \setminus \mathbf{k}_{(1)}$  and a direct computation give  $((\mathfrak{g}_1 \cap \mathfrak{h}_1)^\circ \setminus \mathfrak{i}_1) \cup (\mathfrak{g}_1 \cap \mathbf{k}_{(1)}) \subseteq \mathbf{j}_{(1)}$  and  $((\mathfrak{h}_1 \cap \mathfrak{i}_1)^\circ \setminus \mathfrak{g}_1) \cup (\mathfrak{i}_1 \cap \mathbf{j}_{(1)}) \subseteq \mathbf{k}_{(1)}$ . Notice that  $((\mathfrak{g}_2 \cap \mathfrak{h}_2) \bullet \setminus \mathfrak{i}_2) \cup (\mathfrak{g}_2 \cap \mathbf{k}_{(2)}) \subseteq \mathbf{j}_{(2)}$  and  $((\mathfrak{h}_2 \cap \mathfrak{i}_2) \bullet \setminus \mathfrak{g}_2) \cup (\mathfrak{i}_2 \cap \mathbf{j}_{(2)}) \subseteq \mathbf{k}_{(2)}$  by the equality  $(\mathfrak{g}_2 \cap \mathfrak{h}_2) \bullet \setminus \mathbf{j}_{(2)} = (\mathfrak{h}_2 \cap \mathfrak{i}_2) \bullet \setminus \mathbf{k}_{(2)}$  and a direct computation. Therefore  $(\mathbf{j}_{(3)} \bullet \mathfrak{i}_2) \cup (\mathbf{j}_{(3)} \cap \mathbf{k}_{(2)}) \subseteq \mathbf{j}_{(2)}$  and  $(\mathbf{k}_{(3)} \bullet \mathfrak{g}_2) \cup (\mathbf{j}_{(2)} \cap \mathbf{k}_{(3)}) \subseteq \mathbf{k}_{(2)}$ . By a direct computation and the equality  $(\mathfrak{g}_2 \cap \mathfrak{h}_2) \setminus \mathbf{j}_{(3)} = (\mathfrak{h}_2 \cap \mathfrak{i}_2) \setminus \mathbf{k}_{(3)}$ , notice that  $((\mathfrak{g}_2 \cap \mathfrak{h}_2) \setminus \mathfrak{i}_2) \cup (\mathfrak{g}_2 \cap \mathbf{k}_{(3)}) \subseteq \mathbf{j}_{(3)}$  and  $((\mathfrak{h}_2 \cap \mathfrak{i}_2) \setminus \mathfrak{g}_2) \cup (\mathfrak{i}_2 \cap \mathbf{j}_{(3)}) \subseteq \mathbf{k}_{(3)}$ . The desired lemma thus follows.  $\square$

We are now ready to deduce the remaining main results of this section.

**Theorem 7.14.** *Assume that  $g, h, i \in [0, d]$ . Assume that  $\mathbf{j} = (\mathbf{j}_{(1)}, \mathbf{j}_{(2)}, \mathbf{j}_{(3)}) \in \mathbb{U}_{g,h}$  and  $\mathbf{k} = (\mathbf{k}_{(1)}, \mathbf{k}_{(2)}, \mathbf{k}_{(3)}) \in \mathbb{U}_{\ell,i}$  for some  $\ell \in [0, d]$ . Assume that  $p \nmid k_{[g,h,g]} k_{[\ell,i,\ell]} k_{\mathbf{j}} k_{\mathbf{k}}$ . Then the inequality  $(D_{g,h,\mathbf{j}} + \text{Rad}(\mathbb{T}))(D_{\ell,i,\mathbf{k}} + \text{Rad}(\mathbb{T})) \neq O + \text{Rad}(\mathbb{T})$  holds only if  $h = \ell$ ,  $(\mathfrak{g}_1 \cap \mathfrak{h}_1)^\circ \setminus \mathbf{j}_{(1)} = (\mathfrak{h}_1 \cap \mathfrak{i}_1)^\circ \setminus \mathbf{k}_{(1)}$ ,  $(\mathfrak{g}_2 \cap \mathfrak{h}_2) \bullet \setminus \mathbf{j}_{(2)} = (\mathfrak{h}_2 \cap \mathfrak{i}_2) \bullet \setminus \mathbf{k}_{(2)}$ ,  $(\mathfrak{g}_2 \cap \mathfrak{h}_2) \setminus \mathbf{j}_{(3)} = (\mathfrak{h}_2 \cap \mathfrak{i}_2) \setminus \mathbf{k}_{(3)}$ .*

*Proof.* Assume that  $h = \ell$  by Equation (2.2). The displayed inequality implies that  $D_{g,h,\mathbf{j}} D_{h,i,\mathbf{k}} \neq O$ . The desired theorem is from combining Lemmas 5.14, 7.1, 7.13.  $\square$

**Theorem 7.15.** *Assume that  $g, h, i \in [0, d]$ . Assume that  $\mathbf{j} = (\mathbf{j}_{(1)}, \mathbf{j}_{(2)}, \mathbf{j}_{(3)}) \in \mathbb{U}_{g,h}$ ,  $\mathbf{k} = (\mathbf{k}_{(1)}, \mathbf{k}_{(2)}, \mathbf{k}_{(3)}) \in \mathbb{U}_{h,i}$ ,  $p \nmid k_{[g,h,g]} k_{[h,g,h]} k_{[h,i,h]} k_{[i,h,i]} k_{\mathbf{j}} k_{\mathbf{k}}$ ,  $(\mathfrak{g}_1 \cap \mathfrak{h}_1)^\circ \setminus \mathbf{j}_{(1)} = (\mathfrak{h}_1 \cap \mathfrak{i}_1)^\circ \setminus \mathbf{k}_{(1)}$ ,  $(\mathfrak{g}_2 \cap \mathfrak{h}_2) \bullet \setminus \mathbf{j}_{(2)} = (\mathfrak{h}_2 \cap \mathfrak{i}_2) \bullet \setminus \mathbf{k}_{(2)}$ ,  $(\mathfrak{g}_2 \cap \mathfrak{h}_2) \setminus \mathbf{j}_{(3)} = (\mathfrak{h}_2 \cap \mathfrak{i}_2) \setminus \mathbf{k}_{(3)}$ . Then  $p \nmid k_{[g,i,g]} k_{[i,g,i]} k_{(g,h,i,\mathbf{j},\mathbf{k})}$  and  $(D_{g,h,\mathbf{j}} + \text{Rad}(\mathbb{T}))(D_{h,i,\mathbf{k}} + \text{Rad}(\mathbb{T})) = D_{g,i,(g,h,i,\mathbf{j},\mathbf{k})} + \text{Rad}(\mathbb{T})$ .*

*Proof.* Since  $p \nmid k_{[g,h,g]} k_{[h,g,h]} k_{[h,i,h]} k_{[i,h,i]} k_{\mathbf{j}} k_{\mathbf{k}}$ , Lemmas 7.6 and 7.11 thus imply that  $p \nmid k_{[g,i,g]} k_{[i,g,i]} k_{(g,h,i,\mathbf{j},\mathbf{k})}$ . Assume that  $\mathbf{l} = (\mathbf{l}_{(1)}, \mathbf{l}_{(2)}, \mathbf{l}_{(3)}) \in \mathbb{U}_{g,h}$ ,  $\mathbf{j} \preceq \mathbf{l} \preceq (g, h; \mathbf{j})$ , and the containments  $((\mathfrak{h}_1 \cap \mathfrak{i}_1)^\circ \setminus \mathfrak{g}_1) \cup (\mathfrak{i}_1 \cap \mathbf{l}_{(1)}) \subseteq \mathbf{k}_{(1)}$ ,  $(\mathbf{k}_{(3)} \bullet \setminus \mathfrak{g}_2) \cup (\mathbf{k}_{(3)} \cap \mathbf{l}_{(2)}) \subseteq \mathbf{k}_{(2)}$ ,  $((\mathfrak{h}_2 \cap \mathfrak{i}_2) \setminus \mathfrak{g}_2) \cup (\mathfrak{i}_2 \cap \mathbf{l}_{(3)}) \subseteq \mathbf{k}_{(3)}$  hold together. Notice that  $\mathfrak{i}_1 \cap \mathbf{l}_{(1)} = \mathfrak{g}_1 \cap \mathbf{k}_{(1)} = \mathfrak{i}_1 \cap \mathbf{j}_{(1)}$  and  $\mathbf{l}_{(1)} \setminus \mathfrak{i}_1 = \mathbf{j}_{(1)} \setminus \mathfrak{i}_1$  as  $((\mathfrak{g}_1 \cap \mathfrak{h}_1)^\circ \setminus \mathfrak{i}_1) \cup (\mathfrak{g}_1 \cap \mathbf{k}_{(1)}) \subseteq \mathbf{j}_{(1)}$  by Lemma 7.13. Notice that  $\mathfrak{i}_2 \cap \mathbf{l}_{(2)} = \mathfrak{g}_2 \cap \mathfrak{i}_2 \cap \mathbf{k}_{(3)} \cap \mathbf{l}_{(2)} = \mathfrak{g}_2 \cap \mathfrak{i}_2 \cap \mathbf{j}_{(3)} \cap \mathbf{k}_{(2)} = \mathfrak{i}_2 \cap \mathbf{j}_{(2)}$  and  $\mathbf{l}_{(2)} \setminus \mathfrak{i}_2 = \mathbf{j}_{(2)} \setminus \mathfrak{i}_2$  as  $(\mathbf{j}_{(3)} \bullet \setminus \mathfrak{i}_2) \cup (\mathbf{j}_{(3)} \cap \mathbf{k}_{(2)}) \subseteq \mathbf{j}_{(2)}$  and  $\mathfrak{g}_2 \cap \mathbf{k}_{(3)} \subseteq \mathbf{j}_{(3)}$  by Lemma 7.13. Notice that  $\mathfrak{i}_2 \cap \mathbf{l}_{(3)} = \mathfrak{g}_2 \cap \mathbf{k}_{(3)} = \mathfrak{i}_2 \cap \mathbf{j}_{(3)}$  and  $\mathbf{l}_{(3)} \setminus \mathfrak{i}_2 = \mathbf{j}_{(3)} \setminus \mathfrak{i}_2$  as  $((\mathfrak{g}_2 \cap \mathfrak{h}_2) \setminus \mathfrak{i}_2) \cup (\mathfrak{g}_2 \cap \mathbf{k}_{(3)}) \subseteq \mathbf{j}_{(3)}$  by Lemma 7.13. The above discussion thus implies that  $\mathbf{j} = \mathbf{l}$ . As  $p \nmid k_{[g,h,g]} k_{\mathbf{j}}$ , notice that  $D_{g,h,\mathbf{j}} \neq O$ . The desired theorem thus follows from Lemmas 7.12 and 5.14.  $\square$

We conclude this section by presenting a remark of Theorems 7.14 and 7.15.

**Remark 7.16.** *The structure constants of the  $\mathbb{F}$ -basis of  $\mathbb{T}/\text{Rad}(\mathbb{T})$  in Theorem 7.4 are read off from Theorems 7.14 and 7.15. Notice that they are contained in  $\{\overline{0}, \overline{1}\}$ .*

8. ALGEBRAIC STRUCTURE OF  $\mathbb{T}$ : WEDDERBURN-ARTIN DECOMPOSITION

In this section, we apply Theorems 7.14 and 7.15 to present the Wedderburn-Artin decomposition of  $\mathbb{T}$ . This means that we determine the algebraic structure of the semisimple  $\mathbb{F}$ -algebra  $\mathbb{T}/\text{Rad}(\mathbb{T})$  up to  $\mathbb{F}$ -algebra isomorphism. For this purpose, we recall Notations 3.9, 3.10, 3.17, 3.18, 3.21, 4.13, 5.6, 5.7 and display another lemma.

**Lemma 8.1.** *Assume that  $g, h, i \in [0, d]$ . Assume that  $\mathbf{j} = (\mathbf{j}_{(1)}, \mathbf{j}_{(2)}, \mathbf{j}_{(3)}) \in \mathbb{U}_{g,h}$  and  $\mathbf{k} = (\mathbf{k}_{(1)}, \mathbf{k}_{(2)}, \mathbf{k}_{(3)}) \in \mathbb{U}_{h,i}$ . Assume that  $\mathbb{l}_{(1)} = ((g_1 \cap i_1)^\circ \setminus h_1) \cup (g_1 \cap i_1 \cap (\mathbf{j}_{(1)} \cup \mathbf{k}_{(1)}))$ ,  $\mathbb{l}_{(2)} = ((g_2 \cap i_2)^\bullet \setminus h_2) \cup (g_2 \cap i_2 \cap (\mathbf{j}_{(2)} \cup \mathbf{k}_{(2)}))$ ,  $\mathbb{l}_{(3)} = ((g_2 \cap i_2) \setminus h_2) \cup (g_2 \cap i_2 \cap (\mathbf{j}_{(3)} \cup \mathbf{k}_{(3)}))$ ,  $(g_1 \cap h_1)^\circ \setminus \mathbf{j}_{(1)} = (h_1 \cap i_1)^\circ \setminus \mathbf{k}_{(1)}$ ,  $(g_2 \cap h_2)^\bullet \setminus \mathbf{j}_{(2)} = (h_2 \cap i_2)^\bullet \setminus \mathbf{k}_{(2)}$ ,  $(g_2 \cap h_2) \setminus \mathbf{j}_{(3)} = (h_2 \cap i_2) \setminus \mathbf{k}_{(3)}$ . Then  $(g_1 \cap h_1)^\circ \setminus \mathbf{j}_{(1)} = (h_1 \cap i_1)^\circ \setminus \mathbf{k}_{(1)} = (g_1 \cap i_1)^\circ \setminus \mathbb{l}_{(1)}$ ,  $(g_2 \cap h_2)^\bullet \setminus \mathbf{j}_{(2)} = (h_2 \cap i_2)^\bullet \setminus \mathbf{k}_{(2)} = (g_2 \cap i_2)^\bullet \setminus \mathbb{l}_{(2)}$ , and  $(g_2 \cap h_2) \setminus \mathbf{j}_{(3)} = (h_2 \cap i_2) \setminus \mathbf{k}_{(3)} = (g_2 \cap i_2) \setminus \mathbb{l}_{(3)}$ .*

*Proof.* By a direct computation, notice that  $(g_1 \cap i_1)^\circ \setminus \mathbb{l}_{(1)} = (g_1 \cap h_1 \cap i_1)^\circ \setminus (\mathbf{j}_{(1)} \cup \mathbf{k}_{(1)})$ ,  $(g_2 \cap i_2)^\bullet \setminus \mathbb{l}_{(2)} = (g_2 \cap h_2 \cap i_2)^\bullet \setminus (\mathbf{j}_{(2)} \cup \mathbf{k}_{(2)})$ , and  $(g_2 \cap i_2) \setminus \mathbb{l}_{(3)} = (g_2 \cap h_2 \cap i_2) \setminus (\mathbf{j}_{(3)} \cup \mathbf{k}_{(3)})$ . Notice that  $(g_1 \cap h_1)^\circ \setminus \mathbf{j}_{(1)} = (g_1 \cap h_1 \cap i_1)^\circ \setminus (\mathbf{j}_{(1)} \cup \mathbf{k}_{(1)})$  as  $((g_1 \cap h_1)^\circ \setminus i_1) \cup (g_1 \cap \mathbf{k}_{(1)}) \subseteq \mathbf{j}_{(1)}$  by Lemma 7.13. Notice that  $(g_2 \cap h_2)^\bullet \setminus i_2 \subseteq \mathbf{j}_{(2)}$  as  $(g_2 \cap h_2)^\bullet \setminus \mathbf{j}_{(2)} = (h_2 \cap i_2)^\bullet \setminus \mathbf{k}_{(2)}$ . As  $g_2 \cap \mathbf{k}_{(3)} \subseteq \mathbf{j}_{(3)}$  and  $\mathbf{j}_{(3)} \cap \mathbf{k}_{(2)} \subseteq \mathbf{j}_{(2)}$  by Lemma 7.13,  $(g_2 \cap h_2)^\bullet \setminus \mathbf{j}_{(2)} = (g_2 \cap h_2 \cap i_2)^\bullet \setminus (\mathbf{j}_{(2)} \cup \mathbf{k}_{(2)})$ . Notice that  $(g_2 \cap h_2) \setminus \mathbf{j}_{(3)} = (g_2 \cap h_2 \cap i_2) \setminus (\mathbf{j}_{(3)} \cup \mathbf{k}_{(3)})$  as  $((g_2 \cap h_2) \setminus i_2) \cup (g_2 \cap \mathbf{k}_{(3)}) \subseteq \mathbf{j}_{(3)}$  by Lemma 7.13. The desired lemma thus follows from the above discussion.  $\square$

Lemma 8.1 motivates us to present the following notations and another lemma.

**Notation 8.2.** *Define  $\mathbb{D} = \{(a, b, \mathbf{c}) : a, b \in [0, d], \mathbf{c} \in \mathbb{U}_{a,b}, p \nmid k_{[a,b,a]} k_{[b,a,b]} k_{\mathbf{c}}\}$ . Notice that  $\mathbb{D} \neq \emptyset$  as  $(0, 0, \mathbf{o}) \in \mathbb{D}$ . Assume that  $(g, h, \mathbf{i}), (j, k, \mathbf{l}) \in \mathbb{D}$ ,  $\mathbf{i} = (\mathbf{i}_{(1)}, \mathbf{i}_{(2)}, \mathbf{i}_{(3)})$ , and  $\mathbf{l} = (\mathbf{l}_{(1)}, \mathbf{l}_{(2)}, \mathbf{l}_{(3)})$ . Then  $D_{g,h,\mathbf{i}}$  is a defined nonzero matrix. Write  $(g, h, \mathbf{i}) \sim (j, k, \mathbf{l})$  if and only if  $(g_1 \cap h_1)^\circ \setminus \mathbf{i}_{(1)} = (\mathbf{j}_{(1)} \cap \mathbf{k}_{(1)})^\circ \setminus \mathbf{l}_{(1)}$ ,  $(g_2 \cap h_2)^\bullet \setminus \mathbf{i}_{(2)} = (\mathbf{j}_{(2)} \cap \mathbf{k}_{(2)})^\bullet \setminus \mathbf{l}_{(2)}$ ,  $(g_2 \cap h_2) \setminus \mathbf{i}_{(3)} = (\mathbf{j}_{(2)} \cap \mathbf{k}_{(2)}) \setminus \mathbf{l}_{(3)}$ . Therefore  $\sim$  is an equivalence relation on  $\mathbb{D}$ . There is  $n_\sim \in \mathbb{N}$  such that  $\mathbb{D}_1, \mathbb{D}_2, \dots, \mathbb{D}_{n_\sim}$  are exactly all equivalence classes of  $\mathbb{D}$  with respect to  $\sim$ . Assume that  $m \in [1, n_\sim]$ . Set  $\mathbb{D}(m) = \{a : a \in [0, d], \exists \mathbf{b} \in \mathbb{U}_{a,a}, (a, a, \mathbf{b}) \in \mathbb{D}_m\}$ .*

**Notation 8.3.** *Assume that  $g \in [1, n_\sim]$ . Set  $\mathbb{I}(g) = \langle \{D_{a,b,\mathbf{c}} + \text{Rad}(\mathbb{T}) : (a, b, \mathbf{c}) \in \mathbb{D}_g\} \rangle$ . According to Theorem 7.4, it is clear that the  $\mathbb{F}$ -dimension of  $\mathbb{I}(g)$  is equal to  $|\mathbb{D}_g|$ .*

**Lemma 8.4.** *Assume that  $g \in [1, n_\sim]$ . Then  $\mathbb{I}(g)$  is a two-sided ideal of  $\mathbb{T}/\text{Rad}(\mathbb{T})$ . Moreover,  $\mathbb{T}/\text{Rad}(\mathbb{T})$  is a direct sum of the  $\mathbb{F}$ -linear subspaces  $\mathbb{I}(1), \mathbb{I}(2), \dots, \mathbb{I}(n_\sim)$ .*

*Proof.* The first statement is from combining Theorems 7.4, 7.14, 7.15, Lemma 8.1. As  $\{\mathbb{D}_1, \mathbb{D}_2, \dots, \mathbb{D}_{n_\sim}\}$  is a partition of  $\mathbb{D}$ , the desired lemma is from Theorem 7.4.  $\square$

The following four lemmas focus on the investigation of objects in Notation 8.2.

**Lemma 8.5.** *Assume that  $g \in [1, n_\sim]$  and  $(h, i, \mathbf{j}), (h, i, \mathbf{k}) \in \mathbb{D}_g$ . Then  $\mathbf{j} = \mathbf{k}$ .*

*Proof.* Assume that  $\mathbf{j} = (\mathbf{j}_{(1)}, \mathbf{j}_{(2)}, \mathbf{j}_{(3)})$  and  $\mathbf{k} = (\mathbf{k}_{(1)}, \mathbf{k}_{(2)}, \mathbf{k}_{(3)})$ . As  $(h, i, \mathbf{j}), (h, i, \mathbf{k}) \in \mathbb{D}_g$ , notice that  $\mathbf{j}, \mathbf{k} \in \mathbb{U}_{h,i}$ ,  $(h_1 \cap i_1)^\circ \setminus \mathbf{j}_{(1)} = (h_1 \cap i_1)^\circ \setminus \mathbf{k}_{(1)}$ ,  $(h_2 \cap i_2)^\bullet \setminus \mathbf{j}_{(2)} = (h_2 \cap i_2)^\bullet \setminus \mathbf{k}_{(2)}$ , and  $(h_2 \cap i_2) \setminus \mathbf{j}_{(3)} = (h_2 \cap i_2) \setminus \mathbf{k}_{(3)}$ . So  $\mathbf{j}_{(1)} = (h_1 \cap i_1)^\circ \cap \mathbf{j}_{(1)} = (h_1 \cap i_1)^\circ \cap \mathbf{k}_{(1)} = \mathbf{k}_{(1)}$ ,  $\mathbf{j}_{(2)} = (h_2 \cap i_2)^\bullet \cap \mathbf{j}_{(2)} = (h_2 \cap i_2)^\bullet \cap \mathbf{k}_{(2)} = \mathbf{k}_{(2)}$ ,  $\mathbf{j}_{(3)} = (h_2 \cap i_2) \cap \mathbf{j}_{(3)} = (h_2 \cap i_2) \cap \mathbf{k}_{(3)} = \mathbf{k}_{(3)}$  since  $\mathbf{j}, \mathbf{k} \in \mathbb{U}_{h,i}$ . The desired lemma thus follows from the above discussion.  $\square$

**Lemma 8.6.** *Assume that  $g \in [1, n_\sim]$  and  $(h, i, j) \in \mathbb{D}_g$ . Then there are  $\mathfrak{k} \in \mathbb{U}_{h,h}$  and  $\mathfrak{l} \in \mathbb{U}_{i,i}$  such that  $(h, h, \mathfrak{k}), (i, i, \mathfrak{l}) \in \mathbb{D}_g$ . In particular,  $h, i \in \mathbb{D}(g)$ .*

*Proof.* Assume that  $\mathbf{j} = (\mathbf{j}_{(1)}, \mathbf{j}_{(2)}, \mathbf{j}_{(3)})$ ,  $\mathfrak{k} = (\mathbb{k}_{(1)}, \mathbb{k}_{(2)}, \mathbb{k}_{(3)})$ ,  $\mathfrak{l} = (\mathbb{l}_{(1)}, \mathbb{l}_{(2)}, \mathbb{l}_{(3)})$ , where  $\mathbb{k}_{(1)} = (\mathbb{h}_1^\circ \setminus \mathbb{i}_1) \cup \mathbf{j}_{(1)}$ ,  $\mathbb{k}_{(2)} = (\mathbb{h}_2^\bullet \setminus \mathbb{i}_2) \cup \mathbf{j}_{(2)}$ ,  $\mathbb{k}_{(3)} = (\mathbb{h}_2 \setminus \mathbb{i}_2) \cup \mathbf{j}_{(3)}$ ,  $\mathbb{l}_{(1)} = (\mathbb{i}_1^\circ \setminus \mathbb{h}_1) \cup \mathbf{j}_{(1)}$ ,  $\mathbb{l}_{(2)} = (\mathbb{i}_2^\bullet \setminus \mathbb{h}_2) \cup \mathbf{j}_{(2)}$ , and  $\mathbb{l}_{(3)} = (\mathbb{i}_2 \setminus \mathbb{h}_2) \cup \mathbf{j}_{(3)}$ . As  $(h, i, j) \in \mathbb{D}_g$ , notice that  $\mathbf{j} \in \mathbb{U}_{h,i}$  and  $p \nmid k_{[h,i,h]}k_{[i,h,i]}k_j$ . Hence  $\mathfrak{k} \in \mathbb{U}_{h,h}$ ,  $\mathfrak{l} \in \mathbb{U}_{i,i}$ , and  $p \nmid k_{\mathfrak{k}}k_{\mathfrak{l}}$  by a direct computation.

Notice that  $(\mathbb{h}_1 \cap \mathbb{i}_1)^\circ \setminus \mathbf{j}_{(1)} = \mathbb{h}_1^\circ \setminus \mathbb{k}_{(1)} = \mathbb{i}_1^\circ \setminus \mathbb{l}_{(1)}$ ,  $(\mathbb{h}_2 \cap \mathbb{i}_2)^\bullet \setminus \mathbf{j}_{(2)} = \mathbb{h}_2^\bullet \setminus \mathbb{k}_{(2)} = \mathbb{i}_2^\bullet \setminus \mathbb{l}_{(2)}$ , and  $(\mathbb{h}_2 \cap \mathbb{i}_2) \setminus \mathbf{j}_{(3)} = \mathbb{h}_2 \setminus \mathbb{k}_{(3)} = \mathbb{i}_2 \setminus \mathbb{l}_{(3)}$ . The desired lemma thus follows.  $\square$

**Lemma 8.7.** *Assume that  $g \in [1, n_\sim]$  and  $(h, h, i), (j, j, \mathfrak{k}) \in \mathbb{D}_g$ . Then there exists  $\mathfrak{l} \in \mathbb{U}_{h,j}$  such that  $(h, j, \mathfrak{l}) \in \mathbb{D}_g$ .*

*Proof.* Assume that  $\mathbf{i} = (\mathbf{i}_{(1)}, \mathbf{i}_{(2)}, \mathbf{i}_{(3)})$  and  $\mathfrak{k} = (\mathbb{k}_{(1)}, \mathbb{k}_{(2)}, \mathbb{k}_{(3)})$ . As  $(h, h, i), (j, j, \mathfrak{k}) \in \mathbb{D}_g$ , notice that  $p \nmid k_i k_{\mathfrak{k}}$ ,  $\mathbb{h}_1^\circ \setminus \mathbf{i}_{(1)} = \mathbf{j}_1^\circ \setminus \mathbb{k}_{(1)}$ ,  $\mathbb{h}_2^\bullet \setminus \mathbf{i}_{(2)} = \mathbf{j}_2^\bullet \setminus \mathbb{k}_{(2)}$ , and  $\mathbb{h}_2 \setminus \mathbf{i}_{(3)} = \mathbf{j}_2 \setminus \mathbb{k}_{(3)}$ . Therefore  $\mathbb{h}_1^\circ \setminus \mathbf{j}_1 \subseteq \mathbf{i}_{(1)}$ ,  $\mathbb{h}_2^\bullet \setminus \mathbf{j}_2 \subseteq \mathbf{i}_{(2)}$ ,  $\mathbb{h}_2 \setminus (\mathbb{h}_2^\bullet \cup \mathbf{j}_2) \subseteq \mathbf{i}_{(3)} \setminus \mathbf{i}_{(2)}$ ,  $\mathbf{j}_1^\circ \setminus \mathbb{h}_1 \subseteq \mathbb{k}_{(1)}$ ,  $\mathbf{j}_2^\bullet \setminus \mathbb{h}_2 \subseteq \mathbb{k}_{(2)}$ , and  $\mathbf{j}_2 \setminus (\mathbb{h}_2 \cup \mathbf{j}_2^\bullet) \subseteq \mathbb{k}_{(3)} \setminus \mathbb{k}_{(2)}$ . Therefore  $p \nmid k_{[h,j,h]}k_{[j,h,j]}$  as  $p \nmid k_i k_{\mathfrak{k}}$  and Lemma 3.22 holds. Set  $\mathfrak{l} = h \cap \mathfrak{k}$ . So  $\mathfrak{l} \in \mathbb{U}_{h,j}$  and  $p \nmid k_{[h,j,h]}k_{[j,h,j]}k_{\mathfrak{l}}$  by Lemma 4.5.

Assume that  $\mathfrak{l} = (\mathbb{l}_{(1)}, \mathbb{l}_{(2)}, \mathbb{l}_{(3)})$ . Notice that  $\mathbb{h}_1^\circ \setminus \mathbf{i}_{(1)} = \mathbf{j}_1^\circ \setminus \mathbb{k}_{(1)} = (\mathbb{h}_1 \cap \mathbf{j}_1)^\circ \setminus \mathbb{l}_{(1)}$ ,  $\mathbb{h}_2^\bullet \setminus \mathbf{i}_{(2)} = \mathbf{j}_2^\bullet \setminus \mathbb{k}_{(2)} = (\mathbb{h}_2 \cap \mathbf{j}_2)^\bullet \setminus \mathbb{l}_{(2)}$ , and  $\mathbb{h}_2 \setminus \mathbf{i}_{(3)} = \mathbf{j}_2 \setminus \mathbb{k}_{(3)} = (\mathbb{h}_2 \cap \mathbf{j}_2) \setminus \mathbb{l}_{(3)}$  by a direct computation. The desired lemma thus follows from the above discussion.  $\square$

Lemmas 8.5, 8.6, 8.7 allow us to present the following lemma and another notation.

**Lemma 8.8.** *Assume that  $g \in [1, n_\sim]$ . Then the cartesian product  $\mathbb{D}(g) \times \mathbb{D}(g) \neq \emptyset$ . Moreover, the map that sends  $(h, i, j)$  to  $(h, i)$  is bijective from  $\mathbb{D}_g$  to  $\mathbb{D}(g) \times \mathbb{D}(g)$ . In particular, the  $\mathbb{F}$ -dimension of  $\mathbb{I}(g)$  is equal to  $|\mathbb{D}(g)|^2$ .*

*Proof.* The first statement follows since  $\mathbb{D}_g \neq \emptyset$  and Lemma 8.6 holds. The second statement thus follows from combining Lemmas 8.5, 8.6, and 8.7. As  $|\mathbb{D}_g| = |\mathbb{D}(g)|^2$  by the above discussion, the desired lemma thus follows from Notation 8.3.  $\square$

**Notation 8.9.** *Assume that  $g \in [1, n_\sim]$ . According to Lemma 8.8 and Theorem 7.4, there is a unique  $D_{h,i,j} + \text{Rad}(\mathbb{T}) \in \{D_{a,b,c} + \text{Rad}(\mathbb{T}) : (a, b, c) \in \mathbb{D}_g\}$  for any  $h, i \in \mathbb{D}(g)$ . This nonzero element in  $\mathbb{T}/\text{Rad}(\mathbb{T})$  is denoted by  $D_{h,i}(g)$  for any  $h, i \in \mathbb{D}(g)$ . For completeness, define  $D_{h,i}(g) = O + \text{Rad}(\mathbb{T})$  for any  $h, i \in [0, d]$  and  $\{h, i\} \not\subseteq \mathbb{D}(g)$ . Then  $\mathbb{I}(g)$  has an  $\mathbb{F}$ -basis  $\{D_{a,b}(g) : a, b \in \mathbb{D}(g)\}$  by Notation 8.3 and Lemma 8.8.*

The following lemma gives the computational rule of the objects in Notation 8.9.

**Lemma 8.10.**  *$\mathbb{T}/\text{Rad}(\mathbb{T})$  has an  $\mathbb{F}$ -basis  $\{D_{b,c}(a) : a \in [1, n_\sim], b, c \in \mathbb{D}(a)\}$ . Moreover, if  $g, h \in [1, n_\sim]$ ,  $i, j \in \mathbb{D}(g)$ , and  $k, \ell \in \mathbb{D}(h)$ , then  $D_{i,j}(g)D_{k,\ell}(h) = \delta_{g,h}\delta_{j,k}D_{i,\ell}(g)$ .*

*Proof.* The first statement is from Lemma 8.4 and Notation 8.9. Assume that  $g = h$  by Lemma 8.4. By Notation 8.9, there are unique  $\mathbf{m} \in \mathbb{U}_{i,j}$ ,  $\mathbf{q} \in \mathbb{U}_{k,\ell}$ ,  $\mathbf{r} \in \mathbb{U}_{i,\ell}$  such that  $(i, j, \mathbf{m}), (k, \ell, \mathbf{q}), (i, \ell, \mathbf{r}) \in \mathbb{D}_g$ ,  $D_{i,j}(g) = D_{i,j,\mathbf{m}} + \text{Rad}(\mathbb{T})$ ,  $D_{k,\ell}(g) = D_{k,\ell,\mathbf{q}} + \text{Rad}(\mathbb{T})$ , and  $D_{i,\ell}(g) = D_{i,\ell,\mathbf{r}} + \text{Rad}(\mathbb{T})$ . If  $j \neq k$ , then the desired equality is from the fact  $(i, j, \mathbf{m}), (k, \ell, \mathbf{q}) \in \mathbb{D}_g$  and Theorem 7.14. If  $j = k$ , the desired equality is from the fact  $(i, j, \mathbf{m}), (j, \ell, \mathbf{q}), (i, \ell, \mathbf{r}) \in \mathbb{D}_g$  and Theorem 7.15. The desired lemma follows.  $\square$

The computational rule in Lemma 8.10 allows us to prove the following lemmas.

**Lemma 8.11.** *Assume that  $g \in [1, n_\sim]$  and  $h, i \in \mathbb{D}(g)$ . Then  $\langle \{D_{a,h}(g) : a \in \mathbb{D}(g)\} \rangle$  is an irreducible  $\mathbb{T}$ -module and  $\langle \{D_{a,h}(g) : a \in \mathbb{D}(g)\} \rangle \cong \langle \{D_{a,i}(g) : a \in \mathbb{D}(g)\} \rangle$  as irreducible  $\mathbb{T}$ -modules.*

*Proof.* The first statement follows from Lemma 8.10. By Lemma 8.10, the map that sends  $D_{j,h}(g)$  to  $D_{j,i}(g)$  is a  $\mathbb{T}$ -module isomorphism from  $\langle \{D_{a,h}(g) : a \in \mathbb{D}(g)\} \rangle$  to  $\langle \{D_{a,i}(g) : a \in \mathbb{D}(g)\} \rangle$ . The desired lemma is thus from the above discussion.  $\square$

**Lemma 8.12.** *Assume that  $g \in [1, n_\sim]$ . Then  $\mathbb{I}(g) \cong M_{|\mathbb{D}(g)|}(\mathbb{F})$  as  $\mathbb{F}$ -algebras.*

*Proof.* It suffices to check that  $\mathbb{I}(g) \cong M_{|\mathbb{D}(g)|}(\mathbb{F})$  as  $\mathbb{F}$ -algebras. For any  $h, i \in \mathbb{D}(g)$ , let  $E_{(h,i)}$  be the  $\{\overline{0}, \overline{1}\}$ -matrix in  $M_{|\mathbb{D}(g)|}(\mathbb{F})$  whose unique nonzero entry is the  $(h, i)$ -entry. In particular,  $E_{(h,i)}E_{(j,k)} = \delta_{i,j}E_{(h,k)}$  for any  $h, i, j, k \in \mathbb{D}(g)$ . By Notation 8.9 and Lemma 8.10, the map that sends  $D_{h,i}(g)$  to  $E_{(h,i)}$  is an  $\mathbb{F}$ -algebra isomorphism from  $\mathbb{I}(g)$  to  $M_{|\mathbb{D}(g)|}(\mathbb{F})$ . The desired lemma thus follows from the above discussion.  $\square$

We are now ready to deduce the final main result of this paper and two corollaries.

**Theorem 8.13.** *The number of all pairwise nonisomorphic irreducible  $\mathbb{T}$ -modules equals  $n_\sim$ . Furthermore,*

$$\mathbb{T}/\text{Rad}(\mathbb{T}) \cong \bigoplus_{g=1}^{n_\sim} M_{|\mathbb{D}(g)|}(\mathbb{F}) \text{ as } \mathbb{F}\text{-algebras.}$$

*Proof.* The desired theorem follows from combining Lemmas 8.4, 8.12, and 2.2.  $\square$

**Corollary 8.14.** *Assume that  $g, h \in [1, n_\sim]$  and  $\mathbb{Irr}(g)$  denotes the unique irreducible  $\mathbb{T}$ -module up to  $\mathbb{T}$ -module isomorphism whose definition is given in Lemma 8.11. Then  $\mathbb{Irr}(g) \cong \mathbb{Irr}(h)$  as irreducible  $\mathbb{T}$ -modules if and only if  $g = h$ . Furthermore,  $\{\mathbb{Irr}(a) : a \in [1, n_\sim]\}$  contains exactly the representatives of all isomorphic classes of irreducible  $\mathbb{T}$ -modules and all irreducible  $\mathbb{T}$ -modules are precisely all absolutely irreducible  $\mathbb{T}$ -modules.*

*Proof.* The first statement follows from Lemmas 8.11 and 8.10. The desired corollary thus follows from combining the first statement, Theorem 8.13, and Lemma 2.2.  $\square$

**Corollary 8.15.** *Assume that  $\mathbb{S}$  is a  $p'$ -valenced scheme. Then*

$$\mathbb{T} \cong \bigoplus_{g=1}^{n_\sim} M_{|\mathbb{D}(g)|}(\mathbb{F}) \text{ as } \mathbb{F}\text{-algebras.}$$

*Proof.* The desired corollary is a direct application of Theorems 5.21 and 8.13.  $\square$

We close the whole paper by an example of Theorem 8.13 and Corollary 8.15.

**Example 8.16.** *Assume that  $n = \ell_1 = 2$ ,  $\ell_2 = m_1 = 3$ , and  $m_2 = 4$ . Then  $d = 8$ . If  $p = 2$ , Theorem 8.13 gives  $\mathbb{T}/\text{Rad}(\mathbb{T}) \cong M_4(\mathbb{F}) \oplus 4M_2(\mathbb{F}) \oplus 4M_1(\mathbb{F})$  as  $\mathbb{F}$ -algebras. If  $p = 3$ , Theorem 8.13 gives  $\mathbb{T}/\text{Rad}(\mathbb{T}) \cong M_4(\mathbb{F}) \oplus 5M_2(\mathbb{F}) \oplus 6M_1(\mathbb{F})$  as  $\mathbb{F}$ -algebras. Assume that  $p \notin [2, 3]$ . Theorem 5.21 thus suggests that  $\mathbb{S}$  is a  $p'$ -valenced scheme. Therefore Corollary 8.15 implies that  $\mathbb{T} \cong M_9(\mathbb{F}) \oplus 5M_3(\mathbb{F}) \oplus 6M_1(\mathbb{F})$  as  $\mathbb{F}$ -algebras.*

**Disclosure statement.** No relevant financial or nonfinancial interests are reported.

## REFERENCES

- [1] R. A. Bailey, Association Schemes: Designed Experiments, Algebra and Combinatorics, Cambridge Stud. Adv. Math., vol **84**, Cambridge University Press, Cambridge, 2004.
- [2] G. Bhattacharyya, S. Y. Song, R. Tanaka, Terwilliger algebras of wreath products of one-class association schemes, J. Algebraic Combin. **31** (2010), 455-466.
- [3] Z. Chen, C. Xi, Structure of Terwilliger algebras of quasi-thin association schemes, Preprint (2024), <https://doi.org/10.48550/arXiv.2410.21600>.
- [4] B. Curtin, I. Daqqa, The subconstituent algebra of a Latin square, European J. Combin. **30** (2009), 447-457.
- [5] Y.A. Drozd, V.V. Kirichenko, Finite Dimensional Algebras, Springer-Verlag, Berlin, 1994.
- [6] A. Hanaki, Modular Terwilliger algebras of association schemes, Graphs Combin. **37** (2021), 1521-1529.
- [7] A. Herman, A survey of semisimple algebras in algebraic combinatorics, Indian J. Pure Appl. Math. **52** (2021), 631-642.
- [8] Y. Jiang, A note on modular Terwilliger algebras of association schemes, Beitr. Algebra Geom. **63** (2022), 829-851.
- [9] Y. Jiang, On Terwilliger  $\mathbb{F}$ -algebras of quasi-thin association schemes, J. Algebraic Combin. **57** (2023), 1219-1251.
- [10] Y. Jiang, On Terwilliger  $\mathbb{F}$ -algebras of factorial association schemes, Preprint (2024), <https://doi.org/10.48550/arXiv.2403.08537>.
- [11] G. Karpilovsky, The Jacobson Radical of Group Algebras, North-Holland Publishing Co., Amsterdam, 1987.
- [12] S. Kageyama, Y. Miao, Two classes of  $q$ -ary codes based on group divisible association schemes, Discrete Math. **195** (1999), 269-276.
- [13] F. Levstein, C. Maldonado, D. Penazzi, The Terwilliger algebra of a Hamming scheme  $H(d, q)$ , European J. Combin. **27** (2006), 1-10.
- [14] F. Levstein, C. Maldonado, The Terwilliger algebra of the Johnson schemes, Discrete Math. **307** (2007), 1621-1635.
- [15] B. Lv, C. Maldonado, K. Wang, More on the Terwilliger algebra of Johnson schemes, Discrete Math. **328** (2014), 54-62.
- [16] A. Munemasa, An application of Terwilliger's algebra, Unpublished preprint (1993), <http://www.math.is.tohoku.ac.jp/~munemasa/unpublished.html>.
- [17] P. Terwilliger, The subconstituent algebra of an association scheme. I, J. Algebraic Combin. **1** (1992), 363-388.
- [18] P. Terwilliger, The subconstituent algebra of an association scheme. II, J. Algebraic Combin. **2** (1993), 73-103.
- [19] P. Terwilliger, The subconstituent algebra of an association scheme. III, J. Algebraic Combin. **2** (1993), 177-210.
- [20] M. Tomiyama, N. Yamazaki, The subconstituent algebra of a strongly regular graph, Kyushu J. Math. **48** (1994), 323-334.
- [21] Y. Watanabe, The generalized wreath product of triply-regular association schemes, The 35th Symposium on Algebraic Combinatorics (Proceedings) (2019), 109-114.
- [22] P.-H. Zieschang, Theory of Association Schemes, Springer Monogr. Math., Springer-Verlag, Berlin, 2005.

(Y. Jiang) SCHOOL OF MATHEMATICAL SCIENCES, ANHUI UNIVERSITY (QINGYUAN CAMPUS), NO. 111, JIULONG ROAD, HEFEI, 230601, CHINA

*Email address*, Y. Jiang: [jiangyu@ahu.edu.cn](mailto:jiangyu@ahu.edu.cn)