

**LAWVERE’S FOURTH OPEN PROBLEM:
LEVELS IN THE TOPOS OF SYMMETRIC SIMPLICIAL SETS**

RYUYA HORA AND YUKI MAEHARA

ABSTRACT. In the topos of simplicial sets, it makes sense to ask the following question about a given natural number n : *what is the minimum value m such that n -skeletality implies m -coskeletality?* This is an instance of the *Aufhebung* relation in the sense of Lawvere, who introduced this notion for an arbitrary Grothendieck topos \mathcal{E} in place of \mathbf{sSet} , and levels/essential subtopoi in place of dimensions.

We compute this *Aufhebung* relation for the topos of symmetric simplicial sets. In particular, we show that it is given by $2l - 1$ for the level labelled by $l \geq 3$, which coincides with the previously known case of simplicial sets. This result provides a solution to the fourth of the seven open problems in topos theory posed by Lawvere in 2009.

CONTENTS

1. Introduction	1
1.1. Levels and <i>Aufhebung</i> in a topos	1
1.2. The open problem and its history	2
1.3. Contribution	3
Acknowledgement	3
2. Symmetric sets	3
2.1. Definition and examples of symmetric sets	3
2.2. EZ-decomposition and EZ-congruence	4
2.3. Levels in the topos of symmetric sets	6
2.4. Lower bound for the <i>Aufhebung</i>	10
3. Computing the <i>Aufhebung</i>	12
3.1. Preparation: Propagative graphs	12
3.2. Reduction at points	14
3.3. Reduction graph	17
3.4. Candidate filler and its compatibility with \overline{P}	18
3.5. Compatibility with \underline{P}	22
3.6. Conclusion	22
References	23

1. INTRODUCTION

1.1. **Levels and *Aufhebung* in a topos.** As outlined in [Law06], the concept of dimension in a “topos of spaces” may be captured by a special class of its subtopoi called *levels*. A level of a Grothendieck topos \mathcal{E} is an adjoint

Date: March 6, 2025.

2020 Mathematics Subject Classification. 18F10, 18B25.

Key words and phrases. Topos, *Aufhebung*, level, symmetric simplicial sets, the Boolean algebra classifier, graph.

triple $l_! \dashv l^* \dashv l_* : \mathcal{L} \rightarrow \mathcal{E}$ with fully faithful $l_!$ and l_* . In other words, a level of \mathcal{E} is an essential subtopos of \mathcal{E} . In [KL89], Kelly and Lawvere showed that the levels of a fixed Grothendieck topos form a complete lattice under the natural ordering of subtopoi. In the case of simplicial sets $\mathcal{E} = \mathbf{sSet}$, the notion of levels coincides with the usual notion of dimensions $-\infty < 0 < 1 < \dots < \infty$. The familiar notion of n -skeletality can be extended to an arbitrary level $l_! \dashv l^* \dashv l_* : \mathcal{L} \rightarrow \mathcal{E}$, where we call an object X *l-skeletal* if the counit $l_! l^* X \rightarrow X$ is an isomorphism. Similarly, an object X is *l-coskeletal* if the unit $X \rightarrow l^* l_* X$ is an isomorphism.

This paper is primarily concerned with the *way above* relation between levels. Given two levels $l_1 \geq l_0$ of a topos \mathcal{E} , we say that l_1 is way above l_0 if every l_0 -skeletal object is l_1 -coskeletal. The *Aufhebung* of a level l , if it exists, is the minimum level way above l . The *Aufhebung* of levels has been computed for some topoi, including graphic topoi [Law89, Law91], the topos of ball complexes [Roy97], and the topoi of simplicial and cubical sets [Zak86, KRRZ11].

The purpose of this paper is to compute the *Aufhebung* for the presheaf topos $\mathbf{PSh}(\mathbf{F})$, where \mathbf{F} is the category of non-empty finite sets, which has long been an open problem in topos theory as we now review.

1.2. The open problem and its history. The topos $\mathbf{PSh}(\mathbf{F})$ has attracted interest under two contrasting names that reflect its logical and geometric aspects respectively: *the Boolean algebra classifier/the classifier of non-trivial Boolean algebras* [Law88, Roy97, Law09, Men19, Men24], a name derived from categorical logic, and the category of *symmetric simplicial sets/ensembles simpliciaux symétriques* [Gra01, RT03, Cis06, RRV09, HL25], a term rooted in combinatorial topology. Being two aspects of the same topos, of course these notions are interconnected; their relationship is described in [Law88] through a non-trivial Boolean algebra structure on the infinite-dimensional sphere.

The *Aufhebung* of this topos $\mathbf{PSh}(\mathbf{F})$ has been of interest at least since 1988 [Law88, Roy97, Men19, Men24] from the viewpoint of dimension theory. Its facet as a problem concerning “complexity of automata” was pointed out in [Law04]. This paper adopts the latter viewpoint, but instead of automata, we will think of it as a problem concerning complexity of a variant of Joyal’s *combinatorial species* [Joy81]. Accordingly, the notions of skeletality and coskeletality are interpreted as follows.

- Skeletality of M measures the extent to which an arbitrary M -structure may be recovered from small *quotient structures*. (See Figure 1.)
- Coskeletality of M measures the extent to which an arbitrary M -structure may be recovered from small *substructures*. (See Figure 2.)

Thus, the *Aufhebung* is the quantitative interplay between these dual complexity measures.

In 2009, Lawvere re-emphasised the significance of the *Aufhebung* of this topos by including it as Problem 4 in his list of seven open problems in topos theory:

[Law09] [...] There is another fundamental topos related to classical constructions and combinatorial topology, namely the Boolean algebra classifier that consists of presheaves on the category of finite non-empty sets. [...] What is, in combinatorial or number-theoretic terms, the way below relation for this basic topos?

For more explanations on related topics, see the early article [Law88], the recent paper [Men24], and the papers cited therein.

1.3. **Contribution.** The present paper proves that, in the topos $\mathbf{PSh}(\mathbf{F})$, the Aufhebung of l is given by

$$\left\{ \begin{array}{ll} 0 & (l = -\infty) \\ 1 & (l = 0) \\ 3 & (l = 1) \\ 4 & (l = 2) \\ 2l - 1 & (l \geq 3) \\ \infty & (l = \infty), \end{array} \right.$$

which coincides with the case of simplicial sets proven in [KRRZ11] for $l \geq 3$. In fact, our proof, and in particular the way we utilise the Pigeonhole Principle, is inspired by this simplicial counterpart too.

A notable difference between our approach and that in [KRRZ11] is that we frame some of the combinatorics as a purely graph-theoretic phenomenon (Proposition 3.1.4), which provides an insight from a new angle on why the Aufhebung doubles the dimension.

This use of graph theory is our response to the following phenomenon, which forced us to pay close attention to relationships between points in the underlying set A of a given M -structure $x \in M(A)$: when proving coskeletality of the (suitably skeletal) symmetric simplicial set M , we need *two* substructures to recover the given M -structure (see the construction of f in Lemma 3.4.1) whereas only *one* was needed in the simplicial case (see [KRRZ11, Proposition 3.14] where the filler is simply given by $c_m \sigma^m$). The reason for this contrast can be traced back to the fact that, given a morphism $\alpha: [k] \rightarrow [n]$ in the simplex category Δ and $m \in [k]$, we have

$$m = \min \left\{ i \in [k] \mid \alpha^{-1}(\alpha(i)) \neq \{i\} \right\} \implies \alpha(m) = \alpha(m+1)$$

while we have no such control over functions in \mathbf{F} .

Acknowledgement. We would like to thank Matias Menni for his encouraging advice on this problem. We are also grateful to the Kanda Satellite Lab at the National Institute of Informatics for providing a space for research discussions.

The first-named author would like to thank his supervisor Ryu Hasegawa for helpful discussions and suggestions. He was supported by JSPS KAKENHI Grant Number JP24KJ0837 and FoPM, WINGS Program, the University of Tokyo.

The second-named author gratefully acknowledges the support of JSPS Research Fellowship for Young Scientists and JSPS KAKENHI Grant Number JP23KJ1365.

2. SYMMETRIC SETS

The main result of this paper concerns the *Aufhebung* of the topos of *symmetric sets*. The primary purpose of this section is to recall these two notions. We also construct a family of rather simple symmetric sets which provides a (sharp) lower bound for the Aufhebung.

2.1. Definition and examples of symmetric sets.

Definition 2.1.1. We will write \mathbf{F} for the category of non-empty finite sets and all functions. The category of *symmetric (simplicial) sets* and *symmetric (simplicial) maps* is its presheaf category $\mathbf{PSh}(\mathbf{F}) = [\mathbf{F}^{\text{op}}, \mathbf{Set}]$.

One can think of symmetric sets either

- as a variant of simplicial sets equipped with an action of the symmetric groups, or
- as a variant of Joyal's combinatorial species [Joy81].

Although we will employ some key concepts motivated by the former viewpoint (such as the EZ-decomposition: Definition 2.2.1), our arguments will be framed according to the latter. In particular, given a symmetric set M and a non-empty finite set A , we will refer to elements of $M(A)$ as M -structures on A . Each M -structure $x \in M(B)$ can be pulled back along an arbitrary function $\alpha: A \rightarrow B$, and we denote the resulting M -structure by $x\alpha \in M(A)$.

Below are some typical examples of symmetric sets.

Example 2.1.2 (Graphs). The symmetric set of *graphs* is the contravariant functor $\text{Graph}: \mathbf{F}^{\text{op}} \rightarrow \mathbf{Set}$ that sends a non-empty finite set A to the set of all undirected graphs whose vertex set is A , i.e., the set of all pairs $(A, E \subset \mathcal{P}_2(A))$ where $\mathcal{P}_2(A)$ is the set of all 2-element subsets of A . For a graph $x = (B, E \subset \mathcal{P}_2(B)) \in \text{Graph}(B)$ and a function $\alpha: A \rightarrow B$, the pullback $x\alpha \in \text{Graph}(A)$ is defined to be the graph $(A, \{e \in \mathcal{P}_2(A) \mid \alpha(e) \in E\})$. See Figure 1 for instance. (In what follows, Graph will be our default example when visualising concepts around symmetric sets.)

Example 2.1.3 (Equivalence relations). The symmetric set of *equivalence relations* Eq is defined by

$$\text{Eq}(A) := \{\sim \subset A^2 \mid \sim \text{ is an equivalence relation}\}.$$

A variant of this example will be used to give a lower bound for the Aufhebung (Lemma 2.4.5).

Example 2.1.4 (Representable symmetric set = colouring). For a non-negative integer k , we define the symmetric set Δ^k to be the presheaf $\mathbf{F}(-, S)$ represented by an $(n+1)$ -element set S . Geometrically, one can think of Δ^k as a standard k -simplex. Combinatorially, a Δ^k -structure on a non-empty finite set A is simply a function $A \rightarrow S$, which may be thought of as an S -colouring of A .

2.2. EZ-decomposition and EZ-congruence. Either by direct combinatorics or using [Cam23, Proposition 4.2 and Theorem 5.6], one can verify that the category \mathbf{F} is an *EZ-category* in the sense of [BM11, Definition 6.7] (which allows non-trivial automorphisms, unlike e.g. [Cis19, Definition 1.3.1]). Thus, we are naturally led to the following notions.

Definition 2.2.1 (EZ-decomposition). Let M be a symmetric set, A be a non-empty finite set, and $x \in M(A)$ be an M -structure.

- A *decomposition* of x is a pair (α, y) consisting of a morphism $\alpha: A \rightarrow B$ in \mathbf{F} and $y \in M(B)$ such that $x = y\alpha$.
- We say x is *degenerate* if it admits a decomposition (α, y) where α is a non-invertible surjection. Otherwise x is said to be *non-degenerate*.
- A decomposition (α, y) of x is called an *EZ-decomposition* if α is a surjection and y is non-degenerate.

Example 2.2.2. An EZ-decomposition of a graph (regarded as an M -structure on a seven-element set for $M = \text{Graph}$ of Example 2.1.2) is visualised in Figure 1. (The colours indicate the associated *EZ-congruence*, which will be defined in Definition 2.2.4 together with the term *mass*.)

Each M -structure admits an essentially unique EZ-decomposition in the following sense.

Proposition 2.2.3 (Uniqueness of EZ-decomposition [BM11, Proposition 6.9.]). *Let M be a symmetric set, A be a non-empty finite set, and $x \in M(A)$ be an M -structure. Then x admits an EZ-decomposition. Moreover, for any two EZ-decompositions $(\alpha: A \twoheadrightarrow B, y)$ and $(\alpha': A \twoheadrightarrow B', y')$ of x , there exists a (necessarily unique) bijection*

Then we have $x = y\alpha = y\mu\epsilon = z(\beta\epsilon)$ and the rightmost expression provides an EZ-decomposition of x . Since β is a surjection and μ is an injection, we have

$$|B| \geq |C| \geq |D| = \text{mass}(x).$$

Moreover, the two equalities hold if and only if both μ and β are invertible, or equivalently, if and only if α is a surjection (so that we may take $\mu = \text{id}$) and y is non-degenerate. \square

Lemma 2.2.6. *Let M be a symmetric set, $\phi: A \rightarrow A'$ be a morphism in \mathbf{F} , $a, b \in A$, and x be an M -structure on A' . If $\phi(a) \sim_x \phi(b)$ then we have $a \sim_{x\phi} b$.*

Proof. Let $x = y\alpha$ be an EZ-decomposition of x . Assume $\phi(a) \sim_x \phi(b)$, which is equivalent to the equation $\alpha\phi(a) = \alpha\phi(b)$. We wish to prove that we have $\beta(a) = \beta(b)$ for some (and equivalently all) EZ-decomposition $x\phi = z\beta$ of $x\phi$.

Let $\alpha\phi = \mu\epsilon$ be an epi-mono factorisation of $\alpha\phi$, and let $y\mu = z\gamma$ be an EZ-decomposition of $y\mu$.

$$\begin{array}{ccc}
 A & \xrightarrow{\phi} & A' \\
 \downarrow \epsilon & & \downarrow \alpha \\
 B & \xrightarrow{\mu} & B' \\
 \downarrow \gamma & & \\
 C & & \\
 \beta \curvearrowright & & \\
 & &
 \end{array}
 \quad
 \begin{array}{ccc}
 x\phi & \longleftarrow & x \\
 \uparrow & & \uparrow \\
 y\mu & \longleftarrow & y \\
 \uparrow & & \\
 z & &
 \end{array}$$

Then we have

$$x\phi = y\alpha\phi = y\mu\epsilon = z(\gamma\epsilon)$$

and the rightmost expression is an EZ-decomposition of $x\phi$. It thus suffices to prove $\gamma\epsilon(a) = \gamma\epsilon(b)$, which indeed holds because

$$\mu\epsilon(a) = \alpha\phi(a) = \alpha\phi(b) = \mu\epsilon(b)$$

and μ is monic. \square

Notice that the converse of the above lemma does not hold in general. The lower half of Figure 6 provides a counterexample.

Lemma 2.2.7. *Let M be a symmetric set, $\phi: A \rightarrow A'$ be a morphism in \mathbf{F} , and x be an M -structure on A' . Then we have $\text{mass}(x) \geq \text{mass}(x\phi)$.*

Proof. This is an immediate corollary of Lemma 2.2.6. \square

2.3. Levels in the topos of symmetric sets. For a non-negative integer $l \geq 0$, we write \mathbf{F}_l for the full subcategory of \mathbf{F} consisting of the sets of size less than or equal to $l+1$. Geometrically, \mathbf{F}_l is the category of standard symmetric simplices of dimension less than or equal to l . We also write $\mathbf{F}_{-\infty}$ for the empty category, and \mathbf{F}_{∞} for \mathbf{F} itself.

The following proposition is mentioned in [Law88]. See also [Men24, Corollary 3.3, Example 6.3].

Proposition 2.3.1 (Classification of levels [Law88]). *The complete lattice of levels in the topos $\mathbf{PSh}(\mathbf{F})$ is isomorphic to the total order*

$$-\infty < 0 < 1 < 2 < \dots < \infty,$$

where the level labelled by ¹ l is the adjoint triple

$$\mathbf{PSh}(\mathbf{F}_l) \begin{array}{c} \xrightarrow{\text{Lan}_{\iota_l}} \\ \xleftarrow{\iota_l^*} \\ \xrightarrow{\text{Ran}_{\iota_l}} \end{array} \mathbf{PSh}(\mathbf{F})$$

induced by the inclusion $\iota_l: \mathbf{F}_l \rightarrow \mathbf{F}$.

Remark 2.3.2 (All subtopoi are levels). As mentioned in [Law88], in this special topos $\mathbf{PSh}(\mathbf{F})$, all subtopoi are in fact of the form of Proposition 2.3.1 and thus levels. This fact can be seen as an instance of either [Men24, Corollary 3.3] or [Mar24, Corollary 4].

Let us now recall the notions of *skeleton* and *coskeleton*.

Definition 2.3.3 (Skeleton and coskeleton). For a level of $\mathbf{PSh}(\mathbf{F})$ labelled by $l \in \{-\infty, 0, 1, \dots, \infty\}$, the induced comonad on $\mathbf{PSh}(\mathbf{F})$ is denoted by sk_l , and the induced monad is denoted by cosk_l . A symmetric set M is said to be *l-skeletal* if the counit $\text{sk}_l M \rightarrow M$ is invertible, and is said to be *l-coskeletal* if the unit $M \rightarrow \text{cosk}_l M$ is invertible.

Proposition 2.3.4 (Skeletality). For a symmetric set M , its *l-skeleton* $\text{sk}_l M$ is the symmetric subset $\epsilon_M: \text{sk}_l M \rightarrow M$ consisting of the M -structures whose mass is less than or equal to $l + 1$. In other words, for any non-empty finite set A , we have

$$(\text{sk}_l M)(A) = \{x \in M(A) \mid \text{mass}(x) \leq l + 1\} \subset M(A).$$

In particular, a symmetric set M is *l-skeletal* if and only if $\text{mass}(x) \leq l + 1$ for any non-empty finite set A and any M -structure $x \in M(A)$.

Proof. This is a special case of [BM11, Corollary 6.10]. See also [Men19, Proposition 2.1, Example 2.2]. \square

To provide a concrete description of *l-coskeletality*, we need the notion of *cycle*, whose definition in turn requires the following notation. For a finite set A with $|A| > 1$ and an element $a \in A$, we write δ^a for the canonical embedding

$$\delta^a: A \setminus \{a\} \rightarrow A.$$

We will not notationally distinguish δ^a 's with different codomains. For example, we write $\delta^a \delta^b = \delta^b \delta^a$ to express the commutativity of the square

$$\begin{array}{ccc} A \setminus \{a, b\} & \xrightarrow{\delta^a} & A \setminus \{b\} \\ \downarrow \delta^b & & \downarrow \delta^b \\ A \setminus \{a\} & \xrightarrow{\delta^a} & A. \end{array}$$

Definition 2.3.5 (Cycle). For a positive integer $k > 0$, a *k-cycle* in a symmetric set M is a pair $(P, \{c_p\}_{p \in P})$ of a $(k + 1)$ -element set P and a family of M -structures $c_p \in M(P \setminus \{p\})$ that satisfies *the cycle equation*

$$c_p \delta^q = c_q \delta^p \in M(P \setminus \{p, q\})$$

for any distinct elements $p, q \in P$ (except for the case $k = 1$, in which case we do not enforce the cycle equation because $P \setminus \{p, q\}$ would be empty). A *filler* of a cycle $(P, \{c_p\}_{p \in P})$ is an M -structure $f \in M(P)$ that satisfies $f \delta^p = c_p$ for every $p \in P$. We say $(P, \{c_p\}_{p \in P})$ is *unfillable* if it admits no fillers.

See Figure 2 for a visualisation of cycle filling in the case where $M = \text{Graph}$.

¹We avoid simply calling it “level l ” because the terms “level 0” and “level 1” have specific meanings defined by the Aufhebung relation.




FIGURE 2. A filler of a 6-cycle in Graph

Lemma 2.3.6 (Universality of $\text{sk}_l \Delta^k$). *Let k and l be non-negative integers, and let P be a $(k+1)$ -element set representing $\Delta^k \cong \mathbf{F}(-, P)$. Then for each symmetric set M , there is a bijection between*

- symmetric maps $f: \text{sk}_l \Delta^k \rightarrow M$, and

- families of M -structures $\{x_S \in M(S)\}_{S \in \mathcal{P}_{\leq l+1}(P)}$ such that $x_S * (S \leftarrow S') = x_{S'}$ for all pairs $S' \subset S$ in $\mathcal{P}_{\leq l+1}(P)$ (where $\mathcal{P}_{\leq l+1}(P)$ is the set of non-empty subsets of P of size at most $l+1$)

which is natural in M . Consequently, for a positive integer $k > 0$, the symmetric maps $\text{sk}_{k-1}\Delta^k \rightarrow M$ are in bijection with the k -cycles in M .

Proof. Due to Proposition 2.3.4, $\text{sk}_l(\Delta^k)(A)$ may be identified with the set of functions $P \leftarrow^\alpha A$ with $|\text{Im}(\alpha)| \leq l+1$. Given a symmetric map $f: \text{sk}_l\Delta^k \rightarrow M$, we can construct a family $\{x_S \in M(S)\}_{S \in \mathcal{P}_{\leq l+1}(P)}$ by

$$x_S := f_S(P \leftarrow S) \in M(S).$$

The naturality of f implies that the resulting family satisfies the required compatibility condition.

Conversely, given an arbitrary family $\{x_S \in M(S)\}_{S \in \mathcal{P}_{\leq l+1}(P)}$ that satisfies the compatibility condition, we define a symmetric map $f: \text{sk}_l\Delta^k \rightarrow M$ by

$$f_A(P \leftarrow^\alpha A) := x_{\text{Im}(\alpha)} * (\text{Im}(\alpha) \leftarrow^\alpha A).$$

It is straightforward to verify that this indeed yields a symmetric map, and that it is moreover inverse to the construction in the previous paragraph. The naturality is also clear.

The last statement is a consequence of the observation that, in the case $l = k - 1$, the compatible families $\{x_S \in M(S)\}_{S \in \mathcal{P}_{\leq k}(P)}$ are in bijection with the k -cycles via $c_P := x_{P \setminus \{p\}}$. \square

Proposition 2.3.7 (Coskeletality). *Let M be a symmetric set and l be a non-negative integer. Then the following are equivalent.*

- (1) M is l -coskeletal.
- (2) For any positive integer k greater than l , every k -cycle in M has a unique filler.

Proof. We consider two additional conditions (1'), (2') as follows.

- (1') M is right orthogonal to the counit $\epsilon_{\Delta^n}: \text{sk}_l\Delta^n \rightarrow \Delta^n$ for any non-negative integer $n \geq 0$.
- (2') M is right orthogonal to the counit $\epsilon_{\Delta^k}: \text{sk}_{k-1}\Delta^k \rightarrow \Delta^k$ for any $k > l$.

We have (1) \iff (1') because

$$\begin{aligned} (1) &\iff \text{the unit } \eta_M: M \rightarrow \text{cosk}_l M \text{ is an isomorphism} \\ &\iff \text{for any } n \geq 0, (\eta_M)_*: \mathbf{PSh}(\mathbf{F})(\Delta^n, M) \rightarrow \mathbf{PSh}(\mathbf{F})(\Delta^n, \text{cosk}_l M) \text{ is a bijection} \\ &\iff \text{for any } n \geq 0, (\epsilon_{\Delta^n})^*: \mathbf{PSh}(\mathbf{F})(\Delta^n, M) \rightarrow \mathbf{PSh}(\mathbf{F})(\text{sk}_l\Delta^n, M) \text{ is a bijection} \\ &\iff (1'). \end{aligned}$$

The equivalence (2) \iff (2') follows from Lemma 2.3.6.

Now we prove (1) \implies (2'). Assume (1) and fix $k > l$. Note that, since we have the inclusion $\mathbf{F}_l \subset \mathbf{F}_{k-1}$, the l -coskeletal symmetric set M is also $(k-1)$ -coskeletal. In particular, using an instance of the equivalence (1) \iff (1'), we can deduce that M is right orthogonal to the counit $\epsilon_{\Delta^k}: \text{sk}_{k-1}\Delta^k \rightarrow \Delta^k$ as desired.

It remains to prove (2') \implies (1'). So assume (2') and consider a symmetric map $\text{sk}_l\Delta^n \rightarrow M$ with $n > l$ (note that (1') is trivial in the case $n \leq l$). By Lemma 2.3.6, we may identify this symmetric map with a compatible family $\{x_S\}_{S \in \mathcal{P}_{\leq l+1}(P)}$ of M -structures where P is some $(n+1)$ -element set. The assumption (2') in the case of $k = l+1$ implies that there is a unique extension of this compatible family to $\{x_S\}_{S \in \mathcal{P}_{\leq l+2}(P)}$ indexed by the non-empty subsets of P of size at most $l+2$ (as opposed to $l+1$). By repeating this process, we eventually obtain a unique extension $\{x_S\}_{S \in \mathcal{P}_{\leq n+1}(P)}$, which corresponds to a symmetric map $\Delta^n = \text{sk}_n\Delta^n \rightarrow M$. \square

Remark 2.3.8 (Combinatorial intuition of coskeletality). To clarify the intuition behind coskeletality, which may be somewhat difficult to grasp, let us give an informal explanation. A symmetric set M is l -coskeletal if, for

- any non-empty finite set A and

- any compatible family of M -structures on all subsets of A with at most $l + 1$ elements,

there exists a unique way to glue them together to an M -structure on the entire set A . This is an example of the sheaf condition. In other words, the l -coskeletality of M means that (both the existence and the equality of) M -structures on A are completely determined by their *substructures* on subsets $B \subset A$ of size at most $l + 1$, while the l -skeletality means that they are determined by their *quotient structures*.

For certain examples, it makes sense to think of the coskeletality as measuring how many variables are needed in order to describe what an M -structure is. For instance, the symmetric set Eq of equivalence relations is 2-coskeletal (but not 1-coskeletal), reflecting the fact that the transitivity condition $x \sim y \wedge y \sim z \implies x \sim z$ involves $(2 + 1) = 3$ variables. Another such example is the representable Δ^n , which is the symmetric set of $\{0, 1, \dots, n\}$ -valued functions (without any additional structures nor conditions). Since a function is completely determined by its values at individual points, this symmetric set is 0-coskeletal.

2.4. Lower bound for the Aufhebung.

Definition 2.4.1 (Aufhebung). For a level of the topos $\mathbf{PSh}(\mathbf{F})$ labelled by $l \in \{-\infty, 0, 1, \dots, \infty\}$, its *Aufhebung* is the least level $l \leq a_l \in \{-\infty, 0, 1, \dots, \infty\}$ such that every l -skeletal symmetric set is a_l -coskeletal.

Let us recall the first two Aufhebung $a_{-\infty}$ and a_0 , which are already known. See [Law06, Section 2] for more explanations and motivating ideas around the Aufhebung $a_{-\infty}$ and a_0 .

Proposition 2.4.2. $a_{-\infty} = 0$.

Proof. A symmetric set M is $(-\infty)$ -skeletal if and only if M is the initial object, i.e., the empty symmetric set. Since the empty symmetric set is 0-coskeletal but not $(-\infty)$ -coskeletal, we have $a_{-\infty} = 0$. \square

More generally, if $\mathcal{E}_{\rightarrow}$ is a level of \mathcal{E} , then it coincides with the Aufhebung $a_{-\infty}$ since it is the least dense subtopos (see [Joh02, A.4.5]).

The value of the Aufhebung a_0 was first established in [Law88]. The paper [Men19] contains a detailed explanation on “one-dimensionality” and also on the particular example of the topos $\mathbf{PSh}(\mathbf{F})$.

Proposition 2.4.3 ([Law88], [Men19]). $a_0 = 1$.

Proof. A symmetric set M is 0-skeletal if and only if M is a small coproduct of copies of the terminal object, i.e., if and only if M is discrete. We can easily see that the two-point discrete symmetric set is not 0-coskeletal, which implies $a_0 \geq 1$. It suffices to prove that any discrete M is 1-coskeletal. Since any symmetric map with a discrete codomain must be constant on each connected component, the required orthogonality follows from the fact that $\text{sk}_1 \Delta^n \rightarrow \Delta^n$ induces a bijection between their connected components. \square

In the rest of the present subsection, we will give a lower bound for the Aufhebung.

Definition 2.4.4 ($\text{Eq}_{\leq l}$ and $\text{Eq}_{=l+1}$). For $l \geq 1$, we define the symmetric set $\text{Eq}_{\leq l}$ to be the $(l - 1)$ -skeleton of Eq (See Example 2.1.3)

$$\text{Eq}_{\leq l} := \text{sk}_{l-1} \text{Eq},$$

and the symmetric set $\text{Eq}_{=l+1}$ by the pushout diagram

$$\begin{array}{ccc} \text{Eq}_{\leq l} & \xrightarrow{\quad} & \text{Eq}_{\leq l+1} \\ \downarrow ! & \lrcorner & \downarrow \\ 1 & \longrightarrow & \text{Eq}_{=l+1}. \end{array}$$

More concretely, $\text{Eq}_{\leq l}(A)$ is the set of all partitions of A into at most l non-empty parts,

$$\text{Eq}_{\leq l}(A) = \{\sim \in \text{Eq}(A) \mid |A/\sim| \leq l\}.$$

Similarly, $\text{Eq}_{=l+1}(A)$ is the set of all partitions of A into exactly $l+1$ non-empty parts, with an “error message” $*$:

$$\text{Eq}_{=l+1}(A) = \{\sim \in \text{Eq}(A) \mid |A/\sim| = l+1\} \sqcup \{*\}.$$

If pulling back an equivalence relation results in fewer than $l+1$ parts then it instead returns an error message, and pulling back an error message also results in an error message.

It is easy to see that $\text{Eq}_{=l+1}$ is l -skeletal, and the following unfillable cycle provides the desired lower bound.

Lemma 2.4.5. *Let $l \geq 1$. Then $\text{Eq}_{=l+1}$ has an unfillable k -cycle where*

$$k = \begin{cases} 3 & (l = 1) \\ 4 & (l = 2) \\ 2l - 1 & (l \geq 3). \end{cases}$$

Proof. First, we construct an unfillable $(l+2)$ -cycle on the $(l+3)$ -element set $P = \mathbb{Z}/(l+3)\mathbb{Z}$. (This proves the cases $l = 1, 2$.) For each $i \in \mathbb{Z}/(l+3)\mathbb{Z}$, define an equivalence relation c_i on $P \setminus \{i\}$ by

$$ac_i b \iff (a+1 = i = b-1) \vee (b+1 = i = a-1) \vee (a = b).$$

In other words, c_i partitions $P \setminus \{i\}$ into a single two-element set $\{i-1, i+1\}$ and l singletons. Since there are exactly $l+1$ equivalence classes, we may regard c_i as an element of $\text{Eq}_{=l+1}(P \setminus \{i\})$. These $\text{Eq}_{=l+1}$ -structures satisfy the cycle equations because, for distinct $i, j \in \mathbb{Z}/(l+3)\mathbb{Z}$,

- if $i = j \pm 1$ then both $c_i \delta^j$ and $c_j \delta^i$ partition $P \setminus \{i, j\}$ into $l+1$ singletons, and
- otherwise both $c_i \delta^j$ and $c_j \delta^i$ are the error message $*$.

It is straightforward to see that this cycle is unfillable.

Next, we construct an unfillable $(2l-1)$ -cycle on a $2l$ -element set P for $l \geq 3$. Let us fix a decomposition $P = A \sqcup B$ with $|A| = |B| = l$. For $p \in P$, define an equivalence relation c_p on $P \setminus \{p\}$ by

$$xc_p y \iff (x \in A \wedge y \in A \wedge p \in A) \vee (x \in B \wedge y \in B \wedge p \in B) \vee (x = y).$$

In other words, c_p partitions $P \setminus \{p\}$ into

- a single $(l-1)$ -element set $A \setminus \{p\}$ and l singletons if $p \in A$, and
- a single $(l-1)$ -element set $B \setminus \{p\}$ and l singletons if $p \in B$.

These $\text{Eq}_{=l+1}$ -structures satisfy the cycle equations because, for distinct $p, q \in P$,

- if $p, q \in A$ then both $c_p \delta^q$ and $c_q \delta^p$ partition $P \setminus \{p, q\}$ into a single $(l-2)$ -element set $A \setminus \{p, q\}$ and l singletons,
- similarly in the case $p, q \in B$, and
- otherwise (that is, if either $p \in A$ and $q \in B$, or $p \in B$ and $q \in A$) both $c_p \delta^q$ and $c_q \delta^p$ are the error message $*$.

Again, it is straightforward to see that this cycle is unfillable. □

These cycles will be visualised in Figure 8 after defining the notion of reduction graph (Definition 3.3.2).

Remark 2.4.6 (Comparison with [KRRZ11]). The second family of unfillable cycles in the proof of Lemma 2.4.5 is inspired by [KRRZ11, Example 3.20]. More precisely, our construction essentially extracts the (simplicial version of the) EZ-congruence from that example.

Proposition 2.4.7 (Lower bound). *For $l \geq 1$, we have*

$$a_l \geq \begin{cases} 3 & (l = 1) \\ 4 & (l = 2) \\ 2l - 1 & (l \geq 3). \end{cases}$$

3. COMPUTING THE AUFHEBUNG

We gave a lower bound for the Aufhebung in Proposition 2.4.7. The aim of this section is to prove that the Aufhebung is bounded above by, and hence equal to, those values.

3.1. Preparation: Propagative graphs. In this subsection, we introduce a purely graph-theoretic phenomenon (Proposition 3.1.4), which will eventually provide a sharp upper bound for the Aufhebung (Proposition 3.4.4). Let us start by considering the following notion, which may be visualised as Figure 3.

Definition 3.1.1. Given an undirected graph $G = (V, E \subset \mathcal{P}_2(V))$, we write $\Phi = \Phi_G: \mathcal{P}(V) \rightarrow \mathcal{P}(V)$ for the operation given by

$$\Phi S = S \cup \{v \in V \mid \exists s_0, s_1 \in S (s_0 \neq s_1 \wedge \{s_0, v\}, \{s_1, v\} \in E)\}.$$

We call G *propagative* if, for any $S \subset V$ with $|S| = 2$, there exists $k \geq 1$ such that $\Phi^k S = V$.




FIGURE 3. An example of a propagative graph

Puzzle 3.1.2. Fix $n > 2$. Find the minimum m such that any undirected graph $G = (V, E)$ with

- $|V| = n$, and
- each $v \in V$ has degree at least m

is necessarily propagative.

The following example demonstrates that such m must be strictly greater than $\lfloor \frac{n}{2} \rfloor$.

Example 3.1.3. No bipartite graph (with more than 2 vertices) is propagative because taking one vertex from each part yields a two-element subset S with $\Phi S = S$. In particular, for any $n > 2$, there is a non-propagative n -vertex graph such that each vertex has degree at least $\lfloor \frac{n}{2} \rfloor$ (Figure 4).

This bound is in fact sharp; that is, the answer to Puzzle 3.1.2 is $m = \lfloor \frac{n}{2} \rfloor + 1$.

Proposition 3.1.4. *Let $n > 2$ and let $G = (V, E)$ be an undirected graph with n vertices. If every vertex in G has degree greater than $\frac{n}{2}$, then G is propagative.*




FIGURE 4. Bipartite graphs are never propagative.

Proof. (This proof is visualised in Figure 5.) Let m be the minimum degree of the vertices in G . We will prove the contrapositive of the proposition. More precisely, we will assume that the graph G is not propagative, and prove $2m \leq n$.

Since G is not propagative, we can find a proper subset $S \subsetneq V$ such that $|S| \geq 2$ and $\Phi S = S$. Observe that the equation $\Phi S = S$ is equivalent to the condition

(†) each vertex $v \in V \setminus S$ has at most one neighbour in S .

We claim that the cardinality $x := |S|$ satisfies

- (i) $2 \leq x \leq n - m$, and
- (ii) $x(x - 1) + (n - x) \geq xm$.

For the second inequality in (i), pick $v \in V \setminus S$. Then its degree d_v satisfies $d_v \geq m$ by the minimality of m , and it also satisfies

$$d_v \leq |V \setminus (S \cup \{v\})| + 1 = n - x$$

by (†). Combining these inequalities yields $x \leq n - m$. To obtain (ii), consider the sum of degrees of all elements in S . On the one hand, we can deduce from (†) that it is bounded above by

$$|S|(|S| - 1) + |V \setminus S| = x(x - 1) + (n - x).$$

On the other hand, the minimality of m provides a lower bound $|S|m = xm$. This completes the proof of the two inequalities (i) and (ii).

Since the function $f(t) = t(t-1) + (n-t) - tm$ is downward convex, the existence of a solution x to the inequalities (i) and (ii) implies the satisfaction of (ii) at one of the boundaries of (i). If it is satisfied at $x = 2$ then we have

$$n = 2(2 - 1) + (n - 2) \geq 2m.$$


In the other case, rearranging the resulting inequality

$$(n - m)(n - m - 1) + (n - (n - m)) \geq (n - m)m$$

yields

$$(n - 2m)(n - m - 1) \geq 0.$$

Since (i) implies $n - m - 1 > 0$, we conclude that $n \geq 2m$ holds in either case. \square

FIGURE 5. Illustration for $n = 7, m = 3$.

3.2. Reduction at points. In later subsections, where we will be dealing with highly degenerate M -structures x on a finite set A , it will be useful to keep track of which points of A may be discarded without losing essential information about x . This will be done using the following notion of *reduction*, and this subsection is devoted to proving various technical lemmas around it.

Lemma 3.2.1. *Let M be a symmetric set, A be a non-empty finite set, $a \in A$ be an element, and $x \in M(A)$ be an M -structure. Then the following conditions are equivalent.*

- (1) $\text{mass}(x\delta^a) = \text{mass}(x)$.
- (2) For any EZ-decomposition $x = y\alpha$ of x , the decomposition $x\delta^a = y(\alpha\delta^a)$ of $x\delta^a$ is an EZ-decomposition.
- (3) For some EZ-decomposition $x = y\alpha$ of x , the decomposition $x\delta^a = y(\alpha\delta^a)$ of $x\delta^a$ is an EZ-decomposition.
- (4) The \sim_x -class containing the element $a \in A$ is not a singleton.

Proof. Assume (1), and let $x = y\alpha$ be an EZ-decomposition of x . Then the decomposition $x\delta^a = y(\alpha\delta^a)$ realizes the mass $\text{mass}(x\delta^a) = \text{mass}(x)$, which implies (2) by Lemma 2.2.5.

Clearly (2) implies (3).

The condition (3) implies that the composite $\alpha\delta^a$ is surjective, which in turn implies (4).

Assuming (4), we can take $a \neq a' \in A$ such that $a \sim_x a'$. Fix an EZ-decomposition $x = y\alpha$, and write $\rho: A \rightarrow A \setminus \{a\}$ for the retraction of δ^a that maps a to a' . Then we have $\alpha(a) = \alpha(a')$, which implies $\alpha\delta^a\rho = \alpha$, which in turn implies $x\delta^a\rho = y\alpha\delta^a\rho = y\alpha = x$. Thus by Lemma 2.2.7, we have

$$\text{mass}(x) \geq \text{mass}(x\delta^a) \geq \text{mass}(x\delta^a\rho) = \text{mass}(x).$$

This completes the proof. □

We will borrow the following terminology from [KRRZ11].

Definition 3.2.2 (Reduction). For a symmetric set M , a non-empty finite set A , and an M -structure $x \in M(A)$, we say that $a \in A$ *reduces* x if a and x satisfy the equivalent conditions in Lemma 3.2.1.

Intuitively, a point $a \in A$ reduces an M -structure $x \in M(A)$ if $x\delta^a$ retains enough information about x ; Figure 6 provides an example of reduction/non-reduction in the case where $M = \text{Graph}$. In the rest of this subsection, we will prove lemmas which make this intuition precise.

Lemma 3.2.3 (Congruence lifting). *Let M be a symmetric set, A be a non-empty finite set, $a \in A$ be an element, and $x \in M(A)$ be an M -structure. If a reduces x , then the EZ-congruence associated to $x\delta^a$ is the restriction of that associated to x .*

Proof. This immediately follows from the second (or third) condition in Lemma 3.2.1. \square

Lemma 3.2.4 (Equality lifting: one point). *Let M be a symmetric set, A be a non-empty finite set, $a \in A$ be an element, and $x, y \in M(A)$ be M -structures. Suppose that*

- x and y have the same EZ-congruence,
- a reduces either (hence both) of x and y , and
- $x\delta^a = y\delta^a$.

Then we have $x = y$.

Proof. Let us write \sim for the common EZ-congruence of x and y . Then we can take EZ-decompositions $x = z\pi$ and $y = w\pi$ involving the same quotient map $\pi: A \rightarrow A/\sim$. The second assumption implies that $\pi\delta^a$ is surjective, whereas the third implies

$$z(\pi\delta^a) = x\delta^a = y\delta^a = w(\pi\delta^a).$$

We have thus obtained two EZ-decompositions of the same M -structure, so applying Proposition 2.2.3 yields an automorphism σ on A/\sim such that $\sigma\pi\delta^a = \pi\delta^a$ and $z = w\sigma$. But since $\pi\delta^a$ is surjective, the first of these two equalities implies that σ is the identity, which in turn implies $z = w$. This completes the proof. \square

Lemma 3.2.5 (Equality lifting: two points). *Let M be a symmetric set, A be a non-empty finite set, $a, b \in A$ be elements, and $x, y \in M(A)$ be M -structures. Suppose that*

- each of a and b reduces both x and y ,
- $x\delta^a = y\delta^a$, and
- $x\delta^b = y\delta^b$.

Then we have $x = y$.

Proof. By Lemma 3.2.4, it suffices to prove the equality between their EZ-congruences $\sim_x = \sim_y$. Combining the second and third assumptions with Lemma 3.2.3, we can deduce $c \sim_x d \iff c \sim_y d$ for all $c, d \in A$ except for the pair $\{c, d\} = \{a, b\}$. It thus remains to prove $a \sim_x b \iff a \sim_y b$.

Suppose $a \sim_x b$. If the \sim_x -class containing a and b contains a third element c , then we have

$$a \sim_{x\delta^b} c \sim_{x\delta^a} b \quad \text{or equivalently} \quad a \sim_{y\delta^b} c \sim_{y\delta^a} b,$$

which implies $a \sim_y b$ by Lemma 3.2.3. If this \sim_x -class is just $\{a, b\}$, then a is isolated with respect to $\sim_{x\delta^b} = \sim_{y\delta^b}$. However, a cannot be isolated with respect to $\sim_y \subset A^2$ because a reduces y . This implies $a \sim_y b$ as desired. \square

The following lemma is visualised in Figure 6 in the case of $M = \text{Graph}$.

Lemma 3.2.6 (Decomposition lifting). *Let M be a symmetric set, A be a non-empty finite set, $a \in A$ be an element, and $x \in M(A)$ be an M -structure. Suppose that a reduces x , and fix an EZ-decomposition $x\delta^a = y\alpha$ of $x\delta^a$ where $\alpha: A \setminus \{a\} \rightarrow B$. Then there exists a unique surjection $\beta: A \rightarrow B$ such that*

- $x = y\beta$ is an EZ-decomposition of x , and
- β is an extension of α (that is, $\beta\delta^a = \alpha$).

Proof. We will first prove the uniqueness of such β , assuming that it exists, by applying Lemma 3.2.4 to the representable symmetric set $\mathbf{F}(-, B)$. Observe that, when β is regarded as an $\mathbf{F}(-, B)$ -structure on A , the associated EZ-congruence \sim_β is given by $b \sim_\beta c \iff \beta(b) = \beta(c)$ for $b, c \in A$. Hence \sim_β coincides with \sim_x , and in particular a reduces β . Moreover, pulling back the M -structure β along δ^a simply yields the composite $\beta\delta^a$, which is required to be α . This completes the proof of the uniqueness.

Now we construct the desired β . Let $x = z\gamma$ be an EZ-decomposition of x . Since a reduces x , we obtain another EZ-decomposition $x\delta^a = z(\gamma\delta^a)$ of $x\delta^a$. By Proposition 2.2.3, there exists a bijection σ such that $\sigma\gamma\delta^a = \alpha$ and $z = y\sigma$. It is easy to see that $\beta := \sigma\gamma$ satisfies the required conditions. \square





FIGURE 6. Decomposition lifting

FIGURE 7. Reduction graph of the cycle Figure 2 with black \overline{P} and white \underline{P}

3.3. Reduction graph. Armed with all the preparation, we are now ready to start proving that any cycle that comprises sufficiently degenerate M -structures admits a unique degenerate filler (Theorem 3.5.2). This will provide the desired upper bound for the Aufhebung.

Notation 3.3.1. Throughout Subsections 3.3, 3.4, and 3.5, we fix

- a symmetric set M ,
- a positive integer $k > 0$,
- a $(k + 1)$ -element set P , and
- a k -cycle $\{c_p\}_{p \in P}$ (satisfying the cycle equations $c_p \delta^q = c_q \delta^p$),

and write

- $n := \max\{\text{mass}(c_p) \mid p \in P\} - 1$,
- $d := k - n$,
- $\overline{P} := \{p \in P \mid \text{mass}(c_p) = n + 1\}$, and
- $\underline{P} := \{p \in P \mid \text{mass}(c_p) \leq n\}$.

Note that we have $P = \overline{P} \sqcup \underline{P}$ and $\overline{P} \neq \emptyset$.

The rest of this paper is devoted to proving that, if the following inequalities hold, then the k -cycle $\{c_p\}_{p \in P}$ admits a unique degenerate filler:

$$(\star) \quad \begin{aligned} d &\geq 3 \quad (\iff k > n + 2) \\ k &> 2n - 1. \end{aligned}$$

We emphasise that these inequalities are NOT assumed by default in what follows; when they are assumed, we will explicitly state so.

An example of $\{c_p\}_{p \in P}$ satisfying Inequalities \star in the case where $M = \text{Graph}$ and $(n, k, d) = (3, 6, 3)$ is visualised in Figure 2.

Definition 3.3.2. By the *reduction graph* G of the k -cycle $\{c_p\}_{p \in P}$, we mean the directed graph such that

- its vertex set is P , and
- there is a (unique) edge $p \rightarrow q$ if and only if p reduces c_q .

We write \overline{G} for the subgraph of G consisting of the vertices in \overline{P} and all edges between them.

Proposition 3.3.3. *The reduction graph G has the following properties.*

- (1) *There is no edge from a vertex in \underline{P} to one in \overline{P} .*
- (2) *In the subgraph \overline{G} , there is an edge $p \rightarrow q$ if and only if there is an edge $p \leftarrow q$.*

In the case $d \geq 2$, it further enjoys the following properties.

- (3) The indegree d_p of each vertex $p \in P$ is at least d .
- (4) The subgraph \overline{G} contains at least $d + 1$ vertices.

Proof. To see (1) and (2), consider an edge $p \rightarrow q$ with $q \in \overline{P}$. Since p reduces c_q , we have

$$\text{mass}(c_p) \geq \text{mass}(c_p \delta^q) = \text{mass}(c_q \delta^p) = \text{mass}(c_q) = n + 1 \geq \text{mass}(c_p).$$

In particular, we can deduce $\text{mass}(c_p) = n + 1$ and $\text{mass}(c_p \delta^q) = \text{mass}(c_p)$.

Now suppose $d \geq 2$. For each vertex $p \in P$, its indegree d_p is equal to

$$d_p = k - (\text{the number of isolated points with respect to } \sim_{c_p}).$$

Recall that \sim_{c_p} is an equivalence relation on the k -element set $P \setminus \{p\}$, and the number of \sim_{c_p} -classes is precisely $\text{mass}(c_p) \leq n + 1$. Since we are assuming $k = n + d \geq n + 2$, there must be at least one \sim_{c_p} -class which is not a singleton. Hence the number of isolated points in the above equation is at most n , which establishes (3).

For (4), fix $q \in \overline{P}$, whose existence follows from the definition of n . Then there are at least d edges of the form $p \rightarrow q$ by (3), and each such p belongs to \overline{P} by (1). This completes the proof. \square

Notice that we can regard the graph \overline{G} , which is a priori directed, as an undirected graph thanks to Proposition 3.3.3(2).

Remark 3.3.4 (The reduction graphs of the unfillable cycles in Lemma 2.4.5). It turns out that, assuming Inequalities \star , the undirected graph \overline{G} must be propagative (Proposition 3.4.4), and this property will play a central role when constructing a filler of $\{c_p\}_{p \in P}$ (Proposition 3.4.2). Hence, in order to find an unfillable k -cycle $\{c_p\}_{p \in P}$, it is natural to look for one with a non-propagative reduction graph. Indeed, the cycles appearing in Lemma 2.4.5 (for which $P = \overline{P}$ holds) have non-propagative reduction graphs, as visualised in Figure 8. (Those not achieving the sharp lower bound are de-emphasised.)

3.4. Candidate filler and its compatibility with \overline{P} . We wish to show that, assuming Inequalities \star , we may construct a filler f of the cycle $\{c_p\}_{p \in P}$. In this subsection, we construct a candidate for such f , and prove that it satisfies $f \delta^p = c_p$ for at least all $p \in \overline{P}$. Our strategy is to prove that

- (1) we can find two points $p, q \in \overline{G}$ and $f \in M(P)$ such that $f \delta^p = c_p$ and $f \delta^q = c_q$ (Lemma 3.4.1),
- (2) the property “ $f \delta^r = c_r$ ” propagates along the edges of \overline{G} (Proposition 3.4.2), and
- (3) \overline{G} is propagative (Proposition 3.4.4).

Moreover, such f turns out to be unique if we impose $\text{mass}(f) = n + 1$.

Lemma 3.4.1 (Seed of propagation). *Let $p \leftrightarrow q$ be an edge in \overline{G} . Then there exists a unique $f \in M(P)$ such that $f \delta^p = c_p$, $f \delta^q = c_q$, and $\text{mass}(f) = n + 1$.*

Proof. Observe that, since $p, q \in \overline{G}$ implies $\text{mass}(c_p) = \text{mass}(c_q) = n + 1$, any solution f to these three equations is necessarily reduced by both p and q . It follows by Lemma 3.2.5 that such f is unique if it exists.

Let $c_p \delta^q = c_q \delta^p = x\alpha$ be an EZ-decomposition where $\alpha: P \setminus \{p, q\} \rightarrow S$. Since p reduces c_q and q reduces c_p , Lemma 3.2.6 provides surjections $\beta_p: P \setminus \{p\} \rightarrow S$ and $\beta_q: P \setminus \{q\} \rightarrow S$ such that

- $c_p = x\beta_p$,
- $\beta_p \delta^q = \alpha$,
- $c_q = x\beta_q$, and
- $\beta_q \delta^p = \alpha$.

We define $\beta: P \rightarrow S$ to be the unique common extension of β_p and β_q . Then $f := x\beta$ satisfies $f \delta^p = c_p$, $f \delta^q = c_q$, and $\text{mass}(f) = n + 1$. \square




FIGURE 8. The reduction graphs of the cycles in Lemma 2.4.5, which are not propagative.

In the following proposition, note that we are allowing the case $r \in \underline{P}$. This extra generality will be utilised in the next subsection.

Proposition 3.4.2 (Propagation step). *Let $p, q, r \in P$ be distinct elements and $f \in M(P)$ with $\text{mass}(f) = n + 1$. Suppose that*

- $p, q \in \overline{P}$,
- both p and q reduce c_r ,
- $f\delta^p = c_p$, and
- $f\delta^q = c_q$.

Then we have $f\delta^r = c_r$.

Proof. We wish to apply Lemma 3.2.5 to $x = f\delta^r$ and $y = c_r$. Since we already know $f\delta^r\delta^p = f\delta^p\delta^r = c_p\delta^r = c_r\delta^p$ and similarly $f\delta^r\delta^q = c_r\delta^q$, it remains to prove that p and q reduce $f\delta^r$. Note that, since we already know $\text{mass}(f\delta^r\delta^p) = \text{mass}(c_r\delta^p) = \text{mass}(c_r)$ and similarly $\text{mass}(f\delta^r\delta^q) = \text{mass}(c_r)$, p reduces $f\delta^r$ if and only if $\text{mass}(f\delta^r) = \text{mass}(c_r)$ if and only if q reduces $f\delta^r$.

Suppose for contradiction that neither p nor q reduces $f\delta^r$. Then p is isolated with respect to $\sim_{f\delta^r}$. However, it is not isolated with respect to \sim_f as it reduces f (because $\text{mass}(f) = n + 1 = \text{mass}(c_p) = \text{mass}(f\delta^p)$). Therefore we have $p \sim_f r$ by Lemma 2.2.6, and similarly $q \sim_f r$, which implies $p \sim_f q$. This in turn implies $p \sim_{f\delta^r} q$ again by Lemma 2.2.6, which contradicts the second sentence of this paragraph. \square

Lemma 3.4.3. *Assuming Inequalities \star , at least one of the following conditions holds.*

- (1) For every vertex $p \in P$, we have $d_p > d$.
- (2) The set \underline{P} contains at least $n - 1$ elements.

Proof. We may assume $n \geq 2$ because the condition (2) trivially holds in the case $n \leq 1$. We will further assume

- (-1) there exists $p \in P$ with $d_p = d$, and
- (-2) the set \underline{P} contains less than $n - 1$ elements,

and derive a contradiction. For fix $p \in P$ with $d_p = d$, and define

$$Q := \{q \in P \mid q \rightarrow p\}.$$

We will write $R := P \setminus (Q \cup \{p\})$. Note that this set R contains exactly $(k + 1) - (d + 1) = n$ elements.

Claim 3.4.3.a. *The EZ-congruence \sim_{c_p} partitions $P \setminus \{p\}$ as*

$$P \setminus \{p\} = Q \sqcup \{r_1\} \sqcup \cdots \sqcup \{r_n\}$$

where $r_i \in R$, so in particular $p \in \overline{P}$. Consequently, for any $q \in Q$, the EZ-congruence $\sim_{c_q\delta^p}$ partitions $P \setminus \{p, q\}$ as

$$P \setminus \{p, q\} = Q \setminus \{q\} \sqcup \{r_1\} \sqcup \cdots \sqcup \{r_n\}.$$

Proof of Claim. We know that none of the n elements $r_i \in R$ reduces c_p , so Lemma 3.2.1 implies that each $\{r_i\}$ is a singleton \sim_{c_p} -class. Since there can only be $\text{mass}(c_p) \leq n + 1$ many \sim_{c_p} -classes, it follows that the whole Q forms a single \sim_{c_p} -class. The last assertion follows from the cycle equation $c_q\delta^p = c_p\delta^q$ and Lemma 3.2.3. \square

As a consequence of $p \in \overline{P}$, we obtain another description of the set Q :

$$Q = \{q \in \overline{P} \mid q \leftrightarrow p\}$$

due to Proposition 3.3.3.

Claim 3.4.3.b. *Let $q, q' \in Q$ be distinct elements. Then q' reduces c_q .*

Proof of Claim. Since p reduces c_q , the EZ-congruence $\sim_{c_q\delta^p}$ of Claim 3.4.3.a is the restriction of \sim_{c_q} onto $P \setminus \{p, q\}$. In particular, the \sim_{c_q} -class containing q' must subsume $Q \setminus \{q\}$. It therefore follows from $|Q \setminus \{q\}| = d-1 \geq 3-1 = 2$ that q' reduces c_q . \square

Claim 3.4.3.c. *The set R contains at least $n-1$ elements that reduce no c_q for $q \in Q$.*

Proof of Claim. First, let us fix $q \in Q$ and consider which $r_i \in R$ does not reduce c_q . Since p reduces c_q , the EZ-congruence $\sim_{c_q\delta^p}$ of Claim 3.4.3.a is the restriction of \sim_{c_q} onto $P \setminus \{p, q\}$. It follows that $\{r_i\}$ is a singleton \sim_{c_q} -class for all $1 \leq i \leq n$ except for possibly one i satisfying $r_i \sim_{c_q} p$.

It remains to prove that if we have $r_i \sim_{c_q} p$ and $r_{i'} \sim_{c_{q'}} p$ for distinct $q, q' \in Q$ then $i = i'$. Observe that, since q reduces $c_{q'}$ and q' reduces c_q by Claim 3.4.3.b, the EZ-congruences \sim_{c_q} and $\sim_{c_{q'}}$ restrict to the same equivalence relation \sim on $P \setminus \{q, q'\}$. So $r_i \sim_{c_q} p$ and $r_{i'} \sim_{c_{q'}} p$ would imply $r_i \sim p \sim r_{i'}$, which in turn implies $r_i \sim_{c_q} r_{i'}$, and consequently $r_i \sim_{c_q\delta^p} r_{i'}$. We can thus conclude $i = i'$ by Claim 3.4.3.a. \square

Claim 3.4.3.d. *There exists $r \in \overline{P} \cap R$ such that neither p nor any $q \in Q$ reduces c_r .*

Proof of Claim. Since we are assuming that $\underline{P} = P \setminus \overline{P}$ contains less than $n-1$ elements, Claim 3.4.3.c and the Pigeonhole Principle imply that there exists $r \in \overline{P} \cap R$ that reduces no c_q for $q \in Q$. We also know that r does not reduce c_p because $r \in R$. By Proposition 3.3.3(2), this r has the desired property. \square

We can finally derive the desired contradiction using r of Claim 3.4.3.d as follows. By Lemma 3.2.1, each element of $Q \cup \{p\}$ gives rise to a singleton \sim_{c_r} -class. In addition, since we have

$$|P \setminus \{r\}| = k = n + d > n + 1 = \text{mass}(c_r),$$

there must be at least one non-singleton \sim_{c_r} -class. Hence by counting the number of \sim_{c_r} -classes, we obtain the inequality

$$n + 1 = \text{mass}(c_r) \geq |Q \cup \{p\}| + 1 = d + 2 = k - n + 2,$$

which is equivalent to $k \leq 2n - 1$. This contradicts Inequalities \star . \square

Proposition 3.4.4. *Assuming Inequalities \star , the undirected graph \overline{G} is propagative.*

Proof. Let m be the minimum among the degrees of all vertices in the undirected graph \overline{G} . By Proposition 3.1.4, it suffices to prove $|\overline{P}| < 2m$. We will divide our proof into two cases according to Lemma 3.4.3.

- (1) Suppose that we have $d_p > d$ for all $p \in P$. Then we have $m \geq d + 1$. Combining this inequality with the assumption $k > 2n - 1$, or equivalently $k - 2n + 1 > 0$, we obtain

$$|\overline{P}| \leq |P| = k + 1 < 2k - 2n + 2 = 2(d + 1) \leq 2m.$$

- (2) Suppose that $\underline{P} = P \setminus \overline{P}$ contains at least $n-1$ elements. Then, since we are assuming $d \geq 3$, we have

$$|\overline{P}| = |P| - |\underline{P}| \leq (k + 1) - (n - 1) = d + 2 < 2d \leq 2m$$

where the last inequality follows from Proposition 3.3.3(3).

This completes the proof. \square

Combining the results in this subsection, we obtain the following.

Proposition 3.4.5. *Assuming Inequalities \star , there exists a unique² $f \in M(P)$ such that $\text{mass}(f) = n + 1$ and $f\delta^p = c_p$ for all $p \in \overline{P}$.*

²If we omit $\text{mass}(f) = n + 1$, we lose the uniqueness. For example, consider $\text{Eq}_{=j+1}$ for a large j .

Proof. Take an edge $p \leftrightarrow q$ in \overline{G} , whose existence is guaranteed By Proposition 3.3.3. By Lemma 3.4.1, there exists $f \in M(P)$ with $\text{mass}(f) = n + 1$ such that $f\delta^p = c_p$ and $f\delta^q = c_q$. We define $S \subset \overline{P}$ by $S := \{r \in \overline{P} \mid f\delta^r = c_r\}$. By Proposition 3.4.2 and Proposition 3.4.4, we have $S = \overline{P}$. The uniqueness also follows from Lemma 3.4.1. \square

3.5. **Compatibility with \underline{P} .** In the previous subsection, we saw that the property “ $f\delta^r = c_r$ ” (for the judiciously chosen f) propagates to the whole of \overline{P} . It thus remains to prove that this property propagates to \underline{P} too.

Proposition 3.5.1. *Assume Inequalities \star and let $p \in \underline{P}$. Then there exist distinct points $p_0, p_1 \in \overline{P}$ that reduce c_p .*

Proof. Since $d_p = d$ would imply $p \in \overline{P}$ by Claim 3.4.3.a, we have $d_p \geq d + 1$. We also have $|\overline{P}| \geq d + 1$ by Proposition 3.3.3, so we obtain

$$|\overline{P}| + d_p \geq 2d + 2 = 2(k - n) + 2 = k + 1 + (k - (2n - 1)) > k + 1 = |P \setminus \{p\}| + 1.$$

Thus the subsets \overline{P} and $\{q \in P \mid q \rightarrow p\}$ of $P \setminus \{p\}$ share at least two elements p_0 and p_1 . \square

Theorem 3.5.2. *Assuming Inequalities \star , there exists a unique degenerate $f \in M(P)$ such that $f\delta^p = c_p$ for all $p \in P$.*

Proof. Let $f \in M(P)$ be the M -structure of Proposition 3.4.5. We already know that $f\delta^p = c_p$ holds for all $p \in \overline{P}$. This property then propagates to all $p \in P$ by Proposition 3.4.2 and Proposition 3.5.1. Thus f is a filler of the cycle $\{c_p\}_{p \in P}$.

It remains to prove that any degenerate filler f' of $\{c_p\}_{p \in P}$ must coincide with f . Note that, because of the uniqueness part of Proposition 3.4.5, it suffices to show that such f' necessarily has mass $n + 1$. This is indeed the case because f' being degenerate implies

$$\text{mass}(f') = \max\{\text{mass}(f'\delta^p) \mid p \in P\} = \max\{\text{mass}(c_p) \mid p \in P\} = n + 1.$$

This completes the proof. \square

3.6. Conclusion.

Theorem 3.6.1 (Main theorem). *The Aufhebung of the level labelled by l is given by*

$$a_l = \begin{cases} 0 & (l = -\infty) \\ 1 & (l = 0) \\ 3 & (l = 1) \\ 4 & (l = 2) \\ 2l - 1 & (l \geq 3). \end{cases}$$

Proof. The first two values $a_{-\infty}, a_0$ are already established in Proposition 2.4.2 and Proposition 2.4.3. So fix $l \geq 1$ and let M be an l -skeletal symmetric set. Fix

$$k > \begin{cases} 3 & (l = 1) \\ 4 & (l = 2) \\ 2l - 1 & (l \geq 3), \end{cases}$$

and take an arbitrary k -cycle $\{c_p\}_{p \in P}$ in M . Defining n as in Notation 3.3.1, we obtain Inequalities \star , since the l -skeletality of M implies $n \leq l$. Thus the k -cycle $\{c_p\}_{p \in P}$ admits a unique degenerate filler by Theorem 3.5.2. In fact, removing the degeneracy assumption does not spoil the uniqueness because the l -skeletality forces every filler to be degenerate.

By Proposition 2.3.7, we can conclude that M is a_l -coskeletal where $a_l = \max(l + 2, 2l - 1)$. Combining this observation with the lower bound given in Proposition 2.4.7, we obtain the Aufhebung relation as stated. \square

REFERENCES

- [BM11] Clemens Berger and Ieke Moerdijk. On an extension of the notion of reedy category. *Mathematische Zeitschrift*, 269(3):977–1004, 2011.
- [Cam23] Timothy Champion. Cubical sites as Eilenberg-Zilber categories. *arXiv:2303.06206*, 2023.
- [Cis06] Denis-Charles Cisinski. *Les préfaisceaux comme modèles des types d'homotopie*. Société mathématique de France, 2006.
- [Cis19] Denis-Charles Cisinski. *Higher categories and homotopical algebra*, volume 180. Cambridge University Press, 2019.
- [Gra01] Marco Grandis. Finite sets and symmetric simplicial sets. *Theory and Applications of Categories [electronic only]*, 8:244–252, 2001.
- [HL25] Philip Hackney and Justin Lynd. Partial groups as symmetric simplicial sets. *Journal of Pure and Applied Algebra*, page 107864, 2025.
- [Joh02] Peter T. Johnstone. *Sketches of an Elephant: A Topos Theory Compendium, Volume 1*. Oxford University Press, 2002.
- [Joy81] André Joyal. Une théorie combinatoire des séries formelles. *Advances in mathematics*, 42(1):1–82, 1981.
- [KL89] G. Max Kelly and F. William Lawvere. *On the complete lattice of essential localizations*. University of Sydney. Department of Pure Mathematics, 1989.
- [KRRZ11] Carolyn Kennett, Emily Riehl, Michael Roy, and Michael Zaks. Levels in the toposes of simplicial sets and cubical sets. *Journal of Pure and Applied Algebra*, 215(5):949–961, 2011.
- [Law88] F. William Lawvere. Toposes generated by codiscrete objects, in combinatorial topology and functional analysis. In *Notes for Colloquium lectures given at North Ryde, NSW, Aus*, 1988.
- [Law89] F. William Lawvere. Display of graphics and their applications, as exemplified by 2-categories and the hegelian “taco”. In *Proceedings of the first international conference on algebraic methodology and software technology*, pages 51–75, 1989.
- [Law91] F. William Lawvere. More on graphic toposes. *Cahiers de topologie et géométrie différentielle catégoriques*, 32(1):5–10, 1991.
- [Law04] F. William Lawvere. Functorial concepts of complexity for finite automata. *Theory and Applications of Categories*, 13(10):164–168, 2004.
- [Law06] F. William Lawvere. Some thoughts on the future of category theory. In *Category Theory: Proceedings of the International Conference held in Como, Italy, July 22–28, 1990*, pages 1–13. Springer, 2006.
- [Law09] F. William Lawvere. Open problems in topos theory. Can be found via "<https://lawverearchives.com>", 2009.
- [Mar24] Jérémie Marquès. A criterion for categories on which every grothendieck topology is rigid. *arXiv:2407.18417*, 2024.
- [Men19] Matías Menni. Monic skeleta, boundaries, aufhebung, and the meaning of 'one-dimensionality'. 2019.
- [Men24] Matías Menni. The successive dimension, without elegance. *Proceedings of the American Mathematical Society*, 152(03):1337–1354, 2024.
- [Roy97] Michael John Roy. *The topos of ball complexes*. State University of New York at Buffalo, 1997.
- [RRV09] John E. Roberts, Giuseppe Ruzzi, and Ezio Vasselli. A theory of bundles over posets. *Advances in Mathematics*, 220(1):125–153, 2009.
- [RT03] Jiří Rosický and Walter Tholen. Left-determined model categories and universal homotopy theories. *Transactions of the American Mathematical Society*, 355(9):3611–3623, 2003.
- [Zak86] Michael Zaks. Does n -skeletal imply $n+1$ -coskeletal. *preprint*, 1986.

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF TOKYO, TOKYO, JAPAN
 Email address: hora@ms.u-tokyo

RESEARCH INSTITUTE FOR MATHEMATICAL SCIENCES, KYOTO UNIVERSITY, KYOTO, JAPAN
 Email address: ymaehar@kurims.kyoto-u.ac.jp