A NOTE ON THE TWISTED DEGREE 6 *L*-FUNCTION FOR HERMITIAN CUSP FORMS OF DEGREE 2

RAFAIL PSYROUKIS

ABSTRACT. Let F be a cuspidal Hermitian eigenform of degree two over $\mathbb{Q}(i)$, with first Fourier-Jacobi coefficient not identically zero. Building on a paper by Das and Jha, we prove the meromorphic continuation to \mathbb{C} and the functional equation of a degree six L-function attached to F by Gritsenko, twisted by a Dirichlet character.

1. INTRODUCTION

And rianov initiated the study of the spinor L-function attached to a cuspidal Siegel eigenform of degree two. In [1] he proved its analytic continuation to the complex plane and its functional equation.

A major simplification of the proof of the analytic properties of the spinor L-function was given by Kohnen and Skoruppa, in their paper [7]. In particular, the authors considered the Dirichlet series

$$D_{F,G}(s) := \zeta(2s - 2k + 4) \sum_{N=1}^{\infty} \langle \phi_N, \psi_N \rangle N^{-s}, \qquad (1)$$

where ϕ_N, ψ_N are the Nth Fourier-Jacobi coefficients of two degree two Siegel cusp forms F, G of integral weight $k \ge 0$ and \langle , \rangle denotes the Petersson inner product on the space of Jacobi forms of index N and weight k.

They prove two main results: Firstly, this series admits a meromorphic continuation to \mathbb{C} and satisfies a functional equation. Secondly, when F, G are Hecke eigenforms and G is in the Maass space, this series is proportional to the spinor *L*-function attached to F. We should mention here that their results are only valid when the first Fourier-Jacobi coefficient of F is not identically zero. It turns out that this is not a restriction, as Manickam has shown in [10].

Gritsenko, in his paper [5], considered an analogue of the spinor *L*-function but for cuspidal Hermitian eigenforms of degree two over $\mathbb{Q}(i)$. By using methods similar to Andrianov, he managed to show its meromorphic continuation to \mathbb{C} as well as to establish a functional equation. Moreover, in his later paper [3], he considered a same type Dirichlet series as (1) and proved completely analogous results to the ones of Kohnen of Skoruppa for the Hermitian case, with the same restriction on the first Fourier-Jacobi coefficient.

It should be noted here that the analytic properties of the Hermitian analogue of (1) were also proved by Raghavan and Sengupta in [11]. The authors, however, could not obtain the relation of the Dirichlet series with Gritsenko's *L*-function in the general case, but only when both F, Gbelonged in the Maass space.

In another direction, twists of (1) by Dirichlet characters were considered, initially by Kohnen in [8] for the degree two case (along with some restrictions on the character) and later by [6] for the arbitrary degree case. The authors proved the analytic properties of these twisted Dirichlet series and through these, obtained the analytic properties of the twisted spinor L-function of Andrianov.

Recently, Das and Jha in [2], proved the analytic properties of a twisted Dirichlet series, analogous to (1), attached to two cuspidal Hermitian eigenforms of degree two over $\mathbb{Q}(i)$ (here

the Dirichlet character is defined on $\mathbb{Z}[i]$).

In this short note, we combine the results of Das and Jha with the methods of Gritsenko in [3], in order to obtain the analytic properties of the twisted degree six *L*-function attached to a Hermitian cuspidal eigenform of degree two over $\mathbb{Q}(i)$, with first Fourier-Jacobi coefficient not identically zero. Our main contribution is finding the correct zeta and gamma factors in order to define a **completed** twisted *L*-function and appropriately generalise [3, Theorem 4.1, 5.1].

The structure of the paper is as follows: In Section 2, we give the main definitions for Hermitian modular forms and their Fourier-Jacobi expansions. In Section 3, we present the relevant Hecke theory we need. Finally, in Section 4, we give the main Proposition 4.1, from which the main Theorem 4.1 follows, proving the analytic properties of the twisted degree six L-function considered.

Notation. We denote the space of $n \times n$ matrices with coefficients in a ring R with $M_n(R)$. For any matrix $M \in Mat_n(R)$, we denote by det(M), tr(M) the determinant and trace of M respectively. By $GL_n(R)$, we denote the matrices in $M_n(R)$ with non-zero determinant. For any vector v, we denote by v^t its transpose. For a complex number z let \overline{z} denote its complex conjugate and $e(z) := e^{2\pi i z}$. We also use the bracket notation $A[B] := \overline{B}^t AB$ for suitably sized complex matrices A, B. Finally, we write $A \geq B$ for two matrices A, B if A - B is semi-positive definite.

2. Preliminaries

Let $K = \mathbb{Q}(i)$ denote the Gaussian field and $\mathcal{O}_K = \mathbb{Z}[i]$ denote its ring of integers.

Definition 2.1. Let R be either K, \mathcal{O}_K or \mathbb{C} and fix an embedding $R \hookrightarrow \mathbb{C}$. We write U(2,2)(R) for the *R*-points of the **unitary group** of degree two. That is,

$$U(2,2)(R) := \{g \in \mathrm{GL}_4(R) \mid J_2[g] = J_2\},\$$

where $J_2 := \begin{pmatrix} 0_2 & -1_2 \\ 1_2 & 0_2 \end{pmatrix}$.

Hence, for an element $\begin{pmatrix} A & B \\ C & D \end{pmatrix} \in U(2,2)(R)$ with 2×2 matrices A, B, C, D, these satisfy the relations

$$\overline{A}^t C = \overline{C}^t A, \ \overline{D}^t B = \overline{B}^t D, \ A \overline{D}^t - \overline{B}^t C = 1_2.$$

Definition 2.2. The Hermitian upper half-plane of degree two is defined by

$$\mathbb{I}_2 := \{ Z = X + iY \in \mathcal{M}_2(\mathbb{C}) | \overline{X}^t = X, \overline{Y}^t = Y > 0 \}$$

We also denote the usual upper half plane by \mathbb{H} .

Π

We fix an embedding $K \hookrightarrow \mathbb{C}$. Then, an element $g = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in U(2,2)(K) \hookrightarrow U(2,2)(\mathbb{C})$ of the unitary group acts on the above upper half plane via the action

$$Z \longmapsto g\langle Z \rangle := (AZ + B)(CZ + D)^{-1}.$$

The usual factor of automorphy is defined by $j(g, Z) := \det(CZ + D)$.

Let now Γ_2 denote the **Hermitian modular group**, that is $\Gamma_2 := U(2,2)(\mathcal{O}_K)$.

Definition 2.3. A function $F : \mathbb{H}_2 \longrightarrow \mathbb{C}$ is called a **Hermitian modular form** of degree two and integer weight $k \ge 0$ if

- F is holomorphic,
- $\bullet~F$ satisfies

$$F(g\langle Z\rangle) = j(g,Z)^k F(Z),$$

for all $g \in \Gamma_2$ and $Z \in \mathbb{H}_2$.

It is well-known ([9, Chapter III.2]) that the set of all such forms constitutes a finite dimensional complex vector space, which we denote by M_2^k . Each such F admits a Fourier expansion

$$F(Z) = \sum_{N} a(N)e(\operatorname{tr}(NZ)), \qquad (2)$$

where $a(N) \in \mathbb{C}$ and N runs through all the semi-integral non-negative Hermitian matrices

$$N \in \left\{ (n_{ij})_{i,j=1}^n \ge 0 \mid n_{ii} \in \mathbb{Z}, n_{ij} = \overline{n_{ji}} \in \frac{1}{2}\mathcal{O}_K \right\}$$

F is called a **cusp form** if $a(N) \neq 0$ only for N positive definite. We denote the space of cusp forms by S_2^k . Finally, we have a notion of a Petersson inner product.

Definition 2.4. The **Petersson inner product** for two Hermitian modular forms F, G, where at least one of them is a cusp form, is given by

$$\langle F, G \rangle := \int_{\Gamma_n \setminus \mathbb{H}_n} F(Z) \overline{G(Z)} (\det Y)^k \mathrm{d}^* Z,$$

where $d^*Z = (\det Y)^{-2n} dX dY$ with Z = X + iY.

Let now $F \in S_2^k$ and $Z \in \mathbb{H}_2$. If we partition $Z = \begin{pmatrix} \tau & z_1 \\ z_2 & \omega \end{pmatrix}$, with $\tau, \omega \in \mathbb{H}$ and $z_1, z_2 \in \mathbb{C}$, we can rewrite Fourier expansion of F with respect to the variable ω as

$$F(Z) = \sum_{m=1}^{\infty} \phi_m(\tau, z_1, z_2) e(m\omega),$$

where $\phi_m : \mathbb{H} \times \mathbb{C}^2 \longrightarrow \mathbb{C}$ are holomorphic functions, called the **Fourier-Jacobi coefficients** of F. These are a special kind of modular forms, with respect to the action of the so-called **parabolic subgroups**. Let us make this more precise. Let

$$\Gamma_{1,1} := \left\{ \begin{pmatrix} * & 0 & * & * \\ * & * & * & * \\ * & 0 & * & * \\ 0 & 0 & 0 & * \end{pmatrix} \in \Gamma_2 \right\},\tag{3}$$

the Klingen parabolic of Γ_2 . We now give the definition of the slash operator.

Definition 2.5. Let $k \in \mathbb{Z}$. Then, for a function F on \mathbb{H}_2 and a matrix $g = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in U(2,2)(K)$, we define

$$F|_k g)(Z) := \det(CZ + D)^{-k} F(g\langle Z \rangle).$$

We then have the following Definition.

Definition 2.6. A holomorphic function F on \mathbb{H}_2 is a modular form of weight k with respect to the parabolic subgroup $\Gamma_{1,1}$ if the following conditions hold:

- $F \mid_k M = F$ for all $M \in \Gamma_{1,1}$,
- The function F(Z) is bounded in the domain $\text{Im}(Z) \ge c1_2$, for all c > 0.

Again, each such F has a Fourier expansion as in equation (2) and we call F a **cusp form** if $a(N) \neq 0$ only for positive definite matrices N. Finally, we can define the (Hermitian) Fourier-Jacobi forms as Gritsenko does in [3, p. 2887].

Definition 2.7. A complex-valued, holomorphic function ϕ on $\mathbb{H} \times \mathbb{C}^2$ is said to be a Fourier-Jacobi form of degree two, weight k and index m if the function

$$\tilde{\phi}\left(\begin{pmatrix} \tau & z_1\\ z_2 & \omega \end{pmatrix}\right) := \phi(\tau, z_1, z_2)e(m\omega),$$

where $\omega \in \mathbb{H}$ is chosen so that $\begin{pmatrix} \tau & z_1 \\ z_2 & \omega \end{pmatrix} \in \mathbb{H}_2$, is a modular form with respect to the group $\Gamma_{1,1}$. The space of such forms is denoted by $J_{k,m}$.

3. Hecke Rings and the degree 6 L-function

In this section, we give an account of the Hermitian Hecke theory we will need. We follow Gritsenko in [3]. We start by defining the groups of similitude:

$$S^{2} := \{g \in M_{4}(K) \mid J_{2}[g] = \mu(g)J_{2}, \text{ for some } \mu(g) > 0\},\$$
$$S^{2}_{p} := \{g \in S^{2} \cap M_{4}(\mathcal{O}_{K}[p^{-1}]) \mid \mu(g) = p^{\delta}, \delta \in \mathbb{Z}\},\$$

for each rational prime p. It is then well-known that the pairs $(\Gamma_2, S^2), (\Gamma_2, S_p^2)$ are Hecke pairs and we can define the corresponding Hecke rings, which we will denote by H^2 and H_p^2 respectively (see [4, p. 2869, 2870] for definitions of Hecke pairs and Hecke rings). From [3, Corollary 2.2], we can decompose the global Hecke ring into the tensor product of p-rings as follows:

$$H(\Gamma_2, S^2) = \bigotimes_p H(\Gamma_2, S_p^2).$$

We now want to define a Hecke ring which acts on the Fourier-Jacobi forms. To this end, we define the so-called parabolic Hecke rings. Let $S^{1,1}$ and $S_p^{1,1}$ denote the intersection of the groups S^2 and S_p^2 with $\Gamma_{1,1}$ of equation (3) respectively. Again, the pairs $(\Gamma_{1,1}, S^{1,1})$ and $(\Gamma_{1,1}, S_p^{1,1})$ are Hecke pairs (cf. [3, Section 3]) and we can then define the Hecke rings

$$H^{1,1} := H(\Gamma_{1,1}, S^{1,1}), \ H^{1,1}_p := H(\Gamma_{1,1}, S^{1,1}_p).$$

We can now view the original Hecke rings H^2 and H_p^2 as subrings of the corresponding parabolic Hecke rings, by virtue of the following Lemma.

Lemma 3.1. Let (Γ_0, S_0) and (Γ, S) be two Hecke pairs. We assume that

$$\Gamma_0 \subset \Gamma, \ \Gamma S_0 = S, \ \Gamma \cap S_0 S_0^{-1} \subset \Gamma_0.$$

Then, given an arbitrary element $X \in H(\Gamma, S)$, according to the second condition, we can write it as

$$X = \sum_{i} a_i(\Gamma g_i),$$

with $g_i \in S_0$. Then, if we set

$$\epsilon(X) := \sum_{i} a_i(\Gamma_0 g_i),$$

then ϵ does not depend on the selection of the elements $g_i \in S_0$ and is an embedding (as a ring homomorphism) of the Hecke algebra $H(\Gamma, S)$ to $H(\Gamma_0, S_0)$.

Proof. See [3, page 2890].

Since $\Gamma_2 S_p^{1,1} = S_p^2$ and after writing an element $X \in H_p^2$ as

$$X = \sum_{i} a_i \Gamma_2 g_i,$$

with $g_i \in S_p^2$, we can define an embedding

$$X \longmapsto \epsilon(X) = \sum_{i} a_i \Gamma_{1,1} g_i$$

using Lemma 3.1.

Finally, we can embed the classical Hecke algebra $H(\mathrm{SL}_2(\mathbb{Z}), \mathrm{GL}_2(\mathbb{Q})) \hookrightarrow H(\Gamma_{1,1}, S^{1,1})$ in two ways, as follows: If $X = \mathrm{SL}_2(\mathbb{Z})g\mathrm{SL}_2(\mathbb{Z})$ with $g = \mathrm{diag}(a, d)$, where $a, d \in \mathbb{Z}$, we define

$$j_{-}(X) := \Gamma_{1,1} \operatorname{diag}(a, \det(g), d, 1) \Gamma_{1,1}, \ j_{+}(X) := \Gamma_{1,1} \operatorname{diag}(a, 1, d, \det(g)) \Gamma_{1,1}.$$
(4)

These are related by an anti-homomorphism $*: H^{1,1}_p \longrightarrow H^{1,1}_p,$ given by

$$\sum_{i} a_i \Gamma_{1,1} M_i \Gamma_{1,1} \longmapsto \sum_{i} a_i \Gamma_{1,1} \det(M_i) M_i^{-1} \Gamma_{1,1}, \tag{5}$$

as in [3, Lemma 3.1]. Finally, we define how the elements of the parabolic Hecke algebras act on Fourier-Jacobi forms. Let now

$$X = \Gamma_{1,1} \begin{pmatrix} * & 0 & * & * \\ * & a & * & * \\ * & 0 & * & * \\ 0 & 0 & 0 & b \end{pmatrix} \Gamma_{1,1} = \sum_{i} \Gamma_{1,1} g_i \in H^{1,1},$$

for some $g_i \in S^{1,1}$. Then, we define

$$(F \mid_k X)(Z) := \sum_i (F \mid_k g_i)(Z).$$
(6)

Gritsenko gave the following very convenient definition of the signature.

Definition 3.1. The signature of X is defined as s(X) := b/a.

Using the signature s := s(X) of X, we can now define its action on Fourier-Jacobi forms.

Proposition 3.1. Let $\phi \in J_{k,m}$ denote a Fourier-Jacobi form of weight k and index m. Then, for $Z = \begin{pmatrix} \tau & z_1 \\ z_2 & \omega \end{pmatrix} \in \mathbb{H}_2$, we define the action of X on ϕ via

$$(\phi \mid_k X) (\tau, z_1, z_2) := \left(\widetilde{\phi} \mid_k X \right) (Z) e \left(-\frac{m}{s} \omega \right),$$

with $\widetilde{\phi}(Z) := \phi(\tau, z_1, z_2) e(m\omega)$. Then $\phi \mid_k X$ belongs to $J_{k,m/s}$ if m/s is an integer and is 0 otherwise.

Proof. See [3, Lemma 4.1].

Finally, we define the degree 6 L-function associated to a Hermitian modular form of degree two. For each prime p, we define the following standard elements of $H(\Gamma_2, S_n^2)$:

• $T_p := \Gamma_2 \operatorname{diag}(1, 1, p, p) \Gamma_2$. • $T_{1,p} := \Gamma_2 \operatorname{diag}(1, p, p^2, p)\Gamma_2.$ • $T_{\pi} := \Gamma_2 \operatorname{diag}(1, \pi, p, \pi)\Gamma_2, \ T_{\overline{\pi}} := \Gamma_2(1, \overline{\pi}, p, \overline{\pi})\Gamma_2 \text{ (if } p = \pi\overline{\pi}, \text{ including the case } p = 2).$ • $\Delta_q = \Gamma_2 \operatorname{diag}(q, q, q, q)\Gamma_2, \text{ for } q \in \{p, \pi, \overline{\pi}\}, \text{ depending on the prime } p.$

Let now $F \in S_2^k$ be a Hecke eigenform for H^2 , i.e. an eigenfunction for all Hecke elements in H^2 . We use the following notation: For any polynomial $U[X] \in H^2[X]$, we write U_F for the polynomial obtained when we substitute the elements of the Hecke ring with their corresponding eigenvalues. We then have the following Definition.

Definition 3.2. ([5, Lemma 2.1]) The degree 6 L-function attached to F is defined as:

$$Z_F(s) := \prod_{p \text{ prime in } \mathcal{O}_K} (1 + p^{k-s-2})^{-2} Q_{p,F}^{(2)}(p^{-s})^{-1} \prod_{p \text{ splits}} Q_{p,F}^{(2)}(p^{-s})^{-1},$$

where the polynomials $Q_p^{(2)}(t) \in H^2[t]$ are defined as follows:

• p inert in \mathcal{O}_K :

$$Q_p^{(2)}(t) := 1 - T_p t + (pT_{1,p} + p(p^3 + p^2 - p + 1)\Delta_p)t^2 - p^4\Delta_p T_p t^3 + p^8\Delta_p^2 t^4.$$
• $p = 2$:

$$Q_2^{(2)}(t) := 1 - (T_2 - 3\Delta_{i+1})t + (2T_{1+i}^2 - 8\Delta_{1+i}(T_2 + \Delta_{1+i}))t^2 - (4\Delta_{1+i})^2(T_2 - 3\Delta_{1+i})t^3 + (4\Delta_{1+i})^4t^4$$

• p splits in \mathcal{O}_K :

$$\begin{aligned} Q_p^{(2)}(t) &:= 1 - T_p t + (pT_{\pi}T_{\overline{\pi}} - p^4\Delta_p)t^2 - (p^3(T_{\pi}^2\Delta_{\overline{\pi}} + T_{\overline{\pi}}^2\Delta_{\pi}) - 2p^4\Delta_p T_p)t^3 + p^4\Delta_p (pT_{\pi}T_{\overline{\pi}} - p^4\Delta_p)t^4 - p^8\Delta_p^2 T_p t^5 + p^{12}\Delta_p^3 t^6. \end{aligned}$$

This is defined for $\operatorname{Re}(s)$ large enough (cf. [5, p. 2546]).

We now define the **twisted degree** 6 *L*-function attached to *F* as follows. Let $N \ge 1$ be an integer. Once and for all, we fix a **Dirichlet character** $\chi : (\mathcal{O}_K/N\mathcal{O}_K)^{\times} \to \mathbb{C}^{\times}$.

Definition 3.3. Let $F \in S_2^k$ be an eigenform for H^2 . For $\operatorname{Re}(s)$ large enough, we define

$$Z_F^{(2)}(s,\chi) := \prod_{p \text{ prime in } \mathcal{O}_K} (1+\chi(p)p^{k-2-s})^{-2} Q_{p,F}^{(2)}(\chi(p)p^{-s})^{-1} \prod_{p \text{ splits}} Q_{p,F}^{(2)}(\chi(p)p^{-s})^{-1}.$$

4. Main Theorem

In this section, we prove the meromorphic continuation to \mathbb{C} and functional equation for the twisted degree six L-function. We first make some definitions.

Let $\theta : (\mathbb{Z}/4\mathbb{Z})^* \longrightarrow \mathbb{C}^{\times}$ denote the character with $\theta(3) = -1$.

For any Dirichlet character $\psi : (\mathbb{Z}/m\mathbb{Z})^* \longrightarrow \mathbb{C}^{\times}$, where $m \geq 1$ an integer, we define

$$L(s,\psi) := \sum_{n=1}^{\infty} \psi(n) n^{-s} = \prod_{p \text{ prime}} (1 - \psi(p) p^{-s})^{-1}, \text{ Re}(s) > 1.$$

For a Dirichlet character $\omega : (\mathcal{O}_K/M\mathcal{O}_K)^{\times} \longrightarrow \mathbb{C}^{\times}$, where $M \geq 1$ an integer, we define

$$\zeta_K(s,\omega) := \frac{1}{4} \sum_{0 \neq \lambda \in \mathcal{O}_K} \omega(\lambda) N(\lambda)^{-s}, \ \operatorname{Re}(s) > 1.$$

where $N(\lambda) := \lambda \overline{\lambda}$ denotes the **norm** of λ . We note that we also consider $L(s, \omega)$ by implicitly considering the restriction of ω to \mathbb{Z} .

For any $m \ge 1$, let T(m) denote the standard Hecke element of $SL_2(\mathbb{Z})$. We then define $T_{\pm}(m) := j_{\pm}(T(m))$, where j_{\pm} are the embeddings of equation (4). Let also $\tilde{\chi}$ denote the character given by $\tilde{\chi}(\alpha) := \chi(\overline{\alpha})$. We then have the following Theorem.

Proposition 4.1. Let F be a Hermitian cusp form of degree 2 and weight k for the group Γ_2 , which is an eigenfunction for all Hecke operators of the ring $H(\Gamma_2, S^2)$. Assume ϕ_m are its Fourier-Jacobi coefficients with $\phi_1 \neq 0$. Then, for $\operatorname{Re}(s)$ large enough, we have

$$L^{2}(s-k+2,\theta\chi)L(2s-2k+4,\chi^{2})\zeta_{K}(s-k+3,\chi\tilde{\chi})\sum_{m=1}^{\infty}\phi_{m}|_{k}T_{+}(m)\chi(m)m^{-s} = \zeta_{K}(s-k+2,\chi\tilde{\chi})\phi_{1}Z_{F}^{(2)}(s,\chi).$$

If N = 1 and χ is principal, this coincides with the result of Gritsenko in [3, Theorem 4.1].

Proof. We will use [3, Proposition 4.2], which states that in the ring of formal power series $J_{k,1}[[t]]$, we have:

$$Q_{p,F}^{(2)}(t) \sum_{\delta \ge 0} (\phi_{mp^{\delta}}|_{k}T_{+}(mp^{\delta}))t^{\delta} = \begin{cases} (1-p^{2k-6}t^{2})\phi_{m}|_{k}T_{+}(m) & \text{if } p \text{ is prime in } \mathcal{O}_{K} \\ (1-p^{k-3}t)^{2}(1-p^{2k-4}t^{2})\phi_{m}|_{k}T_{+}(m) & \text{if } p \text{ is decomposable in } \mathcal{O}_{K} \\ (1-2^{k-3}t)(1+2^{k-2}t)\phi_{m}|_{k}T_{+}(m) & \text{if } p = 2 \end{cases}$$

if gcd(m, p) = 1.

Now, $T_+(mn) = T_+(m)T_+(n)$ if gcd(m, n) = 1, because of the multiplicativity of the standard Hecke elements in $SL_2(\mathbb{Z})$ and the fact that j_+ of equation (4) is a ring homomorphism. By then applying the above result successively, have that the expression

$$\frac{1}{\phi_1} \sum_{m=1}^{\infty} \phi_m |_k T_+(m) \chi(m) m^{-s}$$

has an Euler product with p-factor as follows:

$$\begin{cases} (1-\chi^2(p)p^{-2s+2k-6})Q_{p,F}^{(2)}(\chi(p)p^{-s})^{-1} & \text{if } p \text{ is prime in } \mathcal{O}_K\\ (1-\chi(p)p^{-s+k-3})^2(1-\chi^2(p)p^{-2s+2k-4})Q_{p,F}^{(2)}(\chi(p)p^{-s})^{-1} & \text{if } p \text{ is decomposable in } \mathcal{O}_K \\ (1-\chi(2)2^{-s+k-3})(1+\chi(2)2^{-s+k-2})Q_{2,F}^{(2)}(\chi(2)2^{-s})^{-1} & \text{if } p = 2 \end{cases}$$

The result now follows by comparing the local factors for each rational prime p, because of the following Euler products expressions:

•
$$L(s, \theta\chi) = \prod_{\substack{p \text{ prime in } \mathcal{O}_K}} (1 + \chi(p)p^{-s})^{-1} \prod_{\substack{2 \neq p \text{ splits}}} (1 - \chi(p)p^{-s})^{-1}.$$

• $L(s, \chi^2) = \prod_{p} (1 - \chi^2(p)p^{-s})^{-1}.$
• $\zeta_K(s, \chi\tilde{\chi}) = (1 - \chi^2(2)2^{-s})^{-1} \prod_{p \text{ prime in } \mathcal{O}_K} (1 - \chi^2(p)p^{-2s})^{-1} \prod_{\substack{2 \neq p \text{ splits}}} (1 - \chi(p)p^{-s})^{-2}.$

In order to now deduce the analytic properties of Z_F , we consider a specific Dirichlet series, as Das and Jha did in [2]. For any $F, G \in S_2^k$, with Fourier-Jacobi expansions

$$F(Z) = \sum_{m=1}^{\infty} \phi_m(\tau, z_1, z_2) e(m\omega), \ G(Z) = \sum_{m=1}^{\infty} \psi_m(\tau, z_1, z_2) e(m\omega),$$

we define a **twisted Dirichlet series** attached to them via

$$D_{F,G}(s,\chi) := \zeta(2s - 2k + 4)\zeta_K(s - k + 3, \chi\widetilde{\chi}) \sum_{m=1}^{\infty} \chi(m) \langle \phi_m, \psi_m \rangle m^{-s},$$

where $\zeta(s)$ denotes the usual Riemann zeta function. This series converges absolutely and uniformly in compact subsets for $\operatorname{Re}(s) > k + 1$ (cf. [2, Section 4]) and also has good analytic properties ([2, Theorem 4.1]).

We also define the **completion** of Z_F and the **completed twisted Dirichlet series** via $Z_F^*(s,\chi) := \frac{N^{-4s}(4\pi^3)^{-s}\Gamma(s)\Gamma(s-k+3)\Gamma(s-k+2)\zeta(2s-2k+4)\zeta_K(s-k+2,\chi\tilde{\chi})}{L^2(s-k+2,\theta\chi)L(2s-2k+4,\chi^2)}Z_F^{(2)}(s,\chi)$

and

$$D_{F,G}^{*}(s,\chi) := N^{-4s}(4\pi^{3})^{-s}\Gamma(s)\Gamma(s-k+3)\Gamma(s-k+2)\zeta(2s-2k+4)\zeta_{K}(s-k+3,\chi\tilde{\chi})\sum_{m=1}^{\infty}\chi(m)\langle\phi_{m},\psi_{m}\rangle m^{-s}.$$

Here, $\Gamma(s) := \int_0^\infty t^{s-1} e^{-t} dt$ denotes the usual Gamma function.

We then have the following main Theorem.

Theorem 4.1. For a Hermitian cusp form $F \in S_2^k$, which is an eigenfunction for all Hecke operators of the ring $H(\Gamma_2, S^2)$ and has first Fourier-Jacobi coefficient $\phi_1 \neq 0$, the function $Z_F^*(s, \chi)$ is holomorphic for all $s \in \mathbb{C}$, except possibly for some simple poles at s = k, k - 1, k - 2, k - 3 (when χ is principal) and satisfies the functional equation

$$Z_F^*(2k - 3 - s, \chi) = Z_F^*(s, \chi).$$

Proof. Let ψ_1 be a Jacobi cusp form of weight k and index 1. From [3, Proposition 5.1] we have that the adjoint operator of $T_+(m)$ with respect to the inner product of Jacobi forms is $m^{3-k}T_-(m)$. By now taking the inner product of the result of Proposition 4.1 with ψ_1 , we obtain:

$$L^{2}(s-k+2,\theta\chi)L(2s-2k+4,\chi^{2})\zeta_{K}(s-k+3,\chi\tilde{\chi})\sum_{m=1}^{\infty}\langle\phi_{m},\psi_{1}|_{k}m^{3-k}T_{-}(m)\rangle\chi(m)m^{-s}$$
$$=\zeta_{K}(s-k+2,\chi\tilde{\chi})\phi_{1}Z_{F}^{(2)}(s,\chi).$$

But by [3, Proposition 5.2], we have that the function

$$G_{\psi_1}\left(\begin{pmatrix} \tau & z_1\\ z_2 & \omega \end{pmatrix}\right) := \sum_{m \ge 1} m^{3-k} (\psi_1|_k T_-(m))(\tau, z_1, z_2) e(m\omega)$$

is a Hermitian cusp form of degree 2 and weight k (and actually belongs in the Maass space, see [3, Appendix]). Hence, the above relation can be written as

$$D^*_{F,G_{\psi_1}}(s,\chi) = \langle \phi_1, \psi_1 \rangle Z^*_F(s,\chi).$$

The Theorem now follows immediately from the result of Das and Jha in [2, Theorem 4.1].

ACKNOWLEDGEMENTS. This work was supported by the Additional Funding Programme for Mathematical Sciences, delivered by EPSRC (EP/V521917/1) and the Heilbronn Institute for Mathematical Research as well as by a scholarship by Onassis Foundation. The author would also like to thank Prof. Thanasis Bouganis for his continuous guidance and support.

References

- A. N. Andrianov. "Euler Products corresponding to Siegel modular forms of genus 2". In: Russian Mathematical Surveys 29.3 (1974), pp. 45–116.
- [2] S. Das and A.K. Jha. "Analytic properties of twisted real-analytic Hermitian Klingen type Eisenstein series and applications." In: Abh. Math. Semin. Univ. Hambg. 89 (2019), pp. 105–116.
- [3] V.A. Gritsenko. "Jacobi functions and Euler products for Hermitian modular forms." In: J Math Sci 62 (1992), pp. 2883–2914.
- [4] V.A. Gritsenko. "Parabolic extensions of the Hecke ring of the general linear group. II". In: J Math Sci 62 (1992), pp. 2869–2882.
- [5] V.A. Gritsenko. "Zeta function of degree six for Hermite modular forms of genus two". In: J Math Sci 43 (1988), pp. 2540–2553.
- [6] W. Kohnen, J. Sengupta, and A. Krieg. "Characteristic Twists of a Dirichlet series for Siegel Cusp Forms." In: *Manuscripta mathematica* 87.4 (1995), pp. 489–500.
- [7] W. Kohnen and N. P. Skoruppa. "A certain Dirichlet series attached to Siegel modular forms of degree two." In: *Invent Math* 95 (1989), pp. 541–558.
- [8] Winfried Kohnen. "On characteristic twists of certain Dirichlet series". In: Memoirs of the Faculty of Science, Kyushu University. Series A, Mathematics 47.1 (1993), pp. 103–117.
- [9] A. Krieg. Modular forms on half-spaces of quaternions. Vol. 1143. Lecture Notes in Mathematics. Springer-Verlag, Berlin, 1985, pp. xiii+203.
- [10] M. Manickam. "On the first Fourier-Jacobi coefficient of Siegel modular forms of degree two". In: Journal of Number Theory 219 (2021), pp. 404–411.
- [11] S. Raghavan and J. Sengupta. "A Dirichlet series for Hermitian modular forms of degree 2". In: Acta Arithmetica 58.2 (1991), pp. 181–201.

DEPARTMENT OF MATHEMATICAL SCIENCES, DURHAM UNIVERSITY, SOUTH. RD., DURHAM, DH1 3LE, U.K. *Email address*: rafail.psyroukis@durham.ac.uk