

ESO–SKAO Synergies for the Epoch of Reionisation and Cosmic Dawn

Andrei Mesinger¹
 Benedetta Ciardi²
 James E. Davies¹
 Samuel Gagnon-Hartman¹
 Valentina D’Odorico^{3,1}

¹ Scuola Normale Superiore, Pisa, Italy

² Max Planck Institute for Astrophysics, Garching, Germany

³ INAF–Trieste Astronomical Observatory, Italy

Mapping out the first billion years using the 21-cm line with the Square Kilometre Array (SKA) will revolutionise our understanding of the cosmic dawn, reionisation and the galaxies that drove these milestones. However, synergies with other telescopes in the form of cross-correlations will be fundamental in making and confirming initial, low signal-to-noise claims of a detection. Participants in the 2023 ESO–SKAO workshop discussed such synergies for Epoch of Reionisation (EoR) and Cosmic Dawn (CD) science. Here we highlight some of the most promising candidates for cross-correlating SKA EoR/CD observations with ESO instruments such as the Multi-Object Optical and Near-infrared Spectrograph (MOONS), the MOSAIC multi-object spectrograph, and the ArmazoNes high Dispersion Echelle Spectrograph (ANDES).

Introduction

The final phase-change of our Universe, the so-called Epoch of Reionisation (EoR), remains at the forefront of modern cosmology. After decades of studies, we are starting to home-in on the timing of the bulk of reionisation (for example, Qin et al., 2021; Bosman et al., 2022). However, we still do not really know how to connect this final phase-change of our Universe to the populations of stars and black holes that drive it.


The ultimate probe of this fundamental milestone is arguably the 21-cm line of neutral hydrogen, which is sensitive to the ionisation, temperature and density fluctuations of the intergalactic medium (IGM). Current 21-cm interferometers are aiming for a statistical detection of the

21-cm power spectrum (for example, Mertens et al., 2020; Trott et al., 2020; Abdurashidova et al., 2022). However, over the next decade(s), the Square Kilometre Array being built in Australia (SKA-Low) should provide the ultimate dataset: a 3D map of the barely-explored first half of our observable Universe. Such a worthy dataset contains precious insights into the astrophysics of galaxies and the IGM, as well as physical cosmology (see, for example, the review by Mesinger, 2020).

However, it will be a long time until we have a high-signal-to-noise (S/N) map of the EoR and the preceding Cosmic Dawn (CD). Initial claims of a detection will come from a handful of power-spectrum wave modes with low S/N. Given the novel, ground-breaking nature of the observation, it will be challenging to convince ourselves and the broader community that these preliminary ‘detections’ are genuinely cosmological. The best way of doing this is to cross-correlate the 21-cm signal with another signal of known cosmic origin.

In addition to providing an invaluable sanity check on preliminary claims of a 21-cm detection, cross-correlations can also improve the S/N. The cross power spectrum could provide a cleaner probe of the cosmological signal, since the foregrounds and systematics of different datasets are typically not correlated (for example, Amiri et al., 2024). Eventually, with SKA phase 2 we should be able to correlate images (i.e. including phases) of different datasets. This would allow us to study individual ionised or heated regions, comparing their tomography (obtained with SKA) to the brightest galaxies they contain (obtained with optical/IR telescopes like JWST).

Figure 1. An example 21-cm brightness temperature map (see colour bar) and corresponding galaxy map at $z = 10, 8, 7.5$ and 7 , obtained from a 3D multi-frequency radiative transfer code (Eide et al., 2020). The black and red points represent galaxies with [OIII] luminosity $L_{\text{OIII}} > 10^{41}$ erg s^{-1} and $L_{\text{OIII}} > 10^{42}$ erg s^{-1} , respectively, while grey points denote all galaxies within the slice. Galaxy locations are clearly seen to correlate with the 21-cm brightness temperature, since their radiation ionises and heats the surrounding IGM. Figure from Moriwaki et al. (2019).




EoR signals that can be cross-correlated with the 21-cm line

The 21-cm signal from the EoR/CD is determined by the IGM density, temperature, and ionised fraction. During the dark ages, it is a fairly clean probe of density fluctuations. After the CD, radiation from the first galaxies heats and ionises the IGM, effectively coupling modes over a range of scales. However, galaxies themselves are biased tracers of the matter field. Therefore any tracer of the large-scale matter field should in principle correlate with the 21-cm signal from the EoR/CD. The quantitative details of this correlation encode a wealth of information about the first galaxies (see references below).

In practice, however, it is difficult to cross-correlate signals because we need to match the ‘footprints’ of different probes. Foregrounds for 21-cm interferometry live in a ‘wedge’ region in Fourier space (for example, Morales et al., 2012). As such, large-scale and transverse (on-sky) modes might either be lost during foreground cleaning or be subject to a very large cosmic variance in the cross-correlation (i.e. uncorrelated cross terms only vanish in the limit of infinite samples). Therefore, the ideal candidates for cross-correlation with 21-cm interferometry are large-volume surveys with good redshift localisation.

We list some potential candidates here, briefly mentioning their pros and cons.

1. Cosmic backgrounds: (integral) radiation backgrounds like the cosmic microwave background (CMB; for example, Ma et al., 2018a; La Plante, Sippl & Lidz, 2022), near-infrared background (NIR; for example, Mao, 2014), and the X-ray background (XRB; Ma et al., 2018b), contain a contribution from $z > 5.5$, which should correlate with the EoR/CD 21-cm signal. Some radiation background analyses, like those for the CMB, are very mature. However, the overlap with the 21-cm signal is small, given that backgrounds are integrated over all redshifts, while the 21-cm signal loses many on-sky modes to foregrounds.
2. Resolved galaxies: as shown in Figure 1, galaxy maps would be obvious choices



for cross-correlation with the 21-cm signal. Mapping galaxies, at least at lower redshifts, is also a mature field. However, the need for accurate redshift determinations, a reasonably wide field of view, and a sufficient number density of $z > 5.5$ objects, makes it challenging to design wide and deep surveys. Narrow-band dropouts, grism, or spectroscopic follow-up are the most promising options (for example, Wiersma et al., 2013; Vrbanec et al., 2020; Sobacchi, Mesinger & Greig, 2016; Hutter et al., 2017; Moriawaki et al., 2019; Kubota et al., 2020; Heneka & Mesinger, 2020; Hutter et al., 2023; La Plante et al., 2023).

3. Intensity mapping (i.e. unresolved galaxies): other lines such as [CII], [OIII], CO, Lyman- α are mostly sourced by the combined emission from unresolved galaxies. In principle, line intensity maps (LIMs) could have wide fields and good redshift localisation, thus providing an excellent complementary probe to 21-cm observations (for example, Kovetz et al., 2017). However, LIMs have not yet been made at high redshifts, and many line luminosities are expected to be faint (for example, Crites et al., 2014; Yue et al., 2015; Lagache, Cousin & Chatzikos, 2018; Heneka & Cooray, 2021). A reported

cross-correlation between two untested probes might be met with skepticism. However, credible detections of (sub)millimetre lines from high- z galaxies with the Atacama Large Millimeter/submillimeter Array (ALMA) should help demonstrate the viability of LIM at higher redshifts in the near future.

4. Quasar spectra: the Lyman- α forest in quasar spectra provides a 1D skewer through the IGM which could be cross-correlated with the corresponding 21-cm forest. These UV and radio absorption lines in the same background object would provide complimentary information about the intervening IGM. Such a study requires the presence of radio-loud quasars at high redshifts, as well as a sufficiently cold IGM so as to detect the 21-cm forest (Bhagwat et al., 2022).

Bhagwat et al. (2022) have used hydrodynamic cosmological simulations to test how simultaneous absorption from metals (observed with ANDES) and HI 21-cm (with SKA) in the spectrum of a bright, radio-loud background quasar can complement each other as probes of the underlying gas properties. In Figure 4 we show maps of HI column density overlaid with 21-cm optical depth $\tau_{21\text{cm}}$, OI, CII, SiII and FeII column density at various redshifts, resulting from post-processing of the radiation hydrodynamic simulation Aurora (Pawlik et al., 2017). From the figure, the correspondence between a high HI and metal column density is clear.

Such correspondence is better quantified in Figure 5, where we show synthetic spectra from ANDES and SKA-Low. Here we assume a background source with flux of 10 mJy at 150 MHz and a power law index of -1.31 . The shaded regions highlight the locations of aligned and cospatial absorbers (ACA), i.e. those in which 21-cm and metal absorption fall in

the same window in velocity space and absorption originates from the same underlying gas, while for non-cospatial systems the absorption appears aligned only because of peculiar velocity effects and thus does not effectively probe the same region in space. These observations would probe the physical state of the IGM, including its temperature, metallicity and ionisation state, as well as the nature of nearby sources.

Conclusions


The first, low-S/N 21-cm EoR detection should come in the near future. The best way of convincing ourselves and the community that this detection is genuine would be to cross-correlate it with a known $z > 5$ signal (that has completely different foregrounds and systematics).

All high-redshift probes ultimately correlate with the matter field, albeit non-trivially in the case of 21 cm owing to its sensitivity

to large-scale radiation fields. Thus, there are many candidates for cross-correlations, including: (i) cosmic radiation backgrounds (CMB, CXB, CIB); (ii) (resolved) galaxy maps; (iii) line-intensity maps; and (iv) quasar spectra.

Here we identify promising ESO instruments that will synergise well with SKA-Low EoR surveys: (i) MOONS on the VLT; (ii) MOSAIC on the ELT; (iii) ANDES on the ELT. We show that a Lyman- α galaxy map obtained with MOONS or MOSAIC can be used together with the 21-cm interferometric signal to detect the cross-power spectrum at high S/N. Line-intensity maps obtained with a single-dish telescope with spectroscopic capabilities in the submillimetre (like the

Figure 4. Maps of HI column density overlaid with (from left to right): 21-cm optical depth $\tau_{21\text{cm}}$, OI, CII, SiII and FeII column density at $z = 11.32$ (top panels), 7.96 (middle) and 6 (bottom). Each projection is $3 h^{-1} \text{ c Mpc}$ on a side and $1 h^{-1} \text{ c Mpc}$ deep. Figure from Bhagwat et al. (2022).






Figure 5. Panel A: synthetic (noiseless) spectra at $z = 7.96$ showing the 21-cm forest (black lines) and the O I $\lambda 1302$ absorption spectra (orange). Shaded regions (pink and green) highlight the location of ACA, but their extent does not coincide with that of the corresponding absorber. The green shaded regions correspond to the same absorber in panels A and E. Panel B: zoom into the portion of the 21-cm forest spectrum with the ACA shown in panels A and E. The blue line is the noiseless spectrum ($e^{-\tau_{21\text{cm}}}$), while the black one is the simulated spectrum ($S_{\text{obs}} = S_0 e^{-\tau_{21\text{cm}}}$) for a background radio-loud source of flux (S_0) of 10 mJy at 150 MHz. Panel C: zoom into the portion of the O I spectrum of Panel A with the ACA. The orange line shows the noiseless spectrum, while the red one shows the simulated spectrum as observed by the ELT-ANDES instrument assuming an exposure time of 10 hours, two readouts per hour and spectral resolution $R = 10^5$ for a $m_{AB} = 21.0$ mag source. The purple line shows sky transmission at these wavelengths. Panel D: simulated normalised spectra as observed by the ELT and SKA-Low for O I and 21 cm overlaid in velocity space showing the green shaded region from Panel A. Panel E: full extent of the simulated 21-cm forest. The blue line shows the noiseless spectrum (as in panel A) multiplied with the radio continuum of the background source, while the black line shows the simulated spectrum as observed by SKA-Low. Figure from Bhagwat et al. (2022).

proposed concept for the Atacama Large Aperture Submillimeter Telescope [AtLAST; see Klaassen et al., 2020; Mroczkowski et al., 2024]) could also contribute to this. We also demonstrate that correlating Lyman- α and 21-cm absorption as seen in the spectrum of a putative radio-loud high- z quasar can provide complimentary information about the physical state of the IGM.

References

- Abdurashidova, Z. et al. 2022, *ApJ*, 925, 221
 Amiri, M. et al. 2024, *ApJ*, 963, 23
 Bhagwat, A. et al. 2022, *MNRAS*, 517, 2331
 Bosman, S. E. I. et al. 2022, *MNRAS*, 514, 55
 Crites, A. T. et al. 2014, *Proc. SPIE*, 9153, 91531W
 Eide, M. B. et al. 2020, *MNRAS*, 498, 6083
 Heneka, C. & Cooray, A. 2021, *MNRAS*, 506, 1573
 Heneka, C. & Mesinger, A. 2020, *MNRAS*, 496, 581
 Hutter, A. et al. 2017, *ApJ*, 836, 176
 Hutter, A. et al. 2023, *MNRAS*, 525, 1664
 Klaassen, P. D. et al. 2020, *Proc. SPIE*, 11445, 114452F
 Kovetz, E. D. et al. 2017, arXiv:1709.09066

- Kubota, K. et al. 2020, *MNRAS*, 494, 3131
 Lagache, G., Cousin, M. & Chatzikos, M. 2018, *A&A*, 609, A130
 La Plante, P., Sippl, J. & Lidz, A. 2022, *ApJ*, 928, 162
 La Plante, P. et al. 2023, *ApJ*, 944, 59
 Ma, Q. et al. 2018a, *MNRAS*, 476, 4025
 Ma, Q. et al. 2018b, *MNRAS*, 480, 26
 Maiolino, R. et al. 2020, *The Messenger*, 180, 24
 Mao, X.-C. 2014, *ApJ*, 790, 148
 Mertens, F. G. et al. 2020, *MNRAS*, 493, 1662
 Mesinger A. (ed.) 2020, *The cosmic 21-cm revolution: charting the first billion years of our universe*, (Bristol: IOP Publishing)
 Morales M. F. et al. 2012, *ApJ*, 752, 137
 Moriwaki, K. et al. 2019, *MNRAS*, 489, 2471
 Mroczkowski, T. et al. 2024, arXiv:2402.18645
 Pawlik, A. H. et al. 2017, *MNRAS*, 466, 960
 Prelogović, D. & Mesinger, A. 2023, *MNRAS*, 524, 4239
 Qin, Y. et al. 2021, *MNRAS*, 506, 2390
 Sobacchi, E., Mesinger, A. & Greig, B. 2016, *MNRAS*, 459, 2741
 Trott, C. M. et al. 2020, *MNRAS*, 493, 4711
 van Kampen, E. et al. 2024, arXiv:2403.02806
 Vrbancic, D. et al. 2020, *MNRAS*, 492, 4952
 Wiersma, R. P. C. et al. 2013, *MNRAS*, 432, 2615
 Yue, B. et al. 2015, *MNRAS*, 450, 3829



An SKA-Low antenna at the site of the SKA-Low telescope in Western Australia.