

ON THE CLASS OF BENSON'S COFIBRANT MODULES

IOANNIS EMMANOUIL AND WEI REN

ABSTRACT. In this paper, we examine the class of cofibrant modules over a group algebra kG , that were defined by Benson in [2]. We show that this class is always the left-hand side of a complete hereditary cotorsion pair in the category of kG -modules. It follows that the class of Gorenstein projective kG -modules is special precovering in the category of kG -modules, if G is contained in the class $\mathbf{LH}\mathfrak{F}$ of hierarchically decomposable groups defined by Kropholler in [20] and k has finite weak global dimension. It also follows that the obstruction to the equality between the classes of cofibrant and Gorenstein projective kG -modules can be described, over any group algebra kG , in terms of a suitable subcategory of the stable category of Gorenstein projective kG -modules.

CONTENTS

0. Introduction	1
1. Preliminaries	3
2. Cofibrant and Gorenstein projective modules	5
3. The cotorsion pair $(\mathrm{Co}\mathfrak{f}(kG), \mathrm{Co}\mathfrak{f}(kG)^\perp)$	10
4. Some consequences	14
References	19

0. INTRODUCTION

Gorenstein homological algebra is the relative homological theory, which is based on the classes of Gorenstein projective, Gorenstein injective and Gorenstein flat modules. It has developed rapidly during the past several years and has found interesting applications in representation theory. Gorenstein homological algebra has found applications in cohomological group theory as well. If k is a commutative ring and G is a group, the Gorenstein cohomological dimension of G over k generalizes the ordinary cohomological dimension of G over k , as well as the virtual cohomological dimension of G over k , whenever the latter is defined. The Gorenstein cohomological dimension of G over \mathbb{Z} coincides for certain classes of groups with the geometric Bredon dimension, which is defined to be the minimal dimension of a model for the classifying space $\underline{E}G$ for proper actions of G ; the reader is referred to the Introduction of [11] for more details on this relation. In fact, the Gorenstein cohomological dimension of groups over \mathbb{Z} is proposed in [1] to serve as an algebraic invariant, whose finiteness characterizes the groups G that admit a finite dimensional model for $\underline{E}G$. The point here is that the augmented cellular chain complex of such a model for $\underline{E}G$ is a resolution of finite length of the trivial $\mathbb{Z}G$ -module \mathbb{Z} by Gorenstein projective $\mathbb{Z}G$ -modules.

Building upon the relation between the class of finite groups and the class of those groups that admit a cellular action on a finite dimensional contractible CW-complex with finite cell stabilizers, Kropholler defined in [20] the class $\mathbf{LH}\mathfrak{F}$ of those groups that admit a hierarchical

decomposition starting from the class \mathfrak{F} of finite groups. This is a very big class of groups and it does require some effort to construct groups which are not contained therein. A useful tool for the construction of examples of Gorenstein projective $\mathbb{Z}G$ -modules is the module $B(G, \mathbb{Z})$ of bounded functions on G with values in \mathbb{Z} , which is introduced in [21]. Cornick and Kropholler have proved in [5] that the class of Gorenstein projective $\mathbb{Z}G$ -modules contains any $\mathbb{Z}G$ -module M , for which the $\mathbb{Z}G$ -module $M \otimes_{\mathbb{Z}} B(G, \mathbb{Z})$ is projective. Benson [2] realized the importance of the latter condition for the construction of model structures in the category of modules over the group algebras of certain groups and termed the $\mathbb{Z}G$ -modules satisfying that condition as cofibrant. Conversely, it is shown by Dembegioti and Talelli in [8] that any Gorenstein projective $\mathbb{Z}G$ -module is cofibrant if G is an $\mathbf{LH}\mathfrak{F}$ -group. The equality between the classes $\mathbf{Cof}(\mathbb{Z}G)$ and $\mathbf{GProj}(\mathbb{Z}G)$ of cofibrant and Gorenstein projective $\mathbb{Z}G$ -modules respectively, which is conjectured to hold over any group in [loc.cit.], is an important problem in cohomological group theory. Having shown that equality, the study of Gorenstein projective $\mathbb{Z}G$ -modules would become considerably simpler; it would be essentially reduced to certain questions involving classical homological algebra notions. For example, it would follow that being Gorenstein projective is a module property, which is closed under subgroups.

Having in mind the applications to group actions on spaces, the most important ring of coefficients to work with is the ring \mathbb{Z} of integers. The equality between the classes $\mathbf{Cof}(kG)$ and $\mathbf{GProj}(kG)$ of cofibrant and Gorenstein projective kG -modules has been also studied for general coefficient rings though. It is shown in [3] that the equality between these classes holds if G is an $\mathbf{LH}\mathfrak{F}$ -group and k has finite global dimension. Using certain contractibility criteria for acyclic complexes of projective k -modules, we can relax that assumption on k and prove that $\mathbf{Cof}(kG) = \mathbf{GProj}(kG)$ for any $\mathbf{LH}\mathfrak{F}$ -group G , if k satisfies one of the following conditions:

- (i) k has finite weak global dimension or
- (ii) there exists an integer n , such that $\mathrm{pd}_k C \leq n$ for any finitely presented k -module C .

We also prove that, over any group G , the equality $\mathbf{Cof}(kG) = \mathbf{GProj}(kG)$ holds if there exists an integer n , such that $\mathrm{pd}_{kG}(C \otimes_{\mathbb{Z}} B(G, \mathbb{Z})) \leq n$ for any finitely presented kG -module C .

Another example that illustrates the importance of knowing that Gorenstein projective kG -modules are cofibrant is provided by the question regarding the existence of Gorenstein projective precovers. As shown in [6], the pair $(\mathbf{GProj}(R), \mathbf{GProj}(R)^\perp)$ is a hereditary cotorsion pair in the category of R -modules over any unital ring R . An important problem in Gorenstein homological algebra is to show that this cotorsion pair is complete. We show that the pair $(\mathbf{Cof}(kG), \mathbf{Cof}(kG)^\perp)$ is a hereditary cotorsion pair in the category of kG -modules, which is cogenerated by a set for any group algebra kG ; in particular, that cotorsion pair is complete. In this way, we obtain the existence of Gorenstein projective precovers for modules over a big class of group rings. For example, we have the following result:

Theorem A. *Let G be an $\mathbf{LH}\mathfrak{F}$ -group and k a ring of finite weak global dimension. Then, the cotorsion pair $(\mathbf{GProj}(kG), \mathbf{GProj}(kG)^\perp)$ is complete.*

The completeness of the cotorsion pair $(\mathbf{Cof}(kG), \mathbf{Cof}(kG)^\perp)$ can be also used to shed some light to the equality between the classes $\mathbf{Cof}(kG)$ and $\mathbf{GProj}(kG)$, providing us with a quantitative measure for its failure. More precisely, the embedding $\mathbf{Cof}(kG) \hookrightarrow \mathbf{GProj}(kG)$ induces an embedding between the corresponding stable categories $\underline{\mathbf{Cof}}(kG) \hookrightarrow \underline{\mathbf{GProj}}(kG)$. We view the Verdier quotient $\underline{\mathbf{GProj}}(kG)/\underline{\mathbf{Cof}}(kG)$ as a measure for the obstruction to the equality between cofibrant and Gorenstein projective kG -modules. We also consider the full subcategory

\mathcal{T} of the stable category $\underline{\mathbf{GProj}}(kG)$, consisting of those Gorenstein projective kG -modules which are right orthogonal to cofibrant modules.¹ We prove the following result:

Theorem B. *There is an equivalence of triangulated categories $\underline{\mathbf{GProj}}(kG)/\underline{\mathbf{Cof}}(kG) \simeq \mathcal{T}$ for any group algebra kG .*

Here is a description of the contents of the paper: Following the preliminary section, in Section 2, we elaborate on the relation between cofibrant and Gorenstein projective modules over group algebras and obtain some extensions of known results, regarding the equality between these classes. In Section 3, we prove that the class of cofibrant modules is a Kaplansky class and conclude that $(\mathbf{Cof}(kG), \mathbf{Cof}(kG)^\perp)$ is always a complete cotorsion pair. Finally, in Section 4, we describe the application to the completeness of the Gorenstein projective cotorsion pair and identify the Verdier quotient that measures the failure to the equality between cofibrant and Gorenstein projective modules with a suitable subcategory of the stable category of Gorenstein projective modules.

Notations and terminology. Unless otherwise specified, we work with modules over the group algebra kG of a group G with coefficients in a commutative unital ring k .

1. PRELIMINARIES

In this section, we record certain prerequisite notions that are used throughout the paper. These notions concern basic definitions regarding modules over group algebras, cotorsion pairs and stable categories.

I. MODULES OVER GROUP ALGEBRAS. We consider a commutative ring k and let G be a group. Using the diagonal action of G , the tensor product $M \otimes_k N$ of two kG -modules M, N is also a kG -module; we let $g \cdot (x \otimes y) = gx \otimes gy \in M \otimes_k N$ for any $g \in G$, $x \in M$ and $y \in N$. If the kG -module M is projective and N is projective as a k -module, then the diagonal kG -module $M \otimes_k N$ is projective as well; cf. [4, Chapter III, Corollary 5.7]. We also note that for any two kG -modules M, N the k -module $\mathrm{Hom}_k(N, M)$ admits the structure of a kG -module, with the group G acting diagonally; we now let $(g \cdot f)(x) = gf(g^{-1}x) \in M$ for any $g \in G$, $f \in \mathrm{Hom}_k(N, M)$ and $x \in N$. For any three kG -modules L, M, N the natural isomorphism of k -modules

$$\mathrm{Hom}_k(L \otimes_k N, M) \simeq \mathrm{Hom}_k(N, \mathrm{Hom}_k(L, M))$$

is actually an isomorphism of kG -modules, where G acts diagonally on the tensor product and the Hom-groups.

Let $B(G, \mathbb{Z})$ be the $\mathbb{Z}G$ -module consisting of all bounded functions from G to \mathbb{Z} , introduced in [21]. The $\mathbb{Z}G$ -module $B(G, \mathbb{Z})$ is \mathbb{Z} -free; in fact, $B(G, \mathbb{Z})$ is $\mathbb{Z}H$ -free for any finite subgroup $H \subseteq G$. If n is an integer, then the constant function $\iota(n) \in B(G, \mathbb{Z})$ with value n is invariant under the action of G . The map $\iota : \mathbb{Z} \rightarrow B(G, \mathbb{Z})$ which is defined in this way is therefore $\mathbb{Z}G$ -linear. Moreover, ι is a \mathbb{Z} -split monomorphism; an additive splitting for ι may be obtained by evaluating functions at the identity element of G . The kG -module $B(G, k) = B(G, \mathbb{Z}) \otimes_{\mathbb{Z}} k$ may be identified with the kG -module of all functions from G to k that admit finitely many values. It is free as a kH -module for any finite subgroup $H \subseteq G$. We also note that ι induces a k -split kG -linear monomorphism $\iota \otimes 1 : k \rightarrow B(G, k)$. For simplicity of notation, we shall denote $B(G, k)$ by B and let $\overline{B} = \mathrm{coker}(\iota \otimes 1)$.

¹Here, *orthogonality* is understood either in terms of the Ext^1 -pairing in the module category or, equivalently, in terms of the $\underline{\mathbf{Hom}}$ -pairing in the stable category.

The analysis of the homological properties of the kG -module B defined above is manageable over the class $\mathbf{LH}\mathfrak{F}$ of hierarchically decomposable groups, that were introduced by Kropholler in [20]. The class $\mathbf{H}\mathfrak{F}$ is the smallest class of groups which contains the class \mathfrak{F} of finite groups and is such that whenever a group G admits a finite dimensional contractible G -CW-complex with stabilizers in $\mathbf{H}\mathfrak{F}$, then we also have $G \in \mathbf{H}\mathfrak{F}$. The class $\mathbf{LH}\mathfrak{F}$ of locally $\mathbf{H}\mathfrak{F}$ -groups consists of those groups, whose finitely generated subgroups are in $\mathbf{H}\mathfrak{F}$. All soluble groups, all groups of finite virtual cohomological dimension, all one-relator groups and all automorphism groups of Noetherian modules over commutative rings are $\mathbf{LH}\mathfrak{F}$ -groups. The class $\mathbf{LH}\mathfrak{F}$ is closed under extensions, ascending unions, free products with amalgamation and HNN-extensions. Another class of groups over which the homological behaviour of B is tame consists of the groups of type Φ , which were introduced in [26], in connection with the existence of a finite dimensional model for the classifying space for proper actions. We say that a group G is of type Φ (over k) if the class of kG -modules of finite projective dimension coincides with the class of those kG -modules that have finite projective dimension over any finite subgroup of G . Since the kG -module B is kH -projective over any finite subgroup $H \subseteq G$, it is clear that $\text{pd}_{kG} B < \infty$ if the group G is of type Φ over k .

II. COTORSION PAIRS. The Ext^1 -pairing induces an orthogonality relation between modules over any ring R . If \mathbf{S} is a class of modules, then the left orthogonal ${}^\perp\mathbf{S}$ of \mathbf{S} is the class consisting of those modules M , which are such that $\text{Ext}_R^1(M, S) = 0$ for all $S \in \mathbf{S}$. The right orthogonal \mathbf{S}^\perp of \mathbf{S} is the class consisting of those modules N , which are such that $\text{Ext}_R^1(S, N) = 0$ for all $S \in \mathbf{S}$. Any left orthogonal class is closed under continuous ascending filtrations, in the sense that we shall now explain.

Let \mathbf{T} be any class of modules. A continuous ascending filtration of a module M by modules in \mathbf{T} is a family of submodules $(M_\alpha)_{\alpha < \gamma}$ of M , which is indexed by an ordinal γ , such that:

- (i) $M_0 = 0$ and $M = \bigcup_{\alpha < \gamma} M_\alpha$,
- (ii) $M_\alpha \subseteq M_\beta$ whenever α, β are two ordinals with $\alpha < \beta < \gamma$,
- (iii) $M_\beta = \bigcup_{\alpha < \beta} M_\alpha$ for any limit ordinal $\beta < \gamma$ and
- (iv) $M_{\alpha+1}/M_\alpha \in \mathbf{T}$ for any ordinal α with $\alpha + 1 < \gamma$.

We denote by $\text{Filt-}\mathbf{T}$ the class of those modules which admit a continuous ascending filtration by modules in \mathbf{T} . Then, Eklof's lemma [13, Theorem 7.3.4] asserts that $\text{Filt-}({}^\perp\mathbf{S}) \subseteq {}^\perp\mathbf{S}$ for any class of modules \mathbf{S} .

If \mathbf{U} and \mathbf{V} are two classes of modules, then we say that the pair (\mathbf{U}, \mathbf{V}) is a cotorsion pair in the category of modules (cf. [13, Definition 7.1.2]) if $\mathbf{U} = {}^\perp\mathbf{V}$ and $\mathbf{U}^\perp = \mathbf{V}$. In that case, the class \mathbf{U} is closed under extensions, direct summands and (in view of Eklof's lemma) continuous ascending filtrations. The cotorsion pair (\mathbf{U}, \mathbf{V}) is called hereditary if $\text{Ext}_R^i(U, V) = 0$ for all $i > 0$ and all modules $U \in \mathbf{U}$ and $V \in \mathbf{V}$. Equivalently, the cotorsion pair is hereditary if \mathbf{U} is closed under kernels of epimorphisms or else if \mathbf{V} is closed under cokernels of monomorphisms. The cotorsion pair is complete if for any module M there exist short exact sequences

$$0 \longrightarrow V \longrightarrow U \longrightarrow M \longrightarrow 0 \quad \text{and} \quad 0 \longrightarrow M \longrightarrow V' \longrightarrow U' \longrightarrow 0,$$

where $U, U' \in \mathbf{U}$ and $V, V' \in \mathbf{V}$. The existence of these short exact sequences is also referred to by saying that any module admits a special \mathbf{U} -precover and a special \mathbf{V} -preenvelope respectively. The kernel of the cotorsion pair (\mathbf{U}, \mathbf{V}) is the intersection $\mathbf{U} \cap \mathbf{V}$; we say that the cotorsion pair is projective if its kernel coincides with the class of projective modules. The cotorsion pair (\mathbf{U}, \mathbf{V}) is cogenerated by a set of modules \mathbf{S} if $\mathbf{V} = \mathbf{S}^\perp$ (so that $\mathbf{U} = {}^\perp(\mathbf{S}^\perp)$). In that case, the left-hand class \mathbf{U} of the pair consists precisely of the direct summands of $\text{Filt-}(\mathbf{S} \cup \{R\})$ modules; cf.

[13, Corollary 7.3.5]. Eklof and Trlifaj have established in [9, Theorem 10] the completeness of any cotorsion pair which is cogenerated by a set of modules.

III. STABLE CATEGORIES AND LOCALIZATION. An exact additive category \mathcal{E} is said to be Frobenius if it has enough projective and enough injective objects and the classes of projective and injective objects coincide. For example, the category of all kG -modules is Frobenius, if G is a finite group and k is a field. The stable category $\underline{\mathcal{E}}$ of a Frobenius category \mathcal{E} has the same objects as \mathcal{E} , whereas its Hom-groups $\underline{\text{Hom}}(E, E')$ are the quotients of the corresponding Hom-groups $\text{Hom}(E, E')$ of \mathcal{E} , modulo the subgroup of those morphisms that factor through projective-injective objects. We note that the stable category $\underline{\mathcal{E}}$ of a Frobenius exact category \mathcal{E} is canonically a triangulated category; cf. [15, Theorem 2.6].

Following [23] and [27], we say that a sequence of triangulated functors between triangulated categories (in particular, between stable categories of Frobenius categories)

$$\mathcal{T}' \xrightarrow{f} \mathcal{T} \xrightarrow{g} \mathcal{T}''$$

is a localization sequence if the following hold:

- (i) f has a right adjoint $f_\rho : \mathcal{T} \rightarrow \mathcal{T}'$ and the composition $f_\rho \circ f$ is the identity on \mathcal{T}' ,
- (ii) g has a right adjoint $g_\rho : \mathcal{T}'' \rightarrow \mathcal{T}$ and the composition $g \circ g_\rho$ is the identity on \mathcal{T}'' and
- (iii) if X is an object of \mathcal{T} , then $gX = 0 \in \mathcal{T}''$ if and only if $X = fX'$ for some $X' \in \mathcal{T}'$.

In that case, we identify \mathcal{T}' with $\text{im } f$ via f and use g to obtain an equivalence of triangulated categories between the Verdier quotient \mathcal{T}/\mathcal{T}' and \mathcal{T}'' .

2. COFIBRANT AND GORENSTEIN PROJECTIVE MODULES

In this section, we describe a few properties of Benson's cofibrant modules, regarding their relation to Gorenstein projective modules. We also present a few criteria for these two classes of modules to coincide.

Following Benson [2], we say that a kG -module M is cofibrant if the (diagonal) kG -module $M \otimes_k B$ is projective; we denote by $\text{Cof}(kG)$ the class of cofibrant modules. Since all projective kG -modules are cofibrant, the following two conditions are easily seen to be equivalent for a kG -module N :

- (i) There is a non-negative integer n and an exact sequence of kG -modules

$$0 \longrightarrow M_n \longrightarrow \dots \longrightarrow M_1 \longrightarrow M_0 \longrightarrow N \longrightarrow 0,$$

such that $M_0, M_1, \dots, M_n \in \text{Cof}(kG)$.

- (ii) $\text{pd}_{kG}(N \otimes_k B) < \infty$.

If these conditions are satisfied, we say that the kG -module N has finite cofibrant dimension; we denote by $\overline{\text{Cof}}(kG)$ the class of these kG -modules. It is clear that $\text{Cof}(kG) \subseteq \overline{\text{Cof}}(kG)$. An interesting criterion for a $\overline{\text{Cof}}(kG)$ -module to be cofibrant was obtained by Benson in [loc.cit.]. For the reader's convenience, we provide a detailed account of Benson's argument below.

Lemma 2.1. (cf. [2, Lemma 4.5(ii)]) *Let $M \in \overline{\text{Cof}}(kG)$ and assume that $\text{Ext}_{kG}^1(M, N) = 0$ for any kG -module N of finite projective dimension. Then, $M \in \text{Cof}(kG)$.*

Proof. Assuming that M is not cofibrant, let $n = \text{pd}_{kG}(M \otimes_k B) > 0$ and consider a short exact sequence of kG -modules

$$(1) \quad 0 \longrightarrow K \longrightarrow P \longrightarrow M \longrightarrow 0,$$

where P is projective. Tensoring that exact sequence with the k -free kG -module B , it follows that $\text{pd}_{kG}(K \otimes_k B) = n - 1$. We note that there is a commutative diagram with exact rows

$$\begin{array}{ccccccccc} 0 & \longrightarrow & K \otimes_k \overline{B} & \xrightarrow{i} & P \otimes_k \overline{B} & \longrightarrow & M \otimes_k \overline{B} & \longrightarrow & 0 \\ & & \varphi \downarrow & & f \downarrow & & \parallel & & \\ 0 & \longrightarrow & M & \xrightarrow{j} & M \otimes_k B & \longrightarrow & M \otimes_k \overline{B} & \longrightarrow & 0 \end{array}$$

Here, the top row is obtained by tensoring (1) with the k -free kG -module \overline{B} , the bottom row is obtained by tensoring with M the k -split short exact sequence of kG -modules

$$0 \longrightarrow k \longrightarrow B \longrightarrow \overline{B} \longrightarrow 0,$$

f is obtained from the projectivity of the kG -module $P \otimes_k \overline{B}$ and φ is the restriction of f . We also consider a projective kG -module Q admitting a surjective kG -linear map $\pi : Q \longrightarrow M$ and the commutative diagram with exact rows

$$\begin{array}{ccccccccc} 0 & \longrightarrow & Q \oplus (K \otimes_k \overline{B}) & \xrightarrow{1 \oplus \varphi} & Q \oplus (P \otimes_k \overline{B}) & \longrightarrow & M \otimes_k \overline{B} & \longrightarrow & 0 \\ & & \varphi' \downarrow & & f' \downarrow & & \parallel & & \\ 0 & \longrightarrow & M & \xrightarrow{j} & M \otimes_k B & \longrightarrow & M \otimes_k \overline{B} & \longrightarrow & 0 \end{array}$$

where $\varphi' = (\pi, \varphi)$ and $f' = (j \circ \pi, f)$. Since φ' is surjective, an application of the snake lemma shows that f' is also surjective and $\ker \varphi' \simeq \ker f'$. Our assumption that $M \in \overline{\text{Cof}}(kG)$ and the projectivity of $Q \oplus (P \otimes_k \overline{B})$ imply that the kG -module $\ker \varphi' \simeq \ker f'$ has finite projective dimension. Invoking the additional hypothesis on M , we conclude that the epimorphism φ' splits, so that M is a direct summand of $Q \oplus (K \otimes_k \overline{B})$. Hence, $M \otimes_k B$ is a direct summand of $[Q \oplus (K \otimes_k \overline{B})] \otimes_k B = (Q \otimes_k B) \oplus (K \otimes_k \overline{B} \otimes_k B)$. This is a contradiction, since $M \otimes_k B$ has projective dimension equal to n , whereas $(Q \otimes_k B) \oplus (K \otimes_k \overline{B} \otimes_k B)$ has projective dimension $\leq n - 1$. \square

Cofibrant modules are closely related to Gorenstein projective modules. Recall that an acyclic complex of projective kG -modules is called totally acyclic if it remains acyclic after applying the functor $\text{Hom}_{kG}(_, P)$ for any projective kG -module P . A kG -module M is called Gorenstein projective if it is a cokernel of a totally acyclic complex of projective kG -modules. Let $\text{GProj}(kG)$ denote the class of Gorenstein projective kG -modules. We refer to [13, 16] for this notion (which may be, of course, developed over any associative ring). Using an elegant construction, Cornick and Kropholler have shown in [5, Theorem 3.5] that any cofibrant module is Gorenstein projective, so that we always have $\text{Cof}(kG) \subseteq \text{GProj}(kG)$. We now list some basic properties of the class $\text{Cof}(kG)$ of cofibrant modules.

Proposition 2.2. (i) $\text{Cof}(kG) = \overline{\text{Cof}}(kG) \cap \text{GProj}(kG)$.

(ii) The class $\text{Cof}(kG)$ is closed under extensions, kernels of epimorphisms, direct summands and continuous ascending filtrations.

(iii) Any cofibrant module is a cokernel of an acyclic complex of projective kG -modules, all of whose cokernels are cofibrant.

Proof. (i) Since $\text{Cof}(kG) \subseteq \overline{\text{Cof}}(kG)$ and $\text{Cof}(kG) \subseteq \text{GProj}(kG)$, it is clear that $\text{Cof}(kG) \subseteq \overline{\text{Cof}}(kG) \cap \text{GProj}(kG)$. Conversely, if $M \in \overline{\text{Cof}}(kG) \cap \text{GProj}(kG)$, then $\text{Ext}_{kG}^1(M, N) = 0$ for any kG -module N of finite projective dimension; cf. [16, Theorem 2.20]. Hence, Lemma 2.1 implies that M is cofibrant, as needed.

(ii) We consider a short exact sequence of kG -modules

$$0 \longrightarrow M' \longrightarrow M \longrightarrow M'' \longrightarrow 0$$

and assume that M'' is cofibrant. Then, the kG -module $M'' \otimes_k B$ is projective and hence the associated short exact sequence of kG -modules

$$0 \longrightarrow M' \otimes_k B \longrightarrow M \otimes_k B \longrightarrow M'' \otimes_k B \longrightarrow 0$$

splits. We conclude that the kG -module $M' \otimes_k B$ is projective if and only if the kG -module $M \otimes_k B$ is projective. In other words, M' is cofibrant if and only if M is cofibrant.

If the kG -module N is a direct summand of a cofibrant module N' , then $N \otimes_k B$ is a direct summand of the projective kG -module $N' \otimes_k B$. It follows that the kG -module $N \otimes_k B$ is itself projective, so that N is cofibrant. Hence, $\mathbf{Cof}(kG)$ is closed under direct summands.

Finally, let $(L_\alpha)_{\alpha < \tau}$ be a continuous ascending filtration of a kG -module L by cofibrant modules. Then, $(L_\alpha \otimes_k B)_{\alpha < \tau}$ is a continuous ascending filtration of the kG -module $L \otimes_k B$ by projective kG -modules. Since the class $\mathbf{Proj}(kG)$ of projective kG -modules is closed under continuous ascending filtrations, it follows that $L \otimes_k B \in \mathbf{Proj}(kG)$, so that L is cofibrant.

(iii) Let M be a cofibrant module. Since all projective kG -modules are cofibrant, it follows from (ii) above that the cokernels of any projective resolution of M are cofibrant. This provides us with the left half of the required acyclic complex of projective kG -modules. On the other hand, being cofibrant, the kG -module M is Gorenstein projective. Hence, there exists a short exact sequence of kG -modules

$$0 \longrightarrow M \longrightarrow P \longrightarrow N \longrightarrow 0,$$

where P is projective (and hence cofibrant) and N is Gorenstein projective. It follows readily that $N \in \overline{\mathbf{Cof}}(kG)$, so that (i) above implies that the kG -module N is actually cofibrant. We may repeat the same argument with N in the place of M and construct inductively the right half of the required acyclic complex of projective kG -modules, by splicing together short exact sequences with cofibrant end terms and projective middle term. \square

Let us denote by $\mathcal{P}(kG)$ the class of those kG -modules that appear as cokernels of acyclic complexes of projective kG -modules. It is clear that $\mathbf{GProj}(kG) \subseteq \mathcal{P}(kG)$ and equality holds if and only if any acyclic complex of projective kG -modules is totally acyclic. It is conjectured in [8] that $\mathcal{P}(\mathbb{Z}G) \subseteq \mathbf{Cof}(\mathbb{Z}G)$, so that $\mathbf{Cof}(\mathbb{Z}G) = \mathbf{GProj}(\mathbb{Z}G) = \mathcal{P}(\mathbb{Z}G)$. If these equalities hold, then the study of Gorenstein projective $\mathbb{Z}G$ -modules is reduced to questions involving classical homological algebra notions; in particular, Gorenstein projectivity of modules is then a subgroup-closed property. It is proved in [loc.cit., Corollary C] that the equalities above do hold for any group in Kropholler's class $\mathbf{LH}\mathfrak{F}$ of hierarchically decomposable groups. Working over more general rings of coefficients, it is proved in [3, Corollary 5.5] that $\mathcal{P}(kG) \subseteq \mathbf{Cof}(kG)$, if the ring k has finite global dimension and G is either an $\mathbf{LH}\mathfrak{F}$ -group or a group of type Φ . The latter assumption on k can be slightly relaxed, as we shall explain below.

We recall that a k -module is pure-projective if it is a direct summand of a suitable direct sum of finitely presented k -modules. The following two conditions are easily seen to be equivalent:

- (i) All pure-projective k -modules have finite projective dimension.
- (ii) All finitely presented k -modules have finite projective dimension and there is an upper bound on the projective dimension of these finitely presented k -modules.

Lemma 2.3. *Assume that k satisfies one of the following two conditions:*

- (i) *k has finite weak global dimension,*
- (ii) *all pure-projective k -modules have finite projective dimension.*

Then, any acyclic complex of projective k -modules is contractible.

Proof. The conclusion under hypothesis (i) is an immediate consequence of Neeman's result [24, Remark 2.15] on the contractibility of pure acyclic complexes of projective modules; cf.

[11, Proposition 1.1]. The conclusion holds under hypothesis (ii), by applying [12, Remark 2.11(i)] to the cotorsion pair $(\mathbf{Proj}(k), k\text{-Mod})$. \square

Proposition 2.4. *Assume that any acyclic complex of projective k -modules is contractible and let G be either an $\mathbf{LH}\mathfrak{F}$ -group or a group of type Φ over k . Then, $\mathcal{P}(kG) \subseteq \mathbf{Cof}(kG)$ and hence $\mathbf{Cof}(kG) = \mathbf{GProj}(kG) = \mathcal{P}(kG)$.*

Proof. Let M be a cokernel of an acyclic complex of projective kG -modules. Since projective kG -modules are k -projective, our hypothesis on k implies that M is k -projective.

We assume that G is an $\mathbf{LH}\mathfrak{F}$ -group. In order to show that M is cofibrant, i.e. that the kG -module $M \otimes_k B$ is projective, we follow verbatim the arguments in the proof of [8, Theorem B] and prove that $M \otimes_k B$ is projective over any $\mathbf{LH}\mathfrak{F}$ -subgroup of G . The k -projectivity of M is needed for the first step of the transfinite induction therein, showing that $M \otimes_k B$ is projective over any $\mathbf{H}\mathfrak{F}$ -subgroup of G , as it enables us to conclude that $M \otimes_k B$ is kH -projective for any finite subgroup $H \subseteq G$.

We now assume that G is a group of type Φ over k and note that the kG -module B has finite projective dimension. Since the kG -module M is k -projective, it follows that $\mathrm{pd}_{kG}(M \otimes_k B) \leq \mathrm{pd}_{kG} B$. This inequality holds for any kG -module M that appears as a cokernel of an acyclic complex of projective kG -modules and hence a simple argument shows that the kG -module $M \otimes_k B$ is actually projective; cf. [8, Lemma 2.1(c)]. \square

Corollary 2.5. *Assume that k satisfies one of the following two conditions:*

- (i) *k has finite weak global dimension,*
- (ii) *all pure-projective k -modules have finite projective dimension.*

If G is an $\mathbf{LH}\mathfrak{F}$ -group or a group of type Φ over k , then $\mathbf{Cof}(kG) = \mathbf{GProj}(kG) = \mathcal{P}(kG)$.

Remark 2.6. If k is \aleph_n -Noetherian for some integer $n \geq 0$, then the hypothesis in Corollary 2.5 that all pure-projective k -modules have finite projective dimension implies that the global dimension of k is finite (so that the corresponding assertion in Corollary 2.5 can be recovered by [3, Corollary 5.5]). To verify this claim, we may fix a positive integer s , such that $\mathrm{pd}_k C \leq s$ for any finitely presented k -module C . Considering cyclic finitely presented k -modules, it follows that $\mathrm{pd}_k I \leq s - 1$ for any finitely generated ideal $I \subseteq k$. Using the \aleph_n -Noetherian hypothesis on k and invoking [12, Corollary 2.7], we conclude that $\mathrm{id}_k N \leq s + n + 1$ for any k -module N and hence $\mathrm{gl.dim} k \leq s + n + 1$.

We can describe a condition implying that $\mathcal{P}(kG) \subseteq \mathbf{Cof}(kG)$, in terms of the class $\mathbf{PProj}(kG)$ of pure-projective kG -modules. As before, we note that the following conditions are equivalent:

(i) Pure-projective kG -modules have finite cofibrant dimension, i.e. $\mathbf{PProj}(kG) \subseteq \overline{\mathbf{Cof}}(kG)$.

(ii) Finitely presented kG -modules have finite cofibrant dimension and there is an upper bound on $\mathrm{pd}_{kG}(C \otimes_k B)$, where C runs through the finitely presented kG -modules.

Our goal now is to show that these equivalent conditions imply that $\mathcal{P}(kG) \subseteq \mathbf{Cof}(kG)$. To that end, we introduce some notations. We fix a set \mathbf{C} of representatives of the isomorphism classes of finitely presented kG -modules and choose for any kG -module $C \in \mathbf{C}$ a projective resolution of C with syzygy modules $(\Omega_i C)_{i \geq 0}$. We define the set $\mathbf{S} = \{\Omega_i C : C \in \mathbf{C}, i \geq 0\}$ and let $\mathbf{S}_n = \Omega_n \mathbf{S} = \{\Omega_i C : C \in \mathbf{C}, i \geq n\}$ for any $n \geq 0$. Then, the right Ext^1 -orthogonal \mathbf{S}^\perp is the class $\mathbf{SfpInj}(kG)$ of strongly fp-injective kG -modules, introduced in [22]. More generally, for any $n \geq 0$, the right Ext^1 -orthogonal \mathbf{S}_n^\perp is the class $\mathbf{SfpInj}_n(kG)$ of those kG -modules that admit an injective resolution whose n -th cosyzygy module is strongly fp-injective. We are interested in the cotorsion pair cogenerated by \mathbf{S}_n

$$(\perp(\mathbf{S}_n^\perp), \mathbf{S}_n^\perp) = (\mathbf{C}_n, \mathbf{SfpInj}_n(kG)).$$

In view of [13, Corollary 7.3.5], the class \mathcal{C}_n consists precisely of the direct summands of $\text{Filt-}\mathbf{S}_n$ modules. The relevance of the classes $(\mathcal{C}_n)_n$ to our problem stems from the fact that

$$\mathcal{P}(kG) \subseteq \bigcap_n \mathcal{C}_n;$$

this is proved in [10, Corollary 4.9(ii)].

Theorem 2.7. *If $\text{PProj}(kG) \subseteq \overline{\text{Cof}}(kG)$, then $\text{Cof}(kG) = \text{GProj}(kG) = \mathcal{P}(kG)$.*

Proof. In view of the discussion above, it suffices to prove that $\mathcal{P}(kG) \subseteq \text{Cof}(kG)$. We note that our assumption implies the existence of an integer $n \geq 0$, such that $\text{pd}_{kG}(C \otimes_k B) \leq n$ for any finitely presented kG -module C . It follows readily from the definition of the set \mathbf{S}_n that the kG -module $S \otimes_k B$ is projective for all $S \in \mathbf{S}_n$, so that $\mathbf{S}_n \subseteq \text{Cof}(kG)$. Since the class of cofibrant modules is closed under direct summands and continuous ascending filtrations (cf. Proposition 2.2(ii)) and \mathcal{C}_n consists of direct summands of $\text{Filt-}\mathbf{S}_n$ modules (cf. [13, Corollary 7.3.5]), it follows that $\mathcal{C}_n \subseteq \text{Cof}(kG)$. We may now invoke the fact that $\mathcal{P}(kG) \subseteq \mathcal{C}_n$ (cf. [10, Corollary 4.9(ii)]) and conclude that $\mathcal{P}(kG) \subseteq \text{Cof}(kG)$, as needed. \square

We briefly elaborate on the hypothesis of Theorem 2.7. If all kG -modules have finite cofibrant dimension, then there is clearly an upper bound on the projective dimension of the kG -modules $M \otimes_k B$, as M runs through the class of all kG -modules, and the inclusion $\mathcal{P}(kG) \subseteq \text{Cof}(kG)$ is immediate.² As in Remark 2.6, we show that the hypothesis of Theorem 2.7 is equivalent, in some cases, to the apparently stronger assumption that all kG -modules have finite cofibrant dimension.

Lemma 2.8. *If kG is left \aleph_n -Noetherian for some integer $n \geq 0$, then the following conditions are equivalent:*

- (i) *All pure-projective kG -modules have finite cofibrant dimension.*
- (ii) *All kG -modules have finite cofibrant dimension.*

Proof. It suffices to prove that (i) \rightarrow (ii). To that end, assume that (i) holds and fix a positive integer s , such that $\text{pd}_{kG}(C \otimes_k B) \leq s$ for any finitely presented kG -module C . In particular, by considering cyclic finitely presented kG -modules, we conclude that $\text{pd}_{kG}(I \otimes_k B) \leq s - 1$ for any finitely generated left ideal $I \subseteq kG$. Using the standard Hom-tensor adjunction for kG -modules with diagonal action, it follows that the abelian groups

$$\text{Ext}_{kG}^i(I, \text{Hom}_k(B, N)) \simeq \text{Ext}_{kG}^i(I \otimes_k B, N)$$

are trivial for any $i > s - 1$, any finitely generated left ideal $I \subseteq kG$ and any kG -module N . Using the left \aleph_n -Noetherian hypothesis on kG and invoking [12, Corollary 2.7], we conclude that $\text{id}_{kG} \text{Hom}_k(B, N) \leq s + n + 1$ for any kG -module N . It follows that the abelian groups

$$\text{Ext}_{kG}^i(M \otimes_k B, N) \simeq \text{Ext}_{kG}^i(M, \text{Hom}_k(B, N))$$

are trivial for any $i > s + n + 1$ and any kG -modules M, N . Hence, $\text{pd}_{kG}(M \otimes_k B) \leq s + n + 1$ for all kG -modules M , so that all kG -modules have finite cofibrant dimension. \square

Remarks 2.9. (i) Assuming that $\text{Cof}(kG) = \text{GProj}(kG)$, the assertions stated in Proposition 2.2(ii) represent well-known properties of Gorenstein projective modules, whereas Proposition 2.2(iii) is essentially a consequence of the very definition of Gorenstein projective modules. Since all kG -modules that have finite cofibrant dimension have also finite Gorenstein projective

²If X is an acyclic complex of projective kG -modules with cokernels $(C_i X)_i$, then $X \otimes_k B$ is an acyclic complex of projective kG -modules with cokernels $(C_i X \otimes_k B)_i$ and hence $C_i X \otimes_k B$ is a projective kG -module, being the n -th syzygy of $C_{i-n} X \otimes_k B$ for all i and all $n \geq 1$.

dimension, the equality $\mathbf{Cof}(kG) = \mathbf{GProj}(kG)$ reduces Lemma 2.1 to the assertion that, within the class of kG -modules of finite Gorenstein projective dimension, the Gorenstein projective kG -modules are precisely those kG -modules which are left Ext^1 -orthogonal to the kG -modules of finite projective dimension; see [16, Theorem 2.10].

ii) If all pure-projective kG -modules have finite cofibrant dimension, then all pure-projective kG -modules have finite Gorenstein projective dimension. Hence, [12, Theorem 2.3(ii)] implies that any acyclic complex of projective kG -modules is totally acyclic, i.e. $\mathcal{P}(kG) = \mathbf{GProj}(kG)$. In Theorem 2.7, we proved the stronger statement that $\mathcal{P}(kG) = \mathbf{GProj}(kG) = \mathbf{Cof}(kG)$.

iii) As Olympia Talelli has pointed out, the only properties of the kG -module B that are needed for the arguments above to go through are the following:

- (iii1) B is k -projective,
- (iii2) there exists a k -split kG -linear monomorphism $k \rightarrow B$ and
- (iii3) B is kH -projective over any finite subgroup $H \subseteq G$.

Indeed, properties (iii1) and (iii2) are used to show that all cofibrant modules are Gorenstein projective, whereas property (iii3) is the key ingredient to showing that $\mathcal{P}(kG) \subseteq \mathbf{Cof}(kG)$, under various additional assumptions on the pair (k, G) ; cf. the proof of Proposition 2.4.

3. THE COTORSION PAIR $(\mathbf{Cof}(kG), \mathbf{Cof}(kG)^\perp)$

In this section, we show that the class of cofibrant modules is a Kaplansky class and conclude that $(\mathbf{Cof}(kG), \mathbf{Cof}(kG)^\perp)$ is a cotorsion pair, which is cogenerated by a set of modules. We also describe some basic properties of the orthogonal class $\mathbf{Cof}(kG)^\perp$.

We recall from [14, Definition 2.1] that a class \mathcal{K} of kG -modules is called a Kaplansky class if there exists a cardinal number λ , such that for any module $M \in \mathcal{K}$ and any element $x \in M$ there exists a submodule $N \subseteq M$ with $x \in N$, $\text{card } N \leq \lambda$ and $N, M/N \in \mathcal{K}$. Given a cardinal number μ , we say that a kG -module is μ -generated if it can be generated by a set of cardinality $\leq \mu$.

Proposition 3.1. *The class $\mathbf{Cof}(kG)$ is a Kaplansky class.*

Proof. Let $\kappa = \text{card } k$, $\gamma = \text{card } G$ and $\lambda = \max\{\aleph_0, \kappa, 2^\gamma\}$. Then, λ is an infinite cardinal, which is greater than or equal to both $\text{card } kG$ and $\text{card } B$.³ Moreover, it is easily seen that for any λ -generated kG -module K (i.e. for any kG -module K with $\text{card } K \leq \lambda$) the kG -module $K \otimes_k B$ is also λ -generated (i.e. $\text{card}(K \otimes_k B) \leq \lambda$). We prove that this choice of λ works to establish that $\mathbf{Cof}(kG)$ is a Kaplansky class.

To that end, we consider a cofibrant module M and fix an element $x \in M$. Since $M \otimes_k B$ is a projective kG -module, a result by Kaplansky [19] implies that it can be expressed as the direct sum of a family of countably generated projective kG -modules. We fix such a decomposition and write $M \otimes_k B = \bigoplus_{i \in I} P_i$, for a suitable family $(P_i)_{i \in I}$ of countably generated projective kG -modules. For any subset $J \subseteq I$ we define $P(J) = \bigoplus_{i \in J} P_i$; in particular, $P(\emptyset) = 0$ and $P(I) = M \otimes_k B$. Using induction, we construct a sequence of pairs $(I_n, M_n)_n$, such that:

- (i) $(I_n)_n$ is an increasing sequence of subsets of I , with $\text{card } I_n \leq \lambda$ for all $n \geq 0$,
- (ii) $(M_n)_n$ is an increasing sequence of λ -generated kG -submodules of M ,
- (iii) $I_0 = \emptyset$ and $M_0 = kG \cdot x$,
- (iv) $P(I_n) \subseteq M_n \otimes_k B \subseteq P(I_{n+1}) \subseteq P(I) = M \otimes_k B$ for all $n \geq 0$.

Of course, we begin the induction by letting $I_0 = \emptyset$ and $M_0 = kG \cdot x$. Having defined the pairs $(I_0, M_0), (I_1, M_1), \dots, (I_n, M_n)$, we proceed with the inductive step and define (I_{n+1}, M_{n+1}) as

³The inequality $\text{card } B \leq \lambda$ follows since any element of B is a k -linear combination of suitable characteristic functions on subsets of G .

follows: Since the kG -module M_n is λ -generated, the kG -module $M_n \otimes_k B$ is also λ -generated. Expressing each generator of $M_n \otimes_k B \subseteq M \otimes_k B = \bigoplus_{i \in I} P_i$ as a finite sum of elements of the P_i 's, we conclude that there exists a subset $I_{n+1} \subseteq I$ which contains I_n and has cardinality $\leq \lambda$, such that $M_n \otimes_k B \subseteq P(I_{n+1})$. Since $\text{card } I_{n+1} \leq \lambda$ and the kG -module P_i is countably generated for all $i \in I$, the kG -module $P(I_{n+1})$ is λ -generated. Expressing each generator of $P(I_{n+1}) \subseteq P(I) = M \otimes_k B$ as a finite sum of elementary tensors, we conclude that there is a λ -generated kG -submodule $M_{n+1} \subseteq M$ containing M_n , such that $P(I_{n+1}) \subseteq M_{n+1} \otimes_k B$. This completes the inductive step of the construction.

Having constructed the sequence $(I_n, M_n)_n$ with properties (i)-(iv) above, we let $I' = \bigcup_n I_n$ and $M' = \bigcup_n M_n$. Then, $I' \subseteq I$ has cardinality $\leq \lambda$ and M' is a λ -generated kG -submodule of M , which contains M_0 (and hence x), such that

$$M' \otimes_k B = \bigcup_n (M_n \otimes_k B) = \bigcup_n P(I_n) = P(I') \subseteq P(I) = M \otimes_k B.$$

Here, the second equality follows from property (iv). In particular, the kG -module $M' \otimes_k B$ is projective and hence M' is cofibrant. The quotient module M/M' is also cofibrant, since

$$(M/M') \otimes_k B = (M \otimes_k B)/(M' \otimes_k B) = P(I)/P(I') = P(I \setminus I')$$

is a projective kG -module. We have therefore completed the proof that $\mathbf{Cof}(kG)$ is a Kaplansky class. \square

We shall use Proposition 3.1 (and its proof) to construct filtrations of cofibrant modules with successive quotients cofibrant modules of controlled size.

Proposition 3.2. *There is a cardinal number λ , such that any cofibrant module admits a continuous ascending filtration by λ -generated cofibrant modules.*

Proof. Let λ be the cardinal number defined in the proof of Proposition 3.1. We consider a cofibrant module M and fix a decomposition $M \otimes_k B = \bigoplus_{i \in I} P_i$ of the projective kG -module $M \otimes_k B$ into the direct sum of a family $(P_i)_{i \in I}$ of countably generated projective kG -modules, as in the proof of Proposition 3.1. As before, for any subset $J \subseteq I$ we define $P(J) = \bigoplus_{i \in J} P_i$. We also choose a well-ordered set of generators $(x_\alpha)_{\alpha < \tau}$ of the kG -module M , where τ is a suitable cardinal number. It only suffices to consider the case where $\tau > \lambda$, since otherwise M is already λ -generated. Using transfinite induction, we construct a family of pairs $(I_\alpha, M_\alpha)_{\alpha \leq \tau}$, such that:

- (i) $(I_\alpha)_{\alpha \leq \tau}$ is a continuous ascending family of subsets of I ,
- (ii) $\text{card}(I_{\alpha+1} \setminus I_\alpha) \leq \lambda$ for any $\alpha < \tau$,
- (iii) $(M_\alpha)_{\alpha \leq \tau}$ is a continuous ascending family of kG -submodules of M ,
- (iv) the kG -module $M_{\alpha+1}/M_\alpha$ is λ -generated for any $\alpha < \tau$,
- (v) $x_\alpha \in M_{\alpha+1}$ for any $\alpha < \tau$,
- (vi) $I_0 = \emptyset$ and $M_0 = 0$,
- (vii) $M_\alpha \otimes_k B = P(I_\alpha) \subseteq P(I) = M \otimes_k B$ for all $\alpha \leq \tau$.

Of course, we begin the induction, by letting $I_0 = \emptyset$ and $M_0 = 0$. For the inductive step of the proof, we assume that β is an ordinal and the pairs (I_α, M_α) have been constructed for all ordinals $\alpha < \beta$, so that properties (i)-(vii) above hold. In the case where β is a limit ordinal, we define $I_\beta = \bigcup_{\alpha < \beta} I_\alpha$ and $M_\beta = \bigcup_{\alpha < \beta} M_\alpha$. Then, properties (i)-(vii) still hold, since

$$M_\beta \otimes_k B = \bigcup_{\alpha < \beta} (M_\alpha \otimes_k B) = \bigcup_{\alpha < \beta} P(I_\alpha) = P(I_\beta) \subseteq P(I) = M \otimes_k B.$$

We now consider the case where $\beta = \alpha + 1$ is a successor ordinal and note that the quotient M/M_α is a cofibrant module, since

$$(2) \quad (M/M_\alpha) \otimes_k B = (M \otimes_k B)/(M_\alpha \otimes_k B) = P(I)/P(I_\alpha) = P(I \setminus I_\alpha)$$

is a projective kG -module. It follows from the proof of Proposition 3.1, applied to the cofibrant module M/M_α , its element $x_\alpha + M_\alpha$ and the decomposition of $(M/M_\alpha) \otimes_k B$ displayed in (2) above, that there exist a subset $I_{\alpha+1} \subseteq I$ containing I_α and a kG -submodule $M_{\alpha+1} \subseteq M$ containing M_α and x_α , such that $\text{card}(I_{\alpha+1} \setminus I_\alpha) \leq \lambda$, the kG -submodule $M_{\alpha+1}/M_\alpha \subseteq M/M_\alpha$ is λ -generated and $(M_{\alpha+1}/M_\alpha) \otimes_k B = P(I_{\alpha+1} \setminus I_\alpha)$. Since $M_\alpha \otimes_k B = P(I_\alpha)$ and

$$(M_{\alpha+1} \otimes_k B)/(M_\alpha \otimes_k B) = (M_{\alpha+1}/M_\alpha) \otimes_k B = P(I_{\alpha+1} \setminus I_\alpha),$$

it is easily seen that $M_{\alpha+1} \otimes_k B = P(I_{\alpha+1}) \subseteq P(I) = M \otimes_k B$. Indeed, $M_{\alpha+1} \otimes_k B$ is clearly contained in $P(I_{\alpha+1})$ and the embedding $\iota : M_{\alpha+1} \otimes_k B \hookrightarrow P(I_{\alpha+1})$ fits into the commutative diagram with exact rows

$$\begin{array}{ccccccc} 0 & \longrightarrow & M_\alpha \otimes_k B & \longrightarrow & M_{\alpha+1} \otimes_k B & \longrightarrow & (M_{\alpha+1} \otimes_k B)/(M_\alpha \otimes_k B) \longrightarrow 0 \\ & & \downarrow & & \downarrow \iota & & \downarrow \\ 0 & \longrightarrow & P(I_\alpha) & \longrightarrow & P(I_{\alpha+1}) & \longrightarrow & P(I_{\alpha+1} \setminus I_\alpha) \longrightarrow 0 \end{array}$$

whose end vertical arrows are both bijective. The inductive step of the construction is therefore complete.

Having constructed the family of pairs $(I_\alpha, M_\alpha)_{\alpha \leq \tau}$ as above, we note that the kG -submodule $M_\tau = \bigcup_{\alpha < \tau} M_\alpha$ contains all of the generators $(x_\alpha)_{\alpha < \tau}$ of M and hence $M_\tau = M$. Moreover, for any $\alpha < \tau$ the kG -module $(M_{\alpha+1}/M_\alpha) \otimes_k B = P(I_{\alpha+1} \setminus I_\alpha)$ is projective, so that $M_{\alpha+1}/M_\alpha$ is cofibrant. Hence, $(M_\alpha)_{\alpha \leq \tau}$ is the required continuous ascending filtration of M by λ -generated cofibrant modules. \square

We can now state and prove the main result of this section.

Theorem 3.3. *The pair $(\mathbf{Cof}(kG), \mathbf{Cof}(kG)^\perp)$ is a projective hereditary cotorsion pair in the category of kG -modules, which is cogenerated by a set. In particular, it is complete.*

Proof. In view of Proposition 3.2, there exists a cardinal number λ , such that any cofibrant module admits a continuous ascending filtration by λ -generated cofibrant modules. We consider a set \mathcal{S} of representatives of the isomorphism classes of all λ -generated cofibrant modules and prove that

$$(3) \quad (\mathbf{Cof}(kG), \mathbf{Cof}(kG)^\perp) = (\perp(\mathcal{S}^\perp), \mathcal{S}^\perp)$$

is the cotorsion pair cogenerated by \mathcal{S} . First of all, we shall prove that $\mathbf{Cof}(kG)^\perp = \mathcal{S}^\perp$. Since $\mathcal{S} \subseteq \mathbf{Cof}(kG)$, it follows that $\mathbf{Cof}(kG)^\perp \subseteq \mathcal{S}^\perp$. In order to prove the reverse inclusion, we consider a module $N \in \mathcal{S}^\perp$. Then, the functor $\text{Ext}_{kG}^1(-, N)$ vanishes on all modules in \mathcal{S} and hence on all λ -generated cofibrant modules. Since any cofibrant module admits a continuous ascending filtration by λ -generated cofibrant modules, Eklof's lemma [13, Theorem 7.3.4] implies that the functor $\text{Ext}_{kG}^1(-, N)$ vanishes on all cofibrant modules; hence, $N \in \mathbf{Cof}(kG)^\perp$. We have thus proved the equality $\mathbf{Cof}(kG)^\perp = \mathcal{S}^\perp$. We shall now prove that $\mathbf{Cof}(kG) = \perp(\mathcal{S}^\perp)$. We note that the equality $\mathbf{Cof}(kG)^\perp = \mathcal{S}^\perp$ implies that

$$\mathbf{Cof}(kG) \subseteq \perp(\mathbf{Cof}(kG)^\perp) = \perp(\mathcal{S}^\perp).$$

Conversely, \mathcal{S} is contained in $\mathbf{Cof}(kG)$ and the latter class is closed under direct summands and continuous ascending filtrations; cf. Proposition 2.2(ii). Then, [13, Corollary 7.3.5] shows that we also have $\perp(\mathcal{S}^\perp) \subseteq \mathbf{Cof}(kG)$.

Having proved (3), it follows from [9, Theorem 10] that $(\mathbf{Cof}(kG), \mathbf{Cof}(kG)^\perp)$ is a complete cotorsion pair. The cotorsion pair is hereditary, since the left-hand side class $\mathbf{Cof}(kG)$ is closed under kernels of epimorphisms; cf. Proposition 2.2(ii). It remains to show that the cotorsion pair is projective, i.e. that its kernel $\mathbf{Cof}(kG) \cap \mathbf{Cof}(kG)^\perp$ is the class of projective kG -modules. We note that all projective kG -modules are cofibrant and contained in $\mathbf{Cof}(kG)^\perp$; indeed, all projective kG -modules are contained in $\mathbf{Gproj}(kG)^\perp$ and $\mathbf{Cof}(kG) \subseteq \mathbf{GProj}(kG)$. Conversely, assume that $M \in \mathbf{Cof}(kG) \cap \mathbf{Cof}(kG)^\perp$. Since M is cofibrant, Proposition 2.2(iii) implies the existence of a short exact sequence of kG -modules

$$0 \longrightarrow M \longrightarrow P \longrightarrow N \longrightarrow 0,$$

where P is projective and N is cofibrant. Since $M \in \mathbf{Cof}(kG)^\perp$, the group $\mathrm{Ext}_{kG}^1(N, M)$ is trivial and hence the short exact sequence splits. In particular, M is a direct summand of the projective kG -module P , so that M is itself projective, as needed. \square

It follows from Theorem 3.3 that the class $\mathbf{Cof}(kG)^\perp$ is of some interest, as it determines the class of cofibrant modules as its left Ext^1 -orthogonal. We end this section by listing a few properties of that class.

Proposition 3.4. *The class $\mathbf{Cof}(kG)^\perp$ has the 2-out-of-3 property for short exact sequences. It is closed under direct summands and contains all kG -modules of finite projective or injective dimension and all (diagonal) kG -modules of the form $\mathrm{Hom}_k(B, N)$, where N is a kG -module.*

Proof. Since the cotorsion pair $(\mathbf{Cof}(kG), \mathbf{Cof}(kG)^\perp)$ is hereditary, $\mathbf{Cof}(kG)^\perp$ is closed under extensions and cokernels of monomorphisms. We prove that it is also closed under kernels of epimorphisms. To that end, let

$$(4) \quad 0 \longrightarrow N' \longrightarrow N \longrightarrow N'' \longrightarrow 0$$

be a short exact sequence of kG -modules and assume that $N, N'' \in \mathbf{Cof}(kG)^\perp$. In order to show that $N' \in \mathbf{Cof}(kG)^\perp$, we consider a cofibrant module M and invoke Proposition 2.2(iii) to find a short exact sequence of kG -modules

$$0 \longrightarrow M \longrightarrow P \longrightarrow M' \longrightarrow 0,$$

where P is projective and M' is cofibrant. Then, $\mathrm{Ext}_{kG}^1(M, N') = \mathrm{Ext}_{kG}^2(M', N')$ and hence it suffices to prove that the abelian group $\mathrm{Ext}_{kG}^2(M', N')$ is trivial. This follows from the long exact sequence associated with (4)

$$\cdots \longrightarrow \mathrm{Ext}_{kG}^1(M', N'') \longrightarrow \mathrm{Ext}_{kG}^2(M', N') \longrightarrow \mathrm{Ext}_{kG}^2(M', N) \longrightarrow \cdots,$$

since both groups $\mathrm{Ext}_{kG}^1(M', N'')$ and $\mathrm{Ext}_{kG}^2(M', N)$ are trivial.

The class $\mathbf{Cof}(kG)^\perp$ is clearly closed under direct summands. Since $\mathbf{Cof}(kG)^\perp$ contains all projective and all injective kG -modules, an inductive argument and the 2-out-of-3 property for short exact sequences established above show that $\mathbf{Cof}(kG)^\perp$ also contains all kG -modules of finite projective dimension and all kG -modules of finite injective dimension. Finally, if N is any kG -module, the Hom-tensor adjunction for diagonal kG -modules implies that the functor

$$\mathrm{Ext}_{kG}^1(-, \mathrm{Hom}_k(B, N)) = \mathrm{Ext}_{kG}^1(- \otimes_k B, N)$$

vanishes on all cofibrant modules and hence $\mathrm{Hom}_k(B, N) \in \mathbf{Cof}(kG)^\perp$. \square

4. SOME CONSEQUENCES

In this section, we present some applications of the results obtained earlier in the paper. In particular, we show that the class of Gorenstein projective kG -modules is precovering for a big class of group algebras. We also describe, over any group algebra, the obstruction to the equality between the classes of cofibrant and Gorenstein projective kG -modules, in terms of the size of a suitable subcategory of the stable category of Gorenstein projective kG -modules. Finally, we present an interpretation of Theorem 3.3 in the language of model categories.

I. THE COTORSION PAIR $(\mathbf{GProj}(kG), \mathbf{GProj}(kG)^\perp)$. As shown in [6], for any ring R the pair $(\mathbf{GProj}(R), \mathbf{GProj}(R)^\perp)$ is a hereditary cotorsion pair in the category of R -modules. A major problem in Gorenstein homological algebra is to show that this cotorsion pair is complete. In the terminology of [13], the problem is to show that any module admits special Gorenstein projective precovers. Using the results obtained in Section 3, we can show that this is indeed the case for a big class of group algebras.

Theorem 4.1. *Assume that k satisfies one of the following two conditions:*

- (i) *k has finite weak global dimension,*
- (ii) *all pure-projective k -modules have finite projective dimension.*

If G is an $\mathbf{LH}\mathfrak{S}$ -group or a group of type Φ over k , then the Gorenstein projective cotorsion pair $(\mathbf{GProj}(kG), \mathbf{GProj}(kG)^\perp)$ is cogenerated by a set; in particular, it is complete.

Proof. (i) In view of Corollary 2.5, there is an equality $\mathbf{GProj}(kG) = \mathbf{Cof}(kG)$ and hence

$$(\mathbf{GProj}(kG), \mathbf{GProj}(kG)^\perp) = (\mathbf{Cof}(kG), \mathbf{Cof}(kG)^\perp).$$

Therefore, the result follows from Theorem 3.3. \square

Remark 4.2. Assume that all pure-projective kG -modules have finite cofibrant dimension (i.e. that the hypothesis of Theorem 2.7 is satisfied). Since cofibrant modules are Gorenstein projective, it then follows that all pure-projective kG -modules have finite Gorenstein projective dimension. Hence, it follows from [12, Theorem 2.3(i)] that the Gorenstein projective cotorsion pair $(\mathbf{GProj}(kG), \mathbf{GProj}(kG)^\perp)$ is cogenerated by a set.

II. THE EQUALITY $\mathbf{GProj}(kG) = \mathbf{Cof}(kG)$. We show that the existence of special $\mathbf{Cof}(kG)$ precovers for Gorenstein projective kG -modules reduces the proof of the equality $\mathbf{GProj}(kG) = \mathbf{Cof}(kG)$ to proving that Gorenstein projective kG -modules which are contained in $\mathbf{Cof}(kG)^\perp$ are cofibrant.

Proposition 4.3. *The following conditions are equivalent:*

- (i) $\mathbf{GProj}(kG) = \mathbf{Cof}(kG)$,
- (ii) $\mathbf{GProj}(kG) \cap \mathbf{Cof}(kG)^\perp = \mathbf{Proj}(kG)$,
- (iii) $\mathbf{GProj}(kG) \cap \mathbf{Cof}(kG)^\perp \subseteq \mathbf{Cof}(kG)$,
- (iv) $\mathbf{GProj}(kG) \cap \mathbf{Cof}(kG)^\perp \subseteq \mathbf{GProj}(kG)^\perp$.

Proof. (i) \leftrightarrow (ii): If (i) holds, then

$$\mathbf{GProj}(kG) \cap \mathbf{Cof}(kG)^\perp = \mathbf{Cof}(kG) \cap \mathbf{Cof}(kG)^\perp = \mathbf{Proj}(kG),$$

where the last equality follows from Theorem 3.3. Conversely, assume that (ii) holds and let M be a Gorenstein projective kG -module. In view of the completeness of the cotorsion pair $(\mathbf{Cof}(kG), \mathbf{Cof}(kG)^\perp)$, there exists a short exact sequence of kG -modules

$$(5) \quad 0 \longrightarrow K \longrightarrow N \longrightarrow M \longrightarrow 0,$$

where N is cofibrant and $K \in \mathbf{Cof}(kG)^\perp$. Since M, N are Gorenstein projective and the class $\mathbf{GProj}(kG)$ is closed under kernels of epimorphisms (cf. [16, Theorem 2.5]), the kG -module K is Gorenstein projective as well. Then, $K \in \mathbf{GProj}(kG) \cap \mathbf{Cof}(kG)^\perp$ and hence K is projective. It follows that the short exact sequence (5) splits, so that M is a direct summand of N ; in particular, M is cofibrant.

It is clear that (ii) \rightarrow (iii) and (ii) \rightarrow (iv). The converse implications (iii) \rightarrow (ii) and (iv) \rightarrow (ii) follow easily, since both intersections $\mathbf{Cof}(kG) \cap \mathbf{Cof}(kG)^\perp$ and $\mathbf{GProj}(kG) \cap \mathbf{GProj}(kG)^\perp$ coincide with the class $\mathbf{Proj}(kG)$ of projective kG -modules. \square

Remarks 4.4. (i) The classes appearing in Proposition 4.3 may be schematically presented in the following diagram of subcategories of the module category $kG\text{-Mod}$

$$\begin{array}{ccc}
 \mathbf{GProj}(kG) & & \mathbf{Cof}(kG)^\perp \\
 \uparrow & & \uparrow \\
 \mathbf{Cof}(kG) & & \mathbf{GProj}(kG)^\perp \\
 & \swarrow \quad \searrow & \\
 & \mathbf{Proj}(kG) &
 \end{array}$$

Here, all arrows are inclusions.

(ii) Any kG -module $M \in \mathbf{GProj}(kG) \cap \mathbf{Cof}(kG)^\perp$ is a cokernel of an acyclic complex of projective kG -modules, all of whose cokernels are contained in $\mathbf{GProj}(kG) \cap \mathbf{Cof}(kG)^\perp$. Indeed, any totally acyclic complex of projective modules having M as a cokernel has all of its cokernels contained in $\mathbf{GProj}(kG) \cap \mathbf{Cof}(kG)^\perp$; this follows since $\mathbf{Proj}(kG) \subseteq \mathbf{Cof}(kG)^\perp$ and the class $\mathbf{Cof}(kG)^\perp$ has the 2-out-of-3 property for short exact sequences; cf. Proposition 3.4.

(iii) Let M be a Gorenstein projective kG -module. The existence of the short exact sequence (5), where $N \in \mathbf{Cof}(kG)$ and $K \in \mathbf{GProj}(kG) \cap \mathbf{Cof}(kG)^\perp$, shows that M is cofibrant if and only if the functor $\text{Ext}_{kG}^1(M, -)$ vanishes on all modules in $\mathbf{GProj}(kG) \cap \mathbf{Cof}(kG)^\perp$.

III. A LOCALIZATION SEQUENCE. We may elaborate on the completeness of the cotorsion pair $(\mathbf{Cof}(kG), \mathbf{Cof}(kG)^\perp)$ and obtain a measure for the obstruction to the equality $\mathbf{GProj}(kG) = \mathbf{Cof}(kG)$, in terms of the size of a suitable subcategory of the stable category of Gorenstein projective kG -modules.

Lemma 4.5. *The exact categories $\mathbf{Cof}(kG)$, $\mathbf{GProj}(kG)$ and $\mathbf{GProj}(kG) \cap \mathbf{Cof}(kG)^\perp$ (with the canonical exact structures inherited from the full category of kG -modules) are Frobenius, with projective-injective objects the projective kG -modules.*

Proof. The result for the exact category $\mathbf{GProj}(kG)$ is well-known; see, for example [7]. We prove that the exact category $\mathbf{Cof}(kG)$ is Frobenius. Since $\mathbf{Cof}(kG) \cap \mathbf{Cof}(kG)^\perp = \mathbf{Proj}(kG)$, all projective kG -modules are injective objects in $\mathbf{Cof}(kG)$. Proposition 2.2(iii) then implies that the exact category $\mathbf{Cof}(kG)$ has enough injective objects and all of these objects are projective kG -modules. Of course, all projective kG -modules are projective objects in $\mathbf{Cof}(kG)$. Using again Proposition 2.2(iii), it follows that $\mathbf{Cof}(kG)$ has enough projective objects and all of these objects are projective kG -modules. This completes the proof that the exact category $\mathbf{Cof}(kG)$ is Frobenius. An analogous argument, using Remark 4.4(ii) instead of Proposition 2.2(iii), shows that the exact category $\mathbf{GProj}(kG) \cap \mathbf{Cof}(kG)^\perp$ is Frobenius. \square

For simplicity of notation, we denote by \mathcal{T} the stable category of the Frobenius exact category $\mathbf{GProj}(kG) \cap \mathbf{Cof}(kG)^\perp$. Both \mathcal{T} and the stable category $\underline{\mathbf{Cof}}(kG)$ of cofibrant modules are full subcategories of the stable category $\underline{\mathbf{GProj}}(kG)$ of Gorenstein projective kG -modules.

Lemma 4.6. *Let $M \in \mathbf{Cof}(kG)$ and $N \in \mathbf{Cof}(kG)^\perp$. Then, any kG -linear map $M \rightarrow N$ factors through a projective kG -module, i.e. $\underline{\mathbf{Hom}}_{kG}(M, N) = 0$.⁴*

Proof. We fix a short exact sequence of kG -modules

$$0 \longrightarrow K \longrightarrow P \xrightarrow{p} N \longrightarrow 0,$$

where $K \in \mathbf{Cof}(kG)^\perp$ and P is a projective kG -module; cf. Proposition 3.4. Since the abelian group $\mathrm{Ext}_{kG}^1(M, K)$ is trivial, the additive map $p_* : \mathrm{Hom}_{kG}(M, P) \rightarrow \mathrm{Hom}_{kG}(M, N)$ is surjective. Then, any kG -linear map $M \rightarrow N$ factors through P . \square

Lemma 4.7. *Let $f : M \rightarrow M'$ be a kG -linear map, where M, M' are Gorenstein projective kG -modules. We also consider two short exact sequences of kG -modules*

$$0 \longrightarrow K \longrightarrow N \xrightarrow{p} M \longrightarrow 0 \quad \text{and} \quad 0 \longrightarrow K' \xrightarrow{p'} N' \xrightarrow{p'} M' \longrightarrow 0,$$

where $K, K' \in \mathbf{GProj}(kG) \cap \mathbf{Cof}(kG)^\perp$ and $N, N' \in \mathbf{Cof}(kG)$. Then:

- (i) *There exists a kG -linear map $g : N \rightarrow N'$, such that $p'g = fp$.*
- (ii) *If $g, g' : N \rightarrow N'$ are two kG -linear maps with $p'g = fp$ and $p'g' = fp$, then $[g] = [g'] \in \underline{\mathbf{Hom}}_{kG}(N, N')$.*
- (iii) *If $[f] = [0] \in \underline{\mathbf{Hom}}_{kG}(M, M')$ and $g : N \rightarrow N'$ is a kG -linear map with $p'g = fp$, then $[g] = [0] \in \underline{\mathbf{Hom}}_{kG}(N, N')$.*

Proof. (i) The additive map $p'_* : \mathrm{Hom}_{kG}(N, N') \rightarrow \mathrm{Hom}_{kG}(N, M')$ is surjective, since the abelian group $\mathrm{Ext}_{kG}^1(N, K')$ is trivial. Therefore, there exists a kG -linear map $g : N \rightarrow N'$ such that $fp = p'_*(g) = p'g$, as needed.

(ii) Let $g, g' : N \rightarrow N'$ be two kG -linear maps with $p'g = fp$ and $p'g' = fp$.

$$\begin{array}{ccccccccc} 0 & \longrightarrow & K & \longrightarrow & N & \xrightarrow{p} & M & \longrightarrow & 0 \\ & & & & & & & & \\ & & & & & g \downarrow \downarrow g' & & f \downarrow & \\ 0 & \longrightarrow & K' & \xrightarrow{p'} & N' & \xrightarrow{p'} & M' & \longrightarrow & 0 \end{array}$$

Then, $p'(g' - g) = p'g' - p'g = fp - fp = 0$ and hence there exists a kG -linear map $h : N \rightarrow K'$, such that $g' - g = t'h$. In view of Lemma 4.6, the kG -linear map h factors through a projective kG -module. Hence, this is also the case for $g' - g$; it follows that $[g] = [g'] \in \underline{\mathbf{Hom}}_{kG}(N, N')$.

(iii) Assume that f factors as the composition of two kG -linear maps $M \xrightarrow{a} Q \xrightarrow{b} M'$, where the kG -module Q is projective. If $\beta : Q \rightarrow N'$ is a kG -linear map with $p'\beta = b$, then the composition $\beta ap : N \rightarrow N'$ is such that $p'(\beta ap) = (p'\beta)ap = bap = fp$. It follows from (ii) above that $[g] = [\beta ap] \in \underline{\mathbf{Hom}}_{kG}(N, N')$. This finishes the proof, since we obviously have $[\beta ap] = [0] \in \underline{\mathbf{Hom}}_{kG}(N, N')$. \square

Lemma 4.7 implies that for any Gorenstein projective kG -module M the cofibrant module N that appears in a short exact sequence of kG -modules

$$0 \longrightarrow K \longrightarrow N \xrightarrow{p} M \longrightarrow 0$$

where $K \in \mathbf{GProj}(kG) \cap \mathbf{Cof}(kG)^\perp$ is uniquely determined up to a canonical isomorphism in the stable category $\underline{\mathbf{Cof}}(kG)$. Indeed, if

$$0 \longrightarrow K' \longrightarrow N' \xrightarrow{p'} M \longrightarrow 0$$

⁴Conversely, one can show that if N is a kG -module, such that $\underline{\mathbf{Hom}}_{kG}(M, N) = 0$ for any cofibrant module M , then $N \in \mathbf{Cof}(kG)^\perp$.

is another short exact sequence of kG -modules, where $K' \in \mathbf{GProj}(kG) \cap \mathbf{Cof}(kG)^\perp$ and N' is cofibrant, then there are unique morphisms $[g] \in \underline{\mathbf{Hom}}_{kG}(N, N')$ and $[h] \in \underline{\mathbf{Hom}}_{kG}(N', N)$, such that $p'g = p$ and $ph = p'$; cf. Lemma 4.7(i),(ii). The same uniqueness assertion shows that the compositions $[h] \cdot [g]$ and $[g] \cdot [h]$ are the identity morphisms of N and N' respectively. Moreover, Lemma 4.7(iii) implies that the assignment $M \mapsto N$ factors through the stable category $\underline{\mathbf{GProj}}(kG)$ and defines a functor

$$i^! : \underline{\mathbf{GProj}}(kG) \longrightarrow \underline{\mathbf{Cof}}(kG),$$

which is clearly additive.

Proposition 4.8. *The additive functor $i^! : \underline{\mathbf{GProj}}(kG) \longrightarrow \underline{\mathbf{Cof}}(kG)$ defined above is right adjoint to the inclusion functor $i_* : \underline{\mathbf{Cof}}(kG) \longrightarrow \underline{\mathbf{GProj}}(kG)$ and hence it is triangulated. In addition, the composition $i^! \circ i_*$ is the identity on $\underline{\mathbf{Cof}}(kG)$.*

Proof. We fix a cofibrant module M and let $L \in \mathbf{GProj}(kG)$. We also consider a short exact sequence of kG -modules

$$0 \longrightarrow K \longrightarrow N \xrightarrow{p} L \longrightarrow 0,$$

where $K \in \mathbf{GProj}(kG) \cap \mathbf{Cof}(kG)^\perp$ and $N \in \mathbf{Cof}(kG)$. We note that the additive map

$$(6) \quad [p]_* : \underline{\mathbf{Hom}}_{kG}(M, N) \longrightarrow \underline{\mathbf{Hom}}_{kG}(M, L)$$

is natural in both M (this is obvious) and L (this follows from Lemma 4.7(ii)). We establish the adjunction in the statement, by proving that the additive map (6) is bijective. Indeed, since the group $\text{Ext}_{kG}^1(M, K)$ is trivial, the additive map

$$p_* : \text{Hom}_{kG}(M, N) \longrightarrow \text{Hom}_{kG}(M, L)$$

is surjective, whence the surjectivity of (6). Regarding now the injectivity of (6), assume that $f : M \longrightarrow N$ is a kG -linear map, such that $[pf] = [p] \cdot [f] = [p]_*[f] = [0] \in \underline{\mathbf{Hom}}_{kG}(M, L)$. Then, we may consider the commutative diagram

$$\begin{array}{ccccccccc} 0 & \longrightarrow & 0 & \longrightarrow & M & \xrightarrow{1_M} & M & \longrightarrow & 0 \\ & & & & & f \downarrow & & pf \downarrow & \\ 0 & \longrightarrow & K & \longrightarrow & N & \xrightarrow{p} & L & \longrightarrow & 0 \end{array}$$

and invoke Lemma 4.7(iii), in order to conclude that $[f] = [0] \in \underline{\mathbf{Hom}}_{kG}(M, N)$.

Being right adjoint to the triangulated functor i_* , the functor $i^!$ is also triangulated; cf. [23, Lemma 5.3.6]. In order to verify that the composition $i^! \circ i_*$ is the identity on $\underline{\mathbf{Cof}}(kG)$, we simply note that for any cofibrant module M we can choose the approximation sequence

$$0 \longrightarrow 0 \longrightarrow M \xrightarrow{1_M} M \longrightarrow 0,$$

so that $i^!M = M$. □

Let M be a Gorenstein projective kG -module. Then, the completeness of the cotorsion pair $(\mathbf{Cof}(kG), \mathbf{Cof}(kG)^\perp)$ implies the existence of a short exact sequence of kG -modules

$$(7) \quad 0 \longrightarrow M \longrightarrow K \longrightarrow N \longrightarrow 0,$$

where $K \in \mathbf{Cof}(kG)^\perp$ and $N \in \mathbf{Cof}(kG)$. Since both M, N are Gorenstein projective, it follows that K is Gorenstein projective as well and hence $K \in \mathbf{GProj}(kG) \cap \mathbf{Cof}(kG)^\perp$. Working as above, we can show that K is uniquely determined by M up to a canonical isomorphism in the stable category \mathcal{T} and the assignment $M \mapsto K$ defines an additive functor

$$j^* : \underline{\mathbf{GProj}}(kG) \longrightarrow \mathcal{T},$$

which is left adjoint to the inclusion functor $j_* : \mathcal{T} \longrightarrow \underline{\mathbf{GProj}}(kG)$. In particular, the functor j^* is triangulated. Moreover, the composition $j^* \circ j_*$ is the identity on \mathcal{T} .

Lemma 4.9. *Let M be a Gorenstein projective kG -module. Then, $j^*M = 0 \in \mathcal{T}$ if and only if M is cofibrant.*

Proof. If $K \in \underline{\mathbf{GProj}}(kG) \cap \mathbf{Cof}(kG)^\perp$, then $K = 0 \in \mathcal{T}$ if and only if the identity map of K factors through a projective kG -module, i.e. if and only if $K \in \mathbf{Proj}(kG)$. Hence, we have to show that the Gorenstein projective kG -module M is cofibrant if and only if the kG -module K in the short exact sequence (7) is projective.

Indeed, if the kG -module K in (7) is projective, then it is cofibrant. Since the class $\mathbf{Cof}(kG)$ is closed under kernels of epimorphisms, we conclude that M is cofibrant as well. Conversely, if the kG -module M is cofibrant, then we may invoke Proposition 2.2(iii) and obtain a short exact sequence of kG -modules

$$0 \longrightarrow M \longrightarrow P \longrightarrow N \longrightarrow 0,$$

where P is projective and N is cofibrant. Since $\mathbf{Proj}(kG) \subseteq \underline{\mathbf{GProj}}(kG) \cap \mathbf{Cof}(kG)^\perp$, we may use the exact sequence above to compute j^*M . We conclude that $j^*M = P = 0 \in \mathcal{T}$. \square

We may summarize the discussion above in the form of the following result, establishing the existence of a localization sequence of triangulated categories; cf. [27].

Theorem 4.10. *The functors defined above induce a localization sequence*

$$\underline{\mathbf{Cof}}(kG) \xrightarrow{i_*} \underline{\mathbf{GProj}}(kG) \xrightarrow{j^*} \mathcal{T}.$$

The right adjoint of the inclusion i_ is $i^! : \underline{\mathbf{GProj}}(kG) \longrightarrow \underline{\mathbf{Cof}}(kG)$ and the right adjoint of j^* is the inclusion $j_* : \mathcal{T} \longrightarrow \underline{\mathbf{GProj}}(kG)$.*

The following result is a formal consequence of the existence of the localization sequence above; cf. [27].

Corollary 4.11. (i) *The functor j^* induces an equivalence of triangulated categories*

$$\underline{\mathbf{GProj}}(kG)/\underline{\mathbf{Cof}}(kG) \xrightarrow{\sim} \mathcal{T}.$$

(ii) *For any Gorenstein projective kG -module M there is an exact triangle in $\underline{\mathbf{GProj}}(kG)$*

$$(i_* \circ i^!)M \longrightarrow M \longrightarrow (j_* \circ j^*)M \rightsquigarrow$$

which depends functorially on M .

(iii) *A Gorenstein projective kG -module M is cofibrant if and only if $\underline{\mathbf{Hom}}_{kG}(M, K) = 0$ for any $K \in \underline{\mathbf{GProj}}(kG) \cap \mathbf{Cof}(kG)^\perp$.*

IV. A COFIBRANTLY GENERATED MODEL CATEGORY. We may also interpret Theorem 3.3 in the language of model categories. Following Quillen [25], a model category is a category with three distinguished classes of morphisms, called fibrations, cofibrations and weak equivalences, which satisfy a few axioms (which are reminiscent of certain properties of continuous maps on topological spaces). The homotopy category associated with a model category is obtained by formally inverting the weak equivalences, while keeping the same objects. We refer to [17] for the basic definitions and facts on model categories. The bijective correspondence between certain model structures and cotorsion pairs in an abelian category, which is established in [18, Theorem 2.2], provides us with a plethora of examples of model structures. In particular, Theorem 3.3 and Proposition 3.4 imply that the triple $(\mathbf{Cof}(kG), \mathbf{Cof}(kG)^\perp, kG\text{-Mod})$ may

be used to define a model structure in the category $kG\text{-Mod}$ of kG -modules, whose homotopy category is equivalent to the stable category $\underline{\mathbf{Cof}}(kG)$ of cofibrant kG -modules. The next (and final) result follows from general properties of model structures, using standard arguments.

Proposition 4.12. *There is a cofibrantly generated model structure in the category of kG -modules, in which the cofibrations (resp. the trivial cofibrations) are the monomorphisms whose cokernel is cofibrant (resp. the monomorphisms whose cokernel is projective), the fibrations (resp. the trivial fibrations) are the epimorphisms (resp. the epimorphisms with kernel in $\mathbf{Cof}(kG)^\perp$) and the weak equivalences are the kG -linear maps which factor as a trivial cofibration followed by a trivial fibration. The homotopy category of that model structure is equivalent to the stable category $\underline{\mathbf{Cof}}(kG)$ of cofibrant kG -modules.*

Remark 4.13. The model structure defined in Proposition 4.12 is different than the model structure defined by Benson in [2, Theorem 10.6]. The latter model structure is defined, in the special case where the coefficient ring k is Noetherian, on the category \mathcal{A} of countably presented kG -modules with finite cofibrant dimension. The fibrations therein are the epimorphisms in \mathcal{A} and the trivial fibrations are those fibrations whose kernel has finite projective dimension.

Acknowledgments. Wei Ren was supported by the National Natural Science Foundation of China (No. 11871125).

REFERENCES

- [1] Bahlekeh, A., Dembegiotti, F., Talelli, O.: Gorenstein dimension and proper actions. *Bull. London Math. Soc.* **41**, 859-871 (2009)
- [2] Benson, D.: Complexity and varieties for infinite groups I, II. *J. Algebra* **193**, 260-287, 288-317 (1997)
- [3] Biswas, R.: Benson's cofibrants, Gorenstein projectives and a related conjecture. *Proc. Edinb. Math. Soc.* **64**, 779-799 (2021)
- [4] Brown, K.S.: *Cohomology of Groups*, Graduate Texts in Mathematics 87, Springer, Berlin-Heidelberg-New York, 1982.
- [5] Cornick, J., Kropholler, P.H.: On complete resolutions, *Topology Appl.* **78**, 235-250 (1997)
- [6] Cortés-Izurdiaga, M., Saroch, J.: The cotorsion pair generated by the Gorenstein projective modules and λ -pure injective modules. *arXiv:2104.08602v2*
- [7] Dalezios, G., Estrada, S., Holm, H.: Quillen equivalences for stable categories. *J. Algebra* **501**, 130-149 (2018)
- [8] Dembegiotti, F., Talelli, O.: A note on complete resolutions, *Proc. Amer. Math. Soc.* **138**, 3815-3820 (2010).
- [9] Eklof, P., Trlifaj, J.: How to make Ext vanish. *Bull. London Math. Soc.* **23**, 41-51 (2001)
- [10] Emmanouil, I., Kaperonis, I.: On K-absolutely pure complexes. *J. Algebra* **640**, 274-299 (2024)
- [11] Emmanouil, I., Talelli, O.: Gorenstein dimension and group cohomology with group ring coefficients. *J. London Math. Soc.* **97**, 306-324 (2018)
- [12] Emmanouil, I., Talelli, O.: Triviality criteria for unbounded complexes. *J. Algebra* **663**, 786-814 (2025)
- [13] Enochs, E.E., Jenda, O.M.G.: *Relative Homological Algebra*. De Gruyter Expositions in Mathematics no. 30, New York: Walter De Gruyter, 2000
- [14] Enochs, E.E., Lopez-Ramos, J.A.: Kaplansky classes. *Rend. Sem. Mat. Univ. Padova* **107**, 67-79 (2002)
- [15] Happel, D.: *Triangulated Categories in the Representation Theory of Finite Dimensional Algebras*. Cambridge University Press, 1988
- [16] Holm, H.: Gorenstein homological dimensions. *J. Pure Appl. Algebra* **189**, 167-193 (2004)
- [17] Hovey, M.: *Model categories*. Mathematical Surveys and Monographs vol. 63, American Mathematical Society, 1999
- [18] Hovey, M.: Cotorsion pairs, model category structures and representation theory. *Math. Z.* **241**, 553-592 (2002)
- [19] Kaplansky, I.: Projective modules. *Ann. Math.* **68**, 372-377 (1958)
- [20] Kropholler, P.H.: On groups of type FP_∞ . *J. Pure Appl. Algebra* **90**, 55-67 (1993)

- [21] Kropholler, P.H., Talelli, O.: On a property of fundamental groups of graphs of finite groups. *J. Pure Appl. Algebra* **74**, 57-59 (1991)
- [22] Li, W., Guan, J., Ouyang, B.: Strongly FP-injective modules. *Comm. Algebra* **45**, 3816-3824 (2017)
- [23] Neeman, A.: *Triangulated Categories*. *Annals of Mathematics Studies* **148**, Princeton University Press, Princeton, NJ, 2001
- [24] Neeman, A.: The homotopy category of flat modules, and Grothendieck duality. *Invent. Math.* **174**, 255-308 (2008)
- [25] Quillen, D.: *Homotopical Algebra*. *Lecture Notes in Mathematics* no. **43**, Springer-Verlag, 1967
- [26] Talelli, O.: On groups of type Φ . *Arch. Math.* **89**, 24-32 (2007)
- [27] Verdier, J.L.: *Des catégories dérivées abéliennes*. *Asterisque* **239** (1996), xii+253 pp. (1997), with a preface by L. Illusie, edited and with a note by G. Maltsiniotis.

Ioannis Emmanouil

Department of Mathematics, University of Athens, Athens 15784, Greece

E-mail: emmanoui@math.uoa.gr

Wei Ren

School of Mathematical Sciences, Chongqing Normal University, Chongqing 401331, PR China

E-mail: wren@cqnu.edu.cn