POISSON HOMOGENEOUS SPACES OF POISSON 2-GROUPS

HONGLEI LANG AND ZHANGJU LIU

ABSTRACT. Drinfeld classified Poisson homogeneous spaces of a Poisson Lie group in terms of Dirac structures of the Lie bialgebra. In this paper, we study homogeneous spaces of a 2-group and develop Drinfeld theorem in the Poisson 2-group context.

CONTENTS

1.	Introduction	1
2.	Lie 2-group actions and representations	2
3.	Homogeneous spaces of Lie 2-groups	10
4.	Poisson homogeneous space of Poisson 2-groups	14
References		19

1. INTRODUCTION

Poisson Lie groups were introduced by Drinfeld as the semi-classical limit of quantum groups. A Poisson Lie group is a Lie group with a Poisson structure such that the group multiplication is a Poisson map. For the Lie algebra of the Lie group, its dual space is also endowed with a Lie algebra from the Poisson structure. This pair composes a Lie bialgebra. Drinfeld proved that there is a one-one correspondence between connected, simply connected Poisson Lie groups and Lie bialgebras.

A Poisson homogeneous space of a Poisson Lie group is a homogeneous space with a Poisson structure such that the action is a Poisson map. Every Poisson homogeneous space admits an integration to a symplectic groupoid [5]. The classification of Poisson homogeneous spaces was given by Drinfeld [8] in terms of Dirac structures (Lagrangian subalgebras) of the corresponding Lie bialgebra.

Drinfeld's classification theorem on homogeneous spaces was generalized from Poisson Lie groups to Poisson groupoids in [13]. A Poisson groupoid is a groupoid Γ with a Poisson structure Π such that the graph of groupoid multiplication $\{(g, h, gh)|(g, h) \in \Gamma^{(2)}\} \subset \Gamma \times \Gamma \times \overline{\Gamma}$ is coisotropic, where $\overline{\Gamma}$ denotes Γ with the opposite Poisson structure $-\Pi$. The infinitesimals of Poisson groupoids are Lie bialgebroids. In fact, there is a bijection between Poisson Lie groupoid (Γ, Π) and Lie bialgebroid (A, A^*) , where $\Gamma \Rightarrow M$ is an *s*-connected and simply connected Lie groupoid integrating the Lie algebroid A. The double $A \oplus A^*$ of a Lie bialgebroid is a Courant algebroid. In [13], a Dirac structure L of a Lie bialgebroid (A, A^*) was introduced as a subbundle of $A \oplus A^*$ which is maximally isotropic and integrable with respect to the natural pairing and Courant bracket. And it was proved that there is a bijection between Poisson homogeneous spaces of Poisson groupoids and regular Dirac structures of Lie bialgebroids. Here a homogeneous space of a Lie groupoid Γ is of the form $\Gamma/\!/\Gamma_1$, where Γ_1 is a wide (containing all the identities) subgroupoid of Γ . See [10] for homogeneous spaces of Dirac groupoids.

To describe symmetries of symmetries, the concept of (strict) Lie 2-groups was introduced, which is a Lie groupoid such that the total and base manifolds are Lie groups and the groupoid structure maps are Lie group homomorphisms; see [2]. Lie 2-groups can be identified with Lie group crossed modules. A Lie group crossed module \mathbb{G} is a pair of Lie groups G_0 and G_1 together with a Lie group homomorphism $G_1 \xrightarrow{\Phi} G_0$ and an action of G_0 on G_1 by automorphisms satisfying some compatibility conditions. A Poisson 2-group [7] is a Lie 2-group with a Poisson structure on the total space such that it is simultaneously a Poisson groupoid and a Poisson Lie group.

Keywords: Lie 2-group action, Poisson 2-group, homogeneous space

Lie 2-algebras are categorification of Lie algebras [1]. (Strict) Lie 2-algebras can be identified with Lie algebra crossed modules, which are the infinitesimal of Lie group crossed modules. Likewise, a Lie 2-bialgebra can be viewed as a pair of Lie algebra crossed modules $(\mathfrak{g}_1 \xrightarrow{\phi} \mathfrak{g}_0)$ and $(\mathfrak{g}_0^* \xrightarrow{-\phi^*} \mathfrak{g}_1^*)$ such that $(\mathfrak{g}_0 \ltimes \mathfrak{g}_1, \mathfrak{g}_1^* \ltimes \mathfrak{g}_0^*)$ is a Lie bialgebra [6]. See also [3]. In [7], the authors proved that there is a one-to-one correspondence between connected and simply connected Poisson 2-groups and Lie 2-bialgebras.

The goal of this paper is to prove the 2-categorical analogue of the Drinfeld classification theorem for Poisson homogeneous spaces. A Poisson homogeneous space of a Poisson 2-group is expected to be a homogeneous space of the Poisson Lie group and Poisson groupoid at the same time. However, note that a closed Lie 2-subgroup generally does not share the same base manifold with the original Lie 2-group. This leads to the inapplicability of the framework in [13] for homogeneous spaces of Poisson groupoids. This problem is tackled by considering a quotient Poisson groupoid. For a Lie 2-group $\mathbb{G} : G_1 \to G_0$ with a closed 2-subgroup $\mathbb{H} : H_1 \to H_0$, the group quotient \mathbb{G}/\mathbb{H} can also be viewed as a groupoid homogeneous space of the groupoid \mathbb{G}/H_0 . Suppose further that \mathbb{G} is a Poisson 2-group and $H_0 \subset \mathbb{G}$ is a Poisson subgroup so that \mathbb{G}/H_0 is a quotient Poisson groupoid. Then we are able to introduce Poisson homogeneous spaces of Poisson 2-groups and prove that there is a one-one correspondence between such Poisson homogeneous spaces and H_0 -invariant Dirac structures L of the Lie bialgebra $(\mathfrak{g}_1, \mathfrak{g}_1^*)$ such that $L \cap \mathfrak{g}_1 = \mathfrak{h}_1$.

On the way to our main result, we develop the theory of Lie 2-group actions on Lie groupoids, the tangent and cotangent Lie 2-groups of a 2-group and homogeneous spaces of Lie 2-groups. These topics are of independent interest.

Here is the outline of this paper. In Section 2, we interpret a Lie 2-group action on a Lie groupoid as a homomorphism of group crossed modules. Applications include representations of Lie 2-groups on 2-vector spaces. Focusing on the adjoint and coadjoint representations, we constructed the tangent and cotangent Lie 2-groups of a 2-group; Section 3 delves into a comprehensive study of homogeneous spaces of Lie 2-groups. A homogeneous space not only has a quotient Lie groupoid structure, but also a fiber bundle structure. Finally, in Section 4, we establish a correspondence between Poisson homogeneous spaces of Poisson 2-groups and invariant Dirac structures of some Lie bialgebras.

2. LIE 2-GROUP ACTIONS AND REPRESENTATIONS

In [20], an action of a (strict) Lie 2-algebra on a Lie algebroid is introduced as a homomorphism of Lie algebra crossed modules. It is then integrated to an action of a Lie 2-group on a Lie groupoid by using Mackenzie's double Lie algebroids and groupoids. Later in [9], a Lie 2-group action is defined directly as a group action in the category of Lie groupoids. Here we shall show the automorphism group of a Lie groupoid together with its bisection group gives rise to a natural group crossed module and a Lie 2-group action can be interpreted as a homomorphism of group crossed modules.

In particular, for a linear Lie 2-group action, we recover the representation of a Lie 2-group on a 2-vector space. The dual of a linear Lie 2-group action is also studied. As applications, we consider the adjoint and coadjoint actions of a Lie 2-group on its Lie 2-algebra and the dual. The tangent and cotangent Lie 2-groups are thus obtained.

Let us first recall the notions of Lie group (algebra) crossed modules and Lie 2-groups following [1, 2]. A Lie group crossed module is a pair of Lie groups G_1 and G_0 with a Lie group homomorphism $\Phi: G_1 \to G_0$ and an action of G_0 on G_1 by automorphisms such that

$$\Phi(g_0 \triangleright g_1) = g_0 \Phi(g_1) \mathfrak{g}_0^{-1}, \qquad \Phi(g_1) \triangleright g_1' = g_1 g_1' g_1^{-1}, \qquad \forall g_0 \in G_0, g_1, g_1' \in G_1$$

A Lie 2-group is a Lie groupoid $\Gamma \rightrightarrows \Gamma_0$, such that Γ and Γ_0 are both Lie groups and all the groupoid structure maps are group homomorphisms.

There is a bijection between Lie group crossed modules and Lie 2-groups. Given a Lie group crossed module $\mathbb{G}: G_1 \to G_0$, the associated Lie 2-group is $G_0 \ltimes G_1 \rightrightarrows G_0$, or simply denoted by $G_0 \ltimes G_1$. The group multiplication on $G_0 \ltimes G_1$ is given by

(1)
$$(g_0, g_1) \diamond (l_0, l_1) = (g_0 l_0, (l_0^{-1} \triangleright g_1) l_1).$$

The source, target maps of the Lie groupoid are $s(g_0, g_1) = g_0$ and $t(g_0, g_1) = g_0 \Phi(g_1)$, and the groupoid multiplication is

(2)
$$(g_0, g_1) * (l_0, l_1) = (g_0, g_1 l_1), \quad \text{if} \quad l_0 = g_0 \Phi(g_1).$$

A Lie algebra crossed module is a pair of Lie algebras \mathfrak{g}_0 and \mathfrak{g}_1 with a Lie algebra homomorphism $\phi : \mathfrak{g}_1 \to \mathfrak{g}_0$ and an action of \mathfrak{g}_0 on \mathfrak{g}_1 by derivations such that

$$\phi(x \triangleright u) = [x, \phi(u)], \qquad [u, v] = \phi(u) \triangleright v, \qquad \forall x \in \mathfrak{g}_0, u, v \in \mathfrak{g}_1$$

Lie algebra crossed modules are equivalent to strict Lie 2-algebras [1].

Throughout this paper, we use Lie 2-groups (algebras) and Lie group (algebra) crossed modules interchangeably.

Definition 2.1. [9] Let $\mathbb{G} : G_1 \xrightarrow{\Phi} G_0$ be a Lie 2-group and $P \rightrightarrows P_0$ a Lie groupoid. A Lie 2-group action of \mathbb{G} on P, or a \mathbb{G} -action, is a Lie group action $\triangleright : \mathbb{G} \times P \to P$ such that it is also a Lie groupoid homomorphism, where $\mathbb{G} \times P \rightrightarrows G_0 \times P_0$ is the direct product Lie groupoid of \mathbb{G} and P.

For a \mathbb{G} -action, we have

 $\iota(g_0 \triangleright p_0) := \iota(g_0) \triangleright \iota(p_0), \quad s(g \triangleright p) = s(g) \triangleright s(p), \quad t(g \triangleright p) = t(g) \triangleright t(p), \qquad \forall g \in \mathbb{G}, p \in P,$

and

(3)
$$(g *_{\mathbb{G}} g') \triangleright (p * p') = (g \triangleright p) * (g' \triangleright p'), \quad \forall (g, g') \in \mathbb{G}^{(2)}, (p, p') \in P^{(2)}.$$

Here $\mathbb{G}^{(2)} := \{(g, g') \in \mathbb{G} \times \mathbb{G}; t(g) = s(g')\}$ is the set of composable pairs.

The left action of a Lie 2-group on itself by left group translation is obviously a Lie 2-group action on a Lie groupoid, as the group multiplication is a groupoid homomorphism.

Suppose that a Lie 2-group $\mathbb{G}: G_1 \xrightarrow{\Phi} G_0$ acts on a Lie groupoid $P \Rightarrow P_0$ freely and properly. Then the quotient space $P/\mathbb{G} \Rightarrow P_0/G_0$ is a Lie groupoid such that the projection $P \to P/\mathbb{G}$ is a Lie groupoid homomorphism.

A Lie 2-group action can also be characterized as a homomorphism of group crossed modules. Let $P \Rightarrow P_0$ be a Lie groupoid. A *bisection* of P is a splitting $\gamma : P_0 \to P$ of the source map $s (s \circ \gamma = id)$ such that $\phi_{\gamma} := t \circ \gamma : P_0 \to P_0$ is a diffeomorphism. All bisections form a group Bis(P) with the multiplication and inverse given by

(4)
$$\gamma \cdot \gamma'(x) = \gamma'(x) * \gamma(\phi_{\gamma'}(x)), \qquad \gamma^{-1}(x) = \gamma(\phi_{\gamma}^{-1}x)^{-1}, \qquad \forall x \in P_0.$$

Note that $\phi_{\gamma \cdot \gamma'} = \phi_{\gamma} \circ \phi_{\gamma'}$.

Proposition 2.2. Let $P \rightrightarrows P_0$ be a Lie groupoid and Aut(P) the group of Lie groupoid automorphisms. Then we have a group crossed module

$$\Psi : \operatorname{Bis}(P) \to \operatorname{Aut}(P), \qquad \Psi(\gamma)(p) = \gamma(s(p))^{-1} * p * \gamma(t(p)),$$

where the action of Aut(P) on Bis(P) is

(5)
$$(D \triangleright \gamma)(x) = D(\gamma D_0^{-1}(x)), \quad D_0 = D|_{P_0} \in \text{Diff}(P_0).$$

Proof. First it is direct to see

$$(D' \triangleright (D \triangleright \gamma))(x) = D'((D \triangleright \gamma)({D'_0}^{-1}(x))) = D'D(\gamma({D_0}^{-1}{D'_0}^{-1}(x))) = ((D' \circ D) \triangleright \gamma)(x),$$

so \triangleright is a group action. Then we check that Ψ is a group homomorphism. For $p \in P$ with x = s(p) and y = t(p), we have

$$\begin{aligned} \Psi(\gamma \cdot \gamma')(p) &= (\gamma \cdot \gamma')(x)^{-1} * p * (\gamma \cdot \gamma')(y) \\ &= \gamma(\phi_{\gamma'}(x))^{-1} * \gamma'(x)^{-1} * p * \gamma'(y) * \gamma(\phi_{\gamma'}(y)) \\ &= (\Psi(\gamma) \circ \Psi(\gamma'))(p), \end{aligned}$$

which implies $\Psi(\gamma\cdot\gamma')=\Psi(\gamma)\circ\Psi(\gamma').$ It is left to show

$$\Psi(D \triangleright \gamma) = D \circ \Psi(\gamma) \circ D^{-1}, \qquad \Psi(\gamma) \triangleright \gamma' = \gamma \cdot \gamma' \cdot \gamma^{-1}.$$

Actually, since D is a Lie groupoid automorphism, we have

$$\begin{aligned} (D \circ \Psi(\gamma) \circ D^{-1})(p) &= D \big(\gamma (D_0^{-1} x)^{-1} * D^{-1}(p) * \gamma (D_0^{-1} y) \big) \\ &= D (\gamma (D_0^{-1} x))^{-1} * p * D (\gamma (D_0^{-1} y)) \\ &= (D \triangleright \gamma)(x)^{-1} * p * (D \triangleright \gamma)(y) \\ &= \Psi (D \triangleright \gamma)(p). \end{aligned}$$

Also, observing that $\Psi(\gamma)(x) = \phi_{\gamma}(x)$, we have

$$\begin{aligned} \left(\Psi(\gamma) \triangleright \gamma'\right)(x) &= \Psi(\gamma)(\gamma'\phi_{\gamma}^{-1}(x)) \\ &= \gamma(\phi_{\gamma}^{-1}x)^{-1} * \gamma'(\phi_{\gamma}^{-1}(x)) * \gamma(\phi_{\gamma'}\phi_{\gamma}^{-1}x) \\ &= \gamma^{-1}(x) * (\gamma \cdot \gamma')(\phi_{\gamma}^{-1}(x)) \\ &= (\gamma \cdot \gamma' \cdot \gamma^{-1})(x). \end{aligned}$$

Theorem 2.3. Given a Lie 2-group \mathbb{G} and a Lie groupoid P, there is a one-one correspondence between Lie 2-group actions \triangleright of \mathbb{G} on P and homomorphisms (F_0, F_1) of group crossed modules:

$$G_1 \xrightarrow{F_1} \operatorname{Bis}(P)$$

$$\downarrow \Phi \qquad \qquad \downarrow \Psi$$

$$G_0 \xrightarrow{F_0} \operatorname{Aut}(P)$$

They are related by

(6) $F_0(g_0)(p) = g_0 \triangleright p, \quad F_1(g_1)(x) = g_1 \triangleright x, \quad \forall x \in P_0, g_0 \in G_0, g_1 \in G_1, p \in P.$

Proof. Given a Lie 2-group action \triangleright , we show that (F_0, F_1) defined in (6) is a crossed module homomorphism. First, since the \mathbb{G} -action is a Lie group action, we have

$$F_0(g_0g'_0)(p) = (g_0g'_0) \triangleright p = g_0 \triangleright (g'_0 \triangleright p) = (F_0(g_0) \circ F_0(g'_0))(p).$$

Also using the fact that $\phi_{F_1(g_1)}(x) = t(g_1 \triangleright x) = \Phi(g_1) \triangleright x$, we have

$$(F_1(g_1) \cdot F_1(g'_1))(x) = F_1(g'_1)(x) * F_1(g_1)(\Phi(g'_1) \triangleright x) = (g'_1 \triangleright x) * ((g_1 \diamond \Phi(g'_1)) \triangleright x) = ((e, g'_1) *_{\mathbb{G}} (\Phi(g'_1), \Phi(g'_1^{-1}) \triangleright g_1)) \triangleright x = (g_1g'_1) \triangleright x = F_1(g_1g'_1)(x),$$

where we used (3) in the third equation. So F_0 and F_1 are group homomorphisms. Then we show that the diagram is commutative, i.e. $F_0 \circ \Phi = \Psi \circ F_1$. Indeed, by using (3), we have

$$\begin{split} \Psi(F_1(g_1))(p) &= (F_1(g_1)(x))^{-1} * p * F_1(g_1)(y) & x = s(p), y = t(p) \\ &= (g_1 \triangleright x)^{-1} * p * (g_1 \triangleright y) \\ &= ((\Phi(g_1), g_1^{-1}) \triangleright x) * p * (g_1 \triangleright y) \\ &= (\Phi(g_1), e) \triangleright p = F_0(\Phi(g_1))(p). \end{split}$$

It is left to check $F_1(g_0 \triangleright g_1) = F_0(g_0) \triangleright F_1(g_1)$. Indeed, we have

$$(F_0(g_0) \triangleright F_1(g_1))(x) = F_0(g_0)(F_1(g_1)(g_0^{-1} \triangleright x)) = F_0(g_0)((g_1 \diamond g_0^{-1}) \triangleright x)$$

= $(g_0 \diamond g_1 \diamond g_0^{-1}) \triangleright x = (g_0 \triangleright g_1) \triangleright x = F_1(g_0 \triangleright g_1)(x).$

This shows that (F_0, F_1) is a homomorphism of crossed modules.

Conversely, given a crossed module homomorphism (F_0, F_1) , define

(7)
$$\triangleright : \mathbb{G} \times P \to P, \qquad (g_0, g_1) \triangleright p := F_0(g_0) \big(p * F_1(g_1)(y) \big), \quad y = t(p).$$

We claim that \triangleright is a Lie 2-group action. Namely, it is a Lie group action and also a Lie groupoid homomorphism (only check (3)). For $(g_0, g_1), (l_0, l_1) \in G_0 \times G_1$, since F_0, F_1 are group homomorphisms and

$$F_1(l_0^{-1} \triangleright g_1) = F_0(l_0^{-1}) \triangleright F_1(g_1), \qquad \phi_{F_1(l_1)}(x) = \Psi(F_1(l_1))x = F_0(\Phi(l_1))x, \qquad \forall x \in P_0,$$

we have

$$\begin{split} \left((g_0, g_1) \diamond (l_0, l_1) \right) \triangleright p &= F_0(g_0 l_0) \left(p * F_1((l_0^{-1} \triangleright g_1) l_1)(y) \right) \\ &= F_0(g_0 l_0) \left(p * (F_0(l_0^{-1}) \triangleright F_1(g_1)) \cdot F_1(l_1)(y) \right) \\ &= F_0(g_0 l_0) \left(p * F_1(l_1)(y) * (F_0(l_0^{-1}) \triangleright F_1(g_1)) (F_0(\Phi(l_1))y) \right) \\ &= F_0(g_0 l_0) \left(p * F_1(l_1)(y) * F_0(l_0^{-1}) (F_1(g_1) (F_0(l_0 \Phi(l_1))y) \right) \\ &= F_0(g_0) \left((F_0(l_0)(p * F_1(l_1)y)) * (F_1(g_1) F_0(l_0) F_0(\Phi(l_1))y) \right) \\ &= (g_0, g_1) \triangleright \left(F_0(l_0)(p * F_1(l_1)y) \right) = (g_0, g_1) \triangleright \left((l_0, l_1) \triangleright p \right), \end{split}$$

where we have also used (4),(5) and the fact that $F_0(g_0) \in \operatorname{Aut}(P)$. Thus \triangleright is a group action. Then, for a composable pair $(p, p') \in P^{(2)}$, letting y = t(p) = s(p'), z = t(p'), we find

$$\begin{pmatrix} (g_0, g_1) \triangleright p \end{pmatrix} * ((g_0 \Phi(g_1), l_1) \triangleright p') \\ = F_0(g_0) (p * F_1(g_1)(y)) * F_0(g_0 \Phi(g_1)) (p' * F_1(l_1)z) \\ = F_0(g_0) (p * F_1(g_1)(y) * \Psi(F_1(g_1))(p') * \Psi(F_1(g_1))(F_1(l_1)z)) \\ = F_0(g_0) (p * p' * F_1(g_1)z * (F_1(g_1)z)^{-1} * F_1(l_1)z * F_1(g_1)(\phi_{F_1(l_1)}z)) \\ = F_0(g_0) (p * p' * (F_1(g_1) \cdot F_1(l_1))z) \\ = (g_0, g_1 l_1) \triangleright (p * p') = ((g_0, g_1) *_{\mathbb{G}} (g_0 \Phi(g_1), l_1)) \triangleright (p * p'),$$

which implies (3). Hence we obtain a Lie 2-group action from a crossed module homomorphism.

It suffices to show the above two processes are inverse to each other. Given an action \triangleright , we define a crossed module homomorphism (F_0, F_1) by (6), which then gives rise to a Lie 2-group action \triangleright^F by (7). We claim $\triangleright^F = \triangleright$. In fact, we have

$$(g_0, g_1) \triangleright^F p = F_0(g_0) (p * F_1(g_1)(y)) = g_0 \triangleright (p * (g_1 \triangleright y)) = g_0 \triangleright ((e, e) *_{\mathbb{G}} (e, g_1)) \triangleright (p * y)) = g_0 \triangleright (g_1 \triangleright p) = (g_0, g_1) \triangleright p,$$

where we have used the fact that \triangleright is a Lie group action and also a Lie groupoid homomorphism. This completes the proof.

Denote by A the Lie algebroid of the Lie groupoid P. The infinitesimal of the group crossed module $\operatorname{Bis}(P) \xrightarrow{\Psi} \operatorname{Aut}(P)$ in Proposition 2.2 is the Lie algebra crossed module

$$\psi: \Gamma(A) \to \operatorname{Der}(A), \qquad \psi(u) = [u, \cdot]_A,$$

where Der(A) acts on $\Gamma(A)$ naturally.

Given a \mathbb{G} -action on P, taking the differential of the homomorphism in Theorem 2.3, we obtain a homomorphism of Lie algebra crossed modules

$$\begin{array}{c} \mathfrak{g}_1 \longrightarrow \Gamma(A) \\ \downarrow^{\phi} \qquad \qquad \downarrow^{\psi} \\ \mathfrak{g}_0 \longrightarrow \operatorname{Der}(A) \end{array}$$

This gives an action of a Lie 2-algebra \mathfrak{g} on a Lie algebroid A as introduced in [20].


Example 2.4. If the Lie groupoid P is the pair groupoid $M \times M \rightrightarrows M$ for a manifold M, the crossed module $\operatorname{Bis}(P) \xrightarrow{\Psi} \operatorname{Aut}(P)$ becomes $\operatorname{Diff}(M) \xrightarrow{\operatorname{Id}} \operatorname{Diff}(M)$, where $\operatorname{Diff}(M)$ adjoint acts on itself. A crossed module homomorphism from \mathbb{G} to $\operatorname{Diff}(M) \xrightarrow{\operatorname{Id}} \operatorname{Diff}(M)$ can only be of the form

$$\begin{array}{ccc} G_1 & \stackrel{\sigma \circ \Phi}{\longrightarrow} \operatorname{Diff}(M) \\ & & & & \downarrow_{\operatorname{Id}} \\ G_0 & \stackrel{\sigma}{\longrightarrow} \operatorname{Diff}(M) \end{array}$$

where $\sigma : G_0 \to \text{Diff}(M)$ is a Lie group action of G_0 on M. Therefore, by (7), a Lie 2-group action of \mathbb{G} on the pair groupoid $M \times M \rightrightarrows M$ is determined by such a σ in the way:

$$(G_0 \ltimes G_1) \times (M \times M) \to M \times M, \qquad (g_0, g_1) \triangleright (m_1, m_2) = (\sigma(g_0)m_1, \sigma(g_0 \Phi(g_1))m_2)$$

Example 2.5. When the Lie groupoid P is a Lie group H, the group crossed module $Bis(P) \xrightarrow{\Psi} Aut(P)$ turns out to be the well-known Lie group crossed module $H \xrightarrow{AD} Aut(H)$ (after a minor adjustment). A Lie 2-group action of \mathbb{G} on a Lie group H is determined by a pair (F_0, F_1) :



where F_0 and F_1 are Lie group homomorphisms satisfying that

$$F_0(\Phi(g_1))(h) = F_1(g_1)hF_1(g_1)^{-1}, \qquad F_1(g_0 \triangleright g_1) = F_0(g_0)(F_1(g_1)).$$

The Lie 2-group action is by (7) given by

$$\triangleright : (G_0 \ltimes G_1) \times H \to H, \qquad (g_0, g_1) \triangleright h = F_0(g_0) (hF_1(g_1)).$$

Example 2.6. Let the Lie groupoid P be a Lie 2-group $\mathbb{H} : H_0 \ltimes H_1 \Rightarrow H_0$. Consider bisections of \mathbb{H} that are also group homomorphisms and groupoid automorphisms of \mathbb{H} that are also crossed module automorphisms, denoted by $D(H_0, H_1)$ and $\operatorname{Aut}(\mathbb{H})$, respectively. Then we obtain a subcrossed module $D(H_0, H_1) \to \operatorname{Aut}(\mathbb{H})$ of $\operatorname{Bis}(P) \xrightarrow{\Psi} \operatorname{Aut}(P)$. This subcrossed module is called the actor crossed module of \mathbb{H} and was introduced to define actions of a Lie 2-group on another Lie 2-group in [17].

Let us focus on the particular case of Example 2.6 when \mathbb{H} is a 2-vector space \mathbb{V} . We shall see that linear Lie 2-group actions are exactly representations of this Lie 2-group on 2-vector spaces.

A 2-vector space $\mathbb{V}: V_1 \xrightarrow{\phi} V_0$ gives an action Lie groupoid $\mathbb{V}: V_0 \times V_1 \rightrightarrows V_0$, whose source, target and multiplication are

(8)
$$s(x,u) = x, \quad t(x,u) = x + \phi(u), \quad (x,u) * (x + \phi(u), v) = (x, u + v)$$

for $x \in V_0, u, v \in V_1$. With the abelian group structure, it is a linear Lie 2-group. A G-action on \mathbb{V} is **linear** if $g \triangleright (\cdot) : \mathbb{V} \to \mathbb{V}$ is a linear map for all $g \in \mathbb{G}$.

Proposition 2.7. *Given a linear action* \triangleright *of a Lie* 2*-group* \mathbb{G} *on* \mathbb{V} *, then*

(i) there is a semi-direct product Lie 2-group $\mathbb{G} \ltimes \mathbb{V} \rightrightarrows G_0 \ltimes V_0$, with the direct product groupoid structure and the semi-direct product group structure defined by

$$\begin{array}{lll} (g,\omega)*(l,\delta) &=& (g*_{\mathbb{G}}l,\omega*\delta), \qquad (g,l)\in\mathbb{G}^{(2)}, (\omega,\delta)\in\mathbb{V}^{(2)}, \\ (g,\omega)\diamond(g',\omega') &=& (g\diamond g',\omega+g\triangleright\omega'), \qquad \forall g,g'\in\mathbb{G}, \omega,\omega'\in\mathbb{V}. \end{array}$$

(ii) the Lie group crossed module associated to $\mathbb{G} \ltimes \mathbb{V}$ is $G_1 \ltimes V_1 \xrightarrow{\Phi \times \phi} G_0 \ltimes V_0$, where the group structures and the action are defined by

$$(g_1, u) \cdot (g'_1, u') = (g_1g'_1, u + g_1 \triangleright u'),
(g_0, x) \cdot (g'_0, x') = (g_0g'_0, x + g_0 \triangleright x'),
(g_0, x) \triangleright (g_1, u) = (g_0 \triangleright g_1, g_0 \triangleright u - (g_0 \triangleright g_1) \triangleright x + x),$$

for $g_0, g'_0 \in G_0, g_1, g'_1 \in G_1, x, x' \in V_0$ and $u, u' \in V_1$.

Proof. For (i), we only verify that the groupoid multiplication is a group homomorphism, i.e.

(9)
$$((g,\omega)\diamond(g',\omega'))\ast((l,\delta)\diamond(l',\delta')) = ((g,\omega)\ast(l,\delta))\diamond((g',\omega')\ast(l',\delta')),$$

for composable pairs $(g, l), (g', l') \in \mathbb{G}^{(2)}$ and $(\omega, \delta), (\omega', \delta') \in \mathbb{V}^{(2)}$. By (3) and the fact that \mathbb{G}, \mathbb{V} are Lie 2-groups, the left hand side of (9) equals to

$$\begin{aligned} (gg', \omega + g \triangleright \omega') * (ll', \delta + l \triangleright \delta') &= ((gg') *_{\mathbb{G}} (ll'), (\omega + g \triangleright \omega') * (\delta + l \triangleright \delta')) \\ &= ((g *_{\mathbb{G}} l)(g' *_{\mathbb{G}} l'), \omega * \delta + (g \triangleright \omega') * (l \triangleright \delta')) \\ &= ((g *_{\mathbb{G}} l)(g' *_{\mathbb{G}} l'), \omega * \delta + (g *_{\mathbb{G}} l) \triangleright (\omega' * \delta')) \\ &= ((g, \omega) * (l, \delta)) \diamond ((g', \omega') * (l', \delta')), \end{aligned}$$

which matches with the right hand side of (9).

For (ii), from a Lie 2-group $\Gamma \rightrightarrows \Gamma_0$, a crossed module ker $s \xrightarrow{t} \Gamma_0$ can be constructed with the group action of Γ_0 on ker s defined by $g_0 \triangleright g_1 := g_0 \diamond g_1 \diamond g_0^{-1}$, where \diamond is the group multiplication in Γ . Applying this construction to our Lie 2-group $\mathbb{G} \ltimes \mathbb{V}$, we have ker $s = G_1 \ltimes V_1$ and $t|_{\ker s} = \Phi \times \phi$.

Applying this construction to our Lie 2-group $\mathbb{G} \ltimes \mathbb{V}$, we have ker $s = G_1 \ltimes V_1$ and $t|_{\ker s} = \Phi \times \phi$. The group multiplications on $G_i \ltimes V_i$ for i = 0, 1 is direct to obtain. For the action of $G_0 \ltimes V_0$ on $G_1 \ltimes V_1$, we have

$$\begin{aligned} (g_0, x) \triangleright (g_1, u) &:= (g_0, x) \diamond (g_1, u) \diamond (g_0^{-1}, -g_0^{-1} \triangleright x) = (g_0 \diamond g_1, x + g_0 \triangleright u) \diamond (g_0^{-1}, -g_0^{-1} \triangleright x) \\ &= (g_0 \triangleright g_1, x + g_0 \triangleright u - (g_0 \triangleright g_1) \triangleright x) \in G_1 \times V_1, \end{aligned}$$

as desired. The rest is obvious.

Associated to a 2-vector space $\mathbb{V}: V_1 \xrightarrow{\phi} V_0$, we have the general linear Lie 2-group $\operatorname{GL}(\mathbb{V}) : \operatorname{GL}_1(\mathbb{V}) \xrightarrow{\partial} \operatorname{GL}_0(\mathbb{V})$, where

$$\begin{aligned} \operatorname{GL}_1(\mathbb{V}) &:= & \{\gamma \in \operatorname{Hom}(V_0, V_1); \operatorname{Id} + \phi \circ \gamma \in \operatorname{GL}(V_0)\}, \\ \operatorname{GL}_0(\mathbb{V}) &:= & \{(A_0, A_1) \in \operatorname{GL}(V_0) \oplus \operatorname{GL}(V_1); A_0 \circ \phi = \phi \circ A_1\}, \\ & \partial(\gamma) &:= & (\operatorname{Id} + \phi \circ \gamma, \operatorname{Id} + \gamma \circ \phi). \end{aligned}$$

Here the group structures on $GL_i(\mathbb{V})$ are

$$\gamma \cdot \gamma' = \gamma + \gamma' + \gamma \circ \phi \circ \gamma',$$
 $(A_0, A_1) \circ (A'_0, A'_1) = (A_0 \circ A'_0, A_1 \circ A'_1),$

and the action of $GL_0(\mathbb{V})$ of $GL_1(\mathbb{V})$ is given by

$$(A_0, A_1) \triangleright \gamma = A_1 \circ \gamma \circ A_0^{-1}.$$

The Lie 2-algebra of $GL(\mathbb{V})$ is $gl(\mathbb{V})$. See [18] for details. A **representation** of a Lie 2-group \mathbb{G} on \mathbb{V} is a crossed module homomorphism $\sigma : \mathbb{G} \to \mathbb{GL}(\mathbb{V})$.

Proposition 2.8. *There is a one-one correspondence between linear Lie 2-group actions of* \mathbb{G} *on* \mathbb{V} *and representations of* \mathbb{G} *on the 2-vector space* \mathbb{V} .

Proof. We first show that $\operatorname{GL}(\mathbb{V})$ is a subcrossed module of $\operatorname{Bis}(\mathbb{V}) \xrightarrow{\Psi} \operatorname{Aut}(\mathbb{V})$. It is obvious that $\operatorname{GL}_0(\mathbb{V})$ is the set of linear groupoid automorphisms, which is a subgroup of $\operatorname{Aut}(\mathbb{V})$. Then by identifying $\gamma \in \operatorname{GL}_1(\mathbb{V})$ with the bisection $x \mapsto (x, \gamma(x))$, we see that $\operatorname{GL}_1(\mathbb{V})$ can be viewed as the set of linear bisections of \mathbb{V} . For two linear bisections γ and γ' , by Proposition 2.2, we have $\phi_{\gamma} = \operatorname{Id} + \phi \circ \gamma$ and

$$\gamma \cdot \gamma'(x) = (x, \gamma'(x)) * (\phi_{\gamma'}(x), \gamma(\phi_{\gamma'}(x))) = (x, \gamma(x) + \gamma'(x) + (\gamma \circ \phi \circ \gamma')(x))$$

which implies that $\gamma \cdot \gamma' = \gamma + \gamma' + \gamma \circ \phi \circ \gamma'$. Hence one obtains that $\operatorname{GL}_1(\mathbb{V}) \subset \operatorname{Bis}(\mathbb{V})$ is a Lie subgroup. Then by the fact that $(x, u)^{-1} = (x + \phi(u), -u)$, we compute

$$\begin{split} \Psi(\gamma)(x,u) &= (x + \phi\gamma(x), -\gamma(x)) * (x,u) * (x + \phi(u), \gamma(x + \phi(u))) \\ &= ((\mathrm{Id} + \phi \circ \gamma)x, (\mathrm{Id} + \gamma \circ \phi)u) = \partial(\gamma)(x,u), \end{split}$$

which shows $\Psi = \partial$. Then we have

$$((A_0, A_1) \triangleright \gamma)(x) = ((A_0, A_1) \circ \gamma \circ A_0^{-1})x = (x, A_1 \gamma A_0^{-1} x).$$

Hence we see $GL(\mathbb{V})$ is a subcrossed module.

If the Lie 2-group action is linear, the image of (F_0, F_1) from \mathbb{G} to $\operatorname{Bis}(\mathbb{V}) \xrightarrow{\Psi} \operatorname{Aut}(\mathbb{V})$ in Theorem 2.3 falls into the subcrossed module $\operatorname{GL}(\mathbb{V})$, which gives a representation. It is moreover a one-one correspondence.

Explicitly, given a linear Lie 2-group action $\triangleright : \mathbb{G} \times \mathbb{V} \to \mathbb{V}$, by (6), a representation $(\sigma^0, \sigma^1) : \mathbb{G} \to \mathrm{GL}(\mathbb{V})$ is given by

(10)
$$\sigma_{g_0}^0(x,u) := (g_0 \triangleright x, g_0 \triangleright u), \qquad \sigma_{g_1}^1(x) := \operatorname{pr}_{V_1}(g_1 \triangleright x) = g_1 \triangleright x - x,$$

for all $x \in V_0, u \in V_1$. Conversely, by (7), for a representation $(\sigma^0, \sigma^1) : \mathbb{G} \to \mathrm{GL}(\mathbb{V})$, a linear \mathbb{G} -action $\triangleright : \mathbb{G} \times \mathbb{V} \to \mathbb{V}$ is given by

$$(g_0,g_1) \triangleright (x,u) = \left(\sigma_{g_0}^0(x), \sigma_{g_0}^0(u + \sigma_{g_1}^1(x + \phi u))\right) = \left(\sigma_{g_0}^0 x, \sigma_{g_0}^0 \sigma_{g_1}^1 x + \sigma_{g_0 \Phi(g_1)}^0 u\right).$$

For instance, the Lie 2-group $GL(\mathbb{V})$ represents naturally on \mathbb{V} with $(Id, Id) : GL(\mathbb{V}) \to GL(\mathbb{V})$. The associated linear Lie 2-group action $\triangleright : GL(\mathbb{V}) \times \mathbb{V} \to \mathbb{V}$ is

$$(A,\gamma) \triangleright (x,u) = (A_0x, A_1(u + \gamma x + \gamma \phi u)),$$

for all $A_0 \in GL(V_0), A_1 \in GL(V_1)$ and $\gamma \in Hom(V_0, V_1)$.

We remark that the crossed module in Proposition 2.7 is the semi-direct product crossed module induced from the representation (10); see [17].

One approach to obtain a dual Lie 2-group action of \mathbb{G} on \mathbb{V}^* is by taking the dual of a representation and then applying Theorem 2.3.

Proposition 2.9. Let $\sigma = (\sigma^0, \sigma^1) : \mathbb{G} \to \operatorname{GL}(\mathbb{V})$ be a representation of a Lie 2-group \mathbb{G} on a 2-vector space \mathbb{V} .

(i) Taking the dual, we obtain a representation σ^* of \mathbb{G} on \mathbb{V}^* :

$$\begin{array}{ccc}
G_1 & \xrightarrow{\sigma^{*1}} \operatorname{GL}_1(\mathbb{V}^*) \\
\downarrow \Phi & & \downarrow \partial \\
G_0 & \xrightarrow{\sigma^{*0}} \operatorname{GL}_0(\mathbb{V}^*)
\end{array},$$

where

$$\sigma_{g_0}^{*0} := (\sigma_{g_0^{-1}}^0)^*, \qquad \sigma_{g_1}^{*1} := (\sigma_{g_1^{-1}}^1)^*.$$

This representation is called the **dual representation** of σ .

(ii) We have a dual Lie 2-group action $\sigma^{2gp*} : \mathbb{G} \times \mathbb{V}^* \to \mathbb{V}^*$, which is given by

(11)
$$\sigma_{(g_0,g_1)}^{2gp*}(\xi,\alpha) = (\sigma_{g_0}^{*0}\xi,\sigma_{g_0}^{*0}\sigma_{g_1}^{*1}\xi + \sigma_{g_0\Phi(g_1)}^{*0}\alpha), \qquad \forall \xi \in \mathfrak{g}_1^*, \alpha \in \mathfrak{g}_0^*.$$

Proof. (i) follows from a straightforward verification. (ii) follows from Theorem 2.3 and Formula (7). \Box

Remark 2.10. In order to obtain a dual Lie 2-group action, we should take the duals of functors $(GL_0(\mathbb{V}))$ and natural transformations $(GL_1(\mathbb{V}))$ as in Proposition 2.9, instead of taking the dual of linear maps $(GL(\mathbb{V}))$ on the whole vector space \mathbb{V} .

In fact, considering the dual of the group action, we obtain $\triangleright^* : \mathbb{G} \times \mathbb{V}^* \to \mathbb{V}^*$, which is given by

$$\langle g \triangleright^* \gamma, \omega \rangle = \langle \gamma, g^{-1} \triangleright \omega \rangle, \qquad \forall g \in \mathbb{G}, \gamma \in \mathbb{V}^*, \omega \in \mathbb{V}.$$

This is in general not a Lie 2-group action. Actually, \triangleright^* is a Lie 2-group action if and only if

(12)
$$g_1 \triangleright u = u, \qquad \Phi(g_1) \triangleright x = x, \qquad \forall g_1 \in G_1, x \in V_0, u \in V_1.$$

For a Lie 2-group \mathbb{G} with its Lie 2-algebra \mathfrak{g} , denote by Ad and Ad^{*} the adjoint and coadjoint actions of the Lie group \mathbb{G} on \mathfrak{g} and \mathfrak{g}^* , respectively.

Proposition 2.11. (i) For any $(g_0, g_1) \in G_0 \ltimes G_1$ and $(x, u) \in \mathfrak{g}_0 \oplus \mathfrak{g}_1$, we have

(13)
$$\operatorname{Ad}_{(g_0,g_1)}(x,u) = \left(\operatorname{Ad}_{g_0}^{G_0} x, g_0 \triangleright \left(\operatorname{Ad}_{g_1}^{G_1} u - R_{g_1^{-1}*} \widehat{x}_{g_1}\right)\right)$$

where $\hat{x}_{g_1} \in T_{g_1}G_1$ is defined by $\hat{x}_{g_1} := \frac{d}{dt}|_{t=0}(\exp^{tx} \triangleright g_1)$. (ii) The adjoint action $\operatorname{Ad} : \mathbb{G} \times \mathfrak{g} \to \mathfrak{g}$ is a Lie 2-group action. *Proof.* (i) holds by the calculation

$$\begin{aligned} \operatorname{Ad}_{(g_0,g_1)}(x,u) &= \frac{d}{dt}|_{t=0}(g_0,g_1) \diamond (\exp^{tx}, \exp^{tu}) \diamond (g_0^{-1},g_0 \triangleright g_1^{-1}) \\ &= \frac{d}{dt}|_{t=0} \left(g_0 \exp^{tx} g_0^{-1}, g_0 \triangleright ((\exp^{-tx} \triangleright g_1) \exp^{tu} g_1^{-1}) \right) \\ &= \left(\operatorname{Ad}_{g_0}^{G_0} x, g_0 \triangleright (\operatorname{Ad}_{g_1}^{G_1} u - R_{g_1^{-1}*} \widehat{x}_{g_1}) \right). \end{aligned}$$

For (ii), we show the adjoint action Ad is a Lie groupoid homomorphism. We only check (14) $(\operatorname{Ad}_{(q_0,q_1)}(x,u)) * (\operatorname{Ad}_{(q_0\Phi(q_1),h_1)}(x+\phi(u),v)) = \operatorname{Ad}_{(q_0,q_1h_1)}(x,u+v),$

for $g_0 \in G_0, g_1, h_1 \in G_1$ and $x \in \mathfrak{g}_0$ and $u, v \in \mathfrak{g}_1$. By (13), the left hand side of (14) equals $\left(\operatorname{Ad}_{g_0}^{G_0} x, g_0 \triangleright \left(\operatorname{Ad}_{g_1}^{G_1} u - R_{a^{-1}*}\widehat{x}_{g_1}\right)\right) *$

$$(\operatorname{Ad}_{g_0\Phi(g_1)}^{G_0}(x+\phi(u)), g_0\Phi(g_1) \triangleright (\operatorname{Ad}_{h_1}^{G_1}v - R_{h_1^{-1}*}(\widehat{x+\phi(u)})_{h_1}))$$

$$= (\operatorname{Ad}_{g_0}^{G_0}x, g_0 \triangleright (\operatorname{Ad}_{g_1}^{G_1}u - R_{g_1^{-1}*}\widehat{x}_{g_1}) + g_0\Phi(g_1) \triangleright (\operatorname{Ad}_{h_1}^{G_1}v - R_{h_1^{-1}*}(\widehat{x+\phi(u)})_{h_1})),$$

and the right hand side of (14) is

$$\left(\operatorname{Ad}_{g_0}^{G_0} x, g_0 \triangleright \left(\operatorname{Ad}_{g_1h_1}^{G_1}(u+v) - R_{(g_1h_1)^{-1}*}\widehat{x}_{g_1h_1}\right)\right).$$

Using the fact that $\Phi(g_1) \triangleright h_1 = g_1 h_1 g_1^{-1}$ and $g_0 \triangleright (g_1 h_1) = (g_0 \triangleright g_1)(g_0 \triangleright h_1)$, we have $g_0 \Phi(g_1) \triangleright \operatorname{Ad}_{h_1}^{G_1} v = g_0 \triangleright (\operatorname{Ad}_{g_1 h_1}^{G_1} v),$

and

$$g_{0}\Phi(g_{1}) \triangleright (R_{h_{1}^{-1}*}\widehat{\phi(u)}_{h_{1}}) = \frac{d}{dt}|_{t=0}g_{0} \triangleright \left(g_{1}(\Phi(\exp^{tu}) \triangleright h_{1})h_{1}^{-1}g_{1}^{-1}\right)$$
$$= \frac{d}{dt}|_{t=0}g_{0} \triangleright \left(g_{1}\exp^{tu}h_{1}\exp^{-tu}h_{1}^{-1}g_{1}^{-1}\right)$$
$$= g_{0} \triangleright \left(\operatorname{Ad}_{g_{1}}^{G_{1}}u - \operatorname{Ad}_{g_{1}h_{1}}^{G_{1}}u\right),$$

and also

$$\begin{array}{lll} g_0 \triangleright (R_{(g_1h_1)^{-1}*}\widehat{x}_{g_1h_1}) & = & \frac{d}{dt}|_{t=0}g_0 \triangleright \left((\exp^{tx} \triangleright g_1)(\exp^{tx} \triangleright h_1)h_1^{-1}g_1^{-1}\right) \\ & = & g_0 \triangleright (R_{g_1^{-1}*}\widehat{x}_{g_1}) + g_0 \Phi(g_1) \triangleright (R_{h_1^{-1}*}\widehat{x}_{h_1}). \end{array}$$

These three equations show that the terms involving v, u and x of both sides of (14) are equal respectively. Hence (14) holds by linearity.

By Propositions 2.7, associated to the adjoint action Ad, we have a semi-direct product Lie 2-group $\mathbb{G} \ltimes_{\mathrm{Ad}} \mathfrak{g}$. We call it the **tangent 2-group** of a Lie 2-group \mathbb{G} . On the other hand, the tangent bundle $T\mathbb{G} \rightrightarrows TG_0$ of a Lie 2-group \mathbb{G} is naturally a Lie 2-group with the tangent Lie group and Lie groupoid structures. In fact, we have an isomorphism $T\mathbb{G} \cong \mathbb{G} \ltimes \mathfrak{g}$ of Lie 2-groups.

Applying Proposition 2.8 to the adjoint action of \mathbb{G} on its Lie 2-algebra \mathfrak{g} , we have

Corollary 2.12. *The representation* $Ad : \mathbb{G} \to \mathbb{GL}(\mathfrak{g})$ *associated to the adjoint action* Ad *is*

$$G_{1} \xrightarrow{\operatorname{Ad}^{1}} \operatorname{GL}_{1}(\mathfrak{g})$$

$$\downarrow_{\Phi} \qquad \qquad \qquad \downarrow_{\Psi}$$

$$G_{0} \xrightarrow{\operatorname{Ad}^{0}} \operatorname{GL}_{0}(\mathfrak{g})$$

where Ad^0 and Ad^1 are given by

(15)
$$\operatorname{Ad}_{g_0}^0(x,u) = (\operatorname{Ad}_{g_0}^{G_0} x, g_0 \triangleright u), \qquad \operatorname{Ad}_{g_1}^1(x) = -R_{g_1^{-1}*} \widehat{x}_{g_1}.$$

It is called the adjoint representation of \mathbb{G} on \mathfrak{g} .

Taking the Lie 2-group dual as in Proposition 2.9, we shall obtain the coadjoint representation of \mathbb{G} on \mathfrak{g}^* and furthermore a dual Lie 2-group action of \mathbb{G} on \mathfrak{g}^* .

Corollary 2.13. (i) The coadjoint representation $Ad^* : \mathbb{G} \to \mathbb{GL}(\mathfrak{g})$ of a Lie 2-group \mathbb{G} on \mathfrak{g}^* is

$$G_{1} \xrightarrow{\operatorname{Ad}^{*1}} \operatorname{GL}_{1}(\mathfrak{g}^{*})$$

$$\downarrow_{\Phi} \qquad \qquad \qquad \downarrow_{\Psi} ,$$

$$G_{0} \xrightarrow{\operatorname{Ad}^{*0}} \operatorname{GL}_{0}(\mathfrak{g}^{*})$$

where Ad^{*0} and Ad^{*1} are given by

$$\operatorname{Ad}_{g_0}^{*0}(\xi,\alpha) = (g_0 \triangleright^* \xi, \operatorname{Ad}_{g_0}^{*G_0} \alpha), \qquad \operatorname{Ad}_{g_1}^{*1}(\xi) = -\rho_{g_1}^*\xi, \qquad \forall \alpha \in \mathfrak{g}_0^*, \xi \in \mathfrak{g}_1^*.$$

Here $\rho_{g_1}^* : \mathfrak{g}_1^* \to \mathfrak{g}_0^*$ is given by $\langle \rho_{g_1}^*(\xi), x \rangle = \langle \xi, R_{g_1*} \widehat{x}_{g_1^{-1}} \rangle$ for $x \in \mathfrak{g}_0$.

(ii) There is a Lie 2-group action $\mathbb{A}\mathfrak{d}^* = \mathrm{Ad}^{2gp*} : \mathbb{G} \times \mathfrak{g}^* \to \mathfrak{g}^*$ of \mathbb{G} on \mathfrak{g}^* given by

(16)
$$\mathbb{A}\mathfrak{d}^*_{(g_0,g_1)}(\xi,\alpha) = (g_0 \triangleright^* \xi, -\operatorname{Ad}^{*G_0}_{g_0} \rho^*_{g_1}\xi + \operatorname{Ad}^{*G_0}_{g_0\Phi(g_1)} \alpha).$$

By Propositions 2.7, associated to $\mathbb{A}\mathfrak{d}^*$, we have a semi-direct product Lie 2-group $\mathbb{G} \ltimes_{\mathbb{A}\mathfrak{d}^*} \mathfrak{g}^*$, which is called the **cotangent 2-group** of a Lie 2-group.

Remark 2.14. Unlike the coadjoint orbits of a Lie group, the coadjoint orbits of a Lie 2-group defined by (16) in general do not have natural symplectic structures. In fact, the coadjoint action $Ad^* : \mathbb{G} \times \mathfrak{g}^* \to \mathfrak{g}^*$ of the Lie group \mathbb{G} takes the formula

$$\mathrm{Ad}^*_{(g_0,g_1)}(\xi,\alpha) = (g_0 \triangleright^* (\mathrm{Ad}^*_{g_1}\xi), -\mathrm{Ad}^*_{g_0} \rho^*_{g_1}\xi + \mathrm{Ad}^*_{g_0} \alpha),$$

for any $(\alpha, \xi) \in \mathfrak{g}_0^* \oplus \mathfrak{g}_1^*$. It is different from (16) and in general not a Lie 2-group action. By Remark 2.10, it is a Lie 2-group action if and only if

$$\operatorname{Ad}_{g_1}^{G_1} u = u, \qquad \operatorname{Ad}_{\Phi(g_1)}^{G_0} x = x, \qquad \forall g_1 \in G_1, u \in \mathfrak{g}_1, x \in \mathfrak{g}_0$$

So if G_0 and G_1 are connected, Ad^* is a Lie 2-group action if and only if G_1 is abelian and $\operatorname{Im}(\Phi) \subset Z(G_0)$.

3. Homogeneous spaces of Lie 2-groups

In this section, we give a thorough discussion of homogeneous spaces of Lie 2-groups. First, we characterize the algebraic and geometric structures of the homogeneous \mathbb{G} -space \mathbb{G}/\mathbb{H} , where \mathbb{H} is a closed 2-subgroup. It not only inherits a canonical quotient Lie groupoid structure, but also is equipped with a fiber bundle structure. Then it is shown that a homogeneous \mathbb{G} -space is always of the quotient of \mathbb{G} by a closed 2-subgroup. In the end, we discover that \mathbb{G}/\mathbb{H} is not just a homogeneous space of the Lie group \mathbb{G} , but also a homogeneous space of some groupoid.

Given a Lie 2-group \mathbb{G} , we shall use \diamond and * to denote the group and groupoid structures on \mathbb{G} , respectively.

Definition 3.1. Let $\mathbb{G} : G_1 \xrightarrow{\Phi} G_0$ be a Lie 2-group. A homogeneous space of \mathbb{G} , or a homogeneous \mathbb{G} -space, is a Lie groupoid $P \rightrightarrows P_0$ with a transitive \mathbb{G} -action.

Proposition 3.2. Given a Lie 2-group $\mathbb{G} : G_1 \xrightarrow{\Phi} G_0$ and a closed Lie 2-subgroup $\mathbb{H} : H_1 \xrightarrow{\Phi} H_0$, the quotient space \mathbb{G}/\mathbb{H} is a homogeneous \mathbb{G} -space with the \mathbb{G} -action $g \triangleright [g'] = [gg']$, where the Lie groupoid structure on

$$G_0 \ltimes G_1/H_0 \ltimes H_1 \rightrightarrows G_0/H_0$$

is given as follows: the source, target maps are

 $s([g_0, g_1]) = [g_0], \qquad t([g_0, g_1]) = [g_0 \Phi(g_1)],$

and the multiplication is

(17)

$$[g_0,g_1] * [g_0 \Phi(g_1),g_1'] = [g_0,g_1g_1']$$

Proof. To see \mathbb{G}/\mathbb{H} is a Lie groupoid, we only verify that the multiplication is well-defined. For any composable pair $[g_0, g_1], [l_0, l_1] \in \mathbb{G}/\mathbb{H}$, the condition $t([g_0, g_1]) = s([l_0, l_1])$ implies that $[g_0\Phi(g_1)] = [l_0]$. So there exists $h_0 \in H_0$ such that $g_0\Phi(g_1) = l_0h_0$. We can then write

$$[l_0, l_1] = [l_0 h_0, h_0^{-1} \triangleright l_1] = [g_0 \Phi(g_1), h_0^{-1} \triangleright l_1].$$

This shows that to determine * on \mathbb{G}/\mathbb{H} , it is enough to define multiplications of elements of $[g_0, g_1]$ and $[g_0\Phi(g_1), g'_1]$. Then for $(h_0, h_1), (h'_0, h'_1) \in H_0 \ltimes H_1$, we have

$$\begin{split} & [(g_0,g_1)\diamond(h_0,h_1)]\ast[(g_0\Phi(g_1),g_1')\diamond(h_0',h_1')]\\ &= \ [g_0h_0,(h_0^{-1}\rhd g_1)h_1]\ast[g_0\Phi(g_1)h_0',(h_0'^{-1}\rhd g_1')h_1']\\ &= \ [g_0h_0,(h_0^{-1}\rhd g_1)h_1]\ast[g_0\Phi(g_1)h_0\Phi(h_1),(\Phi(h_1)^{-1}h_0^{-1}h_0')\rhd((h_0'^{-1}\rhd g_1')h_1')]\\ &= \ [g_0h_0,(h_0^{-1}\rhd g_1)h_1h_1^{-1}(h_0^{-1}\rhd g_1')((h_0^{-1}h_0')\rhd h_1')h_1]\\ &= \ [(g_0,g_1g_1')\diamond(h_0,((h_0^{-1}h_0')\rhd h_1')h_1)]\\ &= \ [g_0,g_1g_1']. \end{split}$$

This implies that the multiplication (17) does not depend on the choices of representatives in each equivalence class, so it is well-defined. It is direct to show that \mathbb{G}/\mathbb{H} is a homogeneous \mathbb{G} -space, among which (3) holds for the reason that the groupoid multiplication on \mathbb{G} is a group homomorphism.

Given a Lie 2-group \mathbb{G} with a closed Lie 2-subgroup \mathbb{H} , there is a natural Lie group action of H_0 on G_1/H_1 by

$$H_0 \times (G_1/H_1) \to G_1/H_1, \qquad h_0 \triangleright [g_1] = [h_0 \triangleright g_1].$$

We thus obtain an associated bundle

$$G_0 \times_{H_0} (G_1/H_1) \to G_0/H_0, \qquad [g_0, [g_1]] \mapsto [g_0]$$

of the principal H_0 -bundle $G_0 \to G_0/H_0$ with fiber G_1/H_1 .

Lemma 3.3. The above associated bundle is a Lie groupoid $G_0 \times_{H_0} (G_1/H_1) \rightrightarrows G_0/H_0$, where the source, target maps are

$$s([g_0, [g_1]]) = [g_0], \quad t([g_0, [g_1]]) = [g_0 \Phi(g_1)],$$

and the multiplication is given by

$$[g_0, [g_1]] * [g_0 \Phi(g_1), [g'_1]] = [g_0, [g_1g'_1]].$$

Proof. We first show the source and target maps are well-defined. Indeed, taking $[g_0, [g_1]] = [g_0h_0, [(h_0^{-1} \triangleright g_1)h_1]]$ for any $(h_0, h_1) \in H_0 \times H_1$, we have $s([g_0h_0, [(h_0^{-1} \triangleright g_1)h_1]]) = [g_0h_0] = [g_0]$ and

$$t([g_0h_0, [(h_0^{-1} \triangleright g_1)h_1]]) = [g_0h_0h_0^{-1}\Phi(g_1)h_0\Phi(h_1)] = [g_0\Phi(g_1)h_0\Phi(h_1)] = [g_0\Phi(g_1)].$$

Then we show the multiplication is well-defined. For any $[g_0, [g_1]], [l_0, [l_1]] \in G_0 \times_{H_0} (G_1/H_1)$ such that $t([g_0, [g_1]]) = s([l_0, [l_1]])$, there exists some $h_0 \in H_0$ such that $g_0\Phi(g_1) = l_0h_0$. So we have $[l_0, [l_1]] = [g_0\Phi(g_1), [h_0^{-1} \triangleright l_1]]$. This justifies that it is reasonable to only define the multiplication of elements $[g_0, [g_1]]$ and $[g_0\Phi(g_1), [g'_1]]$. It is straightforward to show the structures maps s, t, * define a Lie groupoid structure.

Proposition 3.4. Let \mathbb{G} be a Lie 2-group and \mathbb{H} a closed Lie 2-subgroup. Then

(i) the Lie groupoid $G_0 \times_{H_0} (G_1/H_1) \rightrightarrows G_0/H_0$ is a homogeneous \mathbb{G} -space with the \mathbb{G} -action

 $(l_0, l_1) \triangleright [g_0, [g_1]] = [l_0 g_0, [(g_0^{-1} \triangleright l_1) g_1]];$

(ii) we have an isomorphism of homogeneous G-spaces

$$\theta: G_0 \ltimes G_1/H_0 \ltimes H_1 \to G_0 \times_{H_0} (G_1/H_1), \qquad [g_0, g_1] \mapsto [g_0, [g_1]].$$

Proof. (i) is direct to check. For (ii), we first show that θ is well-defined. Actually,

$$\theta \left([(g_0, g_1) \diamond (h_0, h_1)] \right) = \theta \left([g_0 h_0, (h_0^{-1} \triangleright g_1) h_1] \right) = [g_0 h_0, [(h_0^{-1} \triangleright g_1) h_1]]$$

= $[g_0 h_0, h_0^{-1} \triangleright [g_1]] = [g_0, [g_1]] = \theta \left([g_0, g_1] \right).$

Then following from the calculation

$$\begin{array}{rcl} \theta([g_0,g_1]*[g_0\Phi(g_1),g_1']) &=& \theta([g_0,g_1g_1']) = [g_0,[g_1g_1']] \\ &=& [g_0,[g_1]]*[g_0\Phi(g_1),[g_1']] = \theta([g_0,g_1])*\theta([g_0\Phi(g_1),g_1']), \end{array}$$

we see that θ is a Lie groupoid homomorphism. Next, we check that θ is equivariant with the two \mathbb{G} -actions. Indeed, we have

$$\theta((l_0, l_1) \triangleright [g_0, g_1]) = \theta([l_0 g_0, (g_0^{-1} \triangleright l_1)g_1]) = [l_0 g_0, [(g_0^{-1} \triangleright l_1)g_1]] = (l_0, l_1) \triangleright \theta([g_0, g_1]).$$

At last, define a map

$$\tau: G_0 \times_{H_0} (G_1/H_1) \to G_0 \ltimes G_1/H_0 \ltimes H_1, \qquad [g_0, [g_1]] \mapsto [g_0, g_1].$$

It is straightforward to check that τ is well-defined and it is the inverse of θ .

The Lie groupoid $\mathbb{G} : G_0 \ltimes G_{-1} \rightrightarrows G_0$ is an action groupoid relative to the right action of G_1 on G_0 : $g_0 \triangleleft g_1 = g_0 \Phi(g_1)$. As its quotient, the groupoid $G_0 \times_{H_0} (G_1/H_1) \cong \mathbb{G}/\mathbb{H}$ over G_0/H_0 is in general not an action groupoid any more. We give an example when it is still an action Lie groupoid. Recall from [17] that the subcrossed module \mathbb{H} of a crossed module \mathbb{G} is **normal** if (1) $H_0 \subset G_0$ is a normal subgroup; (2) H_1 is a G_0 -submodule; (3) $(h_0 \bowtie g_1)g_1^{-1} \in H_1$ for all $h_0 \in H_0, g_1 \in G_1$. If $\mathbb{H} \subset \mathbb{G}$ is a normal subcrossed module, then the quotient $\mathbb{G}/\mathbb{H} : G_1/H_1 \xrightarrow{\Phi} G_0/H_0$ inherits a crossed module structure such that the projection $\mathbb{G} \to \mathbb{G}/\mathbb{H}$ is a crossed module homomorphism.

Example 3.5. Let \mathbb{G} be Lie 2-group and \mathbb{H} a closed normal 2-subgroup (normal subcrossed module). Then

- (i) the quotient Lie groupoid \mathbb{G}/\mathbb{H} is naturally isomorphic to the action Lie groupoid (Lie 2-group) $(G_0/H_0) \ltimes (G_1/H_1) \rightrightarrows G_0/H_0.$
- (ii) With the action $\triangleright : (G_0 \ltimes G_1) \times (G_0/H_0 \times G_1/H_1) \to G_0/H_0 \times G_1/H_1:$

$$(g_0, g_1) \triangleright ([g'_0], [g'_1]) = ([g_0g'_0], [(g'_0)^{-1} \triangleright g_1)g'_1]),$$

the Lie groupoid $(G_0/H_0) \ltimes (G_1/H_1) \rightrightarrows G_0/H_0$ is a homogeneous \mathbb{G} -space such that the isomorphism in (i) is an isomomorphism of homogeneous \mathbb{G} -spaces.

Remark 3.6. As for the quotient Lie groupoid $G_0 \times_{H_0} (G_1/H_1) \Rightarrow G_0/H_0$, there are two similar constructions:

- (1) The first one is the quotient of an action Lie groupoid by Lu [15] and Blohmann-Weinstein [4]. A quotient of an action Lie groupoid by a subgroup is of the form $M \times_H G/H \rightrightarrows M/H$, where M is a manifold with a Lie group G-action and $H \subset G$ is a subgroup. Taking $G_0 = G_1$ and $H_0 = H_1$, our quotient Lie groupoid $G_0 \times_{H_0} (G_0/H_0) \rightrightarrows G_0/H_0$ is an example of this case;
- (2) The second one is the quotient of a PBG-Lie groupoid by Mackenzie [16]. A PBG-Lie groupoid is a locally trivial Lie groupoid Γ ⇒ P(M, G) over a principal bundle P(M, G) together with a right action of G on Γ by groupoid automorphisms over the principal action P × G → P. Then the quotient Γ/G ⇒ M is a Lie groupoid. If H₁ is trivial, our quotient Lie groupoid G₀ ×_{H₀} G₁ ⇒ G₀/H₀ is an example of Mackenzie's case.

The Lie algebroid $G_0 \ltimes_{H_0} (\mathfrak{g}_1/\mathfrak{h}_1) \to G_0/H_0$ of the quotient Lie groupoid $G_0 \times_{H_0} (G_1/H_1) \rightrightarrows G_0/H_0$ can also be written down following from the construction in [15]. We omit the details.

It is from Proposition 3.2 that the quotient of a Lie 2-group \mathbb{G} by a closed Lie 2-subgroup is a homogeneous \mathbb{G} -space. We shall see that a homogeneous \mathbb{G} -space $P \Rightarrow P_0$ is always of this form. Given an action of a Lie 2-group \mathbb{G} on a Lie groupoid $P \Rightarrow P_0$, for $x \in P_0$, denote by $\mathcal{O}_x^{\mathbb{G}} \subset P$ and $\mathcal{O}_x^{G_0} \subset P_0$ the \mathbb{G} -orbit and G_0 -orbit passing through x, respectively. Let $G_0^x \subset G_0$ be the isotropy subgroup of xrelative to the G_0 -action on P_0 and let $G_1^x \subset G_1$ and $\mathbb{G}^x \subset \mathbb{G}$ be the isotropy subgroups of x with respect to the G_1 and \mathbb{G} -actions on P, respectively.

Theorem 3.7. With the above notations,

(i) $\mathcal{O}_x^{\mathbb{G}} \rightrightarrows \mathcal{O}_x^{G_0}$ is a Lie subgroupoid of *P*. With the induced \mathbb{G} -action, it is a homogeneous \mathbb{G} -space. Denote it by $\mathcal{O}_x^{\mathbb{G}}$.

- (ii) $G_1^x \xrightarrow{\Phi} G_0^x$ is a subcrossed module of $G_1 \xrightarrow{\Phi} G_0$ and $\mathbb{G}^x = G_0^x \ltimes G_1^x$. And the two homogeneous \mathbb{G} -spaces \mathbb{G}/\mathbb{G}^x and $\mathcal{O}_x^{\mathbb{G}}$ are isomorphic.
- (iii) If P is further a homogeneous \mathbb{G} -space, we have an isomorphism $P \cong \mathbb{G}/\mathbb{G}^x$ of homogeneous \mathbb{G} -spaces.

Proof. For (i), for $g \triangleright x, l \triangleright x \in \mathcal{O}_x^{\mathbb{G}}$ with $t(g) \triangleright x = s(l) \triangleright x$, there exists $k \in G_0^x$ such that $t(g) = s(l)k = s(l \diamond k)$. Then by (3), we have

$$(g \triangleright x) * (l \triangleright x) := (g *_{\mathbb{G}} (l \diamond k)) \triangleright x \in \mathcal{O}_x^{\mathbb{G}},$$

which implies that $\mathcal{O}_x^{\mathbb{G}}$ is a Lie subgroupoid of P.

For (ii), first we show that $G_1^x \xrightarrow{\Phi} G_0^x$ is a subcrossed module of \mathbb{G} . For $g_1 \in G_1^x$, since $\Phi(g_1) \triangleright x = t(g_1) \triangleright x = t_P(g_1 \triangleright x) = t_P(x) = x$, we have $\Phi(g_1) \in G_0^x$. Also, for $g_0 \in G_0^x$ and $g_1 \in G_1^x$, we have

$$(g_0 \triangleright g_1) \triangleright x = (g_0 \diamond g_1 \diamond g_0^{-1}) \triangleright x = x,$$

where \diamond is the group multiplication in \mathbb{G} . Hence $g_0 \triangleright g_1 \in G_1^x$. Thus $G_1^x \xrightarrow{\Phi} G_0^x$ is a subcrossed module and gives a Lie 2-subgroup $G_0^x \ltimes G_1^x \rightrightarrows G_0^x$. To see $\mathbb{G}^x = G_0^x \ltimes G_1^x$, we first notice that $G_0^x \ltimes G_1^x \subset \mathbb{G}^x$. Then for an arbitrary $(g_0, g_1) \in \mathbb{G}^x$, we have $(g_0, g_1) \triangleright x = (g_0 \diamond g_1) \triangleright x = x$. Applying the source map on both sides, we obtain $g_0 \triangleright x = x$ and further $g_1 \triangleright x = g_0^{-1} \triangleright x = x$. Hence $g_0 \in G_0^x, g_1 \in G_1^x$. Thus we proved $\mathbb{G}^x \subset G_0^x \ltimes G_1^x$. So they are equal as Lie groups.

Then we show the map

$$F: \mathcal{O}_x^{\mathbb{G}} \to \mathbb{G}/\mathbb{G}^x, \qquad g \triangleright x \mapsto [g]$$

is an isomorphism of homogeneous \mathbb{G} -spaces. Indeed, it is obviously well-defined. To see it is a groupoid morphism, we find

$$F\big((g \triangleright x) * (l \triangleright x)\big) = F\big((g *_{\mathbb{G}} (l \diamond k)) \triangleright x\big) = [(g *_{\mathbb{G}} (l \diamond k)] = [g] * [l] = F(g \triangleright x) * F(l \triangleright x),$$

where we have used the notations in (i). Also, $F(l \triangleright (g \triangleright x)) = [l \diamond g] = l \triangleright [g] = l \triangleright F(g \triangleright x)$, we see F is an isomorphism of homogeneous spaces with inverse $[g] \mapsto g \triangleright x$.

For (iii), by the transitivity condition, we have $P = \mathcal{O}_x$ for any $x \in P_0$. Then (iii) follows from (ii).

Remark 3.8. From this theorem, we see that for a Lie 2-group action, orbits passing through points in the base manifold are always homogeneous \mathbb{G} -spaces. While the orbit passing through a general point in \mathbb{P} is not necessarily a homogeneous \mathbb{G} -space, since the isotropy group is in general not a Lie 2-subgroup.

In this end of this section, we point out that the quotient space \mathbb{G}/\mathbb{H} is not only a homogeneous space of a Lie group, but also a homogeneous space of a Lie groupoid. This fact will be used in the next section.

Let $\mathbb{G}: G_1 \xrightarrow{\Phi} G_0$ be a Lie 2-group and $\mathbb{H}: H_1 \xrightarrow{\Phi} H_0$ a Lie 2-subgroup. Then the Lie group H_0 acts on G_1 and H_1 , respectively, which induce two associated bundles

$$G_0 \times_{H_0} G_1 \to G_0/H_0, \qquad G_0 \times_{H_0} H_1 \to G_0/H_0$$

Taking H_1 as $\{e\}$ in Propositions 3.2 and 3.4, we get that the associated bundle $G_0 \times_{H_0} G_1$ is a Lie groupoid

$$G_0 \times_{H_0} G_1 \rightrightarrows G_0/H_0$$

with source, target and multiplication

$$s([g_0, g_1]) = [g_0], \qquad t([g_0, g_1]) = [g_0 \Phi(g_1)], \qquad [g_0, g_1] * [g_0 \Phi(g_1), l_1] = [g_0, g_1 l_1].$$

Moreover, it is isomorphic to the Lie groupoid $(G_0 \ltimes G_1)/H_0 \rightrightarrows G_0/H_0$.

Proposition 3.9. With the notations above, we have

- (i) $G_0 \times_{H_0} H_1 \rightrightarrows G_0/H_0$ is a Lie group bundle and a wide normal subgroupoid of $G_0 \times_{H_0} G_1$. Thus, the groupoid quotient $(G_0 \times_{H_0} G_1) / (G_0 \times_{H_0} H_1)$ is a Lie groupoid;
- (ii) there is a Lie groupoid isomorphism

$$\theta: (G_0 \times_{H_0} G_1) /\!\!/ (G_0 \times_{H_0} H_1) \to \mathbb{G}/\mathbb{H}, \qquad [\![g_0, g_1]\!] \to [g_0, g_1].$$

Therefore, \mathbb{G}/\mathbb{H} is a homogeneous space of the Lie group \mathbb{G} and also a homogeneous space of the Lie groupoid $\mathbb{G}/H_0 \rightrightarrows G_0/H_0$.

Proof. For (i), it is obvious that $G_0 \times_{H_0} H_1 \subset G_0 \times_{H_0} G_1$ is a wide subgroupoid. Following from $s([g_0, h_1]) = t([g_0, h_1]) = [g_0]$, we have $G_0 \times_{H_0} H_1$ is a Lie group bundle over G_0/H_0 . For $[g_0, h_1] \in G_0 \times_{H_0} H_1$ and $[g_0, g_1] \in G_0 \times_{H_0} G_1$, we have

$$[g_0, h_1] * [g_0, g_1] = [g_0, h_1] * [g_0 \Phi(h_1), \Phi(h_1)^{-1} \triangleright g_1] = [g_0, g_1 h_1] = [g_0, g_1] * [g_0 \Phi(g_1), h_1] = [g_0, g_1] * [g_0 \Phi(g_1), h_1] = [g_0, g_1] * [g_0 \Phi(g_1), g_1] = [g_0, g$$

which implies that

$$[g_0,g_1]^{\dagger} * [g_0,h_1] * [g_0,g_1] = [g_0 \Phi(g_1),h_1] \in G_0 \times_{H_0} H_1$$

where $[g_0, g_1]^{\dagger}$ is the groupoid inverse of $[g_0, g_1]$. Thus $G_0 \times_{H_0} H_1 \subset G_0 \times_{H_0} G_1$ is a wide normal subgroupoid. Therefore, $(G_0 \times_{H_0} G_1) / (G_0 \times_{H_0} H_1) \rightrightarrows G_0 / H_0$ with the multiplication

$$\llbracket \gamma \rrbracket * \llbracket \gamma' \rrbracket := \llbracket \gamma * \gamma' \rrbracket, \qquad \forall (\gamma, \gamma') \in (G_0 \times_{H_0} G_1)^{(2)}$$

is a Lie groupoid. For (ii), as

$$\begin{aligned} \theta \left(\llbracket [g_0 h_0, h_0^{-1} \triangleright g_1] * [g_0 \Phi(g_1), h_1] \rrbracket \right) &= \theta \left(\llbracket [g_0 h_0, (h_0^{-1} \triangleright g_1) h_1] \rrbracket \right) \\ &= [g_0 h_0, (h_0^{-1} \triangleright g_1) h_1] \\ &= [(g_0, g_1) \diamond (h_0, h_1)] \\ &= [g_0, g_1] = \theta \left(\llbracket [g_0, g_1] \rrbracket \right), \end{aligned}$$

we see that θ is well-defined. Next, for $[g_0, g_1], [g_0 \Phi(g_1), l_1] \in G_0 \times_{H_0} G_1$, we have

$$\theta\big([\![g_0,g_1]]\!]*[\![g_0\Phi(g_1),l_1]]\!]\big) = \theta([\![g_0,g_1l_1]]\!]) = [g_0,g_1l_1] = [g_0,g_1]*[g_0\Phi(g_1),l_1]$$

which implies that θ is a homomorphism. It is moreover an isomorphism with the inverse given by $[g_0, g_1] \mapsto [\![g_0, g_1]]\!]$.

4. POISSON HOMOGENEOUS SPACE OF POISSON 2-GROUPS

Drinfeld classified Poisson homogeneous spaces of a Poisson Lie group by Dirac structures of its Lie bialgebra [8], and then Liu-Weinstein-Xu generalized this result to Poisson groupoids in [13]. Here we present the classification theorem for Poisson homogeneous spaces of Poisson 2-groups. Let us first collect some standard definitions and facts regarding Lie bialgebras and Poisson homogeneous spaces.

A **Lie bialgebra** is a pair $(\mathfrak{g}, \mathfrak{g}^*)$ of Lie algebras such that

$$d_*[x,y] = [d_*(x),y] + [x,d_*(y)], \qquad \forall x,y \in \mathfrak{g}$$

where $d_* : \mathfrak{g} \to \wedge^2 \mathfrak{g}$ is defined by $\langle d_*(x), \xi \wedge \eta \rangle := -\langle x, [\xi, \eta]_{\mathfrak{g}^*} \rangle$. The double $\mathfrak{g} \oplus \mathfrak{g}^*$ of a Lie bialgebra is associated with a natural pairing:

$$\langle x+\xi, y+\eta \rangle_{\bowtie} = \langle x,\eta \rangle + \langle y,\xi \rangle, \qquad \forall x,y \in \mathfrak{g}, \xi,\eta \in \mathfrak{g}^*,$$

and a Lie bracket:

(18)
$$[x + \xi, y + \eta]_{\bowtie} = [x, y]_{\mathfrak{g}} + \mathrm{ad}_{\xi}^* y - \mathrm{ad}_{\eta}^* x + [\xi, \eta]_{\mathfrak{g}^*} + \mathrm{ad}_x^* \eta - \mathrm{ad}_y^* \xi$$

Denote the double Lie algebra by $\mathfrak{g} \bowtie \mathfrak{g}^*$. A **Dirac structure** of a Lie bialgebra $(\mathfrak{g}, \mathfrak{g}^*)$ is a subspace $L \subset \mathfrak{g} \oplus \mathfrak{g}^*$ which is maximal isotropic with respect to $\langle \cdot, \cdot \rangle_{\bowtie}$ and closed under the Lie bracket $[\cdot, \cdot]_{\bowtie}$. The notion of characteristic pairs was introduced in [12] as a useful tool to characterize Dirac structures. Indeed, a maximal isotropic subspace of $\mathfrak{g} \bowtie \mathfrak{g}^*$ is always of the form

(19)
$$L = \{x + \Omega^{\sharp} \xi + \xi; x \in \mathfrak{h}, \xi \in \mathfrak{h}^{\perp}\} = \mathfrak{h} \oplus \operatorname{Graph}(\Omega|_{\mathfrak{h}^{\perp}}),$$

where $\mathfrak{h} \subset \mathfrak{g}$ is a subspace, $\Omega \in \wedge^2 \mathfrak{g}$ and $\Omega^{\sharp}(\xi) := \Omega(\xi, \cdot)$. Note that Ω is actually determined by $\operatorname{pr}(\Omega) \in \wedge^2 \mathfrak{g}/\mathfrak{h}$. We call the pair (\mathfrak{h}, Ω) a **characteristic pair** of *L*. A maximal isotropic subspace *L* as in (19) is a Dirac structure of $(\mathfrak{g}, \mathfrak{g}^*)$ if and only if

- (1) \mathfrak{h} is a Lie subalgebra of \mathfrak{g} ;
- (2) $d_*\Omega + \frac{1}{2}[\Omega,\Omega]_{\mathfrak{g}} \equiv 0 \pmod{\mathfrak{h}};$
- (3) $[\xi,\eta]_{\mathfrak{g}^*} + [\xi,\eta]_{\Omega} \in \mathfrak{h}^{\perp}$ for all $\xi,\eta \in \mathfrak{h}^{\perp}$, where $[\xi,\eta]_{\Omega} = \operatorname{ad}_{\Omega^{\sharp}\varepsilon}^* \eta \operatorname{ad}_{\Omega^{\sharp}\eta}^* \xi$.

Here, the second condition means that $d_*\Omega + \frac{1}{2}[\Omega,\Omega]_{\mathfrak{g}} \in \mathfrak{h} \wedge \mathfrak{g} \wedge \mathfrak{g}$.

Let (G, π) be a Poisson Lie group. A **Poisson homogeneous** G-space is a Poisson manifold (M, π_M) such that M is a homogeneous G-space and the action $G \times M \to M$ is a Poisson map. It is by Drinfeld that there is a one-one correspondence between regular Dirac structures $L \subset \mathfrak{g} \oplus \mathfrak{g}^*$ and Poisson homogeneous spaces G/H, where H is a connected closed subgroup of G with Lie algebra $L \cap \mathfrak{g}$. A Dirac structure L is regular if the left translation of $\mathfrak{h} = L \cap \mathfrak{g}$ defines a simple foliation on G.

Let (P, Π) be a Poisson groupoid. A *P*-space is homogeneous if and only if it is isomorphic to the groupoid quotient $P/\!\!/ P_1$ for some wide (containing all the identities) subgroupoid $P_1 \subset P$. If $P/\!\!/ P_1$ is equipped with a Poisson structure such that the groupoid action is a Poisson map, it is called a **Poisson homogeneous space** of (P, Π) . Liu-Weinstein-Xu [13] proved that there is a one-one correspondence between Poisson homogeneous spaces $P/\!\!/ P_1$ and regular Dirac structures *L* of the Lie bialgebroid (A, A^*) , where P_1 is the *s*-connected closed subgroupoid of *P* corresponding to the subalgebroid $L \cap A$.

A Poisson 2-group is both a Poisson Lie group and a Poisson groupoid. So a Poisson homogeneous space of a Poisson 2-group is naturally expected to be a Poisson homogeneous space of the Poisson Lie group and Poisson groupoid simultaneously. However, given a Lie 2-group \mathbb{G} and a closed Lie 2-subgroup \mathbb{H} , the subgroupoid $H_0 \ltimes H_1 \rightrightarrows H_0$ is not a wide subgroupoid of $G_0 \ltimes G_1 \rightrightarrows G_0$ if $H_0 \neq G_0$. To adjust to the framework in [13], we consider \mathbb{G}/\mathbb{H} as a groupoid homogeneous space of \mathbb{G}/H_0 as in Proposition 3.9.

Lemma 4.1. Let $(\mathbb{G}, \pi_{\mathbb{G}})$ be a Poisson 2-group. Suppose $H_0 \subset G_0$ is a closed subgroup. Then $(\mathbb{G}/H_0, \operatorname{pr}_*\pi_{\mathbb{G}})$ is a quotient Poisson groupoid if and only if $H_0 \subset \mathbb{G}$ is a Poisson subgroup.

Proof. We first recall two facts: the quotient space \mathbb{G}/H_0 inherits a Poisson structure such that the projection $\mathbb{G} \to \mathbb{G}/H_0$ is a Poisson map if and only if the annihilators $\mathfrak{h}_0^{\perp} \oplus \mathfrak{g}_1^*$ of \mathfrak{h}_0 in \mathfrak{g} is a Lie subalgebra of \mathfrak{g}^* . That is, \mathfrak{h}_0 is a coideal in \mathfrak{g} ; The subgroup $H_0 \subset \mathbb{G}$ is a Poisson submanifold of \mathbb{G} if and only if $\mathfrak{h}_0^{\perp} \oplus \mathfrak{g}_1^*$ is an ideal of \mathfrak{g}^* .

Considering the Lie 2-algebra structure on \mathfrak{g}^* , we note that $\mathfrak{h}_0^{\perp} \oplus \mathfrak{g}_1^* \subset \mathfrak{g}^*$ is a Lie subalgebra if and only if it is an ideal, if and only if $\mathfrak{h}_0^{\perp} \subset \mathfrak{g}_0^*$ is a \mathfrak{g}_1^* -submodule. So with the quotient groupoid structure, \mathbb{G}/H_0 is a quotient Poisson groupoid.

Definition 4.2. Let $(\mathbb{G}, \Pi_{\mathbb{G}})$ be a Poisson Lie 2-group and $\mathbb{H} \subset \mathbb{G}$ a closed 2-subgroup such that $H_0 \subset \mathbb{G}$ is a Poisson subgroup. The homogeneous \mathbb{G} -space \mathbb{G}/\mathbb{H} is called a **Poisson homogeneous** \mathbb{G} -space if there is a Poisson structure Π such that $(\mathbb{G}/\mathbb{H}, \Pi)$ is both a Poisson homogeneous space of the Poisson Lie group \mathbb{G} and a Poisson homogeneous space of the Poisson groupoid \mathbb{G}/H_0 .

For a Poisson 2-group $(\mathbb{G}, \pi_{\mathbb{G}})$ with Lie 2-bialgebra $(\mathfrak{g}, \mathfrak{g}^*)$, we see that $(\mathfrak{g}, \mathfrak{g}^*) = (\mathfrak{g}_0 \ltimes \mathfrak{g}_1, \mathfrak{g}_1^* \ltimes \mathfrak{g}_0^*)$ is a Lie bialgebra and $(\mathfrak{g}_0, \mathfrak{g}_0^*), (\mathfrak{g}_1, \mathfrak{g}_1^*)$ are both Lie bialgebras; see [6, Proposition 3.6]. But we emphasize that the doubles $\mathfrak{g}_i \Join \mathfrak{g}_i^* \subset \mathfrak{g} \Join \mathfrak{g}^*$ for i = 0, 1 are not Lie subalgebras. For example, for $x \in \mathfrak{g}_0$ and $\alpha \in \mathfrak{g}_0^*$, the bracket $[x, \alpha]_{\mathfrak{g} \bowtie \mathfrak{g}^*}$ also has component in \mathfrak{g}_1 , which is determined by $\langle [x, \alpha]_{\mathfrak{g} \Join \mathfrak{g}^*}, \xi \rangle = -\langle x, \xi \triangleright \alpha \rangle$ for $\xi \in \mathfrak{g}_1^*$.

A Dirac structure L of the Lie bialgebra $(\mathfrak{g}_1, \mathfrak{g}_1^*)$ is H_0 -invariant if $H_0 \triangleright L = L$, where the action is the natural action of H_0 on $\mathfrak{g}_1 \oplus \mathfrak{g}_1^*$.

Theorem 4.3. Let $(\mathbb{G}, \Pi_{\mathbb{G}})$ be a Poisson Lie 2-group and $\mathbb{H} \subset \mathbb{G}$ a closed 2-subgroup such that $H_0 \subset \mathbb{G}$ is a Poisson subgroup. Then there is a one-one correspondence between Poisson homogeneous \mathbb{G} -spaces \mathbb{G}/\mathbb{H} and H_0 -invariant Dirac structures L of the Lie bialgebra $(\mathfrak{g}_1, \mathfrak{g}_1^*)$ such that $L \cap \mathfrak{g}_1 = \mathfrak{h}_1$.

We first characterize Dirac structures of the Lie bialgebra $(\mathfrak{g}, \mathfrak{g}^*)$. Let us convent that $x, y \in \mathfrak{g}_0, u, v \in \mathfrak{g}_1$ and $\xi, \eta \in \mathfrak{g}_1^*, \alpha, \beta \in \mathfrak{g}_0^*$. A maximal isotropic subspace $L \subset \mathfrak{g} \oplus \mathfrak{g}^*$ such that $L \cap \mathfrak{g}_i$ is a subspace of \mathfrak{g}_i for i = 0, 1 is always of the form:

$$L = \{x + u + r^{\sharp}(\xi) + K^{\sharp}(\xi) + \pi^{\sharp}(\alpha) + K^{\sharp}(\alpha) + \xi + \alpha | x \in \mathfrak{h}_{0}, u \in \mathfrak{h}_{1}, \xi \in \mathfrak{h}_{1}^{\perp}, \alpha \in \mathfrak{h}_{0}^{\perp} \}$$

(20)
$$= \mathfrak{h} \oplus \operatorname{Graph}(\Omega|_{\mathfrak{h}^{\perp}}),$$

where $\mathfrak{h}_i \subset \mathfrak{g}_i, i = 0, 1$ are subspaces and

 $\Omega = \pi + K + r \in \wedge^2 \mathfrak{g}_0 \oplus \mathfrak{g}_0 \wedge \mathfrak{g}_1 \oplus \wedge^2 \mathfrak{g}_1.$

Here $K^{\sharp} : \mathfrak{g}_0^* \to \mathfrak{g}_1$ and $K^{\sharp} : \mathfrak{g}_1^* \to \mathfrak{g}_0$ are defined by $K^{\sharp}(\alpha) = K(\alpha, \cdot)$ and $K^{\sharp}(\xi) = -K(\cdot, \xi)$. Such a pair (\mathfrak{h}, Ω) with $\mathfrak{h} \subset \mathfrak{g}$ and $\Omega \in \wedge^2 \mathfrak{g}$ is a characteristic pair of L.

Lemma 4.4. Let $L \subset \mathfrak{g}_1 \oplus \mathfrak{g}_1^*$ be a Dirac structure of the Lie bialgebra $(\mathfrak{g}_1, \mathfrak{g}_1^*)$ with the characteristic pair $(\mathfrak{h}_1, r \in \wedge^2 \mathfrak{g}_1)$. Then L is H_0 -invariant if and only if $H_0 \triangleright \mathfrak{h}_1 \subset \mathfrak{h}_1$ and $H_0 \triangleright r \equiv r \pmod{\mathfrak{h}_1}$.

Proof. Note that L is H_0 -invariant if $h_0 \triangleright (x + r^{\sharp}\xi + \xi) \in L$ for $h_0 \in H_0, x \in \mathfrak{h}_1$ and $\xi \in \mathfrak{h}_1^{\perp}$, which is equivalent to

$$h_0 \triangleright \mathfrak{h}_1 \subset \mathfrak{h}_1, \qquad h_0 \triangleright (r^{\sharp}\xi) - r^{\sharp}(h_0 \triangleright^* \xi) \in \mathfrak{h}_1.$$

The second equation amounts to the fact that $h_0 \triangleright r - r \in \mathfrak{h}_1 \land \mathfrak{g}_1$.

Lemma 4.5. Let $(\mathfrak{g}, \mathfrak{g}^*)$ be a Lie 2-bialgebra and assume $\mathfrak{h} \subset \mathfrak{g}$ is a Lie 2-subalgebra. Then a maximal isotropic subspace $L \subset \mathfrak{g} \oplus \mathfrak{g}^*$ with characteristic pair $(\mathfrak{h}, r \in \wedge^2 \mathfrak{g}_1)$ is a Dirac structure of the Lie bialgebra $(\mathfrak{g}, \mathfrak{g}^*)$ if and only if the characteristic pair (\mathfrak{h}, r) defines an \mathfrak{h}_0 -invariant Dirac structure L_1 of $(\mathfrak{g}_1, \mathfrak{g}_1^*)$ such that $\mathfrak{h}_1^{\perp} \triangleright \mathfrak{h}_0^{\perp} \subset \mathfrak{h}_0^{\perp}$.

Proof. First, following from the general framework for Dirac structures of Lie bialgebras, we see that $L \subset \mathfrak{g} \oplus \mathfrak{g}^*$ with characteristic pair $(\mathfrak{h}, r \in \wedge^2 \mathfrak{g}_1)$ is a Dirac structure of $(\mathfrak{g}, \mathfrak{g}^*)$ if and only if

- (i) $\mathfrak{h} \subset \mathfrak{g}$ is a Lie subalgebra;
- (ii) $d_*r + \frac{1}{2}[r, r] \equiv 0 \pmod{\mathfrak{h}_1};$
- (iii) $[a,b] + [a,b]_r^g \in \mathfrak{h}^{\perp}$ for $a, b \in \mathfrak{h}^{\perp}$, where $[a,b]_r^g = \operatorname{ad}_{r^{\sharp}a}^* b \operatorname{ad}_{r^{\sharp}b}^* a$ and ad^* is the coadjoint action of \mathfrak{g} on \mathfrak{g}^* .

Note that $d_*r = d_{\mathfrak{a}_1^*}r$ in (ii). Then, we claim that (iii) is equivalent to the following conditions:

$$[\mathfrak{h}_0^{\perp},\mathfrak{h}_0^{\perp}] \subset \mathfrak{h}_0^{\perp}, \qquad \mathfrak{h}_1^{\perp} \triangleright \mathfrak{h}_0^{\perp} \subset \mathfrak{h}_0^{\perp}, \qquad [\xi,\eta] + [\xi,\eta]_r \in \mathfrak{h}_1^{\perp}, \qquad x \triangleright r \equiv 0 \pmod{\mathfrak{h}_1},$$

for all $x \in \mathfrak{h}_0$ and $\xi, \eta \in \mathfrak{h}_1^{\perp}$, where $[\xi, \eta]_r = \mathfrak{a} \mathfrak{d}_{r\sharp\xi}^* \eta - \mathfrak{a} \mathfrak{d}_{r\sharp\eta}^* \xi$ and $\mathfrak{a} \mathfrak{d}^*$ is the coadjoint action of \mathfrak{g}_1 on \mathfrak{g}_1^* . In fact, for $a, b \in \mathfrak{h}_0^{\perp}$ and $a \in \mathfrak{h}_0^{\perp}, b \in \mathfrak{h}_1^{\perp}$, we obtain the first two conditions. For $\xi, \eta \in \mathfrak{h}_1^{\perp}$, we find that

$$\operatorname{pr}_{\mathfrak{g}_1^*}([\xi,\eta] + [\xi,\eta]_r^{\mathfrak{g}}) = [\xi,\eta] + [\xi,\eta]_r$$

and the \mathfrak{g}_0^* -part is determined by

$$\langle \mathrm{pr}_{\mathfrak{g}_0^*}([\xi,\eta] + [\xi,\eta]_r^{\mathfrak{g}}), x \rangle = \langle \mathrm{pr}_{\mathfrak{g}_0^*}([\xi,\eta]_r^{\mathfrak{g}}), x \rangle = \langle \eta, x \triangleright (r^{\sharp}\xi) \rangle - \langle \xi, x \triangleright (r^{\sharp}\eta) \rangle = \langle x \triangleright r, \xi \land \eta \rangle,$$

which implies $x \triangleright r \in \mathfrak{h}_1 \land \mathfrak{g}_1$ for any $x \in \mathfrak{h}_0$. Therefore, for $a, b \in \mathfrak{h}_1^{\perp}$, the condition $[a, b] + [a, b]_r^{\mathfrak{g}} \in \mathfrak{h}^{\perp}$ infers the last two conditions.

Next, on the condition that $\mathfrak{h} \subset \mathfrak{g}$ is a Lie 2-subalgebra, $\mathfrak{h}_1^{\perp} \triangleright \mathfrak{h}_0^{\perp} \subset \mathfrak{h}_0^{\perp}$ implies that $\mathfrak{h}_0^{\perp} \subset \mathfrak{g}_0^*$ is a Lie subalgebra. Thus, *L* is a Dirac structure of $(\mathfrak{g}, \mathfrak{g}^*)$ if and only if

$$\mathfrak{h}_1^{\perp} \triangleright \mathfrak{h}_0^{\perp} \subset \mathfrak{h}_0^{\perp}, \quad d_{\mathfrak{g}_1^*}r + \frac{1}{2}[r,r] \equiv 0 \pmod{\mathfrak{h}_1}, \quad [\xi,\eta] + [\xi,\eta]_r \in \mathfrak{h}_1^{\perp}, \quad x \triangleright r \equiv 0 \pmod{\mathfrak{h}_1},$$

for $\xi, \eta \in \mathfrak{h}_1^{\perp}$ and $x \in \mathfrak{h}_0$, which implies that L_1 is an \mathfrak{h}_0 -invariant Dirac structure of $(\mathfrak{g}_1, \mathfrak{g}_1^*)$ such that $\mathfrak{h}_1^{\perp} \triangleright \mathfrak{h}_0^{\perp} \subset \mathfrak{h}_0^{\perp}$.

This Lemma demonstrates that a Dirac structure $L \subset \mathfrak{g} \Join \mathfrak{g}^*$ with characteristic pair (\mathfrak{h}, r) for $r \in \wedge^2 \mathfrak{g}_1$ induces two Dirac structures $L_1 \subset \mathfrak{g}_1 \Join \mathfrak{g}_1^*$ and $\mathfrak{h}_0 \oplus \mathfrak{h}_0^{\perp} \subset \mathfrak{g}_0 \Join \mathfrak{g}_0^*$, respectively. Moreover, we have $L = L_1 \oplus (\mathfrak{h}_0 \oplus \mathfrak{h}_0^{\perp})$ as vector spaces.

Lemma 4.6. Given a Poisson 2-group $(\mathbb{G}, \Pi_{\mathbb{G}})$, let L be a Dirac structure of the Lie bialgebra $(\mathfrak{g}, \mathfrak{g}^*)$ whose characteristic pair is $(\mathfrak{h}, \Omega = \pi + K + r)$, then $(\mathbb{G}/\mathbb{H}, \Pi)$ is a Poisson homogeneous space of the Poisson group \mathbb{G} with the action given by the group left translation and

$$\Pi = \operatorname{pr}_*(\Pi_{\mathbb{G}} + \overline{\Omega}^{gp}),$$

where $\overleftarrow{\cdot}^{gp}$ denotes the left translation relative to the Lie group structure on \mathbb{G} . In fact, suppose

(21)
$$\pi = \omega^{ij} e_i \wedge e_j \in \wedge^2 \mathfrak{g}_0, \quad K = \theta^{ik} e_i \wedge \xi_k \in \mathfrak{g}_0 \wedge \mathfrak{g}_1, \quad r = \sigma^{kl} \xi_k \wedge \xi_l \in \wedge^2 \mathfrak{g}_1,$$

where $\{e_i\}$ and $\{\xi_k\}$ are bases of \mathfrak{g}_0 and \mathfrak{g}_1 , respectively. Then

$$\overleftarrow{\Omega}^{gp} = \Omega^{2,0} + \Omega^{1,1} + \Omega^{0,2} \in \mathfrak{X}^2(G_0 \ltimes G_1)$$

is given by

$$\begin{split} \Omega^{2,0} &= \omega^{ij} \overleftarrow{e_i} \wedge \overleftarrow{e_j} &\in \mathfrak{X}^2(G_0); \\ \Omega^{1,1} &= -2\omega^{ij} \overleftarrow{e_i} \wedge \widehat{e_j} + \theta^{ik} \overleftarrow{e_i} \wedge \overleftarrow{\xi_k} &\in \mathfrak{X}(G_0) \otimes \mathfrak{X}(G_1); \\ \Omega^{0,2} &= \omega^{ij} \widehat{e_i} \wedge \widehat{e_j} - \theta^{ik} \widehat{e_i} \wedge \overleftarrow{\xi_k} + \sigma^{kl} \overleftarrow{\xi_k} \wedge \overleftarrow{\xi_l} &\in \mathfrak{X}^2(G_1), \end{split}$$

where $\overleftarrow{e_i} \in \mathfrak{X}(G_0)$ and $\overleftarrow{\xi} \in \mathfrak{X}(G_1)$ are vector fields generated by left translations and $\widehat{e_i} \in \mathfrak{X}(G_1)$ is the fundamental vector field coming from the action of G_0 on G_1 .

Proof. The result follows from Drinfeld's classification theorem for Poisson homogeneous spaces in [8]. Here we compute the formula of $\widehat{\Omega}^{gp}$. For $x \in \mathfrak{g}_0, u \in \mathfrak{g}_1$, with respect to the group multiplication on G, we have

$$\overleftarrow{x}^{gp}(g_0,g_1) = (L_{g_0*}x, \frac{d}{dt}|_{t=0}(e^{-tx} \triangleright g_1) = (L_{g_0*}x, -\widehat{x}_{g_1}), \qquad \overleftarrow{u}^{gp}(g_0,g_1) = L_{g_1*}u.$$

This shows that $\overleftarrow{e_i}^{gp} = \overleftarrow{e_i} - \widehat{e_i} \in TG_0 \oplus TG_1$ and $\overleftarrow{\xi_k}^{gp} = \overleftarrow{\xi_k}$.

With the notations and assumptions in Lemma 4.1, denote by (A, A^*) the Lie bialgebroid of the quotient Poisson groupoid $(\mathbb{G}/H_0, \mathrm{pr}_*\pi_{\mathbb{G}}) = (G_0 \times_{H_0} G_1, \mathrm{pr}_*\pi_{\mathbb{G}})$. Note that $A = G_0 \times_{H_0} \mathfrak{g}_1 \to G_0/H_0$ is a quotient of an action Lie algebroid and

 $\Gamma(A) = \{ \Omega' \in C^{\infty}(G_0, \mathfrak{g}_1); \Omega'(g_0 h_0) = h_0^{-1} \triangleright \Omega'(g_0) \}.$

A maximal isotropic subbundle $L \subset A \oplus A^*$ is of the form:

(22)
$$L = \{X + \Omega^{\sharp} \xi + \xi | X \in D, \xi \in D^{\perp}\} = D \oplus \operatorname{Graph}(\Omega|_{D^{\perp}}),$$

where $D \subset A$ is a subbundle, $D^{\perp} \subset A^*$ is the conormal bundle of D and $\Omega \in \Gamma(\wedge^2 A)$. In fact, Ω is determined by its projection $pr(\Omega) \in \Gamma(\wedge^2 A/D)$. Such a pair (D, Ω) is called a characteristic pair of L. For example, for any $r \in \wedge^2 \mathfrak{g}_1$ such that $H_0 \triangleright r \equiv r \pmod{\mathfrak{h}_1}$, the projection

$$\operatorname{pr}(r) \in \wedge^2 (\mathfrak{g}_{-1}/\mathfrak{h}_{-1})^{H_0} \subset \Gamma(\wedge^2 A/D)$$

is a 2-section of A/D. Then any lifting $pr(r) \in \Gamma(\wedge^2 A)$ gives rise to a 2-section of A.

Lemma 4.7. Let (A, A^*) be the above Lie bialgebroid, $L \subset A \oplus A^*$ a maximal isotropic subbundle with characteristic pair $(D = G_0 \times_{H_0} \mathfrak{h}_1, \operatorname{pr}(r))$, where $r \in \wedge^2 \mathfrak{g}_1$ such that $H_0 \triangleright r \equiv r \pmod{\mathfrak{h}_1}$. Then L is a Dirac structure of the Lie bialgebroid (A, A^*) if and only if the following conditions hold:

- (1) $\mathfrak{h}_1 \subset \mathfrak{g}_1$ is a Lie subalgebra;
- (2) $d_{\mathfrak{g}_1^*}r + \frac{1}{2}[r,r] \equiv 0 \pmod{\mathfrak{h}_1};$
- (3) $[\xi, \eta] + [\xi, \eta]_r \in \mathfrak{h}_1^\perp$ for any $\xi, \eta \in \mathfrak{h}_1^\perp$.

Proof. Note that $(D = G_0 \times_{H_0} \mathfrak{h}_1, \operatorname{pr}(r))$ is a Dirac structure of the Lie bialgebroid (A, A^*) if and only if $(G_0 \times \mathfrak{h}_1, r)$ is an H_0 -invariant Dirac structure of the Lie bialgebroid $(G_0 \times \mathfrak{g}_1, G_0 \times \mathfrak{g}_1^*)$, if and only if (\mathfrak{h}_1, r) is a Dirac structure of the Lie bialgebra $(\mathfrak{g}_1, \mathfrak{g}_1^*)$. This proves the lemma.

Now we are ready to prove Theorem 4.3.

Proof. Given a Poisson homogeneous space $(\mathbb{G}/\mathbb{H},\Pi)$ of the Poisson 2-group \mathbb{G} , since it is a Poisson homogeneous space of the Poisson group \mathbb{G} , we have

$$\Pi - \mathrm{pr}_*(\Pi_{\mathbb{G}}) = \mathrm{pr}_* \overleftarrow{\Omega}^{gp}$$

for some $\Omega \in \wedge^2 \mathfrak{g} = (\wedge^2 \mathfrak{g}_0) \oplus (\mathfrak{g}_0 \wedge \mathfrak{g}_1) \oplus (\wedge^2 \mathfrak{g}_1)$. Here we use $\overleftarrow{\cdot} gp$ to denote the left translation with respect to the group multiplication. As $(\mathbb{G}/\mathbb{H},\Pi)$ is also a Poisson homogeneous space of the Poisson groupoid $(\mathbb{G}/H_0, \mathrm{pr}_*\pi_{\mathbb{G}})$, we have

$$\Pi - \mathrm{pr}_*(\Pi_{\mathbb{G}}) = \mathrm{pr}_* \overleftarrow{\Omega'}^{gpd}$$

for

$$\Omega' \in \Gamma(\wedge^2 A) = \{ \Omega' \in C^{\infty}(G_0, \wedge^2 \mathfrak{g}_1); \Omega'(g_0 h_0) = h_0^{-1} \triangleright \Omega'(g_0) \},\$$

where $\overleftarrow{\cdot}^{gpd}$ denotes the left translation with respect to the groupoid multiplication. By Lemma 4.6, we have

$$\overleftarrow{\Omega}^{gp} = \Omega^{2,0} + \Omega^{1,1} + \Omega^{0,2} \in \mathfrak{X}^2(G_0 \ltimes G_1), \quad \overleftarrow{\Omega}^{'gpd} = \Omega^{'0,2} \in \mathfrak{X}^2(G_0 \ltimes G_1).$$

So in order to make sure $pr_*(\overline{\Omega}^{gp} - \overline{\Omega}^{gpd}) = 0$ holds, we must have

 $\Omega = r \in \wedge^2 \mathfrak{g}_1 \pmod{\mathfrak{h}_1}, \qquad \Omega' = \widehat{\operatorname{pr}(r)},$ and $H_0 \triangleright r \equiv r \pmod{\mathfrak{h}_1}$.

Here observe that $\overleftarrow{r}^{gpd} = \overleftarrow{r}^{gp}$.

Then applying Drinfeld theorems for Poisson homogeneous spaces of Poisson groups and groupoids, and by Lemmas 4.4, 4.5 and 4.7, we obtain the one-one correspondence. To be precise, the characteristic pairs of the Dirac structures of the bialgebra $(\mathfrak{g}, \mathfrak{g}^*)$ and the Lie bialgebroid (A, A^*) are $(\mathfrak{h}_0 \oplus \mathfrak{h}_1, r)$ and $(G_0 \times_{H_0} \mathfrak{h}_1, \operatorname{pr}(r))$, respectively. \square

Recall that a bivector field Π on a Lie groupoid P is **multiplicative** if and only if the submanifold

$$\Gamma = \{(g, h, gh); t(g) = s(h)\} \subset P \times P \times P$$

is coisotropic with respect to $\Pi \oplus \Pi \oplus (-\Pi)$. It is called **affine** if and only if the submanifold

$$\Lambda = \{(g, h, l, hg^{-1}l); t(h) = t(g), s(g) = s(l)\} \subset P \times P \times P \times P$$

is coisotropic with respect to $\Pi \oplus (-\Pi) \oplus (-\Pi) \oplus \Pi$. See [19] for affine Poisson structures and [11] for the general case. A Lie groupoid with an affine Poisson structure is called an affine Poisson groupoid.

Corollary 4.8. With the assumptions in Theorem 4.3, if $\mathfrak{h}_1^{\perp} \subset \mathfrak{g}_1^*$ is a Lie subalgebra, then Poisson homogeneous \mathbb{G} -spaces \mathbb{G}/\mathbb{H} are affine Poisson groupoids.

Proof. If $H_0 \subset \mathbb{G}$ is a Poisson subgroup, we see $\mathfrak{h}_0^{\perp} \subset \mathfrak{g}_0^*$ is a \mathfrak{g}_1^* -submodule. Together with the condition $[\mathfrak{h}_1^{\perp}, \mathfrak{h}_1^{\perp}] \subset \mathfrak{h}_1^{\perp}$, we have $\mathfrak{h}^{\perp} = \mathfrak{h}_0^{\perp} \oplus \mathfrak{h}_1^{\perp}$ is a Lie subalgebra of \mathfrak{g}^* , which leads to the fact that $(\mathbb{G}/\mathbb{H},\mathrm{pr}_*\Pi_\mathbb{G})$ is a Poisson manifold and it is obviously a Poisson groupoid.

It is known from [11] that $\Pi \in \mathfrak{X}^2(P)$ is multiplicative if and only if $\Pi + \overleftarrow{\lambda}^{gpd}$ for all $\lambda \in \Gamma(\wedge^2 A_P)$ is affine, where A_P is the Lie algebroid of the Lie groupoid P. By the proof of Theorem 4.3, the Poisson structure on the Poisson homogeneous space \mathbb{G}/\mathbb{H} is

$$\Pi = \mathrm{pr}_*(\Pi_{\mathbb{G}} + \overleftarrow{r}^{gp}) = \mathrm{pr}_*\Pi_{\mathbb{G}} + \widecheck{\mathrm{pr}(r)}^{gpd}, \qquad r \in \wedge^2 \mathfrak{g}_1,$$

which is affine.

By [6], a Poisson 2-group $(G_0 \ltimes G_1 \rightrightarrows G_0, \Pi_{\mathbb{G}})$ has a mixed product Poisson structure (see [14]). Namely, both G_0 and G_1 are quotient Poisson manifolds. Moreover, (G_1, Π_{G_1}) is a Poisson subgroup of \mathbb{G} , where $\Pi_{G_1} = \Pi_{\mathbb{G}}|_{\mathbb{G}_1}$.

Corollary 4.9. Let \mathbb{G}/\mathbb{H} be a Poisson homogeneous space of the Poisson 2-group \mathbb{G} . Then G_1/H_1 is a Poisson submanifold of \mathbb{G}/\mathbb{H} and it is a Poisson homogeneous space of the Poisson group G_1 .

Proof. Suppose \mathbb{G}/\mathbb{H} with the Poisson structure $\Pi = \mathrm{pr}_*(\Pi_{\mathbb{G}} + \overleftarrow{r})$ for $r \in \wedge^2 \mathfrak{g}_1$ is a Poisson homogeneous space of the Poisson 2-group ($\mathbb{G}, \Pi_{\mathbb{G}}$). Since G_1 is a Poisson subgroup of \mathbb{G} and $r \in \wedge^2 \mathfrak{g}_1$, we see that (\mathfrak{h}_1, r) is a Dirac structure of the Lie bialgebra $(\mathfrak{g}_1, \mathfrak{g}_1^*)$. By Drinfeld's theorem, we have that $(G_1/H_1, \operatorname{pr}_*(\Pi_{G_1} + \overleftarrow{r}))$ is a Poisson homogeneous space of the Poisson group (G_1, Π_{G_1}) . Moreover, it is obvious that the inclusion map

$$G_1/H_1 \hookrightarrow \mathbb{G}/\mathbb{H}, \qquad [g_1] \mapsto [e, g_1]$$

is a Poisson map.

Let us recall a class of Lie 2-bialgebras coming from an r-matrix, called a coboundary Lie 2**bialgebra**. Let \mathfrak{g} be a Lie 2-algebra. An element $\mu \in \wedge^2 \mathfrak{g}_1$ is called an *r*-matrix of \mathfrak{g} if $x \triangleright [\mu, \mu] = 0$ for all $x \in \mathfrak{g}_0$. Such a μ gives rise to a Lie 2-bialgebra $(\mathfrak{g}, \mathfrak{g}^*)$, where the Lie bracket on \mathfrak{g}_1^* is defined by

$$\langle [\xi,\eta]_{\mu},u\rangle := \langle \xi \wedge \eta, [\mu,u]\rangle, \qquad \forall \xi,\eta \in \mathfrak{g}_1^*, u \in \mathfrak{g}_1$$

and the action of \mathfrak{g}_1^* on \mathfrak{g}_0^* is given by

(23)
$$\langle \xi \triangleright \alpha, x \rangle := \langle \xi \land \phi^* \alpha, x \triangleright \mu \rangle, \qquad \xi \in \mathfrak{g}_1^*, \alpha \in \mathfrak{g}_0^*, x \in \mathfrak{g}_0.$$

If $[\mu, \mu] = 0$, the coboundary Lie 2-bialgebra is called a **triangular Lie 2-bialgebra**. If the Lie 2-algebra \mathfrak{g} is integrated to a Lie 2-group \mathbb{G} , we further obtain the triangular Poisson 2-group $(\mathbb{G}, \lambda_{\mu}^{gpd} - \overline{\lambda_{\mu}}^{gpd})$, where $\lambda_{\mu} : G_0 \to \wedge^2 \mathfrak{g}_1$ is a 2-section of the Lie algebroid $G_0 \ltimes \mathfrak{g}_1 \to G_0$ defined by

$$\lambda_{\mu}(g_0) = g_0^{-1} \triangleright \mu - \mu, \qquad \forall g_0 \in G_0.$$

See [7, Theorem 3.11] for more details.

Corollary 4.10. Let $(\mathbb{G}, \overleftarrow{\lambda_{\mu}}^{gpd} - \overrightarrow{\lambda_{\mu}}^{gpd})$ be a triangular Poisson 2-group. Then a maximal isotropic subspace L with characteristic pair (\mathfrak{h}_1, r) for $r \in \wedge^2 \mathfrak{g}_1$ defines a Dirac structure of $(\mathfrak{g}_1, \mathfrak{g}_1^*)$ if and only if the following conditions holds

- (1) $\mathfrak{h}_1 \subset \mathfrak{g}_1$ is a Lie subalgebra;
- (2) $[\mu + r, \mu + r] \equiv 0 \pmod{\mathfrak{h}_1};$
- (3) $[u, \mu + r] \equiv 0 \pmod{\mathfrak{h}_1}$ for all $u \in \mathfrak{h}_1$.

In this case, $H_0 \subset \mathbb{G}$ is a Poisson subgroup if and only if $(\phi \wedge 1)(x \triangleright \mu) \in \mathfrak{h}_0 \wedge \mathfrak{g}_{-1}$ for all $x \in \mathfrak{h}_0$. On this condition, Poisson homogenous spaces \mathbb{G}/\mathbb{H} correspond to H_0 -invariant Dirac structures of $(\mathfrak{g}_1, \mathfrak{g}_1^*)$.

Proof. The first part of the corollary follows directly from [12, Corollary 3.2]. Then $H_0 \subset \mathbb{G}$ is a Poisson subgroup if and only if $\mathfrak{h}_0^{\perp} \subset \mathfrak{g}_0^*$ is a \mathfrak{g}_{-1}^* -submodule. By (23), taking $\xi \in \mathfrak{g}_1^*, \alpha \in \mathfrak{h}_0^{\perp}$, we have $\xi \triangleright \alpha \in \mathfrak{h}_0^{\perp}$ if and only if $\langle \xi \land \alpha, (1 \land \phi)(x \triangleright \mu) \rangle = 0, \forall x \in \mathfrak{h}_0$, which amounts to say $(1 \land \phi)(x \triangleright \mu) \in \mathfrak{h}_0 \land \mathfrak{g}_1$ for all $x \in \mathfrak{h}_0$. The rest of this result holds by Theorem 4.3.

Example 4.11. When $\mathbb{H} = \{e\}$ in Theorem 4.3, the Poisson homogeneous space of the Poisson 2-group $(\mathbb{G}, \Pi_{\mathbb{G}})$ is \mathbb{G} itself with the affine Poisson structure $\Pi_{\mathbb{G}} + \overleftarrow{r}$ ([11, 19]), where $r \in \wedge^2 \mathfrak{g}_1$ satisfies that $d_{\mathfrak{g}_1^*}r + \frac{1}{2}[r, r] = 0$. The corresponding Dirac structure of $(\mathfrak{g}_1, \mathfrak{g}_1^*)$ is $\operatorname{Graph}(r^{\sharp})$.

Example 4.12. Let $(\mathbb{G}, \Pi_{\mathbb{G}})$ be a Poisson 2-group with a closed 2-subgroup \mathbb{H} satisfying that $H_0 \subset \mathbb{G}$ is a Poisson subgroup. Then $(\mathbb{G}/\mathbb{H}, \mathrm{pr}_*\Pi_{\mathbb{G}})$ is a Poisson homogeneous space of the Poisson 2-group \mathbb{G} if and only if $\mathfrak{h}_1^{\perp} \subset \mathfrak{g}_1^*$ is a Lie subalgebra. The corresponding Dirac structure of $(\mathfrak{g}_1, \mathfrak{g}_1^*)$ is $\mathfrak{h}_1 \oplus \mathfrak{h}_1^{\perp}$. In this case, $(\mathbb{G}/\mathbb{H}, \mathrm{pr}_*\Pi_{\mathbb{G}})$ is a quotient Poisson groupoid.

Example 4.13. Let \mathbb{G} be a Poisson 2-group with trivial Poisson structure and assume that $\mathbb{H} \subset \mathbb{G}$ is a closed 2-subgroup. Then there is a one-one correspondence between Poisson homogeneous spaces $(\mathbb{G}/\mathbb{H}, \mathrm{pr}_* \overleftarrow{r})$ of \mathbb{G} and H_0 -invariant Dirac structures L of $(\mathfrak{g}_1, \mathfrak{g}_1^*)$ with characteristic pairs (\mathfrak{h}_1, r) , in which $r \in \wedge^2 \mathfrak{g}_1$ satisfies that

$$[r,r] \equiv 0 \pmod{\mathfrak{h}_1}, \qquad [\mathfrak{h}_1^{\perp},\mathfrak{h}_1^{\perp}]_r \subset \mathfrak{h}_1^{\perp}, \qquad H_0 \triangleright r \equiv r \pmod{\mathfrak{h}_1}.$$

Example 4.14. Let $(\mathfrak{g}, \mathfrak{g}^*)$ be a Lie 2-bialgebra with trivial Lie 2-algebra structure on \mathfrak{g} . Then the abelian 2-group $\mathbb{G} = \mathfrak{g}$ with the linear Poisson structure Π_{KKS} is a Poisson 2-group. Let $\mathfrak{h} : \mathfrak{h}_1 \to \mathfrak{h}_0$ be a 2-subspace of \mathfrak{g} such that $\mathfrak{g}_1^* \triangleright \mathfrak{h}_0^\perp \subset \mathfrak{h}_0^\perp$. Then there is a one-one correspondence between Poisson homogeneous spaces $(\mathfrak{g}/\mathfrak{h}, \mathrm{pr}_*(\Pi_{KKS} + r))$ and Dirac structures of $(\mathfrak{g}_1, \mathfrak{g}_1^*)$ with characteristic pair (\mathfrak{h}_1, r) . Here r is an element in $\wedge^2 \mathfrak{g}_1$ satisfies that $d_{\mathfrak{g}_1^*}r \equiv 0 \pmod{\mathfrak{h}_1}$ and $[\mathfrak{h}_1^\perp, \mathfrak{h}_1^\perp] \subset \mathfrak{h}_1^\perp$.

REFERENCES

- [1] J. Baez and A. Crans, Higher dimensional algebra VI: Lie 2-algebras, Theory Appl. Categ. 12 (2014), 492-538.
- [2] J. Baez and A. Lauda, Higher dimensional algebra VI: Lie 2-groups, Theory Appl. Categ. 12 (2014), 423-491.
- [3] C. Bai, Y. Sheng and C. Zhu, Lie 2-bialgebras, Commun. Math. Phys. 320 (2013), no. 1, 70-90.
- [4] C. Blohamnn and A. Weinstein, Hamiltonian Lie algebroids, Mem. Am. Math. Soc. 295, vi+99 pp., Providence, RI: American Mathematical Society (AMS) 2024.
- [5] H. Bursztyn, D. Iglesias-Ponte and J. Lu, Dirac geometry and integration of Poisson homogeneous spaces, J. Differential Geom. 126 (2024), 939-1000.
- [6] Z. Chen, M. Stienon and P. Xu, Weak Lie 2-bialgebras, J. Geom. Phys. 68 (2013), 59-68.
- [7] Z. Chen, M. Stienon and P. Xu, Poisson 2-groups, J. Differential Geom. 94 (2013), no. 2, 209-240.
- [8] V. G. Drinfel'd, On Poisson homogeneous spaces of Poisson-Lie groups, Theo. Math. Phys. 95 (1993), 226-227.
- [9] A. Garmendia and M. Zambon, Quotients of singular foliations and Lie 2-group actions, J. Noncommut. Geom. 15 (2021), no. 4, pp. 1251-1283.
- [10] M. Jotz Lean, Homogeneous spaces of Dirac groupoids, J. Geom. Phys. 104 (2016), 89-111.
- [11] H. Lang, Z. Liu and Y. Sheng, Affine structures on Lie groupoids, Pacific J. Math. 307 (2020), 353-382.
- [12] Z. Liu, Some remarks on Dirac structures and Poisson reductions, Banach Center Publications 51 (2000), 165-173.

- [13] Z. Liu, A. Weinstein and P. Xu, Dirac structures and Poisson homogeneous spaces, *Commun. Math. Phys.* 192 (1998), 121-144.
- [14] J. Lu and V. Mouquin, Mixed product Poisson structures associated to Poisson Lie groups and Lie bialgebras, Int. Math. Res. Notices 19 (2017), 5919-5976.
- [15] J. Lu, A note on Poisson homogeneous spaces, Poisson geometry in mathematics and physics, 173-198, Contemp. Math., 450, Amer. Math. Soc., Providence, RI, 2008.
- [16] K. Mackenzie, General Theory of Lie Groupoids and Lie Algebroids, London Mathematical Society Lecture Note Series, vol. 213, Cambridge University Press.
- [17] K. Norrie, Actions and automorphisms of crossed modules, Bull. Soc. Math. France 118 (1990), 129-146.
- [18] Y. Sheng and C. Zhu, Integration of semidirect product Lie 2-algebras, *Int. J. Geom. Methods Mod. Phys.* 9 (2012), no. 5, 1250043, 1-31.
- [19] A. Weinstein, Affine Poisson structure, Internat. J. Math. 1 (1990), 343-360.
- [20] M. Zambon and C. Zhu, Higher Lie algebra actions on Lie algebroids, J. Geom. Phys. 64 (2013), 155-173.

COLLEGE OF SCIENCE, CHINA AGRICULTURAL UNIVERSITY, BEIJING 100083, CHINA *Email address*: hllang@cau.edu.cn

SCHOOL OF MATHEMATICAL AND STATISTICAL SCIENCES, HENAN UNIVERSITY, KAIFENG 475004, CHINA *Email address*: zhangju@henu.edu.cn