

SHRINKING GRADIENT KÄHLER–RICCI SOLITONS ARE SIMPLY-CONNECTED

CARLOS ESPARZA

ABSTRACT. We show that smooth polarized Fano fibrations have no nontrivial finite covers. Using results by Sun–Zhang and Wylie, it follows that shrinking Kähler–Ricci solitons are simply-connected.

1. INTRODUCTION

A shrinking Kähler–Ricci soliton is a complex manifold M with a complete Kähler metric ω and $f \in C^\infty(M)$ such that

$$\text{Ric}(\omega) + i\partial\bar{\partial}f = \omega .$$

Sun–Zhang have shown [SZ24, Thm. 3.1] that a manifold M admitting such a metric is a quasi-projective variety with additional structure, called a *polarized Fano fibration*:

Definition 1.1 ([SZ24, Def. 2.5]). A smooth complex variety M is a Fano fibration if its algebra of regular functions $R := \mathcal{O}_M^{\text{alg}}(M)$ is finitely generated, the canonical map $\pi: M \rightarrow \text{Spec } R$ is a proper surjection and K_M^{-1} is π -ample.¹

A smooth *polarized* Fano fibration consists of a Fano fibration M equipped with the algebraic action of a torus T and an element $X \in \text{Lie } T$ such that under the weight decomposition

$$R = \bigoplus_{\alpha \in (\text{Lie } T)^*} R_\alpha$$

of $T \curvearrowright R$ we have $\langle \alpha, X \rangle > 0$ for all non-zero α with $R_\alpha \neq 0$, and $T \curvearrowright \text{Spec } R$ only has one fixed point (this last condition is equivalent to $R_0 = \mathbb{C}$).

The purpose of this note is to establish

Proposition 1.2. *Let (M, T, X) be a smooth polarized Fano fibration and let $p: \widetilde{M} \rightarrow M$ be a finite (étale) cover. Then $\widetilde{M} = M$.*

It is known that (Riemannian) shrinking gradient Ricci solitons have finite fundamental group [Wyl08], so we conclude:

Corollary 1.3. *Any shrinking gradient Kähler–Ricci soliton is simply-connected.*

Our proof is inspired by the classical proof that Fano manifolds have no nontrivial finite covers: If $\widetilde{M} \rightarrow M$ is a degree m cover of a Fano manifold then \widetilde{M} is also Fano. By the Hirzebruch–Riemann–Roch formula, the holomorphic Euler characteristic is multiplicative along covers, meaning $\chi(\widetilde{M}, \mathcal{O}) = m\chi(M, \mathcal{O})$. But the Hodge numbers $h^{0,p}(M)$, $p \geq 1$ vanish for any Fano manifold by Serre duality and Kodaira vanishing so $\chi(M, \mathcal{O}) = \chi(\widetilde{M}, \mathcal{O}) = 1$. One concludes that $m = 1$.

To adapt this proof to the noncompact setting of polarized Fano fibrations, we use Wu’s equivariant HRR formula [Wu03, Thm. 3.14]. Sun–Zhang have independently found a different proof of Corollary 1.3, which instead uses Kähler reduction and results from birational geometry. It will appear in a future version of [SZ24].

Date: March 11, 2025.

¹The definition in [SZ24] is worded slightly differently. It is easy to see that the affine variety Y in their definition is necessarily $\text{Spec } R$. Furthermore, the ring of regular functions of a smooth (and thus normal) variety is automatically normal.

Acknowledgements. I would like to thank Song Sun for his encouragement and guidance, and him and Junsheng Zhang for communicating their proof to me. I am grateful to thank Charles Cifarelli for conversations that led me to this question and to Zachary Stier for helpful discussions.

2. PROOF OF PROPOSITION 1.2

Let m be the order of the cover. We can replace T by its m -fold cover in every direction (i.e. we replace $\mathbb{R}^k/\mathbb{Z}^k$ by $\mathbb{R}^k/m\mathbb{Z}^k$), so that T acts on \widetilde{M} and p is T -equivariant.

Since K_M^{-1} is relatively ample over a map $M \rightarrow Y$ to an affine variety, there exist integers $m, N_1, N_2 > 0$, linear diagonal T -actions on \mathbb{C}^{N_1} and \mathbb{P}^{N_2} , and a T -equivariant embedding $(j_1, j_2): M \rightarrow \mathbb{C}^{N_1} \times \mathbb{P}^{N_2}$ such that $j_2^* \mathcal{O}(1) = mK_M^{-1}$ (cf. [SZ24, Def. 2.5]). From this embedding we obtain a hermitian metric $h := e^{-|j_1|^2/2} j_2^* h_{\text{FS}}^{1/m}$ on K_M^{-1} with positive curvature form $\omega := (\omega_{\text{Euc}} \times \omega_{\text{FS}})|_M$. By the definition of the Reeb vector field X , we have $w_i(X) > 0$ for all weights $w_i \in (\text{Lie } T)^*$ of the action on \mathbb{C}^{N_1} . The actions of T on \mathbb{C}^{N_1} and \mathbb{P}^{N_2} are both Hamiltonian, and the ω_{Euc} -moment map for the first action is given by $\sum_i w_i |z_i|^2$. Therefore the action $T \zeta M$ is Hamiltonian for ω , and its moment map $\mu: M \rightarrow \text{Lie } T$ has the property that $\langle \mu, X \rangle$ is proper (for any Reeb vector field X). Since p is finite, this properness is inherited by $\mu \circ p: \widetilde{M} \rightarrow \text{Lie } T$, which is a moment map for $T \zeta \widetilde{M}$. Thus, both actions $T \zeta M$ and $T \zeta \widetilde{M}$ satisfy Assumption 2.15 in [Wu03], if we pick an action chamber C that contains a Reeb vector field.

Let $Z = \text{Fix}_T(M)$, with connected components $\{Z_\alpha\}_{\alpha \in F}$. The covering map p will restrict to (not necessarily connected) covering maps $\widetilde{Z}_\alpha := p^{-1}(Z_\alpha) \rightarrow Z_\alpha$, and $\text{Fix}_T(\widetilde{M}) = \bigcup_{\alpha \in F} \widetilde{Z}_\alpha$. Applying [Wu03, Thm. 3.14] to the trivial line bundle \mathcal{O} on M we obtain

$$\sum_q (-1)^q \text{char } H^q(M, \mathcal{O}) = \sum_{\alpha \in F} \int_{Z_\alpha} \Omega_{Z_\alpha} \quad (2.1)$$

$$\text{where } \Omega_{Z_\alpha} = (-1)^{n-n_\alpha - \text{rk } \mathcal{N}_{Z_\alpha/M}^C} \text{ch}_T \left(\frac{\det(\mathcal{N}_{Z_\alpha/M}^{-C})}{\det(1 - (\mathcal{N}_{Z_\alpha/M}^C)^*) \det(1 - \mathcal{N}_{Z_\alpha/M}^{-C})} \right) \text{td}(Z_\alpha)$$

and an analogous formula for \widetilde{M} and $\{\widetilde{Z}_\alpha\}$. Here $\mathcal{N}_{Z_\alpha/M}^{\pm C} \leq \mathcal{N}_{Z_\alpha/M}$ is the subbundle on which elements of $C \subseteq \text{Lie } T$ act with positive/negative weight and $n_\alpha = \dim Z_\alpha = \dim \widetilde{Z}_\alpha$. Evidently $\mathcal{N}_{\widetilde{Z}_\alpha/\widetilde{M}} \cong p^* \mathcal{N}_{Z_\alpha/M}^C$, T -equivariantly, so the same is true for the subbundles $\mathcal{N}_{\widetilde{Z}_\alpha/\widetilde{M}}^{\pm C}$.

The specific form of the equivariant cohomology classes Ω_{Z_α} is unimportant. The crucial point is that the Todd class and the equivariant Chern character ch_T are functorial along T -equivariant maps, so $\Omega_{\widetilde{Z}_\alpha} = p^* \Omega_{Z_\alpha}$. Thus the RHS of (2.1) is *multiplicative* along the cover $\widetilde{M} \rightarrow M$, meaning

$$\sum_{\alpha \in F} \int_{\widetilde{Z}_\alpha} \Omega_{\widetilde{Z}_\alpha} = m \sum_{\alpha \in F} \int_{Z_\alpha} \Omega_{Z_\alpha}.$$

Consequently the same is true of the LHS, in particular the constant term of the character on the LHS, i.e.

$$\chi^T(M) := \sum_q (-1)^q \dim H^q(M, \mathcal{O})^T = \sum_q (-1)^q \dim H^{n,q}(M, K_M^{-1})^T$$

is multiplicative.

However, since h is a smooth positive metric on K_M^{-1} , the cohomology groups $H^{n,q}(N, K_N^{-1})$ vanish for $q \geq 1$ and $N = M, \widetilde{M}$. This is a direct application of [Dem, Thm. VIII.5.6], since $|j_2|^2$ and $|j_2 \circ p|^2$ are weakly plurisubharmonic exhaustion functions on M and \widetilde{M} respectively. Furthermore $H^0(N, \mathcal{O})^T = H^0(N, \mathcal{O})^{T_c} = \mathbb{C}$ because any JX -invariant holomorphic function f on $N = M, \widetilde{M}$ is constant. This is because f is invariant under the gradient flow of the Hamiltonian $H = \langle \mu, X \rangle$, and since H is Morse–Bott with even indices, a generic gradient flow line approaches the (compact and connected) minimizing submanifold of H , on which f must be constant. Thus $\chi^T(M) = \chi^T(\widetilde{M}) = 1$, contradicting the multiplicativity of χ^T unless $m = 1$. \square

REFERENCES

- [Dem] Jean-Pierre Demailly, *Complex analytic and differential geometry*. [2](#)
- [SZ24] Song Sun and Junsheng Zhang, *Kähler-Ricci shrinkers and Fano fibrations*, arXiv preprint arXiv:2410.09661 (2024). [1](#), [1.1](#), [1](#), [1](#), [2](#)
- [Wu03] Siye Wu, *On the instanton complex of holomorphic Morse theory*, *Comm. Anal. Geom.* **11** (2003), no. 4, 775–807. MR 2015176 [1](#), [2](#)
- [Wyl08] William Wylie, *Complete shrinking Ricci solitons have finite fundamental group*, *Proc. Amer. Math. Soc.* **136** (2008), no. 5, 1803–1806. MR 2373611 [1](#)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA BERKELEY, CA 94720, USA
Email address: esparzac@berkeley.edu