

# A conjecture on monomial realizations and polyhedral realizations for crystal bases

YUKI KANAKUBO\*

## Abstract

Crystal bases are powerful combinatorial tools in the representation theory of quantum groups  $U_q(\mathfrak{g})$  for a symmetrizable Kac-Moody algebras  $\mathfrak{g}$ . The polyhedral realizations are combinatorial descriptions of the crystal base  $B(\infty)$  for Verma modules in terms of the set of integer points of a polyhedral cone, which equals the string cone when  $\mathfrak{g}$  is finite dimensional simple. It is a fundamental and natural problem to find explicit forms of the polyhedral cone. The monomial realization expresses crystal bases  $B(\lambda)$  of integrable highest weight representations as Laurent monomials with double indexed variables. In this paper, we give a conjecture between explicit forms of the polyhedral cones and monomial realizations. We prove the conjecture is true when  $\mathfrak{g}$  is a classical Lie algebra, a rank 2 Kac-Moody algebra or a classical affine Lie algebra.

## 1 Introduction

Crystal bases are introduced for combinatorial study of representations of quantum groups  $U_q(\mathfrak{g})$  for symmetrizable Kac-Moody algebras  $\mathfrak{g}$  over  $\mathbb{C}$  and we can express them by using a bunch of combinatorial objects such as Young tableaux, Young walls, path models and so on [JMMO, KN, Kang, KL, Lit94, Lit95]. In this paper, we focus on two expressions, *monomial realizations* and *polyhedral realizations*. The monomial realization is introduced in [Kas03, Nak], which expresses crystal bases  $B(\lambda)$  of integrable highest weight representations as Laurent monomials with double indexed variables. The following is an example for  $\mathfrak{g} = \mathfrak{sl}_3(\mathbb{C})$ . The crystal base  $B(\Lambda_1)$  is expressed as

$$X_{s,1} \rightarrow \frac{X_{s,2}}{X_{s+1,1}} \rightarrow \frac{1}{X_{s+1,2}} \quad (1.1)$$

and  $B(\Lambda_2)$  is expressed as


$$X_{s,2} \rightarrow \frac{X_{s+1,1}}{X_{s+1,2}} \rightarrow \frac{1}{X_{s+2,1}}. \quad (1.2)$$

The polyhedral realization is invented in [NZ], which expresses elements in crystal bases  $B(\infty)$  of Verma modules (or the negative part  $U_q^-(\mathfrak{g}) \subset U_q(\mathfrak{g})$ ) in terms of the set of integer points of a polyhedral cone. It is defined as the image  $\text{Im}(\Psi_\iota)$  of an embedding  $\Psi_\iota : B(\infty) \rightarrow \mathbb{Z}^\infty$  of crystals associated with an infinite sequence  $\iota = (\cdots, i_3, i_2, i_1)$  of indices from  $I$ . Here,  $I = \{1, 2, \cdots, n\}$  is an index set for simple roots of  $\mathfrak{g}$ . The following is an example of  $\text{Im}(\Psi_\iota) (\cong B(\infty))$  for  $\mathfrak{g} = \mathfrak{sl}_3(\mathbb{C})$  and

---

\*Faculty of Basic Natural Science, Ibaraki University : j\_chi\_sen\_you\_ky@eagle.sophia.ac.jp

$\iota = (\cdots, 2, 1, 2, 1)$ :



Note that the set of elements appearing in the above graph coincides with the set of integer points in a polyhedral cone:

$$\text{Im}(\Psi_\iota) = \{(\cdots, a_3, a_2, a_1) \in \mathbb{Z}^\infty \mid a_1 \geq 0, a_2 \geq a_3 \geq 0, a_k = 0 \ (k > 3)\}. \quad (1.3)$$

It is natural problem to find an explicit form of the polyhedral cone. A lot of researchers are working on this problem. It is known that if  $\mathfrak{g}$  is a finite dimensional simple Lie algebra then the cone coincides with a *string cone* in [Lit98]. When  $\mathfrak{g}$  is a finite dimensional simple Lie algebra and a specific sequence  $\iota = (\cdots, n, \cdots, 2, 1, n, \cdots, 2, 1)$ , explicit forms of cones are given in [Hos05, KS, NZ]. For classical affine Lie algebras  $\mathfrak{g}$  and same  $\iota$ , explicit forms are provided in [Hos13, NZ]. When  $\mathfrak{g}$  is of type  $A_n$  and  $\iota$  is a specific one, it is shown that the inequalities can be obtained from monomial realizations for a system of Demazure crystals [Na]. In our previous paper [KaN20], we combinatorially express inequalities defining the polyhedral cone in terms of column tableaux when  $\mathfrak{g}$  is of type  $A_n, B_n, C_n$  or  $D_n$  and  $\iota$  is adapted (see Definition 2.6). For a classical affine type  $X = A_{n-1}^{(1)}, B_{n-1}^{(1)}, C_{n-1}^{(1)}, D_{n-1}^{(1)}, A_{2n-2}^{(2)}, A_{2n-3}^{(2)}$  or  $D_n^{(2)}$  and an adapted sequence  $\iota$ , we give a combinatorial description of the inequalities in terms of extended Young diagrams and Young walls of type  $X^L$  [Ka23a, Ka23b, Ka24a]. The notation  $X^L$  implies the Langlands dual type for  $X$  (see (2.1)). Note that column tableaux, extended Young diagrams and Young walls are introduced for the combinatorial study of fundamental representations of quantum groups  $U_q(\mathfrak{g})$  [JMMO, KN, Kang]. From these results, one can expect that the inequalities defining the polyhedral cone are expressed by some combinatorial objects deeply related to fundamental representations of quantum groups for general symmetrizable Kac-Moody algebras  $\mathfrak{g}$ . Based on this philosophy, we focus on the monomial realizations as the combinatorial objects.

The main purpose of this paper is to give a conjecture between explicit forms of polyhedral realizations and monomial realizations (Conjecture 4.1). The detail is as follows: We fix an adapted sequence  $\iota$  and consider the union of monomial realizations  $\mathcal{M}_{s,k,\iota}$  with  $s \in \mathbb{Z}_{\geq 1}, k \in I$  for the Langlands dual algebra  $\mathfrak{g}^L$ . Using the tropicalization map (subsection 4.1), one considers the subset of  $\mathbb{Z}^\infty$ :

$$\{\mathbf{a} = (a_{m,j})_{m \in \mathbb{Z}_{\geq 1}, j \in I} \in \mathbb{Z}^\infty \mid \text{Trop}(M)(\mathbf{a}) \geq 0 \text{ for all } M \in \bigcup_{s \in \mathbb{Z}_{\geq 1}, k \in I} \mathcal{M}_{s,k,\iota}\}. \quad (1.4)$$

Here, we identify  $(a_j)_{j \in \mathbb{Z}_{\geq 1}} \in \mathbb{Z}^\infty$  with  $(a_{m,j})_{m \in \mathbb{Z}_{\geq 1}, j \in I}$  by a rule of subsection 2.7. The conjecture claims the above set coincides with  $\text{Im}(\Psi_\iota)$ . For instance, by tropicalizing monomials in (1.1), (1.2), one obtains a system of inequalities

$$a_{s,1} \geq 0, a_{s,2} - a_{s+1,1} \geq 0, -a_{s+1,2} \geq 0, a_{s,2} \geq 0, a_{s+1,1} - a_{s+1,2} \geq 0, -a_{s+2,1} \geq 0 \quad (s \in \mathbb{Z}_{\geq 1}).$$

By  $a_{s,1} \geq 0$ ,  $-a_{s+2,1} \geq 0$  and  $a_{s,2} \geq 0$ ,  $-a_{s+1,2} \geq 0$ , we see that  $a_{m+2,1} = a_{m+1,2} = 0$  for all  $m \geq 1$ . Simplifying other inequalities, the set (1.4) is equal to

$$\{\mathbf{a} = (a_{m,j})_{m \in \mathbb{Z}_{\geq 1}, j \in I} \in \mathbb{Z}^{\infty} \mid a_{1,2} \geq a_{2,1} \geq 0, a_{1,1} \geq 0, a_{m+2,1} = a_{m+1,2} = 0 (m \in \mathbb{Z}_{\geq 1})\},$$

which coincides with  $\text{Im}(\Psi_{\iota})$  in (1.3) as set.

We will prove the conjecture is true when

- $\mathfrak{g}$  is a finite dimensional simple Lie algebra of type  $A_n, B_n, C_n$  or  $D_n$ ,
- $\mathfrak{g}$  is a rank 2 Kac-Moody algebra.
- $\mathfrak{g}$  is an affine Lie algebra of type  $A_{n-1}^{(1)}, B_{n-1}^{(1)}, C_{n-1}^{(1)}, D_{n-1}^{(1)}, A_{2n-2}^{(2)}, A_{2n-3}^{(2)}$  or  $D_n^{(2)}$ .

For the proof, we use our previous expression of inequalities given in [KaN20, Ka23a, Ka24a, Ka24b].

The organization of paper is as follows: In Sect.2, we recall the definition of crystals and polyhedral realizations. In particular, we recall procedures to compute the explicit forms of  $\text{Im}(\Psi_{\iota})$ . In Sect.3, we review monomial realizations of crystal bases. Sect.4 is devoted to give our conjecture and main theorems. In Sect.5, we prove one of theorems, which gives a sufficient condition for the conjecture. We show Theorem 4.3 in Sect.6, 7 and 8 that claims the conjecture is true for specific types.

**Acknowledgements** This work was supported by JSPS KAKENHI Grant Number JP24K22825.

## 2 Crystals and polyhedral realizations for $B(\infty)$

### 2.1 Notation

We set  $I := \{1, 2, \dots, n\}$  with  $n \in \mathbb{Z}_{\geq 1}$ . Let  $\mathfrak{g}$  be a symmetrizable Kac-Moody algebra over  $\mathbb{C}$  with a generalized Cartan matrix  $(a_{i,j})_{i,j \in I}$ , Cartan subalgebra  $\mathfrak{h}$ , weight lattice  $P \subset \mathfrak{h}^*$ , set of simple roots  $\{\alpha_i\}_{i \in I}$  and set of simple coroots  $\{h_i\}_{i \in I}$ , Weyl group  $W$ . Let  $P^+ := \{\lambda \in P \mid \langle h_i, \lambda \rangle \in \mathbb{Z}_{\geq 0}, \text{ for all } i \in I\}$  be the set of dominant integral weights,  $\langle \cdot, \cdot \rangle : \mathfrak{h} \times \mathfrak{h}^* \rightarrow \mathbb{C}$  denote the canonical pairing and  $\Lambda_i \in P^+$  be the  $i$ -th fundamental weight for  $i \in I$ . We obtain  $\langle h_i, \alpha_j \rangle = a_{i,j}$  and  $\langle h_i, \Lambda_j \rangle = \delta_{i,j}$ . Let  $U_q(\mathfrak{g})$  be the quantized universal enveloping algebra of  $\mathfrak{g}$  with indeterminant  $q$ , which is an associative  $\mathbb{C}(q)$ -algebra generated by  $e_i, f_i$  ( $i \in I$ ) and  $q^h$  ( $h \in P^* = \{h \in \mathfrak{h} \mid \langle h, P \rangle \subset \mathbb{Z}\}$ ). It has a subalgebra  $U_q^-(\mathfrak{g})$  generated by  $f_i$  ( $i \in I$ ).

It is known that for  $\lambda \in P^+$ , every integrable highest weight representation  $V(\lambda)$  has a crystal base  $(L(\lambda), B(\lambda))$  and  $U_q^-(\mathfrak{g})$  also has a crystal base  $(L(\infty), B(\infty))$  ([Kas90, Kas91, Lus]). For  $X = A_n, B_n, C_n$  or  $D_n$ , we define the Langlands dual type  $X^L$  as  $X^L = X$  ( $X = A_n, D_n$ ) and  $(B_n)^L = C_n$ ,  $(C_n)^L = B_n$ . We define


$$X^L = X \quad (X = A_{n-1}^{(1)}, D_{n-1}^{(1)}), \quad (C_{n-1}^{(1)})^L = D_n^{(2)}, \quad (D_n^{(2)})^L = C_{n-1}^{(1)}, \quad (A_{2n-3}^{(2)})^L = B_{n-1}^{(1)}, \quad (B_{n-1}^{(1)})^L = A_{2n-3}^{(2)}. \quad (2.1)$$

We also define  $(A_{2n-2}^{(2)})^L = A_{2n-2}^{(2)\dagger}$  and  $(A_{2n-2}^{(2)\dagger})^L = A_{2n-2}^{(2)}$ . Note that  $(X^L)^L = X$ . We often omit the rank and simply write  $A_n, B_n, C_n, D_n$  as  $A, B, C, D$  for simplicity. Let  $\mathfrak{g}^L$  denote the Kac-Moody algebra whose generalized Cartan matrix is the transposed matrix of  $(a_{i,j})_{i,j \in I}$ . For two integers  $m, l$  with  $m \leq l$ , we set  $[m, l] := \{m, m+1, \dots, l-1, l\}$ .


The numbering of Dynkin diagrams and affine Dynkin diagrams are as follows:

$$A_n : \bullet \text{---} \bullet \text{---} \cdots \text{---} \bullet \text{---} \bullet \quad B_n : \bullet \text{---} \bullet \text{---} \cdots \text{---} \bullet \rightleftarrows \bullet$$

1      2                      n-1      n                      1      2                      n-1      n



Replacing our numbering  $1, 2, \dots, n-1, n, n+1$  of  $A_{2n}^{(2)}$  with  $n, n-1, \dots, 1, 0$ , one obtains the numbering in [Kang]. We define the type  $A_{2n-2}^{(2)\dagger}$  by the following diagram whose numbering is same as [Kang]:



## 2.2 Crystals

We briefly review the crystals.

**Definition 2.1.** Let  $B$  be a set and we suppose that there are maps  $\text{wt} : B \rightarrow P$ ,  $\varepsilon_i : B \rightarrow \mathbb{Z} \sqcup \{-\infty\}$ ,  $\varphi_i : B \rightarrow \mathbb{Z} \sqcup \{-\infty\}$ ,  $\tilde{f}_i : B \rightarrow B \sqcup \{0\}$  and  $\tilde{e}_i : B \rightarrow B \sqcup \{0\}$  for  $i \in I$ . Here,  $0$  and  $-\infty$  are additional elements. When the following conditions hold, the set  $B$  together with these maps is called a *crystal*: For  $b, b' \in B$ ,

- (1)  $\text{wt}(\tilde{e}_k b) = \text{wt}(b) + \alpha_k$  if  $\tilde{e}_k(b) \in B$ ,  $\text{wt}(\tilde{f}_k b) = \text{wt}(b) - \alpha_k$  if  $\tilde{f}_k(b) \in B$ ,
- (2)  $\varphi_k(b) = \varepsilon_k(b) + \langle h_k, \text{wt}(b) \rangle$ ,
- (3)  $\varepsilon_k(\tilde{e}_k(b)) = \varepsilon_k(b) - 1$ ,  $\varphi_k(\tilde{e}_k(b)) = \varphi_k(b) + 1$  if  $\tilde{e}_k(b) \in B$ ,
- (4)  $\varepsilon_k(\tilde{f}_k(b)) = \varepsilon_k(b) + 1$ ,  $\varphi_k(\tilde{f}_k(b)) = \varphi_k(b) - 1$  if  $\tilde{f}_k(b) \in B$ ,
- (5)  $\tilde{f}_k(b) = b'$  if and only if  $b = \tilde{e}_k(b')$ ,

(6) if  $\varphi_k(b) = -\infty$  then  $\tilde{e}_k(b) = \tilde{f}_k(b) = 0$ .

It is well known that the sets  $B(\infty)$ ,  $B(\lambda)$  have crystal structures.

**Definition 2.2.** (1) For two crystals  $B_1, B_2$ , a map  $\psi : B_1 \sqcup \{0\} \rightarrow B_2 \sqcup \{0\}$  is said to be a *strict morphism* and denoted by  $\psi : B_1 \rightarrow B_2$  if  $\psi(0) = 0$  and the following conditions hold: For  $k \in I$  and  $b \in B_1$ ,

- if  $\psi(b) \in B_2$  then for  $i \in I$ , it holds

$$\text{wt}(\psi(b)) = \text{wt}(b), \quad \varepsilon_i(\psi(b)) = \varepsilon_i(b), \quad \varphi_i(\psi(b)) = \varphi_i(b),$$

- it holds  $\tilde{e}_i(\psi(b)) = \psi(\tilde{e}_i(b))$  and  $\tilde{f}_i(\psi(b)) = \psi(\tilde{f}_i(b))$  for  $i \in I$ , where we understand  $\tilde{e}_i(0) = \tilde{f}_i(0) = 0$ .

(2) If a strict morphism  $\psi : B_1 \sqcup \{0\} \rightarrow B_2 \sqcup \{0\}$  is injective then  $\psi$  is said to be a *strict embedding* and denoted by  $\psi : B_1 \hookrightarrow B_2$ . If  $\psi$  is bijective then  $\psi$  is said to be an *isomorphism*.

### 2.3 An embedding

Let  $\iota = (\dots, i_r, \dots, i_2, i_1)$  be a sequence of indices from  $I$  such that

$$i_r \neq i_{r+1} \text{ for } r \in \mathbb{Z}_{\geq 1} \text{ and } \#\{r \in \mathbb{Z}_{\geq 1} \mid i_r = k\} = \infty \text{ for all } k \in I. \quad (2.2)$$

We can define a crystal structure on the set

$$\mathbb{Z}^\infty := \{(\dots, a_r, \dots, a_2, a_1) \mid a_r \in \mathbb{Z} \text{ for } r \in \mathbb{Z}_{\geq 1} \text{ and it holds } a_r = 0 \text{ for } r \gg 0\},$$

associated with  $\iota$  (see subsection 2.4 of [NZ]) and denote it by  $\mathbb{Z}_\iota^\infty$ .

**Theorem 2.3.** [Kas93, NZ] *There is the unique strict embedding of crystals*

$$\Psi_\iota : B(\infty) \hookrightarrow \mathbb{Z}_\iota^\infty \quad (2.3)$$

such that the highest weight vector  $u_\infty \in B(\infty)$  is mapped to  $\mathbf{0} := (\dots, 0, \dots, 0, 0) \in \mathbb{Z}_\iota^\infty$ :  $\Psi_\iota(u_\infty) = \mathbf{0}$ .

### 2.4 Nakashima-Zelevinsky's procedure

Following [NZ], let us recall a procedure to compute an explicit form of  $\text{Im}(\Psi_\iota)$ . We set

$$\mathbb{Q}^\infty := \{\mathbf{a} = (\dots, a_r, \dots, a_2, a_1) \mid a_r \in \mathbb{Q} \text{ for } r \in \mathbb{Z}_{\geq 1} \text{ and it holds } a_r = 0 \text{ for } r \gg 0\}.$$

For  $r \in \mathbb{Z}_{\geq 1}$ , one defines  $x_r \in (\mathbb{Q}^\infty)^*$  as  $x_r(\dots, a_r, \dots, a_2, a_1) = a_r$ . We understand  $x_r := 0$  when  $r \in \mathbb{Z}_{< 1}$ . For  $r \in \mathbb{Z}_{\geq 1}$ , we set

$$r^{(+)} := \min\{l \in \mathbb{Z}_{\geq 1} \mid l > r \text{ and } i_r = i_l\}, \quad r^{(-)} := \max\{l \in \mathbb{Z}_{\geq 1} \mid l < r \text{ and } i_r = i_l\} \cup \{0\},$$

and

$$\beta_r := x_r + \sum_{r < j < r^{(+)}} \langle h_{i_r}, \alpha_{i_j} \rangle x_j + x_{r^{(+)}} \in (\mathbb{Q}^\infty)^*, \quad \beta_0 := 0 \in (\mathbb{Q}^\infty)^*. \quad (2.4)$$

We define a piecewise-linear operator  $S_r = S_{r, \iota}$  on  $(\mathbb{Q}^\infty)^*$  by

$$S_r(\varphi) := \begin{cases} \varphi - \varphi_r \beta_r & \text{if } \varphi_r > 0, \\ \varphi - \varphi_r \beta_{r^{(-)}} & \text{if } \varphi_r \leq 0. \end{cases} \quad (2.5)$$

We often simply write the map  $S_r(\varphi)$  as  $S_r\varphi$ . Let us define

$$\Xi_\iota := \{S_{j_l} \cdots S_{j_2} S_{j_1} x_{j_0} \mid l \in \mathbb{Z}_{\geq 0}, j_0, j_1, \dots, j_l \geq 1\}, \quad (2.6)$$

$$\Sigma_\iota := \{\mathbf{x} \in \mathbb{Z}^\infty \subset \mathbb{Q}^\infty \mid \varphi(\mathbf{x}) \geq 0 \text{ for any } \varphi \in \Xi_\iota\}. \quad (2.7)$$

We consider the following *positivity condition* on  $\iota$ :

$$\text{for any } \varphi = \sum_{k \in \mathbb{Z}_{\geq 1}} \varphi_k x_k \in \Xi_\iota, \text{ if } k^{(-)} = 0 \text{ then } \varphi_k \geq 0. \quad (2.8)$$

**Theorem 2.4.** [NZ] *For a sequence  $\iota$  of indices satisfying (2.2) and (2.8), we have  $\text{Im}(\Psi_\iota) = \Sigma_\iota$ .*

## 2.5 Modified procedure

Modifying the procedure in the previous subsection, we get another procedure to compute  $\text{Im}(\Psi_\iota)$ . Let  $\iota = (\dots, i_r, \dots, i_2, i_1)$  be a sequence of indices from  $I$  satisfying (2.2). For each  $r \in \mathbb{Z}_{\geq 1}$ , a map  $S'_r = S'_{r,\iota} : (\mathbb{Q}^\infty)^* \rightarrow (\mathbb{Q}^\infty)^*$  is defined as follows: For  $\varphi = \sum_{l \in \mathbb{Z}_{\geq 1}} c_l x_l \in (\mathbb{Q}^\infty)^*$ ,

$$S'_r(\varphi) := \begin{cases} \varphi - \beta_r & \text{if } c_r > 0, \\ \varphi + \beta_{r^{(-)}} & \text{if } c_r < 0, \\ \varphi & \text{if } c_r = 0. \end{cases} \quad (2.9)$$

Here,  $\beta_r$  is defined in (2.4). When  $c_r > 0$ , we call  $S'_r$  a *positive action*. Just as in the previous subsection, one defines

$$\begin{aligned} \Xi'_\iota &:= \{S'_{j_l} \cdots S'_{j_2} S'_{j_1} x_{j_0} \mid l \in \mathbb{Z}_{\geq 0}, j_0, j_1, \dots, j_l \in \mathbb{Z}_{\geq 1}\}, \\ \Sigma'_\iota &:= \{\mathbf{a} \in \mathbb{Z}^\infty \subset \mathbb{Q}^\infty \mid \varphi(\mathbf{a}) \geq 0 \text{ for any } \varphi \in \Xi'_\iota\}. \end{aligned} \quad (2.10)$$

We impose the following  $\Xi'$ -positivity condition on  $\iota$ :

$$\text{for any } \varphi = \sum_{l \in \mathbb{Z}_{\geq 1}} c_l x_l \in \Xi'_\iota, \text{ if } l^{(-)} = 0 \text{ then } c_l \geq 0. \quad (2.11)$$

**Theorem 2.5.** [Ka23a] *Let  $\iota = (\dots, i_2, i_1)$  be a sequence satisfying (2.2) and (2.11). Then we get  $\text{Im}(\Psi_\iota) = \Sigma'_\iota$ .*

## 2.6 Adapted sequences

In this paper, we will consider only *adapted* sequences.

**Definition 2.6.** [KaN20] Let  $A = (a_{i,j})_{i,j \in I}$  be the symmetrizable generalized Cartan matrix of  $\mathfrak{g}$  and  $\iota = (\dots, i_3, i_2, i_1)$  be a sequence that satisfies (2.2). If the following condition holds then the sequence  $\iota$  is said to be *adapted* to  $A$ : For each  $i, j \in I$  such that  $a_{i,j} < 0$ , the subsequence of  $\iota$  consisting of all  $i, j$  is either

$$(\dots, i, j, i, j, i, j, i, j) \text{ or } (\dots, j, i, j, i, j, i, j, i).$$

When the matrix  $A$  is fixed, we shortly say  $\iota$  is *adapted*.

For an adapted sequence  $\iota = (\dots, i_3, i_2, i_1)$ , we define a set of integers  $(p_{i,j})_{i,j \in I; a_{i,j} < 0}$  by

$$p_{i,j} = \begin{cases} 1 & \text{if the subsequence of } \iota \text{ consisting of } i, j \text{ is } (\dots, j, i, j, i, j, i), \\ 0 & \text{if the subsequence of } \iota \text{ consisting of } i, j \text{ is } (\dots, i, j, i, j, i, j). \end{cases} \quad (2.12)$$

We can verify that if  $a_{i,j} < 0$  then

$$p_{i,j} + p_{j,i} = 1. \quad (2.13)$$

## 2.7 An identification of single indices with double indices

For a fixed sequence  $\iota = (\cdots, i_2, i_1)$  satisfying (2.2), we identify the set of single indices  $\mathbb{Z}_{\geq 1}$  with the set of double indices  $\mathbb{Z}_{\geq 1} \times I$  as follows: We identify each single index  $r \in \mathbb{Z}_{\geq 1}$  with a double index  $(s, k) \in \mathbb{Z}_{\geq 1} \times I$  when  $i_r = k$  and  $k$  appears  $s$  times in  $i_r, i_{r-1}, \cdots, i_1$ . For example, when  $\iota = (\cdots, 2, 1, 3, 2, 1, 3, 2, 1, 3)$ , single indices  $\cdots, 6, 5, 4, 3, 2, 1$  are identified with double indices

$$\cdots, (2, 2), (2, 1), (2, 3), (1, 2), (1, 1), (1, 3).$$

The notation  $x_r, \beta_r, S_r$  and  $S'_r$  in Sect.2 are also written as

$$x_r = x_{s,k}, \quad \beta_r = \beta_{s,k}, \quad S_r = S_{s,k}, \quad \text{and} \quad S'_r = S'_{s,k}.$$

When  $(s, k) \notin \mathbb{Z}_{\geq 1} \times I$ , we understand  $x_{s,k} := 0$ . By this identification and the ordinary order on  $\mathbb{Z}_{\geq 1}$  ( $1 < 2 < 3 < 4 < 5 < 6 < \cdots$ ), we can naturally define an order on  $\mathbb{Z}_{\geq 1} \times I$ . For  $\iota = (\cdots, 2, 1, 3, 2, 1, 3, 2, 1, 3)$ , the order is  $\cdots > (2, 2) > (2, 1) > (2, 3) > (1, 2) > (1, 1) > (1, 3)$ . Using the notation in (2.12), if  $\iota$  is adapted then  $\beta_{s,k}$  is in the form

$$\beta_{s,k} = x_{s,k} + x_{s+1,k} + \sum_{j \in I; a_{k,j} < 0} a_{k,j} x_{s+p_{j,k}, j}. \quad (2.14)$$

One can verify that  $\{\beta_{t,i}\}_{t \in \mathbb{Z}_{\geq 1}, i \in I}$  is  $\mathbb{Z}$ -linearly independent.

## 3 Monomial realizations

Let us review the monomial realizations for crystal bases of highest weight representations. We consider the set of Laurent monomials as follows:

$$\mathcal{Y} := \left\{ X = \prod_{s \in \mathbb{Z}, i \in I} X_{s,i}^{\zeta_{s,i}} \mid \zeta_{s,i} \in \mathbb{Z}, \zeta_{s,i} = 0 \text{ except for finitely many } (s, i) \right\}. \quad (3.1)$$

For  $X = \prod_{s \in \mathbb{Z}, i \in I} X_{s,i}^{\zeta_{s,i}} \in \mathcal{Y}$ , one sets  $\text{wt}(X) := \sum_{s,i} \zeta_{s,i} \Lambda_i$  and

$$\varphi_i(X) := \max \left\{ \sum_{k \leq s} \zeta_{k,i} \mid s \in \mathbb{Z} \right\}, \quad \varepsilon_i(X) := \varphi_i(X) - \text{wt}(X)(h_i) = \max \left\{ - \sum_{k > s} \zeta_{k,i} \mid s \in \mathbb{Z} \right\}.$$

We fix an adapted sequence  $\iota = (\cdots, i_3, i_2, i_1)$  and take  $p_{i,j}$  as in (2.12) and put

$$A_{s,k} := X_{s,k} X_{s+1,k} \prod_{j \in I; a_{j,k} < 0} X_{s+p_{j,k}, j}^{a_{j,k}} \quad (s \in \mathbb{Z}, k \in I). \quad (3.2)$$

For  $i \in I$ , let us define actions of Kashiwara operators as follows:

$$\tilde{f}_i X := \begin{cases} A_{n_{f_i}, i}^{-1} X & \text{if } \varphi_i(X) > 0, \\ 0 & \text{if } \varphi_i(X) = 0, \end{cases} \quad \tilde{e}_i X := \begin{cases} A_{n_{e_i}, i} X & \text{if } \varepsilon_i(X) > 0, \\ 0 & \text{if } \varepsilon_i(X) = 0, \end{cases}$$

where we set

$$n_{f_i} := \min \left\{ r \in \mathbb{Z} \mid \varphi_i(X) = \sum_{k \leq r} \zeta_{k,i} \right\}, \quad n_{e_i} := \max \left\{ r \in \mathbb{Z} \mid \varepsilon_i(X) = \sum_{k \leq r} \zeta_{k,i} \right\}.$$

**Theorem 3.1.** [Kas03, Nak]

(i) The set  $\mathcal{Y}$  together with the above maps  $\text{wt}$ ,  $\varepsilon_i$ ,  $\varphi_i$  and  $\tilde{e}_i$ ,  $\tilde{f}_i$  ( $i \in I$ ) is a crystal.

(ii) Taking  $X \in \mathcal{Y}$  as  $\tilde{e}_i(X) = 0$  for all  $i \in I$ , the set

$$\{\tilde{f}_{j_m} \cdots \tilde{f}_{j_1} X \mid m \in \mathbb{Z}_{\geq 0}, j_1, \dots, j_m \in I\} \setminus \{0\}$$

is isomorphic to  $B(\text{wt}(X))$ . We denote  $\{\tilde{f}_{j_m} \cdots \tilde{f}_{j_1} X_{s,k} \mid m \in \mathbb{Z}_{\geq 0}, j_1, \dots, j_m \in I\} \setminus \{0\}$  by  $\mathcal{M}_{s,k,\iota}$  for  $s \in \mathbb{Z}$  and  $k \in I$ , which is isomorphic to  $B(\Lambda_k)$ .

The following is straightforward from the definitions as above:

**Lemma 3.2.** Let  $X = \prod_{s \in \mathbb{Z}, i \in I} X_{s,i}^{\zeta_{s,i}} \in \mathcal{Y}$  and  $j \in I$ .

- (1) If  $\tilde{f}_j X \neq 0$  and  $\tilde{f}_j X = A_{s,j}^{-1} X$  with some  $s \in \mathbb{Z}$  then  $\zeta_{s,j} > 0$ .
- (2) If  $\tilde{e}_j X \neq 0$  and  $\tilde{e}_j X = A_{s,j} X$  with some  $s \in \mathbb{Z}$  then  $\zeta_{s+1,j} < 0$ .
- (3) If  $\zeta_{s,j} < 0$  with some  $s \in \mathbb{Z}$  and  $\zeta_{s',j} = 0$  for all  $s' > s$  then  $\varepsilon_j(X) > 0$  so that  $\tilde{e}_j X \neq 0$ .


**Example 3.3.** Let  $\mathfrak{g}$  be of type  $C_2$  and  $\iota = (\dots, 2, 1, 2, 1, 2, 1)$ . Then  $\mathcal{M}_{s,1,\iota}$  is

$$X_{s,1} \xrightarrow{1} \frac{X_{s,2}}{X_{s+1,1}} \xrightarrow{2} \frac{X_{s+1,1}}{X_{s+1,2}} \xrightarrow{1} \frac{1}{X_{s+2,1}}$$

and  $\mathcal{M}_{s,2,\iota}$  is


$$X_{s,2} \xrightarrow{2} \frac{X_{s+1,1}^2}{X_{s+1,2}} \xrightarrow{1} \frac{X_{s+1,1}}{X_{s+2,1}} \xrightarrow{1} \frac{X_{s+1,2}}{X_{s+2,1}^2} \xrightarrow{2} \frac{1}{X_{s+2,2}}.$$

**Example 3.4.** Next, we set  $\mathfrak{g}$  is of type  $A_1^{(1)}$  and  $\iota = (\dots, 2, 1, 2, 1)$ . Then the partial crystal graph of  $\mathcal{M}_{s,1,\iota}$  is as follows:





Similarly, the partial crystal graph of  $\mathcal{M}_{s,2,\iota}$  is as follows:



## 4 Main results

We fix an adapted sequence  $\iota = (\cdots, i_2, i_1)$ .

### 4.1 Tropicalizations

We set

$$\mathcal{Y}' := \left\{ X = \prod_{s \in \mathbb{Z}_{\geq 1}, k \in I} X_{s,k}^{\zeta_{s,k}} \mid \zeta_{s,k} \in \mathbb{Z}, \zeta_{s,k} = 0 \text{ except for finitely many } (s,k) \right\}$$

as a subset of  $\mathcal{Y}$  in (3.1) and define

$$\mathcal{H} := \left\{ \sum_{s \in \mathbb{Z}_{\geq 1}, k \in I} \zeta_{s,k} x_{s,k} \mid \zeta_{s,k} \in \mathbb{Z}, \zeta_{s,k} = 0 \text{ except for finitely many } (s,k) \right\}. \quad (4.1)$$

Here,  $x_{s,k} \in (\mathbb{Q}^\infty)$  is the notation as in subsection 2.7. We define a bijection

$$Trop : \mathcal{Y}' \rightarrow \mathcal{H}$$

as

$$Trop \left( \prod_{s \in \mathbb{Z}_{\geq 1}, k \in I} X_{s,k}^{\zeta_{s,k}} \right) = \sum_{s \in \mathbb{Z}_{\geq 1}, k \in I} \zeta_{s,k} x_{s,k}.$$

We define  $DeTrop : \mathcal{H} \rightarrow \mathcal{Y}'$  as its inverse map:

$$DeTrop \left( \sum_{s \in \mathbb{Z}_{\geq 1}, k \in I} \zeta_{s,k} x_{s,k} \right) = \prod_{s \in \mathbb{Z}_{\geq 1}, k \in I} X_{s,k}^{\zeta_{s,k}}.$$

Note that  $Trop(A_{s,k})$  for  $\mathfrak{g}$  coincides with  $\beta_{s,k}$  for  $\mathfrak{g}^L$  by (2.14) and (3.2) when  $s \geq 1$ .

## 4.2 A conjecture and theorems

**Conjecture 4.1.** *Let  $\iota$  be an adapted sequence and  $\Psi_\iota$  be the map in Theorem 2.3 for  $\mathfrak{g}$ . Let  $\mathcal{M}_{s,k,\iota}$  be the set of monomials in Theorem 3.1 (ii) for  $\mathfrak{g}^L$ . Then*

$$\text{Im}(\Psi_\iota) = \left\{ \mathbf{a} \in \mathbb{Z}^\infty \mid \varphi(\mathbf{a}) \geq 0 \text{ for all } \varphi \in \bigcup_{s \in \mathbb{Z}_{\geq 1}, k \in I} \text{Trop}(\mathcal{M}_{s,k,\iota}) \right\}$$

Using maps in (2.9), we set

$$\Xi'_{s,k,\iota} := \{ S'_{j_1} \cdots S'_{j_m} x_{s,k} \mid m \in \mathbb{Z}_{\geq 0}, j_1, \dots, j_m \in \mathbb{Z}_{\geq 1} \}$$

for  $s \in \mathbb{Z}_{\geq 1}, k \in I$ . Let  $\Xi'^+_{s,k,\iota}$  be the set consisting of  $S'_{j_1} \cdots S'_{j_m} x_{s,k}$  such that  $S'_j$  acts by positive actions ( $j = j_1, \dots, j_m$ ). Clearly,  $\Xi'^+_{s,k,\iota} \subset \Xi'_{s,k,\iota}$  holds.

We consider the condition

$$\Xi'^+_{s,k,\iota} = \Xi'_{s,k,\iota} \quad \text{for all } s \in \mathbb{Z}_{\geq 1}, k \in I. \quad (4.2)$$

Note that  $\Xi'$ -positivity condition follows from the assumption (4.2).

**Theorem 4.2.** *If the condition (4.2) holds then Conjecture 4.1 is true.*

**Theorem 4.3.** (1) *When  $\mathfrak{g}$  is a finite dimensional simple Lie algebra of type  $A_n, B_n, C_n$  or  $D_n$ , the Conjecture 4.1 is true.*

(2) *When  $\mathfrak{g}$  is a Kac-Moody algebra of rank 2, the Conjecture 4.1 is true.*

(3) *When  $\mathfrak{g}$  is a classical affine Lie algebra of type  $A_{n-1}^{(1)}, B_{n-1}^{(1)}, C_{n-1}^{(1)}, D_{n-1}^{(1)}, A_{2n-2}^{(2)}, A_{2n-3}^{(2)}$  or  $D_n^{(2)}$ , the Conjecture 4.1 is true.*

**Example 4.4.** We consider the setting as in Example 3.3. By the map Trop, we get the following homomorphisms from  $\mathcal{M}_{s,k,\iota}$  ( $k = 1, 2$ ):

$$x_{s,1}, x_{s,2} - x_{s+1,1}, x_{s+1,1} - x_{s+1,2}, -x_{s+2,1}, \quad x_{s,2}, 2x_{s+1,1} - x_{s+1,2}, x_{s+1,1} - x_{s+2,1}, x_{s+1,2} - 2x_{s+2,1}, -x_{s+2,2}.$$

According to Theorem 4.3 (1), an explicit form of  $\text{Im}(\Psi_\iota)$  of type  $B_2 (= (C_2)^L)$  is obtained from these homomorphisms. By  $a_{s,1} \geq 0, -a_{s+2,1} \geq 0$  for all  $s \in \mathbb{Z}_{\geq 1}$ , we get  $a_{m,1} = 0$  when  $m \geq 3$  and by  $a_{s,2} \geq 0, -a_{s+2,2} \geq 0$  for all  $s \in \mathbb{Z}_{\geq 1}$ , we also get  $a_{m,2} = 0$  when  $m \geq 3$ . Simplifying the inequalities, one obtains

$$\text{Im}(\Psi_\iota) = \{ \mathbf{a} = (a_{m,j}) \in \mathbb{Z}^\infty \mid a_{1,2} \geq a_{2,1} \geq a_{2,2} \geq 0, a_{1,1} \geq 0, a_{m,1} = a_{m,2} = 0 (m \geq 3) \}.$$

**Example 4.5.** We consider the same setting as in Example 3.4, that is,  $\mathfrak{g}$  is of type  $A_1^{(1)}$  and  $\iota = (\dots, 2, 1, 2, 1)$ . Using Theorem 4.3 (2), one obtains several inequalities defining  $\text{Im}(\Psi_\iota)$  of type  $A_1^{(1)} (= (A_1^{(1)})^L)$  from  $\mathcal{M}_{s,k,\iota}$  ( $k = 1, 2$ ):

$$\text{Im}(\Psi_\iota) = \left\{ \mathbf{a} = (a_{m,j}) \in \mathbb{Z}^\infty \mid \begin{array}{l} a_{s,1} \geq 0, 2a_{s,2} - a_{s+1,1} \geq 0, a_{s,2} + a_{s+1,1} - a_{s+1,2} \geq 0, 3a_{s+1,1} - 2a_{s+1,2} \geq 0, \\ a_{s,2} + a_{s+1,2} - a_{s+2,1} \geq 0, 2a_{s+1,1} - a_{s+2,1} \geq 0, a_{s,2} + a_{s+2,1} - a_{s+2,2} \geq 0, \\ a_{s+1,1} + 2a_{s+1,2} - 2a_{s+2,1} \geq 0, 2a_{s+1,1} + a_{s+2,1} - a_{s+1,2} - a_{s+2,2} \geq 0, \\ a_{s,2} + a_{s+2,2} - a_{s+3,1} \geq 0, \dots \\ a_{s,2} \geq 0, 2a_{s+1,1} - a_{s+1,2} \geq 0, a_{s+1,1} + a_{s+1,2} - a_{s+2,1} \geq 0, 3a_{s+1,2} - 2a_{s+2,1} \geq 0, \\ a_{s+1,1} + a_{s+2,1} - a_{s+2,2} \geq 0, 2a_{s+1,2} - a_{s+2,2} \geq 0, a_{s+1,1} + a_{s+2,2} - a_{s+3,1} \geq 0, \\ a_{s+1,2} + 2a_{s+2,1} - 2a_{s+2,2} \geq 0, 2a_{s+1,2} + a_{s+2,2} - a_{s+2,1} - a_{s+3,1} \geq 0, \\ a_{s+1,1} + a_{s+3,1} - a_{s+3,2} \geq 0 \end{array} \right\}.$$

## 5 Proof of Theorem 4.2

By the assumption (4.2), we get  $\Xi'_{s,k,\iota} = \Xi'_{s,k,\iota}$ . For any  $\varphi = S'_{j_1} \cdots S'_{j_r} x_{s,k} \in \Xi'_{s,k,\iota}$  with positive actions  $S'_{j_1}, \dots, S'_{j_r}$ , we can define

$$|\varphi| = r. \quad (5.1)$$

This is well-defined by the linear independence of  $\{\beta_{t,i}\}_{t \in \mathbb{Z}_{\geq 1}, i \in I}$ . If we can prove  $\Xi'_{s,k,\iota} = \text{Trop}(\mathcal{M}_{s,k,\iota})$  then our claim follows from Theorem 2.5. The inclusion  $\Xi'_{s,k,\iota} \supset \text{Trop}(\mathcal{M}_{s,k,\iota})$  is clear from Lemma 3.2 (1),(2) so that we show  $\Xi'_{s,k,\iota} \subset \text{Trop}(\mathcal{M}_{s,k,\iota})$ . For any  $\varphi \in \Xi'_{s,k,\iota}$ , let us show  $\varphi \in \text{Trop}(\mathcal{M}_{s,k,\iota})$  via induction on  $|\varphi|$ . When  $|\varphi| = 0$  then  $\varphi = x_{s,k}$  so that the claim is clear so we assume  $|\varphi| > 0$ . Note that by  $\varphi \in \Xi'_{s,k,\iota} = \Xi'_{s,k,\iota}$ , one can write  $\varphi$  as

$$\varphi = x_{s,k} - \sum_{(t,i) \in \mathbb{Z}_{\geq s} \times I} c_{t,i} \beta_{t,i}$$

with non-negative coefficients  $c_{t,i} \in \mathbb{Z}_{\geq 0}$  and  $c_{t,i} = 0$  except for finitely many  $(t,i)$ . We take

$$(r,j) := \max\{(t,i) \in \mathbb{Z}_{\geq s} \times I | c_{t,i} > 0\}$$

in the order defined in the subsection 2.7. It follows from the definition of  $\beta_{t,i}$  and adaptedness of  $\iota$  that the coefficient of  $x_{r+1,j}$  in  $\varphi$  is negative. By the maximality of  $(r,j)$  and adaptedness of  $\iota$ , we see that the coefficients of  $x_{l,j}$  ( $l \geq r+2$ ) in  $\varphi$  are equal to 0. Thus, writing  $\varphi = \text{Trop}(M)$  with  $M = \text{DeTrop}(\varphi)$ , we get  $\tilde{e}_j M \neq 0$  from Lemma 3.2 (3) and

$$\text{Trop}(\tilde{e}_j M) = \text{Trop}(M) + \beta_{m,j}$$

with some  $m \in \mathbb{Z}_{\geq 0}$ . Note that since the coefficient of  $x_{m+1,j}$  in  $\varphi$  is negative by Lemma 3.2 (2) so it holds  $\text{Trop}(\tilde{e}_j M) = S'_{m+1,j} \varphi \in \Xi'_{s,k,\iota}$ . It follows by (4.2) that  $|\text{Trop}(\tilde{e}_j M)| < |\varphi|$ . Thus, the induction assumption implies  $\text{Trop}(\tilde{e}_j M) \in \text{Trop}(\mathcal{M}_{s,k,\iota})$ . By  $\tilde{e}_j M \in \mathcal{M}_{s,k,\iota}$ , we have  $M = \tilde{f}_j \tilde{e}_j M \in \mathcal{M}_{s,k,\iota}$ . Thus, we get  $\varphi = \text{Trop}(M) \in \text{Trop}(\mathcal{M}_{s,k,\iota})$ .  $\square$

## 6 Finite case

Let  $\mathfrak{g}$  be of type  $X = A_n, B_n, C_n$  or  $D_n$  and  $\iota$  be adapted.

### 6.1 Boxes and tableaux

For  $k$  ( $2 \leq k \leq n$ ), we put

$$P^X(k) := \begin{cases} p_{2,1} + p_{3,2} + \cdots + p_{n-2,n-3} + p_{n,n-2} & \text{if } k = n \text{ and } X = D_n, \\ p_{2,1} + p_{3,2} + p_{4,3} + \cdots + p_{k,k-1} & \text{if otherwise} \end{cases}$$

and  $P^X(0) = P^X(1) = P^X(n+1) = 0$ . We often write  $P^X(k) = P(k)$  when  $X$  is fixed. By  $p_{i,j} \in \{0,1\}$ , it holds for  $k, l \in I$  such that  $k \geq l$ ,

$$P(k) \geq P(l), \quad (6.1)$$

$$(k-l) + P(l) \geq P(k), \quad (6.2)$$

except for the case  $X = D_n, k = n$  and  $l = n-1$ .

We define the following (partial) ordered sets  $J_A, J_B, J_C$  and  $J_D$ :

(i)  $J_A := \{1, 2, \dots, n, n+1\}$  with the order  $1 < 2 < \dots < n < n+1$ .

(ii)  $J_B = J_C := \{1, 2, \dots, n, \bar{n}, \dots, \bar{2}, \bar{1}\}$  with the order

$$1 < 2 < \dots < n < \bar{n} < \dots < \bar{2} < \bar{1}.$$

(iii)  $J_D := \{1, 2, \dots, n, \bar{n}, \dots, \bar{2}, \bar{1}\}$  with the partial order

$$1 < 2 < \dots < n-1 < \frac{n}{\bar{n}} < \overline{n-1} < \dots < \bar{2} < \bar{1}.$$

For  $j \in \{1, 2, \dots, n\}$ , we set  $|j| = |\bar{j}| = j$ .

Next, we define the boxes and tableaux as homomorphisms. Note that the notation is slightly different from [KaN20].

**Definition 6.1.** [KaN20]

(i) For  $1 \leq j \leq n+1$  and  $s \in \mathbb{Z}$ , we define

$$\boxed{j}_s^A := x_{s+PA(j),j} - x_{s+PA(j-1)+1,j-1} \in (\mathbb{Q}^\infty)^*.$$

(ii) For  $1 \leq j \leq n$  and  $s \in \mathbb{Z}$ , we define

$$\boxed{j}_s^C := x_{s+PC(j),j} - x_{s+PC(j-1)+1,j-1} \in (\mathbb{Q}^\infty)^*,$$

$$\boxed{\bar{j}}_s^C := x_{s+PC(j-1)+n-j+1,j-1} - x_{s+PC(j)+n-j+1,j} \in (\mathbb{Q}^\infty)^*.$$

(iii) For  $1 \leq j \leq n-1$  and  $s \in \mathbb{Z}$ , we set

$$\boxed{j}_s^B := x_{s+PB(j),j} - x_{s+PB(j-1)+1,j-1}, \quad \boxed{n}_s^B := 2x_{s+PB(n),n} - x_{s+PB(n-1)+1,n-1} \in (\mathbb{Q}^\infty)^*,$$

$$\boxed{0}_s^B := x_{s+PB(n),n} - x_{s+PB(n)+1,n} \in (\mathbb{Q}^\infty)^*,$$

$$\boxed{\bar{n}}_s^B := x_{s+PB(n-1)+1,n-1} - 2x_{s+PB(n)+1,n}, \quad \boxed{\bar{j}}_s^B := x_{s+PB(j-1)+n-j+1,j-1} - x_{s+PB(j)+n-j+1,j} \in (\mathbb{Q}^\infty)^*,$$

$$\boxed{\overline{n+1}}_s^B := x_{s+PB(n),n} \in (\mathbb{Q}^\infty)^*.$$

(iv) For  $s \in \mathbb{Z}$ , we set

$$\boxed{j}_s^D := x_{s+PD(j),j} - x_{s+PD(j-1)+1,j-1} \in (\mathbb{Q}^\infty)^*, \quad (1 \leq j \leq n-2, j = n),$$

$$\boxed{n-1}_s^D := x_{s+PD(n-1),n-1} + x_{s+PD(n),n} - x_{s+PD(n-2)+1,n-2} \in (\mathbb{Q}^\infty)^*,$$

$$\boxed{\bar{n}}_s^D := x_{s+PD(n-1),n-1} - x_{s+PD(n)+1,n} \in (\mathbb{Q}^\infty)^*,$$

$$\boxed{\overline{n-1}}_s^D := x_{s+PD(n-2)+1,n-2} - x_{s+PD(n-1)+1,n-1} - x_{s+PD(n)+1,n} \in (\mathbb{Q}^\infty)^*,$$

$$\boxed{\bar{j}}_s^D := x_{s+PD(j-1)+n-j,j-1} - x_{s+PD(j)+n-j,j} \in (\mathbb{Q}^\infty)^*, \quad (1 \leq j \leq n-2),$$

$$\boxed{\overline{n+1}}_s^D := x_{s+PD(n),n} \in (\mathbb{Q}^\infty)^*.$$

(v) For  $X = A, B, C$  or  $D$ , we set

$$\begin{array}{c} \boxed{j_1} \\ \boxed{j_2} \\ \vdots \\ \boxed{j_k} \end{array}^X := \boxed{j_k}_s^X + \boxed{j_{k-1}}_{s+1}^X + \cdots + \boxed{j_2}_{s+k-2}^X + \boxed{j_1}_{s+k-1}^X \in (\mathbb{Q}^\infty)^*.$$

**Lemma 6.2.** [KaN20]

(i) When  $\mathfrak{g}$  is of type  $A_n$ , it holds

$$\boxed{j+1}_s^A = \boxed{j}_s^A - \beta_{s+P^A(j),j} \quad (1 \leq j \leq n, s \geq 1 - P^A(j)). \quad (6.3)$$

(ii) When  $\mathfrak{g}$  is of type  $B_n$ , it holds

$$\boxed{j+1}_s^C = \boxed{j}_s^C - \beta_{s+P^C(j),j} \quad (1 \leq j \leq n-1, s \geq 1 - P^C(j)), \quad (6.4)$$

$$\overline{n}_s^C = \overline{n}_s^C - \beta_{s+P^C(n),n} \quad (s \geq 1 - P^C(n)), \quad (6.5)$$

$$\overline{j-1}_s^C = \overline{j}_s^C - \beta_{s+P^C(j-1)+n-j+1,j-1} \quad (2 \leq j \leq n, s \geq j - P^C(j-1) - n). \quad (6.6)$$

(iii) When  $\mathfrak{g}$  is of type  $C_n$ , it holds

$$\boxed{j+1}_s^B = \boxed{j}_s^B - \beta_{s+P^B(j),j} \quad (1 \leq j \leq n-1, s \geq 1 - P^B(j)), \quad (6.7)$$

$$\boxed{0}_s^B = \overline{n}_s^B - \beta_{s+P^B(n),n} \quad (s \geq 1 - P^B(n)), \quad (6.8)$$

$$\overline{n}_s^B = \boxed{0}_s^B - \beta_{s+P^B(n),n} \quad (s \geq 1 - P^B(n)), \quad (6.9)$$

$$\overline{j-1}_s^B = \overline{j}_s^B - \beta_{s+P^B(j-1)+n-j+1,j-1} \quad (2 \leq j \leq n, s \geq j - P^B(j-1) - n) \quad (6.10)$$

$$\overline{n+1}_{l+1}^B + \overline{n}_l^B = \overline{n+1}_l^B - \beta_{l+P^B(n),n} \quad (l \geq 1 - P^B(n)). \quad (6.11)$$

(iv) When  $\mathfrak{g}$  is of type  $D_n$ , it holds

$$\boxed{j+1}_s^D = \boxed{j}_s^D - \beta_{s+P^D(j),j} \quad (1 \leq j \leq n-1, s \geq 1 - P^D(j)), \quad (6.12)$$

$$\overline{n}_s^D = \overline{n-1}_s^D - \beta_{s+P^D(n),n} \quad (s \geq 1 - P^D(n)), \quad (6.13)$$

$$\overline{n-1}_s^D = \overline{n}_s^D - \beta_{s+P^D(n),n} \quad (s \geq 1 - P^D(n)), \quad (6.14)$$

$$\overline{j-1}_s^D = \overline{j}_s^D - \beta_{s+P^D(j-1)+n-j,j-1} \quad (2 \leq j \leq n, s \geq 1 + j - P^D(j-1) - n), \quad (6.15)$$

$$\overline{n+1}_{l+2}^D + \overline{n}_{l+1}^D + \overline{n-1}_l^D = \overline{n+1}_l^D - \beta_{l+P^D(n),n} \quad (l \geq 1 - P^D(n)). \quad (6.16)$$

**Definition 6.3.** (1) For  $X = A_n$  or  $C_n$  and  $k \in I$ , we define

$$\text{Tab}_{k,l}^X := \left\{ \begin{array}{c} \boxed{j_1} \\ \boxed{j_2} \\ \vdots \\ \boxed{j_k} \end{array}^X \mid j_i \in J_X, s \geq 1 - P^X(k), j_1 < \cdots < j_k \right\}.$$

(2) For  $k \in I$  with  $k < n$ , we define

$$\text{Tab}_{k,t}^{B_n} := \left\{ \begin{array}{c} \boxed{j_1} \\ \boxed{j_2} \\ \vdots \\ \boxed{j_k} \end{array} \middle| j_i \in J_B, s \geq 1 - P^B(k), j_1 \leq \cdots \leq j_k, \text{ if } j_i = j_{i+1} \text{ then } j_i = 0 \right\},$$

$$\text{Tab}_{n,t}^{B_n} := \left\{ \begin{array}{c} \boxed{n+1} \\ \boxed{j_1} \\ \vdots \\ \boxed{j_r} \end{array} \middle| r \in [0, n], j_1, \dots, j_r \in J_B, \bar{n} \leq j_1 < \cdots < j_r \leq \bar{1}, s \geq 1 - P^B(n). \right\}$$

(3) For  $k \in I$  with  $k < n - 1$ , we define

$$\text{Tab}_{k,t}^{D_n} := \left\{ \begin{array}{c} \boxed{j_1} \\ \boxed{j_2} \\ \vdots \\ \boxed{j_k} \end{array} \middle| j_1, \dots, j_k \in J_D, j_1 \not\leq j_2 \not\leq \cdots \not\leq j_k, s \geq 1 - P^D(k). \right\},$$

$$\text{Tab}_{n-1,t}^{D_n} := \left\{ \begin{array}{c} \boxed{n+1} \\ \boxed{j_1} \\ \vdots \\ \boxed{j_r} \end{array} \middle| r \in [0, n] \text{ is odd}, j_1, \dots, j_r \in J_D, \bar{n} \leq j_1 < \cdots < j_r \leq \bar{1}, s \geq 1 - P^D(n-1). \right\}$$

$$\text{Tab}_{n,t}^{D_n} := \left\{ \begin{array}{c} \boxed{n+1} \\ \boxed{j_1} \\ \vdots \\ \boxed{j_r} \end{array} \middle| r \in [0, n] \text{ is even}, j_1, \dots, j_r \in J_D, \bar{n} \leq j_1 < \cdots < j_r \leq \bar{1}, s \geq 1 - P^D(n). \right\}$$

For simplicity, we write each tableau  $\begin{array}{c} \boxed{j_1} \\ \vdots \\ \boxed{j_r} \end{array}^X$  as  $[j_1, \dots, j_r]_s^X$ . For  $s \geq 1$ , we define

$$\text{Tab}_{s,k,t}^X := \{[j_1, \dots, j_r]_{s-P^X(k)}^X \in \text{Tab}_{k,t}^X\}.$$

## 6.2 Closedness of $\text{Tab}_{s,k,t}^X$

**Proposition 6.4.** *Let  $\mathfrak{g}$  be of type  $X = A_n, B_n, C_n$  or  $D_n$ . Then  $\text{Tab}_{s,k,t}^{X^L}$  is closed under the action of  $S'_{m,j}$  for all  $m \in \mathbb{Z}_{\geq 1}, j \in I$ .*

*Proof* For  $s \in \mathbb{Z}_{\geq 1 - P^{X^L}(k)}$ , we need to show  $\text{Tab}_{s+P^{X^L}(k),k,t}^{X^L}$  is closed under the action of  $S'_{m,j}$ .

Step1

First, we show that for any  $T = [j_1, \dots, j_r]_s^{X^L} \in \text{Tab}_{s+P^{X^L}(k),k,\ell}^{X^L}$ , if the coefficient of  $x_{t,j}$  in  $T$  is positive (resp. negative) then  $t \geq 1$  (resp.  $t \geq 2$ ). In this proof, we simply write  $P^{X^L}(j)$  as  $P(j)$ .

Recall that for  $X = A, B$  or  $C$ , we have  $T = [j_1, \dots, j_r]_s^{X^L} = \sum_{i=1}^r \boxed{j_i}_{s+r-i}^{X^L}$  and by Definition 6.1, one obtains

$$\boxed{j_i}_{s+r-i}^{X^L} = \begin{cases} c(j_i)x_{s+r-i+P(j_i),j_i} - x_{s+r-i+P(j_i-1)+1,j_i-1} & \text{if } j_i \in \{1, 2, \dots, n\}, \\ x_{s+r-i+P(n),n} - x_{s+r-i+P(n)+1,n} & \text{if } j_i = 0, \\ x_{s+r-i+P(n),n} & \text{if } j_i = \overline{n+1}, \\ x_{s+r-i+P(|j_i|-1)+n-|j_i|+1,|j_i|-1} - c(j_i)x_{s+r-i+P(|j_i|)+n-|j_i|+1,|j_i|} & \text{if } j_i \in \{\overline{n}, \dots, \overline{1}\}, \end{cases} \quad (6.17)$$

where if  $X^L = B_n$  and  $j_i \in \{n, \overline{n}\}$  then  $c(j_i) = 2$ , otherwise  $c(j_i) = 1$ . When  $j_i \in \{1, 2, \dots, n\}$ , it holds  $r = k$  and by  $p_{l,l-1} = 0$  or  $1$  for  $2 \leq l \leq n$ ,  $j_i \geq i$  and  $s \geq 1 - P(k)$ , we have

$$\begin{aligned} s + r - i + P(j_i) &= s + r - i + p_{2,1} + p_{3,2} + \dots + p_{j_i,j_i-1} \\ &\geq s + r - i + p_{2,1} + p_{3,2} + \dots + p_{i,i-1} \\ &\geq s + p_{2,1} + p_{3,2} + \dots + p_{i,i-1} + p_{i+1,i} + \dots + p_{r,r-1} \\ &= s + P(r) = s + P(k) \geq 1. \end{aligned} \quad (6.18)$$

Similarly, if  $j_i > i$  then we get

$$s + r - i + P(j_i - 1) + 1 \geq 2 \quad (6.19)$$

and if  $j_i = i$  then we have  $i = 1$  or  $j_{i-1} = i - 1$  and  $-x_{s+r-i+P(j_i-1)+1,j_i-1}$  in  $\boxed{j_i}_{s+r-i}^{X^L}$  is cancelled in  $\boxed{j_{i-1}}_{s+r-i+1}^{X^L} + \boxed{j_i}_{s+r-i}^{X^L}$ . When  $j_i = 0$  or  $j_i = \overline{n+1}$ , it is easy to see

$$s + r - i + P(n) \geq s + P(k) \geq 1, \quad s + r - i + P(n) + 1 \geq 2 \quad (6.20)$$

from  $r \geq i$ ,  $P(k) \leq P(n)$  ((6.1)) and  $s \geq 1 - P(k)$ . When  $j_i \in \{\overline{n}, \dots, \overline{1}\}$ , it holds  $P(|j_i| - 1) + n - |j_i| + 1 \geq P(n)$ ,  $P(|j_i|) + n - |j_i| \geq P(n)$  so that

$$s + r - i + P(|j_i| - 1) + n - |j_i| + 1 \geq s + P(k) \geq 1, \quad s + r - i + P(|j_i|) + n - |j_i| + 1 \geq 2. \quad (6.21)$$

For  $X = D$ , we see that  $T = [j_1, \dots, j_r]_s^D = \sum_{i=1}^r \boxed{j_i}_{s+r-i}^D$ , and by Definition 6.1, we obtain

$$\boxed{j_i}_{s+r-i}^D = \begin{cases} x_{s+r-i+P(j_i),j_i} - x_{s+r-i+P(j_i-1)+1,j_i-1} & \text{if } j_i \in [1, n-2] \cup \{n\}, \\ x_{s+r-i+P(n-1),n-1} + x_{s+r-i+P(n),n} - x_{s+r-i+P(n-2)+1,n-2} & \text{if } j_i = n-1, \\ x_{s+r-i+P(n),n} & \text{if } j_i = \overline{n+1}, \\ x_{s+r-i+P(n-1),n-1} - x_{s+r-i+P(n)+1,n} & \text{if } j_i = \overline{n}, \\ x_{s+r-i+P(n-2)+1,n-2} - x_{s+r-i+P(n-1)+1,n-1} - x_{s+r-i+P(n)+1,n} & \text{if } j_i = \overline{n-1}, \\ x_{s+r-i+P(|j_i|-1)+n-|j_i|,|j_i|-1} - x_{s+r-i+P(|j_i|)+n-|j_i|,|j_i|} & \text{if } j_i \geq \overline{n-2}. \end{cases} \quad (6.22)$$

When  $j_i \in \{1, 2, \dots, n\}$ , we have  $r = k \leq n - 2$  so that  $P(k) \leq P(n)$ ,  $P(n-1)$  and just as in (6.18), (6.19), one can show

$$s + r - i + P(j_i) \geq s + P(k) \geq 1, \quad \text{if } j_i > i \text{ then } s + r - i + P(j_i - 1) + 1 \geq 2. \quad (6.23)$$

When  $j_i = \overline{n+1}$ , if  $k = n$  then  $s + r - i + P(n) \geq 1$  is clear by  $s \geq 1 - P(k) = 1 - P(n)$ . If  $k = n - 1$  then we have  $i = 1 < r$  so that

$$s + r - i + P(n) \geq s + P(k) \geq 1. \quad (6.24)$$

When  $j_i = \bar{n}$ , if  $k \leq n-1$  then  $s+r-i+P(n-1) \geq 1$  is clear. If  $k = n$  then  $r \geq 3$  and  $r-i \geq 1$  so that

$$s+r-i+P(n-1) \geq s+P(k) \geq 1. \quad (6.25)$$

Similarly, the relation

$$s+r-i+P(n)+1 \geq 2 \quad (6.26)$$

is clear except for the case  $r = i$  and  $k = n-1$ . In this case, we have  $T = \overline{[n+1, \bar{n}]}_s^D = x_{s+P(n-1), n-1}$  and the term  $-x_{s+r-i+P(n)+1, n}$  is cancelled. Similarly, when  $j_i = \overline{n-1}$ , we have

$$s+r-i+P(n-2)+1 \geq s+P(k) \geq 1 \quad (6.27)$$

and it holds

$$s+r-i+P(n-1)+1 \geq 2, \quad s+r-i+P(n)+1 \geq 2 \quad (6.28)$$

or terms  $-x_{s+r-i+P(n-1)+1, n-1}$  or  $-x_{s+r-i+P(n)+1, n}$  are cancelled.

When  $j_i \geq \overline{n-2}$ , by  $P(|j_i|-1)+n-|j_i| \geq P(n), P(n-1)$  and  $P(|j_i|)+n-|j_i|-1 \geq P(n), P(n-1)$ , one obtains

$$s+r-i+P(|j_i|-1)+n-|j_i| \geq s+P(k) \geq 1, \quad s+r-i+P(|j_i|)+n-|j_i| \geq 2. \quad (6.29)$$

Therefore, considering (6.18)-(6.21) and (6.23)-(6.29), if the coefficient of  $x_{t,j}$  in  $T$  is positive (resp. negative) then  $t \geq 1$  (resp.  $t \geq 2$ ).

### Step2

Next, for any  $T = [j_1, \dots, j_r]_s^{X^L} \in \text{Tab}_{s+P(k), k, \iota}^{X^L}$ , let us show  $S'_{m,j}T \in \text{Tab}_{s+P(k), k, \iota}^{X^L}$ . When the coefficient of  $x_{m,j}$  in  $T$  equals 0, the claim is clear so that we assume the coefficient is non-zero. First, we consider the case the coefficient of  $x_{m,j}$  in  $T$  is positive.

#### Case 1 : $j < n$

There is  $i \in [1, r]$  such that  $j_i = j$  or  $j_i = \overline{j+1}$  and  $\boxed{j_i}_{s+k-i}^{X^L}$  has a term  $x_{m,j}$  with a positive coefficient and it holds either  $i = r$  or  $(j_i, j_{i+1}) \neq (j, j+1), (\overline{j+1}, \overline{j}), (\bar{n}, n)$ . When  $j_i = j$  (resp.  $j_i = \overline{j+1}$ ), we see that

$$S'_{m,j}T = T - \beta_{m,j} = [j_1, \dots, j_{i-1}, j+1, j_{i+1}, \dots, j_r]_s^{X^L} \quad (\text{resp. } [j_1, \dots, j_{i-1}, \overline{j}, j_{i+1}, \dots, j_r]_s^{X^L}) \in \text{Tab}_{s+P(k), k, \iota}^{X^L}$$

by Lemma 6.2.

#### Case 2 : $j = n$

We see that there is  $i \in [1, r]$  such that  $\boxed{j_i}_{s+k-i}^{X^L}$  has a term  $x_{m,n}$  with a positive coefficient and the triple  $(X^L, j_i, j_{i+1})$  satisfies one of the following:

$X^L$	$A_n$	$C_n$	$B_n$	$B_n$	$B_n$	$D_n$	$D_n$	$D_n$
$j_i$	$= n$	$= n$	$= n$	$= 0$	$= \overline{n+1}$	$= n-1$	$= n$	$= \overline{n+1}$
$j_{i+1}$	$\neq n+1$	$\neq \bar{n}$	$\neq \bar{n}$	$\neq 0, \bar{n}$	$\neq \bar{n}$	$\neq \bar{n}, n-1$	$\neq \bar{n}, n-1$	$\neq \bar{n}, n-1$

Here, when  $i = r$ , we understand  $j_{i+1}$  satisfies the above condition. When  $(X^L, j_i) = (A_n, n)$  (resp.  $(C_n, n), (B_n, n), (B_n, 0), (B_n, \overline{n+1}), (D_n, n-1), (D_n, n), (D_n, \overline{n+1})$ ), let  $T'$  be a tableau  $[j_1, \dots, j_{i-1}, n+1]$  (resp.  $[j_1, \dots, j_{i-1}, \bar{n}, j_{i+1}, \dots, j_k], [j_1, \dots, j_{i-1}, 0, j_{i+1}, \dots, j_k], [j_1, \dots, j_{i-1}, \bar{n}, j_{i+1}, \dots, j_k]$ ),



$[\overline{n+1}, \overline{n}, j_2, \dots, j_k], [j_1, \dots, j_{i-1}, \overline{n}, j_{i+1}, \dots, j_k], [j_1, \dots, j_{i-1}, \overline{n-1}, j_{i+1}, \dots, j_k], [\overline{n+1}, \overline{n}, \overline{n-1}, j_2, \dots, j_k]$  in  $\text{Tab}_{s+P(k),k,\iota}^{X^L}$ . Here, we simply write  $[j_1, \dots, j_r]_s^{X^L}$  as  $[j_1, \dots, j_r]$ . One obtains

$$S'_{m,n}T = T - \beta_{m,n} = T'$$

by Lemma 6.2.

We can similarly show  $S'_{m,j}T \in \text{Tab}_{s+P(k),k,\iota}^{X^L}$  when the coefficient of  $x_{m,j}$  is negative. Thus, we get our claim.  $\square$

### 6.3 Proof of Theorem 4.3 (1)

Let  $\mathfrak{g}$  of type  $X$  and we show  $\text{Tab}_{s+P(k),k,\iota}^{X^L} = \Xi'_{s+P(k),k,\iota} = \Xi'^+_{s+P(k),k,\iota}$  for  $s \in \mathbb{Z}_{\geq 1-P(k)}$ , which yields our claim by Theorem 4.2. Let  $P_{X^L} = \bigoplus_{j \in I} \mathbb{Z}\Lambda_j$  be the weight lattice of type  $X^L$  with the following partial order : For  $\lambda, \mu \in P_{X^L}$ ,  $\lambda \geq \mu$  if and only if  $\lambda - \mu \in \bigoplus_{j \in I} \mathbb{Z}_{\geq 0}\alpha_j$ .

For the set  $\mathcal{H}$  in (4.1), one considers the  $\mathbb{Z}$ -linear map  $\text{wt} : \mathcal{H} \rightarrow P_{X^L}$  defined as  $\text{wt}(x_{r,j}) := \Lambda_j$  for any  $r \in \mathbb{Z}_{\geq 1}$  and  $j \in I$ . The explicit form (2.14) means

$$\text{wt}(\beta_{r,j}) = \alpha_j \in P_{X^L}. \quad (6.30)$$

First, putting  $s' := s + P(k)$ , we prove  $\text{Tab}_{s',k,\iota}^{X^L} \supset \Xi'_{s',k,\iota} \supset \Xi'^+_{s',k,\iota}$ . Note that

$$x_{s',k} = \begin{cases} [\overline{n+1}]_s^{X^L} & \text{if } k = n, X^L = B_n \text{ or } D_n, \\ [\overline{n+1}, \overline{n}]_s^{X^L} & \text{if } k = n-1, X^L = D_n, \\ [1, 2, \dots, k]_s^{X^L} & \text{otherwise.} \end{cases}$$

Hence,  $x_{s',k} \in \text{Tab}_{s',k,\iota}^{X^L}$ . By the definition of  $\Xi'_{s',k,\iota}$  and Proposition 6.4, we get  $\text{Tab}_{s',k,\iota}^{X^L} \supset \Xi'_{s',k,\iota}$ .

Next, let us show  $\text{Tab}_{s',k,\iota}^{X^L} \subset \Xi'^+_{s',k,\iota}$ . For any  $T = [j_1, \dots, j_r]_s \in \text{Tab}_{s',k,\iota}^{X^L}$ , let us show  $T \in \Xi'^+_{s',k,\iota}$  by the induction on the weight  $\text{wt}(T)$  of  $T$ . Note that it follows from Lemma 6.2 and (6.30) that  $\text{wt}(T) \leq \Lambda_k = \text{wt}(x_{s',k})$ . When

$$T = \begin{cases} [\overline{n+1}]_s^{X^L} & \text{if } k = n, X^L = B_n \text{ or } D_n, \\ [\overline{n+1}, \overline{n}]_s^{X^L} & \text{if } k = n-1, X^L = D_n, \\ [1, 2, \dots, k]_s^{X^L} & \text{otherwise,} \end{cases}$$

it holds  $T = x_{s',k} \in \Xi'^+_{s',k,\iota}$ . Hence, we may assume that  $T \neq x_{s',k}$ . Using Lemma 6.2, (6.18), (6.20), (6.21), (6.23), (6.24), (6.25), (6.27) and (6.29), we can write

$$T = x_{s',k} - \sum_{(t,j) \in \mathbb{Z}_{\geq s'} \times I} c_{t,j} \beta_{t,j}$$

with non-negative coefficients  $c_{t,j} \in \mathbb{Z}_{\geq 0}$ . Except for finitely many  $(t,j)$ , it holds  $c_{t,j} = 0$ . We take

$$(t', j') := \max\{(t,j) \in \mathbb{Z}_{\geq s'} \times I \mid c_{t,j} > 0\}$$

in the order defined in the subsection 2.7. It follows from the definition of  $\beta_{t,j}$  and adaptedness of  $\iota$  that the coefficient of  $x_{t'+1,j'}$  in  $T$  is negative. We can take

$$(t'', j'') := \min\{(t,j) \in \mathbb{Z}_{\geq s'} \times I \mid \text{the coefficient of } x_{t,j} \text{ in } T \text{ is negative}\}.$$

Case 1 :  $j'' < n$

There exists  $i \in [1, r]$  such that  $\boxed{j_i}_{s+r-i}^{X^L}$  has a term  $x_{t'', j''}$  with a negative coefficient and it holds either  $j_i = j'' + 1$  and  $j_{i-1} \neq j'', \bar{n}$  or  $j_i = \bar{j}''$  and  $j_{i-1} \neq \overline{j'' + 1}$ . Let  $T' = [j_1, \dots, j_{i-1}, j'', j_{i+1}, \dots, j_r]_s^{X^L} \in \text{Tab}_{s', k, \iota}^{X^L}$  or  $T' = [j_1, \dots, j_{i-1}, \overline{j'' + 1}, j_{i+1}, \dots, j_r]_s^{X^L} \in \text{Tab}_{s', k, \iota}^{X^L}$ . The minimality of  $(t'', j'')$  implies the coefficient of  $x_{t''-1, j''}$  in  $T'$  is positive. One obtains

$$T = T' - \beta_{t''-1, j''} = S'_{t''-1, j''} T'$$

by Lemma 6.2. Note that by Step1 in the proof of Proposition 6.4, we get  $t'' > 1$ . By the induction assumption, it holds  $T' \in \Xi_{s', k, \iota}^{\prime+}$ , which yields our claim  $T \in \Xi_{s', k, \iota}^{\prime+}$ .

Case 2 :  $j'' = n, (k, X^L) \neq (n, B_n), (n-1, D_n), (n, D_n)$

There exists  $i \in [1, r]$  such that  $\boxed{j_i}_{s+r-i}^{X^L}$  has a term  $x_{t'', n}$  with a negative coefficient and the triple  $(X^L, j_i, j_{i-1})$  is one of the following:

$X^L$	$A_n$	$C_n$	$B_n$	$B_n$	$D_n$	$D_n$
$j_{i-1}$	$\neq n$	$\neq n$	$\neq n, 0$	$\neq n$	$\neq n-1, n$	$\neq n-1, n$
$j_i$	$= n+1$	$= \bar{n}$	$= 0$	$= \bar{n}$	$= \bar{n}$	$= \overline{n-1}$

Here, if  $i = 1$  then we understand  $j_{i-1}$  satisfies the above condition. When  $(j_i, X^L) = (n+1, A_n)$  (resp.  $(j_i, X^L) = (\bar{n}, C_n), (0, B_n), (\bar{n}, B_n), (\bar{n}, D_n), (\overline{n-1}, D_n)$ ), putting  $T' = [j_1, \dots, j_{i-1}, n]$  (resp.  $[j_1, \dots, j_{i-1}, n, j_{i+1}, \dots, j_r], [j_1, \dots, j_{i-1}, n, j_{i+1}, \dots, j_r], [j_1, \dots, j_{i-1}, 0, j_{i+1}, \dots, j_r], [j_1, \dots, j_{i-1}, n-1, j_{i+1}, \dots, j_r], [j_1, \dots, j_{i-1}, n, j_{i+1}, \dots, j_r]$ ), one obtains  $T' \in \text{Tab}_{s', k, \iota}^{X^L}$ . Here, we simply write  $[j_1, \dots, j_r]_s^{X^L}$  as  $[j_1, \dots, j_r]$ . It holds

$$T = T' - \beta_{t''-1, n} = S'_{t''-1, n} T'$$

by Lemma 6.2. Step1 in the proof of Proposition 6.4 yields  $t'' > 1$ . The induction assumption yields  $T' \in \Xi_{s', k, \iota}^{\prime+}$  so that  $T \in \Xi_{s', k, \iota}^{\prime+}$ .

Case 3 :  $j'' = n, (k, X^L) = (n, B_n), (n-1, D_n)$  or  $(n, D_n)$

It holds either  $X^L = B_n$  and  $j_1 = \overline{n+1}, j_2 = \bar{n}$  or  $X^L = D_n$  and  $j_1 = \overline{n+1}, j_2 = \bar{n}, j_3 = \overline{n-1}$ . In the former case, we set  $T' := [\overline{n+1}, j_3, \dots, j_r]_s$  and in the latter case, we set  $T' := [\overline{n+1}, j_4, \dots, j_r]_s$ . Then we have  $T' \in \text{Tab}_{s', k, \iota}^{X^L}$  and

$$T = T' - \beta_{t''-1, n} = S'_{t''-1, n} T'$$

by Lemma 6.2. Using Step1 in the proof of Proposition 6.4, we have  $t'' > 1$ . From the induction assumption, it holds  $T' \in \Xi_{s', k, \iota}^{\prime+}$  so that  $T \in \Xi_{s', k, \iota}^{\prime+}$ . □

## 7 Rank 2 case

We take  $\mathfrak{g}$  as a rank 2 Kac-Moody algebra with a generalized Cartan matrix  $A = \begin{pmatrix} 2 & -a \\ -b & 2 \end{pmatrix}$  ( $a, b \in \mathbb{Z}_{\geq 0}$ ). Let

$$\iota = (\dots, 2, 1, 2, 1, 2, 1).$$

We fix  $s \in \mathbb{Z}$ ,  $i \in I$  and define  $i' \in I$  as  $\{i, i'\} = \{1, 2\} = I$ . One defines  $N := |\langle s_2 s_1 \rangle|$  for  $s_2 s_1 \in W$ . Note that  $N = \infty$  if  $\mathfrak{g}$  is not finite dimensional. For  $m \in \mathbb{Z}_{\geq 0}$  such that  $2m < N$ , let  $P_{m,i}^{(i)}$ ,  $P_{m,i'}^{(i)}$  be integers defined by

$$(s_{i'} s_i)^m \Lambda_i = P_{m,i}^{(i)} \Lambda_i - P_{m,i'}^{(i)} \Lambda_{i'}.$$

Note that when  $2m + 1 < N$ , we have

$$s_i (s_{i'} s_i)^m \Lambda_i = P_{m+1,i'}^{(i)} \Lambda_{i'} - P_{m,i}^{(i)} \Lambda_i,$$

where if  $2m + 2 \geq N$  then we set  $P_{m+1,i'}^{(i)} = 0$ .

Then  $\Xi_{s,i,\ell} := \{S_{j_1} \cdots S_{j_m} x_{s,i} | m \in \mathbb{Z}_{\geq 0}\}$  for  $\mathfrak{g}^L$  is given by

$$\Xi_{s,1,\ell} = \{P_{m,1}^{(1)} x_{s+m,1} - P_{m,2}^{(1)} x_{s+m,2} | m \in \mathbb{Z}_{\geq 0}, 2m < N\} \cup \{P_{m+1,2}^{(1)} x_{s+m,2} - P_{m,1}^{(1)} x_{s+m+1,1} | m \in \mathbb{Z}_{\geq 0}, 2m+1 < N\}$$

and

$$\Xi_{s,2,\ell} = \{P_{m,2}^{(2)} x_{s+m,2} - P_{m,1}^{(2)} x_{s+m+1,1} | m \in \mathbb{Z}_{\geq 0}, 2m < N\} \cup \{P_{m+1,1}^{(2)} x_{s+m+1,1} - P_{m,2}^{(2)} x_{s+m+2,2} | m \in \mathbb{Z}_{\geq 0}, 2m+1 < N\}$$

by the definition of  $S_r$  in (2.5).

**Theorem 7.1.** [Ka24b] Let  $\mathcal{M}_{s,i,\ell}$  be the monomial realization for  $\mathfrak{g}$  in Theorem 3.1 (ii).

(1) The set  $\mathcal{M}_{s,1,\ell}$  includes Laurent monomials in the form

$$\frac{X_{s+m,1}^{P_{m,1}^{(1)}}}{X_{s+m,2}^{P_{m,2}^{(1)}}} \left(1 \leq m < \frac{1}{2}N\right), \quad \frac{X_{s+m,2}^{P_{m+1,2}^{(1)}}}{X_{s+m+1,1}^{P_{m,1}^{(1)}}} \left(1 \leq m < \frac{1}{2}(N-1)\right).$$

Ohter Laurent monomials in  $\mathcal{M}_{s,1,\ell}$  are expressed by a product of above monomials with exponents of positive rational numbers.

(2) The set  $\mathcal{M}_{s,2,\ell}$  includes Laurent monomials in the form

$$\frac{X_{s+m,2}^{P_{m,2}^{(2)}}}{X_{s+m+1,1}^{P_{m,1}^{(2)}}} \left(1 \leq m < \frac{1}{2}N\right), \quad \frac{X_{s+m+1,1}^{P_{m+1,1}^{(2)}}}{X_{s+m+1,2}^{P_{m,2}^{(2)}}} \left(1 \leq m < \frac{1}{2}(N-1)\right).$$

Ohter Laurent monomials in  $\mathcal{M}_{s,2,\ell}$  are expressed by a product of above monomials with exponents of positive rational numbers.

[Proof of Theorem 4.3 (2)]

For  $s \in \mathbb{Z}_{\geq 1}$ , let  $\mathcal{M}'_{s,1,\ell}$  and  $\mathcal{M}'_{s,2,\ell}$  be the sets of monomials in Theorem 7.1 (1) and (2), respectively:

$$\mathcal{M}'_{s,1,\ell} := \left\{ \frac{X_{s+m,1}^{P_{m,1}^{(1)}}}{X_{s+m,2}^{P_{m,2}^{(1)}}} \mid 1 \leq m < \frac{1}{2}N \right\} \cup \left\{ \frac{X_{s+m,2}^{P_{m+1,2}^{(1)}}}{X_{s+m+1,1}^{P_{m,1}^{(1)}}} \mid 1 \leq m < \frac{1}{2}(N-1) \right\},$$

$$\mathcal{M}'_{s,2,\ell} := \left\{ \frac{X_{s+m,2}^{P_{m,2}^{(2)}}}{X_{s+m+1,1}^{P_{m,1}^{(2)}}} \mid 1 \leq m < \frac{1}{2}N \right\} \cup \left\{ \frac{X_{s+m+1,1}^{P_{m+1,1}^{(2)}}}{X_{s+m+1,2}^{P_{m,2}^{(2)}}} \mid 1 \leq m < \frac{1}{2}(N-1) \right\}.$$

Then it is clear that the map  $Trop$  induces a bijection between  $\mathcal{M}'_{s,i,\ell}$  and  $\Xi_{s,i,\ell}$  for  $i = 1, 2$ . For  $M \in \mathcal{M}_{s,i,\ell} \setminus \mathcal{M}'_{s,i,\ell}$  and  $\mathbf{a} = (a_{t,j})_{t \in \mathbb{Z}_{\geq 1}, j \in I} \in \mathbb{Z}^{\infty}$ , if  $Trop(M')(\mathbf{a}) \geq 0$  for all  $M' \in \mathcal{M}'_{s,i,\ell}$  then the inequality  $Trop(M)(\mathbf{a}) \geq 0$  follows by Theorem 7.1. Therefore, our claim is a consequence of Theorem 2.4.  $\square$

## 8 Classical affine case

Let  $X = A_{n-1}^{(1)}, B_{n-1}^{(1)}, C_{n-1}^{(1)}, D_{n-1}^{(1)}, A_{2n-2}^{(2)\dagger}, A_{2n-3}^{(2)}$  or  $D_n^{(2)}$  and  $\iota$  be adapted.

### 8.1 Young walls and truncated walls

Let us review Young walls and truncated walls following [Kang] and [Ka24a]. Each wall consists of  $I$ -colored blocks of three different shapes:

- (1) a block with unit width, unit height and unit thickness:



- (2) a block with unit width, unit height and half-unit thickness:



- (3) a block with unit width, half-unit height and unit thickness:



we simply express the block (1) with color  $j \in I$  as

$$\boxed{j} \tag{8.1}$$

block (2) with color  $j \in I$  as

$$\begin{array}{|} \hline j \\ \hline \end{array} \tag{8.2}$$

or

$$\begin{array}{|} \hline \hline j \\ \hline \end{array} \tag{8.3}$$

and (3) with color  $j \in I$  as

$$\boxed{j} \tag{8.4}$$

We call the blocks (8.1), (8.2), (8.3) and (8.4)  $j$ -blocks.

**Definition 8.1.** For  $k \in I$  and  $X$  other than  $C_{n-1}^{(1)}$ , the index  $k$  is said to be in class 1 if the fundamental weight  $\Lambda_k$  is level 1. The index  $k \in I$  is said to be in class 2 if  $k$  is not in class 1. When  $X = C_{n-1}^{(1)}$ , we understand all indices  $k \in I$  are in class 2. The list of indices  $k \in I$  in class 1 is as follows:

$X$	$A_{n-1}^{(1)}$	$B_{n-1}^{(1)}$	$D_{n-1}^{(1)}$	$A_{2n-2}^{(2)\dagger}$	$A_{2n-3}^{(2)}$	$D_n^{(2)}$
$k$	$1, 2, \dots, n$	$1, 2, n$	$1, 2, n-1, n$	1	1, 2	1, $n$

Let us recall *ground state walls*  $Y_{\Lambda_k}$  of type  $X$  for  $k \in I$  in class 1 [Kang]:

- For  $X = A_{n-1}^{(1)}$  and  $\lambda = \Lambda_k$  ( $k \in I$ ), let  $Y_\lambda$  be the wall that has no blocks.
- For  $X = A_{2n-2}^{(2)\dagger}$  and  $\lambda = \Lambda_1$ , we define

$$Y_{\Lambda_1} := \cdots \boxed{1 \mid 1 \mid 1 \mid 1}$$

The wall  $Y_{\Lambda_1}$  has infinitely many 1-blocks with half-unit height and extends infinitely to the left.

- For  $X = D_n^{(2)}$  and  $\lambda = \Lambda_1, \Lambda_n$ , we define

$$Y_{\Lambda_1} := \cdots \boxed{1 \mid 1 \mid 1}$$

and

$$Y_{\Lambda_n} := \cdots \boxed{n \mid n \mid n}$$

- For  $X = A_{2n-3}^{(2)}$  or  $B_{n-1}^{(1)}$  and  $\lambda = \Lambda_1, \Lambda_2$ ,

$$Y_{\Lambda_1} := \cdots \begin{array}{|c|c|c|c|} \hline \triangle & \triangle & \triangle & \triangle \\ \hline 1 & 2 & 1 & 2 \\ \hline \end{array} \quad Y_{\Lambda_2} := \cdots \begin{array}{|c|c|c|c|} \hline \triangle & \triangle & \triangle & \triangle \\ \hline 2 & 1 & 2 & 1 \\ \hline \end{array}$$

The walls  $Y_{\Lambda_1}, Y_{\Lambda_2}$  extend infinitely to the left. For  $X = B_{n-1}^{(1)}$  and  $\lambda = \Lambda_n$ , we define

$$Y_{\Lambda_n} := \cdots \boxed{n \mid n \mid n}$$

- For  $X = D_{n-1}^{(1)}$  and  $\lambda = \Lambda_1, \Lambda_2, \Lambda_{n-1}$  or  $\Lambda_n$ ,

$$Y_{\Lambda_1} := \cdots \begin{array}{|c|c|c|c|} \hline \triangle & \triangle & \triangle & \triangle \\ \hline 1 & 2 & 1 & 2 \\ \hline \end{array} \quad Y_{\Lambda_2} := \cdots \begin{array}{|c|c|c|c|} \hline \triangle & \triangle & \triangle & \triangle \\ \hline 2 & 1 & 2 & 1 \\ \hline \end{array}$$

$$Y_{\Lambda_{n-1}} := \cdots \begin{array}{|c|c|c|c|} \hline \triangle & \triangle & \triangle & \triangle \\ \hline n-1 & n & n-1 & n \\ \hline \end{array} \quad Y_{\Lambda_n} := \cdots \begin{array}{|c|c|c|c|} \hline \triangle & \triangle & \triangle & \triangle \\ \hline n & n-1 & n & n-1 \\ \hline \end{array}$$

**Definition 8.2.** [Kang] Let  $k \in I$  in class 1. If the following holds then a wall  $Y$  is said to be *Young wall* of ground state  $\Lambda_k$  of type  $X$ :

- (i) The wall  $Y$  is obtained from  $Y_{\Lambda_k}$  by stacking finitely many colored blocks on the ground-state wall  $Y_{\Lambda_k}$ .
- (ii) There are no blocks on the top of a single block with half-unit thickness.
- (iii) Except for the right-most column, there are no free spaces to the right of all blocks.
- (iv) The blocks are stacked following the patterns we give below.

For each type  $X$  and  $\Lambda_k$  with  $k \in I$  in class 1, the patterns of (iv) are as follows:

$$A_{n-1}^{(1)}, \Lambda_k :$$

$n-3$	$n-2$	$n-1$	$n$
$\vdots$	$\vdots$	$\vdots$	$\vdots$
$n-1$	$n$	1	2
$n-2$	$n-1$	$n$	1
$n-3$	$n-2$	$n-1$	$n$
$\vdots$	$\vdots$	$\vdots$	$\vdots$
$k-1$	$k$	$k+1$	$k+2$
$k-2$	$k-1$	$k$	$k+1$
$k-3$	$k-2$	$k-1$	$k$

For  $j_1, j_2$  such that  $\{j_1, j_2\} = \{1, 2\}$ ,

$$A_{2n-3}^{(2)}, \Lambda_{j_1} :$$

3	3	3	3
$j_1/j_2$	$j_2/j_1$	$j_1/j_2$	$j_2/j_1$
3	3	3	3
$\vdots$	$\vdots$	$\vdots$	$\vdots$
$n-1$	$n-1$	$n-1$	$n-1$
$n$	$n$	$n$	$n$
$n-1$	$n-1$	$n-1$	$n-1$
$\vdots$	$\vdots$	$\vdots$	$\vdots$
4	4	4	4
3	3	3	3
$j_1/j_2$	$j_2/j_1$	$j_1/j_2$	$j_2/j_1$

$$B_{n-1}^{(1)}, \Lambda_{j_1} :$$

3	3	3	3
$j_1/j_2$	$j_2/j_1$	$j_1/j_2$	$j_2/j_1$
3	3	3	3
$\vdots$	$\vdots$	$\vdots$	$\vdots$
$n-1$	$n-1$	$n-1$	$n-1$
$n$	$n$	$n$	$n$
$n$	$n$	$n$	$n$
$n-1$	$n-1$	$n-1$	$n-1$
$\vdots$	$\vdots$	$\vdots$	$\vdots$
4	4	4	4
3	3	3	3
$j_1/j_2$	$j_2/j_1$	$j_1/j_2$	$j_2/j_1$

$$B_{n-1}^{(1)}, \Lambda_n :$$

$n-1$	$n-1$	$n-1$	$n-1$
$n$	$n$	$n$	$n$
$n$	$n$	$n$	$n$
$n-1$	$n-1$	$n-1$	$n-1$
$\vdots$	$\vdots$	$\vdots$	$\vdots$
3	3	3	3
$j_1/j_2$	$j_2/j_1$	$j_1/j_2$	$j_2/j_1$
3	3	3	3
$\vdots$	$\vdots$	$\vdots$	$\vdots$
$n-2$	$n-2$	$n-2$	$n-2$
$n-1$	$n-1$	$n-1$	$n-1$
$n$	$n$	$n$	$n$
$n$	$n$	$n$	$n$

Let  $\{j_1, j_2\} = \{1, 2\}$  or  $\{n-1, n\}$ ,  $\{j_3, j_4\} = \{n-1, n\}$  or  $\{1, 2\}$ ,

$$D_{n-1}^{(1)}, \Lambda_{j_1} :$$

	3	3	3	3
$j_1$	$j_2$	$j_1$	$j_2$	
$j_2$	$j_1$	$j_2$	$j_1$	
	3	3	3	3
$\vdots$	$\vdots$	$\vdots$	$\vdots$	
$n-2$	$n-2$	$n-2$	$n-2$	
$j_3$	$j_4$	$j_3$	$j_4$	
$j_4$	$j_3$	$j_4$	$j_3$	
	$n-2$	$n-2$	$n-2$	$n-2$
$\vdots$	$\vdots$	$\vdots$	$\vdots$	
	4	4	4	4
	3	3	3	3
$j_1$	$j_2$	$j_1$	$j_2$	
$j_2$	$j_1$	$j_2$	$j_1$	

$$A_{2n-2}^{(2)\dagger}, \Lambda_1 :$$

2	2	2	2
1	1	1	1
1	1	1	1
2	2	2	2
$\vdots$	$\vdots$	$\vdots$	$\vdots$
$n-1$	$n-1$	$n-1$	$n-1$
$n$	$n$	$n$	$n$
$n-1$	$n-1$	$n-1$	$n-1$
$\vdots$	$\vdots$	$\vdots$	$\vdots$
3	3	3	3
2	2	2	2
1	1	1	1
1	1	1	1

$$D_n^{(2)}, \Lambda_1 :$$

2	2	2	2
1	1	1	1
1	1	1	1
2	2	2	2
$\vdots$	$\vdots$	$\vdots$	$\vdots$
$n-1$	$n-1$	$n-1$	$n-1$
$n$	$n$	$n$	$n$
$n$	$n$	$n$	$n$
$n-1$	$n-1$	$n-1$	$n-1$
$\vdots$	$\vdots$	$\vdots$	$\vdots$
3	3	3	3
2	2	2	2
1	1	1	1
1	1	1	1

$$D_n^{(2)}, \Lambda_n :$$

$n-1$	$n-1$	$n-1$	$n-1$
$n$	$n$	$n$	$n$
$n$	$n$	$n$	$n$
$n-1$	$n-1$	$n-1$	$n-1$
$\vdots$	$\vdots$	$\vdots$	$\vdots$
2	2	2	2
1	1	1	1
1	1	1	1
2	2	2	2
$\vdots$	$\vdots$	$\vdots$	$\vdots$
$n-2$	$n-2$	$n-2$	$n-2$
$n-1$	$n-1$	$n-1$	$n-1$
$n$	$n$	$n$	$n$
$n$	$n$	$n$	$n$

The following is an example of Young wall of ground state  $\Lambda_1$  of type  $A_4^{(2)\dagger}$ .

				2
				3
			2	2
		1	1	1
...	1	1	1	1

Although, when  $X$  is of type  $C_{n-1}^{(1)}$ , the index  $1 \in I$  is not in class 1, we define a pattern as follows:

$$C_{n-1}^{(1)}, \Lambda_1 :$$

2	2	2	2
1	1	1	1
2	2	2	2
$\vdots$	$\vdots$	$\vdots$	$\vdots$
$n-1$	$n-1$	$n-1$	$n-1$
$n$	$n$	$n$	$n$
$n-1$	$n-1$	$n-1$	$n-1$
$\vdots$	$\vdots$	$\vdots$	$\vdots$
3	3	3	3
2	2	2	2
1	1	1	1

Let  $k \in I$  be in class 2. The pattern obtained by the following procedure is called a supporting pattern (resp. covering pattern) for  $\Lambda_k$  of type  $X$ :

- (1) Truncating all boxes of a Young wall pattern for  $\Lambda_1$  of type  $X$  below first (resp. second)  $k$ -boxes from the bottom.
- (2) Splitting the bottom  $k$ -boxes into two half-unit height boxes.

The following is an example of procedure to obtain the supporting pattern for  $\Lambda_k$  of type  $A_{2n-2}^{(2)\dagger}$ :

$A_{2n-2}^{(2)\dagger}, \Lambda_1 :$	$\xrightarrow{(1)}$	$\xrightarrow{(2)}$	$A_{2n-2}^{(2)\dagger}, \Lambda_k :$																																																																																																																																																							
<table border="1" style="border-collapse: collapse; text-align: center; width: 100px; height: 150px;"> <tr><td>2</td><td>2</td><td>2</td><td>2</td></tr> <tr><td>1</td><td>1</td><td>1</td><td>1</td></tr> <tr><td>1</td><td>1</td><td>1</td><td>1</td></tr> <tr><td>2</td><td>2</td><td>2</td><td>2</td></tr> <tr><td><math>\vdots</math></td><td><math>\vdots</math></td><td><math>\vdots</math></td><td><math>\vdots</math></td></tr> <tr><td><math>n-1</math></td><td><math>n-1</math></td><td><math>n-1</math></td><td><math>n-1</math></td></tr> <tr><td><math>n</math></td><td><math>n</math></td><td><math>n</math></td><td><math>n</math></td></tr> <tr><td><math>n-1</math></td><td><math>n-1</math></td><td><math>n-1</math></td><td><math>n-1</math></td></tr> <tr><td><math>\vdots</math></td><td><math>\vdots</math></td><td><math>\vdots</math></td><td><math>\vdots</math></td></tr> <tr><td><math>k</math></td><td><math>k</math></td><td><math>k</math></td><td><math>k</math></td></tr> <tr><td><math>\vdots</math></td><td><math>\vdots</math></td><td><math>\vdots</math></td><td><math>\vdots</math></td></tr> <tr><td>1</td><td>1</td><td>1</td><td>1</td></tr> <tr><td>1</td><td>1</td><td>1</td><td>1</td></tr> </table>	2	2	2	2	1	1	1	1	1	1	1	1	2	2	2	2	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$n-1$	$n-1$	$n-1$	$n-1$	$n$	$n$	$n$	$n$	$n-1$	$n-1$	$n-1$	$n-1$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$k$	$k$	$k$	$k$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	1	1	1	1	1	1	1	1	<table border="1" style="border-collapse: collapse; text-align: center; width: 100px; height: 150px;"> <tr><td>2</td><td>2</td><td>2</td><td>2</td></tr> <tr><td>1</td><td>1</td><td>1</td><td>1</td></tr> <tr><td>1</td><td>1</td><td>1</td><td>1</td></tr> <tr><td>2</td><td>2</td><td>2</td><td>2</td></tr> <tr><td><math>\vdots</math></td><td><math>\vdots</math></td><td><math>\vdots</math></td><td><math>\vdots</math></td></tr> <tr><td><math>n-1</math></td><td><math>n-1</math></td><td><math>n-1</math></td><td><math>n-1</math></td></tr> <tr><td><math>n</math></td><td><math>n</math></td><td><math>n</math></td><td><math>n</math></td></tr> <tr><td><math>n-1</math></td><td><math>n-1</math></td><td><math>n-1</math></td><td><math>n-1</math></td></tr> <tr><td><math>\vdots</math></td><td><math>\vdots</math></td><td><math>\vdots</math></td><td><math>\vdots</math></td></tr> <tr><td><math>k+2</math></td><td><math>k+2</math></td><td><math>k+2</math></td><td><math>k+2</math></td></tr> <tr><td><math>k+1</math></td><td><math>k+1</math></td><td><math>k+1</math></td><td><math>k+1</math></td></tr> <tr><td><math>k</math></td><td><math>k</math></td><td><math>k</math></td><td><math>k</math></td></tr> </table>	2	2	2	2	1	1	1	1	1	1	1	1	2	2	2	2	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$n-1$	$n-1$	$n-1$	$n-1$	$n$	$n$	$n$	$n$	$n-1$	$n-1$	$n-1$	$n-1$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$k+2$	$k+2$	$k+2$	$k+2$	$k+1$	$k+1$	$k+1$	$k+1$	$k$	$k$	$k$	$k$	<table border="1" style="border-collapse: collapse; text-align: center; width: 100px; height: 150px;"> <tr><td>2</td><td>2</td><td>2</td><td>2</td></tr> <tr><td>1</td><td>1</td><td>1</td><td>1</td></tr> <tr><td>1</td><td>1</td><td>1</td><td>1</td></tr> <tr><td>2</td><td>2</td><td>2</td><td>2</td></tr> <tr><td><math>\vdots</math></td><td><math>\vdots</math></td><td><math>\vdots</math></td><td><math>\vdots</math></td></tr> <tr><td><math>n-1</math></td><td><math>n-1</math></td><td><math>n-1</math></td><td><math>n-1</math></td></tr> <tr><td><math>n</math></td><td><math>n</math></td><td><math>n</math></td><td><math>n</math></td></tr> <tr><td><math>n-1</math></td><td><math>n-1</math></td><td><math>n-1</math></td><td><math>n-1</math></td></tr> <tr><td><math>\vdots</math></td><td><math>\vdots</math></td><td><math>\vdots</math></td><td><math>\vdots</math></td></tr> <tr><td><math>k+2</math></td><td><math>k+2</math></td><td><math>k+2</math></td><td><math>k+2</math></td></tr> <tr><td><math>k+1</math></td><td><math>k+1</math></td><td><math>k+1</math></td><td><math>k+1</math></td></tr> <tr><td><math>k</math></td><td><math>k</math></td><td><math>k</math></td><td><math>k</math></td></tr> <tr><td><math>k</math></td><td><math>k</math></td><td><math>k</math></td><td><math>k</math></td></tr> </table>	2	2	2	2	1	1	1	1	1	1	1	1	2	2	2	2	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$n-1$	$n-1$	$n-1$	$n-1$	$n$	$n$	$n$	$n$	$n-1$	$n-1$	$n-1$	$n-1$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$k+2$	$k+2$	$k+2$	$k+2$	$k+1$	$k+1$	$k+1$	$k+1$	$k$	$k$	$k$	$k$	$k$	$k$	$k$	$k$
2	2	2	2																																																																																																																																																							
1	1	1	1																																																																																																																																																							
1	1	1	1																																																																																																																																																							
2	2	2	2																																																																																																																																																							
$\vdots$	$\vdots$	$\vdots$	$\vdots$																																																																																																																																																							
$n-1$	$n-1$	$n-1$	$n-1$																																																																																																																																																							
$n$	$n$	$n$	$n$																																																																																																																																																							
$n-1$	$n-1$	$n-1$	$n-1$																																																																																																																																																							
$\vdots$	$\vdots$	$\vdots$	$\vdots$																																																																																																																																																							
$k$	$k$	$k$	$k$																																																																																																																																																							
$\vdots$	$\vdots$	$\vdots$	$\vdots$																																																																																																																																																							
1	1	1	1																																																																																																																																																							
1	1	1	1																																																																																																																																																							
2	2	2	2																																																																																																																																																							
1	1	1	1																																																																																																																																																							
1	1	1	1																																																																																																																																																							
2	2	2	2																																																																																																																																																							
$\vdots$	$\vdots$	$\vdots$	$\vdots$																																																																																																																																																							
$n-1$	$n-1$	$n-1$	$n-1$																																																																																																																																																							
$n$	$n$	$n$	$n$																																																																																																																																																							
$n-1$	$n-1$	$n-1$	$n-1$																																																																																																																																																							
$\vdots$	$\vdots$	$\vdots$	$\vdots$																																																																																																																																																							
$k+2$	$k+2$	$k+2$	$k+2$																																																																																																																																																							
$k+1$	$k+1$	$k+1$	$k+1$																																																																																																																																																							
$k$	$k$	$k$	$k$																																																																																																																																																							
2	2	2	2																																																																																																																																																							
1	1	1	1																																																																																																																																																							
1	1	1	1																																																																																																																																																							
2	2	2	2																																																																																																																																																							
$\vdots$	$\vdots$	$\vdots$	$\vdots$																																																																																																																																																							
$n-1$	$n-1$	$n-1$	$n-1$																																																																																																																																																							
$n$	$n$	$n$	$n$																																																																																																																																																							
$n-1$	$n-1$	$n-1$	$n-1$																																																																																																																																																							
$\vdots$	$\vdots$	$\vdots$	$\vdots$																																																																																																																																																							
$k+2$	$k+2$	$k+2$	$k+2$																																																																																																																																																							
$k+1$	$k+1$	$k+1$	$k+1$																																																																																																																																																							
$k$	$k$	$k$	$k$																																																																																																																																																							
$k$	$k$	$k$	$k$																																																																																																																																																							



**Definition 8.3.** [Ka24a] Let  $k \in I$  be in class 2. A wall  $Y$  is called a *truncated wall* of supporting (resp. covering) ground state  $\Lambda_k$  of type  $X$  if it satisfies the following:

- (i) The wall  $Y$  is obtained from  $Y_{\Lambda_k}$  by stacking finitely many colored blocks on the ground-state wall  $Y_{\Lambda_k}$ .
- (ii) There are no blocks on the top of a single block with half-unit thickness.
- (iii) Except for the right-most column, there are no free spaces to the right of all blocks.
- (iv) The colored blocks are stacked following the supporting (resp. covering) pattern for  $\Lambda_k$  of type  $X$ .

Here, the ground state wall  $Y_{\Lambda_k}$  is as follows:

$$\cdots \boxed{k \mid k \mid k} \cdots$$

The conditions (i), (ii) and (iii) are same as in the Definition 8.2.

**Definition 8.4.** [Kang] Let  $Y$  be a Young wall or truncated wall of ground state  $\lambda$  of type  $X$ .

- (i) A column of  $Y$  is called a *full column* if its top has unit thickness and its height is a multiple of the unit length.
- (ii) For  $X = A_{n-1}^{(1)}$ , we understand all Young walls are proper.
- (iii) For  $X = A_{2n-3}^{(2)}, A_{2n-2}^{(2)\dagger}, B_{n-1}^{(1)}, D_{n-1}^{(1)}$  and  $D_n^{(2)}$ , the Young wall  $Y$  is said to be *proper* if none of two full columns of  $Y$  have the same height.

**Definition 8.5.** Let  $k \in I$  be in class 2 and  $Y$  be a proper truncated wall of ground state  $\Lambda_k$ . A  $k$ -block with half-unit height put on the ground state wall is called a *second half  $k$ -block*.

**Definition 8.6.** [Kang] Let  $Y$  be a proper Young wall or proper truncated wall of ground state  $\Lambda_k$  and  $i \in I$ .

- (i) An  $i$ -block in  $Y$  other than second half  $k$ -block is said to be a *removable  $i$ -block* if the wall obtained from  $Y$  by removing this  $i$ -block remains proper.
- (ii) A place is said to be an  *$i$ -admissible slot* if the wall obtained from  $Y$  by adding an  $i$ -block other than second half  $k$ -block to this place remains proper.


**Definition 8.7.** [Ka23a] Let  $Y$  be a proper Young wall or proper truncated wall and  $t = 1$  or  $t = n$ .

- (i) Let  $Y'$  be a wall obtained from  $Y$  by putting two  $t$ -blocks of shape (8.4) on the top of a column of  $Y$ :

$$Y = \cdots \begin{array}{c} \text{---} \\ \text{---} \\ \boxed{\phantom{t}} \\ \text{---} \\ \vdots \end{array} \cdots \leftarrow A \qquad Y' = \cdots \begin{array}{c} \boxed{t} \\ \boxed{t} \\ \boxed{\phantom{t}} \\ \text{---} \\ \vdots \end{array} \cdots$$

In  $Y$ , we give a name to a slot as above. The slot  $A$  in  $Y$  is said to be *double  $t$ -admissible* if  $Y'$  is proper.

- (ii) We assume that we obtain a wall  $Y''$  by removing two  $t$ -blocks of shape (8.4) from the top of a column of  $Y$ :



A block in  $Y$  is named  $B$  as above. The block  $B$  in  $Y$  is said to be *double  $t$ -removable* if  $Y''$  is proper.


- (iii) For  $i \in I$ , other  $i$ -admissible slots (resp. removable  $i$ -blocks) are said to be single admissible (resp. single removable).

## 8.2 Assignment of homomorphisms to walls

Following [Ka24a], we assign some homomorphisms to Young walls and truncated walls. As for the things we will not use in this paper, we just cited from the previous paper.

For  $k \in I$ , we define  $T_k^X$ ,  $\overline{T}_k^X$  and  $\overline{\overline{T}}_k^X \in \mathbb{Z}_{\geq 1}$  just as in subsection 3.1 of [Ka24a]. From now on, we will draw proper Young walls of ground state  $\Lambda_k$  of type  $X$  in  $\mathbb{R}_{\leq 0} \times \mathbb{R}_{\geq T_k^X}$  for each  $k \in I$  in class 1. We draw truncated walls of supporting (resp. covering) ground state  $\Lambda_k$  of type  $X$  in  $\mathbb{R}_{\leq 0} \times \mathbb{R}_{\geq \overline{T}_k^X}$  (resp. in  $\mathbb{R}_{\leq 0} \times \mathbb{R}_{\geq \overline{\overline{T}}_k^X}$ ) for each  $k \in I$  in class 2.

Let  $S$  be one of the following slots or blocks in a proper Young wall or truncated wall with  $i \in \mathbb{Z}_{\geq 0}$ ,  $l \in \mathbb{Z}_{\geq 1}$ :



Then we say the position of  $S$  is  $(-i, l)$ . If  $S$  is colored by  $t \in I$  then we assign a homomorphism

$$L_{s,k,\text{ad}}^X(S) := \begin{cases} x_{r,t} & \text{if } S \text{ is admissible,} \\ 0 & \text{otherwise,} \end{cases} \quad L_{s,k,\text{re}}^X(S) := \begin{cases} x_{r+1,t} & \text{if } S \text{ is removable,} \\ 0 & \text{otherwise.} \end{cases} \quad (8.5)$$

Here,  $r \in \mathbb{Z}_{\geq s}$  is determined from  $i, l, s$  and  $\iota$ . The precise definition of  $r$  is in subsection 4.3 of [Ka24a].

Let  $Y$  be a proper Young wall or proper truncated wall of ground state  $\Lambda_k$  of type  $X = A_{n-1}^{(1)}$ ,  $B_{n-1}^{(1)}$ ,  $C_{n-1}^{(1)}$ ,  $D_{n-1}^{(1)}$ ,  $A_{2n-2}^{(2)\dagger}$ ,  $A_{2n-3}^{(2)}$  or  $D_n^{(2)}$ . For  $s \in \mathbb{Z}$ , we define

$$L_{s,k,\iota}^X(Y) := \sum_{t \in I} \left( \sum_{P: \text{single } t\text{-admissible slot}} L_{s,k,\text{ad}}^X(P) - \sum_{P: \text{single removable } t\text{-block}} L_{s,k,\text{re}}^X(P) \right) + \sum_{P: \text{double } t\text{-admissible slot}} 2L_{s,k,\text{ad}}^X(P) - \sum_{P: \text{double removable } t\text{-block}} 2L_{s,k,\text{re}}^X(P). \quad (8.6)$$

### 8.3 Set of walls

**Definition 8.8.** [Ka24a]

- (i) For  $k \in I$  in class 1, we define  $\text{YW}_k^X$  as the set of all proper Young walls of ground state  $\Lambda_k$  of type  $X$ .
- (ii) For  $k \in I$  in class 2, one defines  $\overline{\text{YW}}_k^X$  (resp.  $\overline{\overline{\text{YW}}}_k^X$ ) as the set of all proper truncated walls of supporting (resp. covering) ground state  $\Lambda_k$  of type  $X$ .
- (iii) Let  $k \in I$  be in class 2. We define

$$\text{YW}_k^X := \left\{ (\overline{Y}, \overline{\overline{Y}}) \in \overline{\text{YW}}_k^X \times \overline{\overline{\text{YW}}}_k^X \mid h_j = \frac{1}{2} \text{ if and only if } h'_j = \frac{1}{2} \text{ for } j \in \mathbb{Z}_{\geq 1} \right\},$$

where  $h_j$  (resp.  $h'_j$ ) is the height of  $j$ -th column of  $\overline{Y}$  (resp.  $\overline{\overline{Y}}$ ) from the right. We also define

$$Y_{\Lambda_k} := ( \cdots \boxed{k \mid k \mid k}, \cdots \boxed{k \mid k \mid k} ) \in \text{YW}_k^X.$$

**Definition 8.9.** [Ka24a] Let  $k \in I$  be in class 2.

- (i) For a pair  $Y = (\overline{Y}, \overline{\overline{Y}}) \in \text{YW}_k^X$ , we assume that  $Y' = (\overline{Y}', \overline{\overline{Y}}')$  is obtained from  $Y$  by removing one second half  $k$ -block

$$\boxed{k}$$

from the top of  $j$ -th column of  $\overline{Y}$  and one from the top of  $j$ -th column of  $\overline{\overline{Y}}$  with some  $j \in \mathbb{Z}_{\geq 1}$ . Then the pair of these two  $k$ -blocks are said to be a *removable  $k$ -pair*.

- (ii) For  $Y = (\overline{Y}, \overline{\overline{Y}}) \in \text{YW}_k^X$ , we assume that  $Y'' = (\overline{Y}'', \overline{\overline{Y}}'')$  is obtained from  $Y$  by adding one second half  $k$ -block

$$\boxed{k}$$

to the top of  $j$ -th column of  $\overline{Y}$  and one to the top of  $j$ -th column of  $\overline{\overline{Y}}$  with some  $j \in \mathbb{Z}_{\geq 1}$ . Then the pair of these two  $k$ -slots are said to be a  *$k$ -admissible pair*.

For  $k \in I$  in class 2, let  $P$  be a pair of  $k$ -blocks in some  $Y \in \text{YW}_k^X$  with the following coordinate:

$$P = \begin{array}{cccc} (-i-1, \overline{T}_k+1) & (-i, \overline{T}_k+1) & (-i-1, \overline{\overline{T}}_k+1) & (-i, \overline{\overline{T}}_k+1) \\ & \boxed{k} & & \boxed{k} \\ (-i-1, \overline{T}_k+\frac{1}{2}) & (-i, \overline{T}_k+\frac{1}{2}) & (-i-1, \overline{\overline{T}}_k+\frac{1}{2}) & (-i, \overline{\overline{T}}_k+\frac{1}{2}) \end{array}$$

Then we define

$$L_{s,k,\text{ad}}^X(P) := \begin{cases} x_{s+i,k} & \text{if } P \text{ is a } k\text{-admissible pair,} \\ 0 & \text{otherwise,} \end{cases} \quad L_{s,k,\text{re}}^X(P) := \begin{cases} x_{s+i+1,k} & \text{if } P \text{ is a removable } k\text{-pair,} \\ 0 & \text{otherwise.} \end{cases} \quad (8.7)$$

For above pair  $P$ , we say the *position* of  $P$  is  $(-i, \bar{T}_k)$ . For  $k \in I$  in class 2 and  $Y = (\bar{Y}, \overline{\bar{Y}}) \in \text{YW}_k^X$ , using (8.6) and (8.7), one defines

$$L_{s,k,\iota}^X(Y) := L_{s,k,\iota}^X(\bar{Y}) + L_{s,k,\iota}^X(\overline{\bar{Y}}) + \sum_{P:k\text{-admissible pair in } Y} L_{s,k,\text{ad}}^X(P) - \sum_{P:\text{removable } k\text{-pair in } Y} L_{s,k,\text{re}}^X(P). \quad (8.8)$$

For each  $k \in I$  and  $s \in \mathbb{Z}_{\geq 1}$ , we set  $\text{COMB}_{s,k,\iota}^X[\infty] := \{L_{s,k,\iota}^X(Y) | Y \in \text{YW}_k^X\}$ .

**Theorem 8.10.** [Ka24a] Let  $\mathfrak{g}$  be of type  $X^L$  ( $= A_{n-1}^{(1)}, B_{n-1}^{(1)}, C_{n-1}^{(1)}, D_{n-1}^{(1)}, A_{2n-2}^{(2)}, A_{2n-3}^{(2)}$  or  $D_n^{(2)}$ ), where  $X^L$  is defined in (2.1). Let  $\iota$  be an adapted sequence. Then  $\iota$  satisfies the  $\Xi'$ -positivity condition and we have

$$\Xi'_{s,k,\iota} = \text{COMB}_{s,k,\iota}^X[\infty].$$

**Proposition 8.11.** [Ka24a] Let  $\mathfrak{g}$  be of type  $X^L$  ( $= A_{n-1}^{(1)}, B_{n-1}^{(1)}, C_{n-1}^{(1)}, D_{n-1}^{(1)}, A_{2n-2}^{(2)}, A_{2n-3}^{(2)}$  or  $D_n^{(2)}$ ) and  $s \in \mathbb{Z}_{\geq 1}$ ,  $k \in I$ .

(1) For  $Y_{\Lambda_k} \in \text{YW}_k^X$ , we have  $L_{s,k,\iota}^X(Y_{\Lambda_k}) = x_{s,k}$ .

(2) Let  $Y, Y' \in \text{YW}_k^X$  and we assume that  $Y'$  is obtained from  $Y$  by adding a  $t$ -block or pair of  $t$ -blocks to a  $t$ -admissible slot or  $t$ -admissible pair  $P$  for  $t \in I$ . Then we have

$$L_{s,k,\iota}^X(Y') = L_{s,k,\iota}^X(Y) - \beta_{r,t},$$

where  $r \in \mathbb{Z}_{\geq s}$  and  $L_{s,k,\text{ad}}^X(P) = x_{r,t}$ .

## 8.4 Proof of Theorem 4.3 (3)

We assume  $\mathfrak{g}$  is of type  $X^L$ . Let us show Theorem 4.3 (3). By Theorem 4.2 and 8.10, we need to show  $\text{COMB}_{s,k,\iota}^X[\infty] \subset \Xi'_{s,k,\iota}$ . We take any  $Y \in \text{YW}_k^X$  and let us prove  $L_{s,k,\iota}^X(Y) \in \Xi'_{s,k,\iota}$  by the induction on the number of blocks. Here, we say the number of blocks in  $Y$  is  $m$  if we get the wall  $Y$  by adding  $m$  blocks to  $Y_{\Lambda_k}$ .

If the number of blocks is 0 then we have  $Y = Y_{\Lambda_k}$  and  $L_{s,k,\iota}^X(Y) = x_{s,k} \in \Xi'_{s,k,\iota}$  (Proposition 8.11 (1)) so that we may assume  $Y$  has at least one block. It follows from Proposition 8.11 (2) that  $L_{s,k,\iota}^X(Y)$  can be expressed as

$$L_{s,k,\iota}^X(Y) = x_{s,k} - \sum_{(r,j) \in \mathbb{Z}_{\geq s} \times I} c_{r,j} \beta_{r,j}$$

with non-negative integers  $\{c_{r,j}\}$ . It holds  $c_{r,j} = 0$  except for finitely many  $(r,j)$ . By  $Y \neq Y_{\Lambda_k}$ , there is a pair  $(m', t') \in \mathbb{Z}_{\geq s} \times I$  such as  $(m', t') = \max\{(r,j) \in \mathbb{Z}_{\geq s} \times I | c_{r,j} > 0\}$ . Recall that we defined the order on  $\mathbb{Z}_{\geq 1} \times I$  in the subsection 2.7.

Taking (2.14) into the account, it is easy to see the coefficient of  $x_{m'+1, t'}$  is negative in  $L_{s,k,\iota}^X(Y)$ . Thus, one can also take  $(m'', t'')$  as

$$(m'', t'') = \min\{(r,j) \in \mathbb{Z}_{\geq 1} \times I | \text{the coefficient of } x_{r,j} \text{ in } L_{s,k,\iota}^X(Y) \text{ is negative}\}.$$

The definitions (8.6), (8.8) of  $L_{s,k,t}^X(Y)$  yield that there is a removable  $t''$ -block or removable  $t''$ -pair  $B$  in  $Y$  such that  $L_{s,k, \text{re}}^X(B) = x_{m'',t''}$  and  $m'' \geq 2$ . Let  $Y'' \in \text{YW}_k^X$  be the wall obtained from  $Y$  by removing  $B$ . Using the induction assumption, it follows  $L_{s,k,t}^X(Y'') \in \Xi_{s,k,t}^{\prime+}$ . Proposition 8.11 (2) yields

$$L_{s,k,t}^X(Y) = L_{s,k,t}^X(Y'') - \beta_{m''-1,t''}$$

and by the minimality of  $(m'', t'')$ , the coefficient of  $x_{m''-1,t''}$  in  $L_{s,k,t}^X(Y'')$  is positive. Hence,  $L_{s,k,t}^X(Y) = S'_{m''-1,t''} L_{s,k,t}^X(Y'')$ . Combining with  $L_{s,k,t}^X(Y'') \in \Xi_{s,k,t}^{\prime+}$ , one obtains  $L_{s,k,t}^X(Y) \in \Xi_{s,k,t}^{\prime+}$ .  $\square$

## References

- [Hos05] A.Hoshino, Polyhedral realizations of crystal bases for quantum algebras of finite types, J. Math. Phys. 46, no. 11, 113514, 31 pp, (2005).
- [Hos13] A.Hoshino, Polyhedral realizations of crystal bases for quantum algebras of classical affine types, J. Math. Phys. 54, no. 5, 053511, 28 pp, (2013).
- [JMMO] M.Jimbo, K. C. Misra, T.Miwa, M.Okado, Combinatorics of representations of  $U_q(\widehat{\mathfrak{sl}}(n))$  at  $q = 0$ , Comm. Math. Phys. 136, no. 3, 543–566 (1991).
- [Kang] S.-J. Kang, Crystal bases for quantum affine algebras and combinatorics of Young walls, Proc. London Math. Soc. (3) 86, no. 1, 29–69 (2003).
- [KL] S.-J. Kang, H. Lee, Crystal bases for quantum affine algebras and Young walls. J. Algebra 322, no. 6, 1979–1999 (2009).
- [KMM] S.-J. Kang, K. C. Misra, T.Miwa, Fock space representations of the quantized universal enveloping algebras  $U_q(C_l^{(1)})$ ,  $U_q(A_{2l}^{(2)})$ , and  $U_q(D_{l+1}^{(2)})$ , J. Algebra 155, no. 1, 238–251 (1993).
- [Ka23a] Y.Kanakubo, Polyhedral realizations for  $B(\infty)$  and extended Young diagrams, Young walls of type  $A_{n-1}^{(1)}$ ,  $C_{n-1}^{(1)}$ ,  $A_{2n-2}^{(2)}$ ,  $D_n^{(2)}$ , Algebr. Represent. Theory 26, no.5, 2181–2233 (2023).
- [Ka23b] Y.Kanakubo, Polyhedral realizations for crystal bases of integrable highest weight modules and combinatorial objects of type  $A_{n-1}^{(1)}$ ,  $C_{n-1}^{(1)}$ ,  $A_{2n-2}^{(2)}$ ,  $D_n^{(2)}$ , Lett. Math. Phys. 113 : 60, no.3, 1–50 (2023).
- [Ka24a] Y.Kanakubo, Polyhedral realizations for crystal bases and Young walls of classical affine types, arXiv:2403.01190.
- [Ka24b] Y.Kanakubo, Monomial realizations and LS paths of fundamental representations for rank 2 Kac-Moody algebras, arXiv:2407.19622 (to appear in J. Algebra).
- [KaN20] Y.Kanakubo, T.Nakashima, Adapted sequence for polyhedral realization of crystal bases, Communications in Algebra 48, Issue 11, 4732–4766 (2020).
- [Kas90] M.Kashiwara, Crystallizing the  $q$ -analogue of universal enveloping algebras, Comm. Math. Phys., 133, 249–260 (1990).
- [Kas91] M.Kashiwara, On crystal bases of the  $q$ -analogue of universal enveloping algebras, Duke Math. J., 63 (2), 465–516 (1991).
- [Kas93] M.Kashiwara, The crystal base and Littelmann’s refined Demazure character formula, Duke Math. J., 71, no 3, 839–858 (1993).

- [Kas03] M.Kashiwara, Realizations of crystals, Combinatorial and geometric representation theory (Seoul, 2001), Contemp. Math., 325, Amer. Math. Soc., Providence, RI, pp.133–139 (2003).
- [KN] M.Kashiwara, T.Nakashima, Crystal graphs for representations of the  $q$ -analogue of classical Lie algebras, J. Algebra 165, no. 2, 295–345 (1994).
- [KS] Kim, J.-A., Shin, D.-U., Monomial realization of crystal bases  $B(\infty)$  for the quantum finite algebras, Algebr. Represent. Theory 11, no. 1, 93–105 (2008).
- [Lit94] P.Littelmann, A Littlewood-Richardson rule for symmetrizable Kac-Moody algebras, Invent. Math. 116, no.1-3, pp.329–346 (1994).
- [Lit95] P.Littelmann, Paths and root operators in representation theory, Ann. of Math. (2)142, no.3, pp.499–525 (1995).
- [Lit98] P.Littelmann, Cones, crystals, and patterns, Transform. Groups 3, no. 2, 145–179 (1998).
- [Lus] G.Lusztig, Canonical bases arising from quantized enveloping algebras, J. Amer. Math. Soc. 3, no. 2, 447–498 (1990).
- [MM] K. C. Misra, T.Miwa, Crystal base for the basic representation of  $U_q(\widehat{\mathfrak{sl}}(n))$ , Comm. Math. Phys. 134, no. 1, 79–88 (1990).
- [Nak] H.Nakajima,  $t$ -analogs of  $q$ -characters of quantum affine algebras of type  $A_n, D_n$ , Combinatorial and geometric representation theory (Seoul, 2001), Contemp. Math., 325, Amer. Math. Soc., Providence, RI, pp.141–160 (2003).
- [Na] T.Nakashima, Decorated Geometric Crystals, Polyhedral and Monomial Realizations of Crystal Bases, Contemporary Mathematics 602, 143–163 (2013).
- [NZ] T.Nakashima, A.Zelevinsky, Polyhedral realizations of crystal bases for quantized Kac-Moody algebras, Adv. Math. **131**, no. 1, 253–278 (1997).