Categories meet semigroups in various ways

Bertalan Pécsi

2025

Abstract

This paper touches on several interaction points of semigroups and constructions from category theory.

Introduction

(...)

Throughout the paper we write compositions of arrows of a category in the order of application left to right, and consequently apply functions and functors to the *right* to their arguments, often in exponents, so that x^f (or xf) denotes the image of element x under mapping f.

Accordingly, epimorphisms, monomorphisms, split epi, (...)

1 Wired categories

Definition 1.1. We call a category C equipped with a chosen arrow $w_{a,b} : a \to b$ for each pair of objects $a, b \in Ob C$ a wired category if $w_{a,a} = 1_a$ for each $a \in Ob C$.

Proposition 1.2. Any small wired category (\mathcal{C}, w) induces a semigroup structure on its set of arrows.

Proof. For arrows $\varphi : a \to b$ and $\psi : c \to d$ define

$$\varphi \bullet \psi := \varphi \cdot w_{b,c} \cdot \psi.$$

This is an associative operation.

Recall that the Karoubi envelope (alternatively, idempotent or Cauchy completion) of a semigroup S consists of the idempotents of S as objects and triples $(e, x, f) \in S^3$ as arrows $e \to f$ where e, f are idempotents and exf = x. The composition is defined as follows:

$$(e, x, f) \cdot (f, y, g) := (e, xy, g).$$

Definition 1.3. Let (\mathcal{C}, w) and (\mathcal{D}, v) be wired categories. A functor $F : \mathcal{C} \to \mathcal{D}$ is called a *wired functor* if for all objects $a, b \in \text{Ob}\mathcal{C}$ we have $(w_{a,b})^F = v_{a^F, b^F}$. We denote by $\mathcal{WC}at$ the category of small wired categories and wired functors.

Note that there would be no further restriction on natural transformations between wired functors.

Proposition 1.4. The Karoubi envelope functor $Sgr \to Cat$ factors through $\mathcal{WC}at$, and it is right adjoint to the construction $S : \mathcal{WC}at \to Sgr$ in 1.2.

Proof. Define the wires of the Karoubi envelope by $w_{e,f} := ef$ for idempotents $e, f \in S$.

The elements of both homsets in the adjunction to be proved, naturally correspond to maps from the set of arrows of a wired category (\mathcal{C}, w) to the semigroup S that converts composition in \mathcal{C} to the operation of S.

Definition 1.5. The principal identifier relation of a small wired category (\mathcal{C}, w) is the binary relation Θ on the set of arrows of \mathcal{C} , for which $\underset{a \to x}{\alpha} \Theta \underset{b \to w}{\beta}$ holds if

and only if the following diagram commutes.



It is clear that Θ is reflexive and symmetric, but in general it is not transitive. Also, if α, β are parallel arrows $a \to x$ (i.e. a = b, x = y above) then $\alpha \Theta \beta$ implies $\alpha = \beta$.

2 Regular semigroups

Theorem 2.1. A semigroup S is regular if and only if it has enough idempotents and its Karoubi envelope admits a *split epi* – *split mono* factorization.

3 Lax spiders

Another approach in studying the category-like structure determined by a semigroup S is that instead of a single object we enable a whole set of objects as the "domain" / "codomain" of an element of S.

Here we describe one way to formulate this point of view.

Definition 3.1. Let S be a semigroup and let T be an ordered semigroup. We call a function $f: S \to T$ lax morphism or subhomomorphism if for all $x, y \in S$ it satisfies

$$x^f \cdot y^f \le (x \cdot y)^f$$
.

Note that lax morphisms in the above sense correspond to lax functors in a suitable setup when both S and T are turned into monoidal categories (by adding an identity element and other necessary structure).

Let E denote the set of idempotents in S, $\mathbf{P}(E)$ the set of its subsets, and consider the rectangular band on $\mathbf{P}(E) \times \mathbf{P}(E)$ (so that its operation is $(A, B) \cdot (C, D) = (A, D)$) with partial order $(A, B) \leq (A', B')$ whenever $A \subseteq A'$ and $B \subseteq B'$.

Then we get a lax morphism $S \to \mathbf{P}(E) \times \mathbf{P}(E)$ by assigning

$$x \mapsto (\{e \in E : ex = x\}, \{e \in E : xe = x\})$$

4 Semigroupads

Definition 4.1. A semigroupad or semimonad on a category C is a semigroup object in the category of endofunctors on C, i.e. a monad $T : C \to C$ without the requirement of the unit transformation $\eta : 1_C \to T$ and its defining identities. So, a semigroupad on C is a functor $T : C \to C$ equipped with a natural transformation $\mu : TT \to T$ which is 'associative' in the sense that $T\mu \cdot \mu = \mu T \cdot \mu$, i.e. for each object $a \in Ob C$ the following diagram commutes:



Examples 4.2 (on Set).

• Let T be the functor $A \mapsto A \times S$ for any fixed semigroup S, and let μ be the free S-action, i.e.,

$$A^{\mu} = ((a, s_1), s_2) \mapsto (a, s_1 s_2) : (A \times S) \times S \to A \times S$$

- (Finite) lists or subsets of size greater than a given minimum.
- Nonprincipal ultrafilters.

Theorem 4.3. For a given functor $T : \mathcal{C} \to \mathcal{C}$, a semigroupad structure $\mu : TT \to T$ determines and is determined by a *Kleisli lifting* operation which assigns to each arrow f of the form $a \to b^T$ an arrow $*f : a^T \to b^T$ so that the following conditions hold:

K1. If
$$f: a \to b$$
 and $g: b \to c^T$ then $*(fg) = f^T * g$
K2. If $f: a \to b^T$ and $g: b \to c$ then $*(fg^T) = *fg^T$

K3. If $f: a \to b^T$ and $g: b \to c^T$ then *(f*g) = *f*g

Proof. Assume first that the natural transformation $\mu : TT \to T$ defines a semigroupad structure, that is, $T\mu \cdot \mu = \mu T \cdot \mu$. The same way as for the Kleisli lifting for monads, for an arrow $f : a \to b^T$ we can define

$$f := f^T \cdot b^{\mu}$$
.

K1. If $f: a \to b$ and $g: b \to c^T$ then $*(fg) = f^T g^T c^\mu = f^T *g$.

K2. If $f: a \to b^T$ and $g: b \to c$ then, by naturality of μ ,

$${}^{*}(f g^{T}) = f^{T} g^{TT} c^{\mu} = f^{T} b^{\mu} g^{T} = {}^{*}f g^{T}.$$

K3. If $f: a \to b^T$ and $g: b \to c^T$ then, substituting $*g = g^T c^{\mu}$ and applying $\mu T \cdot \mu = T \mu \cdot \mu$ and naturality of μ , we obtain

$${}^{*}(f \, {}^{*}g) = f^{T} g^{TT} c^{\mu T} c^{\mu} = f^{T} g^{TT} c^{T\mu} c^{\mu} = f^{T} b^{\mu} g^{T} c^{\mu} = {}^{*}f \, {}^{*}g$$

For the converse direction, assume a Kleisli lifting is given. For any object $a \in \operatorname{Ob} \mathcal{C}$ define

$$a^{\mu} := (1_{a^T}) : a^{TT} \to a^T.$$

This is a natural transformation $TT \to T$ because for any arrow $f: a \to b$ in $\mathcal C$ we have

$$f^{TT} b^{\mu} = f^{TT} * (1_{b^{T}}) \stackrel{\mathbf{K1.}}{=} * (f^{T} 1_{b^{T}}) = * (1_{a^{T}} f^{T}) \stackrel{\mathbf{K2.}}{=} * (1_{a^{T}}) f^{T} = a^{\mu} f^{T}.$$

Associativity of μ follows by

$$a^{T\mu} a^{\mu} = {}^{*}(1_{a^{TT}}) {}^{*}(1_{a^{T}}) \stackrel{\mathbf{K3.}}{=} {}^{*}(1_{a^{TT}} {}^{*}(1_{a^{T}})) = {}^{*}(a^{\mu})$$
$$a^{\mu T} a^{\mu} = a^{\mu T} {}^{*}(1_{a^{T}}) \stackrel{\mathbf{K1.}}{=} {}^{*}(a^{\mu} 1_{a^{T}}) = {}^{*}(a^{\mu}).$$

Finally, it is also straightforward to verify that these procedures are inverses to each other. $\hfill \Box$

Remark 4.4. As we can read from this proof, actually there is a one-to-one correspondence between all natural transformations $\mu : TT \to T$ and Kleisli * operations satisfying **K1**. and **K2**.

Remark 4.5. The Kleisli category construction for a monad still gives rise to a *semicategory* structure (i.e., a category without the requirement of identity arrows), when the monad is replaced by a semigroupad. The proof relies on **K3**.

The following theorem, inspired by (...), is a generalization of the celebrated property of the ultrafilter monad $\beta : Set \to Set$ that if S is a semigroup, then S^{β} also admits a semigroup structure, extending the original one. It turns out that this property extends to all monads on Set and even to semigroupads.

Theorem 4.6. Let T be a semigroup d on Set and let S be a semigroup, then there is an induced semigroup operation on S^T .

To achieve this, we introduce the following notation.

For an element $y \in S$, let $(\cdot y)$ denote the map $S \to S$ sending $x \mapsto xy$, and for any $p \in S^T$ we define

$$\bar{p}: S \to S^T, x \mapsto p(\cdot x)^T$$

so that $x \bar{p}$, the image of $x \in S$ under \bar{p} , is defined to be the image of $p \in S^T$ under the map $(\cdot x)^T : S^T \to S^T$. Finally, set

$$\widetilde{p} := \ ^* \! \bar{p} \ : \ S^T \to S^T$$

and define the operation for $p, q \in S^T$ as

$$p \bullet q := q \widetilde{p}.$$

Lemma 4.7. Under the hypothesis of the theorem, for any $y \in S$ and $p \in S^T$ we have

$$(\cdot y) \,\overline{p} = \overline{p} \, (\cdot y)^T \\ (\cdot y)^T \,\widetilde{p} = \widetilde{p} \, (\cdot y)^T$$

Proof. Applying both sides of the first equation to any element $x \in S$ and using associativity of the operation of S, we get

$$x(\cdot y)\bar{p} = (xy)\bar{p} = p(\cdot (xy))^T = p(\cdot x)^T(\cdot y)^T = x\bar{p}(\cdot y)^T.$$

Then the second equation is obtained by Kleisli lifting the first one:

$$(\cdot y)^T \widetilde{p} = (\cdot y)^T * \overline{p} \stackrel{\mathbf{K1.}}{=} * ((\cdot y) \overline{p}) = * (\overline{p} (\cdot y)^T) \stackrel{\mathbf{K2.}}{=} * \overline{p} (\cdot y)^T = \widetilde{p} (\cdot y)^T.$$

Lemma 4.8. Under the hypothesis of the theorem, for any $p, q \in S^T$ we have

$$p \bullet q = \widetilde{q} \, \widetilde{p}$$

Proof. Using the previous lemma, for any $y \in S$ we have

$$y\left(\overline{p \bullet q}\right) = (p \bullet q) (\cdot y)^T = q \widetilde{p} (\cdot y)^T = q (\cdot y)^T \widetilde{p} = y \overline{q} \widetilde{p}$$

yielding $\overline{p \bullet q} = \overline{q} \, \widetilde{p}$, so by **K3.** we arrive to $\widetilde{p \bullet q} = \widetilde{q} \, \widetilde{p}$.

Proof of theorem 4.6. We show that the operation \bullet on S^T is associative. For any $p, q, r \in S^T$, by lemma 4.8 we have

$$(p \bullet q) \bullet r = r\left(\widetilde{p \bullet q}\right) = r \widetilde{q} \widetilde{p} = (q \bullet r) \widetilde{p} = p \bullet (q \bullet r).$$

Definition 4.9. Let (T, μ) be a semigroupad on a category \mathcal{C} . A natural transformation $\xi : 1_{\mathcal{C}} \to T$ is called a

- left identity if $\xi T \cdot \mu = 1_T$,
- right identity if $T\xi \cdot \mu = 1_T$,
- central constant if $\xi T \cdot \mu = T \xi \cdot \mu$,
- idempotent constant if $\xi \cdot \xi T \cdot \mu = \xi$.

We note that any natural transformation $\xi : 1_{\mathcal{C}} \to T$ satisfies $\xi \cdot \xi T = \xi \cdot T\xi$, so ξ is an idempotent constant if and only if $\xi \cdot T\xi \cdot \mu = \xi$.

Proposition 4.10. Let (T, μ) be a semigroupad on a category \mathcal{C} with Kleisli lifting $f \mapsto {}^*f$, and let $\xi : 1_{\mathcal{C}} \to T$ be a natural transformation. Then the following statements hold:

1. ξ is a left identity if and only if for each $a \in Ob \mathcal{C}$ we have

$$(a^{\xi}) = 1_{a^{T}}$$

2. ξ is a right identity if and only if for each $f: a \to b^T$ we have

$$a^{\xi} \cdot {}^*\!f = f.$$

3. ξ is a central constant if and only if for each $f:a\to b^T$ we have

$$a^{\xi} \cdot {}^*f = f \cdot {}^*(b^{\xi}).$$

4. ξ is an idempotent constant if and only if for each $a \in Ob \mathcal{C}$ we have

$$a^{\xi} \cdot {}^*\!(a^{\xi}) = a^{\xi}.$$

Proof. Recall that for an arrow $f: a \to b^T$ we have $*f = f^T b^{\mu}$.

- 1. Obvious.
- 2. If ξ is a right identity, then by naturality of ξ we get

$$a^{\xi} \cdot {}^*f = a^{\xi} \cdot f^T \cdot b^{\mu} = f \cdot b^{T\xi} \cdot b^{\mu} = f.$$

Conversely, apply the hypothesis to $f = 1_{a^T} : a^T \to a^T$ to get

$$1_{a^T} = a^{T\xi} \cdot {}^*\!(1_{a^T}) = a^{T\xi} \cdot a^{\mu}.$$

3. If ξ is central, then similarly as in 2., we have

$$a^{\xi} \cdot {}^*f = f \cdot b^{T\xi} \cdot b^{\mu} = f \cdot b^{\xi T} \cdot b^{\mu} = f \cdot {}^*(b^{\xi}).$$

Conversely, applying again the hypothesis to $f = 1_{a^T}$, we arrive at

$$a^{T\xi} \cdot a^{\mu} = a^{T\xi} \cdot {}^{*}(1_{a^{T}}) = 1_{a^{T}} \cdot {}^{*}(a^{\xi}) = a^{\xi T} \cdot a^{\mu}.$$

4. This readily follows from

$$a^{\xi} \cdot {}^*\!(a^{\xi}) = a^{\xi} \cdot a^{\xi T} \cdot a^{\mu}$$

Theorem 4.11. Assume a semigroupad T on Set is given with Kleisli lifting $f \mapsto {}^*f$, let $\xi : 1_{Set} \to T$ be a natural transformation, and let (S, \cdot) be a semigroup with a distinguished element $e \in S$. As a shorthand, we write $\boldsymbol{\xi}$ for $S^{\boldsymbol{\xi}} : S \to S^T$ below.

Then the following statements hold:

- 1. If ξ is a left identity for T and e is a left identity in S, then e^{ξ} is a left identity in the induced semigroup on S^T .
- 2. If ξ is a right identity for T and e is a right identity in S, then e^{ξ} is a right identity in the induced semigroup on S^T .
- 3. If ξ is a central constant for T and e is a central element in S, then e^{ξ} is a central element in the induced semigroup on S^T .
- 4. If ξ is an idempotent constant for T and e is an idempotent element in S, then e^{ξ} is an idempotent element in the induced semigroup on S^{T} .

Proof. Let $p \in S^T$ be an arbitrary element. Note that by naturality we have $f \boldsymbol{\xi} = \boldsymbol{\xi} f^T$ for any $f: S \to S$.

1. $e^{\boldsymbol{\xi}} \bullet p = p \ \widetilde{e^{\boldsymbol{\xi}}} = p \ ^{\boldsymbol{\xi}} (x \mapsto e^{\boldsymbol{\xi}} (\cdot x)^T) = p \ ^{\boldsymbol{\xi}} (x \mapsto e (\cdot x) \boldsymbol{\xi}) = p \ ^{\boldsymbol{\xi}} \boldsymbol{\xi} = p.$ 2. $p \bullet e^{\boldsymbol{\xi}} = e^{\boldsymbol{\xi}} \ \widetilde{p} = e^{\boldsymbol{\xi}} \ ^{\boldsymbol{\xi}} (x \mapsto p (\cdot x)^T) = e \ (x \mapsto p (\cdot x)^T) = p \ (\cdot e)^T = p.$

3.
$$e^{\boldsymbol{\xi}} \bullet p = p^{\ast}(x \mapsto (ex)^{\boldsymbol{\xi}}) = p^{\ast}(x \mapsto (xe)^{\boldsymbol{\xi}}) = p^{\ast}((\cdot e)\,\boldsymbol{\xi}),$$

 $p \bullet e^{\boldsymbol{\xi}} = e^{\boldsymbol{\xi}}^{\ast}(x \mapsto p(\cdot x)^{T}) = e(x \mapsto p(\cdot x)^{T})^{\ast}\boldsymbol{\xi} = p(\cdot e)^{T}^{\ast}\boldsymbol{\xi} = p^{\ast}((\cdot e)\,\boldsymbol{\xi}).$

4.
$$e^{\boldsymbol{\xi}} \bullet e^{\boldsymbol{\xi}} = e^{\boldsymbol{\xi}} * (x \mapsto e^{\boldsymbol{\xi}} (\cdot x)^T) = e^{\boldsymbol{\xi}} * (x \mapsto (ex)^{\boldsymbol{\xi}}) = e^{\boldsymbol{\xi}} * ((e \cdot) \boldsymbol{\xi}) = e^{\boldsymbol{\xi}} * (\boldsymbol{\xi} (e \cdot)^T) = e^{\boldsymbol{\xi}} * \boldsymbol{\xi} (e \cdot)^T = e^{\boldsymbol{\xi}} (e \cdot)^T = e^{\boldsymbol{\xi}} (e \cdot)^T = e^{\boldsymbol{\xi}}.$$

References

- [1] [A. Costa, B. Steinberg, 2014] The Schützenberger category of a semigroup, arXiv:1408.1615
- [2] [S. Hayashi, 1985] Adjunction of semifunctors: categorical structures in nonextensional lambda calculus, *Theoretical Computer Science* 41