

Exceptional Topology of Non-Hermitian Brillouin Klein Bottles

W. B. Rui^{1,*} and Z. D. Wang^{1,†}

¹*Department of Physics and HK Institute of Quantum Science & Technology,
The University of Hong Kong, Pokfulam Road, Hong Kong, China*

(Dated: March 11, 2025)

Exceptional points (EPs) are prominent non-Hermitian band degeneracies that give rise to a variety of intriguing and unconventional phenomena. Similar to Weyl and Dirac points, EPs carry topological charges and comply with the celebrated fermion doubling theorems in lattices. Beyond these characteristics, EPs exhibit more exotic topological properties, particularly non-Abelian braiding topologies not seen in conventional degeneracies. Here, we investigate these foundational concepts of EPs in two-dimensional non-Hermitian lattices where the fundamental domain of the Brillouin zone is a Klein bottle, rather than a torus assumed in previous studies. We find that EPs do not necessarily appear in pairs with opposite topological charges in the Brillouin Klein bottle, thus violating the fermion doubling theorem. The violation occurs because, without crossing the boundary, the sum of the topological charges of EPs is in fact an even number rather than zero. Moreover, we uncover unique braiding topologies of EPs that cannot be captured by existing theories. Specifically, the composite braidings around all EPs equals the braiding along the boundary of the Brillouin Klein bottle. This novel braiding topology further confirms the failure of the fermion doubling theorem, and allows us to explore the non-Abelian braidings of EPs beyond the scope of topological charges. Our work highlights the fundamental role of Brillouin-zone topology in non-Hermitian systems.

Introduction.— Exceptional points (EPs) represent one of the most intriguing aspects of non-Hermitian physics, drawing significant attention across diverse fields such as optics, photonics, acoustics, electronics, and condensed matters [1–16]. At EPs, both eigenvalues and eigenvectors coalesce [17–19], leading to a variety of unconventional physical phenomena, including enhanced sensitivity [9–12] and unidirectional invisibility [13–16]. In the rapidly growing field of non-Hermitian topology [20–40], the importance of EPs has been further highlighted. These unique non-Hermitian degeneracies serve as defining characteristics of a large class of non-Hermitian topological semimetals, known as exceptional semimetals [41–51].

Like Weyl and Dirac points, EPs carry topological charges and are subject to the celebrated fermion doubling theorem, a universal no-go theorem governing both Hermitian and non-Hermitian lattice systems [52–55]. This theorem dictates that EPs must appear in pairs with opposite topological charges in lattices. The underlying proof relies on the periodic boundary conditions of the Brillouin torus [55], which ensure that the total topological charge of EPs—equal to a line integral along the torus boundary—sums to zero.

While EPs share these fundamental features with conventional band degeneracies, they also exhibit unique and extraordinary topological characteristics, particularly non-Abelian braiding topologies [56–64]. These braiding topologies arise because the complex eigenenergies in the vicinity of EPs can braid around each other, and become non-Abelian when there are more than two eigenenergies. The braiding topology of EPs in lattices follows specific rules [57, 59]: the composite braidings around all EPs must match the braiding along the bound-

ary of the Brillouin torus.

These foundational concepts serve as the cornerstone for studying EPs in lattices. However, they have been developed under the assumption that the Brillouin zone is topologically a torus. Recent discoveries have shown that, under momentum-space nonsymmorphic symmetries, the fundamental domain of the Brillouin zone forms a Klein bottle rather than a torus [65–72]. This raises an important question: Do the fermion doubling theorem and the established braiding topology of EPs, both of which depend on torus boundary conditions, still hold true in the Brillouin Klein bottle?

In this work, we demonstrate that the fermion doubling theorem breaks down, and a unique braiding topology of EPs emerges in the non-Hermitian Brillouin Klein bottle. We focus on the region without crossing the Klein bottle boundary, denoted as K^2 , where a local orientation can be defined to avoid the sign ambiguity of topological charges due to the global non-orientability of the Klein bottle. In this region, the topological charges of EPs maintain definite signs and behave similarly to conventional ones. We find that the sum of all topological charges within K^2 equals a line integral along the Klein bottle boundary ∂K^2 , and satisfies

$$\sum_{\mathbf{k}_i \in K^2} v(\mathbf{k}_i) = \oint_{\partial K^2} d\mathbf{k} \cdot \nabla_{\mathbf{k}} \log \Delta(\mathbf{k}) \in 2\mathbb{Z}, \quad (1)$$

where $v(\mathbf{k}_i)$ is the topological charge of the EP located at \mathbf{k}_i , and $\Delta(\mathbf{k})$ is the discriminant defined below Eq. (4). Since the total charge is an even number rather than zero, EPs do not necessarily appear in pairs with opposite charges, leading to the breakdown of the fermion doubling theorem. Furthermore, we establish the braiding topology of EPs in the Brillouin Klein bottle, and

find that the previously established braiding rules do not apply. Instead, the composite of the braidings around all EPs in K^2 , denoted as $(\mathbf{b}_1 \mathbf{b}_2 \cdots \mathbf{b}_n)$, equals to the braiding along the boundary ∂K^2 , expressed as $(\mathbf{b}_a \mathbf{b}_b \mathbf{b}_a \mathbf{b}_b^{-1})$. That is,

$$\mathbf{b}_1 \mathbf{b}_2 \cdots \mathbf{b}_n = \mathbf{b}_a \mathbf{b}_b \mathbf{b}_a \mathbf{b}_b^{-1}, \quad (2)$$

which can further confirm the breakdown of the fermion doubling theorem. Finally, we explore the non-Abelian braidings associated with EPs, revealing rich topologies that cannot be characterized by topological charges alone.

Brillouin Klein Bottle in non-Hermitian systems.—

Let us consider a non-Hermitian Hamiltonian $\mathcal{H}(\mathbf{k})$ in two dimensions that respects the momentum-space glide reflection symmetry as [65]

$$U\mathcal{H}(k_x, k_y)U^{-1} = \mathcal{H}(-k_x, k_y + \pi), \quad (3)$$

where U is a unitary operator. The symmetry maps (k_x, k_y) to $(-k_x, k_y + \pi)$, partitioning the first Brillouin zone into two equivalent regions: $(k_x, k_y) \in [-\pi, \pi) \times [-\pi, 0)$ and $[-\pi, \pi) \times [0, \pi)$. Hence, it suffices to consider one of these regions, e.g., $[-\pi, \pi) \times [-\pi, 0)$ as plotted in Fig. 1(a).

Examining the two horizontal edges (red lines) of this region at $k_y = -\pi$ and 0 , we find that they must be glued together in opposite directions due to the symmetry $U\mathcal{H}(k_x, -\pi)U^{-1} = \mathcal{H}(-k_x, 0)$. In contrast, the two vertical edges (blue lines) should be glued together in the same direction. This specific edge identification results in a Klein bottle rather than a torus, which is termed the Brillouin Klein bottle.

EPs in Brillouin Klein Bottle.— EPs carry topological charges known as discriminant numbers, which are defined as [22, 55],

$$v(\mathbf{k}_i) = \frac{i}{2\pi} \oint_{\Gamma(\mathbf{k}_i)} d\mathbf{k} \cdot \nabla_{\mathbf{k}} \log \Delta(\mathbf{k}). \quad (4)$$

where $\Delta(\mathbf{k}) = \prod_{j < k} [E_j(\mathbf{k}) - E_k(\mathbf{k})]^2$ is the discriminant, and $E_j(\mathbf{k})$ is the j th eigenvalue of $\mathcal{H}(\mathbf{k})$. It is important to emphasize that $\Gamma(\mathbf{k}_i)$ is a small loop that encircles the EP at \mathbf{k}_i in a *counterclockwise* orientation.

One might try to directly use the discriminant number (4) for EPs in the Brillouin Klein bottle, as shown in Fig. 1(a). However, a fundamental challenge arises: the Klein bottle is globally *non-orientable*, meaning that a consistent counterclockwise or clockwise orientation cannot be maintained over the entire surface. As highlighted by process "I" in Fig. 1(b), when an EP and its associated loop traverse the twist at $k_y = 0 (-\pi)$ (red arrow) and return to its original position, the counterclockwise orientation is reversed to a clockwise orientation. Consequently, the sign of the invariant becomes ambiguous; i.e., $v(\mathbf{k}_i)$ cannot be distinguished from $-v(\mathbf{k}_i)$ over the entire




Fig. 1. (a) Depiction of the Brillouin Klein Bottle, showing EPs (red and blue dots) and the oriented loop Γ used to define the topological charge. (b) Illustration of the global non-orientability of the Brillouin Klein Bottle, where the twist is introduced after gluing the red edges in (a). (c) In the gray region, the loop Γ can be continuously deformed to two lines L_1 and L_2 . (d1-d3) represent the continuous deformation of integration paths to the boundary of Brillouin Klein bottle.

surface. This phenomenon is also observed for the Chern number in Brillouin Klein bottles [65, 68]. Therefore, the global non-orientability poses a challenge for verifying the fermion doubling theorem regarding the pairing of EPs with opposite topological charges.

Local versus global orientability.— In non-orientable manifolds, while a global orientation cannot be established, it is actually feasible to define a local orientation [68, 73]. Here, we focus on the region without crossing the Klein bottle boundary, denoted as K^2 as shown by the gray area in Fig. 1, so that the twist at $k_y = 0 (-\pi)$ can be avoided. Within this region, a counterclockwise orientation can still be meaningfully defined, and the sign of topological charges does not change when moving EPs, as illustrated by process "II" in Fig. 1(b). Thus, the discriminant number from Eq. (4) has definite signs within K^2 , and behaves just like those in a Brillouin torus.

Furthermore, as shown in Fig. 1(c), within the region K^2 , the oriented loop Γ can be deformed to $\tilde{\Gamma}$ and further splitted into two 1D closed paths denoted as L_1 and

$-L_2$, due to topological robustness. Consequently, the topological charge carried by EPs can be computed by the difference between the discriminant number of the two paths at different k_y values, which is given by

$$v(k_y) = \frac{i}{2\pi} \int_{-\pi}^{\pi} dk_x \cdot \partial_{k_x} \log \Delta(k_x, k_y). \quad (5)$$

Failure of fermion doubling theorem.— After specifying the local orientation within the region K^2 , the summation of topological charges for all EPs in K^2 can be unambiguously computed as

$$\sum_{\mathbf{k}_i \in K^2} v(\mathbf{k}_i) = \oint_{\partial K^2} d\mathbf{k} \cdot \nabla_{\mathbf{k}} \log \Delta(\mathbf{k}), \quad (6)$$

as the integration paths can be continuously deformed to the Klein bottle boundary ∂K^2 , a process illustrated in Figs. 1(d₁-d₃) similar to the torus case [55].

Given that the boundary $\partial K^2 = abab^{-1}$, shown in Fig. 1(d₃), the integration cancels out on the two b edges, while it adds up on the two a edges. Hence, we obtain

$$\sum_{\mathbf{k}_i \in K^2} v(\mathbf{k}_i) = 2 \oint_a d\mathbf{k} \cdot \nabla_{\mathbf{k}} \log \Delta(\mathbf{k}) \in 2\mathbb{Z}, \quad (7)$$

since the integration along loop a takes \mathbb{Z} values. By combining the above two equations, we arrive at Eq. (1) in the introduction. Therefore, we can see that the fermion doubling theorem for EPs fails in the Brillouin Klein bottle, as the summation equals an even number. Note that a similar proof can be done for Fermi points, as discussed in the Supplemental Material (SM) [74].

A concrete example violating the fermion doubling theorem.— Let us now investigate a two-band model in the non-Hermitian Brillouin Klein bottle, characterized by the Hamiltonian

$$\mathcal{H}_2(\mathbf{k}) = (\alpha \cos k_x - \beta) \sigma_1 + (\alpha \sin k_x - i\gamma \cos k_y) \sigma_2, \quad (8)$$

where α , β , and γ are model parameters, and σ_i 's are Pauli matrices. The model respects the symmetry of Eq. (3) with $U = \sigma_1$. We propose the experimental realization of this model in the SM [74].

As depicted in Fig. 2(a), there are two EPs located at the position where both $\text{Re} \Delta(\mathbf{k})$ and $\text{Im} \Delta(\mathbf{k})$ vanish [74]. The two EPs both have a discriminant number of $+1$, thereby violating the fermion doubling theorem. This is evident by computing the k_y -resolved topological invariant $v(k_y)$ of Eq. (5). As shown in Fig. 2(b), $v(k_y)$ both increases by $+1$ when crossing both EPs by decreasing k_y , due to their identical topological charge of $+1$.

Braiding topology of EPs in Brillouin Klein Bottles.— Recently, it has been revealed that EPs exhibit a richer braiding topology that cannot be fully captured by the discriminant number [56–62]. To explore this, we start




Fig. 2. (a) A two-band model where two EPs both having positive topological charges. EPs are located at the position where $\text{Re} \Delta(\mathbf{k}) = \text{Im} \Delta(\mathbf{k}) = 0$. (b) The k_y -resolved topological invariants calculated using Eq. (5). (c) The braiding topology of EPs in the Brillouin Klein Bottle. The parameters are $\alpha = 0.5$, $\beta = 1.0 - 0.25i$, and $\gamma = 1.0$. The numerical calculations are given in the SM [74].

by treating EPs as punctures in the Brillouin Klein bottle K^2 [75]. The set of a total of n EPs, denoted as $P = \{\mathbf{k}_1, \mathbf{k}_2, \dots, \mathbf{k}_n\}$, is isolated from K^2 to ensure there is no eigenvalue degeneracy on the n -punctured Klein bottle $K^2 - P$. In this configuration, the N complex eigenvalues $\{E_1, \dots, E_N\}$ are distinct and belong to the unordered configuration space $\text{UConf}_N(\mathbb{C})$ [56, 59]. To reveal the braiding topology, we consider the set of homotopy classes of maps from $K^2 - P$ to $\text{UConf}_N(\mathbb{C})$, denoted by $[K^2 - P, \text{UConf}_N(\mathbb{C})]$. For $\text{UConf}_N(\mathbb{C})$, only its fundamental group is non-trivial and is equal to the braid group, $\pi_1(\text{UConf}_N(\mathbb{C})) = B_N$, while all higher homotopy groups are trivial. Such a space is called an Eilenberg-MacLane space of type $K(G, 1)$, with $G = B_N$. An important property of this space is that there exists a natural bijection [59, 73]

$$[K^2 - P, \text{UConf}_N(\mathbb{C})] = \text{Hom}(\pi_1(K^2 - P), B_N). \quad (9)$$

Hence, we need to compute the set $\text{Hom}(\pi_1(K^2 - P), B_N)$ of the group homomorphisms from $\pi_1(K^2 - P)$ to B_N .

The key difference from the Brillouin torus case is that the base manifold is now the n -punctured Klein bottle $K^2 - P$. Its fundamental group is $\pi_1(K^2 - P) =$

$\{a, b, \Gamma_1, \Gamma_2, \dots, \Gamma_n | abab^{-1} = \Gamma_1 \Gamma_2 \dots \Gamma_n\}$, as calculated in the SM [74]. Here, two generators are from \mathcal{K}^2 (specifically, loops a and b on the edges) and n generators are from the loops surrounding the EPs, denoted as $\Gamma_i = \Gamma(\mathbf{k}_i)$ around the i th EP. There are $(n+2)$ generators constrained by the relation

$$abab^{-1} = \Gamma_1 \Gamma_2 \dots \Gamma_n, \quad (10)$$

which can be understood as the continuous deformation of the loops $\Gamma_1, \dots, \Gamma_n$ to the boundary of \mathcal{K}^2 , a process illustrated in Figs. 1(d₁-d₃). Thus, $\pi_1(\mathcal{K}^2 - P)$ is a free group of $n+1$ generators $\pi_1(\mathcal{K}^2 - P) = *_{n+1}\mathbb{Z}$ [76].

Next, we compute the set of group homomorphisms $\text{Hom}(\pi_1(\mathcal{K}^2 - P), B_N)$. A group homomorphism is a function $f: \pi_1(\mathcal{K}^2 - P) \rightarrow B_N$, which maps loops γ_i in $\mathcal{K}^2 - P$ to braid elements b_i in the braid group B_N . By definition, the group homomorphism preserves the group structure, i.e., $f(\gamma_1 \gamma_2) = f(\gamma_1) f(\gamma_2)$, and is determined by its values on the generators of $\pi_1(\mathcal{K}^2 - P)$. We denote the values of the homomorphism on the generators a and b as $f(a) = \mathbf{b}_a$ and $f(b) = \mathbf{b}_b$, respectively, and on the generators Γ_i as $f(\Gamma_i) = \mathbf{b}_i$. Furthermore, the homomorphism must preserve the relation given by Eq. (10). Thus, we obtain the braiding topology as

$$\text{Hom}(\pi_1(\mathcal{K}^2 - P), B_N) = \{\mathbf{b}_a, \mathbf{b}_b, \mathbf{b}_1, \dots, \mathbf{b}_n \in B_N | \mathbf{b}_1 \dots \mathbf{b}_n = \mathbf{b}_a \mathbf{b}_b \mathbf{b}_a \mathbf{b}_b^{-1}\}, \quad (11)$$

which endorses a no-go theorem on the braiding patterns of EPs in the Brillouin Klein bottle. That is, the composite braidings around all EPs $\mathbf{b}_1 \mathbf{b}_2 \dots \mathbf{b}_n$ must equal to the braiding $\mathbf{b}_a \mathbf{b}_b \mathbf{b}_a \mathbf{b}_b^{-1}$ along the Klein bottle boundary.

As shown in Fig. 2(c) and Fig. 3(c), the braiding patterns can be represented by a sequence of over and under crossings, after sorting the eigenenergies by their real parts. Specifically, τ_i denotes where the i th eigenenergy crosses under the $(i+1)$ th eigenenergy, while τ_i^{-1} denotes where it crosses over.

Revisiting the failure of fermion doubling theorem.— The failure of the fermion doubling theorem can also be proven from the braiding topology. This is because the discriminant number is determined by braid crossings: an over/under crossing contributes $+1/-1$ to the discriminant numbers [57]. As a result, the sum of the discriminant numbers of all EPs equals those along the Klein bottle boundary, using the relation $\mathbf{b}_1 \mathbf{b}_2 \dots \mathbf{b}_n = \mathbf{b}_a \mathbf{b}_b \mathbf{b}_a \mathbf{b}_b^{-1}$ in Eq. (11). Since the discriminant numbers of \mathbf{b}_b and \mathbf{b}_b^{-1} cancel out, we obtain the relation in Eq. (7), where the sum equals twice the discriminant numbers of \mathbf{b}_a .

Returning to the two-band model in Eq. (8), the braid group is B_2 for $N=2$, which is the abelian group \mathbb{Z} and all braid elements commute. As shown in Fig. 2(c), the braiding patterns on the loops Γ_1 and Γ_2 are $\mathbf{b}_1 = \mathbf{b}_2 = \tau_1$, while on the boundary, they are $\mathbf{b}_a = \tau_1$ and $\mathbf{b}_b = 1$. It can be verified that $\mathbf{b}_1 \mathbf{b}_2 = \mathbf{b}_a^2 = \tau_1^2$, in accordance with the braiding theory. The discriminant number for all EPs




Fig. 3. Non-abelian braidings. (a) A three-band model where four EPs all having positive topological charges. (b) The k_y -resolved topological invariants calculated using Eq. (5). (c) The braiding of EPs in the Brillouin Klein bottle. The parameters are $\alpha = 2.0$, $\beta = 1.0$, $\gamma = 1.0$, $\delta = 0.5$ and $\epsilon = 1.0$. The numerical calculations are given in the SM [74].

sums up to $+2$, with each τ_1 contributing $+1$, further confirming the failure of the fermion doubling theorem.

Non-abelian braidings in Brillouin Klein Bottle.— The intriguing aspect of braiding topology is that it becomes intrinsically non-abelian in multi-band cases with $N \geq 3$, which cannot be described by topological charges. In this case, the τ_i 's satisfy the braid relations $\tau_i \tau_j = \tau_j \tau_i$ for $|j - i| > 1$ and $\tau_i \tau_{i+1} \tau_i = \tau_{i+1} \tau_i \tau_{i+1}$ for $1 \leq i \leq N - 1$. Consider a three-band model, whose Hamiltonian is given by

$$\mathcal{H}_3(\mathbf{k}) = \begin{pmatrix} F(\mathbf{k}) & -1 & 0 \\ -1 & 0 & -1 \\ 0 & -1 & G(\mathbf{k}) \end{pmatrix}, \quad (12)$$

where $F(\mathbf{k}) = \alpha \cos k_x + i\beta \sin k_x \cos k_y + i\epsilon$ and $G(\mathbf{k}) = \gamma \cos 2k_y + i\delta \sin 2k_y - \epsilon$. The model satisfies the symmetry of Eq. (3) with $U = 1$. As shown in Fig. 3(a), there are four EPs all having topological charges of $+1$, thereby violating the fermion doubling theorem. This can be verified by the k_y -resolved topological invariant $v(k_y)$ of Eq. (5), as shown in Fig. 3(b).

As depicted in Fig. 3(c), the braiding patterns on the edges are $\mathbf{b}_a = \tau_1 \tau_2$ and $\mathbf{b}_b = \tau_1^{-1}$. They do not com-

mute, i.e., $\mathbf{b}_a\mathbf{b}_b \neq \mathbf{b}_b\mathbf{b}_a$, because $\mathbf{b}_a\mathbf{b}_b = \tau_1\tau_2\tau_1^{-1}$ is not equivalent to $\mathbf{b}_b\mathbf{b}_a = \tau_2$ using the aforementioned braid relations. Thus, the braiding topology cannot be reduced to $\mathbf{b}_1\mathbf{b}_2 \cdots \mathbf{b}_n = \mathbf{b}_a\mathbf{b}_b\mathbf{b}_a\mathbf{b}_b^{-1} \neq \mathbf{b}_a^2$, in contrast to the two-band case. On the other hand, the four EPs have the braidings $\mathbf{b}_1 = \tau_1$, $\mathbf{b}_2 = \tau_2$, $\mathbf{b}_3 = \tau_2$, and $\mathbf{b}_4 = \tau_1$ along the loops Γ_1 , Γ_2 , Γ_3 , and Γ_4 , respectively. It can be checked that $\mathbf{b}_a\mathbf{b}_b\mathbf{b}_a\mathbf{b}_b^{-1} = \mathbf{b}_1\mathbf{b}_2\mathbf{b}_3\mathbf{b}_4 = \tau_1\tau_2\tau_2\tau_1$ as shown in Fig. 3(c), in accordance with Eq. (11). The non-commutative braiding relations highlight the necessity of considering the full braid group structure to understand the topological properties of EPs in the Brillouin Klein bottle.

Conclusions and discussions. — We have demonstrated that the well-established fermion doubling theorem for EPs breaks down in the non-Hermitian Brillouin Klein bottle. Specifically, we have proven that the sum of the total charges is an even number rather than zero for the region without crossing the boundary. Moreover, we have uncovered a novel braiding topology for EPs, particularly the non-Abelian ones, which is distinct from that in the Brillouin torus. While our study has primarily focused on gapless non-Hermitian topologies, exploring gapped topological phases [29, 30] would be a promising direction for future research. Beyond bulk topologies, it would also be valuable to investigate boundary effects, such as non-Hermitian skin effects and topological boundary states [77, 78], in the Brillouin Klein bottle.

This work was supported by the Quantum Science Center of Guangdong-Hong Kong-Macau Greater Bay Area, the NSFC/RGC JRS grant (RGC Grant No. N_HKU774/21, NSFC Grant No. 12161160315), and the GRF of Hong Kong (Grant Nos. 17310622 and 17303023). W.B.R. was supported by the RGC Post-doctoral Fellowship (Ref. No. PDFS2223-7S05).

* wbrui@hku.hk

† zwang@hku.hk

- [1] M.-A. Miri and A. Alu, Exceptional points in optics and photonics, *Science* **363**, eaar7709 (2019).
- [2] Ş. K. Özdemir, S. Rotter, F. Nori, and L. Yang, Parity-time symmetry and exceptional points in photonics, *Nature Materials* **18**, 783 (2019).
- [3] T. Gao, E. Estrecho, K. Y. Bliokh, T. C. H. Liew, M. D. Fraser, S. Brodbeck, M. Kamp, C. Schneider, S. Höfling, Y. Yamamoto, F. Nori, Y. S. Kivshar, A. G. Truscott, R. G. Dall, and E. A. Ostrovskaya, Observation of non-Hermitian degeneracies in a chaotic exciton-polariton billiard, *Nature* **526**, 554 (2015).
- [4] J. Schindler, A. Li, M. C. Zheng, F. M. Ellis, and T. Kottos, Experimental study of active lrc circuits with \mathcal{PT} symmetries, *Phys. Rev. A* **84**, 040101 (2011).
- [5] A. Li, H. Wei, M. Cotrufo, W. Chen, S. Mann, X. Ni, B. Xu, J. Chen, J. Wang, S. Fan, C.-W. Qiu, A. Alù, and L. Chen, Exceptional points and non-Hermitian photonics at the nanoscale, *Nature Nanotechnology* **18**, 706 (2023).
- [6] L. Feng, Z. J. Wong, R.-M. Ma, Y. Wang, and X. Zhang, Single-mode laser by parity-time symmetry breaking, *Science* **346**, 972 (2014).
- [7] H. Hodaei, M.-A. Miri, M. Heinrich, D. N. Christodoulides, and M. Khajavikhan, Parity-time-symmetric microring lasers, *Science* **346**, 975 (2014).
- [8] B. Peng, Ş. K. Özdemir, S. Rotter, H. Yilmaz, M. Liertzer, F. Monifi, C. M. Bender, F. Nori, and L. Yang, Loss-induced suppression and revival of lasing, *Science* **346**, 328 (2014).
- [9] J. Wiersig, Enhancing the Sensitivity of Frequency and Energy Splitting Detection by Using Exceptional Points: Application to Microcavity Sensors for Single-Particle Detection, *Physical Review Letters* **112**, 203901 (2014).
- [10] W. Chen, Ş. Kaya Özdemir, G. Zhao, J. Wiersig, and L. Yang, Exceptional points enhance sensing in an optical microcavity, *Nature* **548**, 192 (2017).
- [11] H. Hodaei, A. U. Hassan, S. Wittek, H. Garcia-Gracia, R. El-Ganainy, D. N. Christodoulides, and M. Khajavikhan, Enhanced sensitivity at higher-order exceptional points, *Nature* **548**, 187 (2017).
- [12] Z.-P. Liu, J. Zhang, Ş. K. Özdemir, B. Peng, H. Jing, X.-Y. Lü, C.-W. Li, L. Yang, F. Nori, and Y.-x. Liu, Metrology with \mathcal{PT} -Symmetric Cavities: Enhanced Sensitivity near the \mathcal{PT} -Phase Transition, *Phys. Rev. Lett.* **117**, 110802 (2016).
- [13] A. Guo, G. J. Salamo, D. Duchesne, R. Morandotti, M. Volatier-Ravat, V. Aimez, G. A. Siviloglou, and D. N. Christodoulides, Observation of \mathcal{PT} -symmetry breaking in complex optical potentials, *Phys. Rev. Lett.* **103**, 093902 (2009).
- [14] Z. Lin, H. Ramezani, T. Eichelkraut, T. Kottos, H. Cao, and D. N. Christodoulides, Unidirectional Invisibility Induced by \mathcal{PT} -Symmetric Periodic Structures, *Phys. Rev. Lett.* **106**, 213901 (2011).
- [15] A. Regensburger, C. Bersch, M.-A. Miri, G. Onishchukov, D. N. Christodoulides, and U. Peschel, Parity-time synthetic photonic lattices, *Nature* **488**, 167 (2012).
- [16] B. Peng, Ş. K. Özdemir, F. Lei, F. Monifi, M. Gianfreda, G. L. Long, S. Fan, F. Nori, C. M. Bender, and L. Yang, Parity-time-symmetric whispering-gallery microcavities, *Nature Physics* **10**, 394 (2014).
- [17] T. Kato, Perturbation theory in a finite-dimensional space, in *Perturbation Theory for Linear Operators*, Classics in Mathematics, edited by T. Kato (Springer, Berlin, Heidelberg, 1995) pp. 62–126.
- [18] M. Berry, Physics of Nonhermitian Degeneracies, *Czechoslovak Journal of Physics* **54**, 1039 (2004).
- [19] W. D. Heiss, The physics of exceptional points, *J. Phys. A: Math. Theor.* **45**, 444016 (2012).
- [20] D. Leykam, K. Y. Bliokh, C. Huang, Y. D. Chong, and F. Nori, Edge modes, degeneracies, and topological numbers in non-hermitian systems, *Phys. Rev. Lett.* **118**, 040401 (2017).
- [21] Y. Xu, S.-T. Wang, and L.-M. Duan, Weyl exceptional rings in a three-dimensional dissipative cold atomic gas, *Phys. Rev. Lett.* **118**, 045701 (2017).
- [22] H. Shen, B. Zhen, and L. Fu, Topological band theory for non-hermitian hamiltonians, *Phys. Rev. Lett.* **120**,

- 146402 (2018).
- [23] H. Zhou, C. Peng, Y. Yoon, C. W. Hsu, K. A. Nelson, L. Fu, J. D. Joannopoulos, M. Soljacic, and B. Zhen, Observation of bulk fermi arc and polarization half charge from paired exceptional points, *Science* **359**, 1009 (2018).
- [24] S. Yao and Z. Wang, Edge states and topological invariants of non-hermitian systems, *Phys. Rev. Lett.* **121**, 086803 (2018).
- [25] F. K. Kunst, E. Edvardsson, J. C. Budich, and E. J. Bergholtz, Biorthogonal bulk-boundary correspondence in non-hermitian systems, *Phys. Rev. Lett.* **121**, 026808 (2018).
- [26] K. Zhang, Z. Yang, and C. Fang, Correspondence between winding numbers and skin modes in non-hermitian systems, *Phys. Rev. Lett.* **125**, 126402 (2020).
- [27] N. Okuma, K. Kawabata, K. Shiozaki, and M. Sato, Topological origin of non-hermitian skin effects, *Phys. Rev. Lett.* **124**, 086801 (2020).
- [28] W. B. Rui, M. M. Hirschmann, and A. P. Schnyder, Ptsymmetric non-hermitian dirac semimetals, *Phys. Rev. B* **100**, 245116 (2019).
- [29] Z. Gong, Y. Ashida, K. Kawabata, K. Takasan, S. Higashikawa, and M. Ueda, Topological phases of non-hermitian systems, *Phys. Rev. X* **8**, 031079 (2018).
- [30] K. Kawabata, K. Shiozaki, M. Ueda, and M. Sato, Symmetry and topology in non-hermitian physics, *Phys. Rev. X* **9**, 041015 (2019).
- [31] L. Li, C. H. Lee, S. Mu, and J. Gong, Critical non-Hermitian skin effect, *Nature Communications* **11**, 5491 (2020).
- [32] W. B. Rui, Y. X. Zhao, and A. P. Schnyder, Topology and exceptional points of massive dirac models with generic non-hermitian perturbations, *Phys. Rev. B* **99**, 241110 (2019).
- [33] W. B. Rui, Y. X. Zhao, and Z. D. Wang, Hermitian topologies originating from non-hermitian braidings, *Phys. Rev. B* **108**, 165105 (2023).
- [34] K. Ding, C. Fang, and G. Ma, Non-Hermitian topology and exceptional-point geometries, *Nature Reviews Physics* **4**, 745 (2022).
- [35] W. B. Rui, Y. X. Zhao, and Z. D. Wang, Making topologically trivial non-hermitian systems nontrivial via gauge fields, *Phys. Rev. Lett.* **131**, 176402 (2023).
- [36] H. Hu and E. Zhao, Knots and non-hermitian bloch bands, *Phys. Rev. Lett.* **126**, 010401 (2021).
- [37] K. Yokomizo and S. Murakami, Non-bloch band theory of non-hermitian systems, *Phys. Rev. Lett.* **123**, 066404 (2019).
- [38] T. Helbig, T. Hofmann, S. Imhof, M. Abdelghany, T. Kiessling, L. W. Molenkamp, C. H. Lee, A. Szameit, M. Greiter, and R. Thomale, Generalized bulk-boundary correspondence in non-Hermitian topoelectrical circuits, *Nature Physics* **16**, 747 (2020).
- [39] L. Xiao, T. Deng, K. Wang, G. Zhu, Z. Wang, W. Yi, and P. Xue, Non-Hermitian bulk-boundary correspondence in quantum dynamics, *Nature Physics* **16**, 761 (2020).
- [40] Y. Ashida, Z. Gong, and M. Ueda, Non-Hermitian physics, *Advances in Physics* **69**, 249 (2020).
- [41] K. Kawabata, T. Bessho, and M. Sato, Classification of exceptional points and non-hermitian topological semimetals, *Phys. Rev. Lett.* **123**, 066405 (2019).
- [42] E. J. Bergholtz, J. C. Budich, and F. K. Kunst, Exceptional topology of non-hermitian systems, *Rev. Mod. Phys.* **93**, 015005 (2021).
- [43] J. Carlström and E. J. Bergholtz, Exceptional links and twisted fermi ribbons in non-hermitian systems, *Phys. Rev. A* **98**, 042114 (2018).
- [44] T. Yoshida, R. Peters, N. Kawakami, and Y. Hatsugai, Symmetry-protected exceptional rings in two-dimensional correlated systems with chiral symmetry, *Phys. Rev. B* **99**, 121101 (2019).
- [45] X. Zhang, K. Ding, X. Zhou, J. Xu, and D. Jin, Experimental observation of an exceptional surface in synthetic dimensions with magnon polaritons, *Phys. Rev. Lett.* **123**, 237202 (2019).
- [46] Z. Yang, C.-K. Chiu, C. Fang, and J. Hu, Jones polynomial and knot transitions in hermitian and non-hermitian topological semimetals, *Phys. Rev. Lett.* **124**, 186402 (2020).
- [47] W. Tang, X. Jiang, K. Ding, Y.-X. Xiao, Z.-Q. Zhang, C. T. Chan, and G. Ma, Exceptional nexus with a hybrid topological invariant, *Science* **370**, 1077 (2020).
- [48] I. Mandal and E. J. Bergholtz, Symmetry and higher-order exceptional points, *Phys. Rev. Lett.* **127**, 186601 (2021).
- [49] T. Liu, J. J. He, Z. Yang, and F. Nori, Higher-order weyl-exceptional-ring semimetals, *Phys. Rev. Lett.* **127**, 196801 (2021).
- [50] W. Tang, K. Ding, and G. Ma, Realization and topological properties of third-order exceptional lines embedded in exceptional surfaces, *Nature Communications* **14**, 6660 (2023).
- [51] Y. Wu, Y. Wang, X. Ye, W. Liu, Z. Niu, C.-K. Duan, Y. Wang, X. Rong, and J. Du, Third-order exceptional line in a nitrogen-vacancy spin system, *Nature Nanotechnology* **19**, 160 (2024).
- [52] H. B. Nielsen and M. Ninomiya, A no-go theorem for regularizing chiral fermions, *Physics Letters B* **105**, 219 (1981).
- [53] H. B. Nielsen and M. Ninomiya, Absence of neutrinos on a lattice: (I). Proof by homotopy theory, *Nuclear Physics B* **185**, 20 (1981).
- [54] H. B. Nielsen and M. Ninomiya, Absence of neutrinos on a lattice: (II). Intuitive topological proof, *Nuclear Physics B* **193**, 173 (1981).
- [55] Z. Yang, A. P. Schnyder, J. Hu, and C.-K. Chiu, Fermion doubling theorems in two-dimensional non-hermitian systems for fermi points and exceptional points, *Phys. Rev. Lett.* **126**, 086401 (2021).
- [56] Z. Li and R. S. K. Mong, Homotopical characterization of non-hermitian band structures, *Phys. Rev. B* **103**, 155129 (2021).
- [57] H. Hu, S. Sun, and S. Chen, Knot topology of exceptional point and non-hermitian no-go theorem, *Phys. Rev. Res.* **4**, L022064 (2022).
- [58] C.-X. Guo, S. Chen, K. Ding, and H. Hu, Exceptional non-abelian topology in multiband non-hermitian systems, *Phys. Rev. Lett.* **130**, 157201 (2023).
- [59] C. C. Wojcik, K. Wang, A. Dutt, J. Zhong, and S. Fan, Eigenvalue topology of non-hermitian band structures in two and three dimensions, *Phys. Rev. B* **106**, L161401 (2022).
- [60] C. C. Wojcik, X.-Q. Sun, T. c. v. Bzdušek, and S. Fan, Homotopy characterization of non-hermitian hamiltonians, *Phys. Rev. B* **101**, 205417 (2020).
- [61] Y. S. S. Patil, J. Höller, P. A. Henry, C. Guria, Y. Zhang, L. Jiang, N. Kralj, N. Read, and J. G. E. Harris, Measuring the knot of non-Hermitian degeneracies and non-

- commuting braids, [Nature](#) **607**, 271 (2022).
- [62] J. L. K. König, K. Yang, J. C. Budich, and E. J. Bergholtz, Braid-protected topological band structures with unpaired exceptional points, [Phys. Rev. Res.](#) **5**, L042010 (2023).
- [63] K. Wang, A. Dutt, C. C. Wojcik, and S. Fan, Topological complex-energy braiding of non-Hermitian bands, [Nature](#) **598**, 59 (2021).
- [64] Q. Zhang, Y. Li, H. Sun, X. Liu, L. Zhao, X. Feng, X. Fan, and C. Qiu, Observation of acoustic non-hermitian bloch braids and associated topological phase transitions, [Phys. Rev. Lett.](#) **130**, 017201 (2023).
- [65] Z. Y. Chen, S. A. Yang, and Y. X. Zhao, Brillouin Klein bottle from artificial gauge fields, [Nature Communications](#) **13**, 2215 (2022).
- [66] C. Zhang, Z. Y. Chen, Z. Zhang, and Y. X. Zhao, General theory of momentum-space nonsymmorphic symmetry, [Phys. Rev. Lett.](#) **130**, 256601 (2023).
- [67] Z. Y. Chen, Z. Zhang, S. A. Yang, and Y. X. Zhao, Classification of time-reversal-invariant crystals with gauge structures, [Nature Communications](#) **14**, 743 (2023).
- [68] A. G. Fonseca, S. Vaidya, T. Christensen, M. C. Rechtsman, T. L. Hughes, and M. Soljačić, Weyl points on nonorientable manifolds, [Phys. Rev. Lett.](#) **132**, 266601 (2024).
- [69] J. Hu, S. Zhuang, and Y. Yang, Higher-order topological insulators via momentum-space nonsymmorphic symmetries, [Phys. Rev. Lett.](#) **132**, 213801 (2024).
- [70] C.-A. Li, J. Sun, S.-B. Zhang, H. Guo, and B. Trauzettel, Klein-bottle quadrupole insulators and dirac semimetals, [Phys. Rev. B](#) **108**, 235412 (2023).
- [71] P. Lai, J. Wu, Z. Pu, Q. Zhou, J. Lu, H. Liu, W. Deng, H. Cheng, S. Chen, and Z. Liu, Real-projective-plane hybrid-order topological insulator realized in phononic crystals, [Phys. Rev. Appl.](#) **21**, 044002 (2024).
- [72] Y.-L. Tao, M. Yan, M. Peng, Q. Wei, Z. Cui, S. A. Yang, G. Chen, and Y. Xu, Higher-order klein bottle topological insulator in three-dimensional acoustic crystals, [Phys. Rev. B](#) **109**, 134107 (2024).
- [73] A. Hatcher, *Algebraic Topology* (Cambridge University Press, 2001).
- [74] See the Supplemental Material for details.
- [75] Note that a different font of K^2 is used to denote the full Klein bottle, including its boundary.
- [76] The fundamental group $\pi_1(K^2 - P) = *_{n+1}\mathbb{Z}$ as a free group of $n + 1$ generators can also be seen by the deformation retraction of $K^2 - P$: it reduces to the wedge product of $n + 1$ circles $\vee_{n+1}S^1$, i.e., $n + 1$ circles joined at a point. Here, the free group of $n + 1$ generators is denoted by $*_{n+1}\mathbb{Z} = \underbrace{\mathbb{Z} * \mathbb{Z} * \cdots * \mathbb{Z}}_{n+1}$, and the wedge product is $\vee_{n+1}S^1 = \underbrace{S^1 \vee S^1 \vee \cdots \vee S^1}_{n+1}$.
- [77] H.-Y. Wang, F. Song, and Z. Wang, Amoeba formulation of non-bloch band theory in arbitrary dimensions, [Phys. Rev. X](#) **14**, 021011 (2024).
- [78] D. Nakamura, T. Bessho, and M. Sato, Bulk-boundary correspondence in point-gap topological phases, [Phys. Rev. Lett.](#) **132**, 136401 (2024).