#### TANGENT CURRENTS, KING'S RESIDUE FORMULA AND INTERSECTION THEORY

#### TAEYONG AHN

ABSTRACT. In this work, we study the intersection of positive closed currents on domains. We use the theory of tangent currents in connection with King's residue formula. We find a sufficient condition for the local existence of tangent currents, and express the shadow of tangent currents and the *h*-dimension of tangent currents in terms of the complex Monge-Ampère type current. Further, a reasonable integrability condition for the existence of the unique tangent current with minimal *h*-dimension is introduced. We apply it to the study of the intersection of positive closed currents, find a sufficient condition for the intersection of positive closed currents on domains and describe the intersection in terms of the complex Monge-Ampère type current. At the same time, we obtain regularizations of positive closed currents that work well with the suggested intersection of positive closed currents. In particular, the standard regularization of currents by convolution actually produces the convergence towards the intersection of positive closed currents. In this sense, our approach generalizes King's work on currents defined by analytic varieties, which is obtained from Federer's slicing theory. Some classical examples are computed. Our work is applicable to general complex manifolds not necessarily compact or Kähler.

Keywords: tangent current, complex Monge-Ampère operator, King's residue formula, intersection of positive closed currents

# 1. INTRODUCTION

One of the fundamental problems in pluripotential theory is to find a reasonable definition for the intersection of positive closed currents. (For instance, see [1].) Due to the availability of (quasi-)plurisubharmonic functions, the case of bidegree (1,1) has been intensively studied. To name a few, see Demailly ([2]), Hyunh-Kaufmann-Vu ([12]) and references therein. There had not been much known about the higher bidegree case until Dinh-Sibony brought in remarkable ideas: the theory of superpotentials in [7] and the theory of tangent currents in [9]. In [7], Dinh-Sibony used the theory of superpotentials to answer Demailly's question raised in [1] in the case of complex projective space. From the perspective of intersection theory, the theory of superpotentials has been further studied in Dinh-Sibony ([8]), Vu ([20]), and very recently, Luo-Zhou ([16]) and the theory of tangent currents has been investigated in Vu ([21]), Hyunh-Kaufmann-Vu ([12]), Kaufmann-Vu ([13]) and very recently, Nguyen ([17]) and Nguyen-Truong ([18]). In [17], Nguyen systematically studied tangent currents from the perspective of generalized Lelong numbers and introduced a reasonable sufficient condition for the definition of the intersection of positive closed currents in the case of compact Kähler manifolds.

The primary purpose of this article is to study the intersection of positive closed currents of arbitrary bidegree in quite a general case including compact Kähler manifolds. To this end, we use King's residue formula to explore and investigate the relationship between tangent currents and complex Monge-Ampère type currents.

To give a flavor of our approach, we consider the intersection of two positive closed currents on a domain rather than a general tangent current. The residue formula of the following type was proved in [14]. There are various versions, but we refer the reader to [11], [19] for our purpose.

Date: March 11, 2025.

**Theorem 1.1** (Theorem 5.2.1 in [19]). Let  $U \subseteq \mathbb{C}^N$  be a domain. Let  $1 \le n \le N$  be an integer. If the zero set  $A_f$  of a holomorphic mapping  $f: U \to \mathbb{C}^n$  has codimension n, then the currents  $(dd^c \log |f|)^i$  and  $\log |f|(dd^c \log |f|)^i$  with i < n have locally integrable coefficients, and

$$(dd^c \log |f|)^n = \sum_j m_i [A_j] = Z_j$$

where  $A_j$ 's are the irreducible components of  $A_f$ , the summation is made only for (N-n)-dimensional components  $A_j$ 's of the variety  $A_f$  and  $m_j$ 's are the generic multiplicities of f at  $A_j$  and  $Z_f$  is the corresponding holomorphic chain.

Let *S* and *R* be two smooth positive closed currents of bidegree (s, s) and (r, r) on a bounded simply connected domain  $D \in \mathbb{C}^n$  with smooth boundary, where  $1 \le s + r \le n$ . Let  $\pi_i : D \times D \to D$ be the canonical projection onto the *i*-th factor for i = 1, 2. Due to Theorem 1.1, we have

$$\langle S \wedge R, \varphi \rangle = \int_{D \times D} \pi_1^* S \wedge \pi_2^* (R \wedge \varphi) \wedge [\Delta] = \int_{D \times D} \pi_1^* S \wedge \pi_2^* (R \wedge \varphi) \wedge (dd^c u)^* dd^c u \wedge (dd^c u)^*$$

where  $\varphi$  is a smooth test form of bidegree (n - s - r, n - s - r) on D and  $u(x, y) = \log |x - y|$ for  $(x, y) \in D \times D$ . Using the notion of double currents in [10], the current  $\pi_1^*S \wedge \pi_2^*R$  can be defined for the general positive closed currents S and R. So, if the product of the currents  $(dd^c u)^n$  and  $\pi_1^*S \wedge \pi_2^*R$  makes sense, one can expect the resulting product to have the desired properties of the intersection of positive closed currents. For this, we introduce the horizontal condition in Definition 3.5. Under the horizontal condition, we first define  $(dd^c u) \wedge (\pi_1^*S \wedge \pi_2^*R) :=$  $dd^c (u\pi_1^*S \wedge \pi_2^*R)$  and once  $(dd^c u)^{i-1} \wedge (\pi_1^*S \wedge \pi_2^*R)$  is defined, we inductively define  $(dd^c u)^i \wedge$  $(\pi_1^*S \wedge \pi_2^*R) := dd^c (u((dd^c u)^{i-1} \wedge \pi_1^*S \wedge \pi_2^*R))$  for  $1 \le i \le n$ . Inductively given integrability conditions may feel somewhat too much to require but actually it turns out that each integrability condition reduces the h-dimension by 1 as shown in Proposition 3.7.

We relate the above intersection and tangent currents. We approximate the above intersection. We may assume  $D \times D$  is a subset of  $E := \Delta \times \mathbb{C}^n$ , which is the normal bundle of  $\Delta$  in  $D \times D$ . Let  $\overline{E}$  be the projective compactification of E. Let  $\pi_{\Delta} : \overline{E} \to \Delta$  and  $\pi_F : \overline{E} \to \mathbb{P}^n$  denote the canonical projection onto  $\Delta$  and the fiber space  $\mathbb{P}^n$ , respectively. Let  $A_{\lambda} : \overline{E} \to \overline{E}$  be the multiplication by  $\lambda$  on fibers of E. We regularize  $(dd^c u)^n$  by the family  $(\mathcal{K}^n_{\theta})_{0 < |\theta| \ll 1}$  of smooth positive closed forms of bidegree (n, n). For the definition, see Section 2. From the integrability condition, we have the following convergence

$$\lim_{\theta \to 0} \int_{D \times D} \left( \pi_1^* S \wedge \pi_2^* \left( R \wedge \varphi \right) \right) \wedge \mathcal{K}_{\theta}^n = \int_{D \times D} \left( \pi_1^* S \wedge \pi_2^* \left( R \wedge \varphi \right) \right) \wedge \left( dd^c u \right)^n$$

On the other hand, a tangent current  $(\pi_1^* S \wedge \pi_2^* R)_{\infty}$  of  $\pi_1^* S \wedge \pi_2^* R$  along  $\Delta$  is a limit current of the family  $((A_{\lambda})_* (\pi_1 S \wedge \pi_2^* R))_{\lambda \in \mathbb{C}^*}$  of currents in E. For the definition of the tangent current and its shadow, see Definition 2.2. Under the horizontal condition, there exists a unique tangent current of minimal h-dimension. (See Theorem 1.3.) The key point is that we utilize Proposition 3.9 together with the form  $\Omega_n$  as in Lemma 2.7, which is a smooth positive closed form of maximal bidegree with compact support in  $\mathbb{C}^n \subset \mathbb{P}^n$  cohomologous to a linear subspace. There exists a squence  $(\lambda_k) \subset \mathbb{C}^*$  such that  $(\pi_1^* S \wedge \pi_2^* R)_{\infty} = \lim_{k \to \infty} (A_{\lambda_k})_* (\pi_1^* S \wedge \pi_2^* R)_{\infty}$  in the sense of currents on E. Then, the shadow  $(\pi_1^* S \wedge \pi_2^* R)_{\infty}^h$  of the tangent current  $(\pi_1^* S \wedge \pi_2^* R)_{\infty}$  is exactly the same as the suggested intersection as follows.

$$\left( \left( \pi_1^* S \wedge \pi_2^* R \right)_{\infty}^h, \varphi \right) \coloneqq \left( \left( \pi_\Delta \right)_* \left( \pi_1^* S \wedge \pi_2^* R \right)_{\infty} \wedge \pi_F^* \Omega_n, \varphi \right) = \lim_{k \to \infty} \left\langle \left( A_{\lambda_k} \right)_* \left( \pi_1^* S \wedge \pi_2^* R \right) \wedge \pi_F^* \Omega_n, \pi_\Delta^* \varphi \right\rangle$$

$$= \lim_{k \to \infty} \left\langle \pi_1^* S \wedge \pi_2^* R \wedge \left( A_{\lambda_k} \right)^* \left( \pi_F^* \Omega_n \right), \pi_\Delta^* \varphi \right\rangle = \lim_{k \to \infty} \left\langle \pi_1^* S \wedge \pi_2^* R \wedge \mathcal{K}_{e^{-M} |\lambda_k|^{-1}}^n, \pi_2^* \varphi \right\rangle.$$

Here, since  $(D \times D) \cap \Delta$  is isomorphic to D, after proper change of coordinates, we can switch  $\pi_{\Delta}^* \varphi$  to  $\pi_2^* \varphi$ . Clearly, we see that the last limit equals the suggested intersection. One remark is

that the last equality is obtained via change of variables as below:

$$(A_{\lambda_{k}})^{*} (\pi_{F}^{*}\Omega_{m}) = (A_{\lambda_{k}})^{*} \left[ dd^{c}\chi ((\log|x-y|)+M) \wedge (dd^{c}\log|x-y|)^{i-1} \right] = dd^{c}\chi ((\log|\lambda_{k}(x-y)|)+M) \wedge (dd^{c}\log|\lambda_{k}(x-y)|)^{i-1} = dd^{c} \left(\chi ((\log|x-y|) + \log(e^{M}|\lambda_{k}|)) - \log(|\lambda_{k}|e^{M})) \wedge (dd^{c}\log|x-y|)^{i-1} = \mathcal{K}_{e^{-M}|\lambda_{k}|^{-1}}^{m}.$$

Actually, this kind of relationship also holds for other products such as  $\pi_1^* S \wedge \pi_2^* R \wedge (dd^c u)^j$  for  $1 \le j \le n - 1$ . The change of coordinates in the fiber space in this way is the key ingredient of this work. Applying this viewpoint to general tangent currents, we obtain the following theorems linking tangent currents to complex Monge-Ampère type currents. For the definitions, see Section 2 and Section 3.

**Theorem 1.2.** Let  $U \,\subset\, \mathbb{C}^N$  be a bounded simply connected domain with smooth boundary and V a complex submanifold of codimension n. Let T be a positive closed current of bidegree (p,p) on U with a tangent current  $T_{\infty}$  along V. Let m be an integer such that  $n - p \leq m \leq n$ . Suppose that the family  $\left(T \wedge \mathcal{K}^m_{\theta} \wedge \pi^*_V \omega_V^{N-p-m}\right)_{0 < |\theta| \ll 1}$  has locally uniformly bounded mass in U. Then, for every smooth test form  $\varphi$  of bidegree  $\left(N - p - m, N - p - m\right)$  on V, there exists a sequence  $\left(\theta_k\right)_{k \in \mathbb{N}}$  such that the sequence  $\left(T \wedge \mathcal{K}^m_{\theta_k} \wedge \pi^*_V \varphi\right)_{k \in \mathbb{N}}$  of currents of maximal bidegree converges to a current  $L_{\varphi}$  of maximal bidegree in the sense of currents on U and

$$\langle \mathbf{1}_V L_{\varphi}, 1 \rangle = \langle (\pi_V)_* (T_{\infty} \wedge \pi_F^* \Omega_m), \varphi \rangle,$$

where  $\Omega_m$  is the smooth positive closed form of bidegree (m,m) as in Lemma 2.7.

With slightly more restrictive conditions, we can describe it more concretely. See Subsection 3.2 and Corollary 2.12. Another important corollary is Proposition 2.13, which expresses the *h*-dimension of tangent currents in terms of limit currents of the families in Theorem 1.2. See also Corollary 2.15 for a practical criterion. If another family  $(\mathcal{T}^m_{\theta})_{0<|\theta|\ll 1}$  are used instead of the family  $(\mathcal{K}^m_{\theta})_{0<|\theta|\ll 1}$ , we can find a sufficient condition for the local existence of tangent currents as in Proposition 2.17.

Under the horizontal condition in Section 3, the above properties are put together and the existence of a unique tangent current with minimal *h*-dimension follows.

**Theorem 1.3.** Let  $u = \log |x''|$  for  $(x', x'') \in U$  with  $V = \{x'' = 0\}$ . Suppose that the current T satisfies the horizontal condition. Then, there exists a unique tangent current  $T_{\infty}$  of T along V, and the h-dimension of the tangent current  $T_{\infty}$  is minimal. For a smooth test form  $\varphi$  of bidegree (N - p - n, N - p - n) on V, its shadow  $T_{\infty}^h$  is computed as

$$\langle T^h_{\infty}, \varphi \rangle = \int_{U \smallsetminus V} T \wedge (u (dd^c u)^{n-1}) \wedge dd^c \Phi,$$

where  $\Phi$  is a smooth form of bidegree (N - p - n, N - p - n) on U with compact support such that  $\Phi = \pi_V^* \varphi$  in a neighborhood of V. The value of the integral is independent of the choice of  $\Phi$ .

In particular, we obtain a sufficient condition for the definition of the intersection of positive closed currents. For the definition of the product  $\langle \cdot \rangle_{\kappa}$ , see Section 3.

**Theorem 1.4.** Let  $D \in \mathbb{C}^n$  be a bounded simply connected domain with smooth boundary. Let s and r be integers such that  $1 \leq s + r \leq n$ . Let S and R be two positive closed currents of bidegree (s, s) and (r, r), respectively. Let  $u = \log |x - y|$  for  $(x, y) \in D \times D$ . Suppose that  $u \in L^1_{loc}(\langle \pi_1^*S \wedge \pi_2^*(R \wedge \omega_y^{2n-s-r-i+1}) \wedge (dd^c u)^{i-1} \rangle_{\mathcal{K}})$  inductively from i = 1 through n. Then, the current  $\pi_1^*S \wedge \pi_2^*R$  has a unique tangent current along  $\Delta$  and its h-dimension is minimal. Moreover, its shadow is exactly  $(S \wedge R)_{\mathcal{K}}$ , where we have

(1.1) 
$$\langle (S \wedge R)_K, \varphi \rangle \coloneqq \int_{D \times D \setminus \Delta} \pi_1^* S \wedge \pi_2^* R \wedge (u(dd^c u)^{n-1}) \wedge dd^c \Phi,$$

where  $\Phi$  is smooth test form with compact support of bidegree (n - s - r, n - s - r) on  $D \times D$  such that  $\Phi = \pi_2^* \varphi$  in a neighborhood of  $\Delta$ . The value of the integral is independent of the choice of  $\Phi$ . Namely, the Dinh-Sibony product of S and R in [9] is well defined and equal to  $(S \wedge R)_K$ .

In the case of compact Kähler manifolds, Nguyen's work in [17] and our work use different approaches but both works arrive at the essentially exactly same conclusion about the wedgability condition ([17, Theorem 2.18]) and the characterization of the *h*-dimension.

We also obtain a regularization of positive closed currents that fits in with the intersection.

**Theorem 1.5.** Let S and R be two currents as in Theorem 1.4. Then, we have

$$\mathscr{L}^{\mathcal{K}^n_{\theta}}(S) \wedge R = S \wedge \mathscr{L}^{\mathcal{K}^n_{\theta}}(R) \to (S \wedge R)_K$$

in the sense of currents as  $|\theta| \rightarrow 0$ .

We can do the same to the intersection of many positive closed currents. See Subsection 5.3. Due to the invariant nature of the tangent currents, our results can be carried over to general complex manifolds. See Section 6.

An interesting question is whether the standard regularization of positive closed currents by convolution would give the intersection of positive closed currents. Actually, it is so under the wedgability condition as in Theorem 1.4.

**Theorem 1.6.** Let *S* and *R* be two currents as in Theorem 1.4. Let  $(S_{\epsilon})_{0 < \epsilon \ll 1}$  and  $(R_{\epsilon})_{0 < \epsilon \ll 1}$  be standard regularizations of *S* and *R* by convolution, respectively. Then, we have

$$S_{\epsilon} \wedge R, S \wedge R_{\epsilon} \rightarrow (S \wedge R)_K$$

in the sense of currents as  $\epsilon \rightarrow 0$ .

At the same time, this convergence implies that our work generalizes King's work in [15], where Federer's slicing theory is used.

This article is organized as follows. In Section 2, we establish the relationship between tangent currents and complex Monge-Ampère type currents; in Section 3, we introduce integrability conditions and prove the existence of the unique tangent current with minimal *h*-dimension under the integrability conditions; in Section 4, we consider semi-regular transforms on domains and discuss regularizations of positive closed currents on domains; in Section 5, we apply our results on tangent currents and regularizations to the study of the intersection of positive closed currents; in Section 6, complex manifolds are considered and we define a family of superfunctions similar to superpotentials; in Section 7, we study some classical examples; in Section 8, we investigate the relationship between our intersection and the standard regularization by convolution.

**Notations.** We frequently use a smooth convex increasing function  $\chi : \mathbb{R} \to \mathbb{R}_{\geq 0}$  such that  $\chi(t) = 0$ if  $t \leq -1$  and  $\chi(t) = t$  if  $t \geq 1$  in this work. We denote by  $|\cdot|$  the standard Euclidean norm on an Euclidean space or its subset. We use  $||\cdot||_{\infty}$  (or  $||\cdot||_{\infty,U}$ ) and  $||\cdot||_{C^{\alpha}}$  (or  $||\cdot||_{C^{\alpha},U}$ ) for the uniform norm of a given function (or on the set U) and for the  $C^{\alpha}$  norm of a given function (or on the set U), respectively. By each norm applied to a form, we mean the supremum of the norms of the coefficients of the form. For a domain or a complex manifold X, we denote by  $\mathscr{C}_k(X)$  the set of positive closed currents of bidegree (k, k) on X. We write [V] for the current of integration on the regular part of V for an analytic subset V. For a positive current S, we denote by ||S|| either the mass of the current S or the trace measure of S with respect to the standard Kähler form of a given Kähler manifold; in the case of  $\mathbb{P}^n$ , it is with respect to the standard Fubini-Study form  $\omega_{\mathbb{P}^n}$ such that  $\int_{\mathbb{P}^n} \omega_{\mathbb{P}^n}^n = 1$ .

# 2. TANGENT CURRENTS

The theory of tangent currents was introduced in [9] by Dinh-Sibony and further studied by Vu in [21] and by Nguyen in [17]. The following definition is the version in [21].

Let X be a complex manifold of dimension N and V a smooth complex submanifold of X of codimension n. We suppose that n < N. Let T be a positive closed current of bidegree (p,p) on X, where  $1 \le p \le N - n$ . We will consider the other cases later in Section 3. For our purpose, it suffices to consider T that has no mass on V. Let  $\pi_V : E \to V$  denote the normal bundle of V in X and  $\overline{E} := \mathbb{P}(E \oplus \mathbb{C})$  the projective compactification of E. The hypersurface at infinity  $H_{\infty} := \overline{E} \setminus E$  of  $\overline{E}$  is isomorphic to  $\mathbb{P}(E)$  as a fiber bundle over V. We also have a canonical projection  $\pi_{\infty} : E \setminus V \to H_{\infty}$ .

**Definition 2.1.** Let U be an open subset of X with  $U \cap V \neq \emptyset$ . A (local) holomorphic admissible map is a biholomorphism  $\tau$  from U to an open neighborhood of  $U \cap V$  in E, which is the identity on  $U \cap V$ , and the restriction of whose differential  $d\tau$  to  $U \cap V$  is the identity.

The adjective "local" means that it may not be defined in a neighborhood of the entire submanifold V but is defined in the neighborhood U of  $U \cap V$ . The admissible maps in [9] are global and may not be holomorphic. In this work, we use local holomorphic admissible maps. When we need to emphasize the difference, we will add "local". Otherwise, for notational convenience, we will drop "local" and simply write a "holomorphic admissible map."

Suppose that U sits inside a coordinate domain. In local coordinates  $x = (x', x'') \in U$  such that  $U \cap V = \{x'' = 0\}$ , a holomorphic admissible map  $\tau$  can be described as follows:

$$\tau(x) = \left(x' + O\left(|x''|\right), x'' + O\left(|x''|^2\right)\right),$$

and

$$d\tau(x) = \left(dx' + \widetilde{O}\left(\left|x''\right|\right), dx'' + \widetilde{O}\left(\left|x''\right|^2\right)\right).$$

Here, for a positive integer k,  $\tilde{O}(|x''|^k)$  denotes a sum of 1-forms with  $O(|x''|^k)$  coefficients and 1 forms dx'',  $d\overline{x''}$  with  $O(|x''|^{k-1})$  coefficients.

For  $\lambda \in \mathbb{C}^*$ , let  $A_{\lambda} : E \to E$  be the multiplication by  $\lambda$  on fibers of E.

**Definition 2.2.** A tangent current  $T_{\infty}$  of T along V is a positive closed current on E such that there exists a sequence  $(\lambda_k)_{k \in \mathbb{N}} \subset \mathbb{C}^*$  converging to  $\infty$  and a collection of holomorphic admissible maps  $\tau_i : U_i \to E$  for  $i \in I$  satisfying the following two properties:

$$(i)V \subset \bigcup_{i \in I} U_i$$
  
$$(ii)T_{\infty} \coloneqq \lim_{k \to \infty} (A_{\lambda_k})_* (\tau_i)_*T \text{ on } \pi_V^{-1}(U_i \cap V) \text{ for every } i \in I$$

For an open subset  $V_0$  of V, the horizontal dimension (or the h-dimension for short) of  $T_{\infty}$  over  $V_0$  is the largest integer  $h_T$  such that  $T_{\infty} \wedge \pi_V^*(\omega_V^{h_T}) \neq 0$  on  $\pi_V^{-1}(V_0)$ , where  $\omega_V$  is a Kähler form on V.

**Remark 2.3.** A tangent current  $T_{\infty}$  trivially extends to  $\overline{E}$ . We still denote it by  $T_{\infty}$ .

**Remark 2.4.** Thanks to [13, Proposition 2.5], the tangent current is independent of the choice of holomorphic admissible maps  $(\tau_i)_{i \in I}$  and in particular, is well defined on any intersection  $\pi_V^{-1}(U_i \cap V) \cap \pi_V^{-1}(U_i \cap V)$  whenever the limits exist.

**Definition 2.5** (Definition 3.6 in [9]). Let  $T_{\infty}$  be a tangent current of T along V. Let  $h_T$  be the *h*-dimension of  $T_{\infty}$  over an open subset  $V_0$  in V. Let  $\Omega$  be a smooth closed form of bidegree  $(N - p - h_T, N - p - h_T)$  on  $\pi_V^{-1}(V_0)$  whose restriction to each fiber of  $\pi_V$  is cohomologous to a linear

subspace in this fiber. The shadow of  $T_{\infty}$  on  $V_0$  is the postive closed current  $T_{\infty}^h := (\pi_V)_* (T_{\infty} \wedge \Omega)$  of bidegree  $(N - n - h_T, N - n - h_T)$  with support in  $\pi_V (\text{supp} (T_{\infty})) \cap V_0$ . The shadow of T is defined to be its shadow on V.

**Remark 2.6.** According to [9, Proposition 3.5], the shadow  $T^h_{\infty}$  of a tangent current T is independent of the choice of  $\Omega$ .

In this section, we consider the following local case. A bounded simply connected domain  $U \subseteq \mathbb{C}^N$  with smooth boundary near the origin is considered instead of X and V is an N - n-dimensional complex submanifold of U such that  $0 \in V$ . Indeed, due to Remark 2.4, the global case can be easily deduced from the local case by use of a partition of unity.

We will use the coordinates  $(x', x'') \in U$  such that  $x' = (x_1, \ldots, x_{N-n})$ ,  $x'' = (x_{N-n+1}, \ldots, x_N)$ and  $U \cap V = \{x'' = 0\}$ . The projective compactification of the normal bundle of V in U can be written as  $\overline{E} = V \times \mathbb{P}^n$  with the canonical projection map  $\pi_V : \overline{E} \to V$  of  $\overline{E}$  to V. Also, in our case, it makes sense to consider the projection to the fiber space  $\mathbb{P}^n$ , which will be denoted by  $\pi_F : \overline{E} \to \mathbb{P}^n$ . We can extend the coordinates (x', x'') to  $V \times \mathbb{P}^n$  in the following way. We use the coordinates (x', [x'':t]) for a point in  $V \times \mathbb{P}^n$ . We identify  $(x', x'') \in V \times \mathbb{C}^n = E$  with  $(x', [x'':1]) \in V \times \mathbb{P}^n = \overline{E}$ . Hence, we may consider U as a subset of  $\overline{E}$ . Thanks to Remark 2.4, shrinking U and V if necessary, we may assume that  $\{(x', x'') : (x', 0) \in V, |x''| < \varepsilon\} \subset U$  for some  $\varepsilon > 0$  and that the holomorphic admissible map  $\tau : U \to \overline{E}$  is the inclusion map.

We will use the standard Kähler form  $\omega_V = dd^c |x'|$  on V and on each fiber  $\mathbb{P}^n$  of  $\pi_V : \overline{E} \to V$ , we take the Fubini-Study form  $\omega_F = \frac{1}{2}dd^c \log(|x''|^2 + |t|^2)$ . Then,  $\omega := \pi_V^* \omega_V + \pi_F^* \omega_F$  defines a Kähler form on  $\overline{E}$ .

Together with  $\omega_F$ , we will also use the forms  $\Omega_i$  for i = 1, ..., n defined on the fiber space in the following lemma. The lemma below is elementary, but is one of the main ingredients of this work.

**Lemma 2.7.** Let M > 0 be a sufficiently large real number. Let [x'' : t] denote the homogeneous coordinates for  $\mathbb{P}^n$ . We identify  $x'' \in \mathbb{C}^n$  with  $[x'' : 1] \in \mathbb{P}^n$ . Let  $\alpha$  be a smooth form of bidegree (1,1) on  $\mathbb{P}^n$  of bidegree (1,1) defined by

$$\alpha \coloneqq \begin{cases} dd^{c}\chi\left(\left(\log|x''|\right) + M\right) & \text{for } x \in \mathbb{C}^{n} \\ dd^{c}\log|x''| & \text{for } x \text{ in a neighborhood of } \mathbb{P}^{n} \smallsetminus \mathbb{C}^{n} \text{ in } \mathbb{P}^{n}, \end{cases}$$

where  $\chi : \mathbb{R} \to \mathbb{R}_{\geq 0}$  is the convex increasing function in Introduction. Let  $1 \leq i \leq n$  be an integer. Let  $\Omega_i$  be a smooth form defined by

$$\Omega_i \coloneqq \alpha \land \left( dd^c \log |x''| \right)^{i-1}.$$

Then,  $\Omega_i$  is a smooth positive closed form of bidegree (i, i), has support away from the set  $\{x'' = 0\}$ , and is in the same cohomology class as  $\omega_F^i$ .

*Proof.* For  $x'' \in \mathbb{C}^n$  with  $|x''| > e^{1-M}$ ,  $\alpha = dd^c \log |x''|$ . Hence,  $\alpha$  is a well-defined smooth form of bidegree (1,1). Positivity and closedness are local properties and therefore,  $\Omega_i$  is positive closed and of bidegree (*i*, *i*). Notice that  $\alpha = 0$  if  $x'' \in \mathbb{C}^n$  satisfies  $|x''| < e^{-1-M}$ . So,  $\Omega_i$  has support away from the set  $\{x'' = 0\}$ .

Now, we compute the cohomology class of  $\Omega_i$ . Due to the cohomology group  $H^{i,i}(\mathbb{P}^n, \mathbb{R})$  of  $\mathbb{P}^n$  is of dimension 1, it is enough to compute the mass  $\langle \Omega_i, \omega_F^{n-i} \rangle$  and to show that it equals 1. The function  $\chi(\log |x''| + N) - \log |x''|$  is integrable and has compact support in  $\mathbb{C}^n \subset \mathbb{P}^n$  and therefore,

it is defined over the entire  $\mathbb{P}^n$ . So, we have

$$\begin{split} \left\langle \Omega_{i}, \omega_{F}^{n-i} \right\rangle &= \int_{\mathbb{P}^{n}} dd^{c} \log |x''| \wedge \left( dd^{c} \log |x''| \right)^{i-1} \wedge \left( \frac{1}{2} dd^{c} \log \left( |x''|^{2} + |t|^{2} \right) \right)^{n-i} \\ &+ \int_{\mathbb{P}^{k}} dd^{c} \left( \chi (\log |x''| + N) - \log |x''| \right) \wedge \left( dd^{c} \log |x''| \right)^{i-1} \wedge \left( \frac{1}{2} dd^{c} \log \left( |x''|^{2} + |t|^{2} \right) \right)^{n-i} \\ &= \int_{\mathbb{P}^{n}} \left( dd^{c} \log |x''| \right)^{i} \wedge \left( \frac{1}{2} dd^{c} \log \left( |x''|^{2} + |t|^{2} \right) \right)^{n-i} . \end{split}$$

Then, since  $\frac{1}{2}\log\left(\frac{|x''|^2}{|t|^2+|x''|^2}\right)$  is well defined over  $\mathbb{P}^n$ , we have

$$\begin{split} \left\langle \Omega_{i}, \omega_{F}^{n-i} \right\rangle &= \int_{\mathbb{P}^{n}} \left( dd^{c} \log |x''| \right)^{i} \wedge \left( \frac{1}{2} dd^{c} \log \left( |x''|^{2} + |t|^{2} \right) \right)^{n-i} \\ &= \int_{\mathbb{P}^{n}} \left( dd^{c} \log |x''| \right)^{i-1} \wedge \left( \frac{1}{2} dd^{c} \log \left( |x''|^{2} + |t|^{2} \right) \right)^{n-i+1} \\ &+ \int_{\mathbb{P}^{n}} \left( dd^{c} \log |x''| \right)^{i-1} \wedge dd^{c} \frac{1}{2} \log \left( \frac{|x''|^{2}}{|x''|^{2} + |t|^{2}} \right) \wedge \left( \frac{1}{2} dd^{c} \log \left( |x''|^{2} + |t|^{2} \right) \right)^{n-i} \\ &= \int_{\mathbb{P}^{n}} \left( dd^{c} \log |x''| \right)^{n-1} \wedge \left( \frac{1}{2} dd^{c} \log \left( |x''|^{2} + |t|^{2} \right) \right)^{n-i+1} . \end{split}$$

Repeating the argument, we reach

$$\left\langle\Omega_i,\omega_F^{n-i}\right\rangle = \int_{\mathbb{P}^n} \left(dd^c \frac{1}{2}\log\left(|x''|^2 + |t|^2\right)\right)^n = 1.$$

We introduce some more functions and forms. We define

$$u(x) = \log |x''| = \frac{1}{2} \log \left( \sum_{i=N-n+1}^{N} |x_i|^2 \right).$$

on U. Then, notice that from Theorem 1.1, we have

 $(dd^c u)^n = [V]$ 

in the sense of currents. We introduce two different approximations of u. Let  $\theta \in \mathbb{C}^*$  denote a non-zero complex number such that  $|\theta| < 1$ . We may assume that  $|\theta|$  is sufficiently small. Let  $\chi: \mathbb{R} \to \mathbb{R}_{\geq 0}$  be the smooth convex increasing function as in Introduction. We define

$$u_{\theta}^{\mathcal{K}} \coloneqq \chi(u - \log |\theta|) + \log |\theta|$$
 and  $u_{\theta}^{\mathcal{T}} \coloneqq \frac{1}{2} \log \left( |x''|^2 + |\theta|^2 \right).$ 

Both  $u_{\theta}^{\mathcal{K}}$  and  $u_{\theta}^{\mathcal{T}}$  decreasingly converge to u as  $|\theta| \to 0$ . With these approximating functions, we will consider related approximations of the current [V].

- (1)  $\mathcal{K}^{i} := (dd^{c}u)^{i}$  for  $i = 1, \dots, n$ ;
- (2)  $\mathcal{K}^i_{\theta} \coloneqq (dd^c u^{\mathcal{K}}_{\theta}) \land (dd^c u)^{i-1} \text{ for } i = 1, \cdots, n;$ (3)  $\mathcal{T}^i_{\theta} \coloneqq (dd^c u^{\mathcal{T}}_{\theta})^i \text{ for } i = 1, \cdots, n.$

We will use these forms to study tangent currents. In particular, we will use the properties below to investigate tangent currents in connection with Condition (I) in Section 3. The following theorem describes our approach to tangent currents best.

**Theorem 1.2.** Let T be a positive closed current of bidegree (p,p) on U with a tangent current  $T_{\infty}$ along V. Let m be an integer such that  $n-p \le m \le n$ . Suppose that the family  $\left(T \land \mathcal{K}^m_{\theta} \land \pi^*_V \omega_V^{N-p-m}\right)_{0 < |\theta| \ll 1}$ has locally uniformly bounded mass in U. Then, for every smooth test form  $\varphi$  of bidegree (N – p – m, N - p - m) on V, there exists a sequence  $(\theta_k)_{k \in \mathbb{N}}$  such that the sequence  $(T \wedge \mathcal{K}^m_{\theta_k} \wedge \pi^*_V \varphi)_{k \in \mathbb{N}}$  of

currents of maximal bidegree converges to a current  $L_{\varphi}$  of maximal bidegree in the sense of currents on U and

$$\langle \mathbf{1}_V L_{\varphi}, \mathbf{1} \rangle = \langle (\pi_V)_* (T_{\infty} \wedge \pi_F^* \Omega_m), \varphi \rangle$$

where  $\Omega_m$  is the smooth positive closed form of bidegree (m,m) as in Lemma 2.7.

**Remark 2.8.** When  $m = N - p - h_T$ , the form  $\Omega_{N-p-h_T}$  can be replaced any smooth closed  $(N - p - h_T, N - p - h_T)$ -form in the de Rham cohomology class  $\{\omega_F^{N-p-h_T}\}$  as in Definition 2.5. However, this may not be true if  $m < N - p - h_T$ .

*Proof.* We may suppose that  $\varphi$  is positive in the sense of forms as any smooth test form can be written as a difference of two positive smooth test forms. Let  $(\lambda_k)_{k\in\mathbb{N}} \subset \mathbb{C}^*$  be a sequence such that  $(A_{\lambda_k})_* T \to T_\infty$  in E. Let M > 0 be a constant as in Lemma 2.7. From the hypothesis, the sequence  $\left(T \wedge \mathcal{K}^m_{e^{-M}|\lambda_k|^{-1}} \wedge \pi^*_V \varphi\right)_{k\in\mathbb{N}}$  of currents of maximal bidegree has locally uniformly bounded mass, and therefore, there exists a subsequence  $(\lambda_{k_l})$  such that  $\left(T \wedge \mathcal{K}^m_{e^{-M}|\lambda_{k_l}|^{-1}} \wedge \pi^*_V \varphi\right)_{l\in\mathbb{N}}$  converges in U in the sense of currents. By passing to a convergent subsequence, we assume that  $\left(T \wedge \mathcal{K}^m_{e^{-M}|\lambda_k|^{-1}} \wedge \pi^*_V \varphi\right)_{k\in\mathbb{N}}$  converges and let  $L_{\varphi}$  denote the limit current. Notice that if f is a smooth test function on U such that  $\operatorname{supp} f \cap V = \emptyset$ , then  $\lim_{k\to\infty} \left(T \wedge \mathcal{K}^m_{e^{-M}|\lambda_k|^{-1}} \wedge \pi^*_V \varphi, f\right) = \langle T \wedge \mathcal{K}^m \wedge \pi^*_V \varphi, f\rangle$ . So, we have  $\mathbf{1}_{U \smallsetminus V} L_{\varphi} = \mathbf{1}_{U \smallsetminus V} (T \wedge \mathcal{K}^m \wedge \pi^*_V \varphi)$ .

Let  $W \,\subset V$  an open subset of V with compact closure such that  $\operatorname{supp}\varphi \subseteq W$ . Let  $\chi_W : \mathbb{C}^n \to [0,1]$ be a smooth function with compact support defined on the fiber space such that  $\chi_W \equiv 1$  in a neighborhood of  $0 \in \mathbb{C}^n$  and that  $W \times \operatorname{supp}\chi_W \in U$ . Notice that it is not difficult to see from Definition 2.2 that in  $E \cap \pi_V^{-1}(W)$ , the sequence  $((A_{\lambda_k})_*((\pi_F^*\chi_W)T))_{k\in\mathbb{N}}$  converges to the same tangent current  $T_\infty$ . Be aware that on  $\overline{E} \cap \pi_V^{-1}(W)$ , the sequence  $((A_{\lambda_k})_*((\pi_F^*\chi_W)T))_{k\in\mathbb{N}}$  may not converge.

We prove that the convergence of  $\left(T \wedge \mathcal{K}^m_{e^{-M}|\lambda_k|^{-1}} \wedge \pi^*_V \varphi\right)_{k \in \mathbb{N}}$  on U implies the convergence of

$$\left(\int_{\overline{E}} (A_{\lambda_k})_* \left( (\pi_F^* \chi_W) T \right) \wedge \pi_F^* \Omega_m \wedge \pi_V^* \varphi \right)_{k \in \mathbb{N}}\right)$$

in  $\overline{E}$ . We have

$$\begin{aligned} \int_{\overline{E}} (A_{\lambda_k})_* \left( (\pi_F^* \chi_W) T \right) \wedge \pi_F^* \Omega_m \wedge \pi_V^* \varphi &= \int_{\overline{E}} (\pi_F^* \chi_W) T \wedge (A_{\lambda_k})^* (\pi_F^* \Omega_m) \wedge \pi_V^* \varphi \\ &= \int_U (\pi_F^* \chi_W) T \wedge (A_{\lambda_k})^* (\pi_F^* \Omega_m) \wedge \pi_V^* \varphi. \end{aligned}$$

For the second equality, recall that U is assumed to be a subset of  $\overline{E}$  and that T is supported in U. Observe that over U, we have

$$(2.1) \qquad (A_{\lambda_k})^* (\pi_F^* \Omega_m) = (A_{\lambda_k})^* \left[ dd^c \chi \left( \left( \log |x''| \right) + M \right) \wedge \left( dd^c \log |x''| \right)^{i-1} \right] \\ = dd^c \chi \left( \left( \log |\lambda_k x''| \right) + M \right) \wedge \left( dd^c \log |\lambda_k x''| \right)^{i-1} \\ = dd^c \chi \left( \left( \log |x''| \right) + \log \left( e^M |\lambda_k| \right) \right) \wedge \left( dd^c \left( \log |x''| + \log |\lambda_k| \right) \right)^{i-1} \\ = dd^c \left( \chi \left( \left( \log |x''| \right) + \log \left( e^M |\lambda_k| \right) \right) - \log \left( |\lambda_k| e^M \right) \right) \wedge \left( dd^c \log |x''| \right)^{i-1} \\ = \mathcal{K}_{e^{-M} |\lambda_k|^{-1}}^m.$$

Hence, the above integral becomes

$$\int_{\overline{E}} (A_{\lambda_k})_* \left( (\pi_F^* \chi_W) T \right) \wedge \pi_F^* \Omega_m \wedge \pi_V^* \varphi = \int_U (\pi_F^* \chi_W) T \wedge (A_{\lambda_k})^* (\pi_F^* \Omega_m) \wedge \pi_V^* \varphi$$
$$= \int_U T \wedge \mathcal{K}_{e^{-M} |\lambda_k|^{-1}}^m \wedge (\pi_F^* \chi_W) (\pi_V^* \varphi).$$

Since the support of  $(\pi_F^*\chi_W)(\pi_V^*\varphi)$  is compact support in U, the desired convergence follows from our hypothesis.

Let  $\chi_{\varepsilon}^{\infty} : \mathbb{P}^n \to [0,1]$  be a smooth function such that  $\chi_{\varepsilon}^{\infty} \equiv 1$  in a neighborhood of the hyperplane at infinity and the support sits inside the  $\varepsilon$ -neighborhood of the hyperplane at infinity, where the distance is measured with respect to the norm associated with the form  $\frac{1}{2}dd^c \log (|x''|^2 + |t|^2)$ . Then, we have

(2.2) 
$$\int_{U} T \wedge \mathcal{K}^{m}_{e^{-M}|\lambda_{k}|^{-1}} \wedge (\pi^{*}_{F}\chi_{W}) (\pi^{*}_{V}\varphi) = \int_{\overline{E}} (A_{\lambda_{k}})_{*} ((\pi^{*}_{F}\chi_{W})T) \wedge \pi^{*}_{F}\Omega_{m} \wedge \pi^{*}_{V}\varphi$$

(2.3) 
$$= \int_{\overline{E}} \pi_F^* \chi_{\varepsilon}^{\infty} (A_{\lambda_k})_* ((\pi_F^* \chi_W) T) \wedge \pi_F^* \Omega_m \wedge \pi_V^* \varphi$$

(2.4) 
$$+ \int_{\overline{E}} (1 - \pi_F^* \chi_{\varepsilon}^{\infty}) (A_{\lambda_k})_* ((\pi_F^* \chi_W) T) \wedge \pi_F^* \Omega_m \wedge \pi_V^* \varphi$$

For (2.3), we have

$$\int_{\overline{E}} \pi_F^* \chi_{\varepsilon}^{\infty} \left( A_{\lambda_k} \right)_* \left( \left( \pi_F^* \chi_W \right) T \right) \wedge \pi_F^* \Omega_m \wedge \pi_V^* \varphi = \int_{\overline{E}} T \wedge \left( \pi_F^* \chi_W \right) \left( A_{\lambda_k} \right)_* \left( \pi_F^* \left( \chi_{\varepsilon}^{\infty} \Omega_m \right) \right) \wedge \pi_V^* \varphi.$$

For a given sufficiently small  $\varepsilon > 0$ , we have  $\pi_F^* \Omega_m = \mathcal{K}^m$  on the support of  $\pi_F^* \chi_{\varepsilon}^{\infty}$  and  $(A_{\lambda_k})^* \mathcal{K}^m = \mathcal{K}^m$  on  $\overline{E} \smallsetminus V$ . Hence, we get

$$\begin{split} \int_{\overline{E}} \pi_F^* \chi_{\varepsilon}^{\infty} \left( A_{\lambda_k} \right)_* \left( \left( \pi_F^* \chi_W \right) T \right) \wedge \pi_F^* \Omega_m \wedge \pi_V^* \varphi \\ &= \int_{\mathrm{supp} T \cap A_{\lambda_k} \left( \mathrm{supp} \left( \pi_F^* \chi_{\varepsilon}^{\infty} \right) \right)} \left( \left( \pi_F^* \chi_{\varepsilon}^{\infty} \right) \circ A_{\lambda_k}^{-1} \right) T \wedge \mathcal{K}^m \wedge \left( \pi_F^* \chi_W \right) \left( \pi_V^* \varphi \right). \end{split}$$

As we have  $0 \le (\pi_F^* \chi_{\varepsilon}^{\infty}) \circ A_{\lambda_k}^{-1} \le 1$  and  $A_{\lambda_k} (\operatorname{supp} (\pi_F^* \chi_{\varepsilon}^{\infty})) \cap V = \emptyset$  for all  $k \in \mathbb{N}$ , we obtain

$$\int_{\overline{E}} \pi_F^* \chi_{\varepsilon}^{\infty} \left( A_{\lambda_k} \right)_* \left( \left( \pi_F^* \chi_W \right) T \right) \wedge \pi_F^* \Omega_m \wedge \pi_V^* \varphi \le \int_{U \smallsetminus V} \left( \pi_F^* \chi_W \right) T \wedge \mathcal{K}^m \wedge \pi_V^* \varphi$$

Since  $\left(T \wedge \mathcal{K}_{e^{-M}|\lambda_k|^{-1}}^m \wedge \pi_V^* \varphi\right)_{k \in \mathbb{N}}$  converges in U and the support of  $(\pi_F^* \chi_W)(\pi_V^* \varphi)$  is compact, we let  $k \to \infty$  to get from (2.2) that

$$\int_{U} (\pi_{F}^{*}\chi_{W}) L_{\varphi} \leq \int_{U \setminus V} (\pi_{F}^{*}\chi_{W}) T \wedge \mathcal{K}^{m} \wedge \pi_{V}^{*}\varphi + \int_{\overline{E}} (1 - \chi_{\varepsilon}^{\infty}) T_{\infty} \wedge \pi_{F}^{*}\Omega_{m} \wedge \pi_{V}^{*}\varphi.$$

The convergence of  $((A_{\lambda_k})_*((\pi_F^*\chi_W)T))_{k\in\mathbb{N}}$  to  $T_{\infty}$  in the second integral is from the discussion in the second paragraph of the proof. Letting  $\varepsilon \to 0$  and applying the argument in the beginning of the proof, we have

(2.5) 
$$\int_{U} \mathbf{1}_{V} L_{\varphi} \leq \int_{E} T_{\infty} \wedge \pi_{F}^{*} \Omega_{m} \wedge \pi_{V}^{*} \varphi$$

In the last integral, it does not actually matter whether the region of integration is E or  $\overline{E}$  as the current  $T_{\infty}$  trivially extends to  $\overline{E}$ .

We prove the other direction. As the tangent current  $T_{\infty}$  trivially extends to  $H_{\infty} = \overline{E} \setminus E$ , we have

$$\int_{E} T_{\infty} \wedge \pi_{F}^{*} \Omega_{m} \wedge \pi_{V}^{*} \varphi = \lim_{\varepsilon \to 0} \int_{E} (1 - \pi_{F}^{*} \chi_{\varepsilon}^{\infty}) T_{\infty} \wedge \pi_{F}^{*} \Omega_{m} \wedge \pi_{V}^{*} \varphi$$
$$= \lim_{\varepsilon \to 0} \lim_{k \to \infty} \int_{E} (1 - \pi_{F}^{*} \chi_{\varepsilon}^{\infty}) (A_{\lambda_{k}})_{*} T \wedge \pi_{F}^{*} \Omega_{m} \wedge \pi_{V}^{*} \varphi$$

Let  $\chi_{\delta}^{0} : U \to [0,1]$  be another smooth function with compact support such that  $\chi_{\delta}^{0} \equiv 1$  in a neighborhood of  $\overline{W}$  in U and that supp $\chi_{\delta}^{0}$  lies inside the  $\delta$ -neighborhood of V, where the distance

is measured with respect to the Euclidean norm on U. Let  $\delta > 0$  be given. Then, arguing as previously, we have

$$\int_{\overline{E}} (1 - \pi_F^* \chi_{\varepsilon}^{\infty}) (A_{\lambda_k})_* T \wedge \pi_F^* \Omega_m \wedge \pi_V^* \varphi = \int_{\overline{E}} (1 - (\pi_F^* \chi_{\varepsilon}^{\infty}) \circ A_{\lambda_k}) T \wedge \mathcal{K}_{e^{-M} |\lambda_k|^{-1}}^m \wedge \pi_V^* \varphi$$
$$\leq \int_{\overline{E}} \chi_{\delta}^0 T \wedge \mathcal{K}_{e^{-M} |\lambda_k|^{-1}}^m \wedge \pi_V^* \varphi = \int_{U} \chi_{\delta}^0 T \wedge \mathcal{K}_{e^{-M} |\lambda_k|^{-1}}^m \wedge \pi_V^* \varphi$$

for all sufficiently large  $k \in \mathbb{N}$  as the support of  $(1 - (\pi_F^* \chi_{\varepsilon}^{\infty}) \circ A_{\lambda_k}) \pi_V^* \varphi$  shrinks to W. Letting  $k \to \infty$  and  $\varepsilon \to 0$  in this order, we obtain

$$\int_E T_\infty \wedge \pi_F^* \Omega_m \wedge \pi_V^* \varphi \le \int_U \chi_\delta^0 L_\varphi$$

for all sufficiently small  $\delta > 0$ . Finally, we let  $\delta \rightarrow 0$  and we get

(2.6) 
$$\int_E T_{\infty} \wedge \pi_F^* \Omega_m \wedge \pi_V^* \varphi \leq \int_U \mathbf{1}_V L_{\varphi}$$

Take  $\theta_k = e^{-M} |\lambda_k|^{-1}$  for  $k \in \mathbb{N}$  and the inequalities (2.5) and (2.6) conclude the desired equality.

Consider the case where m = n. Intuitively, in the above approximation  $(A_{\lambda_k})^* \Omega_n$  of the current of integration [V], we concentrate all the mass to the submanifold V to get some information of the current on the submanifold V. On the other hand, for the tangent current, we push all the mass outside the submanifold V to the hyperplane  $H_{\infty}$  at infitnity so that only the derivative type information of the current along the submanifold V remains over E. So, if we sum the information up over the set E not  $\overline{E}$ , then we get the same information as in the case of the approximation of the current of integration [V].

**Remark 2.9.** In the definition of the tangent current, the convergence is not in  $\overline{E}$  but in E. So, tangent currents are the restriction to E of limit currents of  $((A_{\lambda_k})_* T)_{k\in\mathbb{N}}$  in  $\overline{E}$ , which is always possible to define due to the El Mir-Skoda theorem. When the family  $(T \wedge \mathcal{K}^m_{e^{-M}|\lambda_k|^{-1}} \wedge \pi^*_V \varphi)_{k\in\mathbb{N}}$  is considered, such difference between the convergence of  $((A_{\lambda_k})_* T)_{k\in\mathbb{N}}$  in E and that in  $\overline{E}$  appears as a limit value of integrals with the region of integration in  $U \setminus V$ . Actually, the proof of Theorem 1.2 is about this.

If we are given a slightly stronger condition, by a smilar proof to the proof of Theorem 1.2, we obtain the following characterization. Because of the similarity, we omit the proof.

**Corollary 2.10.** Let T be a positive closed current of bidegree (p, p) on U with a tangent current  $T_{\infty}$  along V. Let m be an integer such that  $n - p \le m \le n$ . Suppose that the family  $(T \land \mathcal{K}^m_{\theta})_{0 < |\theta| \ll 1}$  has locally uniformly bounded mass in U. Then, there exists a positive closed current  $T_{\mathcal{K}}^{p+m}$  of bidegree (p+m, p+m) on U such that  $T_{\mathcal{K}}^{p+m}$  is a limit current of the family  $(T \land \mathcal{K}^m_{\theta})_{0 < |\theta| \ll 1}$  and that we have

$$(\pi_V)_* \left( \mathbf{1}_V T_{\mathcal{K}}^{p+m} \right) = (\pi_V)_* \left( T_{\infty} \wedge \pi_F^* \Omega_m \right),$$

where  $\Omega_m$  is the smooth positive closed form of bidegree (m,m) as in Lemma 2.7. In particular, when N - p - m is the horizontal dimension  $h_T$ , that is,  $m = N - p - h_T$ , then  $(\pi_V)_* (\mathbf{1}_V T_{\mathcal{K}}^{p+m})$  equals the shadow  $T_{\infty}^h$  of  $T_{\infty}$ .

**Remark 2.11.** The push-forward  $(\pi_V)_*$  in the two sides of the equality has slightly different meanings although they look alike in our settings. The left hand side is actually  $\pi_V : U \to V$  while the right hand side is  $\pi_V : \overline{E} \to V$ . Also, the set U is not compact, but since the support of  $\mathbf{1}_V T_{\mathcal{K}}^{p+m}$  sits inside V, the left hand side is well defined. **Corollary 2.12.** Let T be a positive closed current of bidegree (p, p) on U with a tangent current  $T_{\infty}$  along V. Suppose that the family  $\left(T \wedge \mathcal{K}_{\theta}^{n} \wedge \pi_{V}^{*} \omega_{V}^{N-p-n}\right)_{0 < |\theta| \ll 1}$  has locally uniformly bounded mass in U. Then, the family  $\left(T \wedge \mathcal{K}_{\theta}^{n}\right)_{0 < |\theta| \ll 1}$  has locally uniformly bounded mass in U and there exists a limit current  $L_{T_{\infty}}$  of  $\left(T \wedge \mathcal{K}_{\theta}^{n}\right)_{0 < |\theta| \ll 1}$  such that

$$(\pi_V)_*L_{T_{\infty}} = (\pi_V)_* (T_{\infty} \wedge \pi_F^* \Omega_n),$$

where  $\Omega_n$  is the smooth positive closed form of bidegree (n,n) as in Lemma 2.7. In particular, when T has the minimal horizontal dimension, then  $(\pi_V)_* L_{T_{\infty}}$  equals the shadow  $T^h_{\infty}$  of  $T_{\infty}$ .

Proof. For the first assertion, it is enough to observe that for bidegree reason, we have

$$T \wedge \mathcal{K}^n_{\theta} \wedge \omega^{N-p-n} = T \wedge \mathcal{K}^n_{\theta} \wedge \pi^*_V \omega_V^{N-p-n}$$

For the second assertion, it suffices to notice that the limit currents of  $(T \wedge \mathcal{K}^n_\theta)_{0 < |\theta| \ll 1}$  are supported in *V*.

Since the *h*-dimension is actually independent of tangent currents of the current T along V, the following theorem is a direct consequence of Theorem 1.2.

**Proposition 2.13.** Let T be a positive closed current of bidegree (p,p) on U with a tangent current  $T_{\infty}$  along V. Suppose that for each integer m with  $N - p - n \le m \le N - p$ , the family  $\left(T \wedge \mathcal{K}_{\theta}^{N-p-m} \wedge \pi_{V}^{*} \omega_{V}^{m}\right)_{0 < |\theta| \ll 1}$  of currents of maximal bidegree has locally uniformly bounded mass. The h-dimension of  $T_{\infty}$  is the largest integer m such that there exists a limit current of the family  $\left(T \wedge \mathcal{K}_{\theta}^{N-p-m} \wedge \pi_{V}^{*} \omega_{V}^{m}\right)_{0 < |\theta| \ll 1}$  having mass on V.

If we use  $\omega_F^m$  instead of  $\Omega_m$ , then it does not seem easy to obtain the same result as Theorem 1.2, but we can still get a partial estimate and a useful corollary.

**Proposition 2.14.** Let T be a positive closed current of bidegree (p, p) on U with a tangent current  $T_{\infty}$  along V. Suppose that the family  $\left(T \wedge \mathcal{T}_{\theta}^{m} \wedge \pi_{V}^{*} \omega_{V}^{N-p-m}\right)_{0 < |\theta| \ll 1}$  has locally uniformly bounded mass in U for some integer m with  $n - p \le m \le n$ . Then, there exists a positive closed current  $L_{T}^{\mathcal{T}}$  of maximal bidegree on U such that  $L_{T}^{\mathcal{T}}$  is a limit current of the family  $\left(T \wedge \mathcal{T}_{\theta}^{m} \wedge \pi_{V}^{*} \omega_{V}^{N-p-m}\right)_{0 < |\theta| \ll 1}$  and that we have

$$(\pi_V)_* (T_{\infty} \wedge \pi_F^* \omega_F^m) \wedge \omega_V^{N-p-m} \le (\pi_V)_* (\mathbf{1}_V L_T^{\mathcal{T}})$$

in the sense of currents on V.

Proof. We apply the exactly same argument in the proof of Theorem 1.2 for the inequality

(2.7) 
$$\int_{E} T_{\infty} \wedge \pi_{F}^{*} \omega_{F}^{m} \wedge \pi_{V}^{*} \left( f \omega_{V}^{N-p-m} \right) \leq \int_{U} \mathbf{1}_{V} \pi_{V}^{*} f L_{T}^{\mathcal{T}},$$

where  $f: V \to \mathbb{R}$  is a positive smooth test function.

The following corollary is straightforward from Definition 2.2 and Proposition 2.14.

**Corollary 2.15.** Let T be a positive closed current of bidegree (p,p) on U with a tangent current  $T_{\infty}$  along V. Let m be an integer such that  $n - p \le m \le n - 1$ . Suppose that the family  $(T \wedge \mathcal{T}_{\theta}^{i} \wedge \pi_{V}^{*} \omega_{V}^{N-p-i})_{0 < |\theta| \ll 1}$  has locally uniformly bounded mass in U for  $n - p \le i \le m$  and that for each  $i = n - p, \ldots, m$ , there exists a limit current of  $(T \wedge \mathcal{T}_{\theta}^{i} \wedge \pi_{V}^{*} \omega_{V}^{N-p-i})_{0 < |\theta| \ll 1}$  having no mass on V. Then, the h-dimension of  $T_{\infty}$  is at most N - p - m - 1.

The local existence of tangent currents is obtained as a corollary of the following proposition.

**Proposition 2.16.** Let *T* be a positive closed current of bidegree (p, p) on *U*. Let  $\varphi : U \to \mathbb{R}_{\geq 0}$  be a non-negative smooth function with compact support. Then, the mass for the family  $((A_{\lambda})_* (\varphi T))_{\lambda \in \mathbb{C}^*}$  is uniformly bounded in  $\overline{E}$  if and only if the mass for the family  $(\varphi T \wedge \mathcal{T}_{\lambda^{-1}}^i \wedge \pi_V^* \omega_V^{N-p-i})_{\lambda \in \mathbb{C}^*}$  is uniformly bounded on *U* for i = 1, ..., n. If the current *T* has bounded mass on *U*, then the mass for the family  $((A_{\lambda})_* T)_{\lambda \in \mathbb{C}^*}$  is uniformly bounded in  $\overline{E}$  if and only if the mass of the family  $(T \wedge \mathcal{T}_{\lambda^{-1}}^i \wedge \pi_V^* \omega_V^{N-p-i})_{\lambda \in \mathbb{C}^*}$  is uniformly bounded on *U* for i = 1, ..., n.

*Proof.* Since the support of the current  $(A_{\lambda})_*(\varphi T)$  stays in E, we have

$$\left\langle (A_{\lambda})_{*} (\varphi T), \omega^{N-p} \right\rangle = \int_{E} (A_{\lambda})_{*} (\varphi T) \wedge \omega^{N-p}$$

$$= \sum_{l=0}^{n} c_{l} \int_{E} (A_{\lambda})_{*} (\varphi T) \wedge (\pi_{F}^{*} \omega_{F})^{l} \wedge (\pi_{V}^{*} \omega_{V})^{N-p-l}$$

$$= \sum_{l=0}^{n} c_{l} \int_{U} \varphi T \wedge \left[ (A_{\lambda})_{*} (\pi_{F}^{*} \omega_{F})^{l} \right] \wedge (\pi_{V}^{*} \omega_{V})^{N-p-l}$$

$$= \sum_{l=0}^{n} c_{l} \int_{U} \varphi T \wedge \mathcal{T}_{\lambda^{-1}}^{l} \wedge (\pi_{V}^{*} \omega_{V})^{N-p-l} ,$$

where  $c_l$ 's are some nonnegative constants independent of  $\lambda \in \mathbb{C}^*$ . The last equality is from the following observation. Notice that the restriction of  $\omega_F$  to  $\mathbb{C}^n$  can be written as  $\frac{1}{2}dd^c \log(1+|x''|^2)$ . Then, we have

$$(A_{\lambda})^{*} (\pi_{F}^{*}\omega) = (A_{\lambda})^{*} \left[\frac{1}{2} dd^{c} \log(1 + |x''|^{2})\right]$$
$$= \frac{1}{2} dd^{c} \log\left(1 + |\lambda x''|^{2}\right) = \frac{1}{2} dd^{c} \log\left(|x''|^{2} + |\lambda|^{-2}\right) = dd^{c} u_{\lambda^{-1}}^{\mathcal{T}}$$

For the second assertion, the same arguments work. We just remove the smooth function  $\varphi$  in the proof.

**Proposition 2.17.** If the family  $\left(T \wedge \mathcal{T}_{\lambda^{-1}}^{i} \wedge \pi_{V}^{*} \omega_{V}^{N-p-i}\right)_{\lambda \in \mathbb{C}^{*}}$  has locally uniformly bounded mass on U for i = 1, ..., n, then tangent currents of T along V exist.

*Proof.* Let  $(V_i)$  be a sequence of open subsets with compact closure in V such that  $\overline{V_i} \in V_{i+1}$  for every  $i \in \mathbb{N}$ . Let  $\chi_{V_i} : U \to [0,1]$  be a smooth function with compact support such that  $\chi_{V_i} \equiv 1$  in a neighborhood of  $\overline{V_i}$  in U. Then, by Proposition 2.16, the sequence  $((A_\lambda)_*(\chi_{V_i}T))$  is uniformly bounded in  $\overline{E}$  for each  $i \in \mathbb{N}$ . Let K be a compact subset of E. Then, there exists an  $i \in \mathbb{N}$  such that  $K \in \pi_V^{-1}(V_i)$ . Also, for all  $\lambda \in \mathbb{C}^*$  sufficiently large  $|\lambda|$ ,  $(A_\lambda)_* \chi_{V_i} \equiv 1$  on K. Hence, we can find a sequence  $(\lambda_k)$  such that  $(A_{\lambda_k})_* T$  converges on K.

We find a sequence of open subset  $E_j$  with compact closure in E such that  $\bigcup_{j \in \mathbb{N}} E_j = E$ . We apply the above arguments to each  $E_j$  and then, we use the classical diagonal argument to obtain a tangent current.

The following lemma will be used several times later. The point is that  $dd^c u^{\mathcal{T}}_{\theta}$  is strictly positive, but  $dd^c u$  lacks strict positivity in one direction. However, due to the properties of the exterior algebra, multiplying a single  $dd^c u^{\mathcal{T}}_{\theta}$  makes up the missing direction and we obtain the estimate.

**Lemma 2.18.** There exists a constant C > 0 such that for  $1 \le i \le n$ , we have

$$\mathcal{T}^{i}_{\theta} \leq C dd^{c} u_{\theta}^{\mathcal{T}} \wedge \mathcal{K}^{i-1}$$

outside V for all  $\theta$  with  $0 < |\theta| \ll 1$ , where the positivity is the strong positivity for forms.

*Proof.* For notational convenience, in this proof, we will write w instead of x'', that is,  $w_1 = x_{N-n+1}, \ldots, w_j = x_{N-n+j}, \ldots, w_n = x_N$ . On  $\{w \neq 0\}$ , we have

$$dd^{c}u = \frac{\sum_{j=1}^{n} \sqrt{-1}dw_{j} \wedge d\overline{w_{j}}}{|w|^{2}} - \frac{\sqrt{-1}\left(\sum_{j=1}^{n} \overline{w_{j}}dw_{j}\right) \wedge \left(\sum_{j=1}^{n} w_{j}d\overline{w_{j}}\right)}{|w|^{4}} \quad \text{and}$$
$$dd^{c}u_{\theta}^{\mathcal{T}} = \frac{\sum_{j=1}^{n} \sqrt{-1}dw_{j} \wedge d\overline{w_{j}}}{|w|^{2} + |\theta|^{2}} - \frac{\sqrt{-1}\left(\sum_{j=1}^{n} \overline{w_{j}}dw_{j}\right) \wedge \left(\sum_{j=1}^{n} w_{j}d\overline{w_{j}}\right)}{(|w|^{2} + |\theta|^{2})^{2}}$$
$$= \frac{|\theta|^{2}}{(|w|^{2} + |\theta|^{2})^{2}} \sum_{j=1}^{n} \sqrt{-1}dw_{j} \wedge d\overline{w_{j}} + \frac{|w|^{4}}{(|w|^{2} + |\theta|^{2})^{2}} dd^{c}u.$$

Since  $dd^c u$  is positive, we have

$$0 \le dd^c u_{\theta}^{\mathcal{T}} \le \frac{|\theta|^2}{\left(|w|^2 + |\theta|^2\right)^2} \sum_{j=1}^n \sqrt{-1} dw_j \wedge d\overline{w_j} + dd^c u$$

on the set  $\{w \neq 0\}$ , where the positivity is the strong positivity for forms. Due to the inequality, in order to prove the claim, it is enough to show that for each i = 1, ..., n, there exists a constant  $C_i > 0$  such that

$$\left(\frac{|\theta|^2}{(|w|^2 + |\theta|^2)^2} \sum_{j=1}^n \sqrt{-1} dw_j \wedge d\overline{w_j}\right)^i \le C_i \left(\frac{|\theta|^2}{(|w|^2 + |\theta|^2)^2} \sum_{j=1}^n \sqrt{-1} dw_j \wedge d\overline{w_j}\right) \wedge (dd^c u)^{i-1}$$

on  $\{w \neq 0\}$ , where the positivity is the strong positivity for forms.

Without loss of generality, we consider  $0 < |w_1| < 1$ ,  $|w_j| < 2|w_1|$  for j = 2, ..., n and use the coordinates  $(v_1, ..., v_n)$  where  $v_1 = w_1$  and  $v_j = w_j/w_1$  for j = 2, ..., n. For k = 2, ..., n, we have

$$dw_k = d(v_1v_k) = v_1dv_k + v_kdv_1$$
 and  $d\overline{w_k} = d(\overline{v_1v_k}) = \overline{v_1}d\overline{v_k} + \overline{v_k}d\overline{v_1}$ 

Since  $\sqrt{-1}(v_1 dv_k - v_k dv_1) \wedge (\overline{v_1} d\overline{v_k} - \overline{v_k} d\overline{v_1})$  is a positive form, for k = 2, ..., n, we have

$$\frac{|\theta|^{2}}{(|w|^{2}+|\theta|^{2})^{2}} \left(\sqrt{-1}dw_{k} \wedge d\overline{w_{k}}\right) \wedge \left(\frac{|\theta|^{2}}{(|w|^{2}+|\theta|^{2})^{2}} \sum_{j=1}^{n} \sqrt{-1}dw_{j} \wedge d\overline{w_{j}}\right)^{i-1} \\
= \frac{|\theta|^{2}}{(|w|^{2}+|\theta|^{2})^{2}} \sqrt{-1} \left(v_{1}dv_{k}+v_{k}dv_{1}\right) \wedge \left(\overline{v_{1}}d\overline{v_{k}}+\overline{v_{k}}d\overline{v_{1}}\right) \wedge \left(\frac{|\theta|^{2}}{(|w|^{2}+|\theta|^{2})^{2}} \sum_{j=1}^{n} \sqrt{-1}dw_{j} \wedge d\overline{w_{j}}\right)^{i-1} \\
\leq \frac{|\theta|^{2}}{(|w|^{2}+|\theta|^{2})^{2}} \sqrt{-1} \left(v_{1}dv_{k}+v_{k}dv_{1}\right) \wedge \left(\overline{v_{1}}d\overline{v_{k}}+\overline{v_{k}}d\overline{v_{1}}\right) \wedge \left(\frac{|\theta|^{2}}{(|w|^{2}+|\theta|^{2})^{2}} \sum_{j=1}^{n} \sqrt{-1}dw_{j} \wedge d\overline{w_{j}}\right)^{i-1} \\
+ \frac{|\theta|^{2}}{(|w|^{2}+|\theta|^{2})^{2}} \sqrt{-1} \left(v_{1}dv_{k}-v_{k}dv_{1}\right) \wedge \left(\overline{v_{1}}d\overline{v_{k}}-\overline{v_{k}}d\overline{v_{1}}\right) \wedge \left(\frac{|\theta|^{2}}{(|w|^{2}+|\theta|^{2})^{2}} \sum_{j=1}^{n} \sqrt{-1}dw_{j} \wedge d\overline{w_{j}}\right)^{i-1} \\
= \frac{2|\theta|^{2}}{\left(|w|^{2}+|\theta|^{2}\right)^{2}} \sqrt{-1} \left(|v_{1}|^{2}dv_{k} \wedge d\overline{v_{k}}+|v_{k}|^{2}dv_{1} \wedge d\overline{v_{1}}\right) \wedge \left(\frac{|\theta|^{2}}{(|w|^{2}+|\theta|^{2})^{2}} \sum_{j=1}^{n} \sqrt{-1}dw_{j} \wedge d\overline{w_{j}}\right)^{i-1}.$$

Note that with respect to the coordinates in our consideration,  $|v_k|$ 's are bounded by 2 for k = 2, ..., n. Inductively applying the above inequality, we see that on  $\{w \neq 0\}$ , the form  $\left(\frac{|\theta|^2}{(|w|^2 + |\theta|^2)^2} \sum_{j=1}^n \sqrt{-1} dw_j \wedge d\overline{w_j}\right)^i$  is bounded by a linear combination with positive coefficients of forms of bidegree (i, i) of the following types:

(1) 
$$\left(\sqrt{-1}\right)^{i} \left(\frac{|\theta|^{2}}{\left(|v_{1}|^{2}+|\theta|^{2}\right)^{2}}\right)^{i} |v_{1}|^{2i-2} dv_{1} \wedge d\overline{v_{1}} \wedge dv_{j_{1}} \wedge d\overline{v_{j_{1}}} \wedge \cdots \wedge dv_{j_{i-1}} \wedge d\overline{v_{j_{i-1}}} \quad \text{or}$$
  
(2)  $\left(\sqrt{-1}\right)^{i} \left(\frac{|\theta|^{2}}{\left(|v_{1}|^{2}+|\theta|^{2}\right)^{2}}\right)^{i} |v_{1}|^{2i} dv_{j_{1}} \wedge d\overline{v_{j_{1}}} \wedge \cdots \wedge dv_{j_{i}} \wedge d\overline{v_{j_{i}}},$   
pre  $i_{1} = i_{i} \geq 2$ 

where  $j_1, \ldots, j_i \ge 2$ .

In the case of (1), by the relationship between the arithmetic mean and the geometric mean, we have

$$\left(\sqrt{-1}\right)^{i} \left(\frac{|\theta|^{2}}{\left(|v_{1}|^{2}+|\theta|^{2}\right)^{2}}\right)^{i} |v_{1}|^{2i-2} dv_{1} \wedge d\overline{v_{1}} \wedge dv_{j_{1}} \wedge d\overline{v_{j_{1}}} \wedge \cdots \wedge dv_{j_{i-1}} \wedge d\overline{v_{j_{i-1}}}$$

$$\leq \left(\sqrt{-1}\right)^{i} \left(\frac{|\theta|^{2}}{\left(|v_{1}|^{2}+|\theta|^{2}\right)^{2}}\right) dv_{1} \wedge d\overline{v_{1}} \wedge dv_{j_{1}} \wedge d\overline{v_{j_{1}}} \wedge \cdots \wedge dv_{j_{i-1}} \wedge d\overline{v_{j_{i-1}}}$$

Since  $v_1 = w_1$ , we have

$$\sqrt{-1} \left( \frac{|\theta|^2}{(|v_1|^2 + |\theta|^2)^2} \right) dv_1 \wedge d\overline{v_1} \leq \sqrt{-1} \frac{4n|\theta|^2}{(|w|^2 + |\theta|^2)^2} \sum_{j=1}^n dw_j \wedge d\overline{w_j}$$

and therefore, we have

$$\left(\sqrt{-1}\right)^{i} \left(\frac{|\theta|^{2}}{\left(|v_{1}|^{2}+|\theta|^{2}\right)^{2}}\right)^{i} |v_{1}|^{2i-2} dv_{1} \wedge d\overline{v_{1}} \wedge dv_{j_{1}} \wedge d\overline{v_{j_{1}}} \wedge \cdots \wedge dv_{j_{i-1}} \wedge d\overline{v_{j_{i-1}}}$$

$$\leq \left(\sqrt{-1}\right)^{i} \left(\frac{4n|\theta|^{2}}{\left(|w|^{2}+|\theta|^{2}\right)^{2}} \sum_{j=1}^{n} dw_{j} \wedge d\overline{w_{j}}\right) \wedge dv_{j_{1}} \wedge d\overline{v_{j_{1}}} \wedge \cdots \wedge dv_{j_{i-1}} \wedge d\overline{v_{j_{i-1}}}$$

With respect to  $v_1, \ldots, v_n$ ,  $dd^c u$  is written as

$$dd^{c}u = \frac{1}{2}dd^{c}\log\left(1 + \sum_{j=2}^{n} |v_{j}|^{2}\right)$$

and on a bounded domain,  $\frac{1}{2}dd^c \log \left(1 + \sum_{j=2}^n |v_j|^2\right)$  is equivalent to  $dd^c \left(\sum_{j=2}^n |v_j|^2\right)$ . Hence, we see that there exists a constant  $C_1 > 0$  such that

$$\left(\sqrt{-1}\right)^{i} \left(\frac{|\theta|^{2}}{\left(|v_{1}|^{2}+|\theta|^{2}\right)^{2}}\right)^{i} |v_{1}|^{2i-2} dv_{1} \wedge d\overline{v_{1}} \wedge dv_{k_{1}} \wedge d\overline{v_{k_{1}}} \wedge \cdots dv_{k_{i-1}} \wedge d\overline{v_{k_{i-1}}}$$
$$\leq C_{1} \left(\frac{|\theta|^{2}}{\left(|w|^{2}+|\theta|^{2}\right)^{2}} \sum_{j=1}^{n} \sqrt{-1} dw_{j} \wedge d\overline{w_{j}}\right) \wedge (dd^{c}u)^{i-1}.$$

In the case of (2), we have the following inequality. For k = 2, ..., n, we have

$$2\sqrt{-1}dw_j \wedge d\overline{w_j} + 2\sqrt{-1}|v_j|^2 dv_1 \wedge d\overline{v_1}$$
  
=  $\sqrt{-1}|v_1|^2 dv_j \wedge d\overline{v_j} + \sqrt{-1} \left(|v_1|^2 dv_j \wedge d\overline{v_j} + 2v_1\overline{v_j}dv_j \wedge d\overline{v_1} + 2v_j\overline{v_1}dv_1 \wedge d\overline{v_j} + 4|v_j|^2 dv_1 \wedge dv_1\right)$   
=  $\sqrt{-1}|v_1|^2 dv_j \wedge d\overline{v_j} + \sqrt{-1}(v_1dv_j + 2v_jdv_1) \wedge (\overline{v_1}d\overline{v_j} + 2\overline{v_j}d\overline{v_1}) \geq \sqrt{-1}|v_1|^2 dv_j \wedge d\overline{v_j},$ 

where the positivity is the strong positivity for forms. We have  $|v_k| < 2$  for k = 2, ..., n. Arguing as in the case of (1), we obtain from the above inequality that there exists  $C_2 > 0$  such that

$$\left(\sqrt{-1}\right)^{i} \left(\frac{|\theta|^{2}}{\left(|v_{1}|^{2}+|\theta|^{2}\right)^{2}}\right)^{i} |v_{1}|^{2i} dv_{j_{1}} \wedge d\overline{v_{k_{1}}} \wedge \cdots dv_{j_{i}} \wedge d\overline{v_{j_{i}}}$$

$$\leq \left(\sqrt{-1}\right)^{i} \left(\frac{|\theta|^{2}}{\left(|v_{1}|^{2}+|\theta|^{2}\right)^{2}}\right) |v_{1}|^{2} dv_{j_{1}} \wedge d\overline{v_{j_{1}}} \wedge \cdots dv_{j_{i}} \wedge d\overline{v_{j_{i}}}$$

$$\leq C_{2} \left(\frac{|\theta|^{2}}{\left(|w|^{2}+|\theta|^{2}\right)^{2}} \sum_{j=1}^{n} \sqrt{-1} dw_{j} \wedge d\overline{w_{j}}\right) \wedge (dd^{c}u)^{i-1}.$$

Combining the two inequalities proves the claim. Notice that the constant  $C_1$  and  $C_2$  are independent of  $\theta$ .

# 3. INTEGRABILITY CONDITIONS AND TANGENT CURRENTS

In this section, we introduce integrability conditions, which are related to the existence, uniqueness and *h*-dimension of tangent currents. Let U be a bounded simply connected domain with smooth boundary in  $\mathbb{C}^N$  and V a complex submanifold of U of codimension n.

3.1. The case of n < N. We continue to employ the same notations as in Section 2. Let *T* be a positive closed current of bidegree (p, p) on *U*. We continue to assume n < N and  $1 \le p \le N - n$ .

Let  $1 \leq i \leq n$  be an integer and  $\varphi_{N-p-i}$  a real smooth form of bidegree (N-p-i, N-p-i) on V. We define two products of  $\mathcal{K}^i$ , T and  $\pi_V^* \varphi_{N-p-i}$ :  $\langle T \wedge \mathcal{K}^i \wedge \pi_V^* \varphi_{N-p-i} \rangle_{\mathcal{K}}$  and  $\langle T \wedge \mathcal{K}^i \wedge \pi_V^* \varphi_{N-p-i} \rangle_{\mathcal{T}}$  below.

**Definition 3.1.** For a positive smooth form  $\varphi_{N-p-i}$  of bidegree (N - p - i, N - p - i), the current  $\langle T \wedge \mathcal{K}^i \wedge \pi_V^* \varphi_{N-p-i} \rangle_{\mathcal{K}}$  of maximal bidegree on U is defined to be the limit

$$\langle T \wedge \mathcal{K}^i \wedge \pi_V^* \varphi_{N-p-i} \rangle_{\mathcal{K}} \coloneqq \lim_{|\theta| \to 0} T \wedge \mathcal{K}^i_{\theta} \wedge \pi_V^* \varphi_{N-p-i}$$

in the sense of currents, provided that the limit exists. For general  $\varphi_{N-p-i}$ , we express  $\varphi_{N-p-i} = \varphi_{N-p-i}^+ - \varphi_{N-p-i}^-$ , where  $\varphi_{N-p-i}^{\pm}$  are positive smooth forms of bidegree (N - p - i, N - p - i) on V. We define

$$\left\langle T \wedge \mathcal{K}^{i} \wedge \pi_{V}^{*} \varphi_{N-p-i} \right\rangle_{\mathcal{K}} \coloneqq \left\langle T \wedge \mathcal{K}^{i} \wedge \pi_{V}^{*} \varphi_{N-p-i}^{+} \right\rangle_{\mathcal{K}} - \left\langle T \wedge \mathcal{K}^{i} \wedge \pi_{V}^{*} \varphi_{N-p-i}^{-} \right\rangle_{\mathcal{K}},$$

provided that both limits  $\langle T \wedge \mathcal{K}^i \wedge \pi_V^* \varphi_{N-p-i}^{\pm} \rangle_{\mathcal{K}}$  exist.

Note that since the support of  $dd^c u_{\theta}^{\mathcal{K}}$  is away from the set V, the product inside the limit is well defined. Observe that outside V, we have

$$\langle T \wedge \mathcal{K}^i \wedge \pi_V^* \varphi_{N-p-i} \rangle_{\mathcal{K}} = T \wedge \mathcal{K}^i \wedge \pi_V^* \varphi_{N-p-i}.$$

In the lemma below, we define the product  $\langle T \wedge \mathcal{K}^i \wedge \pi_V^* \varphi_{N-p-i} \rangle_{\mathcal{T}}$  under a certain integrability condition.

**Lemma 3.2.** Let  $1 \le i \le n$  be an integer. Suppose that the current  $\langle T \land \mathcal{K}^{i-1} \land \pi_V^* \omega_V^{N-p-i+1} \rangle_{\mathcal{K}}$  is well defined and that

$$u \in L^1_{loc}\left(\left\langle T \wedge \mathcal{K}^{i-1} \wedge \pi_V^* \omega_V^{N-p-i+1} \right\rangle_{\mathcal{K}}\right).$$

Then,  $(T \wedge \mathcal{K}^i \wedge \pi_V^* \varphi_{N-p-i})_{\mathcal{K}}$  is well defined for every real smooth form  $\varphi_{N-p-i}$  of bidegree (N - p - i, N - p - i). Also, for  $i \ge 2$  and for each  $\theta \in \mathbb{C}^*$ , the limit

$$\left(\left(T \wedge dd^{c}u_{\theta}^{\mathcal{T}}\right) \wedge \mathcal{K}^{i-1} \wedge \pi_{V}^{*}\varphi_{N-p-i}\right)_{\mathcal{K}} = \lim_{|\theta'| \to 0} \left(T \wedge dd^{c}u_{\theta}^{\mathcal{T}}\right) \wedge \mathcal{K}_{\theta'}^{i-1} \wedge \pi_{V}^{*}\varphi_{N-p-i}$$

exists and furthermore, we have

$$\left\langle T \wedge \mathcal{K}^{i} \wedge \pi_{V}^{*} \varphi_{N-p-i} \right\rangle_{\mathcal{T}} \coloneqq \lim_{|\theta| \to 0} \left\langle \left( T \wedge dd^{c} u_{\theta}^{\mathcal{T}} \right) \wedge \mathcal{K}^{k-1} \wedge \pi_{V}^{*} \varphi_{N-p-i} \right\rangle_{\mathcal{K}} \\ = \left\langle T \wedge \mathcal{K}^{i} \wedge \pi_{V}^{*} \varphi_{N-p-i} \right\rangle_{\mathcal{K}}.$$

Remark 3.3. We also have

$$\left(\left(T \wedge dd^{c} u_{\theta}^{\mathcal{T}}\right) \wedge \mathcal{K}^{i-1} \wedge \pi_{V}^{*} \varphi_{N-p-i}\right)_{\mathcal{K}} = T \wedge dd^{c} u_{\theta}^{\mathcal{T}} \wedge \mathcal{K}^{i-1} \wedge \pi_{V}^{*} \varphi_{N-p-i}$$

on  $U \smallsetminus V$ .

We need the following lemma.

**Lemma 3.4.** Let the sets U and V and the map  $\pi_V$  be as in Section 2. Let T' be a positive closed current of maximal bidegree on U. Namely, T' is a Radon measure. Let  $T'|_V := \mathbf{1}_V T'$  be the restriction of T' to V. Then, for every test function f on U, we have

$$\left\langle T'|_V, f \right\rangle = \left\langle T'|_V, \pi_V^* f|_V \right\rangle.$$

*Proof.* Let  $\varepsilon > 0$  be given. Then, we have

$$\left|\int_{V_{\varepsilon}} (f - \pi_V^* f|_V) T'\right| \leq \sup_{V_{\varepsilon}} |f - \pi_V^* f|_V |\int_{V_{\varepsilon} \cap \operatorname{supp} f} T' \leq \varepsilon \|f\|_{C^1} \int_{V_{\varepsilon} \cap \operatorname{supp} f} T',$$

where  $V_{\varepsilon}$  is the  $\varepsilon$ -neighborhood of V in U. Hence, as  $\varepsilon$  shrinks to 0, we obtain the equality.  $\Box$ 

Proof of Lemma 3.2. We first prove that the sequence

$$(T \wedge \mathcal{K}^{i}_{\theta} \wedge \pi^{*}_{V} \varphi_{N-p-i})_{0 < |\theta| \ll 1}$$

has locally uniformly bounded mass. Let  $\chi_V : \mathbb{C}^{N-n} \to [0,1]$  and  $\chi_F : \mathbb{C}^n \to [0,1]$  be smooth function with compact support such that  $\chi_V \equiv 1$  and  $\chi_F \equiv 1$  in a neighborhood of 0 and that  $\operatorname{supp} \pi_V^* \chi_V \cap \operatorname{supp} \pi_F^* \chi_F$  is a compact subset of U. Here, for  $\delta > 0$ , the set  $\{\pi_F^* \chi_F < \delta\}$  is a neighborhood of V. Due to the positivity, we may assume that  $\varphi_{N-p-i} = \omega_V^{N-p-i}$ . When i = 1, our claim is straightforward. Assume that  $i \geq 2$ . For  $\theta' \in \mathbb{C}^*$  with  $|\theta'| \ll |\theta|$ , We have

$$\int (\pi_V^* \chi_V) (\pi_F^* \chi_F) T \wedge \mathcal{K}_{\theta}^i \wedge \pi_V^* \omega_V^{N-p-i} = \int (\pi_V^* \chi_V) (\pi_F^* \chi_F) T \wedge dd^c u_{\theta}^{\mathcal{K}} \wedge \mathcal{K}_{\theta'}^{i-1} \wedge \pi_V^* \omega_V^{N-p-i}$$

$$= \int u_{\theta}^{\mathcal{K}} (\pi_F^* \chi_F) dd^c (\pi_V^* \chi_V) \wedge T \wedge \mathcal{K}_{\theta'}^{i-1} \wedge \pi_V^* \omega_V^{N-p-i}$$

$$+ \int u_{\theta}^{\mathcal{K}} d(\pi_F^* \chi_F) \wedge d^c (\pi_F^* \chi_F) \wedge T \wedge \mathcal{K}_{\theta'}^{i-1} \wedge \pi_V^* \omega_V^{N-p-i}$$

$$+ \int u_{\theta}^{\mathcal{K}} d(\pi_F^* \chi_F) \wedge d^c (\pi_F^* \chi_F) \wedge T \wedge \mathcal{K}_{\theta'}^{i-1} \wedge \pi_V^* \omega_V^{N-p-i}$$

$$+ \int u_{\theta}^{\mathcal{K}} (\pi_V^* \chi_V) dd^c (\pi_F^* \chi_F) \wedge T \wedge \mathcal{K}_{\theta'}^{i-1} \wedge \pi_V^* \omega_V^{N-p-i}.$$

Every factor except *T* in the last three integrals is smooth and the support of each current is a definite distance away from the set *V* independently of  $\theta$ . We first let  $|\theta'| \to 0$  and then  $|\theta| \to 0$ . We see the last three integrals converge. Hence, for all  $\theta$  with  $0 < |\theta| \ll 1$ , the last three integrals are uniformly bounded. We look into the first integral. There exists a constant  $c_{\chi_1} > 0$  such that  $c_{\chi_1}\omega_V \ge dd^c \pi_V^* \chi_V$  over the set  $\sup p \pi_V^* \chi_V \cap \pi_F^* \chi_F$ . Hence, the measure

$$u_{ heta}^{\mathcal{K}}(\pi_F^*\chi_F)dd^c(\pi_V^*\chi_V)\wedge T\wedge \mathcal{K}_{ heta'}^{i-1}\wedge \pi_V^*\omega_V^{N-p-i}$$

Let  $\varphi_{N-p-i}$  be a real smooth form of bidegree (N - p - i, N - p - i). Without loss of generality, we may assume that  $\varphi_{N-p-i}$  is positive. We show that there exists a unique limit current for the family  $(T \wedge \mathcal{K}^i_{\theta} \wedge \pi^*_V \varphi_{N-p-i})_{0 < |\theta| \ll 1}$ . When i = 1, it is obvious. So, we may assume that  $i \ge 2$ . Observe that every limit current coincides on  $U \setminus V$ . Hence, we are interested in the restriction to the set V of the limit currents. The limit currents are positive and therfore, currents of order 0, which means the current depends only on the value of the test function on the set. More precisely, we have the following. Let L be a limit current and  $(\theta_j)_{j \in \mathbb{N}}$  a sequence corresponding to the limit current L. Let  $f: U \to \mathbb{R}$  be a smooth test function with compact support. Let  $\varepsilon > 0$  be given and  $V_{\varepsilon}$  a  $\varepsilon$ -neighborhood of V. Then, for some C > 0, we have

$$\left|\int_{V_{\varepsilon}} \left(f - \pi_{V}^{*}f|_{V}\right) T \wedge \mathcal{K}_{\theta_{j}}^{i} \wedge \pi_{V}^{*}\varphi_{N-p-i}\right| \leq \varepsilon \|f\|_{C^{1}} \int_{V_{\varepsilon}} T \wedge \mathcal{K}_{\theta_{j}}^{i} \wedge \pi_{V}^{*}\varphi_{N-p-i} \leq C\varepsilon.$$

The last inequality comes from the uniform boundedness of the mass proven previously. By letting  $\varepsilon \to 0$ , we see that the restrictions of fL and  $\pi_V^* f_V L$  to V are the same. Hence, it suffices to check whether the sequence  $(\pi_V)_* \left( \mathbf{1}_{V_{\varepsilon_0}} T \wedge \mathcal{K}^i_{\theta_j} \wedge \pi_V^* \varphi_{N-p-i} \right)_{j \in \mathbb{N}}$  of currents of maximal bidegree is uniquely determined as  $n \to \infty$  for some  $\varepsilon_0 > 0$ .

Let  $g: V \to \mathbb{R}$  be a smooth test function with compact support. Since we are dealing with the neighborhood  $V_{\varepsilon_0}$  of V, we may assume that T has finite mass over the intersection of the support of  $\pi_V^*g$  and  $V_{\varepsilon_0}$ . Choose  $\chi_F: \mathbb{C}^n \to [0,1]$  so that the support of  $\pi_F^*\chi_F \subset V_{\varepsilon_0}$ . We have

(3.1)  

$$\int_{V_{\varepsilon_0}} (\pi_V^*g) T \wedge \mathcal{K}^i_{\theta_j} \wedge \pi_V^*\varphi_{N-p-i}$$

$$= \int (\pi_F^*\chi_F) (\pi_V^*g) T \wedge \mathcal{K}^i_{\theta_j} \wedge \pi_V^*\varphi_{N-p-i}$$

$$(3.2) \qquad \qquad + \int_{V_{\varepsilon_0}} \left(1 - (\pi_F^* \chi_F)\right) (\pi_V^* g) T \wedge \mathcal{K}^i_{\theta_j} \wedge \pi_V^* \varphi_{N-p-i}$$

As previously, for  $\theta'$  with  $0 < |\theta'| \ll |\theta|$ , we have

$$(3.1) = \int (\pi_F^* \chi_F) (\pi_V^* g) T \wedge \mathcal{K}_{\theta_j}^i \wedge \pi_V^* \varphi_{N-p-i} = \int (\pi_F^* \chi_F) (\pi_V^* g) T \wedge dd^c u_{\theta_j}^{\mathcal{K}} \wedge \mathcal{K}_{\theta'}^{i-1} \wedge \pi_V^* \varphi_{N-p-i}$$

$$= \int u_{\theta_j}^{\mathcal{K}} (\pi_F^* \chi_F) dd^c (\pi_V^* g) \wedge T \wedge \mathcal{K}_{\theta'}^{i-1} \wedge \pi_V^* \varphi_{N-p-i}$$

$$+ \int u_{\theta_j}^{\mathcal{K}} d (\pi_V^* g) \wedge d^c (\pi_F^* \chi_F) \wedge T \wedge \mathcal{K}_{\theta'}^{i-1} \wedge \pi_V^* \varphi_{N-p-i}$$

$$+ \int u_{\theta_j}^{\mathcal{K}} d (\pi_V^* g) dd^c (\pi_F^* \chi_F) \wedge T \wedge \mathcal{K}_{\theta'}^{i-1} \wedge \pi_V^* \varphi_{N-p-i}$$

As previously, since the support of every partial derivative of  $\pi_F^*\chi_F$  is a definite distance away from V independently from  $\theta'$  and j, the last three integrals converges as  $|\theta'| \to 0$  and  $j \to \infty$  in this order. By the integrability condition, the first integral is also converges as  $|\theta'| \to 0$ . Then we let  $j \to \infty$  and we see the first integral also converges.

In (3.2), the support of  $1 - (\pi_F^* \chi_F)$  is a definite distance away from V independently of  $\theta$ . Hence, over the support of g, the integral converges. Indeed,  $u - u_{\theta_j}^{\mathcal{K}}$  converges to 0 locally uniformly on  $U \setminus V$  and T is assumed to have finite mass over the support of  $\pi_V^* g$ .

The assertions related to  $u^{\mathcal{T}}_{\theta}$  can be proved exactly in the same way as above.

Below we introduce integrability conditions, which we will refer to as Condition (I).

**Definition 3.5** (Condition (I)). Let T be a positive closed current of bidegree (p,p) on U. Let  $1 \le m \le n$  be an integer. We say that the current T satisfies Condition (I) with respect to V up to codimension m if

$$u \in L^1_{loc}\left(\left\langle T \wedge (dd^c u)^{i-1} \wedge \pi_V^* \omega_V^{N-p-i+1} \right\rangle_{\mathcal{K}}\right)$$

inductively holds from i = 1 through m. When m = n, the current T is said to satisfy the horizontal condition with respect to V.

The following proposition is straightforward.

**Proposition 3.6.** Let T be a positive closed current of bidegree (p, p) on U satisfying Condition (I) with respect to V up to codimension m for some  $1 \le m \le n$ . Let  $\varphi$  be a real smooth closed form of bidegree (q,q) on V for some  $1 \le q \le N - p - n$ . Then,  $T \land \pi_V^* \varphi$  satisfies Condition (I) with respect to V up to codimension m.

The next theorem finds a relationship between the *h*-dimension and Condition (I).

**Proposition 3.7.** Let  $1 \le m \le n$  be an integer. Let T be a positive closed current of bidegree (p, p) on U. Suppose that a tangent current  $T_{\infty}$  of T along V exists. Then, if the current T satisfies Condition (1) with respect to V up to codimension m then the h-dimension of the tangent current  $T_{\infty}$  along V is at most N - p - m. In particular, if T satisfies the horizontal condition with respect to V, then the h-dimension of  $T_{\infty}$  is minimal.

*Proof.* Lemma 2.18 and Lemma 3.2 imply that the family  $\left(T \wedge \mathcal{T}_{\theta}^{i} \wedge \pi_{V}^{*} \omega_{V}^{N-p-i}\right)_{0 < |\theta| \ll 1}$  has locally uniformly bounded mass for i = 1, ..., m. Moreover, for any limit current  $T_{\mathcal{T}}^{p+i}$  of the family  $\left(T \wedge \mathcal{T}_{\theta}^{i}\right)_{0 < |\theta| \ll 1}$ , we have

$$T_{\mathcal{T}}^{p+i} \wedge \pi_{V}^{*} \omega_{V}^{N-p-i} \leq C \left\langle T \wedge \mathcal{K}^{i} \wedge \pi_{V}^{*} \omega_{V}^{N-p-i} \right\rangle_{\mathcal{K}}$$

in the sense of currents, where C > 0 is the constant as in Lemma 2.18. From the integrability condition, we see that  $\left(T \wedge \mathcal{K}^i \wedge \pi_V^* \omega_V^{N-p-i}\right)_{\mathcal{K}}$  has no mass on V for  $i = 1, \ldots, m-1$ , and so does  $T_{\mathcal{T}}^{p+i} \wedge \pi_V^* \omega_V^{N-p-i}$  for  $i = 1, \ldots, m-1$ . Then, by Corollary 2.15, the *h*-dimension of  $T_{\infty}$  is at most N - p - m.

We consider the relationship between the horizontal condition and the existence of tangent currents.

**Proposition 3.8.** Let T be a positive closed current of bidegree (p, p) on U. Suppose that the current T satisfies the horizontal condition. Then, tangent currents exist and their h-dimension is minimal.

*Proof.* Since T satisfies the horizontal condition, the family  $(\langle (T \wedge dd^c u_{\theta}^T) \wedge \mathcal{K}^{i-1} \wedge \pi_V^* \omega^{N-p-i} \rangle_{\mathcal{K}})$  has locally uniformly bounded mass for i = 1, ..., n. By Lemma 2.18, we see that the family  $(T \wedge \mathcal{T}_{\theta}^i \wedge \pi_V^* \omega^{N-p-i})$  has uniformly bounded mass for i = 1, ..., n. By Proposition 2.17, T has a tangent current  $T_{\infty}$  along V. Proposition 3.7 implies that their h-dimension is minimal.

Now, we investigate the relationship between the Condition (I) and the uniqueness of tangent currents. We prove Theorem 1.3.

We first recall the precise statement of [9, Proposition 3.5] in our case. Note that the h-dimension in Definition 2.2 and the shadow in Definition 2.5 are actually defined for V-conic positive closed currents defined on projective bundles. For the V-conic positive closed current, see [9, Definition 3.9].

**Proposition 3.9** (Proposition 3.5 in [9]). Let T' be a positive closed current of bidegree (p, p) on  $\overline{E}$ and  $h_{T'}$  the h-dimension of T' over an open subset  $V_0 \subset V$ . Let  $\Omega$  be a smooth closed form of bidegree  $(N-p-h_{T'}, N-p-h_{T'})$  on  $\pi_V^{-1}(V_0)$  whose restriction to each fiber of  $\pi_V$  is cohomologous to a linear subspace in this fiber. Then the current  $(T')^h := (\pi_V)_* (T' \land \Omega)$  on  $V_0$  is positive closed of bidegree  $(N-n-h_{T'}, N-n-h_{T'})$  with support in  $\pi_V(\operatorname{supp}(T')) \cap V_0$ . Moreover, it does not depend on the choice of  $\Omega$ .

**Theorem 1.3.** Let T be a positive closed current of bidegree (p,p) on U. Suppose that the current T satisfies the horizontal condition. Then, there exists a unique tangent current  $T_{\infty}$  of T along V, and the h-dimension of the tangent current  $T_{\infty}$  is minimal. For a smooth test form  $\varphi$  of bidegree (N - p - n, N - p - n) on V, its shadow  $T_{\infty}^h$  is computed as

$$\langle T_{\infty}^{h},\varphi\rangle = \langle \langle T \wedge \mathcal{K}^{n} \wedge \pi_{V}^{*}\varphi \rangle_{\mathcal{K}}, \pi_{F}^{*}\chi_{F}\rangle = \langle \langle T \wedge \mathcal{K}^{n} \wedge \pi_{V}^{*}\varphi \rangle_{\mathcal{K}}, 1\rangle = \int_{U \smallsetminus V} T \wedge \left( u \left( dd^{c}u \right)^{n-1} \right) \wedge dd^{c}\Phi,$$

where  $\chi_F : \mathbb{C}^n \to [0,1]$  is a smooth function with compact support such that  $\operatorname{supp} \pi_V^* \varphi \cap \operatorname{supp} \pi_F^* \chi_F \in U$  and  $\chi_F \equiv 1$  in a neighborhood of 0, and  $\Phi$  is a smooth form of bidegree (N - p - n, N - p - n) on U with compact support such that  $\Phi = \pi_V^* \varphi$  in a neighborhood of V. The value of the pairing is independent of  $\chi_F$  and the value of the integral is independent of the choice of  $\Phi$ .

We first prove the following proposition.

**Proposition 3.10.** Let T and  $\varphi$  be as in Theorem 1.3. Let  $\chi_F : \mathbb{C}^n \to [0,1]$  be a smooth function with compact support such that  $\operatorname{supp} \pi_V^* \varphi \cap \operatorname{supp} \pi_F^* \chi_F \in U$  and  $\chi_F \equiv 1$  in a neighborhood of 0. Then, the current  $\varphi \to \langle \langle T \wedge \mathcal{K}^n \wedge \pi_V^* \varphi \rangle_{\mathcal{K}}, \pi_F^* \chi_F \rangle$  for a smooth test form  $\varphi$  of bidegree (N - p - n, N - p - n)on V is a well defined positive closed current of bidegree (p, p) on V. Moreover, we have

$$\langle\langle T \wedge \mathcal{K}^n \wedge \pi_V^* \varphi \rangle_{\mathcal{K}}, \pi_F^* \chi_F \rangle = \langle\langle T \wedge \mathcal{K}^n \wedge \pi_V^* \varphi \rangle_{\mathcal{K}}, 1 \rangle = \int_{U \smallsetminus V} T \wedge \left( u \left( dd^c u \right)^{n-1} \right) \wedge dd^c \Phi,$$

where  $\Phi$  is a smooth form of bidegree (N - p - n, N - p - n) on U with compact support such that  $\Phi = \pi_V^* \varphi$  in a neighborhood of V. The integral is independent of the choice of  $\Phi$ .

*Proof.* By Lemma 3.2, for each smooth test form  $\varphi$  of bidegree (N - p - n, N - p - n), the current  $\langle T \wedge \mathcal{K}^n \wedge \pi_V^* \varphi \rangle_{\mathcal{K}}$  of maximal bidegree is well defined. Observe that the support of the current is inside  $\operatorname{supp} \pi_V^* \varphi \cap V$ , which is a proper subset of V and is a compact set. Hence, for such  $\chi_F$ , we have

$$\langle \langle T \wedge \mathcal{K}^n \wedge \pi_V^* \varphi \rangle_{\mathcal{K}}, \pi_F^* \chi_F \rangle = \langle \langle T \wedge \mathcal{K}^n \wedge \pi_V^* \varphi \rangle_{\mathcal{K}}, 1 \rangle$$

and it is a well defined definite number. Obviously, it is linear. The continuity of the function  $\varphi \rightarrow \langle \langle T \wedge \mathcal{K}^n \wedge \pi_V^* \varphi \rangle_{\mathcal{K}}, \pi_F^* \chi_F \rangle$  with respect to the  $\| \cdot \|_{\infty}$  norm is also quite straightforward. Hence, it is a well defined current.

By definition, we have

$$\left\langle \left\langle T \wedge \mathcal{K}^n \wedge \pi_V^* \varphi \right\rangle_{\mathcal{K}}, \pi_F^* \chi_F \right\rangle = \lim_{|\theta| \to 0} \int T \wedge \mathcal{K}_{\theta}^n \wedge \left(\pi_F^* \chi_F\right) \left(\pi_V^* \varphi\right),$$

from which we can see that the current is positive and closed.

Further, for  $\theta' \in \mathbb{C}^*$  with  $|\theta'| \ll |\theta|$ , we get

$$\int T \wedge \mathcal{K}_{\theta}^{n} \wedge (\pi_{F}^{*}\chi_{F}) (\pi_{V}^{*}\varphi) = \int T \wedge dd^{c}u_{\theta}^{\mathcal{K}} \wedge \mathcal{K}_{\theta'}^{n-1} \wedge (\pi_{F}^{*}\chi_{F}) (\pi_{V}^{*}\varphi)$$

$$= \int u_{\theta}^{\mathcal{K}} dd^{c} ((\pi_{F}^{*}\chi_{F}) (\pi_{V}^{*}\varphi)) \wedge T \wedge \mathcal{K}_{\theta'}^{n-1}$$

$$= \int u_{\theta}^{\mathcal{K}} (\pi_{F}^{*}\chi_{F}) dd^{c} (\pi_{V}^{*}\varphi) \wedge T \wedge \mathcal{K}_{\theta'}^{n-1} + \int u_{\theta}^{\mathcal{K}} d(\pi_{F}^{*}\chi_{F}) \wedge d^{c} (\pi_{V}^{*}\varphi) \wedge T \wedge \mathcal{K}_{\theta'}^{n-1}$$

$$+ \int u_{\theta}^{\mathcal{K}} d(\pi_{V}^{*}\varphi) \wedge d^{c} (\pi_{F}^{*}\chi_{F}) \wedge T \wedge \mathcal{K}_{\theta'}^{n-1} + \int u_{\theta}^{\mathcal{K}} (\pi_{V}^{*}\varphi) dd^{c} (\pi_{F}^{*}\chi_{F}) \wedge T \wedge \mathcal{K}_{\theta'}^{n-1}.$$

As discussed previously in the proof of Lemma 3.2, the last three integrals converges. For the first integral, from the horizontal condition, the measure  $\langle T \wedge \mathcal{K}^{n-1} \wedge dd^c(\pi_V^*\varphi) \rangle_{\mathcal{K}}$  has no mass on V. So, as  $|\theta'| \to 0$  and  $|\theta| \to 0$  in this order, the integral converges and we obtain

$$\langle \langle T \wedge \mathcal{K}^n \wedge \pi_V^* \varphi \rangle_{\mathcal{K}}, \pi_F^* \chi_F \rangle = \int_{U \setminus V} T \wedge \left( u \left( dd^c u \right)^{n-1} \right) \wedge dd^c \left( \pi_F^* \chi_F \pi_V^* \varphi \right).$$

We prove the independence of the value of the integral from the choice of  $\Phi$ . Let  $\Phi'$  denote another smooth form of bidegree (N - p - n, N - p - n) on U with compact support such that  $\Phi' = \pi_V^* \varphi$  in a neighborhood of V. So, in a neighborhood of V,  $\Phi = \Phi'$ . Also, from the horizontal condition, we have the following convergence:

$$\int_{U} T \wedge \mathcal{K}^{n}_{\theta} \wedge \Phi = \int_{U \setminus V} T \wedge \left( u^{\mathcal{K}}_{\theta} (dd^{c}u)^{n-1} \right) \wedge dd^{c} \Phi \to \int_{U \setminus V} T \wedge \left( u(dd^{c}u)^{n-1} \right) \wedge dd^{c} \Phi$$

as  $|\theta| \to 0$ . The support of  $T \wedge \mathcal{K}^n_{\theta}$  shrinks to a subset of V. Therefore, for all  $\theta$  with sufficiently small  $|\theta|$ , we have

$$\int_{U} T \wedge \mathcal{K}_{\theta}^{n} \wedge \Phi = \int_{U} T \wedge \mathcal{K}_{\theta}^{n} \wedge \Phi'$$

Simply letting  $|\theta| \to 0$  completes the proof.

We extract from the above proof the following. Since the support of  $\mathcal{K}_{\theta}^{n}$  is very close to a subset of V, the support of  $T \wedge \mathcal{K}_{\theta}^{n} \wedge \pi_{V}^{*} \varphi$  is compact in U and lies inside a small neighborhood of V. So, we can write

$$(3.3) \qquad \langle \langle T \wedge \mathcal{K}^n \wedge \pi_V^* \varphi \rangle_{\mathcal{K}}, 1 \rangle = \lim_{|\theta| \to 0} \langle T \wedge \mathcal{K}^n_{\theta} \wedge \pi_V^* \varphi, 1 \rangle = \lim_{|\theta| \to 0} \int_U T \wedge \mathcal{K}^n_{\theta} \wedge \pi_V^* \varphi$$
$$= \lim_{|\theta| \to 0} \int_V (\pi_V)_* (T \wedge \mathcal{K}^n_{\theta}) \wedge \varphi.$$

Observe that for bidegree reason, we have  $(T \wedge \mathcal{K}^n_{\theta}) \wedge \psi = 0$  if  $\psi$  is not a linear combination of smooth forms of the form  $f\pi^*_V \varphi$ , where f is a smooth function on U and  $\varphi$  is a smooth test form of bidegree (N - p - n, N - p - n) on V. So, (3.3) implies that the family  $(T \wedge \mathcal{K}^n_{\theta})_{0 < |\theta| \ll 1}$  has locally uniformly bounded mass. The limit currents of the family  $(T \wedge \mathcal{K}^n_{\theta})_{0 < |\theta| \ll 1}$  has support in V. Hence, Lemma 3.4 implies that (3.3) proves the convergence of  $(\langle T \wedge \mathcal{K}^n_{\theta}, f\pi^*_V \varphi \rangle)_{0 < |\theta| \ll 1}$  as  $|\theta| \to 0$ , where f is a smooth function on U and  $\varphi$  is a smooth test form of bidegree (N - p - n, N - p - n) on V. Hence, summarizing the discussion, we obtain the following proposition.

**Proposition 3.11.** Let T be a positive closed current of bidegree (p,p) on U. Suppose that the current T satisfies the horizontal condition. Then, the family  $(T \wedge \mathcal{K}^n_{\theta})_{0 < |\theta| \ll 1}$  converges as  $|\theta| \to 0$ .

When T is smooth, we know that  $T \wedge \mathcal{K}^n_{\theta}$  converges to  $T \wedge [V]$ . Inspired by this, we define the following notation.

**Definition 3.12.** Let T be a positive closed current of bidegree (p,p) on U satisfying the horizontal condition. We define the current  $T \wedge_C [V]$  on U by

(3.4) 
$$T \wedge_C [V] := \langle T \wedge \mathcal{K}^n \rangle_{\mathcal{K}} = \lim_{|\theta| \to 0} T \wedge \mathcal{K}^n_{\theta}$$

in the sense of currents.

For the proof of Theorem 1.3, we also need the following proposition, which can be deduced from [9, Lemma 3.4].

**Proposition 3.13.** Suppose that the set of tangent currents of T along V is not empty and the h-dimension of every tangent current over V is minimal. The tangent current of T along V is unique if and only if there is only one shadow for tangent currents.

Proof of Theorem 1.3. Proposition 3.8 proves the existence of tangent currents and the minimality of the *h*-dimension. Let  $T_{\infty}$  be a tangent current and let  $(\lambda_k)_{k\in\mathbb{N}}$  be a sequence of non-zero complex numbers diverging to  $\infty$  such that  $((A_{\lambda_k})_* T)_{k\in\mathbb{N}}$  converges to  $T_{\infty}$ .

From Proposition 3.11, we have

$$T \wedge \mathcal{K}^n_\theta \to T \wedge_C [V]$$

as  $|\theta| \to 0$  on U in the sense of currents. Hence, from Corollary 2.12, we have

$$T_{\infty}^{h} = (\pi_{V})_{*} \left( T_{\infty} \wedge \pi_{F}^{*} \Omega_{n} \right) = (\pi_{V})_{*} \left( L_{T_{\infty}} \right) = (\pi_{V})_{*} \left( T \wedge_{C} \left[ V \right] \right),$$

which implies that there exists only one shadow for tangent currents. By Proposition 3.13, we prove the uniqueness of the tangent current and by Proposition 3.10, we obtain the desired expression.  $\Box$ 

Note that  $T_{\infty} \wedge [V]$  can be understood as a double current in de Rham's language ([3]).

**Proposition 3.14.** Suppose that T is a positive closed current of bidegree (p, p) with bounded mass on U satisfying the horizontal condition. Then, we have

$$T_{\infty} \wedge [V] = T \wedge_C [V]$$

in the sense of currents defined on U. Here, we identify currents in U with their trivially extended currents in E.

*Proof.* The current  $T_{\infty}$  is a current in x' where (x', x'') is a coordinate system of E. So, we have

$$(\pi_V)_* (T_{\infty} \wedge [V]) = (\pi_V)_* (T_{\infty} \wedge \Omega_n) = \lim_{|\lambda| \to \infty} (\pi_V)_* ((A_{\lambda})_* T \wedge \Omega_n)$$
$$= \lim_{|\lambda| \to \infty} (\pi_V)_* (T \wedge (A_{\lambda})^* \Omega_n) = \lim_{|\lambda| \to \infty} (\pi_V)_* (T \wedge \mathcal{K}_{e^{-M}|\lambda|^{-1}}^n) = (\pi_V)_* (T \wedge_C [V]),$$

where  $\Omega_n$  is the form as in Lemma 2.7. The last equality comes from the horizontal condition. Then, as in the arguments for Proposition 3.11, Lemma 3.4 implies the desired equality.

The following can be considered as a version of [9, Proposition 3.5 and Remark 4.9]. We will need this later for the study of the relationship between the horizontal condition and slicing theory. The proof itself is almost straightforward and so omitted.

**Proposition 3.15.** Let T be a positive closed current of bidegree (p, p) on U, which satisfies the horizontal condition with respect to V. Let  $\varphi$  be a smooth positive closed form of bidegree (N - p - n, N - p - n) on U such that  $\varphi|_V \neq 0$ . Then, the family  $((A_\lambda)_*(T \wedge \varphi))$  admits a tangent current  $(T \wedge \varphi)_{\infty}$  along V and is unique. The h-dimension of  $(T \wedge \varphi)_{\infty}$  is minimal. Furthermore, its shadow is a Radon measure that satisfies

$$(T \wedge \varphi)^h_\infty = T^h_\infty \wedge \varphi|_V.$$

3.2. **General case.** We introduce Condition (I'), which is slightly more restrictive but works more generally than Condition (I). Let *T* be a positive closed current of bidegree (p,p) on *U*. Now, we allow N = n and also N - n < p.

Here, we are considering families  $(T \wedge \mathcal{K}^m_{\theta})_{0 < |\theta| \ll 1}$  and  $(T \wedge \mathcal{T}^m_{\theta})_{0 < |\theta| \ll 1}$  for  $1 \le m \le \min\{N - p, n\}$  instead of the families  $(T \wedge \mathcal{K}^m_{\theta} \wedge \pi^*_V \omega_V^{N-p-m})_{0 < |\theta| \ll 1}$  and  $(T \wedge \mathcal{T}^m_{\theta} \wedge \pi^*_V \omega_V^{N-p-m})_{0 < |\theta| \ll 1}$  for  $1 \le m \le n \le N - p$ .

Corollary 2.10 actually works in this case as well.

**Proposition 3.16** (Corollary 2.10). Let T be a positive closed current of bidegree (p,p) on U with a tangent current  $T_{\infty}$  along V. Let m be an integer such that  $1 \le m \le \min\{N - p, n\}$ . Suppose that the family  $(T \land \mathcal{K}^m_{\theta})_{0 < |\theta| \ll 1}$  has locally uniformly bounded mass in U. Then, there exists a positive closed current  $T_{\mathcal{K}}^{p+m}$  of bidegree (p + m, p + m) on U such that  $T_{\mathcal{K}}^{p+m}$  is a limit current of the family  $(T \land \mathcal{K}^m_{\theta})_{0 < |\theta| \ll 1}$  and that we have

$$(\pi_V)_* \left( \mathbf{1}_V T_{\mathcal{K}}^{p+m} \right) = (\pi_V)_* \left( T_{\infty} \wedge \pi_F^* \Omega_m \right),$$

where  $\Omega_m$  is the smooth positive closed form of bidegree (m,m) as in Lemma 2.7. In particular, when N - p - m is the horizontal dimension  $h_T$ , that is,  $m = N - p - h_T$ , then  $(\pi_V)_* (\mathbf{1}_V T_{\mathcal{K}}^{p+m})$  equals the shadow  $T_{\infty}^h$  of  $T_{\infty}$ .

Other corresponding statements are as follows. Each proof is the same as that for the corresponding statement. The only difference is that we can use the limit currents of the families  $(T \wedge \mathcal{K}_{\theta}^m)_{0 < |\theta| \ll 1}$  and  $(T \wedge \mathcal{T}_{\theta}^m)_{0 < |\theta| \ll 1}$  for  $1 \le m \le \min\{N - p, n\}$ .

**Proposition 3.17** (Corollary 2.15). Let T be a positive closed current of bidegree (p, p) on U with a tangent current  $T_{\infty}$  along V. Let m be an integer such that  $1 \le m \le \min\{N - p, n\} - 1$ . Suppose that the family  $(T \land \mathcal{T}^i_{\theta})_{0 < |\theta| \ll 1}$  has locally uniformly bounded mass in U for  $1 \le i \le m$  and that for each  $i = n - p, \ldots, m$ , there exists a limit current of  $(T \land \mathcal{T}^i_{\theta})_{0 < |\theta| \ll 1}$  having no mass on V. Then, the h-dimension of  $T_{\infty}$  is at most N - p - m - 1.

**Proposition 3.18** (Proposition 2.17). If the family  $(T \wedge \mathcal{T}^i_{\lambda^{-1}})_{\lambda \in \mathbb{C}^*}$  has locally uniformly bounded mass on U for  $i = 1, ..., \min\{N - p, n\}$ , then tangent currents of T along V exist.

**Definition 3.19.** Let T be a positive closed (p, p)-current on U. If  $u \in L^1_{loc}(||T||)$ , we define

$$\langle dd^c u \wedge T \rangle_C \coloneqq dd^c (uT).$$

Let  $1 \le i \le \min\{N - p, n\}$  be an integer. We inductively define

$$\left\langle \left(dd^{c}u\right)^{i}\wedge T\right\rangle _{C}\coloneqq dd^{c}\left(u\left\langle \left(dd^{c}u\right)^{i-1}\wedge T\right\rangle _{C}\right)$$

provided that  $\langle (dd^c u)^k \wedge T \rangle_C$  is well defined and  $u \in L^1_{loc} \left( \left\| \langle (dd^c u)^k \wedge T \rangle_C \right\| \right)$ for k = 0 through i-1. When  $n \leq N - p$  and m = n, we define the product denoted by  $\langle T \wedge [V] \rangle_C$  to be

$$\langle T \wedge [V] \rangle_C \coloneqq \langle (dd^c u)^n \wedge T \rangle_C$$

Notice that we have  $\langle T \wedge [V] \rangle_C = T \wedge_C [V]$  whenever both are defined. It can be shown from the regularization of [V].

**Definition 3.20** (Condition (I')). Let *T* be a positive closed current of bidegree (p, p) on *U*. Let  $1 \le m \le \min\{N - p, n\}$  be an integer. We say that the current *T* satisfies Condition (I') with respect to *V* up to codimension *m* if

$$u \in L^1_{loc}\left(\left\langle T \wedge (dd^c u)^{i-1} \right\rangle_C\right)$$

inductively holds from i = 1 through m. When  $m = \min\{N - p, n\}$ , the current T is said to satisfy the minimally vertical condition with respect to V.

**Theorem 3.21** (Theorem 1.3). Let  $u = \log |x''|$  for  $(x', x'') \in U$  with  $V = \{x'' = 0\}$ . Suppose that the current T satisfies the minimally vertical condition. Then, there exists a unique tangent current  $T_{\infty}$  of T along V, and the h-dimension of the tangent current  $T_{\infty}$  is minimal. Its shadow is computed as in Proposition 3.16.

# 4. REGULARIZATIONS OF POSITIVE CLOSED CURRENTS

In this section, we introduce regularizations of positive closed currents. Throughout this section, we consider a bounded simply connected domain D with smooth boundary in  $\mathbb{C}^n$ . We use x for the coordinates of D and  $\omega_D$  for the standard Kähler form  $dd^c |x|^2$ . For an integer  $1 \le k \le n$ ,  $\mathscr{C}_k(D)$  denotes the set of positive closed currents of bidegree (k, k) defined on D and  $\mathscr{C}_k(\overline{D})$  the set of positive closed currents of bidegree (k, k) defined on D and  $\mathscr{C}_k(\overline{D})$  the set of positive closed currents of (k, k) defined in a neighborhood of  $\overline{D}$ . Let  $\pi_i : D \times D \to D$  be the two canonical projections onto the *i*-th factor for each i = 1, 2. We will use (x, y) for the coordinates of  $D \times D$  so that we have  $\pi_1(x, y) = x$  and  $\pi_2(x, y) = y$ . We will also use the Kähler form  $\omega_{x,y} = \pi_1^* \omega_D + \pi_2^* \omega_D$ .

4.1. **Semi-regular transformations on a domain.** Semi-regular transforms on compact Kähler manifolds were introduced and used by Dinh-Sibony in [8]. Here, we consider the essentially same notion on domains. For our purpose, instead of defining the notion using blown-up, we define the notion in terms of the regularity property. We start by justifying the notions to be used.

Let  $0 \le q \le n$  be an integer. Let Q be a form of bidegree (q,q) on  $\mathbb{C}^n \times \mathbb{C}^n$  smooth outside  $\Delta$  such that for some constant  $M_Q > 0$ , we have

$$(4.1) \qquad Q \le M_Q \omega_{x,y}, \ |Q| \le -M_Q [\log \operatorname{dist}(\cdot, \Delta)] \operatorname{dist}(\cdot, \Delta)^{2-2n} \text{ and } |\nabla Q| \le M_Q \operatorname{dist}(\cdot, \Delta)^{1-2n}$$

where |Q| means the sum of the absolute values of the coefficients and  $|\nabla Q|$  the sum of the absolute values of the derivatives of the coefficients, and the first inequality is understood in the sense of currents.

The following proposition is quite straightforward.

**Proposition 4.1.** Let  $n - q \le s \le n$  be an integer. Let S be an  $L^{\infty}$ -form of bidegree (s,s) (not necessarily with compact support) in D. The form  $(\pi_2)_*(\pi_1^*S \land Q)$  is a well defined form of bidegree (s+q-n, s+q-n) and has  $C^1$  coefficients. If Q can be written as a form of  $x-y = (x_1-y_1, \ldots, x_n-y_n)$  and S has compact support, then  $(\pi_2)_*(\pi_1^*S \land Q)$  has  $C^{\infty}$  coefficients.

**Definition 4.2.** Let  $n - q \le s \le n$  be an integer. Let S be an  $L^{\infty}$ -form of bidegree (s,s) (not necessarily with compact support) in D. The semi-regular transform  $\mathscr{L}^Q(S)$  of S is a form of bidegree (s + q - n, s + q - n) with  $C^1$ -coefficients on  $\mathbb{C}^n$  defined by

$$\mathscr{L}^Q(S) \coloneqq (\pi_2)_*(\pi_1^* S \wedge Q).$$

Notice that we can easily see that for any smooth test form  $\varphi$  of bidegree (2n - s - q, 2n - s - q) on D, we have

$$\langle \mathscr{L}^Q(S), \varphi \rangle = \langle \mathscr{L}^Q(S), \varphi \rangle = \langle S, \mathbf{1}_D \mathscr{L}^{\rho^* Q}(\varphi) \rangle$$

where  $\rho : \mathbb{C}^n \times \mathbb{C}^n \to \mathbb{C}^n \times \mathbb{C}^n$  is the involution defined by  $\rho(x, y) = (y, x)$ . So, for a general current *S*, we define the semi-regular transform as below:

**Definition 4.3.** Let  $n - q \le s \le n$  be an integer. Let *S* be a positive current of bidegree (s, s) with bounded mass on *D*. The semi-regular transform  $\mathscr{L}^Q(S)$  of *S* is a current of bidegree (s + q - n, s + q - n) defined by

$$\left\langle \mathscr{L}^{Q}(S), \varphi \right\rangle \coloneqq \left\langle S, \mathbf{1}_{D} \mathscr{L}^{\rho^{*}Q}(\varphi) \right\rangle$$

for a smooth test form  $\varphi$  of bidegree (2n - s - q, 2n - s - q), where  $\rho : \mathbb{C}^n \times \mathbb{C}^n \to \mathbb{C}^n \times \mathbb{C}^n$  is the involution defined by  $\rho(x, y) = (y, x)$ . The bidegree of the transform is defined to be (q - n, q - n).

If Q is a smooth form, then the transform is said to be smooth. If Q is positive (or closed), then the transform is said to be positive (or closed, respectively).

Note that the pairing on the right hand side makes sense since  $\mathscr{L}^{\rho^*Q}(\varphi)$  has  $C^1$  coefficients on  $\mathbb{C}^n$  and S is a positive current with bounded mass. When Q is smooth and has compact support in  $D \times D$ , one can easily see that the current  $\mathscr{L}^Q(S)$  is well defined and consistent with Definition 4.2 and Definition 4.3. In this case, the current  $\mathscr{L}^Q(S)$  is actually a smooth form with compact support.

Now, we want to justify that Definition 4.3 is consistent with Definition 4.2 in general and also with intuition. The main problem in defining the transform  $\mathscr{L}^Q(S)$  to be  $(\pi_2)_*(\pi_1^*S \wedge Q)$  is as follows: since Q is not smooth, it is not clear whether  $\pi_1^*S \wedge Q$  is well defined or not for now. We present some formulations equivalent to our definition. As a result, we see that the expression  $(\pi_2)_*(\pi_1^*S \wedge Q)$  is well defined as a current and also, we express  $\mathscr{L}^Q(S)$  as an integral formula. In particular, after this section, one may use any of the equivalent formulations for the definition of the transform  $\mathscr{L}^Q(S)$ .

Observe that the semi-regular transform is linear with respect to the form Q. Any smooth form of bidegree (q,q) can be written as a linear combination of positive smooth form of bidegree (q,q). Due to the first condition in (4.1), we may assume that Q is of the form

$$Q(x,y) = f(x,y) \prod_{i=1}^{q} \sqrt{-1} \psi_i \wedge \overline{\psi_i}$$

where  $\psi_i$ 's are smooth (1,0)-forms on  $\mathbb{C}^n \times \mathbb{C}^n$ , and f(x, y) is an  $L^1$  function on  $D \times D$ , which is smooth outside  $\Delta$  and bounded above, and satisfies

$$f \ge [\log \operatorname{dist}(\cdot, \Delta)] \operatorname{dist}(\cdot, \Delta)^{2-2n}$$
 and  $|\nabla f| \le \operatorname{dist}(\cdot, \Delta)^{1-2n}$ ,

where  $\nabla f$  means the gradient of f.

Let  $(\chi_k)$  be a sequence of smooth functions  $\chi_k : D \times D \to [0,1]$  with compact support such that  $\chi_k \equiv 1$  on the set  $\{(x,y) \in D \times D : \operatorname{dist}(x,\partial D) \ge k^{-1} \text{ and } \operatorname{dist}(y,\partial D) \ge k^{-1}\}$ ,  $\chi_k \le \chi_{k+1}$  and  $\lim_{k\to\infty} \chi_k = \mathbf{1}_{D\times D}$ .

Let  $(Q_k)_{k \in \mathbb{N}}$  denote a sequence of smooth forms with compact support in  $D \times D$  defined by

$$Q_k = \chi_k \cdot (\chi \circ (f+k) - k) \prod_{i=1}^q \sqrt{-1} \psi_i \wedge \overline{\psi_i}$$

where the function  $\chi : \mathbb{R} \to \mathbb{R}_{\geq 0}$  is as in Introduction. Then, observe that  $\chi_k(\chi \circ (f+k) - k) \searrow f$  as  $k \to \infty$ . In this sense, the Lebesgue dominated convergence theorem implies that the sequence  $(Q_k)$  approximates the form Q in the sense of currents.

**Proposition 4.4.** For a smooth test form  $\varphi$  of bidegree (2n - s - q, 2n - s - q), we have

$$\left\langle \mathscr{L}^{Q_k}(S), \varphi \right\rangle \to \left\langle \mathscr{L}^Q(S), \varphi \right\rangle$$

in the sense of currents.

*Proof.* Since  $Q_k$  is smooth and has compact support in  $D \times D$ , we have

$$\left\langle \mathscr{L}^{Q_k}(S),\varphi\right\rangle = \int_{D\times D} \pi_1^*S \wedge \pi_2^*\varphi \wedge Q_k = \left\langle S, \mathscr{L}^{\rho^*Q_k}(\varphi)\right\rangle.$$

We claim that

$$\left(S, \mathscr{L}^{\rho^*Q_k}(\varphi) - \mathscr{L}^{\rho^*Q}(\varphi)\right) \to 0 \quad \text{as } k \to \infty.$$

For each  $x \in D$ , direct computations give us

$$\begin{aligned} \mathscr{L}^{\rho^*Q_k}(\varphi) - \mathscr{L}^{\rho^*Q}(\varphi) &= \int_{y \in D \setminus \{x\}} \varphi(y) \wedge [Q_k(y, x) - Q(y, x)] \\ &= \int_{y \in D \setminus \{x\}} [\chi_k(\chi \circ (f+k) - k) - f(x, y)] \varphi(y) \wedge \prod_{i=1}^q \sqrt{-1} \psi_i \wedge \overline{\psi_i} \\ &= \int_{y \in \text{supp } \varphi \setminus \{x\}} [\chi_k(\chi \circ (f+k) - k) - f(x, y)] \varphi(y) \wedge \prod_{i=1}^q \sqrt{-1} \psi_i \wedge \overline{\psi_i} \end{aligned}$$

Since  $y \in \operatorname{supp} \varphi$  and  $\operatorname{supp} \varphi$  is compact, for all sufficiently large  $k \in \mathbb{N}$ , the sets  $\{x \in D : \operatorname{dist}(x, \operatorname{supp} \varphi) < k^{-1/2n}\}$  and  $\{x \in D : \operatorname{dist}(x, \partial D) < 1/k\}$  are disjoint. If  $|x - y| \ge k^{-1/2n}$  and  $\operatorname{dist}(x, \partial D) \ge 1/k$ , then  $\chi_k \equiv 1$  and  $\chi \circ (f + k) - k = f$ . Hence,  $\chi_k(\chi \circ (f + k) - k) - f(x, y) = 0$ , which means that the form  $\mathscr{L}^{\rho^*Q_k}(\varphi) - \mathscr{L}^{\rho^*Q}(\varphi)$  does not vanish only when x satisfies either  $\operatorname{dist}(x, \operatorname{supp}, \varphi) < k^{-1/2n}$  or  $\operatorname{dist}(x, \partial D) < 1/k$ .

If x satisfies dist $(x, \operatorname{supp}, \varphi) < k^{-1/2n}$ , the integral is bounded by  $c_1 k^{-1/2n}$ , where  $c_1$  is a constant independent of x and k. If x satisfies dist $(x, \partial D) < 1/k$ , the coefficients of  $\mathscr{L}^{\rho^*Q_k}(\varphi) - \mathscr{L}^{\rho^*Q}(\varphi)$  is bounded by  $-c_2[\log \operatorname{dist}(\operatorname{supp}\varphi, \partial D)]\operatorname{dist}(\operatorname{supp}\varphi, \partial D)^{2-2n}$ , where  $c_2$  is a constant independent of x and k. So, we can estimate

$$\left| \left( S, \mathscr{L}^{\rho^* Q_k}(\varphi) - \mathscr{L}^{\rho^* Q}(\varphi) \right) \right| \le c \left( k^{-1/2n} \|S\| + \int_{\{x \in D: \operatorname{dist}(x, \partial D) < 1/k\}} S \wedge \omega_x^{n-s} \right),$$

where c > 0 is a constant independent of k. Obviously, the first term converges to 0 as  $k \to \infty$ . Since  $S \wedge \omega_x^{n-s}$  is a Radon measure, S is assumed to have a finite mass over D, and D is bounded, the second integral converges to 0 as  $k \to \infty$ .

Let  $\varphi$  be a smooth test form of bidegree (2n - s - q, 2n - s - q). Without loss of generality, we may assume that  $\varphi$  is positive. We first consider the expression  $\int_{D \times D} \pi_1^* S \wedge \pi_2^* \varphi \wedge Q$ . Due to the main result in [6],  $\pi_1^* S$  is a positive closed current of bidegree (s, s). Since  $\pi_2^* \varphi \wedge \prod_{i=1}^q \sqrt{-1}\psi_i \wedge \overline{\psi_i}$  is smooth, the measure  $\pi_1^* S \wedge \pi_2^* \varphi \wedge \prod_{i=1}^q \sqrt{-1}\psi_i \wedge \overline{\psi_i}$  is well defined. So, the expression can be understood as an integration

$$\int_{D\times D} \pi_1^* S \wedge \pi_2^* \varphi \wedge Q \coloneqq \int_{D\times D} f(x, y) \pi_1^* S \wedge \pi_2^* \varphi \wedge \prod_{i=1}^q \sqrt{-1} \psi_i \wedge \overline{\psi_i}.$$

**Proposition 4.5.** Let  $\varphi$  be a smooth test form of bidegree (2n - s - q, 2n - s - q). The integrals

$$\int_{D \times D} \pi_1^* S \wedge \pi_2^* \varphi \wedge Q \quad and \quad \int_{D \times D \setminus \Delta} \pi_1^* S \wedge \pi_2^* \varphi \wedge Q$$

are well defined and equal to  $\langle \mathscr{L}^Q(S), \varphi \rangle$ . In particular, we can write

$$\mathscr{L}^Q(S) = \int_{x \in D \setminus \{y\}} S(x) \wedge Q(x,y) = (\pi_2)_* (\pi_1^* S \wedge Q).$$

Proof. As in the proof of Proposition 4.4, we can write

$$\langle \mathscr{L}^{Q_k}(S), \varphi \rangle = \int_{D \times D} \pi_1^* S \wedge \pi_2^* \varphi \wedge Q_k.$$

Then, by the monotone convergence theorem in measure theory and Proposition 4.4, we have

$$\int_{D\times D} \pi_1^* S \wedge \pi_2^* \varphi \wedge Q = \lim_{k \to \infty} \int_{D\times D} \pi_1^* S \wedge \pi_2^* \varphi \wedge Q_k = \lim_{k \to \infty} \left\langle \mathscr{L}^{Q_k}(S), \varphi \right\rangle = \left\langle \mathscr{L}^Q(S), \varphi \right\rangle$$

Hence, the integral is well defined as a real value and equal to  $\langle \mathscr{L}^Q(S), \varphi \rangle$ .

On the other hand,  $\pi_1^* S \wedge \pi_2^* \varphi \wedge Q$  is well defined as a current on  $D \times D \setminus \Delta$  since Q is smooth outside  $\Delta$ . So, if we allow  $-\infty$  as a value, the integral  $\int_{D \times D \setminus \Delta} \pi_1^* S \wedge \pi_2^* \varphi \wedge Q$  is well defined. Moreover, since the current  $\pi_1^* S$  has no mass on  $\Delta$  for bidegree reason, we have

$$\int_{D \times D} \pi_1^* S \wedge \pi_2^* \varphi \wedge Q = \lim_{k \to \infty} \int_{D \times D} \pi_1^* S \wedge \pi_2^* \varphi \wedge Q_k$$
$$= \lim_{k \to \infty} \int_{D \times D \setminus \Delta} \pi_1^* S \wedge \pi_2^* \varphi \wedge Q_k = \int_{D \times D \setminus \Delta} \pi_1^* S \wedge \pi_2^* \varphi \wedge Q_k$$

The last equality comes from the monotone convergence theorem in measure theory applied in the space  $D \times D \setminus \Delta$ . Hence, we see the desired equalities.

$$\left\langle \mathscr{L}^Q(S),\varphi\right\rangle = \int_{D\times D} \pi_1^* S \wedge \pi_2^* \varphi \wedge Q = \int_{D\times D\smallsetminus \Delta} \pi_1^* S \wedge \pi_2^* \varphi \wedge Q,$$

which means that the last integral is also well defined.

By Fubini's theorem, we can write

$$\int_{D \times D \setminus \Delta} \pi_1^* S \wedge \pi_2^* \varphi \wedge Q = \int_{y \in D} \left[ \int_{x \in D \setminus \{y\}} S(x) \wedge Q(x, y) \right] \wedge \varphi(y).$$

Also, the finiteness of the integral  $\int_{D \times D} \pi_1^* S \wedge \pi_2^* \varphi \wedge Q$  means that the expression  $\pi_1^* S \wedge Q$  is a current of order 0 and of finite mass. Hence, the formal definition  $\mathscr{L}^Q(S) = (\pi_2)_*(\pi_1^* S \wedge Q)$  is actually well defined. Therefore, we obtain the expressions of  $\mathscr{L}^Q(S)$  as

$$\mathscr{L}^Q(S) = \int_{x \in D \setminus \{y\}} S(x) \wedge Q(x,y) = (\pi_2)_* (\pi_1^* S \wedge Q).$$

**Remark 4.6.** By Fubini's theorem, when S is positive,  $\mathscr{L}^Q(S)$  is a form with  $L^{1+1/(n-1)}$ -coefficients. For instance, see [8, Theorem 2.3.1].

The following is quite straightforward.

**Proposition 4.7.** When Q is smooth, then  $\mathscr{L}^Q(S)$  is smooth.

**Proposition 4.8.** Let S be a current in  $\mathscr{C}_s(\overline{D})$ . Let  $S_{\epsilon}$  be the current on D obtained from the regularization of S in a neighborhood of  $\overline{D}$  by the standard convolution method in  $\mathbb{C}^n$ . Then, the transform  $\mathscr{L}^Q(S_{\epsilon})$  converges to  $\mathscr{L}^Q(S)$  in D as  $\epsilon \to 0$  in the sense of currents.

*Proof.* Let  $g : \mathbb{C}^n \to \mathbb{R}_{\geq 0}$  be a smooth function with compact support in the unit ball such that  $\int_{\mathbb{C}^n} g \, d\mu = 1$  and that for  $z_1, z_2 \in \mathbb{C}^n$  with  $|z_1| = |z_2|$ , we have  $g(z_1) = g(z_2)$ , where  $\mu$  denotes the standard Lebesgue measure. For  $z \in \mathbb{C}^n$ , we define  $\tau_z(x) = x - z$ . Let  $\epsilon > 0$  be sufficiently small so that in the  $\epsilon$ -neighborhood of  $\overline{D}$ , the current S is well defined. The standard convolution formula gives us

$$S_{\epsilon}(x) = \int_{z \in \mathbb{C}^n} \tau_z^* S(x) \left( \frac{1}{\epsilon^n} g\left( \frac{z}{\epsilon} \right) \right) d\mu(z).$$

Let  $\varphi$  be a smooth test form of bidegree (2n - s - q, 2n - s - q) in D. Then, we have

$$\left\langle \mathscr{L}^{Q}(S_{\epsilon}) - \mathscr{L}^{Q}(S), \varphi \right\rangle = \left\langle S_{\epsilon} - S, \mathbf{1}_{D} \mathscr{L}^{\rho^{*}Q}(\varphi) \right\rangle = \int_{z \in \mathbb{C}^{n}} \left\langle \tau_{z}^{*} S - S, \mathbf{1}_{D} \mathscr{L}^{\rho^{*}Q}(\varphi) \right\rangle \left( \frac{1}{\epsilon^{n}} g\left( \frac{z}{\epsilon} \right) \right) d\mu(z) = \int_{z \in \mathbb{C}^{n}} \left\langle S, \tau_{z}^{*}(\mathbf{1}_{D} \mathscr{L}^{\rho^{*}Q}(\varphi)) - \mathscr{L}^{\rho^{*}Q}(\varphi) \right\rangle \left( \frac{1}{\epsilon^{n}} g\left( \frac{z}{\epsilon} \right) \right) d\mu(z) = \int_{z \in \mathbb{C}^{n}} \left\langle S, \mathbf{1}_{D} \tau_{z}^{*}(\mathscr{L}^{\rho^{*}Q}(\varphi)) - \mathscr{L}^{\rho^{*}Q}(\varphi) \right\rangle \left( \frac{1}{\epsilon^{n}} g\left( \frac{z}{\epsilon} \right) \right) d\mu(z)$$

$$(4.2)$$

(4.3) 
$$+ \int_{z \in \mathbb{C}^n} \left\langle S, (\tau_z^*(\mathbf{1}_D) - \mathbf{1}_D) \tau_z^*(\mathscr{L}^{\rho^*Q}(\varphi)) \right\rangle \left(\frac{1}{\epsilon^n} g\left(\frac{z}{\epsilon}\right)\right) d\mu(z)$$

Since  $\mathscr{L}^{\rho^*Q}(\varphi)$  is defined on  $\mathbb{C}^n$  and has  $C^1$ -coefficients. Recall that D is bounded and so the  $\epsilon$ -neighborhood of  $\overline{D}$  has compact closure  $\overline{(D)_{\epsilon}}$ . So, the uniform continuity theorem applied on the set  $(\overline{D})_{\epsilon}$  implies that

$$\left| \tau_z^* (\mathscr{L}^{\rho^* Q}(\varphi))(x) - \mathscr{L}^{\rho^* Q}(\varphi)(x) \right| \lesssim \epsilon \quad \text{ for all } x \in D \text{ and for } z \text{ with } |z| < \epsilon$$

where  $|\cdot|$  means the sum of the absolute values of the coefficients of the given form and  $\leq$  means  $\leq$  up to a multiplicative constant independent of x, z and  $\epsilon$ . Hence, we have  $|(4.2)| \leq \epsilon$ , where  $\leq$  means  $\leq$  up to a multiplicative constant independent of x, z and  $\epsilon$ . Since D is bounded and D has smooth boundary, from the well known tube formula, |(4.3)| is also bounded by  $\epsilon$  up to a multiplicative constant. Thus, by the triangle inequality, we can draw the desired conclusion.

4.2. Regularizations of positive closed currents. Applying Theorem 1.1 with  $U = \mathbb{C}^n \times \mathbb{C}^n$  and  $f: \mathbb{C}^n \times \mathbb{C}^n \to \mathbb{C}^n$  defined by  $f(x_1, \dots, x_n, y_1, \dots, y_n) = (x_1 - y_1, \dots, x_n - y_n)$ , we have

$$(4.4) \qquad \qquad (dd^c \log |x-y|)^n = [\Delta].$$

We set  $u = \log |x - y|$  and we have  $(dd^c u)^n = [\Delta]$  in the sense of currents.

We consider two approximations of u introduced in Section 2 for this setup:

$$u_{\theta}^{\mathcal{K}} \coloneqq \chi(u - \log |\theta|) + \log |\theta| \quad \text{and} \quad u_{\theta}^{\mathcal{T}} \coloneqq \frac{1}{2} \log \left(|x - y|^2 + |\theta|^2\right).$$

Also, the following forms are defined there:

- (1)  $\mathcal{K}^{i} \coloneqq (dd^{c}u)^{i}$  for  $i = 1, \dots, n-1$ ; (2)  $\mathcal{K}^{i}_{\theta} \coloneqq (dd^{c}u^{\mathcal{K}}_{\theta}) \land (dd^{c}u)^{i-1}$  for  $i = 1, \dots, n$ ; (3)  $\mathcal{T}^{i}_{\theta} \coloneqq (dd^{c}u^{\mathcal{T}}_{\theta})^{i}$  for  $i = 1, \dots, n$ .

Observe that the forms from (1) through (6) are all invariant under the involution  $\rho(x, y) = (y, x)$ . Also, notice that all the kernels define semi-regular transforms of currents due to their regularity properties. We also write  $\mathcal{K}^n := (dd^c u)^n$ , which is equal to the current of integration  $[\Delta]$  on the diagonal submanifold  $\Delta$  of  $\mathbb{C}^n \times \mathbb{C}^n$  due to King's formula.

**Proposition 4.9.** Let  $1 \leq s \leq n$  be an integer. Let  $S \in \mathscr{C}_s(D)$ . Then, the current  $\mathscr{L}^{\mathcal{K}^n_{\theta}}(S)$  is positive and the family  $(\mathscr{L}^{\mathcal{K}^n_{\theta}}(S))$  converge to S in the sense of currents as  $|\theta| > 0$ . Let D' be an open subset of D with compact closure. Then, for all  $\theta \in \mathbb{C}^*$  with  $|\theta| \ll 1$ , the restriction of  $\mathscr{L}^{\mathcal{K}^n_{\theta}}(S)$  is a positive closed current in D'.

*Proof.* We first consider a smooth current  $S \in \mathscr{C}_s(D)$ . Let D' be a open subset of D with compact closure. It suffices to prove that the restriction of  $\mathscr{L}^{\mathcal{K}^n_{\theta}}(S)$  to D' is a smooth positive closed current converging to S in D' as  $|\theta| \to 0$ . We assume that  $|\theta|$  be sufficiently small that  $\operatorname{supp} \mathcal{K}_{\theta}^n \subset$  $\{(x,y) \in D \times D : e^{-1}|\theta| \le \operatorname{dist}(x,y) \le e|\theta|\} \subset \overline{\Delta_{e|\theta|}}$ . Let  $\varphi$  be a smooth test form of bidegree (n-s, n-s) such that  $\operatorname{supp} \varphi \in D'$ . Let  $\chi_{\varphi} : D \to [0, 1]$  be a smooth function with compact support such that  $\operatorname{supp}\chi_{\varphi} \in D'$  and that the  $\{\chi_{\varphi} \equiv 1\}$  contains the set  $\operatorname{supp}\varphi$ . We have

$$\left\langle \mathscr{L}^{\mathcal{K}^{n}_{\theta}}(S),\varphi\right\rangle \coloneqq \left\langle S,\mathscr{L}^{\mathcal{K}^{n}_{\theta}}(\varphi)\right\rangle = \int_{D\times D} \pi_{1}^{*}S \wedge \pi_{2}^{*}(\varphi) \wedge \mathcal{K}^{n}_{\theta} = \int_{D\times D} \pi_{1}^{*}(\chi_{\varphi}S) \wedge \pi_{2}^{*}(\varphi) \wedge \mathcal{K}^{n}_{\theta}$$
$$= \int_{D\times D} \pi_{1}^{*}S \wedge [\pi_{1}^{*}(\chi_{\varphi})\pi_{2}^{*}(\varphi)] \wedge (dd^{c}u_{\theta}^{\mathcal{K}}) \wedge (dd^{c}u)^{n-1}.$$

Since  $\pi_1^*(\chi_{\varphi})\pi_2^*(\varphi)$  is a smooth form with compact support in  $D \times D$ . The above becomes

$$= \int_{D \times D} u_{\theta}^{\mathcal{K}} \pi_1^* S \wedge dd^c [\pi_1^*(\chi_{\varphi}) \pi_2^*(\varphi)] \wedge (dd^c u)^{n-1}$$

In the same way, we have

$$S,\varphi\rangle = \int_{D\times D} u\pi_1^* S \wedge dd^c [\pi_1^*(\chi_\varphi)\pi_2^*(\varphi)] \wedge (dd^c u)^{n-1}$$

and therefore, we have

$$\begin{split} \left| \left\langle \mathscr{L}^{\mathcal{K}^{n}_{\theta}}(S), \varphi \right\rangle - \left\langle S, \varphi \right\rangle \right| &= \left| \int_{D \times D} (u_{\theta}^{\mathcal{K}} - u) \pi_{1}^{*} S \wedge dd^{c} [\pi_{1}^{*}(\chi_{\varphi}) \pi_{2}^{*}(\varphi)] \wedge (dd^{c}u)^{n-1} \right| \\ &\leq \|\pi_{1}^{*}(\chi_{\varphi}) \pi_{2}^{*}(\varphi)\|_{C^{2}} \int_{D \times D} (u_{\theta}^{\mathcal{K}} - u) \pi_{1}^{*} S \wedge \pi_{1}^{*}(\chi_{D'}) \pi_{2}^{*}(\chi_{D'}) \omega_{x,y}^{n-s} \wedge (dd^{c}u)^{n-1}, \end{split}$$

where  $\chi_{D'}: D \to [0,1]$  is a smooth function with compact support such that  $\chi_{D'} \equiv 1$  on  $\overline{D'}$ . Since the singularity of  $u \wedge (dd^c u)^{n-1}$  is of the form  $\operatorname{dist}(\cdot, \Delta)^{2-2n} \log \operatorname{dist}(\cdot, \Delta)$  and the support of  $(u_{\theta}^{\mathcal{K}} - u)$  is a tubular neighborhood of  $\Delta$  whose radius is proportional to  $|\theta| > 0$ , the form  $(\pi_1)_*[(u_{\theta}^{\mathcal{K}} - u)\pi_1^*(\chi_{D'})\pi_2^*(\chi_{D'})\omega_{x,y}^{n-s} \wedge (dd^c u)^{n-1}]$  is a  $C^1$ -form whose  $C^1$ -norm is bounded by  $|\theta|$  up to a multiplicative constant independent of  $\theta$ . The support of the form the form  $(\pi_1)_*[(u_{\theta}^{\mathcal{K}} - u)\chi_{D'\times D'}\omega^{n-s} \wedge (dd^c u)^{n-1}]$  sits inside  $\operatorname{supp}\chi_{D'}$ . Hence, we have

$$\left|\left\langle \mathscr{L}^{\mathcal{K}^{n}_{\theta}}(S),\varphi\right\rangle-\left\langle S,\varphi\right\rangle\right|\lesssim\theta\left\|\pi_{1}^{*}(\chi_{\varphi})\pi_{2}^{*}(\varphi)\right\|_{C^{2}}\|S\|_{\mathrm{supp}\chi_{D'}}.$$

For a general  $S \in \mathscr{C}_s(D)$ , we use the regularization  $S_{\epsilon}$  of S obtained from the standard convolution method in  $\mathbb{C}^n$ . We have  $\|S_{\epsilon}\|_{\operatorname{supp}\chi_{D'}} \leq c\|S\|_{\operatorname{supp}\chi_{D'}}$  for some constant c > 0 independent of  $\epsilon$ . So, the convergence of  $\mathscr{L}^{\mathcal{K}^n_{\theta}}(S_{\epsilon}) \to S_{\epsilon}$  in D' is uniform with respect to  $\epsilon$  and therefore, Proposition 4.8 implies that the convergence in D' is true for  $S \in \mathscr{C}_s(D)$ .

We prove the closedness of  $\mathscr{L}^{\mathcal{K}^n_{\theta}}(S)$ . Assume that  $\varphi = d\psi$  for some smooth form  $\psi$  with compact support in D'. By Proposition 4.5, we have

$$\left\langle \mathscr{L}^{\mathcal{K}^n_{\theta}}(S), d\psi \right\rangle = \int_{D \times D} \pi_1^* S \wedge \pi_2^*(d\psi) \wedge \mathcal{K}^n_{\theta} = \int_{D \times D} \pi_1^* S \wedge d\left(\pi_2^*(\psi) \wedge \mathcal{K}^n_{\theta}\right).$$

Notice that from our choice of  $\theta$ , the support of  $\pi_2^*(\psi) \wedge \mathcal{K}_{\theta}^n$  is compact in  $D \times D$ . So, the Stokes theorem conclude that  $\mathscr{L}^{\mathcal{K}_{\theta}^n}(S)$  is closed when restricted to D'.

**Remark 4.10.** If S is just a current in  $\mathscr{C}_s(D)$ , it is not clear whether  $\mathscr{L}^{\mathcal{K}^n_{\theta}}(S)$  is closed at all. However, Proposition 4.9 implies that if S is a current in  $\mathscr{C}_s(\overline{D})$ , then for all  $\theta \in \mathbb{C}^*$  with sufficiently small  $|\theta| > 0$ ,  $\mathscr{L}^{\mathcal{K}^n_{\theta}}(S)$  is a smooth positive closed current in  $\mathscr{C}_s(D)$ .

Below are the estimates of the  $C^{\alpha}$ -norms of  $\mathscr{L}^{\mathcal{K}^{n}_{\theta}}(S)$ .

**Proposition 4.11.** Let  $\theta$  be such that  $0 < |\theta| \ll 1$ . Let  $1 \le s \le n$  be an integer. Let  $S \in C_s(D)$ . Then, for each compact subset  $K \in D$ , we have

$$\|\mathscr{L}^{\mathcal{K}^n_{\theta}}(S)\|_{C^{\alpha},K} \le c_{\alpha,K} |\theta|^{-2n-c}$$

for some constant  $c_{\alpha,K} > 0$  independent of  $\theta$ .

*Proof.* Notice that from the choice of the function  $\chi$  in Introduction,  $\operatorname{supp} \mathcal{K}_{\theta}^{n} \subset \{(x,y) \in D \times D : e^{-1}|\theta| \leq \operatorname{dist}(x,y) \leq e|\theta|\} =: W_{|\theta|}$ . Over the region  $W_{|\theta|}$ ,  $dd^{c}u$  is smooth and we have  $\|(dd^{c}u)^{n-1}\|_{C^{\alpha},W_{\theta}\cap\pi_{2}^{-1}(K)} \leq c'_{\alpha,K}|\theta|^{2-2n-\alpha}$  and also  $\|dd^{c}u_{\theta}^{\mathcal{K}}\|_{C^{\alpha},W_{\theta}\cap\pi_{2}^{-1}(K)} \leq c'_{\alpha,K}|\theta|^{-2}$  for some constant  $c'_{\alpha,K} > 0$  independent of  $\theta$ . Hence, we can conclude that

$$\|\mathscr{L}^{\mathcal{K}^n_{\theta}}(S)\|_{C^{\alpha},K} \le c_{\alpha,K}|\theta|^{-2n-\epsilon}$$

for some constant  $c_{\alpha,K} > 0$  independent of  $\theta$ .

We consider another regularization of currents:  $(\mathscr{L}^{\mathcal{T}^n_{\theta}}(S))_{0 < |\theta| \ll 1}$ .

**Proposition 4.12.** Let  $S \in \mathscr{C}_s(\overline{D})$ . Then, the sequence  $\mathscr{L}^{\mathcal{T}^n_{\theta}}(S)$  converges to S in the sense of currents as  $|\theta| \leq 0$ .

*Proof.* Let  $\varphi$  be a smooth test form of bidegree (n - s, n - s) in D. We define  $\chi_{\varphi} : D \times D \to [0, 1]$  to be a smooth function such that  $\chi_{\varphi} \equiv 1$  in a neighborhood of  $\Delta$  and  $\operatorname{supp} \chi_{\varphi} \cap \pi_2^{-1}(\operatorname{supp} \varphi)$  is compact.

For now, we assume that S is smooth. Then, due to Theorem 1.1, we have

$$\left\langle \mathscr{L}^{\mathcal{T}^{n}_{\theta}}(S),\varphi\right\rangle - \left\langle S,\varphi\right\rangle = \left\langle (\pi_{2})_{*}(\pi_{1}^{*}S \wedge \mathcal{T}^{n}_{\theta}) - (\pi_{2})_{*}(\pi_{1}^{*}S \wedge \mathcal{K}^{n}),\varphi\right\rangle$$

$$= \left\langle (\pi_{2})_{*}(\pi_{1}^{*}S \wedge \mathcal{T}^{n}_{\theta}) - (\pi_{2})_{*}(\pi_{1}^{*}S \wedge \mathcal{K}^{n}),\varphi\right\rangle = \int_{D \times D} S(x) \wedge \varphi(y) \wedge (\mathcal{T}^{n}_{\theta} - \mathcal{K}^{n})(x,y)$$

$$= \int_{D \times D} \chi_{\varphi}(x,y)S(x) \wedge \varphi(y) \wedge (\mathcal{T}^{n}_{\theta} - \mathcal{K}^{n})(x,y)$$

$$+ \int_{D \times D} (1 - \chi_{\varphi}(x,y))S(x) \wedge \varphi(y) \wedge (\mathcal{T}^{n}_{\theta} - \mathcal{K}^{n})(x,y)$$

$$5$$

(4.5)

$$= \frac{1}{2^{n}} \sum_{i=0}^{n-1} \int_{\mathbb{C}^{n} \times \mathbb{C}^{n}} \chi_{\varphi}(x, y) S(x) \wedge \varphi(y) \wedge dd^{c} [\log(|x - y|^{2} + |\theta|^{2}) - \log|x - y|^{2}] \\ \wedge (dd^{c} \log(|x - y|^{2} + |\theta|^{2}))^{i} \wedge (dd^{c} \log|x - y|^{2})^{k-i-1}$$

(4.6)

$$= \frac{1}{2^n} \sum_{i=0}^{n-1} \int_{D \times D} (1 - \chi_{\varphi}(x, y)) S(x) \wedge \varphi(y) \wedge dd^c [\log(|x - y|^2 + |\theta|^2) - \log|x - y|^2] \\ \wedge (dd^c \log(|x - y|^2 + |\theta|^2))^i \wedge (dd^c \log|x - y|^2)^{k-i-1}.$$

Since  $\operatorname{supp}\chi_{\varphi} \cap \pi_2^{-1}(\operatorname{supp}\varphi)$  is compact, (4.5) can be written as

$$(4.5) = \frac{1}{2^n} \sum_{i=0}^{n-1} \int_{\mathbb{C}^n \times \mathbb{C}^n} \left[ \log(|x-y|^2 + |\theta|^2) - \log|x-y|^2 \right] S(x) \wedge (dd^c \log(|x-y|^2 + |\theta|^2))^i \\ \wedge (dd^c \log|x-y|^2)^{k-i-1} \wedge dd^c [\chi_{\varphi}(x,y)\varphi(y)]$$

In the last inequality, we use the fact that the current  $(dd^c \log ||x-y||^2)^{k-i-1}$  has no mass on the diagonal submanifold  $\Delta$  for  $i = 0, \dots, k-1$ . We use the change of coordinates  $(x, y) \rightarrow (x, x-y)$  for  $\mathbb{C}^n \times \mathbb{C}^n$  and denote by w = x - y. Then, the above sum becomes

$$(4.5) = \frac{1}{2^n} \sum_{i=0}^{n-1} \int_{\mathbb{C}^n \times \mathbb{C}^n} [\log(|w|^2 + |\theta|^2) - \log|w|^2] S(x) \wedge (dd^c \log(|w|^2 + |\theta|^2))^i \\ \wedge (dd^c \log|w|^2)^{k-i-1} \wedge dd^c [\chi_{\varphi}(x, x - w)\varphi(x - w)]$$

and we get

$$\begin{aligned} |(4.5)| \lesssim \|\chi_{\varphi}\|_{C^{2}} \|\varphi\|_{C^{2}} \|S\| \sum_{i=0}^{n-1} \int_{|w| < R} [\log(|w|^{2} + |\theta|^{2}) - \log|w|^{2}] (dd^{c} \log(|w|^{2} + |\theta|^{2}))^{i} \\ & \wedge (dd^{c} \log|w|^{2})^{k-i-1} \wedge \omega(w), \end{aligned}$$

where R > 0 is a sufficiently large number so that the image of  $\operatorname{supp}\chi_{\varphi} \cap \pi_2^{-1}(\operatorname{supp}\varphi)$  under the projection  $(x, w) \to w$  is contained in  $\{w \in \mathbb{C}^n : |w| < R\}$ . Hereafter in this proof, the inequality  $\leq x \leq w$  is  $\leq w$  to a multiplicative constant independent of S and  $\theta$ .

We have

$$dd^{c} \log(|w|^{2} + |\theta|^{2}) = \sqrt{-1}\partial\bar{\partial} \log(|w|^{2} + |\theta|^{2}) = \sqrt{-1}\partial\left(\frac{\sum_{i=1}^{n} w_{i}d\bar{w}_{i}}{|w|^{2} + |\theta|^{2}}\right)$$
$$= \frac{\sum_{i=1}^{n} \sqrt{-1}dw_{i} \wedge d\bar{w}_{i}}{|w|^{2} + |\theta|^{2}} - \frac{\sqrt{-1}(\sum_{i=1}^{n} \bar{w}_{i}dw_{i}) \wedge (\sum_{i=1}^{n} w_{i}d\bar{w}_{i})}{(|w|^{2} + |\theta|^{2})^{2}}.$$

So, for all  $i = 0, \ldots, n-1$ , the integral

$$\int_{|w| < R} \left[ \log(|w|^2 + |\theta|^2) - \log|w|^2 \right] (dd^c \log(|w|^2 + |\theta|^2))^i \wedge (dd^c \log|w|^2)^{k-i-1} \wedge \omega(w)$$

is bounded by a constant times

$$\int_{|w| < R} \frac{\log(|w|^2 + |\theta|^2) - \log|w|^2}{|w|^{2k-2}} d\mu(w),$$

where  $\mu$  denotes the standard Lebesgue measure on  $\mathbb{C}^k$ .

If we use the polar coordinate system, we can easily get

$$\int_{|w| \le t} \frac{1}{|w|^{2k-2}} d\mu(w) \le c_1 t^2 \quad \text{and} \quad \int_{|w| \le t} \frac{1}{|w|^{2k-1}} d\mu(w) \le c_2 t,$$

where 0 < t < 1 and the constants  $c_1, c_2 > 0$  are independent of t.

So, we have

$$\begin{split} \left| \int_{|w| < R} \frac{\log(|w|^2 + |\theta|^2) - \log|w|^2}{|w|^{2k-2}} d\mu(w) \right| \\ \leq \left| \int_{|\theta|^{1/2} \le |w| < R} \frac{\log(|w|^2 + |\theta|^2) - \log|w|^2}{|w|^{2k-2}} d\mu(w) \right| \\ + \left| \int_{|w| < |\theta|^{1/2}} \frac{\log(|w|^2 + |\theta|^2) - \log|w|^2}{|w|^{2k-2}} d\mu(w) \right| \\ \leq \left| \int_{|\theta|^{1/2} \le |w| < R} \frac{\log(|w|^2 + |\theta|^2) - \log|w|^2}{|w|^{2k-2}} d\mu(w) \right| \\ + \left| \int_{|w| < |\theta|^{1/2}} \frac{\log(|w|^2 + |\theta|^2)}{|w|^{2k-2}} d\mu(w) \right| + \left| \int_{0 < |w| < |\theta|^{1/2}} \frac{\log|w|^2}{|w|^{2k-2}} d\mu(w) \right| \\ \leq C_1 |\theta| + C_2 |\theta| \log |\theta| + C_3 |\theta|^{1/2} \end{split}$$

where  $C_1, C_2, C_3 > 0$  are constants independent of  $\theta$  and S. Hence, we have  $|(4.5)| \leq |\theta|^{1/2}$ , where the inequality  $\leq \text{mean} \leq \text{up to a multiplicative constant independent of } \theta$  and S.

Concerning (4.6), we directly compute the form

$$dd^{c} \log(|w|^{2} + |\theta|^{2}) - dd^{c} \log|w|^{2}$$
  
=  $|\theta|^{2} \frac{\sqrt{-1}(2|w|^{2} + |\theta|^{2})(\sum_{i=1}^{n} \bar{w}_{i}dw_{i}) \wedge (\sum_{i=1}^{n} w_{i}d\bar{w}_{i})}{|w|^{4}(|w|^{2} + |\theta|^{2})^{2}} - |\theta|^{2} \frac{\sum_{i=1}^{n} \sqrt{-1}dw_{i} \wedge d\bar{w}_{i}}{|w|^{2}(|w|^{2} + |\theta|^{2})}$ 

as previously. Due to the choice of our function  $\chi_{\varphi}$ , there exists an r > 0 such that  $r < \text{dist}(\text{supp}(1-\chi_{\text{supp}\varphi}), \Delta)$ . We see that

$$\begin{aligned} |(4.6)| &= \left| \frac{1}{2^n} \sum_{i=0}^{n-1} \int_{D \times D \cap \{|w| > r\}} (1 - \chi_{\varphi}(x, y)) S(x) \wedge \varphi(y) \wedge dd^c [\log(|x - y|^2 + |\theta|^2) - \log|x - y|^2] \right| \\ &\wedge (dd^c \log(|x - y|^2 + |\theta|^2))^i \wedge (dd^c \log|x - y|^2)^{k-i-1} \right| \lesssim |\theta|^2 \|\varphi\|_{\infty} \|S\|. \end{aligned}$$

Hence, for a given smooth test form  $\varphi$  of bidegree (n - s, n - s), when S is smooth, we see that

$$\left|\left\langle \mathscr{L}^{\mathcal{T}^n_{\theta}}(S),\varphi\right\rangle - \left\langle S,\varphi\right\rangle\right| \lesssim \|S\| \|\theta\|^{1/2},$$

where the inequality  $\leq$  mean  $\leq$  up to a multiplicative constant independent of  $\theta$  and *S*.

For a general  $S \in \mathscr{C}_s(\overline{D})$ , we use the regularization of  $\mathbf{1}_D S$  obtained from the standard convolution method in  $\mathbb{C}^n$ . The mass of  $||S_{\epsilon}|| \leq c ||S||$  for some constant c > 0 independent of  $\epsilon$ . So, the

convergence of  $\mathscr{L}_{\theta}^{\mathcal{T}_{\theta}^{n}}(S_{\epsilon}) \to S_{\epsilon}$  is uniform with respect to  $\epsilon$  and therefore, Proposition 4.8 implies that the convergence is true for  $S \in \mathscr{C}_{s}(\overline{D})$ .

**Remark 4.13.** The approximating currents  $(\mathscr{L}^{\mathcal{T}_{\theta}^{n}}(S))_{0 < |\theta| \ll 1}$  may not be closed on any open subset with compact closure in general as the support of  $\mathcal{T}_{\theta}^{n}$  does not shrink to  $\Delta$ .

# 5. INTERSECTION OF POSITIVE CLOSED CURRENTS

In this section, we apply the existence of the unique tangent current with minimal h-dimension to study the intersection of positive closed currents on domains. Throughout this section, we continue to assume D to be a bounded simply connected domain with smooth boundary in  $\mathbb{C}^n$ . We continue to use the same notations as in Section 4.

5.1. Intersection of two positive closed currents. Let *s* and *r* be two positive integers such that  $1 < s + r \le n$ . Let *S* and *R* be currents in  $\mathscr{C}_s(D)$  and  $\mathscr{C}_r(D)$ , respectively.

We take  $U = D \times D$ , N = 2n. We take as V the diagonal submanifold  $\Delta$  of  $D \times D$ , which is defined by  $\Delta = \{(x, y) \in D \times D : x = y\}$ . We consider  $D \times D$  as a subset of  $\mathbb{C}^n \times \Delta$ . We take  $E = \mathbb{C}^n \times \Delta$ ,  $\overline{E} = \mathbb{P}^n \times \Delta$  and  $\pi_\Delta : \mathbb{P}^n \times \Delta \to \Delta$  to be the natural extension of the projection of  $D \times D$  onto  $\Delta$  and  $\pi_F : \overline{E} \to \mathbb{P}^n$  the complementary projection of  $\overline{E}$  onto the fiber space  $\mathbb{P}^n$ . We will use coordinates  $([w:t], y) \in \overline{E}$  and identify (w, y) with  $([w:1], y) \in E$ . Then, we have  $\pi_\Delta([w:t], y) = y$  and  $\pi_F([w:t], y) = [w:t]$ . The inclusion map  $\iota : D \times D \to E$  can be written as  $\iota(x, y) = (x - y, y)$ . We will study the case of  $T = \pi_1^* S \wedge \pi_2^* R$ . We will use  $\omega_\Delta = dd^c |y|^2$  on  $\Delta$ ,  $\omega_F = \frac{1}{2}dd^c \log(|w|^2 + |t|^2)$  on  $\mathbb{P}^n$  and  $\omega = \pi_\Delta^* \omega_\Delta + \pi_F^* \omega_F$  on  $\overline{E}$ . In these coordinates, we may say  $\pi_2^* \omega_D = \pi_\Delta^* \omega_\Delta$  on  $D \times D$ . We will also continue to use  $\omega_{x,y} = \pi_1^* \omega_D + \pi_2^* \omega_D$ . We take  $u = \log |x - y|$ .

We start by an observation below, which is actually the main motivation of this work.

Let  $\varphi$  be a smooth test form of bidegree (n - s - r, n - s - r) with compact support in D. From Theorem 1.1, we have  $[\Delta] = (dd^c u)^n$  in the sense of currents. Let  $\chi_{\varphi} : D \to [0, 1]$  denote a smooth function with compact support such that  $\chi_{\varphi}$  is smooth,  $\chi_{\varphi} \equiv 1$  on the support of  $\varphi$  and  $\chi_{\varphi}$  has compact support in U. Then, if S and R are smooth, we can write

$$\langle S \wedge R, \varphi \rangle = \langle S, R \wedge \varphi \rangle = \langle \chi_{\varphi}S, R \wedge \varphi \rangle = \int_{D \times D} \pi_1^*(\chi_{\varphi}S) \wedge \pi_2^*(R \wedge \varphi) \wedge [\Delta]$$
  
= 
$$\int_{D \times D} \pi_1^*(\chi_{\varphi}S) \wedge \pi_2^*(R \wedge \varphi) \wedge (dd^c u)^n$$
  
= 
$$\int_{D \times D} \pi_1^*S \wedge \pi_2^*R \wedge \pi_1^*(\chi_{\varphi})\pi_2^*(\varphi) \wedge (dd^c u)^n.$$

Since  $\pi_1^*(\chi_{\varphi})\pi_2^*(\psi)$  has compact support, by Stoke's theorem, we can write the above integral as follows:

$$= \int_{D \times D} \pi_1^* S \wedge \pi_2^* R \wedge \left( u (dd^c u)^{n-1} \right) \wedge dd^c (\pi_1^*(\chi_\varphi) \pi_2^*(\varphi)).$$

Observe that the form  $dd^c(\pi_1^*(\chi_{\varphi})\pi_2^*(\varphi))$  is  $dd^c\pi_2^*(\varphi)$  in a neighborhood of  $\Delta$ . Recall that we previously had this type of representation in Theorem 1.3. Hence, we will extend this description of the wedge product to general positive closed currents.

Now, we adapt a sufficient condition for the existence of the unique tangent current with the minimal *h*-dimension to introduce a sufficient condition for the definition of the wedge product of  $S \in \mathscr{C}_s(D)$  and  $R \in \mathscr{C}_r(D)$ . Note that the current  $\pi_1^*S \wedge \pi_2^*R$  is well defined as  $\pi_1^*S \otimes \pi_2^*R$ . For a rigorous treatment of this, see the double current in [3].

**Definition 5.1.** We say that the currents S and R are wedgable if  $\pi_1^* S \wedge \pi_2^* R$  satisfies the horizontal condition as in Definition 3.5.

Remark 5.2. In our coordinates, the horizontal condition can be written as

$$u \in L^1_{loc}\left(\left\langle \pi_1^* S \land \pi_2^* \left( R \land \omega_D^{2n-s-r-i+1} \right) \land \mathcal{K}^{i-1} \right\rangle_{\mathcal{K}} \right)$$

inductively from i = 1 through n. By symmetry, the wedgability condition can also be written as

$$u \in L^{1}_{loc}\left(\left\langle \pi_{1}^{*}\left(S \wedge \omega_{D}^{2n-s-r-i+1}\right) \wedge \pi_{2}^{*}R \wedge \mathcal{K}^{i-1}\right\rangle_{\mathcal{K}}\right)$$

inductively from i = 1 through n.

Proposition 3.10 can be interpreted as follows:

**Proposition 5.3.** Let  $S \in \mathscr{C}_s(D)$  and  $R \in \mathscr{C}_r(D)$  be two wedgable currents. Let  $\varphi$  be a smooth test form with compact support of bidegree (n - s - r, n - s - r) on D. Let  $\Phi$  be a smooth test form with compact support of bidegree (n - s - r, n - s - r) on  $D \times D$  such that  $\Phi = \pi_2^*(\varphi)$  in a neighborhood of  $\Delta$ . Then, the following integral is well-defined and independent of the choice of  $\Phi$ .

(5.1) 
$$\langle (S \wedge R)_K, \varphi \rangle \coloneqq \int_{D \times D \setminus \Delta} \pi_1^* S \wedge \pi_2^* R \wedge (u(dd^c u)^{n-1}) \wedge dd^c \Phi,$$

which defines a positive closed current of bidegree (s + r, s + r) on D.

**Definition 5.4.** Let  $S \in \mathscr{C}_s(D)$  and  $R \in \mathscr{C}_r(D)$  be two currents such that the currents S and R are wedgable. The wedge product of S and R in the sense of King's formula is defined to be as in (5.1) in Proposition 5.3 and denoted by  $(S \land R)_K$ .

Theorem 1.3 gives us the following theorem on the intersection of positive closed currents, which shows that Definition 5.4 is actually intrinsic.

**Theorem 1.4.** Let  $S \in \mathscr{C}_s(D)$  and  $R \in \mathscr{C}_r(D)$  be two currents such that the currents S and R are wedgable. Then, the current  $\pi_1^*S \wedge \pi_2^*R$  has a unique tangent current along  $\Delta$  and its h-dimension is minimal. Moreover, its shadow is exactly  $(S \wedge R)_K$ . Namely, the Dinh-Sibony product of S and R in [9] is well defined and equal to  $(S \wedge R)_K$ .

The following propositions are quite straightforward.

**Proposition 5.5.** If S is a current in  $\mathscr{C}_s(D)$  and R is a smooth current in  $\in \mathscr{C}_r(D)$ , then S and R are wedgable and we have  $(S \land R)_K = S \land R$ .

**Proposition 5.6.** Suppose that two currents  $S \in \mathscr{C}_s(D)$  and  $R \in \mathscr{C}_r(D)$  are wedgable. Then, we have  $(S \wedge R)_K = (R \wedge S)_K$ .

The regularization by the kernel  $\mathcal{K}_{\theta}^{n}$  works well with the wedge product in the sense of King. The following is straightforward from Theorem 1.3 and the arguments near (3.4).

**Theorem 1.5.** Let  $S \in \mathscr{C}_s(D)$  and  $R \in \mathscr{C}_r(D)$  be two currents such that S and R are wedgable. Then, we have

$$\mathscr{L}^{\mathcal{K}^n_\theta}(S) \wedge R = S \wedge \mathscr{L}^{\mathcal{K}^n_\theta}(R) \to (S \wedge R)_K$$

in the sense of currents as  $|\theta| \rightarrow 0$ .

**Remark 5.7.** It is not clear whether the same is true for the kernel  $\mathcal{T}_{\theta}^{n}$ . The difference is that the kernel  $\mathcal{K}_{\theta}^{n}$  has compact support in a neighborhood of the origin while  $\mathcal{T}_{\theta}^{n}$  does not. This may or may not lead to concentration of mass of  $(A_{\lambda})_{*}(\pi_{1}^{*}S \wedge \pi_{2}^{*}R) \wedge \pi_{F}^{*}\omega_{F}^{n}$  in  $H_{\infty}$  as  $|\lambda| \to \infty$ .

**Remark 5.8.** In this work, in the case of an analytic subset Z of pure codimension m in D, we have two notions of the wedge product:  $S \wedge_C [Z]$  and  $(S \wedge [Z])_K$  for  $S \in \mathscr{C}_s(D)$  provided that both of them are well defined. (Previously,  $\wedge_C$  was defined for smooth submanifolds of D, but the notion can actually be defined any analytic subsets satisfying Theorem 1.1.) Then, one can relate  $S \wedge_C [Z]$  to the tangent current of S along Z on the normal bundle of Z in D while  $(S \wedge [Z])_K$  is related to the tangent current of  $\pi_1^*S \wedge \pi_2^*[Z]$  along the diagonal submanifold  $\Delta \subset D \times D$  on the normal bundle of  $\Delta$  in  $D \times D$ .By [21, Lemma 2.3], we see that they are identical. See also [9, Lemma 5.4]. Let  $S \in \mathscr{C}_s(D)$  be a current on D. We consider the following superfunctions, by which we mean functions whose domain and range are a subset of the space of positive closed currents and the set of extended real numbers, respectively. For each current  $R \in \mathscr{C}_{n-s+1}(D)$ , we define

$$\mathscr{F}^{1}_{S,j,\theta}(R) \coloneqq \int_{D_{j+1}} u_{\theta}^{\mathcal{K}} \chi_{j} \pi_{1}^{*} S \wedge \pi_{2}^{*}(R \wedge \omega_{D}^{n-1})$$

This sequence of functions is decreasing as  $|\theta|$  decreases to 0. We allow  $-\infty$  as value and we take the limit as  $|\theta| \to 0$ . We denote by  $\mathscr{F}_{S,j}^1$  the limit function and its domain is written by  $\mathscr{D}_{S,j}^1 := \{R \in \mathscr{C}_{n-s} : \mathscr{F}_{S,j}^1(R) > -\infty\}$ . Further, we define

$$\mathscr{D}^1_S \coloneqq \bigcap_{j \in \mathbb{N}} \mathscr{D}^1_{S,j}.$$

For every  $R \in \mathscr{D}_S^1$ , according to Lemma 3.2, the current  $\langle \pi_1^* S \wedge \pi_2^* (R \wedge \omega_D^{n-2}) \wedge dd^c u \rangle_{\mathcal{K}}$  is well defined. Then, we consider a function defined not on the whole space of  $\mathscr{C}_{n-s}(D)$  but on  $\mathscr{D}_S^1$ . We define

$$\mathscr{F}_{S,j,\theta}^{2}(R) \coloneqq \int_{D_{j+1}} u_{\theta}^{\mathcal{K}} \chi_{j} \left\langle \pi_{1}^{*} S \wedge \pi_{2}^{*} \left( R \wedge \omega_{D}^{n-2} \right) \wedge dd^{c} u \right\rangle_{\mathcal{K}},$$

which is again decreasing in  $|\theta|$ . We take the limit and denote by  $\mathscr{F}_{S,j}^2$  the limit function on  $\mathscr{D}_S^1$ . In the same way, we define its associated sets  $\mathscr{D}_{S,j}^2$  and  $\mathscr{D}_S^2$ . Inductively, we can define  $\mathscr{F}_{S,j}^i, \mathscr{D}_{S,j}^i$ and  $\mathscr{D}_S^i$  for i = 1, ..., n. By construction, we have  $\mathscr{D}_S^{i+1} \subset \mathscr{D}_S^i$ . So, the set  $\mathscr{D}_S^n$  can be used to define a reasonable collection of currents wedgable with the current S. The following proposition is straightforward.

**Proposition 5.9.** For an integer  $1 \le r \le n-s$ , a current  $R \in \mathscr{C}_r(D)$  is wedgable with S if

$$R \wedge \omega_D^{n-s-r+1} \in \mathscr{D}_S^n$$

In this case, for a smooth test form  $\varphi$  of bidegree (n - s - r, n - s - r) on D, we have

$$\langle (S \wedge R)_K, \varphi \rangle = \mathscr{F}_{S,j}^n (R \wedge (M_\varphi \omega_D^{n-s-r+1} + dd^c \varphi)) - \mathscr{F}_{S,j}^n (R \wedge (M_\varphi \omega_D^{n-s-r+1})),$$

where j is an integer such that  $\operatorname{supp} \varphi \in D_j$ , and  $M_{\varphi} > 0$  is a constant such that  $M_{\varphi} \omega_D^{n-s-r+1} + dd^c \varphi$  is positive in the sense of forms.

5.3. Intersection of more than two positive closed currents. In the same spirit, we can define the intersection of many positive closed currents. Let  $D \,\subset \,\mathbb{C}^n$  be a bounded simply connected domain with smooth boundary. We use the coordinates x for D and take a Kähler form  $\omega_D = dd^c |x|^2$ . Let  $k \geq 2$  be an integer. Let  $s_1, \ldots, s_k$  be positive integers such that  $1 \leq s := s_1 + \cdots + s_k \leq n$  and  $S_i \in \mathscr{C}_{s_i}(D)$  currents for  $i = 1, \ldots, k$ . We take as  $U \, D^k = D_1 \times \cdots \times D_k$  and N = kn, where  $D_i$ 's are copies of D. Let  $\pi_i : D^k \to D_i$  denote the canonical projection onto the *i*-th factor for each  $1 \leq i \leq k$ . We use the coordinates  $(x^1, \ldots, x^k) \in D^k$  for  $D^k$  so that  $\pi_i(x^1, \ldots, x^k) = x^i \in D_i$  for each  $1 \leq i \leq k$ . We take as V the diagonal submanifold  $\Delta_{D^k}$  of  $D^k$  defined by  $\Delta_{D^k} := \{(x, \ldots, x) \in D^k : x \in D\}$ . We consider  $D^k$  as a subset of  $\mathbb{C}^{(k-1)n} \times \Delta_{D^k}$ . We take  $E = \mathbb{C}^{(k-1)n} \times \Delta_{D^k}, \overline{E} = \mathbb{P}^{(k-1)n} \times \Delta_{D^k}$  and  $\pi_{\Delta_{D^k}} : \mathbb{P}^{(k-1)n}$  the complementary projection of  $\overline{E}$  onto the fiber space  $\mathbb{P}^{(k-1)n}$ . We will use coordinates  $([w^1 : \cdots : w^{k-1} : t], x) \in \overline{E}$  and identify  $(w_1, \ldots, w^{k-1}, x)$  with  $([w_1 : \cdots : w^{k-1} : 1], x) \in E$ . Then, we have  $\pi_{\Delta_{D^k}}([w^1 : \cdots : w^{k-1} : t], x) = x$ 

and  $\pi_F\left([w^1:\dots:w^{k-1}:t],x\right) = [w^1:\dots:w^{k-1}:t]$ . The inclusion map  $\iota:D^k \to E$  can be written as  $\iota(x^1,\dots,x^k) = (x^1 - x^k,\dots,x^{k-1} - x^k,x^k)$ . We will keep the notations  $\omega_{\Delta_{D^k}} = dd^c |x|^2$  on  $\Delta_{D^k}$ ,  $\omega_F = \frac{1}{2}dd^c \log(|w^1|^2 + \dots + |w^{k-1}|^2 + |t|^2)$  on  $\mathbb{P}^{(k-1)n}$  and  $\omega = \pi^*_{\Delta_{D^k}}\omega_{\Delta_{D^k}} + \pi^*_F\omega_F$  on  $\overline{E}$ . We will also continue to use  $\omega_{D^k} = \pi^*_1\omega_D + \dots + \pi^*_k\omega_D$ . In these coordinates, we may say  $\pi^*_{\Delta_{D^k}}\omega_{\Delta_{D^k}} = \pi^*_k\omega_D$ . We take  $u = \frac{1}{2}\log\sum_{i=1}^{k-1} |x^i - x^k|^2$ .

**Definition 5.10.** We say that  $S_1, \ldots, S_k$  are wedgable if the current  $\pi_1^* S_1 \wedge \cdots \wedge \pi_k^* S_k$  satisfies the horizontal condition with respect to  $\Delta_{D^k}$ .

Remark 5.11. In our coordinates, the horizontal condition can be written as

$$u \in L^1_{loc}\left(\left\langle \pi_1^* S_1 \wedge \dots \wedge \pi_{k-1}^* S_{k-1} \wedge \pi_k^* \left(S_k \wedge \omega_D^{kn-s-i+1}\right) \wedge \mathcal{K}^{i-1} \right\rangle_{\mathcal{K}}\right)$$

inductively from i = 1 through (k - 1)n.

In the same way as in Proposition 5.3, we obtain the following from Proposition 3.10:

**Proposition 5.12.** Let  $s_1, \ldots, s_k$  be positive integers such that  $1 \le s := s_1 + \cdots + s_k \le n$ . Let  $S_i \in \mathscr{C}_{s_i}(D)$  be wedgable currents for  $i = 1, \ldots, k$ . Let  $\varphi$  be a smooth test form with compact support of bidegree (n - s, n - s) on D. Let  $\Phi$  be a smooth test form with compact support of bidegree (n - s, n - s) on  $D^k$  such that  $\Phi = \pi_k^* \varphi$  in a neighborhood of  $\Delta_{D^k}$ . Then, the following integral is well-defined and independent of the choice of  $\Phi$ .

(5.2) 
$$\langle (S_1 \wedge \dots \wedge S_k)_K, \varphi \rangle \coloneqq \int_{D^k \smallsetminus \Delta_{D^k}} \pi_1^* S \wedge \dots \wedge \pi_k^* S_k \wedge \left( u (dd^c u)^{(k-1)n-1} \right) \wedge dd^c \Phi,$$

which defines a positive closed current of bidegree (s, s) on D.

**Definition 5.13.** Let  $S_i \in \mathscr{C}_{s_i}(D)$  be wedgable currents defined on D for i = 1, ..., k, where  $s_i$ 's are positive integers such that  $1 \le s_1 + \dots + s_k \le n$ . We define the wedge product of the currents  $S_1, ..., S_k$  in the sense of King to be the positive closed current (5.2) of bidegree (s, s) in Proposition 5.12 and denote it by  $(S_1 \land \dots \land S_k)_K$ .

Then, we again obtain the following theorem from Theorem 1.2.

**Theorem 5.14.** Let  $S_i \in \mathscr{C}_{s_i}(D)$  be wedgable currents defined on D for i = 1, ..., k, where  $s_i$ 's are positive integers such that  $1 \le s_1 + \cdots + s_k \le n$ . Then, there exists a unique tangent current for  $\pi_1^*S_1 \land \cdots \land \pi_k^*S_k$  along  $\Delta_{D^k}$  and its h-dimension is minimal. Moreover, the shadow of the unique tangent current, which is called the Dinh-Sibony product, coincides with the positive closed currents  $(S_1 \land \cdots \land S_k)_K$  of bidegree (s, s) on D.

We show that the above intersection of many positive closed currents can be approximated in terms of the regularization operator  $\mathscr{L}^{\mathcal{K}^n_{\theta}}(\cdot)$ . For simplicity, we consider the case of k = 3. The general case can be treated in the same way.

**Theorem 5.15.** Let  $S_i \in \mathscr{C}_{s_i}(D)$  be wedgable currents for i = 1, 2, 3, where  $s_i$ 's are positive integers such that  $1 \le s := s_1 + s_2 + s_3 \le n$ . Then, we have

$$S_{1} \wedge \mathscr{L}^{\mathcal{K}_{\theta}^{n}}(S_{2}) \wedge \mathscr{L}^{\mathcal{K}_{\theta}^{n}}(S_{3}) \rightarrow (S_{1} \wedge S_{2} \wedge S_{3})_{K},$$
  
$$\mathscr{L}^{\mathcal{K}_{\theta}^{n}}(S_{1}) \wedge S_{2} \wedge \mathscr{L}^{\mathcal{K}_{\theta}^{n}}(S_{3}) \rightarrow (S_{1} \wedge S_{2} \wedge S_{3})_{K} \quad and$$
  
$$\mathscr{L}^{\mathcal{K}_{\theta}^{n}}(S_{1}) \wedge \mathscr{L}^{\mathcal{K}_{\theta}^{n}}(S_{2}) \wedge S_{3} \rightarrow (S_{1} \wedge S_{2} \wedge S_{3})_{K}$$

in the sense of currents, as  $|\theta| \rightarrow 0$ .

Proof. Let E be the normal bundle of  $\Delta_{D^3}$  in  $D^3$  and  $\overline{E}$  its projective compactification. We continue to assume that  $D^3$  is a subset of E. We use the change of coordinates  $(x^1, x^2, x^3) \rightarrow (x, z, w) = (x^1 - x^3, x^2 - x^3, x^3)$  for E. Then, the coordinates for  $\overline{E}$  can be written as ([x:z:t], w) and (x, z, w) is identified with  $([x:z:1], w) \in E$ . Also, let  $\pi_V : \overline{E} \rightarrow \Delta_{D^3}$  be the canonical projection onto the diagonal submanifold  $\Delta_{D^3}$  and that  $\pi_F : \overline{E} \rightarrow \mathbb{P}^{2n}$ . Namely,  $\pi_V([x:z:t], w) = w$  and  $\pi_F([x:z:t], w) = [x:z:t]$ . We denote by the forms  $\Omega_n^1$  and  $\Omega_n^2$  in the fiber space  $\mathbb{P}^{2n}$  the form  $\Omega_n$  as in Lemma 2.7 with x'' replaced by x and z, respectively.

From the assumption, the current  $\pi_1^*S_1 \wedge \pi_2^*S_2 \wedge \pi_3^*S_3$  satisfies the horizontal condition with respect to  $\Delta_{D^3}$ . Let  $\varphi$  be a smooth test form of bidegree (n - s, n - s) on D. With respect to our coordinates,  $\varphi$  can be understood as a form on  $\Delta_{D^3}$  and so, we may say  $\pi_{\Delta_{D^3}}^*\varphi = \pi_3^*\varphi$ . We have  $A_\lambda(x, z, w) = (\lambda x, \lambda z, w)$ . Then, as in the proof of Theorem 1.2, from Theorem 5.14 together with the independence of the shadow of the tangent current in Definition 2.2, we get

$$\langle (S_1 \wedge S_2 \wedge S_3)_K, \varphi \rangle = \lim_{\theta \to 0} \langle (\pi_{\Delta_{D^3}})_* \left( (A_{\theta^{-1}})_* (\pi_1^* S_1 \wedge \pi_2^* S_2 \wedge \pi_3^* S_3) \wedge \pi_F^* \Omega_n^1 \wedge \pi_F^* \Omega_n^2 \right), \varphi \rangle$$

$$= \lim_{\theta \to 0} \langle (A_{\theta^{-1}})_* (\pi_1^* S_1 \wedge \pi_2^* S_2 \wedge \pi_3^* S_3) \wedge \pi_F^* \Omega_n^1 \wedge \pi_F^* \Omega_n^2, \pi_3^* \varphi \rangle$$

$$= \lim_{\theta \to 0} \langle \pi_1^* S_1 \wedge \pi_2^* S_2 \wedge \pi_3^* S_3 \wedge (A_{\theta^{-1}})^* (\pi_F^* \Omega_n^1 \wedge \pi_F^* \Omega_n^2), \pi_3^* \varphi \rangle$$

$$= \lim_{\theta \to 0} \langle \pi_1^* S_1 \wedge \pi_2^* S_2 \wedge \pi_3^* S_3 \wedge \mathcal{K}_{e^{-M}\theta}^{n,1} \wedge \mathcal{K}_{e^{-M}\theta}^{n,2}, \pi_3^* \varphi \rangle,$$

where M > 0 is a constant as in Lemma 2.7, and  $\mathcal{K}_{e^{-M}\theta}^{n,i}$  denotes the form  $\mathcal{K}_{e^{-M}\theta}^{n}$  with respect to the variable x if i = 1, and z if i = 2. In the above, we need to use the same arguments as in the proof of Theorem 1.2 in order to guarantee that there is no mass concentration in  $H_{\infty} = \overline{E} \times E$ . The current  $\pi_1^* S_1 \wedge \pi_2^* S_2 \wedge \mathcal{K}_{e^{-M}\theta}^{n,1} \wedge \mathcal{K}_{e^{-M}\theta}^{n,2}$  can be considered as a smooth form with respect to  $x^3$  because of the support of  $\mathcal{K}_{e^{-M}\theta}^{n,1} \wedge \mathcal{K}_{e^{-M}\theta}^{n,2}$ . Hence, the above limit can be written as

$$\left\langle \left(S_1 \wedge S_2 \wedge S_3\right)_K, \varphi \right\rangle = \lim_{\theta \to 0} \left\langle (\pi_3)_* \left( \pi_1^* S_1 \wedge \pi_2^* S_2 \wedge \mathcal{K}_{e^{-M}\theta}^{n,1} \wedge \mathcal{K}_{e^{-M}\theta}^{n,2} \right) \wedge S_3, \varphi \right\rangle.$$

We observe that for each  $x^3 \in D_3$ ,  $(\pi_3)_*$  is the integration over the fiber  $D_1 \times D_2$ . So, for each  $x^3 \in D_3$ , since  $\pi_1^*S_1 \wedge \mathcal{K}_{e^{-M}\theta}^{n,1}$  is a form with measures in  $x_1$  as coefficients and  $\pi_2^*S_2 \wedge \mathcal{K}_{e^{-M}\theta}^{n,2}$  is a form with measures in  $x_2$  as coefficients, the Fubini theorem or the notion of double currents implies that for each  $x^3 \in D_3$ , we can split the form  $\pi_1^*S_1 \wedge \pi_2^*S_2 \wedge \mathcal{K}_{e^{-M}\theta}^{n,1} \wedge \mathcal{K}_{e^{-M}\theta}^{n,2}$  into the product of a form in  $x_1$  and another form in  $x_2$ . So, we get

$$\langle (S_1 \wedge S_2 \wedge S_3)_K, \varphi \rangle = \lim_{\theta \to 0} \langle (\pi_3)_* \left( \pi_1^* S_1 \wedge \mathcal{K}_{e^{-M}\theta}^{n,1} \right) \wedge (\pi_3)_* \left( \pi_2^* S_2 \wedge \mathcal{K}_{e^{-M}\theta}^{n,2} \right) \wedge S_3, \varphi \rangle$$
$$= \lim_{\theta \to 0} \left\langle \mathscr{L}_{e^{-M}\theta}^{\mathcal{K}_{e^{-M}\theta}^n}(S_1) \wedge \mathscr{L}_{e^{-M}\theta}^{\mathcal{K}_{e^{-M}\theta}^n}(S_2) \wedge S_3, \varphi \right\rangle.$$

In the above, we abuse notation in the sense that the same notation  $\pi_1$  is used for projections  $\pi_3: D^3 \to D_3$ ,  $\pi_3: D_2 \times D_3 \to D_3$  and  $\pi_3: D_3 \times D_1 \to D_3$ . In the same way, we used  $\pi_1$  and  $\pi_2$ . Or one can say that we identified  $D_2 \times D_3$  with  $\{x_0^1\} \times D_2 \times D_3$  for some  $x_0^1 \in D_1$  and  $D_3 \times D_1$  with  $D_1 \times \{x_0^2\} \times D_3$  for some  $x_0^2 \in D_2$ .

The other cases can be proved exactly in the same way.

The following seems to be intuitively obvious and the proof is quite straightforward but we may need to go through lengthy and tedious computations for the proof. For this reason, we skip the proof of the proposition. For now, we do no know whether there is a more efficient way to obtain it. The case where  $S_3$  is not smooth needs to be further investigated.

**Proposition 5.16.** Suppose that  $S_1 \in \mathscr{C}_{s_1}(D)$  and  $S_2 \in \mathscr{C}_{s_2}(D)$  are wedgable and  $S_3 \in \mathscr{C}_{s_3}(D)$  is smooth. Then, the currents  $S_1$ ,  $S_2$ ,  $S_3$  are wedgable,  $S_1$  and  $(S_2 \wedge S_3)_K$  are wedgable, and we have

$$(S_1 \land S_2 \land S_3)_K = (S_1 \land (S_2 \land S_3)_K)_K = ((S_1 \land S_2)_K \land S_3)_K$$

# 6. TANGENT CURRENTS AND INTERSECTION OF POSITIVE CLOSED CURRENTS ON COMPLEX MANIFOLDS

In this section, we apply our results to compact complex manifolds. Due to Remark 2.4, we can carry the results over to general complex manifolds.

6.1. Tangent currents and intersection of positive closed currents. Let X be a complex manifold of complex dimension N. Let  $V \subset X$  be a smooth complex submanifold of codimension n. Let T be a positive closed current of bidegree (p, p) on X.

**Definition 6.1.** We define a T-cover of V in X to be a countable collection  $(U_j)_{j \in \mathbb{N}}$  of open subsets of X such that each  $U_j$  is biholomorphic to a bounded simply connected domain with smooth boundary sitting inside a single coordinate chart of X, that  $V \subset \bigcup_{j \in \mathbb{N}} U_j$  and that for each  $x \in V$ , there exist a finitely many  $j \in \mathbb{N}$  with  $x \in U_j$ . For simplicity, a T-cover of X in X is called a T-cover of X.

Due to Remark 2.4, we obtain the following theorems, whose proof is quite straightforward. So, we skip proofs.

**Proposition 6.2.** Suppose that there exists a T-cover  $(U_j)_{j \in \mathbb{N}}$  of V in X such that on each  $U_j$  with  $U_j \cap V \neq \emptyset$ , Proposition 2.17 holds. Then, T admits a tangent current along V.

We can also estimate the h-dimension in terms of the local language. Below is an example induced from Proposition 3.7.

**Proposition 6.3.** Suppose that T admits a tangent current  $T_{\infty}$  along V. Let  $1 \le m \le n$  be an integer. Further assume that there exists a T-cover  $(U_j)_{j \in \mathbb{N}}$  of V in X such that on each  $U_j$  with  $U_j \cap V \ne \emptyset$ , T satisfies Condition (I) with respect to V up to codimension m. Then, the h-dimension of the tangent current  $T_{\infty}$  along V is at most N - p - m.

From Theorem 1.3, we obtain the following.

**Theorem 6.4.** Suppose that there exists a T-cover  $(U_j)_{j \in \mathbb{N}}$  of V in X such that on each  $U_j$  with  $U_j \cap V \neq \emptyset$ , T satisfies the horizontal condition. Then, the tangent current  $T_{\infty}$  of T along V exists and is unique. Its local representation is as in Theorem 1.3.

In the same way, the intersection of positive closed currents can be also stated. Let *Y* be a complex manifold of complex dimension  $n_Y$ . Let  $Y^k = Y_1 \times \cdots \times Y_k$  denote the *k*-fold product of the copies of *Y* and  $\pi_i : Y^k \to Y_i$  the canonical projection onto the *i*-th factor. Let  $\Delta_{Y^k} := \{(y, \ldots, y) \in Y^k : y \in Y\}$  be the diagonal submanifold of  $Y^k$ .

**Theorem 6.5.** Let  $s_1, \ldots, s_k$  be positive integers such that  $1 \le s := s_1 + \cdots + s_k \le m$ . Let  $S_1, \ldots, S_k$  be positive closed currents of bidegree  $(s_1, s_1), \ldots, (s_k, s_k)$  on Y, respectively. Let  $\{U_i\}_{i \in I}$  be a T-cover  $(U_j)_{j \in \mathbb{N}}$  of X such that the current  $\pi_1^*(S_1) \land \cdots \land \pi_k^*(S_k)$  on  $U_j \times \cdots \times U_j$  satisfies the horizontal condition with respect to the diagonal submanifold  $\Delta_{Y^k} \cap U_j \times \cdots \times U_j$  for each  $j \in \mathbb{N}$ . Then, the Dinh-Sibony product of the currents  $S_1, \ldots, S_k$  is well defined. Its local representation is as in Theorem 1.4.

Due to the local nature of our approach, Theorem 1.5, which is about regularization of positive closed currents, is not completely carried over to compact complex manifolds. We may be able to use a partition of unity but the regularization obtained by use of a partition of unity is not closed in the global sense.

6.2. **Superfunctions.** We can use a similar idea to the one in Subsection 5.2 to show that on a compact Kähler manifold, a positive closed current with continuous superpotentials is wedgable with every positive closed currents.

Let  $(X, \omega_X)$  be a compact Kähler manifold of dimension n. We will use the coordinates x for X. Let  $\mathfrak{X} = X_1 \times X_2$  denote the product of two copies of X and  $\pi_i : \mathfrak{X} \to X_i$  the canonical projection onto its *i*-th factor for each i = 1, 2. We will use the coordinates (x, y) for  $\mathfrak{X}$ . In this subsection, we use a Kähler form  $\omega_{\mathfrak{X}} := \pi_1^* \omega_X + \pi_2^* \omega_X$  on  $\mathfrak{X}$ . Let  $\Delta$  denotes the diagonal submanifold of  $\mathfrak{X}$ , which is defined by  $\{(x, x) \in \mathfrak{X} : x \in X\}$ . Let  $\pi : \mathfrak{X} \to \mathfrak{X}$  denote the blow-up of  $\mathfrak{X}$  along  $\Delta$ . Then, by a theorem of Blanchard,  $\mathfrak{X}$  is a compact Kähler manifold and let  $\omega_{\mathfrak{X}}$  denote a Kähler form on  $\mathfrak{X}$ . Let  $\widehat{\Delta}$  denote the exceptional divisor. Let  $\Pi_i := \pi_i \circ \pi$  for each i = 1, 2. Let  $\alpha_{\widehat{\Delta}}$  be a smooth real closed form of bidegree (1, 1) cohomologous to  $[\widehat{\Delta}]$ . Then, there exists a quasi-plurisubharmonic function  $u_{\widehat{\Delta}}$  on  $\mathfrak{X}$  such that  $[\widehat{\Delta}] - \alpha_{\widehat{\Delta}} = dd^c u_{\widehat{\Delta}}$ . For local treatment, we follow the notations used in Subsection 5.1.

Let  $(U_j^i)_{j \in J}$  for i = 0, ..., n be finite coverings of X such that each  $U_j^i$  is biholomorphic to a bounded simply connected domain with smooth boundary, and that  $\overline{U_j^i} \in U_j^{i-1}$  for each  $j \in J$  and for each i = 1, ..., n. For each  $j \in J$  and for each i = 1, ..., n, let  $\chi_j^i : U_j^{i-1} \to [0, 1]$  be a smooth function with compact support with  $\chi_j^i \equiv 1$  on the set  $\overline{U_j^i}$ .

Let s and r be positive integers such that  $s + r \le n$ . Let  $S \in \mathscr{C}_s(X)$  and  $R \in \mathscr{C}_{n-s+1}(X)$  be two currents. As in Subsection 5.2, we can inductively define

$$\mathcal{F}_{S,j}^{i}(R) \coloneqq \lim_{|\theta| \to 0} \int_{U_{j}^{i-1}}^{t} u_{\theta}^{\mathcal{K}} \chi_{j}^{i} \left\langle \pi_{1}^{*}S \wedge \pi_{2}^{*} \left( R \wedge \omega_{X}^{n-i} \right) \wedge \mathcal{K}^{i-1} \right\rangle_{\mathcal{K}},$$
  
$$\mathcal{D}_{S,j}^{i} \coloneqq \left\{ R \in \mathcal{D}_{S}^{i-1} : \mathcal{F}_{S,j}^{i} > -\infty \right\} \quad \text{and} \quad \mathcal{D}_{S}^{i} \coloneqq \bigcap_{j \in J} \mathcal{D}_{S,j}^{i} \quad \text{for each} \quad i = 1, \dots, n.$$

The each function  $\mathscr{F}_{S,j}^i$  is defined on  $\mathscr{D}_S^{i-1}$ . By convention,  $\mathscr{D}_S^0 = \mathscr{C}_{n-s+1}(X)$ . Here, for each  $j \in J$ , we use the same coordinate chart for  $U_j^0, \ldots, U_j^n$  and u is the function as in Subsection 5.1 on this coordinate chart. Associated to  $\mathscr{F}_{S,j}^i$ , we define . Again, the following proposition is straightforward.

**Proposition 6.6.** For an integer  $1 \le r \le n - s$ , a current  $R \in \mathscr{C}_r(X)$  is wedgable with S if

$$R \wedge \omega_X^{n-s-r+1} \in \mathscr{D}_S^n$$

In this case, for a smooth test form  $\varphi$  of bidegree (n - s - r, n - s - r) on X, we have

$$\langle (S \wedge R)_K, \varphi \rangle = \sum_{j \in J} \mathscr{F}_{S,j}^n (R \wedge (M_{\varphi,j} \omega_X^{n-s-r+1} + dd^c(\chi_j \varphi))) - \mathscr{F}_{S,j}^n (R \wedge (M_{\varphi,j} \omega_X^{n-s-r+1})),$$

where the collection  $(\chi_j)_{j \in J}$  is a partition of unity subordinate to the cover  $(U_j^n)_{j \in J}$  of X, and for each  $j \in J$ ,  $M_{\varphi,j} > 0$  is a constant such that  $M_{\varphi,j} \omega_X^{n-s-r+1} + dd^c (\chi_j \varphi)$  is positive in the sense of forms.

Now, we consider positive closed currents with continuous superpotentials on compact Kähler manifolds. For definitions and related properties of superpotentials on compact Kähler manifolds, we refer the reader to [8] and [4]. See also [7] for the theory on complex projective spaces.

We claim that if  $S \in \mathscr{C}_s(X)$  admits continuous superpotentials, each function  $\mathscr{F}_{S,j}^i : \mathscr{D}_S^{i-1} \to \mathbb{R} \cup \{-\infty\}$  is upper-semicontinuous in a certain sense for  $i = 1, \ldots, n$ .

**Proposition 6.7.** Suppose that  $S \in \mathscr{C}_s(X)$  admits continuous superpotentials. Let  $R \in \mathscr{D}_S^{i-1}$  be a current. Let  $(R_k)_{k \in \mathbb{N}}$  be a sequence of smooth currents in  $\mathscr{D}_S^{i-1}$  such that  $R_k \to R$  as  $k \to \infty$  and that

there exists an M > 0 such that  $\mathscr{F}_{S,j}^{i-1}(R_k) \ge -M$  for some  $j \in J$ . Then, we have

$$\limsup_{k \to \infty} \mathscr{F}^{i}_{S,j}(R_k) \le \mathscr{F}^{i}_{S,j}(R)$$

**Lemma 6.8.** Let  $(R_k)_{k \in \mathbb{N}}$  and R be as above in Proposition 6.7. Then, we have

$$\lim_{k \to \infty} \left\langle \pi_1^* S \wedge \pi_2^* \left( R_n \wedge \omega_X^{n-i} \right) \wedge \mathcal{K}^{i-1} \right\rangle_{\mathcal{K}} = \left\langle \pi_1^* S \wedge \pi_2^* \left( R \wedge \omega_X^{n-i} \right) \wedge \mathcal{K}^{i-1} \right\rangle_{\mathcal{K}}$$

*Proof.* When i = 1, this is obvious. So, we may assume that  $i \ge 2$ . Let R be a current in  $\mathscr{D}_S^{i-1,M}$  and  $(R_k)_{k\in\mathbb{N}}$  a sequence of currents in  $\mathscr{D}_S^{i-1,M}$  converging to R in the sense of currents. We first claim that the sequence  $\langle \pi_1^*S \wedge \pi_2^* (R_k \wedge \omega_X^{n-i}) \wedge \mathcal{K}^{i-1} \rangle_{\mathcal{K}}$  has locally uniformly bounded mass with respect to  $k \in \mathbb{N}$ .

Let  $\chi_1: U_j^{i-1} \to [0,1]$  be a smooth function with compact support and  $\chi_2: U_j^{i-1} \to [0,1]$  another smooth function with compact support such that  $\chi_1 \equiv 1$  on a neighborhood of supp $\chi_2$ . We have

$$\begin{split} \int_{U_{j}^{i-1} \times U_{j}^{i-1}} \pi_{1}^{*} \chi_{1} \pi_{2}^{*} \chi_{2} \left\langle \pi_{1}^{*} S \wedge \pi_{2}^{*} \left( R_{k} \wedge \omega_{X}^{n-i} \right) \wedge \mathcal{K}^{i-1} \right\rangle_{\mathcal{K}} \\ &= \lim_{|\theta| \to 0} \int_{U_{j}^{i-1} \times U_{j}^{i-1}} \pi_{1}^{*} \chi_{1} \pi_{2}^{*} \chi_{2} \pi_{1}^{*} S \wedge \pi_{2}^{*} \left( R_{k} \wedge \omega_{X}^{n-i} \right) \wedge \left( dd^{c} u_{\theta}^{\mathcal{K}} \right) \wedge \mathcal{K}^{i-2} \\ &= \lim_{|\theta| \to 0} \int_{U_{j}^{i-1} \times U_{j}^{i-1}} u_{\theta}^{\mathcal{K}} \pi_{1}^{*} \chi_{1} dd^{c} \left( \pi_{2}^{*} \chi_{2} \right) \wedge \pi_{1}^{*} S \wedge \pi_{2}^{*} \left( R_{k} \wedge \omega_{X}^{n-i} \right) \wedge \mathcal{K}^{i-2} \\ &+ \int_{U_{j}^{i-1} \times U_{j}^{i-1}} u_{\theta}^{\mathcal{K}} d \left( \pi_{1}^{*} \chi_{1} \right) \wedge d^{c} \left( \pi_{2}^{*} \chi_{2} \right) \wedge \pi_{1}^{*} S \wedge \pi_{2}^{*} \left( R_{k} \wedge \omega_{X}^{n-i} \right) \wedge \mathcal{K}^{i-2} \\ &+ \int_{U_{j}^{i-1} \times U_{j}^{i-1}} u_{\theta}^{\mathcal{K}} d \left( \pi_{2}^{*} \chi_{2} \right) \wedge d^{c} \left( \pi_{1}^{*} \chi_{1} \right) \wedge \pi_{1}^{*} S \wedge \pi_{2}^{*} \left( R_{k} \wedge \omega_{X}^{n-i} \right) \wedge \mathcal{K}^{i-2} \\ &+ \int_{U_{j}^{i-1} \times U_{j}^{i-1}} u_{\theta}^{\mathcal{K}} \pi_{2}^{*} \chi_{2} \left( \pi_{1}^{*} \chi_{1} \right) \wedge \pi_{1}^{*} S \wedge \pi_{2}^{*} \left( R_{k} \wedge \omega_{X}^{n-i} \right) \wedge \mathcal{K}^{i-2} \end{split}$$

The first integral is uniformly bounded with respect to k due to the assumption that  $R_k$ 's and R belong to  $\mathscr{D}_S^{i-1,M}$ . For the last three integrals, the region of integration does not intersection with  $\Delta$ . So, it is actually a definite distance away from  $\Delta$ . Hence, the function u and the form  $dd^c u$  are smooth on the region of integration. So, it is easy to see that the last three integrals are uniformly bounded with respect to k.

Now, we look into limit currents of the sequence  $\left(\left\langle \pi_1^*S \land \pi_2^*\left(R_k \land \omega_X^{n-i}\right) \land \mathcal{K}^{i-1}\right\rangle_{\mathcal{K}}\right)$ . Notice that over  $U_j^{i-1} \times U_j^{i-1} \smallsetminus \Delta$ , every limit current equals  $\pi_1^*S \land \pi_2^*\left(R \land \omega_X^{n-i}\right) \land \mathcal{K}^{i-1}$ . So, we are interested in the restrictions of the limit currents. By replacing  $R_k$  by its subsequence, we may assume that the integrals and the currents both converge. Due to Lemma 3.4, it suffices to consider a test form of the type  $\pi_F^*\chi_{\varepsilon}\pi_{\Delta}^*f$ , where  $f: \left(U_j^{i-1} \times U_j^{i-1}\right) \cap \Delta \to \mathbb{R}$  is a smooth test function on  $\left(U_j^{i-1} \times U_j^{i-1}\right) \cap \Delta$ and  $\chi_{\varepsilon}: \mathbb{C}^n \to [0,1]$  is a smooth function with compact support in the  $\varepsilon$ -neighborhood of the origin. We may assume that  $0 < \varepsilon \ll 1$ . We have

(6.1) 
$$\lim_{k \to \infty} \int \pi_F^* \chi_{\varepsilon} \pi_{\Delta}^* f \left( \pi_1^* S \wedge \pi_2^* \left( R_k \wedge \omega_X^{n-i} \right) \wedge \mathcal{K}^{i-1} \right)_{\mathcal{K}}$$
$$= \lim_{k \to \infty} \lim_{|\theta| \to 0} \int \pi_F^* \chi_{\varepsilon} \pi_{\Delta}^* f \pi_1^* S \wedge \pi_2^* \left( R_k \wedge \omega_X^{n-i} \right) \wedge dd^c u_{\theta}^{\mathcal{K}} \wedge \mathcal{K}^{i-2}$$
$$= \lim_{k \to \infty} \lim_{|\theta| \to 0} \int u_{\theta}^{\mathcal{K}} \pi_F^* \chi_{\varepsilon} dd^c \left( \pi_{\Delta}^* f \right) \wedge \pi_1^* S \wedge \pi_2^* \left( R_k \wedge \omega_X^{n-i} \right) \wedge \mathcal{K}^{i-2}$$
$$+ \int u_{\theta}^{\mathcal{K}} d \left( \pi_F^* \chi_{\varepsilon} \right) \wedge d^c \left( \pi_{\Delta}^* f \right) \wedge \pi_1^* S \wedge \pi_2^* \left( R_k \wedge \omega_X^{n-i} \right) \wedge \mathcal{K}^{i-2}$$

(6.3) 
$$+ \int u_{\theta}^{\mathcal{K}} d\left(\pi_{\Delta}^{*}f\right) \wedge d^{c}\left(\pi_{F}^{*}\chi_{\varepsilon}\right) \wedge \pi_{1}^{*}S \wedge \pi_{2}^{*}\left(R_{k} \wedge \omega_{X}^{n-i}\right) \wedge \mathcal{K}^{i-2}$$

(6.4) 
$$+ \int u_{\theta}^{\mathcal{K}} \pi_{\Delta}^{*} f dd^{c} \left(\pi_{F}^{*} \chi_{\varepsilon}\right) \wedge \pi_{1}^{*} S \wedge \pi_{2}^{*} \left(R_{k} \wedge \omega_{X}^{n-i}\right) \wedge \mathcal{K}^{i-2}.$$

In the same way, we see that

(6.5) 
$$\int \pi_F^* \chi_{\varepsilon} \pi_{\Delta}^* f \left\langle \pi_1^* S \wedge \pi_2^* \left( R_k \wedge \omega_X^{n-i} \right) \wedge \mathcal{K}^{i-1} \right\rangle_{\mathcal{K}}$$
$$= \lim_{|\theta| \to 0} \int u_{\theta}^{\mathcal{K}} \pi_F^* \chi_{\varepsilon} dd^c \left( \pi_{\Delta}^* f \right) \wedge \pi_1^* S \wedge \pi_2^* \left( R_k \wedge \omega_X^{n-i} \right) \wedge \mathcal{K}^{i-2}$$
$$(6.6)$$

(6.6) 
$$+ \int u_{\theta}^{\mathcal{K}} d\left(\pi_{F}^{*}\chi_{\varepsilon}\right) \wedge d^{c}\left(\pi_{\Delta}^{*}f\right) \wedge \pi_{1}^{*}S \wedge \pi_{2}^{*}\left(R_{k} \wedge \omega_{X}^{n-i}\right) \wedge \mathcal{K}^{i-2}$$

(6.7) 
$$+ \int u_{\theta}^{\mathcal{K}} d(\pi_{\Delta}^{*}f) \wedge d^{c}(\pi_{F}^{*}\chi_{\varepsilon}) \wedge \pi_{1}^{*}S \wedge \pi_{2}^{*}(R_{k} \wedge \omega_{X}^{n-i}) \wedge \mathcal{K}^{i-2}$$

(6.8) 
$$+ \int u_{\theta}^{\mathcal{K}} \pi_{\Delta}^* f dd^c \left(\pi_F^* \chi_{\varepsilon}\right) \wedge \pi_1^* S \wedge \pi_2^* (R_k \wedge \omega_X^{n-i}) \wedge \mathcal{K}^{i-1}$$

Because of the region of integration being away from  $\Delta$ , it follows that (6.2), (6.3), (6.4) converge to (6.6), (6.7), (6.8), respectively.

We claim that the limit currents of  $u\pi_F^*\chi_{\varepsilon}\pi_1^*S \wedge \pi_2^*(R_k \wedge \omega_X^{n-i+1}) \wedge \mathcal{K}^{i-2}$  have no mass on  $\Delta$ . Since  $R_k$  is smooth, it satisfies the horizontal condition and the current  $\langle \pi_1^*S \wedge \pi_2^*(R_k \wedge \omega_X^{n-i+1}) \wedge \mathcal{K}^{i-2} \rangle_{\mathcal{K}}$  is well defined and has no mass on  $\Delta$ . So, we have

$$u\left(\pi_1^*S \wedge \pi_2^*(R_k \wedge \omega_X^{n-i+1}) \wedge \mathcal{K}^{i-2}\right)_{\mathcal{K}} = u\left(\pi_1^*S \wedge \pi_2^*(R_k \wedge \omega_2^{n-i+1})\right) \wedge \mathcal{K}^{i-2}\Big|_{\left(U_j^{i-1} \times U_j^{i-1}\right) \setminus \Delta}$$

By change of cooridnates, which is the restriction  $\pi_{\widehat{\mathfrak{X}}\setminus\widehat{\Delta}}$  of the blow-up map, we can compare the currents  $(\pi^*u) (dd^c(\pi^*u))^{i-2} \wedge \prod_1^* S \wedge \prod_2^* (R_k \wedge \omega_X^{n-i+1}) \text{ and } u_{\widehat{\Delta}} \prod_1^* S \wedge \prod_2^* (R_k \wedge \omega_X^{n-i+1}) \wedge \omega_{\widehat{\mathfrak{X}}}^{i-2}$  over the set  $U_j^{i-1} \times U_j^{i-1} \setminus \widehat{\Delta}$ . Writing out the integral in the blown-up space, the form  $dd^c(\pi^*u)$  becomes a bounded smooth form. Since the singularity of  $\pi^*u$  is the same as  $\log \operatorname{dist}(\cdot, \widehat{\Delta})$ , we see that the current  $u \langle \pi_1^* S \wedge \pi_2^* (R_k \wedge \omega_X^{n-i+1}) \wedge \mathcal{K}^{i-2} \rangle_{\mathcal{K}}$  is dominated by  $\pi_* \left( u_{\widehat{\Delta}} \prod_1^* S \wedge \prod_2^* (R_n \wedge \omega_X^{n-i+1}) \wedge \omega_{\widehat{\mathfrak{X}}}^{j-2} \right)$ . Then, since  $R_k \to R$  as  $k \to \infty$ , the mass of  $R_k$  is uniformly bounded independently of k. Therefore, [4, Proposition 2.7] implies the desired conclusion.

As we are interested in the measures restricted to  $\Delta$ , we may shrink  $\varepsilon > 0$  to 0. If we use proper coordinates for  $\overline{E}$  as in Subsection 5.1, we see that  $\pi_{\Delta}^* f$  is a form in  $x^2$  without  $x^1$ . Hence, as  $\varepsilon$  shrinks to 0, the above claim implies that the difference between (6.1) and (6.5) converges to 0. Hence, all together, we just obtained the desired convergence.

Proof of Proposition 6.7. By Lemma 6.8, the function  $R \to \langle \pi_1^* S \land \pi_2^* (R \land \omega_X^{n-i}) \land \mathcal{K}^{i-1} \rangle_{\mathcal{K}}$  is continuous on the space  $\{R, R_1, R_2, \ldots\}$ . So, each function  $\mathscr{F}_{S,j}^i$  can actually be written as a decreasing limit of continuous functions for  $i = 1, \ldots, n$  and  $j \in J$ . Hence it is upper-semicontinuous on  $\{R, R_1, R_2, \ldots\}$  and we obtain the inequality.

**Proposition 6.9.** Let  $S \in \mathscr{C}_s(X)$  be a current with continuous superpotentials. Let  $1 \le r \le n - s$  be an integer. Then, for every  $R \in \mathscr{C}_r(X)$ , S and R are wedgable. Namely, the Dinh-Sibony product of S and R is well defined.

*Proof.* According to [5], every  $R \in \mathscr{C}_r(X)$  can be written as  $R = R^+ - R^-$ , where  $R^{\pm} \in \mathscr{C}_r(X)$  are currents that can be approximated by smooth currents in  $\mathscr{C}_r(X)$  in the same cohomology class as  $R^{\pm}$ , respectively. Let  $(R_k^+)_{k \in \mathbb{N}}$  be a sequence of smooth currents in  $\mathscr{C}_r(X)$  such that  $\{R_k^+\} = \{R^+\}$  and  $\lim_{k \to \infty} R_k^+ = R^+$ . By [4, Proposition 2.5], there exists a constant  $M_{R^+} > 0$  such that  $\int_{\widehat{\mathfrak{X}}} u_{\widehat{\mathfrak{A}}} \prod_1^* S \wedge \prod_2^* (R_k^+ \wedge \omega_X^{n-i}) \wedge \omega_{\widehat{\mathfrak{X}}}^{i-1} > -M_{R^+}$  for  $k \in \mathbb{N}$ . Then, the negativity of the current and the upper-semicontinuity in Proposition 6.7, we see that  $\mathscr{F}_{S,j}^i(R^+)$  is finite for every  $i = 1, \ldots, n$  and  $j \in J$ . The same is true for  $R^-$ . Hence, we can say that S and R are wedgable.

# 7. EXAMPLES

In this section, we consider examples of tangent currents and wedgable positive closed currents on domains or on complex manifolds. We see that the horizontal condition is natural in that the horizontal condition is satisfied in many known cases.

7.1. The classical Lelong number. As discussed in [9] and [21], the Lelong number corresponds to the case where V is a single point. We check that Theorem 3.21 is applicable in this case. Let U be an open subset of  $\mathbb{C}^N$  containing 0. We may assume that  $V = \{0\}$  and  $u = \log |x|$ , where the coordinates x are used for U. Since the dimension of V is 0, we have dim L(u) = 0. By [2, Theorem III.4.5] and [2, Proposition III.4.9], the following integral is always finite

$$\int_U u (dd^c u)^i \wedge T \wedge \omega^{N-p-i} > -\infty$$

for  $0 \le i \le N - p - 1$ . Hence, *T* satisfies the minimally vertical condition and Theorem 3.21 implies that there exists a unique tangent current with *h*-dimension 0.

7.2. **Intersection of analytic subsets.** We can apply our theorems to the intersection of analytic subsets in general position in both domains and complex manifolds. Here, complex manifolds are not limited to compact complex manifolds or Kähler manifolds, but general complex manifolds. Without loss of generality, we only consider domains.

We continue to use the same notations as in Subsection 5.3.

Let  $H_1, \ldots, H_k$  be irreducible analytic subsets of pure codimension  $h_1, \ldots, h_k$  defined on D, respectively, where  $h_1, \ldots, h_k$  are positive integers such that  $1 \le h := h_1 + \cdots + h_k \le n$ . We suppose that the intersection  $H := H_1 \cap \cdots \cap H_k$  is generic in the sense that  $\operatorname{codim} H = h$ .

We take  $T = \pi_1^*[H_1] \land \ldots \land \pi_k^*[H_k]$ . We consider the formal integrals:

$$\int_{D^k} \left( \pi_1^*[H_1] \wedge \ldots \wedge \pi_k^*[H_k] \right) \wedge \left( u (dd^c u)^i \right) \wedge \omega_{\Delta_{D^k}}^{kn-h-i} \quad \text{for } i = 0, \ldots, (k-1)n.$$

The unbounded locus L(u) of u is  $\Delta_{D^k}$  and therefore  $L(u) \cap \text{supp}\left(\pi_1^*[H_1] \wedge \ldots \wedge \pi_k^*[H_k]\right)$  is essentially  $H \coloneqq \bigcap_{i=1}^k H_i$ , which is of complex dimension n-h. So,  $\mathscr{H}_{2(kn-h)-2(k-1)n+1}(L(u) \cap H) =$ 0. By [2, Theorem III.4.5] and [2, Proposition III.4.9], the current  $\pi_1^*[H_1] \wedge \ldots \wedge \pi_k^*[H_k]$  satisfies the horizontal condition with respect to  $\Delta_{D^k}$ . Hence, there exists a unique tangent current, which is denoted by  $[H]_{\infty}$ , along  $\Delta_{D^k}$  and its *h*-dimension is minimal.

Now, we compute the shadow  $[H]^h_{\infty}$  of the unique tangent current  $[H]_{\infty}$ . Let  $\varphi$  be a smooth test form on  $\Delta_{D^k}$  (or equivalently on  $D_k$ ) of bidegree ((k-1)n - h, (k-1)n - h) and  $\Phi$  a smooth form with compact support such that  $\Phi = \pi^*_{\Delta_{D^k}} \varphi$  in a neighborhood of  $\Delta_{D^k}$ .

The support of the current  $(\pi_1^*[H_1] \land \ldots \land \pi_k^*[H_k]) \land_C [\Delta_{D^k}]$  is an analytic subset of dimension n - h and the current is positive and closed. By restricting ourselves to an open subset, we may assume that analytic subsets is irreducible. So, by the support theorem by Federer or Siu, the current should be a constant multiple of [H]. We can find this constant by looking at the regular part of H. So, we may further shrink the support of  $\varphi$  and may assume that each  $H_i$  is a smooth submanifold of codimension  $h_i$  for each  $i = 1, \ldots, k$  and their pairwise intersection is transversal. Let  $\widehat{H} := \pi_1^{-1}H_1 \cap \cdots \cap \pi_k^{-1}H_k$ . Then, from [2, Proposition III.4.12], we see that  $\pi_1^*[H_1] \land \ldots \land \pi_k^*[H_k] = [\widehat{H}]$ .

Then, the desired shadow is

$$\left\langle \left[H\right]_{\infty}^{h},\varphi\right\rangle = \int_{D^{k}}\left[\widehat{H}\right]\wedge\left(u(dd^{c}u)^{(k-1)n-1}\right)\wedge dd^{c}\Phi$$

where  $\Phi$  is a smooth test form on  $D^k$  as above. The set  $L(u) \cap \widehat{H}$  is of dimension n - h and the dimension of the current  $u(dd^c u)^{(k-1)n-1} \wedge [\widehat{H}]$  is n - h + 1. The support theorem for flat currents tells us that the current  $u(dd^c u)^{(k-1)n-1} \wedge [\widehat{H}]$  does not charge any mass on  $L(u) \cap \widehat{H}$ . Also, the current  $u|_{\widehat{H}}(dd^c u|_{\widehat{H}})^{(k-1)n-1}$  has dimension n - h + 1 in  $\widehat{H}$  while the dimension of  $L(u) \cap \widehat{H} = n - h$ . So, again, by the support theorem for flat currents,  $u|_{\widehat{H}}(dd^c u|_{\widehat{H}})^{(k-1)n-1}$  does not charge any mass on  $L(u) \cap \widehat{H}$ . Hence, we have

$$\begin{split} \int_{D^k} u (dd^c u)^{(k-1)n-1} \wedge \left[\widehat{H}\right] \wedge dd^c \Phi &= \int_{\widehat{H}} u|_{\widehat{H}} \left( dd^c u|_{\widehat{H}} \right)^{(k-1)n-1} \wedge dd^c \Phi|_{\widehat{H}} = \int_{\widehat{H}} \left( dd^c u|_{\widehat{H}} \right)^{(k-1)n} \wedge \Phi|_{\widehat{H}} \\ &= \int_{\widehat{H}} \left[ \Delta_{D^k} \cap \widehat{H} \right] \wedge \Phi|_{\widehat{H}} = \int_{\widehat{H}} \left[ \Delta_{D^k} \cap \widehat{H} \right] \wedge \left( \pi^*_{\Delta_{D^k}} \varphi \right) \Big|_{\widehat{H}} = \langle [H], \varphi \rangle \,. \end{split}$$

The third to last equality comes from King's residue formula. The last equality comes from the biholomorphism of  $\Delta_{D^k}$  and D. Indeed, in the coordinates as in Subsection 5.3, we have the desired equality. Hence, we see that the constant equals 1 and we conclude that the shadow is [H] and the tangent current is simply  $\pi^*_{\Delta_{D^k}}[H]$ .

In Section 8, we see that our approach also extends King's work in [15] and gives a reasoanble partial answer to the question in [10, 4.3.20] in the holomorphic category.

**Remark 7.1.** Slightly more generally, we can consider an analytic subset T and T is in a generic position with respect to V, we can apply the same argument as above. Then, we see that its tangent current along V exists and its h-dimension is minimal. Moreover, the unique tangent current is simply the inverse image of the intersection of T and V under the projection of E onto V.

7.3. Intersection with a positive closed current of bidegree (1,1). We continue to use the same notations as in Subsection 5.1. Let D be a simply connected domain in  $\mathbb{C}^n$  with smooth boundary. We use x for the coordinates of D. Let  $1 \le r \le n-1$  be an integer. Let  $S \in \mathscr{C}_1(D)$  and  $R \in \mathscr{C}_r(D)$  be currents. Let f be a plurisubharmonic function such that  $S = dd^c f$ . Suppose that f is locally integrable with respect to the trace measure of R. That means the measure  $|f|(R \land \omega_D^{n-r})$  is locally integrable. We prove that in this classical case, the currents S and R are wedgable.

**Proposition 7.2.** Let S and R be as above. Then, S and R are wedgable and the Dinh-Sibony product is equal to  $dd^c(fR)$ .

**Lemma 7.3.** Let S and R be as above. Then, the current  $\langle \pi_1^* S \wedge \pi_2^* (R \wedge \omega_D^{n-r}) \wedge \mathcal{K}^{n-1} \rangle_{\mathcal{K}}$  is well defined and the map  $S \to \langle \pi_1^* S \wedge \pi_2^* (R \wedge \omega_D^{n-r}) \wedge \mathcal{K}^{n-1} \rangle_{\mathcal{K}}$  is continuous in the sense of currents.

*Proof.* Let  $S \in \mathscr{C}_1(D)$  and  $R \in \mathscr{C}_{n-1}(D)$  be currents. We consider the integrals

$$\int_{D\times D} u \left\langle \pi_1^* S \wedge \pi_2^* \left( R \wedge \omega_D^{2n-r-1-i} \right) \wedge \mathcal{K}^i \right\rangle_{\mathcal{K}}$$

for i = 0, ..., n - 2. Since the bidegree of  $R \wedge \omega_D^{2n-r-1-i}$  is 2n - 1 - i, this integral is 0 for i = 0, 1, ..., n - 2. We only consider the case i = n - 1. By Lemma 3.2, we see that the current  $\langle \pi_1^* S \wedge \pi_2^* (R \wedge \omega_D^{n-r}) \wedge \mathcal{K}^{n-1} \rangle_{\mathcal{K}}$  is well defined for each  $S \in \mathscr{C}_1(D)$  and for each  $R \in \mathscr{C}_{n-1}(D)$ .

We fix currents  $S \in \mathscr{C}_1(D)$  and  $R \in \mathscr{C}_{n-1}(D)$ . Let  $(S_k)_{k \in \mathbb{N}}$  be a sequence of currents in  $\mathscr{C}_1(D)$ such that  $S_k \to S$  as  $k \to \infty$  in the sense of currents. Let D' be an open subset of D with compact closure. Let  $\chi_1 : D \to [0,1]$  and  $\chi_2 : D \to [0,1]$  be smooth functions with compact support such that  $\chi_2 \equiv 1$  on D' and that  $\chi_1 \equiv 1$  on a neighborhood of the support of  $\chi_2$ . We first prove that the family  $(\langle \pi_1^* S_k \land \pi_2^* (R \land \omega_D^{n-r}) \land \mathcal{K}^{n-1} \rangle)_{n \in \mathbb{N}}$  has locally uniformly bounded mass. We basically follow the arguments used in the proof of Lemma 3.2. Then, for  $\theta$  with  $0 < |\theta| \ll 1$ , we have

$$\begin{split} \int_{D\times D} (\pi_{1}^{*}\chi_{1}) (\pi_{2}^{*}\chi_{2}) \left\langle \pi_{1}^{*}S_{k} \wedge \pi_{2}^{*} (R \wedge \omega_{D}^{n-r}) \wedge \mathcal{K}^{n-1} \right\rangle_{\mathcal{K}} \\ &= \lim_{|\theta| \to 0} \int_{D\times D} (\pi_{1}^{*}\chi_{1}) (\pi_{2}^{*}\chi_{2}) \pi_{1}^{*}S_{k} \wedge \pi_{2}^{*} (R \wedge \omega_{D}^{n-r}) \wedge dd^{c}u_{\theta}^{\mathcal{K}} \wedge \mathcal{K}^{n-2} \\ &= \lim_{|\theta| \to 0} \int_{D\times D} u_{\theta}^{\mathcal{K}} (\pi_{1}^{*}\chi_{1}) dd^{c} (\pi_{2}^{*}\chi_{2}) \wedge \pi_{1}^{*}S_{k} \wedge \pi_{2}^{*} (R \wedge \omega_{D}^{n-r}) \wedge \mathcal{K}^{n-2} \\ &+ \int_{D\times D} u_{\theta}^{\mathcal{K}} d (\pi_{1}^{*}\chi_{1}) \wedge d^{c} (\pi_{2}^{*}\chi_{2}) \wedge \pi_{1}^{*}S_{k} \wedge \pi_{2}^{*} (R \wedge \omega_{D}^{n-r}) \wedge \mathcal{K}^{n-2} \\ &+ \int_{D\times D} u_{\theta}^{\mathcal{K}} d (\pi_{2}^{*}\chi_{2}) \wedge d^{c} (\pi_{1}^{*}\chi_{1}) \wedge \pi_{1}^{*}S_{k} \wedge \pi_{2}^{*} (R \wedge \omega_{D}^{n-r}) \wedge \mathcal{K}^{n-2} \\ &+ \int_{D\times D} u_{\theta}^{\mathcal{K}} (\pi_{2}^{*}\chi_{2}) dd^{c} (\pi_{1}^{*}\chi_{1}) \wedge \pi_{1}^{*}S_{k} \wedge \pi_{2}^{*} (R \wedge \omega_{D}^{n-r}) \wedge \mathcal{K}^{n-2}. \end{split}$$

The first integral is 0 for bidegree reason. Since the support of any derivative of  $\chi_1$  is a definite distance away from  $\Delta$ , the last thre integrals converges as  $|\theta| \rightarrow 0$  and  $k \rightarrow \infty$  in this order. Hence, the sequence has uniformly bounded mass.

We show that the sequence  $\left(\left\langle \pi_1^* S_k \wedge \pi_2^* \left(R \wedge \omega_D^{n-r}\right) \wedge \mathcal{K}^{n-1}\right\rangle_{\mathcal{K}}\right)_{n \in \mathbb{N}}$  of currents converges to the current  $\left\langle \pi_1^* S \wedge \pi_2^* \left(R \wedge \omega_D^{n-r}\right) \wedge \mathcal{K}^{n-1}\right\rangle_{\mathcal{K}}$ . Observe that outside the set  $\Delta$ , the limit currents of the sequence  $\left(\left\langle \pi_1^* S_k \wedge \pi_2^* \left(R \wedge \omega_D^{n-r}\right) \wedge \mathcal{K}^{n-1}\right\rangle_{\mathcal{K}}\right)_{n \in \mathbb{N}}$  are equal to  $\left\langle \pi_1^* S \wedge \pi_2^* \left(R \wedge \omega_D^{n-r}\right) \wedge \mathcal{K}^{n-1}\right\rangle_{\mathcal{K}}$ . So, we only need to check the limit currents on  $\Delta$ . As in the proof of Lemma 3.2, by Lemma 3.4, in order to investigate the measure restricted to  $\Delta$ , we only need to apply a test function of the type  $\pi_V^* g$  where g is a smooth test function on  $\Delta$ .

Hence, we claim that for a smooth test function g defined on  $\Delta$ , we have

$$\lim_{k \to \infty} \int_{D \times D} (\pi_F^* \chi_F) (\pi_V^* g) \langle \pi_1^* S_k \wedge \pi_2^* (R \wedge \omega_D^{n-r}) \wedge \mathcal{K}^{n-1} \rangle_{\mathcal{K}}$$
$$= \int_{D \times D} (\pi_F^* \chi_F) (\pi_V^* g) \langle \pi_1^* S \wedge \pi_2^* (R \wedge \omega_D^{n-r}) \wedge \mathcal{K}^{n-1} \rangle_{\mathcal{K}}$$

Here,  $\chi_F : \mathbb{C}^n \to [0,1]$  is a smooth function with compact support such that  $\chi_F \equiv 1$  in a neighborhood of 0. We have

$$\begin{split} \lim_{k \to \infty} \int_{D \times D} \left( \pi_F^* \chi_F \right) \left( \pi_V^* g \right) \left\langle \pi_1^* S_k \wedge \pi_2^* \left( R \wedge \omega_D^{n-r} \right) \wedge \mathcal{K}^{n-1} \right\rangle_{\mathcal{K}} \\ &= \lim_{k \to \infty} \lim_{|\theta| \to 0} \int_{D \times D} \left( \pi_F^* \chi_F \right) \left( \pi_V^* g \right) \left( dd^c u_{\theta}^{\mathcal{K}} \right) \wedge \pi_1^* S_k \wedge \pi_2^* \left( R \wedge \omega_D^{n-r} \right) \wedge \mathcal{K}^{n-2} \\ &= \lim_{k \to \infty} \lim_{|\theta| \to 0} \int_{D \times D} u_{\theta}^{\mathcal{K}} dd^c \left( \left( \pi_F^* \chi_F \right) \left( \pi_V^* g \right) \right) \wedge \pi_1^* S_k \wedge \pi_2^* \left( R \wedge \omega_D^{n-r} \right) \wedge \mathcal{K}^{n-2} \\ &= \lim_{k \to \infty} \lim_{|\theta| \to 0} \int_{D \times D} u_{\theta}^{\mathcal{K}} \left( \pi_V^* g \right) dd^c \left( \pi_F^* \chi_F \right) \wedge \pi_1^* S_k \wedge \pi_2^* \left( R \wedge \omega_D^{n-r} \right) \wedge \mathcal{K}^{n-2} \\ &= u \left( \pi_V^* g \right) dd^c \left( \pi_F^* \chi_F \right) \wedge \pi_1^* S \wedge \pi_2^* \left( R \wedge \omega_D^{n-r} \right) \wedge \mathcal{K}^{n-2}. \end{split}$$

$$\int_{D\times D} \left(\pi_F^* \chi_F\right) \left(\pi_V^* g\right) \left\langle \pi_1^* S \wedge \pi_2^* \left(R \wedge \omega_D^{n-r}\right) \wedge \mathcal{K}^{n-1} \right\rangle_{\mathcal{K}}$$

in the above form, we can see the desired convergence.

*Proof of Proposition 7.2.* The work in [12] proves that the Dinh-Sibony product of the currents S and R is  $dd^c(fR)$ . See also [13]. So, it suffices to prove that S and R are wedgable.

In the above proof, we have seen that

$$\int_{D \times D} u \left\langle \pi_1^* S \wedge \pi_2^* \left( R \wedge \omega_D^{2n-r-1-i} \right) \wedge \mathcal{K}^i \right\rangle_{\mathcal{K}} = 0$$

for i = 0, ..., n - 2. So, we consider the case of i = n - 1.

Let D' be an open subset of D with compact closure. Let  $\chi_1 : D \to [0,1]$  and  $\chi_2 : D \to [0,1]$  be smooth functions with compact support as in the proof of Lemma 7.3. We may assume that f is negative on the support of  $\chi_2$ . Let  $(f_j)$  be a sequence of plurisubharmonic functions decreasingly converging to f and negative on the support of  $\chi_2$ . Since f is locally integrable with respect to  $(R \land \omega_D^{n-r})$ , we have

$$\int_{D\times D} \pi_1^*(\chi_1 f_j) \pi_2^*(\chi_2 R \wedge \omega_D^{n-r}) \wedge [\Delta] = \int_D \chi_2 f_j R \wedge \omega_D^{n-r} \ge \int_D \chi_2 f R \wedge \omega_D^{n-r} > -\infty$$

The current  $\pi_2^*(\chi_2 R \wedge \omega_D^{n-r})$  is closed since it is of maximal bidegree with respect to y. Hence, we have

$$\begin{split} &\int_{D\times D} \pi_1^*(\chi_1 f_j) \pi_2^*(\chi_2 R \wedge \omega_D^{n-r}) \wedge [\Delta] = \int_{D\times D} dd^c \pi_1^*(\chi_1 dd^c f_j) \wedge \pi_2^*(\chi_2 R \wedge \omega_D^{n-r}) \wedge u(dd^c u)^{n-1} \\ &= \int_{D\times D} \pi_1^*(\chi_1 dd^c f_j) \wedge \pi_2^*(\chi_2 R \wedge \omega_D^{n-r}) \wedge u(dd^c u)^{n-1} \\ &+ \int_{D\times D} \pi_1^*(df_j \wedge d^c \chi_1) \wedge \pi_2^*(\chi_2 R \wedge \omega_D^{n-r}) \wedge u(dd^c u)^{n-1} \\ &+ \int_{D\times D} \pi_1^*(d\chi_1 \wedge d^c f_j) \wedge \pi_2^*(\chi_2 R \wedge \omega_D^{n-r}) \wedge u(dd^c u)^{n-1} \\ &+ \int_{D\times D} \pi_1^*(f_j dd^c \chi_1) \wedge \pi_2^*(\chi_2 R \wedge \omega_D^{n-r}) \wedge u(dd^c u)^{n-1}. \end{split}$$

The first equality holds as R is the only current that is not smooth. Observe that the support of any derivative of  $\chi_1$  is disjoint from the support of  $\chi_2$ . So, over the support of  $\pi_2^*(\chi_2 R \wedge \omega_D^{n-r})$ , if the three last integrals are first integrated with respect to y, then they are smooth forms. We consider the second integral, which is of the form

$$\int_D df_j \wedge d^c \chi_1 \wedge \psi$$

where  $\psi = \int_{y \in D} \pi_2^*(\chi_2 R \wedge \omega_D^{n-r}) \wedge u(dd^c u)^{n-1}$  is a smooth (n-1, n-1)-form. The other two can be dealt in the same way. Since the support of  $\chi_1$  is compact, we have

$$\int_D df_j \wedge d^c \chi_1 \wedge \psi = \int_D f_j \wedge dd^c \chi_1 \wedge \psi + \int_D f_j d^c \chi_1 \wedge d\psi.$$

It is bounded by  $||f_j||_{L^1} ||\psi_1||_{C^1}$  up to a multiplicative constant independent of j. Since the support of  $d\chi_1$  and the support of  $\chi_2$  are a definite distance away from each other, the integral is uniformly bounded. Hence, what we have obtained so far is that the first integral

$$\int_{D\times D} \pi_1^*(\chi_1 dd^c f_j) \wedge \pi_2^*(\chi_2 R \wedge \omega_D^{n-r}) \wedge u(dd^c u)^{n-1} > -M_1$$

for some constant  $M_1 > 0$  independent of j. The function  $S \to \pi_1^*(\chi_1 S) \land \pi_2^*(\chi_2 R \land \omega_D^{n-r}) \land u(dd^c u)^{n-1}$  is upper-semicontinuous as Lemma 7.3 says that  $S \to \pi_1^*(\chi_1 S) \land \pi_2^*(\chi_2 R \land \omega_D^{n-r}) \land u_{\theta}^{\mathcal{K}}(dd^c u)^{n-1}$  is continuous and  $S \to \pi_1^*(\chi_1 S) \land \pi_2^*(\chi_2 R \land \omega_D^{n-r}) \land u(dd^c u)^{n-1}$  can be written as

a decreasing limit of continuous functions  $(\pi_1^*(\chi_1 S) \wedge \pi_2^*(\chi_2 R \wedge \omega_D^{n-r}) \wedge u_{\theta}^{\mathcal{K}}(dd^c u)^{n-1})_{0 < |\theta| \ll 1}$ . By the upper-semicontinuity, we conclude that the integral

$$\int_{D\times D} \pi_1^*(\chi_1 dd^c f) \wedge \pi_2^*(\chi_2 R \wedge \omega_D^{n-r}) \wedge u(dd^c u)^{n-1} > -M_1$$

is finite. For every compact subset of  $D \times D$ , we can always find valid  $\chi_1$  and  $\chi_2$  satisfying the support condition. Due to the negativity, the current  $\pi_1^*(dd^c f) \wedge \pi_2^*(R \wedge \omega_D^{n-r}) \wedge u(dd^c u)^{n-1}$  is locally integrable, which means  $dd^c f$  and S are wedgable.

# 8. SLICING THEORY AND THE CURRENTS DEFINED BY ANALYTIC VARIETIES

In this section, we present another type of application of our approach. The basic idea is to exploit [9, Proposition 3.5] together with (2.1). It can be used to create many other regularizations. We again continue to use the same spaces, coordinates, maps, forms and currents as in Subsection 5.1 except that in this section, we take  $\pi_{\Delta} : E \to \Delta$  to be the orthogonal projection onto  $\Delta$  with respect to the usual Euclidean metric.

Let  $S \in \mathscr{C}_s(D)$  and  $R \in \mathscr{C}_r(D)$  be two wedgable currents. Recall from Proposition 4.8 that  $g : \mathbb{C}^n \to \mathbb{R}_{\geq 0}$  is a smooth function with compact support in the unit ball such that  $\int_{\mathbb{C}^n} g \, d\mu = 1$  and that for  $z_1, z_2 \in \mathbb{C}^n$  with  $|z_1| = |z_2|$ , we have  $g(z_1) = g(z_2)$ , where  $\mu$  denotes the standard Lebesgue measure. We also used  $\tau_z(x) = x - z$  for  $z \in \mathbb{C}^n$ .

Let  $\varphi$  be a smooth test form of bidegree (n - s - r, n - s - r) on D. Further, we may assume that  $\varphi = f\Theta$ , where f is a positive smooth function with compact support in D and  $\Theta$  is a smooth positive closed form of bidegree (n - s - r, n - s - r) on D. Indeed, any smooth test form of bidegree (n - s - r, n - s - r) can be written as a finite sum of such forms of the same bidegree.

Since currents *S* and *R* are wedgable, the current  $\pi_1^* S \wedge \pi_2^* (R \wedge \Theta)$  on  $D \times D$  admits tangent current  $(S \wedge R \wedge \Theta)_{\infty}$  along  $\Delta$  and its *h*-dimension is minimal due to Proposition 3.15. We choose  $\Omega = gd\mu = \frac{1}{n!}g(z)(dd^c|z|^2)^n$ . Observe that it has compact support in  $\mathbb{C}^n$ , belongs to the same cohomology class as [z = 0] does, and therefore, it can trivially be extended to  $\mathbb{P}^n$ .

Notice that we have  $\pi_{\Delta}^*((\pi_2^*f)|_{\Delta}) = f(\frac{x+y}{2})$  and that with respect to the coordinates  $(x, y) \in E$ , we have  $A(\lambda)(x, y) = (\frac{x+y}{2} + \lambda \frac{x-y}{2}, \frac{x+y}{2} - \lambda \frac{x-y}{2})$ . We take z = x - y. Then, arguing as in the proof of Theorem 1.2, we have

$$\langle (S \wedge R)_{K}, f\Theta \rangle = \langle (S \wedge R \wedge \Theta)_{K}, f \rangle = \langle (S \wedge R \wedge \Theta)_{\infty}^{h}, f \rangle = \langle (\pi_{\Delta})_{*} ((S \wedge R \wedge \Theta)_{\infty} \wedge \pi_{F}^{*}\Omega), (\pi_{2}^{*}f)|_{\Delta} \rangle$$

$$= \int_{\overline{E}} f\left(\frac{x+y}{2}\right) (S \wedge R \wedge \Theta)_{\infty} \wedge \left(\frac{1}{n!}g(z)(dd^{c}|z|^{2})^{n}\right)$$

$$= \lim_{\epsilon \to 0} \int_{\overline{E}} f\left(\frac{x+y}{2}\right) (A_{\epsilon^{-1}})_{*} (\pi_{1}^{*}S \wedge \pi_{2}^{*}(R \wedge \Theta)) \wedge \left(\frac{1}{n!}g(z)(dd^{c}|z|^{2})^{n}\right)$$

$$= \lim_{\epsilon \to 0} \int_{\overline{E}} f\left(\frac{x+y}{2}\right) \pi_{1}^{*}S \wedge \pi_{2}^{*}(R \wedge \Theta) \wedge (A_{\epsilon^{-1}})^{*} \left(\frac{1}{n!}g(z)(dd^{c}|z|^{2})^{n}\right)$$

$$= \lim_{\epsilon \to 0} \left[\int_{\overline{E}} f(y)\pi_{1}^{*}S \wedge \pi_{2}^{*}(R \wedge \Theta) \wedge (A_{\epsilon^{-1}})^{*} \left(\frac{1}{n!}g(z)(dd^{c}|z|^{2})^{n}\right) \right]$$

$$+ \int_{\overline{E}} \left(f\left(\frac{x+y}{2}\right) - f(y)\right) \pi_{1}^{*}S \wedge \pi_{2}^{*}(R \wedge \Theta) \wedge (A_{\epsilon^{-1}})^{*} \left(\frac{1}{n!}g(z)(dd^{c}|z|^{2})^{n}\right)$$

$$= \lim_{\epsilon \to 0} \left[\int_{\overline{E}} f(y)\pi_{1}^{*}S \wedge \pi_{2}^{*}(R \wedge \Theta) \wedge (A_{\epsilon^{-1}})^{*} \left(\frac{1}{n!}g(z)(dd^{c}|z|^{2})^{n}\right)\right]$$

$$(8.1)$$

(8.2) 
$$+ \int_{\overline{E}} \left( f\left(y + \frac{x - y}{2}\right) - f(y) \right) \pi_1^* S \wedge \pi_2^* (R \wedge \Theta) \wedge (A_{\epsilon^{-1}})^* \left(\frac{1}{n!} g(z) (dd^c |z|^2)^n \right) \right]$$

For sufficiently small  $\epsilon > 0$ , the support of  $\sup \left( f\left(y + \frac{x-y}{2}\right) - f(y) \right) \cap \Delta_{\epsilon}$  lies inside the set  $\{\pi_{\Delta}^{*}((\pi_{2}^{*}\chi_{f})|_{\Delta}) \equiv 1\}$  for some smooth function  $\chi_{f} : D \to [0,1]$  with compact support. Hence, for

all sufficiently small  $\epsilon > 0$ , we have

$$|(\mathbf{8.2})| = \left| \int_{\overline{E}} \left( f\left(y + \frac{x - y}{2}\right) - f(y) \right) \pi_1^* S \wedge \pi_2^* (R \wedge \Theta) \wedge (A_{\epsilon^{-1}})^* \left( \frac{1}{n!} g(z) (dd^c |z|^2)^n \right) \right| \\ \leq \epsilon \|f\|_{C^1} \left| \left\langle (S \wedge R \wedge \Theta)_K, \chi_f \right\rangle \right|$$

Hence, the second term converges to 0. The first term can be written as below:

$$(8.1) = \int_{\overline{E}} f(y) \pi_1^* S \wedge \pi_2^* (R \wedge \Theta) \wedge (A_{\epsilon^{-1}})^* \left(\frac{1}{n!} g(z) (dd^c |z|^2)^n\right)$$
$$= \int_{\overline{E}} \pi_1^* S \wedge \pi_2^* (R \wedge f\Theta) \wedge \left(\frac{1}{n! \epsilon^{2n}} g\left(\frac{z}{\epsilon}\right) (dd^c |z|^2)^n\right)$$
$$= \int_{y \in D} \left(\int_{z \in \mathbb{C}^n} S(y+z) \wedge \left(\frac{1}{\epsilon^{2n}} g\left(\frac{z}{\epsilon}\right) dV(z)\right)\right) \wedge R \wedge f\Theta$$
$$= \langle S_{\epsilon} \wedge R, f\Theta \rangle.$$

Summarizing the above arguments, we obtain

**Theorem 1.6.** Let D be a bounded simply connected domain with smooth boundary. Let  $S \in \mathscr{C}_s(D)$  and  $R \in \mathscr{C}_r(D)$  be two wedgable currents. Let  $(S_{\epsilon})_{0 < \epsilon \ll 1}$  be a standard regularization by convolution. Then, we have

$$S_{\epsilon} \wedge R, S \wedge R_{\epsilon} \rightarrow (S \wedge R)_K$$

in the sense of currents as  $\epsilon \to 0$ . Also, this exactly means that the slice of  $\pi_1^* S \wedge \pi_2^* R$  along  $\Delta$  with respect to the submersion  $D \times D \to \Delta^{\perp}$  exists, where  $\Delta^{\perp}$  is the orthogonal complement of  $\Delta$  in  $D \times D$ . Together with Subsection 7.2, it shows that our approach extends the result of King's in [15] and also, our approach partially answers a question in [10, 4.3.20] in the holomorphic category.

Notice that the same applies to the intersection of many positive closed currents. In general, as in Subsection 5.3, we have

$$(S_1)_{\epsilon} \wedge \dots \wedge (S_{k-1})_{\epsilon} \wedge S_k \to (S_1 \wedge \dots \wedge S_k)_K$$

in the sense of currents, as  $\epsilon \rightarrow 0$ .

**Acknowledgments.** The research of the author was supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government (MSIT) (No. RS-2023-00250685). The paper was partially prepared during the visit of the author at the University of Michigan, Ann Arbor. He would like to express his gratitude to the organization for hospitality. The author would like to thank Dan Burns and Mattias Jonsson for constructive discussions and for their support.

# REFERENCES

- [1] DEMAILLY, J. -P.: Courants positifs et théorie de l'intersection, Gaz. Math., 53 (1992), 131-159.
- [2] DEMAILLY, J. -P.: Complex Analytic and Differential Geometry
- [3] DE RHAM, G.: Differentiable manifolds, Grundlehren Math. Wiss., 266 [Fundamental Principles of Mathematical Sciences] Springer-Verlag, Berlin, 1984, x+167 pp.
- [4] DINH, T. -C.; NGUYEN, V. -A.; VU, D. V.: Super-potentials, densities of currents and number of periodic points for holomorphic maps, Adv. Math. 331 (2018), 874–907
- [5] DINH, T. -C.; SIBONY, N.: Regularization of currents and entropy, Ann. Sci. Éc. Norm. Supér. 37 (2004), 959–971
- [6] DINH, T. -C.; SIBONY, N.: Pull-back of currents by holomorphic maps, Manuscripta Math. 123 (2007), no. 3, 357–371
- [7] DINH, T.-C.; SIBONY, N.: Super-potentials of positive closed currents, intersection theory and dynamics, Acta Math., 203 (2009), 1–82
- [8] DINH, T.-C.; SIBONY, N.: Super-potentials for currents on compact Kähler manifolds and dynamics of automorphisms, J. Algebraic Geom. 19 (2010), no. 3, 473–529

- [9] DINH, T.-C.; SIBONY, N.: Density of positive closed currents: A theory of non-generic intersections, J. Algebraic Geom. 27 (2018), no. 3, 497–551.
- [10] FEDERER, H.: Geometric measure theory, Die Grundlehren der mathematischen Wissenschaften, Band 153 Springer-Verlag New York, Inc., New York, 1969. xiv+676 pp.
- [11] GRIFFITHS, P. A.; KING, J.: Nevanlinna theory and holomorphic mappings between algebraic varieties, Acta Math. 130 (1973), 145–220
- [12] HUYNH, D. T.; KAUFMANN, L.; Vu, D. -V.: Intersection of (1,1)-currents and the domain of definition of the Monge-AmpÈre operator, Indiana Univ. Math. J. 72 (2023), no. 1, 239–261.
- [13] KAUFFMAN, L.; VU, D.-V: Density and intersection of (1,1)-currents, J. Funct. Anal. 277 (2019), 392–417.
- [14] KING, J. R.: A residue formula for complex subvarieties Proc. Carolina conf. on holomorphic mappings and minimal surfaces, Univ. of North Carolina, Chapel Hill , (1970) 43–56.
- [15] KING, J. R.: The currents defined by analytic varieties, Acta Math. 127 (1971), no. 3-4, 185-220.
- [16] LUO, M.; ZHOU, Q.: Equidistribution of saddle periodic points for Hénon-like maps, arXiv:2502.20103
- [17] NGUYEN, V. -A.: The generalized Lelong numbers and intersection theory, arXiv:2501.02150
- [18] NGUYEN, V. -A.; TRUONG, T. T.: Uniqueness of tangent currents for positive closed currents, arXiv:2502.06532
- [19] RASHKOVSKII, A.: Singularities of plurisubharmonic functions and positive closed currents, arXiv:math/0108159
- [20] VU, D.-V.: Intersection of positive closed currents of higher bidegrees, Michigan Math. J. 65 (2016), no. 4, 863–872.
- [21] VU, D.-V.: Densities of currents on non-Kählermanifolds, Int. Math. Res. Not. IMRN 2021, no. 17, 13282–13304.

(Ahn) Department of Mathematics Education, Inha University, 100 Inha-ro, Michuhol-gu, Incheon 22212, Republic of Korea

Email address: t.ahn@inha.ac.kr