

A SPLITTING THEOREM FOR MANIFOLDS WITH SPECTRAL NONNEGATIVE RICCI CURVATURE AND MEAN-CONVEX BOUNDARY

HAN HONG AND GAOMING WANG

ABSTRACT. We prove a splitting theorem for a smooth noncompact manifold with (possibly noncompact) boundary. We show that if a noncompact manifold of dimension $n \geq 2$ has $\lambda_1(-\alpha\Delta + \text{Ric}) \geq 0$ for some $\alpha < \frac{4}{n-1}$ and mean-convex boundary, then it is either isometric to $\Sigma \times \mathbb{R}_{\geq 0}$ for a closed manifold Σ with nonnegative Ricci curvature or it has no interior ends.

1. INTRODUCTION

The classical Cheeger-Gromoll splitting theorem [CG72] states that a complete noncompact manifold with at least two ends and nonnegative Ricci curvature must be isometrical to $\Sigma \times \mathbb{R}_+$ where Σ is a closed manifold with nonnegative Ricci curvature. This result has been extended to manifolds with compact boundary by Kasue [Kas83] and Croke-Kleiner [CK92]. More specifically, a noncompact manifold with a compact mean-convex boundary and nonnegative Ricci curvature is isometric to $\mathbb{R}_+ \times \Sigma$. The compactness of the boundary is essential due to the example that the catenoid bounds a domain in \mathbb{R}^n .

Recently, Antonelli-Pozzetta-Xu [APX24] extended the Cheeger-Gromoll splitting theorem to manifolds with nonnegative Ricci curvature in the spectral sense (to be defined below). Catino-Mari-Mastrolia-Roncoroni [CMMR24] proved the same theorem using a different approach. Many classical and important theorems in Riemannian geometry have been generalized under this framework, for example, [AX24]. In particular, these extensions have played a significant role in the resolution of the stable Bernstein problem [CL23, CLS24, Maz24]. The goal of this paper is to establish a spectral splitting theorem for manifolds with boundary.

Let (M, g) be a non-compact n -dimensional Riemannian manifold with boundary ∂M . Denote the outward unit normal vector of ∂M by η . One can define the Lipschitz function $\text{Ric} : M \rightarrow \mathbb{R}$ by

$$\text{Ric}(x) = \inf\{\text{Ric}(v, v) : g(v, v) = 1 \text{ and } v \in T_x M\}.$$

For a constant $\alpha \geq 0$, one says that M has nonnegative α -Ricci curvature in the spectral sense if $\lambda_1(-\alpha\Delta + \text{Ric}) \geq 0$.

Equivalently, this can be expressed as

$$\int_M \alpha |\nabla \varphi|^2 + \text{Ric} \cdot \varphi^2 \geq 0$$

for any compactly supported function φ , or there exists a positive function $u \in C^{2,\alpha}(M)$ satisfying

$$(1.1) \quad -\alpha \Delta u + \text{Ric} \cdot u = 0 \quad \text{on } M,$$

$$(1.2) \quad \frac{\partial u}{\partial \eta} = 0 \quad \text{on } \partial M.$$

This equivalence follows from arguments similar to those of Fischer-Colbrie and Schoen [FCS80].

The boundary ∂M consists of components that may be compact or noncompact. As in the case of geodesically complete manifolds, one can define the notion of ends. Fixing a point $p \in \partial M$ and choosing a sufficiently large subset containing p , such as the geodesic ball $B_R(p)$ for large R , we consider the connected unbounded components of $M \setminus B_R(p)$. These are called the ends of M with respect to $B_R(p)$. Fix a point $p \in \partial M$, and choose a large subset containing p , e.g., $B_R(p)$ with large R , such that the complement $M \setminus B_R(p)$ has finitely many connected unbounded components. These components are called ends with respect to $B_R(p)$. There are two types of ends: some have a boundary portion belonging to ∂M , while others only have a boundary belonging to $\partial B_R(p)$. The former are called boundary ends, and the latter are called interior ends. In dimension two, these concepts have been introduced to study noncompact capillary surfaces by the author [HS23]. An example is that one can imagine attaching very tiny half-cylinders to a long strip at integer interior points. See also [Wu23, Hon24] for some discussions in higher dimensions.

The main result of this paper is the following.

Theorem 1.1. *Let (M, g) be a smooth noncompact n -dimensional Riemannian manifold with mean-convex boundary and nonnegative α -Ricci curvature in the spectral sense. Assume that $\alpha < \frac{4}{n-1}$. Then, either*

1. M is isometric to a product space $\Sigma \times \mathbb{R}_{\geq 0}$ with a compact manifold Σ satisfying $\text{Ric}_\Sigma \geq 0$, or
2. M has no interior ends.

Notice that the compactness of ∂M is not assumed a priori but rather emerges as a consequence of the splitting result. The proof of the theorem relies on two key techniques: the second variation of the weighted length functional and the curve-capture argument.

First, we analyze the second variation of the weighted length functional. By computing the second variation, we establish that the associated weighted function remains constant along any weighted minimizing line or ray. Additionally, it implies that the Ricci curvature is non-negative along such curves.

Next, we employ the curve-capture technique to construct a minimizing ray or line passing through any given point $p \in M$. This construction is analogous to the prescribed surface method used to resolve the Milnor conjecture in dimension 3 [Liu13]. Further applications of this technique in splitting theorems can be found in [CEM19, CLZ24, HW25]. In particular, our approach provides a distinct strategy compared to those in [APX24, CMMR24]. Finally, we conclude that the Ricci curvature is non-negative on M and finish the proof by the result due to Sakurai [Sak17].

We recall that, given two orthonormal vectors $x, y \in T_p N$, the α -biRic curvature operator is defined as

$$\text{biRic}_\alpha(x, y) := \text{Ric}(x, x) + \alpha \text{Ric}(y, y) - R(x, y, x, y).$$

One can define the α -biRic curvature as

$$\text{biRic}_\alpha(p) = \min\{\text{biRic}_\alpha(x, y) : x, y \text{ are orthonormal vectors in } T_p N\}.$$

When $\alpha = 1$, the α -biRic curvature is just the biRic curvature firstly raised by Shen-Ye[SY96] to study stable minimal hypersurfaces. The α -biRic curvature has shown its importance in the resolution of stable Bernstein problem in \mathbb{R}^{n+1} (see [CLMS24, Maz24]). We say that M has nonnegative biRic curvature if $\text{biRic}(p) \geq 0$ for any point p on N . This condition is weaker than nonnegative sectional curvature but stronger than nonnegative scalar curvature. We show two interesting consequences of the main theorem.

Corollary 1.2. *Let $(N, \partial N, g)$ be a smooth $(n+1)$ -dimensional Riemannian manifold with boundary and $n \leq 4$. Assume that $\text{biRic} \geq 0$ and ∂N is weakly mean-convex. Then, for any two-sided noncompact stable free boundary minimal immersion $(M, \partial M) \rightarrow (N, \partial N, g)$, either*

- 1, M splits isometrically as $\Sigma \times \mathbb{R}_{\geq 0}$, where $\text{Ric}_\Sigma \geq 0$. Moreover, M is totally geodesic in N , and $\text{Ric}_N(\nu_M, \nu_M) = 0$ along M , or
- 2, M has no interior ends.

This corollary can be compared to [Wu23, Theorem 5.3], where the author estimates the number of nonparabolic ends under stronger curvature conditions. In general, we have a more general result.

Corollary 1.3. *Let $(N, \partial N, g)$ be a smooth $(n+1)$ -dimensional Riemannian manifold with boundary and $n \leq 4$. Assume that for some $\alpha \in (\frac{n-1}{n}, \frac{4}{n-1})$,*

$$\inf_N \text{biRic}_\alpha + \frac{4\alpha^2 - n + 1}{4\alpha((\alpha - 1)n + 1)} H^2 \geq 0,$$

and ∂N is weakly mean-convex. Then, for any two-sided noncompact stable free boundary CMC H -hypersurface M in N , either

- 1, M splits isometrically as $\Sigma \times \mathbb{R}_{\geq 0}$, where $\text{Ric}_\Sigma \geq 0$. Moreover, M is totally geodesic in N , and $\text{Ric}_N(\nu_M, \nu_M) = 0$ along M , or
- 2, M has no interior ends.

In Section 2, we prepare some calculations about the weighted length functional L_u^α . In Section 3, we show how to construct a L_u^α -minimizing ray passing through any given point. In Section 4, we prove a splitting theorem for manifolds with boundary and pointwise nonnegative Ricci curvature. In the last Section 5, we prove our main result and its corollary.

1.1. Acknowledgements. The first author is supported by NSFC No. 1240-1058 and the Talent Fund of Beijing Jiaotong University No. 2024XKRC008. The second author is supported by the China Postdoctoral Science Foundation (No. 2024M751604).

2. A WEIGHTED LENGTH FUNCTIONAL

Assume that (M, g) is a noncompact manifold with boundary ∂M and the unit outward normal η . Let u be a positive smooth function on M . Fix a positive constant α which will be determined in later sections. For any curve $\gamma(s)$ parametrized by arclength s that is proper in the sense that $\gamma \cap \partial M = \partial\gamma$, consider the following functional

$$(2.1) \quad L_u^\alpha(\gamma) = \int_\gamma u^\alpha ds.$$

Definition 2.1. A curve $\gamma : I \subset \mathbb{R} \rightarrow M$ is said to be a L_u^α -minimizing curve in the free boundary sense if it satisfies the following two conditions for any $s_1, s_2 \in I$:

- (i) For any curve $\tilde{\gamma}$ connecting $\gamma(s_1)$ and $\gamma(s_2)$, we have

$$L_u^\alpha(\tilde{\gamma}) \geq L_u^\alpha(\gamma([s_1, s_2])).$$

- (ii) For any two curves $\tilde{\gamma}_1$ and $\tilde{\gamma}_2$, each connecting $\gamma(s_i)$ ($i = 1, 2$) to ∂M , we have

$$L_u^\alpha(\tilde{\gamma}_1) + L_u^\alpha(\tilde{\gamma}_2) \geq L_u^\alpha(\gamma([s_1, s_2])).$$

In particular, we say γ is L_u^α -minimizing if γ satisfies only the first condition.

Remark 2.2. If $\gamma : I \rightarrow M$, parametrized by arc length, is an L_u^α -minimizing curve in the free boundary sense and is a complete curve, then I can only be $[0, +\infty)$ or $(-\infty, +\infty)$ after a reparametrization.

- If $I = (-\infty, +\infty)$, we call γ a L_u^α -minimizing line in the free boundary sense.
- If $I = [0, +\infty)$, we call γ a L_u^α -minimizing ray in the free boundary sense.

2.1. Weighted minimizing ray. Let $\gamma(s) : [0, b) \rightarrow M$ be a curve on M with $\gamma(0) \in \partial M$ parametrized by arclength s and it is proper in the sense that $\gamma \cap \partial M = \gamma(0)$. We allow that $b = +\infty$ and in this case γ is a ray.

We denote $\{e_1, \dots, e_n\}$ to be a system of orthonormal vector fields parallel along γ such that $e_n = \frac{\partial}{\partial \gamma_s}$. We simply denote e_n by ∂s . Notice that $\partial s|_{\gamma(0)} = -\eta$.

Lemma 2.3. *If γ_t is a smooth proper 1-parameter family of curves with $\gamma_0 = \gamma$, $\gamma_t(0) \in \partial M$ and $\gamma_t([a, b]) = \gamma([a, b])$, denote the normal speed along the direction e_i at $t = 0$ by $\phi = \langle \frac{\partial \gamma_t}{\partial t} |_{t=0}, e_i \rangle$, then*

$$(2.2) \quad \left. \frac{dL_u^\alpha(\gamma_t)}{dt} \right|_{t=0} = \int_\gamma \kappa_i u^\alpha \phi + \langle \nabla u^\alpha, e_i \rangle \phi ds + \langle \frac{\partial \gamma_t}{\partial t}, \eta \rangle |_{\gamma(0)} u^\alpha(\gamma(0))$$

In particular, the critical curve γ of L_u^α satisfies

$$(2.3) \quad \kappa_i = -\langle \nabla_{\partial s} \partial s, e_i \rangle = -\alpha \nabla_{e_i} \log u$$

along γ and it is perpendicular to ∂M at $\gamma(0)$.

Lemma 2.4. *Let γ be a L_u^α -minimizing curve in the free boundary sense. Let γ_t be a smooth proper 1-parameter family of curves with $\gamma_0 = \gamma$, $\gamma_t(0) \in \partial M$, and $\gamma_t([a, b]) = \gamma([a, b])$, denote the normal speed by ϕ at $t = 0$ along the direction e_i . Then the quadratic form $Q_u(\phi) := \left. \frac{d^2 L_u^\alpha(\gamma_t)}{dt^2} \right|_{t=0} \geq 0$ where*

$$\begin{aligned} Q_u(\phi) &= \int_\gamma \phi_s^2 u^\alpha - R(\partial s, e_i, \partial s, e_i) u^\alpha \phi^2 - (\alpha^2 + \alpha) \phi^2 u^\alpha (\nabla_{e_i} \log u)^2 ds \\ &\quad + \int_\gamma \alpha \phi^2 u^{\alpha-1} (\nabla^2 u)(e_i, e_i) ds - h_{\partial M}(e_i, e_i) u^\alpha \phi^2 |_{\gamma(0)} \end{aligned}$$

where $R(\partial s, e_i, \partial s, e_i)$ is the sectional curvature of the plane spanned by $\{\partial s, e_i\}$ and $h_{\partial M}(\cdot, \cdot) = \langle \nabla \cdot, \eta \rangle$ is the second fundamental form of ∂M with respect to the outer unit normal η .

Proof. Differentiating (2.2) and evaluating at $t = 0$ yields

$$\begin{aligned} Q_u(\phi) &= \int_\gamma u^{\alpha-1} \phi ((-\phi_{ss} - R(\partial s, e_i, \partial s, e_i) \phi - \kappa_i^2 \phi) u + \kappa_i \phi \nabla_{e_i} u) ds \\ &\quad + \int_\gamma \alpha \phi u^{\alpha-1} (\nabla^2 u)(e_i, e_i) + \alpha \phi u^{\alpha-1} \langle \nabla u, -\phi_s \partial s \rangle ds. \end{aligned}$$

Notice that we can write, at the point $\gamma(0)$,

$$\phi \phi_s = \langle V, \nabla_\eta V \rangle = \langle V, \nabla_V \eta \rangle = -\phi^2 h_{\partial M}(e_i, e_i)$$

where V is the associated variational vector field. Then using integration by parts and the critical equation (2.3) completes the proof. \square

Summing over all i we have

Lemma 2.5. *Let γ be a L_u^α -minimizing curve in the free boundary sense. Then for any compact supported function ϕ on γ , we obtain*

$$\begin{aligned} (n-1) \int_\gamma \phi_s^2 u^\alpha ds &\geq \int_\gamma \left(\text{Ric}(\partial s, \partial s) - \alpha \frac{\Delta u}{u} \right) u^\alpha \phi^2 ds - 2\alpha \int_\gamma \phi u^{\alpha-1} u_s \phi_s ds \\ &\quad + \alpha \int_\gamma \phi^2 u^\alpha |\nabla^\perp \log u|^2 ds - \alpha(\alpha-1) \int_\gamma u^{\alpha-2} \phi^2 u_s^2 \\ &\quad - \alpha \phi^2 u^{\alpha-1} u_s |_{\gamma(0)} + H_{\partial M} u^\alpha \phi^2 |_{\gamma(0)}. \end{aligned}$$

Proof. Directly add the result in Lemma 2.4 over i , then we have

$$\begin{aligned} (n-1) \int_{\gamma} \phi_s^2 u^\alpha ds &\geq \int_{\gamma} \text{Ric}(\partial s, \partial s) u^\alpha \phi^2 ds + h_{\partial M}(e_i, e_i) u^\alpha \phi^2|_{\gamma(0)} \\ &+ \int_{\gamma} (\alpha^2 + \alpha) \phi^2 u^\alpha \sum_{i=1}^{n-1} (\nabla_{e_i} \log u)^2 - \alpha \phi^2 u^{\alpha-1} \sum_{i=1}^{n-1} (\nabla^2 u)(e_i, e_i) ds. \end{aligned}$$

Note that

$$\begin{aligned} \Delta u &= \sum_{i=1}^{n-1} (\nabla^2 u)(e_i, e_i) + (\nabla^2 u)(\partial s, \partial s) \\ &= \sum_{i=1}^{n-1} (\nabla^2 u)(e_i, e_i) + u_{ss} + \sum_{i=1}^{n-1} \kappa_i \nabla_{e_i} u \\ &= \sum_{i=1}^{n-1} (\nabla^2 u)(e_i, e_i) + u_{ss} - \alpha u \sum_{i=1}^{n-1} (\nabla_{e_i} \log u)^2. \end{aligned}$$

Simplify the inequality and do an integration by parts to u_{ss} . This leads to the inequality in the lemma. \square

As a corollary, we have

Proposition 2.6. *Let γ be a L_u^α -minimizing curve in the free boundary sense. Then for any compactly supported function ψ on γ , we obtain*

$$\begin{aligned} (2.4) \quad c_1(n, \alpha, \varepsilon) \int_{\gamma} \psi_s^2 ds &\geq c_2(n, \alpha, \varepsilon) \int_{\gamma} u^{-2} \psi^2 u_s^2 ds \\ &+ \int_{\gamma} \left(\text{Ric}(\partial s, \partial s) - \alpha \frac{\Delta u}{u} \right) \psi^2 ds + \alpha \int_{\gamma} \psi^2 |\nabla^\perp \log u|^2 ds \\ &+ \psi^2 \left(H_{\partial M} - \alpha \frac{u_s}{u} \right) \Big|_{\gamma(0)} \end{aligned}$$

where

$$c_1(n, \alpha, \varepsilon) = n - 1 + \frac{\alpha(n-3)}{4\varepsilon} > 0$$

and

$$c_2(n, \alpha, \varepsilon) = \alpha \left(1 - \frac{n-1}{4} \alpha - (n-3)\varepsilon \right)$$

which is positive if $\alpha < \frac{4}{n-1}$ and ε is small, or $n = 3$ and $\alpha \leq \frac{4}{n-1}$.

Proof. Plug $\phi = u^{-\alpha/2} \psi$ into Lemma 2.5 gives

$$\begin{aligned} (n-1) \int_{\gamma} \psi_s^2 ds &\geq \alpha \left(1 - \frac{n-1}{4} \alpha \right) \int_{\gamma} u^{-2} \psi^2 u_s^2 ds + \alpha(n-3) \int_{\gamma} u^{-1} \psi u_s \psi_s ds \\ &+ \int_{\gamma} \left(\text{Ric}(\partial s, \partial s) - \alpha \frac{\Delta u}{u} \right) \psi^2 ds + \alpha \int_{\gamma} \psi^2 |\nabla^\perp \log u|^2 ds \\ &+ \psi^2 \left(H_{\partial M} - \alpha \frac{u_s}{u} \right) \Big|_{\gamma(0)}. \end{aligned}$$

Using Cauchy-Schwarz inequality we have

$$\int_{\gamma} u^{-1} \psi u_s \psi_s ds \leq \varepsilon \int_{\gamma} u^{-2} \psi^2 u_s^2 ds + \frac{1}{4\varepsilon} \int_{\gamma} \psi_s^2 ds.$$

Plugging it into the above inequality completes the proof. \square

Remark 2.7. If we have information on the Neumann derivative of u along the boundary ∂M , the above boundary term can be expressed as

$$(n-1)\psi\psi_s - \frac{n+1}{2}\alpha u^{-1}u_s\psi^2 \Big|_{\gamma(0)}.$$

To make this term vanish, it suffices to choose ψ constant near $\gamma(0)$ and assume $u_s|_{\gamma(0)} = 0$. This condition does not depend on the mean convexity of the boundary ∂M .

2.2. Weighted minimizing line. In this part, we adapt some results from the previous subsection to curves entirely contained in $M \setminus \partial M$. Let $\gamma(s) : (-b, b) \rightarrow M$ be a curve on M parametrized by arc length s , and assume it is proper in the sense that $\gamma \cap \partial M = \emptyset$. We allow $b = +\infty$, in which case γ is a line.

Using calculations similar to those above, we obtain the following two results.

Lemma 2.8. *If γ_t is a smooth proper 1-parameter family of curves with $\gamma_0 = \gamma$, $\gamma_t(0) \in \partial M$ and $\gamma_t((-b, -a]) \cup \gamma_t([a, b]) = \gamma((-b, -a]) \cup \gamma([a, b])$. Denote the normal speed along the direction e_i at $t = 0$ by $\phi = \langle \frac{\partial \gamma_t}{\partial t} |_{t=0}, e_i \rangle$. Then*

$$(2.5) \quad \frac{dL_u^\alpha(\gamma_t)}{dt} \Big|_{t=0} = \int_{\gamma} \kappa_i u^\alpha \phi + \langle \nabla u^\alpha, e_i \rangle \phi ds.$$

In particular, the critical curve γ of L_u^α satisfies

$$(2.6) \quad \kappa_i = -\langle \nabla_{\partial_s} \partial_s, e_i \rangle = -\alpha \nabla_{e_i} \log u$$

along γ .

Proposition 2.9. *Let γ be a L_u^α -minimizing curve in the free boundary sense. Then for any compact supported function ψ on γ , we obtain*

$$(2.7) \quad c_1(n, \alpha, \varepsilon) \int_{\gamma} \psi_s^2 ds \geq c_2(n, \alpha, \varepsilon) \int_{\gamma} u^{-2} \psi^2 u_s^2 ds + \int_{\gamma} \left(\text{Ric}(\partial_s, \partial_s) - \alpha \frac{\Delta u}{u} \right) \psi^2 ds + \alpha \int_{\gamma} \psi^2 |\nabla^\perp \log u|^2 ds$$

where $c_1(n, \alpha, \varepsilon)$ and $c_2(n, \alpha, \varepsilon)$ are the same as the constants in Proposition 2.6.

2.3. Rigidity of L_u^α -minimizing curve in the free boundary sense.

When the function u and the curvatures of M satisfy certain conditions, we can derive a rigidity result. Specifically, we prove the following.

Proposition 2.10. *Let (M, g) be a Riemannian manifold with boundary. Suppose that γ is a complete L_u^α -minimizing curve in the free boundary sense in M . If*

$$\text{Ric} - \alpha u^{-1} \Delta u \geq 0$$

and

$$u_s|_{\partial M} \leq 0, \text{ or, } (H_{\partial M} - \alpha u^{-1} u_s)|_{\partial M} \geq 0.$$

Then $|\nabla u| = 0$ and $\text{Ric} - \alpha u^{-1} \Delta u = 0$ along γ .

Proof. The proof directly follows from Proposition 2.6 and Proposition 2.9. In fact, if γ is a complete ray, we let

$$\psi = \begin{cases} 1, & s \in [0, R] \\ \frac{2R-s}{R}, & s \in [R, 2R] \\ 0, & \text{otherwise.} \end{cases}$$

Plug it into (2.4), together with Remark 2.7, we obtain

$$\frac{c_1}{R} \geq c_2 \int_0^R u^{-2} u_s^2 ds + \int_0^R \text{Ric}(\partial s, \partial s) - \alpha \frac{\Delta u}{u} ds + \alpha \int_0^R |\nabla^\perp \log u|^2 ds.$$

By letting $R \rightarrow \infty$, we have that $u_s = 0$, $|\nabla^\perp u| = 0$ and $\text{Ric} - \alpha u^{-1} \Delta u = 0$ along γ . The other case if γ is a complete line, we have the same conclusion by plugging the following ψ into (2.7)

$$\psi = \begin{cases} 1, & s \in [-R, R] \\ \frac{2R-s}{R}, & |s| \in [R, 2R] \\ 0, & \text{otherwise.} \end{cases}$$

The proof is complete. \square

3. CONSTRUCTION OF WEIGHTED MINIMIZING RAY OR LINE

Suppose that M has nonnegative α -Ricci curvature in the spectral sense. Then, there exists a positive function $u \in C^{2,\alpha}(M)$ satisfying (1.1) and (1.2). Note that we do not assume the mean convexity of the boundary ∂M in this section.

The main theorem of this section is

Theorem 3.1. *If M contains a complete L_u^α -minimizing curve γ in the free boundary sense, then for any $p \in M$, there exists a complete L_u^α -minimizing curve in the free boundary sense, denoted by γ_p , that passes through p*

We first prove the following lemma.

Lemma 3.2. *Suppose γ is a L_u^α -minimizing curve in the free boundary sense in a compact set $K \subset M$. Then, the total length of γ is uniformly bounded, and the weighted length of γ is uniformly bounded by a constant C , where C depends on the diameter of K , α , and the infimum and supremum of u on K .*

Proof. We establish the following inequality:

$$\inf_K u^\alpha \int_\gamma d\gamma \leq \int_\gamma u^\alpha d\gamma \leq \int_{\tilde{\gamma}} u^\alpha d\tilde{\gamma} \leq \sup_K u^\alpha \int_{\tilde{\gamma}} d\tilde{\gamma} \leq \sup_K u^\alpha (\text{diam}(K) + \varepsilon),$$

where $\tilde{\gamma}$ is a curve connecting the endpoints of γ , and the total length of γ is bounded by $\text{diam}(K) + \varepsilon$. Taking $\varepsilon \rightarrow 0^+$, we obtain the desired result. \square

We will need the following lemma to construct a perturbed weighted function $u_{r,t}$ near an existing L_u^α -minimizing curve in the free boundary sense.

Lemma 3.3. *For any $p_0 \in M$, there exist positive constants $r_0 = r_0(n, M, p_0)$ and $t_0 = t_0(n, M, p_0)$ such that for any $r \leq r_0$ and any p in the closure of $B_{2r}(p_0)$, there exists a positive function $u_{r,t}$ on M for any $0 < t \leq t_0$ satisfying the following conditions:*

- (i) $u_{r,t} \rightarrow u$ as $r, t \rightarrow 0^+$ in the C^2 sense, and $u_{r,t} \rightarrow u$ smoothly for any fixed $r > 0$.
- (ii) $u_{r,t} = u$ on $M \setminus B_{3r}(p)$, and $u_{r,t} < u$ in $B_{3r}(p)$.
- (iii) $-\alpha \Delta u_{r,t} + \text{Ric} \cdot u_{r,t} > 0$ in $B_{3r}(p) \setminus B_r(p)$.
- (iv) $\frac{\partial u_{r,t}}{\partial \eta} \geq 0$ on ∂M .

Proof. This is a purely local construction. First, we consider the case $p_0 \in \partial M$. We choose the standard Fermi coordinate near p_0 such that $p_0 = 0$ and M is locally given by $\{x_1 \leq 0\}$ near 0, and the metric g_{ij} at 0 is δ_{ij} . In particular, we can write $\eta = \frac{\partial}{\partial x_1}$.

Given any $p \in B_{2r_0}(p_0)$ for r_0 small enough, we choose the function d_p by

$$d_p(\cdot) = \text{dist}_\delta(p, \cdot),$$

where dist_δ denotes the distance function with respect to the metric δ (which is the Euclidean metric using the Fermi coordinate). It is easy to see that

$$(3.1) \quad \frac{\partial d_p^2}{\partial \eta} \geq 0.$$

By choosing r_0 small enough (not depending on p), we know $g - \delta$ is sufficiently small in $B_{5r_0}(p_0)$ in the C^2 sense and hence

$$(3.2) \quad \frac{1}{2} \leq |\nabla d_p|_g \leq 2, \quad \Delta_g d_p^2 \leq 2n + \frac{1}{2} \text{ in } B_{5r_0}(p_0).$$

Note that this implies

$$d_p \Delta_g d_p \leq n.$$

Next, we choose a function $f(s)$ defined on \mathbb{R} such that

- (i) $f(s)$ is a negative constant on $(-\infty, \frac{1}{2}]$,

- (ii) $f(s) = -\exp(k/(s-3))$ on $[1, 3)$,
- (iii) $f(s) = 0$ when $s \geq 3$.

Then, for any $l > 0$ and $s \in [1, 3)$, we have

$$sf''(s) + lf'(s) = -\exp\left(\frac{k}{s-3}\right) \frac{k}{(s-3)^4} (ks + 2s(s-3) - l(s-3)^2).$$

If we choose $k > 4 + 4l$, then

$$(3.3) \quad sf''(s) + lf'(s) < 0 \quad \text{for } s \in [1, 3).$$

Note that $f'(s) > 0$ and $f''(s) < 0$ for $s \in [1, 3)$.

We consider

$$v_r = r^4 f\left(\frac{d_p}{r}\right).$$

Then, we know

$$v_r < 0 \text{ in } B_{3r}(p),$$

and define

$$u_{r,t} = (1 + tv_r)u.$$

In particular, for t_0 small enough, we know $u_{r,t} > 0$ for any $0 < r \leq r_0$ and $t \in (0, t_0]$. In $B_{3r}(p) \setminus B_r(p)$, we compute

$$\begin{aligned} \alpha \Delta u_{r,t} &= \alpha(1 + tv_r) \Delta u + \alpha t u \Delta v + 2\alpha t \nabla u \cdot \nabla v \\ &= \text{Ric} \cdot u_{r,t} + \alpha t (u \Delta v + 2 \nabla u \cdot \nabla v) \\ &\leq \text{Ric} \cdot u_{r,t} + \alpha t \left(ur^2 f''\left(\frac{d_p}{r}\right) |\nabla d_p|_g^2 + ur^3 f'\left(\frac{d_p}{r}\right) \Delta d_p \right. \\ &\quad \left. + 2 |\nabla u|_g r^3 f'\left(\frac{d_p}{r}\right) |\nabla d_p|_g \right) \\ &\leq \text{Ric} \cdot u_{r,t} + \frac{\alpha t u r^3}{d_p} \left(\frac{d_p}{4r} f''\left(\frac{d_p}{r}\right) + n f'\left(\frac{d_p}{r}\right) \right. \\ &\quad \left. + \frac{12 \|\nabla u\|_g \|L^\infty(B_{5r_0}(p_0))\} r_0}{\inf_{B_{5r_0}(p_0)} u} f'\left(\frac{d_p}{r}\right) \right). \end{aligned}$$

Now, we select $l = 4n + \frac{48 \|\nabla u\|_g \|L^\infty(B_{5r_0}(p_0))\} r_0}{\inf_{B_{5r_0}(p_0)} u}$ and $k = 4l + 4 + 1$. It follows from (3.3) that

$$\alpha \Delta u_{r,t} < \text{Ric} \cdot u_{r,t}$$

in $B_{3r}(p) \setminus B_r(p)$. Now, we compute

$$\frac{\partial u_{r,t}}{\partial \eta} = (1 + tv_r) \frac{\partial u}{\partial \eta} + tu \frac{\partial v_r}{\partial \eta} = t u r^3 f'\left(\frac{d_p}{r}\right) \frac{\partial d_p}{\partial \eta} \geq 0$$

by (3.1). It is easy to verify that other conditions are also satisfied by the construction of $u_{r,t}$.

If $p_0 \in M \setminus \partial M$, we can choose r_0 sufficiently small such that $5r_0 < \text{dist}_g(p_0, \partial M)$, and then we do not need to consider the boundary condition. The construction of $u_{r,t}$ is the same as above. \square

Theorem 3.4. *Let $\gamma : I \rightarrow M$ be a complete L_u^α -minimizing curve in the free boundary sense in M . Then, for any $s \in I$, there exists a positive constant $r_0 = r_0(s, \gamma) < \frac{1}{3}$ such that for any $p \in M \setminus \gamma$ with $\text{dist}_g(p, \gamma) = \text{dist}_g(p, \gamma(s)) = 2r$ and $0 < r \leq r_0$, we can construct another complete L_u^α -minimizing curve in the free boundary sense, denoted by γ_r , which passes through the closure of $B_r(p)$.*

Proof. First, we consider the case that $I = [0, +\infty)$. We fix any $s \in [0, +\infty)$ and let $p_0 = \gamma(s)$. Let r_0, t_0 be the positive constants given by Lemma 3.3 for p_0 . Let p be the point such that $\text{dist}_g(p, p_0) = \text{dist}_g(p, \gamma) = 2r$ and $u_{r,t}$ be the function constructed in Lemma 3.3 for p and $t \in (0, t_0)$.

We construct new smooth weighted functions $u_{r,t,h}$ for any $h > 0$ by requiring

- (i) $u_{r,t,h} = u_{r,t}$ in $\{q \in M : \text{dist}_g(\gamma([0, h + s + 1]), q) \leq 2h\}$,
- (ii) $u_{r,t,h} \geq u_{r,t}$ in $M \setminus \{q \in M : \text{dist}_g(\gamma([0, h + s + 1]), q) \leq 2h\}$,
- (iii) $u_{r,t,h} \geq \max\{1, u_{r,t}\}$ in $\{q \in M : \text{dist}_g(\gamma([0, h + s + 1]), q) \geq 2h + 1\}$.

Let $\gamma_{r,t,h}$ to be the curve from $\gamma(h + s + 1)$ to ∂M which minimizes the functional $L_{u_{r,t,h}}^\alpha$. Note that since M is complete and $u_{r,t,h} \geq 1$ outside a compact set, the curve $\gamma_{r,t,h}$ is well-defined and has finite length by Lemma 3.2. It is straightforward to verify that $\gamma_{r,t,h}$ is a $L_{u_{r,t,h}}^\alpha$ -minimizing curve in the free boundary sense.

We claim that $\gamma_{r,t,h}$ has a non-empty intersection with $B_{3r-\varepsilon}(p)$ for $\varepsilon > 0$ small enough, which does not depend on h . Note that by the construction of $u_{r,t}$, we have

$$u_{r,t}^\alpha \geq (1 - C\varepsilon)u^\alpha$$

in $B_{3r}(p) \setminus B_{3r-\varepsilon}(p)$ for some positive constant C .

If $\gamma_{r,t,h} \cap B_{3r-\varepsilon}(p) = \emptyset$, then we have

$$\begin{aligned}
(3.4) \quad & 0 < \int_0^{s+h+1} u^\alpha d\gamma - \int_0^{s+h+1} u_{r,t}^\alpha d\gamma \\
& = \int_0^{s+h+1} u^\alpha d\gamma - \int_0^{s+h+1} u_{r,t,h}^\alpha d\gamma \\
& \leq \int_0^{s+h+1} u^\alpha d\gamma - \int_{\gamma_{r,t,h}} u_{r,t,h}^\alpha d\gamma_{r,t,h} \\
& \leq \int_0^{s+h+1} u^\alpha d\gamma - \int_{\gamma_{r,t,h} \cap B_{3r}(p)} (1 - C\varepsilon)u^\alpha d\gamma_{r,t,h} - \int_{\gamma_{r,t,h} \setminus B_{3r}(p)} u^\alpha d\gamma_{r,t,h} \\
& = \int_0^{s+h+1} u^\alpha d\gamma - \int_{\gamma_{r,t,h}} u^\alpha d\gamma_{r,t,h} + C\varepsilon \int_{\gamma_{r,t,h} \cap B_{3r}(p)} u^\alpha d\gamma_{r,t,h} \\
& \leq \int_0^{s+h+1} u^\alpha d\gamma - \int_0^{s+h+1} u^\alpha d\gamma + C\varepsilon \int_{\gamma_{r,t,h} \cap B_{3r}(p)} u^\alpha d\gamma_{r,t,h}
\end{aligned}$$

$$= C\varepsilon \int_{\gamma_{r,t,h} \cap B_{3r}(p)} u^\alpha d\gamma_{r,t,h}.$$

By Lemma 3.2, we can choose ε small enough (independent of h) to obtain a contradiction, since the right-hand side of the first line is a constant that does not depend on h by the definition of $u_{r,t}$.

For any compact $K \subset M$, the total length of $\gamma_{r,t,h}$ is uniformly bounded by Lemma 3.2. The critical equation (2.3) also implies the geodesic curvature of $\gamma_{r,t,h}$ is uniformly bounded in K . Then, for fixed $r, t > 0$, we can choose a subsequence of $\gamma_{r,t,h}$ that converges smoothly to $\tilde{\gamma}_{r,t}$ as $h \rightarrow +\infty$. Note that $\tilde{\gamma}_{r,t}$ may have multiple connected components. Since $u_{r,t,h} \rightarrow u_{r,t}$ in any compact set, each component of $\tilde{\gamma}_{r,t}$ is a complete $L_{u_{r,t}}^\alpha$ -minimizing curve in the free boundary sense.

Moreover, $\tilde{\gamma}_{r,t}$ has non-empty intersection with the closure of $B_{3r-\varepsilon}(p)$. We define $\gamma_{r,t}$ to be the component of $\tilde{\gamma}_{r,t}$ which passes through the closure of $B_{3r-\varepsilon}(p)$.

Claim: $\gamma_{r,t} \cap B_r(p) \neq \emptyset$.

This follows from Proposition 2.10. Lemma 3.3 ensures that $u_{r,t}$ satisfies the assumption of Proposition 2.10. Thus $\text{Ric} - \alpha u_{r,t}^{-1} \Delta u_{r,t} = 0$ along γ , contradicting (iii) of Lemma 3.3.

Now, we may further choose $t \rightarrow 0^+$ and, up to a subsequence, have $\gamma_{r,t}$ converging smoothly and locally to $\tilde{\gamma}_r$. We choose γ_r to be the component of $\tilde{\gamma}_r$ that passes through the closure of $B_r(p)$. Then, γ_r is a complete L_u^α -minimizing curve in the free boundary sense and passes through the closure of $B_r(p)$. We remark that it is not known whether γ_r is a ray or line. See Figure 1.




FIGURE 1. The case where γ_r is a line

Now, we consider the case that $I = (-\infty, +\infty)$. For any $p_0 \in \gamma$, we assume $\gamma(0) = p_0$ after a reparametrization. The choice of $u_{r,t}$, $u_{r,t,h}$ is the same as above. The difference lies in the construction of $\gamma_{r,t,h}$. For a fixed

$h > 0$, we define the class of curves A_h as follows. A curve β belongs to A_h if β satisfies one of the following conditions:

- (i) β is a curve connecting $\gamma(-h-1)$ and $\gamma(h+1)$,
- (ii) β has two components β_1 and β_2 , where each β_i is a curve connecting $\gamma((-1)^i(h+1))$ and ∂M .

We define $\gamma_{r,t,h}$ to be the curve in A_h that minimizes $L_{u_{r,t,h}}^\alpha$. Such a curve exists because M is complete and $u_{r,t,h} \geq 1$ outside a compact set. The computation in (3.4) still holds, and we can choose ε small enough to ensure that $\gamma_{r,t,h}$ has non-empty intersection with $B_{3r-\varepsilon}(p)$. Then, we can repeat the argument from the case $I = [0, +\infty)$ to obtain a complete L_u^α -minimizing curve in the free boundary sense, denoted by γ_r , which passes through the closure of $B_r(p)$. \square

Now, we are ready to prove the main theorem stated at the beginning of this section.

Proof of theorem 3.1. We define the set $\Omega \subset M$ by

$$\Omega := \{p \in M : \text{there exists a complete } L_u^\alpha\text{-minimizing curve in the free boundary sense through } p\}.$$

It is easy to see that Ω is a non-empty closed set by the assumption of the theorem and the compactness of L_u^α -minimizing curves in the free boundary sense.

If $\Omega \neq M$, we pick any p_1 in $M \setminus \Omega$. Let $p_0 \in \Omega$ be the nearest point to p_1 , and let r_0 be the positive constant given by Proposition 3.4 for p_0 . We choose γ to be any L_u^α -minimizing curve in the free boundary sense through p_0 . Let β be a shortest curve connecting p_0 and p_1 , with $\beta(0) = p_0$ parametrized by arclength. It is easy to see that $B_{2r}(\beta(2r)) \cap \Omega = \emptyset$ for any $r > 0$ sufficiently small. It implies

$$\text{dist}_g(\beta(2r), \gamma) = \text{dist}_g(\beta(2r), p_0) = 2r.$$

Then, we can apply Theorem 3.4 for $r < r_0$ to conclude that $\overline{B_r(\beta(2r))} \cap \Omega \neq \emptyset$. This is a contradiction. Hence, $\Omega = M$. \square

4. SPLITTING THEOREM WITH INTERIOR END

We need the following proposition, which can be proved by a standard argument.

Proposition 4.1. *Let (M, g) be a smooth noncompact Riemannian manifold with boundary. If M has an interior end, then there exists an L_u^α -minimizing curve in the free boundary sense in M .*

Proof. This part is similar to the construction in the proof of Theorem 3.4. Fix a point $p \in \partial M$. Since M has an interior end by assumption, we can choose a sufficiently large R such that $M \setminus B_R(p)$ has at least one unbounded component, denoted it by E , that does not intersect ∂M .

Now, consider a family of exhaustions $B_{R+n}(p)$ of M for $n = 1, 2, \dots$. We also choose a perturbation of the weighted function u_n such that:

- $u_n = u$ in $B_{R+n}(p)$,
- $u_n \geq u$ on M ,
- $u_n > \max\{1, u\}$ in $M \setminus B_{R+n+1}(p)$.

Since $\partial B_{R+n}(p) \cap E$ is compact, there exists a curve γ_n connecting $\partial B_{R+n}(p) \cap E$ and ∂M that minimizes the weighted length $L_{u_n}^\alpha$. Note that $\gamma_n \cap B_R(p) \neq \emptyset$ for all n .

Taking $n \rightarrow \infty$ and passing to a subsequence, γ_n converges locally and smoothly to $\tilde{\gamma}$, which passes through the closure of $B_R(p)$. Each component of $\tilde{\gamma}$ is L_u^α -minimizing in the free boundary sense. We then select one of the connected components of $\tilde{\gamma}$ that passes through the closure of $B_R(p)$. This is the desired curve. \square

Remark 4.2. Note that if we choose $u \equiv 1$, then each γ_n is the shortest geodesic connecting $\partial B_{R+n}(p) \cap E$ and ∂M , and the limit γ is a minimizing geodesic ray starting from ∂M since M is complete.

Theorem 4.3. *Let (M, g) be a smooth noncompact Riemannian manifold with mean-convex boundary. Assume that the Ricci curvature of M is non-negative and M has an interior end. Then, M is isometric to $\Sigma \times \mathbb{R}_{\geq 0}$, where Σ is closed manifold with nonnegative Ricci curvature.*

If we replace the existence of an interior end with the assumption that the boundary of M is compact, this result is due to Kasue[Kas83] and Croke-Kleiner[CK92]. The proof essentially follows from Kasue[Kas83]. The interesting point is that the existence of an interior end forces the boundary of the manifold to be compact and the number of interior ends to be exactly one.

Proof. By choosing $u \equiv 1$, Proposition 4.1 provides us with a free boundary minimizing geodesic ray $\gamma : [0, \infty) \rightarrow M$ such that $\text{dist}(\gamma(t), \partial M) = t$ for any $t \geq 0$. Then result follows from [Sak17, Theorem 1.8]. \square

5. PROOF OF THE MAIN THEOREM

In this section, we provide a proof of the main theorem.

Proof of Theorem 1.1. First, we claim that if M has an interior end, it must split. By the assumption, let u satisfy

$$\begin{aligned} -\alpha \Delta u + \text{Ric} \cdot u &= 0 \text{ on } M, \\ \frac{\partial u}{\partial \eta} &= 0 \text{ on } \partial M. \end{aligned}$$

Recall the weighted length function

$$L_u^\alpha(\gamma) = \int_\gamma u^\alpha.$$

By Proposition 4.1 we can find a ray τ starting from a curve in ∂M that is L_u^α -minimizing in the free boundary sense. Moreover, by Lemma 2.2, τ is perpendicular to ∂M at $\tau(0)$.

With this ray as a starting point, by Theorem 3.1, for any point on M there exists a complete L_u^α minimizing curve passing through it. It follows from Proposition 2.10 that $|\nabla u| = 0$ on M . Thus, u is a positive constant, so $\text{Ric} \geq 0$ on M , i.e., the Ricci curvature is pointwise nonnegative. Hence, the claim follows from Theorem 4.3. \square

Proof of Corollary 1.2 and Corollary 1.3. The stability inequality is

$$\int_M |\nabla \varphi|^2 - (\text{Ric}(\nu, \nu) + |A|^2)\varphi^2 - \int_{\partial M} h_{\partial N}(\nu, \nu)\varphi^2 \geq 0.$$

Let $\{e_i\}_{i=1}^n$ be a local orthonormal frame of M . We assume that $\text{Ric} = \text{Ric}(e_1, e_1)$. By the Gauss equation, it implies

$$\begin{aligned} & \int_M |\nabla \varphi|^2 + \beta \text{Ric} \varphi^2 - \int_{\partial M} h_{\partial N}(\nu, \nu)\varphi^2 \\ & \geq \int_M (\text{Ric}(\nu, \nu) + \beta \sum_{i=1}^n R_{1i1i} + |A|^2 - \beta h_{11}H + \beta \sum_{i=1}^n h_{1i}^2)\varphi^2 \\ & \geq \int_M \left(\text{biRic}_\beta(e_1, \nu) + \frac{4\beta^2 - n + 1}{4\beta((\beta - 1)n + 1)} H^2 \right) \varphi^2 \end{aligned}$$

for $\beta < \frac{n}{n-1}$. In the above second inequality we have used the following inequality ([Den11, Lemma 2.1]): for any $a > \frac{n-1}{n}$,

$$a|A|^2 + h_{11}H - \sum_{i=1}^n h_{1i}^2 \geq \frac{4a^2 - n + 1}{4((a-1)n + 1)} H^2.$$

It follows from assumptions of the theorem that for any compactly supported function $\varphi \in C_c(M)$, it holds

$$\int_M |\nabla \varphi|^2 + \beta \text{Ric} \varphi^2 - \int_{\partial M} h_{\partial N}(\nu, \nu)\varphi^2 \geq 0.$$

Then there exists a positive function u on M satisfying

$$\begin{aligned} -\Delta u + \beta \text{Ric} \cdot u &= 0 \text{ on } M, \\ \frac{\partial u}{\partial \eta} &= h_{\partial N}(\nu, \nu)u \text{ on } \partial M. \end{aligned}$$

The boundary term of Proposition 2.6 now becomes by letting $\alpha = \frac{1}{\beta}$,

$$\phi^2 \left(H_{\partial M} + \frac{1}{\beta} h_{\partial N}(\nu, \nu) \right) \Big|_{\gamma(0)}.$$

Since M is free boundary hypersurface in $(N, \partial N)$, then the above term is nonnegative if we choose $\beta = 1$ and assume the weakly mean-convexity of ∂N . Then all the analysis in the proof of Theorem 1.1 follows if $n \leq 4$. That

is, if we assume the existence of an interior end, M is isometric to $\Sigma \times \mathbb{R}_{\geq 0}$ where Σ is a closed manifold.

Let r be the distance function to the boundary ∂M and let $\phi(r)$ be a Lipschitz function that is 1 when $r < R$, and is 0 when $r > 2R$. Moreover, $|\phi_r| \leq 1/R$. Then by the stability inequality

$$0 \leq \int_{r \leq R} \text{Ric}(\nu, \nu) + |A|^2 \leq \frac{1}{R^2} \int_{r \leq 2R} 1 \rightarrow 0, \text{ as } R \rightarrow \infty.$$

Thus $|A| \equiv 0$ and $\text{Ric}(\nu, \nu) = 0$. The proof of the Corollary is complete. \square

Remark 5.1. In the proof of Corollary 1.2 and Corollary 1.3, $n \leq 4$ can be relaxed to $n \leq 5$ if we assume weakly convexity of ∂M .

REFERENCES

- [APX24] Gioacchino Antonelli, Marco Pozzetta, and Kai Xu. A sharp spectral splitting theorem. *arXiv:2412.12707*, 2024.
- [AX24] Gioacchino Antonelli and Kai Xu. New spectral bishop-gromov and bonnetmyers theorems and applications to isoperimetry. *arXiv:2405.08918*, 2024.
- [CEM19] Otis Chodosh, Michael Eichmair, and Vlad Moraru. A splitting theorem for scalar curvature. *Comm. Pure Appl. Math.*, 72(6):1231–1242, 2019.
- [CG72] Jeff Cheeger and Detlef Gromoll. The splitting theorem for manifolds of non-negative Ricci curvature. *J. Differential Geometry*, 6:119–128, 1971/72.
- [CK92] Christopher B. Croke and Bruce Kleiner. A warped product splitting theorem. *Duke Math. J.*, 67(3):571–574, 1992.
- [CL23] Otis Chodosh and Chao Li. Stable anisotropic minimal hypersurfaces in \mathbb{R}^4 . *Forum Math. Pi*, 11(e3):1–22, 2023.
- [CLMS24] Otis Chodosh, Chao Li, Paul Minter, and Douglas Stryker. Stable minimal hypersurfaces in \mathbb{R}^5 . *arXiv: Differential Geometry*, 2024.
- [CLS24] Otis Chodosh, Chao Li, and Douglas Stryker. Complete stable minimal hypersurfaces in positively curved 4-manifolds. *to appear in J. Eur. Math. Soc.*, 2024.
- [CLZ24] Jianchun Chu, Man-Chun Lee, and Jintian Zhu. On kähler manifolds with non-negative mixed curvature. *arXiv:2408.14043*, 2024.
- [CMMR24] Giovanni Catino, Luciano Mari, Paolo Mastrolia, and Alberto Roncoroni. Criticality, splitting theorems under spectral ricci bounds and the topology of stable minimal hypersurfaces. *arXiv:2412.12631*, 2024.
- [Den11] Qintao Deng. Complete hypersurfaces with constant mean curvature and finite index in hyperbolic spaces. *Acta Math. Sci. Ser. B (Engl. Ed.)*, 31(1):353–360, 2011.
- [FCS80] Doris Fischer-Colbrie and Richard Schoen. The structure of complete stable minimal surfaces in 3-manifolds of nonnegative scalar curvature. *Comm. Pure Appl. Math.*, 33(2):199–211, 1980.
- [Hon24] Han Hong. CMC hypersurface with finite index in hyperbolic space \mathbb{H}^4 . *arXiv:2404.10276*, 2024.
- [HS23] Han Hong and Artur B. Saturnino. Capillary surfaces: stability, index and curvature estimates. *J. Reine Angew. Math.*, 803:233–265, 2023.
- [HW25] Han Hong and Gaoming Wang. A splitting theorem for 3-manifold with nonnegative scalar curvature and mean-convex boundary. *arXiv:2501.08677*, 2025.

- [Kas83] Atsushi Kasue. Ricci curvature, geodesics and some geometric properties of Riemannian manifolds with boundary. *J. Math. Soc. Japan*, 35(1):117–131, 1983.
- [Liu13] Gang Liu. 3-manifolds with nonnegative Ricci curvature. *Invent. Math.*, 193(2):367–375, 2013.
- [Maz24] Laurent Mazet. Stable minimal hypersurfaces in \mathbb{R}^6 . *arXiv:2405.14676*, 2024.
- [Sak17] Yohei Sakurai. Rigidity of manifolds with boundary under a lower Ricci curvature bound. *Osaka J. Math.*, 54(1):85–119, 2017.
- [SY96] Ying Shen and Rugang Ye. On stable minimal surfaces in manifolds of positive bi-Ricci curvatures. *Duke Math. J.*, 85(1):109–116, 1996.
- [Wu23] Yujie Wu. Free boundary stable minimal hypersurfaces in positively curved 4-manifolds. *arXiv:2308.08103*, 2023.

DEPARTMENT OF MATHEMATICS AND STATISTICS, BEIJING JIAOTONG UNIVERSITY,
BEIJING, CHINA, 100044

Email address: hanhong@bjtu.edu.cn

YAU MATHEMATICAL SCIENCES CENTER, TSINGHUA UNIVERSITY, 100084, BEIJING,
CHINA

Email address: gmwang@tsinghua.edu.cn