arXiv:2503.07054v1 [math.DG] 10 Mar 2025

A REMARK ON THE REACH AND UPPER BOUNDS ON SOME EXTRINSIC GEOMETRY INVARIANTS OF SUBMANIFOLDS

R. MIRZAIE

ABSTRACT. We consider a compact submanifold M of a Riemannian manifold N and we use the second variation formula as a tool to drive some geometric results on reach(M, N) the reach of M in N, including some useful relations between the extrinsic geometry of M in N and reach(M, N). Our results generalize some theorems previously proved for the special case where N is Euclidean space.

Key words: Riemannian manifold, Reach, Submanifold. MSC: : 53C70, 53C22, 53B25, 62C20, 68U05.

1. INTRODUCTION

The reach of a subset M of Euclidean space was introduced by Federer [9] as the largest number τ such that any point at distance less than τ from M has a unique nearest point on M (foot point on M). The reach appears under various names in the literature. It was called the condition number in topological data analysis. In computational geometry it is used under the name of local feature size. It is an important geometric invariant of subsets of Euclidean space which plays effective roles in the field of manifold learning and topological data analysis (see [4, 8, 13]) and has been widely used by the researches in geometric measure theory, integral geometry ([10, 15]), image processing arguments and biomathematics. If M is a submanifold of Euclidean space, it is proved that the reach quantifies the maximum ratio between the geodesic distance on M and Euclidean distance for pairs of points in M, and it is a measure of a manifolds departure from convexity. In fact, the reach is a global invariant incorporating informations about the second

1

R. Mirzaie, Department of Pure Mathematics, Faculty of Science, Imam Khomeini International University, Qazvin, Iran

r.mirzaei@sci.ikiu.ac.ir.

fundamental form of the embedding and the location of the first critical point of the distance from the submanifold. Thus, it incorporates the local and global topology and geometry of M. Positivity of the reach for subsets of Euclidean space is the minimal regularity assumption on sets. If M is a smooth compact submanifold of Euclidean space, it has a positive reach and if the reach is bounded, then strong topological and geometric restrictions on M will appear. For instance, it is proved that for compact submanifolds of \mathbb{R}^n with the bounded reach, the angle between tangent spaces at different points are bounded. There are many papers about the relations between the reach of M in N and the geometry and topology of M when N is a Euclidean space. We refer to [7] and [2] for some results about the relations between the reach and the geometry of M when N is Euclidean space or it is a sphere. Authors of [11] and [5] extended the reach from subsets of Euclidean space to the reach of subsets of Riemannian manifolds. Although some results about the reach obtained for the submanifolds of Euclidean space can directly be extended to the general case where the ambient space is a Riemannian manifold, but in many cases we need more calculations and differential geometric tools. In the present article, we consider a compact differentiable submanifold M of a Riemannian manifold N. In Theorem 3.2, we find an upper bound for the exterior part of the acceleration vector of the unit speed geodesics of M, which is a calculation of the impact of the reach on the exterior geometry of M. Then, we drive some more results about the geometry of M and as application we use our theorem in the special case where N has constant curvature.

2. Preliminaries

The main tool in our proofs is the second variation formula which measures the change in the arch length under the small displacements. We refer to [14] for definitions and further results.

In what follows, N is a smooth and complete Riemannian manifold and M is a connected compact submanifold of N. All maps are considered to be smooth.

Remark 2.1 (Second variation formula). Let $\sigma : [0,1] \to N$ be a geodesic segment and $\chi : [0,1] \times (-\delta,\delta) \to N$ be a smooth map such that $\chi(t,0) = \sigma(t)$. Then, χ is called a variation of σ . Denote by L(s) the length of the curve $\sigma_s : [0,1] \to N$, $t \to \chi(t,s)$ ($\sigma_0 = \sigma$) and let

 $V(t)=\frac{d}{ds}\chi(t,s)_{|s=0}=\chi_s(t,0), \ A(t)=\chi_{ss}(t,0).$ By the first and second variation formulas we have:

$$L'(0) = \frac{1}{L(0)} \int_0^1 \langle \sigma'(t), V'(t) \rangle dt$$

and

$$L''(0) = \frac{1}{L(0)} \int_0^1 \{\langle V'^{\perp}, V'^{\perp} \rangle - \langle R_{V\sigma'}V, \sigma' \rangle \} dt$$

+ $\frac{1}{L(0)} (\langle \sigma'(1), A(1) \rangle - \langle \sigma'(0), A(0) \rangle \rangle).$

Where, $(V')^{\perp}$ is the component of V' normal to σ' .

Similarly, if for all $s \in (-\delta, \delta)$, the curve σ_s is a geodesic in N, then

$$L'(s) = \frac{1}{L(s)} \int_0^1 \langle \sigma'_s(t), V'_s(t) \rangle dt$$
$$L''(s) = \frac{1}{L(s)} \int_0^1 \{\langle V'_s^\perp, V'_s^\perp \rangle - \langle R_{V_s \sigma'_s} V_s, \sigma'_s \rangle \} dt$$
$$+ \frac{1}{L(s)} (\langle \sigma'_s(1), A_s(1) \rangle - \langle \sigma'_s(0), A_s(0) \rangle).$$

where, $V_{s}(t) = \chi_{s}(t, s), A_{s}(t) = \chi_{ss}(t, s).$

Definition 2.2 (medial axes and reach).

Let M be a compact submanifold of a Riemannian manifold N. **1.** The medial axes of M in N which we denote by med(M, N) is the closure of the following subset of N.

$$A = \{x \in N : \exists p \neq p' \in M \text{ such that } d(x, M) = d(x, p) = d(x, p')\}$$

2. The reach of M in N which we denote by τ , is defined as follows:

$$\tau = \inf\{d(x, M) : x \in med(M, N)\}.$$

3. If $q \in N$, then a point $p \in M$, with d(q, M) = d(q, p) is called a foot point of q.

4. A point $q \in med(M, N)$ is called a reach assigning point of Med(M, N) if $d(q, M) = \tau$. A foot point of a reach assigning point of Med(M, N) is called a reach assigning point on M.

Remark 2.3. If N is compact then there is a reach assigning point on med(M, N). In fact, by definition of the reach, for each positive integer n, there is a point $q_n \in Med(M, N)$ such that $\tau \leq d(q_n, M) < \tau + \frac{1}{n}$. By the compactness of N, $\{q_n\}$ has a convergent subsequence which we denote it again by $\{q_n\}$. If $q = \lim_{n \to \infty} q_n$, then $d(q, M) = \tau$.

If $p \in M$, denote by $(T_p M)^{\perp}$ the subspace of all vectors normal to M at p. Consider the usual exponential map $exp_p : T_p N \to N$, and let $B_r(p) = exp_p(\{v \in (T_p M)^{\perp} : |v| \leq r\}).$

Remark 2.4. If $reach(M, N) = \tau$, then for all positive numbers $r < \tau$, the *r*-tube around M, defined by $\{x \in N : d(x, M) < r\}$ is an embedded submanifold of N (a consequence of the tubular neighborhood theorem). Consequently, the reach of M in N can also be defined as:

 $sup\{\eta > 0 : B_{\eta}(p) \cap B_{\eta}(p') = \emptyset, for all p, p' \in M with p \neq p'\}.$

3. Some inequalities describing the impact of the reach on the extrinsic geometry of submanifolds

It is proved in [7] that the reach gives an upper bound for the length of the acceleration of the arc length parametrized geodesics in the compact submanifolds of the Euclidean space. Also we refer to [1] for more similar calculations relating the reach and the extrinsic geometry of the submanifolds of Euclidean space. The theorems of the present section can be used as a tool to obtain some similar results for the compact submanifolds of a Riemannian manifold.

In what follows, we will denote by ∇ the Levi Civita connection of N. ∇ will denote to the induced connection on M. If $\alpha : I \to M$ is a differentiable curve in M and $t \in I$, we will denote by $\ddot{\alpha}(t)$ and $\alpha''(t)$ the accerelations of α in N and M, respectively. If v, w are linearly in depended vectors in T_pN , then $\kappa(v, w)$ will denote the sectional curvature of N at the point p along the plan generated by v, w.

If $p \in M$ and Π is the second fundamental form at p and η is a unit vector normal to M at p, we will denote by A_{η} the shape operator in direction of η , which is a linear map $A_{\eta}: T_pM \to T_pM$ defined by

$$< A_{\eta}(w), v > = < \Pi(v, w), \eta >, v, w \in T_p M.$$

We will use the following lemma.

Lemma 3.1. Let M be a compact submanifold of a Riemannian manifold $N, p \in M$ and q be a point in N with $d(q,p) = d(q,M) = \tau > 0$ and let $\sigma : [0,1] \to N$ be a geodesic with the length τ joining p to q and $\eta \in T_pN$ be the unit vector in direction of $\sigma'(0)$. If α is a unit speed geodesic in M with $\alpha(0) = p$, then there is a vector field V along σ such that

$$<\eta, \ddot{lpha}(0)> \leq \ rac{1}{ au}- au\int_0^1\kappa(V,\sigma')(1-t)^2dt.$$

Proof. Since $d(q, M) = \tau$, then for all points $x \in M$, $d(q, x) \geq \tau$ and $\sigma'(0)$ is perpendicular to M at p. Let $u = \alpha'(0)$ and denote by U(t) the vector field along σ obtained by the parallel transformation of u along σ . Put V(t) = (1 - t)U(t) and keeping the symbols used in Remark 2.1, consider a variation $\chi(t,s)$, $(t,s) \in [0,1] \times (-\epsilon,\epsilon)$ of σ such that $\chi_s(t,0) = V(t)$ and $\chi(0,s) = \alpha(s)$ (such a variation exists for sufficiently small ϵ). Since V(1) = 0, then $\chi(1,s)$ is a constant point which must be q and A(1) = 0. Also, we get from $A(t) = \chi_{ss}(t,s)|_{s=0}$ that

$$A(0) = \ddot{\alpha}(0) \quad (1)$$

Denote by L(s) the length of the curve $t \to \chi(t, s)$. Thus, L(0) is equal to the length of σ (which is equal to $\tau = |\sigma'(0)|$). Then, by the second variation formula we have

$$L''(0) = \frac{1}{\tau} \int_0^1 \{ \langle V'^{\perp}, V'^{\perp} \rangle - \langle R_{V\sigma'}V, \sigma' \rangle \} dt$$
$$-\frac{1}{\tau} \langle \sigma'(0), \ddot{\alpha}(0) \rangle \quad (2)$$

Since U is parallel along σ , then $\overline{\nabla}_{\sigma'} U = 0$. Thus,

$$V'(t) = \bar{\nabla}_{\sigma'} V = \bar{\nabla}_{\sigma'} (1-t)U(t) = -U(t) + (1-t)\bar{\nabla}_{\sigma'} U = -U(t) \quad (3)$$

Since U(0)(=u) is unit and normal to σ at $\sigma(0)(=p)$, U(t) is unit and normal to σ at $\sigma(t)$. Thus,

$$\langle V'^{\perp}, V'^{\perp} \rangle = \langle U, U \rangle = 1$$
 (4)

We have also

$$\langle R_{V\sigma'}V, \sigma' \rangle = \kappa(V, \sigma') \langle V, V \rangle \langle \sigma', \sigma' \rangle =$$

$$\kappa(V, \sigma') \langle (1-t)U, (1-t)U \rangle \langle \sigma', \sigma' \rangle = \kappa(V, \sigma')(1-t)^2 \tau^2 \quad (5)$$

R. Mirzaie

Since η is supposed to be a unit vector in direction of σ' , then $\eta = \frac{1}{\tau}\sigma'(0)$. Now, we get from (1)-(5) that

$$L''(0) = \frac{1}{\tau} - \tau \int_0^1 \kappa(V, \sigma') (1 - t)^2 dt - \langle \eta, \ddot{\alpha}(0) \rangle$$
 (6)

L(s) is minimum at s = 0, then $L''(0) \ge 0$ and we get the result from (6).

Theorem 3.2. If $reach(M, N) = \tau > 0$ and $p \in M$, then for each unit speed geodesic α in M and each unit vector η normal to M at $\alpha(0) = p$, there is a geodesic $\sigma : [0, 1] \to N$ and a vector field V along σ such that

$$\langle \ddot{\alpha}, \eta \rangle \leq \frac{1}{\tau} - \tau \int_0^1 \kappa(V, \sigma')(1-t)^2 dt.$$

Proof. Let n > 1 be a positive integer such that $\tau > \frac{1}{n}$ and put $\tau_n = \tau - \frac{1}{n}$. let $\sigma_n(t) = exp_p(t\tau_n\eta)$ and put $q_n = \sigma_n(1)$. The length of σ_n between $\sigma_n(0) = p$ and $\sigma_n(1) = q_n$ is equal to τ_n . Thus, $q_n \in B_{\tau_n}(p)$. We show that $d(q_n, M) = \tau_n$. Clearly, $d(q_n, M) \leq d(q_n, p) = \tau_n < \tau$. Since $d(q_n, M) < \tau$, then $q_n \notin med(M, N)$. Consequently, there is a unique point $p' \in M$ such that $d(q_n, M) = d(q_n, p')$. Thus, $q_n \in B_{\tau_n}(p')$, $r_n = d(q_n, M)$. Since $r_n \leq \tau_n$, then $B_{r_n}(p') \subset B_{\tau_n}(p')$. Thus, $q_n \in B_{\tau_n}(p) \cap B_{\tau_n}(p')$. Now, by Remark 2.4, we get that p = p'. Therefore,

$$d(q_n, M) = d(q_n, p) = \tau_n \quad (1)$$

The set $E = \{w \in T_p N : |w| \leq \tau\}$ is compact in $T_p N$. Since the sequence $\{q_n\}$ is a subset of the compact set $exp_p(E)$, it has a convergent subsequence which we denote it again by $\{q_n\}$. Let $q_n \to q$ and take limits in (1) to get $d(q, M) = d(q, p) = \tau$. It is clear that for all n, $\sigma'_n(0) = |\sigma'_n(0)|\eta$. Then, for the minimal geodesic $\sigma : [0, 1] \to N$, joining p to q, we have $\sigma'(0) = |\sigma'(0)|\eta$. Therefore, we get the result by Lemma 3.1.

Remark 3.3. In Theorem 3.2, If $M \alpha$ is a unit speed geodesic in M, then we have $\ddot{\alpha} = \bar{\nabla}_{\alpha'} \alpha' = \alpha'' + nor(\bar{\nabla}_{\alpha'} \alpha') = nor(\bar{\nabla}_{\alpha'} \alpha')$. Then, if Mis not totally geodesic in N, we can put $\eta = \frac{1}{|\vec{\alpha}(0)|} \ddot{\alpha}(0)$, which leads to

$$|\ddot{\alpha}(0)| \le \frac{1}{\tau} - \tau \int_0^1 \kappa(V, \sigma') (1-t)^2 dt.$$

We recall that if W is a vector space with an inner product, and $T: W \to W$ is a linear map, then the norm of T is defined by:

$$|T| = \sup\{|T(v)|: v \text{ is a unit vector in } W\}.$$

Keeping the symbols used in Theorem 3.2, since A_{η} is self adjoint, then by a little linear algebra computations we can show that

$$|A_{\eta}| = \sup_{|w|=1} | < A_{\eta}(w), w > |.$$

Since the set of all unit vectors of T_pM is compact, then there is a unit vector w in T_pM such that $|A_\eta| = | < A_\eta(w), w > |$. We can suppose $< A_\eta(w), w >$ is positive and $|A_\eta| = < A_\eta(w), w >$ (if not, use $-\eta$ instead of η). Let α be a geodesic in M with $\alpha'(0) = w$. Since $\Pi(\alpha', \alpha') = \ddot{\alpha}$, then $|A_\eta| = < A_\eta(\alpha'(0)), \alpha'(0) > = < \Pi(\alpha'(0), \alpha'(0)), \eta > = < \ddot{\alpha}(0), \eta >$. Now, by Theorem 3.2, we get the following corollary.

Corollary 3.4. Under the assumptions of Theorem 3.2,

$$|A_{\eta}| \le \frac{1}{\tau} - \tau \int_0^1 \kappa(V, \sigma') (1-t)^2 dt$$

If N has lower bounded curvature $\kappa \geq c$, then we have

$$\frac{1}{\tau} - \tau \int_0^1 \kappa(V, \sigma') (1-t)^2 dt \le \frac{1}{\tau} - \tau \int_0^1 c(1-t)^2 dt = \frac{1}{\tau} - \frac{c\tau}{3}.$$

If M is a compact submanifold of the Euclidean space with the positive reach and α is unit speed geodesic in M, then it is proved that $|\ddot{\alpha}| \leq \frac{1}{reach(M,R^n)}$ (see [7]). The following theorem is a generalization which easily comes from from Remark 3.3 and Corollary 3.4.

Theorem 3.5. Let N be a Riemannian manifold of curvature bounded from below to c, M be a compact submanifold of N with the reach $(M, N) = \tau > 0$ and $p \in M$. Then, for each unit speed geodesic α in M, with $\alpha(0) = p$ and each unit vector $\eta \in (T_p M)^{\perp}$ we have:

$$<\eta, \ddot{\alpha}(0)> \leq \frac{3-\tau^2 c}{3\tau}, \quad |\ddot{\alpha}(0)| \leq \frac{3-\tau^2 c}{3\tau}, \quad |A_{\eta}| \leq \frac{3-\tau^2 c}{3\tau}.$$

Note that in the previous theorem we need only the curvature of N be bounded along geodesics. This is a special case of a general category of Riemannian manifolds which are called manifolds with bounded radial curvature (see [12]). There are many useful topological and geometric results about manifolds with the bounded radial geodesics, which can be used in the subject of the present article.

4. Existence of the reach assigning points and the geometry of M

Existence of the reach assigning points imposes important geometric restrictions on the manifold. For example, it is proved that if $M \subset \mathbb{R}^n$ and there exists a reach assigning point on M, then either M has bottleneck or $|\ddot{\alpha}|$ is equal to $\frac{1}{reach(M,\mathbb{R}^n)}$ at some points of a unit speed geodesic α of M (see [1]). In the present section we show that similar results are true when the ambient space is an arbitrary Riemannian manifold. Note that in the terminology used in the texts of topological data analysis or computational geometry a point in med(M, N) with more than

Lemma 4.1. Let N be a Riemannian manifold and M be a compact submanifold of N with reach $(M, N) = \tau > 0$. If there is a reach assigning point q in med(M, N) with at least two foot points in M, then there is a unit speed geodesic α in M and a point $p = \alpha(s_0)$ on α such that

one foot point in N is called a bottleneck point.

$$<\ddot{\alpha}(s_0), \sigma'(0)>= 1-L^2 \int_0^1 \kappa(V, \sigma')(1-t)^2 dt.$$

Where, $\sigma : [0,1] \to N$ is a geodesic in N joining $\alpha(s_0)$ to q, V(t) = (1-t)U(t) such that U is a vector field along σ obtained by the parallel translation of $\alpha'(s_0)$ along σ , and $L = d(q, \alpha(s_0))$.

Proof. Let p and p' be different points in M such that $d(q, M) = d(q, p) = d(q, p') = \tau$, and let $\alpha : [0, b] \to M$, $b = d_M(p, p')$, be the unit speed minimizing geodesic in M joining p to p'. Extend α to a geodesic $\alpha : (-\epsilon, b + \epsilon) \to M$ (at least for a small number $\epsilon > 0$) and put $L(s) = d(q, \alpha(s))$. L is minimum at s = 0 and s = b ($L(0) = L(b) = \tau$). Consequently,

$$L''(0) \ge 0, \quad L''(b) \ge 0.$$

In other way, since L(0) = L(b) are minimum points for L, then L will be maximum at a point $d \in [0, b]$, which implies $L''(d) \ge 0$. Then, we

get from $L''(0) \leq 0$ and $L''(d) \geq 0$ that $L''(s_0) = 0$, for some number $s_0 \in [0, d]$. By the arguments similar to the proof of Lemma 3.1, we get that

$$L''(s_0) = \frac{1}{L} - \langle \ddot{\alpha}(s_0), \eta \rangle - L \int_0^1 \kappa(V, \sigma') (1-t)^2 \quad (1)$$

Where, $\sigma : [0,1] \to N$ is a geodesic with the length $L = d(\alpha((s_0),q))$ joining $\alpha(s_0)$ to q, $\eta = \frac{\sigma'(0)}{|\sigma'(0)|}$, V = (1-t)U(t) such that U(t) is the parallel translation of $\alpha'(s_0)$ along σ . Since $|\sigma'(0)| = L$ and $L''(s_0) = 0$, then we get from (1) that

$$\langle \ddot{\alpha}(s_0), \sigma'(0) \rangle = 1 - L^2 \int_0^1 \kappa(V, \sigma') (1-t)^2 dt.$$

Remark 4.2. Keeping the symbols used in Definition 2.2, suppose that there is a reach assigning point q in med(M, N) with only one foot point in M. Then, q belongs to the closure of A but $q \notin A$. Thus, there is a sequence $\{q_n\}$ in A such that $q_n \to q$. There are different points p_n, p'_n in M such that

$$d(q_n, p_n) = d(q_n, p'_n) = d(q_n, M)$$
 (1)

Since M is compact, we can consider convergent subsequences of $\{p_n\}$ and $\{p'_n\}$. Then, without lose of generality, we can suppose that

$$p_n \to p, p'_n \to p', p, p' \in M.$$

Taking limits $(n \to \infty)$ in (1), we get that $d(q, p) = d(q, p') = d(q, M) = \tau$. Since by the assumption, q has only one base point in M, we have p = p'. Consider a neghibourhood W in M for p and a number $\delta > 0$, such that all unit speed geodesics of M with a point in W can be defined in $[0, \delta]$ (such a neghibourhood exist for sufficiently small number $\delta > 0$). Put $d_n = d_M(p_n, p'_n)$ and let $\alpha_n : [0, d_n] \to M$ be the unit speed geodesic in M joining p_n to p'_n . If n is sufficiently large, then $p_n, p'_n \in W$ and $d_n < \delta$. Then, we can extend $\alpha_n : [0, d_n] \to M$ to the unite speed geodesic $\alpha_n : [0, \delta] \to M$. Extend $\alpha_n : [0, \delta] \to M$ to a geodesic $\alpha_n : (-\epsilon, \delta + \epsilon)$ for a small number $\epsilon > 0$ and put $L_n(s) = d(q_n, \alpha_n(s))$. By the arguments similar to the proof of Lemma 4.1, we get that there is a point $s_n \in [0, d_n]$ such that

$$<\ddot{\alpha_n}(s_n), \sigma'_n(0)>= 1 - L_n^2 \int_0^1 \kappa(V_n, \sigma'_n)(1-t)^2 dt$$
 (2).

Where, $\sigma_n : [0,1] \to N$ is a geodesic joining $\alpha_n(s_n)$ to q_n , L_n is equal to the length of σ_n , $V_n(t) = (1-t)U_n(t)$ and U_n is the vector field along σ_n obtained by the parallel transformation of $\alpha'_n(s_n)$ along σ_n .

Since $d_n \to 0$, then $s_n \to 0$. Consequently, $L_n \to d(q, p) = \tau$. From the compactness of M and the fact that the geodesics α_n are unit speed, we get that the sequence $\{(\alpha_n(s_n), \alpha'(s_n))\}$ has convergent subsequence in TM, which allows us to suppose that $(\alpha_n(s_n), \alpha'_n(s_n)) \to (p, Z), Z \in T_p M$, and Z must be a unit vector. Let $\alpha : (-\epsilon, \delta + \epsilon) \to M$ be the unit speed geodesic in M with the initial conditions $\alpha(0) = p, \alpha'(0) = Z$. Let $\sigma : [0,1] \to N$ be the minimizing geodesic in N joining $\alpha(0)(=p)$ to q, and let U(t) be the vector field along σ obtained by the parallel transformation of $\alpha'(0)$ along σ and put V(t) = (1-t)U(t). It is easy to show that

$$<\ddot{\alpha_n}(s_n), \sigma'_n(0) > \rightarrow <\ddot{\alpha}(0), \sigma'(0) >$$

and

$$\kappa(V_n(t), \sigma'_n(t)) \to \kappa(V(t), \sigma'(t)).$$

Taking limits in (2) leads to

$$<\ddot{\alpha}(0),\sigma'(0)>=1-\tau^2\int_0^1\kappa(V,\sigma')(1-t)^2dt.$$

Note that since the length of σ is equal to d(q, M), then $\sigma'(0)$ is perpendicular to M at p. Thus,

Now, putting together Lemma 4.1 and Remark 4.2, yields to the following theorem.

Theorem 4.3. Let N be a Riemannian manifold and M be a compact submanifold of N with reach $(M, N) = \tau > 0$. If there is a reach assigning point q in med(M, N), then there is a unit speed geodesic α in M and a point $p = \alpha(0)$ on α , a geodesic $\sigma : [0, 1] \rightarrow N$ in N joining p to q and a vector field V along σ such that:

$$<\ddot{\alpha}(0), \sigma'(0)>= 1 - L^2 \int_0^1 \kappa(V, \sigma')(1-t)^2 dt, \quad L = d(p,q).$$

If q has only one foot point in M, then

$$<\ddot{\alpha}(0),\sigma'(0)>=1-\tau^2\int_0^1\kappa(V,\sigma')(1-t)^2dt.$$

If in the previous theorem, N has constant curvature c and q has two foot points, then $\langle \ddot{\alpha}(0), \sigma'(0) \rangle = 1 - L^2 \frac{c}{3}$, L = d(p,q). Consider the

unit speed geodesic $\beta(t) = \sigma(L^{-1}t)$. We have $\beta'(0) = \frac{1}{L}\sigma'(0)$. Thus, $\langle \ddot{\alpha}(0), \beta'(0) \rangle = \frac{1}{L} - \frac{cL}{3}$, L = d(p,q). Similarly, If q has a unique foot point in M, $\langle \ddot{\alpha}(0), \beta'(0) \rangle = \frac{1}{\tau} - \frac{c\tau}{3}$. Since β is of unit speed, then

$$\langle \ddot{\alpha}(0), \beta'(0) \rangle \leq |\alpha(0)|$$

But, by Theorem 3.5 we have $|\ddot{\alpha(0)}| \leq \frac{1}{\tau} - \frac{c\tau}{3}$. Thus, $|\ddot{\alpha(0)}| = \frac{1}{\tau} - \frac{c\tau}{3}$. Putting c = 0 yields to $|\ddot{\alpha}(0)| = \frac{1}{\tau}$, which is the criterion previously proved for the compact submanifolds of Euclidean space (see [1]).

5. Comparing the extrinsic and intrinsic geometries of $M_{\rm ~~IN}$ N

The reach of M in N is a useful tool for comparing the intrinsic and extrinsic geometries of M in N. For instance, the authors of [7] find a bound for the angle between $\gamma'(0)$ and $\gamma'(1)$ for an arc length parametrized geodesic γ in $M \subset \mathbb{R}^d$. Also they give a bound for the angle between T_pM and T_qM related to $d_N(p,q)$ and the reach. In a more general case, we compare the parallel translations of a vector with respect to the geometries of M and N, when N has curvature bounded from below.

Remark 5.1. Suppose that $\tau = reac(M, N) > 0$ and N has curvature bounded from below to c. Let $v_0 \in T_pM$ be a unit vector and consider a point $q \in M$ and let $\alpha : [0,1] \to M$ be a path joining p to q (for simplicity we suppose that α is a unit speed geodesic in M). Denote by $v^M(t)$ and $v^N(t)$ the parallel translations of v_0 along α to the point $\alpha(t)$, in M and N respectively. We compute the following number, which compares the parallel transformations in M and N.

$$D = \langle v^N(1), v^M(1) \rangle - \langle v_0, v_0 \rangle$$
.

Put

$$f(t) = \langle v^M(t), v^N(t) \rangle - \langle v_0, v_0 \rangle, \ t \in [0, 1].$$

We have

$$f'(t) = \frac{d}{dt} < v^M, v^N > = <\bar{\nabla}_{\alpha'}(v^M), v^N > + < v^M, \bar{\nabla}_{\alpha'}(v^N) > .$$
(1)

Since v^N is parallel in N and v^M is parallel in M then $\bar{\nabla}_{\alpha'}(v^N) = 0$ and

$$\bar{\nabla}_{\alpha'}(v^M) = \nabla_{\alpha'}(v^M) + \Pi(\alpha', v^M) = \Pi(\alpha', v^M).$$

R. Mirzaie

Thus, we get from (1) that

$$f'(t) = <\Pi(\alpha', v^M), v^N > \qquad (2)$$

Let $(v^N)^{\perp}$ be the normal (to M) component of v^N and put $\eta = \frac{(v^N)^{\perp}}{|(v^N)^{\perp}|}$ (we consider the typical case where M is not totally geodesic in N and $|(v^N)^{\perp}| \neq 0$). Then,

$$f'(t) = <\Pi(\alpha', v^M), (v^N)^{\perp} > = |(v^N)^{\perp}| < \Pi(\alpha', v^M), \eta >$$

= $|(v^N)^{\perp}| < v^M, A_{\eta}(\alpha') >$ (3)

Thus,

$$|f'(t)| \le |(v^N)^{\perp}| |v^M| |A_{\eta}| |(\alpha')| \quad (4)$$

Since v^M , v^N and α' are unit vectors, then $|f'(t)| \leq |A_\eta|$, which by Theorem 3.5, yields to $|f'(t)| \leq \frac{3-\tau c}{3\tau}$. By mean value theorem, we have f(1) = f(0) + f'(t) for some t. Since f(0) = 0, then

$$|D| = |f(1)| \le \frac{3 - \tau c}{3\tau}.$$

This means that when the reach is positive, the intrinsic geometry of M is not very far away from the extrinsic geometry of M in N.

References

- E. Aamari, J. Kim, F. chazal, B. Michel, A. Rinaldo, and L. Wasserman, *Estimating the Reach of a Manifold*, Electron. J. Stat., **13(1)**, (2019) 1359-1399.
- [2] R. J. Adler, S. R. Krishnan, J. E. Taylor, S. Weinberger, Convergence of the reach for a sequence of Gaussian-embedded manifolds, Probab. Theory Relat. Fields, 71, (2018) 1045-1091.
- [3] N. Amenta, M. W. Bern, Surface reconstruction by Voronoi filtering, Discrete Comput. Geom., 22(4), (1999) 481-504.
- [4] D. Attali, A. Lieutier, Geometry driven collapses for converting a Cech complex into a triangulation of a ` nicely triangulable shape, Discrete Comput. Geom., 54(4), (2015) 798-825.
- [5] V. Bangert, Sets with positive reach, Archiv der Mathematik, 38(1), (1982) 54-57.
- [6] M. Belkin, J. Sun, Y. Wang, Constructing Laplace operator from point clouds in R^d, Proceedings of the Twentieth Annual ACM-SIAM Symposium on Discrete Algorithms, SIAM, (2009) 1031-1040.
- [7] J. D. Boissonnat, A. Lieutier, M. Wintraecken, The reach, metric distortion, geodesic convexity and the variation of tangent spaces, Journal of Applied and Computational Topology, 3, (2019) 29-58.

12

- [8] T. Dey, J. Giesen, E. Ramos, B. Sadri, Critical points of distance to an εsampling of a surface and flow-complex-based surface reconstruction, Int. J. Comput. Geom. Appl., 18, (2008) 29-61.
- [9] H. Federer, Curvature measures, Trans. Amer. Math. Soc., 93, (1959) 418-491.
- [10] H. Federer, Geometric measure theory, Series Classics in Mathematics, Springer-Verlag New York, (1969).
- [11] N. Kleinjohann, Convexity and the unique footpoint property in Riemannian geometry, Archiv der Mathematik, 35(1), (1980) 574-582.
- [12] K. Kondo, Topology of complete manifolds with radial curvature bounded from below, Geometric and Functional analysis, 17, (2007) 1237-1247.
- [13] P. Niyogi, S. Smale, S. Weinberger, Finding the homology of submanifolds with high confidence from random samples, Discrete Comput. Geom., 39(1-3), (2008) 419-441.
- [14] B. O'Neill, Semi-Riemannian geometry with applications to Relativity, Academic Press, New York, Berkeley, (1983).
- [15] C. Thale, 50 years sets with positive reach a survey, Surv. Math. Appl., 3, (2008) 123-165.