Magic dynamics in many-body localized systems

Pedro R. Nicácio Falcão,^{1,2,*} Piotr Sierant,³ Jakub Zakrzewski,^{2,4} and Emanuele Tirrito^{5,6,†}

¹Szkoła Doktorska Nauk Ścisłych i Przyrodniczych,

Uniwersytet Jagielloński, Łojasiewicza 11, PL-30-348 Kraków, Poland

²Instytut Fizyki Teoretycznej, Wydział Fizyki, Astronomii i Informatyki Stosowanej,

Uniwersytet Jagielloński, Łojasiewicza 11, PL-30-348 Kraków, Poland

³Barcelona Supercomputing Center, Barcelona 08034, Spain

⁴Mark Kac Complex Systems Research Center, Uniwersytet Jagielloński, Kraków, Poland

⁵ The Abdus Salam International Centre for Theoretical Physics (ICTP), Strada Costiera 11, 34151 Trieste, Italy

⁶Dipartimento di Fisica "E. Pancini", Università di Napoli "Federico II", Monte S. Angelo, 80126 Napoli, Italy

(Dated: March 11, 2025)

Nonstabilizerness, also known as quantum magic, characterizes the beyond-Clifford operations needed to prepare a quantum state and constitutes an essential resource, alongside entanglement, for achieving quantum advantage. This work investigates how nonstabilizerness spreads under the dynamics of disordered quantum many-body systems. Using the ℓ -bit model, a phenomenological model of many-body localization (MBL), we present an analytical description of the nonstabilizerness growth in MBL systems. We demonstrate that our analytical formulas describe the nonstabilizerness growth in strongly disordered quantum spin chains. Our findings establish a new facet of MBL phenomenology and identify the vital role of the disorder in slowing down the growth of the complexity of quantum states, important for our understanding of quantum advantage.

Quantum state $|\psi\rangle$ of L qubits is spec-Introduction. ified by a state vector in 2^L dimensional Hilbert space [1]. The *exponential* growth of many-body Hilbert space implies that quantum states may become intractable for classical computers for sufficiently large L [2], motivating the development of quantum simulators [3-5] and quantum computers [6-9]. However, certain quantum states possess a structure that enables their efficient representation on classical computers. For instance, when $|\psi\rangle$ is weakly entangled [10], it can be simulated at cost increasing *polynomially* with L using tensor network approaches [11–16]. Hence, extensive entanglement [17, 18] is necessary for quantum devices to reach computational advantage over classical computers [19]. Nevertheless, stabilizer states [20, 21], may host an extensive entanglement, and still be simulated with classical resources scaling *polynomially* with L [22, 23]. Therefore, nonstabilizerness, commonly referred to as "magic", quantifying the extent to which $|\psi\rangle$ departs from the set of stabilizer states [24-29], is a quantum resource [30] essential for characterizing the complexity of quantum states.

Understanding the generation of magic resources in many-body systems is fundamental for assessing their classical simulability. Generically, quantum many-body systems prepared in an out-of-equilibrium state are expected to follow eigenstate thermalization hypothesis (**ETH**) [31–34] and to thermalize reaching an equilibrium state described by appropriate ensembles of statistical mechanics [35–38]. Thermalization is accompanied by fast, ballistic [39, 40] or sub-ballistic [41, 42], growth of entanglement entropy, and a rapid saturation of nonstabilizerness measures [43, 44] to their maximal values [45]. The process of thermalization slows down in the presence of disorder [46–49]. Sufficiently strong disorder leads to a phenomenon of the many-body localization (**MBL**) [50– 53] which prevents the thermalization [54–68] at any experimentally relevant time scale [69, 70]. The absence of thermalization starkly affects the dynamics of MBL systems, leading to a logarithmic in time growth of entanglement entropy [71–75] and the memory of the initial state due to the presence of an emergent set of local integrals of motion, dubbed localized bits or ℓ -bits [56, 76]. This raises the question: how does magic spread in MBL systems?

In this work, we address this question by investigating the dynamics of nonstabilizerness in strongly disor-



Figure 1. Magic spreading in disordered quantum systems. In the absence of interactions (III), \mathcal{M}_2 is described solely by spin precession and saturates to $L \log_2(8/7)$. In the interacting case, slow spin dephasing leads to a power-law growth of \mathcal{M}_2 . For the initial X-polarized state, magic grows as described by Eq. 5, c.f. (I), and saturates on a timescale much shorter than that required for the entanglement entropy to saturate. For a generic random product state, (II), the growth of nonstabilizerness is limited.

dered spin chains that exhibit MBL. Leveraging the ℓ bits, we develop a theoretical framework that describes how magic evolves under MBL dynamics. Specifically, we analyze the stabilizer Rényi entropy (SRE) [28] as a measure of magic, characterizing its growth in both non-interacting and interacting disorder localized sys-In non-interacting localized systems, the SRE tems. remains limited, while in MBL systems, the slow spin dephasing leads to a power-law growth of nonstabilizerness before its eventual saturation. To corroborate our theoretical predictions, we perform numerical simulations on a microscopic MBL model, the disordered transverse-field Ising model (TFIM). We track the evolution of magic across different disorder strengths, initial states, and system sizes, showing remarkable agreement with the ℓ -bits-based predictions. Additionally, we demonstrate that nonstabilizerness enables a clear distinction between MBL and ergodic regimes.

Quantifying nonstabilizerness. As a measure of nonstabilizerness, we consider stabilizer Rényi entropy (SRE) [28], which quantifies the spread of a state in the basis of Pauli string operators, and is defined as

$$\mathcal{M}_k(|\Psi\rangle) = \frac{1}{1-k} \log_2\left[\sum_{P \in \mathcal{P}_L} \frac{\langle \Psi|P|\Psi\rangle^{2k}}{D}\right], \quad (1)$$

where L is the number of qubits, k is the Rényi index, and P is a Pauli string that belongs to the Pauli group \mathcal{P}_L . In particular, \mathcal{M}_1 is defined by the limit $k \to 1$ in (1), and $\mathcal{M}_k \geq 0$, with the equality holding if and only if $|\Psi\rangle$ is a stabilizer state [77, 78]. In our study, we fix k = 2 and evaluate the SRE using the algorithm of [79], which allowed us to obtain numerically exact results for $L \leq 22$. One advantage of the SRE over many other proposed measures of magic [26] is that it allows an efficient computation even for a large L [80–85]. Moreover, $\mathcal{M}_k(|\Psi\rangle)$ is also experimentally measurable [85–89].

Quantifying nonstabilizerness and understanding how these resources grow is a current topic of interest in quantum many-body physics [27]. Recent works have addressed this question for ergodic many-body systems [44, 45], where initial state information is rapidly lost and $|\Psi\rangle$ behaves similarly to a random vector [34]. For example, in random unitary circuits, the SRE saturates to the Haar-random state value [90]

$$\mathcal{M}_2^{\text{Haar}} = \log_2(D+3) - 2,$$
 (2)

where $D = 2^{L}$ is the Hilbert space dimension, at times scaling *logarithmically* with system size L [43]. Generic many-body systems exhibit more intricate behavior. Floquet systems behave similarly to random circuits, while for Hamiltonian dynamics, the time required to approach the Haar value scales linearly in L, and the SRE may not reach the Haar value [44, 91]. The ℓ -bit model. A characteristic hallmark of MBL is the emergent integrability that microscopic models acquire at sufficiently strong disorder [56, 76, 92, 93]. In the MBL regime, the system is described by a set of ℓ -bits, $\hat{\tau}_i^z$, and its Hamiltonian reads

$$\hat{\mathcal{H}}_{\ell-\text{bit}} = \sum_{i} h_i \hat{\tau}_i^z + \sum_{i < j} J_{ij} \hat{\tau}_i^z \hat{\tau}_j^z + \sum_{i < j < k} J_{ijk} \hat{\tau}_i^z \hat{\tau}_j^z \hat{\tau}_k^z + \dots$$
(3)

where h_i are random on-site fields, $J_{ij...}$ are interaction terms that decay exponentially with the distance between the spins, and $\hat{\tau}^z$ are quasilocal operators that mutually commute [94]. The ℓ -bit model, (3), captures many phenomenological properties of MBL, including eigenvalue statistics [95, 96], entanglement [97–99], and other aspects of the dynamics [100–102]. In the following, we utilize (3) to understand nonstabilizerness dynamics in an MBL system.

We start by analyzing how the SRE grows when different terms are included in (3) for an X-polarized initial state $|\Psi_X^+\rangle = \bigotimes_{k=1}^L (|\downarrow\rangle + |\uparrow\rangle)/\sqrt{2}$. We first focus on the case where the ℓ -bits do not interact, $J_{ij...} = 0$. In this case, (3) describes an Anderson insulator, and the dynamics of the Pauli strings are governed solely by the spin precession. The SRE exhibits rapid initial growth at $t \sim 1$ and saturates fast to a nearly constant value (after averaging over disorder realizations). This dynamics can be accurately captured by decomposing the Pauli strings as the product of individual single-spin observables, yielding [94]

$$\mathcal{M}_2 = -\frac{L}{W} \int_0^W dh \log_2 \left[1 - \frac{1}{4} \sin^2(4ht) \right] \qquad (4)$$

for the initial $|\Psi_X^+\rangle$ state. In the limit of $t \to \infty$, this expression yields $\mathcal{M}_2 \approx L \log_2(8/7)$, see Fig. 1. For a generic product initial state $|\Psi_R\rangle$, our numerical results show that the SRE rapidly saturates to a constant value not bigger than $\mathcal{M}_2(|\Psi\rangle) + L \log_2(8/7)$, see [94].

The behavior of SRE significantly changes when the ℓ bits interact and $J_{ij...} \neq 0$. Similar to the non-interacting case, the spin precession terms induce a fast growth of the SRE at $t \sim 1$. Subsequently, in the presence of interactions, \mathcal{M}_2 continues to grow until it reaches a saturation value after sufficiently long times.

To understand the dynamics of SRE, it is essential to examine the impact of the spin dephasing terms on the time evolution of Pauli strings expectation values [103]. The expectation values of Pauli strings composed solely of $\hat{\tau}^z$ operators remain constant under the dynamics of $\hat{\mathcal{H}}_{\ell-\text{bit}}$. On the other hand, Pauli strings containing $\hat{\tau}^x$ or $\hat{\tau}^y$ exhibit distinctly different behavior. The slow dephasing of the spins causes a power-law decay of the single-spin expectation values $|\hat{\tau}_j^{\alpha}|$, with $\alpha \in \{x, y\}$; for multi-spin observables, the situation depends on whether the spins corresponding to $\hat{\tau}^{\alpha}$ operators are entangled.



Figure 2. Dynamics of nonstabilizerness in disordered TFIM. Evolution of SRE for initial states (a) $|\Psi_Z\rangle$, (b) $|\Psi_X\rangle$, (c) $|\Psi_Y\rangle$, and (d) $|\Psi_R\rangle$ (see text). The results are for L = 16 and averaged over 1000 realizations, considering the product state close to the middle of the spectrum. To demonstrate the validity of (5), we performed a numerical fit at W = 5 for all states. The saturation value $\mathcal{M}_2^{\text{sat}}$ depends on the initial state: $\mathcal{M}_2^{\text{sat}} \approx 7.3$ for $|\Psi_Z\rangle$, $\mathcal{M}_2^{\text{sat}} \approx 11.53$ for $|\Psi_R\rangle$, while the SRE for $|\Psi_X\rangle$ and $|\Psi_Y\rangle$ states saturates to the Haar value $\mathcal{M}_2^{\text{Haar}}$. The power-law growth exponents are $\beta \approx 0.16$ for $|\Psi_Z\rangle$, $\beta \approx 0.29$ for $|\Psi_X\rangle$, $\beta \approx 0.39$ for $|\Psi_Y\rangle$, and $\beta \approx 0.19$ for $|\Psi_R\rangle$. Similar behavior is obtained for other disorder strengths within the MBL regime.

If these spins are entangled, the expectation value decays in the same power-law fashion as for the single-spin observables. However, if these spins are not entangled, the expectation value decays as a product of individual single-spin observables and, therefore, decays much faster. Before the spins get entangled, the sum of all Pauli strings results in a stretched exponential behavior of the sum in (1), leading to a power-law growth of \mathcal{M}_2 with a certain exponent β . As the particles gradually become entangled, this exponent decreases, and the growth of the SRE slows down with time. This results in the following dynamics of SRE in the MBL regime

$$\mathcal{M}_2^{\text{MBL}} = \mathcal{M}_2^{\text{sat}} - c/t^\beta, \qquad (5)$$

where $\mathcal{M}_{2}^{\text{sat}}$, c and β are constants dependent on the initial state. In particular, for the X-polarized initial state $|\Psi_X^+\rangle$, the saturation value $\mathcal{M}_2^{\text{sat}} = \mathcal{M}_2^{\text{Haar}}$ is the same as for the ergodic dynamics, $c = \ln(D)$, and $\beta = \xi \ln(2)$ is the localization length of ℓ -bits [94], see the orange dashed line in Fig. 1. For a generic initial product state, the saturation value is smaller, $\mathcal{M}_2^{\text{sat}} < \mathcal{M}_2^{\text{Haar}}$ and $\beta < \xi \ln(2)$ [94].

Microscopic model. To assess the accuracy of the ℓ bit in capturing the magic dynamics of strongly interacting disordered systems, we analyze a microscopic model expected to exhibit an MBL phase at sufficiently strong disorder, the disordered TFIM, with Hamiltonian

$$\hat{\mathcal{H}}_{\text{TFIM}} = \sum_{i=1}^{L-1} J_{i,i+1} \hat{Z}_i \hat{Z}_{i+1} + \sum_{i=1}^{L} h_i \hat{Z}_i + g \sum_{i=1}^{L} \hat{X}_i \quad (6)$$

where $h_i \in [-W, W]$ are random on-site fields that are drawn from a uniform distribution, g is the transverse field, and $J_{i,i+1}$ are the interactions between neighboring spins. Building on [104], we consider nearest-neighbor couplings drawn from a uniform distribution $J_{i,i+1} \in$ [0.8, 1.2], and fix the transverse field at g = 1. Mathematical arguments [104, 105] suggest that this model hosts an MBL phase for sufficiently large disorder strength W. Finite-size scaling analysis [66, 106] places the critical disorder threshold at $W_c \approx 3.5$. However, similar to the XXZ chain [65], the model exhibits finite-size drifts [107]. We study the quench dynamics of an initial state $|\Psi\rangle$ using the Chebyshev polynomial expansion [69, 108] up to $t = 2 \times 10^4 J$. We consider chains of $L \in [8, 20]$ spins, with results averaged over 1000 disorder realizations. The chosen initial state $|\Psi\rangle$ is a product state with energy near the middle of the spectrum.

In Fig. 2(a), we show the time evolution of a Z polarized initial state $|\Psi_Z\rangle$. This state belongs to the computational basis and is a conventional choice in MBL studies. For weak disorder, $W \sim 1$, \mathcal{M}_2 quickly grows towards the Haar value in the weak disorder regime, consistent with the ergodic dynamics results [43, 44]. For strong disorder $W \sim 5$, $|\Psi_Z\rangle$ is close to an eigenstate of (6) and, therefore, SRE increases very slowly. In the MBL regime, the behavior of SRE is accurately captured by the phenomenological formula (5), with the saturation value $\mathcal{M}_2^{\text{sat}}$ considerably smaller than $\mathcal{M}_2^{\text{Haar}}$.

In Fig. 2(b), we present the SRE evolution of initial X-polarized state $|\Psi_X\rangle$, a random state in the X basis. The SRE, \mathcal{M}_2 , initially grows rapidly before slowing down at longer times. Even for strong disorder, the SRE approaches the Haar value at late times. A similar trend is observed for a randomly selected state of Y basis, $|\Psi_Y\rangle$, as shown in Fig. 2(c). This choice leads to the smallest contribution of the Pauli strings involving only \hat{Z}_i operators, and, consequently, the saturation value in (5) is $\mathcal{M}_2^{\text{Haar}}$.

Additionally, we examine product states constructed by random rotations on the Bloch sphere, denoted as $|\Psi_B\rangle$. Since such states cannot be constructed by Clifford



Figure 3. Nonstabilizerness versus entanglement in the MBL regime. The SRE \mathcal{M}_2 is plotted as a function of the halfchain entanglement entropy \mathcal{S} for different disorder strengths W and system size L. For the Y-polarized state $|\Psi_Y\rangle$, (**a**), $\mathcal{M}_2(\mathcal{S})$ collapse, without any fitting parameters, on a single master curve both for ℓ -bit model and TFIM. For the random product state $|\Psi_R\rangle$, (**b**), the collapse occurs when $\mathcal{M}_2(\mathcal{S})$ are rescaled by an L-independent factor f(W).

gates, $\mathcal{M}_2 \neq 0$ at t = 0. However, this initial value is significantly below the Haar limit, as non-entangled states typically possess low nonstabilizerness [109]. Therefore, under Hamiltonian dynamics, the magic resources spread over time, consistently with (5), and the SRE growth is intermediate between the dynamics for the $|\Psi_Z\rangle$ and $|\Psi_Y\rangle$ states.

Collapsing of stabilizer entropy. To understand the interplay of nonstabilizerness and entanglement in the MBL regime, we analyze in Fig. 3 the growth of \mathcal{M}_2 as a function of the half-chain entanglement entropy $\mathcal{S} = -\text{tr}(\rho_{L/2} \ln \rho_{L/2})$, where $\rho_{L/2}$ is the reduced density matrix obtained by tracing out the degrees of freedom of the first half of the system. The entanglement entropy has been proposed as an "internal clock" for disordered localized systems, providing a natural way to compare the dynamic evolution at different values of W [49].

We consider $|\Psi_Y\rangle$ as our initial state. As shown in Fig. 3 (a), both ℓ -bit model and TFIM exhibit similar dynamics, with $\mathcal{M}_2(S)$ collapsing onto a single master curve without any fitting parameters. For an initial random product state $|\Psi_R\rangle$ state the ℓ -bit model predictions are aligned with the TFIM results when a disorderdependent rescaling function, f(W), is introduced. In Fig. 3(b) we show $\mathcal{M}_2/f(W)$ for $|\Psi_R\rangle$, where f(W) is found by minimizing deviations of TFIM results from the ℓ -bit model predictions. Importantly, f(W) is independent of system size L. These results demonstrate a close connection between the SRE and entanglement growth in the MBL regime.

Finally, we investigate whether \mathcal{M}_2 can serve as a diagnostic for the crossover between ETH and MBL regimes. The SRE has been linked to phase transitions in many ground-state problems [81, 110–113] and, therefore, one may expect that it is sensitive to the ETH-MBL crossover



Figure 4. Nonstabilizerness across the ETH-MBL crossover in TFIM. (a) Deviation from the Haar value ΔM_2 as a function of system size L and for different disorder strengths. (b) ΔM_2 as a function of disorder strength W at the latest time available for the $|\Psi_Z\rangle$ distinguishes the ETH and MBL regimes.

observed for $L \approx 20$ at $W_c \sim 3.5$ [66]. We consider time evolution of $|\Psi_Z\rangle$ and compute $\Delta \mathcal{M}_2 = \mathcal{M}_2^{\text{Haar}} - \mathcal{M}_2$ for different size systems L at the longest available time $(t = 2 \times 10^4)$, as shown in Fig. (4)(a). For weak disorder strength, $\Delta \mathcal{M}_2 \rightarrow 0$ as L increases, as expected for the ETH regime. On the other hand, $\Delta \mathcal{M}_2$ grows linearly with L for strong disorder. The crossover between these two regimes occurs near $W_c \approx 3.5$, showing how the SRE distinguishes the ergodic and non-ergodic dynamical regimes.

Conclusions and outlook. In this work, we have explored the dynamics of the nonstabilizerness in disordered many-body systems exhibiting MBL. By developing a theoretical framework founded on an ℓ -bit phenomenology, we have shown that magic spreading in MBL systems is fundamentally constrained by the slow dynamics characteristic of the MBL regime. Unlike ergodic systems, where SRE rapidly saturates to its maximal value, MBL systems display a much slower power-law relaxation towards the saturation value, which exhibits a strong variability with respect to the choice of initial state. Through numerical simulations of the disordered TFIM, we have verified our analytical formulas for the SRE growth and demonstrated a strong connection between the SRE and entanglement entropy growth in the MBL regime. In particular, our findings show that the disorder suppresses the rapid spread of magic resources, and interactions play a crucial role in enabling its slow but sustained growth. Furthermore, we have shown that SRE can serve as a diagnostic tool distinguishing ergodic and non-ergodic dynamics.

Our results open up several intriguing directions for future research. One promising avenue is investigating the interplay between magic and other forms of ergodicity breaking, such as quantum many-body scars [114– 116] and disorder-free localization [117–123]. In particular, extending our analysis to gauge theories [124, 125] and constrained quantum systems [126–130] could shed further light on the role of magic in non-thermalizing quantum dynamics. Moreover, this also motivates further explorations of Clifford-augmented matrix product states [131–135] in scenarios where continuous Hamiltonians govern the dynamics. Such investigations may provide a broader understanding of the interplay of disorder and magic spreading for quantum error correction and fault-tolerant quantum computing.

Acknowledgments. E.T. and P.S. acknowledge collaboration with X. Turkeshi and M. Dalmonte on related subjects. P.S. acknowledges insightful discussions with P. Stornati, S. Masot-Llima, A. Garcia-Saez. The work of P.R.N.F. was funded by the National Science Centre, Poland, project 2021/03/Y/ST2/00186 within the QuantERA II Programme that has received funding from the European Union Horizon 2020 research and innovation programme under Grant agreement No 101017733. P.S. acknowledges fellowship within the "Generación D" initiative, Red.es, Ministerio para la Transformación Digital y de la Función Pública, for talent atraction (C005/24-ED CV1), funded by the European Union NextGenerationEU funds, through PRTR. The work of J.Z. was funded by the National Science Centre, Poland, project 2021/43/I/ST3/01142 - OPUS call within the WEAVE programme. E.T. acknowledges support from ERC under grant agreement n.101053159 (RAVE), and CINECA (Consorzio Interuniversitario per il Calcolo Automatico) award, under the ISCRA initiative and Leonardo early access program, for the availability of high-performance computing resources and support. The study was also partially funded by the "Research Support Module" as part of the "Excellence Initiative - Research University" program at the Jagiellonian University in Kraków. We gratefully acknowledge Polish high-performance computing infrastructure PLGrid (HPC Centers: ACK Cyfronet AGH) for providing computer facilities and support within computational grant no. PLG/2024/017289

- * pedro.nicaciofalcao@doctoral.uj.edu.pl
- [†] etirrito@sissa.it
- M. A. Nielsen and I. L. Chuang, *Quantum Computa*tion and *Quantum Information* (Cambridge University Press, 2000).
- [2] 40 years of quantum computing, Nature Reviews Physics 4, 1 (2022).
- [3] M. Lewenstein, A. Sanpera, V. Ahufinger, B. Damski, S. Adithi, and U. Sen, Ultracold atomic gases in optical lattices: mimicking condensed matter physics and beyond, Advances in Physics 56, 243 (2007).
- [4] J. I. Cirac and P. Zoller, Goals and opportunities in quantum simulation, Nature Physics 8, 264 (2012).
- [5] E. Altman *et al.*, Quantum simulators: Architectures and opportunities, PRX Quantum 2, 017003 (2021).
- [6] D. P. DiVincenzo, The physical implementation of

quantum computation, Fortschritte der Physik 48, 771 (2000).

- [7] A. Montanaro, Quantum algorithms: an overview, npj Quantum Information 2, 15023 (2016).
- [8] M. Cerezo, A. Arrasmith, R. Babbush, S. C. Benjamin, S. Endo, K. Fujii, J. R. McClean, K. Mitarai, X. Yuan, L. Cincio, and P. J. Coles, Variational quantum algorithms, Nature Reviews Physics 3, 625 (2021).
- [9] K. Bharti, A. Cervera-Lierta, T. H. Kyaw, T. Haug, S. Alperin-Lea, A. Anand, M. Degroote, H. Heimonen, J. S. Kottmann, T. Menke, W.-K. Mok, S. Sim, L.-C. Kwek, and A. Aspuru-Guzik, Noisy intermediatescale quantum algorithms, Rev. Mod. Phys. 94, 015004 (2022).
- [10] G. Vidal, Efficient classical simulation of slightly entangled quantum computations, Phys. Rev. Lett. 91, 147902 (2003).
- [11] F. Verstraete, V. Murg, and J. Cirac, Matrix product states, projected entangled pair states, and variational renormalization group methods for quantum spin systems, Advances in Physics 57, 143 (2008).
- [12] N. Schuch, M. M. Wolf, F. Verstraete, and J. I. Cirac, Entropy scaling and simulability by matrix product states, Phys. Rev. Lett. **100**, 030504 (2008).
- [13] U. Schollwöck, The density-matrix renormalization group in the age of matrix product states, Annals of Physics **326**, 96 (2011).
- [14] J. Haegeman, C. Lubich, I. Oseledets, B. Vandereycken, and F. Verstraete, Unifying time evolution and optimization with matrix product states, Phys. Rev. B 94, 165116 (2016).
- [15] R. Orús, Tensor networks for complex quantum systems, Nature Reviews Physics 1, 538 (2019).
- [16] M. C. Bañuls, Tensor network algorithms: A route map, Annual Review of Condensed Matter Physics 14, 173–191 (2023).
- [17] L. Amico, R. Fazio, A. Osterloh, and V. Vedral, Entanglement in many-body systems, Rev. Mod. Phys. 80, 517 (2008).
- [18] R. Horodecki, P. Horodecki, M. Horodecki, and K. Horodecki, Quantum entanglement, Rev. Mod. Phys. 81, 865 (2009).
- [19] J. Preskill, Quantum computing and the entanglement frontier (2012), arXiv:1203.5813 [quant-ph].
- [20] D. Gottesman, The heisenberg representation of quantum computers (1998), arXiv:quant-ph/9807006 [quantph].
- [21] D. Gottesman, Theory of fault-tolerant quantum computation, Phys. Rev. A 57, 127 (1998).
- [22] S. Aaronson and D. Gottesman, Improved simulation of stabilizer circuits, Phys. Rev. A 70, 052328 (2004).
- [23] C. Gidney, Stim: a fast stabilizer circuit simulator, Quantum 5, 497 (2021).
- [24] S. Bravyi and A. Kitaev, Universal quantum computation with ideal clifford gates and noisy ancillas, Phys. Rev. A 71, 022316 (2005).
- [25] D. Gross, Hudson's theorem for finite-dimensional quantum systems, Journal of Mathematical Physics 47, 122107 (2006).
- [26] V. Veitch, S. A. Hamed Mousavian, D. Gottesman, and J. Emerson, The resource theory of stabilizer quantum computation, New Journal of Physics 16, 013009 (2014).
- [27] Z.-W. Liu and A. Winter, Many-body quantum magic, PRX Quantum 3, 020333 (2022).

- [28] L. Leone, S. F. Oliviero, and A. Hamma, Stabilizer rényi entropy, Phys. Rev. Lett. 128, 050402 (2022).
- [29] L. Leone and L. Bittel, Stabilizer entropies are monotones for magic-state resource theory, Phys. Rev. A 110, L040403 (2024).
- [30] E. Chitambar and G. Gour, Quantum resource theories, Rev. Mod. Phys. 91, 025001 (2019).
- [31] J. M. Deutsch, Quantum statistical mechanics in a closed system, Phys. Rev. A 43, 2046 (1991).
- [32] M. Srednicki, Chaos and quantum thermalization, Phys. Rev. E 50, 888 (1994).
- [33] M. Rigol, V. Dunjko, and M. Olshanii, Thermalization and its mechanism for generic isolated quantum systems, Nature 452, 854 (2008).
- [34] L. D'Alessio, Y. Kafri, A. Polkovnikov, and M. Rigol, From quantum chaos and eigenstate thermalization to statistical mechanics and thermodynamics, Advances in Physics 65, 239 (2016).
- [35] L. Foini and J. Kurchan, Eigenstate thermalization hypothesis and out of time order correlators, Phys. Rev. E 99, 042139 (2019).
- [36] S. Pappalardi, L. Foini, and J. Kurchan, Eigenstate thermalization hypothesis and free probability, Phys. Rev. Lett. **129**, 170603 (2022).
- [37] S. Pappalardi, F. Fritzsch, and T. Prosen, General eigenstate thermalization via free cumulants in quantum lattice systems (2023), arXiv:2303.00713 [condmat.stat-mech].
- [38] S. Pappalardi, L. Foini, and J. Kurchan, Microcanonical windows on quantum operators, Quantum 8, 1227 (2024).
- [39] A. M. Läuchli and C. Kollath, Spreading of correlations and entanglement after a quench in the one-dimensional bose-hubbard model, Journal of Statistical Mechanics: Theory and Experiment 2008, P05018 (2008).
- [40] H. Kim and D. A. Huse, Ballistic spreading of entanglement in a diffusive nonintegrable system, Phys. Rev. Lett. 111, 127205 (2013).
- [41] T. Rakovszky, F. Pollmann, and C. W. von Keyserlingk, Sub-ballistic growth of rényi entropies due to diffusion, Phys. Rev. Lett. **122**, 250602 (2019).
- [42] M. Žnidarič, Entanglement growth in diffusive systems, Communications Physics 3, 100 (2020).
- [43] X. Turkeshi, E. Tirrito, and P. Sierant, Magic spreading in random quantum circuits (2025), arXiv:2407.03929 [quant-ph].
- [44] E. Tirrito, X. Turkeshi, and P. Sierant, Anticoncentration and magic spreading under ergodic quantum dynamics (2024), arXiv:2412.10229 [quant-ph].
- [45] X. Turkeshi, A. Dymarsky, and P. Sierant, Pauli spectrum and nonstabilizerness of typical quantum manybody states, Physical Review B 111, 054301 (2025).
- [46] D. J. Luitz and Y. Bar Lev, Anomalous thermalization in ergodic systems, Phys. Rev. Lett. 117, 170404 (2016).
- [47] S. Bera, G. De Tomasi, F. Weiner, and F. Evers, Density propagator for many-body localization: Finite-size effects, transient subdiffusion, and exponential decay, Phys. Rev. Lett. 118, 196801 (2017).
- [48] D. Sels and A. Polkovnikov, Thermalization of dilute impurities in one-dimensional spin chains, Phys. Rev. X 13, 011041 (2023).
- [49] F. Evers, I. Modak, and S. Bera, Internal clock of manybody delocalization, Phys. Rev. B 108, 134204 (2023).
- [50] R. Nandkishore and D. A. Huse, Many-Body Localiza-

tion and Thermalization in Quantum Statistical Mechanics, Annual Review of Condensed Matter Physics 6, 15 (2015).

- [51] F. Alet and N. Laflorencie, Many-body localization: An introduction and selected topics, Comptes Rendus Physique 19, 498 (2018).
- [52] D. A. Abanin, E. Altman, I. Bloch, and M. Serbyn, Colloquium: Many-body localization, thermalization, and entanglement, Rev. Mod. Phys. 91, 021001 (2019).
- [53] P. Sierant, M. Lewenstein, A. Scardicchio, L. Vidmar, and J. Zakrzewski, Many-body localization in the age of classical computing, Reports on Progress in Physics 88, 026502 (2025).
- [54] V. Oganesyan and D. A. Huse, Localization of interacting fermions at high temperature, Phys. Rev. B 75, 155111 (2007).
- [55] A. Pal and D. A. Huse, Many-body localization phase transition, Phys. Rev. B 82, 174411 (2010).
- [56] D. A. Huse, R. Nandkishore, and V. Oganesyan, Phenomenology of fully many-body-localized systems, Phys. Rev. B 90, 174202 (2014).
- [57] D. J. Luitz, N. Laflorencie, and F. Alet, Many-body localization edge in the random-field Heisenberg chain, Phys. Rev. B **91**, 081103 (2015).
- [58] T. B. Wahl, A. Pal, and S. H. Simon, Efficient representation of fully many-body localized systems using tensor networks, Phys. Rev. X 7, 021018 (2017).
- [59] M. Mierzejewski, M. Kozarzewski, and P. Prelovšek, Counting local integrals of motion in disordered spinless-fermion and Hubbard chains, Phys. Rev. B 97, 064204 (2018).
- [60] S. J. Thomson and M. Schiró, Time evolution of manybody localized systems with the flow equation approach, Phys. Rev. B 97, 060201 (2018).
- [61] J. Šuntajs, J. Bonča, T. Prosen, and L. Vidmar, Quantum chaos challenges many-body localization, Phys. Rev. E 102, 062144 (2020).
- [62] P. Sierant, D. Delande, and J. Zakrzewski, Thouless Time Analysis of Anderson and Many-Body Localization Transitions, Phys. Rev. Lett. **124**, 186601 (2020).
- [63] R. K. Panda, A. Scardicchio, M. Schulz, S. R. Taylor, and M. Znidaric, Can we study the many-body localisation transition?, Europhysics Letters 128, 67003 (2020).
- [64] D. Sels and A. Polkovnikov, Dynamical obstruction to localization in a disordered spin chain, Phys. Rev. E 104, 054105 (2021).
- [65] P. Sierant, M. Lewenstein, and J. Zakrzewski, Polynomially filtered exact diagonalization approach to manybody localization, Phys. Rev. Lett. **125**, 156601 (2020).
- [66] D. Abanin, J. H. Bardarson, G. De Tomasi, S. Gopalakrishnan, V. Khemani, S. Parameswaran, F. Pollmann, A. Potter, M. Serbyn, and R. Vasseur, Distinguishing localization from chaos: Challenges in finite-size systems, Annals of Physics 427, 168415 (2021).
- [67] M. Kiefer-Emmanouilidis, R. Unanyan, M. Fleischhauer, and J. Sirker, Unlimited growth of particle fluctuations in many-body localized phases, Annals of Physics 435, 168481 (2021).
- [68] R. Ghosh and M. Žnidarič, Resonance-induced growth of number entropy in strongly disordered systems, Phys. Rev. B 105, 144203 (2022).
- [69] P. Sierant and J. Zakrzewski, Challenges to observation of many-body localization, Phys. Rev. B 105, 224203 (2022).

- [70] A. Morningstar, L. Colmenarez, V. Khemani, D. J. Luitz, and D. A. Huse, Avalanches and many-body resonances in many-body localized systems, Phys. Rev. B 105, 174205 (2022).
- [71] G. D. Chiara, S. Montangero, P. Calabrese, and R. Fazio, Entanglement entropy dynamics of Heisenberg chains, Journal of Statistical Mechanics: Theory and Experiment **2006**, P03001 (2006).
- [72] M. Žnidarič, T. Prosen, and P. Prelovšek, Many-body localization in the Heisenberg XXZ magnet in a random field, Phys. Rev. B 77, 064426 (2008).
- [73] J. H. Bardarson, F. Pollmann, and J. E. Moore, Unbounded growth of entanglement in models of manybody localization, Phys. Rev. Lett. 109, 017202 (2012).
- [74] M. Serbyn, Z. Papić, and D. A. Abanin, Universal slow growth of entanglement in interacting strongly disordered systems, Phys. Rev. Lett. **110**, 260601 (2013).
- [75] F. Iemini, A. Russomanno, D. Rossini, A. Scardicchio, and R. Fazio, Signatures of many-body localization in the dynamics of two-site entanglement, Phys. Rev. B 94, 214206 (2016).
- [76] M. Serbyn, Z. Papić, and D. A. Abanin, Local conservation laws and the structure of the many-body localized states, Phys. Rev. Lett. 111, 127201 (2013).
- [77] T. Haug and L. Piroli, Quantum 7, 1092 (2023).
- [78] D. Gross, S. Nezami, and M. Walter, Commun. Math. Phys. 385, 1325 (2021).
- [79] P. Sierant *et al.*, To appear.
- [80] T. Haug and L. Piroli, Quantifying nonstabilizerness of matrix product states, Phys. Rev. B 107, 035148 (2023).
- [81] P. S. Tarabunga, E. Tirrito, T. Chanda, and M. Dalmonte, Many-body magic via pauli-markov chains—from criticality to gauge theories, PRX Quantum 4, 040317 (2023).
- [82] G. Lami and M. Collura, Nonstabilizerness via perfect pauli sampling of matrix product states, Phys. Rev. Lett 131, 180401 (2023).
- [83] P. S. Tarabunga, E. Tirrito, M. C. Bañuls, and M. Dalmonte, Nonstabilizerness via matrix product states in the pauli basis, Phys. Rev. Lett. 133, 010601 (2024).
- [84] M. Frau, P. S. Tarabunga, M. Collura, M. Dalmonte, and E. Tirrito, Nonstabilizerness versus entanglement in matrix product states, Phys. Rev. B 110, 045101 (2024).
- [85] E. Tirrito, P. S. Tarabunga, G. Lami, T. Chanda, L. Leone, S. F. Oliviero, M. Dalmonte, M. Collura, and A. Hamma, Quantifying nonstabilizerness through entanglement spectrum flatness, Phys. Rev. A 109, L040401 (2024).
- [86] X. Turkeshi, M. Schirò, and P. Sierant, Measuring nonstabilizerness via multifractal flatness, Phys. Rev. A 108, 042408 (2023).
- [87] S. F. E. Oliviero, L. Leone, A. Hamma, and S. Lloyd, Measuring magic on a quantum processor, npj Quantum Information 8, 148 (2022).
- [88] T. Haug and M. Kim, Scalable measures of magic resource for quantum computers, PRX Quantum 4, 010301 (2023).
- [89] P. Niroula, C. D. White, Q. Wang, S. Johri, D. Zhu, C. Monroe, C. Noel, and M. J. Gullans, Phase transition in magic with random quantum circuits, Nat. Phys. 20, 1786 (2024).
- [90] X. Turkeshi, A. Dymarsky, and P. Sierant, Pauli spectrum and nonstabilizerness of typical quantum manybody states, Phys. Rev. B 111, 054301 (2025).

- [91] J. Odavić, M. Viscardi, and A. Hamma, Stabilizer entropy in non-integrable quantum evolutions (2024), arXiv:2412.10228 [quant-ph].
- [92] V. Ros, M. Mueller, and A. Scardicchio, Integrals of motion in the many-body localized phase, Nuclear Physics B 891, 420 (2015).
- [93] J. Z. Imbrie, V. Ros, and A. Scardicchio, Local integrals of motion in many-body localized systems, Annalen der Physik 529, 1600278 (2017).
- [94] See Supplementary Material at [URL will be inserted by publisher] for further details on the analytical calculation of \mathcal{M}_2 for Anderson systems, details on the ℓ -bit model, dependence of the initial state and entanglement based collapse of \mathcal{M}_2 .
- [95] P. Sierant and J. Zakrzewski, Model of level statistics for disordered interacting quantum many-body systems, Phys. Rev. B 101, 104201 (2020).
- [96] A. Prakash, J. Pixley, and M. Kulkarni, Universal spectral form factor for many-body localization, Phys. Rev. Res. 3, L012019 (2021).
- [97] M. Serbyn, Z. Papić, and D. A. Abanin, Quantum quenches in the many-body localized phase, Phys. Rev. B 90, 174302 (2014).
- [98] M. Žnidarič, Entanglement in a dephasing model and many-body localization, Phys. Rev. B 97, 214202 (2018).
- [99] D. Aceituno Chávez, C. Artiaco, T. Klein Kvorning, L. Herviou, and J. H. Bardarson, Ultraslow growth of number entropy in an l-bit model of many-body localization, Phys. Rev. Lett. **133**, 126502 (2024).
- [100] V. Berger, A. Nava, J. H. Bardarson, and C. Artiaco, Numerical study of disordered noninteracting chains coupled to a local lindblad bath (2024), arXiv:2412.03233 [cond-mat.dis-nn].
- [101] T. Szołdra, P. Sierant, M. Lewenstein, and J. Zakrzewski, Catching thermal avalanches in the disordered xxz model, Phys. Rev. B 109, 134202 (2024).
- [102] A. Scocco, G. Passarelli, M. Collura, P. Lucignano, and A. Russomanno, Thermalization propagation front and robustness against avalanches in localized systems, Phys. Rev. B 110, 134204 (2024).
- [103] M. Serbyn, Z. Papić, and D. A. Abanin, Quantum quenches in the many-body localized phase, Phys. Rev. B 90, 174302 (2014).
- [104] J. Z. Imbrie, On many-body localization for quantum spin chains, Journal of Statistical Physics 163, 998 (2016).
- [105] W. D. Roeck, L. Giacomin, F. Huveneers, and O. Prosniak, Absence of normal heat conduction in strongly disordered interacting quantum chains (2024), arXiv:2408.04338 [math-ph].
- [106] G. De Tomasi, I. M. Khaymovich, F. Pollmann, and S. Warzel, Rare thermal bubbles at the many-body localization transition from the fock space point of view, Phys. Rev. B **104**, 024202 (2021).
- [107] P. Sierant, M. Lewenstein, A. Scardicchio, and J. Zakrzewski, Stability of many-body localization in floquet systems, Phys. Rev. B 107, 115132 (2023).
- [108] H. Tal-Ezer and R. Kosloff, An accurate and efficient scheme for propagating the time dependent schrödinger equation, The Journal of chemical physics 81, 3967 (1984).
- [109] S. F. E. Oliviero, L. Leone, and A. Hamma, Magic-state resource theory for the ground state of the transverse-

field ising model, Phys. Rev.A 106, 042426 (2022).

- [110] P. S. Tarabunga and C. Castelnovo, Magic in generalized Rokhsar-Kivelson wavefunctions, Quantum 8, 1347 (2024).
- [111] P. R. N. Falcão, P. S. Tarabunga, M. Frau, E. Tirrito, J. Zakrzewski, and M. Dalmonte, Nonstabilizerness in U(1) lattice gauge theory, Phys. Rev. B 111, L081102 (2025).
- [112] P. S. Tarabunga, Critical behaviors of non-stabilizerness in quantum spin chains, Quantum 8, 1413 (2024).
- [113] Y.-M. Ding, Z. Wang, and Z. Yan, Evaluating manybody stabilizer r\'enyi entropy by sampling reduced pauli strings: singularities, volume law, and nonlocal magic (2025), arXiv:2501.12146 [quant-ph].
- [114] M. Serbyn, D. A. Abanin, and Z. Papić, Quantum many-body scars and weak breaking of ergodicity, Nature Physics 17, 675 (2021).
- [115] R. Smith, A. Hallam, et al., Non-stabilizerness in kinetically-constrained rydberg atom arrays (2024), arXiv:2406.14348 [quant-ph].
- [116] J. Hartse, L. Fidkowski, and N. Mueller, Stabilizer scars (2024), arXiv:2411.12797 [quant-ph].
- [117] A. Smith, J. Knolle, D. L. Kovrizhin, and R. Moessner, Disorder-free localization, Phys. Rev. Lett. 118, 266601 (2017).
- [118] M. Brenes, M. Dalmonte, M. Heyl, and A. Scardicchio, Many-body localization dynamics from gauge invariance, Phys. Rev. Lett. **120**, 030601 (2018).
- [119] E. van Nieuwenburg, Y. Baum, and G. Refael, From Bloch oscillations to many-body localization in clean interacting systems, Proceedings of the National Academy of Sciences 116, 9269 (2019).
- [120] M. Schulz, C. A. Hooley, R. Moessner, and F. Pollmann, Stark many-body localization, Phys. Rev. Lett. 122, 040606 (2019).
- [121] R. Yao and J. Zakrzewski, Many-body localization of bosons in an optical lattice: Dynamics in disorder-free potentials, Phys. Rev. B 102, 104203 (2020).
- [122] R. Yao, T. Chanda, and J. Zakrzewski, Nonergodic dynamics in disorder-free potentials, Annals of Physics 435, 168540 (2021), special Issue on Localisation 2020.
- [123] R. Yao, T. Chanda, and J. Zakrzewski, Many-body localization in tilted and harmonic potentials, Phys. Rev. B 104, 014201 (2021).
- [124] M. C. Bañuls *et al.*, Simulating lattice gauge theories within quantum technologies, The European Physical Journal D 74, 165 (2020).
- [125] M. Aidelsburger, L. Barbiero, A. Bermudez, T. Chanda, A. Dauphin, D. González-Cuadra, P. R. Grzybowski, S. Hands, F. Jendrzejewski, J. Jünemann, G. Juzeliūnas, V. Kasper, A. Piga, S.-J. Ran, M. Rizzi, G. Sierra, L. Tagliacozzo, E. Tirrito, T. V. Zache, J. Zakrzewski, E. Zohar, and M. Lewenstein, Cold atoms meet lattice gauge theory, Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences **380**, 10.1098/rsta.2021.0064 (2021).
- [126] C. Chen, F. Burnell, and A. Chandran, How does a locally constrained quantum system localize?, Phys. Rev. Lett. 121, 085701 (2018).
- [127] P. Sierant, E. G. Lazo, M. Dalmonte, A. Scardicchio, and J. Zakrzewski, Constraint-induced delocalization, Phys. Rev. Lett. **127**, 126603 (2021).
- [128] H. Théveniaut, Z. Lan, G. Meyer, and F. Alet, Transition to a many-body localized regime in a two-

dimensional disordered quantum dimer model, Phys. Rev. Res. **2**, 033154 (2020).

- [129] F. Pietracaprina and F. Alet, Probing many-body localization in a disordered quantum dimer model on the honeycomb lattice, SciPost Phys. 10, 044 (2021).
- [130] K. Royen, S. Mondal, F. Pollmann, and F. Heidrich-Meisner, Enhanced many-body localization in a kinetically constrained model, Phys. Rev. E 109, 024136 (2024).
- [131] S. Masot-Llima and A. Garcia-Saez, Stabilizer tensor networks: Universal quantum simulator on a basis of stabilizer states, Phys. Rev. Lett. 133, 230601 (2024).
- [132] X. Qian, J. Huang, and M. Qin, Augmenting density matrix renormalization group with clifford circuits, Phys. Rev. Lett. 133, 190402 (2024).
- [133] G. Lami, T. Haug, and J. D. Nardis, Quantum state designs with clifford enhanced matrix product states (2024), arXiv:2404.18751 [quant-ph].
- [134] X. Qian, J. Huang, and M. Qin, Clifford circuits augmented time-dependent variational principle (2024), arXiv:2407.03202 [cond-mat.str-el].
- [135] A. C. Nakhl, B. Harper, M. West, N. Dowling, M. Sevior, T. Quella, and M. Usman, Stabilizer tensor networks with magic state injection (2024), arXiv:2411.12482 [quant-ph].
- [136] A. S. Aramthottil, P. Sierant, M. Lewenstein, and J. Zakrzewski, Phenomenology of many-body localization in bond-disordered spin chains, Phys. Rev. Lett. 133, 196302 (2024).
- [137] If the disorder is not coupled to on-site operators, the standard ℓ -bit description may fail to capture the system's physics. In such cases, a more sophisticated RSRG-X approach is required to correctly describe the model's properties [136].

SUPPLEMENTARY MATERIAL: MAGIC DYNAMICS IN MANY-BODY LOCALIZED SYSTEMS

NONSTABILIZERNESS OF ANDERSON STATES

In this section, we demonstrate how to obtain an analytical expression for the stabilizer Rényi entropy (SRE) for a perfect Anderson insulator. In this case, the ℓ -bits do not interact with each other and, therefore, only the first term of $\hat{\mathcal{H}}_{\ell-\mathrm{bit}}$ remains relevant. The SRE can be rewritten as

$$\mathcal{M}_2 = \mathbb{E}_{\{h\}} \left(-\log_2 \left[\sum_{P \in \mathcal{P}_L} \frac{|\mathrm{Tr}(\rho P)|^4}{D} \right] \right)$$
(S.1)

where $\mathbb{E}_{\{h\}}$ denotes the average over different disorder realizations and ρ is the density matrix of the state $|\Psi(t)\rangle$. Since the system is non-interacting, the density matrix can be written as $\rho = \bigotimes_{k=1}^{L} \rho_k$, where ρ_k is the reduced density matrix of the k-th spin. For any initial state $|\Psi\rangle = \bigotimes_{k=1}^{L} (\cos(\theta_k/2)| \uparrow\rangle + e^{i\phi_k} \sin(\theta_k/2)| \downarrow\rangle)$, the reduced density matrix ρ_k at time t is given by [97]:

$$\rho_k(t) = \begin{pmatrix} \cos^2(\theta_k/2) & \frac{\sin\theta_k}{2}e^{-i(2h_k t + \phi_k)} \\ \frac{\sin\theta_k}{2}e^{i(2h_k t + \phi_k)} & \sin^2(\theta_k/2) \end{pmatrix}$$
(S.2)

where h_k is the local magnetic field at site k. In this case, the sum of expectation values of all Pauli strings can be rewritten as

$$\sum_{P \in \mathcal{P}_L} |\mathrm{Tr}(\rho P)|^4 = \prod_{k=1}^L \sum_{\sigma \in \{I, X, Y, Z\}} |\mathrm{Tr}(\rho_k \sigma_k)|^4 \qquad (S.3)$$

implying that to understand how the SRE grows in the non-interacting picture, it suffices to understand the dynamics of single-spin observables. The single qubit SRE can be easily obtained through the reduced density matrix ρ_k , yielding

$$\sum_{\sigma \in \{I, X, Y, Z\}} |\operatorname{Tr}(\rho_k \sigma_k)|^4 = \frac{1}{2} (4 - \sin^2(2\theta_k) - \sin^4 \theta_k \sin^2 \delta_k)$$
(S.4)

where $\delta_k = 4h_k t + 2\phi_k$. Substituting this expression into Eq. S.3, and assuming that the on-site fields h_k are drawn independently and identically from a uniform distribution in the interval [-W, W], we obtain

$$\mathcal{M}_{2} = -\sum_{k=1}^{L} \frac{1}{W} \int_{0}^{W} dh \, \log_{2} \left[1 - \frac{1}{4} \sin^{2}(2\theta_{k}) - \frac{1}{4} \sin^{4}\theta_{k} \sin^{2}\delta_{k} \right]. \quad (S.5)$$

where the first term inside the bracket comes from the identity contribution, the second to the expectation value



Figure S.1. (a) Nonstabilizerness gain (see text) for a single qubit SRE throughout the whole parameter space (θ, ϕ) . The greatest gain occur for an initial $|+\rangle$ state, while the smallest occur for an initial $|T\rangle$ state. Comparison between the analytical solution given by Eq. S.5 (dashed line) and the numerical calculations of the ℓ -bit model (solid line) for an initial (b) $|\Psi_X^+\rangle$ and (c) $|\Psi_T\rangle$ state. The results were obtained for L = 16 and 10^3 disorder realizations.

of τ^z operators and the third term is obtained from the $\tau^{x/y}$ operators. At $\theta = \pi/2$, the second term vanishes and the single-qubit SRE significantly differ from its initial value. To quantify this difference, we define the non-stabilizerness gain

$$\mathcal{M}_2^{g} = \mathcal{M}_2^{sat} - \mathcal{M}_2(|\Psi\rangle) \tag{S.6}$$

where $\mathcal{M}_2^{\text{sat}}$ is the asymptotic value of the SRE and $\mathcal{M}_2(|\Psi\rangle)$ its value at t = 0. In Fig S.1(a), we show \mathcal{M}_2^{g} throughout the parameter space (θ, ϕ) , confirming that the highest variance of the SRE is around $\theta = \pi/2$.

In particular, the X polarized state gives us the greatest gain. To confirm our analytical result, we consider $|\Psi_X^+\rangle = \bigotimes_{k=1}^L |+\rangle$ as the initial state, where $|+\rangle$ is an eigenstate of the \hat{X} operator. In this particular case, Eq. S.5 can be written as

$$\mathcal{M}_{2} = -\frac{L}{W} \int_{0}^{W} dh \log_{2} \left[1 - \frac{1}{4} \sin^{2}(4ht) \right]$$
(S.7)

which, in the limit $t \to \infty$, yields $\mathcal{M}_2 \approx L \log_2(8/7)$. In Fig. S.1(b), we show a comparison between the analytical solution (dashed red line) and the numerical result (solid black line). The analytical solution perfectly describes the behavior of \mathcal{M}_2 , showing the accuracy of our analytical calculation. In a similar spirit, we compare the exact and numerical results for an initial state



10⁻² 10² t 10⁶ 10¹⁰ 10⁻² 10² t 10⁶ 10¹⁰ Figure S.2. (a) Half chain entanglement entropy (S) and (b) SRE dynamics in the ℓ -bit model for a chain with L = 16 spins and localization length $\xi = 0.5$. The simulation was performed assuming an initial state $|\Psi(0)\rangle = |+\rangle^{\otimes L}$ and the results were averaged over several disorder realizations.

2

0

 \mathcal{S}

5

4

3

 $\mathbf{2}$

1

0

Page

 $|\Psi_T\rangle = \bigotimes_{k=1}^L |T\rangle$, where $|T\rangle = (1/\sqrt{2})(|0\rangle + e^{i\pi/4}|1\rangle)$ is the magic state. In the disorderless case, $|\Psi_T\rangle$ has the highest SRE of a unentangled state [28]. Here, however, the initial spin precession quickly decreases its SRE value, leading to a saturation value that significantly differs from its initial value, as shown in Fig. S.1(c).

DETAILS ON THE *l*-BIT MODEL

In the MBL regime, when the disorder is coupled to on-site operators [137], the system is simply described by the emergence of an exponentially large set of integrals of motion, also known as localized bits (ℓ -bits). The quasilocal structure of the ℓ -bits encapsulates key signatures of MBL, making the Hamiltonian $\hat{\mathcal{H}}_{\ell-\text{bit}}$ (see Eq. 3) of the main letter) a natural framework to probe the nonstabilizerness dynamics in MBL. The physical degrees of freedom are related to the ℓ -bits by guasilocal unitary transformations U, satisfying $\hat{\tau}_i^z = U \hat{Z}_i U^{\dagger}$. Ideally, U reflects the system's localized nature, with its matrix elements decaying exponentially with the distance between *l*-bits. However, for practical purposes, a good approximation is to take U = I, where I is the identity operator, significantly simplifying the analysis while retaining the essential features of MBL dynamics.

In our simulations, we model interactions between ℓ bits as random variables drawn from a Gaussian distribution with zero mean, where the variance decays exponentially with the distance between spins, following the approach in [98]. Specifically, the variance of the twobody interaction term is given by

$$\langle (J_{i,j})^2 \rangle = e^{-|j-i|/\xi} \tag{S.8}$$

where ξ is the localization length of the system. A similar exponential decay is assumed for higher-order interactions, considering the maximum separation between spins.



Figure S.3. (a) Nonstabilizerness spread in the ℓ -bit model for different system sizes, with $|\Psi_X^+\rangle$ as the initial state. The dashed lines show the analytical solution (Eq. 5 of the main text), which accurately describes the SRE growth after $t \sim 1$. (b) Time evolution of $\Delta \mathcal{M}_2$ for different system sizes L. It decays polynomially in t until it eventually saturates, except for L = 18, for which the Heisenberg time is beyond the considered here. The saturation value exhibits an exponential decay with exponent $\lambda \approx -\ln 2$, as shown in the inset.

To investigate how different interaction terms in the ℓ -bit Hamiltonian influence key observables, we first analyze the dynamics of the half-chain entanglement entropy, defined as

$$\mathcal{S} = -\mathrm{tr}(\rho_{L/2} \ln \rho_{L/2}) \tag{S.9}$$

where $\rho_{L/2}$ is the reduced density matrix obtained by tracing out the degrees of freedom of the first half of spins. Fig. S.2(a) shows the growth of S starting from the initial state $|\Psi_{x}^{+}\rangle$. In the Anderson insulating case (solid black line), where interactions are absent, correlations cannot spread, and therefore $\mathcal{S} = 0$. However, in interacting cases, \mathcal{S} grows logarithmically in time, a hallmark of MBL. The rate of entanglement growth increases slightly as additional interaction terms are included in $\mathcal{H}_{\ell-\text{bit}}$ as illustrated in Fig. S.2(a). The differences between the interaction terms are more striking in the SRE behavior, as shown in Fig. S.2(b). Assuming only twobody interactions between the ℓ -bits does not lead to the SRE reaching the Haar value, at least for the time scales analyzed here. However, as we add corrections to the ℓ bit Hamiltonian, the dynamics of the SRE approximates the semi-analytical prediction described in Eq. 5.

To evaluate the accuracy of our semi-analytical description (Eq. 5 of the main letter), we examine in Fig. S.3(a) the growth of the SRE in the ℓ -bit model for different system sizes L, using $|\Psi_X^+\rangle$ as the initial state. For small chains, the analytical solution (dashed lines) deviates from the behavior obtained by the numerics. However, as the system size increases, the analytical expression accurately describes the behavior of \mathcal{M}_2 at time scales where entanglement is relevant, and it saturates close to the Haar value (dotted lines) after sufficiently long times. Fig. S.3(b) further explores this



Figure S.4. (a) Entanglement entropy and (b) SRE dynamics for a slightly perturbed eigenstate of the system. We consider $L = 16, \xi = 0.5$, and averaged the results over 500 disorder realizations. Despite the initial value, the dynamics of each state resembles the dynamics of the $|\Psi_Z\rangle$ in microscopic models.

convergence by showing the time evolution of the deviation from the Haar value, defined as ΔM_2 . ΔM_2 decreases polynomially with time before stabilizing at a finite asymptotic value ΔM_2^{∞} . As illustrated in the inset of Fig. S.3(b), this residual deviation decreases exponentially with system size, following a decay rate of approximately $\lambda \approx -\ln 2$. This exponential suppression suggests that in the thermodynamic limit M_2 converges to the Haar value, confirming that nonstabilizerness reaches its maximal extent in large-scale systems.

We also verify how the SRE will spread for an initial state that is close to an eigenstate of the system. To this end, we prepare a random product state in the computational basis and apply a unitary operator that is exponentially localized in the Fock space. Concretely, we select a random product state $|i'\rangle$ among $D = 2^L$ basis states and construct the state

$$|\Psi\rangle = \frac{1}{\sqrt{Z}} \sum_{i=1}^{D} e^{-\alpha d(i,i')} |i\rangle, \quad Z = \sum_{i=1}^{D} e^{-2\alpha d(i,i')} \quad (S.10)$$

where d(i, i') denotes the Hamming distance between two different strings of the same length. The parameter α controls the degree of localization in the computational basis and, therefore, the dynamics of this state in the ℓ -bit model should resemble the quench dynamics of an initial $|\Psi_Z\rangle$ in microscopic models, with α playing the role of an effective disorder strength.

In Fig. S.4(a), we present the time evolution of the entanglement entropy for different localization strengths α . Although S exhibits the expected logarithmic growth over time, its saturation value remains well below the Page value (black dotted line), indicating that the spins are not fully dephased. The extent of this saturation depends on α . A similar dependence on α is observed in the saturation value of \mathcal{M}_2 . However, as shown in Fig. S.4(b), the overall time-scaling behavior of \mathcal{M}_2 remains unchanged, suggesting that while localization



Figure S.5. W_Z dynamics for two different states, $|\Psi_Y\rangle$ and $|\Psi_Z\rangle$, in the (a) ergodic and (b) localized regimes. In the simulation, we consider L = 16 and average the results over, at least, 700 realizations. In the ergodic regime, the two different states evolve similarly due to the lack of integrals of motion. In the localized regime, however, the weight of Z gates (W_Z) depend on the initial state, leading to different saturation values of the SRE.

strength influences the final magic content, the fundamental growth dynamics are robust to variations in α .

DEPENDENCE ON THE INITIAL STATE AND WEIGHT OF Z GATES

In the previous section, we checked the choice of initial state strongly influences the saturation value of the SRE. This behavior arises from the system's integrals of motion: for an initial $|\Psi_X^+\rangle$ state, the expectation values of any Pauli operator containing Z gates vanish, allowing the SRE to reach its maximum value. On the other hand, for a slightly perturbed eigenstate, the presence of Z gates contributes nontrivially to the SRE calculation, leading to a lower saturation value. To assess how well the transverse-field Ising model (TFIM) aligns with this expected behavior, we compute the contribution of \hat{Z} gates, quantified by the weight function

$$\mathcal{W}_Z(|\Psi\rangle) = \sum_{P \in \mathcal{P}_{IZ}} \frac{|\langle \Psi|P|\Psi\rangle|^2}{D}$$
(S.11)

where \mathcal{P}_{IZ} is a subgroup of the Pauli group \mathcal{P}_L that contains only \hat{I} and \hat{Z} operators. In the following analysis, we study the dynamics of \mathcal{W}_Z for two different choices of initial state, $|\Psi_Y\rangle$ and $|\Psi_Z\rangle$, across the ergodic (W = 2.0) and localized (W = 8.0) regimes.

We begin by analyzing the dynamics of \mathcal{W}_Z in the ergodic regime. In this case, the system lacks an extensive set of integrals of motion, and beyond an initial transient period, the evolution of \mathcal{W}_Z becomes independent of the initial state, as shown in Fig. S.5(a). Here, Pauli strings containing only Z gates behave similarly to other Pauli strings, allowing the SRE to reach the Haar value at sufficiently long times. The behavior changes significantly in the localized regime, as depicted in Fig. S.5(b).



Figure S.6. Rescaled SRE $(\mathcal{M}_2/f(W))$ versus entanglement entropy (\mathcal{S}) for the initial state $|\Psi_R\rangle$ and (a) L = 12, (b) L =14, (c) L = 16 and (d) L = 20. The data collapse improves with increasing L, without altering the rescaling parameter f(W). For L = 20, the data were obtained via Monte-Carlo sampling using 15000 samples, and the shaded black region indicate the sampling errors.

For an initial $|\Psi_Y\rangle$ state, \mathcal{W}_Z saturates to a value close to that expected when only the identity operator contributes, indicated by the orange dashed line in the inset. This result is consistent with ℓ -bit predictions, where expectation values of Pauli operators containing Z gates should vanish for this specific state. On the other hand, for an initial $|\Psi_Z\rangle$ state, the weight of Z gates is considerable, and it reaches a value that is significantly higher than other gates. Its behavior is almost constant in time, as expected when the system is described by a set of constants of motion. However, by fitting with a power-law decay ($W_Z = at^{-\lambda}$), one obtains a small decay in time ($\lambda = 0.005 \pm 0.001$) suggesting an extremely slow delocalization at this particular disorder strength.

ENTANGLEMENT AS THE "INTERNAL-CLOCK" FOR THE NONSTABILIZERNESS DYNAMICS

In the main letter, we show that the entanglement entropy can be seen as an "internal-clock" for the SRE growth in the MBL regime, similarly to what occurs with other observables [49]. Here, however, we show that this result is fully consistent with the ℓ -bit framework after a proper rescaling by a function f(W) that is independent of the system size L. In this section, we explore this idea further by considering the $|\Psi_Y\rangle$ and $|\Psi_R\rangle$ as the initial state.

For the former choice, as shown in Fig 3(a) of the main text, both models agree without any need of rescaling by a function f(W). The situation differs if we set $|\Psi_R\rangle$ as the initial state. In this case, in order to achieve the collapse between the curves, we minimize the distance between the results obtained for the TFIM with respect to the one obtained for the $\hat{\mathcal{H}}_{\ell-\text{bit}}$. As shown in Fig S.6, the quality of the collapse improves as we increase the system size. Most importantly, the rescaling parameter f(W) is independent of L, as depicted in the inset of Fig 3(b) of the main text. Hence, our phenomenological description of the SRE growth based on the integrals of motion is fully consistent with the more intricate structure of microscopic Hamiltonians.