

Remarks on the Gluing Theorems for Compact Special Lagrangian Submanifolds with Isolated Conical Singularities

Yohsuke Imagi

Abstract

We make two improvements upon the gluing theorems of [6, 7] for compact special Lagrangian submanifolds with isolated conical singularities. Firstly, we get rid of a few technical hypotheses of them. Secondly, we replace another hypothesis by a weaker one and prove that the latter is a necessary condition for the gluing process to be possible.

1 Introduction

We begin by recalling briefly the set-up of [6, 7] (summarized in [8]) in which we glue together special Lagrangian submanifolds. We use nearly the same notation as in [6, 7]. Let M be a Kähler manifold of complex dimension $m > 2$, whose canonical bundle is holomorphically trivial. Let M be given a C^∞ family $(\omega^s, J^s, \Omega^s)_{s \in \mathbb{R}^n}$ of almost Calabi–Yau structures (in the sense of Definition 2.1). Let $X \subseteq (M; \omega^0, J^0, \Omega^0)$ be a special Lagrangian submanifold whose closure \bar{X} is compact and of the form $X \sqcup X^{\text{sing}}$ for some finite set X^{sing} . Suppose that for each $x \in X^{\text{sing}}$ there exists a unique tangent cone C_x to X at x (which is smooth outside vertex and of multiplicity one). More precisely, we suppose that X approaches with power order at x (as in Definition 2.6) the cone C_x . Let L_x be a smoothing model for C_x , which is a special Lagrangian submanifold of $T_x M \cong \mathbb{C}^m$ which approaches (with order ≤ 0) at infinity the cone C_x . Recall that there are two cohomology classes $Y(L_x) \in H^1(C_x, \mathbb{R})$ and $Z(L_x) \in H^{m-1}(C_x, \mathbb{R})$ relevant to the gluing process.

We prove that we can get rid of the hypothesis $m < 6$ in [8, Theorem 7.3], [7, Theorem 6.12] and [7, Theorem 8.9] (in the ascending order of generality). These are the results of [6, 7] in which there is at least one smoothing model L_x with the cohomology class $Y(L_x) \neq 0$.

There is also a technical hypothesis in the most general version [7, Theorem 8.9] of the gluing theorems in [6, 7]. This has to do with an equation for the cohomology classes $[\text{Im } \Omega^s]$ and $\bigoplus_{x \in X^{\text{sing}}} Z(L_x)$; for the more precise expression see [7, (108)]. It is supposed in [7, (135)–(136)] that both the sides of the equation vanish rather than that the equation itself holds. We prove that we can

get rid of this restriction; that is, the equation itself is sufficient for [7, Theorem 8.9].

Our version of the equation, (2.2) of the present paper, is in fact slightly more general than [7, (108)]. Following the proofs of [6, 7] we see immediately that we can make such a generalization. But we prove also that the generalized equation is a necessary condition for the gluing process to be possible.

We explain now how we prove our results. We begin with the same approximate solutions as in [6, 7] which we make from X and $\bigsqcup_{x \in X^{\text{sing}}} L_x$ using partitions of unity (not doing any analysis yet). We then perturb these to the true solutions, not by the method of [6, 7] but by the method of [3, 11]. The advantage of the latter is that the two building blocks X and $\bigsqcup_{x \in X^{\text{sing}}} L_x$ play symmetric rôles. This happens to other gluing methods including the well-known case of Yang–Mills gauge theory. For instance, if we want to glue together the model ASD connections over \mathbb{R}^4 and the flat connection over a compact 4-manifold M then we can take the actual building blocks to be $S^4 = \mathbb{R}^4 \sqcup \{\infty\}$ and M (as in [1, §7.2]) whose rôles are symmetric.

For X connected, using the method of [3, 11] we can perturb the approximate solutions directly to the true solutions. Otherwise, there is a non-trivial obstruction space and the results we get are still approximate solutions. Using the cohomology class equation we see however that the second approximate solutions are so improved that we can apply to them the original perturbation result [6, Theorem 5.3] (or Theorem 9.1 below). Note that as we work under the weaker hypotheses, the first approximate solutions need not satisfy the hypotheses of [6, Theorem 5.3]. Having the second approximate solutions is therefore crucial.

The proof that the cohomology class equation is a necessary condition for the gluing process is similar to the proof that the second approximate solutions satisfy the better estimates.

We begin in §2 with the more precise statements of our results. In §3 we prove that the cohomology class equation is strictly necessary, which is independent of proving the gluing theorem. The latter is done in §§4–9.

In §4 we produce the first approximate solutions, which is essentially the same as in [6, 7]. But the estimates we shall need for these are weaker than those in [6, 7]; for more details see Remark 4.8. The weaker estimates are proved in §5. In §6 we prove the quadratic estimates for the non-linear part of the special Lagrangian equations, which is not very different from the estimates in [6, 7]. In §7 we prove that the linearized operators are uniformly invertible, which is essentially the same as in [3, 11]. In §8 we produce the second approximate solutions, which we perturb in §9 to the true solutions.

Acknowledgements. The author was supported by the grant 21K13788 of the Japan Society for the Promotion of Science.

2 Statements of the Results

We begin by making the definitions we will use to state our results. We define almost Calabi–Yau m -folds in the first place.

Definition 2.1. Let M be a manifold of even dimension $2m$. An *almost Calabi–Yau* structure on M is the data (ω, J, Ω) such that $(M; \omega, J)$ is a Kähler manifold with Kähler form ω and complex structure J , and such that Ω is a nowhere-vanishing holomorphic $(m, 0)$ form on the complex manifold (M, J) . We call $(M; \omega, J, \Omega)$ an *almost Calabi–Yau m -fold*. Notice that there is a C^∞ function $\psi : M \rightarrow (0, \infty)$ defined by

$$\psi^{2m} \omega^m / m! = (i/2)^m (-1)^{m(m-1)/2} \Omega \wedge \bar{\Omega}. \quad (2.1)$$

We call (ω, J, Ω) a *Calabi–Yau* structure, without the word almost, if $\psi : M \rightarrow (0, \infty)$ is constant. In this case ω is a Ricci-flat Kähler form on (M, J) .

We define special Lagrangian submanifolds of almost Calabi–Yau m -folds.

Definition 2.2. Let $(M; \omega, J, \Omega)$ be an almost Calabi–Yau m -fold and $L \subseteq (M, \omega)$ a Lagrangian submanifold. We say that L is *special* if $\text{Im } \Omega|_L = 0$.

We define a Calabi–Yau structure (ω', J', Ω') on \mathbb{C}^m and special Lagrangian cones in $(\mathbb{C}^m; \omega', J', \Omega')$.

Definition 2.3. Let $m \geq 1$ be an integer and z_1, \dots, z_m the complex coordinates of \mathbb{C}^m . Denote by J' the complex structure of \mathbb{C}^m and define on \mathbb{C}^m a Calabi–Yau structure (ω', J', Ω') by $\omega' := \frac{i}{2}(dz_1 \wedge d\bar{z}_1 + \dots + dz_m \wedge d\bar{z}_m)$ and $\Omega' := dz_1 \wedge \dots \wedge dz_m$.

Define $(0, \infty) \times \mathbb{C}^m \rightarrow \mathbb{C}^m$ by $(t, z) \mapsto tz$, by which the multiplicative group $(0, \infty)$ acts upon \mathbb{C}^m . For $t \in (0, \infty)$ we denote for short by $t : \mathbb{C}^m \rightarrow \mathbb{C}^m$ the map $z \mapsto tz$. A *special Lagrangian cone* in $(\mathbb{C}^m; \omega', J', \Omega')$ is a special Lagrangian submanifold $C \subseteq (\mathbb{C}^m; \omega', J', \Omega')$ invariant under the $(0, \infty)$ action and such that $C \cap S^{2m-1}$ is a compact $(m-1)$ dimensional submanifold of the unit sphere $S^{2m-1} \subseteq \mathbb{C}^m$. The *cone metric* on C is the Riemannian metric induced from the embedding $C \subseteq \mathbb{C}^m$.

Remark 2.4. Our C corresponds to Joyce’s C' in [4–8], and our $\bar{C} = C \sqcup \{0\}$ to Joyce’s C . We change the notation because we refer more often to C than to \bar{C} .

We define an asymptotically conical special Lagrangian submanifold $L \subseteq \mathbb{C}^m$ and its cohomology classes $Y(L), Z(L)$.

Definition 2.5. Let $C \subseteq (\mathbb{C}^m; \omega', J', \Omega')$ be a special Lagrangian cone and $L \subseteq (\mathbb{C}^m; \omega', J', \Omega')$ a properly-embedded special Lagrangian submanifold. We say that L approaches *with order $\lambda < 2$ at infinity* the cone C if there exist a compact set $K \subseteq \mathbb{C}^m$ and a diffeomorphism $\phi : C \setminus K \cong L \setminus K$ such that if we denote by $\iota : C \rightarrow \mathbb{C}^m$ the inclusion map then for $k = 0, 1, 2, \dots$ we have

$|\nabla^k(\phi - \iota)| = O(r^{\lambda-1-k})$ where $|\cdot|$ and ∇ are computed with respect to the cone metric on C .

Let $C \subseteq (\mathbb{C}^m; \omega', J', \Omega')$ be a special Lagrangian cone and $L \subseteq (\mathbb{C}^m; \omega', J', \Omega')$ a properly-embedded special Lagrangian submanifold which approaches with order < 2 at infinity the cone C . Using the homology groups with coefficient field \mathbb{R} we make the following definition: $Y(L) \in H^1(C)$ and $Z(L) \in H^{m-1}(C)$ are the respective images of the relative de Rham classes

$$[\omega'|_L] \in H^2(\mathbb{C}^m, L) \cong H^1(L) \text{ and } [\text{Im } \Omega'|_L] \in H^m(\mathbb{C}^m, L) \cong H^{m-1}(L)$$

under the natural maps $H^1(L) \rightarrow H^1(C)$ and $H^{m-1}(L) \rightarrow H^{m-1}(C)$.

We define compact special Lagrangian submanifolds with isolated conical singularities.

Definition 2.6. Let $(M; \omega, J, \Omega)$ be an almost Calabi–Yau m -fold and define $\psi : M \rightarrow (0, \infty)$ by (2.1). Let $X \subseteq (M; \omega, J, \Omega)$ be a special Lagrangian submanifold with compact closure $\bar{X} = X \sqcup X^{\text{sing}}$ for some finite set X^{sing} . For each $x \in X^{\text{sing}}$ choose a \mathbb{C} -vector space isomorphism $(T_x M, J|_x) \cong (\mathbb{C}^m, J')$ under which $(\omega|_x, \Omega|_x)$ corresponds to (ω', Ω') . We call \bar{X} a compact special Lagrangian submanifold *with isolated conical singularities* if for each $x \in X^{\text{sing}}$ there exist a special Lagrangian cone $C_x \subseteq (\mathbb{C}^m, \omega', J', \Omega')$, an open neighbourhood U_x of $0 \in \mathbb{C}^m$, and a Darboux embedding $\Upsilon_x : U_x \rightarrow M$ such that $\Upsilon_x(0) = x$, $\Upsilon_x^* \Omega = \psi(x)^m \Omega'$, $\Upsilon_x^* \omega = \omega'$ and the following holds: there exist $\mu > 2$ and a diffeomorphism $\phi : C_x \cap U_x \rightarrow \Upsilon_x^{-1}(X)$ such that if we denote by $\iota : C_x \rightarrow \mathbb{C}^m$ the inclusion map then for $k = 0, 1, 2, \dots$ we have $|\nabla^k(\phi - \iota)| = O(r^{\mu-1-k})$ where $|\cdot|, \nabla$ are computed with respect to the cone metric on C_x .

We call C_x the *tangent cone* to X at x . We introduce the following notation for short: for each connected component $Y \subseteq X$ and each connected component $C_{xj} \subseteq C_x$ we write $C_{xj} \subseteq Y$ if $\Upsilon_x(C_{xj} \cap U_x) \subseteq Y$.

Remark 2.7. Denote by $H_c^*(X) := H_c^*(X, \mathbb{R})$ the compact-support cohomology group. Since X is outside a compact set diffeomorphic to $\bigsqcup_{x \in X^{\text{sing}}} C_x$ we get a long exact sequence $\dots \rightarrow H_c^*(X) \rightarrow H^*(X) \rightarrow \bigoplus_{x \in X^{\text{sing}}} H^*(C_x) \rightarrow \dots$. Since $m \geq 3$ it follows also that for $q = 2, 3, 4, \dots$ there are natural \mathbb{R} -vector space isomorphisms $H_c^q(X) \cong H^q(\bar{X}, X^{\text{sing}}) \cong H^q(\bar{X})$.

We state now the hypothesis under which we carry out the gluing process.

Hypothesis 2.8. Let $m \geq 3$ be an integer and M a C^∞ manifold of dimension $2m$. Let $n \geq 0$ be an integer and $(\omega^s, J^s, \Omega^s)_{s \in \mathbb{R}^n}$ a C^∞ family of almost Calabi–Yau structures on M . Let $\bar{X} \subseteq (M; \omega^0, J^0, \Omega^0)$ be a compact special Lagrangian submanifold with isolated conical singularities. Suppose that for each $x \in X^{\text{sing}}$ there exists a properly-embedded special Lagrangian submanifold $L_x \subseteq (\mathbb{C}^m; \omega', J', \Omega')$ which approaches with order ≤ 0 at infinity the cone C_x .

We make a definition we will use shortly.

Definition 2.9. Let $n \geq 0$ be an integer and $\mathcal{G} \subseteq \mathbb{R}^n \times (0, \infty)$ a subset whose closure in $\mathbb{R}^n \times [0, \infty)$ contains $(0, 0)$. We say then that a function $\beta : \mathcal{G} \rightarrow \mathbb{R}$ is *smooth* if every point of \mathcal{G} has an open neighbourhood in $\mathbb{R}^n \times (0, \infty)$ to which β extends smoothly. In the same way we define a smooth function $\mathcal{G} \times N \rightarrow M$ for M, N manifolds. Suppose now that $(N^{st} \subseteq M)_{(s,t) \in \mathcal{G}}$ a family of submanifolds, each N^{st} diffeomorphic to a common manifold N . We say then that the family is *smooth* if there exists a smooth function $\mathcal{G} \times N \rightarrow M$ whose restriction to $\{(s, t)\} \times N$ defines the submanifold $N^{st} \subseteq M$.

We state the condition which will play the central rôle in stating our main results.

Condition 2.10. Under Hypothesis 2.8 there exist a closed subset $\mathcal{G} \subseteq \mathbb{R}^n \times (0, \infty)$ whose closure in $\mathbb{R}^n \times [0, \infty)$ contains $(0, 0)$, and a smooth family $(N^{st} \subseteq M)_{(s,t) \in \mathcal{G}}$ of compact submanifolds with the following three properties.

- (a) Each $N^{st} \subseteq (M; \omega^s, J^s, \Omega^s)$ is a special Lagrangian submanifold.
- (b) As (s, t) tends to $(0, 0)$ the family N^{st} converges in the sense of geometric measure theory to \bar{X} . More precisely, the multiplicity-one varifolds supported on N^{st} converge to the multiplicity-one varifold supported on \bar{X} .
- (c) There exist constants $c, \epsilon > 0$ independent of s, t and such that the following holds. For $x \in X^{\text{sing}}$ let $U_x \subseteq M$ be as in Definition 2.6, which we identify with the corresponding neighbourhood of $0 \in \mathbb{C}^m$. Let $K_x \subseteq \mathbb{C}^m$ be a compact neighbourhood of 0 and denote by $\iota : L_x \cap K_x \rightarrow \mathbb{C}^m$ the inclusion map. Then for every (s, t) close enough to $(0, 0)$ there exists a diffeomorphism $\phi : L_x \cap K_x \rightarrow t^{-1}(N^{st} \cap tK_x)$ such that for $k = 0, 1, 2, \dots$ we have $|\nabla^k(\phi - \iota)| \leq c(|s|^\epsilon + t^\epsilon)$ where $|\cdot|, \nabla$ are computed pointwise with respect to the metric on $L_x \cap K_x$ induced from the Euclidean metric of $K_x \subseteq \mathbb{C}^m$.

Remark 2.11. Suppose that these (a)–(c) hold. Combining them with known results we can control N^{st} as follows. By Allard’s regularity theorem it follows from (a),(b) above that the family $N^{st} \subseteq M$ converges smoothly in every compact set to X as (s, t) tends to $(0, 0)$. So there exists a neighbourhood $\bigsqcup_{x \in X^{\text{sing}}} U_x$ of $X^{\text{sing}} \subseteq M$ such that $N^{st} \setminus \bigsqcup_{x \in X^{\text{sing}}} U_x$ is C^∞ close to $X \setminus \bigsqcup_{x \in X^{\text{sing}}} U_x$. On the other hand, by (c) above we can control N^{st} also in $\bigsqcup_{x \in X^{\text{sing}}} tK_x$. By the proof of [2, Theorem 2.2] we can in fact control N^{st} everywhere; that is, the annulus region $\bigsqcup_{x \in X^{\text{sing}}}(N^{st} \cap U_x \setminus tK_x)$ is C^∞ close to $\bigsqcup_{x \in X^{\text{sing}}}(C_x \cap U_x \setminus tK_x)$ where each C_x is embedded in X as in Definition 2.6.

We make now a definition and state then the result that the cohomology class equations are strictly necessary for the gluing process to be possible.

Definition 2.12. Let $n \geq 0$ be an integer and $\mathcal{G} \subseteq \mathbb{R}^n \times (0, \infty)$ a subset whose closure in $\mathbb{R}^n \times [0, \infty)$ contains $(0, 0)$. We say then that a function $\beta : \mathcal{G} \rightarrow \mathbb{R}$ *decays with power decay at $(0, 0) \in \bar{\mathcal{G}}$* if there exist $\epsilon, c > 0$ such that for every $(s, t) \in \mathcal{G}$ we have $|\beta(s, t)| \leq c(|s|^\epsilon + t^\epsilon)$.

Theorem 2.13. *If Hypothesis 2.8 and Condition 2.10 hold then there exist smooth functions $\beta_x, \gamma_x : \mathcal{G} \rightarrow \mathbb{R}$ decaying with power order at $(0, 0) \in \overline{\mathcal{G}}$ and such that for every $(s, t) \in \mathcal{G}$ the following hold.*

- (i) *The image of $[\omega^s] \in H^2(M)$ under the natural map $H^2(M) \rightarrow H^2(\overline{X}) \cong H_c^2(X)$ agrees with the image of $\bigoplus_{x \in X^{\text{sing}}} t^2(1 + \beta_x)Y(L_x) \in \bigoplus_{x \in X^{\text{sing}}} H^1(C_x)$ under the natural map $\bigoplus_{x \in X^{\text{sing}}} H^1(C_x) \rightarrow H_c^2(X)$.*
- (ii) *Define $\psi^s : M \rightarrow (0, \infty)$ by (2.1) with $(\omega^s, J^s, \Omega^s)$ in place of (ω, J, Ω) . Then for each connected component $Y \subseteq X$ we have*

$$\sum_{C_{xj} \subseteq Y} t^m \psi^s(x)^m (1 + \gamma_x) [C_{xj} \cap S^{2m-1}] \cdot Z(L_x) = [\overline{Y}] \cdot [\text{Im } \Omega^s] \quad (2.2)$$

where the sum on the left-hand side is taken over connected components $C_{xj} \subseteq C_x$ lying in Y after embedded in M , and the right-hand side is the pairing of $[\overline{Y}] \in H_m(M)$ and $[\text{Im } \Omega^s] \in H^m(M)$.

We finally state the gluing theorem under the weaker hypotheses.

Theorem 2.14. *Let Hypothesis 2.8 hold. Denote by N the compact manifold without boundary which we obtain from X and $\bigsqcup_{x \in X^{\text{sing}}} L_x$ after we identify their common ends. Let N be connected.*

Let $\mathcal{G} \subseteq \mathbb{R}^n \times (0, \infty)$ be a subset whose closure in $\mathbb{R}^n \times [0, \infty)$ contains $(0, 0)$. Suppose that for each $x \in X^{\text{sing}}$ we are given two smooth functions $\beta_x, \gamma_x : \mathcal{G} \rightarrow \mathbb{R}$ decaying with power order at $(0, 0) \in \overline{\mathcal{G}}$ and such that every $(s, t) \in \mathcal{G}$ satisfies (i), (ii) of Theorem 2.13. Then for every $\theta > 2$ there exist $t_0 > 0$ and a smooth family $(N^{st} \subseteq M : (s, t) \in \mathcal{G}, |s| \leq t^\theta < t_0^\theta)$ of compact submanifolds satisfying (a)–(c) of Condition 2.10.

3 Proof of Theorem 2.13

Let Hypothesis 2.8 and Condition 2.10 hold. For $x \in X^{\text{sing}}$ choose as in [4, Theorem 3.8] a Weinstein neighbourhood of the tangent cone $C_x \subseteq \mathbb{C}^m$ and a symplectic embedding of it into the cotangent bundle T^*C_x . The difference from the ordinary Weinstein theorem is the group action by the multiplicative group $(0, \infty)$. This acts on C_x and on \mathbb{C}^m , under which the inclusion map $C_x \rightarrow \mathbb{C}^m$ is equivariant. The action $(0, \infty) \curvearrowright C_x$ induces also an $(0, \infty)$ action on T^*C_x ; and in particular, if α is a 1-form on C_x then $t(\text{Graph } \alpha) = \text{Graph}(t^2 t_* \alpha) \subseteq T^*C_x$. We require that the neighbourhoods of $C_x \in \mathbb{C}^m$ and $C_x \subseteq T^*C_x$ are both $(0, \infty)$ invariant and that the diffeomorphism between these is $(0, \infty)$ equivariant.

For $x \in X^{\text{sing}}$ let $\Upsilon_x : U_x \rightarrow M$ be as in Definition 2.6 with $(\omega^0, J^0, \Omega^0)$ in place of (ω, J, Ω) . Following [4, Theorem 4.4] write $X \cap U_x$ as the graph of a 1-form η_x over $C_x \cap U_x$ in its Weinstein neighbourhood. It is proved in [4, Lemma 4.5] that the 1-form η_x is exact. Take a Weinstein neighbourhood of $X \cap U_x$

such that the following holds.

Denote by $\phi : C_x \cap U_x \rightarrow X \cap U_x$ the diffeomorphism assigning to each point $y \in C_x \cap U_x$ the unique point of $X \cap U_x = \text{Graph } \eta_x$ in the fibre over y (in the Weinstein neighbourhood of $C_x \cap U_x$). Then for every 1-form α on $X \cap U_x$ its graph in the Weinstein neighbourhood of $X \cap U_x$ is equal to the graph of $\phi^*\alpha + \eta_x$ in the Weinstein neighbourhood of $C_x \cap U_x$. (3.1)

Following [4, Theorem 4.8] we define a Weinstein neighbourhood of $X \subseteq M$.

Definition 3.1. Define $\psi^s : M \rightarrow (0, \infty)$ by (2.1) with $(\omega^s, J^s, \Omega^s)$ in place of (ω, J, Ω) (which is the same as in Theorem 2.13 (ii)). For each $x \in X^{\text{sing}}$ choose a C^∞ family of \mathbb{C} -vector space isomorphism $(T_x M, J^s|_x) \cong (\mathbb{C}^m, J')$ under which $(\omega^s|_x, \Omega^s|_x)$ corresponds to (ω', Ω') . By a family version of Darboux's theorem we can find, making U_x smaller if we need, a Darboux embedding $\Upsilon_x^s : U_x \rightarrow M$ such that $\Upsilon_x^s(0) = x$, $\Upsilon_x^{s*}\Omega^s = \psi^s(x)^m \Omega'$ and $\Upsilon_x^{s*}\omega^s = \omega'$. We often identify $U_x \subseteq \mathbb{C}^m$ with its image $\Upsilon_x^s(U_x) \subseteq M$.

Choose a finite-dimensional \mathbb{R} -vector space V consisting of closed 2-forms on X supported in the compact set $X \setminus \bigsqcup_{x \in X^{\text{sing}}} U_x$ and such that the natural projection $V \rightarrow H_c^2(X)$ is an \mathbb{R} -vector space isomorphism. Denote by $\nu^s \in V$ the element whose compact-support cohomology class is equal to the image of $[\omega^s]$ under $H^2(M) \rightarrow H^2(\tilde{X}) \cong H_c^2(X)$. Recall then from [4, Theorem 4.8] that there exists an s -dependent neighbourhood W of $X \subseteq M$ such that for each $x \in X^{\text{sing}}$ the pre-image $(\Upsilon_x^s)^{-1}(W)$ agrees with the Weinstein neighbourhood of $X \cap U_x \subseteq \mathbb{C}^m$, and W is diffeomorphic to a neighbourhood of the zero-section $X \subseteq T^*X$ so that if we denote by $\hat{\omega}$ the standard symplectic form of the cotangent bundle T^*X then the Kähler form ω^s corresponds to $\hat{\omega} + \nu^s$. We call W the *Weinstein neighbourhood* of $X \subseteq (M, \omega^s)$.

Consider now the smooth model $L_x \subseteq \mathbb{C}^m$ for $x \in X^{\text{sing}}$.

Definition 3.2. Following [4, Theorem 7.4] let $K_x \subseteq \mathbb{C}^m$ be a compact neighbourhood of $0 \in \mathbb{C}^m$ and write $L_x \setminus K_x$ as the graph of a 1-form χ_x over $C_x \setminus K_x$ in the Weinstein neighbourhood of C_x . It is proved in [4, Proposition 7.6] that the cohomology class $[\chi_x]$ is equal to $Y(L_x)$. Take a Weinstein neighbourhood of $L_x \cap K_x$ such that (3.1) holds with $L_x \setminus K_x$ in place of $X \cap U_x$ and with χ_x in place of η_x . Following [4, Theorem 7.5] we take a Weinstein neighbourhood of $L_x \subseteq (\mathbb{C}^m, \omega')$ which extends that of $L_x \cap K_x$ which we have just defined.

Denote by K_x° the interior of the compact subset $K_x \subseteq \mathbb{C}^m$. This is by definition an open set in \mathbb{C}^m . Making K_x larger if we need, suppose that K_x° is contractible and $L_x \cap K_x^\circ$ diffeomorphic to L_x . Define then the cohomology class $Y(L_x \cap K_x) \in H^1(C_x)$ to be image of the relative de Rham cohomology class $[\omega'] \in H^2(K_x^\circ, L_x \cap K_x^\circ)$ under the composite of the natural maps $H^2(K_x^\circ, L_x \cap K_x^\circ) \cong H^1(L_x \cap K_x^\circ) \cong H^1(L_x) \rightarrow H^1(C_x \cap S^{2m-1}) \cong H^1(C_x)$.

By hypothesis we are given N^{st} as in Condition 2.10. As $t^{-1}(N^{st} \cap tK_x^\circ) \subseteq K_x^\circ$ is a Lagrangian submanifold we can assign to it in the same way as above

the cohomology class $Y(t^{-1}(N^{st} \cap tK_x^\circ))$. Part (c) of Condition 2.10 implies then that $t^{-1}(N^{st} \cap tK_x)$ converges with power orders to $L_x \cap K_x$ so that

$$Y(t^{-1}(N^{st} \cap tK_x)) = (1 + \beta_x)Y(L_x \cap K_x) \quad (3.2)$$

for some $\beta_x = \beta_x(s, t)$ decaying with power orders. But by definition $Y(L_x \cap K_x)$ and $Y(L_x)$ are equal. The equation (3.2) implies therefore (i) of Theorem 2.13.

For each connected component $Y \subseteq X$ there is a unique connected component of $N^{st} \setminus \bigsqcup_{x \in X^{\text{sing}}} tK_x$ that is C^1 close to $Y \setminus \bigsqcup_{x \in X^{\text{sing}}} tK_x$, which we denote by Y^{st} . This Y^{st} is a manifold with boundary, with $\partial Y^{st} \subseteq \bigsqcup_{x \in X^{\text{sing}}} tK_x$. For $x \in X^{\text{sing}}$ let $\Sigma_{xj} \subseteq t^{-1}(\partial Y^{st} \cap tK_x)$ be a connected component, which is a compact $m-1$ manifold diffeomorphic to some connected component $C_{xj} \cap S^{2m-1} \subseteq C_x \cap S^{2m-1}$. Denote by $0^* \Sigma_{xj} \subseteq \mathbb{C}^m$ the union of the line segments between the origin 0 and the points of Σ_{xj} . The union $Y^{st} \cup \bigsqcup_{C_{xj} \subseteq Y} t(0^* \Sigma_{xj})$ defines in M an m -cycle homologous to \bar{Y} . Integrating $\text{Im } \Omega^s$ over this and using $\text{Im } \Omega^s|_{Y^{st}} = 0$ we get

$$[\bar{Y}] \cdot [\text{Im } \Omega^s] = \sum_{C_{xj} \subseteq Y} \int_{t(0^* \Sigma_{xj})} \text{Im } \Omega^s. \quad (3.3)$$

On the other hand, for each $x \in X^{\text{sing}}$ we have

$$\int_{t(0^* \Sigma_{xj})} \text{Im } \Omega^s = t^m \psi^s(x)^m \int_{0^* \Sigma_{xj}} \text{Im } t^{-m} \psi^s(x)^{-m} t^* \Omega^s. \quad (3.4)$$

Notice that $t^{-m} \psi^s(x)^{-m} t^* \Omega^s$ depends smoothly on (s, t) including $(0, 0)$, at which it is equal to Ω' , so that

$$\int_{0^* \Sigma_{xj}} \text{Im } t^{-m} \psi^s(x)^{-m} t^* \Omega^s = \int_{0^* \Sigma_{xj}} \text{Im } \Omega' + O(|s| + t). \quad (3.5)$$

Let Ψ be an $m-1$ form on \mathbb{C}^m with $d\Psi = \text{Im } \Omega'$. Then

$$\int_{0^* \Sigma_{xj}} \text{Im } \Omega' = \int_{\Sigma_{xj}} \Psi. \quad (3.6)$$

Notice that Σ_{xj} converges, as (s, t) tends to $(0, 0)$, to some connected component of $\partial(L_x \cap K_x)$ which we denote by Σ_{xj}^0 . Since Σ_{xj} converges with power orders (as in (c) of Condition 2.10) to Σ_{xj}^0 it follows that

$$\int_{\Sigma_{xj}} \Psi = \int_{\Sigma_{xj}^0} \Psi + O(|s|^\epsilon + t^\epsilon) \quad (3.7)$$

for some $\epsilon > 0$ independent of s, t . More precisely, making K_x larger if we need, we can suppose that the 1-form χ_x in Definition 3.2 is defined also near $\partial(C_x \cap K_x)$. Define a diffeomorphism $\phi : \partial(C_x \cap K_x) \cong \partial(L_x \cap K_x)$ by assigning to each point $y \in \partial(C_x \cap K_x)$ the unique point at which the graph of χ_x intersects the fibre over y . Then $C_{xj} = \phi^{-1}(\Sigma_{xj}^0)$ and

$$\int_{\Sigma_{xj}^0} \Psi = \int_{C_{xj}} \phi^* \Psi. \quad (3.8)$$

Finally, by the definition of $Z(L_x) \in H^{m-1}(C_x) \cong H^{m-1}(C_x \cap S^{2m-1})$ the right-hand side of (3.8) may be identified with the pairing $\sum_{C_{x_j} \subseteq Y} [C_{x_j} \cap S^{2m-1}] \cdot Z(L_x)$. This with (3.3)–(3.8) implies (ii) of Condition 2.10. \square

4 First Approximate Solutions

Let the hypotheses of Theorem 2.14 hold. We work in the circumstances of Definitions 3.1 and 3.2. We begin by introducing the cylinder metric on the cone $C_x \subseteq \mathbb{C}^m$ for $x \in X^{\text{sing}}$.

Definition 4.1. Denote by $r : C_x \rightarrow (0, \infty)$ the Euclidean distance from $0 \in \mathbb{C}^m$, which we call the *radius function* on C_x . The *cylinder metric* on C_x is then the conformal change by r^{-2} of the cone metric on C_x . This is the product metric $(d \log r)^2 + g_x$ where g_x is the induced metric on $C_x \cap S^{2m-1}$ (on which the cone and cylinder metrics induce the same metric).

We say that a smooth function $T_x : C_x \cap U_x \rightarrow \mathbb{R}$ is *of order* $\delta > 0$ if for $k = 0, 1, 2, \dots$ we have $|\nabla^k T_x| = O(r^\delta)$ where $|\cdot|, \nabla$ are computed pointwise on $C_x \cap U_x$ with respect to its cylinder metric. This is equivalent to saying that $|\nabla^k T_x| = O(r^{\delta-k})$ with respect to the cone metric on C_x .

Fix $(s, t) \in \mathcal{G}$. Recall from [4, Theorem 2.19(b)] that there exists on X a 1-form α satisfying the following condition.

- Condition 4.2.**
- (i) There exists $\delta > 0$ independent of s, t and such that for each $x \in X^{\text{sing}}$ there exists on $X \cap U_x \cong C_x \cap U_x$ a 0-form T_x of order $\delta > 0$ and with $\alpha = (1 + \beta_x)Y(L_x) + dT_x$.
 - (ii) Give $(M; \omega, J)$ the Kähler metric and $X \subseteq M$ the induced metric; then $d^*\alpha = 0$.
 - (iii) $d\alpha$ is a compactly supported 2-form on X , whose compact-support cohomology class is the image under the natural map $\bigoplus_{x \in X^{\text{sing}}} H^1(C_x) \rightarrow H_c^2(X)$ of the element $(1 + \beta_x)Y(L_x) \in \bigoplus_{x \in X^{\text{sing}}} H^1(C_x)$.

We prove a lemma we will use shortly.

Lemma 4.3. *For $(s, t) \in \mathcal{G}$ the graph of $t^2\alpha$ in the Weinstein neighbourhood of $X \subseteq (M, \omega^s)$ is a Lagrangian submanifold.*

Proof. Let ν^s be as in Definition 3.1. Since the condition (i) of Theorem 2.13 holds for $(s, t) \in \mathcal{G}$ (as supposed in Theorem 2.14) it follows that the compact-support cohomology class $[\nu^s]$ is the image under the natural map $\bigoplus_{x \in X^{\text{sing}}} H^1(C_x) \rightarrow H_c^2(X)$ of the element $t_x^2 Y(L_x) \in \bigoplus_{x \in X^{\text{sing}}} H^1(C_x)$. The latter is represented by $t^2 d\alpha$, by Condition 4.2 (iii). Since the natural projection $V \rightarrow H_c^2(X)$ is an isomorphism it follows therefore that $\nu^s = t^2 d\alpha$. But the symplectic form in the Weinstein neighbourhood of $X \subseteq (M, \omega^s)$ is equal to $\widehat{\omega} + \nu^s$, which thus vanishes on the graph of $t^2\alpha$. \square

We define a compact connected Lagrangian submanifold $N \subseteq (M, \omega^s)$.

Definition 4.4. We define N by gluing together $\bigsqcup_{x \in X^{\text{sing}}} (L_x \cap K_x)$, $\bigsqcup_{x \in X^{\text{sing}}} (C_x \cap U_x \setminus t_x K_x)$ and $X \setminus \bigsqcup_{x \in X^{\text{sing}}} U_x$. For $x \in X^{\text{sing}}$ put $t_x := \sqrt{1 + \beta_x} t$. Notice that $t_x = O(t)$ because β_x decays to 0. Identify each $L_x \cap K_x$ with $t_x(L_x \cap K_x) \subseteq U_x$ and its image under $\Upsilon_x^s : U_x \rightarrow M$. This is then a Lagrangian submanifold of (M, ω^s) .

We interpolate between $t_x(L_x \cap K_x) \subseteq U_x$ and the graph of $t^2\alpha$ in the Weinstein neighbourhood of $X \setminus \bigsqcup_{x \in X^{\text{sing}}} U_x$. Recall from Hypothesis 2.8 that each L_x approaches with order ≤ 0 at infinity the cone C_x . This implies that for $x \in X^{\text{sing}}$ and $k = 0, 1, 2, \dots$ we have $|\nabla^k \chi_x| = O(r^{-1})$ on $L_x \setminus K_x \cong C_x \setminus K_x$ with respect to the cylinder metric. Identify $Y(L_x) \in H^1(C_x)$ with the harmonic 1-form on the link $C_x \cap S^{2m-1}$ and identify this further with its pull-back under the projection $C_x \rightarrow C_x \cap S^{2m-1}$. We thus regard $Y(L_x)$ as a closed 1-form on C_x . Recall then from [4, Proposition 7.6] that the difference $\chi_x - Y(L_x)$ is an exact 1-form on $C_x \setminus K_x$. Write this as dE_x with E_x a C^∞ function $C_x \setminus K_x \rightarrow \mathbb{R}$ decaying at infinity to 0. By [4, Theorem 7.11(b)] there is in fact a constant $\lambda < 0$ such that for $k = 0, 1, 2, \dots$ we have $|\nabla^k \chi_x| = O(r^{\lambda-k})$ with respect to the cylinder metric on $C_x \setminus K_x$. As X^{sing} is finite, we can suppose that λ is independent of x after we make it smaller if we need. Thus $L_x \setminus K_x$ is the graph of $Y(L_x) + dE_x$, and $t_x(L_x \setminus K_x)$ the graph of $t_x^2 Y(L_x) + t_x^2 dt_{x*} E_x$.

On the other hand, recall from Definition 2.6 that X approaches with order > 2 at each $x \in X^{\text{sing}}$ the cone C_x . So there exists $\mu > 2$ such that for $k = 0, 1, 2, \dots$ we have $|\nabla^k \eta_x| = O(r^{\mu-1})$ where $|\cdot|, \nabla$ are computed pointwise on $X \cap U_x \cong C_x \cap U_x$ with respect to its cylinder metric. As X^{sing} is finite, we can suppose that μ is independent of x after we make it smaller if we need. It is easy to show (as in [4, Lemma 4.5]) that η_x is an exact 1-form dA_x such that for $k = 0, 1, 2, \dots$ we have $|\nabla^k A_x| = O(r^\mu)$ with respect to the cylinder metric on $C_x \cap U_x$. Condition 4.2 (i) and Definition 3.1 imply now that the graph of $t^2\alpha$ in the Weinstein neighbourhood of $X \cap U_x$ is the graph of $t^2(1 + \beta_x)Y(L_x) + t^2 dT_x + dA_x$.

Let $\tau \in (\frac{2}{3}, 1)$ be a constant independent of s, t . For $x \in X^{\text{sing}}$ and $\epsilon > 0$ denote by $B_x(\epsilon) \subseteq U_x$ the ball of Euclidean distance $< \epsilon$ from 0 in \mathbb{C}^m . Let $f_x : U_x \rightarrow [0, 1]$ be a C^∞ function with $f_x = 0$ on $B_x(t^\tau)$ and with $f_x = 1$ on $B_x(2t^\tau)$. Consider the graph of

$$t_x^2 d[(1 - f_x)t_{x*} E_x] + t_x^2 Y(L_x) + t^2 d(f_x T_x) + d(f_x A_x) \quad (4.1)$$

on $C_x \cap U_x \setminus K_x$. This joins smoothly to $t_x(L_x \cap K_x) \subseteq U_x$ and the graph of $t^2\alpha$ in the Weinstein neighbourhood of $X \cap U_x$. As a result we get in (M, ω^s) a compact Lagrangian submanifold

$$N := \bigsqcup_{x \in X^{\text{sing}}} (L_x \cap K_x) \sqcup \bigsqcup_{x \in X^{\text{sing}}} (C_x \cap U_x \setminus t_x K_x) \sqcup \left(X \setminus \bigsqcup_{x \in X^{\text{sing}}} U_x \right). \quad (4.2)$$

The hypothesis in the first paragraph of Theorem 2.14 implies that N is connected.

We define radius functions $r : X \rightarrow (0, \infty)$, $r : \bigsqcup_{x \in X^{\text{sing}}} : L_x \rightarrow (0, \infty)$ and

$r : N \rightarrow (0, \infty)$. We use the same symbol r for short although these are of course different functions.

Definition 4.5. Let $r : X \rightarrow (0, \infty)$ be a smooth function which is nearly constant on $X \setminus \bigsqcup_{x \in X^{\text{sing}}} U_x$ and such that for each $x \in X^{\text{sing}}$ the restriction of $r : X \rightarrow (0, \infty)$ to $X \cap U_x$ is equal to the restriction of $r : C_x \rightarrow (0, \infty)$ to $C_x \cap U_x$. under the diffeomorphism $X \cap U_x \cong C_x \cap U_x$ (corresponding to the 1-form η_x).

For $x \in X^{\text{sing}}$ let $r : L_x \rightarrow (0, \infty)$ be a smooth function which is nearly constant on K_x and agrees outside K_x with the radius function on $C_x \setminus K_x$ under the diffeomorphism $L_x \cong K_x \cong C_x \setminus K_x$ corresponding to the 1-form χ_x . Note that $r : \bigsqcup_{x \in X^{\text{sing}}} C_x \rightarrow (0, \infty)$, $r : X \rightarrow (0, \infty)$ and $r : \bigsqcup_{x \in X^{\text{sing}}} L_x \rightarrow (0, \infty)$ are all independent of s, t .

There is now a unique smooth function $r : N \rightarrow (0, \infty)$, depending smoothly on s, t and such that the following holds: for $x \in X^{\text{sing}}$ the restriction of $r : N \rightarrow (0, \infty)$ to the first piece $t_x(L_x \cap K_x)$ in 4.2 is equal to the restriction of $r : L_x \rightarrow (0, \infty)$ to the same region $t_x(L_x \cap K_x) \subseteq t_x L_x$; for $x \in X^{\text{sing}}$ the restriction of $r : N \rightarrow (0, \infty)$ to the second piece $C_x \cap U_x \setminus t_x K_x$ in (4.2) is equal to the restriction of $r : C_x \rightarrow (0, \infty)$ to the same region $C_x \cap U_x \setminus t_x K_x \subseteq C_x$; and the restriction of $r : N \rightarrow (0, \infty)$ to the last piece $(X \setminus \bigsqcup_{x \in X^{\text{sing}}} U_x)$ in (4.2) is equal to the restriction of $r : X \rightarrow (0, \infty)$ to the same region $(X \setminus \bigsqcup_{x \in X^{\text{sing}}} U_x) \subseteq X$.

We define conical and cylindrical metrics on $X, \bigsqcup_{x \in X^{\text{sing}}} L_x$ and N .

Definition 4.6. For $x \in X^{\text{sing}}$ the *conical metric* on L_x is the Riemannian metric induced from the Euclidean metric on \mathbb{C}^m . The *cylindrical metric* on L_x is its conformal change by $r^{-2} : L_x \rightarrow (0, \infty)$. The *conical metric* on X is the Riemannian metric induced from the Kähler metric of $(M; \omega, J)$. The *cylindrical metric* on X is its conformal change by $r^{-2} : X \rightarrow (0, \infty)$. For each s, t the *conical metric* on N is the Riemannian metric induced from the Kähler metric of $(M; \omega^s, J^s)$. The *cylindrical metric* on N is its conformal change by $r^{-2} : N \rightarrow (0, \infty)$.

We use Hölder norms with exponent $\frac{1}{2} \in (0, 1)$; the choice of the exponent is unimportant and any other constant in $(0, 1)$ will do. Put

$$\kappa := \min\{4 - 4\tau, (\mu - 2)\tau, (2 - \lambda)(1 - \tau), 2 + \tau(\delta - 2)\} > 0. \quad (4.3)$$

Since $\tau < 1$, $\mu > 2$, $\lambda < 0$ and $\nu > 0$ it follows that $\kappa > 2 - 2\tau$. We state now the key estimates for $\text{Im } \Omega^s|_N$.

Proposition 4.7. *There exists $c > 0$ independent of s, t and such that*

$$|r^{-m} \text{Im } \Omega|_N|_{C^{1/2}} \leq cr \text{ on } \bigsqcup_{x \in X^{\text{sing}}} (N \cap B_x(t^\tau)), \quad (4.4)$$

$$|r^{-m} \text{Im } \Omega|_N|_{C^{1/2}} \leq c(|s|r + t^\kappa) \text{ on } \bigsqcup_{x \in X^{\text{sing}}} (B_x(2t^\tau) \setminus B_x(t^\tau)) \text{ and} \quad (4.5)$$

$$|r^{-m} \text{Im } \Omega|_N|_{C^{1/2}} \leq c(|s|r + t^4 r^{-4}) \text{ on } N \setminus \bigsqcup_{x \in X^{\text{sing}}} B_x(2t^\tau) \quad (4.6)$$

where the Hölder norms are computed with respect to the cylindrical metric of N .

Remark 4.8. Proposition 4.7 with $s = 0$ corresponds to [7, Theorem 6.7] (our κ corresponding to Joyce's $\kappa - 1$) except the $L^{\frac{2m}{m+2}}$ estimate in [7, Theorem 6.7]. For the latter estimate we need $m < 6$, as in the computation after [7, (57)]. We do not need such a restriction in Proposition 4.7.

The matter is not whether to use the Hölder norms or the Sobolev norms but whether to use the cylindrical metrics or the conical metrics. Although these two are conformally equivalent, they are in fact parametrized by s, t and the estimates we prove ought to be uniform with respect to them. Using the cylindrical metrics or the conical metrics therefore causes the significant difference.

5 Proof of (4.4)–(4.6)

We use the following lemma which we can prove by straightforward computation:

Lemma 5.1. *Let $(\Psi^s)_{s \in \mathbb{R}^n}$ be a C^∞ family of p -forms on U_x , with $\Psi^s|_0 = 0$. Then there exist $\epsilon, c > 0$ independent of s and such that the following holds: let ξ be a closed 1-form on an open set in $C_x \cap U_x$ with $|\xi|_{C^{3/2}} < \epsilon$ with respect to the cylinder metric on C_x ; then the graph of ξ lies in U_x , and for $s \in \mathbb{R}^n$ with $|s| < \epsilon$ we have*

$$|r^{-p}\Psi^s|_{\text{Graph}\xi}|_{C^{1/2}} \leq cr \text{ at every point of Graph}\xi \quad (5.1)$$

where $|\cdot|_{C^{1/2}}$ is computed with respect to the cylinder metric under the diffeomorphism $\text{Graph}\xi \cong C_x$. \square

Remark 5.2. Here $|\xi|_{C^{3/2}}$ is taken into account because $\Psi^s|_{\text{Graph}\xi}$ is over a fixed point of C' a smooth function of $\xi, \nabla\xi$ whose C^β norms are bounded by $|\xi|_{C^{3/2}}$.

Corollary 5.3. *Let $(\Psi^s)_{s \in \mathbb{R}^n}$ be as in Lemma 5.1 and suppose moreover that $\Psi^0 = 0$ at every point of U_x . Then there exist again $\epsilon, c > 0$ such that the same statement holds with the upper bound $c|s|r$ in place of cr in (5.1).*

Proof. Putting $s = (s_1, \dots, s_n) \in \mathbb{R}^n$ and using Taylor's formula, write $\phi^s = s_1\psi_1^s + \dots + s_n\psi_n^s$. We can then apply Lemma 5.1 to $\psi_1^s, \dots, \psi_n^s$ and the rest is easy. \square

We prove now the estimates (4.4)–(4.6).

Proof of (4.4). Fix $x \in X^{\text{sing}}$. Notice that there is on $K_x \subseteq \mathbb{C}^m$ a C^∞ family $(\Xi^{ts})_{t \in [-1, 1], s \in \mathbb{R}^n}$ of m -forms defined by saying that for $t \in [-1, 1] \setminus \{0\}$ we have $\Xi^{ts} := \text{Im}(t_x^{-m} t_x^s \Omega^s - \psi^s(x)^m \Omega')$ and for $t = 0$ we have $\Xi^{0s} := 0$. This is then smooth with respect to t and there exists a constant $c > 0$ independent of s, t and such that for s, t near $(0, 0)$ we have

$$|r^{-m}\Xi^{ts}|_{L_x}|_{C^{1/2}} \leq c|t| \text{ at every point of } L_x \cap K_x \quad (5.2)$$

On the other hand, since $L_x \subseteq (\mathbb{C}^m; \omega', J', \Omega')$ is a special Lagrangian submanifold it follows that $\Xi^{ts}|_{L_x} = \text{Im}(t_x^{-m} t_x^s \Omega^s)|_{L_x}$. Simple computation shows that

$r^{-m} \operatorname{Im}(t_x^{-m} t_x^* \Omega^s)|_{L_x} = t_x^*(r^{-m} \operatorname{Im} \Omega^s|_{t_x L_x})$. On the other hand, $t_x L_x = N$ in $t_x K_x$. Combining these with (5.2) we see that

$$|t_x^*(r^{-m} \operatorname{Im} \Omega^s|_N)|_{C^{1/2}} \leq c|t| \text{ at every point of } t^{-1}N \cap K_x. \quad (5.3)$$

This rescaled by t_x implies $|r^{-m} \operatorname{Im} \Omega^s|_N|_{C^{1/2}} \leq c|t|$ at every point of $N \cap t_x K_x$. But the radius function r is, at every point of $N \cap t_x K_x$, bounded below by t up to constant. Making c larger if we need, therefore, we have

$$|r^{-m} \operatorname{Im} \Omega^s|_N|_{C^{1/2}} \leq cr \text{ at every point of } N \cap t_x K_x. \quad (5.4)$$

We prove that the same estimate holds also on $N \cap B_x(t^\tau) \setminus t_x K_x$. Applying Lemma 5.1 to $\Psi^s := \operatorname{Im}(\Omega^s - \psi^s(x)^m \Omega')$ we see that making c larger if we need, we have

$$|r^{-m} \Psi^s|_N|_{C^{1/2}} \leq cr \text{ at every point of } N \cap B_x(t^\tau) \setminus t_x K_x. \quad (5.5)$$

Since $N \cap B_x(t^\tau) = t_x L_x \cap B_x(t^\tau) \subseteq (\mathbb{C}^m; \omega', J', \Omega')$ is a special Lagrangian submanifold it follows that $\Psi^s|_N = \operatorname{Im} \Omega^s|_N$. The estimate (5.5) implies therefore that $|r^{-m} \operatorname{Im} \Omega^s|_N|_{C^{1/2}} \leq cr$ at every point of $N \cap B_x(t^\tau) \setminus t_x K_x$, completing the proof. \square

Proof of (4.5). If ξ is a C^1 small 1-form on X then we can write $\operatorname{Im} \Omega|_{\operatorname{Graph} \xi} =: P_X(\xi, \nabla \xi) \operatorname{dvol}$ where P_X is a ξ -independent smooth function $T^*X \oplus (T^*X)^{\otimes 2} \rightarrow \mathbb{R}$, and dvol is the volume form of the conical metric on X . Recall from [5, Proposition 6.3] that

$$P_X(\xi, \nabla \xi) = -d^*(\psi^m \xi) + O(r^{-2}|\xi|^2) + O(|\nabla \xi|^2) \quad (5.6)$$

where $|\cdot|, \nabla$ and the O terms are computed with respect to the conical metric on X . If we replace the conical metric by the cylindrical metric then $|\xi|$ becomes $r^{-1}|\xi|$, $|\nabla \xi|$ becomes $r^{-2}|\nabla \xi|$ and the formula becomes

$$P_X(\xi, \nabla \xi) = -d^*(\psi^m \xi) + r^{-4}O(|\xi|^2) + r^{-4}O(|\nabla \xi|^2) \quad (5.7)$$

where $|\cdot|, \nabla$ and the O terms are computed now with respect to the cylindrical metric on X . The same result [5, Proposition 6.3] computes also the derivatives of (5.6) and in particular we can estimate the Hölder norm of $P_X(\xi, \nabla \xi)$. If we use again the cylindrical metric then we have

$$|P_X(\xi, \nabla \xi)|_{C^{1/2}} = |d^*(\psi^m \xi)|_{C^{1/2}} + r^{-4}O(|\xi|^2) + r^{-4}O(|\nabla \xi|^2); \quad (5.8)$$

that is, these $|\cdot|, \nabla$ and O terms are computed with respect to the cylindrical metric on X .

Fix $x \in X^{\operatorname{sing}}$. The estimate (4.5) is concerned with the region $B_x(2t^\tau) \setminus B_x(t^\tau)$ in which $N \cap B_x(2t^\tau) \setminus B_x(t^\tau)$ is the graph of a 1-form ζ on $C_x \cap B_x(2t^\tau) \setminus B_x(t^\tau)$ such that $\zeta - t^2 \alpha$ is an exact 1-form dF . The 0-form F is made from $A_x, t_x^2 t_x^* E_x$ and $t^2 T_x$ using cut-off functions.

We compute the right-hand side of (5.8). For $\xi = t^2\alpha$ the d^* term vanishes and the last two quadratic terms on (5.8) contribute $r^{-4}t^4$. But $r \geq t^\tau$ so this is bounded above by $t^{4-4\tau} \leq t^\kappa$. For $\xi = dA_x, t_x^2 t_{x*} dE_x$ and $t^2 dT_x$ the dominant term on the right-hand side of (5.8) is the d^* term. Since d^* is computed with respect to the conical metric it follows that there exists $c > 0$ independent of s, t and such that

$$|d^*(\psi^m \xi)|_{C^{1/2}} \leq cr^{-2} |\xi|_{C^{3/2}} \text{ at every point of } C_x \cap B_x(2t^\tau) \setminus B_x(t^\tau). \quad (5.9)$$

where the Hölder norms are computed with respect to the cylinder metric. Notice that

$$\begin{aligned} \xi = dA_x = O(r^\mu) &\Rightarrow r^{-2} |\xi|_{C^{3/2}} = r^{\mu-2} \leq 2^{\mu-2} t^{\tau(\mu-2)} \leq 2^{\mu-2} t^\kappa, \\ \xi = t_x^2 t_{x*} dE_x = t^{2-\lambda} O(r^\lambda) &\Rightarrow r^{-2} |\xi|_{C^{3/2}} = t^{2-\lambda} r^{\lambda-2} \leq t^{(2-\lambda)(1-\tau)} \leq t^\kappa, \\ \xi = t^2 dT_x = t^2 O(r^\delta) &\Rightarrow r^{-2} |\xi|_{C^{3/2}} = t^2 r^{\delta-2} \leq t^{2+\tau(\delta-2)} \leq t^\kappa. \end{aligned}$$

It is easy to show that the cut-off functions which piece together $A_x, t_x^2 t_{x*} E_x$ and $t^2 T_x$ have only negligible contributions, so the 1-form dF_x contributes t^κ up to constant. We have thus

$$|P_X(\zeta, \nabla \zeta)|_{C^{1/2}} \leq ct^\kappa \text{ at every point of } B_x(2t^\tau) \setminus B_x(t^\tau) \quad (5.10)$$

after we make c larger if we need.

Since $r^{-m} \text{Im} \Omega|_N = P_X(\zeta, \nabla \zeta) r^{-m} \text{dvol}$ and since $r^{-m} \text{dvol}$ is the volume form of the cylinder metric it follows that

$$|r^{-m} \text{Im} \Omega|_N|_{C^{1/2}} = |P_X(\zeta, \nabla \zeta) r^{-m} \text{dvol}|_{C^{1/2}} = |P_X(\zeta, \nabla \zeta)|_{C^{1/2}}. \quad (5.11)$$

Applying Corollary 5.3 to $\text{Im}(\Omega^s - \Omega)$ we get

$$|r^{-m} \text{Im}(\Omega^s - \Omega)|_N|_{C^{1/2}} \leq |s|r \text{ at every point of } B_x(2t^\tau) \setminus B_x(t^\tau). \quad (5.12)$$

Combining (5.10)–(5.12) we get (4.5). \square

Proof of (4.6). The same computation as in (5.11) shows that

$$|r^{-m} \text{Im} \Omega|_N|_{C^{1/2}} = |P_X(t^2 \alpha, t^2 \nabla \alpha)| \text{ at every point of } N \setminus \bigsqcup_{x \in X^{\text{sing}}} B_x(2t^\tau). \quad (5.13)$$

Applying (5.8) to $\xi = t^2 \alpha$ we get

$$|P_X(t^2 \alpha, t^2 \nabla \alpha)| = r^{-4} O(t^4 |\alpha|^2) + r^{-4} O(t^4 |\nabla \alpha|^2) = t^4 O(r^{-4}). \quad (5.14)$$

On the other hand, we show now that

$$|r^{-m} \text{Im}(\Omega^s - \Omega)|_N|_{C^{1/2}} \leq |s|r \text{ at every point of } N \setminus \bigsqcup_{x \in X^{\text{sing}}} B_x(2t^\tau). \quad (5.15)$$

As the proof of (5.12) is valid also on $N \cap \bigsqcup_{x \in X^{\text{sing}}} (U_x \setminus B_x(2t^\tau))$, the estimate (5.15) does hold on this region. Since $N \setminus \bigsqcup_{x \in X^{\text{sing}}} U_x$ is compact and t -independent it follows that

$$|r^{-m} \text{Im}(\Omega^s - \Omega^0)|_N|_{C^{1/2}} \leq |s| \text{ at every point of } N \setminus \bigsqcup_{x \in X^{\text{sing}}} U_x. \quad (5.16)$$

But $r : X \rightarrow (0, \infty)$ is bounded below on this region, so (5.16) implies that (5.15) holds on the same region. Thus (5.15) holds everywhere on $N \setminus \bigsqcup_{x \in X^{\text{sing}}} B_x(2t^\tau)$. This combined with (5.13) and (5.14) implies (4.6). \square

6 Quadratic Estimates

Since $\kappa > 2 - 2\tau$ and $\tau \in (\frac{2}{3}, 1)$ it follows that

$$\nu' := \max \left\{ \frac{3\tau - 2}{\tau - 2}, \frac{2 - 2\tau - \kappa}{2 - \tau}, \frac{2\tau - 2}{2 - \tau} \right\} < 0. \quad (6.1)$$

Fix $\nu \in (\nu', 0)$. Making ν larger if we need, suppose also that $2 - 2\nu < \theta$ where θ is the constant > 2 given in Theorem 2.14. Give $C^{5/2}(N, \mathbb{R})$ a weighted Hölder norm $|\bullet|_\nu$ with weight $r^{-\nu}$; that is, for $u \in C^{5/2}(N, \mathbb{R})$ define $|u|_\nu := |r^{-\nu}u|_{C^{5/2}}$ where the right-hand side is computed with respect to the cylindrical metric on N . In the same way, give $C^{1/2}(N, \mathbb{R})$ the weighted Hölder norm $|\bullet|_{\nu-2}$ with weight $r^{\nu-2}$. We prove then a corollary of Proposition 4.7.

Corollary 6.1. *There exists $c > 0$ independent of s, t and such that*

$$|r^{-n} \operatorname{Im} \Omega|_N|_{\nu-2} \leq c|s| + t^\theta \text{ at every point of } N. \quad (6.2)$$

Proof. From (4.4)–(4.6) we get

$$|r^{-n} \operatorname{Im} \Omega|_N|_{\nu-2} \leq cr^{2-\nu}r = r^{3-\nu} \leq ct^{\tau(3-\nu)}, \quad (6.3)$$

$$|r^{-n} \operatorname{Im} \Omega|_N|_{\nu-2} \leq cr^{2-\nu}(|s|t^\tau + t^\kappa) \leq c|s| + ct^{\tau(2-\nu)+\kappa} \text{ and} \quad (6.4)$$

$$|r^{-n} \operatorname{Im} \Omega|_N|_{\nu-2} \leq cr^{2-\nu}(|s|r + t^4r^{-4}) \leq c|s| + ct^{4-\tau(2+\nu)} \quad (6.5)$$

respectively. As $\nu \in (\nu', 0)$ it follows from (6.1) that the respective powers of t in the last terms of (6.3)–(6.5) are all $> 2 - 2\nu$. \square

We define now a neighbourhood of $N \subseteq (M, \omega^s)$ by gluing together the respective Weinstein neighbourhoods of $\bigsqcup_{x \in X^{\text{sing}}} t(L_x \cap K_x)$, $\bigsqcup_{x \in X^{\text{sing}}} (C_x \cap U_x \setminus t_x K_x)$ and $X \setminus \bigsqcup_{x \in X^{\text{sing}}} U_x$ in (4.2). We call this the *Weinstein* neighbourhood of $N \subseteq (M, \omega^s)$. Using it we define the graph of a C^1 small 1-form on N . If ξ is such a 1-form then we define $P\xi \in C^0(N, \mathbb{R})$ by $(P\xi)d\text{vol} = \operatorname{Im} \Omega^s|_{\text{Graph } \xi}$, using the diffeomorphism $\text{Graph } \xi \cong N$, where $d\text{vol}$ is the volume form of the conical metric on N . If ξ is moreover $C^{3/2}$ then $P\xi \in C^{1/2}(N, \mathbb{R})$.

Thus P is an operator from an open subset of $C^{3/2}(T^*N)$ to the whole space $C^{1/2}(N, \mathbb{R})$. This is a smooth map and its linearization at $0 \in C^{3/2}(T^*N)$ is well defined. Denote the latter by Δ and put $Q := P - P0 - \Delta$. We prove then a quadratic estimate for Q . Give $C^{3/2}(T^*N)$ the weighted Hölder norm $|\bullet|_\nu$ with weight $r^{-\nu}$. Notice that if $|\xi|_\nu = O(t^{\theta+\nu})$ then $|\xi|_{C^{3/2}} = O(t^\theta)$ so the 1-form ξ and the rescaled 1-form $t^{-2}\xi$ are both small enough.

Proposition 6.2. *For $c_0 > 0$ independent of s, t there exists $c_1 > 0$ independent of s, t and such that for $\phi, \phi' \in C^{3/2}(T^*N)$ with $|\phi|_\nu + |\phi'|_\nu \leq c_0 t^{\theta+\nu}$ we have*

$$|Q\phi' - Q\phi|_{C^{1/2}} \leq c_1 r^{-4} |\phi' - \phi|_{C^{3/2}} (|\phi|_{C^{3/2}} + |\phi'|_{C^{3/2}}) \text{ at every point of } N \quad (6.6)$$

where r is the radius function on N .

Proof. Put $V := T^*N \oplus T^*N^{\otimes 2}$, a vector bundle over N . Re-write $P\phi$ as $P(\phi, \nabla\phi)$ where the latter P is a ϕ -independent smooth function $V \rightarrow \mathbb{R}$. Denote then by $\partial P : V^* \otimes V \rightarrow \mathbb{R}$ the partial derivative of P in the fibres of V (for which we do not have to choose a connection on V). Define also the second derivative $\partial^2 P : V^* \otimes V^* \otimes V \rightarrow \mathbb{R}$. For $a \in \mathbb{R}$ put $\phi^a := \phi + a(\phi' - \phi)$. Put also $\Phi^a := (\phi^a, \nabla\phi^a)$, which is a section of V . Put $\Phi = \Phi^0$ and $\Phi' := \Phi^1$. Taylor's theorem implies then

$$Q\phi' - Q\phi = (\Phi' - \Phi) \lrcorner \int_0^1 \int_0^1 \Phi^a \lrcorner \partial^2 P(ba\Phi^a) da db. \quad (6.7)$$

Taking the $C^{1/2}$ norms we get

$$|Q\phi' - Q\phi|_{C^{1/2}} \leq |\Phi' - \Phi|_{C^{1/2}} \int_0^1 \int_0^1 |\Phi^a|_{C^{1/2}} |\partial^2 P(ba\Phi^a)|_{C^{1/2}} da db. \quad (6.8)$$

On the other hand,

$$|\Phi' - \Phi|_{C^{1/2}} \leq |\phi' - \phi|_{C^{1/2}} + |\nabla\phi' - \nabla\phi|_{C^{1/2}} \leq |\phi' - \phi| \quad (6.9)$$

and we can compute $|\Phi^a|_{C^{1/2}}$ in the same way. In a way similar to proving (5.8) we can show that $|\partial^2 P(ba\Phi^a)|_{C^{1/2}} = O(r^{-4})$. So (6.8) implies (6.6). \square

Passing to the weighted norms we get

Corollary 6.3. *For $c_0 > 0$ independent of s, t there exists $c_1 > 0$ independent of s, t and such that for $\phi, \phi' \in C^{3/2}(T^*N)$ with $|\phi|_\nu + |\phi'|_\nu \leq c_0 t^{\theta+\nu}$ we have*

$$|Q\phi' - Q\phi|_{\nu-2} \leq c_1 t^{\nu-2} |\phi' - \phi|_\nu (|\phi|_\nu + |\phi'|_\nu). \quad (6.10)$$

Proof. After we pass to the weighted norms, the factor r^{-4} in (6.6) becomes $r^{-4} r^\nu r^\nu r^{-\nu+2} = r^{\nu-2}$. But the radius function $r : N \rightarrow (0, \infty)$ is bounded below by t up to positive constant, so we have the factor $t^{\nu-2}$ as above. \square

7 Uniform Invertibility

For each connected component $Y \subseteq X$ choose a C^∞ function $E_Y : X \rightarrow \mathbb{R}$ supported in $Y \setminus \bigsqcup_{x \in X^{\text{sing}}} U_x$ and with $\int_Y E_Y d\text{vol} \neq 0$ where $d\text{vol}$ is the volume form of the conical metric on Y . Regard this as an \mathbb{R} -linear map $\mathbb{R} \rightarrow C^{1/2}(N, \mathbb{R})$ which maps 1 to E_Y . Varying $Y \in \pi_0 X$ we get a basis of the \mathbb{R} -vector space $H^0(X, \mathbb{R})$ and the family $(E_Y)_{Y \in \pi_0 X}$ defines then an \mathbb{R} -linear map $H^0(X, \mathbb{R}) \rightarrow C^{1/2}(N, \mathbb{R})$ which we denote by E . Consider now the operator

$$\Delta \oplus E : C^{5/2}(N, \mathbb{R}) \oplus H^0(X, \mathbb{R}) \rightarrow C^{1/2}(N, \mathbb{R}). \quad (7.1)$$

The E factor compensates for the one-dimensional cokernel of $\Delta : C^{5/2}(N, \mathbb{R}) \rightarrow C^{1/2}(N, \mathbb{R})$ so that $\Delta \oplus E$ is surjective. We prove

Proposition 7.1. *There exists an (s, t) uniformly bounded right inverse to (7.1) with respect to the weighted norms on the Hölder spaces.*

The idea of the proof is that we can decompose the operator (7.1) into two pieces. One depends only on $(L_x)_{x \in X^{\text{sing}}}$ and the other only on X . Both are uniformly invertible, from which we deduce that the original operator is invertible. The heart of the proof is therefore to make a reasonable definition of the decomposition. This is rather long but more or less straightforward. We do it following [1, §7.2] in outline, although the background geometry is different. We begin by introducing the cut-off functions with which we will glue together the operators we have defined.

Definition 7.2. Let $F_X : N \rightarrow [0, 1]$ and $F_L : N \rightarrow [0, 1]$ be a partition of unity on N subordinate to $N \setminus \bigsqcup_{x \in X^{\text{sing}}} B_x(t^\tau)$ and $N \cap \bigsqcup_{x \in X^{\text{sing}}} B_x(2t^\tau)$, F_X supported on the former and F_L on the latter. For each connected component $Y \subseteq X$ put $F_Y := F_X|_Y$.

Choose a constant $\rho < \tau$ independent of s, t . Put $L := \bigsqcup_{x \in X^{\text{sing}}} L_x$ and let $G_L : L \rightarrow [0, 1]$ be a cut-off function supported on $L \cap \bigsqcup_{x \in X^{\text{sing}}} B_x(t^\rho)$, with $G_L = 1$ on $L_x \cap \bigsqcup_{x \in X^{\text{sing}}} B_x(2t^\tau)$ and satisfying the following condition: there exists $c > 0$ independent of s, t and such that at every point of L we have $|G_L|_{C^{5/2}} \leq c(-\log t)^{-1}$ where the Hölder norm is computed with respect to the cylindrical metric on L .

We do something similar on X . Choose a constant $\sigma > \tau$ independent of s, t . Let $G_X : X \rightarrow [0, 1]$ be a cut-off function supported on $X \setminus \bigsqcup_{x \in X^{\text{sing}}} B_x(t^\sigma)$, with $G_X = 1$ on $X \setminus \bigsqcup_{x \in X^{\text{sing}}} B_x(t^\tau)$ and satisfying the following condition: there exists $c > 0$ independent of s, t and such that at every point of X we have $|G_X|_{C^{5/2}} \leq c(-\log t)^{-1}$ where the Hölder norm is computed with respect to the cylindrical metric on X . For each connected component $Y \subseteq X$ put $G_Y := G_X|_Y$.

We define two operators $\Delta_L, D_L : C_\nu^{5/2}(L, \mathbb{R}) \rightarrow C_{\nu-2}^{1/2}(L, \mathbb{R})$.

Definition 7.3. Denote by g' the Euclidean metric of \mathbb{C}^m , and by the same g' the induced metric on $L = \bigsqcup_{x \in X^{\text{sing}}} L_x \subseteq \bigsqcup_{x \in X^{\text{sing}}} \mathbb{C}^m$. Define $\Delta_L : C_\nu^{5/2}(L, \mathbb{R}) \rightarrow C_{\nu-2}^{1/2}(L, \mathbb{R})$ by $\Delta_L := -d^{*'}d$ where $d^{*'}$ is the formal adjoint of d computed with respect to g' . This is a bounded operator with respect to the weighted Hölder norms. It is invertible, by [10, Theorem 6.2.15].

We choose on L a Riemannian metric g^{st} depending smoothly on s, t and such that the restriction to $L_x \setminus B_x(t_x^{-1}t^\rho)$ of g^{st} agrees with g' , and the restriction to each $L_x \cap B_x(2t_x^{-1}t^\tau)$ of g^{st} agrees with the metric induced by the embedding $L_x \cap B_x(2t_x^{-1}t^\tau) \cong t_x L_x \cap B_x(2t^\tau) \subseteq U_x$ where $U_x \subseteq M$ is given the Kähler metric corresponding to (ω^s, J^s) . Choose a smooth function $\psi^{st} : L \rightarrow (0, \infty)$ depending smoothly on s, t , with $\psi^{st} = 1$ on each $L_x \setminus B_x(t_x^{-1}t^\rho)$, and with $\psi^{st} = t_x^* \psi^s$ on each $L_x \cap B_x(2t_x^{-1}t^\tau)$. Define $D_L : C_\nu^{5/2}(L, \mathbb{R}) \rightarrow C_{\nu-2}^{1/2}(L, \mathbb{R})$ by $D_L := -d^{*st}[(\psi^{st})^m d]$ where d^{*st} is the formal adjoint of d computed with respect to g^{st} .

We show that D_L is uniformly invertible.

Lemma 7.4. $D_L : C_\nu^{5/2}(L, \mathbb{R}) \rightarrow C_{\nu-2}^{1/2}(L, \mathbb{R})$ is invertible and its inverse is bounded uniformly with respect s, t .

Proof. Computation similar to the proof of (4.4) show that there exists $c > 0$ independent of s, t and such that

$$|g^{st} - g'|_{C^{3/2}} \leq ctr \leq 2ct^\rho \text{ at every point of } L \quad (7.2)$$

where the $C^{3/2}$ norm is computed with respect to the cylindrical metric on L . Denote by ∇^{st}, ∇' the respective Levi-Civita connections of g^{st}, g' . The estimate (7.2) implies then that there exists $c > 0$ independent of s, t and such that for $\phi \in C^{3/2}(T^*L)$ we have

$$|\nabla^{st}\phi - \nabla'\phi|_{C^{1/2}} \leq ct^\rho|\phi|_{C^{3/2}} \text{ at every point of } L. \quad (7.3)$$

In general, the operator d^* of a Riemannian metric g_{ab} is of the form $-g^{ab}\nabla_b$ in the index notation. In the current circumstances the g^{ab} term contributes r^{-2} so that

$$|d^{*st}\phi - d^{*t}\phi|_{C^{1/2}} \leq cr^{-2}t^\rho|\phi|_{C^{3/2}} \text{ at every point of } L. \quad (7.4)$$

Hence making c larger if we need, we find that for $u \in C^{5/2}(L, \mathbb{R})$ we have

$$|d^{*st}[(\psi^{st})^m du] - d^{*t}[(\psi^{st})^m du]|_{C^{1/2}} \leq cr^{-2}t^\rho|u|_{C^{5/2}}. \quad (7.5)$$

Since $\psi^{00} = 1$ it follows that making c larger if we need, we have

$$|D_L u - \Delta_L u|_{C^{1/2}} \leq cr^{-2}(|s| + t^\rho)|u|_{C^{5/2}}. \quad (7.6)$$

Passing to the weighted norms we get

$$|D_L u - \Delta_L u|_{\nu-2} \leq c(|s| + t^\rho)|u|_\nu. \quad (7.7)$$

Recall now from [10, Theorem 6.2.15] that the operator $\Delta_L : C_\nu^{5/2}(L, \mathbb{R}) \rightarrow C_{\nu-2}^{1/2}(L, \mathbb{R})$ is invertible. Since g^{st} and ψ^{st} depend smoothly on s, t it follows that the operator $D_L : C_\nu^{5/2}(L, \mathbb{R}) \rightarrow C_{\nu-2}^{1/2}(L, \mathbb{R})$ is bounded uniformly with respect to s, t (where the operator norm is computed with respect to the weighted norms). The estimate (7.7) implies therefore that that the operator $D_L \Delta_L^{-1} - \text{id} : C_{\nu-2}^{1/2}(L, \mathbb{R}) \rightarrow C_{\nu-2}^{1/2}(L, \mathbb{R})$ has operator norm $\leq c(|s| + t^\rho) \ll 1$ with respect to the $C_{\nu-2}^{1/2}$ norm. Thus $D_L \Delta_L^{-1}$ is invertible and its inverse is bounded uniformly with respect to s, t . On the other hand, Δ_L has nothing to do with s, t and is therefore bounded uniformly with respect to s, t . Accordingly so is $\Delta_L^{-1}(D_L \Delta_L^{-1})^{-1}$, which is a right inverse to D_L . Changing the order of D_L and Δ_L^{-1} we see also that $\Delta_L^{-1} D_L$ is invertible and that D_L has a left inverse. The latter must agree with the right inverse, which completes the proof. \square

We define two other operators $\Delta_Y \oplus E_Y$ and $D_Y \oplus E_Y$.

Definition 7.5. Recall that $N \setminus \bigsqcup_{x \in X^{\text{sing}}} t_x K_x$ lies in the Weinstein neighbourhood of $X \subseteq (M, \omega^s)$ and write this as the graph of a 1-form ξ on $X \setminus \bigsqcup_{x \in X^{\text{sing}}} t_x K_x$. Introduce on X a 1-form ξ'' which agrees with ξ on $X \setminus \bigsqcup_{x \in X^{\text{sing}}} B_x(t^\sigma)$ and vanishes on $\bigsqcup_{x \in X^{\text{sing}}} B_x(2t^\sigma)$. Fix a connected component $Y \subseteq X$. Define $\Delta_Y : C_\nu^{5/2}(Y, \mathbb{R}) \rightarrow C_{\nu-2}^{1/2}(Y, \mathbb{R})$ by $\Delta_Y = -d^*[(\psi^0)^m d]$ where d^* is computed with respect to the conical metric on $Y \subseteq X$. Since $E_Y : Y \rightarrow \mathbb{R}$ is compactly supported it follows that this is an element of the weighted space $C_{\nu-2}^{1/2}(N, \mathbb{R})$ and defines therefore an \mathbb{R} -linear map $\mathbb{R} \rightarrow C_{\nu-2}^{1/2}(N, \mathbb{R})$ which maps 1 to E_Y . We denote the latter also by E_Y . Combining it with Δ_Y we get an operator

$$\Delta_Y \oplus E_Y : C_\nu^{5/2}(Y, \mathbb{R}) \oplus \mathbb{R} \rightarrow C_{\nu-2}^{1/2}(Y, \mathbb{R}). \quad (7.8)$$

Denote by g'' the Riemannian metric on X induced by the embedding $X \cong \text{Graph } \xi'' \subseteq M$ where M is given the Kähler metric corresponding to (ω^s, J^s) . Define $D_Y : C_\nu^{5/2}(Y, \mathbb{R}) \rightarrow C_{\nu-2}^{1/2}(Y, \mathbb{R})$ by $D_Y := -d^{*''}[(\psi^s)^m d]$ where $d^{*''}$ is computed with respect to g'' .

We show that (7.8) has a uniformly bounded right inverse.

Lemma 7.6. *Using the $C_\nu^{5/2}$ norm on $C_\nu^{5/2}(Y, \mathbb{R})$ and the Euclidean norm on \mathbb{R} give $C_\nu^{5/2}(Y, \mathbb{R}) \oplus \mathbb{R}$ the product norm. The operator (7.8) has then a right inverse which is uniformly bounded with respect to s, t .*

Proof. Denote by g the conical metric on $Y \subseteq X$. We show that there exists $c > 0$ independent of s, t and for which

$$|g'' - g|_{C^{3/2}} \leq ct^{2-2\sigma} \text{ at every point of } X \quad (7.9)$$

where the $C^{3/2}$ norm is computed with respect to the cylindrical metric on X . It is clear that this holds on $\bigsqcup_{x \in X^{\text{sing}}} B_x(t^\sigma)$ because $g'' = g$ there. On $X \setminus \bigsqcup_{x \in X^{\text{sing}}} B_x(2t^\sigma)$ we have $\xi = t^2 \alpha$ which implies the stronger estimate $|g'' - g|_{C^{3/2}} = O(t^2)$. On each $B_x(2t^\sigma) \setminus B_x(t^\sigma)$ the 1-form ξ is made from $dA_x, t_x^2 t_{x*} dE_x, dT_x$ and $t^2 \alpha$. Computation similar to the proof of (4.5) shows that the dominant contribution is that of $t^2 \alpha$, which is $r^{-2} O(t^2)$, including the rescale factor r^{-2} . This with $r \geq t^\sigma$ implies (7.9).

If we denote by ∇'', ∇ the Levi-Civita connection of g'' then we find from (7.9) a constant $c > 0$ independent of s, t and such that for $\phi \in C^{3/2}(T^*Y)$ we have

$$|\nabla'' \phi - \nabla \phi|_{C^{1/2}} \leq ct^{2-2\sigma} |\phi|_{C^{3/2}} \text{ at every point of } Y. \quad (7.10)$$

Computing the d^* operators as in (7.4) we get

$$|d^{*''} \phi - d^* \phi|_{C^{1/2}} \leq cr^{-2} t^{2-2\sigma} |\phi|_{C^{3/2}} \text{ at every point of } Y. \quad (7.11)$$

Hence making c larger if we need, we find that for $u \in C^{5/2}(Y, \mathbb{R})$ we have

$$|d^{*''}[(\psi^s)^m du] - d^*[(\psi^s)^m du]|_{C^{1/2}} \leq cr^{-2} t^{2-2\sigma} |u|_{C^{5/2}}. \quad (7.12)$$

Since $\psi^0 = \psi$ it follows that making c larger if we need, we have

$$|D_Y u - \Delta_Y u|_{C^{1/2}} \leq cr^{-2}(|s| + t^{2-2\sigma})|u|_{C^{5/2}}. \quad (7.13)$$

Passing to the weighted norms we get $|D_Y u - \Delta_Y u|_{\nu-2} \leq c(|s| + t^{2-2\sigma})|u|_{\nu}$; that is,

$$D_Y - \Delta_Y : C_{\nu}^{5/2}(Y, \mathbb{R}) \rightarrow C_{\nu-2}^{1/2}(Y, \mathbb{R}) \text{ has operator norm } \leq c(|s| + t^{2-2\sigma}) \quad (7.14)$$

with respect to the weighted Hölder norms.

Recall now from [4, Theorems 2.14 and 2.16(b)] that the Sobolev space version of (7.8) is surjective, and from [10, Theorem 6.9] that the Hölder version holds too; that is, (7.8) itself is surjective. So there exists a bounded linear operator $\Theta : C_{\nu-2}^{1/2}(Y, \mathbb{R}) \rightarrow C_{\nu}^{5/2}(Y, \mathbb{R}) \oplus \mathbb{R}$ which is a right inverse to (7.8). Hence it follows by (7.14) that the operator $(D_Y \oplus E_Y)\Theta - \text{id} : C_{\nu-2}^{1/2}(Y, \mathbb{R}) \rightarrow C_{\nu-2}^{1/2}(Y, \mathbb{R})$ has operator norm less than $c(|s| + t^{2-2\sigma}) \ll 1$. So $(D_Y \oplus E_Y)\Theta$ is invertible and its inverse is bounded uniformly. As Θ has nothing to do with s, t it is also bounded uniformly with respect to s, t . Accordingly so is $\Theta[(D_Y \oplus E_Y)\Theta]^{-1}$, which is a right inverse to $D_Y \oplus E_Y$. This completes the proof. \square

We prove Proposition 7.1 now.

Proof of Proposition 7.1. For each connected component $Y \subseteq X$ denote by $R_Y \oplus \pi_Y : C_{\nu-2}^{1/2}(Y, \mathbb{R}) \rightarrow C_{\nu}^{5/2}(Y, \mathbb{R}) \oplus \mathbb{R}$ the uniformly bounded right inverse produced in Lemma 7.6. Define $S : C^{1/2}(N, \mathbb{R}) \rightarrow C^{5/2}(N, \mathbb{R})$ by $S := G_L D_L^{-1} F_L + \sum_{Y \in \pi_0 X} G_Y R_Y F_Y$. Define $\varpi : C^{1/2}(N, \mathbb{R}) \rightarrow H^0(X, \mathbb{R})$ by $v \mapsto \sum_{Y \in \pi_0 X} \pi_Y F_Y v$. Recall from Definitions 7.3 and 7.5 that $D_L = \Delta$ on the support of G_L and that for each connected component $Y \subseteq X$ we have $D_Y = \Delta$ on the support on G_Y . So for $v \in C^{1/2}(N, \mathbb{R})$ we have

$$\begin{aligned} & D_L G_L D_L^{-1} F_L v + \sum_{Y \in \pi_0 X} (D_Y G_Y R_Y F_Y v) + \sum_{Y \in \pi_0 X} G_Y E_Y \pi_Y F_Y v \\ &= \Delta G_L R_L F_L v + \sum_{Y \in \pi_0 X} (\Delta G_Y R_Y F_Y v) + \sum_{Y \in \pi_0 X} E_Y \pi_Y F_Y v \\ &= \Delta S v + E \varpi v = (\Delta \oplus E)(S \oplus \varpi)v. \end{aligned} \quad (7.15)$$

On the other hand, since $G_L F_L = F_L$ it follows that

$$D_L G_L D_L^{-1} F_L v - [D_L, G_L] F_L v = G_L F_L v = F_L v. \quad (7.16)$$

For $Y \in \pi_0 X$, since $(D_Y \oplus E_Y)(R_Y \oplus \pi_Y) = \text{id}$ and $G_Y F_Y = F_Y$ it follows that

$$\sum_{Y \in \pi_0 X} G_Y D_Y R_Y F_Y v + \sum_{Y \in \pi_0 X} G_Y E_Y \pi_Y F_Y v = G_Y F_Y v = \sum_{Y \in \pi_0 X} F_Y v. \quad (7.17)$$

Since F_L and $(F_Y)_{Y \in \pi_0 X}$ form a partition of unity it follows that the rightmost term $F_L v$ on (7.16) and the rightmost term $\sum_{Y \in \pi_0 X} F_Y v$ on (7.17) sum up to v . This with (7.15)–(7.17) implies

$$(\Delta \oplus E)(S \oplus \varpi)v - v = [D_L, G_L]R_L F_L v + \sum_{Y \in \pi_0 X} [D_Y, G_Y]R_Y F_Y v. \quad (7.18)$$

Since the bracket terms contains derivatives of G_L, G_Y , which are $O((-\log t)^{-1})$, we get a constant $c > 0$ independent of s, t, v and satisfying

$$|(\Delta \oplus E)(S \oplus \varpi)v - v|_{C^{1/2}} \leq c(-\log t)^{-1}|v|_{C^{1/2}} \quad (7.19)$$

where the $C^{1/2}$ norms are computed with respect to the cylindrical metric on N . Passing to the weighted norms we get

$$|(\Delta \oplus E)(S \oplus \varpi)v - v|_{\nu-2} \leq c(-\log t)^{-1}|v|_{\nu-2}. \quad (7.20)$$

The operator $(\Delta \oplus E)(S \oplus \varpi) - \text{id} : C^{1/2}(N, \mathbb{R}) \rightarrow C^{1/2}(N, \mathbb{R})$ has thus operator norm $\leq c(-\log t)^{-1} \ll 1$. So $J := (\Delta \oplus E)(S \oplus \varpi)$ is invertible and J^{-1} is bounded uniformly with respect to s, t . Now $J^{-1}(S \oplus \varpi)$ is a right inverse to $\Delta \oplus E$ which is bounded uniformly with respect to s, t , which completes the proof of Proposition 7.1. \square

8 Second Approximate Solutions

We solve the special Lagrangian equation with an additional term.

Corollary 8.1. *There exists $c > 0$ independent of s, t and such that the following holds: for $(s, t) \in \mathcal{G}$ close enough to $(0, 0)$ there exists $u \oplus w \in C_\nu^{5/2}(N, \mathbb{R}) \oplus H^0(X, \mathbb{R})$ such that $Pdu + Ew = 0$ with $|u|_\nu \leq ct^{\theta+\nu}$.*

Proof. Denote by $R \oplus \pi : C_{\nu-2}^{1/2}(N, \mathbb{R}) \rightarrow C_\nu^{5/2}(N, \mathbb{R}) \oplus H^0(X, \mathbb{R})$ the uniformly bounded right inverse to $\Delta \oplus E$ produced in Proposition 7.1. Put $B := \{v \in C_{\nu-2}^{1/2}(N, \mathbb{R}) : |v|_{\nu-2} \leq t^{\theta+\nu}\}$. Since R is uniformly bounded it follows that there exists $c_0 > 0$ independent of s, t and such that for $v \in B$ we have $|Rv|_\nu \leq c_0 t^{\theta+\nu}$, which implies that we can define $QdRv$ as in Proposition 6.2. We can therefore define $T : B \rightarrow C_{\nu-2}^{1/2}(N, \mathbb{R})$ by $v \mapsto -P0 - QdRv$.

We show that T maps B to itself. Recall from Corollary 6.3 that there exists $c_1 > 0$ independent of s, t and such that for $v \in B$ we have

$$|QdRv|_{\nu-2} \leq c_1 t^{\nu-2} |Rv|_{\nu-2}^2 \leq c_1 c_0^2 t^{\nu-2} t^{2(\theta+\nu)}, \quad (8.1)$$

where the last inequality follows from the definition of B . On the other hand, by Corollary 6.1 we have

$$|P0|_{\nu-2} \leq |s| + t^\theta \leq 2t^\theta. \quad (8.2)$$

This and (8.1) imply that

$$|Tv|_{\nu-2} \leq |P0|_{\nu-2} + |QdRv|_{\nu-2} \leq (2t^{-\nu} + c_1 c_0^2 t^{\theta+2\nu-2}) t^{\theta+\nu}. \quad (8.3)$$

Since $\theta > 2 - 2\nu$ and $\nu > 0$ it follows by (8.3) that for t small enough we have $Tv \in B$.

We have thus defined the map $T : B \rightarrow B$. For $v, v' \in B$ we have

$$\begin{aligned} |Tv - Tv'|_{\nu-2} &\leq c_1 t^{\nu-2} |Rv - Rv'|_{\nu} (|Rv|_{\nu} + |Rv'|_{\nu}) \\ &\leq c_1 c_0^2 t^{\nu-2} |v - v'|_{\nu-2} (|v|_{\nu-2} + |v'|_{\nu-2}) \leq c_1 c_0^2 t^{\theta+2\nu-2} |v - v'|_{\nu-2}. \end{aligned} \quad (8.4)$$

Hence we see, using again $\theta + 2\nu - 2 > 0$, that for t small enough the map $T : B \rightarrow B$ is a contraction map. As B is closed we can apply to it the fixed point theorem; that is, there exists $v \in B$ with $Tv = v$.

Since $R \oplus \pi$ is a right inverse to $\Delta \oplus E$ it follows that putting $u \oplus w := (R \oplus \pi)v$ we have $(\Delta \oplus E)(u \oplus w) = v$. Now

$$Pdu + Ew = P0 + Qu + \Delta u + Ew = P0 + QdRv + v = -Tv + v = 0. \quad (8.5)$$

Moreover $|u|_{\nu} \leq c_0 |v|_{\nu-2} \leq c_0 t^{\theta+\nu}$ as we have to prove. \square

We prove now Theorem 2.14 for X connected.

Proof of Theorem 2.14 for X connected. Regard $w \in H^0(X, \mathbb{R}) \cong \mathbb{R}$ as a real constant. Since $E_X : N \rightarrow [0, 1]$ is supported in $N \setminus \bigsqcup_{x \in X^{\text{sing}}} t_x K_x \cong X \setminus \bigsqcup_{x \in X^{\text{sing}}} t_x K_x$ it follows that

$$w \int_X E_X \, \text{dvol} = w \int_N E_X \, \text{dvol} = - \int_N Pdu \, \text{dvol} = - \int_N \text{Im } \Omega^s. \quad (8.6)$$

Here the right-hand side is the pairing of $[N] \in H_m(M, \mathbb{Z})$ and $[\text{Im } \Omega^s] \in H^m(M, \mathbb{R})$. Definition 4.4 shows that $[N]$ converges, as (s, t) tends to $(0, 0)$, to $[X] \in H_m(N, \mathbb{Z})$. But the integer homology classes are discrete, so $[N^{st}] = [X]$ for (s, t) close enough to $(0, 0)$. The right-hand side of (8.6) is therefore equal to the pairing $[X] \cdot [\text{Im } \Omega^s]$. In the current case, as X is connected the left-hand side of (2.2) is the sum over all connected components of $C_{x_j} \subseteq C_x$. But the pairing $\sum_{C_{x_j} \subseteq C_x} [C_{x_j} \cap S^{2m-1}] \cdot Z(L_x)$ is equal to the integral of $\text{Im } \Omega$ over a compact set in L_x , which vanishes automatically. So the left-hand side of (2.2) vanishes, and accordingly $[X] \cdot [\text{Im } \Omega^s] = 0$. Thus $w \int_X E_X \, \text{dvol} = 0$. But E_X is so chosen that $\int_X E_X \, \text{dvol} \neq 0$ and we have therefore $w = 0$. This means that the graph of du is a special Lagrangian submanifold of $(M; \omega^s, J^s, \Omega^s)$. \square

When X need not be connected, we deal with the additional term w as follows.

Lemma 8.2. *Suppose that for each $(s, t) \in \mathcal{G}$ we are given $u \oplus w \in C_{\nu}^{5/2}(N, \mathbb{R}) \oplus H^0(X, \mathbb{R})$ with $Pdu + Ew = 0$. Then there exist $c, \epsilon > 0$ independent of s, t and such that if we write $w = \bigoplus_{Y \in \pi_0 X} w_Y \in H^0(X, \mathbb{R})$ then $\max_{Y \in \pi_0 X} |w_Y| \leq ct^{m+\epsilon}$.*

Proof. We proceed as in the proof of Theorem 2.13. For each connected component $Y \subseteq X$ there is a unique connected component of $(\text{Graph } du) \setminus (\bigsqcup_{x \in X^{\text{sing}}} t_x K_x)$ that is C^1 close to $Y \setminus \bigsqcup_{x \in X^{\text{sing}}} t_x K_x$, which we denote by Y^u . This Y^u is a manifold with boundary, with $\partial Y^u \subseteq \bigsqcup_{x \in X^{\text{sing}}} t_x K_x$. For $x \in X^{\text{sing}}$ let $\Sigma_{xj} \subseteq t_x^{-1}(Y^u \cap t_x K_x)$ be a connected component, which is a compact $m-1$ manifold diffeomorphic to some connected component $C_{xj} \cap S^{2m-1} \subseteq C_x \cap S^{2m-1}$. Denote by $0 * \Sigma_{xj} \subseteq \mathbb{C}^m$ the union of the line segments between the origin 0 and the points of Σ_{xj} . The union $Y^u \cup \bigsqcup_{C_{xj} \subseteq Y} t_x(0 * \Sigma_{xj})$ defines in M an m -cycle homologous to \bar{Y} . Integrating $\text{Im } \Omega^s$ over this we get

$$[\bar{Y}] \cdot [\text{Im } \Omega^s] = \sum_{C_{xj} \subseteq Y} \int_{t_x(0 * \Sigma_{xj})} \text{Im } \Omega^s + \int_{Y^u} \text{Im } \Omega^s. \quad (8.7)$$

Following with modification the proof of Theorem 2.13 we find $\epsilon > 0$ such that

$$\sum_{C_{xj} \subseteq Y} \int_{t_x(0 * \Sigma_{xj})} \text{Im } \Omega^s = \sum_{C_{xj} \subseteq Y} t_x^m \psi^s(x)^m [C_{xj} \cap S^{2m-1}] \cdot Z(L_x) + O(t^{m+\epsilon}). \quad (8.8)$$

But it is supposed in Theorem 2.14 that $(s, t) \in \mathcal{G}$ satisfies (ii) of Theorem 2.13; that is,

$$[\bar{Y}] \cdot [\text{Im } \Omega^s] = \sum_{x \in X^{\text{sing}}} t^m \psi^s(x)^m (1 + \gamma_x) [C_x \cap S^{2m-1}] \cdot Z(L_x). \quad (8.9)$$

Recalling that $t_x = \sqrt{1 + \beta_x} t$ and that β_x, γ_x decay with power orders it follows from (8.7)–(8.9) that

$$\int_{Y^u} \text{Im } \Omega^s = O(t^{m+\epsilon}). \quad (8.10)$$

Notice that $\text{Im } \Omega^s|_{\text{Graph } du} = (Pdu)d\text{vol} = -(Ew)d\text{vol}$. By the definition of E we have $Ew = \sum_{Y \in \pi_0 X} w_Y E_Y$. Since each E_Y is supported in $Y^u \cong Y \setminus \bigsqcup_{x \in X^{\text{sing}}} t_x K_x$ it follows that

$$\int_{Y^u} \text{Im } \Omega^s = - \int_{Y^u} (Ew)d\text{vol} = -w_Y \int_Y E_Y d\text{vol}. \quad (8.11)$$

As E_Y is so chosen that $\int_Y E_Y d\text{vol}$ is a non-zero constant independent of s, t it follows from (8.9)–(8.11) that $w_Y = O(t^{m+\epsilon})$. \square

9 Proof of Theorem 2.14

We begin by recalling from [6] the key result we will use.

Theorem 9.1 (Theorem 5.3 of [6]). *Let $m \geq 3$ be an integer and $(M; \omega, J, \Omega)$ an almost Calabi–Yau m -fold. Define $\psi : M \rightarrow (0, \infty)$ by (2.1). Let $Q \subseteq (M, \omega)$ be a compact oriented Lagrangian submanifold, and g the Riemannian metric on Q induced from the Kähler metric of $(M; \omega, J)$. Suppose given $c, t > 0$ such that*

- (i) the Riemannian manifold (Q, g) has injectivity radius $\geq ct$ and sectional curvature $\leq ct^{-2}$.

Let $U \subseteq (M, \omega)$ be a Weinstein neighbourhood of Q isomorphic to a neighbourhood of the zero-section $Q \subseteq T^*Q$ whose fibres are balls of radius ct around the zero-section. Denote by $\pi : T^*Q \rightarrow Q$ the vector bundle projection. Notice that the Levi-Civita connection of (Q, g) induces over T^*Q a vector bundle isomorphism $T(T^*Q) \cong \pi^*TQ \oplus \pi^*T^*Q$. Since the two vector bundles π^*TQ and π^*T^*Q have metrics and Levi-Civita connections induced from g , we get on $T(T^*Q)$ a metric and a connection. Denote these by \hat{g} and $\hat{\nabla}$. Suppose that

- (ii) for $k = 0, 1, 2, 3$ we have $\sup_U |\hat{\nabla}^k \text{Im } \Omega| \leq ct^{-k}$ where $|\cdot|$ is computed pointwise with respect to \hat{g} .

For $p \in [1, \infty]$ denote by $L^p(Q, \mathbb{R})$ the L^p space of the Riemannian manifold Q . For $p \in [1, \infty)$ and $k = 0, 1, 2, \dots$ denote by $L_k^p(Q, \mathbb{R})$ the Sobolev space of the Riemannian manifold Q . So $L_0^p(Q, \mathbb{R}) = L^p(Q, \mathbb{R})$. Suppose now that $W \subseteq L^2(Q, \mathbb{R})$ is a finite dimensional \mathbb{R} -vector subspace, and $\pi_W : L^2(Q, \mathbb{R}) \rightarrow W$ the L^2 orthogonal projection. Suppose that

- (iii) for $v \in L_1^2(Q, \mathbb{R})$ with $\pi_W v = 0$ we have $\|v\|_{L^{\frac{2m}{m-2}}} \leq c \|dv\|_{L^2}$,

- (iv) for $w \in W$ we have $\|d^*dw\|_{L^{\frac{2m}{m+2}}} \leq \frac{1}{2}c \|dw\|_{L^2}$ and

- (v) for $w \in W$ with $\int_Q w \, d\text{vol} = 0$ we have $\|w\|_{C^0} \leq ct^{1-\frac{m}{2}} \|dw\|_{L^2}$.

Denote by $d\text{vol}$ the volume form of the Riemannian manifold Q and define a smooth function $e^{i\theta} : Q \rightarrow S^1$ by $\Omega|_Q = (\psi|_Q)^m e^{i\theta} d\text{vol}$. Denote by $\cos \theta : Q \rightarrow \mathbb{R}$ its real part and by $\sin \theta : Q \rightarrow \mathbb{R}$ its imaginary part. Suppose that there exists $\kappa > 0$ such that

- (vi) $\|\pi_W(\psi^m \sin \theta)\|_{L^1} \leq ct^{\kappa+m-1}$ and

- (vii) for each $p \in [1, \infty]$ we have $\|\psi^m \sin \theta\|_{L^p} + t \|d(\psi^m \sin \theta)\|_{L^p} \leq ct^{\kappa+\frac{m}{p}-1}$.

Then there exists $t_0 > 0$ depending only on m, c, κ and such that if $t \in (0, t_0)$ then there exists on Q an exact 1-form ξ whose graph of lies in the Weinstein neighbourhood $U \subseteq (M, \omega)$ and defines a special Lagrangian submanifold of $(M; \omega, J, \Omega)$. There is moreover a constant $c' > 0$ depending only on m, c, κ such that $\sup_Q |\xi| \leq c't^\kappa$ where $|\cdot|$ is computed pointwise with respect to g .

Remark 9.2. The idea of using such a uniform Sobolev inequality as in (iii) above goes back to [9].

Let s, t satisfy $|s| \leq t^\theta$ and $(s, t) \in \mathcal{G}$. Let $u \oplus w$ be as in Corollary 8.1 and put $Q := \text{Graph } du$. We show that Q satisfies the hypotheses (i)–(vii) of Theorem 9.1 with $(M; \omega^s, J^s, \Omega^s)$ in place of $(M; \omega, J, \Omega)$. The proof that Q satisfies (i)–(vi) of them are essentially the same as in [6, 7]. More precisely, in the circumstances of [6, 7] it is proved that the first approximate solution N satisfies (i)–(vi) of

Theorem 9.1. The details about (i),(ii) are given in [6, §6.3] in the simplest case (amongst the results of [6, 7]). The details about (iii)–(v) are given in [6, §7.3] in the simplest case. The details about (vi) are given in [6, §7.2] in the simplest case.

Our current circumstances are more complex because the submanifold Q is a further perturbation of N . But Q is in $\bigsqcup_{x \in X^{\text{sing}}} U_x$ as close to $t_x L_x$ as N is, and outside $\bigsqcup_{x \in X^{\text{sing}}} t_x K_x$ as close to X as N is. These properties are all we need for (i)–(v) of Theorem 9.1. For (vi) we use also Lemma 8.2.

The proof of (vii) is the difference from [6, 7] as we show now. Since $u \oplus w$ is as in Corollary 8.1 it follows that $\text{Im } \Omega|_Q = -\sum_{Y \in \pi_0 X} w_Y E_Y$. But as in Lemma 8.2 every constant w_Y is $O(t^{m+\epsilon})$. Since each E_Y is supported in $Y \setminus \bigsqcup_{x \in X^{\text{sing}}} U_x$ it follows that for every $p \in [1, \infty]$ we have

$$\|\psi^m \sin \theta\|_{L^p} + \|d(\psi^m \sin \theta)\|_{L^p} \leq ct^{m+\epsilon}, \quad (9.1)$$

which implies (vii). We can thus apply Theorem 9.1 to Q , which completes the proof of Theorem 2.14. \square

References

- [1] S.K. Donaldson and P.B. Kronheimer ‘The Geometry of Four-Manifolds’ Oxford Science Publications 1990
- [2] Y. Imagi ‘A uniqueness theorem for gluing calibrated submanifolds’ *Comm. Anal. Geom.* 23 (2015) 691–715
- [3] Y. Imagi ‘Example of compact special Lagrangians with a stable singularity’ *Int. Math. Res. Not.* (2020) 7975–8006
- [4] D.D. Joyce ‘Special Lagrangian submanifolds with isolated conical singularities I. Regularity’ *Ann. Global Anal. Geom.* 25 (2003) 201–58
- [5] D.D. Joyce ‘Special Lagrangian submanifolds with isolated conical singularities. II. Deformation theory’ *Ann. Global Anal. Geom.* 25 (2003) 201–58
- [6] D.D. Joyce ‘Special Lagrangian submanifolds with isolated conical singularities. III. Desingularization, the unobstructed case’ *Ann. Global Anal. Geom.* 26 (2004) 1–58
- [7] D.D. Joyce ‘Special Lagrangian submanifolds with isolated conical singularities. IV. Desingularization, obstructions and families’ *Ann. Global Anal. Geom.* 26 (2004) 117–174
- [8] D.D. Joyce ‘Special Lagrangian submanifolds with isolated conical singularities. V. Survey and applications’ *J. Differential Geom.* 63 (2003) 279–347

- [9] Y-I. Lee ‘Embedded special Lagrangian submanifolds in Calabi-Yau manifolds’ *Comm. Anal. Geom.* 11 (2003) 391–423
- [10] S.P. Marshall ‘Deformations of special Lagrangian submanifolds’ Oxford D.Phil. thesis 2002
- [11] T. Pacini ‘Special Lagrangian conifolds, II: gluing constructions in \mathbb{C}^m ’ *Proc. London Math. Soc.* (3) 107 (2013) 225–66

Institute of Mathematical Sciences, ShanghaiTech University, 393 Middle Huaxia Road, Pudong New District, Shanghai, China
e-mail address: yosukeimagi@shanghaitech.edu.cn