

RIEMANN–LIOUVILLE INTEGRALS IN BESOV SPACES

Elena P. Ushakova *

*V.A. Trapeznikov Institute of Control Sciences of Russian Academy of Sciences, 65 Profsoyuznaya str.,
Moscow 117997, Russia*

Steklov Mathematical Institute of Russian Academy of Sciences, 8 Gubkina str., Moscow 119991, Russia

*Computing Center of Far Eastern Branch of Russian Academy of Sciences, 65 Kim Yu Chena str.,
Khabarovsk 680000, Russia*

Key words: Riemann–Liouville operator, Besov space, Muckenhoupt weight, local Muckenhoupt weight, B –spline, Battle–Lemarié wavelet system, decomposition theorem

MSC (2010): 44A15, 41A15, 26A33

Abstract. Criteria for the fulfillment of inequalities in weighted smoothness function spaces of Besov type with Riemann–Liouville operators of natural orders on the real axis and semi–axes are found. The obtained estimates are refined under additional conditions on weights.

1 Introduction

Consider the left and right Riemann–Liouville operators of natural orders α of the forms

$$I_{c+}^{\alpha} f(x) := \frac{1}{\Gamma(\alpha)} \int_c^x (x-y)^{\alpha-1} f(y) dy \quad (x > c) \quad (1.1)$$

and

$$I_{c-}^{\alpha} f(x) := \frac{1}{\Gamma(\alpha)} \int_x^c (x-y)^{\alpha-1} f(y) dy \quad (x < c) \quad (1.2)$$

(see e.g. [18]). Here $c \in [-\infty, +\infty]$. If $c = \mp\infty$ we denote (1.1) and (1.2) by I_{\mp}^{α} and I^{α} , respectively.

It is assumed that images $I_{c\pm}^{\alpha} f$ and pre–images f of our operators $I_{c\pm}^{\alpha}$ are identified with density functions of regular distributions (see e.g. [21]) from spaces of Besov type $B_{pq}^{s,w}(\mathbb{R})$ with summation parameters $p > 0$ and $0 < q \leq \infty$, smoothness indices $s \in \mathbb{R}$ and weights w of Muckenhoupt type $\mathcal{A}_{\infty}^{\text{loc}}$ (see related definitions in Section 2). By weight w we mean a locally integrable almost everywhere positive function on \mathbb{R} . The main problem studied in this paper is characterisation of inequalities connecting the norms of images and pre–images of $I_{c\pm}^{\alpha}$ in $B_{pq}^{s,w}(\mathbb{R})$.

Similar inequalities for some classes of operators acting in Besov type spaces were studied in [23, Theorem 2.3.8], [5], [17, Theorem 2.20], [10, § 4], [25, p. 23]. For $I_{c\pm}^{\alpha}$ the problem was considered in [17, Theorem 2.20], [10, Proposition 4.2], as well as in [30–32] and [33]. In particular, the following results were obtained in [33].

Theorem 1.1 ([33, Theorem 5.5]). *Let $p > 1$, $0 < q \leq \infty$, $s \in \mathbb{R}$, weights $u, v, w \in \mathcal{A}_{\infty}^{\text{loc}}$, $\alpha \in \mathbb{N}$ and $f \in L^1(\mathbb{R})$.*

(i) *Then $I_{\pm}^{\alpha} f \in B_{pq}^{s,u}(\mathbb{R})$ if $f \in B_{pq}^{s+\kappa^*-\alpha,v}(\mathbb{R})$ with some $\kappa^* \in \mathbb{R}$ such that*

$$\mathbf{C}_{\pm}^{\alpha}(\kappa^*) = \mathbf{M}_{\pm}^{\alpha}(1) + \mathbf{M}_{\pm}^{\alpha}(0) + \sup_{d \in \mathbb{N}} \left[\mathcal{M}_{\pm}^{\alpha}(d, \kappa^*, 1) + \mathcal{M}_{\pm}^{\alpha}(d, \kappa^*, 0) \right] < \infty, \quad \text{where} \quad (1.3)$$

$$\mathbf{M}_{\pm}^{\theta}(\varepsilon) := \sup_{\tau \in \mathbb{Z}} \left(\sum_{r \geq \tau} (r - \tau + 1)^{p(\theta-1)\varepsilon} u(Q_{0r}) \right)^{\frac{1}{p}} \left(\sum_{r \leq \tau} (\tau - r + 1)^{p'(\theta-1)(1-\varepsilon)} [v(Q_{0r})]^{1-p'} \right)^{\frac{1}{p'}},$$

*elenau@inbox.ru

$$\begin{aligned}
\mathbf{M}_-^\theta(\varepsilon) &:= \sup_{\tau \in \mathbb{Z}} \left(\sum_{r \leq \tau} (\tau - r + 1)^{p(\theta-1)\varepsilon} u(Q_{0r}) \right)^{\frac{1}{p}} \left(\sum_{r \geq \tau} (r - \tau + 1)^{p'(\theta-1)(1-\varepsilon)} [v(Q_{0r})]^{1-p'} \right)^{\frac{1}{p'}}, \\
\mathcal{M}_+^\theta(d, \kappa, \varepsilon) &:= 2^{-d\kappa} \sup_{\tau \in \mathbb{Z}} \left(\sum_{r \geq \tau} (r - \tau + 1)^{p(2\theta-1)\varepsilon} u(Q_{(d-1)r}) \right)^{\frac{1}{p}} \\
&\quad \times \left(\sum_{r \leq \tau} (\tau - r + 1)^{p'(2\theta-1)(1-\varepsilon)} [v(Q_{(d-1)r})]^{1-p'} \right)^{\frac{1}{p'}}, \\
\mathcal{M}_-^\theta(d, \kappa, \varepsilon) &:= 2^{-d\kappa} \sup_{\tau \in \mathbb{Z}} \left(\sum_{r \leq \tau} (\tau - r + 1)^{p(2\theta-1)\varepsilon} u(Q_{(d-1)r}) \right)^{\frac{1}{p}} \\
&\quad \times \left(\sum_{r \geq \tau} (r - \tau + 1)^{p'(2\theta-1)(1-\varepsilon)} [v(Q_{(d-1)r})]^{1-p'} \right)^{\frac{1}{p'}}.
\end{aligned}$$

Moreover,

$$\|I_\pm^\alpha f\|_{B_{pq}^{s,u}(\mathbb{R})} \lesssim \mathbf{C}_\pm^\alpha(\kappa^*) \|f\|_{B_{pq}^{s+\kappa^*-\alpha,v}(\mathbb{R})}. \quad (1.4)$$

(ii) If $I_\pm^\alpha f \in B_{pq}^{s,u}(\mathbb{R})$ then $f \in B_{pq}^{s-\kappa_*-\alpha,w}(\mathbb{R})$ with $\kappa_* \in \mathbb{R}$ such that there exist $0 \leq \varepsilon_M, \varepsilon_M \leq 1$, satisfying the condition

$$\mathbf{C}_\pm^\alpha(\kappa_*) = M_\pm^\alpha(\varepsilon_M) + \sup_{d \in \mathbb{N}} \mathbf{M}_\pm^\alpha(d, \kappa_*, \varepsilon_M) < \infty, \quad \text{where} \quad (1.5)$$

$$\begin{aligned}
M_+^\theta(\varepsilon) &:= \sup_{\tau \in \mathbb{Z}} \left(\sum_{r=\tau-\theta}^{\tau} (\tau - r + 1)^{-p(\theta+1)\varepsilon} w(Q_{0r}) \right)^{\frac{1}{p}} \left(\sum_{r=\tau}^{\tau+\theta} (r - \tau + 1)^{-p'(\theta+1)(1-\varepsilon)} [u(Q_{0r})]^{1-p'} \right)^{\frac{1}{p'}}, \\
M_-^\theta(\varepsilon) &:= \sup_{\tau \in \mathbb{Z}} \left(\sum_{r=\tau}^{\tau+\theta} (r - \tau + 1)^{-p(\theta+1)\varepsilon} w(Q_{0r}) \right)^{\frac{1}{p}} \left(\sum_{r=\tau-\theta}^{\tau} (\tau - r + 1)^{-p'(\theta+1)(1-\varepsilon)} [u(Q_{0r})]^{1-p'} \right)^{\frac{1}{p'}}, \\
\mathbf{M}_+^\theta(d, \kappa, \varepsilon) &:= 2^{-d\kappa} \sup_{\tau \in \mathbb{Z}} \left(\sum_{r=\tau-\theta}^{\tau} (\tau - r + 1)^{-p(2\theta+1)\varepsilon} w(Q_{(d-1)r}) \right)^{\frac{1}{p}} \\
&\quad \times \left(\sum_{r=\tau}^{\tau+\theta} (r - \tau + 1)^{-p'(2\theta+1)(1-\varepsilon)} [u(Q_{(d-1)r})]^{1-p'} \right)^{\frac{1}{p'}}, \\
\mathbf{M}_-^\theta(d, \kappa, \varepsilon) &:= 2^{-d\kappa} \sup_{\tau \in \mathbb{Z}} \left(\sum_{r=\tau}^{\tau+\theta} (r - \tau + 1)^{-p(2\theta+1)\varepsilon} w(Q_{(d-1)r}) \right)^{\frac{1}{p}} \\
&\quad \times \left(\sum_{r=\tau-\theta}^{\tau} (\tau - r + 1)^{-p'(2\theta+1)(1-\varepsilon)} [u(Q_{(d-1)r})]^{1-p'} \right)^{\frac{1}{p'}}.
\end{aligned}$$

Besides,

$$\|f\|_{B_{pq}^{s-\kappa_*-\alpha,w}(\mathbb{R})} \lesssim \mathbf{C}_\pm^\alpha(\kappa_*) \|I_\pm^\alpha f\|_{B_{pq}^{s,u}(\mathbb{R})}. \quad (1.6)$$

(iii) If $I_\pm^\alpha f \in B_{pq}^{s,u}(\mathbb{R})$ then $f \in B_{pq}^{s-\alpha,u}(\mathbb{R})$, and

$$\|f\|_{B_{pq}^{s-\alpha,u}(\mathbb{R})} \lesssim \|I_\pm^\alpha f\|_{B_{pq}^{s,u}(\mathbb{R})}. \quad (1.7)$$

To further list the results from [33], we introduce the following notations:

$$\tilde{\mathbf{C}}_\pm^\alpha(\kappa^*, c) = \tilde{\mathbf{M}}_\pm^\alpha(1, c) + \tilde{\mathbf{M}}_\pm^\alpha(0, c) + \sup_{d \in \mathbb{N}} \left[\tilde{\mathcal{M}}_\pm^\alpha(d, \kappa^*, 1, c) + \tilde{\mathcal{M}}_\pm^\alpha(d, \kappa^*, 0, c) \right],$$

$$\widetilde{\mathbf{C}}_{\pm}^{\alpha}(\kappa_*, c) = \widetilde{M}_{\pm}^{\alpha}(\varepsilon_{\widetilde{M}}, c) + \sup_{d \in \mathbb{N}} \widetilde{M}_{\pm}^{\alpha}(d, \kappa_*, \varepsilon_{\widetilde{M}}, c),$$

where with $Q_{dr}^{(c)} := \left[\frac{r+c}{2^d}, \frac{r+c+1}{2^d} \right]$ for some $c \in \mathbb{R}$

$$\begin{aligned} \widetilde{M}_{+}^{\theta}(\varepsilon, c) &:= \sup_{\tau \in \mathbb{N}_0} \left(\sum_{r \geq \tau} (r - \tau + 1)^{p(\theta-1)\varepsilon} u(Q_{0r}^{(c)}) \right)^{\frac{1}{p}} \left(\sum_{0 \leq r \leq \tau} (\tau - r + 1)^{p'(\theta-1)(1-\varepsilon)} [v(Q_{0r}^{(c)})]^{1-p'} \right)^{\frac{1}{p'}}, \\ \widetilde{M}_{-}^{\theta}(\varepsilon, c) &:= \sup_{\tau \in -\mathbb{N}_0} \left(\sum_{r \leq \tau} (\tau - r + 1)^{p(\theta-1)\varepsilon} u(Q_{0r}^{(c)}) \right)^{\frac{1}{p}} \left(\sum_{0 \geq r \geq \tau} (r - \tau + 1)^{p'(\theta-1)(1-\varepsilon)} [v(Q_{0r}^{(c)})]^{1-p'} \right)^{\frac{1}{p'}}, \\ \widetilde{\mathcal{M}}_{+}^{\theta}(d, \kappa, \varepsilon, c) &:= 2^{-d\kappa} \sup_{\tau \in \mathbb{N}_0} \left(\sum_{r \geq \tau} (r - \tau + 1)^{p(2\theta-1)\varepsilon} u(Q_{(d-1)r}^{(c)}) \right)^{\frac{1}{p}} \\ &\quad \times \left(\sum_{0 \leq r \leq \tau} (\tau - r + 1)^{p'(2\theta-1)(1-\varepsilon)} [v(Q_{(d-1)r}^{(c)})]^{1-p'} \right)^{\frac{1}{p'}}, \\ \widetilde{\mathcal{M}}_{-}^{\theta}(d, \kappa, \varepsilon, c) &:= 2^{-d\kappa} \sup_{\tau \in -\mathbb{N}_0} \left(\sum_{r \leq \tau} (\tau - r + 1)^{p(2\theta-1)\varepsilon} u(Q_{(d-1)r}^{(c)}) \right)^{\frac{1}{p}} \\ &\quad \times \left(\sum_{0 \geq r \geq \tau} (r - \tau + 1)^{p'(2\theta-1)(1-\varepsilon)} [v(Q_{(d-1)r}^{(c)})]^{1-p'} \right)^{\frac{1}{p'}}, \\ \widetilde{M}_{+}^{\theta}(\varepsilon, c) &:= \sup_{\tau \in \mathbb{N}_0} \left(\sum_{r=\tau-\theta}^{\tau} (\tau - r + 1)^{-p(\theta+1)\varepsilon} w(Q_{0r}^{(c)}) \right)^{\frac{1}{p}} \left(\sum_{r=\tau}^{\tau+\theta} (r - \tau + 1)^{-p'(\theta+1)(1-\varepsilon)} [u(Q_{0r}^{(c)})]^{1-p'} \right)^{\frac{1}{p'}}, \\ \widetilde{M}_{-}^{\theta}(\varepsilon, c) &:= \sup_{\tau \in -\mathbb{N}_0} \left(\sum_{r=\tau}^{\tau+\theta} (r - \tau + 1)^{-p(\theta+1)\varepsilon} w(Q_{0r}^{(c)}) \right)^{\frac{1}{p}} \left(\sum_{r=\tau-\theta}^{\tau} (\tau - r + 1)^{-p'(\theta+1)(1-\varepsilon)} [u(Q_{0r}^{(c)})]^{1-p'} \right)^{\frac{1}{p'}}, \\ \widetilde{M}_{+}^{\theta}(d, \kappa, \varepsilon, c) &:= 2^{-d\kappa} \sup_{\tau \in \mathbb{N}_0} \left(\sum_{r=\tau-\theta}^{\tau} (\tau - r + 1)^{-p(2\theta+1)\varepsilon} w(Q_{(d-1)r}^{(c)}) \right)^{\frac{1}{p}} \\ &\quad \times \left(\sum_{r=\tau}^{\tau+\theta} (r - \tau + 1)^{-p'(2\theta+1)(1-\varepsilon)} [u(Q_{(d-1)r}^{(c)})]^{1-p'} \right)^{\frac{1}{p'}}, \\ \widetilde{M}_{-}^{\theta}(d, \kappa, \varepsilon, c) &:= 2^{-d\kappa} \sup_{\tau \in -\mathbb{N}_0} \left(\sum_{r=\tau}^{\tau+\theta} (r - \tau + 1)^{-p(2\theta+1)\varepsilon} w(Q_{(d-1)r}^{(c)}) \right)^{\frac{1}{p}} \\ &\quad \times \left(\sum_{r=\tau-\theta}^{\tau} (\tau - r + 1)^{-p'(2\theta+1)(1-\varepsilon)} [u(Q_{(d-1)r}^{(c)})]^{1-p'} \right)^{\frac{1}{p'}}. \end{aligned}$$

Theorem 1.2 ([33, Theorem 6.1]). *Let $p > 1$, $0 < q \leq \infty$, $s \in \mathbb{R}$, $u, v \in \mathcal{A}_{\infty}^{\text{loc}}$, $\alpha \in \mathbb{N}$ and $f \in L_{\text{loc}}(\mathbb{R})$. Suppose that $f \equiv 0$ on $(-\infty, c)$ in the case of operator I_{c+}^{α} and, similarly, $f \equiv 0$ on (c, ∞) for I_{c-}^{α} .*

(i) *Then $I_{c\pm}^{\alpha} f \in B_{pq}^{s,u}(\mathbb{R})$ if $f \in B_{pq}^{s+\kappa^*-\alpha,v}(\mathbb{R})$ with some $\kappa^* \in \mathbb{R}$ such that $\widetilde{\mathbf{C}}_{\pm}^{\alpha}(\kappa^*, c) < \infty$. Moreover,*

$$\|I_{c\pm}^{\alpha} f\|_{B_{pq}^{s,u}(\mathbb{R})} \lesssim \widetilde{\mathbf{C}}_{\pm}^{\alpha}(\kappa^*, c) \|f\|_{B_{pq}^{s+\kappa^*-\alpha,v}(\mathbb{R})}. \quad (1.8)$$

(ii) *If $I_{c\pm}^{\alpha} f \in B_{pq}^{s,u}(\mathbb{R})$ then $f \in B_{pq}^{s-\kappa^*-\alpha,w}(\mathbb{R})$ with such $\kappa^* \in \mathbb{R}$ that with some $0 \leq \widetilde{\varepsilon}_{\widetilde{M}}, \widetilde{\varepsilon}_{\widetilde{M}} \leq 1$ $\widetilde{\mathbf{C}}_{\pm}^{\alpha}(\kappa^*, a) < \infty$. Besides,*

$$\|f\|_{B_{pq}^{s-\kappa^*-\alpha,w}(\mathbb{R})} \lesssim \widetilde{\mathbf{C}}_{\pm}^{\alpha}(\kappa^*, a) \|I_{c\pm}^{\alpha} f\|_{B_{pq}^{s,u}(\mathbb{R})}. \quad (1.9)$$

(iii) If $I_{c_{\pm}}^{\alpha} f \in B_{pq}^{s,u}(\mathbb{R})$ then $f \in B_{pq}^{s-\alpha,u}(\mathbb{R})$, and (1.7) is true with $I_{c_{\pm}}^{\alpha}$ instead of I_{\pm}^{α} .

In Theorems 1.1 or 1.2, sufficient conditions of the form (1.3) (or $\widetilde{\mathbf{C}}_{\pm}^{\alpha}(\kappa^*, c) < \infty$) and (1.5) (or $\widetilde{\mathbf{C}}_{\pm}^{\alpha}(\kappa_*, a) < \infty$) for the validity of the inequalities (1.4) (or (1.8)) and (1.6) (or (1.9)), respectively, are obtained. In this paper, we improve these results up to criteria (see Theorems 3.4 and 3.5) by suitably modifying the second component (the mother wavelet) of the spline wavelet systems used for obtaining upper bounds in the decomposition theorems (see Theorem 3.3).

We write $h(\Omega) := \int_{\Omega} h(x) dx$, where $\Omega \subset \mathbb{R}$ is a bounded measurable set. Symbol \mathbb{R} denotes the real axis, \mathbb{N} and \mathbb{Z} – the sets of natural and integer numbers, respectively. Notation \mathbb{N}_0 is used for the set $\mathbb{N} \cup \{0\}$, $[s]$ – for the integer part of $s \in \mathbb{R}$. Relation $A \lesssim B$ means that $A \leq cB$ with some constant $c > 0$, depending only on fixed numerical parameters. Similarly, $A \simeq B$ is used instead of $A = cB$. We write $A \approx B$ in the case of $A \lesssim B \lesssim A$. For $r > 1$, the conjugate (or dual) exponent r' is defined as $r/(r-1)$, with $r' = \infty$ if $r = 1$. Symbols $:=$ and $=:$ are used to denote new quantities. By $L_{\text{loc}}(\mathbb{R})$ we denote the collection of all locally integrable functions on \mathbb{R} . The notation $L^p(\mathbb{R})$, $0 < p \leq \infty$, stands for the Lebesgue spaces of all measurable functions on \mathbb{R} with the (quasi-)norm $\|f\|_{L^p(\mathbb{R})} := (\int_{\mathbb{R}} |f(x)|^p dx)^{1/p}$ admitting standard modification for the case $p = \infty$.

2 Weighted Besov type spaces

2.1 Weights from the Muckenhoupt class A_{∞}

Let Q be a segment in \mathbb{R} and $|Q|$ – its length. Denote

$$A_{\rho}[w(Q)] := \frac{w(Q)}{|Q|} \left(\frac{1}{|Q|} \int_Q w^{1-\rho'} \right)^{\frac{\rho}{\rho'}}, \quad A_1[w(Q)] := \frac{w(Q)}{|Q|} \|1/w\|_{L^{\infty}(Q)}. \quad (2.1)$$

Definition 2.1. (i) Weight w belongs to *Muckenhoupt class* A_{ρ} , $1 < \rho < \infty$, if

$$A_{\rho}(w) := \sup_{Q \subset \mathbb{R}} A_{\rho}[w(Q)] < \infty;$$

(ii) $w \in A_1$ if $A_1(w) := \sup_{Q \subset \mathbb{R}} A_1[w(Q)] < \infty$;

(iii) the *Muckenhoupt class* A_{∞} is $\bigcup_{\rho \geq 1} A_{\rho}$.

Let us mention some properties of weights from the Muckenhoupt class [13, 22]:

- if $w \in A_{\rho}$, $1 < \rho < \infty$, then $w^{-\rho'/\rho} \in A_{\rho'}$;
- if $1 \leq \rho_1 < \rho_2 \leq \infty$ then $A_{\rho_1} \subset A_{\rho_2}$;
- if $w \in A_{\rho}$, $p > 1$, then there exists $r < \rho$ such that $w \in A_r$.

The last property implies the existence of the number

$$r_0 := r_0(w) := \inf\{r \geq 1 : w \in A_r\} < \infty, \quad w \in A_{\infty}.$$

The functions $w \in A_{\infty}$ satisfy two important generalised doubling-type conditions [6].

(I) For $w \in A_{\rho}$, $\rho \geq 1$, there exists a constant $c_I > 0$ such that for all segments B and F satisfying $F \subset B$, the following holds:

$$\left(\frac{|F|}{|B|} \right)^{\rho} \leq c_I \frac{w(F)}{w(B)}. \quad (2.2)$$

(II) If $w \in A_{\infty}$ then there are $\rho^* > 0$ and $c_{II} > 0$ such that for all B and E such that $E \subset B$,

$$\frac{w(E)}{w(B)} \leq c_{II} \left(\frac{|E|}{|B|} \right)^{\rho^*}. \quad (2.3)$$

For a more detailed study of the class \mathcal{A}_{∞} , one can refer to [22, Chapter V] or, for example, to [9, Lemma 1.3], [8, Lemma 2.3] and [9, Lemma 1.4] (and the literature therein).

2.2 Weights from the local Muckenhoupt class A_∞^{loc}

The next class of weights covers admissible [7], locally regular [12, 19, 20] as well as A_∞ -weights.

Definition 2.2. (i) Weight w is of *local Muckenhoupt type* A_ρ^{loc} , $1 < \rho < \infty$, if (see. (2.1))

$$A_\rho^{\text{loc}}(w) := \sup_{|Q| \leq 1} A_\rho[w(Q)] < \infty;$$

(ii) $w \in A_1^{\text{loc}}$ if $A_1^{\text{loc}}(w) := \sup_{|Q| \leq 1} A_1[w(Q)] < \infty$;

(iii) we say that $w \in A_\infty^{\text{loc}}$ if $w \in A_\rho^{\text{loc}}$ for some $1 \leq \rho < \infty$, that is

$$A_\infty^{\text{loc}} := \bigcup_{\rho \geq 1} A_\rho^{\text{loc}}.$$

The weights of class A_∞^{loc} largely repeat the properties of the weights from A_∞ [17]. In particular, by analogy with r_0 in the case of $w \in A_\infty$, there exists a number

$$r_w := \inf\{r \geq 1 : w \in A_r^{\text{loc}}\} < \infty, \quad w \in A_\infty^{\text{loc}}. \quad (2.4)$$

The doubling type property for $w \in A_\infty^{\text{loc}}$ can be formulated as follows: if $w \in \mathcal{A}_\rho^{\text{loc}}$, $1 < \rho \leq \infty$, then there exists a constant $c_w > 0$ such that the estimate

$$w(Q_t) \leq \exp(c_w t/s)w(Q_s)$$

holds for all Q_s with $|Q_s| = s$, $0 < s \leq 1$ and Q_t such that $Q_s \subset Q_t$, where $|Q_t| = t$, $t \geq 1$.

Let us present *basic property* of A_ρ^{loc} -weights with respect to A_ρ -functions: any $w \in A_\rho^{\text{loc}}$ can be continued outside a given cube Q in such a way that a new extended weight \bar{w} belongs to A_ρ .

Lemma 2.1 ([17, Lemma 1.1]). *Let $1 \leq \rho < \infty$, $w \in A_\rho^{\text{loc}}$ and Q be a cube with $|Q| = 1$. Then there is $\bar{w} \in A_\rho$ such that $\bar{w} = w$ on Q and with a constant c independent of Q the following holds:*

$$A_\rho(\bar{w}) \leq cA_\rho^{\text{loc}}(w).$$

More information about local Muckenhoupt weights can be found in [12, 17, 34, 35].

2.3 Weighted Besov type spaces

Let $p > 0$, $0 < q \leq \infty$ and $s \in \mathbb{R}$. Definitions of unweighted Besov type spaces $B_{pq}^s(\mathbb{R})$ are given in [23, 24]. For a given $w \not\equiv 1$, their weighted generalisations can be specified in several ways depending on which class w belongs to [9, 16, 17, 19].

The space $\mathcal{S}'(\mathbb{R})$, which is topologically dual to the space $\mathcal{S}(\mathbb{R})$ of Schwartz functions, is used in unweighted case, as well as in situations when w belongs to the Muckenhoupt class A_∞ . For $\mathcal{A}_\infty^{\text{loc}}$ -weights, a more general space of distributions $\mathcal{S}'_e(\mathbb{R})$ is required than $\mathcal{S}'(\mathbb{R})$.

Definition 2.3. The space $\mathcal{S}_e(\mathbb{R})$ consists of all functions $\phi \in C^\infty(\mathbb{R})$ such that

$$q_N(\phi) := \sup_{x \in \mathbb{R}} e^{N|x|} \sum_{0 \leq \gamma \leq N} |D^\gamma \phi(x)| < \infty \quad \text{for any } N \in \mathbb{N}_0.$$

The set $\mathcal{S}_e(\mathbb{R})$ is equipped with a locally convex topology defined by the system of seminorms q_N . More details on the properties of $\mathcal{S}_e(\mathbb{R})$ can be found in [17, 19] and [12, § 3].

By $\mathcal{S}'_e(\mathbb{R})$ we denote the space of all linear continuous functionals on $\mathcal{S}_e(\mathbb{R})$. The class $\mathcal{S}'_e(\mathbb{R})$ is sometimes conveniently identified with the subset in $\mathcal{D}'(\mathbb{R})$ of all distributions f on $\mathcal{D}(\mathbb{R})$ for which the estimate

$$|\langle f, \phi \rangle| \leq C \sup \left\{ |D^\gamma \phi(x)| e^{N|x|} : x \in \mathbb{R}, 0 \leq \gamma \leq N \right\}$$

is satisfied for any $\phi \in \mathcal{D}(\mathbb{R})$ with some constants C and N depending on f . Each such f can be extended to a continuous functional on $\mathcal{S}_e(\mathbb{R})$.

For $\phi, \psi \in \mathcal{S}_e(\mathbb{R})$, the convolution $\phi * \psi$ belongs to $\mathcal{S}_e(\mathbb{R})$. If $f \in \mathcal{S}'_e(\mathbb{R})$ and $\phi \in \mathcal{S}_e(\mathbb{R})$ then the convolution

$$f * \phi(x) =: \langle f(\cdot), \phi(x - \cdot) \rangle, \quad x \in \mathbb{R},$$

is a function in $C^\infty(\mathbb{R})$ of at most exponential growth.

To define Besov spaces with Muckenhoupt weights of local type, we take $\phi_0 \in \mathcal{D}(\mathbb{R})$ such that

$$\int_{\mathbb{R}} \phi_0(x) dx \neq 0, \quad (2.5)$$

further, set

$$\phi(x) = \phi_0(x) - 2^{-1}\phi_0(x/2) \quad (2.6)$$

and define $\phi_d(x) := 2^{(d-1)}\phi(2^{d-1}x)$ for $d \in \mathbb{N}$. One can choose ϕ_0 satisfying

$$\int_{\mathbb{R}} x^\gamma \phi(x) dx = 0 \quad (2.7)$$

for all $\gamma \in \mathbb{N}_0$, $\gamma \leq \Gamma$, with $\Gamma \in \mathbb{N}$. We write $\Gamma = -1$ if (2.7) does not hold.

Definition 2.4. Let (2.5) hold for some $\phi_0 \in \mathcal{D}(\mathbb{R})$, and let ϕ of the form (2.6) satisfy (2.7) with $0 \leq \gamma \leq \Gamma$, where $\Gamma \geq [s]$. For $w \in A_\infty^{\text{loc}}$, we define the weighted Besov type space $B_{pq}^{s,w}(\mathbb{R})$ of distributions as the set of all $f \in \mathcal{S}'_e(\mathbb{R})$ such that the (quasi-)norm

$$\|f\|_{B_{pq}^{s,w}(\mathbb{R})} := \left(\sum_{d=0}^{\infty} 2^{dsq} \|\phi_d * f\|_{L^p(\mathbb{R},w)}^q \right)^{\frac{1}{q}}$$

(with standard modification for $q = \infty$) is finite. Here for $p > 0$ and a weight w the symbol $L^p(\mathbb{R}, w)$ denotes the weighted Lebesgue space with the norm $\|f\|_{L^p(\mathbb{R},w)} := \|w^{1/p}f\|_{L^p(\mathbb{R})}$.

The definition of $B_{pq}^{s,w}(\mathbb{R})$ does not depend on the choice of the function ϕ_0 . It generalises the notion of Besov type smoothness spaces with $w \in A_\infty$ [17, § 2.2 and Remark 2.23].

The spaces $B_{pq}^{s,w}(\mathbb{R})$ are similar in their properties to unweighted $B_{pq}^s(\mathbb{R})$, as well as to the classes $B_{pq}^s(\mathbb{R}, w)$ with $w \in A_\infty$ (for more details, see [10, 12, 17, 34, 35]).

Next, for $\tau \in \mathbb{Z}$ and $d \in \mathbb{N}_0$ we define dyadic segments $Q_{d\tau} := [2^{-d}\tau, 2^{-d}(\tau + 1)]$ with the origin at the point $x_{Q_{d\tau}} := 2^{-d}\tau$. In what follows, we will need Besov type spaces $b_{pq}^s(w)$ of sequences $\lambda = \{\lambda_{d\tau}\}_{d \in \mathbb{N}_0, \tau \in \mathbb{Z}}$, $\lambda_{d\tau} \in \mathbb{C}$, such that

$$\|\lambda\|_{b_{pq}^s(w)} := \left\| \sum_{\tau \in \mathbb{Z}} |\lambda_{0\tau}| \chi_{Q_{0\tau}} \right\|_{L^p(\mathbb{R},w)} + \left(\sum_{d \in \mathbb{N}_0} 2^{qds} \left\| \sum_{\tau \in \mathbb{Z}} |\lambda_{d\tau}| \chi_{Q_{(d-1)\tau}} \right\|_{L^p(\mathbb{R},w)}^q \right)^{1/q} < \infty.$$

In the definition of $\|\lambda\|_{b_{pq}^s(w)}$, we can take $Q_{(d-1)(\tau \pm 1/2)}$ instead of $Q_{(d-1)\tau}$ in the second term. Due to (2.2) property of Muckenhoupt weights (see also Lemma 2.1), this is allowed in this situation.

3 Criteria for the fulfillment of norm inequalities

The results obtained in this section are based on decomposition theorems those relate elements f of Besov spaces $B_{pq}^{s,w}(\mathbb{R})$ with sequential spaces $b_{pq}^s(w)$, as well as the corresponding to these spaces norms. To prove them, we decompose f with respect to spline wavelet systems of natural orders.

3.1 Decompositions in terms of spline wavelets

Characterization of the spaces $B_{pq}^{s,w}(\mathbb{R})$ in terms of spline wavelets of natural orders were given in [29, 33] (see also [28]). In particular, it was established that

$$\|f\|_{B_{pq}^{s,w}(\mathbb{R})} \approx \|\tilde{\lambda}^{(k,0)}\|_{b_{pq}^s(w)}, \quad (3.1)$$

where elements

$$\tilde{\lambda}_{0\tau} = \langle f, \tilde{\Phi}_\tau \rangle \quad (\tau \in \mathbb{Z}), \quad \tilde{\lambda}_{d\tau}^{(k,0)} = 2^{d/2} \langle f, \tilde{\Psi}_{(d-1)\tau}^{(k,0)} \rangle \quad (d \in \mathbb{N}, \tau \in \mathbb{Z}) \quad (3.2)$$

of sequences $\tilde{\lambda}^{(k,0)} = \{\tilde{\lambda}_{0,\tau}\}_{\tau \in \mathbb{Z}} \cup \{\tilde{\lambda}_{d\tau}^{(k,0)}\}_{d \in \mathbb{N}, \tau \in \mathbb{Z}}$ c $k \in \{0, 1\}$ are defined by $\tilde{\Phi}_\tau$ и $\tilde{\Psi}_{(d-1)\tau}^{(k,0)}$ of the form (5.15) with possible "starting" points $\mathbf{a} \in \{0, \pm 1/2\}$. In [29], the case $\mathbf{a} = 0$ was considered, while in [28] — an even simpler situation $\mathbf{a} = k = 0$.

The result (3.1) allows for some generalization (see Theorem 3.3). Let us start, however, with the base case $\zeta = k = 0$. For this, taking into account (2.4), we denote

$$\sigma_p(w) := r_w / \min\{p, r_w\} - 2 + r_w.$$

We will assume that $\mathbf{s} = 0$ in the definitions of $\Psi^{(\mathbf{a})} \in (5.14)$ and $\tilde{\Psi}^{(k,\zeta)} \in (5.13)$, $k, \zeta \in \{0, 1\}$.

Theorem 3.1 ([29, 33]). *Let $p > 0$, $0 < q \leq \infty$, $s \in \mathbb{R}$ and $w \in A_\infty^{\text{loc}}$. Let $\Phi^{(\mathbf{a})}$ and $\Psi^{(\mathbf{a})}$, where $\mathbf{a} \in \{0, \pm 1/2\}$, be functions satisfying (5.14) with some $n \in \mathbb{N}$ such that*

$$n \geq \max\left\{0, [s] + 1, [(r_w - 1)/p - s] + 1, [\sigma_p(w) - s]\right\} + 1. \quad (3.3)$$

Then $f \in \mathcal{S}'_e(\mathbb{R})$ belongs to $B_{pq}^{s,w}(\mathbb{R})$ if and only if the representation holds

$$f = \sum_{\tau \in \mathbb{Z}} \lambda_{0\tau}^{(\mathbf{a})} \Phi_\tau^{(\mathbf{a})} + \sum_{d \in \mathbb{N}} \sum_{\tau \in \mathbb{Z}} \lambda_{d\tau}^{(\mathbf{a})} 2^{-d/2} \Psi_{(d-1)\tau}^{(\mathbf{a})}, \quad (3.4)$$

where $\{\lambda_{d\tau}^{(\mathbf{a})}\}_{d \in \mathbb{N}_0, \tau \in \mathbb{Z}} =: \lambda^{(\mathbf{a})} \in b_{pq}^s(w)$ and the series converges in $\mathcal{S}'_e(\mathbb{R})$. The representation (3.4) is unique with

$$\lambda_{0\tau}^{(\mathbf{a})} = \langle f, \Phi_\tau^{(\mathbf{a})} \rangle \quad (\tau \in \mathbb{Z}), \quad \lambda_{d\tau}^{(\mathbf{a})} = 2^{d/2} \langle f, \Psi_{(d-1)\tau}^{(\mathbf{a})} \rangle \quad (d \in \mathbb{N}, \tau \in \mathbb{Z}), \quad (3.5)$$

and $I: f \mapsto \lambda^{(\mathbf{a})}$ is a linear isomorphism of $B_{pq}^{s,w}(\mathbb{R})$ onto $b_{pq}^s(w)$. Moreover,

$$\|f\|_{B_{pq}^{s,w}(\mathbb{R})} \approx \|\lambda^{(\mathbf{a})}\|_{b_{pq}^s(w)}, \quad (3.6)$$

where the equivalence constants depend only on the numerical parameters.

Below there is a formulation of an analogue of Theorem 3.1, which uses a dictionary of functions $\tilde{\Phi}$ and $\tilde{\Psi}^{(k,0)}$ of the form (5.13) with $k = 1$ and $\zeta = 0$. It also asserts (3.1) (see [33, Theorem 4.8]), but with different representations for f in its sufficiency and necessity parts.

Theorem 3.2. *Let $p > 0$, $0 < q \leq \infty$, $s \in \mathbb{R}$ and $w \in A_\infty^{\text{loc}}$. Suppose that for some $n \in \mathbb{N}$ the condition (3.3) is satisfied, and there are functions $\Phi^{(\mathbf{a})}$ and $\Psi^{(\mathbf{a})}$ with $\mathbf{a} \in \{0, \pm 1/2\}$, satisfying (5.14), as well as functions $\tilde{\Phi}$ and $\tilde{\Psi}^{(1,0)}$ of the form (5.13), both the pairs of smoothness n .*

(i) *A distribution $f \in \mathcal{S}'_e(\mathbb{R})$ belongs to $B_{pq}^{s,w}(\mathbb{R})$, if there is a representation*

$$f = \sum_{\tau \in \mathbb{Z}} \tilde{\lambda}_{0\tau} \tilde{\Phi}_\tau + \sum_{d \in \mathbb{N}} \sum_{\tau \in \mathbb{Z}} \tilde{\lambda}_{d\tau}^{(1,0)} 2^{-d/2} \tilde{\Psi}_{(d-1)\tau}^{(1,0)}, \quad (3.7)$$

where $\tilde{\lambda}^{(1,0)}$ of the form (3.2) is from $b_{pq}^s(w)$ and the series (3.7) converges in $\mathcal{S}'_e(\mathbb{R})$. Moreover,

$$\|f\|_{B_{pq}^{s,w}(\mathbb{R})} \lesssim \|\tilde{\lambda}^{(1,0)}\|_{b_{pq}^s(w)} \lesssim \sum_{\mathbf{a}=0,\pm 1/2} \|\lambda^{(\mathbf{a})}\|_{b_{pq}^s(w)}$$

with $\lambda^{(\mathbf{a})}$ of the form (3.5).

(ii) Conversely, if $f \in \mathcal{S}'_e(\mathbb{R})$ belongs to $B_{pq}^{s,w}(\mathbb{R})$, then for $\mathbf{a} \in \{0, \pm 1/2\}$ we have (3.4) in the sense of convergence in $\mathcal{S}'_e(\mathbb{R})$, where $\lambda^{(\mathbf{a})} \in b_{pq}^s(w)$ is given by (3.5). Moreover,

$$\|\tilde{\lambda}^{(1,0)}\|_{b_{pq}^s(w)} \leq \sum_{\mathbf{a}=0,1/2} \|\lambda^{(\mathbf{a})}\|_{b_{pq}^s(w)} \lesssim \|f\|_{B_{pq}^{s,w}(\mathbb{R})}.$$

A generalisation of Theorem 3.2 in terms of $\tilde{\Phi}$ and $\tilde{\Psi}^{(1,1)}$ of the form (5.13) is given below.

Theorem 3.3. *Let $p > 0$, $0 < q \leq \infty$, $s \in \mathbb{R}$ and $w \in A_\infty^{\text{loc}}$. Suppose that for some $n \in \mathbb{N}$ the condition (3.3) is satisfied, and there are functions $\Phi^{(\mathbf{a})}$ and $\Psi^{(\mathbf{a})}$ with $\mathbf{a} \in \{0, \pm 1/2\}$, satisfying (5.14), as well as functions $\tilde{\Phi}$ and $\tilde{\Psi}^{(1,1)}$ of the form (5.13), the both pairs of smoothness n .*

(i) *A distribution $f \in \mathcal{S}'_e(\mathbb{R})$ belongs to $B_{pq}^{s,w}(\mathbb{R})$ if there is a representation*

$$f = \sum_{\tau \in \mathbb{Z}} \tilde{\lambda}_{0\tau} \tilde{\Phi}_\tau + \sum_{d \in \mathbb{N}} \sum_{\tau \in \mathbb{Z}} \tilde{\lambda}_{d\tau}^{(1,1)} 2^{-d/2} \tilde{\Psi}_{(d-1)\tau}^{(1,1)},$$

where the sequence $\tilde{\lambda}^{(1,1)} = \{\tilde{\lambda}_{0,\tau}\}_{\tau \in \mathbb{Z}} \cup \{\tilde{\lambda}_{d\tau}^{(1,1)}\}_{d \in \mathbb{N}, \tau \in \mathbb{Z}}$ of the form

$$\tilde{\lambda}_{0\tau} = \langle f, \tilde{\Phi}_\tau \rangle \quad (\tau \in \mathbb{Z}), \quad \tilde{\lambda}_{d\tau}^{(1,1)} = 2^{d/2} \langle f, \tilde{\Psi}_{(d-1)\tau}^{(1,1)} \rangle \quad (d \in \mathbb{N}, \tau \in \mathbb{Z}) \quad (3.8)$$

is from $b_{pq}^s(w)$ with $\tilde{\Phi}_\tau$ and $\tilde{\Psi}_{(d-1)\tau}^{(1,1)}$ given by (5.15) with $\mathbf{a} = \{0, \pm 1/2\}$. In this case, the series (3.7) converges in $\mathcal{S}'_e(\mathbb{R})$. In addition, with $\lambda^{(\mathbf{a})}$ of the form (3.5), we have

$$\|f\|_{B_{pq}^{s,w}(\mathbb{R})} \lesssim \|\tilde{\lambda}^{(1,1)}\|_{b_{pq}^s(w)} \lesssim \sum_{\mathbf{a}=0,\pm 1/2} \|\lambda^{(\mathbf{a})}\|_{b_{pq}^s(w)}. \quad (3.9)$$

(ii) Conversely, if $f \in \mathcal{S}'_e(\mathbb{R})$ belongs to $B_{pq}^{s,w}(\mathbb{R})$, then the (3.4) holds for $\mathbf{a} \in \{0, \pm 1/2\}$ in the sense of convergence in $\mathcal{S}'_e(\mathbb{R})$, where $\lambda^{(\mathbf{a})} \in b_{pq}^s(w)$ is given by (3.5). Besides,

$$\|\tilde{\lambda}^{(1,1)}\|_{b_{pq}^s(w)} \leq \sum_{\mathbf{a}=0,1/2} \|\lambda^{(\mathbf{a})}\|_{b_{pq}^s(w)} \lesssim \|f\|_{B_{pq}^{s,w}(\mathbb{R})}.$$

Observe that in the original decomposition theorem using spline wavelet systems of natural orders (see [29, § 4.3]) the case $\mathbf{a} = 0$ was considered. Later in [33, Theorem 4.8] the cases $\mathbf{a} = \pm 1/2$ were added, they are justified by the (2.2) property of Muckenhoupt weights and Lemma 2.1. The proof of Theorem 3.3 is similar to the proof of Theorem 3.2 (see [33, Theorem 4.8], as well as [29, Theorem 4.3]). The basis for it is the fact that the elements of the sequence $\tilde{\lambda}^{(1,1)}$ form a family of atoms [10, 34] (see also [29, Definition 4.1]).

3.2 The main result

Let us introduce into consideration a functional

$$\mathfrak{M}(d, \kappa; \sigma_1, \sigma_2) := \sup_{\tau \in \mathbb{Z}} \begin{cases} [\sigma_1(Q_{0\tau})]^{-\frac{1}{p}} [\sigma_2(Q_{0\tau})]^{\frac{1}{p}}, & d = 0, \\ 2^{-d\kappa} [\sigma_1(Q_{(d-1)\tau})]^{-\frac{1}{p}} [\sigma_2(Q_{(d-1)\tau})]^{\frac{1}{p}}, & d \in \mathbb{N}. \end{cases}$$

Theorem 3.4. Let $p > 1$, $0 < q \leq \infty$, $s \in \mathbb{R}$, weights $u, v \in \mathcal{A}_\infty^{\text{loc}}$ and $\alpha \in \mathbb{N}$. Suppose that $f \in L^1(-\infty, x)$ for any $x \in \mathbb{R}$ in the case of I_\pm^α and $f \in L^1(x, +\infty)$ for all $x \in \mathbb{R}$ for I_\pm^α .

(i) The operator I_\pm^α is bounded from $B_{pq}^{s+\kappa^*-\alpha, v}(\mathbb{R})$ to $B_{pq}^{s, u}(\mathbb{R})$ with some $\kappa^* \in \mathbb{R}$ if and only if

$$\mathfrak{C}_\pm^\alpha(\kappa^*) = \mathbf{M}_\pm^\alpha(1) + \mathbf{M}_\pm^\alpha(0) + \sup_{d \in \mathbb{N}} \mathfrak{M}(d, \kappa^*; v, u) < \infty. \quad (3.10)$$

In addition, the norm $\|I_\pm^\alpha\|_{B_{pq}^{s+\kappa^*-\alpha, v}(\mathbb{R}) \rightarrow B_{pq}^{s, u}(\mathbb{R})}$ of the operator $I_\pm^\alpha: B_{pq}^{s+\kappa^*-\alpha, v}(\mathbb{R}) \rightarrow B_{pq}^{s, u}(\mathbb{R})$ is equivalent to $\mathfrak{C}_\pm^\alpha(\kappa^*)$ with constants depending only on the numerical parameters.

(ii) Let $\kappa_* \in \mathbb{R}$ and suppose that $0 < \mathfrak{C}$ is the best constant of the inequality

$$\|f\|_{B_{pq}^{s-\kappa_*-\alpha, w}(\mathbb{R})} \lesssim \mathfrak{C} \|I_\pm^\alpha f\|_{B_{pq}^{s, u}(\mathbb{R})} \quad (3.11)$$

independent of f . Inequality (3.11) holds if and only if

$$\mathfrak{C}(\kappa_*) := \sup_{d \in \mathbb{N}_0} \mathfrak{M}(d, \kappa_*; u, w) < \infty,$$

and $\mathfrak{C} \approx \mathfrak{C}(\kappa_*)$. In particular, if $w = u$, then $\mathfrak{C} \approx \mathfrak{C}(0) = 1$.

Proof. (i) *Sufficiency.* Let us turn to the beginning of the proof of assertion (i) from [33, Theorem 5.5] in the sufficiency part and show the validity of

$$\|I_+^\alpha\|_{B_{pq}^{s+\kappa^*-\alpha, v}(\mathbb{R}) \rightarrow B_{pq}^{s, u}(\mathbb{R})} \lesssim \mathfrak{C}_+^\alpha(\kappa^*).$$

To do this, we choose $n^* \in \mathbb{N}$ satisfying the condition (3.3) with respect to parameters of the space $B_{pq}^{s, u}(\mathbb{R})$ including r_u (see (2.4)). Besides, we set $m^* := n^* + \alpha$, which must also satisfy (3.3) with respect to parameters of the space $B_{pq}^{s+\kappa^*-\alpha, v}(\mathbb{R})$ and r_v . Then, by (3.9) (see Theorem 3.3(i)),

$$\|I_+^\alpha f\|_{B_{pq}^{s, u}(\mathbb{R})} \approx \|\tilde{\lambda}^{\langle 1, 1 \rangle}\|_{b_{pq}^s(u)},$$

where $\tilde{\lambda}^{\langle 1, 1 \rangle}$ is defined by (3.8) with $I_+^\alpha f$ instead of f . Besides, $\tilde{\Phi}$ is given by B_{n^*} , and $\tilde{\Psi}^{\langle 1, 1 \rangle}$ (see (5.13)) is specified by $\Psi_{n^*, 0, 0; m^*(1), \alpha(1)}$. This means that we can write:

$$\|I_+^\alpha f\|_{B_{pq}^{s, u}(\mathbb{R})} \approx \left\| \sum_{\tau \in \mathbb{Z}} |\tilde{\lambda}_{0\tau}| \chi_{Q_{0\tau}} \right\|_{L^p(\mathbb{R}, u)} + \left(\sum_{d \in \mathbb{N}} 2^{qd} \left\| \sum_{\tau \in \mathbb{Z}} |\tilde{\lambda}_{d\tau}^{\langle 1, 1 \rangle}| \chi_{Q_{(d-1)\tau}} \right\|_{L^p(\mathbb{R}, u)}^q \right)^{1/q}, \quad (3.12)$$

where

$$\tilde{\lambda}_{0\tau} = \langle I_+^\alpha f, \tilde{\Phi}_\tau \rangle \quad (\tau \in \mathbb{Z}), \quad \tilde{\lambda}_{d\tau}^{\langle 1, 1 \rangle} = 2^{d/2} \langle I_+^\alpha f, \tilde{\Psi}_{(d-1)\tau}^{\langle 1, 1 \rangle} \rangle \quad (d \in \mathbb{N}, \tau \in \mathbb{Z}).$$

The first term on the right-hand side of (3.12) can be estimated by repeating a similar stage of the proof of assertion (i) from [33, Theorem 5.5]. We present it here in a slightly modified form, by relying on the differential property (5.1) of the basic splines B_n .

As in the proof of [33, Theorem 5.5], we denote $\tilde{\Phi}_\tau(\cdot) = B_{n^*}(\cdot - \tau) =: P(\tau)$ and set

$$R(\tau - r) := \sum_{j=0}^{\alpha} (-1)^j \binom{\alpha}{j} P(\tau - r - j).$$

It is true that

$$P(\tau - r) = R(\tau - r) - \sum_{j=1}^{\alpha} (-1)^j \binom{\alpha}{j} P(\tau - r - j). \quad (3.13)$$

Then

$$P(\tau) = R(\tau) - \sum_{j=1}^{\alpha} (-1)^j \binom{\alpha}{j} P(\tau - j) = R(\tau) + \binom{\alpha}{1} P(\tau - 1) - \sum_{j=2}^{\alpha} (-1)^j \binom{\alpha}{j} P(\tau - j).$$

Again on the strength of (3.13),

$$\begin{aligned} P(\tau) - R(\tau) &= \binom{\alpha}{1} \left[R(\tau - 1) - \sum_{j=1}^{\alpha} (-1)^j \binom{\alpha}{j} P(\tau - 1 - j) \right] - \sum_{j=2}^{\alpha} (-1)^j \binom{\alpha}{j} P(\tau - j) \\ &= \binom{\alpha}{1} R(\tau - 1) - \binom{\alpha}{1} \sum_{j=1}^{\alpha} (-1)^j \binom{\alpha}{j} P(\tau - 1 - j) - \sum_{j=2}^{\alpha} (-1)^j \binom{\alpha}{j} P(\tau - j) = \binom{\alpha}{1} R(\tau - 1) \\ &\quad + \left[\binom{\alpha}{1}^2 - \binom{\alpha}{2} \right] P(\tau - 2) - \binom{\alpha}{1} \sum_{j=2}^{\alpha} (-1)^j \binom{\alpha}{j} P(\tau - 1 - j) - \sum_{j=3}^{\alpha} (-1)^j \binom{\alpha}{j} P(\tau - j). \end{aligned}$$

Setting $A_0 := \binom{\alpha}{0} = 1$, $A_1 := A_0 \binom{\alpha}{1}$, $A_2 := A_1 \binom{\alpha}{1} - A_0 \binom{\alpha}{2} = \binom{\alpha}{1}^2 - \binom{\alpha}{2}$, we continue the procedure for forming finite iterated differences as shown below

$$\begin{aligned} P(\tau) &= A_0 R(\tau) + A_1 R(\tau - 1) + A_2 \left[R(\tau - 2) - \sum_{j=1}^{\alpha} (-1)^j \binom{\alpha}{j} P(\tau - 2 - j) \right] \\ &\quad - \binom{\alpha}{1} \sum_{j=2}^{\alpha} (-1)^j \binom{\alpha}{j} P(\tau - 1 - j) - \sum_{j=3}^{\alpha} (-1)^j \binom{\alpha}{j} P(\tau - j) \\ &= A_0 R(\tau) + A_1 R(\tau - 1) + A_2 R(\tau - 2) - \underbrace{\left[A_2 \binom{\alpha}{1} - A_1 \binom{\alpha}{2} + A_0 \binom{\alpha}{3} \right]}_{:=A_3} P(\tau - 3) \\ &\quad - A_2 \sum_{j=2}^{\alpha} (-1)^j \binom{\alpha}{j} P(\tau - 2 - j) - \binom{\alpha}{1} \sum_{j=3}^{\alpha} (-1)^j \binom{\alpha}{j} P(\tau - 1 - j) - \sum_{j=4}^{\alpha} (-1)^j \binom{\alpha}{j} P(\tau - j), \end{aligned}$$

which is resulting in the sum of the form

$$P(\tau) = \sum_{r \geq 0} A_r R(\tau - r), \quad \text{where } A_r = A_r(\alpha) = \sum_{j=1}^r (-1)^{j-1} A_{r-j} \binom{\alpha}{j}.$$

In [33], it was established that $A_r(\alpha) \leq (1+r)^{\alpha-1}$ (see proof of (5.18) in [33]).

Similar procedure of forming differences can be done by "moving to the right side" and assuming

$$R^*(\tau + r) := \sum_{j=0}^{\alpha} (-1)^j \binom{\alpha}{j} P(\tau + r + j).$$

Then

$$P(\tau) = \sum_{r \geq 0} A_r R^*(\tau + r), \quad \text{where } R(\tau + r)^* = B_m^{(\alpha)}(\cdot - \tau - r).$$

Since we have $f \in L^1(-\infty, x)$ for any $x \in \mathbb{R}$, also

$$\text{supp } B_m^{(\alpha)}(\cdot - \tau + r + \alpha) = \text{supp } B_{m^*, \tau - r - \alpha}^{(\alpha)}(\cdot) = [\tau - r - \alpha, \tau - r + n^* + 1]$$

and $B_m^{(k)}(\cdot) = 0$ at the points $\tau - r - \alpha$ and $\tau - r + n^* + 1$ for all $k = 0, \dots, \alpha$ (see §5), then

$$\int_{\mathbb{R}} I_+^{\alpha} f(x) B_m^{(\alpha)}(x - \tau - r) dx = (-1)^{\alpha} \int_{\mathbb{R}} f(y) B_{m^*}(y - \tau + r + \alpha) dy. \quad (3.14)$$

Hence and with respect to the equality $R(\tau - r) = (-1)^\alpha B_{m^*}^{(\alpha)}(\cdot - \tau + r + \alpha)$ (see (5.1)), the first term in (3.12) can be estimated as follows:

$$\begin{aligned}
\left\| \sum_{\tau \in \mathbb{Z}} |\tilde{\lambda}_{0\tau}| \chi_{Q_{0\tau}} \right\|_{L^p(\mathbb{R}, u)}^p &= \sum_{\tau \in \mathbb{Z}} u(Q_{0\tau}) |\langle I_+^\alpha f, \tilde{\Phi}_\tau \rangle|^p \\
&= \sum_{\tau \in \mathbb{Z}} u(Q_{0\tau}) \left[\left| \sum_{r \geq 0} A_r \int_{\mathbb{R}} I_+^\alpha f(x) B_{m^*}^{(\alpha)}(x - \tau + r + \alpha) dx \right|^p \right] \\
&= \Gamma(\alpha) \sum_{\tau \in \mathbb{Z}} u(Q_{0\tau}) \left[\left| \sum_{r \geq 0} A_r \int_{\mathbb{R}} f(y) B_{m^*}(y - \tau + r + \alpha) dy \right|^p \right] \\
&\lesssim \sum_{\tau \in \mathbb{Z}} u(Q_{0\tau}) \left[\sum_{r \geq 0} (r+1)^{\alpha-1} \left| \int_{\mathbb{R}} f(y) B_{m^*}(y - \tau + r + \alpha) dy \right|^p \right] \\
&= \sum_{\tau \in \mathbb{Z}} u(Q_{0\tau}) \left[\sum_{l \leq \tau} (\tau - l + 1)^{\alpha-1} \left| \int_{\mathbb{R}} f(y) B_{m^*}(y + \alpha - l) dy \right|^p \right].
\end{aligned}$$

Further, by virtue of (3.10),

$$\sum_{\tau \in \mathbb{Z}} u(Q_{0\tau}) |\langle I_+^\alpha f, \tilde{\Phi}_\tau \rangle|^p \lesssim [\mathbf{M}_+^\alpha(1) + \mathbf{M}_+^\alpha(0)]^p \sum_{\tau \in \mathbb{Z}} v(Q_{0\tau}) \left| \int_{\mathbb{R}} f(y) B_{m^*}(y + \alpha - \tau) dy \right|^p \quad (3.15)$$

(see [33, Theorem 5.1] or [15, Theorems 1.6 & 1.11]). Using (2.2) and relying on Lemma 2.1, we derive an estimate of the form

$$v(Q_{0\tau}) = \int_{Q_{0\tau}} v \leq \int_{\cup_{i=\tau-\alpha}^{\tau} Q_{0i}} v \leq \alpha^{r_v} \int_{Q_{0(\tau-\alpha)}} v.$$

Then, by substituting $r = \tau - \alpha$, we can transform (3.15) to

$$\begin{aligned}
\left\| \sum_{\tau \in \mathbb{Z}} |\lambda_{0\tau}| \chi_{Q_{0\tau}} \right\|_{L^p(\mathbb{R}, u)} &= \left(\sum_{\tau \in \mathbb{Z}} u(Q_{0\tau}) |\langle I_+^\alpha f, \tilde{\Phi}_\tau \rangle|^p \right)^{1/p} \\
&\lesssim \mathfrak{e}_+^\alpha(\kappa^*) \left(\sum_{r \in \mathbb{Z}} v(Q_{0r}) \left| \int_{\mathbb{R}} f(y) B_{m^*}(y - r) dy \right|^p \right)^{1/p}.
\end{aligned}$$

We set $\int_{\mathbb{R}} f(y) B_{m^*}(y - \tau) dy =: \langle f, \bar{\Phi}_\tau \rangle$ and name $\bar{\Phi}$ as the first component of a new wavelet system forming spline basis of order m^* . The second component of this system, as well as the elements it generates, will be denoted by $\tilde{\Psi}^{(1,0)}$ and $\tilde{\Psi}_{(d-1)\tau}^{(1,0)}$, respectively.

The next step is to estimate $\tilde{\lambda}_{d\tau}^{(1,1)}$ for $d > 0$ on the right-hand side (3.12) via $\langle f, \tilde{\Psi}_{(d-1)\tau}^{(1,0)} \rangle$. To do this, we write out an exact expression for the element $\tilde{\Psi}_{(d-1)\tau}^{(1,1)}$, based on (5.10) and (5.12):

$$\begin{aligned}
2^{-d/2} \gamma_{n^*} \mathbf{\Lambda}_{n^*} \tilde{\Psi}_{(d-1)\tau}^{(1,1)}(\cdot) &= \gamma_{n^*} \Psi_{n^*, 0, 0; m^*(1), \alpha(1)}(2^{d-1} \cdot -\tau) \\
&= \sum_{|l| \leq m^*} \lambda_{|l|}(m^*) (-1)^{|l|} \sum_{|s| \leq n^*} \lambda_{|s|}(n^*) (-1)^{|s|} \\
&\quad \times \sum_{k=0}^{2\alpha+n^*+1} (-1)^k \binom{1+m^*+2\alpha}{k} B_{n^*}(2^d \cdot -2\tau + (n^* - k) - s - l).
\end{aligned}$$

We will also need the Zhu–Vandermonde identity [18, p. 15]

$$\binom{r+s}{k} = \sum_{n=0}^k \binom{r}{n} \binom{s}{k-n}, \quad \text{where } r, s \in \mathbb{R} \text{ and } k \in \mathbb{N}, \quad (3.16)$$

from which it follows for $r = 1 + m^*$ and $s = \alpha$ that

$$\begin{aligned}
& \sum_{k=0}^{2\alpha+n^*+1} (-1)^k \binom{1+n^*+2\alpha}{k} B_{n^*}(2^d x - 2\tau + (m^* - k) - s - l) \\
&= \sum_{k=0}^{2\alpha+n^*+1} (-1)^k \sum_{r=0}^k \binom{1+m^*}{r} \binom{\alpha}{k-r} B_{n^*}(2^d x - 2\tau + (m^* - k) - s - l) \\
&= \sum_{r=0}^{m^*+1} (-1)^r \binom{1+m^*}{r} \sum_{m=0}^{\alpha} (-1)^m \binom{\alpha}{m} B_{n^*}(2^d x - 2\tau + (m^* - r - m) - s - l).
\end{aligned}$$

Observe that

$$\begin{aligned}
& \sum_{m=0}^{\alpha} (-1)^m \binom{\alpha}{m} B_{n^*}(2^d x - 2\tau + (m^* - r - m) - s - l) \\
&= 2^{-d\alpha} B_{m^*}^{(\alpha)}(2^d x - 2\tau + (m^* - r) - s - l). \quad (3.17)
\end{aligned}$$

Also,

$$\begin{aligned}
& \text{supp } B_{m^*}^{(\alpha)}(2^d x - 2\tau + (m^* - r) - s - l) =: \text{supp } B_{m^*, 2\tau+s+l+r-m^*}^{(\alpha)}(2^d \cdot) \\
&= \left[\frac{2\tau + s + l + r - m^*}{2^d}, \frac{2\tau + s + l + r + 1}{2^d} \right]
\end{aligned}$$

and $B_{m^*, 2\tau+s+l+r-m^*}^{(\alpha)}(2^d \cdot) = 0$ at the points $2^{-d}(2\tau + s + l + r - m^*)$ and $2^{-d}(2\tau + s + l + r + 1)$ for all $k = 0, \dots, \alpha$. Therefore, and in view of integrability of f on semiaxes $(-\infty, x)$ for any x ,

$$\begin{aligned}
& \int_{\mathbb{R}} \left(\int_{-\infty}^x \frac{f(y) dy}{(x-y)^{1-\alpha}} \right) \sum_{m=0}^{\alpha} (-1)^m \binom{\alpha}{m} B_{n^*}(2^d x - 2\tau + (m^* - r - m) - s - l) \\
&= \int_{\frac{2\tau+s+l+r+m+1}{2^d}}^{\frac{2\tau+s+l+r+m+1}{2^d}} \left(\int_{-\infty}^x \frac{f(y) dy}{(x-y)^{1-\alpha}} \right) B_{m^*}^{(\alpha)}(2^d x - 2\tau + (m^* - r) - s - l) \\
&= \int_{-\infty}^{\frac{2\tau+s+l+r+m-m^*}{2^d}} f(y) \left(\int_{\frac{2\tau+s+l+r+m-m^*}{2^d}}^{\frac{2\tau+s+l+r+m+1}{2^d}} \frac{B_{m^*}^{(\alpha)}(2^d x - 2\tau + (m^* - r) - s - l) dx}{(x-y)^{1-\alpha}} \right) dy \\
&+ \int_{\frac{2\tau+s+l+r+m-m^*}{2^d}}^{\frac{2\tau+s+l+r+m+1}{2^d}} f(y) \left(\int_{\frac{2\tau+s+l+r+m-m^*}{2^d}}^y \frac{B_{m^*}^{(\alpha)}(2^d x - 2\tau + (m^* - r) - s - l) dx}{(x-y)^{1-\alpha}} \right) dy \\
&= \frac{\Gamma(\alpha)}{2^{d\alpha}} \int_{\frac{2\tau+s+l+r-m^*}{2^d}}^{\frac{2\tau+s+l+r+m+1}{2^d}} f(y) B_{m^*}^{(\alpha)}(2^d y - 2\tau + (m^* - r) - s - l) dy.
\end{aligned}$$

From this we obtain that

$$\begin{aligned}
& \frac{2^{d\alpha} \Lambda_{n^*}}{2^{d/2} \Gamma(\alpha)} \langle I_+^\alpha f, \tilde{\Psi}_{(d-1)\tau}^{(1,1)} \rangle = \frac{2^{d\alpha} \gamma_{n^*}}{\Gamma(\alpha)} \langle I_+^\alpha f, \Psi_{n^*, 0, 0; m^*(1), \alpha(1)}(2^{d-1} \cdot - \tau) \rangle \\
&= \sum_{|l| \leq m^*} \lambda_{|l|}(m^*) (-1)^{|l|} \sum_{|s| \leq n^*} \lambda_{|s|}(n^*) (-1)^{|s|} \sum_{r=0}^{m^*+1} (-1)^r \binom{1+m^*}{r} \\
&\quad \times \int_{\mathbb{R}} f(y) B_{m^*}^{(\alpha)}(2^d y - 2\tau + (m^* - r) - s - l) dy = \frac{\gamma_{m^*} \Lambda_{m^*}}{2^{d/2}} \langle f, \bar{\Psi}_{(d-1)\tau}^{(1,0)} \rangle,
\end{aligned}$$

where $\bar{\Psi}^{(1,0)}$ is specified by $\Psi_{m^*,0,0;n^*(1),\alpha(0)}$, and we can write that

$$2^{pd\alpha} \sum_{\tau \in \mathbb{Z}} u(Q_{(d-1)\tau}) |\langle I_+^\alpha f, \tilde{\Psi}_{(d-1)\tau}^{(1,1)} \rangle|^p = \sum_{\tau \in \mathbb{Z}} u(Q_{(d-1)\tau}) \left| \int_{\mathbb{R}} f(y) \bar{\Psi}_{(d-1)\tau}^{(1,0)}(y) dy \right|^p.$$

Applying the condition (3.10), we obtain

$$\begin{aligned} 2^{-d/2} 2^{d(-\kappa^* + \alpha)} \left\| \sum_{\tau \in \mathbb{Z}} |\tilde{\lambda}_{d\tau}^{(1,1)}| \chi_{Q_{(d-1)\tau}} \right\|_{L^p(\mathbb{R}, u)} &= 2^{d(-\kappa^* + \alpha)} \left(\sum_{\tau \in \mathbb{Z}} u(Q_{(d-1)\tau}) |\langle I_+^\alpha f, \tilde{\Psi}_{(d-1)\tau}^{(1,1)} \rangle|^p \right)^{1/p} \\ &\simeq 2^{-d\kappa^*} \left(\sum_{\tau \in \mathbb{Z}} u(Q_{(d-1)\tau}) |\langle f, \bar{\Psi}_{(d-1)\tau}^{(1,0)} \rangle|^p \right)^{1/p} \\ &\lesssim \sup_{d \in \mathbb{N}} \mathfrak{M}(d, \kappa^*; v, u) \left(\sum_{\tau \in \mathbb{Z}} v(Q_{(d-1)\tau}) |\langle f, \bar{\Psi}_{(d-1)\tau}^{(1,0)} \rangle|^p \right)^{1/p} \\ &\lesssim \mathfrak{C}_+^\alpha(\kappa^*) \left(\sum_{\tau \in \mathbb{Z}} v(Q_{(d-1)\tau}) |\langle f, \bar{\Psi}_{(d-1)\tau}^{(1,0)} \rangle|^p \right)^{1/p} = \mathfrak{C}_+^\alpha(\kappa^*) \left\| \sum_{\tau \in \mathbb{Z}} \chi_{Q_{(d-1)\tau}} |\langle f, \bar{\Psi}_{(d-1)\tau}^{(1,0)} \rangle| \right\|_{L^p(\mathbb{R}, v)}. \end{aligned}$$

Thus, $\bar{\Psi}_{(d-1)\nu}^{(1,0)}$ together with $\bar{\Phi}_\nu$ (see p. 11) forms a spline wavelet system of order m^* . The sequential norms corresponding to this system, by Theorem 3.2, are equivalent to the norm of f in the space $B_{pq}^{s+\kappa^*-\alpha, v}(\mathbb{R})$. This means that assertion (i) is proven.

Necessity. Suppose that the operator I_+^α is bounded from $B_{pq}^{s+\kappa^*-\alpha, v}(\mathbb{R})$ with some $\kappa^* \in \mathbb{R}$ to $B_{pq}^{s, u}(\mathbb{R})$. Accordingly, the following inequality holds

$$\|I_+^\alpha f\|_{B_{pq}^{s, u}(\mathbb{R})} \lesssim \mathfrak{C} \|f\|_{B_{pq}^{s+\kappa^*-\alpha, v}(\mathbb{R})}, \quad (3.18)$$

where $\mathfrak{C} > 0$ is the best constant in (3.18), i.e. $\mathfrak{C} = \|I_\pm^\alpha\|_{B_{pq}^{s+\kappa^*-\alpha, v}(\mathbb{R}) \rightarrow B_{pq}^{s, u}(\mathbb{R})}$. We will show that

$$\mathfrak{C} \gtrsim \mathfrak{C}_+(\kappa^*) \geq \max\{\mathbf{M}_+^\alpha(1), \mathbf{M}_+^\alpha(0), \sup_{d \in \mathbb{N}} \mathfrak{M}(d, \kappa^*; v, u)\}. \quad (3.19)$$

Let $n^* \in \mathbb{N}$ and $m^* \in \mathbb{N}$ satisfy the condition (3.3) with respect to the parameters of the spaces $B_{pq}^{s, u}(\mathbb{R})$ and $B_{pq}^{s+\kappa^*-\alpha, v}(\mathbb{R})$, respectively. By theorem 3.1,

$$\|I_+^\alpha f\|_{B_{pq}^{s, u}(\mathbb{R})} \approx \|\tilde{\lambda}_{(I_+^\alpha f)}^{(\tilde{\mathbf{a}})}\|_{b_{pq}^s(u)}, \quad (3.20)$$

where the sequence $\tilde{\lambda}_{(I_+^\alpha f)}^{(\tilde{\mathbf{a}})}$ is defined by the elements

$$\tilde{\lambda}_{0\tau}^{(\tilde{\mathbf{a}})} = \langle I_+^\alpha f, \tilde{\Phi}_\tau^{(\tilde{\mathbf{a}})} \rangle \quad (\tau \in \mathbb{Z}), \quad \tilde{\lambda}_{d\tau}^{(\tilde{\mathbf{a}})} = 2^{d/2} \langle I_+^\alpha f, \tilde{\Psi}_{(d-1)\tau}^{(\tilde{\mathbf{a}})} \rangle \quad (d \in \mathbb{N}, \tau \in \mathbb{Z}). \quad (3.21)$$

Here, $\tilde{\Phi}^{(\tilde{\mathbf{a}})}$ are defined by $B_{n^*, \tilde{\mathbf{a}}}$ and $\tilde{\Psi}^{(\tilde{\mathbf{a}})}$ (see (5.13)) — via $\Psi_{n^*, \tilde{\mathbf{a}}, 0}$, where $\tilde{\mathbf{a}}$ can take values 0 or $\pm 1/2$. Based on the above, we can write:

$$\|I_+^\alpha f\|_{B_{pq}^{s, u}(\mathbb{R})} \approx \left\| \sum_{\tau \in \mathbb{Z}} |\tilde{\lambda}_{0\tau}^{(\tilde{\mathbf{a}})}| \chi_{Q_{0\tau}} \right\|_{L^p(\mathbb{R}, u)} + \left(\sum_{d \in \mathbb{N}} 2^{qds} \left\| \sum_{\tau \in \mathbb{Z}} |\tilde{\lambda}_{d\tau}^{(\tilde{\mathbf{a}})}| \chi_{Q_{(d-1)\tau}} \right\|_{L^p(\mathbb{R}, u)}^q \right)^{1/q}. \quad (3.22)$$

Moreover, also based on Theorem 3.1,

$$\|f\|_{B_{pq}^{s+\kappa^*-\alpha, v}(\mathbb{R})} \approx \|\lambda_f^{(\mathbf{a})}\|_{b_{pq}^{s+\kappa^*-\alpha}(v)},$$

where the sequence $\lambda_f^{(a)}$ with $a \in \{0, \pm 1/2\}$ is defined by elements

$$\lambda_{0\tau}^{(a)} = \langle f, \Phi_\tau^{(a)} \rangle \quad (\tau \in \mathbb{Z}), \quad \lambda_{d\tau}^{(a)} = 2^{d/2} \langle f, \Psi_{(d-1)\tau}^{(a)} \rangle \quad (d \in \mathbb{N}, \tau \in \mathbb{Z}). \quad (3.23)$$

In this case, $\Phi^{(a)}$ is defined by $B_{m^*, a}$, and $\Psi^{(a)}$ (see (5.14)) – via $\Psi_{m^*, a, 0}$. Thus, for the right-hand side of (3.18) we can write that

$$\|f\|_{B_{pq}^{s+\kappa^*-\alpha, v}(\mathbb{R})} \approx \left\| \sum_{\tau \in \mathbb{Z}} |\lambda_{0\tau}^{(a)}| \chi_{Q_{0\tau}} \right\|_{L^p(\mathbb{R}, v)} + \left(\sum_{d \in \mathbb{N}} 2^{qd(s+\kappa^*-\alpha)} \left\| \sum_{\tau \in \mathbb{Z}} |\lambda_{d\tau}^{(a)}| \chi_{Q_{(d-1)\tau}} \right\|_{L^p(\mathbb{R}, v)}^q \right)^{1/q}. \quad (3.24)$$

To prove (3.19) we first assume that the maximum on the right side of (3.19) is equal to

$$\mathbf{M}_+^\alpha(0) = \sup_{\tau \in \mathbb{Z}} \left(\sum_{r \geq \tau} u(Q_{0r}) \right)^{\frac{1}{p}} \left(\sum_{r \leq \tau} (\tau - r + 1)^{p'(\alpha-1)} [v(Q_{0r})]^{1-p'} \right)^{\frac{1}{p'}}.$$

Then fix n^* and $m^* = n^* + \alpha$, and leave only the first of the two terms on the right side of (3.22). Due to the procedure of forming iterative differences (to the left) of order α with coefficients

$$A_r(\alpha) = \sum_{j=1}^r (-1)^{j-1} \binom{\alpha}{j} A_{r-j}(\alpha), \quad \text{where } A_0(\alpha) = 1,$$

(see e.g. (3.14) and (5.1)) the following equalities are true:

$$\begin{aligned} \frac{m^*!}{n^*!} \left\| \sum_{\tau \in \mathbb{Z}} |\tilde{\lambda}_{0\tau}^{(\tilde{a})}| \chi_{Q_{0\tau}} \right\|_{L^p(\mathbb{R}, u)}^p &= \sum_{\tau \in \mathbb{Z}} u(Q_{0\tau}) |\langle I_+^\alpha f, \tilde{\Phi}_\tau^{(\tilde{a})} \rangle|^p \\ &= \sum_{\tau \in \mathbb{Z}} u(Q_{0\tau}) \left| \sum_{r \geq 0} A_r(\alpha) \int_{\mathbb{R}} f(y) B_{m^*}(y - \tau + \alpha - \tilde{a} + r) dy \right|^p \\ &= \sum_{\tau \in \mathbb{Z}} u(Q_{0\tau}) \left| \sum_{r \geq 0} A_r(\alpha) \langle f, B_{m^*, \tau - \alpha + \tilde{a} - r} \rangle \right|^p = \sum_{\tau \in \mathbb{Z}} u(Q_{0\tau}) \left| \sum_{l \leq \tau} A_{\tau-l}(\alpha) \langle f, B_{m^*, l - \alpha + \tilde{a}} \rangle \right|^p. \end{aligned}$$

Then we obtain from (3.18), (3.22) and (3.24):

$$\begin{aligned} \left(\sum_{\tau \in \mathbb{Z}} u(Q_{0\tau}) \left| \sum_{l \leq \tau} A_{\tau-l}(\alpha) \langle f, B_{m^*, l - \alpha + \tilde{a}} \rangle \right|^p \right)^{1/p} &\simeq \left\| \sum_{\tau \in \mathbb{Z}} |\tilde{\lambda}_{0\tau}^{(\tilde{a})}| \chi_{Q_{0\tau}} \right\|_{L^p(\mathbb{R}, u)} \\ &\lesssim \|I_+^\alpha f\|_{B_{pq}^{s, u}(\mathbb{R})} \lesssim \mathfrak{e} \|f\|_{B_{pq}^{s+\kappa^*-\alpha, v}(\mathbb{R})} \\ &\approx \mathfrak{e} \left[\left\| \sum_{\tau \in \mathbb{Z}} |\lambda_{0\tau}^{(a)}| \chi_{Q_{0\tau}} \right\|_{L^p(\mathbb{R}, v)} + \left(\sum_{d \in \mathbb{N}} 2^{qd(s+\kappa^*-\alpha)} \left\| \sum_{\tau \in \mathbb{Z}} |\lambda_{d\tau}^{(a)}| \chi_{Q_{(d-1)\tau}} \right\|_{L^p(\mathbb{R}, v)}^q \right)^{1/q} \right] \\ &= \mathfrak{e} \left[\left(\sum_{\tau \in \mathbb{Z}} v(Q_{0\tau}) |\langle f, B_{m^*, \tau + a} \rangle|^p \right)^{1/p} + \left(\sum_{d \in \mathbb{N}} 2^{qd(s+\kappa^*-\alpha)} \left(\sum_{\tau \in \mathbb{Z}} v(Q_{(d-1)\tau}) |\langle f, \Psi_{(d-1)\tau}^{(a)} \rangle|^p \right)^{q/p} \right)^{1/q} \right]. \end{aligned}$$

On the strength of Lemma 2.1 and property (2.2),

$$\sum_{\tau \in \mathbb{Z}} v(Q_{0\tau}) |\langle f, B_{m^*, \tau + a} \rangle|^p \lesssim \sum_{l \in \mathbb{Z}} v(Q_{0l}) |\langle f, B_{m^*, l - \alpha + a} \rangle|^p.$$

Thus,

$$\left(\sum_{\tau \in \mathbb{Z}} u(Q_{0\tau}) \left| \sum_{l \leq \tau} A_{\tau-l}(\boldsymbol{\alpha}) \langle f, B_{m^*, l-\boldsymbol{\alpha}+\tilde{\mathbf{a}}} \rangle \right|^p \right)^{1/p} \lesssim \mathfrak{C} \left[\left(\sum_{l \in \mathbb{Z}} v(Q_{0l}) |\langle f, B_{m^*, l-\boldsymbol{\alpha}+\mathbf{a}} \rangle|^p \right)^{1/p} + \left(\sum_{d \in \mathbb{N}} 2^{qd(s+\kappa^*-\boldsymbol{\alpha})} \left(\sum_{\tau \in \mathbb{Z}} v(Q_{(d-1)\tau}) |\langle f, \Psi_{(d-1)\tau}^{(\mathbf{a})} \rangle|^p \right)^{q/p} \right)^{1/q} \right]. \quad (3.25)$$

For $A_r(m)$ with fixed $m \in \mathbb{N}$, the following lower bound holds for all $r \in \mathbb{N}$:

$$(m-1)!A_r(m) \geq (1+r)^{m-1} = \sum_{s=0}^{m-1} \frac{(m-1)!r^s}{s!(m-1-s)!}. \quad (3.26)$$

We will show that this is so, taking into account (see [33, (5.20)]) that $A_1(m) = m$ and

$$A_r(m) = \sum_{1 \leq l \leq m} A_{r-1}(l), \quad (m \in \mathbb{N}). \quad (3.27)$$

For $r = 1$, (3.26) holds by (3.27) for all $m = m_0 \in \mathbb{N}$. Suppose that (3.26) holds for $r = k - 1$ with some $k > 1$ and any $m = l$, where $1 \leq l \leq m_0$. Then it follows from (3.27) that

$$A_k(m_0) = \sum_{1 \leq l \leq m_0} A_{k-1}(l) \geq \sum_{1 \leq l \leq m_0} \frac{k^{l-1}}{(l-1)!} = \sum_{s=0}^{m_0-1} \frac{k^s}{s!} \geq \sum_{s=0}^{m_0-1} \frac{k^s}{s!(m_0-1-s)!} = \frac{(1+k)^{m_0-1}}{(m_0-1)!}.$$

Next, we set $\mathbf{a} = \tilde{\mathbf{a}} = 0$ and denote by \mathfrak{B}_{m^*} the class of functions from $B_{pq}^{s+\kappa^*-\boldsymbol{\alpha}, v}(\mathbb{R})$ which can be represented as a finite linear combination of integer shifts of B -splines of order m^* . Due to the semi-orthogonality of the basis elements on the right-hand side of (3.24) (see (5.16)), the restriction of (3.25) to the sequences $\boldsymbol{\lambda}_f^{(0)}$ with f from the class \mathfrak{B}_{m^*} has the form

$$\left(\sum_{\tau \in \mathbb{Z}} u(Q_{0\tau}) \left| \sum_{l \leq \tau} A_{\tau-l}(\boldsymbol{\alpha}) \langle f, B_{m^*, l-\boldsymbol{\alpha}} \rangle \right|^p \right)^{1/p} \lesssim \mathfrak{C}_{\mathfrak{B}_{m^*}} \left(\sum_{l \in \mathbb{Z}} v(Q_{0l}) |\langle f, B_{m^*, l-\boldsymbol{\alpha}} \rangle|^p \right)^{1/p}. \quad (3.28)$$

Besides, the best constant $\mathfrak{C}_{\mathfrak{B}_{m^*}}$ of this inequality does not exceed \mathfrak{C} .

For a sufficiently large $R \in \mathbb{N}$, we substitute into (3.28) a function f_R^* of the form

$$\begin{aligned} f_R^*(y) &= \sum_{-R \leq \tau \leq \nu} (\nu - \tau + 1)^{(\boldsymbol{\alpha}-1)(p'-1)} [v(Q_{0\tau})]^{1-p'} B_{m^*, \tau-\boldsymbol{\alpha}}(y) \\ &= \sum_{\tau \leq \nu} (\nu - \tau + 1)^{(\boldsymbol{\alpha}-1)(p'-1)} [v(Q_{0\tau})]^{1-p'} B_{m^*}(y - \tau + \boldsymbol{\alpha}) \quad (\nu \in \mathbb{Z}). \end{aligned} \quad (3.29)$$

This will lead us to inequality

$$\left(\sum_{\tau \in \mathbb{Z}} u(Q_{0\tau}) \left| \sum_{l \leq \tau} A_{\tau-l}(\boldsymbol{\alpha}) \langle f_R^*, B_{m^*, l-\boldsymbol{\alpha}} \rangle \right|^p \right)^{1/p} \lesssim \mathfrak{C}_{\mathfrak{B}_{m^*}} \left(\sum_{l \in \mathbb{Z}} v(Q_{0l}) |\langle f_R^*, B_{m^*, l-\boldsymbol{\alpha}} \rangle|^p \right)^{1/p}. \quad (3.30)$$

Due to the non-negativity of f_R^* and B_{m^*} and on the basis of (3.26)

$$\sum_{l \leq \tau} A_{\tau-l}(\boldsymbol{\alpha}) \langle f_R^*, B_{m^*, l-\boldsymbol{\alpha}} \rangle \gtrsim \sum_{l \leq \tau} (\tau - l + 1)^{\boldsymbol{\alpha}-1} \langle f_R^*, B_{m^*, l-\boldsymbol{\alpha}} \rangle.$$

Besides,

$$\begin{aligned} & \left(\sum_{\tau \in \mathbb{Z}} u(Q_{0\tau}) \left| \sum_{l \leq \tau} (\tau - l + 1)^{\alpha-1} \langle f_R^*, B_{m^*, l-\alpha} \rangle \right|^p \right)^{1/p} \\ & \geq \left(\sum_{\tau \geq \nu} u(Q_{0\tau}) \right)^{1/p} \sum_{l \leq \nu} (\nu - l + 1)^{\alpha-1} \langle f_R^*, B_{m^*, l-\alpha} \rangle. \end{aligned}$$

Thus, (3.30) is transformable into

$$\left(\sum_{\tau \geq \nu} u(Q_{0\tau}) \right)^{1/p} \sum_{l \leq \nu} (\nu - l + 1)^{\alpha-1} \langle f_R^*, B_{m^*, l-\alpha} \rangle \lesssim \mathfrak{C}_{\mathfrak{B}_{m^*}} \left(\sum_{l \in \mathbb{Z}} v(Q_{0l}) |\langle f_R^*, B_{m^*, l-\alpha} \rangle|^p \right)^{1/p}. \quad (3.31)$$

Further, we write for the sum depending on f_R^* on the left-hand side (3.31):

$$\begin{aligned} & \sum_{l \leq \nu} (\nu - l + 1)^{\alpha-1} \langle f_R^*, B_{m^*, l-\alpha} \rangle \\ & = \sum_{l \leq \nu} (\nu - l + 1)^{\alpha-1} \sum_{-R \leq \tau \leq \nu} (\nu - \tau + 1)^{(\alpha-1)(p'-1)} [v(Q_{0\tau})]^{1-p'} \langle B_{m^*, \tau-\alpha}, B_{m^*, l-\alpha} \rangle \\ & = \sum_{-R \leq \tau \leq \nu} (\nu - \tau + 1)^{(\alpha-1)(p'-1)} [v(Q_{0\tau})]^{1-p'} \sum_{l \leq \nu} (\nu - l + 1)^{\alpha-1} \langle B_{m^*, \tau-\alpha}, B_{m^*, l-\alpha} \rangle \\ & \geq \sum_{-R \leq \tau \leq \nu} (\nu - \tau + 1)^{(\alpha-1)(p'-1)} [v(Q_{0\tau})]^{1-p'} \sum_{l \leq \tau} (\nu - l + 1)^{\alpha-1} \langle B_{m^*, \tau-\alpha}, B_{m^*, l-\alpha} \rangle \\ & \geq \sum_{-R \leq \tau \leq \nu} (\nu - \tau + 1)^{(\alpha-1)p'} [v(Q_{0\tau})]^{1-p'} \sum_{l \leq \tau} \langle B_{m^*, \tau-\alpha}, B_{m^*, l-\alpha} \rangle. \end{aligned}$$

Due to (5.5) and orthonormality of $\{\phi_{m^*, \tau}\}_{\tau \in \mathbb{Z}}$,

$$\begin{aligned} & \beta_{m^*}^2 \sum_{l \leq \tau} \langle B_{m^*, \tau-\alpha}, B_{m^*, l-\alpha} \rangle \\ & = \sum_{\varkappa'=0}^{m^*} \alpha'_{\varkappa'} \sum_{\varkappa''=0}^{m^*} \alpha'_{\varkappa''} \sum_{l \leq \tau} \int \phi_{m^*}(y + \varkappa' - \tau + \alpha) \phi_{m^*}(y + \varkappa'' - l + \alpha) \\ & = \sum_{\varkappa=0}^{m^*} (\alpha'_{\varkappa})^2 + \sum_{\varkappa''-\varkappa'=1} \alpha'_{\varkappa'} \alpha'_{\varkappa''} + \dots + \sum_{\varkappa''-\varkappa'=m^*} \alpha'_{\varkappa'} \alpha'_{\varkappa''} \\ & \gtrsim \sum_{\varkappa=0}^{m^*} (\alpha'_{\varkappa})^2 + \sum_{|\varkappa'-\varkappa''|=1} \alpha'_{\varkappa'} \alpha'_{\varkappa''} + \dots + \sum_{|\varkappa'-\varkappa''|=m^*} \alpha'_{\varkappa'} \alpha'_{\varkappa''} = \left(\sum_{\varkappa=0}^{m^*} \alpha'_{\varkappa} \right)^2 = \beta_{m^*}^2. \quad (3.32) \end{aligned}$$

Thus, the left-hand side of (3.31) is estimated with some constant from below by

$$\left(\sum_{\tau \geq \nu} u(Q_{0\tau}) \right)^{1/p} \sum_{-R \leq \tau \leq \nu} (\nu - \tau + 1)^{(\alpha-1)p'} [v(Q_{0\tau})]^{1-p'}. \quad (3.33)$$

To estimate the right-hand side of (3.31) from above, we write, using the Hölder inequality,

$$\begin{aligned} & |\langle f_R^*, B_{m^*, l-\alpha} \rangle|^p = \left[\sum_{-R \leq \tau \leq \nu} (\nu - \tau + 1)^{(\alpha-1)(p'-1)} [v(Q_{0\tau})]^{1-p'} \langle B_{m^*}(\cdot - \tau + \alpha), B_{m^*}(\cdot - l + \alpha) \rangle \right]^p \\ & \leq \sum_{-R \leq \tau \leq \nu} (\nu - \tau + 1)^{(\alpha-1)p'} [v(Q_{0\tau})]^{-p'} \langle B_{m^*}(\cdot - \tau + \alpha), B_{m^*}(\cdot - l + \alpha) \rangle \end{aligned}$$

$$\times \left[\sum_{-R \leq r \leq \nu} \langle B_{m^*}(\cdot - r + \alpha), B_{m^*}(\cdot - l + \alpha) \rangle \right]^{p-1}.$$

Hence, due to bounded supports of B_{m^*} ,

$$\begin{aligned} & \sum_{l \in \mathbb{Z}} v(Q_{0l}) |\langle f_R^*, B_{m^*, l - \alpha} \rangle|^p \\ & \leq \sum_{l \in \mathbb{Z}} v(Q_{0l}) \sum_{-R \leq \tau \leq \nu} (\nu - \tau + 1)^{(\alpha-1)p'} [v(Q_{0\tau})]^{-p'} \langle B_{m^*}(\cdot - \tau + \alpha), B_{m^*}(\cdot - l + \alpha) \rangle \\ & \quad \times \left[\sum_{-R \leq r \leq \nu} \langle B_{m^*}(\cdot - r + \alpha), B_{m^*}(\cdot - l + \alpha) \rangle \right]^{p-1} \\ & = \sum_{-R \leq \tau \leq \nu} (\nu - \tau + 1)^{(\alpha-1)p'} [v(Q_{0\tau})]^{-p'} \sum_{l \in \mathbb{Z}} v(Q_{0l}) \langle B_{m^*}(\cdot - \tau + \alpha), B_{m^*}(\cdot - l + \alpha) \rangle \\ & \quad \times \left[\sum_{-R \leq r \leq \nu} \langle B_{m^*}(\cdot - r + \alpha), B_{m^*}(\cdot - l + \alpha) \rangle \right]^{p-1} \\ & \leq \sum_{-R \leq \tau \leq \nu} (\nu - \tau + 1)^{(\alpha-1)p'} [v(Q_{0\tau})]^{-p'} \sum_{|l - \tau| \leq m^*} v(Q_{0l}) \langle B_{m^*}(\cdot - \tau + \alpha), B_{m^*}(\cdot - l + \alpha) \rangle \\ & \quad \times \left[\sum_{r \leq \nu} \langle B_{m^*}(\cdot - r + \alpha), B_{m^*}(\cdot - l + \alpha) \rangle \right]^{p-1}. \end{aligned}$$

For a fixed l , due to (5.5) and orthonormality of $\{\phi_{m^*, \tau}\}_{\tau \in \mathbb{Z}}$ (see (3.32)),

$$\sum_{r \leq \nu} \langle B_{m^*}(\cdot - r + \alpha), B_{m^*}(\cdot - l + \alpha) \rangle \leq \sum_{|r - l| \leq m^*} \langle B_{m^*}(\cdot - r + \alpha), B_{m^*}(\cdot - l + \alpha) \rangle \leq 1.$$

By virtue of Lemma 2.1 and property (2.2) we have $v(Q_{0l}) \lesssim v(Q_{0\tau})$ for any $|l - \tau| \leq m^*$. Therefore, for the right-hand side (3.31) it holds that

$$\begin{aligned} & \sum_{l \in \mathbb{Z}} v(Q_{0l}) |\langle f_R^*, B_{m^*, l - \alpha} \rangle|^p \\ & \lesssim \sum_{-R \leq \tau \leq \nu} (\nu - \tau + 1)^{(\alpha-1)p'} [v(Q_{0\tau})]^{1-p'} \sum_{|l - \tau| \leq m^*} \langle B_{m^*}(\cdot - \tau + \alpha), B_{m^*}(\cdot - l + \alpha) \rangle \\ & \leq \sum_{-R \leq \tau \leq \nu} (\nu - \tau + 1)^{(\alpha-1)p'} [v(Q_{0\tau})]^{1-p'}. \end{aligned}$$

Combining this with (3.31) and (3.33), we obtain from (3.30) that

$$\mathfrak{C}_{\mathfrak{B}_{m^*}} \gtrsim \sup_{\nu \in \mathbb{Z}} \left(\sum_{\tau \geq \nu} u(Q_{0\tau}) \right)^{\frac{1}{p}} \left(\sum_{-R \leq \tau \leq \nu} (\nu - \tau + 1)^{p'(\alpha-1)} [v(Q_{0\tau})]^{1-p'} \right)^{\frac{1}{p'}},$$

from which, tending $R \rightarrow \infty$, we extract the estimate $\mathfrak{C} \geq \mathfrak{C}_{\mathfrak{B}_{m^*}} \gtrsim \mathbf{M}_+^\alpha(0)$ for the best constant of the inequality (3.18). Similarly, it can be shown that $\mathfrak{C} \gtrsim \mathbf{M}_+^\alpha(1)$, where

$$\mathbf{M}_+^\alpha(1) = \sup_{\tau \in \mathbb{Z}} \left(\sum_{r \geq \tau} (r - \tau + 1)^{p(\alpha-1)} u(Q_{0r}) \right)^{\frac{1}{p}} \left(\sum_{r \leq \tau} [v(Q_{0r})]^{1-p'} \right)^{\frac{1}{p'}}.$$

To do this, we need to define a test function dual to f_R^*

$$g_R^*(y) = \sum_{\nu \leq \tau \leq R} (\tau - \nu + 1)^{(\alpha-1)(p-1)} [u(Q_{0\tau})] B_{m^*, \tau - \alpha}(y)$$

and substitute it into the dual to (3.28) inequality of the form

$$\left(\sum_{\tau \in \mathbb{Z}} [v(Q_{0\tau})]^{1-p'} \left| \sum_{l \geq \tau} A_{l-\tau}(\alpha) \langle g, B_{m^*, l-\alpha} \rangle \right|^{p'} \right)^{1/p'} \lesssim \mathfrak{E}_{\mathfrak{B}_{m^*}} \left(\sum_{l \in \mathbb{Z}} [u(Q_{0l})]^{1-p'} \left| \langle g, B_{m^*, l-\alpha} \rangle \right|^{p'} \right)^{1/p'}.$$

Now suppose the maximum on the right-hand side (3.19) is achieved at $\sup_{d \in \mathbb{N}} \mathfrak{M}(d, \kappa^*; v, u)$. Let us show the necessity of such a characteristic. To do this, for some d_0 and τ_0 we define

$$\begin{aligned} h_{\tau_0}^*(y) &= \Lambda_{m^*}^{-1} \Psi_{m^*, \bar{a}, \bar{s}; n^*(1), \alpha(1)}(2^{d_0-1}y - \tau_0) \\ &= \sum_{k=0}^{2\alpha} (-1)^k \binom{2\alpha}{k} h_{\tau_0; k}(y) = \sum_{|\varkappa| \leq n^*} \lambda_{|\varkappa|}(n^*) (-1)^{|\varkappa|} \sum_{k=0}^{2\alpha} (-1)^k \binom{2\alpha}{k} h_{\tau_0; k, \varkappa}(y) \end{aligned} \quad (3.34)$$

with $\bar{s} \in \mathbb{R}$, which we will specified later, and

$$\begin{aligned} h_{\tau_0; k}(y) &= \Psi_{m^*, \bar{a}, \bar{s}; n^*(1), \alpha(0)}(2^{d_0-1}y - \tau_0 - k/2 + \alpha/2), \\ h_{\tau_0; k, \varkappa}(y) &= \Psi_{m^*, \bar{a}, \bar{s}}(2^{d_0-1}y - \tau_0 - k/2 - \varkappa/2 + \alpha/2). \end{aligned}$$

We will need the representation (3.22) for $f = h_{\tau_0}^*$ and (3.24) for $f = h_{\tau_0; k, \varkappa}$ with $k = 0, \dots, 2\alpha$ and $\varkappa = -n^*, \dots, 0, \dots, n^*$. In this situation, we fix n^* and m^* so that $n^* = m^* + \alpha$.

Then (see (5.1) and (3.17))

$$\begin{aligned} \gamma_{m^*} I_+^\alpha h_{\tau_0}^*(x) &= 2^{-\alpha d_0} \gamma_{n^*} \Psi_{n^*, \bar{a}, \bar{s}; m^*(1), \alpha(0)}(2^{d_0-1} \cdot -\tau_0) = 2^{-\alpha d_0} \sum_{|r| \leq m^*} \lambda_{|r|}(m^*) (-1)^{|r|} \\ &\times \sum_{|\kappa| \leq n^*} \lambda_{|\kappa|}(n^*) (-1)^{|\kappa|} \sum_{j=0}^{n^*+1} (-1)^j \binom{1+n^*}{j} B_{n^*}(2^{d_0}(x - \bar{s} - \bar{a}) - 2\tau_0 + n^* - j - r - \kappa). \end{aligned} \quad (3.35)$$

We have $\text{supp}(I_+^\alpha h_{\tau_0}^*) = \left[\frac{2(\tau_0 + \bar{s} + \bar{a}) - 2n^* - m^*}{2^{d_0}}, \frac{2(\tau_0 + \bar{s} + \bar{a}) + 2n^* + m^* + 2}{2^{d_0}} \right]$ (see §5). On the other hand, the supports of the functions $\Psi_{n^*, \bar{a}, 0}(2^{d_0-1} \cdot -\tau)$ forming the coefficients $\tilde{\lambda}_{d\tau}^{(\bar{a})}$ in (3.22), coincide with

$$\left[\frac{2\tau + 2\tilde{a} - 2n^*}{2^d}, \frac{2\tau + 2\tilde{a} + 2n^* + 2}{2^d} \right]$$

(see §5), besides

$$\begin{aligned} \gamma_{n^*} \Psi_{n^*, \bar{a}, 0}(2^{d_0-1} \cdot -\tau) &= \sum_{|l| \leq n^*} \lambda_{|l|}(n^*) (-1)^{|l|} \\ &\times \sum_{s=0}^{n^*+1} (-1)^s \binom{1+n^*}{s} B_{n^*}(2^{d_0}(y - \tilde{a}) - 2\tau + n^* - s - l). \end{aligned} \quad (3.36)$$

Denote $\int_{\mathbb{R}} B_n(t) B_n(t-n) dt =: \mathbf{B}_n$ and observe that $\mathbf{B}_n > 0$ for any $n \in \mathbb{N}$. We will call \mathbf{B}_n as \mathbf{B}_n -intersection, which is minimal in some sense. It is true: (see (3.22))

$$\begin{aligned} \|I_+^\alpha h_{\tau_0}^*\|_{B_{p\bar{q}}^{s,u}(\mathbb{R})}^p &\gtrsim 2^{d_0 s p} |\tilde{\lambda}_{d_0\tau_0}^{(\bar{a})}|^p [u(Q_{(d_0-1)\tau_0})] = 2^{d_0 s p} 2^{d_0 p/2} |\langle I_+^\alpha h_{\tau_0}^*, \tilde{\Psi}_{(d_0-1)\tau_0}^{(\bar{a})} \rangle|^p [u(Q_{(d_0-1)\tau_0})] \\ &= 2^{d_0(s-\alpha)p} 2^{d_0 p} |\langle \Psi_{n^*, \bar{a}, \bar{s}; m^*(1), \alpha(0)}(2^{d_0-1} \cdot -\tau_0), \Psi_{n^*, \bar{a}, 0}(2^{d_0-1} \cdot -\tau_0) \rangle|^p [u(Q_{(d_0-1)\tau_0})] \\ &= 2^{d_0(s-\alpha)p} |\langle \Psi_{n^*, \bar{a}, \bar{s}; m^*(1), \alpha(0)}(\cdot/2 - \tau_0), \Psi_{n^*, \bar{a}, 0}(\cdot/2 - \tau_0) \rangle|^p [u(Q_{(d_0-1)\tau_0})]. \end{aligned} \quad (3.37)$$

If we choose $\bar{s} = -1 - m^* - 4n^*$ and set $\bar{a} = \tilde{a} = 0$, the supports of $\Psi_{n^*, \bar{a}, \bar{s}; m^*(1), \alpha(0)}$ and $\Psi_{n^*, \bar{a}, 0}$ in (3.37) intersect only at intersection of the supports of the leftmost B_{n^*} in the representation (3.36),

which is $B_{n^*}(\cdot - 2\tau_0 + 2n^*)$, and the rightmost B_{n^*} in the representation (3.35), which corresponds to $B_{n^*}(\cdot - 2\tau_0 + 3n^*)$. So, it will be exactly intersection of the type \mathbf{B}_{n^*} . The supports of other B_{n^*} in the specified representations will not intersect each other if $\bar{s} = -1 - m^* - 4n^*$. Therefore, taking into account all the coefficients, we conclude that

$$\langle \Psi_{n^*, \bar{a}, \bar{s}; m^*(1), \alpha(0)}(\cdot/2 - \tau_0), \Psi_{n^*, \bar{a}, 0}(\cdot/2 - \tau_0) \rangle = \frac{16\mathbf{B}_{n^*}}{\gamma_{m^*}\gamma_{n^*}} =: \Theta(n^*). \quad (3.38)$$

From here and (3.37) we obtain that

$$\|I_+^\alpha h_{\tau_0}^*\|_{B_{pq}^{s,u}(\mathbb{R})} \gtrsim 2^{d_0(s-\alpha)} [u(Q_{(d_0-1)\tau_0})]^{1/p} \Theta(n^*).$$

On the other hand,

$$\|h_{\tau_0}^*\|_{B_{pq}^{s+\kappa^*-\alpha, v}(\mathbb{R})} \lesssim \sum_{|\varkappa| \leq n^*} \lambda_{|\varkappa|}(n^*) \sum_{k=1}^{2\alpha} \binom{2\alpha}{k} \|h_{\tau_0; k, \varkappa}\|_{B_{pq}^{s+\kappa^*-\alpha, v}(\mathbb{R})},$$

where $h_{\tau_0; k, \varkappa}$ are finite linear combinations of mutually orthogonal integer shifts of $\psi_{m^*, \bar{a}, \bar{s}}(2^{d_0-1} \cdot -\tau_0 - k/2 - \varkappa/2 + \alpha/2)$ with \bar{a} and \bar{s} as above. For fixed k and \varkappa , this corresponds to $\psi_{m^*, \mathbf{a}, \mathbf{s}}(2^{d_0-1} \cdot -\tau_0)$ with $\mathbf{a} = (k + \varkappa - \alpha)/2 - [(k + \varkappa - \alpha)/2]$ and $\mathbf{s} = \bar{s} + [(k + \varkappa - \alpha)/2]$ (see (5.7)).

From the representation (3.24) we can write for each $k + \varkappa$:

$$\|h_{\tau_0; k, \varkappa}\|_{B_{pq}^{s+\kappa^*-\alpha, v}(\mathbb{R})} \approx \left\| \sum_{\tau \in \mathbb{Z}} |\lambda_{0\tau}^{(\mathbf{a})}| \chi_{Q_{0\tau}} \right\|_{L^p(\mathbb{R}, v)} + \left(\sum_{d \in \mathbb{N}} 2^{qd(s+\kappa^*-\alpha)} \left\| \sum_{\tau \in \mathbb{Z}} |\lambda_{d\tau}^{(\mathbf{a})}| \chi_{Q_{(d-1)\tau}} \right\|_{L^p(\mathbb{R}, v)}^q \right)^{1/q}, \quad (3.39)$$

where the elements $\lambda_{\tau_0; k, \varkappa}^{(\mathbf{a})}$ are given in (3.23) by $B_{m^*, \mathbf{a}}$ and $\Psi_{m^*, \mathbf{a}, 0}$ with \mathbf{a} as above. Due to the semi-orthogonality of the basis generated by $B_{m^*, \mathbf{a}}$ and $\Psi_{m^*, \mathbf{a}, 0}$,

$$\begin{aligned} \|h_{\tau_0; k, \varkappa}\|_{B_{pq}^{s+\kappa^*-\alpha, v}(\mathbb{R})}^p &\approx 2^{d_0(s-\alpha+\kappa^*+1)p} \left\| \sum_{\tau \in \mathbb{Z}} |\langle h_{\tau_0; k, \varkappa}(\cdot), \Psi_{m^*, \mathbf{a}, 0}(2^{d_0-1} \cdot -\tau) \rangle| \chi_{Q_{(d_0-1)\tau}} \right\|_{L^p(\mathbb{R}, v)}^p \\ &= \Lambda_{m^*}^{-2} 2^{d_0(s-\alpha+\kappa^*+1)p} \sum_{\tau \in \mathbb{Z}} |\langle \Psi_{m^*, \mathbf{a}, \mathbf{s}}(2^{d_0-1} \cdot -\tau_0), \Psi_{m^*, \mathbf{a}, 0}(2^{d_0-1} \cdot -\tau) \rangle|^p [v(Q_{(d_0-1)\tau})]. \end{aligned} \quad (3.40)$$

Since the support of $\Psi_{m^*, \mathbf{a}, 0}$ is compact, then the sum over τ is limited to $|\tau_0 + \mathbf{s} - \tau| \leq 2m^*$. In this case, $v(Q_{(d_0-1)\tau}) \lesssim v(Q_{(d_0-1)\tau_0})$ with a constant depending on n^* and α . In view of $\gamma_{m^*} \Psi_{m^*, \mathbf{a}, 0}(2^{d_0-1} \cdot -\tau) = \sum_{|\rho| \leq m^*} \alpha''_\rho \cdot \psi_{m^*, \mathbf{a}, 0}(2^{d_0-1} \cdot -\rho - \tau)$ we have (see [28, § 4.1])

$$\sum_{|\tau_0 + \mathbf{s} - \tau| \leq 2m^*} |\langle \Psi_{m^*, \mathbf{a}, \mathbf{s}}(2^{d_0-1} \cdot -\tau_0), \Psi_{m^*, \mathbf{a}, 0}(2^{d_0-1} \cdot -\tau) \rangle| \lesssim \Lambda_{m^*}^2.$$

From this we obtain that $\mathfrak{C} \gtrsim 2^{-d_0\kappa^*} [v(Q_{(d_0-1)\tau_0})]^{-1/p} [u(Q_{(d_0-1)\tau_0})]^{1/p}$ for arbitrary d_0 and τ_0 . Hence,

$$\mathfrak{C} \gtrsim \sup_{d \in \mathbb{N}} \mathfrak{M}(d, \kappa^*; v, u). \quad (3.41)$$

The proof of statement (i) in the case of I_+^α is similar. In this case, it is necessary to apply the procedure for forming iterative differences to the right side.

(ii) *Sufficiency* of the conditions in this part follows from Theorem 1.1 (iii) and the criteria for embeddings of Besov weighted spaces (see [9, Proposition 2.1] and [12, Proposition 5.1]).

Necessity can be proven similarly to the estimate $\mathfrak{C} \gtrsim \sup_{d \in \mathbb{N}} \mathfrak{M}(d, \kappa^*; v, u)$. Namely, it is first assumed that the inequality (3.11) is valid. Next, one should fix $n_* \in \mathbb{N}$ and $m_* \in \mathbb{N}$ satisfying $n_* = m_* + \alpha$ and the condition (3.3) with respect to parameters of the spaces $B_{pq}^{s,u}(\mathbb{R})$ and $B_{pq}^{s-\kappa_*-\alpha, w}(\mathbb{R})$,

respectively. Then, by Theorem 3.1, the representation (3.20) is true with $\tilde{\lambda}_{(I_+^\alpha f)}^{(\tilde{\mathbf{a}})}$ of the form (3.21), where $\tilde{\Phi}^{(\tilde{\mathbf{a}})}$ are defined by $B_{n^*, \tilde{\mathbf{a}}}$, and $\tilde{\Psi}^{(\tilde{\mathbf{a}})}$ (see (5.13)) – via $\Psi_{n^*, \tilde{\mathbf{a}}, 0}$, with some $\tilde{\mathbf{a}} \in \{0, \pm 1/2\}$. Thus, the representation (3.22) takes place for the right-hand side of (3.11).

Similarly, also based on Theorem 3.1,

$$\|f\|_{B_{pq}^{s-\kappa^*-\alpha, w}(\mathbb{R})} \approx \|\lambda_f^{(\mathbf{a})}\|_{b_{pq}^{s-\kappa^*-\alpha}(w)},$$

where the sequence $\lambda_f^{(\mathbf{a})}$ with $\mathbf{a} \in \{0, \pm 1/2\}$ is defined in (3.23) with $\Phi^{(\mathbf{a})}$ and $\Psi^{(\mathbf{a})}$ (see (5.14)), specified by $B_{m^*, \mathbf{a}}$ and $\Psi_{m^*, \mathbf{a}, 0}$. That is, for the left side of (3.11) the following is true:

$$\|f\|_{B_{pq}^{s-\kappa^*-\alpha, w}(\mathbb{R})} \approx \left\| \sum_{\tau \in \mathbb{Z}} |\lambda_{0\tau}^{(\mathbf{a})}| \chi_{Q_{0\tau}} \right\|_{L^p(\mathbb{R}, w)} + \left(\sum_{d \in \mathbb{N}} 2^{qd(s-\kappa^*-\alpha)} \left\| \sum_{\tau \in \mathbb{Z}} |\lambda_{d\tau}^{(\mathbf{a})}| \chi_{Q_{(d-1)\tau}} \right\|_{L^p(\mathbb{R}, w)}^q \right)^{1/q}. \quad (3.42)$$

Next, for arbitrary d_0 and τ_0 , the function $h_{*, \tau_0} = h_{\tau_0}^*$ is introduced into consideration (see (3.34)). In this case (see (3.35)),

$$\begin{aligned} \gamma_{m^*} 2^{\alpha d_0} I_+^\alpha h_{\tau_0}^*(x) &= \Psi_{n^*, \bar{\mathbf{a}}, \bar{\mathbf{s}}; m^*(1), \alpha(0)}(2^{d_0-1}y - \tau_0) \\ &= \sum_{|l| \leq m^*} \lambda_{|l|}(m^*) (-1)^{|l|} \Psi_{n^*, \bar{\mathbf{a}}, \bar{\mathbf{s}}}(2^{d_0-1}y - \tau_0 - l/2) =: \sum_{|l| \leq m^*} \lambda_{|l|}(m^*) (-1)^{|l|} h_{*, \tau_0, l}(y). \end{aligned}$$

We will use the representation (3.42) for $f = h_{*, \tau_0}$ and (3.22) for $h_{*, \tau_0, l}$ with $|l| \leq m^*$ instead of $I_+^\alpha f$. First, we estimate the left-hand side of (3.11). To do this, we write (see (3.42))

$$\begin{aligned} \|h_{*, \tau_0}\|_{B_{pq}^{s-\kappa^*-\alpha, w}(\mathbb{R})}^p &\gtrsim 2^{d_0(s-\kappa^*-\alpha)p} |\lambda_{d_0\tau_0}^{(\mathbf{a})}|^p [w(Q_{(d_0-1)\tau_0})] \\ &= 2^{d_0(s-\kappa^*-\alpha)p} 2^{d_0p/2} |\langle h_{*, \tau_0}, \Psi_{(d_0-1)\tau_0}^{(\mathbf{a})} \rangle|^p [w(Q_{(d_0-1)\tau_0})] \\ &= 2^{d_0(s-\kappa^*-\alpha)p} 2^{d_0p} |\langle \Psi_{m^*, \bar{\mathbf{a}}, \bar{\mathbf{s}}; n^*(1), \alpha(1)}(2^{d_0-1} \cdot -\tau_0), \Psi_{m^*, \mathbf{a}, 0}(2^{d_0-1} \cdot -\tau_0) \rangle|^p [w(Q_{(d_0-1)\tau_0})]. \end{aligned}$$

Just as in (3.37) here it is sufficient to fix $\bar{\mathbf{s}}$ in such a way to reduce the intersection of the supports $\Psi_{m^*, \bar{\mathbf{a}}, \bar{\mathbf{s}}; n^*(1), \alpha(1)}(2^{d_0-1} \cdot -\tau_0)$ and $\Psi_{m^*, \mathbf{a}, 0}$ with $\bar{\mathbf{a}} = \mathbf{a} = 0$ to \mathbf{B}_{m^*} -intersection of the supports of the rightmost B_{m^*} in the corresponding representation for $\Psi_{m^*, \bar{\mathbf{a}}, \bar{\mathbf{s}}; n^*(1), \alpha(1)}(2^{d_0-1} \cdot -\tau_0)$ and the leftmost B_{m^*} in the representation for $\Psi_{m^*, \mathbf{a}, 0}$. This will lead us to the estimate

$$\|h_{*, \tau_0}\|_{B_{pq}^{s-\kappa^*-\alpha, w}(\mathbb{R})} \gtrsim 2^{d_0(s-\kappa^*-\alpha)} [w(Q_{(d_0-1)\tau_0})]^{1/p} \Theta(m^*). \quad (3.43)$$

On the other hand, we will estimate $\|I_+^\alpha h_{*, \tau_0}\|_{B_{pq}^{s, u}(\mathbb{R})}^p$ from above via $2^{(s-\alpha)d_0} \Lambda_{n^*}^2$ by analogy with (3.39) and (3.40). Together with (3.43) this will lead us to the inequality

$$\mathfrak{C} \gtrsim 2^{-d_0\kappa^*} [u(Q_{(d_0-1)\tau_0})]^{-1/p} [w(Q_{(d_0-1)\tau_0})]^{1/p}, \quad (3.44)$$

and hence, to the required lower bound for the best constant \mathfrak{C} of the inequality (3.11). \square

For the operators $I_{c^\pm}^\alpha$ with $c \in \mathbb{R}$ the following analogous to Theorem 3.4 statement is true.

Theorem 3.5. *Let $p > 1$, $0 < q \leq \infty$, $s \in \mathbb{R}$, weights $u, v \in \mathcal{A}_\infty^{\text{loc}}$, $\alpha \in \mathbb{N}$ and $f \in L_{\text{loc}}(\mathbb{R})$. Suppose that $f \equiv 0$ on $(-\infty, c)$ in the case of the operator $I_{c^+}^\alpha$ and $f \equiv 0$ on (c, ∞) for $I_{c^-}^\alpha$.*

(i) $I_{c^\pm}^\alpha$ is bounded from $B_{pq}^{s+\kappa^*-\alpha, v}(\mathbb{R})$ to $B_{pq}^{s, u}(\mathbb{R})$ with some $\kappa^* \in \mathbb{R}$ if and only if

$$\tilde{\mathfrak{C}}_\pm^\alpha(\kappa^*, c) = \tilde{\mathbf{M}}_\pm^\alpha(1, c) + \tilde{\mathbf{M}}_\pm^\alpha(0, c) + \sup_{d \in \mathbb{N}} \tilde{\mathfrak{M}}_\pm(d, \kappa^*, c; v, u) < \infty,$$

where

$$\widetilde{\mathfrak{M}}_{\pm}(d, \kappa, c; v, u) := \sup_{\tau \in \pm \mathbb{N}_0} 2^{-d\kappa} [v(Q_{(d-1)\tau}^{(c)})]^{-\frac{1}{p}} [u(Q_{(d-1)\tau}^{(c)})]^{\frac{1}{p}}.$$

Moreover, the norm $\|I_{c^{\pm}}^{\alpha}\|_{B_{pq}^{s+\kappa^*-\alpha, v}(\mathbb{R}) \rightarrow B_{pq}^{s, u}(\mathbb{R})}$ of the operator $I_{c^{\pm}}^{\alpha}$ is equivalent to $\widetilde{\mathfrak{C}}_{\pm}^{\alpha}(\kappa^*, c)$.

(ii) Let $\kappa_* \in \mathbb{R}$ and assume that $0 < \widetilde{\mathfrak{C}}$ – the best independent of f constant in

$$\|f\|_{B_{pq}^{s-\kappa_*-\alpha, w}(\mathbb{R})} \lesssim \widetilde{\mathfrak{C}} \|I_{c^+}^{\alpha} f\|_{B_{pq}^{s, u}(\mathbb{R})}. \quad (3.45)$$

The inequality (3.45) holds if and only if

$$\widetilde{\mathfrak{C}}(\kappa_*) := \sup_{d \in \mathbb{N}_0} \widetilde{\mathfrak{M}}_{\pm}(d, \kappa_*, c; u, w) < \infty,$$

where

$$\widetilde{\mathfrak{M}}_{\pm}(d, \kappa, c; u, w) := \sup_{\tau \in \pm \mathbb{N}_0} \begin{cases} [u(Q_{0\tau}^{(c)})]^{-\frac{1}{p}} [w(Q_{0\tau}^{(c)})]^{\frac{1}{p}}, & d = 0, \\ 2^{-d\kappa} [u(Q_{(d-1)\tau}^{(c)})]^{-\frac{1}{p}} [w(Q_{(d-1)\tau}^{(c)})]^{\frac{1}{p}}, & d \in \mathbb{N}. \end{cases}$$

Besides, $\widetilde{\mathfrak{C}} \approx \widetilde{\mathfrak{C}}(\kappa_*)$.

Proof. Details of the proof can be found in [33, §6]. \square

Remark 3.1. The characteristics found in Theorem 1.1 (i) differ from those presented in Theorem 3.4 (i) by components supremal in $d \in \mathbb{N}$, namely,

$$\sup_{d \in \mathbb{N}} [\mathcal{M}_{\pm}^{\alpha}(d, \kappa^*, 1) + \mathcal{M}_{\pm}^{\alpha}(d, \kappa^*, 0)] \geq \sup_{d \in \mathbb{N}} \mathfrak{M}(d, \kappa^*; v, u).$$

The characteristics obtained in Part (ii) of Theorem 1.1 are also redundant compared to the new ones established in Theorem 3.4(ii): namely, $\mathfrak{C}_{\pm}^{\alpha}(\kappa_*) \geq \mathfrak{C}(\kappa_*)$.

The same is true relatively to the results of Theorems 1.2 and 3.5.

At the end of Section, let us give an example.

Example 3.1. Let $u(x) = w(x) = (1 + |x|)^{-t}$ and $v(x) = (1 + |x|)^{-t+sp}$ with $t > \max\{1, p\alpha - (p-1)\}$ and $\min\{1, \alpha\} < \alpha \leq s$. For the operator I_{0+}^{α} from $B_{pq}^{s-\alpha, v}(\mathbb{R})$ to $B_{pq}^{s, u}(\mathbb{R})$ we have

$$\widetilde{\mathfrak{M}}_{+}^{\alpha}(0, 0) \lesssim \begin{cases} \sup_{\tau \in \mathbb{N}_0} (1 + \tau)^{-\frac{t}{p} + \alpha}, & \alpha p \leq t < sp \\ \sup_{\tau \in \mathbb{N}_0} (1 + \tau)^{-s+1}, & t \geq sp \end{cases}$$

and

$$\widetilde{\mathfrak{M}}_{+}^{\alpha}(1, 0) \lesssim \begin{cases} \sup_{\tau \in \mathbb{N}_0} (1 + \tau)^{-\frac{t}{p} + \alpha - \frac{1}{p'}}, & t < sp \\ \sup_{\tau \in \mathbb{N}_0} (1 + \tau)^{-s+\alpha}, & t \geq sp \end{cases},$$

that is the functionals $\widetilde{\mathfrak{M}}_{+}^{\alpha}(i, 0)$, $i = 0, 1$, are finite. The condition

$$\widetilde{\mathfrak{M}}_{+}(d, 0, 0; v, u) < \infty$$

is also satisfied and $\widetilde{\mathfrak{M}}_{+}(d, 0, 0; u, w) < \infty$ (see also [26]). Therefore, under the conditions assumed and by virtue of Theorem 3.5, the operator I_{0+}^{α} is bounded from $B_{pq}^{s-\alpha, v}(\mathbb{R})$ to $B_{pq}^{s, u}(\mathbb{R})$, besides, the $B_{pq}^{s, u}(\mathbb{R}) \rightarrow B_{pq}^{s-\alpha, w}(\mathbb{R})$ norm inequality of the form (3.45) holds.

For I_{+}^{α} , assume that $t + (p-1) + p(\alpha-1) < sp$ and $\alpha p < t$. Then

$$\mathfrak{M}_{+}^{\alpha}(1) \lesssim \sup_{\tau \in \mathbb{N}} (1 + \tau)^{-\frac{t}{p} + \alpha - \frac{1}{p'}} + \sup_{\tau \in -\mathbb{N}_0} (1 - \tau)^{t/p - s + \alpha - 1/p} < \infty,$$

$$\mathfrak{M}_{+}^{\alpha}(0) \lesssim \sup_{\tau \in \mathbb{N}} (1 + \tau)^{-\frac{t}{p} + \alpha} + \sup_{\tau \in -\mathbb{N}_0} (1 - \tau)^0 < \infty.$$

Moreover, $\mathfrak{M}(d, 0; v, u) < \infty$ and $\mathfrak{M}(d, 0; u, w) < \infty$ for all $d \in \mathbb{N}_0$. Hence and by Theorem 3.4, the operator I_{+}^{α} is bounded from $B_{pq}^{s-\alpha, v}(\mathbb{R})$ to $B_{pq}^{s, u}(\mathbb{R})$ and the inequality (3.11) holds with $\kappa_* = 0$.

For the estimates in Example 3.1, the «power rule» for sums was applied (see [4, Lemma 1]).

4 Estimates under additional conditions on weights

To simplify the quantitative characteristics obtained in Theorems 3.4 and 3.5, we can consider them on certain subclasses of weight functions.

4.1 Weights with homogeneous anti-derivatives

Parameters κ_* and κ^* in the definitions of function spaces $B_{pq}^{s+\kappa^*-\alpha, v}(\mathbb{R})$ and $B_{pq}^{s-\kappa_*-\alpha, w}(\mathbb{R})$ (see Theorems 3.4 and 3.5) depend on weight functions v and w , as well as on weight u of $B_{pq}^{s, u}(\mathbb{R})$. That is, their values display themselves for specific u , v , and w only. In general case, κ_* and κ^* can be refined by narrowing the functionals $\sup_{d \in \mathbb{N}} \mathfrak{M}(d, \kappa^*; v, u)$ and $\sup_{d \in \mathbb{N}_0} \mathfrak{M}(d, \kappa_*; u, w)$ from Theorem 3.4, as well as $\sup_{d \in \mathbb{N}} \widetilde{\mathfrak{M}}_{\pm}(d, \kappa^*, c; v, u)$ and $\sup_{d \in \mathbb{N}_0} \widetilde{\mathfrak{M}}_{\pm}(d, \kappa_*, c; u, w)$ to subclasses of weights.

Consider the collection of positively homogeneous weights. Recall that a function $h: \mathbb{R} \rightarrow \mathbb{R}$ is positively homogeneous of degree $s(h)$ if for any $x \in \mathbb{R}$ from the function's domain and any $\lambda > 0$

$$h(\lambda x) = \lambda^{s_h} h(x).$$

The number s_h is called the order of homogeneity.

In our situation, parameter λ will take values 2^{-d} , $d \in \mathbb{N}$. Given that the anti-derivative of each weight σ_i , $i = 1, 2$, is a homogeneous function Σ_i of degree s_{Σ_i} , one can give the functional $\widetilde{\mathfrak{M}}_{\pm}(d, \kappa_*, c; \sigma_1, \sigma_2)$ a more transparent form in the case of $d \in \mathbb{N}$. Namely,

$$\widetilde{\mathfrak{M}}_{\pm}(d, \kappa, c; \sigma_1, \sigma_2) = 2^{-d\kappa} 2^{\frac{d-1}{p}(s_{\Sigma_1} - s_{\Sigma_2})} \widetilde{\mathfrak{M}}_{\pm}(0, \kappa, c; \sigma_1, \sigma_2).$$

Assuming $\kappa = (s_{\Sigma_1} - s_{\Sigma_2})/p$, we obtain that

$$\sup_{d \in \mathbb{N}_0} \widetilde{\mathfrak{M}}_{\pm}(d, \kappa, c; \sigma_1, \sigma_2) \simeq \widetilde{\mathfrak{M}}_{\pm}(0, \kappa, c; \sigma_1, \sigma_2).$$

Examples are $\Sigma_i(x) \simeq |x|^{\zeta}$ на $|x| \geq |c|$ for I_{c+}^{α} with $c > 0$ or I_{c-}^{α} with $c < 0$.

To simplify functionals of the type \mathfrak{M} , one can also use the direct (2.2) and/or inverse (2.3) doubling conditions for Muckenhoupt-type functions in combination with Lemma 2.1.

4.2 Using an average type condition

The functionals $\mathbf{M}_{\pm}^{\alpha}(\varepsilon)$, as well as $\widetilde{\mathbf{M}}_{\pm}^{\alpha}(\varepsilon, c)$, $\varepsilon = 0, 1$, in assertions (i) of Theorems 3.4 and 3.5 can be reduced to integral form if weights σ involved satisfy the condition

$$\int_r^{r+1} \sigma \approx \sigma(\tau) \quad \text{for some } \tau \in [r, r+1] \quad (4.1)$$

(see [30, 31]). The result when $\sigma = u$ and $\sigma = v$ are of the type (4.1) is as follows:

$$\mathbf{M}_{\pm}^{\theta}(\varepsilon) \approx \sup_{\tau \in \mathbb{Z}} \left(\int_{\tau}^{\infty} (x - \tau + 1)^{p(\theta-1)\varepsilon} u(x) dx \right)^{\frac{1}{p}} \left(\int_{-\infty}^{\tau} (\tau - y + 1)^{p'(\theta-1)(1-\varepsilon)} [v(y)]^{1-p'} dy \right)^{\frac{1}{p'}},$$

$$\mathbf{M}_{-}^{\theta}(\varepsilon) \approx \sup_{\tau \in \mathbb{Z}} \left(\int_{-\infty}^{\tau} (\tau - x + 1)^{p(\theta-1)\varepsilon} u(x) dx \right)^{\frac{1}{p}} \left(\int_{\tau}^{\infty} (y - \tau + 1)^{p'(\theta-1)(1-\varepsilon)} [v(y)]^{1-p'} dy \right)^{\frac{1}{p'}},$$

$$\widetilde{\mathbf{M}}_{\pm}^{\theta}(\varepsilon) \approx \sup_{\tau \in \mathbb{N}_0} \left(\int_{\tau+c}^{\infty} (x - \tau - c + 1)^{p(\theta-1)\varepsilon} u(x) dx \right)^{\frac{1}{p}}$$

$$\begin{aligned} & \times \left(\int_c^{\tau+c} (\tau + c - y + 1)^{p'(\theta-1)(1-\varepsilon)} [v(y)]^{1-p'} dy \right)^{\frac{1}{p'}}, \\ \mathbf{M}_-^\theta(\varepsilon) & \approx \sup_{\tau \in -\mathbb{N}_0} \left(\int_{-\infty}^{\tau+c} (\tau + c - x + 1)^{p(\theta-1)\varepsilon} u(x) dx \right)^{\frac{1}{p}} \\ & \times \left(\int_{\tau+c}^c (y - \tau - c + 1)^{p'(\theta-1)(1-\varepsilon)} [v(y)]^{1-p'} dy \right)^{\frac{1}{p'}}. \end{aligned}$$

5 Appendix

Here we give precise formulations of elements of the Battle–Lemarié type wavelet systems, in terms of which theorems of decomposition of Besov spaces $B_{pq}^{s,w}(\mathbb{R})$ are formulated in §3.1.

We start by defining a basic spline of order $n \in \mathbb{N}_0$: namely, $B_0 = \chi_{[0,1]}$ and

$$B_n(x) := (B_{n-1} * B_0)(x) = \int_0^1 B_{n-1}(x-t) dt.$$

The key property of B_n in our case is the property of its derivative $B_n^{(k)}$ of order k :

$$B_n^{(k)}(\cdot) = \sum_{l=0}^k \binom{k}{l} B_{n-k}(\cdot - l) \quad (n \geq k \in \mathbb{N}). \quad (5.1)$$

Other properties of B_n can be seen in [1] and, for example, in [33].

The B_n is a scaling function, i.e. it generates in $L^2(\mathbb{R})$ the so-called multi-resolution analysis KMA_{B_n} in the form of expanding subspaces $V_d \subset V_{d+1}$, $d \in \mathbb{N}_0$, such that $\text{clos}_{L^2(\mathbb{R})} \left(\bigcup_{d \in \mathbb{Z}} V_d \right) = L^2(\mathbb{R})$ and $\bigcap_{d \in \mathbb{Z}} V_d = \{0\}$. Moreover, for each $d \in \mathbb{N}_0$ the system

$$\{B_n(2^d \cdot -\tau)\}_{\tau \in \mathbb{Z}}$$

of integer shifts of dyadic dilations of B-spline B_n , which has compact support $[0, n+1]$, forms a non-orthogonal Riesz basis in $V_d \subset L^2(\mathbb{R})$.

Let W_d denote the orthogonal complement of V_d to V_{d+1} , i.e. $V_{d+1} = V_d \oplus W_d$ for all $d \in \mathbb{N}$. Then $W_i \perp W_j$ for $i \neq j$. From this, in particular, we obtain that

$$L^2(\mathbb{R}) = V_0 \oplus \left(\bigoplus_{l \geq 0} W_l \right).$$

The spaces W_d , $d \in \mathbb{Z}$, are called wavelet spaces. Just as the spaces V_d are generated by integer shifts of dyadic dilations of the scaling function B_n , the spaces W_d are generated by basis functions–wavelets, constructed from B_n . The multiresolution analysis KMA_ϕ with $\phi = B_n$ allows one to construct a wavelet Riesz basis in $L^2(\mathbb{R})$ using integer shifts of B_n and the same type of shifts of dyadic dilations of the corresponding to B_n wavelets. Within the framework of KMA_{B_n} , one can move from the scaling function B_n to another, orthonormal one, satisfying the relation

$$\hat{\phi}_n(\omega) = \frac{\beta_n \hat{B}_n(\omega)}{(e^{i\omega r_1} + 1) \dots (e^{i\omega r_n} + 1)} = \beta_n \sum_{l_1=0}^{\infty} (-r_1 e^{i\omega})^{l_1} \dots \sum_{l_n=0}^{\infty} (-r_n e^{i\omega})^{l_n} \hat{B}_n(\omega). \quad (5.2)$$

Here $\beta_n := (r_1 + 1) \dots (r_n + 1)$, where the sets $\{r_j\}_{j=1}^n \subset (0, 1)$ are unique for each n , and r_j are the roots of the Euler trigonometric polynomial (see, for example, [27]). The function ϕ_n is named after G. Battle [2,3] and P. G. Lemarié–Rieusset [11]. One can write (see e.g. [14,27,33])

$$\phi_n(x) = \beta_n \sum_{l_1=0}^{\infty} (-r_1)^{l_1} \dots \sum_{l_n=0}^{\infty} (-r_n)^{l_n} B_n(x + l_1 \dots + l_n).$$

The Battle–Lemarié wavelets ψ_n corresponding to ϕ_n satisfy the condition (see [27, 33])

$$\begin{aligned} \hat{\psi}_{n,\mathbf{s}}(\omega) &= \frac{\gamma_n}{2e^{i\omega\mathbf{s}}} (1/r_1 - e^{-i\omega/2}) \dots (1/r_n - e^{-i\omega/2}) \sum_{k=0}^{n+1} \frac{(-1)^k (n+1)!}{k!(n+1-k)!} e^{(n-k)i\omega/2} \\ &\quad \times \prod_{j=1}^n \sum_{m_j=0}^{\infty} (-r_j e^{-i\omega})^{m_j} \sum_{l_j=0}^{\infty} (-r_j e^{i\omega/2})^{l_j} \hat{B}_n(\omega/2), \quad (\mathbf{s} \in \mathbb{Z}) \end{aligned} \quad (5.3)$$

where $\gamma_n = 2^{-n} \beta_n(r_1 \dots r_n)$ and \mathbf{s} is fixed. Their integer shifts form an orthonormal basis in $W_0 \subset L^2(\mathbb{R})$. By analogy with ϕ_n , one can write:

$$\begin{aligned} \psi_{n,\mathbf{s}}(\cdot + \mathbf{s}) &= \gamma_n \sum_{m_1=0}^{\infty} (-r_1)^{m_1} \sum_{l_1=0}^{\infty} (-r_1)^{l_1} \dots \sum_{m_n=0}^{\infty} (-r_n)^{m_n} \sum_{l_n=0}^{\infty} (-r_n)^{l_n} \\ &\quad \left[\sum_{k=0}^{n+1} \frac{(-1)^k (n+1)!}{k!(n+1-k)!} B_n(2 \cdot -k - 2m_1 + l_1 - \dots - 2m_n + l_n) \right. \\ &\quad - \left\{ \sum_{j_1 \in \{1, \dots, n\}} \frac{1}{r_{j_1}} \right\} \sum_{k=0}^{n+1} \frac{(-1)^k (n+1)!}{k!(n+1-k)!} B_n(2 \cdot -k + 1 - 2m_1 + l_1 - \dots - 2m_n + l_n) + \\ &\quad \left\{ \sum_{\substack{j_1, j_2 \in \{1, \dots, n\} \\ j_1 \neq j_2}} \frac{1}{r_{j_1} r_{j_2}} \right\} \sum_{k=0}^{n+1} \frac{(-1)^k (n+1)!}{k!(n+1-k)!} B_n(2 \cdot -k + 2 - 2m_1 + l_1 - \dots - 2m_n + l_n) + \dots \\ &\quad \dots + \left\{ \sum_{\substack{j_1, \dots, j_c \in \{1, \dots, n\} \\ j_1 \neq \dots \neq j_c}} \frac{(-1)^c}{r_{j_1} \dots r_{j_c}} \right\} \sum_{k=0}^{n+1} \frac{(-1)^k (n+1)!}{k!(n+1-k)!} B_n(2 \cdot -k + c - 2m_1 + l_1 - \dots - 2m_n + l_n) + \dots \\ &\quad \left. + \dots + \frac{(-1)^n}{r_1 \dots r_n} \sum_{k=0}^{n+1} \frac{(-1)^k (n+1)!}{k!(n+1-k)!} B_n(2 \cdot +(n-k) - 2m_1 + l_1 - \dots - 2m_n + l_n) \right]. \end{aligned}$$

The Fourier transform of $\psi_{n,\mathbf{s}}$ in (5.3) can be represented as (see details in [33, (2.31)])

$$\begin{aligned} \hat{\psi}_{n,\mathbf{s}}(\omega) &= \frac{\gamma_n}{2e^{i\omega\mathbf{s}}} \sum_{k=0}^{n+1} \frac{(-1)^k (n+1)!}{k!(n+1-k)!} e^{(n-k)i\omega/2} \\ &\quad \times \prod_{j=1}^n \frac{|1 - e^{i\omega/2} r_j|^2}{r_j} \prod_{j=1}^n \sum_{m_j=0}^{\infty} (-r_j e^{-i\omega})^{m_j} \sum_{l_j=0}^{\infty} (r_j^2 e^{i\omega})^{l_j} \hat{B}_n(\omega/2). \end{aligned}$$

The orthonormal wavelet basis in $L^2(\mathbb{R})$ generated by the pair $\{\phi_n, \psi_{n,\mathbf{s}}\}$ is formed as follows: it is $\sqrt{2}\phi_n(x - \tau)$ and $2^{d-1}\psi_{n,\mathbf{s}}(2^{d-1}x - \tau)$ for all $\tau \in \mathbb{Z}$ and $d \in \mathbb{N}_0$.

Instead of B_n one can take the function $B_{n,\mathbf{a}}(\cdot) := B_n(\cdot - \mathbf{a})$ with $\mathbf{a} \in \mathbb{R}$. This would form another $KMA_{B_{n,\mathbf{a}}}$ and, generally speaking, some other system $\{\phi_{n,\mathbf{a}}, \psi_{n,\mathbf{a},\mathbf{s}}\}$, generating an orthonormal basis of Battle–Lemarié wavelets in $L^2(\mathbb{R})$.

Since $\phi_{n,\mathbf{a}}$ and $\psi_{n,\mathbf{a},\mathbf{s}}$ have unbounded supports, in [29], with the aim of effectively using spline systems of this type in decomposition theorems, a so-called localisation algorithm was proposed. It consists in forming, within the framework of $KMA_{B_{n,\mathbf{a}}}$, new basis elements which are finite linear combinations of integer shifts of the scaling or wavelet Battle–Lemarié functions.

As a localised analogue of $\phi_{n,\mathbf{a}}$ we will use the function $\Phi_{n,\mathbf{a}} = B_{n,\mathbf{a}}$, which turns out to be a linear combination of $n+1$ integer shifts of $\phi_{n,\mathbf{a}}$. Indeed, on the strength of (5.2),

$$\hat{\phi}_{n,\mathbf{a}}(\omega) \prod_{j=1}^n \frac{1}{r_j + 1} (e^{i\omega} r_j + 1) = \hat{B}_{n,\mathbf{a}}(\omega) =: \hat{\Phi}_{n,\mathbf{a}}(\omega). \quad (5.4)$$

Therefore, we have

$$\text{supp } \Phi_{n,\mathbf{a}} = [\mathbf{a}, \mathbf{a} + n + 1] \quad \text{and} \quad \Phi_{n,\mathbf{a}}(\cdot) = \beta_n^{-1} \sum_{\varkappa=-n}^0 \alpha'_{\varkappa} \cdot \phi_{n,\mathbf{a}}(\cdot - \varkappa), \quad (5.5)$$

where $\alpha'_{\varkappa} > 0$ for all $\varkappa = -n, \dots, 0$ and $\sum_{\varkappa=-n}^0 \alpha'_{\varkappa} = \beta_n$. Recall that integer shifts of $\phi_{n,\mathbf{a}}$ in (5.5) are mutually orthogonal, moreover, $0 < r_j < 1$ for all $j = 1, \dots, n$ in (5.4).

As a localised analogue of $\psi_{n,\mathbf{a},\mathbf{s}}$, we consider the function $\Psi_{n,\mathbf{a},\mathbf{s}}$, satisfying the condition

$$\begin{aligned} \hat{\Psi}_{n,\mathbf{a},\mathbf{s}}(\omega) &= \gamma_n^{-1} \hat{\psi}_{n,\mathbf{a},\mathbf{s}}(\omega) \prod_{j=1}^n (1 + e^{-i\omega} r_j)(1 - e^{i\omega} r_j^2) \\ &= \frac{\hat{B}_{n,\mathbf{a}}(\omega/2)}{2e^{i\omega\mathbf{s}}} \sum_{k=0}^{n+1} \frac{(-1)^k (n+1)!}{k!(n+1-k)!} e^{(n-k)i\omega/2} \prod_{j=1}^n \frac{|1 - e^{i\omega/2} r_j|^2}{r_j} \\ &= \frac{\hat{B}_{n,\mathbf{a}}(\omega/2)}{2e^{i\omega\mathbf{s}}} \sum_{k=0}^{n+1} \frac{(-1)^k (n+1)!}{k!(n+1-k)!} e^{(n-k)i\omega/2} \prod_{j=1}^n [\rho_j - 2 \cos(j\omega/2)], \end{aligned} \quad (5.6)$$

where $\rho_j := r_j + 1/r_j = 2(2\alpha_j - 1)$ with $\alpha_j > 1$, $j = 1, \dots, n$ (see details in [33]). From this it is clear that $\Psi_{n,\mathbf{a},\mathbf{s}}$ is, on the one hand, a linear combination of $2n + 1$ mutually orthogonal integer shifts of $\psi_{n,\mathbf{s},\mathbf{a}}$. On the other hand, $\Psi_{n,\mathbf{a},\mathbf{s}}$ is a finite linear combination of integer and half shifts of $B_{n,\mathbf{a}}(2\cdot)$. Since all r_j lie inside $(0, 1)$, the system $\{\Psi_{n,\mathbf{a},\mathbf{s}}(\cdot - \tau)\}_{\tau \in \mathbb{Z}}$ forms a Riesz basis in $W_0(B_{n,\mathbf{a}})$. Also observe that

$$\text{supp } \Psi_{n,\mathbf{a},\mathbf{s}} = [\mathbf{s} + \mathbf{a} - n, \mathbf{s} + \mathbf{a} + n + 1] \quad \text{and} \quad \Psi_{n,\mathbf{a},\mathbf{s}}(\cdot) = \gamma_n^{-1} \sum_{|\varkappa| \leq n} \alpha''_{\varkappa} \cdot \psi_{n,\mathbf{a},\mathbf{s}}(\cdot - \varkappa), \quad (5.7)$$

where

$$\gamma_n^{-1} \sum_{|\varkappa| \leq n} \alpha''_{\varkappa} = 2^n \prod_{j=1}^n \frac{(1+r_j)(1-r_j^2)}{r_j(1+r_j)} = 2^n \prod_{j=1}^n (1/r_j - r_j) := \Lambda_n > 0.$$

To formulate decomposition theorems, we will also need functions of the form:

$$\hat{\Psi}_{n,\mathbf{a},\mathbf{s};m(\mathbb{k}),\alpha(\zeta)}(\omega) = \hat{\Psi}_{n,\mathbf{a},\mathbf{s}}(\omega) \left[\prod_{j=1}^m [\rho_j(m) - 2 \cos(\omega/2)] \right]^{\mathbb{k}} \left[\sum_{\iota=0}^{2\alpha} \frac{(-1)^\iota (2\alpha)!}{\iota! (2\alpha - \iota)!} e^{(\alpha-\iota)i\omega/2} \right]^{\zeta},$$

where $\mathbb{k}, \zeta \in \{0, 1\}$. Due to the fact that $r_j(m) [\rho_j(m) - 2 \cos(\omega/2)] = |1 - e^{i\omega/2} r_j(m)|^2$ and

$$\sum_{\iota=0}^{2\alpha} \frac{(-1)^\iota (2\alpha)!}{\iota! (2\alpha - \iota)!} e^{(\alpha-\iota)i\omega/2} = e^{\alpha i\omega/2} (1 - e^{-i\omega/2})^{2\alpha},$$

the element $\Psi_{n,\mathbf{a},\mathbf{s};m(1),\alpha(\zeta)}$, $\zeta = 0, 1$, is linear combination of integer and half shifts of $\Psi_{n,\mathbf{a},\mathbf{s}}$. Observe that

$$\text{supp } \Psi_{n,\mathbf{a},\mathbf{s};m(\mathbb{k}),\alpha(\zeta)} = [\mathbf{s} + \mathbf{a} - n - m\mathbb{k}/2 - \alpha\zeta/2, \mathbf{s} + \mathbf{a} + n + 1 + m\mathbb{k}/2 + \alpha\zeta/2] \quad (5.8)$$

and

$$\Psi_{n,\mathbf{a},\mathbf{s};m(1),\alpha(0)}(\cdot) = \sum_{|\varkappa| \leq m} \lambda_{|\varkappa|}(m) (-1)^{|\varkappa|} \Psi_{n,\mathbf{a},\mathbf{s}}(\cdot - \varkappa/2), \quad (5.9)$$

$$\begin{aligned} \Psi_{n,\mathbf{a},\mathbf{s};m(1),\alpha(1)}(\cdot) &= \sum_{|\mathbf{z}|\leq m} \lambda_{|\mathbf{z}|}(m)(-1)^{|\mathbf{z}|} \sum_{\zeta=0}^{2\alpha} \binom{2\alpha}{\zeta} (-1)^\zeta \Psi_{n,\mathbf{a},\mathbf{s}}(\cdot - \zeta/2 + \alpha/2 - \mathbf{z}/2) \\ &\stackrel{c}{=} \sum_{|\mathbf{z}|\leq m} \lambda_{|\mathbf{z}|}(m)(-1)^{|\mathbf{z}|} = \prod_{j=1}^m \frac{1 - r_j^2(m)}{r_j(m)} = 2^{-m} \Lambda_m, \end{aligned} \quad (5.10)$$

(see [33, p. 18]), besides,

$$\prod_{j=1}^m [(r_j + 1/r_j) - 2 \cos(\omega/2)] =: \sum_{j=0}^m (-1)^j \lambda_j \cos(j\omega/2), \quad (5.11)$$

where $\lambda_m = 2$ and $0 < \lambda_j = \lambda(\rho_1, \dots, \rho_m)$ in the case $j \neq m$, and λ_j are even for $j \neq 0$. For completeness, we will also write down that (see (5.6))

$$\Psi_{n,\mathbf{a},\mathbf{s}}(\cdot) = \gamma_n \sum_{|s|\leq n} \lambda_{|s|}(n)(-1)^{|s|} \sum_{k=0}^{n+1} (-1)^k \binom{1+n}{k} B_n(2\cdot + (n-k) - s). \quad (5.12)$$

In conclusion of Section, for $\mathbf{a} \in \mathbb{R}$ we define pairs of functions of the form

$$\tilde{\Phi} = \Phi_{n,\mathbf{a}} \quad \text{and} \quad \tilde{\Psi}^{(\mathbb{k},\zeta)} = \Psi_{n,\mathbf{a},\mathbf{s};m(\mathbb{k}),\alpha(\zeta)}/\Lambda_n, \quad \text{where } \mathbb{k}, \zeta \in \{0, 1\}; \quad (5.13)$$

$$\Phi^{(a)} = \Phi_{n,\mathbf{a}} \quad \text{and} \quad \Psi^{(a)} = \Psi_{n,\mathbf{a},\mathbf{s}}/\Lambda_n. \quad (5.14)$$

In (5.13) and (5.14) the first components $\tilde{\Phi}$ and $\Phi^{(a)}$ (the scaling functions or the father wavelets) are identical to each other. We denote them differently only to distinguish the wavelet systems $\{\tilde{\Phi}, \tilde{\Psi}^{(\mathbb{k},\zeta)}\}$ and $\{\Phi^{(a)}, \Psi^{(a)}\}$ from each other. The difference in the systems is in the second components $\tilde{\Psi}^{(\mathbb{k},\zeta)}$ and $\Psi^{(a)}$ or the wavelet functions (the mother wavelets). Here $\tilde{\Psi}^{(\mathbb{k},\zeta)}$ is a more general function, which, however, coincides with $\Psi^{(a)}$ when $\mathbb{k} = \zeta = 0$.

The smoothness of the functions in (5.13) and (5.14) is n . For $x \in \mathbb{R}$ we denote:

$$\tilde{\Phi}_\tau(x) := \tilde{\Phi}(x - \tau) \quad \text{and} \quad \tilde{\Psi}_{d\tau}^{(\mathbb{k},\zeta)}(x) := 2^{d/2} \tilde{\Psi}^{(\mathbb{k},\zeta)}(2^d x - \tau) \quad (\tau \in \mathbb{Z}, d \in \mathbb{N}_0), \quad (5.15)$$

and also,

$$\Phi_\tau^{(a)}(x) := \Phi^{(a)}(x - \tau) \quad \text{and} \quad \Psi_{d\tau}^{(a)}(x) := 2^{d/2} \Psi^{(a)}(2^d x - \tau) \quad (\tau \in \mathbb{Z}, d \in \mathbb{N}_0). \quad (5.16)$$

As noted in [33, Remark 2.4], the construction of $\Psi_{n,\mathbf{a},\mathbf{s};m(\mathbb{k}),\alpha(\zeta)}(\cdot - \tau)$, $\mathbb{k}, \zeta \in \{0, 1\}$, starts with $B_{n,2(s+\mathbf{a}+\tau)}(\cdot)$. As for observations (5.7) and (5.8) concerning supports of $\Psi_{n,\mathbf{a},\mathbf{s};m(\mathbb{k}),\alpha(\zeta)}$, they are slightly corrected in this paper relatively to the information presented in [33, §2].

The systems (5.16), as well as (5.15), if they are generated by elements (5.13) with $\mathbb{k} = \zeta = 0$, form a semi-orthogonal Riesz basis in $L^2(\mathbb{R})$ (i.e. elements from V_0 are orthogonal to W_l with $l \geq 0$, and functions of the space W_{l_0} with fixed $l_0 \in \mathbb{N}_0$ are orthogonal to W_l if $l \neq l_0$). The special structure of the elements of such systems allows them to be used for obtaining lower bounds (or proofs of necessity) in decomposition theorems (see § 3.1). Upper bounds in such theorems (or proofs of sufficiency) are less demanding and allow representation with respect to combined elements of the type (5.15), when they are generated by functions (5.13) with $\mathbb{k} = 1$ and/or $\zeta = 1$.

Acknowledgment. The research presented in § 4 of the paper was performed at Steklov Mathematical Institute of Russian Academy of Sciences under support of the RSF grant 24-11-00170, <https://rscf.ru/project/24-11-00170/>. The remaining work was done in framework of the state assignments of Ministry of Science and Higher Education of Russia for V. A. Trapeznikov Institute of Control Sciences of Russian Academy of Sciences and Computing Center of Far Eastern Branch of Russian Academy of Sciences.

Bibliography

- [1] Chui C. K. *An Introduction to Wavelets*. NY: Academic Press, 1992.
- [2] Battle G. *A block spin construction of ondelettes, Part I: Lemarie functions*, *Comm. Math. Phys.* **110** (1987), 601–615.
- [3] Battle G. *A block spin construction of ondelettes, Part II: QFT connection*, *Comm. Math. Phys.* **114** (1988), 93–102.
- [4] Bennett G., Grosse-Erdmann K.-G., *On series of positive terms*, *Houston J. Math.* **31**:2 (2005), 541–586.
- [5] Berkolaiko M. Z., Novikov I. Ya. *Images of wavelets under the influence of convolution operators*, *Math. Notes* **55**:5 (1994), 13–24.
- [6] Dyda, B., Ihnatsyeva, L., Lehrbäck, J. et al. *Muckenhoupt A_p -properties of distance functions and applications to Hardy–Sobolev –type inequalities*, *Potential Anal.* **50** (2019), 83–105.
- [7] Haroske D., Triebel H. *Wavelet bases and entropy numbers in weighted function spaces*, *Math. Nachr.* **278** (2005), 108–132.
- [8] Haroske D. D., Piotrowska I. *Atomic decompositions of function spaces with Muckenhoupt weights, and some relation to fractal analysis*, *Math. Nachr.* **281**:10 (2008), 1476–1494.
- [9] Haroske D., Skrzypczak L. *Entropy and approximation numbers of embeddings of function spaces with Muckenhoupt weights, I*, *Rev. Mat. Complut.* **21**:1 (2008), 135–177.
- [10] Izuki I., Sawano Y. *Atomic decomposition for weighted Besov and Triebel–Lizorkin spaces with local class of weights*, *Math. Nachr.* **285** (2012), 103–126.
- [11] Lemarie P. G. *Une nouvelle base d’ondelettes de $L^2(\mathbb{R}^n)$* , *J. de Math. Pures et Appl.* **67** (1988), 227–236.
- [12] Malecka A. *Haar functions in weighted Besov and Triebel–Lizorkin spaces*, *J. Approx. Theory* **200** (2015), 1–27.
- [13] Muckenhoupt B. *Weighted norm inequalities for the Hardy maximal function*, *Trans. Amer. Math. Soc.* **165** (1972), 207–226.
- [14] Novikov I. Ya., Stechkin S. B. *Basic constructions of wavelets*, *Fundam. Prikl. Mat.* **3**:4 (1997), 999–1028.
- [15] Prokhorov D. V., Stepanov V. D., Ushakova E. P. *Hardy–Steklov integral operators. Part I*, *Proc. Steklov Inst. Math. (Supplementary issues)* **300**:2 (2018), 1–112.
- [16] Roudenko S. *Matrix-weighted Besov spaces*, *Trans. Amer. Math. Soc.* **355**:1 (2002), 273–314.
- [17] Rychkov V. S. *Littlewood–Paley theory and function spaces with A_p^{loc} weights*, *Math. Nachr.* **224** (2001), 145–180.
- [18] Samko S. G., Kilbas A. A., Marichev O. I. *Fractional integrals and derivatives: theory and applications*. Gordon and Breach Science Publishers, 1993.
- [19] Schott T. *Function spaces with exponential weights. I*, *Math. Nachr.* **189** (1998), 221–242.
- [20] Schott T. *Function spaces with exponential weights. II*, *Math. Nachr.* **196** (1998), 231–250.

- [21] Sickel W., Triebel H. *Hölder inequalities and sharp embeddings in function spaces of B_{pq}^s and F_{pq}^s type*, Z. Anal. Anwend. **14**:1 (1995), no. 1, 105–140.
- [22] Stein E. M. Harmonic analysis. Princeton University Press, Princeton, 1993.
- [23] Triebel H. Theory of function spaces, Birkhäuser Verlag, Basel, 1983.
- [24] Triebel H. Theory of function spaces II. Birkhäuser Verlag, Basel, 1992.
- [25] Triebel H. Tempered homogeneous function spaces. European Math. Soc. Publishing House, Zurich, 2015.
- [26] Nasyrova M. G., Ushakova E. P. *Wavelet bases and entropy numbers of Hardy operator*, Anal. Math. **44**:4 (2018), 543–576.
- [27] Ushakova E. P., Ushakova K. E. *Localisation property of Battle-Lemarié wavelets' sums*, J. Math. Anal. Appl. **461**:1 (2018), 176–197.
- [28] Ushakova E. P. *Spline wavelet bases in function spaces with Muckenhoupt weights*, Rev. Mat. Complut. **33**:1 (2020), 125–160.
- [29] Ushakova E. P. *Spline wavelet decomposition in weighted function spaces*, Proc. Steklov Inst. Math. **312** (2021), 301–324.
- [30] Ushakova E. P. *The images of integration operators in weighted function spaces*, Siber. Math. J. **63**:6 (2022), 1181–1207.
- [31] Ushakova E. P., Ushakova K. E. *Norm inequalities with fractional integrals*, St. Petersburg Math. J. **35**:3 (2024), 571–595.
- [32] Ushakova E. P. *Boundedness of the Hilbert transform in Besov spaces*, Anal. Math. **49**:4 (2023), 1137–1174.
- [33] Ushakova E. P. *The study by splines of norm inequalities for Riemann–Liouville operators in weighted Besov spaces*, Journal of Mathematical Sciences **480**:5 (2024), 758–820.
- [34] Wojciechowska A. *Local means and wavelets in function spaces with local Muckenhoupt weights*, in: Function spaces IX. Vol. 92. Banach Center Publications, 2011, 399–412.
- [35] Wojciechowska A. Multidimensional wavelet bases in Besov and Triebel–Lizorkin spaces. PhD. thesis, Adam Mickiewicz University Poznań, Poznań, 2012.