

# Inferring diffusivity from killed diffusion

Richard Nickl and Fanny Seizilles

UNIVERSITY OF CAMBRIDGE

March 20, 2025

## Abstract

We consider diffusion of independent molecules in an insulated Euclidean domain with unknown diffusivity parameter. At a random time and position, the molecules may bind and stop diffusing in dependence of a given ‘binding potential’. The binding process can be modeled by an additive random functional corresponding to the canonical construction of a ‘killed’ diffusion Markov process. We study the problem of conducting inference on the infinite-dimensional diffusion parameter from a histogram plot of the ‘killing’ positions of the process. We show first that these positions follow a Poisson point process whose intensity measure is determined by the solution of a certain Schrödinger equation. The inference problem can then be re-cast as a non-linear inverse problem for this PDE, which we show to be consistently solvable in a Bayesian way under natural conditions on the initial state of the diffusion, provided the binding potential is not too ‘aggressive’. In the course of our proofs we obtain novel posterior contraction rate results for high-dimensional Poisson count data that are of independent interest. A numerical illustration of the algorithm by standard MCMC methods is also provided.

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
1.1	Observation model and main results . . . . .	2
1.2	Basic notation . . . . .	3
<b>2</b>	<b>Posterior consistency for high-dimensional Poisson count data</b>	<b>4</b>
2.1	Measurement setup and information inequalities . . . . .	4
2.2	The basic contraction theorem . . . . .	5
2.3	Contraction rates for intensity densities . . . . .	6
<b>3</b>	<b>Derivation of the observation model and proof of Theorem 1.1</b>	<b>7</b>
3.1	Probability density of the binding location $\times$ time . . . . .	8
3.2	Construction of the Poisson point process . . . . .	11
<b>4</b>	<b>Inference on the diffusion parameter from killed diffusion</b>	<b>11</b>
4.1	Construction of the prior and posterior . . . . .	12
4.2	Numerical illustration . . . . .	13
4.3	Posterior consistency theorems . . . . .	14
<b>5</b>	<b>Analytical results from PDE theory</b>	<b>17</b>
5.1	Schrödinger operators and their spectrum . . . . .	17
5.2	Identifiability and Stability. . . . .	20
5.3	Proof of Theorem 5.6 . . . . .	21

arXiv:2503.14978v1 [math.ST] 19 Mar 2025

<b>6 Proofs for Section 2</b>	<b>23</b>
6.1 Proof of Lemma 2.1	23
6.2 Proof of Lemma 2.2	24
6.3 Proof of Theorem 2.3	25
6.4 A concentration inequality for high-dimensional Poisson regression	27

# 1 Introduction

## 1.1 Observation model and main results

Diffusion of particles in an insulated domain  $\Omega \subset \mathbb{R}^d$  is mathematically described by a stochastic differential equation (SDE) with reflection at the boundary  $\partial\Omega$ : we consider the Markov process  $(X_t : t \geq 0)$  starting at initial distribution  $X_0 \sim \phi$  and then evolving according to the SDE

$$dX_t = \nabla D(X_t)dt + \sqrt{2D(X_t)}dW_t + \nu(X_t)dL_t, \quad t > 0, \tag{1}$$

where  $D : \Omega \rightarrow [D_{min}, \infty)$ ,  $D_{min} > 0$ , is a real-valued diffusion parameter with Lipschitz gradient vector field  $\nabla D$ , and  $(W_t)$  is a  $d$ -dimensional Brownian motion. Moreover  $(L_t)$  is a continuous non-decreasing process, non-zero only when  $X_t$  is at  $\partial\Omega$ , and  $\nu(x)$ ,  $x \in \partial\Omega$ , is the inward pointing normal vector. To avoid technicalities we assume that  $\Omega$  is convex with smooth boundary – the existence of a process  $(X_t)$  with almost surely continuous sample paths is then shown in [40, 28]. One can model a further drift term  $\nabla U(X_t)$  in the above SDE (see after (32) for discussion), but the main conceptual challenges already emerge when  $\nabla U = 0$ .

This work is partly motivated by observations arising in biochemistry experiments [23, 4], [29], where molecules diffuse inside a membrane until they suddenly ‘bind’ at an apparently random time and position – see Section 3 for more details. We will introduce (in fact, recall) a Markovian model for such binding events based on the notion of a ‘killed diffusion process’ (see [38, 2, 6, 3]) associated to a potential  $q : \Omega \rightarrow [0, \infty)$ . Larger values of  $q(x)$  indicate a higher probability of the process to bind when diffusing near  $x \in \Omega$ . It induces an additive functional  $\int_0^t q(X_s)ds$  – for example if  $q$  equals the indicator  $1_A$  of a subset  $A$  of  $\Omega$  this measures the occupation time of the  $(X_t)$  process in  $A$  up until  $t$ . Then let  $Y$  be a (standard) exponential random variable independent of  $(X_t)$ . The *binding time*  $S$  is the first time the cumulative binding integral of the potential along the path exceeds  $Y$ ; i.e.  $S$  is defined as

$$S = \inf \left\{ t : \int_0^t q(X_s)ds > Y \right\}. \tag{2}$$

By properties of exponential random variables, this process is ‘memory-less’ in the sense that not having bound by time  $t$  does not affect the distribution of binding events at later times. The killed diffusion process is defined as

$$\tilde{X}_t = \begin{cases} X_t, & t < S, \\ \dagger, & t \geq S; \end{cases}$$

so that  $X_t$  is ‘removed to the cemetery’  $\{\dagger\}$  at binding time  $S$ . The ‘binding’ location is given by  $X_S$  encoding the terminal position of the particle at time  $S$ .

In the abovementioned applications from biochemistry (and elsewhere), the aim is to make statistical inference on the functional parameter  $D$  from observing several molecules diffusing independently according to (1) in the presence of ‘binding’ events. The equilibrium distribution of the Markov process (1) is uniform for all  $D$  and hence the occupation times spent by the process  $X_t$  in particular areas of  $\Omega$  before binding are not informative for this task. If one can track the trajectories of the particles then methods such as those from [33, 24, 19] (see also related contributions [20, 35] for  $d = 1$  and for ‘continuous’ data in [34, 18]) can be used to infer  $D$  from the observed dynamics. This generally requires a high-resolution imaging technique that can take measurements for an extended period, but once a binding process is superimposed (or in any case) it can become difficult or expensive to collect accurate such data; for instance single particle tracking methods only allow for short path measurements [22].

A potentially much simpler observational signature is to take a dissection of  $\Omega$  into finitely many disjoint subsets  $B_1, \dots, B_K$  and to measure the relative frequency of binding events  $\{X_S \in B_i\}$  in each bin  $B_i$ . *The main contribution of this article is to demonstrate that merely observing sufficiently dense histograms of the binding locations, rather than the trajectories of diffusion, is enough to consistently identify  $D$ .* This is shown to be true as long as a) the initial state  $\phi$  of the system satisfies the mild identifiability Condition 5.5 (of being either close to equilibrium, or appropriately prepared by the experimenter), and b) if the binding potential  $q$  is not too ‘aggressive’ (i.e., small enough relative to the parameter of the exponential variable  $Y$ ), but still strictly positive on an open set, so that all molecules eventually bind at a finite time with probability one (see Proposition 3.1). If  $n_{mol}$  molecules  $X_t^m, m = 1, \dots, n_{mol}$ , diffuse independently, denote by

$$N(A) = \sum_{m=1}^{n_{mol}} 1\{X_S^m \in A\}, \quad A \subset \Omega \text{ (measurable)}, \quad (3)$$

the count of the number of molecules having bound in the subregion  $A$  of  $\Omega$  (regardless of the point in time when binding has occurred). It is physically plausible to assume that the counts for two disjoint subsets  $A, A'$  are statistically independent. In this case a consistent model requires  $n_{mol}$  to be randomly chosen from a Poisson distribution since it is a consequence of Rényi’s theorem that such ‘completely random’ point processes are necessarily Poisson processes (see Theorem 6.12 in [26] and also the discussion in Sec. 3.2). Our first main result – whose details and proof will be given in Section 3 under Conditions 4.1 and 4.2 – is:

**Theorem 1.1.** *For  $n \in \mathbb{N}$ , let  $n_{mol} \sim \text{Poisson}(n)$  be the total number of molecules diffusing independently according to (1), each with initial condition  $X_0 \sim \phi$  and binding at random time  $S$  with bounded potential  $q \geq 0$  that is strictly positive on an open subset  $\Omega_{00}$  of  $\Omega$ . Then  $N$  from (3) is a Poisson point process with intensity measure  $\Lambda$  defined by its Lebesgue density  $\lambda(x) = nq(x)u_{D,q}(x), x \in \Omega$ , where  $u = u_{D,q}$  solves the stationary Schrödinger equation  $\nabla \cdot (D\nabla u) - qu = -\phi$  on  $\Omega$  with Neumann boundary conditions.*

The connection between the intensity  $\lambda$  and an elliptic partial differential equation (PDE) revealed by the previous theorem allows to recast the problem within the paradigm of Bayesian non-linear inverse problems [39, 32] and to adapt techniques from [30, 32, 33] to the present setting. We establish necessary analytical properties of the non-linear PDE solution map  $D \mapsto u_{D,q}$ , including its injectivity under certain hypotheses (discussed in detail after Theorem 5.6), to show that this inverse problem can be solved in principle. We further obtain a novel statistical theorem about posterior contraction rates with Poisson count data of independent interest for the theory of non-linear inverse problems, where physical data often arise in this fashion. The details will be developed in the next sections, but in essence we shall show that:

**Theorem 1.2.** *Suppose  $q \geq 0$  is known, strictly positive on an open subset  $\Omega_{00}$  of  $\Omega$ , and that its Sobolev norm  $\|q\|_{H^2}$  is sufficiently small. Assume moreover that the initial condition  $X_0 \sim \phi$  satisfies the identifiability Condition 5.5. Then there exists an estimator  $\hat{D}_n$  (arising from the posterior mean of a Gaussian process prior) based on observations of the Poisson point process from the previous theorem such that any positive  $D_0 \in H^\alpha(\Omega), \alpha > 2 + d$ , that is constant near  $\partial\Omega$  can be recovered as  $n \rightarrow \infty$  at convergence rate*

$$\|\hat{D}_n - D_0\|_{L^2} = O_{P_{D_0}}(n^{-\beta}), \quad \text{for some } \beta > 0.$$

See Theorem 4.6 for full details. The posterior mean can be computed by standard MCMC and numerical PDE methods, see Figure 1 and after (32) for more discussion.

## 1.2 Basic notation

The  $L^p$ , Hölder,  $L^2$ - and  $L^p$ -Sobolev spaces over a bounded domain  $\Omega$  in  $\mathbb{R}^d$  with smooth boundary are denoted by  $L^p(\Omega), C^\alpha(\Omega), H^\alpha(\Omega), W^{a,p}(\Omega)$ , respectively. We denote by  $\|\cdot\|_B$  the norm on the resulting Banach space  $B$ , and  $\langle \cdot, \cdot \rangle_{L^2}$  denotes the inner product on  $L^2$ . By  $\|H\|_\infty$  we always denote the supremum norm over the domain of the map  $H$ , and  $dx$  denotes Lebesgue measure on  $\Omega$ , while  $\nabla, \nabla \cdot, \Delta$  denote the gradient, divergence, and Laplace operator, respectively. We also denote by  $\mathbb{R}^+$  the positive half line  $[0, \infty)$ .

We recall the well known multiplication inequalities

$$\|uv\|_{H^\alpha} \lesssim \|u\|_{B^\alpha} \|v\|_{H^\alpha}, \quad u \in B^\alpha, v \in H^\alpha, \quad (4)$$




Figure 1: Example of density reconstruction through MCMC, for  $n = 10^7$ . The true diffusivity function is displayed on the left, and generates the random observation counts (centre) on each of the  $K = 36$  bins. The posterior mean computed via MCMC is displayed on the right. More details about the implementation can be found in Sec. 4.2.

where  $B^a = C^a$  or, if  $a > d/2$ ,  $B^a = H^a$ . We also make frequent use of the standard interpolation inequalities for Sobolev spaces. First we have for all  $u \in W^{a,p}$  and  $0 \leq j \leq a$ ,  $1 \leq p < \infty$  that

$$\|u\|_{W^{j,p}} \lesssim \|u\|_{W^{a,p}}^{j/a} \|u\|_{L^p}^{(a-j)/a}, \quad (5)$$

see p.135 in [1]. The following inequality will also be useful, cf. p.139 in [1],

$$\|u\|_{L^2} \lesssim \|u\|_{W^{a,1}}^{1-\theta} \|u\|_{L^1}^\theta, \quad u \in W^{a,1} \quad (6)$$

as long as  $a > d/2$  and with  $\theta = (a - d/2)/a$ .

## 2 Posterior consistency for high-dimensional Poisson count data

In order to prove Theorem 1.2, we need a general posterior contraction theorem for Poisson count data with a finite but possibly large number  $K$  of ‘bins’  $B_1, \dots, B_K$ . This is the appropriate measurement setting for various physical experiments in inverse problems (beyond the context of this paper, also, e.g., with  $X$ -ray imaging, [31, 30]). While there are several contributions to posterior consistency with Poisson process data – see [5, 25, 9, 21, 17], Sec. 10.4.4 in [13], as well as references therein – we are not aware of a result that provides what we require. A main challenge arises from the fact that the relevant data is just independent but not iid, that the sample size  $n$  is random, and that the log-likelihood ratios appearing with the Poisson family are not bounded, so that the machinery from [13] via Hellinger testing does not apply in a straightforward way. We develop here some techniques based on the alternative approach of [15] to establish tests by sharp concentration properties of appropriate estimators for high-dimensional Poisson bin count data – these are of independent interest and can be found in Section 6, along with the other proofs for this section.

### 2.1 Measurement setup and information inequalities

We consider a measurable space  $(\mathcal{O}, \mathcal{B})$  and a collection  $\mathcal{L}$  of finite measures  $\Lambda : \mathcal{B} \rightarrow [0, \infty)$ . Let  $B_1, \dots, B_K$  be a measurable partition  $\mathcal{O} = \cup_{i=1}^K B_i$ . We are given observations drawn independently across the ‘bins’  $B_i$  from a Poisson distribution

$$Y_i \sim Poi(n\Lambda(B_i)), \quad i = 1, \dots, K. \quad (7)$$

We can think of  $n\Lambda(\mathcal{O})$  as the ‘average number of observations’, and we assume for simplicity that  $n$  is known. This models the setting where we measure  $K$  batches of count data arising from  $n$  realisations of a Poisson point process over  $\mathcal{O}$  with intensity measure  $\Lambda$ . We emphasise that we consider a high-dimensional setup in the sense that  $K = K_n \rightarrow \infty$  with  $n \rightarrow \infty$ , so that we may recover finer properties of  $\Lambda$  (such as its density w.r.t. some dominating measure) by standard approximation theoretic methods, see Subsection 2.3.

To proceed, consider  $K$  realisations  $Y_1 = y_1, \dots, Y_K = y_K$  of (7). The probability density of each observation  $Y_i$  is given by

$$p_{i,\Lambda,n}(y_i) = e^{-n\Lambda(B_i)} \frac{(n\Lambda(B_i))^{y_i}}{y_i!}, \quad y_i \in \mathbb{N} \cup \{0\} \equiv \mathbb{N}_0,$$

and, by independence the resulting likelihood function is

$$p_\Lambda(y_1, \dots, y_K) \equiv p_{\Lambda,n}^K(y_1, \dots, y_K) = \prod_{i=1}^K p_{i,\Lambda,n}(y_i). \quad (8)$$

Denote by  $P_\Lambda = P_{\Lambda,n}^K$  and  $\mathbb{E}_\Lambda = \mathbb{E}_{n,\Lambda}^K$  the probability measure and expectation operator with respect to the density  $p_{\Lambda,n}^K$ , respectively.

For two finite measures  $\Lambda, \Lambda_0$  on  $\mathcal{O}$ , we define a (semi-)metric in sequence space as

$$\|\Lambda - \Lambda_0\|_{\ell_1} = \sum_{i=1}^K |\Lambda(B_i) - \Lambda_0(B_i)| \quad (9)$$

as well as the following ‘information divergences’

$$\mathcal{D}_{2,K}^2(\Lambda, \Lambda_0) = \sum_{i=1}^K \frac{|\Lambda(B_i) - \Lambda_0(B_i)|^2}{\Lambda_0(B_i)}, \quad \mathcal{D}_{\infty,K} = \max_{1 \leq i \leq K} \frac{|\Lambda(B_i) - \Lambda_0(B_i)|}{\Lambda_0(B_i)} \quad (10)$$

which are understood to equal  $\infty$  if  $\Lambda_0(B_i) = 0$  for some  $i$  for which  $\Lambda(B_i) \neq 0$ .

We first derive a ‘local’ bound for the information distance related to such measurements.

**Lemma 2.1.** *Let  $K \in \mathbb{N}$  and let  $\Lambda, \Lambda_0$  be finite measures on  $\mathcal{O}$  such that  $\mathcal{D}_{\infty,K}(\Lambda, \Lambda_0) < 1/2$ . Then*

$$\mathbb{E}_{\Lambda_0} \left( \log \frac{p_{\Lambda_0,n}^K(Y_1, \dots, Y_K)}{p_{\Lambda,n}^K(Y_1, \dots, Y_K)} \right) \leq 2n\mathcal{D}_{2,K}^2(\Lambda, \Lambda_0)$$

Using the preceding lemma, we can obtain a lower bound on certain integrated likelihood ratios (‘evidence’ lower bound) that will be essential in the posterior contraction rate theorem to follow. For this we consider a collection  $\mathcal{L}$  of finite measures  $\Lambda$  over  $(\mathcal{O}, \mathcal{B})$  and equip  $\mathcal{L}$  with a  $\sigma$ -field  $\mathcal{S}$  such that the maps

$$(\Lambda, y) \rightarrow p_\Lambda(y_1, \dots, y_K) \quad (11)$$

are jointly Borel-measurable from  $\mathcal{L} \times \mathbb{N}_0^K$  to  $\mathbb{R}$ .

**Lemma 2.2.** *Fix  $\Lambda_0 \in \mathcal{L}$  such that  $\Lambda_0(B_i) > 0$  for all  $i$ . Let  $\nu$  be a probability measure on*

$$A := \{\Lambda \in \mathcal{L} : \mathcal{D}_{2,K}^2(\Lambda, \Lambda_0) \leq \epsilon^2, \mathcal{D}_{\infty,K}(\Lambda, \Lambda_0) \leq 1/2\} \in \mathcal{S}, \quad \epsilon > 0.$$

*Then for every  $c > 0$  there exists a constant  $L_c > 0$  such that for all  $n, \epsilon, K$  we have*

$$P_{\Lambda_0} \left( \int_A \frac{p_\Lambda}{p_{\Lambda_0}}(Y_1, \dots, Y_K) d\nu(\Lambda) \leq e^{-(2+c)n\epsilon^2} \right) \leq 2e^{-L_c n\epsilon^2}$$

## 2.2 The basic contraction theorem

We now turn to the posterior contraction theorem, and consider a prior  $\Pi$  defined on the measurable sets  $\mathcal{S}$  (from before (11)) of the collection  $\mathcal{L}$  of ‘intensity’ measures  $\Lambda$  over  $\mathcal{O}$ . The posterior distribution arising from data (7) with likelihood (8) then arises from Bayes’ formula and is given by

$$\Pi(A|Y_1, \dots, Y_K) = \frac{\int_A p_{\Lambda,n}^K(Y_1, \dots, Y_K) d\Pi(\Lambda)}{\int_{\mathcal{L}} p_{\Lambda,n}^K(Y_1, \dots, Y_K) d\Pi(\Lambda)}, \quad A \in \mathcal{S}. \quad (12)$$

Given the preparations from Section 2.1, the following proof follows the standard pattern [13], assuming the existence of sufficiently good ‘tests’ which we will construct in Theorem 6.1 below, using the concentration of measure approach of [15] adapted to high-dimensional Poisson count data.

**Theorem 2.3.** Let  $K = K_n \in \mathbb{N}$  and  $\varepsilon_n > 0$  be sequences such that

$$\frac{K_n}{n} \rightarrow 0, \varepsilon_n \leq \sqrt{\frac{K_n}{n}}, \varepsilon_n \rightarrow 0, \sqrt{n}\varepsilon_n \rightarrow \infty.$$

Suppose a sequence of prior probability measures  $\Pi = \Pi_n$  defined on  $\mathcal{L}$  satisfies, for some  $C > 0$ , some  $\Lambda_0 \in \mathcal{L}$ , and all  $n$ ,

$$\Pi(\Lambda : D_{2,K}(\Lambda, \Lambda_0) \leq \varepsilon_n^2, \mathcal{D}_{\infty,K}(\Lambda, \Lambda_0) \leq 1/2) \geq e^{-Cn\varepsilon_n^2}. \quad (13)$$

For fixed  $\Lambda_{max} < \infty$  define  $\mathcal{R} = \{\Lambda \in \mathcal{L} : \Lambda(\mathbb{O}) \leq \Lambda_{max}\}$  and assume further that there exist measurable sets  $\Theta_n \subset \mathcal{L}$  such that, for all  $n$  large enough and some  $L > 0$ ,

$$\Pi(\mathcal{L} \setminus (\Theta_n \cap \mathcal{R})) \leq Le^{-(C+4)n\varepsilon_n^2}. \quad (14)$$

Then the posterior distribution  $\Pi(\cdot | Y_1, \dots, Y_K)$  from (12) arising from data (7) contracts about the ground truth intensity measure  $\Lambda_0$  in the  $\ell_1$  metric; specifically we have for some  $k > 0$  that

$$\Pi(\Lambda \in \Theta_n \cap \mathcal{R} : \|\Lambda - \Lambda_0\|_{\ell_1} < M\sqrt{K_n/n}|Y_1, \dots, Y_K) = 1 - O_{P_{\Lambda_0}}(e^{-kn\varepsilon_n^2})$$

whenever  $M$  is large enough depending on  $C, \Lambda_{max}, k$ .

In typical applications of Bayesian nonparametrics,  $K \simeq n\varepsilon_n^2$  will be a natural choice that matches the number of bins with the effective smoothness / dimension of the prior for the unknown intensity  $\Lambda$ . However, below, we shall employ priors that arise as pushforwards of PDE solution maps and therefore naturally oversmooth. In this case the contraction rate obtained from ‘Poisson bin count’ data will be dominated by the number of bins used, hence the previous theorem is stated in this more flexible form.

The uniform bound on  $\Lambda(\mathbb{O})$  can be replaced by a condition that the posterior concentrates on uniformly bounded intensities with frequentist probability approaching one.

### 2.3 Contraction rates for intensity densities

The  $\ell_1$ -metric on the bin-counts is a natural metric for regression problems arising from (7), but does not identify the full intensity measure for fixed  $K$ . However, as  $K \rightarrow \infty$ , if the measures  $\Lambda$  have sufficiently regular densities  $\lambda$  with respect to a joint dominating measure, we can recover these consistently as  $K \rightarrow \infty$ . We now discuss this for the case relevant below where  $\mathbb{O}$  is a bounded subset of  $\mathbb{R}^d$  of positive Lebesgue measure  $dx$  and the intensities have Lipschitz densities.

Denote the  $L^2(dx)$ -orthonormal functions arising from the measurable partition of  $\mathbb{O}$  by  $e_i = 1_{B_i}/\sqrt{|B_i|}$  where  $|B|$  denotes the volume of  $B$ . The  $L^2(dx)$  projection of  $g \in L^2$  onto  $E_K = \text{span}\{e_i : i \leq K\}$  is then

$$P_{E_K}g = \sum_{i \leq K} \langle e_i, g \rangle_{L^2} e_i = \sum_{i=1}^K \frac{1}{|B_i|} \int_{B_i} g(z) dz 1_{B_i}$$

and this makes sense for any finite signed measure  $\Lambda$  if we replace  $\int_{B_i} h$  by  $\Lambda(B_i)$ . For such  $\Lambda$  we have

$$\|P_{E_K}\Lambda\|_{L^1} = \int_{\mathbb{O}} \left| \sum_{i=1}^K \frac{1}{|B_i|} \Lambda(B_i) 1_{B_i}(x) \right| dx \leq \sum_{i=1}^K |\Lambda(B_i)| = \|\Lambda\|_{\ell_1}.$$

Now given  $x \in \mathbb{O}$ , denote by  $B_i(x)$  the unique element of the partition containing  $x$ . Then if  $\Lambda$  has a Lebesgue density  $\lambda$  that is  $B$ -Lipschitz we obtain

$$|P_{E_K}\Lambda(x) - \lambda(x)| = \left| \frac{1}{|B_i(x)|} \int_{B_i(x)} (\lambda(z) - \lambda(x)) dz \right| \leq B \sup_{z \in B_i(x)} |z - x|_{\mathbb{R}^d} \leq c(B, d)h$$

if each bin  $B_i$  has Euclidean diameter at most  $h$ . This forces  $K \simeq h^{-d}$  in Euclidean domains by standard volumetric arguments, e.g., p.373 in [14]. The previous bound can be  $dx$ -integrated to give

$$\|P_{E_K}\Lambda - \lambda\|_{L^1} \leq c_{B,\mathbb{O}}h, \quad (15)$$

which controls the  $L^1$ -approximation error from the ‘bin count’ histogram estimator. Using slightly refined approximation theory (as in Ch.4.3 in [14]), the assumption of  $B$ -Lipschitzness can be weakened to  $\lambda$  having a bounded  $B_{1\infty}^1(\mathbb{O})$ -norm, if the latter Besov space is appropriately defined on  $\mathbb{O}$ .

**Theorem 2.4.** *In the setting of Theorem 2.3, suppose that  $\mathbb{O}$  is a bounded subset of  $\mathbb{R}^d$  and that the bins  $(B_i)_{i=1}^K$  have diameter bounded by a constant multiple of  $K^{-1/d}$ . Assume further that*

$$\Theta_n \subseteq \{ \text{the density } \lambda \text{ of } \Lambda \text{ exists and is } B\text{-Lipschitz} \} \quad (16)$$

for all  $n$  and some  $B > 0$ . Then if the density  $\lambda_0$  of the true intensity measure  $\Lambda_0$  exists and is  $B$ -Lipschitz on  $\mathbb{O}$ , we have as  $n \rightarrow \infty$  that

$$\Pi(\lambda : \|\lambda - \lambda_0\|_{L^1} > M(\sqrt{K/n} + K^{-1/d})|Y_1, \dots, Y_K) = O_{P_{\Lambda_0}}(e^{-kn\varepsilon_n^2}) \quad (17)$$

for some large enough constant  $M$  depending on  $B, d, \mathbb{O}, C, \Lambda_{max}$  and some  $k > 0$ .

*Proof.* The result follows from Theorem 2.3, (15) and

$$\|\lambda - \lambda_0\|_{L^1}^2 \leq \|\Lambda - \Lambda_0\|_{\ell_1}^2 + \|P_{E_K}(\Lambda - \Lambda_0) - (\lambda - \lambda_0)\|_{L^1}^2.$$

□

For high-dimensional Poisson regression problems, the  $L^1$ -structure is useful to deploy appropriate concentration of measure techniques in the proof of Theorem 6.1. When we know that the  $W^{\beta,1}$  norms of  $\lambda - \lambda_0$  are uniformly bounded, then we can interpolate (6) and obtain a contraction rate for  $\|\lambda - \lambda_0\|_{L^2}$  as well.

The approach based on histograms presented here is naturally confined to obtain bounds of the order  $O(K^{-1/d})$ , which is sufficient to prove consistency at ‘algebraic’ rates  $n^{-\beta}$  for some  $\beta > 0$ , but which prohibits to attain ‘fast’ convergence rates for intensity measures  $\Lambda$  whose densities are smoother than Lipschitz. But note that for Lipschitz densities  $\lambda$ , the rate of convergence obtained in (17) as well as in Theorem 6.1 is minimax optimal (as can be proved similarly to Ch.6.3 in [14], using Lemma 2.1 above and also Exercise 6.3.2 there). Note also that the assumption of disjoint support of the bins is crucial to retain independence of the  $K$  samples in (7), and obtaining faster rates will require rather different techniques based on general functionals of Poisson point processes that we do not wish to develop here.

### 3 Derivation of the observation model and proof of Theorem 1.1

In a variety of imaging tasks in biochemistry (e.g., [23, 22, 4]) one observes molecules diffusing in a ‘nucleus’ (a domain  $\Omega$  in  $\mathbb{R}^d$ ) that contains genetic material. Inside such nuclei, certain proteins (such as chromatin) may be concentrated in sub-compartments, called ‘foci’. Diffusing molecules can be seen to bind (‘get trapped’) at random times, to then stop diffusing (or to visibly diffuse much slower). The binding probability is higher in regions of high protein concentration, but diffusion is still visible inside such foci, and it is a scientifically relevant question whether diffusive behaviour of unbound molecules inside of foci and outside is different. The presence of binding events makes it difficult to discern such effects with high-resolution microscope measurements, and the main ideas of this article are concerned with the task of inferring the diffusivity  $D$  from such data. To do this, we proposed a Markovian ‘binding’ model in the introduction, where a motion of the molecule evolves according to the basic diffusion equation (1) and where binding is modelled by the random process  $S$  from (2), with a binding potential  $q$  that is given and which models the (observable) density of chromatin inside of  $\Omega$ . This captures the essence of the biological mechanism described above (but is still a simplification as it does not model the fact that molecules can also unbind, and may in fact not stop after binding but just move much slower).

The purpose of this subsection is to rigorously derive the Poisson process measurement model from the axioms sketched above, and to prove Theorem 1.1 under natural regularity Conditions 4.1 and 4.2 (to be introduced in the next subsection). We recall the killed process  $\tilde{X}_t$  from the introduction with ‘binding’ time  $S$  from (2). Some well known properties of this process are explained, e.g., in Section III.18 in [38]: On the enlarged state space  $\Omega \cup \{\dagger\}$  where all measurable functions  $f$  are extended simply by setting  $f(\dagger) = 0$ , one

shows that  $\tilde{X}_t$  is a Markov process with infinitesimal generator equal to the Schrödinger operator  $\mathcal{L}_{D,q}$  from (38). The law of  $\tilde{X}_t$  on  $\Omega$  models the ‘subprobability density’ of the surviving particles at time  $t$ , given by the solution  $v_{D,q}(t, \cdot)$  of the PDE (42). The first fact we prove is that when  $q$  is strictly positive on a set of positive Lebesgue measure, the survival time of a particle is finite almost surely.

**Proposition 3.1.** *Suppose  $q, D, \phi$  satisfy Conditions 4.1 and 4.2. Let  $S$  be the binding time from (2). Then  $P(S < \infty) = 1$ .*

*Proof.* For any  $m \in \mathbb{N}$  we use (44) and the Cauchy-Schwarz inequality to bound

$$P(S > m) \leq P(\tilde{X}_m \notin \{\dagger\}) = P(\tilde{X}_m \in \Omega) = \int_{\Omega} v_{D,q}(m, x) dx \leq |\Omega|^{1/2} \|v_{D,q}(m, \cdot)\|_{L^2} \lesssim e^{-cm}$$

from which we deduce that  $\sum_{m>1} P(S > m) < \infty$  and hence by the Borel-Cantelli lemma we have

$$P(\{S > m\} \text{ i.o.}) = 0,$$

so the result follows. □

In the killed process,  $\tilde{X}_S = \dagger$  is ‘not visible’ any longer at time  $S$ , whereas we observe the actual binding location  $X_S$  of the molecule. This is a well-defined random variable since  $S$  is a stopping time for the process  $(X_t)$  (e.g., Sections 3.3 and 22 in [3]). Since all particles will bind at some random point in time, intuitively, the distribution of their binding locations should equal the time-average  $\int_0^\infty v(t, \cdot) dt$  of the density of their positions, multiplied by the intensity  $q$  at the relevant position. We will now show that this intuition is rigorous and gives rise to Theorem 1.1.

### 3.1 Probability density of the binding location $\times$ time

Define a random variable  $Z$  taking values in  $\hat{\Omega} := \Omega \times [0, \infty]$  by

$$Z = (X_S, S)$$

such that  $Z$  jointly records the location and time of the binding event and is a measurable map for the Borel  $\sigma$ -algebra  $\mathcal{B}_{\hat{\Omega}} := \mathcal{B}_{\Omega} \otimes \mathcal{B}_{[0, \infty]}$ , where  $\mathcal{B}_{[0, \infty]} = \{B \cup \{\infty\} : B \in \mathcal{B}_{\mathbb{R}^+}\}$ . The law  $\Lambda_Z = \text{Law}(Z)$  can be characterised as follows on a generating family of ‘rectangles’ of measurable sets:

**Proposition 3.2.** *Suppose  $q, D, \phi$  satisfy Conditions 4.1 and 4.2. The probability measure  $\Lambda_Z$  has a density function such that for every open set  $A \subset \Omega$  and every interval  $(t_1, t_2], 0 < t_1 < t_2 < \infty$ ,*

$$P(Z \in A \times (t_1, t_2]) = \int_{t_1}^{t_2} \int_A \lambda(t, x) dx dt \tag{18}$$

where

$$\lambda(t, x) = q(x)v_{D,q}(t, x), \quad t > 0, x \in \Omega, \tag{19}$$

for  $v_{D,q}$  the solution of the parabolic Schrödinger equation (42) with initial condition  $\phi$ .

*Proof.* Note first that from the exponential decay estimate (44) and since  $\|q\|_\infty < \infty$ , we know that the density  $\lambda$  is Lebesgue integrable on  $[0, \infty) \times \Omega$ , hence the r.h.s. of (18) is well defined.

We start with the following key lemma.

**Lemma 3.3.** *For  $A$  any open subset of  $\Omega$  and any  $t \geq 0$*

$$P(Z \in A \times (t, t+h]) = h \int_A \lambda(t, x) dx + o(h) \quad \text{as } h \rightarrow 0. \tag{20}$$

*Proof.* By the memory-less property of the exponential random variable  $Y$ , if  $\tilde{X}_t$  binds at time  $S \in (t, t+h]$ , it has not bound before time  $t$  and we can restart the clock and the particle will bind at the first subsequent time  $t + \sigma$  for which  $\int_t^{t+\sigma} q(X_s)ds > Y$ . We also know that if  $\tilde{X}_s$  binds in  $A$  at time  $t + \sigma$ , the variables  $\tilde{X}_s$  and  $X_s$  co-incide for all  $s < t + \sigma$  (since the process  $\tilde{X}_s$  cannot return from the cemetery  $\{\dagger\}$ ), and by continuity of sample paths of the diffusion,

$$X_{t+\sigma} = \lim_{r \rightarrow t+\sigma} X_r = \lim_{r \rightarrow t+\sigma} \tilde{X}_r$$

must hold almost surely in this case.

Thus let us write the probability

$$\begin{aligned} P(Z \in A \times (t, t+h]) &= P\left(\int_t^{t+\sigma} q(X_s)ds > Y \text{ and } \lim_{r \rightarrow t+\sigma} \tilde{X}_r \in A \text{ for some } 0 < \sigma \leq h\right) \\ &= P(\cup_{0 < \sigma \leq h} B_\sigma \cap A_\sigma) \end{aligned} \quad (21)$$

where for  $y > 0$  we have defined measurable subsets

$$B_\sigma = \{w, y : \int_t^{t+\sigma} q(w(s))ds > y\} \quad \text{and} \quad A_\sigma = \{w : w(t+\sigma) \in A\} \times \mathbb{R}^+.$$

Here  $w = (w(s)) \in \Omega$  for  $0 < s \leq t + \sigma$ , are points in the space of possible continuous trajectories of  $\tilde{X}_s$  before binding, extended by continuity to  $s = t + \sigma$ . We prove that the last probability is both upper and lower bounded by  $h \int_A \lambda(t, z)dz$  plus an error that is  $o(h)$ . Let us write  $P^W$  for the law of the process  $(\tilde{X}_s)$  in what follows and  $P^Y$  for the law of  $Y$ .

**Lower bound** We clearly have  $B_h \cap A_h \subset \cup_{0 < \sigma \leq h} B_\sigma \cap A_\sigma$  and therefore the probability in (21) is greater than or equal to  $P(B_h \cap A_h)$ . Also by the remark from earlier,  $B_h \cap A_h \subset C([0, t+h] \rightarrow \Omega) \times \mathbb{R}^+$ , and so by continuity of  $w$  and since  $A$  is open we must have  $1_{A_h} \rightarrow 1_{\{w(t) \in A\}}$  as  $h \rightarrow 0$  for every fixed  $w$ . From the dominated convergence theorem and since the density of the distribution of  $\tilde{X}_t$  is the solution  $v_{D,q}(t)$  at time  $t$  of (42), we see

$$\int \mathbb{1}_{A_h}(w)q(w(t))dP^W(w) \xrightarrow{h \rightarrow 0} \int \mathbb{1}_A(w(t))q(w(t))dP^W(w) = \int_A q(z)v_{D,q}(t, z)dz.$$

Using this, boundedness and continuity of  $q$  combined with the fundamental theorem of calculus and again the dominated convergence theorem we have

$$\begin{aligned} &\int \int_{\mathbb{R}} \mathbb{1}_{A_h}(w, \cdot) \mathbb{1}_{B_h}(w, y) dP^Y(y) dP^W(w) = \int \mathbb{1}_{A_h}(w, \cdot) P^Y\left(\int_t^{t+h} q(w(s))ds > Y\right) dP^W(w) \\ &= \int \mathbb{1}_{A_h}(w, \cdot) (1 - e^{-\int_t^{t+h} q(w(s))ds}) dP^W(w) \\ &= h \int \mathbb{1}_{A_h}(w, \cdot) \frac{1}{h} \int_t^{t+h} q(w(s))ds dP^W(w) + o(h) \\ &= h \int \mathbb{1}_{A_h}(w, \cdot) q(w(t)) dP^W(w) + o(h) = h \int_A q(z)v_{D,q}(t, z)dz + o(h) \\ &= h \int_A \lambda(t, z)dz + o(h), \end{aligned}$$

which gives the desired lower bound.

**Upper bound** Since  $q(\dagger) = 0 \leq q$ , we can regard  $B_\sigma \subset B_h$  and note that

$$P^Y\left(\int_t^{t+h} q(w(s))ds > Y\right) \leq hq(w(t)) + o(h) \quad (22)$$

follows just as above. Hence the probability in question is upper bounded by

$$h \int \mathbb{1}_{\cup_{0 \leq \sigma \leq h} A_\sigma}(w, \cdot) q(w(t)) dP^W(w) + o(h).$$

Now letting  $h = 1/n$  for  $n \in \mathbb{N}$ , we see that

$$\bigcup_{0 \leq \sigma \leq 1/n} A_\sigma \downarrow \bigcap_n \bigcup_{0 \leq \sigma \leq 1/n} A_\sigma = \{w(t + 1/n) \in A \text{ i.o.}\} \times \mathbb{R}_+ \equiv \bar{A} \times \mathbb{R}_+.$$

Using that  $q \geq 0$  and the monotone convergence theorem we can take limits to see that the penultimate expression equals

$$h \int \mathbb{1}_{\bar{A}}(w) q(w(t)) dP^W(w) + o(h) = h \int \mathbb{1}_A(w(t)) q(w(t)) dP^W(w) + o(h)$$

where in the last equality we used that by continuity of  $w$  and for any open set  $A$  we must have

$$\{w(t + 1/n) \in A \text{ i.o.}\} = \{w(t) \in A\}.$$

The lemma is proved.  $\square$

Now fix any open set  $A$  in the domain  $\Omega$  and define  $S_A$  as the binding time in  $A$ , that is  $S_A = S$  if  $X_S \in A$  and  $S_A = \infty$  otherwise. Write  $F_{S_A}(t) = P(S_A \leq t)$ ,  $t > 0$ , for its distribution function and let  $t, h$  be arbitrary. Then as in (21) and above (22), using also  $1 - e^{-x} \leq x$  for all  $x$ , we can estimate

$$\begin{aligned} F_{S_A}(t+h) - F_{S_A}(t) &= P(Z \in A \times (t, t+h]) \leq P(B_h) \\ &= \int \int \mathbb{1}_{B_h}(w, y) dP^Y(y) dP^W(w) \\ &= \int (1 - e^{-\int_t^{t+h} q(w(s)) ds}) dP^W(w) \leq h \|q\|_\infty. \end{aligned}$$

For any finite set of disjoint intervals  $(a_1, b_1), \dots, (a_N, b_N)$  lying in  $[t_1, t_2]$ , we see that

$$\sum_{j=1}^N (b_j - a_j) < \varepsilon / \|q\|_\infty \implies \sum_{j=1}^N |F_{S_A}(b_j) - F_{S_A}(a_j)| < \|q\|_\infty \sum_{j=1}^N (b_j - a_j) < \varepsilon,$$

so the map  $t \mapsto F_{S_A}(t)$  is absolutely continuous on  $[t_1, t_2]$ . By Theorem 3.35 in [11],  $F_{S_A}$  therefore admits an integrable density  $f_{S_A}$  with respect to the Lebesgue measure such that  $P(Z \in A \times \tau) = \int_\tau f_{S_A}(s) ds$  for every subinterval  $\tau \subseteq [t_1, t_2]$  and  $A$  fixed. Moreover, by Lebesgue's differentiation theorem and Lemma 3.3,

$$\begin{aligned} f_{S_A}(t) - \int_A \lambda(t, x) dx &= \lim_{h \rightarrow 0} \frac{1}{h} \left( \int_t^{t+h} f_{S_A}(s) ds - h \int_A \lambda(t, x) dx \right) \\ &= \lim_{h \rightarrow 0} \frac{1}{h} (P(Z \in ((t, t+h], A)) - h \int_A \lambda(t, x) dx) = 0 \end{aligned}$$

for almost every  $t \in (t_1, t_2]$ , so that  $f_{S_A}$  and  $\int_A \lambda(\cdot, x) dx$  coincide almost everywhere. The proposition is proved.  $\square$

We can extend Proposition 3.2 to the  $\sigma$ -field  $\mathcal{B}_{\hat{\Omega}} = \mathcal{B}_\Omega \otimes \mathcal{B}_{[0, \infty]}$ . Denote by  $\Lambda$  the finite measure induced by the application

$$(B, \tau) \mapsto \Lambda(B \times \tau) = \int_B \int_\tau \lambda(t, x) dt dx, \quad (B, \tau) \in \mathcal{B}_\Omega \otimes \mathcal{B}_{\mathbb{R}^+}.$$

By Proposition 3.2 the law  $\Lambda_Z$  of  $Z$  and  $\Lambda$  coincide on the 'rectangles'  $A \times \tau$  where  $\tau = (t_1, t_2]$  and  $A$  is open, and therefore, by standard uniqueness arguments (e.g., Theorem 1.14 in [11]) they coincide on the

Borel  $\sigma$ -field  $\mathcal{B}_\Omega \otimes \mathcal{B}_{\mathbb{R}^+}$ . We can extend  $\Lambda$  to  $\mathcal{B}_{\hat{\Omega}}$  by setting  $\Lambda(B \times \tau \cup \{\infty\}) = \Lambda(B \times \tau)$ , in particular  $\Lambda(\Omega \times \{\infty\}) = 0$ , which is consistent with

$$\Lambda(\Omega \times \mathbb{R}^+) = \lim_{T \rightarrow \infty} \Lambda(\Omega \times [0, T]) = \lim_{T \rightarrow \infty} \Lambda_Z(\Omega \times [0, T]) = \Lambda_Z(\Omega \times \mathbb{R}^+) = P(S < \infty) = 1$$

in view of Proposition 3.1 and countable additivity of measures. In particular  $\lambda$  is a probability density on  $\Omega \times [0, \infty)$ , that is

$$\int_{\Omega} q(x) \int_{\mathbb{R}^+} v_{D,q}(t, x) dt dx = \Lambda(\Omega \times \mathbb{R}^+) = 1. \quad (23)$$

If we record only the binding position (without reporting the time of binding), then the probability to bind in a particular measurable subset  $A$  of  $\Omega$ , at some time, is given by

$$P(\tilde{X}_S \in A, S < \infty) = \Lambda_Z(A \times \mathbb{R}^+) = \int_A \int_0^\infty q(x) v_{D,q}(t, x) dt dx = \int_A q(x) u_{D,q}(x) dx \equiv \Lambda_D(A) \quad (24)$$

where the second identity follows from (45), and where  $u_{D,q}$  is the solution to the steady state Schrödinger equation

$$\begin{aligned} \nabla \cdot (D \nabla u) - qu &= -\phi \quad \text{on } \Omega \\ \frac{\partial u}{\partial \nu} &= 0 \quad \text{on } \partial\Omega, \end{aligned} \quad (25)$$

with  $\phi$  modelling the initial condition of the system.

### 3.2 Construction of the Poisson point process

The measure  $\Lambda_D$  from (24) will serve as the intensity measure for the point process we construct now. We take independently diffusing molecules  $\tilde{X}_s^i, i = 1, \dots$ , and observe their terminal positions  $X_S^i$ . We then record the count of binding locations through the number  $N(A)$  of molecules having bound in a measurable subset  $A$  of  $\Omega$ . Concretely, we take independent and identically distributed random variables  $T_1, T_2, \dots$ ,

$$T_i(A) = \begin{cases} 1, & \text{if molecule } i \text{ has bound in } A \\ 0, & \text{otherwise} \end{cases}$$

such that  $P(T_i(A) = 1) = \Lambda_D(A)$ , and obtain a mixed binomial process

$$N(A) = \sum_{i=1}^{n_{mol}} T_i(A), \quad A \text{ measurable,}$$

where  $n_{mol} \in \mathbb{N}$  is the total number of diffusing molecules. If we assume that the binding positions in disjoint subsets of  $\Omega$  are statistically independent, then such a ‘completely random’ point process is necessarily a Poisson processes, see Theorem 6.12 in [26]. The preceding mixed binomial process is a Poisson process precisely when  $n_{mol}$  is drawn from a Poisson distribution (see Propositions 3.5 and 3.8 in [26]). To model an effective sample size of  $n$ , we therefore assume  $n_{mol} \sim \text{Poisson}(n)$  for some  $n \in \mathbb{N}$ . One then shows as in Section 3.2 of [26] that  $N$  is indeed a Poisson point process with intensity  $n\lambda_D = nqu_{D,q}$  on  $\Omega$ . The expected number  $\mathbb{E}[N(A)]$  of bound molecules in every Borel set  $A$  of  $\Omega$  is precisely given by Campbell’s formula (see Prop. 2.7 in [26]):

$$\mathbb{E}[N(A)] = \int_A nq(x) u_{D,q}(x) dx. \quad (26)$$

## 4 Inference on the diffusion parameter from killed diffusion

We formulate appropriate hypotheses on  $D, q, \phi$  that allow to combine the results from the previous sections. Since boundary reflection dominates the diffusive dynamics of (1) near  $\partial\Omega$ , it is convenient to consider  $q = 0$

near  $\partial\Omega$  so that binding events occur only in the interior of  $\Omega$ . Along with that we assume that  $D$  is known to be constant ( $= 1/2$ , say) near  $\partial\Omega$  and bounded away from zero (by  $1/4$ , for notational convenience). We also assume that the initial condition is strictly positive and satisfies a basic smoothness and boundary compatibility condition.

**Condition 4.1.** *The potential  $q$  lies in  $H^\eta(\Omega)$ ,  $\eta > 1 + d/2$ , with  $\|q\|_\infty \leq Q$  for some  $Q > 0$ . Suppose further that  $\inf_{x \in \Omega_{00}} q(x) \geq q_{\min} > 0$  on some open set  $\Omega_{00} \subset \Omega$  such that  $\text{dist}(\Omega_{00}, \partial\Omega) > 0$  and that  $q = 0$  outside a compact subset  $\Omega_0 \subset \Omega$  that contains  $\Omega_{00} \subset \Omega_0$ .*

**Condition 4.2.** *The diffusivity  $D$  lies in  $H^\alpha(\Omega)$  for some  $\alpha > 2 + d/2$ , equals  $1/2$  outside of  $\Omega_{00}$  and satisfies  $\inf_{x \in \Omega} D(x) > 1/4$ . Moreover  $\phi$  is a probability density that satisfies  $\inf_{x \in \Omega} \phi(x) \geq \phi_{\min}$  for some  $\phi_{\min} > 0$  and  $\phi$  either i) lies in  $H^b(\Omega)$  for some  $b$ ,  $d/2 < b \leq \min(\alpha + 1, \eta + 2)$ , and is constant outside of  $\Omega_{00}$  or ii) lies in  $H^2_\nu(\Omega)$  from (39) below, in which case we set  $b = 2, d < 4$ , in what follows.*

The hypotheses ensure that  $\phi$  belongs to the spectrally defined Sobolev space  $\tilde{H}^b_{D,q}$  from (41) below for any such  $D, q$  and  $b$  in view of Proposition 5.2 (for appropriate  $\alpha, \eta$ ). Lemma 5.4, (46) and the Sobolev imbedding then imply that the unique solution  $u_{D,q}$  to (25) is non-negative and lies in  $\tilde{H}^{b+2} \subset H^{b+2} \subset C^2$ . Since also  $0 \leq q \in H^\eta \subset C^1$  this ensures that the density

$$\frac{d\Lambda_D}{dx} = \lambda_D = qu_{D,q} \quad (27)$$

is Lipschitz and non-negative on  $\Omega$ . We can therefore consider the Poisson point process  $N$  over  $\Omega$  constructed in Section 3.2 with intensity measure  $n\Lambda_D$  where  $\Lambda_D$  is the (probability) measure with Lebesgue density  $\lambda_D$ . From what precedes  $N$  will generate samples lying in  $\Omega_0$  almost surely. Considering that there are on average  $n$  particles diffusing, and in light of (17), we then consider

$$K = K_{n,d} \simeq n^{d/(2+d)} \quad (28)$$

many bins  $B_i$  that partition  $\Omega_{00}$  in a way such that each bin has Euclidean diameter bounded by a multiple of  $K^{-1/d}$ . We measure the count of the killed particles in each bin:

$$Y_i := N(B_i) \quad 1 \leq i \leq K, \quad (29)$$

which for these choices of  $K, \Lambda = \Lambda_D$  and  $\mathbb{O} = \Omega_{00}$  fits into the general measurement setting (7). We denote the resulting law  $P_{\Lambda_D}$  of the observation vector by  $P_D$  in this section.

## 4.1 Construction of the prior and posterior

Our goal is to make inference on  $D$  and hence we construct a prior directly for this parameter which induces a pushforward prior for the intensity measure  $\Lambda_D$  via (27). We start with a Gaussian random field  $(w(x) : x \in \Omega)$  that induces a probability measure  $\text{Law}(w)$  on the (separable) Banach space  $H^a(\Omega) \subset C^1(\Omega)$  with reproducing kernel Hilbert space (RKHS) equal to  $H^a(\Omega)$ , where

$$\alpha > 2 + d, \alpha - d/2 > a > 2 + d/2.$$

For the existence of such a process see Theorem B.1.3 in [32]. We then define the rescaled random function

$$W(x) = \frac{1}{n^{d/(4\alpha+2d)}} \zeta(x)w(x), \quad x \in \Omega, \quad (30)$$

where  $\zeta$  is a smooth cutoff function that is zero outside of  $\Omega_0$  and identically  $\zeta = 1$  on  $\Omega_{00}$ . The prior probability measure for  $D$  is then the law  $\Pi = \Pi_n = \text{Law}(D)$  on  $H^a$  of the random function given by

$$D(x) = \frac{1}{4} + \frac{1}{4}e^{W(x)}, \quad x \in \Omega, \quad (31)$$

which equals  $D = 1/2$  near the boundary  $\partial\Omega$ , specifically on  $\Omega \setminus \Omega_0$ . For given  $q$  this induces a prior probability measure  $\Pi_\Lambda$  supported in  $H^a$  for the intensity measures  $\Lambda_D$  from (27) simply via the image

measure theorem (and since the map  $D \mapsto u_{D,q}$  is appropriately measurable in view of Lemmata 4.3 and 5.4 below). This prior and then also the resulting posterior measures concentrate precisely on the non-linear manifold of intensity functions that arise from observing a killed diffusion process for some  $D$  (and  $q$ ), which will be crucial later to solve the non-linear inverse problem via Theorem 5.6. Note that the posterior measure from (12) based on data (29) is then given by

$$\Pi(A|Y_1, \dots, Y_K) = \frac{\int_A p_{\Lambda_{D,n}}^K(Y_1, \dots, Y_K) d\Pi(D)}{\int_{H^a} p_{\Lambda_{D,n}}^K(Y_1, \dots, Y_K) d\Pi(D)}, \quad A \text{ a Borel set in } H^a(\Omega). \quad (32)$$

Despite the lack of convexity of the negative loglikelihood function  $D \mapsto -\log p_{\Lambda_{D,n}}^K$ , following ideas in [39], approximate computation of this posterior measure is possible by MCMC methods, with each MCMC iterate requiring the numerical evaluation of  $p_{\Lambda_{D,n}}^K$  and hence of the elliptic PDE (25). For instance one can use the pCN, ULA or MALA algorithm, see [7] and also Sec.1.2.4 in [32] as well as references therein. Computational guarantees for such algorithms can be given following ideas in [37], see also Ch.5 in [32]. In concrete applications one may wish to model a further drift  $\nabla U(X_t)dt$  in (1), which corresponds to subtracting a further vector field  $\nabla U \cdot \nabla u$  on the l.h.s. in (25) and hence changes little in the implementation if  $U$  is known or determined independently – see also the introduction of [33] for further discussion of this aspect.

## 4.2 Numerical illustration

To illustrate the theory, we perform numerical experiments where  $\Omega$  is the unit disk ( $d = 2$ ) represented by a triangular mesh with 6161 nodes. We take  $n = 10^7$  as the expected number of molecules, and  $K = 36$  the number of bins. We select parameters  $D, q$  such that  $q$  equals 200 on a small circular ‘focus’, and 100 outside (with a smooth sigmoid transition in between), and such that in the link function (31) we replace the additive constant  $1/4$  by 15. These values were chosen for convenience to amplify the effects of both diffusion and binding in the graphical representation of our results – a more comprehensive numerical investigation of the algorithm proposed here will be undertaken in future work.




Figure 2: Problem setting

**Synthetic data** We take the initial condition  $\phi$  and ground truth  $D_0$  such that  $\phi(x, y) \propto \exp(\cos(3\pi(x^2 + y^2)))$ ,  $W_0(x, y) = 5e^{-10x^2 - 10(y-0.4)^2}$  for  $(x, y)$  in the interior of the mesh and zero elsewhere. The corresponding solution  $u_{D_0, q}$ , is computed on the mesh nodes via FEM, and can be extended to the whole domain via standard interpolation, as needed to compute the integrals  $\Lambda_{D_0}(B_i) = \int_{B_i} q u_{D_0, q}$ ,  $i = 1, \dots, K$ . We then draw the random observation count  $Y_i = N(B_i)$  for each bin as  $Poisson(n\Lambda_{D_0}(B_i))$ . For the sake of speed, a basic trapezoidal method on a fine grid ( $\Delta x = \Delta y = 0.01$ ) is used to compute the integrals over the bins.

**MCMC** Following a standard implementation from [16], we run the preconditioned Crank-Nicolson (pCN) scheme for 15 000 steps, on the posterior density (32) arising from a Matérn Gaussian process prior of smoothness parameter  $\alpha = 2.5$  and length scale  $\ell = 0.15$ . At each step, the function  $W$  is represented by the 6161-dimensional vector of its values on the mesh nodes, and we again use a standard interpolation algorithm to define it on the whole domain when solving the PDE. We initialise the chain uniformly at 0. After running the chain, we thin it by retaining every 200<sup>th</sup> sample after removal of the burn-in (initial 5 000 iterations) - which we then use to compute the posterior mean  $\bar{W}$  giving rise to  $\bar{D} = 15 + \frac{1}{4}e^{\bar{W}}$ .




Figure 3: Posterior mean estimate  $\bar{D}$  for the diffusivity function (left), along with the corresponding PDE solution  $u_{\bar{D}, q}$  (centre), and the true solution  $u_{D_0, q}$  (right) for comparison.




Figure 4: The second and third displays compare, respectively, the ‘fit’ of the histogram bin count  $N(B_i)$  and the estimated posterior intensities  $n\Lambda_{\bar{D}}(B_i)$  with the ground truth  $n\Lambda_{D_0}(B_i)$ . The numerically improved recovery also of the intensity  $\Lambda_{D_0}$  by the Bayesian method when compared to the naive bin count is clearly visible in these images.

### 4.3 Posterior consistency theorems

We now adapt techniques from Bayesian non-linear inverse problems [30, 32, 33] to the present situation to derive some posterior consistency results.

While the probabilistic structure of the prior for  $D$  is well understood via Gaussian process theory, the prior for the intensity measures  $\Lambda_D$  is more complex as it involves the non-linear PDE solution map  $D \mapsto u_{D, q}$ . To verify the hypotheses of Theorem 2.3 we will require the following lemma.

**Lemma 4.3.** *Suppose  $D, q, \phi$  satisfy Conditions 4.1 and 4.2. Let the intensity measure  $\Lambda_D$  be given as in (27) with Lebesgue density  $\lambda_D$ . Then we have*

$$\inf_{x \in \Omega_{00}} \lambda_D(x) \geq \frac{q_{\min} \phi_{\min}}{Q} > 0.$$

Next consider two diffusivities  $D_1, D_2 \geq 1/4$  s.t.  $D_1 = D_2$  on  $\Omega \setminus \Omega_0$  and  $\|D_i\|_{H^a} \leq B$  for some  $B > 0$  and  $a > 2 + d/2$ . Then we have

$$\|\lambda_{D_1} - \lambda_{D_2}\|_{L^2} \leq c_{B, q_{\min}, Q} \|D_1 - D_2\|_{L^2}$$

and for every  $a' > 1 + d/2$  that

$$\|\lambda_{D_1} - \lambda_{D_2}\|_{\infty} \leq c_{B, Q} \|D_1 - D_2\|_{H^{a'}}$$

for finite positive constants  $c_{B, q_{\min}, Q}, c_{B, Q}$ .

*Proof.* We use PDE techniques and notation from Sec. 5.1. The first claim is Lemma 5.3a) with  $\varphi = \phi$ . Next, suppose  $u_{D_i}, i = 1, 2$ , solve the elliptic equation (25) for two distinct diffusivities  $D_i$  s.t.  $\|D_i\|_{H^a} \leq b$ . Set  $\bar{u} = u_{D_1} - u_{D_2}$  and notice that

$$\|u_{D_i}\|_{C^2} \lesssim \|u_{D_i}\|_{\tilde{H}^{b+2}} \leq c_B < \infty$$

in view of Lemma 5.4 for appropriate  $b > d/2$ , Proposition 5.2, the Sobolev imbedding, and our hypotheses which ensure  $\phi \in \tilde{H}_{D_i, q}^b$ . Then we have

$$\mathcal{L}_{D_1, q}(\bar{u}) = -\phi - \mathcal{L}_{D_2, q} u_{D_2} + \nabla \cdot (D_2 - D_1) \nabla u_{D_2} = \nabla \cdot (D_2 - D_1) \nabla u_{D_2}$$

so that  $\bar{u}$  equals  $\mathcal{L}_{D_1, q}^{-1}[\varphi]$  with  $\varphi = \nabla \cdot (D_2 - D_1) \nabla u_{D_2}$  which lies in  $\tilde{H}_{D_1, q}^b, b > d/2$ , by what precedes, Proposition 5.2, and since  $D_1 - D_2$  vanishes near  $\partial\Omega$ . We thus obtain from the second inequality in Part a) of Lemma 5.3 that

$$\|u_{D_1} - u_{D_2}\|_{\infty} \leq \|\nabla \cdot ((D_1 - D_2) \nabla u_{D_2})\|_{\infty} \lesssim \|D_1 - D_2\|_{C^1} \leq \|D_1 - D_2\|_{H^{a'}} \quad (33)$$

which implies the desired  $L^\infty$ -estimate since  $\lambda_D = qu_{D, q}$  for all  $D$ . For the  $L^2 - L^2$ -estimate, we notice that  $D_1 - D_2$  has compact support in  $\Omega$  and so we see from the divergence theorem, again Lemma 5.3 combined with the previous bound for  $\|u_{D_2}\|_{C^1}$  and the Cauchy-Schwarz inequality, that

$$\begin{aligned} \|u_{D_1} - u_{D_2}\|_{L^2} &\lesssim \|\nabla \cdot ((D_1 - D_2) \nabla u_{D_2})\|_{\tilde{H}^{-1}} \\ &= \sup_{\|\psi\|_{\tilde{H}^1} \leq 1} \left| \int_{\Omega} (D_1 - D_2) \nabla \psi \cdot \nabla u_{D_2} \right| \lesssim \|D_1 - D_2\|_{L^2} \end{aligned}$$

completing the proof. □

We now provide an appropriate result for the Gaussian small deviation asymptotics of the base prior.

**Lemma 4.4.** *Consider the Gaussian process  $W$  from (30) with  $\alpha > 2 + d$  and let  $W_0 \in H^\alpha$  be such that  $W_0 = 0$  outside of  $\Omega_{00}$ . Let  $\bar{\varepsilon}_n = n^{-\alpha/(2\alpha+d)}$ . Then for every fixed  $\epsilon > 0, a < \alpha - d/2$ , there exists a positive constant  $C = C(\epsilon, \alpha, a, \Omega, \zeta, \Omega_{00}) > 0$  such that for all  $n$ ,*

$$\Pr(W : \|W - W_0\|_{L^2} < \bar{\varepsilon}_n, \|W - W_0\|_{H^a} < \epsilon) \geq e^{-Cn\bar{\varepsilon}_n^2}. \quad (34)$$

*Proof.* Note that the rescaling factor in (30) equals  $1/\sqrt{n\bar{\varepsilon}_n}$ . Since  $\zeta^{-1}$  is smooth on the support of  $W_0$ , using also Exercise 2.6.5 in [14] and (4), the RKHS-norm of the ‘shift’  $W_0$  can be bounded as

$$\sqrt{n\bar{\varepsilon}_n} \|W_0/\zeta\|_{H^\alpha} \lesssim \sqrt{n\bar{\varepsilon}_n} \|W_0\|_{H^\alpha}.$$

Therefore, using Corollary 2.6.18 in [14], the Gaussian correlation inequality (e.g., Theorem B.1.2 in [32]), and again (4), we can lower bound the probability in question by

$$e^{-n\bar{\varepsilon}_n^2 \|W_0\|_{H^\alpha}^2} \Pr(w : \|w\|_{L^2} < \sqrt{n}\bar{\varepsilon}_n / \|\zeta\|_\infty) \Pr(\|w\|_{H^a} < \epsilon \sqrt{n}\bar{\varepsilon}_n / \|\zeta\|_{C^a}).$$

Since the law  $\text{Law}(w)$  is supported on the separable Banach space  $H^a$ , the second probability is bounded from below by a positive constant independent of  $n$  in view of Anderson's lemma (Theorem 2.4.5 in [14]) and for  $\sqrt{n}\bar{\varepsilon}_n \geq 1$  (in fact the probability converges to one as  $n \rightarrow \infty$ ). For the first probability, the required lower bound follows by the same computation as the one on p.34 in [32], so that the result follows.  $\square$

Collecting these facts we can now show:

**Theorem 4.5.** *Suppose that  $q, D_0, \phi$  satisfy Conditions 4.1 and 4.2 with  $\alpha > 2 + d$ , respectively. The prior  $\Pi_\Lambda$  for the intensities  $\Lambda_D$  from (27) as constructed after (31) satisfies conditions (13) and (14) with  $\Lambda_{D_0}, \Lambda_{max} = 1, \varepsilon_n = \bar{k}n^{-\alpha/(2\alpha+d)}$  for some  $\bar{k} > 0$ , and with regularisation set*

$$\Theta_n = \{D : \|D\|_{H^a} \leq B\}, \quad (35)$$

for any  $B$  large enough and every  $a < \alpha - d/2$ . As a consequence we obtain, for all  $M$  large enough, some  $k > 0$ , and  $K = K_n$  as in (28) that

$$\Pi\left(D \in \Theta_n : \|\lambda_D - \lambda_{D_0}\|_{L^1(\Omega_{00})} \leq Mn^{-1/(2+d)} |Y_1, \dots, Y_K\right) = 1 - O_{P_{D_0}}(e^{-kn\varepsilon_n^2}) \quad (36)$$

*Proof.* To the ground truth  $D_0$  corresponds the potential  $W_0 = \log(4D_0 - 1)$  which lies in  $H^\alpha$  by the chain rule and since  $D_0 > 1/4$  does, and which equals  $W_0 = \log 1 = 0$  outside of  $\Omega_{00}$ , hence satisfies the hypotheses of Lemma 4.4. To verify (13), we work on the event in (34), so that in particular  $\|W\|_{H^a} + \|W_0\|_{H^a}$  is bounded by a fixed constant, which also bounds the  $\|\cdot\|_\infty$ -norms by the Sobolev imbedding with  $a > d/2$ . By Lemma 4.3 and (27), the true intensity function  $\lambda_{D_0}$  is uniformly lower bounded on  $\Omega_{00}$  and hence on any  $B_i$ , by  $\Lambda_{min} = q_{min}\phi_{min}/Q > 0$ . We then have from Jensen's inequality

$$\begin{aligned} \sum_{i=1}^K \frac{|\Lambda_D(B_i) - \Lambda_{D_0}(B_i)|^2}{\Lambda_{D_0}(B_i)} &\leq \Lambda_{min}^{-1} \sum_{i=1}^K \frac{1}{|B_i|} \left( \int_{B_i} (\lambda_D - \lambda_{D_0})(x) dx \right)^2 \\ &= \Lambda_{min}^{-1} \sum_{i=1}^K \left( \int_{B_i} (\lambda_D - \lambda_{D_0}) \frac{dx}{|B_i|} \right)^2 |B_i| \\ &\leq \Lambda_{min}^{-1} \sum_{i=1}^K \int_{B_i} (\lambda_D - \lambda_{D_0})^2 dx \\ &= \Lambda_{min}^{-1} \|\lambda_D - \lambda_{D_0}\|_{L^2(\Omega_{00})}^2 \leq C_0 \|D - D_0\|_{L^2}^2 \leq C_1 \|W - W_0\|_{L^2}^2, \end{aligned}$$

using also Lemma 4.3 in the last line, and for a fixed positive constant  $C_1 > 0$ . Similarly from Lemma 4.3,

$$\begin{aligned} \frac{|\Lambda_D(B_i) - \Lambda_{D_0}(B_i)|}{\Lambda_{D_0}(B_i)} &\leq \Lambda_{min}^{-1} \frac{1}{|B_i|} \left| \int_{B_i} (\lambda_D - \lambda_{D_0})(x) dx \right| \leq \Lambda_{min}^{-1} \|\lambda_D - \lambda_{D_0}\|_\infty \\ &\leq c \|D - D_0\|_{H^{a'}} \leq c' \|W - W_0\|_{H^{a'}} \end{aligned}$$

for some  $c' > 0$ . The condition (13) then follows from Lemma 4.4 with  $\varepsilon_n$  a constant multiple of  $\bar{\varepsilon}_n$  as well as  $a' = a$  and  $\epsilon > 0$  small enough. The condition (14) is verified for the rescaled prior and our  $\Theta_n$  just as on p.33 in [32]. We also note the uniform upper bound on  $\Lambda(\Omega_{00}) \leq 1 = \Lambda_{max}$  in view (23), (24). Therefore, noting also that our choices of  $K_n, \varepsilon_n$  are as required, we can invoke Theorem 2.3 to obtain the contraction rate  $O(\sqrt{K/n})$  for the  $\|\Lambda_D - \Lambda_{D_0}\|_{\ell_1}$  distance over the  $K$  bins  $B_i$  partitioning  $\Omega_{00} = \emptyset$ . Using again Lemma 5.4, Conditions 4.1, 4.2, and the Sobolev imbedding implies that  $\lambda_D = qu_D$  is uniformly bounded in  $\tilde{H}^{2+b} \subset C^1(\Omega)$  for  $D \in \Theta_n \cup \{D_0\}$ . This implies that these densities admit a uniform Lipschitz constant  $B$  and so the final conclusion for the  $L^1$ -norms follows from (17).  $\square$

The last step is to take advantage of the stability estimate Theorem 5.6 to show that the posterior inference for  $D$  is also valid, which in particular implies the proof of Theorem 1.2. To focus on the main statistical ideas we show this for ‘small potentials’  $q$  and  $d \leq 3$ , see Section 5.2 for discussion.

**Theorem 4.6.** *In the setting of Theorem 4.5, assume further that  $\phi$  satisfies the identifiability Condition 5.5, that  $d \leq 3$  and that  $\|q\|_{H^2} < \delta$  for  $\delta$  small enough as in Theorem 5.6. Then if  $K = K_n$  is as in (28) we have*

$$\Pi(D : \|D - D_0\|_{L^2} \leq Mn^{-\beta} | Y_1, \dots, Y_K) = 1 - O_{P_{D_0}}(e^{-kn\varepsilon_n^2}) \quad (37)$$

for some  $k > 0$  and  $\beta = \frac{1}{2+d} \frac{b+2-d/2}{b+2}$ . If  $\bar{D} = (1 + e^{\bar{W}})/4$  for posterior mean  $\bar{W} = E^\Pi[W | Y_1, \dots, Y_K]$ , then we also have

$$\|\bar{D} - D_0\|_{L^2} = O_{P_{D_0}}(n^{-\beta}).$$

*Proof.* For  $D \in \Theta_n$  from (35) and  $D_0 \in H^\alpha$  we know from Lemma 5.4 and our hypothesis on  $\phi$  that  $u_{D,q} - u_{D_0,q}$  is bounded in  $\tilde{H}^{b+2} \subset H^{b+2} \subset W^{b+2,1}$ . We then combine (36) with the stability estimate Theorem 5.6 for  $D_1 = D_0, D_2 = D$  and the interpolation inequality (6) to obtain

$$\begin{aligned} \|W - W_0\|_{L^2} &\lesssim \|D - D_0\|_{L^2} \lesssim \|u_{D,q} - u_{D_0,q}\|_{H^2(\Omega_{00})} \\ &\lesssim \|u_{D,q} - u_{D_0,q}\|_{L^1(\Omega_{00})}^{c(b,d)} \lesssim \|\lambda_D - \lambda_{D_0}\|_{L^1(\Omega_{00})}^{c(b,d)} \\ &\lesssim n^{-\frac{1}{2+d} \frac{b+2-d/2}{b+2}} \end{aligned}$$

from which the first claim of the theorem follows. The convergence of the posterior mean can then be proved as in Theorem 2.3.2 in [32], and is left to the reader.  $\square$

## 5 Analytical results from PDE theory

### 5.1 Schrödinger operators and their spectrum

We now recall some analytical results for Schrödinger type equations that will allow to derive some estimates for solutions of the PDE (25) that are essential in our proofs. Define the Schrödinger operator

$$\mathcal{L}_{D,q}v \equiv \nabla \cdot (D\nabla v) - qv \quad (38)$$

acting on  $v$  belonging to the usual Sobolev space

$$H_\nu^2 = \left\{ u \in H^2(\Omega) : \frac{\partial u}{\partial \nu} = 0 \text{ on } \partial\Omega \right\} \quad (39)$$

equipped with Neumann-boundary conditions. In this subsection we generally assume Conditions 4.1 and 4.2 even though weaker assumptions would be possible too. Adapting standard arguments from spectral theory for the Laplace equation with Neumann boundary conditions in Ch.5.7 in [41] to the presence of  $D, q$  (with  $D = 1/2, q = 0$  near  $\partial\Omega$ ), one shows that the Schrödinger operator  $\mathcal{L}_{D,q}$  has  $L^2(\Omega)$ -orthonormal eigen-pairs

$$(e_{j,D,q}, -\lambda_{j,D,q}) \in H_\nu^2(\Omega) \times [0, \infty), \quad j \geq 1,$$

with eigenvalues satisfying

$$\lambda_{j,D,q} \leq \lambda_{j+1,D,q} \simeq j^{2/d}, \quad (40)$$

where the constants implicit in  $\simeq$  depend on upper bounds for  $\|q\|_\infty, \|D\|_\infty$ . We now prove the following key Poincaré-type inequality.

**Lemma 5.1.** *Suppose  $q \geq q_{min} > 0$  on some subset  $\Omega_{00}$  of  $\Omega$  of positive Lebesgue measure  $|\Omega_{00}| > 0$ . Then we have the spectral gap inequality*

$$\lambda_{1,D,q} \geq c(\Omega, |\Omega_{00}|, q_{min}) > 0.$$

*Proof.* The variational characterisation of eigenvalues (Sec.4.5 in [8]) and  $D \geq 1/4$  imply that

$$\begin{aligned}\lambda_{1,D,q} &= - \sup_{u \in H_\nu^2, \|u\|_{L^2(\Omega)}=1} \langle \mathcal{L}_{D,q} u, u \rangle_{L^2(\Omega)} \\ &= \inf_{u \in H_\nu^2, \|u\|_{L^2(\Omega)}=1} [\langle D \nabla u, \nabla u \rangle_{L^2(\Omega)} + \langle qu, u \rangle_{L^2(\Omega)}] \\ &\geq C_{q_{\min}} \inf_{u \in H_\nu^2, \|u\|_{L^2(\Omega)}=1} [\langle \nabla u, \nabla u \rangle_{L^2(\Omega)} + \|u\|_{L^2(\Omega_{00})}^2]\end{aligned}$$

for some  $C_{q_{\min}} > 0$ . Suppose the last expression equals zero. By the usual Poincaré inequality (p.292 in [10]) this implies first that  $u = \text{const}$  and then that  $|\Omega_{00}|/|\Omega| = 0$ , a contradiction since  $\Omega_{00}$  has positive Lebesgue measure. The result follows.  $\square$

We deduce from Parseval's identity that for any  $\varphi \in H^2$  we have

$$\mathcal{L}_{D,q} \varphi = - \sum_{j=1}^{\infty} \lambda_{j,D,q} e_{j,D,q} \langle e_{j,D,q}, \varphi \rangle_{L^2},$$

with convergence at least in  $L^2(\Omega)$ , and that for any  $\varphi \in L^2$  the functions

$$\mathcal{L}_{D,q}^{-1}[-\varphi] = \sum_{j=1}^{\infty} \lambda_{j,D,q}^{-1} e_{j,D,q} \langle e_{j,D,q}, \varphi \rangle_{L^2}$$

solve the PDE (25) with r.h.s.  $-\phi = -\varphi$  in the  $L^2$ -sense. To see that these solutions satisfy Neumann boundary conditions, let us define the associated spectrally defined Sobolev-type spaces

$$\tilde{H}^b(\Omega) = \tilde{H}_{D,q}^b(\Omega) = \left\{ f : \sum_j \lambda_{j,D,q}^b \langle f, e_{j,D,q} \rangle_{L^2}^2 \equiv \|f\|_{\tilde{H}^b}^2 < \infty \right\}, b \geq -1. \quad (41)$$

By Parseval's identity  $\tilde{H}^0 = L^2$  while  $\tilde{H}^{-1} = (\tilde{H}^1)^*$  is interpreted by duality in the usual way. For  $b > 0$  these are subspaces of  $L^2$  with the following properties:

**Proposition 5.2.** *a) For any  $D \in C^1, 0 \leq q \in L^\infty$ , we have that*

$$\tilde{H}_{D,q}^2 = H_\nu^2$$

*and the graph norm  $\|\mathcal{L}_{D,q} f\|_{L^2} = \|f\|_{\tilde{H}_{D,q}^2}$  is Lipschitz-equivalent to the  $H^2$ -norm on  $H_\nu^2$ , with equivalence constants depending only on upper bounds on  $\|q\|_\infty, \|D\|_{C^1}$ .*

*b) Let  $\|D\|_{H^a} + \|q\|_{H^\eta} \leq B$  for some  $a > 2 + d/2, \eta > 1 + d/2$ , and let  $|b| \leq \min(a + 1, \eta + 2)$ . Then  $\tilde{H}^b \subset H^b$  and  $\|u\|_{H^b} \leq c(B, b) \|u\|_{\tilde{H}^b}$  for all  $u \in \tilde{H}^b$ . Conversely, if  $u \in H^b$  is constant outside of a compact set  $K \subset \Omega$ , then  $u \in \tilde{H}^b$  and  $\|u\|_{\tilde{H}^b} \leq c(B, b, K) \|u\|_{H^b}$ .*

*Proof.* First, let  $f \in \tilde{H}_{D,q}^2$  and define its  $J$ -th partial sum as

$$f_J = \sum_{j \leq J} e_{j,D,q} \langle f, e_{j,D,q} \rangle_{L^2}, \quad J \in \mathbb{N},$$

so that  $f_J \in H_\nu^2$  since  $e_j \in H_\nu^2$  and clearly  $f_J \rightarrow f$  in  $\tilde{H}_{D,q}^2$ . Using standard isomorphism theorems for the Neumann Laplacian (e.g., Theorem II.5.4 in [27]) and  $D \geq 1/4$  we see for any  $u \in H_\nu^2$  that

$$\begin{aligned}\|u\|_{H^2} &\leq \|\Delta u\|_{L^2} \lesssim \|\mathcal{L}_{D,q} u\|_{L^2} + \|D\|_{C^1} \|u\|_{H^1} + \|q\|_\infty \|u\|_{L^2} \\ &\lesssim \|u\|_{\tilde{H}_{D,q}^2} + \langle \mathcal{L}_{D,q} u, u \rangle_{L^2}^{1/2} + \|q\|_\infty \|u\|_{L^2} \lesssim \|u\|_{\tilde{H}_{D,q}^2}.\end{aligned}$$

Applying this to  $u = f_J - f_{J'}$  we see that  $f_J$  is a Cauchy sequence in  $H^2$  and hence  $f \in H^2$ , in fact by the trace Theorem I.9.4 in [27] we must have  $f \in H^2_\nu$ . Conversely, let  $f \in H^2_\nu$ . Then by the divergence theorem and Parseval's identity

$$\|f\|_{\tilde{H}^2}^2 = \sum_j \lambda_{j,D,q}^2 \langle f, e_{j,D,q} \rangle_{L^2}^2 = \sum_j \langle \mathcal{L}_{D,q} f, e_{j,D,q} \rangle_{L^2}^2 = \|\mathcal{L}_{D,q} f\|_{L^2}^2 \lesssim (\|D\|_{C^1} + \|q\|_\infty) \|f\|_{H^2} < \infty,$$

so  $f \in \tilde{H}^2_{D,q}$  and Part a) of the proposition follows. Part b) is proved by induction similar to Proposition 2 in [33] and left to the reader.  $\square$

### Parabolic and steady state equation

We now study the underlying time-dependent equation and recall its relationship to the steady state equation (25). Specifically, consider solutions  $v = v_{D,q}$  to the parabolic PDE

$$\begin{aligned} \frac{\partial}{\partial t} v(t, \cdot) &= \mathcal{L}_{D,q} v \quad \text{on } (0, T] \times \Omega \\ v(0, \cdot) &= \varphi \quad \text{on } \Omega \\ \frac{\partial v}{\partial \nu} &= 0 \quad \text{on } (0, T] \times \partial\Omega. \end{aligned} \tag{42}$$

For any  $\varphi \in L^2$  and  $D, q$  as in Conditions 4.1, 4.2, unique solutions exist by standard PDE arguments (as, e.g., in Sec 7.1.2 in [10]) and can be represented spectrally as the convergent series in  $L^2(\Omega)$  given by

$$v_{D,q}(t, \cdot) = \sum_{j=1}^{\infty} e^{-t\lambda_{j,D,q}} e_{j,D,q} \langle e_{j,D,q}, \varphi \rangle_{L^2}. \tag{43}$$

One shows, using Proposition 5.2, that for  $\varphi \in \tilde{H}^b$ , the solutions  $v_{D,q}(t, \cdot)$  lie in  $C([0, T], \tilde{H}^b) \subset C([0, T], H^b)$ , where the parabolic spaces  $C([0, T], X)$  are defined as the bounded continuous maps from  $[0, T]$  into a Banach space  $X$ . We also have from Lemma 5.1, (40), and Parseval's identity that

$$\|v_{D,q}(t, \cdot)\|_{L^2} \leq e^{-ct} \|\varphi\|_{L^2} \quad \forall t > 0, \tag{44}$$

for some  $c > 0$ , which shows that the energy of the system dissipates as  $t \rightarrow \infty$  (corresponding to the 'killing of all particles' in Proposition 3.1). Integrating the penultimate expression in time and using

$$\int_0^s e^{-t\lambda} dt = -\frac{1}{\lambda} [e^{-s\lambda} - 1], \quad \lambda > 0,$$

we have after letting  $s \rightarrow \infty$ , and using the dominated convergence theorem,

$$\int_0^\infty v_{D,q}(t, \cdot) dt = \sum_j \lambda_{j,D,q}^{-1} e_{j,D,q} \langle e_{j,D,q}, \varphi \rangle_{L^2} = \mathcal{L}_{D,q}^{-1} [-\varphi], \quad \varphi \in L^2. \tag{45}$$

Thus the time average of the solutions to (42) equal the solutions  $u_{D,q}$  to the steady state equation (25) with source  $-\phi = -\varphi$ . From Proposition 5.2 and the Sobolev imbedding we have that  $\varphi \in \tilde{H}^b_{D,q} \subset H^b \subset L^\infty$  for some  $b > d/2$  implies that the solutions  $u_{D,q}$  lie in  $\tilde{H}^{2+b}_{D,q} \subset H^{2+b} \subset C^2$ . Theorem 5.1 (or 5.2) in [12] then allows us to represent the solution via the Feynman-Kac formula

$$u_{D,q}(x) = \int_0^\infty v(t, x) dt = E^x \int_0^\infty \varphi(X_t) \exp \left\{ - \int_0^t q(X_s) ds \right\} dt, \quad x \in \Omega, \tag{46}$$

where  $X_s$  is the Markov process from (1) started at  $X_0 = x$ .

**Lemma 5.3.** *Assume Conditions 4.1, 4.2 and that  $u_{D,q} = \mathcal{L}_{D,q}^{-1}(-\varphi)$  is the solution to (25) with source  $-\varphi = -\phi$ .*

a) Suppose further  $\varphi \in \tilde{H}_{D,q}^b$  for some  $b > d/2$  and that  $\|q\|_\infty \leq Q < \infty$ . Then

$$\inf_{x \in \Omega} u_{D,q}(x) \geq \frac{\inf_{x \in \Omega} \varphi(x)}{Q},$$

as well as

$$\|u_{D,q}\|_\infty \leq \|\varphi\|_\infty.$$

b) We also have for any  $\varphi \in L^2$

$$\|u_{D,q}\|_{L^2} \leq C_{q,\Omega_0,0,\Omega} \|\varphi\|_{\tilde{H}^{-1}} \lesssim \|\varphi\|_{L^2}$$

*Proof.* For Part a) note that (46) and the hypotheses on  $\varphi, q$  imply

$$u(x) \geq \inf_{x \in \Omega} \varphi(x) \int_0^\infty e^{-Qt} dt = \frac{\inf_{x \in \Omega} \varphi(x)}{Q}$$

and similarly

$$|u(x)| \leq \|\varphi\|_\infty E^x \exp \left\{ - \int_0^t q(X_s) ds \right\} dt \leq \|\varphi\|_\infty$$

since  $q \geq 0$ . For Part b) we use (the proof of) Lemma 5.1 and sequence space duality to the effect that for any  $\psi \in \tilde{H}^1$  and testing  $g = -\psi$

$$\|\mathcal{L}_{D,q}\psi\|_{\tilde{H}^{-1}} = \sup_{\|g\|_{\tilde{H}^1} \leq 1} |\langle \mathcal{L}_{D,q}\psi, g \rangle_{L^2}| \geq -\langle \mathcal{L}_{D,q}\psi, \psi \rangle_{L^2} \geq c\|\psi\|_{L^2}.$$

Inserting  $\psi = u = \mathcal{L}_{D,q}^{-1}\varphi \in \tilde{H}^1$  and the elementary embedding  $\tilde{H}^1 \subset L^2$  imply the inequalities in b).  $\square$

We finally record the following lemma on regularity of solutions of (25).

**Lemma 5.4.** *In the setting of Lemma 5.3, suppose that  $\varphi \in \tilde{H}_{D,q}^b$  for some  $b \leq \min(a+1, \eta+2) - 2$ . Then*

$$\sup_{\|\varphi\|_{\tilde{H}^b} + \|D\|_{H^a} + \|q\|_{H^\eta} \leq B} \|\mathcal{L}_{D,q}^{-1}(-\varphi)\|_{\tilde{H}^{b+2}} < \infty. \quad (47)$$

The result remains true if  $q = 0$  as well if we assume in addition that  $\int_\Omega \varphi = 0$ .

*Proof.* Within the spectrally defined  $\tilde{H}^\alpha$  scale, by definition of the  $\tilde{H}^b$  norm we immediatly get

$$\sup_{\|\varphi\|_{\tilde{H}^b} + \|D\|_{H^a} + \|q\|_{H^\eta} \leq B} \|\mathcal{L}_{D,q}^{-1}\varphi\|_{\tilde{H}^{b+2}} = \sup_{\|\varphi\|_{\tilde{H}^b} + \|D\|_{H^a} + \|q\|_{H^\eta} \leq B} \|\varphi\|_{\tilde{H}^b} \leq B < \infty.$$

The proof for  $q = 0$  is similar under the additional constraint on  $\varphi$ , cf. Sec.3.1 in [33].  $\square$

## 5.2 Identifiability and Stability.

We will now show that the the diffusivity  $D$  can be identified from  $u_{D,q}$  on a fixed subset  $\Omega_{00} \subset \Omega$  if  $q$  is small enough in a  $H^2$  sense, and assuming  $d \leq 3$ . These hypotheses can be weakened by the introduction of further technicalities (e.g., replacing energy methods by Schauder estimates), but our focus here is not on obtaining the most general stability estimate but on providing a concept proof that inference on diffusivity from killed diffusion is possible under certain hypotheses on the initial condition  $\phi$ . To ease notation we take  $\Omega$  to have normalised Lebesgue measure equal to one, so that the invariant measure of the process  $(X_t)$  from (1) equals 1 identically on  $\Omega$ .

**Condition 5.5.** *Let  $d \leq 3$ . Assume that  $|\Omega| = 1$  and that for some strict subdomain  $\Omega_{00}$  of  $\Omega$  and some  $\epsilon > 0$  the initial condition  $\phi$  of (1) is a probability density that satisfies*

(a) *either  $\phi = 1 + \epsilon e_1$  where  $e_1 \in H_\nu^2$  is the first non-constant eigenfunction of the generator  $\mathcal{L}_{D_0,0}$  of the diffusion process (1), and  $e_1$  satisfies  $\inf_{x \in \Omega_{00}} |\nabla e_1(x)| \geq c > 0$ ,*

(b) or  $\phi \geq 1 + \epsilon$  on  $\Omega_{00}$ .

Before we discuss this condition, let us state our main stability (inverse continuity) estimate for the ‘forward’ map  $D \mapsto u_{D,q}$  of our inverse problem.

**Theorem 5.6.** *Assume  $D_i, i = 1, 2; q, \phi$  satisfy Conditions 4.1, 4.2 with  $\alpha \geq 2 + d/2$  and moreover that  $\phi$  satisfies Condition 5.5 with  $D_0 = D_1$  in case a). Denote by  $u_{D_i,q}$  the corresponding solutions to (25). Suppose further that  $\|D_1\|_{H^\alpha} + \|D_2\|_{H^\alpha} \leq b$ . Then there exist positive constants  $\delta, C$  depending on  $b, \phi, \epsilon, \Omega, \Omega_{00}, \alpha$  such that if  $\|q\|_{H^2} < \delta$  we have*

$$\|D_1 - D_2\|_{L^2(\Omega)} \leq C \|u_{D_1,q} - u_{D_2,q}\|_{H^2(\Omega_{00})}. \quad (48)$$

The following remark discusses some physical interpretations of our hypotheses.

**Remark 5.7.** Since  $\phi$  is a probability density, Condition 5.5b) requires the initial state to be prepared such that more than average (= 1) particles start diffusing in the sub-domain  $\Omega_{00}$  where we wish to identify  $D$ . This can be thought of as ensuring that we will see sufficiently many binding events in the region  $\Omega_{00}$ .

The hypothesis a) is somewhat more subtle to interpret: It essentially requires the domain  $\Omega$  and operator  $\mathcal{L}_{D_0,0} = \nabla \cdot (D_0 \nabla)$  to satisfy the hotspots conjecture (see Sections 2.2.2 and 3.7 in [33] for concrete examples and relevant references). We also notice that  $e_1 \in H^2_\nu \subset L^\infty$  (cf. Prop.2 in [33]), so Condition 4.2 is satisfied for  $b = 2, d < 4$  and  $\epsilon = \epsilon(\|e_1\|_\infty)$  small enough. Since the distribution  $p_t(\psi)$  of  $X_t$  from (1) for any initial condition  $\psi$  at time  $t$  equals  $1 + \sum_{j \geq 1} e^{-t\lambda_j} e_j \langle e_j, \psi \rangle_{L^2}$  (with  $(e_j, -\lambda_j)$  the eigen-pairs of  $\mathcal{L}_{D_0,0}$ ), the perturbative ( $t \rightarrow \infty$ ) interpretation of hypothesis a) is that  $\phi = 1 + \epsilon e_1$  is already close to its equilibrium distribution. Statistically speaking, the first eigenfunction  $e_1 = e_{1,D_0}$  is the last ‘informative’ part of diffusion that we can see before it reaches its non-informative (uniform) equilibrium distribution. [As  $\phi = e_1$  now depends on  $D_0$ , one needs to replace it by an empirical estimate in the MCMC implementation.]

That  $q$  needs to be small enough can be related to the problem that too large  $q$  may kill diffusing particles too aggressively for them to visit all areas of  $\Omega_{00}$  before they ‘bind’. One could equivalently calibrate the parameter of the exponential random variable  $Y$  featuring in the binding time. It remains an open problem whether this hypothesis is necessary or not. The numerical findings from Sec. 4.2 suggest that the ratio  $D/q$  is relevant for stable recovery, and moreover that recovery of  $D$  is possible also after relaxing the hypothesis Condition 5.5. This presents interesting avenues for future research.

### 5.3 Proof of Theorem 5.6

We first recall a general stability lemma for a transport operator, taken from [33] and inspired by earlier work in [36, 32].

**Condition 5.8.** *Let  $u_0 \in H^2(\Omega)$  be a function such that  $\sup_{x \in \Omega_{00}} |u_0(x)| \leq u < \infty$  and*

$$\frac{1}{2} \Delta u_0(x) + \mu |\nabla u_0(x)|^2 \geq c_0 > 0, \quad \text{a.e. } x \in \Omega_{00} \quad (49)$$

for some subset  $\Omega_{00}$  s.t.  $\text{dist}(\Omega_{00}, \Omega) > 0$ .

**Lemma 5.9.** *(Lemma 2 in [33]) Suppose  $u_0$  (or  $-u_0$ ) satisfies Condition 5.8. Then there exists a constant  $\underline{c} = \underline{c}(u, c_0, \mu) > 0$  such that we have, for any  $h \in C^1$  that vanishes on  $\Omega \setminus \Omega_{00}$ ,*

$$\|\nabla \cdot (h \nabla u_0)\|_{L^2(\Omega)} \geq \underline{c} \|h\|_{L^2(\Omega)}.$$

Given the preceding lemma, the proof of Theorem 5.6 proceeds in several steps by comparing the solution of (25) to the solution of the standard Neumann problem with  $q = 0$ , which exists if the ‘source’ term  $\phi$  integrates to zero.

**Step 1:** For  $\phi$  satisfying Conditions 4.2 and 5.5 we set

$$\phi_0 := \phi - 1 = \phi - \int_{\Omega} \phi \quad \text{and} \quad w_\phi = \mathcal{L}_{D_1,0}^{-1}[-\phi_0].$$

From Lemma 5.4 combined with Proposition 5.2 we know that  $w_\phi \in H^2_\nu \cap H^{\beta+2} \subset C^2$  for some  $2 > \beta > d/2$ . Let us show that  $w_\phi$  further satisfies Condition 5.8 for some  $\mu > 0$  and subset  $\Omega_{00} \subset \Omega$ .

(a) If  $\phi = 1 + \epsilon e_1$  with  $|\nabla e_1| > 0$  on  $\Omega_{00}$ ,

$$w_\phi = \mathcal{L}_{D_1,0}^{-1}[-\epsilon e_1] = \sum_{j=1}^{\infty} \frac{\epsilon \langle e_j, e_1 \rangle}{\lambda_j} e_j = \frac{\epsilon e_1}{\lambda_1}$$

such that  $|\nabla w_\phi| = |\frac{\epsilon \nabla e_1}{\lambda_1}| > 0$  on  $\Omega_{00}$ . Since  $\Delta w_\phi \in L^\infty$  we see that for  $\mu$  large enough, Condition 5.8 is satisfied with  $u_0 = w_\phi$ .

(b) If  $\phi \geq 1 + \epsilon$  with  $\epsilon > 0$  on  $\Omega_{00}$  then  $\phi_0 \geq \epsilon$  on  $\Omega_{00}$ . Let us write  $\bar{w}_\phi = -w_\phi$  then,

$$\phi_0 = \nabla \cdot (D_1 \nabla \bar{w}_\phi) = D_1 \Delta \bar{w}_\phi + \nabla D_1 \cdot \nabla \bar{w}_\phi. \quad (50)$$

This forces either  $D_1 \Delta \bar{w}_\phi(x) > \frac{\epsilon}{2} > 0$  and then  $\Delta \bar{w}_\phi(x) > \frac{\epsilon}{2 \|D_1\|_\infty} > 0$ , or

$$\frac{\epsilon}{2} \leq \nabla D_1(x) \cdot \nabla \bar{w}_\phi(x) \leq |\nabla \bar{w}_\phi(x)| |\nabla D_1(x)|$$

so that  $|\nabla \bar{w}_\phi| > \frac{\epsilon}{2 \|D_1\|_{C^1}}$  on  $\Omega_{00}$ . Conclude that again, Condition 5.8 is verified for  $u_0 = \bar{w}_\phi = -w_\phi$  and  $\mu$  large enough, since  $\Delta \bar{w}_\phi \in L^\infty$ .

**Step 2.** We next consider a constant potential  $q = q_{min} > 0$  on the domain  $\Omega$ . In this case the spectral decomposition of  $\mathcal{L}_{D_1, q_{min}}$  is the same as the one of  $\mathcal{L}_{D_1, 0}$  but with eigenvalues

$$\lambda'_j = \lambda_j + q_{min}, \quad j \in \mathbb{N} \cup \{0\},$$

shifted by  $q_{min}$ , including  $\lambda_0 = 0$  corresponding to  $e_0 = 1$ . The operator  $\mathcal{L}_{D_1, q_{min}}$  thus has inverse  $\mathcal{L}_{D_1, q_{min}}^{-1}$  on  $L^2(\Omega)$  with spectral representation

$$u = \mathcal{L}_{D_1, q_{min}}^{-1}[\varphi] = - \sum_{j=0}^{\infty} \frac{1}{\lambda'_j} e_j \langle e_j, \varphi \rangle, \quad \varphi \in L^2,$$

solving  $\nabla \cdot (D \nabla u) - q_{min} u = \varphi$  subject to Neumann boundary conditions. Now since  $w_\phi$  solves  $\mathcal{L}_{D_1, 0} w_\phi = -\phi_0$  with Neumann boundary conditions, the function

$$v_\phi := w_\phi + \frac{\int_\Omega \phi}{q_{min}}$$

also satisfies Neumann boundary conditions and solves

$$\mathcal{L}_{D_1, q_{min}} v_\phi = \mathcal{L}_{D_1, 0} v_\phi - q_{min} v_\phi = -\phi_0 - \int_\Omega \phi - q_{min} w_\phi = -\phi - q_{min} w_\phi$$

Therefore if  $u = u_{D_1, q_{min}} = \mathcal{L}_{D_1, q_{min}}^{-1}[-\phi]$  is the actual solution of the steady state Schrödinger equation

$$\nabla \cdot (D_1 \nabla u) - q_{min} u = -\phi$$

with Neumann boundary conditions, then

$$\begin{aligned} \|u_{D_1, q_{min}} - v_\phi\|_{C^2} &= \|\mathcal{L}_{D_1, q_{min}}^{-1}[-\phi] - \mathcal{L}_{D_1, q_{min}}^{-1}[-\phi - q_{min} w_\phi]\|_{C^2} \\ &= q_{min} \|\mathcal{L}_{D_1, q_{min}}^{-1}[w_\phi]\|_{C^2} \lesssim q_{min} \|\mathcal{L}_{D_1, q_{min}}^{-1}[w_\phi]\|_{\tilde{H}_{D_1, q_{min}}^{2+\beta}} \lesssim q_{min} \end{aligned}$$

using the Sobolev imbedding ( $d < 4$ ) as well as the elliptic regularity Lemma 5.4. When Condition 5.8 holds for  $w_\phi$  then it also holds for  $v_\phi$  as the constant subtracted vanishes after taking the gradient and the Laplacian. Hence for  $q_{min}$  small enough the function  $u_{D_1, q_{min}}$  satisfies Condition 5.8 too since its  $C^2$ -norm distance to  $v_\phi$  can be made as small as desired.

**Step 3.** Now, let  $u_{D_1,q} = \mathcal{L}_{D_1,q}^{-1}[-\phi]$  be the solution of the steady state Schrödinger equation with non-constant potential  $q$  such that  $\|q - q_{min}\|_{H^n}$  is small. We then see that

$$\mathcal{L}_{D_1,q}(u_{D_1,q} - u_{D_1,q_{min}}) = (q - q_{min})u_{D_1,q_{min}}.$$

We have that  $u_{D_1,q_{min}}$  lies in  $\tilde{H}^2 = H^2_\nu$  and so does  $q - q_{min}$  as it belongs to  $H^2$  and is constant near the boundary. Therefore  $(q - q_{min})u_{D_1,q} \in H^2_\nu$  as well in view of (4) and  $d < 4$ . Hence we obtain from the Sobolev imbedding with  $d/2 < \beta < 2$ , the definition of the  $\tilde{H}^b$ -norms, and again (4)

$$\begin{aligned} \|u_{D_1,q} - u_{D_1,q_{min}}\|_{C^2} &\lesssim \|\mathcal{L}_{D_1,q}^{-1}[(q - q_{min})u_{D_1,q_{min}}]\|_{\tilde{H}_{D_1,q}^{2+\beta}} \\ &\lesssim \|(q - q_{min})u_{D_1,q_{min}}\|_{\tilde{H}_{D_1,q}^2} \\ &\lesssim \|q - q_{min}\|_{H^2} \|u_{D_1,q_{min}}\|_{H^2}, \end{aligned}$$

and the r.h.s. can be made as small as desired for  $\|q - q_{min}\|_{H^2}$  small enough, so that again Condition 5.8 is inherited by  $u_{D_1,q}$ . From what precedes Lemma 5.9 applies to give for any  $h \in C^1$  that vanishes on  $\Omega \setminus \Omega_{00}$

$$\|\nabla \cdot (h \nabla u_{D_1,q})\|_{L^2} \geq c \|h\|_{L^2}$$

and applying this to  $h = D_1 - D_2$  gives

$$\|D_1 - D_2\|_{L^2} \lesssim \|\nabla \cdot ((D_1 - D_2) \nabla u_{D_1,q})\|_{L^2}. \quad (51)$$

Now, since  $u_{D_1,q}$  solves  $\mathcal{L}_{D_1,q} u = -\phi$ , and  $u_{D_2,q}$  solves the same equation with  $D_2$  in place of  $D_1$ , the r.h.s equals, with all  $L^2$ -norms over  $\Omega_{00}$ ,

$$\begin{aligned} \|\nabla \cdot ((D_1 - D_2) \nabla u_{D_1,q})\|_{L^2}^2 &= \|\nabla \cdot (D_1 \nabla u_{D_1,q}) - \nabla \cdot (D_2 \nabla (u_{D_1,q} - u_{D_2,q} + u_{D_2,q}))\|_{L^2}^2 \\ &\leq \|-\phi + q u_{D_1,q} + \phi - q u_{D_2,q}\|_{L^2}^2 + \|\nabla \cdot (D_2 \nabla (u_{D_1,q} - u_{D_2,q}))\|_{L^2}^2 \\ &= \|q(u_{D_1,q} - u_{D_2,q})\|_{L^2}^2 + \|\nabla \cdot (D_2 \nabla (u_{D_1,q} - u_{D_2,q}))\|_{L^2}^2 \\ &\lesssim \|u_{D_1,q} - u_{D_2,q}\|_{H^2(\Omega_{00})}^2. \end{aligned}$$

This completes the proof of Theorem 5.6.

## 6 Proofs for Section 2

### 6.1 Proof of Lemma 2.1

By independence and the definitions, it suffices to prove

$$-\mathbb{E}_{\Lambda_0} \left( \log \frac{p_\Lambda}{p_{\Lambda_0}}(Y) \right) \leq 2n \frac{|\Lambda(B) - \Lambda_0(B)|^2}{\Lambda_0(B)}$$

for every  $Y = Y_i, B = B_i$  and where  $p_\Lambda = p_{\Lambda,n}^1$  in this proof. From the definitions of Poisson densities we have

$$\frac{p_\Lambda}{p_{\Lambda_0}}(y) = e^{-n(\Lambda(B) - \Lambda_0(B))} \left[ \frac{\Lambda(B)}{\Lambda_0(B)} \right]^y \quad (52)$$

and therefore

$$\begin{aligned} \mathbb{E}_{\Lambda_0} \left[ \log \frac{p_\Lambda}{p_{\Lambda_0}}(Y) \right] &= -n[\Lambda(B) - \Lambda_0(B)] + \log[\Lambda(B)/\Lambda_0(B)] \mathbb{E}_{\Lambda_0}[Y] \\ &= -n[\Lambda(B) - \Lambda_0(B)] + n\Lambda_0(B) \log \left[ 1 + \frac{\Lambda(B) - \Lambda_0(B)}{\Lambda_0(B)} \right] \end{aligned}$$

Using a second order Taylor expansion about zero with mean value remainder

$$\log(1+x) = x - \frac{1}{2}x^2 \frac{1}{(1+\zeta x)^2} \quad \text{for some } \zeta \in (0, 1)$$

at  $x = \frac{\Lambda(B) - \Lambda_0(B)}{\Lambda_0(B)} \in (-1/2, 1/2)$ , we deduce

$$\log\left(1 + \frac{\Lambda(B) - \Lambda_0(B)}{\Lambda_0(B)}\right) = \frac{\Lambda(B) - \Lambda_0(B)}{\Lambda_0(B)} - \frac{1}{2} \left(\frac{\Lambda(B) - \Lambda_0(B)}{\Lambda_0(B)}\right)^2 \frac{1}{(1 + \zeta \frac{\Lambda(B) - \Lambda_0(B)}{\Lambda_0(B)})^2}.$$

The last factor is bounded by 4 under our hypothesis and we deduce

$$-\mathbb{E}_{P_{\Lambda_0}} \left[ \log \frac{p_{\Lambda}}{p_{\Lambda_0}}(Y) \right] = n \frac{(\Lambda(B) - \Lambda_0(B))^2}{2\Lambda_0(B)} \frac{1}{(1 + \zeta \frac{\Lambda(B) - \Lambda_0(B)}{\Lambda_0(B)})^2} \leq 2n \frac{(\Lambda(B) - \Lambda_0(B))^2}{\Lambda_0(B)},$$

completing the proof.

## 6.2 Proof of Lemma 2.2

We prove the result for  $c \leq 2$  relevant below, in which case we will show that one can take  $L_c = c^2/8$ . The case  $c > 2$  follows from simple modifications and is left to the reader. By Jensen's inequality,

$$\log \int_A \frac{p_{\Lambda}}{p_{\Lambda_0}}(Y_1, \dots, Y_K) d\nu(\Lambda) \geq \int_A \log \frac{p_{\Lambda}}{p_{\Lambda_0}}(Y_1, \dots, Y_K) d\nu(\Lambda)$$

so using also Lemma 2.1, the probability in question is bounded, for  $s > 0$ , by

$$\begin{aligned} P_{\Lambda_0} \left( \int_A \log \frac{p_{\Lambda}}{p_{\Lambda_0}}(Y_1, \dots, Y_K) d\nu(\Lambda) \leq -(2+c)n\epsilon^2 \right) \\ &= P_{\Lambda_0} \left( \int_A \left[ \log \frac{p_{\Lambda}}{p_{\Lambda_0}}(Y_1, \dots, Y_K) - \mathbb{E}_{\Lambda_0} \log \frac{p_{\Lambda}}{p_{\Lambda_0}}(Y_1, \dots, Y_K) \right] d\nu(\Lambda) \right. \\ &\quad \left. \leq -(2+c)n\epsilon^2 - \int_A \mathbb{E}_{\Lambda_0} \log \frac{p_{\Lambda}}{p_{\Lambda_0}}(Y_1, \dots, Y_K) d\nu(\Lambda) \right) \\ &\leq P_{\Lambda_0} \left( \int_A \left[ \log \frac{p_{\Lambda}}{p_{\Lambda_0}}(Y_1, \dots, Y_K) - \mathbb{E}_{\Lambda_0} \log \frac{p_{\Lambda}}{p_{\Lambda_0}}(Y_1, \dots, Y_K) \right] d\nu(\Lambda) \leq -cn\epsilon^2 \right) \\ &\leq P_{\Lambda_0} \left( \exp s \left| \mathbb{E}_{\nu} \left[ \log \frac{p_{\Lambda}}{p_{\Lambda_0}}(Y_1, \dots, Y_K) - \mathbb{E}_{\Lambda_0} \log \frac{p_{\Lambda}}{p_{\Lambda_0}}(Y_1, \dots, Y_K) \right] \right| \geq e^{snc\epsilon^2} \right) \\ &\leq e^{-snc\epsilon^2} \mathbb{E}_{\Lambda_0} \left[ \exp s \left| \mathbb{E}_{\nu} \left[ \log \frac{p_{\Lambda}}{p_{\Lambda_0}}(Y_1, \dots, Y_K) - \mathbb{E}_{\Lambda_0} \log \frac{p_{\Lambda}}{p_{\Lambda_0}}(Y_1, \dots, Y_K) \right] \right| \right] \\ &\leq e^{-snc\epsilon^2} \mathbb{E}_{\nu} \left[ \mathbb{E}_{\Lambda_0} \left[ \exp s \left| \log \frac{p_{\Lambda}}{p_{\Lambda_0}}(Y_1, \dots, Y_K) - \mathbb{E}_{\Lambda_0} \log \frac{p_{\Lambda}}{p_{\Lambda_0}}(Y_1, \dots, Y_K) \right| \right] \right] \end{aligned}$$

using also Jensen's inequality and Fubini's theorem in the last step. We define random variables

$$Z_i := \log \frac{p_{\Lambda}}{p_{\Lambda_0}}(Y_i) - \mathbb{E}_{\Lambda_0} \log \frac{p_{\Lambda}}{p_{\Lambda_0}}(Y_i) = Y_i \log \frac{\Lambda(B_i)}{\Lambda_0(B_i)} - n\Lambda_0(B_i) \log \frac{\Lambda(B_i)}{\Lambda_0(B_i)}, \quad i = 1, \dots, K,$$

and estimate the expectation of  $e^{s|\sum_i Z_i|} \leq e^{s\sum_i Z_i} + e^{-s\sum_i Z_i}$ . For the first term we have by independence

$$\begin{aligned} \mathbb{E}_{\Lambda_0} [e^{s\sum_i Z_i}] &= \prod_{i=1}^K \mathbb{E}_{\Lambda_0} \left[ \exp \left( s \log \frac{\Lambda(B_i)}{\Lambda_0(B_i)} Y_i - s \log \frac{\Lambda(B_i)}{\Lambda_0(B_i)} n\Lambda_0(B_i) \right) \right] \\ &= \prod_{i=1}^K \exp \left( n\Lambda_0(B_i) \left( e^{s \log \frac{\Lambda(B_i)}{\Lambda_0(B_i)}} - 1 - s \log \frac{\Lambda(B_i)}{\Lambda_0(B_i)} \right) \right). \end{aligned} \quad (**)$$

Using a second order Taylor expansion with mean value remainder,  $0 < \zeta < 1$ , and the hypothesis  $\max_i \left| \frac{\Lambda(B_i) - \Lambda_0(B_i)}{\Lambda_0(B_i)} \right| < 1/2$  we get for  $0 < s \leq 1$  that

$$e^{s \log \frac{\Lambda(B_i)}{\Lambda_0(B_i)}} = 1 + s \log \frac{\Lambda(B_i)}{\Lambda_0(B_i)} + \frac{s^2}{2} \left( \log \frac{\Lambda(B_i)}{\Lambda_0(B_i)} \right)^2 \underbrace{e^{\zeta s \log \frac{\Lambda(B_i)}{\Lambda_0(B_i)}}}_C$$

with  $C \leq 3/2$ . Similarly from the mean value theorem

$$\log \frac{\Lambda(B_i)}{\Lambda_0(B_i)} = \log \left( 1 + \frac{\Lambda(B_i) - \Lambda_0(B_i)}{\Lambda_0(B_i)} \right) = \frac{\Lambda(B_i) - \Lambda_0(B_i)}{\Lambda_0(B_i)} \underbrace{\frac{1}{1 + \zeta \frac{\Lambda(B_i) - \Lambda_0(B_i)}{\Lambda_0(B_i)}}}_{C'}$$

where  $1/C' \leq 2$ . We conclude that (\*\*) is bounded by

$$\begin{aligned} \mathbb{E}_{\Lambda_0} [e^{s \sum_i Z_i}] &\leq \prod_{i=1}^K \exp \left( n \Lambda_0(B_i) s^2 \left( \log \frac{\Lambda(B_i)}{\Lambda_0(B_i)} \right)^2 \right) \\ &\leq \prod_{i=1}^K \exp \left( 2n s^2 \frac{(\Lambda(B_i) - \Lambda_0(B_i))^2}{\Lambda_0(B_i)} \right) = e^{2s^2 n \mathcal{D}_{2,K}^2(\Lambda, \Lambda_0)}, \quad 0 < s \leq 1. \end{aligned}$$

and the same inequality can be established for  $e^{-s \sum_i Z_i}$  by similar arguments. Combining what precedes with the support of the probability measure  $\nu$  we obtain for all  $0 < s \leq 1$  the bound

$$\begin{aligned} P_{\Lambda_0} \left( \int_A \frac{p_{\Lambda}}{p_{\Lambda_0}}(Y_1, \dots, Y_K) d\nu(\Lambda) \leq e^{-(2+c)n\epsilon^2} \right) &\leq 2e^{-snc\epsilon^2} \mathbb{E}_{\nu} e^{2s^2 n \mathcal{D}_{2,K}^2(\Lambda, \Lambda_0)} \\ &\leq 2e^{-snc\epsilon^2 + 2s^2 n \epsilon^2} = 2e^{-n\epsilon^2(sc - 2s^2)} \end{aligned}$$

Choosing  $s = c^2/8 < 1$  (permitted since we assumed  $c \leq 2$ ) gives the overall bound

$$P_{\Lambda_0} \left( \int_A \frac{p_{\Lambda}}{p_{\Lambda_0}}(Y_1, \dots, Y_K) d\nu(\Lambda) \leq e^{-(2+c)n\epsilon^2} \right) \leq 2e^{-c^2 n \epsilon^2 / 8},$$

completing the proof.

### 6.3 Proof of Theorem 2.3

**Step 1.** Let us introduce the sets

$$\mathcal{L}^n = \Theta_n \cap \mathcal{R}.$$

We can find tests (indicator functions)  $\Psi_n \equiv \Psi(Y_1, \dots, Y_K, n)$  such that for every  $n \in \mathbb{N}$  and  $M > 0$  large enough depending on  $C, \Lambda_{min}$ ,

$$(i) \quad \mathbb{E}_{\Lambda_0} \Psi_n \rightarrow 0, \quad (ii) \quad \sup_{\Lambda \in \mathcal{L}^n: \|\Lambda - \Lambda_0\|_{\ell_1} \geq M\sqrt{K/n}} \mathbb{E}_{\Lambda} (1 - \Psi_n) \leq L e^{-(C+4)n\epsilon_n^2}$$

*Proof.* Define  $\Psi_n = \mathbb{1}\{\|\hat{\Lambda} - \Lambda_0\|_{\ell_1} \geq t\sqrt{K/n}\}$ . Part (i) follows from the concentration inequality Theorem 6.1 and so does (ii) in view of the following argument: note that

$$\mathbb{E}_{\Lambda} [1 - \Psi_n] = P_{\Lambda}(\Psi_n = 0) = P_{\Lambda}(\|\hat{\Lambda} - \Lambda_0\|_{\ell_1} \leq t\sqrt{K/n}).$$

But

$$\|\hat{\Lambda} - \Lambda_0\|_{\ell_1} = \|\hat{\Lambda} - \Lambda + \Lambda - \Lambda_0\|_{\ell_1} \geq \|\Lambda - \Lambda_0\|_{\ell_1} - \|\hat{\Lambda} - \Lambda\|_{\ell_1}$$

so that

$$\mathbb{E}_{\Lambda} [1 - \Psi_n] \leq P_{\Lambda}(\|\Lambda - \Lambda_0\|_{\ell_1} - \|\hat{\Lambda} - \Lambda\|_{\ell_1} < t\sqrt{K/n}) = P_{\Lambda}(\|\hat{\Lambda} - \Lambda\|_{\ell_1} > \|\Lambda - \Lambda_0\|_{\ell_1} - t\sqrt{K/n})$$

and hence, using (55), we have for all  $\Lambda \in \mathcal{L}^n$  such that  $\|\Lambda - \Lambda_0\|_{\ell_1} \geq M\sqrt{K/n}$  the bound

$$\begin{aligned}\mathbb{E}_\Lambda(1 - \Psi_n) &\leq P_\Lambda(\|\hat{\Lambda} - \Lambda\|_{\ell_1} > \|\Lambda - \Lambda_0\|_{\ell_1} - t\sqrt{K/n}) \leq P_\Lambda(\|\hat{\Lambda} - \Lambda\|_{\ell_1} > M\varepsilon_n - t\sqrt{K/n}) \\ &\leq P_\Lambda(\|\hat{\Lambda} - \Lambda\|_{\ell_1} > (M-t)\sqrt{K/n}) \leq e^{-c'K} \leq e^{-c'n\varepsilon_n^2}\end{aligned}$$

□

**Step 2.** Notice first

$$\mathbb{E}_{\Lambda_0} \left[ \Pi(\{\Lambda \in (\mathcal{L}^n)^c \cup \|\Lambda^{(K)} - \Lambda_0^{(K)}\|_{\ell_1} \geq M\sqrt{K/n} | Y_1, \dots, Y_K\}) \Psi_n \right] \leq \mathbb{E}_{\Lambda_0} \Psi_n \rightarrow 0$$

by assumption on the tests. Let us write shorthand

$$\bar{L}(K, n) \equiv (\mathcal{L}^n)^c \cup \{\Lambda : \|\Lambda^{(K)} - \Lambda_0^{(K)}\|_{\ell_1} \geq M\sqrt{K/n}\}$$

so we only need to prove convergence in  $P_{\Lambda_0}$  to 0 of

$$\Pi(\bar{L}(K, n) | Y_1, \dots, Y_K)(1 - \Psi_n) = \frac{\int_{\bar{L}(K, n)} (p_\Lambda/p_{\Lambda_0})(Y_1, \dots, Y_K) d\Pi(\Lambda)(1 - \Psi_n)}{\int (p_\Lambda/p_{\Lambda_0})(Y_1, \dots, Y_K) d\Pi(\Lambda)}$$

Lemma 2.2 shows that for all  $c > 0$  and probability measures  $\nu$  with support in

$$B_n := \{\Lambda \in \mathcal{L}^n : D_{2,K}^2(\Lambda, \Lambda_0)^2 \leq \varepsilon_n^2, \mathcal{D}_{\infty, K}(\Lambda, \Lambda_0) \leq 1/2\},$$

we have

$$P_{\Lambda_0} \left( \int \frac{p_\Lambda}{p_{\Lambda_0}}(Y_1, \dots, Y_K) d\nu(\Lambda) \leq e^{-(2+c)n\varepsilon_n^2} \right) \leq 2e^{-L\varepsilon_n^2}$$

We apply this result with  $c = 1$  and  $\nu$  equal to the normalised restriction of the prior  $\Pi$  to  $B_n$ . Together with Step 1, this shows that if  $A_n$  is the event

$$A_n := \left\{ \int_{B_n} \frac{p_\Lambda}{p_{\Lambda_0}}(Y_1, \dots, Y_K) d\Pi(\Lambda) \geq \Pi(B_n) e^{-3n\varepsilon_n^2} \geq e^{-(3+C)n\varepsilon_n^2} \right\},$$

then  $P_{\Lambda_0}(A_n) \geq 1 - 2e^{-L_1 n\varepsilon_n^2} \rightarrow 1$ , and we can write for every  $\epsilon > 0$ ,

$$\begin{aligned}P_{\Lambda_0} \left( \frac{\int_{\bar{L}(K, n)} (p_\Lambda/p_{\Lambda_0})(Y_1, \dots, Y_K) d\Pi(\Lambda)(1 - \Psi_n)}{\int (p_\Lambda/p_{\Lambda_0})(Y_1, \dots, Y_K) d\Pi(\Lambda)} > \epsilon \right) \\ \leq P_{\Lambda_0}(A_n^c) + P_{\Lambda_0} \left( e^{(3+C)n\varepsilon_n^2} (1 - \Psi_n) \int_{\bar{L}(K, n)} \frac{p_\Lambda}{p_{\Lambda_0}}(Y_1, \dots, Y_K) d\Pi(\Lambda) \geq \epsilon \right).\end{aligned}$$

Now, using that

$$\begin{aligned}\mathbb{E}_{\Lambda_0} \left[ \frac{p_\Lambda}{p_{\Lambda_0}}(Y_1, \dots, Y_K) \right] &= \int_{p_{\Lambda_0} > 0} p_\Lambda d\mu \leq 1, \quad (\mu \text{ counting measure on } \mathbb{N}_0^K) \\ \mathbb{E}_{\Lambda_0} \left[ \frac{p_\Lambda}{p_{\Lambda_0}}(Y_1, \dots, Y_K)(1 - \Psi_n) \right] &\leq \mathbb{E}_\Lambda(1 - \Psi_n)\end{aligned}$$

and that  $0 \leq 1 - \Psi_n \leq 1$ , we obtain

$$\mathbb{E}_{\Lambda_0} \left[ (1 - \Psi_n) \int_{\bar{L}(K, n)} \frac{p_\Lambda}{p_{\Lambda_0}}(Y_1, \dots, Y_K) d\Pi(\Lambda) \right] \leq 2\Pi(\mathcal{L} \setminus \mathcal{L}^n) + \sup_{\Lambda \in \mathcal{L}^n : \|\Lambda^{(K)} - \Lambda_0^{(K)}\|_{\ell_1} \geq M\sqrt{K/n}} \mathbb{E}_\Lambda(1 - \Psi_n).$$

Now the assumptions on  $\mathcal{L}^n$  and concentration properties of the tests combined with Markov's inequality give, for every  $\epsilon > 0$ ,

$$P_{\Lambda_0} \left( (1 - \Psi_n) \int_{\bar{L}(K, n)} \frac{p_\Lambda}{p_{\Lambda_0}}(Y_1, \dots, Y_K) d\Pi(\Lambda) > \frac{\epsilon}{e^{(3+C)n\varepsilon_n^2}} \right) \leq (3L/\epsilon) e^{-n\varepsilon_n^2}$$

and the theorem follows by combining the preceding estimates and appropriate choice of  $\epsilon$ .

## 6.4 A concentration inequality for high-dimensional Poisson regression

For the construction of tests in the previous proof, we rely on a nonparametric estimator  $\hat{\Lambda}$  of  $\Lambda$  with good concentration properties. But note that  $\hat{\Lambda}$  does not satisfy the ‘PDE constraint’ (24), hence cannot be used itself in the context of our inverse problem since the stability estimate Theorem 5.6 does not apply to it.

A natural statistic is the normalised bin count, which gives the random vector in  $(\mathbb{R}^+)^K$  with entries

$$\hat{\Lambda}_i = \frac{Y_i}{n}, \quad i = 1, \dots, K, \quad (53)$$

for data from (7). Now consider the average  $\ell_1$ -risk across the bins

$$\|\hat{\Lambda} - \Lambda\|_{\ell_1} = \sum_{i=1}^K |\hat{\Lambda}_i - \Lambda(B_i)| = \frac{1}{n} \sum_{i=1}^K |Y_i - \mathbb{E}_\Lambda Y_i| \quad (54)$$

A basic first inequality is, using also the Cauchy-Schwarz inequality twice,

$$\mathbb{E}_\Lambda \|\hat{\Lambda} - \Lambda\|_{\ell_1} \leq \frac{1}{n} \sum_{i \leq K} \sqrt{\text{Var}(Y_i)} = \frac{1}{n} \sum_{i \leq K} \sqrt{n\Lambda(B_i)} \leq \frac{\sqrt{K}\sqrt{\Lambda(0)}}{\sqrt{n}}.$$

This is upgraded to a probability inequality in the following theorem which for  $K \gtrsim n\varepsilon_n^2$  has a sufficiently rapidly decaying concentration function to combine with the posterior contraction rate proof from above.

**Theorem 6.1.** *If  $K \leq Cn$  for some  $C > 0$ , then for any  $c' > 0$  we can find  $t$  large enough depending on  $c', C$  and on an upper bound  $\Lambda_{\max} \geq \Lambda(0)$ , such that we have for every  $K, n \in \mathbb{N}$ ,*

$$P_\Lambda \left( \|\hat{\Lambda} - \Lambda\|_{\ell_1} \geq t\sqrt{K/n} \right) \leq e^{-c'K} \quad (55)$$

*Proof.* We represent the  $\ell_1$ -norm of the vector  $\hat{\Lambda} - \Lambda$  on  $\mathbb{R}^K$  by duality,

$$\|\hat{\Lambda} - \Lambda\|_{\ell_1} = \sup_{v \in \mathbb{R}^K: \max_i |v_i| \leq 1} |v^T(\hat{\Lambda} - \Lambda)|.$$

Then let  $v_{(m)}, m = 1, \dots, N(1/2)$ , be vectors in the  $\ell_\infty$ -unit ball  $B_\infty(1)$  of  $\mathbb{R}^K$  that form the centers of a radius-1/2 covering of  $B_\infty^1$  for the  $\ell_\infty$  norm  $\|v\|_{\ell_\infty} = \max_{i \leq K} |v_i|$ . By Prop. 4.3.34 in [14] we require at most  $N(1/2) \leq 6^K$  such balls. We deduce from the triangle inequality for every fixed  $v \in B_\infty(1)$  and some  $v_m$  that

$$|v^T(\hat{\Lambda} - \Lambda)| \leq |(v - v_{(m)})^T(\hat{\Lambda} - \Lambda)| + |v_{(m)}^T(\hat{\Lambda} - \Lambda)| \leq \frac{1}{2} \|\hat{\Lambda} - \Lambda\|_{\ell_1} + |v_{(m)}^T(\hat{\Lambda} - \Lambda)|$$

Taking suprema over  $v \in B_\infty(1)$  and subtracting we get

$$\frac{1}{2} \|\hat{\Lambda} - \Lambda\|_{\ell_1} \leq \max_{m=1, \dots, N(1/2)} |v_{(m)}^T(\hat{\Lambda} - \Lambda)|$$

so that the l.h.s. of (55) is bounded, using also a union bound for probabilities, by

$$P_\Lambda \left( \max_{m=1, \dots, N(1/2)} |v_{(m)}^T(\hat{\Lambda} - \Lambda)| \geq t\sqrt{K/n}/2 \right) \leq 6^K \sup_{v \in B_\infty(1)} P_\Lambda \left( |n v^T(\hat{\Lambda} - \Lambda)| \geq t\sqrt{Kn}/2 \right) \quad (56)$$

We now need an appropriate tail bound for the random variables

$$n v^T(\hat{\Lambda} - \Lambda) = \sum_{i=1}^K v_i (Y_i - \mathbb{E}_\Lambda Y_i), \quad v \in B_\infty(1).$$

We consider each summand and  $0 < x < 3$ . First, if  $0 \leq v_i \leq 1$  so that  $0 < xv_i < 3$ , the moment generating function can be bounded as

$$\mathbb{E}_\Lambda \exp\{xv_i(Y_i - \mathbb{E}_\Lambda Y_i)\} = \exp\{n\Lambda(B_i)[e^{xv_i} - 1 - xv_i]\} \leq \exp\left\{\frac{n\Lambda(B_i)(xv_i)^2}{2(1 - |xv_i|/3)}\right\} \leq \exp\left\{\frac{n\Lambda(B_i)x^2}{2(1 - |x|/3)}\right\},$$

using that the  $Y_i \sim Poi(n\Lambda(B_i))$  as well as (3.10) and (3.22) in [14]. Similarly, if  $-1 < v_i < 0$  so that  $0 < -xv_i < 3$  then we have from  $e^{-y} + y - 1 \leq y^2/2$  for  $-xv_i = y \in (0, 3)$  that

$$\mathbb{E}_\Lambda e^{xv_i(Y_i - \mathbb{E}_\Lambda Y_i)} = e^{n\Lambda(B_i)[e^{xv_i} - 1 - xv_i]} \leq e^{n\Lambda(B_i)(xv_i)^2/2},$$

which is also upper bound by the r.h.s. of the penultimate display. In summary, by Markov's inequality and independence of the bin-counts, for any  $z > 0$  and  $0 < x < 3$ ,

$$\begin{aligned} P_\Lambda \left( \sum_{i=1}^K v_i(Y_i - \mathbb{E}_\Lambda Y_i) > z \right) &\leq \frac{\mathbb{E}_\Lambda [e^{\sum_{i=1}^K x v_i(Y_i - \mathbb{E}_\Lambda Y_i)}]}{e^{xz}} \leq e^{-xz} \prod_{i=1}^K \exp \left\{ \frac{n\Lambda(B_i)x^2}{2(1 - |x|/3)} \right\} \\ &\leq e^{-xz + \frac{n\Lambda(0)x^2}{2-2|x|/3}} \leq e^{-\frac{z^2}{2n\Lambda(0)+2z}}, \end{aligned}$$

after choosing  $x = z/(n\Lambda(0) + z) \in (0, 1)$ . Applying this with  $z = t\sqrt{nK}/2$  (and repeating the argument with  $-v$  to deal with  $|\cdot|$ ) gives the following bound for (56)

$$6^K \sup_{v \in B_\infty(1)} P_\Lambda \left( n|v^T(\hat{\Lambda} - \Lambda)| \geq t\sqrt{nK}/2 \right) \leq e^{cK} e^{-\frac{t^2 nK/4}{2n\Lambda(0) + t\sqrt{nK}}},$$

with universal constant  $c > 0$ . Now since  $K \leq Cn$  we obtain that the last tail is bounded by

$$e^{cK} e^{-\frac{1}{4} \frac{t^2 nK}{n(2\Lambda(0) + \sqrt{\sigma t})}} \leq e^{-c'K}$$

for all  $t$  large enough, completing the proof.  $\square$

**Acknowledgements** The authors would like to thank Aleksandra Jartseva, David Klenerman and Ernest Laue for various stimulating discussions and for drawing our attention to the problem of inferring diffusivity in the presence of binding events arising in biochemistry. RN and FS were supported by ERC Advanced Grant (Horizon Europe UKRI G116786) and RN was further supported by EPSRC grant EP/V026259.

## References

- [1] Robert A. Adams and John J. F. Fournier. *Sobolev Spaces*. en. Elsevier, June 2003. ISBN: 978-0-08-054129-7.
- [2] Richard Bass. *Diffusions and Elliptic Operators*. en. Probability and its Applications. New York: Springer-Verlag, 1998. ISBN: 978-0-387-98315-8. DOI: 10.1007/b97611.
- [3] Richard F. Bass. *Stochastic Processes*. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge: Cambridge University Press, 2011. ISBN: 978-1-107-00800-7. DOI: 10.1017/CB09780511997044.
- [4] S. Basu et al. *Live-cell 3D single-molecule tracking reveals how NuRD modulates enhancer dynamics*. en. July 2021. DOI: 10.1101/2020.04.03.003178.
- [5] Eduard Belitser, Paulo Serra, and Harry van Zanten. "Rate-optimal Bayesian intensity smoothing for inhomogeneous Poisson processes". In: *Journal of Statistical Planning and Inference*. Special Issue on Bayesian Nonparametrics 166 (Nov. 2015), pp. 24–35. ISSN: 0378-3758. DOI: 10.1016/j.jspi.2014.03.009.
- [6] Kai Lai Chung and Zhongxin Zhao. *From Brownian Motion to Schrödinger's Equation*. Ed. by M. Artin et al. Vol. 312. Grundlehren der mathematischen Wissenschaften. Berlin, Heidelberg: Springer, 1995. ISBN: 978-3-642-63381-2 978-3-642-57856-4. DOI: 10.1007/978-3-642-57856-4.
- [7] S. L. Cotter, G. O. Roberts, A. M. Stuart, and D. White. "MCMC Methods for Functions: Modifying Old Algorithms to Make Them Faster". In: *Statistical Science* 28.3 (Aug. 2013). Publisher: Institute of Mathematical Statistics, pp. 424–446. ISSN: 0883-4237, 2168-8745. DOI: 10.1214/13-STS421.
- [8] E. Brian Davies. *Spectral Theory and Differential Operators*. Cambridge Studies in Advanced Mathematics. Cambridge: Cambridge University Press, 1995. ISBN: 978-0-521-58710-5. DOI: 10.1017/CB09780511623721.

- [9] Sophie Donnet, Vincent Rivoirard, Judith Rousseau, and Catia Scricciolo. “Posterior Concentration Rates for Counting Processes with Aalen Multiplicative Intensities”. en. In: *Bayesian Analysis* 12.1 (Mar. 2017). ISSN: 1936-0975. DOI: 10.1214/15-BA986.
- [10] Lawrence C. Evans. *Partial Differential Equations*. en. American Mathematical Soc., 2010. ISBN: 978-0-8218-4974-3.
- [11] G. B. Folland. *Real Analysis: Modern Techniques and Their Applications, 2nd Edition — Wiley*. en. Apr. 1999.
- [12] Mark Freidlin. *Functional integration and partial differential equations*. Vol. 109. Annals of Mathematics Studies. Princeton University Press, Princeton, NJ, 1985, pp. x+545. ISBN: 0-691-08354-1; 0-691-08362-2. DOI: 10.1515/9781400881598.
- [13] Subhashis Ghosal and Aad van der Vaart. *Fundamentals of Nonparametric Bayesian Inference*. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge: Cambridge University Press, 2017. ISBN: 978-0-521-87826-5. DOI: 10.1017/9781139029834.
- [14] Evarist Giné and Richard Nickl. *Mathematical foundations of infinite-dimensional statistical models*. en. Cambridge series in statistical and probabilistic mathematics 40. Cambridge New York, NY Melbourne Dehli Singapore: Cambridge University Press, 2016. ISBN: 978-1-107-04316-9.
- [15] Evarist Giné and Richard Nickl. “Rates of contraction for posterior distributions in  $L^r$ -metrics,  $1 \leq r \leq \infty$ ”. In: *Ann. Statist.* 39.6 (2011), pp. 2883–2911. ISSN: 0090-5364,2168-8966. DOI: 10.1214/11-AOS924.
- [16] Matteo Giordano. “Bayesian nonparametric inference in PDE models: asymptotic theory and implementation”. In: *arXiv preprint arXiv:2311.18322* (2023).
- [17] Matteo Giordano, Alisa Kirichenko, and Judith Rousseau. *Nonparametric Bayesian intensity estimation for covariate-driven inhomogeneous point processes*. arXiv:2312.14073 [math]. Dec. 2023. DOI: 10.48550/arXiv.2312.14073.
- [18] Matteo Giordano and Kolyan Ray. “Nonparametric Bayesian inference for reversible multidimensional diffusions”. In: *Ann. Statist.* 50.5 (2022), pp. 2872–2898. ISSN: 0090-5364,2168-8966. DOI: 10.1214/22-aos2213.
- [19] Matteo Giordano and Sven Wang. *Statistical algorithms for low-frequency diffusion data: A PDE approach*. arXiv:2405.01372 [stat]. Jan. 2025. DOI: 10.48550/arXiv.2405.01372.
- [20] Emmanuel Gobet, Marc Hoffmann, and Markus Reiß. “Nonparametric estimation of scalar diffusions based on low frequency data”. In: *Ann. Statist.* 32.5 (2004), pp. 2223–2253. ISSN: 0090-5364,2168-8966. DOI: 10.1214/009053604000000797.
- [21] Shota Gugushvili, Frank van der Meulen, Moritz Schauer, and Peter Spreij. *Fast and scalable nonparametric Bayesian inference for Poisson point processes*. arXiv:1804.03616 [stat]. Mar. 2020.
- [22] Alec Heckert, Liza Dahal, Robert Tjian, and Xavier Darzacq. “Recovering mixtures of fast-diffusing states from short single-particle trajectories”. In: *eLife* 11 (Sept. 2022). Ed. by Ihor Smal, Anna Akhmanova, and Maarten W Paul. Publisher: eLife Sciences Publications, Ltd, e70169. ISSN: 2050-084X. DOI: 10.7554/eLife.70169.
- [23] Mathias Luidor Heltberg, Judith Miné-Hattab, Angela Taddei, Aleksandra M. Walczak, and Thierry Mora. *Physical observables to determine the nature of membrane-less cellular sub-compartments*. en. Pages: 2021.04.01.438041 Section: New Results. Apr. 2021. DOI: 10.1101/2021.04.01.438041.
- [24] Marc Hoffmann and Kolyan Ray. “Nonparametric Bayesian estimation in a multidimensional diffusion model with high frequency data”. en. In: *Probability Theory and Related Fields* (Oct. 2024). ISSN: 1432-2064. DOI: 10.1007/s00440-024-01317-w.
- [25] Alisa Kirichenko and Harry van Zanten. “Optimality of Poisson Processes Intensity Learning with Gaussian Processes”. In: *Journal of Machine Learning Research* 16.91 (2015), pp. 2909–2919. ISSN: 1533-7928.
- [26] Günter Last and Mathew Penrose. *Lectures on the Poisson Process*. Institute of Mathematical Statistics Textbooks. Cambridge: Cambridge University Press, 2017. ISBN: 978-1-107-08801-6. DOI: 10.1017/9781316104477.
- [27] J. L. Lions and E. Magenes. *Non-Homogeneous Boundary Value Problems and Applications*. Berlin, Heidelberg: Springer Berlin Heidelberg, 1972. ISBN: 978-3-642-65163-2 978-3-642-65161-8. DOI: 10.1007/978-3-642-65161-8.

- [28] P. L. Lions and A. S. Sznitman. “Stochastic differential equations with reflecting boundary conditions”. en. In: *Communications on Pure and Applied Mathematics* 37.4 (1984), pp. 511–537. ISSN: 1097-0312. DOI: 10.1002/cpa.3160370408.
- [29] Judith Miné-Hattab and Angela Taddei. “Physical principles and functional consequences of nuclear compartmentalization in budding yeast”. en. In: *Current Opinion in Cell Biology* 58 (June 2019), pp. 105–113. ISSN: 09550674. DOI: 10.1016/j.ceb.2019.02.005.
- [30] François Monard, Richard Nickl, and Gabriel P. Paternain. “Consistent inversion of noisy non-Abelian X-Ray transforms”. en. In: *Communications on Pure and Applied Mathematics* 74.5 (May 2021), pp. 1045–1099. ISSN: 0010-3640, 1097-0312. DOI: 10.1002/cpa.21942.
- [31] François Monard, Richard Nickl, and Gabriel P. Paternain. “Efficient nonparametric Bayesian inference for X-ray transforms”. In: *The Annals of Statistics* 47.2 (Apr. 2019). Publisher: Institute of Mathematical Statistics, pp. 1113–1147. ISSN: 0090-5364, 2168-8966. DOI: 10.1214/18-AOS1708.
- [32] Richard Nickl. *Bayesian Non-linear Statistical Inverse Problems*. en. ISBN: 9783985470532 9783985475537 ISSN: 2943-4963, 2943-4971. June 2023. DOI: 10.4171/zlam/30.
- [33] Richard Nickl. “Consistent inference for diffusions from low frequency measurements”. In: *Ann. Statist.* 52.2 (2024), pp. 519–549. ISSN: 0090-5364,2168-8966. DOI: 10.1214/24-aos2357.
- [34] Richard Nickl and Kolyan Ray. “Nonparametric statistical inference for drift vector fields of multi-dimensional diffusions”. In: *Ann. Statist.* 48.3 (2020), pp. 1383–1408. ISSN: 0090-5364,2168-8966. DOI: 10.1214/19-AOS1851.
- [35] Richard Nickl and Jakob Söhl. “Nonparametric Bayesian posterior contraction rates for discretely observed scalar diffusions”. In: *Ann. Statist.* 45.4 (2017), pp. 1664–1693. ISSN: 0090-5364,2168-8966. DOI: 10.1214/16-AOS1504.
- [36] Richard Nickl, Sara Van De Geer, and Sven Wang. “Convergence Rates for Penalized Least Squares Estimators in PDE Constrained Regression Problems”. en. In: *SIAM/ASA Journal on Uncertainty Quantification* 8.1 (Jan. 2020), pp. 374–413. ISSN: 2166-2525. DOI: 10.1137/18M1236137.
- [37] Richard Nickl and Sven Wang. “On polynomial-time computation of high-dimensional posterior measures by Langevin-type algorithms”. en. In: *Journal of the European Mathematical Society* 26.3 (Dec. 2022), pp. 1031–1112. ISSN: 1435-9855. DOI: 10.4171/jems/1304.
- [38] L. C. G. Rogers and David Williams. *Diffusions, Markov processes, and martingales. Vol. 1*. Cambridge Mathematical Library. Foundations, Reprint of the second (1994) edition. Cambridge University Press, Cambridge, 2000, pp. xx+386. ISBN: 0-521-77594-9. DOI: 10.1017/CB09781107590120.
- [39] A. M. Stuart. “Inverse problems: A Bayesian perspective”. en. In: *Acta Numerica* 19 (May 2010), pp. 451–559. ISSN: 1474-0508, 0962-4929. DOI: 10.1017/S0962492910000061.
- [40] Hiroshi Tanaka. “Stochastic differential equations with reflecting boundary condition in convex regions”. In: *Hiroshima Mathematical Journal* 9.1 (Jan. 1979). Publisher: Hiroshima University, Mathematics Program, pp. 163–177. ISSN: 0018-2079, 2758-9641. DOI: 10.32917/hmj/1206135203.
- [41] Michael E. Taylor. *Partial Differential Equations I: Basic Theory*. en. Springer Science & Business Media, Oct. 2010. ISBN: 978-1-4419-7055-8.

DEPARTMENT OF PURE MATHEMATICS & MATHEMATICAL STATISTICS  
 UNIVERSITY OF CAMBRIDGE, Cambridge, UK  
 Email: nickl@maths.cam.ac.uk, fps25@cam.ac.uk