

THE BETTI NUMBERS OF KUNZ-WALDI SEMIGROUPS

MARIO GONZÁLEZ-SÁNCHEZ, SRISHTI SINGH, AND HEMA SRINIVASAN

ABSTRACT. Given two coprime numbers $p < q$, KW semigroups contain p, q and are contained in $\langle p, q, r \rangle$ where $2r = p, q, p + q$ whichever is even. These semigroups were first introduced by Kunz and Waldi. Kunz and Waldi proved that all KW semigroups of embedding dimension $n \geq 4$ have Cohen-Macaulay type $n - 1$ and first Betti number $\binom{n}{2}$. In this paper, we characterize KW semigroups whose defining ideal is generated by the 2×2 minors of a $2 \times n$ matrix. In addition, we identify all KW semigroups that lie on the interior of the same face of the Kunz cone \mathcal{C}_p as a KW semigroup with determinantal defining ideal. Thus, we provide an explicit formula for the Betti numbers of all those KW semigroups.

INTRODUCTION

Let $3 \leq p < q$ be two relatively prime integers. In [6], Kunz and Waldi build numerical semigroups with multiplicity p by filling in the gaps of the symmetric semigroup $\langle p, q \rangle$ in a systematic way. In particular, they characterize all numerical semigroups $\langle p, q \rangle \subsetneq H \subset \langle p, q, r \rangle$, where $r = p/2$ if p is even, $r = q/2$ if q is even, and $r = (p + q)/2$ otherwise. Such semigroups H are in one-to-one correspondence to the lattice paths, with right and downward steps, in the rectangle $R \subset \mathbb{R}^2$ with vertices $(0, 0)$, $(0, p' - 1)$, $(q' - 1, p' - 1)$, and $(q' - 1, 0)$, where $p' = \lfloor p/2 \rfloor$ and $q' = \lfloor q/2 \rfloor$. We call these semigroups Kunz-Waldi semigroups and denote by $KW(p, q)$ the class formed by all of them. In fact, a semigroup $H = \langle p, q, h_1, \dots, h_{n-2} \rangle \in KW(p, q)$ if and only if $h_i = pq - x_i p - y_i q$ satisfying $0 < x_1 < \dots < x_{n-2} \leq q/2$ and $p/2 \geq y_1 > \dots > y_{n-2} > 0$, as observed in Remark 1 of [8], and the embedding dimension of H is $e(H) = n$.

Let $H = \langle p, q, h_1, \dots, h_{n-2} \rangle \in KW(p, q)$ and fix a field \mathbb{k} . Consider the polynomial ring $T = \mathbb{k}[u, v, u_1, \dots, u_{n-2}]$ graded by H via $\deg(u) = p$, $\deg(v) = q$ and $\deg(u_i) = h_i$ for all $1 \leq i \leq n - 2$, and let t be another unknown. Define $\phi_H : T \rightarrow \mathbb{k}[t]$ by $u \mapsto t^p$, $v \mapsto t^q$, $u_i \mapsto t^{h_i}$, $1 \leq i \leq n - 2$. The *defining ideal of H* is $I_H = \ker \phi_H$, and it is homogeneous, prime,

2020 *Mathematics Subject Classification*. Primary 13D02, 13D05; Secondary 20M14.

The first author has been supported in part by the grant PID2022-137283NB-C22 funded by MICIU/AEI/10.13039/501100011033 and by ERDF/EU, and thanks financial support from European Social Fund, *Programa Operativo de Castilla y León*, and *Consejería de Educación de la Junta de Castilla y León*.

This collaboration is the outcome of the first author's visit to the University of Missouri during Fall 2024.

The third author is supported by a grant from Simons Foundation.

and binomial. In line with the convention of calling a numerical semigroup H Gorenstein or Complete Intersection when I_H is Gorenstein or Complete Intersection, we say H is determinantal if the ideal I_H is a determinantal ideal. We know from the theorem in Appendix A of [7] that I_H is minimally generated by the following $\binom{n}{2}$ homogeneous binomials:

$$\begin{aligned} f_{ij} &= u_i u_j - u^{q-x_i-x_j} v^{p-y_i-y_j}, \quad 1 \leq i \leq j \leq n-2 \\ g_i &= v^{y_i-y_{i+1}} u_i - u^{x_{i+1}-x_i} u_{i+1}, \quad 1 \leq i \leq n-3 \\ \eta_1 &= v^{p-y_1} - u^{x_1} u_1 \\ \eta_2 &= v^{y_{n-2}} u_{n-2} - u^{q-x_{n-2}}. \end{aligned}$$

Since I_H is homogeneous, the semigroup algebra $\mathbb{k}[H] \simeq T/I_H$ is a H -graded module, and we can consider a minimal graded free resolution of T/I_H ,

$$0 \rightarrow T^{\beta_{n-1}} \rightarrow T^{\beta_{n-2}} \rightarrow \dots \rightarrow T^{\beta_1} \rightarrow T \rightarrow T/I_H \rightarrow 0.$$

The numbers β_i , $0 \leq i \leq n-1$, are the *Betti numbers* of T/I_H . Note that $\beta_0 = 1$, $\beta_1 = \binom{n}{2}$ is the number of elements in any minimal generating set of I_H formed by homogeneous polynomials, and $\beta_{n-1} = n-1$ by Corollary 3.1 of [7].

Given a numerical semigroup $S \subset \mathbb{N}$, let $m = \min(S \setminus \{0\})$ be its multiplicity, and write

$$\text{Ap}(S) = \{s \in S \mid s - m \notin S\} = \{0, a_1, \dots, a_{m-1}\}$$

for the *Apéry set* of S with respect to m , where for all $1 \leq i \leq m-1$, $a_i \in S$ is the minimum element in S congruent to i modulo m . The *Apéry coordinate vector* of S with respect to m is the tuple (a_1, \dots, a_{m-1}) . For each $m \in \mathbb{Z}_{>0}$, $m \geq 2$, the *Kunz cone* $\mathcal{C}_m \subset \mathbb{R}_{\geq 0}^{m-1}$ is a pointed cone with defining inequalities $a_i + a_j \geq a_{i+j}$ whenever $i + j \neq 0$, where $i, j \in \mathbb{Z}_m \setminus \{0\}$. If S_1, S_2 are two numerical semigroups of multiplicity m whose Apéry coordinate vectors lie on the interior of the same face of the Kunz cone \mathcal{C}_m , then they have the same Betti numbers by Theorem 2.7 of [1].

The study of determinantal ideals is a well-developed area in commutative algebra, as these ideals provide extensive information about the structure of the ring. In particular, their minimal free resolutions are completely understood. In [4] and [9], the authors give criteria for a numerical semigroup of embedding dimension n to have the defining ideal generated by the 2×2 minors of a $2 \times n$ matrix in terms of Pseudo-Frobenius numbers, in some specific cases. In [8], the authors examine the defining ideals of KW semigroups and show that the ideal can be written as a sum of determinantal ideals (Theorem 4.1, [8]). However, the results did not extend beyond this, and a key motivation was to determine whether any KW semigroups are determinantal.

In this paper, we characterize the semigroups $H \in KW(p, q)$ whose defining ideal is minimally generated by the 2×2 minors of a $2 \times n$ matrix, in terms of the Pseudo-Frobenius numbers of H (Proposition 1.2), and later we use the Kunz cone C_p to compute the Betti numbers of a larger class. These determinantal semigroups form a subclass $KW_D(p, q)$ of $KW(p, q)$, defined as follows: a semigroup $H = \langle p, q, h_1, \dots, h_{n-2} \rangle \in KW(p, q)$ is in $KW_D(p, q)$ if there exist $x, y \in \mathbb{Z}_{>0}$ with $(n-2)x \leq q/2$, $(n-2)y \leq p/2$, and $x_i = ix$, $y_i = (n-1-i)y$ for all $1 \leq i \leq n-2$. In Proposition 1.2 we prove that the 2×2 minors of the matrix

$$\begin{bmatrix} u_{n-2} & u^x & v^{p-(n-1)y} & u_1 & u_2 & \cdots & u_{n-4} & u_{n-3} \\ u^{q-(n-1)x} & v^y & u_1 & u_2 & u_3 & \cdots & u_{n-3} & u_{n-2} \end{bmatrix}$$

generate the ideal I_H for a semigroup $H \in KW_D(p, q)$, and in Theorem 1.4 we prove that the semigroups in $KW_D(p, q)$ are the only ones in $KW(p, q)$ whose defining ideal is determinantal. For any semigroup $H \in KW_D(p, q)$, the Eagon-Northcott complex [3] resolves the ideal I_H . Hence, their Betti numbers are given by $\beta_i = i \binom{n}{i+1}$ for all $1 \leq i \leq n-1$. Moreover, for two semigroups $H, G \in KW(p, q)$, we characterize when G and H lie on the interior of the same face of the Kunz cone C_p in Proposition 2.7. As a consequence, we obtain a larger subclass of semigroups of $KW(p, q)$ whose Betti numbers are given by $\beta_i = i \binom{n}{i+1}$, $1 \leq i \leq n-1$.

1. WHEN IS A $KW(p, q)$ - SEMIGROUP DETERMINANTAL?

The goal of this section is to completely characterize all Kunz-Waldi numerical semigroups whose defining ideals are determinantal. To this end, we give a necessary and sufficient condition for any $H \in KW(p, q)$ to have a determinantal defining ideal I_H (1.4) and provide a matrix A such that $I_H = I_{2 \times 2}(A)$ (1.2), i.e. I_H is generated by the 2×2 minors of A .

Definition 1.1. *Let $3 \leq p < q$ be relatively prime. We define a subclass $KW_D(p, q)$ of $KW(p, q)$ as follows. A semigroup $H = \langle p, q, h_1, \dots, h_{n-2} \rangle \in KW(p, q)$ is in $KW_D(p, q)$ if there exist $x, y \in \mathbb{Z}_{>0}$ such that $(n-2)x \leq q/2$, $(n-2)y \leq p/2$, and $h_i = pq - x_i p - y_i q$, where $x_i = ix$ and $y_i = (n-1-i)y$ for all $1 \leq i \leq n-2$.*

The notation chosen to denote this class will be justified after Theorem 1.4, where we prove that the semigroups in $KW_D(p, q)$ are precisely all the semigroups in $KW(p, q)$ with determinantal defining ideal.

Remark 1. *Let $H \in KW_D(p, q)$ for some $x, y \in \mathbb{Z}_{>0}$. Note that $h_{i+1} - h_i = yq - xp$ for all $1 \leq i \leq n-3$. Thus, the generators h_1, h_2, \dots, h_{n-2} necessarily form an arithmetic sequence with common difference $yq - xp$. Note that this sequence is increasing if $yq > xp$ and decreasing otherwise.*

Proposition 1.2. *If $H \in KW_D(p, q)$ for some $x, y, n \in \mathbb{Z}_{>0}$ then I_H is generated by the 2×2 minors of the following $2 \times n$ matrix*

$$A = \begin{bmatrix} u_{n-2} & u^x & v^{p-(n-1)y} & u_1 & u_2 & \cdots & u_{n-4} & u_{n-3} \\ u^{q-(n-1)x} & v^y & u_1 & u_2 & u_3 & \cdots & u_{n-3} & u_{n-2} \end{bmatrix}.$$

Proof. Let a_{ij} denote the determinant of the submatrix of A obtained from the i - and j -th columns, and $I_2(A)$ be the ideal generated by all the a_{ij} . Each a_{ij} is a H -homogeneous binomial and thus, is in I_H . Therefore, $I_2(A) \subset I_H$. Let us prove the other inclusion. Note that for all $1 \leq i \leq n-3$, $g_i = -a_{2,i+3}$; $\eta_1 = -a_{23}$, and $\eta_2 = a_{12}$, so all of them are in $I_2(A)$. To finish the proof, it suffices to show that $f_{ij} \in I_2(A)$ for all $1 \leq i \leq j \leq n-2$. We have $f_{1,n-2} = a_{13} \in I_2(A)$ and for all $1 \leq j \leq n-3$, one can check that

$$f_{1,j} = -a_{3,j+3} - \sum_{k=0}^{n-j-4} (u^{kx} v^{p-(n+k)y} a_{2,j+4+k}) + u^{(n-j-3)x} v^{p-(2n-j-3)y} a_{12} \in I_2(A).$$

Moreover, for all $2 \leq i \leq n-2$,

$$f_{i,n-2} = a_{1,i+2} + \sum_{k=0}^{i-2} (u^{q-(n+k)x} v^{ky} a_{2,i+1-k}) \in I_2(A).$$

Finally, note that for all $2 \leq i \leq j \leq n-3$, $f_{i,j} = f_{i-1,j+1} - a_{i+2,j+3}$, and hence all the f_{ij} belong to $I_2(A)$. \square

When Kunz and Waldi initially construct these numerical semigroups of multiplicity p obtained by adjoining arbitrary gaps of $\langle p, q \rangle$, no conditions are imposed on the gaps. In this initial work, they discuss the set of Pseudo-Frobenius elements of these semigroups (page 668, [6]). Later, they prove that if the gaps are chosen so that the resulting numerical semigroups belong to $KW(p, q)$, then their type is one less than the embedding dimension (Example 4.6, [7]). From this, one can deduce the complete set of Pseudo-Frobenius elements for any $H \in KW(p, q)$, as we will illustrate below.

Lemma 1.3. *Let $H \in KW(p, q)$ of embedding dimension n . Then $PF(H) = \{g_i := pq - (x_i + 1)p - (y_{i+1} + 1)q \mid 0 \leq i \leq n-2\}$, where $x_0 = y_{n-1} = 0$.*

Proof. Set $x_0 = y_{n-1} = 0$ and let $\mathcal{H} = \{g_i := pq - (x_i + 1)p - (y_{i+1} + 1)q \mid 0 \leq i \leq n-2\}$. Recall from the introduction that $H = \langle p, q, h_1, \dots, h_{n-2} \rangle$ is in one-to-one correspondence to a lattice path l in the rectangle R . In particular, each h_i corresponds to the point $(x_i - 1, y_i - 1)$ under the line $pq - p - q = px + qy$ ([6]). To see that $\mathcal{H} = PF(H)$, first note that $\mathcal{H} = \{h \in \mathbb{N} \setminus H \mid h + p \in H, h + q \in H\}$. This is clear from Figure 1. Hence, $PF(H) \subset \mathcal{H}$, and the equality $PF(H) = \mathcal{H}$ follows from the fact $|\mathcal{H}| = n-1 = |PF(H)|$ (the type of H is $n-1$). \square




FIGURE 1. Pseudo Frobenius Elements of $H \in KW(p, q)$ correspond to the points \times .

Theorem 1.4. *Let $H \in KW(p, q)$ of embedding dimension n . Then the following are equivalent:*

- (1) $H \in KW_D(p, q)$.
- (2) *There exist positive integers x, y such that $(n - 2)y \leq p/2$, $(n - 2)x \leq q/2$ and I_H is generated by the 2×2 minors of the matrix*

$$\begin{bmatrix} u_{n-2} & u^x & v^{p-(n-1)y} & u_1 & u_2 & \cdots & u_{n-4} & u_{n-3} \\ u^{q-(n-1)x} & v^y & u_1 & u_2 & u_3 & \cdots & u_{n-3} & u_{n-2} \end{bmatrix}.$$

- (3) *There exist H -homogeneous polynomials $F_i, G_i \in (u, v, u_1, \dots, u_{n-2})$ such that I_H is generated by the 2×2 minors of the matrix*

$$\begin{bmatrix} F_1 & F_2 & \cdots & F_n \\ G_1 & G_2 & \cdots & G_n \end{bmatrix}.$$

- (4) $\text{PF}(H) = \{z + k, z + 2k, \dots, z + (n - 1)k\}$ for some $z \geq 0$ and $k > 0$.

When this is the case, $\text{PF}(H) = \{pq - (ix + 1)p - ((n - i - 2)y + 1)q \mid 0 \leq i \leq n - 2\}$, for the numbers $x, y \in \mathbb{Z}_{>0}$ defining $H \in KW_D(p, q)$, and hence $k = |xp - yq|$.

Proof. (1) \Rightarrow (2) is Proposition 1.2, (2) \Rightarrow (3) is trivial and (3) \Rightarrow (4) is proved in general for any numerical semigroup in Section 2 of [9].

(4) \Rightarrow (1) By Lemma 1.3, $PF(H) = \{g_i := pq - (x_i + 1)p - (y_{i+1} + 1)q \mid 0 \leq i \leq n - 2\}$, where $x_0 = y_{n-1} = 0$. Since $PF(H)$ satisfies (4) by our hypothesis, we must have that the difference between any two consecutive elements is constant. Firstly, for any $0 \leq i \leq n - 2$,

$$\begin{aligned} g_i - g_{i+1} &= pq - (x_i + 1)p - (y_{i+1} + 1)q - [pq - (x_{i+1} + 1)p - (y_{i+2} + 1)q] \\ &= p(x_{i+1} - x_i) + q(y_{i+2} - y_{i+1}). \end{aligned}$$

Let $\alpha_i = x_{i+1} - x_i$ and $\beta_i = y_{i+2} - y_{i+1}$. Now we must have that for any $0 \leq i < j \leq n - 3$,

$$p\alpha_i + q\beta_i = p\alpha_j + q\beta_j \implies p(\alpha_i - \alpha_j) = q(\beta_j - \beta_i).$$

Since p and q are relatively prime, there exists $l \in \mathbb{Z}$ such that $\alpha_i - \alpha_j = ql$. But $|\alpha_i - \alpha_j| = |x_{i+1} + x_j - x_i - x_{j+1}| \leq q - 2$, so $l = 0$. Thus, since $x_{i+1} > x_i$, there is some $x \in \mathbb{Z}_{>0}$ such that $\alpha_i = \alpha_j = x$. The recursive definition of α_i now gives us $x_i = ix$ for $1 \leq i \leq n - 2$. If $x > \frac{q}{2(n-2)}$, then $x_{n-2} = (n-2)x > q/2$, a contradiction.

Similarly, $\beta_i = \beta_j = -y$, for some $y \in \mathbb{Z}_{>0}$ since $y_{i+2} < y_{i+1}$. This implies $y_{n-2} = y, y_{n-3} = 2y, \dots, y_1 = (n-2)y$. If $y > \frac{p}{2(n-2)}$ then $y_1 > p/2$, a contradiction.

We have shown that H must be as in Definition 1.1. In addition, when (4) holds, we have shown that $PF(H) = \{pq - (ix + 1)p - ((n - i - 2)y + 1)q \mid 0 \leq i \leq n - 2\}$, which implies $k = |xp - yq|$. \square

Remark 2. Let $3 \leq p < q$ relatively prime integers and denote $p' = \lfloor p/2 \rfloor$, and $q' = \lfloor q/2 \rfloor$. By Theorem 1.4, the cardinality of $KW_D(p, q)$ is

$$|KW_D(p, q)| = \sum_{n=3}^{p'+2} \left\lfloor \frac{p'}{n-2} \right\rfloor \left\lfloor \frac{q'}{n-2} \right\rfloor = \sum_{n=1}^{p'} \left\lfloor \frac{p'}{n} \right\rfloor \left\lfloor \frac{q'}{n} \right\rfloor,$$

and the cardinality of $KW(p, q)$ is

$$|KW(p, q)| = \sum_{n=3}^{p'+2} \binom{p'}{n-2} \binom{q'}{n-2} = \sum_{n=1}^{p'} \binom{p'}{n} \binom{q'}{n} = \binom{p' + q'}{p'} - 1.$$

Thus, the proportion of semigroups in $KW(p, q)$ whose defining ideal is determinantal is

$$\rho_D(p, q) := \frac{|KW_D(p, q)|}{|KW(p, q)|} = \frac{\sum_{n=1}^{p'} \left\lfloor \frac{p'}{n} \right\rfloor \left\lfloor \frac{q'}{n} \right\rfloor}{\binom{p' + q'}{p'} - 1}.$$

2. BETTI SEQUENCE OF ANY $H \in KW(p, q)$

Since the ideal $I_H \subset T = \mathbb{k}[u, v, u_1, \dots, u_{n-2}]$ for any $H \in KW_D(p, q)$ is generated by the 2×2 minors of a $2 \times n$ matrix and its height is $n - 1$ ($= n - 2 + 1$), it is resolved by the Eagon-Northcott complex (Theorem 2 of [3]). In particular, if the minimal graded free resolution of T/I_H is

$$0 \rightarrow T^{\beta_{n-1}} \rightarrow T^{\beta_{n-2}} \rightarrow \dots \rightarrow T^{\beta_1} \rightarrow T \rightarrow T/I_H \rightarrow 0,$$

then the Betti numbers are given by the formula $\beta_i = i \binom{n}{i+1}$ for all $1 \leq i \leq n-1$, by the construction of the Eagon-Northcott complex. We want to expand this to compute the Betti sequence of more KW-semigroups. One way to do this is by using the fact that any two numerical semigroups of multiplicity p lying in the interior of the same face of the Kunz cone \mathcal{C}_p have the same Betti sequence.

In this section, we identify all $H \in KW(p, q)$ that lie on the interior of the same face of \mathcal{C}_p as some $H' \in KW_D(p, q)$. To achieve this, we compute the Apéry posets of $H \in KW(p, q)$ and use the fact that two semigroups lie on the same face of \mathcal{C}_p if and only if they have the same Apéry posets.

We first recall how to construct the Apéry set and poset of any semigroup $H \in KW(p, q)$ from the lattice path defining H .

Proposition 2.1. *Let $H = \langle p, q, h_1, \dots, h_{n-2} \rangle \in KW(p, q)$ be the semigroup defined by the sequences $x_1 < x_2 < \dots < x_{n-2}$ and $y_1 > y_2 > \dots > y_{n-2}$, and set $y_{n-1} := 0$. The Apéry set of H is*

$$\text{Ap}(H) = \{\lambda q \mid 0 \leq \lambda < p - y_1\} \cup \left(\bigcup_{i=1}^{n-2} \{h_i + \lambda q \mid 0 \leq \lambda < y_i - y_{i+1}\} \right).$$

Proof. Let A be the Apéry set of the numerical semigroup $\langle p, q \rangle$, $A = \{0, q, 2q, \dots, (p-1)q\}$. We can obtain the Apéry set $\text{Ap}(H)$ from A as follows. Recall that

$$H = \langle p, q \rangle \cup \{pq - xp - yq \mid x \leq x_i, y \leq y_i \text{ for some } 1 \leq i \leq n-2\}.$$

Note that $pq - xp - yq \equiv -yq \pmod{p}$, so the elements that we have to replace in A are the ones congruent to $-yq$ modulo p for $0 < y \leq y_1$. For each one of these congruence classes, we choose the smallest element in H , i.e. the one with the largest x . This corresponds to the element in the lattice path defining H that has Y -coordinate $y-1$. Therefore,

$$\text{Ap}(H) = \{\lambda q \mid 0 \leq \lambda < p - y_1\} \cup \left(\bigcup_{i=1}^{n-3} \{h_i + \lambda q \mid 0 \leq \lambda < y_i - y_{i+1}\} \right) \cup \{h_{n-2} + \lambda q \mid 0 \leq \lambda < y_{n-2}\}.$$

□

Remark 3. *By Corollary 3.10 of [5], $\text{Ap}(H)$ is homogeneous since the variable u appears in all non-homogeneous binomials of a minimal generating set of I_H . This means that for every element $z \in \text{Ap}(H)$ all the factorizations of z have the same length.*

We now consider a poset structure on $\text{Ap}(H)$.

Definition 2.2. *Let S be a numerical semigroup with multiplicity m , and write $\text{Ap}(S) = \{a_0 := 0, a_1, \dots, a_{m-1}\}$ so that $a_i \equiv i \pmod{m}$ for all i . The Apéry poset of S is a poset $\mathcal{P}(S) = (\mathbb{Z}_m, \preceq)$, where $i \preceq j$ if and only if $a_j - a_i \in S$ for $i, j \in \mathbb{Z}_m$. We write $i \prec j$ and say j covers i if $i \preceq j$ and there is no k such that $i \prec k \prec j$.*

Proposition 2.3. For all $i, j \in \mathbb{Z}_m$, $i \prec j$ if and only if $a_j - a_i$ is a minimal generator of S .

Proof. Being (\Leftarrow) trivial, let us prove (\Rightarrow) . Let $i, j \in \mathbb{Z}_m$ such that $i \prec j$ and write $a_j = a_i + \alpha + \beta$ for some $\alpha, \beta \in H$ with α a minimal generator of H . Note that $a_i \prec a_i + \alpha \preceq a_j$. Since $a_j \in \text{Ap}(H)$, then $a_i + \alpha \in \text{Ap}(H)$, so $a_j = a_i + \alpha$ as j covers i . \square

Theorem 2.4. (Theorem 3.10, [2]) Any two numerical semigroups with the same multiplicity m belong to the interior of the same face of \mathcal{C}_m if and only if their Apéry posets are the same.

Example 2.5. Consider the semigroups $H = \langle 8, 17, 60, 69, 78 \rangle$ and $G = \langle 8, 17, 53, 62, 55 \rangle$ with Apéry sets $\text{Ap}(H) = \{0, 17, 34, 51, 60, 69, 78, 95\}$ and $\text{Ap}(G) = \{0, 17, 34, 51, 68, 53, 62, 55\}$. Their Apéry posets are shown in Figure 2.




FIGURE 2. Apéry posets in Example 2.5.

Now consider a semigroup $H \in KW(p, q)$ and let us see how its Apéry poset look like. For $0 \leq \lambda < p - y_1$ set $i_{0,\lambda}$ the label of λq in $\mathcal{P}(H)$, i.e. $0 \leq i_{0,\lambda} < p$ and $i_{0,\lambda} \equiv \lambda q \pmod{p}$. Similarly, for each $1 \leq j \leq n - 2$ and $0 \leq \lambda < y_j - y_{j+1}$, let $i_{j,\lambda}$ be the label of $h_j + \lambda q$ in $\mathcal{P}(H)$.

Proposition 2.6. Let $3 \leq p < q$ be relatively prime and $H \in KW(p, q)$.

(1) For all $i_{j_1, k_1}, i_{j_2, k_2} \in \mathbb{Z}_p$,

$$i_{j_1, k_1} \prec i_{j_2, k_2} \Leftrightarrow \begin{cases} j_2 = j_1 & \text{and } k_2 = k_1 + 1, \text{ or} \\ j_1 = 0 & \text{and } k_2 = k_1. \end{cases}$$

Thus, the Hasse diagram of $\mathcal{P}(H)$ is the one shown in Figure 3.

(2) The poset $\mathcal{P}(H)$ is graded for the rank function $\rho : \text{Ap}(H) \rightarrow \mathbb{N}$ defined by $\rho(i_{j,k}) = j + k$.

Proof. Note that $(\lambda + 1)q - \lambda q = q$, $(h_j + (\lambda + 1)q) - (h_j + \lambda q) = q$, $(h_j + \lambda q) - \lambda q = h_j$, so $i_{j,\lambda} \prec i_{j,\lambda+1}$ and $i_{0,\lambda} \prec i_{j,\lambda}$ for all j, λ by Proposition 2.3. Let us prove that there are no more covering relations. By Remark 3 and Proposition 2.3, it suffices to prove that $(h_i + (\lambda + 1)q) - (h_j + \lambda q) = h_i - h_j + q$ is not a minimal generator of H when $i \neq j$ and $\lambda \geq 0$. Note that $h_i - h_j + q \neq q$ since $i \neq j$, and $h_i - h_j + q \neq p$ because $h_i + q, h_j \in \text{Ap}(H)$. Now suppose that $h_i - h_j + q = h_k$ for some $1 \leq k \leq n - 2$, $k \neq i, j$. Then

$$(x_j + x_k - x_i)p + (y_j + y_k - y_i + 1)q = pq.$$

Thus, q divides $x_j + x_k - x_i$ and since $3 - q/2 \leq x_j + x_k - x_i \leq q - 2$, then $x_j + x_k - x_i = 0$, so $x_i > x_j$ and hence $i > j$. With a similar argument, one can prove that $y_i = y_j + y_k + 1 > y_j$, so $i < j$, a contradiction. This completes the proof of part (1), and part (2) is a direct consequence of part (1) and Remark 3. \square




FIGURE 3. Hasse diagram of the Apéry poset of a semigroup $H \in KW(p, q)$.

Let $H, G \in KW(p, q)$ be of the same embedding dimension n . Write $H = \langle p, q, h_1, \dots, h_{n-2} \rangle$ and $G = \langle p, q, g_1, \dots, g_{n-2} \rangle$. Further, write $h_i = pq - x_i p - y_i q$ and $g_i = pq - z_i p - w_i q$, $1 \leq i \leq n - 2$.

Proposition 2.7. *Let $H, G \in KW(p, q)$. Then H and G belong to the interior of the same face of the Kunz cone \mathcal{C}_p if and only if*

- (1) $e(H) = e(G) = n$, and
- (2) $y_i = w_i$ for all $1 \leq i \leq n - 2$.

Proof. By Proposition 2.6, the Apéry poset of any semigroup $H \in KW(p, q)$ is completely determined by the embedding dimension of H , n , and the sequence $p/2 \geq y_1 > \dots > y_{n-2} > 0$. Thus, the result follows from Theorem 2.4. \square

Corollary 2.8. *Let $H \in KW(p, q)$ of embedding dimension n be such that $y_i = (n - i - 1)y$ for some $y \in \mathbb{N}$ with $(n - 2)y \leq p/2$. Then the Betti numbers of $\mathbb{k}[H]$ are*

$$\beta_i = i \binom{n}{i+1}, \quad 1 \leq i \leq n - 1.$$

Proof. Consider the semigroup $H' \in KW_D(p, q)$ defined by the sequence $x_1 = 1 < x_2 = 2 < \dots < x_{n-2} = n - 2$ and the sequence $y_1 > y_2 > \dots > y_{n-2}$, $y_i = (n - 1 - i)y$. By Proposition 2.7, H and H' lie in the interior of the same face of the Kunz cone. Therefore, $\mathbb{k}[H]$ and $\mathbb{k}[H']$ have the same Betti sequence by Theorem 2.7 of [1], that is, $\beta_i = i \binom{n}{i+1}$, $1 \leq i \leq n - 1$ by the construction of the Eagon-Northcott complex for $I_{H'}$. \square

Example 2.9. *In Example 2.5, G is not in $KW_D(8, 17)$ as 53, 62, 55 are not in an arithmetic sequence. However, G is in the same face of \mathcal{C}_8 as $G' = \langle 8, 17, 69, 70, 71 \rangle$. Note that for both G and G' , $y_1 = 3, y_2 = 2, y_3 = 1$. Their Betti sequence is 4, 15, 20, 10, 1, and following is the Hasse diagram of $\mathcal{P}(G)$ and $\mathcal{P}(G')$:*




FIGURE 4. $\mathcal{P}(G)$ and $\mathcal{P}(G')$

Remark 4. *By Corollary 2.8, the number of numerical semigroups in $KW(p, q)$ whose Betti numbers are $\beta_i = i \binom{n}{i+1}$, $1 \leq i \leq n - 1$, is at least $\sum_{n=1}^{p'} \binom{p'}{n} \binom{q'}{n}$, where $p' = \lfloor p/2 \rfloor$ and $q' = \lfloor q/2 \rfloor$.*

It is interesting to note that even if $H \in KW(p, q)$ does not satisfy the hypothesis of Corollary 2.8, the conclusion still seems to hold. In particular, if the y_1, \dots, y_{n-2} are not in an arithmetic sequence, is it still true that $\beta_i = i \binom{n}{i+1}$ for $1 \leq i \leq n - 1$? We give an example to support a positive answer:

Example 2.10. $H = \langle 8, 17, 36, 45, 63 \rangle \in KW(8, 17) \setminus KW_D(8, 17)$. *The Betti sequence of $\mathbb{k}[H]$ is 4, 15, 20, 10, 1, and yet, H is not even in the same face as some $H' \in KW_D(8, 17)$.*

This is just one of many examples but we have yet to find a KW-semigroup of embedding dimension $n - 1$, whose Betti numbers are not $\beta_i = i \binom{n}{i+1}$, $1 \leq i \leq n - 1$. This is clearly

true for embedding dimension 3. With this in mind, we end this article with the following conjecture:

Conjecture. *The Betti numbers of any numerical semigroup $H \in KW(p, q)$ of embedding dimension n are $\beta_i = i \binom{n}{i+1}$, $1 \leq i \leq n - 1$.*

REFERENCES

- [1] Benjamin Braun et al. “Minimal free resolutions of numerical semigroup algebras via Apéry specialization”. In: *Pacific Journal of Mathematics* 334.2 (2025), pp. 211–231. DOI: <https://doi.org/10.2140/pjm.2025.334.211>.
- [2] Winfried Bruns et al. “Wilf’s conjecture in fixed multiplicity”. In: *International Journal of Algebra and Computation* 30.04 (2020), pp. 861–882. DOI: <https://doi.org/10.1142/S021819672050023X>.
- [3] John A Eagon and Douglas Geoffrey Northcott. “Ideals defined by matrices and a certain complex associated with them”. In: *Proceedings of the Royal Society of London. Series A. Mathematical and Physical Sciences* 269.1337 (1962), pp. 188–204. DOI: <https://doi.org/10.1098/rspa.1962.0170>.
- [4] Shiro Goto et al. “Pseudo-Frobenius numbers versus defining ideals in numerical semigroup rings”. In: *Journal of Algebra* 508 (2018), pp. 1–15. ISSN: 0021-8693. DOI: <https://doi.org/10.1016/j.jalgebra.2018.04.025>.
- [5] Raheleh Jafari and Santiago Zarzuela Armengou. “Homogeneous numerical semigroups”. In: *Semigroup Forum* 97.2 (2018), pp. 278–306. ISSN: 1432-2137. DOI: <https://doi.org/10.1007/s00233-018-9941-6>.
- [6] E. Kunz and R. Waldi. “Geometrical illustration of numerical semigroups and of some of their invariants”. In: *Semigroup Forum* 89.3 (June 2014), pp. 664–691. ISSN: 1432-2137. DOI: <http://dx.doi.org/10.1007/s00233-014-9599-7>.
- [7] E. Kunz and R. Waldi. “On the deviation and the type of certain local Cohen–Macaulay rings and numerical semigroups”. In: *Journal of Algebra* 478 (2017), pp. 397–409. ISSN: 0021-8693. DOI: <https://doi.org/10.1016/j.jalgebra.2017.01.041>.
- [8] Srishti Singh and Hema Srinivasan. “A Class of Numerical Semigroups Defined by Kunz and Waldi - Their Principal Matrices and Structure”. In: *Journal of Algebra and Its Applications* (Feb. 2025). ISSN: 1793-6829. DOI: <http://dx.doi.org/10.1142/S0219498825410130>.
- [9] Do Van Kien and Naoyuki Matsuoka. “Numerical Semigroup Rings of Maximal Embedding Dimension with Determinantal Defining Ideals”. In: *Numerical Semigroups : IMNS 2018*. Ed. by Valentina Barucci et al. Cham: Springer International Publishing, 2020,

pp. 185–196. ISBN: 978-3-030-40822-0. DOI: https://doi.org/10.1007/978-3-030-40822-0_12.

INSTITUTO DE INVESTIGACIÓN EN MATEMÁTICAS DE LA UNIVERSIDAD DE VALLADOLID (IMUVA),
UNIVERSIDAD DE VALLADOLID, 47011 VALLADOLID, SPAIN.

Email address: `mario.gonzalez.sanchez@uva.es`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MISSOURI, COLUMBIA MO 65211, USA.

Email address: `spkdq@missouri.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MISSOURI, COLUMBIA MO 65211, USA.

Email address: `srinivasanh@missouri.edu`