

Martingale property and moment explosions in signature volatility models

Eduardo Abi Jaber^{*1}, Paul Gassiat^{†2}, and Dimitri Sotnikov^{‡1,3}

¹École Polytechnique, CMAP

²CEREMADE, Université Paris Dauphine-PSL & DMA, ENS Paris & Institut Universitaire de France

³Engie Global Markets

March 24, 2025

Abstract

We study the martingale property and moment explosions of a signature volatility model, where the volatility process of the log-price is given by a linear form of the signature of a time-extended Brownian motion. Excluding trivial cases, we demonstrate that the price process is a true martingale if and only if the order of the linear form is odd and a correlation parameter is negative. The proof involves a fine analysis of the explosion time of a signature stochastic differential equation. This result is of key practical relevance, as it highlights that, when used for approximation purposes, the linear combination of signature elements must be taken of odd order to preserve the martingale property. Once martingality is established, we also characterize the existence of higher moments of the price process in terms of a condition on a correlation parameter.

1 Introduction

The martingale property and moment explosions of the asset price process are fundamental considerations in stochastic volatility modeling, particularly due to their implications for derivative pricing and numerical methods. The martingale property is a key criterion for assessing the validity of a stochastic volatility model, as emphasized by [Sepp and Rakhmonov \(2023\)](#). In particular, strict local martingale models exhibit structural deficiencies, such as the failure of put-call parity and deviations in the asymptotic behavior of implied volatility wings ([Jacquier and Keller-Ressel, 2018](#)). Similarly, the existence of moments for the price process plays a crucial role in both theoretical and computational aspects of option pricing. In particular, the existence of finite moments is crucial for ensuring the numerical stability and convergence of Monte Carlo methods used to compute derivative prices and their sensitivities. Moments also play a key role in the proof of asymptotic formulae, allowing one to transition from large deviations of the price process to call price asymptotics, see for instance ([Friz et al., 2021](#), Section 4.2).

Both of these problems have been extensively studied in the literature, dating back to the works of [Girsanov \(1960\)](#) and [Novikov \(1973\)](#). However, Novikov’s criterion is often not applicable to stochastic volatility models. [Sin \(1998\)](#) observed for the first time that in certain stochastic volatility models—such as the Hull–White model—the stock price can be a strict martingale for certain parameter values. In this context, [Lions and Musiela \(2007\)](#) present general sufficient conditions for martingality and strict local martingality when the volatility process follows a stochastic differential equation (SDE), and also address the problem of moment existence. Subsequently, [Mijatović and Urusov \(2012\)](#) provide a more general criterion for martingality in SDE-based models, while [Bernard, Cui, and McLeish \(2017\)](#) extend this work to the general spot-volatility

^{*}eduardo.abi-jaber@polytechnique.edu. The first author is grateful for the financial support from the Chaires FiME-FDD, Financial Risks, Deep Finance & Statistics and Machine Learning and systematic methods in finance at École Polytechnique.

[†]gassiat@ceremade.dauphine.fr

[‡]dmitrii.sotnikov@polytechnique.edu. The author is grateful for the financial support provided by Engie Global Markets.

correlation case $-1 \leq \rho \leq 1$, offering a complete classification of convergence properties for general one-factor Markovian stochastic volatility models.

Most of these works, however, focus on Markovian models where both the asset price and volatility processes are solutions of SDEs. For example, [Jourdain \(2004\)](#) proposes a martingality criterion, along with integrability conditions for prices in SABR-like models. Similarly, [Andersen and Piterbarg \(2007\)](#) address the moment problem and provide sufficient martingality conditions for SABR-like and Heston-like models.

In contrast, results in non-Markovian settings tend to be less general and are often derived on a case-by-case basis. For instance, in the case of volatility processes governed by stochastic Volterra equations, several results have been derived. For the rough Bergomi model, [Gassiat \(2018\)](#) establishes conditions for both martingality and non-martingality, as well as a sufficient condition for moment explosions. [Gerhold, Pachschröll, and Ruf \(2024\)](#) propose an alternative approach to verifying higher-order moment explosions by examining the integrability of the price supremum and apply this method to the rough Bergomi model. In the context of the Volterra Heston model, [Abi Jaber, Larsson, and Pulido \(2019\)](#) establish conditions for the martingale property, while [Gerhold, Gerstenecker, and Pinter \(2018\)](#) study the problem of moment explosions. In the class of so-called path-dependent volatility (PDV) models ([Guyon and Lekeufack, 2023](#)), [Nutz and Riveros Valdevenito \(2024\)](#) analyze the martingale property in a two-factor PDV model. Additionally, [Ruf \(2015\)](#) provides a general martingality criterion for stochastic exponentials defined as non-anticipative path-dependent functionals of solutions to stochastic differential equations. Despite its generality, this result cannot be directly applied to stochastic volatility models and requires a thorough analysis to verify the condition for each particular model.

Recently, signature-based models have gained popularity in non-Markovian modeling. Such models were first studied by [Arribas, Salvi, and Szpruch \(2020\)](#) and further developed by [Cuchiero, Gazzani, and Svaluto-Ferro \(2023\)](#); [Cuchiero, Gazzani, Möller, and Svaluto-Ferro \(2025\)](#), and [Abi Jaber and Gérard](#). Their distinguishing feature is that the volatility process is represented as a (possibly infinite) linear combination of signature elements. These models can serve both for the approximation, see [Cuchiero et al. \(2023\)](#), and the exact representation, see [Abi Jaber and Gérard](#), of a wide range of models, though in both cases, the linear combination of signature elements must be truncated for practical use. As we will demonstrate, the truncation order plays a crucial role in determining whether the price process is a true martingale. However, no results on either martingality or moment explosions have been established for these models so far. The aim of this paper is to fill this gap by providing necessary and sufficient conditions for the martingality of the price process and by studying the finiteness of its moments in signature volatility models. We adopt the multiplicative framework of [Abi Jaber and Gérard](#); [Cuchiero et al. \(2025\)](#), in which the price process is given by a stochastic exponential, making the question of martingality non-trivial.

The Markovian setting provides key insights into the martingality problem. In particular, for truncated exponential Ornstein-Uhlenbeck models with negative spot-volatility correlation, obtained by truncating the power series expansion of the exponential, the price process is a martingale if and only if the truncation order is odd. A similar pattern appears in polynomial volatility models, where volatility is expressed as a polynomial function of a Brownian motion or an Ornstein-Uhlenbeck process, as in the Quintic model of [Abi Jaber, Illand, and Li \(2023\)](#). These results can be established using techniques similar to those in [Lions and Musiela \(2007, Theorem 2.4\(i\)\)](#). Since path signatures play an analogous role to polynomials on path space, we expect similar characterizations to hold beyond the Markovian framework.

In this work, we show that the order of the linear combination of signature elements plays the same role as the order of a polynomial and that a similar result holds for signature volatility models. Specifically, given two Brownian motions, B and W , correlated with coefficient $\rho \in [-1, 1]$, we consider the following signature volatility model for a price process S :

$$\begin{aligned} \frac{dS_t}{S_t} &= \sigma_t dB_t, \quad t \geq 0, \\ \sigma_t &= \langle \boldsymbol{\sigma}, \widehat{\mathbb{W}}_t \rangle, \end{aligned}$$

where the volatility σ_t is a finite linear combination of order N in the signature $\widehat{\mathbb{W}}_t$ of the extended Brownian motion $\widehat{W} = (t, W)$, defined as a sequence of iterated Stratonovich integrals, with the first orders given by

$$\widehat{\mathbb{W}}_t^0 = 1, \quad \widehat{\mathbb{W}}_t^1 = \begin{pmatrix} t \\ W_t \end{pmatrix}, \quad \widehat{\mathbb{W}}_t^2 = \begin{pmatrix} \frac{t^2}{2!} & \int_0^t s dW_s \\ \int_0^t W_s ds & \frac{W_t^2}{2!} \end{pmatrix}, \quad \dots$$

The martingale property. Our main result, Theorem 3.1, establishes a very simple necessary and sufficient condition for the martingality of the price process S under the assumption that the leading coefficient $\sigma^{2^{\otimes N}}$ in front of the signature element $\frac{W_t^N}{N!}$ is nonzero. More precisely, in the nontrivial case where $N \geq 2$ and $\rho \neq 0$, we have that

the price process S is a true martingale if and only if N is odd and $\rho\sigma^{2^{\otimes N}} \leq 0$.

The standard technique for verifying the martingale property reduces to studying the explosion time of a certain stochastic differential equation (SDE) following a change of probability measure. In our case, this yields a signature SDE whose explosion time is intricate to analyze, see Proposition 5.1 and Theorem 5.2. To address this, we establish two key bounds for linear combinations of signature elements in Lemmas 4.1 and 4.2. These bounds exploit the fact that the shuffle algebra forms a polynomial algebra over the Lyndon (1954) words, thanks to the Radford (1979) Theorem. Using these bounds, we then prove the finiteness or explosion of the solution of the signature SDE, depending on the parity of the order N and the sign of ρ , leading to Theorem 3.1. We stress that such result is of key practical relevance, as it highlights that, when used for approximation purposes, the linear combination of signature elements must be taken of odd order to preserve the martingale property. The non-positivity of $\rho\sigma^{2^{\otimes N}}$ does not imply that the spot-volatility correlation is necessarily non-positive, see Remark 3.1.3.

Additionally, we extend the martingale characterization to a volatility process constructed from the signature of a multidimensional Brownian motion, as presented in Theorem 3.3.

Moment explosions. Assuming the conditions for S to be a martingale, we prove in Theorem 3.2 that

*the moment $\mathbb{E}[S_T^m]$ of order m is finite for any $T > 0$ if $|\rho| > \sqrt{1 - \frac{1}{m}}$.
Conversely, if $|\rho| < \sqrt{1 - \frac{1}{m}}$, then $\mathbb{E}[S_T^m] = +\infty$ for all $T > 0$.*

This result aligns well with those obtained in the Markovian case, such as (Lions and Musiela, 2007, Theorem 2.3) and (Jourdain, 2004, Proposition 6). The same condition for moment explosions in the non-Markovian rough Bergomi model was proved in Gassiat (2018), see also Gulisashvili (2020). To prove Theorem 3.2, we rely on the Boué and Dupuis (1998) formula, which reduces the moment existence problem to the finiteness of an optimal control problem value. In the case of infinite moments, we explicitly construct a sequence of controls to show that the problem value is infinite, whereas in the case of finite moments, we establish the finiteness of the value function using the bounds on linear combinations derived earlier.

Outline. Section 2 introduces the necessary definitions and key concepts related to tensor algebra and signatures. In Section 3, we present our main results concerning the martingality criterion and moment explosions for the signature volatility model. Section 4 establishes two essential bounds on signature words, which are extensively used in the subsequent proofs. The proofs of these bounds are postponed to Section 6. Section 5 demonstrates how the martingality problem can be reduced to the explosion problem of a signature SDE. Sections 7 and 8 are dedicated to proving the martingality and strict local martingality criteria, respectively. Finally, Section 9 contains the proof concerning moment explosions.

2 Preliminary concepts of path signatures

2.1 Tensor algebra

In this section, we introduce the notation and definitions necessary for dealing with signatures of continuous semimartingales. We also refer the reader to the introductory sections in Cuchiero et al. (2025); Abi Jaber and Gérard.

Fix $d \in \mathbb{N}$. For $n \in \mathbb{N}$, let $(\mathbb{R}^d)^{\otimes n}$ denote the space of tensors of order n and $(\mathbb{R}^d)^{\otimes 0} = \mathbb{R}$. The path signatures defined below, will be defined as tensor sequences, i.e. the elements of the extended tensor algebra space $T((\mathbb{R}^d))$ over \mathbb{R}^d defined by

$$T((\mathbb{R}^d)) := \{\ell = (\ell^n)_{n=0}^\infty : \ell^n \in (\mathbb{R}^d)^{\otimes n}\}.$$

The truncated tensor algebra $T^N(\mathbb{R}^d)$ is defined as the space of sequences such that all their elements of order greater than $N \in \mathbb{N}$ are equal to zero. We will also denote by $T(\mathbb{R}^d)$ the space of all finite tensor sequences,

i.e. $T(\mathbb{R}^d) := \bigcup_{N \in \mathbb{N}} T^N(\mathbb{R}^d)$, and we note that $T(\mathbb{R}^d) \subset T((\mathbb{R}^d))$.

Let $\{e_1, \dots, e_d\} \subset \mathbb{R}^d$ denote the canonical basis of \mathbb{R}^d . Then, $(e_{i_1} \otimes \dots \otimes e_{i_n})_{(i_1, \dots, i_n) \in \{1, \dots, d\}^n}$ is a basis of $(\mathbb{R}^d)^{\otimes n}$. To simplify the notations, we identify its elements $e_{i_1} \otimes \dots \otimes e_{i_n}$ with the words $\mathbf{i}_1 \cdots \mathbf{i}_n$ of length n in the alphabet $A_d = \{\mathbf{1}, \mathbf{2}, \dots, \mathbf{d}\}$. All the words of length n will be denoted by

$$V_n := \{\mathbf{i}_1 \cdots \mathbf{i}_n : \mathbf{i}_k \in A_d \text{ for } k = 1, 2, \dots, n\}.$$

We adopt the convention of denoting by \emptyset the empty word and by $V_0 = \{\emptyset\}$ identified with the basis of $(\mathbb{R}^d)^{\otimes 0} = \mathbb{R}$. The basis of $T((\mathbb{R}^d))$ can be then identified with $V := \bigcup_{n \geq 0} V_n$, and each sequence $\ell \in T((\mathbb{R}^d))$ can be written as

$$\ell = \sum_{n=0}^{\infty} \sum_{\mathbf{v} \in V_n} \ell^{\mathbf{v}} \mathbf{v},$$

where $\ell^{\mathbf{v}} = (\ell^n)^{\mathbf{v}} \in \mathbb{R}$ is the coefficient of ℓ corresponding to the word \mathbf{v} and $n = |\mathbf{v}|$ stands for its length. The concatenation $\ell \mathbf{v}$ of elements $\ell \in T((\mathbb{R}^d))$ and the word $\mathbf{v} = \mathbf{i}_1 \cdots \mathbf{i}_n$ means $\ell \otimes e_{i_1} \otimes \dots \otimes e_{i_n}$.

We also define the bracket between $\ell \in T(\mathbb{R}^d)$ and $\mathbf{p} \in T((\mathbb{R}^d))$ by

$$\langle \ell, \mathbf{p} \rangle := \sum_{n=0}^{\infty} \sum_{\mathbf{v} \in V_n} \ell^{\mathbf{v}} \mathbf{p}^{\mathbf{v}}.$$

Note that since ℓ is finite, the sum contains only a finite number of terms, so that the bracket is always well-defined.

Another key operation on the space of tensor sequences is the *shuffle product*, which generalizes the integration by parts formula and plays a crucial role in the signature linearization property stated in Proposition 2.3 below.

Definition 2.1 (Shuffle product). *For two words $\mathbf{v}, \mathbf{w} \in V$ and two letters $\mathbf{i}, \mathbf{j} \in A_d$, the shuffle product $\sqcup: V \times V \rightarrow T(\mathbb{R}^d)$ is defined recursively by*

$$\begin{aligned} (\mathbf{vi}) \sqcup (\mathbf{wj}) &= (\mathbf{v} \sqcup (\mathbf{wj}))\mathbf{i} + ((\mathbf{vi}) \sqcup \mathbf{w})\mathbf{j}, \\ \mathbf{w} \sqcup \emptyset &= \emptyset \sqcup \mathbf{w} = \mathbf{w}. \end{aligned}$$

It extends to $T((\mathbb{R}^d))$ by linearity: for $\ell = \sum_{n=0}^{\infty} \sum_{\mathbf{v} \in V_n} \ell^{\mathbf{v}} \mathbf{v}$ and $\mathbf{p} = \sum_{n=0}^{\infty} \sum_{\mathbf{v} \in V_n} \mathbf{p}^{\mathbf{v}} \mathbf{v}$, the shuffle product is given by

$$\ell \sqcup \mathbf{p} = \sum_{n=0}^{\infty} \sum_{\substack{\mathbf{v}, \mathbf{w} \in V \\ |\mathbf{v}| + |\mathbf{w}| = n}} \ell^{\mathbf{v}} \mathbf{p}^{\mathbf{w}} \mathbf{v} \sqcup \mathbf{w}.$$

Note that the shuffle product is commutative.

The shuffle product represents all possible riffle shuffles of two separate decks of cards, where the relative order within each individual deck is preserved. For instance, $\mathbf{12} \sqcup \mathbf{3} = \mathbf{123} + \mathbf{132} + \mathbf{312}$. For more details on the shuffle product, we refer to [Ree \(1958\)](#) and [Gaines \(1994\)](#).

2.2 Signatures

Let X and Y be two semimartingales. Throughout the paper, we denote the Itô integral by $\int_0^\cdot Y_t dX_t$ and the Stratonovich integral by $\int_0^\cdot Y_t \circ dX_t$. The following relation holds:

$$\int_0^\cdot Y_t \circ dX_t = \int_0^\cdot Y_t dX_t + \frac{1}{2}[X, Y].$$

Definition 2.2 (Signature). *Let $(X_t)_{t \geq 0}$ be a continuous \mathbb{R}^d -valued semimartingale on a filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathbb{P})$ and fix $T > 0$. The signature of X is defined by*

$$\begin{aligned} \mathbb{X} : \Omega \times [0, T] &\rightarrow T((\mathbb{R}^d)) \\ (\omega, t) &\mapsto \mathbb{X}_t(\omega) := (1, \mathbb{X}_t^1(\omega), \dots, \mathbb{X}_t^n(\omega), \dots), \end{aligned}$$

where

$$\mathbb{X}_t^n := \int_{0 < u_1 < \dots < u_n < t} \circ dX_{u_1} \otimes \dots \otimes \circ dX_{u_n} \in (\mathbb{R}^d)^{\otimes n}, \quad n \geq 0, \quad t \in [0, T].$$

The signature can be seen as a set of monomials in the path-space, while linear combinations of signature elements $\langle \ell, \mathbb{X}_t \rangle$ play the role of polynomials. For instance, if $d = 1$, the signature of X is exactly the sequence of monomials $(\frac{1}{n!}(X_t - X_0)^n)_{n \in \mathbb{N}}$ and any finite combination $\langle \ell, \mathbb{X}_t \rangle$ for $\ell \in T^N(\mathbb{R}^d)$ is a polynomial of degree N in X_t .

Remark 2.2.1. Each element of the signature $\mathbb{X}_t^n = (\mathbb{X}_t^{i_1 \dots i_n})_{(i_1 \dots i_n) \in V_n}$ can be written coordinate-wise in an iterative form

$$\mathbb{X}_t^{i_1 \dots i_n} = \int_0^t \mathbb{X}_s^{i_1 \dots i_{n-1}} \circ dX_s^{i_n}.$$

2.3 Linear combinations of signature elements

For the signature volatility model, we will consider the signature $\widehat{\mathbb{X}}_t$ of the time-extended path $\widehat{X}_t := (t, X_t) \in \mathbb{R}^{d+1}$. The main object of our study is the finite linear combination in $\widehat{\mathbb{X}}_t$ given by

$$\langle \ell, \widehat{\mathbb{X}}_t \rangle = \sum_{n=0}^N \sum_{\mathbf{v} \in V_n} \ell^{\mathbf{v}} \widehat{\mathbb{X}}_t^{\mathbf{v}}, \quad \ell \in T^N(\mathbb{R}^{d+1}),$$

for some $N \in \mathbb{N}$.

Linear combinations of signature elements have a remarkable property: they allow for the linearization of polynomials formed from these combinations. This property is frequently emphasized in the literature when highlighting the signature's linearization power.

Proposition 2.3 (Shuffle property). *If $\ell_1, \ell_2 \in T(\mathbb{R}^{d+1})$, then*

$$\langle \ell_1, \widehat{\mathbb{X}}_t \rangle \langle \ell_2, \widehat{\mathbb{X}}_t \rangle = \langle \ell_1 \sqcup \ell_2, \widehat{\mathbb{X}}_t \rangle.$$

Proof. See (Lyons, Caruana, and Lévy, 2007, Theorem 2.15). □

3 Main results

Let $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathbb{P})$ be a filtered probability space supporting a two-dimensional correlated Brownian motion (B, W) with correlation parameter $\rho \in [-1, 1]$ and with the filtration generated by this Brownian motion. We set $\bar{\rho} = \sqrt{1 - \rho^2}$ and, for $\rho \in (-1, 1)$, we define

$$B^\perp := \frac{W - \rho B}{\bar{\rho}}, \quad W^\perp := \frac{B - \rho W}{\bar{\rho}},$$

the Brownian motions independent from B and W respectively.

We consider a signature volatility model in the sense of [Abi Jaber and Gérard](#), see also [Cuchiero, Gazzani, Möller, and Svaluto-Ferro \(2025\)](#). In this model, the stock price process S under the risk-neutral probability \mathbb{P} is given by a stochastic volatility model where the volatility process σ_t is a finite linear combination of the elements of signature $\widehat{\mathbb{W}}_t$ of the time extended Brownian motion $\widehat{W}_t := (t, W_t)$:

$$\begin{aligned} \frac{dS_t}{S_t} &= \sigma_t dB_t, \quad t \geq 0, \\ \sigma_t &= \langle \boldsymbol{\sigma}, \widehat{\mathbb{W}}_t \rangle. \end{aligned} \tag{3.1}$$

Here $\boldsymbol{\sigma} \in T^N(\mathbb{R}^2)$ is a tensor sequence of order $N \in \mathbb{N}$ in the form

$$\boldsymbol{\sigma} = \sum_{k \geq 0} \sum_{|\mathbf{v}|=k} \sigma^{\mathbf{v}} \mathbf{v}, \quad \sigma^{\mathbf{v}} \in \mathbb{R},$$

We note that

$$\mathbb{E} \left[\int_0^T \sigma_t^2 dt \right] = \left\langle \sigma^{\perp \perp 2} \mathbf{1}, \mathbb{E}[\widehat{\mathbb{W}}_T] \right\rangle,$$

is finite, as the right-hand side involves a sum of a finite number of terms where the expected signature $\mathbb{E}[\widehat{\mathbb{W}}_T]$ is given by Fawcett's formula [Fawcett \(2003\)](#) extended to the time-augmented Brownian motion in ([Lyons and Victoir, 2004](#), Proposition 4.10). This ensures the existence of a unique strong solution to (3.1) given by the exponential local martingale

$$S_t = S_0 \exp \left\{ -\frac{1}{2} \int_0^t \sigma_s^2 ds + \int_0^t \sigma_s dB_s \right\}, \quad t \geq 0.$$

We note that the volatility dynamics in the model (3.1) are driven by the signature $\widehat{\mathbb{W}}_t$ of a one-dimensional Brownian motion. We will first characterize the martingale property in Subsection 3.1, then study moment explosions in Subsection 3.2, before extending the characterization of the martingale property to volatility dynamics driven by a multidimensional Brownian motion in Subsection 3.3.

3.1 Martingale property

Our main result in this section provides a necessary and sufficient condition for the price process S , defined by (3.1), to be a martingale. This condition is given in terms of the parity of the signature level N and the correlation parameter ρ .

Theorem 3.1. *Let S be given by (3.1). Suppose that the leading coefficient $\sigma^{2^{\otimes N}} \neq 0$.*

- (i) *If $\rho = 0$ or $N = 1$, then S is a martingale.*
- (ii) *If $\rho \neq 0$ and $N \geq 2$, then S is a martingale if and only if the order N is odd and $\rho \sigma^{2^{\otimes N}} \leq 0$.*

Proof. (i) and sufficiency in (ii) are proved in Section 7 and necessity in (ii) is proved in Subsection 8.1. \square

The results of Theorem 3.1 can be illustrated numerically on implied volatility surfaces. Namely, if the spot price S is a strict local martingale, the put-call parity is not verified, and hence, put-option implied volatility is different from the call-option implied volatility which may be not even defined, as shown by [Jacquier and Keller-Ressel \(2018\)](#). More precisely, recall that the Black-Scholes Call and Put option prices $C_{\text{BS}}(T, K, \sigma)$ and $P_{\text{BS}}(T, K, \sigma)$ satisfy the put-call parity

$$C_{\text{BS}}(T, K, \sigma) - P_{\text{BS}}(T, K, \sigma) = S_0 - K,$$

while the model Call and Put option prices $C(T, K)$ and $P(T, K)$ satisfy only

$$C(T, K) - P(T, K) = \mathbb{E}[S_T] - K.$$

Thus, if $P(T, K) = P_{\text{BS}}(T, K, \sigma_{\text{BS}}^P)$, one obtains

$$C(T, K) = C_{\text{BS}}(T, K, \sigma_{\text{BS}}^P) + \mathbb{E}[S_T] - S_0 = C_{\text{BS}}(T, K, \sigma_{\text{BS}}^C).$$

Hence, the call and put implied volatilities coincide if and only if the put-call parity is satisfied, which holds if and only if $\mathbb{E}[S_T] = S_0$. In turn, this is equivalent to the martingality of S , since S is a positive local martingale and therefore a supermartingale.

We compute the put and call implied volatility smiles via the Monte Carlo method using 10^6 trajectories of the spot price given by (3.1) for even ($N = 4$) and odd ($N = 5$) orders and for positive ($\rho = 0.9$) and negative ($\rho = -0.9$) values of the correlation parameter. The maturity is taken equal to $T = 1$. The coefficients σ are chosen randomly (same random numbers were used in all cases) from the uniform distribution on $[-0.5, 0.5]$, except for the coefficient $\sigma^{2^{\otimes N}}$ fixed equal to 1.

Simulation results are provided in Figure 1. As expected, the only case where call and put implied volatilities coincide corresponds to the odd order $N = 5$ and negative correlation $\rho = -0.9$.




Figure 1: The put (brown) and call (dark green) implied volatility smiles for $N = 4$, $\rho = 0.9$ (upper-left), $N = 4$, $\rho = -0.9$ (upper-right), $N = 5$, $\rho = 0.9$ (lower-left), and $N = 5$, $\rho = -0.9$ (lower-right). The confidence intervals with confidence level of 95% do not intersect anywhere except for the forth plot $N = 5$, $\rho = -0.9$, corresponding to the martingale spot price.

Remark 3.1.1. The characterization of Theorem 3.1 still holds if, instead of $\widehat{\mathbb{W}}_t$, we consider the signature of a time-extended Ornstein–Uhlenbeck process $\widehat{Y}_t = (t, Y_t)$ for the volatility process, where Y satisfies

$$dY_t = aY_t dt + b dW_t, \quad t \geq 0,$$

for some $a, b \in \mathbb{R}$. Although slight modifications in the proof of Lemma 4.2 are needed, the proof of this extension follows the same ideas presented later in this paper. The case of the Ornstein–Uhlenbeck process is important for practical applications as it includes the Quintic model of [Abi Jaber, Illand, and Li \(2023\)](#) and can be seen as a particular case of the multi-factor signature volatility model proposed by [Cuchiero, Gazzani, Möller, and Svaluto-Ferro \(2025\)](#).

Remark 3.1.2. Theorem 3.1 provides useful insights into the practical use of the signature volatility model (3.1). Specifically, when performing a linear regression of the volatility process against the driving Brownian motion, using its exact signature representation, or calibrating the model, a suitable truncation order must be chosen. Our result offers clear and simple guidelines for selecting hyperparameters: N should be chosen to be odd, while the condition on $\rho\sigma^{2^{\otimes N}}$ can be easily incorporated as a constraint in the optimization problem. Failing to do so could have dramatic consequences for martingality, as illustrated in Figure 1.

Remark 3.1.3. We note that the condition $\rho\sigma^{2^{\otimes N}} \leq 0$ does not imply that the spot-volatility correlation must be negative. Indeed, as shown in ([Abi Jaber and Gérard](#), Subsection 3.2), the spot-volatility correlation is

given by

$$\frac{d[\log S, |\sigma|]_t}{\sqrt{d[\log S]_t} \sqrt{d[|\sigma|]_t}} = \rho \cdot \text{sign} \left(\left\langle \sigma|_{\mathbf{2}}, \widehat{\mathbb{W}}_t \right\rangle \right),$$

where $\sigma|_{\mathbf{2}} = \sum_{n=0}^N \sum_{|\mathbf{v}| \leq n} \sigma^{\mathbf{v}2} \mathbf{v}$. Since the sign depends on the other coefficients of σ , the spot-volatility correlation can be positive while still ensuring the price martingality. In this way, the model can be applied to markets with a positive implied volatility skew, such as FX or commodity markets.

3.2 Moments

In this subsection, we present our main result concerning the moment explosions of the process S in the signature volatility model (3.1). Under the assumptions ensuring true martingality stated in Theorem 3.1, we identify the threshold value of m for which $\mathbb{E}[S_T^m]$ is finite.

Theorem 3.2. *Let S be given by (3.1). Suppose that the order $N \geq 2$ is odd and $\rho \sigma^{2^{\otimes N}} < 0$.*

(i) *If $|\rho| < \sqrt{1 - \frac{1}{m}}$, then $\mathbb{E}[S_T^m] = +\infty$ for any $T > 0$.*

(ii) *If $|\rho| > \sqrt{1 - \frac{1}{m}}$, then $\mathbb{E}[S_T^m] < +\infty$ for any $T > 0$.*

Proof. The proof is given in Section 9. □

3.3 Extension of the martingality result to multidimensional setting

Theorem 3.1 was stated for the signature volatility model (3.1), where the volatility process was constructed based on the signature of a one-dimensional Brownian motion W correlated with B . Consider now a model

$$\begin{aligned} \frac{dS_t}{S_t} &= \sigma_t dB_t, \quad t \geq 0, \\ \sigma_t &= \left\langle \sigma, \widehat{\mathbb{Y}}_t \right\rangle, \end{aligned} \tag{3.2}$$

where $\widehat{\mathbb{Y}}_t$ denotes the signature of $\widehat{Y} = (t, B, Z^1, \dots, Z^d)$, $Z = (Z^1, \dots, Z^d)$ is a d -dimensional Brownian motion independent from B and $\sigma \in T^N(\mathbb{R}^{d+2})$. Theorem 3.1 can be generalized as follows.

Theorem 3.3. *Let S be given by (3.2). Suppose that the leading coefficient $\sigma^{2^{\otimes N}} \neq 0$.*

(i) *If the order $N = 1$, then S is a martingale.*

(ii) *If the order $N \geq 2$, then S is a martingale if and only if N is odd and $\sigma^{2^{\otimes N}} \leq 0$.*

Proof. (i) and sufficiency in (ii) are proved in Section 7, while necessity in (ii) is proved in Subsection 8.2. □

Remark 3.3.1. The price model (3.1) is indeed a special case of the model (3.2), since the elements of the signature of $\widehat{W} = (t, W) = (t, \rho B + \sqrt{1 - \rho^2} B^\perp)$ can be written as a linear combination of elements of the signature of $\widehat{Y} = (t, B, B^\perp)$, such that $\left\langle \sigma, \widehat{\mathbb{W}}_t \right\rangle = \left\langle \tilde{\sigma}, \widehat{\mathbb{Y}}_t \right\rangle$ for some $\tilde{\sigma} \in T^N(\mathbb{R}^3)$ with the same order N . Moreover, the coefficient of $\tilde{\sigma}$ corresponding to $\mathbf{2}^{\otimes N}$ equals $\rho^N \sigma^{2^{\otimes N}}$.

Remark 3.3.2. Although Theorem 3.1 directly follows from Theorem 3.3 when $\rho \neq 0$, we chose to present the multidimensional framework after the one-dimensional one for several reasons. First, the model in (3.1) is more natural and efficient in practice, as it uses a two-dimensional alphabet instead of the three-dimensional one used in its multidimensional counterpart. Moreover, the link between Theorems 3.1 and 3.2 and their Markovian counterparts, is much clearer when the one-dimensional model is introduced first. Finally, the methods used in the proof of Theorem 3.2 rely heavily on the structure of the equations in model (3.1) and would be difficult to generalize to the multidimensional framework.

4 Key lemmas for bounds on the signatures

In this section, we state some bounds on signature elements which will be of crucial use throughout the rest of the paper. The first result states that terms in the signature of $\widehat{X}_t = (t, X_t)$ may be bounded in terms of the supremum norm of X .

Lemma 4.1. *Let $X: [0, T] \rightarrow \mathbb{R}$ be a continuous path, and let \widehat{X}_t denote the signature of \widehat{X} . Let $N_2(\mathbf{w})$ denote the number of occurrences of letter **2** in \mathbf{w} . Then, for any word \mathbf{w} , there exists a constant $C_{\mathbf{w}} > 0$, such that*

$$\sup_{t \in [0, T]} \left| \left\langle \mathbf{w}, \widehat{X}_t \right\rangle \right| \leq C_{\mathbf{w}} T^{|\mathbf{w}| - N_2(\mathbf{w})} \max_{t \in [0, T]} |X_t|^{N_2(\mathbf{w})}. \quad (4.1)$$

Proof. The proof is given in Subsection 6.2. □

We then consider the signature of the extended process $\widehat{Y}_t^X = (t, X_t, Z_t^1, \dots, Z_t^d)$, where $Z_t = (Z_t^1, \dots, Z_t^d)$ is a d -dimensional Brownian motion and X is an arbitrary Itô process. While the terms of the signature no longer satisfy an almost sure bound as in the previous lemma, we show that a similar result still holds in a probabilistic sense.

Lemma 4.2. *Let $Z = (Z^1, \dots, Z^d)$ be a d -dimensional Brownian motion, and let X be an Itô process defined on the same probability space and such that $\langle Z^i, X \rangle \equiv 0$ for any $i = 1, \dots, d$. Let \widehat{Y}_t^X denote the signature of \widehat{Y}_t^X . Then, for any word \mathbf{w} , any $q \geq 2$, $\epsilon > 0$, and any stopping time τ taking values in $[0, T]$, there exists a constant $C = C_{\mathbf{w}, q, \epsilon, T}$ such that*

$$\mathbb{E} \left[\int_0^\tau \left\langle \mathbf{w}, \widehat{Y}_t^X \right\rangle^q dt \right] \leq C \cdot \mathbb{E} \left[\int_0^\tau X_t^{q(N_2(\mathbf{w}) + \epsilon)} dt \right], \quad (4.2)$$

where $N_2(\mathbf{w})$ denotes the number of occurrences of the letter **2** in \mathbf{w} .

Proof. The proof is given in Subsection 6.3. □

Remark 4.2.1. The statement of Lemma 4.2 remains true if, for some $C_0 > 0$ and for any $i = 1, \dots, d$,

$$|d\langle Z^i, X \rangle_t| \leq C_0 dt.$$

In this case, the same proof is applicable since the cross-variation term appearing in the bounds can be controlled in exactly the same way as the drift term.

5 A general martingality criterion

In this section, we derive a general criterion to establish the martingality of the price process in signature volatility models. Similarly to the Markovian setting, lack of martingality is seen by a Girsanov transform to be equivalent to the explosion of a certain SDE of signature drift form.

We let S be given by (3.2), i.e. the volatility process is given by

$$\sigma_t = \left\langle \boldsymbol{\sigma}, \widehat{Y}_t \right\rangle, \quad t \geq 0$$

where \widehat{Y}_t is the signature of the process $\widehat{Y}_t = (t, B_t, Z_t^1, \dots, Z_t^d)$, and $\boldsymbol{\sigma} \in T^N(\mathbb{R}^{d+2})$ for some $N \geq 1$.

The criterion will be stated, in Theorem 5.2, in terms of the solution to the SDE

$$X_t = \int_0^t \left\langle \boldsymbol{\sigma}, \widehat{Y}_s^X \right\rangle ds + B_t, \quad (5.1)$$

where \widehat{Y}_t^X denotes the signature of $\widehat{Y}_t^X = (t, X_t, Z_t^1, \dots, Z_t^d)$.

5.1 Signature SDE

In this subsection, we show the signature SDE (5.1) is well-defined (up to an explosion time) for arbitrary σ in the truncated tensor algebra $T^N(\mathbb{R}^{d+2})$.

We note that X may explode in finite time, so the solution is understood as a process satisfying equation (5.1) on $[[0, \tau_\infty^X[[$, where the explosion time τ_∞^X is a stopping time defined by

$$\tau_\infty^X := \lim_{n \rightarrow \infty} \tau_n^X, \quad \tau_n^X := \inf\{t \geq 0: |X_t| = n\},$$

such that, on the event $\{\tau_\infty^X < \infty\}$, we have $\lim_{t \rightarrow \tau_\infty^X} |X_t| = +\infty$ a.s.

Proposition 5.1. *The SDE (5.1) admits a unique strong solution. In addition, it holds that*

$$\tau_\infty^X = \inf \left\{ t \geq 0, \int_0^t \langle \sigma, \widehat{Y}_s^X \rangle^2 ds = +\infty \right\}. \quad (5.2)$$

Proof. For the well-posedness of the SDE, we note that it can be equivalently written as a system of classical (Markovian) SDEs. Indeed, setting $\widehat{Y}_t^{X, \mathbf{w}} = \langle \mathbf{w}, \widehat{Y}_t^X \rangle$, the signature SDE (5.1) is equivalent to $X = \widehat{Y}^{X, \mathbf{2}}$, where the $(\widehat{Y}^{X, \mathbf{w}})_{\mathbf{w}: |\mathbf{w}| \leq N}$ satisfy $\widehat{Y}^{X, \emptyset} \equiv 1$ and for $|\mathbf{u}| < N$,

$$\begin{aligned} \widehat{Y}_t^{X, \mathbf{u1}} &= \int_0^t \widehat{Y}_s^{X, \mathbf{u}} ds, \\ \widehat{Y}_t^{X, \mathbf{u2}} &= \int_0^t \widehat{Y}_s^{X, \mathbf{u}} \left(\sum_{|\mathbf{w}| \leq N} \sigma^{\mathbf{w}} \widehat{Y}_s^{X, \mathbf{w}} \right) ds + \int_0^t \widehat{Y}_s^{X, \mathbf{u}} \circ dB_s, \\ \widehat{Y}_t^{X, \mathbf{ui}} &= \int_0^t \widehat{Y}_s^{X, \mathbf{u}} \circ dZ_s^{i-2}, \quad i = 3, \dots, d+2. \end{aligned}$$

This is a Stratonovich SDE with locally Lipschitz coefficients, for which existence and uniqueness up to an explosion time are standard.

It remains to prove (5.2). Since on the event $\{\tau_\infty^X < \infty\}$, we have $\int_0^{\tau_\infty^X} \langle \sigma, \widehat{Y}_s^X \rangle ds = X_{\tau_\infty^X} - B_{\tau_\infty^X} = \pm\infty$, it follows that

$$\int_0^{\tau_\infty^X} \langle \sigma, \widehat{Y}_s^X \rangle^2 ds \geq \left(\int_0^{\tau_\infty^X} \langle \sigma, \widehat{Y}_s^X \rangle ds \right)^2 = +\infty,$$

we have $\int_0^{\tau_\infty^X} \langle \sigma, \widehat{Y}_s^X \rangle^2 ds = +\infty$ on $\{\tau_\infty^X < \infty\}$. Conversely, using Lemma 4.2, we obtain that for each word \mathbf{w} ,

$$\mathbb{E} \left[\int_0^{\tau_n^X \wedge T} \langle \mathbf{w}, \widehat{Y}_t^X \rangle^2 dt \right] \leq C_{\mathbf{w}, T} n^{2|\mathbf{w}|+1} < \infty.$$

Since $\tau_\infty^X = \sup_{n \geq 1} \tau_n^X$, this implies that $\int_0^t \langle \sigma, \widehat{Y}_s^X \rangle^2 ds < \infty$ for each $t < \tau_\infty^X$. \square

5.2 Martingality and explosion

By the results of the previous subsection, equation (5.1) admits a unique strong solution up to an explosion time τ_∞^X , satisfying (5.2).

Theorem 5.2. *For each $T > 0$, for S defined by (3.2), it holds that*

$$S \text{ is a martingale on } [0, T] \Leftrightarrow \mathbb{P}(\tau_\infty^X < T) = 0,$$

where τ_∞^X is the explosion time of X defined by (5.1).

Proof. Since S is a nonnegative local martingale, it is a supermartingale by Fatou's Lemma, so that $\mathbb{E}[S_T] \leq S_0$ for all $T \geq 0$. Hence, for a given $T > 0$, it is a martingale on $[0, T]$ if and only if $\mathbb{E}[S_T] = S_0$. Let

$$\tau_n = \inf \left\{ t > 0: \int_0^t \langle \sigma, \widehat{Y}_s \rangle^2 ds = n \right\}.$$

As the stopped process $\left(\int_0^{t \wedge \tau_n} \langle \sigma, \widehat{Y}_s \rangle^2 ds \right)_{t \geq 0}$ is bounded by n , the stopped price process $S^{\tau_n} = (S_{t \wedge \tau_n})_{t \geq 0}$ is an exponential martingale by Novikov's criterion and it follows that

$$S_0 = \mathbb{E}[S_T^{\tau_n}] = \mathbb{E}[S_{T \wedge \tau_n}] = \mathbb{E}[S_T \mathbf{1}_{\{T \leq \tau_n\}}] + \mathbb{E}[S_{\tau_n} \mathbf{1}_{\{\tau_n < T\}}] = \mathbb{E}[S_T \mathbf{1}_{\{T \leq \tau_n\}}] + S_0 \widehat{\mathbb{P}}_n(\tau_n < T),$$

where $\frac{d\widehat{\mathbb{P}}_n}{d\mathbb{P}} = \frac{S_{T \wedge \tau_n}}{S_0}$.

By the monotone convergence theorem, the term $\mathbb{E}[S_T \mathbf{1}_{\{T \leq \tau_n\}}]$ converges to $\mathbb{E}[S_T]$ as $n \rightarrow \infty$. Thus,

$$S_0 - \mathbb{E}[S_T] = \lim_{n \rightarrow \infty} \mathbb{E}[S_{\tau_n} \mathbf{1}_{\{\tau_n < T\}}] = \lim_{n \rightarrow \infty} S_0 \widehat{\mathbb{P}}_n(\tau_n < T).$$

On the other hand, by Girsanov theorem, for each n , it holds that for each $t \leq T$,

$$B_t = \int_0^{t \wedge \tau_n} \langle \sigma, \widehat{Y}_s \rangle ds + \widehat{B}_t,$$

where \widehat{B} is a Brownian motion under $\widehat{\mathbb{P}}_n$. This implies that, under $\widehat{\mathbb{P}}_n$, B is a solution to (5.1) on $[[0, \tau_n]]$, so that by (weak) uniqueness for (5.1), the law of $(B_{t \wedge \tau_n})_{t \geq 0}$ under $\widehat{\mathbb{P}}_n$ is the same as the law of $(X_{t \wedge \tilde{\tau}_n})_{t \geq 0}$ under \mathbb{P} , with

$$\tilde{\tau}_n = \inf \left\{ t > 0: \int_0^t \langle \sigma, \widehat{Y}_s^X \rangle^2 ds = n \right\},$$

so that

$$\lim_{n \rightarrow \infty} \widehat{\mathbb{P}}_n(\tau_n < T) = \lim_{n \rightarrow \infty} \mathbb{P}(\tilde{\tau}_n < T) = \mathbb{P}(\tau_\infty^X < T),$$

since $\lim_{n \rightarrow \infty} \tilde{\tau}_n = \tau_\infty^X$.

□

Corollary 5.3. *The same proof applied to the one-dimensional model implies that S given by (3.1) is a martingale on $[0, T]$ if and only if $\mathbb{P}(\tau_\infty^X < T)$, where τ_∞^X is the explosion time of X defined by*

$$X_t = \rho \int_0^t \langle \sigma, \widehat{X}_s \rangle ds + W_t, \tag{5.3}$$

and \widehat{X}_t denotes the signature of $\widehat{X}_t = (t, X_t)$, for $t \in [0, T]$.

Remark 5.3.1. Theorem 5.2 is closely related to the result proved in (Kazamaki, 2006, Lemma 1.5) rewritten in terms of new probability measures. A more general criterion was proposed in (Ruf, 2015, Theorem 3.3). It states that the price process is a martingale if and only if

$$\widehat{\mathbb{P}} \left(\int_0^{T \wedge \vartheta} \sigma_s^2 ds < \infty \right) = 1, \quad T \geq 0,$$

where ϑ stands for the explosion time and $\widehat{\mathbb{P}}$ denotes the extension of $(\widehat{\mathbb{P}}_n)_{n \geq 1}$ which is constructed in (Ruf, 2015, Proposition 3.1). However, this approach assumes a specific structure of the underlying probability space as highlighted in (Ruf, 2015, Example 4.3). For this reason, we prefer to work with Theorem 5.2, ensuring that our results are independent of the choice of probability space.

6 Proof of bounds on signature elements

6.1 Polynomial representations in the shuffle algebra

In this subsection, we recall the notion of Lyndon (1954) words which will help us prove the bounds given by Lemma 4.1 and Lemma 4.2. The extended tensor algebra $T((\mathbb{R}^d))$ can be considered as an algebra endowed with a shuffle product. We define a shuffle multivariate polynomial P^{\sqcup} of n arguments as

$$P^{\sqcup} : T((\mathbb{R}^d))^n \rightarrow T((\mathbb{R}^d)), \quad (\mathbf{v}_1, \dots, \mathbf{v}_n) \mapsto \sum_{p_1, \dots, p_n} c_{p_1, \dots, p_n} \mathbf{v}_1^{\sqcup p_1} \sqcup \mathbf{v}_2^{\sqcup p_2} \dots \sqcup \mathbf{v}_n^{\sqcup p_n},$$

where the sum is finite and taken over some set of positive natural numbers p_1, \dots, p_n .

The Lyndon words defined below are useful to construct the polynomial representations of the shuffle algebra elements.

Definition 6.1. *A Lyndon word in an alphabet A_d is a non-empty word that is strictly lexicographically greater than any of its non-trivial cyclic rotations. A set of all Lyndon words in the alphabet A_d will be denoted by $\mathcal{L}(A_d)$, and $\mathcal{L}_N(A_d)$ will denote the Lyndon words of length not greater than N .*

Example 6.1.1. The first Lyndon words in the alphabets A_2 and A_3 are given by

- $\mathcal{L}(A_2) = \{2, 1, 21, 221, 211, 2221, 2211, 2111, \dots\}$.
- $\mathcal{L}(A_3) = \{3, 2, 1, 32, 31, 21, 332, 331, 322, 321, 312, 311, 221, 211, \dots\}$.

An important property of the Lyndon words is given by the Radford (1979) Theorem. It states that the Lyndon words generate the shuffle algebra, i.e. each element $\ell \in T(\mathbb{R}^d)$ can be represented by a shuffle polynomial P_ℓ^{\sqcup} :

$$\ell = P_\ell^{\sqcup}(\mathbf{v}_1, \dots, \mathbf{v}_{n_\ell}), \quad \mathbf{v}_1, \dots, \mathbf{v}_{n_\ell} \in \mathcal{L}.$$

Moreover, if $\ell \in T^N(\mathbb{R}^d)$, then $\mathbf{v}_1, \dots, \mathbf{v}_{n_\ell} \in \mathcal{L}_N$ and $\sum_{i=1}^{n_\ell} p_i |\mathbf{v}_i| \leq N$ for all terms $\mathbf{v}_1^{\sqcup p_1} \sqcup \dots \sqcup \mathbf{v}_{n_\ell}^{\sqcup p_{n_\ell}}$ in the polynomial P_ℓ^{\sqcup} .

The following example provides explicit polynomial representations for the words over alphabet $\{1, 2\}$ of length up to 3 that can be easily verified.

Example 6.1.2. 1. $\mathcal{L}_1 = \{1, 2\}$ contains all the words of length 1.

2. $\mathcal{L}_2 = \{1, 2, 21\}$. The remaining words of length 2 can be obtained as the following polynomials:

$$\begin{aligned} 11 &= \frac{1}{2} \cdot 1^{\sqcup 2}, \\ 12 &= 1 \sqcup 2 - 21, \\ 22 &= \frac{1}{2} \cdot 2^{\sqcup 2}. \end{aligned}$$

3. $\mathcal{L}_3 = \{1, 2, 21, 211, 221\}$. The words of length 2 were constructed above. The remaining words of length 3 can be obtained as polynomials:

$$\begin{aligned} 111 &= \frac{1}{6} \cdot 1^{\sqcup 3}, & 122 &= \frac{1}{2} \cdot 1 \sqcup 2^{\sqcup 2} - 21 \sqcup 2 + 221, \\ 112 &= 11 \sqcup 2 - 21 \sqcup 1 + 211, & 212 &= 21 \sqcup 2 - 2 \cdot 221, \\ 121 &= 21 \sqcup 1 - 2 \cdot 211, & 222 &= \frac{1}{6} \cdot 2^{\sqcup 3}. \end{aligned}$$

We note that no Lyndon word ends with the lexicographically greatest letter \mathbf{d} , except for the word \mathbf{d} itself. Together with Radford's theorem, and noting that the letters in the alphabet can be ordered arbitrarily, this implies the following result.

Proposition 6.2. *Let \mathbf{w} be a word in A_d , and fix $\mathbf{k} \in A_d$. Then \mathbf{w} can be expressed as a shuffle polynomial*

$$\mathbf{w} = P_{\mathbf{w}}^{\sqcup}(\mathbf{v}_1, \dots, \mathbf{v}_k),$$

where each \mathbf{v}_i either does not end with the letter \mathbf{k} or is equal to \mathbf{k} .

Remark 6.2.1. In the subsequent proofs, we will often deal with the signatures of $\widehat{Y}_t^X = (t, X_t, Z_t^1, \dots, Z_t^d)$, where X_t is an arbitrary Itô process and Z_t is a d -dimensional Brownian motion. It is then beneficial, in proofs using induction on word length, to represent an arbitrary word \mathbf{w} as a polynomial of words that do not end with $\mathbf{2}$ (except for $\mathbf{2}^{\otimes k}$), corresponding to the process X . This representation reduces the analysis to standard Riemann integrals and Stratonovich integrals with respect to the Brownian motion.

6.2 Proof of Lemma 4.1

We will prove the result by induction on the length $|\mathbf{w}|$ of the word \mathbf{w} . For $|\mathbf{w}| = 1$, the result holds trivially with $C_{\mathbf{w}} = 1$. For $|\mathbf{w}| > 1$, we assume that (4.1) holds for words of length strictly less than $|\mathbf{w}|$. By Proposition 6.2, the word \mathbf{w} can be written as a linear combination of monomials of the form

$$\mathbf{2}^{\sqcup k} \sqcup \mathbf{u}_1 \mathbf{1} \sqcup \dots \sqcup \mathbf{u}_m \mathbf{1},$$

where

$$k + \sum_{i=1}^m N_2(\mathbf{u}_i) = N_2(\mathbf{w}), \quad k + m + \sum_{i=1}^m |\mathbf{u}_i| = |\mathbf{w}|. \quad (6.1)$$

Hence, it suffices to bound each monomial:

$$\begin{aligned} \left| \left\langle \mathbf{2}^{\sqcup k} \sqcup \mathbf{u}_1 \mathbf{1} \sqcup \dots \sqcup \mathbf{u}_m \mathbf{1}, \widehat{\mathbb{X}}_t \right\rangle \right| &= \left| X_t^k \prod_{i=1}^m \int_0^t \left\langle \mathbf{u}_i, \widehat{\mathbb{X}}_s \right\rangle ds \right| \\ &\leq \max_{t \in [0, T]} |X_t|^k \prod_{i=1}^m \left(T \max_{t \in [0, T]} \left| \left\langle \mathbf{u}_i, \widehat{\mathbb{X}}_t \right\rangle \right| \right) \\ &\leq T^m \max_{t \in [0, T]} |X_t|^k \prod_{i=1}^m \left(C_{\mathbf{u}_i} T^{|\mathbf{u}_i| - N_2(\mathbf{u}_i)} \max_{t \in [0, T]} |X_t|^{N_2(\mathbf{u}_i)} \right) \\ &= \left(\prod_{i=1}^m C_{\mathbf{u}_i} \right) T^{m + \sum_{i=1}^m (|\mathbf{u}_i| - N_2(\mathbf{u}_i))} \max_{t \in [0, T]} |X_t|^{k + \sum_{i=1}^m N_2(\mathbf{u}_i)}, \end{aligned}$$

where we have used for the second inequality the induction hypothesis (4.1) on the words \mathbf{u}_i , since $|\mathbf{u}_i| < |\mathbf{w}|$. It remains to observe that, by (6.1),

$$k + \sum_{i=1}^m N_2(\mathbf{u}_i) = N_2(\mathbf{w}), \quad m + \sum_{i=1}^m (|\mathbf{u}_i| - N_2(\mathbf{u}_i)) = |\mathbf{w}| - N_2(\mathbf{w}),$$

which yields (4.1) for the word \mathbf{w} and ends the proof.

6.3 Proof of Lemma 4.2

We proceed by induction on the length n of the word \mathbf{w} . For $n = 1$, (4.2) holds trivially. For $n \geq 2$, we assume the bound to hold for all words of length strictly less than n .

If $\mathbf{w} = \mathbf{2}^{\otimes n}$, the claim is obvious, as equality holds in (4.2) for $C = (n!)^{-q}$ and $\epsilon = 0$.

Otherwise, by Proposition 6.2, one can write \mathbf{w} as

$$\mathbf{w} = \sum_{k=1}^m \alpha_k \mathbf{w}_{k,1} \sqcup \dots \sqcup \mathbf{w}_{k,i_k}$$

where the words $\mathbf{w}_{k,j}$ contain at least one letter and satisfy

$$|\mathbf{w}_{k,1}| + \dots + |\mathbf{w}_{k,i_k}| = |\mathbf{w}| = n,$$

$$N_2(\mathbf{w}_{k,1}) + \dots + N_2(\mathbf{w}_{k,i_k}) = N_2(\mathbf{w}),$$

and for each k, j , $\mathbf{w}_{k,j}$ is either of length not greater than $n - 1$ or ending with a letter different from $\mathbf{2}$.

By the elementary inequality

$$(x_1 + \dots + x_m)^q \leq C_{q,m} (|x_1|^q + \dots + |x_m|^q),$$

it suffices to bound each term corresponding to the summands separately.

We then distinguish two cases. First, assume that $i_k \geq 2$, meaning that each of the words has length less than n . In this case, we apply Young's inequality to write

$$\prod_{j=1}^{i_k} \left| \langle \mathbf{w}_{k,j}, \widehat{\mathbb{Y}}_t^X \rangle \right|^q \lesssim \sum_{j=1}^k \left| \langle \mathbf{w}_{k,j}, \widehat{\mathbb{Y}}_t^X \rangle \right|^{q \frac{N_2(\mathbf{w}) + \epsilon}{N_2(\mathbf{w}_{k,j}) + \epsilon_j}},$$

where $\epsilon_j > 0$ are arbitrary such that $\sum_j \epsilon_j = \epsilon$. We then conclude by the induction hypothesis applied to each $\mathbf{w}_{k,j}$ since $|\mathbf{w}_{k,j}| < n$.

In the case where $i_k = 1$, we know that $\mathbf{w}_{k,1}$ is of the form $\mathbf{u}\mathbf{1}$ or $\mathbf{u}\mathbf{j}$ for $3 \leq j \leq d+2$. In the first case, this means that

$$\mathbb{E} \left[\int_0^\tau \left| \langle \mathbf{w}_{k,1}, \widehat{\mathbb{Y}}_t^X \rangle \right|^q dt \right] = \mathbb{E} \left[\int_0^\tau \left| \int_0^t \langle \mathbf{u}, \widehat{\mathbb{Y}}_s^X \rangle ds \right|^q dt \right] \leq T^{q-1} \mathbb{E} \left[\int_0^\tau \left| \langle \mathbf{u}, \widehat{\mathbb{Y}}_s^X \rangle \right|^q ds \right],$$

where we have used Hölder's inequality, and we conclude by the induction hypothesis applied to \mathbf{u} since $|\mathbf{u}| = n-1$.

For the case $\mathbf{w}_{k,1} = \mathbf{u}\mathbf{j} = \mathbf{u}'\mathbf{u}_n\mathbf{j}$, $j \geq 3$, this means that

$$\begin{aligned} \langle \mathbf{w}_{k,1}, \widehat{\mathbb{Y}}_t^X \rangle &= \int_0^t \langle \mathbf{u}, \widehat{\mathbb{Y}}_s^X \rangle \circ dZ_s^{j-2} \\ &= \int_0^t \langle \mathbf{u}, \widehat{\mathbb{Y}}_s^X \rangle dZ_s^{j-2} + \mathbf{1}_{\{\mathbf{u}_n = \mathbf{j}+2\}} \int_0^t \langle \mathbf{u}', \widehat{\mathbb{Y}}_s^X \rangle ds + \mathbf{1}_{\{\mathbf{u}_n = 2\}} \int_0^t \langle \mathbf{u}', \widehat{\mathbb{Y}}_s^X \rangle d\langle X_s, Z_s^{j-2} \rangle. \end{aligned}$$

Again, it is enough to bound separately the integrals of each of the three terms. The second one follows exactly as in the case $\mathbf{w}_{k,1} = \mathbf{u}\mathbf{1}$, and the third one vanishes, as follows from the assumption $\langle Z^i, X \rangle \equiv 0$. For the first term, using the Burkholder–Davis–Gundy inequality, we obtain

$$\begin{aligned} \mathbb{E} \left[\int_0^\tau \left| \int_0^t \langle \mathbf{u}, \widehat{\mathbb{Y}}_s^X \rangle dZ_s^{j-2} \right|^q dt \right] &\leq T \mathbb{E} \left[\sup_{0 \leq t \leq \tau} \left| \int_0^t \langle \mathbf{u}, \widehat{\mathbb{Y}}_s^X \rangle dZ_s^{j-2} \right|^q \right] \\ &\leq C_{T,q} \mathbb{E} \left[\left(\int_0^\tau \langle \mathbf{u}, \widehat{\mathbb{Y}}_s^X \rangle^2 ds \right)^{\frac{q}{2}} \right] \\ &\leq C'_{T,q} \mathbb{E} \left[\int_0^\tau \left| \langle \mathbf{u}, \widehat{\mathbb{Y}}_s^X \rangle \right|^q ds \right] \end{aligned}$$

Using Hölder's inequality, recalling that $q \geq 2$, and applying the induction hypothesis to \mathbf{u} since $|\mathbf{u}| = n-1$, we establish (4.2) for the word \mathbf{w} , thereby completing the proof.

7 Proof of martingality in Theorem 3.1 and Theorem 3.3

As noted in Remark 3.3.1, Theorem 3.1 follows immediately from Theorem 3.3 when $\rho \neq 0$. The case $\rho = 0$ is trivial, as X , given by (5.3), is a Brownian motion. Consequently, we have $\tau_\infty^X = +\infty$ almost surely, and S is a martingale by Corollary 5.3. Hence, we turn directly to the more general model (3.2) and proceed with the proof of Theorem 3.3.

If $N = 1$, the equation (5.1) is a linear SDE, which admits a unique strong solution on \mathbb{R}^+ . This ensures that $\tau_\infty^X = +\infty$ almost surely. We conclude that S is a martingale by Theorem 5.2. This establishes (i) of Theorem 3.3.

We now consider the non-trivial case $N \geq 2$, corresponding to point (ii) of Theorem 3.3. By Theorem 5.2, we need to prove that the solution X to (5.1) does not explode in finite time. Specifically, defining

$$\tau_n^X = \inf \{t \geq 0, |X_t| = n\},$$

we will show that

$$\lim_{n \rightarrow \infty} \mathbb{P}(\tau_n^X < T) = \mathbb{P}(\tau_\infty^X < T) = 0.$$

The probabilities on the left-hand side are controlled by

$$\mathbb{P}(\tau_n^X \leq T) \leq \frac{1}{n^{2N}} \mathbb{E} \left[X_{T \wedge \tau_n^X}^{2N} \right],$$

since

$$\mathbb{E} \left[X_{T \wedge \tau_n^X}^{2N} \right] \geq \mathbb{E} \left[n^{2N} \mathbf{1}_{\{\tau_n^X \leq T\}} \right] = n^{2N} \mathbb{P}(\tau_n^X \leq T),$$

so that it is enough to obtain a uniform bound (in n) for $\mathbb{E} \left[X_{T \wedge \tau_n^X}^{2N} \right]$.

By Itô's formula, we have on $t \in [[0, \tau_n^X]]$

$$dX_t^{2N} = 2N X_t^{2N-1} dX_t + N(2N-1) X_t^{2N-2} dt = 2N X_t^{2N-1} \left\langle \sigma, \widehat{Y}_t^X \right\rangle dt + 2N X_t^{2N-1} dB_t + N(2N-1) X_t^{2N-2} dt$$

Integration from 0 to $T \wedge \tau_n^X$, taking the expectation and denoting $\tilde{\sigma} = \sigma - \sigma^{2^{\otimes N}} \mathbf{2}^{\otimes N}$, we obtain

$$\begin{aligned} \mathbb{E} \left[X_{T \wedge \tau_n^X}^{2N} \right] &= \mathbb{E} \left[\int_0^{T \wedge \tau_n^X} \left(2N X_s^{2N-1} \left\langle \sigma, \widehat{Y}_s^X \right\rangle + N(2N-1) X_s^{2N-2} \right) ds \right] \\ &= \mathbb{E} \left[\int_0^{T \wedge \tau_n^X} \left(\frac{2N}{N!} \sigma^{2^{\otimes N}} X_s^{3N-1} + \frac{2N}{N!} X_s^{2N-1} \left\langle \tilde{\sigma}, \widehat{Y}_s^X \right\rangle + N(2N-1) X_s^{2N-2} \right) ds \right] \\ &\leq \mathbb{E} \left[\int_0^{T \wedge \tau_n^X} \left(\frac{2N}{N!} \sigma^{2^{\otimes N}} X_s^{3N-1} + \delta X_s^{3N-1} + C(N, \delta) \left\langle \tilde{\sigma}, \widehat{Y}_s^X \right\rangle^{\frac{3N-1}{N}} + N(2N-1) X_s^{2N-2} \right) ds \right] \\ &\leq \mathbb{E} \left[\int_0^{T \wedge \tau_n^X} \left(\left(\frac{2N}{N!} + \delta \right) \sigma^{2^{\otimes N}} X_s^{3N-1} + \tilde{C}(N, \delta, T) \left(|X_s|^{(3N-1)\frac{N-1}{N} + \epsilon} + 1 \right) + N(2N-1) X_s^{2N-2} \right) ds \right], \end{aligned}$$

where we have applied Young's inequality to obtain the first inequality and Lemma 4.2 for the second one.

Choosing $\delta \in \left(0, -\frac{2N}{N!} \sigma^{2^{\otimes N}} \right)$ and ϵ such that $(3N-1)\frac{N-1}{N} + \epsilon < 3N-1$, we ensure that the integrand is

bounded from above uniformly in n since $3N-1$ is even and $\left(\frac{2N}{N!} + \delta \right) \sigma^{2^{\otimes N}} < 0$. This finishes the proof.

8 Proofs of strict local martingality

In this section, we prove the necessity in Theorem 3.1 (ii) and Theorem 3.3 (ii). Unlike the ‘‘one-shot’’ proof of the martingale property in Section 7, we first prove strict local martingality in the simpler one-dimensional case and then extend it to the multidimensional setting.

8.1 Proof of necessity in Theorem 3.1 (ii)

The proof of the theorem follows the ideas used by Lions and Musiela (2007) in the Markovian setting.

Fix $T > 0$. According to Corollary 5.3, it is enough to prove that $\mathbb{P}(\tau_\infty^X < T) > 0$, where

$$\tau_\infty^X = \lim_{n \rightarrow \infty} \tau_n^X, \quad \tau_n^X := \inf \{t \geq 0: |X_t| = n\},$$

and $X = (X_t)_{t \geq 0}$ satisfies

$$dX_t = dW_t + \langle \rho \sigma, \widehat{\mathbb{X}}_t \rangle dt, \quad t \in [[0, \tau_\infty^X]], \quad X_0 = 0.$$

For the linear form of signature, the following inequality is a direct consequence of Lemma 4.1:

$$\sup_{t \in [0, T]} \left| \langle \sigma - \sigma^{2^{\otimes N}} \mathbf{2}^{\otimes N}, \widehat{\mathbb{X}}_t \rangle \right| \leq C_{\sigma, T} \left(1 + \sup_{t \in [0, T]} |X_t|^{N-1} \right) \quad a.s.$$

for some positive $C_{\sigma, T} > 0$ depending only on T and the coefficients σ . This implies

$$\frac{\sigma^{2^{\otimes N}}}{N!} X_t^N - C_{\sigma, T} \left(1 + \sup_{s \in [0, t]} |X_s|^{N-1} \right) \leq \langle \sigma, \widehat{\mathbb{X}}_t \rangle \leq \frac{\sigma^{2^{\otimes N}}}{N!} X_t^N + C_{\sigma, T} \left(1 + \sup_{s \in [0, t]} |X_s|^{N-1} \right) \quad a.s. \quad (8.1)$$

for all $t \in [0, \tau_\infty^X]$.

In particular, it follows that $\lim_{X_t \rightarrow \infty} \langle \sigma, \widehat{\mathbb{X}}_t \rangle^2 = +\infty$, so it is sufficient to show that $\mathbb{P}(\tau_\infty^X < T) > 0$.

For an arbitrary $\lambda \in \mathbb{R}$, define the event

$$\mathcal{A}_\lambda = \left\{ \omega : \sup_{t \in [0, T]} |W_t - \lambda t| \leq 1 \right\}, \quad \mathbb{P}(\mathcal{A}_\lambda) > 0.$$

If $\langle \sigma_t, \widehat{\mathbb{X}}_t \rangle$ was a deterministic function of X_t , so that X_t was Markovian, we would write on \mathcal{A}_λ

$$X_t \geq \lambda t - 1 + \rho \int_0^t \langle \sigma, \widehat{\mathbb{X}}_s \rangle ds$$

and apply the comparison theorem for ODEs to establish the explosion of X_t . However, in general, it is not true, and the equation becomes path-dependent. Our strategy consists of isolating the term $\frac{\sigma^{2^{\otimes N}}}{N!} X_t^N$

$$\langle \sigma, \widehat{\mathbb{X}}_t \rangle = \frac{\sigma^{2^{\otimes N}}}{N!} X_t^N + \langle \sigma - \sigma^{2^{\otimes N}} \mathbf{2}^{\otimes N}, \widehat{\mathbb{X}}_t \rangle$$

and bounding the remainder using (8.1). Then, we compare the solutions on $[\tau_n^X, \tau_{n+1}^X)$, bounding $\max_{s \in [0, t]} |X_s|^{N-1}$

by $(n+1)^{N-1}$, and apply the comparison principle for ODEs on this interval. This leads to a bound on $(\tau_{n+1}^X - \tau_n^X)$, which is used in its turn to bound τ_∞^X on \mathcal{A}_λ .

Assume first that $\rho \sigma^{2^{\otimes N}} > 0$. By (8.1), for $t \in [\tau_n^X, \tau_{n+1}^X)$, we have on \mathcal{A}_λ

$$\begin{aligned} X_t &\geq X_{\tau_n^X} + (W_t - W_{\tau_n^X}) + \frac{\rho \sigma^{2^{\otimes N}}}{N!} \int_{\tau_n^X}^t X_s^N ds - |\rho| C_{\sigma, T} \int_{\tau_n^X}^t \left(1 + \sup_{u \in [0, s]} |X_u|^{N-1} \right) ds \\ &\geq n + \lambda(t - \tau_n^X) - 2 + \frac{\rho \sigma^{2^{\otimes N}}}{N!} \int_{\tau_n^X}^t X_s^N ds - |\rho| C_{\sigma, T} (1 + (n+1)^{N-1}) (t - \tau_n^X). \end{aligned}$$

This allows us to apply the comparison theorem for ODEs, so that $X_t \geq Y_n(t)$ on $[\tau_n^X, \tau_{n+1}^X)$, where $Y_n(t)$ satisfies

$$\dot{Y}_n(t) = \lambda - |\rho| C_{\sigma, T} (1 + (n+1)^{N-1}) + \frac{\rho \sigma^{2^{\otimes N}}}{N!} Y_n(t)^N, \quad Y(\tau_n^X) = n - 2.$$

Since $\rho \sigma^{2^{\otimes N}} > 0$, we can choose λ large enough to guarantee that for all $n \geq 0$, the right-hand side is positive and $Y_n(t)$ is increasing. Then, since $Y_n(t) \geq n - 2$, for $t \in [\tau_n^X, \tau_{n+1}^X)$,

$$Y_n(t) \geq n - 2 + \left(\lambda + \frac{\rho \sigma^{2^{\otimes N}}}{N!} (n - 2)_+^N - |\rho| C_{\sigma, T} (1 + (n+1)^{N-1}) \right) (t - \tau_n^X).$$

Hence, on \mathcal{A}_λ , the following estimate holds

$$\tau_{n+1}^X - \tau_n^X \leq \tau_{n+1}^Y - \tau_n^X \leq \frac{3}{\lambda + \frac{\rho\sigma^{2^{\otimes N}}}{N!}(n-2)_+^N - |\rho|C_{\sigma,T}(1+(n+1)^{N-1})}$$

and

$$\tau_\infty^X = \sum_{n \geq 0} (\tau_{n+1}^X - \tau_n^X) \leq \sum_{n \geq 0} \frac{3}{\lambda + \frac{\rho\sigma^{2^{\otimes N}}}{N!}(n-2)_+^N - |\rho|C_{\sigma,T}(1+(n+1)^{N-1})}.$$

The series converges since $N \geq 2$ and the its sum can be made arbitrary small by choosing λ large enough.

For the remaining case of even N and $\rho\sigma^{2^{\otimes N}} < 0$, we take $\lambda < 0$ large enough, so that $X_{\tau_n^X} = -n$ on \mathcal{A}_λ , and we have on $[\tau_n^X, \tau_{n+1}^X)$

$$\begin{aligned} X_t &\leq X_{\tau_n^X} + (W_t - W_{\tau_n^X}) + \frac{\rho\sigma^{2^{\otimes N}}}{N!} \int_{\tau_n^X}^t X_s^N ds + |\rho|C_{\sigma,T} \int_{\tau_n^X}^t \left(1 + \sup_{u \in [0, s]} |X_u|^{N-1}\right) ds \\ &\leq -n + \lambda(t - \tau_n^X) + 2 + \frac{\rho\sigma^{2^{\otimes N}}}{N!} \int_{\tau_n^X}^t X_s^N ds + |\rho|C_{\sigma,T}(1+(n+1)^{N-1})(t - \tau_n^X). \end{aligned}$$

As $\frac{\rho\sigma^{2^{\otimes N}}}{N!}X_s^N < 0$, using the same argument as before, we can choose $\lambda(T)$ in order to ensure that X_t blows up to $-\infty$ on \mathcal{A}_λ before T for any $T > 0$.

8.2 Proof of necessity in Theorem 3.3 (ii)

We prove the result for the case where $\sigma^{2^{\otimes N}} > 0$ showing that X blows up to $+\infty$ with positive probability. The case where $\sigma^{2^{\otimes N}} < 0$ and N is even is treated similarly by showing that X blows up to $-\infty$.

The idea of this proof is to extend the proof of strict local martingality in Section 8 by finding a suitable event \mathcal{A} of positive probability such that the solution of (5.1) blows up before $T > 0$ on \mathcal{A} . We achieve this by keeping the elements of the signature containing Z (i.e., those corresponding to words containing the letters $\mathbf{3}, \dots, \mathbf{d} + \mathbf{2}$) small, so that the solution of the SDE remains close to the one considered in the one-dimensional case. More precisely, we define

$$\tau_n^X = \inf\{t \geq t_0 : X_t = n\},$$

and we fix $t_0 > 0$, $M \in \mathbb{N}$, and ϵ such that $N\epsilon < 1$. We require that on \mathcal{A} , the following inequalities hold:

$$X_{t_0} \geq M, \tag{8.2}$$

$$|\langle \mathbf{w}, \widehat{\mathbb{Y}}_{t_0}^X \rangle| \leq 1, \quad \text{for all } \mathbf{w} \text{ containing at least one of the letters } \mathbf{3}, \dots, \mathbf{d} + \mathbf{2}; \tag{8.3}$$

$$\sup_{s \in [\tau_n^X, \tau_{n+1}^X]} |\langle \mathbf{w}, \widehat{\mathbb{Y}}_s^X \rangle| \leq n^{N_2(\mathbf{w}) + \epsilon|\mathbf{w}|}, \quad \text{for } n \geq M \text{ and for all words } \mathbf{w} \text{ such that } |\mathbf{w}| \leq N; \tag{8.4}$$

$$\sup_{s \in [\tau_n^X, \tau_{n+1}^X]} |B_s - B_{\tau_n}| \leq n^{-N/2 + \epsilon}, \quad \text{for } n \geq M. \tag{8.5}$$

Step 1: Proving the explosion on \mathcal{A} . Conditional on this event, we can use ODE comparison on the dynamics of X to obtain that

$$\tau_{n+1}^X - \tau_n^X \leq Cn^{-N}, \quad n \geq M, \tag{8.6}$$

for some positive constant $C = C_{\sigma,N} > 0$ to be determined later. Indeed, for $t \in [\tau_n^X, \tau_n^X + Cn^{-N}]$, we have

$$\begin{aligned} X_t &= X_{\tau_n^X} + \int_{\tau_n^X}^t \langle \boldsymbol{\sigma}, \widehat{\mathbb{Y}}_s^X \rangle ds + B_t - B_{\tau_n^X} \\ &\geq X_{\tau_n^X} + \frac{\sigma^{2^{\otimes N}}}{N!} \int_{\tau_n^X}^t X_s^N ds - C_\sigma n^{(N-1) + \epsilon N} (t - \tau_n^X) - n^{-N/2 + \epsilon} \\ &\geq n - \left(C_\sigma C n^{N\epsilon - 1} + n^{-N/2 + \epsilon}\right) + \frac{\sigma^{2^{\otimes N}}}{N!} \int_{\tau_n^X}^t X_s^N ds, \end{aligned}$$

where we used conditions (8.4) and (8.5) in the first inequality. For n large enough, we have

$$C_\sigma C n^{N\epsilon-1} + n^{-N/2+\epsilon} < 1,$$

since $N\epsilon < 1$, so that

$$X_t \geq (n-1) + \frac{\sigma 2^{\otimes N}}{N!} \int_{\tau_n^X}^t X_s^N ds.$$

By the comparison theorem for ODEs, this implies that $\tau_{n+1}^X - \tau_n^X \leq \tau_{(n-1) \rightarrow (n+1)}^x$, where $\tau_{(n-1) \rightarrow (n+1)}^x$ is the time needed for the ODE

$$\dot{x}(t) = \frac{\sigma 2^{\otimes N}}{N!} x^N(t),$$

to go from $(n-1)$ to $(n+1)$. Since the coefficient $\sigma 2^{\otimes N}$ is positive, a straightforward computation gives

$$\tau_{(n-1) \rightarrow (n+1)}^x = \left(\frac{N!}{(N-1)\sigma 2^{\otimes N}} \right) ((n-1)^{1-N} - (n+1)^{1-N}) \leq C n^{-N}$$

for a suitable choice of $C > 0$. It then follows that, on the event \mathcal{A} , we have

$$\tau_\infty \leq t_0 + C \sum_{n \geq M} n^{-N}.$$

Since t_0 and M in (8.2)–(8.3) are arbitrary, they can be chosen so that $\tau_\infty < T$ on \mathcal{A} . It remains to show that $\mathbb{P}(\mathcal{A}) > 0$.

Step 2: Showing that $\mathbb{P}(\mathcal{A}) > 0$. First, we note that

$$\mathcal{A} = A_M \cap \bigcap_{n \geq M} \left(A_n^B \cap \bigcap_{\mathbf{w}: |\mathbf{w}| \leq N} A_n^{\mathbf{w}} \right),$$

where the event A_M corresponds to the inequalities (8.2)–(8.3), $A_n^{\mathbf{w}}$ corresponds to (8.4) for fixed n and \mathbf{w} , and A_n^B corresponds to (8.5) for a given n .

We first claim that $\mathbb{P}(A_M) > 0$. By the support theorem of [Stroock and Varadhan \(1972\)](#), it suffices to show that, for a suitable choice of (w, z^1, \dots, z^d) (deterministic smooth functions on $[0, T]$), the corresponding deterministic dynamics (i.e. where W, Z^i are replaced by w, z^i in (5.1)) satisfy the conditions. Taking $z^1 \equiv \dots \equiv z^d \equiv 0$ clearly leads to (8.3). Further taking $w_t = \lambda t$ leads to (8.2) by the analysis of Section 8.1. It follows that $\mathbb{P}(A_M) > 0$.

We define

$$A_{n+1} := A_n \cap \left(\bigcap_{|\mathbf{w}| \leq N} A_{n+1}^{\mathbf{w}} \right) \cap A_{n+1}^B, \quad n \geq M.$$

It is enough to prove that, if M is large enough, for each $n \geq M$, we have

$$\mathbb{P}(A_{n+1}^c | A_n) \leq c e^{-cn^\delta}, \tag{8.7}$$

for some $c, \delta > 0$. Indeed, since $A_{n+1} \subset A_n$ and $\mathcal{A} = \bigcap_{n \geq M} A_n$, we write

$$\mathbb{P}(\mathcal{A}) = \mathbb{P}(A_M) \prod_{n \geq M} \mathbb{P}(A_{n+1} | A_n),$$

so that, if (8.7) is verified, we have

$$\log(\mathbb{P}(\mathcal{A})) = \log(\mathbb{P}(A_M)) + \sum_{n \geq M} \log(\mathbb{P}(A_{n+1} | A_n)) \geq \log(\mathbb{P}(A_M)) + \sum_{n \geq M} \log(1 - c e^{-cn^\delta}).$$

As $n \rightarrow \infty$, $\log(1 - c e^{-cn^\delta}) \sim -c e^{-cn^\delta}$, and the series converges. Thus, we obtain $\mathbb{P}(\mathcal{A}) > 0$.

Note that since

$$\mathbb{P}(A_{n+1}^c | A_n) = \mathbb{P} \left(\bigcup_{\mathbf{w}: |\mathbf{w}| \leq N} (A_{n+1}^{\mathbf{w}})^c \cup (A_{n+1}^B)^c \middle| A_n \right) \leq \mathbb{P}((A_{n+1}^B)^c | A_n) + \sum_{\mathbf{w}: |\mathbf{w}| \leq N} \mathbb{P}((A_{n+1}^{\mathbf{w}})^c | A_n),$$

the condition (8.7) is stable under finite intersections, and it suffices to prove (8.7) for $\mathbb{P}((A_{n+1}^B)^c | A_n)$ and for each of the $\mathbb{P}((A_{n+1}^{\mathbf{w}})^c | A_n)$.

The bound (8.7) for $\mathbb{P}(A_{n+1}^B | A_n)$ follows from Gaussian scaling and Gaussian tail estimates for the Brownian motion B . Thus, it remains to establish (8.7) for the events $A_{n+1}^{\mathbf{w}}$, which we proceed to do via induction on the length of the word $|\mathbf{w}|$.

For $|\mathbf{w}| = 1$, we consider three cases. If $\mathbf{w} = \mathbf{1}$, (8.4) holds for sufficiently large n due to the bound (8.6) on τ_n^X . If $\mathbf{w} = \mathbf{2}$, (8.4) follows directly from the definition of τ_n^X for large n . In the remaining case, where $\mathbf{w} \in \{\mathbf{3}, \dots, \mathbf{d} + \mathbf{2}\}$, (8.7) follows from Gaussian tail estimates, as was previously done for the probability $\mathbb{P}(A_{n+1}^B | A_n)$.

Now, suppose that $|\mathbf{w}| > 1$ and that (8.7) holds for all words \mathbf{u} such that $|\mathbf{u}| < |\mathbf{w}|$.

By Proposition 6.2, we can write \mathbf{w} as a linear combination of shuffle products and we consider three cases.

The first case consists of words of the form $\mathbf{w} = \mathbf{w}_1 \sqcup \dots \sqcup \mathbf{w}_k$ with $k \geq 2$. The result then follows from the induction hypothesis for \mathbf{w}_j , since on $\bigcap_{j=1}^k A_{n+1}^{\mathbf{w}_j}$ we have

$$\sup_{s \in [\tau_n^X, \tau_{n+1}^X]} |\langle \mathbf{w}, \widehat{\mathbb{Y}}_s^X \rangle| \leq \prod_{j=1}^k \sup_{s \in [\tau_n^X, \tau_{n+1}^X]} |\langle \mathbf{w}_j, \widehat{\mathbb{Y}}_s^X \rangle| \leq \prod_{j=1}^k n^{N_2(\mathbf{w}_j) + \epsilon |\mathbf{w}_j|} = n^{N_2(\mathbf{w}) + \epsilon |\mathbf{w}|}.$$

This implies that $A_{n+1}^{\mathbf{w}} \subset \bigcap_{j=1}^k A_{n+1}^{\mathbf{w}_j}$.

The second case concerns words of the form $\mathbf{u}\mathbf{1}$, which is again handled by the induction hypothesis for \mathbf{u} . Indeed, conditional on A_n and $A_{n+1}^{\mathbf{u}}$, we have

$$\begin{aligned} \sup_{s \in [\tau_n^X, \tau_{n+1}^X]} |\langle \mathbf{w}, \widehat{\mathbb{Y}}_s^X \rangle| &= \sup_{s \in [\tau_n^X, \tau_{n+1}^X]} \left| \langle \mathbf{w}, \widehat{\mathbb{Y}}_{\tau_n^X}^X \rangle + \int_{\tau_n^X}^s \langle \mathbf{u}, \widehat{\mathbb{Y}}_s^X \rangle ds \right| \\ &\leq (n-1)^{N_2(\mathbf{w}) + \epsilon |\mathbf{w}|} + (\tau_{n+1}^X - \tau_n^X) n^{N_2(\mathbf{u}) + \epsilon |\mathbf{u}|} \\ &\leq (n-1)^{N_2(\mathbf{w}) + \epsilon |\mathbf{w}|} + C n^{-N + N_2(\mathbf{u}) + \epsilon |\mathbf{u}|} \\ &\leq (n-1)^{N_2(\mathbf{w}) + \epsilon |\mathbf{w}|} + C n^{-1 + \epsilon(N-1)}, \end{aligned}$$

which is bounded by $n^{N_2(\mathbf{w}) + \epsilon |\mathbf{w}|}$ for sufficiently large n (recall that M can be chosen large enough).

The last case corresponds to $\mathbf{w} = \mathbf{u}\mathbf{i}$ for $i \in \{3, \dots, d+2\}$. We write

$$\begin{aligned} \mathbb{P}((A_{n+1}^{\mathbf{w}})^c | A_n) &= \mathbb{P}((A_{n+1}^{\mathbf{w}})^c, (A_{n+1}^{\mathbf{u}})^c | A_n) + \mathbb{P}((A_{n+1}^{\mathbf{w}})^c, A_{n+1}^{\mathbf{u}} | A_n) \\ &\leq \mathbb{P}((A_{n+1}^{\mathbf{u}})^c | A_n) + \mathbb{P}((A_{n+1}^{\mathbf{w}})^c, A_{n+1}^{\mathbf{u}} | A_n), \end{aligned}$$

and note that (8.7) holds for the first term by the induction hypothesis for \mathbf{u} . For $s \geq \tau_n$, we have

$$\langle \mathbf{w}, \widehat{\mathbb{Y}}_s^X \rangle = \langle \mathbf{w}, \widehat{\mathbb{Y}}_{\tau_n^X}^X \rangle + \int_{\tau_n^X}^s \langle \mathbf{u}, \widehat{\mathbb{Y}}_s^X \rangle \circ dZ_s^{i-2}.$$

The first term and the Stratonovich correction are bounded on A_n and $A_{n+1}^{\mathbf{u}}$ as in the previous case. Thus, it suffices to prove the bound for the conditional probability of the event

$$\left\{ \sup_{t \in [\tau_n^X, \tau_{n+1}^X]} \int_{\tau_n^X}^t \langle \mathbf{u}, \widehat{\mathbb{Y}}_s^X \rangle dZ_s^{i-2} \geq n^{N_2(\mathbf{w}) + \epsilon |\mathbf{w}| - 1} \right\}.$$

We use the concentration inequality

$$\mathbb{P} \left(\sup_{t \in [\tau, \tau+h]} \left| \int_{\tau}^t \gamma_s dW_s \right| \geq \lambda, \sup_{t \in [\tau, \tau+h]} |\gamma_t| \leq \eta \right) \leq \exp \left(-\frac{\lambda^2}{2\eta^2 h} \right),$$

which follows from the Gaussian tails of Brownian motion and the Dambis—Dubins—Schwarz theorem and holds for any stopping time τ , any $h > 0$, and any progressively measurable γ .

$$\begin{aligned} & \mathbb{P} \left(\sup_{t \in [\tau_n^X, \tau_{n+1}^X]} \int_{\tau_n}^t \langle \mathbf{u}, \widehat{\mathbb{Y}}_s^X \rangle dZ_s^i \geq n^{N_2(\mathbf{w}) + \epsilon|\mathbf{w}| - 1}, A_{n+1}^{\mathbf{u}} \mid A_n \right) \\ & \leq \mathbb{P} \left(\sup_{t \in [\tau_n^X, \tau_{n+1}^X]} \int_{\tau_n}^t \langle \mathbf{u}, \widehat{\mathbb{Y}}_s^X \rangle \circ dZ_s^i \geq n^{N_2(\mathbf{w}) + \epsilon|\mathbf{w}| - 1}, \sup_{t \in [\tau_n^X, \tau_{n+1}^X]} \left| \langle \mathbf{u}, \widehat{\mathbb{Y}}_t^X \rangle \right| \leq n^{N_2(\mathbf{w}) + \epsilon(|\mathbf{w}| - 1)} \mid A_n \right) \\ & \leq \exp \left(-\frac{n^{N-2(1-\epsilon)}}{2} \right). \end{aligned}$$

Thus, (8.7) holds for \mathbf{w} . This completes the proof.

9 Proof of Theorem 3.2: Moments explosion

Without loss of generality, we will assume that N is odd and

$$\sigma^{2^{\otimes N}} > 0 \text{ and } \rho < 0.$$

The proof for the remaining case $\sigma^{2^{\otimes N}} < 0$ and $\rho > 0$ is analogous.

By Itô's formula,

$$\begin{aligned} S_T^m &= S_0^m \exp \left(m \int_0^T \sigma_s dB_s - \frac{m}{2} \int_0^T \sigma_s^2 ds \right) \\ &= S_0^m \exp \left(\rho m \int_0^T \sigma_s dW_s + \bar{\rho} m \int_0^T \sigma_s dW_s^\perp - \frac{m}{2} \int_0^T \sigma_s^2 ds \right), \end{aligned}$$

As $\sigma_t = \langle \boldsymbol{\sigma}, \widehat{\mathbb{W}}_t \rangle$ is adapted to the filtration generated by W , conditioning with respect to W implies

$$S_0^{-m} \mathbb{E} [S_T^m] = \mathbb{E} \left[\exp \left(\rho m \int_0^T \langle \boldsymbol{\sigma}, \widehat{\mathbb{W}}_s \rangle dW_s + \frac{\bar{\rho}^2 m^2 - m}{2} \int_0^T \langle \boldsymbol{\sigma}, \widehat{\mathbb{W}}_s \rangle^2 ds \right) \right].$$

We then proceed as in Gassiat (2018) and apply the Boué and Dupuis (1998) formula (more precisely, a variant allowing unbounded functionals, see Appendix A) to obtain

$$\begin{aligned} \ln \mathbb{E} [S_T^m / S_0^m] &= \sup_{(u_t)_{t \geq 0} \in \mathcal{U}_q} \mathbb{E} \left[\int_0^T \left(-\frac{u_s^2}{2} + \rho m \langle \boldsymbol{\sigma}, \widehat{\mathbb{X}}_s^u \rangle u_s + \frac{\bar{\rho}^2 m^2 - m}{2} \langle \boldsymbol{\sigma}, \widehat{\mathbb{X}}_s^u \rangle^2 \right) ds \right] \\ &= \sup_{(u_t)_{t \geq 0} \in \mathcal{U}_q} \mathbb{E} \left[\int_0^T \left(\frac{m^2 - m}{2} \langle \boldsymbol{\sigma}, \widehat{\mathbb{X}}_s^u \rangle^2 - \frac{(u_s - \rho m \langle \boldsymbol{\sigma}, \widehat{\mathbb{X}}_s^u \rangle)^2}{2} \right) ds \right] \\ &=: \sup_{u \in \mathcal{U}_q} \mathcal{J}(u; \rho, m, T) =: \mathcal{V}(\rho, m, T), \end{aligned} \tag{9.1}$$

where

$$\mathcal{U}_q = \left\{ (u_t)_{t \geq 0} \text{ progressively measurable with } \mathbb{E} \left[\int_0^T |u_t|^q dt \right] < +\infty \right\},$$

for some fixed $q \geq 2N$, and $\widehat{\mathbb{X}}_t^u$ denotes the signature of $\widehat{X}_t^u = (t, X_t^u)$ with X_t^u defined by

$$X_t^u = W_t + \int_0^t u_s ds.$$

The finiteness of the moments is therefore equivalent to the control problem having a finite value. We now proceed to prove that, depending on the position of ρ relative to $-\sqrt{1 - \frac{1}{m}}$, the value will either be finite or infinite.

9.1 Proof of Theorem 3.2 (i): negative result

In this subsection we assume that

$$-\sqrt{1 - \frac{1}{m}} < \rho < 0,$$

and we will show that $\mathcal{V}(\rho, m, T) = +\infty$. The idea is simple: we take u of feedback form

$$u_t = \lambda \left\langle \boldsymbol{\sigma}, \widehat{\mathbb{X}}_t^u \right\rangle$$

for $\lambda \in \mathbb{R}$.

In that case, we denote by X^λ the solution of the ODE

$$dX_t^\lambda = dW_t + \lambda \left\langle \boldsymbol{\sigma}, \widehat{\mathbb{X}}_t^\lambda \right\rangle dt,$$

and we denote its signature by $\widehat{\mathbb{X}}_t^\lambda$. This solution exists (up to a possible explosion time) by Proposition 5.1, the value (9.1) corresponding to the control is simply

$$P(\lambda) \mathbb{E} \left[\int_0^T \left\langle \boldsymbol{\sigma}, \widehat{\mathbb{X}}_s^\lambda \right\rangle^2 ds \right],$$

with

$$P(\lambda) = \frac{m^2 - m}{2} - \frac{(\lambda - \rho m)^2}{2}.$$

The assumption on ρ is exactly equivalent to $P(0) > 0$, which guarantees that there exists λ s.t.

$$\lambda > 0 \text{ and } P(\lambda) > 0.$$

As shown in the proof in Section 8, the first inequality $\lambda > 0$ will imply that X^λ blows up with positive probability, which with the help of the second one implies that the problem value (9.1) is infinite.

The above argument is not completely rigorous, since the feedback control is only defined up to the explosion time and clearly not in \mathcal{U}_q . This can be dealt with by a simple truncation argument : for $R > 0$, let

$$\tau_{R,\lambda} = \inf\{t \geq 0 : |X_t^\lambda| = R\}$$

and take

$$u_t^{\lambda,R} = \begin{cases} \lambda \left\langle \boldsymbol{\sigma}, \widehat{\mathbb{X}}_t^\lambda \right\rangle, & t < \tau_{R,\lambda} \\ 0, & t \geq \tau_{R,\lambda} \end{cases}$$

Then, by Lemma 4.2, $u^{\lambda,R}$ is in \mathcal{U}_q (for any $q \geq 2$), and

$$\mathcal{V}(\rho, m, T) \geq \mathcal{J}(u^{\lambda,R}; \rho, m, T) = P(\lambda) \mathbb{E} \left[\int_0^{T \wedge \tau_{R,\lambda}} \left\langle \boldsymbol{\sigma}, \widehat{\mathbb{X}}_s^\lambda \right\rangle^2 ds \right] + P(0) \mathbb{E} \left[\int_{T \wedge \tau_{R,\lambda}}^T \left\langle \boldsymbol{\sigma}, \widehat{\mathbb{X}}_s^{u^{\lambda,R}} \right\rangle^2 ds \right].$$

Taking the supremum over R , we obtain

$$\mathcal{V}(\rho, m, T) \geq P(\lambda) \mathbb{E} \left[\int_0^{T \wedge \tau_{\infty,\lambda}} \left\langle \boldsymbol{\sigma}, \widehat{\mathbb{X}}_s^\lambda \right\rangle^2 ds \right]$$

where $\tau_{\infty,\lambda}$ is the explosion time of X^λ . However, the r.h.s is infinite since the event $\{\tau_{\infty,\lambda} < T\}$ has positive probability (by the results of Section 8.1), and on this event $\int_0^{\tau_{\infty,\lambda}} \left\langle \boldsymbol{\sigma}, \widehat{\mathbb{X}}_s^\lambda \right\rangle^2 ds = +\infty$ (by Proposition 5.1).

9.2 Proof of Theorem 3.2 (ii): positive result

We assume now that the condition

$$\rho < -\sqrt{1 - \frac{1}{m}},$$

is verified and fix $(u_t)_{t \geq 0} \in \mathcal{U}_q$. We want to bound

$$\mathbb{E} \left[\int_0^T \left(-\frac{u_s^2}{2} + \rho m \langle \boldsymbol{\sigma}, \widehat{\mathbb{X}}_s^u \rangle u_s + \frac{\bar{\rho}^2 m^2 - m}{2} \langle \boldsymbol{\sigma}, \widehat{\mathbb{X}}_s^u \rangle^2 \right) ds \right]. \quad (9.2)$$

Define

$$U_t := \int_0^t u_s ds.$$

We denote by $\widehat{Y}_t^U = (t, U_t, W_t)$ and we denote its signature by \widehat{Y}_t^U . Since this is a 3-dimensional process, this signature is in the dual of the tensor algebra over words with letters **1, 2, 3**. Since $X = U + W$, we can rewrite the linear forms in terms of \widehat{Y}_t^U . Namely, there exists $\tilde{\boldsymbol{\sigma}} \in T^N(\mathbb{R}^3)$, such that $\langle \boldsymbol{\sigma}, \widehat{\mathbb{X}}_t^u \rangle = \langle \tilde{\boldsymbol{\sigma}}, \widehat{Y}_t^U \rangle$.

In the expectation (9.2), the term with the highest degree in U is

$$\frac{\bar{\rho}^2 m^2 - m}{2} \left(\frac{\boldsymbol{\sigma}^{2^{\otimes N}}}{N!} \right)^2 \mathbb{E} \left[\int_0^T U_t^{2N} dt \right].$$

Note that $\bar{\rho}^2 m^2 - m$ is strictly negative by assumption.

We return to the value in (9.2) and bound the three terms in the integrand separately. The first one is non-positive and can therefore be ignored.

Step 1. Bounding the third term. Using the same techniques as in the proof of Theorem 3.3, we can bound the third term $\frac{\bar{\rho}^2 m^2 - m}{2} \mathbb{E} \left[\int_0^T \langle \tilde{\boldsymbol{\sigma}}, \widehat{Y}_s^U \rangle^2 ds \right]$ from above by

$$-c_1 \mathbb{E} \left[\int_0^T U_t^{2N} dt \right] + \sum_{\substack{|\mathbf{w}| \leq N, \\ N_2(\mathbf{w}) < N}} c_{\mathbf{w}} \mathbb{E} \left[\int_0^T \langle \mathbf{w}, \widehat{Y}_t^U \rangle^2 dt \right] \quad (9.3)$$

for some positive c_1 and $c_{\mathbf{w}}$. By Lemma 4.2, it is further bounded from above by

$$\mathbb{E} \left[\int_0^T (-c_1 U_t^{2N} + c_2 U_t^{2N-1}) dt \right] + c_3 < +\infty.$$

Step 2. Bounding the second term. For the second term $\rho m \mathbb{E} \left[\int_0^T \langle \tilde{\boldsymbol{\sigma}}, \widehat{Y}_s^U \rangle u_s ds \right]$, we note that, for suitable coefficients $c_{\mathbf{w}}$,

$$\rho m \mathbb{E} \left[\int_0^T \langle \boldsymbol{\sigma}, \widehat{\mathbb{X}}_s^u \rangle u_s ds \right] = \rho m \mathbb{E} \left[\int_0^T \langle \boldsymbol{\sigma}, \widehat{\mathbb{X}}_s^u \rangle dU_s \right] = -c_4 \mathbb{E}[U_T^{N+1}] + \sum_{\substack{|\mathbf{w}| \leq N, \\ N_2(\mathbf{w}) < N}} c_{\mathbf{w}} \mathbb{E} \left[\langle \mathbf{w}, \widehat{Y}_T^U \rangle \right], \quad (9.4)$$

where $c_4 = -\frac{\rho m \boldsymbol{\sigma}^{2^{\otimes N}}}{(N+1)!}$. We note that the Itô and Stratonovich integrals with respect to U coincide since U has bounded variation. By the assumption of the theorem, $c_4 > 0$, so that the first term in (9.4) is negative.

For the second term, we claim that it holds, for any \mathbf{w} with $|\mathbf{w}| \leq N+1$ and $N_2(\mathbf{w}) \leq N$, for arbitrary $\epsilon > 0$,

$$\mathbb{E} \left[\langle \mathbf{w}, \widehat{Y}_T^U \rangle \right] \lesssim 1 + \mathbb{E}[|U_T|^{N+\epsilon}] + \int_0^T \mathbb{E}[|U_t|^{2(N-1)+\epsilon}] dt \quad (9.5)$$

We first consider separately the case where $|\mathbf{w}| = N + 1$ and $N_2(\mathbf{w}) = N$, i.e., only one letter differs from $\mathbf{2}$, say \mathbf{j} . By Proposition 6.2, we can express \mathbf{w} in the form

$$\mathbf{w} = \sum_{k=0}^N \alpha_k \mathbf{2}^{\sqcup k} \sqcup \mathbf{2}^{\otimes N-k} \mathbf{j} \quad (9.6)$$

If $\mathbf{j} = \mathbf{3}$, then (9.6) takes the form

$$\mathbb{E} \left[\left\langle \mathbf{w}, \widehat{\mathbb{Y}}_T^U \right\rangle \right] = \sum_{k=1}^N \alpha_k \mathbb{E} \left[U_T^k \left(\int_0^T U_t^{N-k} dW_t \right) \right]$$

Note that we omitted the term for $k = 0$, as it has zero expectation. Applying Young's inequality and the Burkholder–Davis–Gundy inequality to each term, we obtain

$$\begin{aligned} \mathbb{E} \left[U_T^k \left(\int_0^T U_t^{N-k} dW_t \right) \right] &\lesssim \mathbb{E} [|U_T|^{N+\epsilon}] + \mathbb{E} \left[\left(\int_0^T U_t^{(N-k)} dW_t \right)^{\frac{N+\epsilon}{(N-k+\epsilon)}} \right] \\ &\lesssim \mathbb{E} [|U_T|^{N+\epsilon}] + \mathbb{E} \left[\left(\int_0^T U_t^{2(N-k)} dt \right)^{\frac{N+\epsilon}{2(N-k+\epsilon)}} \right] \\ &\lesssim 1 + \mathbb{E} [|U_T|^{N+\epsilon}] + \mathbb{E} \left[\left(\int_0^T U_t^{2(N-k)} dt \right)^{\max\{\frac{N+\epsilon}{2(N-k+\epsilon)}, 1\}} \right] \\ &\lesssim 1 + \mathbb{E} [|U_T|^{N+\epsilon}] + \mathbb{E} \left[\int_0^T |U_t|^{\max\{N+\epsilon, 2(N-k)\}} dt \right], \\ &\lesssim 1 + \mathbb{E} [|U_T|^{N+\epsilon}] + \mathbb{E} \left[\int_0^T |U_t|^{2(N-1)+\epsilon} dt \right], \end{aligned}$$

which proves the claim (9.5) in this case.

If $\mathbf{j} = \mathbf{1}$, a similar but simpler computation yields

$$\mathbb{E} \left[U_T^k \left(\int_0^T U_t^{N-k} dt \right) \right] \lesssim \mathbb{E} [|U_T|^N] + \mathbb{E} \left[\int_0^T |U_t|^N dt \right],$$

We then assume that $N_2(\mathbf{w}) \leq N - 1$. In this case, by Proposition 6.2, we can write \mathbf{w} as shuffle polynomial with monomials of the form

$$\mathbf{2}^{\sqcup k} \sqcup \mathbf{u}_1 \mathbf{1} \sqcup \dots \sqcup \mathbf{u}_m \mathbf{1} \sqcup \mathbf{v}_1 \mathbf{3} \sqcup \dots \sqcup \mathbf{v}_n \mathbf{3},$$

so that it remains to prove estimates on

$$\mathbb{E} \left[U_T^k \prod_{i=1}^m \left(\int_0^T \langle \mathbf{u}_i, \widehat{\mathbb{Y}}_t^U \rangle dt \right) \prod_{j=1}^n \left(\int_0^T \langle \mathbf{v}_j, \widehat{\mathbb{Y}}_t^U \rangle \circ dW_t \right) \right].$$

Again, an application of Young's inequality yields that this expectation is bounded by

$$\mathbb{E} [|U_T|^{N_2(\mathbf{w})+\epsilon}] + \sum_{i=1}^m \mathbb{E} \left[\left| \int_0^T \langle \mathbf{u}_i, \widehat{\mathbb{Y}}_t^U \rangle dt \right|^{\frac{N_2(\mathbf{w})+\epsilon}{N_2(\mathbf{u}_i)+\epsilon_{1,i}}} \right] + \sum_{j=1}^n \mathbb{E} \left[\left| \int_0^T \langle \mathbf{v}_j, \widehat{\mathbb{Y}}_t^U \rangle \circ dW_t \right|^{\frac{N_2(\mathbf{w})+\epsilon}{N_2(\mathbf{v}_j)+\epsilon_{2,j}}} \right],$$

where $\epsilon_{1,i}, \epsilon_{2,j} > 0$ are arbitrary, with $\sum_{i=1}^m \epsilon_{1,i} + \sum_{j=1}^n \epsilon_{2,j} = \epsilon$. The first term is of the correct order since $N_2(\mathbf{w}) \leq N$. For the second one, we use Hölder's inequality and Lemma 4.2 to obtain

$$\begin{aligned} \mathbb{E} \left[\left| \int_0^T \langle \mathbf{u}_i, \widehat{\mathbb{Y}}_t^U \rangle dt \right|^{\frac{N_2(\mathbf{w})+\epsilon}{N_2(\mathbf{u}_i)+\epsilon_{1,i}}} \right] &\lesssim \int_0^T \mathbb{E} \left| \langle \mathbf{u}_i, \widehat{\mathbb{Y}}_t^U \rangle \right|^{\frac{N_2(\mathbf{w})+\epsilon}{N_2(\mathbf{u}_i)+\epsilon_{1,i}}} dt \\ &\lesssim 1 + \int_0^T \mathbb{E} \left| \langle \mathbf{u}_i, \widehat{\mathbb{Y}}_t^U \rangle \right|^{\max\left\{\frac{N_2(\mathbf{w})+\epsilon}{N_2(\mathbf{u}_i)+\epsilon_{1,i}}, 2\right\}} dt \\ &\lesssim 1 + \int_0^T \mathbb{E} \left[|U_t|^{\max\{N_2(\mathbf{w})+\epsilon, 2(N_2(\mathbf{u}_i)+\epsilon_{1,i})\}} \right] dt \\ &\lesssim 1 + \int_0^T \mathbb{E} \left[|U_t|^{2(N-1)+\epsilon} \right] dt. \end{aligned}$$

For the second term, the Itô–Stratonovich correction is treated as in the inequality above, and for the Itô integral, we use the Burkholder–Davis–Gundy inequality and Lemma 4.2 to obtain that

$$\begin{aligned} \mathbb{E} \left[\left| \int_0^T \langle \mathbf{v}_j, \widehat{\mathbb{Y}}_t^U \rangle dW_t \right|^{\frac{N_2(\mathbf{w})+\epsilon}{N_2(\mathbf{v}_j)+\epsilon_{2,j}}} \right] &\lesssim \mathbb{E} \left[\left| \int_0^T \langle \mathbf{v}_j, \widehat{\mathbb{Y}}_t^U \rangle^2 dt \right|^{\frac{N_2(\mathbf{w})+\epsilon}{2(N_2(\mathbf{v}_j)+\epsilon_{2,j})}} \right] \\ &\lesssim 1 + \mathbb{E} \left[\left| \int_0^T \langle \mathbf{v}_j, \widehat{\mathbb{Y}}_t^U \rangle^2 dt \right|^{\max\left\{\frac{N_2(\mathbf{w})+\epsilon}{2(N_2(\mathbf{v}_j)+\epsilon_{2,j})}, 1\right\}} \right] \\ &\lesssim 1 + \mathbb{E} \left[\int_0^T \left| \langle \mathbf{v}_j, \widehat{\mathbb{Y}}_t^U \rangle \right|^{\max\left\{\frac{N_2(\mathbf{w})+\epsilon}{N_2(\mathbf{v}_j)+\epsilon_{2,j}}, 2\right\}} dt \right] \\ &\lesssim 1 + \mathbb{E} \left[\int_0^T |U_t|^{\max\{N_2(\mathbf{w})+\epsilon, 2(N_2(\mathbf{v}_j)+\epsilon_{2,j})\}} dt \right] \\ &\lesssim 1 + \int_0^T \mathbb{E} \left[|U_t|^{2(N-1)+\epsilon} \right] dt. \end{aligned}$$

This proves (9.5).

To conclude, combining (9.3), (9.4), and (9.5), we observe that we have proven the existence of polynomial functions P and Q of orders strictly less than $2N$ and $N + 1$, respectively, such that, for any $u \in \mathcal{U}_q$,

$$\mathbb{E} \left[\int_0^T \left(\frac{\bar{\rho}^2 m^2 - m}{2} \langle \boldsymbol{\sigma}, \widehat{\mathbb{X}}_s^u \rangle^2 + \rho m \langle \boldsymbol{\sigma}, \widehat{\mathbb{X}}_s^u \rangle u_s - \frac{u_s^2}{2} \right) ds \right] \leq \mathbb{E} \left[\int_0^T (-c_1 U_t^{2N} + P(|U_t|)) dt + (-c_4 U_T^{N+1} + Q(|U_T|)) \right].$$

Note that the polynomials under the expectation are bounded from above uniformly in u . This implies the finiteness of $\mathcal{V}(\rho, m, T)$, and by (9.1), this proves the finiteness of $\mathbb{E}[S_T^m]$.

A Boué–Dupuis formula and proof of (9.1)

We will use the following variant of the Boué–Dupuis formula, which is a minor extension on recent results of Hariya and Watanabe (2022).

In this section, $W = (W_t)_{0 \leq t \leq T}$ is a (d -dimensional) Brownian motion under the measure \mathbb{P} , $(\mathcal{F}_t)_{t \geq 0}$ is its augmented natural filtration, and for any $q \geq 2$, we let

$$\mathcal{U}_q = \left\{ (u_t)_{t \geq 0}(\mathcal{F}_t) - \text{progressively measurable with } \mathbb{E} \left[\int_0^T |u_t|^q dt \right] < +\infty \right\}.$$

Proposition A.1. *Let $q \geq 2$ be fixed, and $F : C([0, T], \mathbb{R}^d) \rightarrow \mathbb{R}$ be (Borel-)measurable and such that*

$$\mathbb{E} \left[\left| F \left(W + \int_0^\cdot u \right) \right| \right] < \infty, \quad u \in \mathcal{U}_q. \quad (\text{A.1})$$

Then it holds that

$$\ln \mathbb{E} \left[e^{F(W)} \right] = \sup_{u \in \mathcal{U}_q} \mathbb{E} \left[F \left(W + \int_0^\cdot u \right) - \frac{1}{2} \int_0^T u_s^2 ds \right].$$

Proof. We proceed as in (Hariya and Watanabe, 2022, Remark 1.2 (1)) by a truncation argument. For $n \geq 0$, let $F_n = F \wedge n$, then it holds that $\mathbb{E}[|F_n(W)|]$ and $\mathbb{E}[e^{F_n(W)}]$ are finite, so that we can apply the results of Hariya and Watanabe (2022) to obtain

$$\ln \mathbb{E} \left[e^{F_n(W)} \right] = \sup_{u \in \mathcal{U}_q} \mathbb{E} \left[F_n \left(W + \int_0^\cdot u \right) - \frac{1}{2} \int_0^T u_s^2 ds \right]$$

(They prove that the above formula holds for the supremum taken over both \mathcal{U}_2 and the set of bounded controls \mathcal{U}_b , see Theorem 1.1 and Corollary 2.11 therein. This implies the case of arbitrary $q \geq 2$ since $\mathcal{U}_b \subset \mathcal{U}_q \subset \mathcal{U}_2$).

We then have by monotone convergence that

$$\begin{aligned} \ln \mathbb{E} \left[e^{F(W)} \right] &= \sup_n \ln \mathbb{E} \left[e^{F_n(W)} \right] \\ &= \sup_{n \geq 0} \sup_{u \in \mathcal{U}_q} \mathbb{E} \left[F_n \left(W + \int_0^\cdot u \right) - \frac{1}{2} \int_0^T u_s^2 ds \right] \\ &= \sup_{u \in \mathcal{U}_q} \sup_{n \geq 0} \mathbb{E} \left[F_n \left(W + \int_0^\cdot u \right) - \frac{1}{2} \int_0^T u_s^2 ds \right] \\ &= \sup_{u \in \mathcal{U}_q} \mathbb{E} \left[F \left(W + \int_0^\cdot u \right) - \frac{1}{2} \int_0^T u_s^2 ds \right] \end{aligned}$$

where the last equality follows from (A.1) and dominated convergence. \square

Proof of (9.1). We let

$$F(W) = \rho m \int_0^T \langle \sigma, \widehat{\mathbb{W}}_s \rangle dW_s + \frac{\bar{\rho}^2 m^2 - m}{2} \int_0^T \langle \sigma, \widehat{\mathbb{W}}_s \rangle^2 ds$$

and check that it satisfies (A.1) for $q \geq 2N$. For $u \in \mathcal{U}_q$, letting $\widehat{\mathbb{X}}_t^u$ denote the signature of $\widehat{X}_t^u = (t, X_t^u)$ with X_t^u defined by

$$X_t^u = W_t + \int_0^t u_s ds,$$

one has that

$$F \left(W + \int_0^\cdot u \right) = \rho m \int_0^T \langle \sigma, \widehat{\mathbb{X}}_s^u \rangle (dW_s + u_s ds) + \frac{\bar{\rho}^2 m^2 - m}{2} \int_0^T \langle \sigma, \widehat{\mathbb{X}}_s^u \rangle^2 ds$$

As in Section 10.2, there exists $\tilde{\sigma} \in T^N(\mathbb{R}^3)$, such that $\langle \sigma, \widehat{\mathbb{X}}_t^u \rangle = \langle \tilde{\sigma}, \widehat{\mathbb{Y}}_t^U \rangle$ where $\widehat{\mathbb{Y}}_t^U$ is the signature of $Y_t^U = (t, \int_0^t u_s ds, W_t)$. By Lemma 4.2 and Hölder's inequality, this implies that

$$\mathbb{E} \left[\int_0^T \langle \sigma, \widehat{\mathbb{X}}_s^u \rangle^2 ds \right] \leq C \left(1 + \mathbb{E} \left[\int_0^T |u_s|^{2N} ds \right] \right) < \infty \quad (\text{A.2})$$

since $q \geq 2N$. Since it holds that

$$\mathbb{E} \left[\left| F \left(W + \int_0^\cdot u \right) \right| \right] \leq C \left(1 + \mathbb{E} \left[\int_0^T u_s^2 ds \right] + \mathbb{E} \left[\int_0^T \langle \sigma, \widehat{\mathbb{X}}_s^u \rangle^2 ds \right] \right)$$

(using Itô isometry for the term in dW and Young's inequality for the term in $u_s ds$), we see that (A.1) holds.

We then use Proposition A.1 to obtain that

$$\begin{aligned} \ln \mathbb{E} \left[e^{F(W)} \right] &= \sup_{u \in \mathcal{U}_q} \mathbb{E} \left[- \int_0^T \frac{u_s^2}{2} ds + \int_0^T \rho m \langle \boldsymbol{\sigma}, \widehat{\mathbb{X}}_s^u \rangle (dW_s + u_s ds) + \frac{\bar{\rho}^2 m^2 - m}{2} \int_0^T \langle \boldsymbol{\sigma}, \widehat{\mathbb{X}}_s^u \rangle^2 ds \right] \\ &= \sup_{u \in \mathcal{U}_q} \mathbb{E} \left[- \int_0^T \frac{u_s^2}{2} ds + \int_0^T \rho m \langle \boldsymbol{\sigma}, \widehat{\mathbb{X}}_s^u \rangle u_s ds + \frac{\bar{\rho}^2 m^2 - m}{2} \int_0^T \langle \boldsymbol{\sigma}, \widehat{\mathbb{X}}_s^u \rangle^2 ds \right], \end{aligned}$$

where we have used that the stochastic integral has zero expectation by (A.2). \square

References

- Eduardo Abi Jaber and Louis-Amand Gérard. Signature volatility models: Pricing and hedging with fourier. *Siam Journal on Financial Mathematics*, to appear. doi: 10.2139/ssrn.4714535.
- Eduardo Abi Jaber, Martin Larsson, and Sergio Pulido. Affine Volterra processes. *The Annals of Applied Probability*, 29(5):3155–3200, 2019.
- Eduardo Abi Jaber, Camille Illand, and Shaun Xiaoyuan Li. The quintic Ornstein-Uhlenbeck volatility model that jointly calibrates SPX & VIX smiles. *Risk*, Cutting edge section, 2023.
- Leif B. G. Andersen and Vladimir V. Piterbarg. Moment explosions in stochastic volatility models. *Finance and Stochastics*, 11(1):29–50, 2007. ISSN 1432-1122. doi: 10.1007/s00780-006-0011-7. URL <https://doi.org/10.1007/s00780-006-0011-7>.
- Imanol Perez Arribas, Cristopher Salvi, and Lukasz Szpruch. Sig-SDEs model for quantitative finance. In *Proceedings of the First ACM International Conference on AI in Finance*, pages 1–8, 2020.
- Carole Bernard, Zhenyu Cui, and Don McLeish. On the martingale property in stochastic volatility models based on time-homogeneous diffusions. *Mathematical Finance*, 27(1):194–223, 2017.
- Michelle Boué and Paul Dupuis. A variational representation for certain functionals of brownian motion. *The Annals of Probability*, 26(4):1641–1659, 1998.
- Christa Cuchiero, Guido Gazzani, and Sara Svaluto-Ferro. Signature-based models: Theory and calibration. *SIAM Journal on Financial Mathematics*, 14(3):910–957, 2023. doi: 10.1137/22M1512338.
- Christa Cuchiero, Guido Gazzani, Janka Möller, and Sara Svaluto-Ferro. Joint calibration to SPX and VIX options with signature-based models. *Mathematical Finance*, 35(1):161–213, 2025. doi: <https://doi.org/10.1111/mafi.12442>. URL <https://onlinelibrary.wiley.com/doi/abs/10.1111/mafi.12442>.
- Thomas Fawcett. *Problems in stochastic analysis. Connections between rough paths and noncommutative harmonic analysis*. PhD thesis, University of Oxford, 2003.
- PK Friz, P Gassiat, and P Pigato. Precise asymptotics: Robust stochastic volatility models. *The Annals of Applied Probability*, 31(2):896, 2021.
- J. G. Gaines. The algebra of iterated stochastic integrals. *Stochastics and Stochastic Reports*, 49(3-4):169–179, 1994. doi: 10.1080/17442509408833918. URL <https://doi.org/10.1080/17442509408833918>.
- Paul Gassiat. On the martingale property in the rough bergomi model. *Electronic Communications in Probability*, 2018. URL <https://api.semanticscholar.org/CorpusID:140230875>.
- Stefan Gerhold, Christoph Gerstenecker, and Arpad Pinter. Moment explosions in the rough heston model. *Decisions in Economics and Finance*, 42, 01 2018. doi: 10.1007/s10203-019-00267-6.
- Stefan Gerhold, Julian Pachschwöll, and Johannes Ruf. On the integrability of the supremum of stochastic volatility models and other martingales, 2024.
- I. V. Girsanov. On transforming a certain class of stochastic processes by absolutely continuous substitution of measures. *Theory of Probability & Its Applications*, 5(3):285–301, 1960. doi: 10.1137/1105027. URL <https://doi.org/10.1137/1105027>.

- Archil Gulisashvili. Gaussian stochastic volatility models: Scaling regimes, large deviations, and moment explosions. *Stochastic Processes and their Applications*, 130(6):3648–3686, 2020.
- Julien Guyon and Jordan Lekeufack. Volatility is (mostly) path-dependent. *Quantitative Finance*, 23(9):1221–1258, 2023. doi: 10.1080/14697688.2023.2221281.
- Yuu Hariya and Sou Watanabe. The boué–dupuis formula and the exponential hypercontractivity in the gaussian space. *Electronic Communications in Probability*, 27:1–13, 2022.
- Antoine Jacquier and Martin Keller-Ressel. Implied volatility in strict local martingale models. *SIAM Journal on Financial Mathematics*, 9(1):171–189, 2018. doi: 10.1137/16M1069651. URL <https://doi.org/10.1137/16M1069651>.
- Benjamin Jourdain. Loss of martingality in asset price models with lognormal stochastic volatility. *Preprint CERMICS*, 08 2004.
- N. Kazamaki. *Continuous Exponential Martingales and BMO*. Lecture Notes in Mathematics. Springer Berlin Heidelberg, 2006. ISBN 9783540484219.
- P.-L. Lions and M. Musiela. Correlations and bounds for stochastic volatility models. *Annales de l’Institut Henri Poincaré C, Analyse non linéaire*, 24(1):1–16, 2007. ISSN 0294-1449. doi: <https://doi.org/10.1016/j.anihpc.2005.05.007>.
- R. C. Lyndon. On burnside’s problem. *Transactions of the American Mathematical Society*, 77(2):202–215, 1954. ISSN 00029947, 10886850. URL <http://www.jstor.org/stable/1990868>.
- Terry Lyons and Nicolas Victoir. Cubature on Wiener space. *Proceedings: Mathematical, Physical and Engineering Sciences*, 460(2041):169–198, 2004. ISSN 13645021. URL <http://www.jstor.org/stable/4143098>.
- Terry Lyons, Michael Caruana, and Thierry Lévy. *Differential Equations Driven by Rough Paths*. 01 2007. ISBN 978-3-540-71284-8. doi: 10.1007/978-3-540-71285-5.5.
- Aleksandar Mijatović and Mikhail Urusov. On the martingale property of certain local martingales. *Probability Theory and Related Fields*, 152(1):1–30, 2012. ISSN 1432-2064. doi: 10.1007/s00440-010-0314-7. URL <https://doi.org/10.1007/s00440-010-0314-7>.
- A. A. Novikov. On an identity for stochastic integrals. *Theory of Probability & Its Applications*, 17(4):717–720, 1973. doi: 10.1137/1117088. URL <https://doi.org/10.1137/1117088>.
- Marcel Nutz and Andrés Riveros Valdevenito. On the guyon–lekeufack volatility model. *Finance and Stochastics*, 28:1203–1223, 09 2024. doi: 10.1007/s00780-024-00544-2.
- David E. Radford. A natural ring basis for the shuffle algebra and an application to group schemes. *Journal of Algebra*, 58:432–454, 1979. URL <https://api.semanticscholar.org/CorpusID:121992265>.
- Rimhak Ree. Lie elements and an algebra associated with shuffles. *Annals of Mathematics*, 68(2):210–220, 1958. ISSN 0003486X. URL <http://www.jstor.org/stable/1970243>.
- Johannes Ruf. The martingale property in the context of stochastic differential equations. *Electronic Communications in Probability*, 20, January 2015. ISSN 1083-589X. doi: 10.1214/ecp.v20-3449. URL <http://dx.doi.org/10.1214/ECP.v20-3449>.
- Artur Sepp and Parviz Rakhmonov. What is a robust stochastic volatility model. *Available at SSRN*, 2023. doi: <http://dx.doi.org/10.2139/ssrn.4647027>.
- Carlos A Sin. Complications with stochastic volatility models. *Advances in Applied Probability*, 30(1):256–268, 1998.
- Daniel W Stroock and Srinivasa RS Varadhan. On the support of diffusion processes with applications to the strong maximum principle. In *Proceedings of the Sixth Berkeley Symposium on Mathematical Statistics and Probability (Univ. California, Berkeley, Calif., 1970/1971)*, volume 3, pages 333–359, 1972.