

A NOTE ON THE LONG TIME BEHAVIOR OF THE ELEPHANT RANDOM WALK WITH STOPS

TATSUYA AKIMOTO, MASATO TAKEI, AND KEISUKE TANIGUCHI

ABSTRACT. We study the long time behavior of the elephant random walk with stops, introduced by Kumar, Harbola and Lindenberg (2010), and establish the phase transition of the number of visited points up to time n , and the correlation between the position at time n and the number of moves up to time n .

1. INTRODUCTION

Recently there has been considerable interest in the elephant random walk (ERW), introduced by Schütz and Trimper [7] (see Laulin [5] and references therein). In this note we study the elephant random walk with stops (ERWS), which is a crucial generalization introduced by Kumar, Harbola and Lindenberg [4]. Let $s \in [0, 1]$. Assume that $p, q \in [0, 1]$ and $r \in [0, 1]$ satisfy $p + q + r = 1$. The first step X_1 of the walker is $+1$ with probability s , and -1 with probability $1 - s$. For each $n \in \mathbb{N} := \{1, 2, \dots\}$, let U_n be uniformly distributed on $\{1, \dots, n\}$, and

$$X_{n+1} = \begin{cases} X_{U_n} & \text{with probability } p, \\ -X_{U_n} & \text{with probability } q, \\ 0 & \text{with probability } r. \end{cases} \quad (1.1)$$

Each of choices in the above procedure is made independently. A one-dimensional random walk $\{S_n\}$ is defined by $S_0 := 0$, and $S_n = \sum_{i=1}^n X_i$ for $n \in \mathbb{N}$. The standard ERW corresponds to the special case $r = 0$.

It turns out to be convenient to introduce the following new parameters: $a := p - q$, and $b := 1 - r$. Note that $b \in (0, 1]$ and $a \in [-b, b]$.

An important quantity for the ERWS is the number Σ_n of moves up to time n , i.e.,

$$\Sigma_0 := 0, \quad \text{and} \quad \Sigma_n := \sum_{i=1}^n 1_{\{X_i \neq 0\}} = \sum_{i=1}^n X_i^2 \quad \text{for } n \in \mathbb{N}. \quad (1.2)$$

Let \mathcal{F}_n be the σ -algebra generated by X_1, \dots, X_n . Since $X_{n+1}^2 \in \{0, 1\}$ and $P(X_{n+1}^2 = 1 \mid \mathcal{F}_n) = (b\Sigma_n)/n$ for $n \in \mathbb{N}$, the process $\{\Sigma_n\}$ is nothing but the ‘laziest’ minimal random walk model of elephant type with parameter b , studied in detail in Miyazaki and Takei [6]. By Corollary 1 in [6] (see also Lemma 2.1 in [1]),

$$\lim_{n \rightarrow \infty} \frac{\Sigma_n}{n^b} = \Sigma > 0 \quad \text{a.s.}, \quad (1.3)$$

where Σ has a Mittag-Leffler distribution with parameter b .

In a remarkable paper [1], Bercu obtained several limit theorems for the ERWS $\{S_n\}$. Among other results in [1], we quote the central limit theorem and the law of the iterated logarithm. For a fixed $b \in (0, 1]$, the critical value for $a \in [-b, b]$ is given by $a_c(b) := b/2$.

Theorem 1.1 (Bercu [1]). *Let $\phi(x) := \sqrt{2x \log \log x}$.*

(i) (Subcritical regime) *If $a \in [-b, b/2]$ then $\frac{S_n}{\sqrt{\Sigma_n}} \xrightarrow{d} N\left(0, \frac{b}{b-2a}\right)$ as $n \rightarrow \infty$, and $\limsup_{n \rightarrow \infty} \pm \frac{S_n}{\phi(\Sigma_n)} = \sqrt{\frac{b}{b-2a}}$ a.s..*

(ii) (Critical regime) *If $a = b/2$ then $\frac{S_n}{\sqrt{\Sigma_n \log \Sigma_n}} \xrightarrow{d} N(0, 1)$ as $n \rightarrow \infty$, and*

$\limsup_{n \rightarrow \infty} \pm \frac{S_n}{\phi(\Sigma_n \log \Sigma_n)} = 1$ a.s.

(iii) (Supercritical regime) *If $a \in (b/2, b]$ then there exists a random variable L such that $P(L \neq 0) > 0$ and $\lim_{n \rightarrow \infty} \frac{S_n}{n^a} = L$ a.s.*

Remark 1.2. *This theorem implies that $\{S_n\}$ is recurrent [resp. transient] when $a \in [-b, b/2]$ [resp. $a \in (b/2, b]$]: More precisely, $S_n = 0$ for infinitely many n with probability one if $a \in [-b, b/2]$, while $S_n = 0$ for only finitely many n with positive probability if $a \in (b/2, b]$. For the standard ERW (i.e. $b = 1$), it is shown in [3] that $P(L \neq 0) = 1$.*

In this note, we carry out a further study on the phase transition concerning the long time behavior of the ERWS. Our first result in subsection 2.1 shows that the range of the ERWS admits phase transition. To our best knowledge, this is the first result on the range of all versions of elephant random walks. In subsection 2.2, we show that the correlation coefficient of S_n and Σ_n exhibits a non-trivial phase transition when $r \in (0, 1)$ i.e. the walker can stay on his own position with positive probability, and the number Σ_n of moves up to time n is random. Note that when $r = 0$ the model is nothing but the original ERW, and the correlation coefficient of S_n and $\Sigma_n \equiv n$ is zero for all $a \in [-1, 1]$. Proofs are given in Sections 3, 4 and 5.

2. MAIN RESULTS

2.1. Almost sure range of the elephant random walk with stops.

For the ERWS $\{S_n\}$, we define the range $\{R_n\}$ of by $R_0 := 1$ and $R_n := \#\{S_0, S_1, \dots, S_n\}$ for $n \in \mathbb{N}$. Combining the strong law of large numbers for $\{S_n\}$ obtained by [1] and Theorem 2 in [2], we have $\lim_{n \rightarrow \infty} \frac{R_n}{n} = 0$ a.s. for any $b \in (0, 1]$ and $a \in [-b, b]$. The following theorem is a considerable improvement, which shows that the range admits phase transition at $a = a_c(b) = b/2$.

Theorem 2.1. *Assume that $b \in (0, 1]$.*

(i) *If $a \in [-b, b/2]$ then $\sqrt{\frac{b}{b-2a}} \leq \limsup_{n \rightarrow \infty} \frac{R_n}{\phi(\Sigma_n)} \leq 2\sqrt{\frac{b}{b-2a}}$ a.s..*

(ii) *If $a = b/2$ then $1 \leq \limsup_{n \rightarrow \infty} \frac{R_n}{\phi(\Sigma_n \log \Sigma_n)} \leq 2$ a.s..*

(iii) If $a \in (b/2, b]$, then $\lim_{n \rightarrow \infty} \frac{R_n}{n^a} = |L|$ a.s., where L is the limiting random variable in Theorem 1.1 (iii).

2.2. Correlation between the position and the number of moves.

In this subsection, we assume that $b = 1 - r \in (0, 1)$, and thus the number of moves Σ_n up to time n is a nondegenerate random variable. Our interest is how big the correlation between S_n and Σ_n is, and the next theorem implies that it is drastically changed at $a = a_c(b) = b/2$.

The correlation coefficient of X and Y is denoted by $\rho[X, Y]$. The beta function is defined by

$$B(x, y) := \int_0^1 u^{x-1}(1-u)^{y-1} du \quad \text{for } x, y > 0.$$

Theorem 2.2. *Assume that $b \in (0, 1)$. If $a = 0$ or $s = 1/2$, then $\rho[S_n, \Sigma_n] = 0$ for all $n \in \mathbb{N}$. If $a \neq 0$ and $s \neq 1/2$, then as $n \rightarrow \infty$,*

$$\rho[S_n, \Sigma_n] \sim \begin{cases} (2s-1) \cdot P_{a,b} n^{-(b-2a)/2} & \text{if } a \in [-b, b/2), \\ (2s-1) \cdot Q_a (\log n)^{-1/2} & \text{if } a = b/2, \\ (2s-1) \cdot R_{a,b,s} & \text{if } a \in (b/2, b]. \end{cases}$$

Here $P_{a,b}$, Q_a and $R_{a,b,s}$ are constants given by

$$\begin{aligned} P_{a,b} &= \frac{\sqrt{(b-2a)\Gamma(b)} \cdot \{(a+b) \cdot B(a+1, b) - 1\}}{\Gamma(a+1) \cdot \sqrt{b \cdot B(b, b) - 1}}, \\ Q_a &= \frac{\sqrt{\Gamma(2a)} \cdot \{a \cdot B(a, 2a) - 1\}}{\Gamma(a+1) \cdot \sqrt{2a \cdot B(2a, 2a) - 1}}, \\ R_{a,b,s} &= \frac{\sqrt{2a-b} \cdot \{a \cdot B(a, b) - 1\}}{\sqrt{\{a^2 \cdot B(a, a) - (2s-1)^2 \cdot (2a-b)\} \{b \cdot B(b, b) - 1\}}}. \end{aligned}$$

Note that $P_{a,b}$ is positive [resp. negative] for $a \in (0, b/2)$ [resp. $a \in [-b, 0)$] while Q_a and $R_{a,b,s}$ are positive.

By Theorem 2.2, the correlation between S_n and Σ_n vanishes as $n \rightarrow \infty$ when $a \in [-b, b/2]$, while it remains positive when $a \in (b/2, b]$. This is in line with our expectations, in view of the recurrence property of the ERWS, described in Remark 1.2. The (sub)critical ERWS is recurrent and after having completed an excursion (that is when $S_n = 0$) the walk has the same number of +1 and -1 steps in its history. Hence the probability of moving to +1 is the same as going to -1 independently of the past, that is, the signs of the excursions are determined by independent coin flips. It shows that in these regimes, the random sign of S_n causes cancellation in a way which makes S_n asymptotically independent of Σ_n . In the supercritical regime, the walk is transient and after the last return to 0, the more non-zero steps the walk takes the further it gets away from 0.

On the other hand, the next theorem says that S_n^2 and Σ_n has a positive correlation for all $a \in [-b, b]$, which means that conditioning on Σ_n gives non-trivial information on S_n^2 and hence on $|S_n|$.




FIGURE 1. The graphs of $\rho = \lim_{n \rightarrow \infty} \rho[S_n^2, \Sigma_n]$ as a function of $a \in [-b, b]$ for $b = 0.3, 0.6$ and 0.9 . Note that the critical value is $a_c(b) = b/2$.

Theorem 2.3. For $b \in (0, 1)$,

$$\lim_{n \rightarrow \infty} \rho[S_n^2, \Sigma_n] = \begin{cases} Q'_b & \text{if } a \in [-b, b/2], \\ R'_{a,b} & \text{if } a \in (b/2, b]. \end{cases} \quad (2.1)$$

Here Q'_b and $R'_{a,b}$ are positive constants given by

$$Q'_b = \sqrt{\frac{b \cdot B(b, b) - 1}{3b \cdot B(b, b) - 1}}, \quad R'_{a,b} = \frac{2a \cdot B(2a, b) - 1}{\sqrt{\left\{ \frac{6(2a^2 + 2ab - b^2)}{4a - b} \cdot B(2a, 2a) - 1 \right\} \{b \cdot B(b, b) - 1\}}}.$$

The limit in (2.1) can be regarded as a function of $a \in [-b, b]$ for fixed $b \in (0, 1)$: See Figure 1.

3. PROOF OF THEOREM 2.1

Theorem 2.1 is a consequence of Theorem 1.1 and the following lemma, which is an extension of Theorem 2 in Boudabra and Markowsky [2], and is of independent interest.

Lemma 3.1. Let $\{x_n\}$ be an integer valued sequence with $x_n - x_{n-1} \in \{-1, 0, +1\}$ for each $n \in \mathbb{N}$. We define $\{r_n\}$ by $r_0 := 1$ and $r_n := \#\{x_0, x_1, \dots, x_n\}$ for $n \in \mathbb{N}$. Let $f(n)$ be a positive real valued function defined for all sufficiently large $n \in \mathbb{Z}_+ := \{0, 1, 2, \dots\}$, which is increasing and satisfies that $\lim_{n \rightarrow \infty} f(n) = +\infty$.

- (i) If $\lim_{n \rightarrow \infty} \frac{x_n}{f(n)} = c$, then $\lim_{n \rightarrow \infty} \frac{r_n}{f(n)} = |c|$.
- (ii) If $\limsup_{n \rightarrow \infty} \pm \frac{x_n}{f(n)} = 1$, then $1 \leq \limsup_{n \rightarrow \infty} \frac{r_n}{f(n)} \leq 2$.

Proof. Without loss of generality, we can assume that $x_0 = 0$.

(i) Suppose that $c > 0$. For any $\varepsilon \in (0, c)$, there exists N s.t. $(0 <)(c - \varepsilon)f(n) \leq x_n \leq (c + \varepsilon)f(n)$ for all $n \geq N$. If $n \geq N$ then $\{x_0, x_1, \dots, x_n\} \supset$

$\{0, \dots, \lfloor (c - \varepsilon)f(n) \rfloor\}$ and $\{x_N, \dots, x_n\} \subset [0, (c + \varepsilon)f(n)]$, which imply that $\lfloor (c - \varepsilon)f(n) \rfloor \leq r_n \leq r_{N-1} + (c + \varepsilon)f(n)$. Thus we have $c - \varepsilon \leq \liminf_{n \rightarrow \infty} \frac{r_n}{f(n)} \leq \limsup_{n \rightarrow \infty} \frac{r_n}{f(n)} \leq c + \varepsilon$. Letting $\varepsilon \searrow 0$, we obtain the desired conclusion when $c > 0$. For the case $c < 0$, consider $\{-x_n\}$. Next suppose that $c = 0$. For any $\varepsilon > 0$, there exists N such that $-\varepsilon f(n) \leq x_n \leq \varepsilon f(n)$ for all $n \geq N$. Since $\{x_N, \dots, x_n\} \subset [-\varepsilon f(n), \varepsilon f(n)]$ for all $n \geq N$, we have $0 \leq \liminf_{n \rightarrow \infty} \frac{r_n}{f(n)} \leq \limsup_{n \rightarrow \infty} \frac{r_n}{f(n)} \leq 2\varepsilon$. This completes the proof of (i). (ii) For any $\varepsilon > 0$, there exists N such that $-(1 + \varepsilon)f(n) \leq x_n \leq (1 + \varepsilon)f(n)$ for all $n \geq N$. Noting that $\{x_N, \dots, x_n\} \subset [-(1 + \varepsilon)f(n), (1 + \varepsilon)f(n)]$ for all $n \geq N$, we have $r_n \leq r_{N-1} + 2(1 + \varepsilon)f(n)$ for all $n \geq N$, and $\limsup_{n \rightarrow \infty} \frac{r_n}{f(n)} \leq 2(1 + \varepsilon)$. On the other hand, for any $\varepsilon \in (0, 1)$, $x_n \geq (1 - \varepsilon)f(n) (> 0)$ for infinitely many n . As $\{x_0, x_1, \dots, x_n\} \supset \{0, \dots, \lfloor (1 - \varepsilon)f(n) \rfloor\}$ for infinitely many n , we have $r_n \geq \lfloor (1 - \varepsilon)f(n) \rfloor$ for infinitely many n , and $\limsup_{n \rightarrow \infty} \frac{r_n}{f(n)} \geq 1 - \varepsilon$. This completes the proof of (ii). \square

4. PROOF OF THEOREM 2.2

For $x > -1$ and $n \in \mathbb{N}$, we define

$$c_n(x) := \prod_{k=1}^{n-1} \left(1 + \frac{x}{k}\right) = \frac{\Gamma(n+x)}{\Gamma(n)\Gamma(x+1)} \sim \frac{n^x}{\Gamma(x+1)} \quad \text{as } n \rightarrow \infty, \quad (4.1)$$

where we used Stirling's formula.

Lemma 4.1. *Assume that $b \in (0, 1)$ and $a \in [-b, b]$. If $a = 0$ or $s = 1/2$, then $\text{Cov}[S_n, \Sigma_n] = 0$ for all $n \in \mathbb{N}$. If $a \neq 0$ and $s \neq 1/2$, then*

$$\text{Cov}[S_n, \Sigma_n] \sim \frac{2s-1}{\Gamma(a+1)\Gamma(b+1)} \cdot \{(a+b) \cdot B(a+1, b) - 1\} \cdot n^{a+b} \quad \text{as } n \rightarrow \infty.$$

Proof. Eqs. (C.17), (4.20) and (C.19) in [1] can be read as follows:

$$E[S_n] = (2s-1) \cdot c_n(a), \quad (4.2)$$

$$E[\Sigma_n] = c_n(b), \quad (4.3)$$

$$E[S_n \Sigma_n] = \frac{2s-1}{b} \cdot \{(a+b) \cdot c_n(a+b) - a \cdot c_n(a)\}. \quad (4.4)$$

If $a = 0$ then $E[S_n \Sigma_n] = (2s-1) \cdot c_n(b) = E[S_n]E[\Sigma_n]$ for all $n \in \mathbb{N}$. If $s = 1/2$ then $E[S_n \Sigma_n] = 0 = E[S_n]E[\Sigma_n]$ for all $n \in \mathbb{N}$. If $a \neq 0$ and $s \neq 1/2$, then by (4.1),

$$E[S_n \Sigma_n] \sim \frac{2s-1}{b\Gamma(a+b)} \cdot n^{a+b} \quad \text{and} \quad E[S_n]E[\Sigma_n] \sim \frac{2s-1}{\Gamma(a+1)\Gamma(b+1)} \cdot n^{a+b},$$

as $n \rightarrow \infty$. This completes the proof. \square

Lemma 4.2. *Assume that $b \in (0, 1]$. As $n \rightarrow \infty$,*

$$V[S_n] \sim \begin{cases} \frac{1}{(b-2a)\Gamma(b)} \cdot n^b & \text{if } a \in [-b, b/2), \\ \frac{1}{\Gamma(2a)} \cdot n^{2a} \log n & \text{if } a = b/2, \\ \frac{a^2 \cdot B(a, a) - (2s-1)^2 \cdot (2a-b)}{(2a-b)\Gamma(a+1)^2} \cdot n^{2a} & \text{if } a \in (b/2, b]. \end{cases}$$

Proof. From the calculation leading Eq. (A.6) in [1],

$$E[S_n^2] = \begin{cases} \frac{b \cdot c_n(b) - 2a \cdot c_n(2a)}{b - 2a} & \text{if } a \neq b/2, \\ 2a \cdot c_n(2a) \cdot \sum_{k=1}^n \frac{1}{k+2a} & \text{if } a = b/2. \end{cases} \quad (4.5)$$

By (4.1), we have as $n \rightarrow \infty$,

$$E[S_n^2] \sim \begin{cases} \frac{1}{(b-2a)\Gamma(b)} \cdot n^b & \text{if } a \in [-b, b/2), \\ \frac{1}{\Gamma(2a)} \cdot n^{2a} \log n & \text{if } a = b/2. \\ \frac{1}{(2a-b)\Gamma(2a)} \cdot n^{2a} & \text{if } a \in (b/2, b]. \end{cases} \quad (4.6)$$

On the other hand, by (4.2) and (4.1), $E[S_n]^2 \sim (2s-1)^2 n^{2a} / \{\Gamma(a+1)\}^2$ as $n \rightarrow \infty$. Now the desired conclusion follows. \square

Lemma 4.3. *If $b \in (0, 1)$ then $V[\Sigma_n] \sim \frac{b \cdot B(b, b) - 1}{\Gamma(b+1)^2} \cdot n^{2b}$ as $n \rightarrow \infty$.*

Proof. From Eq. (4.20) in [1],

$$E[\Sigma_n(\Sigma_n + 1)] = 2c_n(2b) \quad \text{for } n \in \mathbb{N}. \quad (4.7)$$

We have $V[\Sigma_n] = E[\Sigma_n^2] - E[\Sigma_n]^2 = \{2c_n(2b) - c_n(b)\} - \{c_n(b)\}^2$, which together with (4.1) yields the claim. \square

Combining the above lemmata, we have Theorem 2.2. For $a \in [-b, 0)$, since $a(b-1) > 0$, we have $(a+b) \cdot B(a+1, b) \leq (a+b)/\{(a+1)b\} < 1$, and thus $P_{a,b} < 0$. As for $a \in (0, b/2)$, noting that $(a+b) \cdot B(a+1, b) = a \cdot B(a, b) > a \cdot B(a, 1) = 1$, we have $P_{a,b} > 0$. Similarly we see that $Q_a, R_{a,b,s} > 0$.

5. PROOF OF THEOREM 2.3

Lemma 5.1. *For $b \in (0, 1)$,*

$$\text{Cov}[S_n^2, \Sigma_n] \sim \begin{cases} \frac{b \cdot B(b, b) - 1}{b(b-2a)\Gamma(b)^2} \cdot n^{2b} & \text{if } a \in [-b, b/2), \\ \frac{b \cdot B(b, b) - 1}{b\Gamma(b)^2} \cdot n^{2b} \log n & \text{if } a = b/2, \\ \frac{2a \cdot B(2a, b) - 1}{b(2a-b)\Gamma(2a)\Gamma(b)} \cdot n^{2a+b} & \text{if } a \in (b/2, b]. \end{cases}$$

Proof. First we show that

$$E[S_n^2 \Sigma_n] \sim \begin{cases} \frac{1}{(b-2a)\Gamma(2b)} \cdot n^{2b} & \text{if } a \in [-b, b/2), \\ \frac{1}{\Gamma(2b)} \cdot n^{2b} \log n & \text{if } a = b/2, \\ \frac{2a}{b(2a-b)\Gamma(2a+b)} \cdot n^{2a+b} & \text{if } a \in (b/2, b]. \end{cases} \quad (5.1)$$

Since

$$\begin{aligned} E[S_{n+1}^2 \Sigma_{n+1} | \mathcal{F}_n] &= E[(S_n + X_{n+1})^2 (\Sigma_n + X_{n+1}^2) | \mathcal{F}_n] \\ &= \left(1 + \frac{2a+b}{n}\right) \cdot S_n^2 \Sigma_n + \frac{2a}{n} \cdot S_n^2 + \frac{b}{n} \cdot \Sigma_n (\Sigma_n + 1), \end{aligned}$$

we have

$$E[S_{n+1}^2 \Sigma_{n+1}] = \left(1 + \frac{2a+b}{n}\right) \cdot E[S_n^2 \Sigma_n] + \frac{2a}{n} \cdot E[S_n^2] + \frac{b}{n} \cdot E[\Sigma_n (\Sigma_n + 1)].$$

Setting $J_n := S_n^2 \Sigma_n / c_n(2a+b)$, we have

$$E[J_{n+1}] - E[J_n] = \frac{2a \cdot E[S_n^2]}{n \cdot c_{n+1}(2a+b)} + \frac{b \cdot E[\Sigma_n (\Sigma_n + 1)]}{n \cdot c_{n+1}(2a+b)},$$

and

$$E[J_n] = E[J_1] + \sum_{k=1}^{n-1} \frac{2a \cdot E[S_k^2]}{k \cdot c_{k+1}(2a+b)} + \sum_{k=1}^{n-1} \frac{b \cdot E[\Sigma_k (\Sigma_k + 1)]}{k \cdot c_{k+1}(2a+b)}. \quad (5.2)$$

From the definition of X_1 and (1.2), $E[J_1] = 1$. It is straightforward to prove that

$$\sum_{k=1}^{n-1} \frac{c_k(x)}{k \cdot c_{k+1}(y)} = \frac{1}{x-y} \cdot \left\{ \frac{c_n(x)}{c_n(y)} - 1 \right\} \quad \text{for } x, y > -1 \text{ with } x \neq y. \quad (5.3)$$

For the case $a \neq b/2$, using (4.5), (4.7), (5.2), and (5.3), we see that

$$\begin{aligned} E[S_n^2 \Sigma_n] &= \frac{1}{b-2a} \left\{ 2b \cdot c_n(2b) - b \cdot c_n(b) + \frac{2a}{b} (2a \cdot c_n(2a) - (2a+b) \cdot c_n(2a+b)) \right\}, \end{aligned} \quad (5.4)$$

which coincides with Eq. (C.23) in [1]. In view of (4.1), the first (resp. fourth) term in the right hand side of (5.4) is dominant when $a \in [-b, b/2)$ (resp. $a \in (b/2, b]$). This completes the proof of (5.1) for the case $a \neq b/2$. For the critical case $a = b/2$, by (4.1), (4.6), and (4.7), in the right hand side of (5.2), the third term is dominant. Since $2a = b$, we have

$$E[S_n^2 \Sigma_n] \sim c_n(2b) \cdot \sum_{k=1}^{n-1} \frac{2b \cdot c_k(2b)}{k \cdot c_{k+1}(2b)} \sim \frac{1}{\Gamma(2b)} \cdot n^{2b} \log n \quad \text{as } n \rightarrow \infty,$$

which is (5.1) for $a = b/2$. Finally, by (4.1), (4.3), and (4.6), we have

$$E[S_n^2]E[\Sigma_n] \sim \begin{cases} \frac{1}{b(b-2a)\Gamma(b)^2} \cdot n^{2b} & \text{if } a \in [-b, b/2), \\ \frac{1}{b\Gamma(b)^2} \cdot n^{2b} \log n & \text{if } a = b/2, \\ \frac{1}{b(2a-b)\Gamma(2a)\Gamma(b)} \cdot n^{2a+b} & \text{if } a \in (b/2, b]. \end{cases} \quad (5.5)$$

The conclusion of Lemma 5.1 follows from (5.1) and (5.5). \square

Lemma 5.2. *If $b \in (0, 1]$ then as $n \rightarrow \infty$,*

$$V[S_n^2] \sim \begin{cases} \frac{3b \cdot B(b, b) - 1}{(b-2a)^2 \Gamma(b)^2} \cdot n^{2b} & \text{if } a \in [-b, b/2), \\ \frac{3b \cdot B(b, b) - 1}{\Gamma(b)^2} \cdot n^{2b} (\log n)^2 & \text{if } a = b/2, \\ \frac{1}{(2a-b)^2 \Gamma(2a)^2} \cdot \left\{ \frac{6(2a^2 + 2ab - b^2)}{4a-b} \cdot B(2a, 2a) - 1 \right\} \cdot n^{4a} & \text{if } a \in (b/2, b]. \end{cases}$$

Proof. First we prove

$$E[S_n^4] \sim \begin{cases} \frac{3b}{(b-2a)^2 \Gamma(2b)} \cdot n^{2b} & \text{if } a \in [-b, b/2), \\ \frac{3b}{\Gamma(2b)} \cdot n^{2b} (\log n)^2 & \text{if } a = b/2, \\ \frac{6(2a^2 + 2ab - b^2)}{(2a-b)^2 (4a-b) \Gamma(4a)} \cdot n^{4a} & \text{if } a \in (b/2, b]. \end{cases} \quad (5.6)$$

Our starting point is Eq. (C.22) in [1], i.e.

$$E[S_{n+1}^4] = \left(1 + \frac{4a}{n}\right) \cdot E[S_n^4] + \frac{6b}{n} \cdot E[S_n^2 \Sigma_n] + \frac{4a}{n} \cdot E[S_n^2] + \frac{b}{n} \cdot E[\Sigma_n].$$

Putting $L_n := S_n^4/c_n(4a)$, we see that

$$E[L_n] = 1 + \sum_{k=1}^{n-1} \frac{6b \cdot E[S_k^2 \Sigma_k]}{k \cdot c_{k+1}(4a)} + \sum_{k=1}^{n-1} \frac{4a \cdot E[S_k^2]}{k \cdot c_{k+1}(4a)} + \sum_{k=1}^{n-1} \frac{b \cdot E[\Sigma_k]}{k \cdot c_{k+1}(4a)}. \quad (5.7)$$

For the case $a \neq b/2$, using (5.4), (4.5), (4.3), and (5.3), we have

$$\begin{aligned} E[S_n^4] &= \frac{24a(2a^2 + 2ab - b^2)}{(2a-b)^2(4a-b)} \cdot c_n(4a) - \frac{12a(2a+b)}{(2a-b)^2} \cdot c_n(2a+b) \\ &\quad + \frac{8a}{2a-b} \cdot c_n(2a) + \frac{6b^2}{(2a-b)^2} \cdot c_n(2b) - \frac{b(5b-2a)}{(2a-b)(4a-b)} \cdot c_n(b), \end{aligned} \quad (5.8)$$

which coincides with Eq. (C.24) in [1]. By (4.1), the fourth (resp. first) term in the right hand side of (5.8) is dominant when $a \in [-b, b/2)$ (resp. $a \in (b/2, b]$). This completes the proof of (5.6) for the case $a \neq b/2$. For the critical case $a = b/2$, from (4.1), (5.1), (4.6), and (4.7), the first term is

dominant in the right hand side of (5.7). Noting that $4a = 2b$, we have

$$E[S_n^4] \sim c_n(2b) \cdot \sum_{k=1}^{n-1} \frac{6b \cdot E[S_k^2 \Sigma_k]}{k \cdot c_{k+1}(2b)} \sim \frac{6bn^{2b}}{\Gamma(2b)} \cdot \sum_{k=1}^{n-1} \frac{\log k}{k} \sim \frac{3b}{\Gamma(2b)} \cdot n^{2b} (\log n)^2$$

as $n \rightarrow \infty$, which is (5.6) for $a = b/2$. Lemma 5.2 follows from (5.6) and (4.6). \square

Theorem 2.3 follows from Lemmata 5.1, 5.2, and 4.3.

ACKNOWLEDGEMENT

The authors thank the referee for helpful and constructive comments, in particular on the heuristics behind our results in Section 2.2. M.T. is partially supported by JSPS KAKENHI Grant Numbers JP19H01793, JP19K03514 and JP22K03333.

REFERENCES

- [1] Bercu, B. (2022). On the elephant random walk with stops playing hide and seek with the Mittag-Leffler distribution, *J. Stat. Phys.*, **189**, Paper No. 12.
- [2] Boudabra, M. and Markowsky, G. (2021). Remarks on the speeds of a class of random walks on the integers, *Discrete Math.*, **344**, 112436
- [3] Guérin, H., Laulin, L., and Raschel, K. (2023). A fixed-point equation approach for the superdiffusive elephant random walk, to appear in *Ann. Inst. Henri Poincaré, Probab. Stat.*, arXiv:2308.14630
- [4] Kumar, N., Harbola, U., and Lindenberg, K. (2010). Memory-induced anomalous dynamics: Emergence of diffusion, *Phys. Rev. E*, **82**, 021101
- [5] Laulin, L. (2022). Autour de la marche aléatoire de l'éléphant (About the elephant random walk), thèse, Université de Bordeaux, HAL Id: tel-03708512
- [6] Miyazaki, T. and Takei, M. (2020). Limit theorems for the 'laziest' minimal random walk model of elephant type, *J. Stat. Phys.*, **181**, 587–602.
- [7] Schütz, G. M. and Trimper, S. (2004). Elephants can always remember: Exact long-range memory effects in a non-Markovian random walk, *Phys. Rev. E*, **70**, 045101

GRADUATE SCHOOL OF ENGINEERING SCIENCE, YOKOHAMA NATIONAL UNIVERSITY,
79-5 TOKIWADAI, HODOGAYA-KU, YOKOHAMA, 240-8501, KANAGAWA, JAPAN

DEPARTMENT OF APPLIED MATHEMATICS, FACULTY OF ENGINEERING, YOKOHAMA
NATIONAL UNIVERSITY, 79-5 TOKIWADAI, HODOGAYA-KU, YOKOHAMA, 240-8501, KANA-
GAWA, JAPAN

Email address: takei-masato-fx@ynu.ac.jp

GRADUATE SCHOOL OF ENGINEERING SCIENCE, YOKOHAMA NATIONAL UNIVERSITY,
79-5 TOKIWADAI, HODOGAYA-KU, YOKOHAMA, 240-8501, KANAGAWA, JAPAN