

SMALL EXTENSIONS OF ANALYTIC FIELDS

MICHAEL TEMKIN

ABSTRACT. An extension K/k of analytic (i.e. real valued complete) fields is called small if it is topologically-algebraically generated by finitely many elements. We prove that this property is inherited by subextensions and hence topological generating degree of such extensions is monotonic. Much more detailed results are obtained in the case of degree one. Let k be an analytic algebraically closed field of positive residual characteristic p and $K = \widehat{k(t)^a}$ with a non-trivial valuation. In a previous work it was shown that the set $I_{K/k}$ of intermediate complete algebraically closed subextensions $k \subseteq F \subseteq K$ is totally ordered by inclusion. In this paper we show that $I_{K/k}$ is an interval parameterized by the distance between t and F . Moreover, logarithmic parameterizations induced by other generators differ by PL functions with slopes in $p^{\mathbb{Z}}$ and corners in $|K^\times|$, so $I_{K/k}$ acquires a natural PL structure.

1. INTRODUCTION

1.1. **Recollections.** Let us recall main results about topological degree that were proved in [Tem21]. These results will be used without further referencing throughout the paper. We will use the abbreviation *t.a.* instead of topologically algebraically or topgebraically, as in [Tem21]. Given an extension L/K of analytic fields, a set $S \subset L$ is called *t.a. independent over K* if $\widehat{K(S')^a} \subsetneq \widehat{K(S)^a}$ for any $S' \subsetneq S$, and it is called *t.a. generating* if $\widehat{L^a} = \widehat{k(S)^a}$. The minimum of cardinals of t.a. generating sets of L/K is called the *topological generating degree* $\text{Tr.c}(L/K)$ and the supremum of all cardinals of t.a. independent sets is called the *topological transcendence degree* $\text{tr.c}(L/K)$. For shortness, we use notation similar to Huber's topological transcendence degree tr.c. , see [Hub96, §1.8] and [Sch22, §21], but we warn the reader that our terminology is different – Huber's transcendence degree is the generating degree in our meaning which is denoted Tr.c. Since we consider the two cardinals, our terminology seems more natural. By *small* extensions we mean t.a. finitely generated extensions, that is, extensions L/K with $\text{Tr.c}(L/K) < \infty$. Naturally, the extensions with $\text{Tr.c}(L/K) = \infty$ will be called *large*.

It is almost obvious that for a tower $F/L/K$ one invariant is monotonic and another one is subadditive: $\text{tr.c}(L/K) \leq \text{tr.c}(F/K)$ and $\text{Tr.c}(F/K) \leq \text{Tr.c}(L/K) + \text{Tr.c}(F/L)$, but any other property is not clear at all. In fact, one of the main goals of this paper and my motivation to continue the research of [Tem21] was proving Theorem 2.3.2, which states that Tr.c is monotonic for small extensions and essentially reduces to proving that being a small extension is preserved by passing to subextensions. This answers a natural question in Berkovich and adic geometries which was (somewhat implicitly) open since their foundation, and was

Key words and phrases. Non-archimedean fields, topological transcendence degree.

This research was supported by ERC Consolidator Grant 770922 - BirNonArchGeom, and by the Israel Science Foundation Grant 1203/22.

explicitly stated by Scholze in [Sch22, Question 21.4]. I am very grateful to Peter Scholze for an email exchange and pointing out that the results of [Tem21] were not sufficient to completely answer this question.

It is shown in [Tem21] that a version of exchange lemma holds for t.a. dependency, hence $\text{tr.c}(L/K) \leq \text{Tr.c}(L/K)$, see [Tem21, Theorem 3.2.1] and its proof. Of course, the two cardinals are equal if there exists a topological transcendence basis – a set, which is both t.a. independent and generating, and it always exists for small extensions by [Tem21, Theorem 3.2.3], but large extensions, such as spherical completions, tend not to have such a basis, see [Tem21, Theorem 5.3.2]. If the residual characteristic $p = \text{char}(\tilde{K})$ is positive, the situation is even stranger as there exists the following pathologies, which are essentially equivalent: there exist non-invertible \widehat{K}^a -endomorphisms of $\widehat{K}(x)^a$, there exist towers $\widehat{K}^a \subsetneq L = \widehat{K}(x)^a \subsetneq F = \widehat{K}(y)^a$, there exist large extensions L/K which satisfy $\text{tr.c}(L/K) = 1$. In particular, the topological transcendence degree is not additive in towers even for small extensions. Finally, it was observed in [Tem21, Theorem 5.3.9] that if L and K are algebraically closed and $\text{tr.c}(L/K) = 1$, then the set $I_{L/K} = \{F_i\}$ of analytic algebraically closed intermediate fields $K \subseteq F_i \subseteq L$ is totally ordered.

1.2. Main results. In this paper we answer questions about extensions of finite topological transcendence degree, which were left open or unexplored in [Tem21]. In particular, we prove that if L/K is a small extension of analytic algebraically closed fields, then any intermediate algebraically closed analytic field F is small over K , and hence satisfies $\text{Tr.c}(F/K) \leq \text{Tr.c}(L/K)$, see Theorem 2.3.2. Moreover, the equality holds if and only if F and L are K -isomorphic, see Theorem 2.3.4. In fact, the claim about small extensions follows by applying Baire category theorem, once we prove the following result, which is of its own interest: any chain of algebraically closed analytic intermediate fields has at most countable cofinality, see Theorem 2.2.6. The latter result is reduced to the case of topological transcendence degree one, already covered by Lemma 2.2.3, but later on we prove in Theorem 3.1.13 the following complete and simple description of $I_{L/K}$ which was unanticipated in [Tem21]: any t.a. generator $t \in L$ induces a bijection $\rho_t: I_{L/K} \xrightarrow{\sim} [0, r_K(t)]$ by sending an intermediate field F to $r_F(t)$. In fact, this is a really simple result, which was overlooked in [Tem21] and noticing it recently made all progress of the current paper possible.

Furthermore, we study the following natural question in §3: what is the natural structure of the interval $I_{L/K}$ for $L = \widehat{K}(t)^a$? We identify it with $[-\log(r), \infty)$ by $-\log \circ \rho_t$ and show in Theorem 3.2.12 that any other t.a. generator t' induces a parametrization which differs from this one by a PL-function with slopes in $p^{\mathbb{Z}}$ and corners in $|L^\times|$ (satisfying an addition restriction at the endpoint when $|K^\times| \neq |L^\times|$). Moreover, in Theorem 3.2.16 we show that the converse is also true, obtaining a complete characterization of the set of possible parameterizations of $I_{L/K}$. In particular, $I_{L/K}$ acquires a natural $\mathbb{Z}[\frac{1}{p}]$ -structure.

Finally, a good control on the subfields of small extensions enables us to study in §4 large extensions L/K of finite topological transcendence degree and the set of their algebraically closed subfields. In particular, in this case $\text{Tr.c}(L/K)$ is necessarily the first uncountable cardinal \aleph_1 and if $d = 1$, then $I_{L/K}$ is nothing else but an Alexandroff's long ray.

2. THE MONOTONICITY THEOREM

In this section we present the fastest route (we know) to prove the monotonicity theorem for small extensions.

2.1. Countably generated extensions.

2.1.1. *Completed filtered unions.* We will use completed filtered unions of analytic fields, and the question whether the completion is non-trivial will be essential.

Lemma 2.1.2. *Assume K is an analytic field and $\{F_i\}_{i \in I}$ is a filtered family of analytic subfields, then:*

(i) *If I is of countable cofinality and not all F_i are trivially valued, then $F = \cup_{i \in I} F_i$ is not complete.*

(ii) *Conversely, F is complete in any of the following cases: the cofinality is one, the fields are trivially valued, or any countable subset in I is bounded (for example, the family is a chain of an uncountable cofinality).*

Proof. (i) Choose a cofinal increasing subsequence, which we denote for shortness $F_0 \subsetneq F_1 \subsetneq \dots$. Then $F = \cup_{j=0}^{\infty} F_j$ and each F_j is nowhere dense in \widehat{F} . Therefore $F \neq \widehat{F}$ by Baire category theorem.

(ii) The first two cases are trivial, so assume that any countable subset is bounded. Any Cauchy sequence a_0, a_1, \dots of elements of F lies already in a countable union of F_i 's, and hence is contained in a single field F_j . So, the limit already lies in some F_j . \square

Remark 2.1.3. If $F_0 \subsetneq F_1 \subsetneq \dots$ is a strictly increasing sequence of analytic k -fields such that $F_i = \widehat{k(x_i)^a}$ and F is the completed union, then $\text{tr.c}(F/k) = 1$ because any pair $\{x, y\} \subset F$ which is t.a. independent over k can be slightly deformed by [Tem21, Theorem 3.1.9] but no F_i contains a t.a. independent pair since $\text{tr.c}(F_i/k) = 1$. It follows that any $x \in F \setminus \cup_i F_i$ is a t.a. generator of F over k because for any i the pair x, x_i is t.a. dependent, $x \notin F_i$, and hence $x_i \in \widehat{k(x)^a}$. Already this simple observation was missed in [Tem21]. In particular, it implies that the answer to [Tem21, Question 5.3.7] is negative – for any k -endomorphism ϕ of $K = \widehat{k(t)^a}$ the completed union of $\phi^{-n}(K)$ is again generated by a single element.

The argument from the above remark extends to countable unions of analytic k -fields of bounded topological generating degree.

Theorem 2.1.4. *Let K/k be an extension of analytic fields, $\{F_i\}_{i \in I}$ a filtered family of intermediate fields of at most countable cofinality and F the completion of $\cup_i F_i$. If $d = \max_i(\text{Tr.c}(F_i/k))$ is finite, then $\text{Tr.c}(F/k) = d$.*

Proof. We can replace K and all other fields by $\widehat{K^a}$ and completed algebraic closures in $\widehat{K^a}$ as this does not change the invariant Tr.c . So in the sequel all fields are assumed to be algebraically closed. Note that $\text{tr.c}(F/k) = d$ and hence $\text{tr.c}(L/k) \leq d$ for any intermediate extension $k \subseteq L \subseteq F$. Indeed, otherwise there exist elements $x_0, \dots, x_d \in F$ t.a. independent over k , and by [Tem21, Theorem 3.1.9] a small enough perturbation x'_0, \dots, x'_d of x_0, \dots, x_d is also t.a. independent over k . In particular, we can achieve that x'_0, \dots, x'_d lie in $\cup_i F_i$ and hence also in some F_i , which contradicts that $\text{tr.c}(F_i/k) = \text{Tr.c}(F_i/k) \leq d$.

We will run induction on d with the base $d = 0$ being obvious. Since $\text{tr.c}(F_i/k) = \text{Tr.c}(F_i/k)$, we have that $\text{Tr.c}(F_i/K) \leq \text{Tr.c}(F_j/K)$ for $i \leq j$. So, we can replace the family by a cofinal sequence $F_0 \subseteq F_1 \subseteq \dots$ such that $\text{Tr.c}(F_i/k) = d$ for any i . If the sequence stabilizes, the claim is obvious, so assume that it does not and hence by Lemma 2.1.2 there exists $x \in F \setminus \cup_i F_i$. Set $k' = \widehat{k(x)^a}$ and $F'_i = \widehat{F_i(x)^a}$ and note that F is the completed union of F'_i . We claim that $\text{Tr.c}(F'_i/k') \leq d - 1$, and once we prove this the theorem will follow as follows: by the induction assumption $\text{Tr.c}(F/k') \leq d - 1$ and hence $\text{Tr.c}(F/k) \leq d - 1 + \text{Tr.c}(k'/k) \leq d$. Therefore $\text{tr.c}(F/k) = \text{Tr.c}(F/k)$ and the monotonicity of the invariant tr.c implies that in fact $\text{Tr.c}(F/k) = d$.

To prove the claim we choose a topological transcendence basis x_1, \dots, x_d of the small extension F_i/k . Then F'_i is t.a. generated over k by the set $S = \{x, x_1, \dots, x_d\}$ and since $\text{tr.c}(F'_i/k) \leq d$, there exists a t.a. dependency between the elements. By our construction, $x \notin F_i \supseteq \widehat{k(x_1, \dots, x_d)^a}$ and hence there exists x_m lying in $k(S \setminus \{x_m\})^a$. Therefore $F'_i = \widehat{k(S)^a} = \widehat{k(S \setminus \{x_m\})^a}$ and hence $S \setminus \{x, x_m\}$ t.a. generates F'_i over k' . In particular, $\text{Tr.c}(F'_i/k') \leq d - 1$. \square

As an immediate corollary we can now extend [Tem21, Theorem 3.2.3] to the countable cardinal. As we will later see, this is already optimal, as the result fails for the first uncountable cardinal.

Corollary 2.1.5. *If K/k is an extension of analytic fields such that $\text{Tr.c}(K/k) \leq \aleph_0$, then $\text{tr.c}(K/k) = \text{Tr.c}(K/k)$.*

Proof. Since $\text{tr.c}(K/k) \leq \text{Tr.c}(K/k)$, we should only rule out the possibility that simultaneously $\text{tr.c}(K/k) = d$ and $\text{Tr.c}(K/k) = \aleph_0$. In the latter case $\widehat{K^a}$ is the completion of $\widehat{k(x_0, x_1, \dots)^a}$ and hence K is the completed union of the sequence $K_n = \widehat{k(x_0, \dots, x_n)^a}$. Since $\text{Tr.c}(K_n/k)$ is finite, it coincides with $\text{tr.c}(K_n/k)$ and hence is bounded by d . Thus $\text{Tr.c}(K/k) \leq d$ by Theorem 2.1.4 yielding a contradiction. \square

2.2. Cofinality of subfields of small extensions. Now we will prove that algebraically closed subextensions of small extensions have countable cofinality.

2.2.1. One generator. We start with extensions with a single t.a. generator. The following self-similarity result is hidden in the proof of [Tem21, Theorem 5.2.2], so we formulate and prove it for the sake of completeness. As in [Tem21] given an extension L/K and an element $x \in L$ by $r_K(x) = \inf_{c \in K^a} |x - c|$ we denote the distance between x and K^a evaluated in L^a .

Lemma 2.2.2. *Assume that $L/K/k$ is a tower of algebraically closed analytic fields such that $\text{Tr.c}(K/k) = \text{Tr.c}(L/k) = 1$. Then the analytic k -fields K and L are isomorphic and the extension L/K is immediate.*

Proof. We can assume that the valuation on K is non-trivial as the other case is easily ruled out by classical tools. Choose generators $K = \widehat{k(x)^a}$ and $L = \widehat{k(y)^a}$. Since $k(y)$ is dense in L and $r_k(x) > 0$, we can find an element $y' \in \widehat{k(y)^a}$ such that $|x - y'| < r_k(x)$. Then $k(y)^a$ is algebraic over $k(y')$ and hence $L = \widehat{k(y')^a}$. In addition, $\widehat{k(y')^a} \xrightarrow{\sim} \widehat{k(x)^a}$ by [Tem21, Lemma 3.1.6], and hence the analytic k -fields K and L are also isomorphic. The latter also implies that $E_{L/K} = E_{L/k} - E_{K/k} = 0$

and $F_{L/K} = F_{L/k} - F_{K/k} = 0$, where we use the notation $F_{L/K} = \text{tr.deg.}(\widetilde{L}/\widetilde{K})$ and $E_{L/K}$ is the \mathbb{Q} -dimension of $|L^\times|^\mathbb{Q}/|K^\times|^\mathbb{Q}$. Since \widetilde{k} is algebraically closed and $|K^\times|$ is divisible this implies that $\widetilde{L} = \widetilde{K}$ and $|L^\times| = |K^\times|$, that is, the extension L/K is immediate. \square

Now, we can bound the ordered set $I_{K/k}$.

Lemma 2.2.3. *Let K/k be an extension of algebraically closed analytic fields such that $\text{Tr.c}(K/k) = 1$ and let $I_{K/k} = \{F_i\}$ denote the set of analytic algebraically closed intermediate fields $k \subseteq F_i \subseteq K$. Then any t.a. generator $t \in K$ induces an injective map $\rho_t: I_{K/k} \hookrightarrow [0, r_k(t)]$, which sends $F \in I_{K/k}$ to $r_F(t)$.*

Proof. We should prove that if $l \subsetneq L$ are two different elements of $I_{K/k}$, then the inequality $r_L(t) \leq r_l(t)$ is strict. We will argue by a contradiction, so assume that $r_L(t) = r_l(t)$. Since $K = \widehat{l(t)^a}$, we can replace k by l . Thus, we assume in the sequel that $k = l$ and $r_k(t) = r_L(t)$. In addition, decreasing L can only increase $r_L(t)$, so we can replace L by a subfield of the form $\widehat{k(y)^a}$ with $y \in L \setminus k$. Then we still have that $r_L(t) = r_k(t)$ and by Lemma 2.2.2 there exists a k -isomorphism of analytic fields $\phi: L = \widehat{k(t)^a} \xrightarrow{\sim} K$.

Set $x = \phi(t)$ and note that $|x - t| \leq r_k(x) = r_k(t)$. Since $r_k(t) = r_L(t) \leq |x - t|$ we obtain that in fact $|x - t| = r_L(t)$. This implies that the valuation on $L(x - t)$ is a generalized Gauss valuation. In particular, $L(t)/L$ is not immediate, hence K/L is not immediate and this contradicts the second assertion of Lemma 2.2.2. \square

Remark 2.2.4. Of course, each ρ_t is a bijection, and we will prove this later by an explicit construction taken from [Tem21]. But we even do not need the surjectivity to prove the monotonicity theorem, as already at this stage one can easily show using Theorem 2.1.4 that any intermediate field F_i is of the form $\widehat{k(x_i)^a}$.

2.2.5. *Finitely many generators.* Even more generally than in the above remark, we can now bound subextensions of arbitrary small extensions.

Theorem 2.2.6. *Let K/k be a small extension of algebraically closed analytic fields. Then any chain $\{F_i\}_{i \in I}$ of intermediate algebraically closed analytic fields is of at most countable cofinality.*

Proof. Assume to the contrary that the cofinality is uncountable. Decreasing I we can then assume that it is well-ordered and uncountable. We will run induction on $d = \text{Tr.c}(K/k)$, with the case of $d = 0$ being trivial. So, assume that $d \geq 1$ and choose a topological transcendence basis $x = x_1, \dots, x_d$ of K/k . Set $k' = \widehat{k(x)^a}$ and note that $\text{Tr.c}(K/k') = d - 1$. The fields $F'_i = \widehat{F_i(x)^a}$ form a chain of intermediate subfields of K/k' , hence this chain stabilizes at some i_0 such that there exist at most countably many elements $i < i_0$. Removing these elements and renaming i_0 by 0 we can assume that $F'_i = F'_0$ for any $i \in I$. Therefore, $\{F_i\}_{i \in I}$ is an uncountable well-ordered chain of intermediate fields for the extension $\widehat{F_0(x)^a}/F_0$ and by Corollary 2.2.3, there exists an order reversing embedding of I into the interval $[0, r_{F_0}(x)]$, which is an absurd. \square

2.3. Finite monotonicity of Tr.c.

2.3.1. *The monotonicity theorem.* As a corollary we can prove that the topological generating degree is monotonic in the finite case. In particular, this provides a positive answer to [Sch22, Question 21.4].

Theorem 2.3.2. *Let $K \subseteq E \subseteq F \subseteq L$ be a tower of analytic fields such that the extension L/K is small. Then $\text{Tr.c}(F/E) \leq \text{Tr.c}(L/K)$, in particular, F/E is small.*

Proof. We should prove that $\text{Tr.c}(F/K) \leq \text{Tr.c}(L/K)$, as $\text{Tr.c}(F/E) \leq \text{Tr.c}(F/K)$ in the obvious way. Replacing the fields with their completed algebraic closures does not affect the invariants Tr.c , so we can assume that the fields are algebraically closed. If $\text{Tr.c}(F/K) < \infty$, then the topological generating degrees coincide with the topological transcendence degree, and the assertion follows from monotonicity of tr.c . Thus, we should only rule out the possibility of $\text{Tr.c}(F/K) = \infty$.

By transfinite induction there exists a maximal strictly increasing chain F_i ordered by ordinals $i \leq \omega$, where ω is the maximal ordinal in the chain, such that $F_0 = K$, for any $i < \omega$ one has that $F_{i+1} = \widehat{F_i(x_i)^a}$, and for any limit ordinal $j \leq \omega$ the field F_j is the completion of $\cup_{i < j} F_i$. In particular, $F_\omega = F$ by the maximality assumption. Then the cardinal ω is countable by Theorem 2.2.6, hence F is topologically generated over k by countably many elements x_i , and therefore by Corollary 2.1.5 $\text{Tr.c}(F/K) = \text{tr.c}(F/K) \leq \text{tr.c}(L/K) < \infty$. \square

2.3.3. *The self-similarity theorem.* Also, one can now strengthen the self-similarity lemma by extending it to any finite topological degree.

Theorem 2.3.4. *Let L/K be a small extension of algebraically closed analytic fields and F an intermediate algebraically closed analytic field. Then L and F are isomorphic as analytic K -fields if and only if the inequality $\text{tr.c}(F/K) \leq \text{tr.c}(L/K)$ is an equality. In addition, in this case the extension L/F is immediate.*

Proof. Only the inverse implication is non-trivial, so assume that both topological transcendence degrees are equal to d . We claim that for an element $y \in L \setminus F$ the K -field $E = \widehat{F(y)^a}$ is isomorphic to F and the extension E/F is immediate. Since L is t.a. finitely generated over F (even over K), the theorem will follow by induction on the number of t.a. generators.

Now, let us prove the claim. By Theorem 2.3.2 F/K is small, hence possesses a topological transcendence basis x_1, \dots, x_d . The set $\{y, x_1, \dots, x_d\}$ is t.a. dependent over K , and up to renumbering the elements $x_1 \in K(y, \widehat{x_2, \dots, x_d})^a$. In particular, the latter field coincides with E , and setting $K' = K(x_2, \dots, x_d)^a$ we obtain that $F = \widehat{K'(x_1)^a} \subset E = \widehat{K'(y)^a}$. So, by Lemma 2.2.2 F and E are isomorphic even as analytic K' -fields and the extension E/F is immediate. \square

3. SUBFIELDS OF $\widehat{k(t)^a}$

Throughout this section K/k is an extension of algebraically closed analytic fields such that $p = \text{char}(\tilde{k}) > 0$, $|K^\times| \neq 1$ and $\text{Tr.c}(K/k) = 1$. Often we will also fix a t.a. generator t and then $K = \widehat{k(t)^a}$ and $r = r_k(t) > 0$. Our goal is to study the set $I_{K/k}$ of intermediate algebraically closed analytic fields.

3.1. Ordered structure of $I_{K/k}$. We already know that $\rho_t: I_{K/k} \hookrightarrow [0, r_k(t)]$ by Lemma 2.2.3, so our goal is prove that it is onto by constructing an intermediate field F of a prescribed radius $r_F(t)$. The trick which was already used in [Tem21] is to construct an extension L/K with $\text{Tr.c}(L/k) = 1$ instead, and then use an isomorphism $L \xrightarrow{\sim} K$ which sends K to a subfield F .

3.1.1. Endomorphisms. We start with recalling how endomorphisms of $\widehat{k(t)^a}$ are constructed. It is easy to see that if K/k is an extension of analytic fields and $x, t \in K$ satisfy $|x - t| \leq r_k(t)$, then $k(x) \xrightarrow{\sim} k(t)$ as valued fields (in particular, if $x \in k$, then $t = x$) and hence one obtains an isomorphism of analytic k -fields $\widehat{k(x)} \xrightarrow{\sim} \widehat{k(t)}$ sending x to t . This is proved in [Tem10, Lemma 6.3.2] when $|x - t| < r_k(t)$ but the same argument applies when the equality holds. If K is algebraically closed, then this extends further to an isomorphism of subfields $\widehat{k(x)^a} \xrightarrow{\sim} \widehat{k(t)^a}$ of K . In particular, we obtain the following result:

Lemma 3.1.2. *Assume that k is an algebraically closed analytic field and $K = \widehat{k(t)^a}$. Then for any k -endomorphism $\psi: K \rightarrow K$ the element $x = \psi(t)$ satisfies $|x - t| \leq r_k(t) = r_k(x)$. Conversely, for any x with $|x - t| \leq r_k(t)$ there exists a K -endomorphism ψ taking t to x , and such ψ is unique up to an action of $\text{Gal}(\widehat{k(t)})$.*

3.1.3. Recollections on completed differentials. Next, we recall the following relation between differentials and t.a. dependence.

Lemma 3.1.4. *Let K/k be an extension of analytic field and $t \in K$ such that $K = \widehat{k(t)}$. Then the following conditions are equivalent:*

- (i) $k^s \cap K$ is dense in K (in particular, $t \in \widehat{k^a}$),
- (ii) $\widehat{d}_{K/k}(t) = 0$,
- (iii) $d_{K^\circ/k^\circ}(t)$ is infinitely divisible.

Proof. The equivalence of (i) and (ii) is covered by [KT18, Lemma 2.4], and the equivalence of (ii) and (iii) follows from [Tem21, Lemma 4.1.11]. \square

3.1.5. Deeply ramified points of geometric type 4. As in [Tem21], fancy extensions are constructed in this paper using deeply ramified extensions which force some differentials to vanish and hence lead to a t.a. dependence. More specifically, let k be a non-trivially valued analytic field. It is proved in [Tem21, Theorem 5.1.12] that if k is not perfectoid, then there exist points z of type 4 with deeply ramified extension $\mathcal{H}(z)/k$. Moreover, we are going to show that the proof in loc.cit. also yields the following refinement, which provides a control on the geometric radius of z in a k -split disc.

Lemma 3.1.6. *Let k be an analytic field such that $p = \text{char}(\widetilde{k}) > 0$ and the group $|k^\times|$ is dense. Assume that $t \in k^\circ$ is an element such that $d_{k^\circ}(t)$ is not infinitely divisible in Ω_{k° and $r \in (0, 1)$. Then there exists a primitive extension $K = \widehat{k(x)}$ of type 4 and an algebraic subextension l/k of K/k such that $d_{l^\circ}(t)$ is infinitely divisible, $|x| = 1$ and $r_{\widehat{k^a}}(x) = r$.*

Proof. Fix $\pi \in k$ such that $|p| < |\pi| < 1$, and if the characteristic is mixed, also replace t by $p^{-n}t$ achieving that $|p| < |t| < 1$. Choose a strictly increasing sequence of natural numbers $0 < d_0 < d_1 < \dots$ such that $\prod_{n=0}^{\infty} |\pi|^{1/(p^{d_n}-1)} > r$. Now, we will construct an extension l/k precisely as in the proof of [Tem21, 5.1.12]: set

$t_0 = t$, take t_{n+1} to be a root of $f_n(x) = x^{p^{d_n}} - \pi x - t_n$ and define $k_0 = k$, $k_{n+1} = k_n(t_{n+1})$ and $l = \cup_n k_n$. A simple computation in the proof [Tem21, 5.1.12] shows that $d_{l^\circ} t$, is infinitely divisible in Ω_{l° and $r_{\text{spl}}(t_{n+1}/k_n) = |\pi|^{1/(p^{d_n}-1)}$.

Since $\prod_{n=0}^{\infty} r_{\text{spl}}(t_{n+1}/k_n) > r$ and $|k^\times|$ is dense, there exists a sequence $r_0 = 1, r_1, r_2, \dots$ of elements of $|k^\times|$ which converges to r and satisfies $r_{n+1}/r_n < r_{\text{spl}}(t_{n+1}/k_n)$ for any $n \geq 0$. It then follows from [Tem21, Lemma 5.1.7] that starting with the unit disc $E_0 = \mathcal{M}(k\{x\})$ one can construct a sequence of discs $E_0 \supset E_1 \supset E_2 \supset \dots$ such that E_n is k_n -split and is of geometric radius r_n in E_0 . The intersection of these discs does not contain k^a -points and hence $\bigcap_n E_n = \{z\}$ is a single point of type 4 of geometric radius r . Thus, $K = \mathcal{H}(z) = \widehat{k(x)}$ contains l and $r_{\widehat{k^a}}(x) = \inf_{c \in k^a} |c - x|_z = r$, as required. \square

3.1.7. Completed tame extensions. In order to deal with discretely valued fields we will have to pass to an appropriate completed tame extension. This makes the group of values dense, and the following result shows that the divisibility properties of the differentials are not changed much.

Lemma 3.1.8. *Assume that k is an analytic field, $\omega \in \Omega_{k^\circ}$ is an element, l/k a tame algebraic extension and $K = \widehat{l}$. Then ω is infinitely divisible if and only if the image of $\omega \otimes 1$ in Ω_{K° is infinitely divisible.*

Proof. Since K/k is separable, $H_1(\mathbb{L}_{K^\circ/k^\circ}) = 0$ by [Tem16, Theorem 5.2.3(ii)] and hence the sequence

$$0 \rightarrow \Omega_{k^\circ} \otimes_{k^\circ} K^\circ \rightarrow \Omega_{K^\circ} \rightarrow \Omega_{K^\circ/k^\circ} \rightarrow 0$$

is exact. In the same vein we have the exact sequence

$$0 \rightarrow \Omega_{l^\circ/k^\circ} \otimes_{k^\circ} l^\circ \rightarrow \Omega_{K^\circ/k^\circ} \rightarrow \Omega_{K^\circ/l^\circ} \rightarrow 0,$$

whose first term is a torsion module annihilated by $k^{\circ\circ}$, and whose third term is torsion free by [Tem16, Lemma 5.2.9]. Thus, from the second sequence we obtain that Ω_{K°/k° contains no non-zero infinitely divisible torsion elements, and this combined with the first sequence implies the lemma. \square

3.1.9. Construction of towers with tr.c defect. Now we can construct towers of simple extensions in which the topological transcendence degree is non-additive. This refines [Tem21, Theorems 5.2.2] by providing control on radii of generators, but we use essentially the same argument.

Lemma 3.1.10. *Assume that F/k is a non-trivial extension of algebraically closed analytic fields such that $F = \widehat{k(t)^a}$ is non-trivially valued and $\text{char}(\widehat{k}) > 0$. Then for any positive $r < r_k(t)$ there exists an extension $K = \widehat{F(x)^a}$ of F such that $|x - t| < r_k(t) = r_k(x)$, $r = r_F(x)$ and $K = \widehat{k(x)^a}$.*

Proof. Set $L = \widehat{k(t)}$ and recall that $d_{L^\circ/k^\circ}(t)$, and hence also $d_{L^\circ}(t)$, is not infinitely divisible by Lemma 3.1.4. Since L can be discretely valued (when k is trivially valued) we also consider the completed tame closure $L' = \widehat{L^t}$, which has dense group of absolute values, and note that $d_{L'^\circ}(t)$ is not infinitely divisible by Lemma 3.1.8. Choose $\pi \in F$ such that $r < |\pi| < r_k(t)$. By Lemma 3.1.6 there exists an extension of analytic fields $E' = \widehat{L'(y)}$ such that $d_{E'^\circ}(t)$ is infinitely divisible, $|y| = 1$ and $r_F(y) = r/|\pi|$. Since E' is a completed tame extension of $E = \widehat{k(y, t)}$, we obtain

by Lemma 3.1.8 that already $d_{E^\circ}(t)$ is infinitely divisible. Set $H = \widehat{k(y)}$. Then $d_{E^\circ/H^\circ}(t)$ is infinitely divisible, and hence $t \in H$ by Lemma 3.1.4.

We claim that $K = \widehat{k(y)^a}$ and $x = t + \pi y$ are as required. First, $|x - t| = |\pi| < r_k(t)$, and hence $r_k(x) = r_k(t)$. Second, $r_F(x) = r_F(\pi y) = |\pi| r_F(y) = r$. Finally, $\widehat{k(t, x)^a} = \widehat{k(t, y)^a} = K$. Since $\text{tr.c}(K/k) = 1$ and $K \neq \widehat{k(t)^a}$, this implies that $t \in \widehat{k(x)^a}$ and hence $K = \widehat{k(x)^a}$. \square

Corollary 3.1.11. *Assume that K/k is a non-trivial extension of algebraically closed analytic fields such that $K = \widehat{k(t)^a}$ and $\text{char}(\tilde{k}) > 0$. Then for any positive $r < r_k(t)$ there exists a k -subfield $F = \widehat{k(x)^a} \subsetneq K$ such that $r_F(t) = r$.*

Proof. By Lemma 3.1.10 there exists an extension $L = \widehat{K(y)^a}$ such that $r_K(y) = r$, $|y - t| < r_k(t)$ and $L = \widehat{k(y)^a}$. By Lemma 3.1.2 there exists a k -isomorphism $\psi: L \xrightarrow{\sim} K$ taking y to t . Then for $F = \psi(K)$ and $x = \psi(t)$ we have that $F = \widehat{k(x)^a} \subsetneq K$ and $r_F(t) = r_K(x) = r$. \square

3.1.12. *Applications.* Summarizing our results so far we obtain a surprisingly simple answer to [Tem21, Question 5.3.10], namely the map ρ_t from Lemma 2.2.3 is a bijection.

Theorem 3.1.13. *Let K/k be an extension of algebraically closed analytic fields such that $\text{Tr.c}(K/k) = 1$, $\text{char}(\tilde{k}) > 0$ and $|K^\times| \neq 1$. Then any $t \in K$ with $\widehat{k(t)^a} = K$ induces a bijection $\rho_t: I_{K/k} \xrightarrow{\sim} [0, r_k(t)]$ sending an intermediate algebraically closed analytic field F to $r_F(t)$. The bijection ρ_t is monotonic and respects limits: $F \subseteq F'$ if and only if $r_F(t) \geq r_{F'}(t)$, and for any family $\{F_i\} \subset I_{K/k}$ with $F' = \bigcap_i F_i$ and F'' the completion of $\bigcup_i F_i$ one has that $r_{F'}(t) = \sup_i r_{F_i}(t)$ and $r_{F''}(t) = \inf_i r_{F_i}(t)$.*

Proof. The map ρ_t is injective by Lemma 2.2.3 and surjective by Corollary 3.1.11. Clearly, ρ_t is monotonic and reverses the order, and hence it takes infima to suprema and vice versa. It remains to note that infima and suprema in $I_{K/k}$ are intersections and completed unions. \square

3.2. **The PL structure of $I_{K/k}$.** Throughout §3.2 we assume that k is algebraically closed, $p = \text{char}(\tilde{k}) > 0$, $k \subsetneq K = \widehat{k(t)^a}$, $r = r_k(t) > 0$, and we will study the natural structure the interval $I_{K/k}$ acquires.

3.2.1. *Log radius parameterizations.* Instead of the parametrization $\rho_t: I_{K/k} \xrightarrow{\sim} [0, r]$ by the radius function, it will be convenient to use its logarithm, which will be denoted by $\rho_t^{\log}: I_{K/k} \xrightarrow{\sim} [-\log(r), \infty]$, where $\rho_t^{\log}(F) = -\log(r_F(t))$. In particular, this parametrization is order preserving. Such functions also make sense for t which is not a t.a. generator: if $F = \widehat{k(t)^a}$, then $\rho_t^{\log}: [k, F] \xrightarrow{\sim} [-\log(r), \infty]$ and $\rho_t^{\log}([F, K]) = \infty$.

3.2.2. *Transition functions.* One can wonder to which extent the metric structure ρ_t^{\log} induces on $I_{K/k}$ is canonical. Since the parametrization is given by the log radius function ρ_t^{\log} of the t.a. generator t , an equivalent question is to describe the *transition functions*, that is, functions $[-\log r, \infty] \xrightarrow{\sim} [-\log r', \infty]$ of the form $\rho_{t,t'}^{\log} = \rho_{t'}^{\log} \circ (\rho_t^{\log})^{-1}$, where t' is another t.a. generator and $r' = -\log r_k(t')$.

A tightly related question is as follows: what is the image of the restriction

$$\Lambda_t: \text{Aut}_k(K) \rightarrow \text{Aut}(I_{K/k}) \xrightarrow{\sim} \text{Aut}([-\log r, \infty])?$$

Of course, this restriction takes an automorphism $\psi \in \text{Aut}_k(K)$ to $\rho_{t, \psi(t)}^{\log}$.

3.2.3. Immediate simple extensions. The main tool of our study of the radii and transition functions will be the following results about simple immediate extensions $L = \widehat{k(x)}$. They are not well recorded in the literature (though maybe known to experts), so we provide all needed details. Assume that k is algebraically closed and L/k is immediate. In particular, $r = r_k(x) = \inf_{a \in k} |x - a|$ is not attained. For shortness, set $x_a = x - a$ and $r_a = |x_a|$ for $a \in k$. For any element $f \in k[x]$ of degree d let $f = \sum_{i=0}^d c_{a,i} x_a^i$ be its presentation with respect to the coordinate x_a .

Lemma 3.2.4. *Keep the above notation, then*

(i) $L = \widehat{k[x]}$ and, moreover, L° is the completion of $\cup_a k^\circ[x_a/\pi_a]$, where $\pi_a \in k$ is such that $|\pi_a| = r_a$.

(ii) There exists $r_0 > r$ such that for any $a \in k$ with $r_a < r_0$ the numbers $s_i = |c_{a,i}|$ are independent of a and satisfy the inequalities $s_j > \binom{i}{j} s_i r_a^{i-j}$ for any $0 \leq j < i \leq d$ such that $s_j > 0$.

(iii) One has that $r_k(f) = \max_{i>0} |s_i| r^i$ and $r_k(f) > |s_i| r^i$ whenever $i \notin \{0\} \cup p^{\mathbb{N}}$. In particular, if r_a is sufficiently close to r and $i_1 = p^N$ and $i_2 = p^{N'}$ are the maximal and the minimal values of i such that $r_k(f) = s_i |r^i|$, then

$$f - c_{a,0} = g + \sum_{n=N'}^N c_{a,p^n} x_a^{p^n},$$

where $|g| < r_k(f)$.

(iv) If $F = \widehat{k(f)}$, then $[L : F] = p^N$.

Proof. (i) The valuation of K corresponds to a point $z \in \mathbb{A}_k^1$ on the Berkovich affine line and by Berkovich's classification of points on the affine line, $\{z\} = \cap_a D(a, r_a)$ is the intersection of discs given by $|x - a| \leq r_a$. In particular, the valuation $|\cdot|_z$ on $k[x]$ is the infimum of the translated Gauss valuations $|\cdot|_{a,r_a}$ around a of radius r_a , and the claim of (i) follows easily.

(ii) Let ∂^j denote the j -th divided power derivative given by $\partial^j(f) = \sum_{i=j}^d \binom{i}{j} c_j x^{i-j}$ and choose r_0 so close to r that the disc $D_0 = D(a, r_0)$ of radius r_0 around z does not contain zeros of any non-zero $\partial^j(f)$. Then each non-zero $\partial^j(f)$ is invertible in any disc contained in D_0 and hence its dominant term is the free one, that is $|c_{a,j}| > \binom{i}{j} |c_{a,i}| r_a^{i-j}$ for any $i > j$ (we use that $\partial^j(f) = \sum_{i=i}^d \binom{i}{j} c_{a,i} x_a^{i-j}$). Moreover, for any $b \in k$ we have that $c_{b,j} = \sum_{i=j}^d \binom{i}{j} c_{a,i} (b-a)^{i-j}$, hence whenever $|b-a| < r_0$ we have that $|c_{a,j}| = |c_{b,j}|$ is independent of a .

(iii) It follows from (i) that $r_k(f)$ is the infimum of the distance between f and k with respect to the Gauss valuations $|\cdot|_{a,r}$, that is,

$$r_k(f) = \inf_{a \in k} \max_{i>0} |c_{a,i}| r_a^i.$$

Since $r = \inf_a r_a$ and the values of $|c_{a,i}|$ stabilise when r_a is close enough to r , we obtain that $r_k(f) = \max_{i>0} |s_i| r^i$. If $i > 0$ is not a p -th power, then $i = p^n m$ with $m > 1$ and $(m, p) = 1$. It follows that $j = p^n$ satisfies $p \nmid \binom{i}{j}$ and hence $s_i r^{i-j} < s_j$ by (ii). In particular, $s_i r^i < s_j r^i \leq r_f(k)$. The last claim of (iii) follows.

(iv) Consider the morphism $h: \mathbb{A}_k^1 \rightarrow \mathbb{A}_k^1$ induced by f and let $y = h(z)$. Then $\mathcal{H}(z) = L$ and $\mathcal{H}(y) = F$, so the claim reduces to showing that the local degree $\deg_h(z)$ of h at z is p^N . For $r' > r$ let $D_{r'}$ be the disc around z of radius r' and let $z_{r'}$ be its maximal point. By definition of N in (iii), decreasing r_0 we can achieve that $s_i r'^i < s_{p^N} r'^{p^N}$ for any $r < r' < r_0$ and $0 < i \neq p^N$. For any such r' , after translating the coordinates by a and $c_{a,0}$ the morphism $D_{r'} \rightarrow h(D_{r'})$ becomes the morphism between discs with center at 0 given by a polynomial of order p^N (recall that the order is the largest degree of a dominant term). By Weierstrass division theorem this morphism is of degree p^N , hence $\deg_h(z) \leq p^N$. In addition, this morphism is precisely of degree p^N at the maximal point of the disc. So, $\deg_h(z_{r'}) = p^N$ and by semicontinuity of the degree we obtain the equality $\deg_h(z) = p^N$. \square

It seems that a basic theory of almost tame extension is missing in the literature. Of course, this gap should be filled out, but perhaps in another venue. So, we will just state what we need without proof and will use in side remarks in the sequel.

Remark 3.2.5. (i) A finite extension of valued fields l/k is *almost tame* if $\Omega_{l^\circ/k^\circ}^{\log} = 0$. Otherwise we say that l/k is *wildest*. Also, we say that l/k is *purely wildest* if for any $k \subseteq k_1 \subsetneq l_1 \subseteq l$ the extension l_1/k_1 is wildest.

(ii) It is easy to see that the class of almost tame extensions is closed under compositions and taking subextensions. It includes the class of tame extensions and “mildest” wild extension and is, in fact, the first step of the ramification filtration beyond the tame class. A defectless extension l/k is almost tame if and only if it is tame. A purely wild almost tame extension is immediate. Finally, l/k contains a maximal almost tame subextension l_{at} . A bit subtler fact is that l/l_{at} is purely wildest. We call $[l : l_{at}]$ the *wildest degree* of l/k . This is the analogue of the *wild degree* of l/k , which is the degree of l over the maximal tame subfield l_t . Naturally, the wild and the wildest degrees are multiplicative in towers.

(iii) The meaningful interpretation of N' in Lemma 3.2.4 is that $p^{N'}$ is the wildest degree of $L/\widehat{k(f)}$, while p^N is its usual degree, which is also the wild degree.

(iv) If $L = \widehat{k(t)}$ is not immediate over k , then L is stable by [Tem10, Theorem 6.3.1(iii)] and hence any finite extension F/L is defectless and its wild and wildest degrees coincide. In addition, the wild degree can be read off from the simple invariants – it is the p -primary part of $e_{F/L}$ if $|L^\times| \neq |k^\times|$, and it is the inseparable degree of $\widetilde{F}/\widetilde{L}$ otherwise.

3.2.6. Explicit examples in positive characteristic. Now we can construct a large family of explicit examples, which completely illustrate the general case. Technically, it is easier to deal with the case of positive characteristic, so assume in the next example that $\text{char}(k) = p$.

Example 3.2.7. (i) For any $t' = \sum_{i=-\infty}^N c_i t^{p^i}$ with $c_i \in k$, we have that $r_F(t') = \max_i (|c_i| r_F(t)^{p^i})$ for any $F \in I_{K/k}$. Indeed, to check this we can remove all terms of absolute value smaller than $r_F(t')$, and we can replace t' by t'^{p^m} because $r_F(t'^{p^m}) = (r_F(t'))^{p^m}$ (just apply Fr^m). This reduces us to the case when $t' = \sum_{i=0}^N c_i t^{p^i}$, and then we can compute $r_F(t')$ by Lemma 3.2.4(iii) because $c_i \in F$ and the coefficients c_i do not change under any coordinate change $t'_a = t' - a$. Passing to the log scale

we obtain that

$$\rho_{t,t'}^{\log}(s) = \min_i(p^i s - \log |c_i|),$$

so the functions realized in this way as $\rho_{t,t'}^{\log}$ are convex PL functions on $[-\log r, \infty)$ with slopes in $p^{\mathbb{Z}}$ and corners in $\log |k^\times|$ such that the sequence $-p^i \log |t| - \log |c_i|$ converges to infinity.

(ii) The same example with t being the dominant term of $t' = t + \sum_{i=-\infty}^{-1} a_i t^p$ can be used to construct explicit elements in the image of Λ_t . Also, any PL function can be realized as a composition of a convex one and an inverse of a convex one, leading to a much richer family of examples of the form $(\rho_{t',t'}^{\log})^{-1} \circ \rho_{t,t'}^{\log} = \rho_{t,t''}^{\log}$.

3.2.8. Explicit examples in mixed characteristic. In mixed characteristic computations are more difficult, so we consider only the simplest case we will use later.

Example 3.2.9. (i) Let us study $\rho_{t,t'}^{\log}$, where $t' = t^p + ct$ with $c \in k$. Using the presentations $t' - ca - a^p = ct_a + \sum_{i=1}^p \binom{p}{i} a^{p-i} t_a^i$ with $t_a = t - a$ and $a \in k$ such that $|a| = |t|$, one obtains that unless there is a cancellation $|c + pa^{p-1}| < |c|$, which does not happen when $|c| \neq |pa^{p-1}|$ and hence can be easily avoided in the sequel by translating t , the dominant term on the left is either t_a^p or the linear term $(c + pa^{p-1})t_a$. So, in this case $\rho_{t,t^p+ct}^{\log}(s) = \min(ps, s + \gamma)$, where $\gamma = \min(-\log |c|, -\log |pt^{p-1}|)$. This function is linear if $s_1 = \gamma/(p-1)$ is smaller than the endpoint $s_0 = -\log r$ and otherwise has a single corner at s_1 and slopes p and 1 . The latter happens whenever $|c| < r^{p-1}$ and $|t|$ is sufficiently close to r (see below). A technical complication in the mixed characteristic is that the corner cannot be too far from the endpoint:

$$s_1 - s_0 \leq -\frac{1}{p-1} \log |p| - \log |t| + \log r \leq -\frac{1}{p-1} \log |p|$$

and the equality is obtained when $|c| < |pt^{p-1}|$, including the case of $c = 0$ and $t' = t^p$. It follows easily that any PL function ρ with a single corner $s_1 \in |k^\times|$ such that $s_1 - s_0 < -\frac{1}{p-1} \log |p|$, $\rho(s) = ps$ for $s \in (s_0, s_1)$ and $\rho(s)$ is linear for $s > s_1$ can be realized as $\rho = \rho_{t_a, t_a + ct_a^p}^{\log}$ for appropriate $c \in k$ and $t_a = t - a$ with $|t_a|$ close enough to r .

(ii) Let now ρ be any PL function on $[s_0, \infty)$ with two corners $s_1, s_2 \in |k^\times|$ such that $s_1 < s_2 < s_0 - \frac{1}{p-1} \log |p|$, and $\rho(s) = s$ for $s < s_1$, ρ has slope p^{-1} for $s \in (s_1, s_2)$ and is linear for $s > s_2$. It follows from (i) that choosing c and a appropriately we can find a realization

$$\rho = \rho_{t_a^{1/p}, t_a + ct_a^{1/p}}^{\log} \circ (\rho_{t_a^{1/p}, t_a}^{\log})^{-1} = \rho_{t_a, t_a + ct_a^{1/p}}^{\log}.$$

Furthermore, when $|t_a|$ is close enough to r the inequality $|c| < r^{\frac{p-1}{p}}$ strengthens to $|ct_a^{1/p}| < r$ and hence there exists an automorphism $\psi \in \text{Aut}_k(K)$ taking t_a to $t_a + ct_a^{1/p}$. By the construction $\rho = \Lambda_t(\psi)$.

3.2.10. PL functions. Let us formalize the properties of transition functions observed in the above examples. Assume that Γ is a divisible subgroup of \mathbb{R} . By a $(p^{\mathbb{Z}}, \Gamma)$ -PL function on an interval $I \subseteq \mathbb{R}$ we mean a continuous function $f: I \rightarrow \mathbb{R}$ which is locally of the form $f(t) = p^n t + \gamma$ with $n \in \mathbb{Z}$ and $\gamma \in \Gamma$ outside of a discrete set of *corner points*. In particular, f is strictly increasing and each corner point t_0 lies in Γ because it satisfies $p^n t_0 + \gamma = p^{n_1} t_0 + \gamma_1$ with $n \neq n_1$. By a $(p^{\mathbb{Z}}, \Gamma)$ -PL

function on an interval $[t_0, \infty]$ we mean an unbounded $(p^{\mathbb{Z}}, \Gamma)$ -PL function f on the ray $[t_0, \infty)$ augmented by the rule $f(\infty) = \infty$. In particular, the set of corner points is discrete in $[t_0, \infty)$ but can accumulate at infinity. Note that the set of $(p^{\mathbb{Z}}, \Gamma)$ -PL functions is closed under compositions and taking inverses.

3.2.11. *The piecewise linearity theorem.* Now, let us establish the main result of §3.2 that any transition function is, indeed, of this form.

Theorem 3.2.12. *Let $K = \widehat{k(t)^a}/k$ be an extension of algebraically closed analytic fields with $r = r_k(t) > 0$, and assume that $|K^\times| \neq 1$ and $\text{char}(\tilde{k}) = p > 0$. Then:*

- (i) *For any other t.a. generator t' the function $\rho = \rho_{t,t'}^{\log}$ is $(p^{\mathbb{Z}}, |K^\times|)$ -PL.*
- (ii) *If $|K^\times| \neq |k^\times|$, then $|K^\times| = |k^\times| \oplus r^{\mathbb{Q}}$ and $\rho(-\log r) = -p^n b \log r - \log |a|$, where $a \in k^\times$, $b \in \mathbb{Z} \setminus p\mathbb{Z}$ and p^n is the slope of ρ at the endpoint $-\log r$.*

Proof. We will use the observation that the set of PL functions satisfying conditions of the theorem is closed under compositions and taking the inverse function. Also, the claim of the theorem is local at a field $F \in I_{K/k} \setminus \{K\}$. So set $r_F := r_F(t)$ and $s = -\log r_F$, and let us study ρ locally at s . It will be convenient to work with a neighborhood $I' = [F_1, F_2]$ of F in $I_{K/k}$ and shrink it when needed. In particular, we can assume that there exists $\varepsilon > 0$ such that $\rho_{t'} \geq \varepsilon$ on I' , and hence t' can be replaced by $t' - c$ with $c \in K$ and $|c| < \varepsilon$ without affecting $\rho_{t'}$ and hence also ρ on I' . In particular, we can assume that $t' \in k(t)^a$, and then $L = \widehat{k(t, t')}$ is finite over $\widehat{k(t)}$. Since L is topologically finitely generated over \widehat{k} , by [Tem10, Theorem 6.3.1] there exists $t'' \in L$ such that L is unramified over $\widehat{k(t'')}$. Since ρ is the composition of $\rho' = \rho_{t'', t'}^{\log}$ and $\rho'' = \rho_{t, t''}^{\log}$, it suffices to prove the claim for ρ' and $(\rho'')^{-1}$. Replacing t by t'' for shortness of notation we can assume in the sequel that L is unramified over $\widehat{k(t)}$ itself. If $\tilde{F} = \tilde{K}$ this simply means that $L = \widehat{k(t)}$. Finally, we will use that we can freely replace t by $t_a = t - a$ with $a \in F$ without affecting ρ .

In the sequel we will have to separately consider three cases due to the type of the extension L/k . Recall that by Lemma 2.2.2 if K/F is not immediate, then $F = k$ is the endpoint of $I_{K/k}$.

Type 4. Assume first that K/F is immediate. If $F \neq k$, then shrinking I' we can assume that $k \notin I'$. In either case K/F_1 is immediate too, so $F_1[t]$ is dense in $L = \widehat{F_1(t)}$ by Lemma 3.2.4(i). Thus, moving t' slightly we can even assume that $t' = \sum_i c_i t^i \in F_1[t]$. By claim (iii) of the same lemma, there exists $r' > r_F$ such that for any $a \in F$ with $|t_a| < r'$ we have that $t' = c_{a,0} + g + \sum_{n=N}^{N'} c_{a,p^n} t_a^{p^n}$ with $|g| < r_F(t')$. Since ρ_t and $\rho_{t'}$ are continuous, shrinking I' we can assume that $|g| < \rho_{t'}$ and $\rho_t < r'$ on I' . In particular, there exists $a \in F_1 \subseteq F$ such that $|t_a| < r'$ and for this a we have that $c_{a,0} \in F_1$. Therefore this presentation of t' and Lemma 3.2.4(iii) can be used to compute $r_E(t')$ for any $E \in I'$, and we obtain that $\rho_{t'}(s) = \max_n |c_{a,p^n}| \rho_t^{p^n}(s)$ for any $s \in I'$. Thus, $\rho(s) = \min_n (p^n s - \log |c_{a,p^n}|)$ on I' , as required.

It remains to consider the case when K/k is not immediate and automatically $F = k$. In this case, the k -norm on finite-dimensional k -subspaces of K is Cartesian, hence translating t and t' by elements of k we can achieve that $|t| = r = r_k(t)$ and $|t'| = r' = r_k(t')$. Now we again have to split to cases.

Type 2. Assume that $|k^\times| = |K^\times|$. Then we can also rescale t and t' by elements of $|k^\times|$ so that $r = |t| = |t'| = 1$, while $\rho_{t,t'}^{\log}$ shifts by a value from $\log |k^\times|$. Thus

we assume in the sequel that \tilde{t}, \tilde{t}' are well defined and are not contained in \tilde{k} . Let us start with two particular cases: (1) if $\tilde{t}' = \tilde{t}^{p^n}$, then $r' = |t' - t^{p^n}| < 1$, hence $\rho_{t, t'}^{\log} = \rho_{t, t^{p^n}}^{\log}$ on $[0, -\log(r')]$, and the same computation as earlier shows that $\rho_{t, t^{p^n}}^{\log}$ has slope p^n at 0. (2) If \tilde{L} is separable over $\tilde{k}(\tilde{t}')$, then $L = \widehat{k(t)}$ is unramified over $\widehat{k(t')}$ because this extension is defectless by the stability theorem (e.g. see [Tem10, Theorem 6.3.1(iii)]). Therefore $\widehat{F(t)}/\widehat{F(t')}$ is unramified for any $F \in I_{K/k}$, and for $F \neq k$ this implies that $\widehat{F(t)} = \widehat{F(t')}$. As earlier, moving t' a bit we can assume that $t' \in F[t]$, and then Lemma 3.2.4(iv) implies that $t' = \sum_i a_i t^i$ with $|a_1| r_F(t) > |a_i| r_F(t)^i$ for $i > 1$. Thus, by the already established case of extensions of type 4, $\rho_{t, t'}^{\log}$ has slope 1 on $(-\log r, \infty)$ and hence also at the endpoint $-\log r$.

Now, assume that t' is arbitrary. Since \tilde{L} is a one-dimensional function field over \tilde{k} , the p -rank of \tilde{L} equals one and hence \tilde{L} is separable over a subfield of the form $\tilde{k}(\tilde{t}^{1/p^n})$. Take any $t'' \in L$ with $\tilde{t}'' = t^{1/p^n}$, then $\rho_{t'', t'}^{\log}$ and $\rho_{t, t''}^{\log}$ are $(p^{\mathbb{Z}}, |k^\times|)$ -PL at $s = 0$ by the cases (1) and (2) above, and hence the same is true for $\rho_{t, t'}^{\log}$.

Type 3. Finally, assume that $|k^\times| \neq |K^\times|$. Then $k\{t, t^{-1}\}_{r, r^{-1}}$ is already a field and hence coincides with L . Thus, $|L^\times| = |k^\times| \oplus r^{\mathbb{Z}}$ and moving t' a bit we can assume that $t' = \sum_i c_i t^i \in k[t^{\pm 1}]$. Then $|t'| = |c_m t^m|$ for a single m , and $m \neq 0$ because $|t'| = r_k(t')$. Thus $|t' - c_m t^m| < r'$ and we can replace t' by $c_m t^m$ without affecting $\rho_{t'}$ on a small enough neighborhood I' of k in $I_{K/k}$. Furthermore, $\rho_{t'} = |c_m| \rho_{t^m}$, so it suffices to study ρ_{t^m} on $I_{K/k}$. Write $m = bp^n$ with $(p, b) = 1$. For any $E \in I_{K/k}$ the value of $r_E(t^m)$ is computed using the presentations $t^m = \sum_{i=0}^{\infty} \binom{m}{i} a^{m-i} t_a^i$ with $a \in E$, $|a| = |t|$ and $r_a = |t - a|$ tending to $r_E(t)$. Note that this formula covers the case of $m < 0$ as well, and the minimal $i > 0$ with $p \nmid \binom{m}{i}$ is $i = p^n$. Therefore, the p^n -th term is the dominant non-free one when $r_E(t)$ is close enough to $|t| = r_k(t)$, and in a small neighborhood of the endpoint of $I_{K/k}$ we have that $\rho_{t^m}(E) = |t|^{(b-1)p^n} \rho_t(E)^{p^n}$. That is, $\rho_{t, t^m}^{\log}(s) = p^n s - (b-1)p^n \log(r)$ as (i) asserts and (ii) follows by substituting $s = -\log(r)$. \square

Remark 3.2.13. Let t, t' be as in the above theorem and assume that $t' \in k(t)^a$. We define the wild (resp. the wildest) degree of t' over t with respect to $F \in I_{K/k}$ as the ratio $n_{F, t/t'}^w$ (resp. $n_{F, t/t'}^W$) of the wild (resp. wildest) degrees of $\widehat{F(t, t')}/\widehat{F(t')}$ and $\widehat{F(t, t')}/\widehat{F(t)}$. Using Remark 3.2.5 it is easy to see that the arguments above in fact establish the following more precise result: the slope of $\rho_{t, t'}^{\log}$ to the right (resp. left) of $s = r_t^{\log}(F)$ equals $n_{F, t/t'}^w$ (resp. $n_{F, t/t'}^W$). In particular, even though at the endpoint $F = k$ the degrees $[L : \widehat{k(t)}]$ and $[L : \widehat{k(t')}]$ do not have to be a power of p , the slope only takes the wild degrees into account and hence lies in $p^{\mathbb{Z}}$.

In a sense, one can view $\widehat{F(t)}$ as a “lattice” in $\widehat{F(t)}^a$ and the transition functions measure the index between the lattices of t and t' . If $\widehat{F(t')} \subseteq \widehat{F(t)}$, then the function is convex, and a corner point occurs at F when this extension is not purely wildest. In general, we can only locally compare $\widehat{F(t')}$ and $\widehat{F(t)}$ by embedding them into a larger “lattice”, so the transition function is the ratio of two convex $(p^{\mathbb{Z}}, |K^\times|)$ -PL functions (that is, just a $(p^{\mathbb{Z}}, |K^\times|)$ -PL function).

3.2.14. *The PL structure.* Theorem 3.2.12 implies that the ray $I_{K/k} \setminus \{K\}$ possesses a natural structure of a $(\mathbb{Z}[\frac{1}{p}], |K^\times|)$ -PL space. In fact, the structure is even more

rigid since any change of coordinates has slopes in $p^{\mathbb{Z}}$. It is an interesting question if this non-typical restriction shows up elsewhere in the theory of PL spaces or their applications.

3.2.15. Full characterization of transition functions. Example 3.2.7 indicates that in the positive characteristic the PL structure we have just described is the finest natural structure on $I_{K/k}$. This holds in mixed characteristic too and can be formulated in the very precise form that the conditions of Theorem 3.2.12 are the only restrictions functions $\rho_{t,t'}^{\log}$ satisfy.

Theorem 3.2.16. *Let $K = \widehat{k(t)^a}/k$ be as in Theorem 3.2.12, then*

(i) *Any $(p^{\mathbb{Z}}, |K^\times|)$ -PL function ρ satisfying condition (ii) of Theorem 3.2.12 is of the form $\rho_{t,t'}^{\log}$ for an appropriate choice of t' .*

(ii) *The image of the map*

$$\Lambda_t: \text{Aut}_k(K) \rightarrow \text{Aut}(I_{K/k}) \xrightarrow{\sim} \text{Aut}([-\log r, \infty])$$

contains any $(p^{\mathbb{Z}}, \log |k^\times|)$ -PL function ρ on $[-\log r, \infty]$ which is the identity in a neighborhood of the endpoint $-\log r$.

Proof. The idea is to compose PL functions with a single corner. First we observe that computations in the proof of Theorem 3.2.12 easily imply that for any ρ as in (i) there exists t'' such that $\rho = \rho_{t'',t'}^{\log}$ locally at the endpoint $s_0 = -\log(r)$. Indeed, if $|k^\times| = |K^\times|$, then $\rho(s) = p^n s - \log |a|$ locally at s_0 , and one can take $t' = at^{p^n}$. Otherwise locally at s_0 we have that $\rho(s) = p^n s - (b-1)p^n \log(r) - \log |a|$ with $b \in \mathbb{Z} \setminus p\mathbb{Z}$ and $a \in k^\times$, and one can take $t' = at^{bp^n}$. Now we can deduce (i) from (ii) as follows: $\rho \circ (\rho_{t'',t'}^{\log})^{-1}$ is a $(p^{\mathbb{Z}}, \log |K^\times|)$ -PL function, which is the identity in a neighborhood of s_0 , hence if (ii) holds, then it comes from an automorphism ψ and hence is of the form $\rho_{t'',t'}^{\log}$, where $t'' = \psi(t')$. Thus, $\rho = \rho_{t'',t'}^{\log} \circ \rho_{t'',t'}^{\log} = \rho_{t,t'}^{\log}$.

The rest is devoted to the proof of (ii), so in the sequel we assume that ρ is the identity in a neighborhood of s_0 . First, suppose that the claim is proved in the particular case when ρ has a single corner and let us deduce the general claim. By induction one easily finds functions ρ_1, ρ_2, \dots such that ρ_n has a single break at s_n , $\rho_n(s) = s$ for $s < s_n$ and $\rho_n \circ \dots \circ \rho_1$ coincides with ρ on $[s_0, s_{n+1}]$. Choose any F_i corresponding to a point $s'_i \in (s_{i-1}, s_i)$, then by our assumption there exists $\psi_i \in \text{Aut}_{F_i}(K) \subset \text{Aut}_k(K)$ such that $\Lambda_t(\psi_i) = \rho_i$. Therefore $\Lambda_t(\psi_n \circ \dots \circ \psi_1)$ coincides with ρ on $[s_0, s_n]$, and it remains to show that if the number of corners is infinite, then the infinite composition $\dots \psi_2 \circ \psi_1$ converges. In this case s'_i tend to infinity, hence $\rho_t(F_i)$ monotonically decrease to 0. In particular, the infinite composition converges to a (valuation preserving) k -automorphism of $\cup_i F_i$, and hence also of its completion K .

It remains to consider the case when ρ has a single corner at a point $s_1 \in \log |K^\times|$ with $s_1 > s_0$. Let p^{-n} be the slope of ρ for $s > s_1$. Then ρ is the $-n$ -th composition power of the function ρ' which is the identity on $[s_0, s_1]$ and has slope p^{-1} for $s > s_1$. Therefore, it suffices to show that ρ' is in the image of Λ_t and for simplicity of notation we assume that ρ itself has slope p^{-1} for $s > s_1$.

We start with the simpler case when $\text{char}(k) = p$. Then by Example 3.2.7 $\rho = \rho_{t,t'}^{\log}$, where $t' = t + ct^{1/p}$ and $c \in k$ is chosen so that $\log |c| = (p^{-1} - 1)s_1$. Furthermore, $r > |cr^{1/p}|$ because $s_1 > s_0$, hence translating t by an element of k

and making $|t|$ close enough to r we can also achieve that $r > |ct^{1/p}|$. Then there exists an automorphism ψ taking t to t' , and we have that $\Lambda_t(\psi) = \rho$.

Finally, in the mixed characteristic case we will construct ψ by composing functions $\psi_i \in \text{Aut}_{F_i}(K)$ which take appropriate t_a (with $a \in F_i$) to $t_a + ct_a^{1/p}$. Let $x_i = \rho_i^{\log}(F_i) \geq s_0$ denote the endpoint. The behaviour of a function $\rho_i = \Lambda_t(\psi_i)$ was described in Example 3.2.9: it has slopes $1, p^{-1}, 1$, and corners $x'_i, x''_i \in \log |K^\times|$ such that $x_i < x'_i < x''_i < x_i - \frac{1}{p-1} \log |p|$. We claim that ψ_1, ψ_2, \dots can be chosen by induction so that each composition $\rho_n \circ \dots \circ \rho_1$ has slopes $1, p^{-1}, 1$, and corners at s_1 and y_n so that $\lim_n y_n = \infty$. Indeed, one should make the choices so that the first corner is given by $x'_1 = s_1$ and $x'_{n+1} = y_n$ for $n \geq 1$, the series $\sum_n (x''_n - x'_n)$ diverges (one can just take $x''_n = x'_n - \frac{1}{p} \log |p|$), and the endpoint x_n satisfies $x''_n + \frac{1}{p-1} \log |p| < x_n < x'_n$. In this case $\lim_n x_n = \infty$, and hence the composition $\psi = \dots \circ \psi_2 \circ \psi_1$ converges and satisfies $\Lambda_t(\psi) = \dots \circ \rho_2 \circ \rho_1 = \rho$. \square

4. LARGE EXTENSIONS

4.1. Large extensions of finite topological transcendence degree. So far we have studied small extensions, which always have a finite topological transcendence basis x_1, \dots, d_d and satisfy $d = \text{tr.c}(K/k) = \text{Tr.c}(K/k)$. In this subsection we will study the remaining class of extensions of finite topological transcendence degree, the large ones. Recall that these are extensions satisfying $\text{tr.c}(K/k) < \infty = \text{Tr.c}(K/k)$, they do not possess a topological transcendence basis, and we already know from Corollary 2.1.5 that they necessarily satisfy $\aleph_0 < \text{Tr.c}(K/k)$. Surprisingly, one can say quite a bit about such extensions and the mechanism preventing them from having a topological transcendence basis.

4.1.1. Large extensions of topological degree one. The one-dimensional case is very concrete:

Theorem 4.1.2. *Let K/k be an extension of algebraically closed analytic fields such that $\text{tr.c}(K/k) = 1$ and $\text{char}(\tilde{k}) > 0$. Then the following conditions are equivalent: (1) the extension is large, (2) $\text{Tr.c}(K/k) \geq \aleph_1$, (3) $\text{Tr.c}(K/k) = \aleph_1$, (4) the family $\{F_i\}_{i \in I}$ of intermediate algebraically closed analytic subextensions $k \subsetneq F_i \subsetneq K$ has uncountable cofinality, (5) the family $\{F_i\}$ has cofinality \aleph_1 , (6) ω_1 is the only uncountable ordinal that can be embedded into I , (7) K is the (uncompleted) union of F_i , (8) K is a maximal extension of k of topological degree one, (9) any k -endomorphism of K is an automorphism.*

Proof. First, assume that the extension is not large, say, $K = \widehat{k(x)^a}$, and let us show that all other conditions are not satisfied too. This is obvious for (2) and (3), conditions (4), (5), (6) fail by Theorem 2.2.6, (7) fails by Lemma 2.1.2, (8) fails by [Tem21, Theorem 5.2.2], and (9) fails by [Tem21, Theorem 5.2.5].

Conversely, assume that K/k is large and let us deduce all other conditions. First, if (8) fails, then there exists an extension $K \subsetneq L$ such that $\text{tr.c}(L/k) = 1$ and then any $x \in L \setminus K$ is t.a. independent of K and hence $K \subset \widehat{k(x)^a}$, which contradicts Theorem 2.3.2. By maximality, any k -endomorphism of K maps it onto itself, hence (9) holds too. By Theorem 2.1.4 (4) holds, and hence (7) holds by Lemma 2.1.2. To deduce (6), and hence also (5), from (4) we should show that if J is an uncountable well ordered subset of I , then J is cofinal. Indeed, $\cup_{j \in J} F_j$ is a large extension of k by the above paragraph, hence it is maximal by (8), and thus $\cup_{j \in J} F_j = K$.

Finally, by (6) there exists an uncountable cofinal family $\{F_j\}_{j \in J}$ of intermediate fields and J is of cardinality \aleph_1 . Since, each F_j is not large, $F_j = \widehat{k(x_j)^a}$, and hence the set $\{x_j\}_{j \in J}$ t.a. generates K over k , and we obtain that $\text{Tr.c}(K/k) \leq \aleph_1$. Since $\text{Tr.c}(K/k)$ is uncountable, (3) holds, and hence also (2). \square

Corollary 4.1.3. *Assume that K/k is a large extension of topological transcendence degree one. Then the ordered set $I_{K/k}$ of intermediate algebraically closed analytic fields is naturally homeomorphic to an Alexandroff ray (also known as a long ray) compactified by adding a point at infinity.*

Proof. By the above Theorem there exists a cofinal chain of fields $\{F_i\}_{i \in \omega_1} \subset I_{K/k}$. Hence $I_{K/k} \setminus \{K\}$ is the filtered union of ω_1 closed intervals, which implies that $I_{K/k} \setminus \{K\}$ is a long ray and adding the maximal point K compactifies it at infinity. \square

4.1.4. *Cofinality.* It is less clear, what can be said about large extensions in general, but at least one can use the same argument to bound the cofinality of $I_{K/k}$.

Theorem 4.1.5. *Let K/k be an extension of algebraically closed analytic fields such that $\text{tr.c}(K/k) = d < \infty$ and $\text{char}(\tilde{k}) > 0$. Then the cofinality of the set $I_{K/k}$ does not exceed the ordinal $d\omega_1$.*

Proof. If the cofinality is larger, then we can find subfields $k = F_0, F_1, \dots, F_d \in I_{K/k}$ such that $F_d \neq K$ and each interval $[F_i, F_{i+1}] = I_{F_{i+1}/F_i}$ in $I_{K/k}$ has uncountable cofinality. We claim that $\text{tr.c}(K/F_{i+1}) < \text{tr.c}(K/F_i)$ for any i . Once this is proved, one obtains that $\text{tr.c}(K/F_d) = 0$ and hence $K = F_d$, giving a contradiction.

Without restriction of generality it suffices to show that $\text{tr.c}(K/F) < d$ for $F = F_1$. Assume that this is not so, and hence there exists $x_1, \dots, x_d \in K$ which are t.a. independent over F . For any $x_0 \in F$ the elements x_0, \dots, x_d are t.a. dependent over k and this leaves only one possibility: $x_0 \in k(\widehat{x_1, \dots, x_d})^a$. Thus, the small extension $L = k(\widehat{x_1, \dots, x_d})^a$ of k contains F and hence $I_{L/k}$ has uncountable cofinality, which contradicts Theorem 2.2.6. \square

Remark 4.1.6. Assume that K/k is a maximal extension of k of topological transcendence degree d . Then the theorem easily implies that the cofinality of the set $I_{K/k}$ is $d'\omega_1$, where $1 \leq d' \leq d$. Of course it can happen that $d' = d$. For example, this is the case when K/k splits into a tower of d large extensions of topological transcendence degree one. It is a natural question, if any positive $d' < d$ can occur as well.

4.2. **Further questions.** I do not know about possible applications, but the natural further question to study is to which extent the same results hold for countably generated analytic extensions. In particular, this includes the following

Question 4.2.1. (i) *Assume that $\text{Tr.c}(K/k) = \aleph_0$. What is the maximal cofinality of a chain of intermediate algebraically closed analytic fields? Can it be uncountable?*

(ii) *Assume that $\text{tr.c}(K/k) = \aleph_0$. What is the maximal cardinality of $\text{Tr.c}(K/k)$? Can it exceed \aleph_1 ?*

(iii) *Can it happen that $\aleph_1 \leq \text{tr.c}(K/k) < \text{Tr.c}(K/k)$?*

REFERENCES

- [Hub96] Roland Huber, *Étale cohomology of rigid analytic varieties and adic spaces*, Aspects of Mathematics, E30, Friedr. Vieweg & Sohn, Braunschweig, 1996. MR 1734903 (2001c:14046)
- [KT18] Kiran Kedlaya and Michael Temkin, *Endomorphisms of power series fields and residue fields of Fargues-Fontaine curves*, Proceedings of AMS **146** (2018), 489–495.
- [Sch22] Peter Scholze, *Étale cohomology of diamonds*, 2022.
- [Tem10] Michael Temkin, *Stable modification of relative curves*, J. Algebraic Geom. **19** (2010), no. 4, 603–677. MR 2669727 (2011j:14064)
- [Tem16] ———, *Metrization of differential pluriforms on Berkovich analytic spaces*, Nonarchimedean and Tropical Geometry, Simons Symposia, Springer, 2016, pp. 195–285.
- [Tem21] ———, *Topological transcendence degree*, J. Algebra **568** (2021), 35–60. MR 4166051

EINSTEIN INSTITUTE OF MATHEMATICS, THE HEBREW UNIVERSITY OF JERUSALEM, GIV'AT RAM,
JERUSALEM, 91904, ISRAEL

Email address: `temkin@math.huji.ac.il`