

# NEW GOSPER'S LAMBERT SERIES IDENTITIES OF LEVELS 12 AND 16

RUSSELLE GUADALUPE

ABSTRACT. We derive new Gosper's Lambert series identities of levels 12 and 16 using certain sums of generalized  $\eta$ -quotients on the genus zero congruence subgroups  $\Gamma_0(12)$  and  $\Gamma_0(16)$ .

## 1. INTRODUCTION

For an element  $\tau$  in the complex upper half-plane  $\mathbb{H}$ , we define  $q := e^{2\pi i\tau}$ , and for any complex number  $a$ , we define  $(a; q)_\infty := \prod_{n=0}^\infty (1 - aq^n)$  and

$$(a_1, a_2, \dots, a_m; q)_\infty := (a_1; q)_\infty (a_2; q)_\infty \cdots (a_m; q)_\infty.$$

For any complex numbers  $a$  and  $b$  with  $|ab| < 1$ , the Ramanujan's general theta function is defined by

$$f(a, b) = \sum_{n=-\infty}^\infty a^{n(n+1)/2} b^{n(n-1)/2} = (-a, -b, ab; ab)_\infty,$$

where the last equality follows from the Jacobi triple product identity [4, p. 35, Entry 19]. We have the following special case given by  $\psi(q) := f(q, q^3) = (q^2; q^2)_\infty / (q; q^2)_\infty$ . In his study on the properties of the  $q$ -analogues of trigonometric functions, Gosper [6] introduced the function

$$\Pi_q := q^{1/4} \frac{(q^2; q^2)_\infty^2}{(q; q^2)_\infty^2} = q^{1/4} \psi^2(q)$$

and conjectured the following Lambert series identities

- (1) 
$$\sum_{n \geq 1} \frac{q^n}{(1 - q^n)^2} - 2 \sum_{n \geq 1} \frac{q^{2n}}{(1 - q^{2n})^2} = \frac{1}{24} \left( \frac{\Pi_q^4}{\Pi_{q^2}^2} - 1 \right) + \frac{2}{3} \Pi_{q^2}^2,$$
- (2) 
$$\frac{1}{\Pi_{q^5}^2} \left( \sum_{n \geq 1} \frac{q^{2n-1}}{(1 - q^{2n-1})^2} - 5 \sum_{n \geq 1} \frac{q^{10n-5}}{(1 - q^{10n-5})^2} \right) = \sqrt{\frac{\Pi_q^3}{\Pi_{q^5}^3} - 2 \frac{\Pi_q^2}{\Pi_{q^5}^2} + 5 \frac{\Pi_q}{\Pi_{q^5}}},$$

$$6 \left( \sum_{n \geq 1} \frac{q^n}{(1 - q^n)^2} - 5 \sum_{n \geq 1} \frac{q^{5n}}{(1 - q^{5n})^2} \right) + 1$$
- (3) 
$$= \left( \frac{\Pi_q}{\Pi_{q^5}} + 2 + 5 \frac{\Pi_{q^5}}{\Pi_q} \right) \left( \sum_{n \geq 1} \frac{q^{2n-1}}{(1 - q^{2n-1})^2} - 5 \sum_{n \geq 1} \frac{q^{10n-5}}{(1 - q^{10n-5})^2} \right),$$
- (4) 
$$\frac{1}{\Pi_{q^6}^2} \left( \sum_{n \geq 1} \frac{q^{2n-1}}{(1 - q^{2n-1})^2} - 6 \sum_{n \geq 1} \frac{q^{12n-6}}{(1 - q^{12n-6})^2} \right) = \frac{\Pi_{q^2}^2}{\Pi_{q^6}^2} + 2 \frac{\Pi_{q^2}}{\Pi_{q^6}},$$

$$\frac{1}{\Pi_{q^9}^2} \left( \sum_{n \geq 1} \frac{q^{2n-1}}{(1 - q^{2n-1})^2} - 9 \sum_{n \geq 1} \frac{q^{18n-9}}{(1 - q^{18n-9})^2} \right)$$
- (5) 
$$= \left( \frac{\Pi_q}{\Pi_{q^9}} + 3 \right) \sqrt{\left( \frac{\Pi_q}{\Pi_{q^9}} \right)^{3/2} - 3 \frac{\Pi_q}{\Pi_{q^9}} + 3 \left( \frac{\Pi_q}{\Pi_{q^9}} \right)^{1/2}},$$

---

2020 *Mathematics Subject Classification*: Primary 11F03, 11F11, 33D15.

*Key words and phrases*: Gosper's identities, Lambert series,  $\eta$ -quotients, generalized  $\eta$ -quotients

$$\begin{aligned}
& 3 \left( \sum_{n \geq 1} \frac{q^n}{(1-q^n)^2} - 9 \sum_{n \geq 1} \frac{q^{9n}}{(1-q^{9n})^2} \right) + 1 \\
(6) \quad & = \left( \sqrt{\frac{\Pi_q}{\Pi_{q^9}}} + 3\sqrt{\frac{\Pi_{q^9}}{\Pi_q}} \right) \left( \sum_{n \geq 1} \frac{q^{2n-1}}{(1-q^{2n-1})^2} - 9 \sum_{n \geq 1} \frac{q^{18n-9}}{(1-q^{18n-9})^2} \right)
\end{aligned}$$

involving  $\Pi_q$ . Using the methods of Gosper and integrals of Ramanujan's theta functions, El Bachraoui [1] proved (1). He [8, 9] confirmed (2) and (3) using Ramanujan's modular equations of degree five and proved (5) via modular forms. Wang [13] observed that the above identities can be seen as equalities of certain modular forms on the congruence subgroup  $\Gamma_0(N)$  and called such equality an identity of level  $N$ . Wang also proved these identities and devised a method of discovering Lambert series identities of level  $N$  via Eisenstein series and a basis for the vector space of modular forms on  $\Gamma_0(N)$  of some positive weight. Yathirajsharma, Harshitha, and Vasuki [15] deduced the above identities except (4) by applying Ramanujan's theta function identities.

In this paper, we offer another proof of the identity (4) of level 12 and derive new Lambert series identities of levels 12 and 16 using sums of generalized  $\eta$ -quotients on  $\Gamma_0(N)$  for  $N \in \{12, 16\}$ .

**Theorem 1.1** (level 12 identity). *We have*

$$\begin{aligned}
& \frac{1}{\Pi_{q^6}^2} \left[ 24 \left( \sum_{n \geq 1} \frac{q^n}{(1-q^n)^2} - 6 \sum_{n \geq 1} \frac{q^{6n}}{(1-q^{6n})^2} \right) + 5 \right] \\
(7) \quad & = 5 \frac{\Pi_{q^2}^3}{\Pi_{q^6}^3} + 24 \frac{\Pi_{q^2}^2}{\Pi_{q^6}^2} + 42 \frac{\Pi_{q^2}}{\Pi_{q^6}} + 9 \frac{\Pi_{q^6}}{\Pi_{q^2}}.
\end{aligned}$$

**Theorem 1.2** (level 16 identities). *We have*

$$\begin{aligned}
(8) \quad & \frac{1}{\Pi_{q^8}^2} \left( \sum_{n \geq 1} \frac{q^{2n-1}}{(1-q^{2n-1})^2} - 8 \sum_{n \geq 1} \frac{q^{16n-8}}{(1-q^{16n-8})^2} \right) = \frac{\Pi_{q^4}^3}{\Pi_{q^8}^3} + 2 \frac{\Pi_{q^4}^2}{\Pi_{q^8}^2} + 4 \frac{\Pi_{q^4}}{\Pi_{q^8}} + 4, \\
(9) \quad & \frac{1}{\Pi_{q^8}^2} \left[ 24 \left( \sum_{n \geq 1} \frac{q^n}{(1-q^n)^2} - 8 \sum_{n \geq 1} \frac{q^{8n}}{(1-q^{8n})^2} \right) + 7 \right] \\
& = 7 \frac{\Pi_{q^4}^4}{\Pi_{q^8}^4} + 24 \frac{\Pi_{q^4}^3}{\Pi_{q^8}^3} + 72 \frac{\Pi_{q^4}^2}{\Pi_{q^8}^2} + 96 \frac{\Pi_{q^4}}{\Pi_{q^8}} + 112.
\end{aligned}$$

Our main tool for proving (4) and Theorems 1.1 and 1.2 is the following identity of Bailey [3] given by

$$(10) \quad \sum_{n=-\infty}^{\infty} \left[ \frac{aq^n}{(1-aq^n)^2} - \frac{bq^n}{(1-bq^n)^2} \right] = af^6(-q) \frac{f(-ab, -\frac{q}{ab})f(-\frac{b}{a}, -\frac{aq}{b})}{f^2(-a, -\frac{q}{a})f^2(-b, -\frac{q}{b})}$$

for  $|ab| < 1$ , which is a special case of his very well-poised  ${}_6\psi_6$  summation formula (see [2, (4.7)]). This formula transforms a given Lambert series identity of level  $N$  into sums of products of theta functions, which in turn can be written as sums of generalized  $\eta$ -quotients on  $\Gamma_0(N)$ . We then exploit the fact that  $\Gamma_0(N)$  has genus zero for  $N \in \{12, 16\}$ , which will be essential in establishing these identities by expressing them in terms of quotients of  $\Pi_q$ . We note that our method presented in this paper is different from that of He [9] and Wang [13] in that we only apply modular functions instead of modular forms of positive weight to derive the above identities. In addition, the author [7] quite recently obtained two new Gosper's Lambert series identities of level 14 using this method.

We organize the rest of the paper as follows. We give a short background on modular functions, particularly  $\eta$ -quotients and generalized  $\eta$ -quotients, on the congruence subgroups

$\Gamma_1(N)$  and  $\Gamma_0(N)$  in Section 2. Applying Bailey's formula (10) and some properties of sums of generalized  $\eta$ -quotients on  $\Gamma_0(N)$  for  $N \in \{12, 16\}$ , we establish identity (4) and Theorem 1.1 in Section 3, and Theorem 1.2 in Section 4. In the proofs of these identities, we express the corresponding sum of generalized  $\eta$ -quotients on  $\Gamma_0(N)$  in terms of the generator of the field of modular functions on  $\Gamma_0(N)$  involving quotients of  $\Pi_q$ . We have executed most of our computations using *Mathematica*.

## 2. MODULAR FUNCTIONS ON $\Gamma_1(N)$ AND $\Gamma_0(N)$

We consider in this paper modular functions on congruence subgroups

$$\Gamma_1(N) := \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \mathrm{SL}_2(\mathbb{Z}) : a - 1 \equiv c \equiv d - 1 \equiv 0 \pmod{N} \right\},$$

$$\Gamma_0(N) := \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \mathrm{SL}_2(\mathbb{Z}) : c \equiv 0 \pmod{N} \right\}.$$

The action an element of a congruence subgroup  $\Gamma$  on the extended upper half-plane  $\mathbb{H}^* := \mathbb{H} \cup \mathbb{Q} \cup \{\infty\}$  can be described by a linear fractional transformation. We define the cusps of  $\Gamma$  as the equivalence classes of  $\mathbb{Q} \cup \{\infty\}$  under this action. We define a modular function on  $\Gamma$  as a meromorphic function  $f : \mathbb{H}^* \rightarrow \mathbb{C}$  such that  $f(\gamma\tau) = f(\tau)$  for all  $\gamma \in \Gamma$  and the  $q$ -expansion of  $f$  at each cusp  $r$  of  $\Gamma$  is

$$f(\gamma\tau) = \sum_{n \geq n_0} a_n q^{n/h}$$

for some  $\gamma \in \mathrm{SL}_2(\mathbb{Z})$  with  $\gamma(\infty) = r$  and integers  $h$  and  $n_0$  with  $a_{n_0} \neq 0$ . We call  $n_0$  the order of  $f(\tau)$  at  $r$ , denoted by  $\mathrm{ord}_r f(\tau)$ , and  $h$  the width of  $r$ , which is the smallest positive integer such that  $\gamma \begin{bmatrix} 1 & h \\ 0 & 1 \end{bmatrix} \gamma^{-1} \in \pm\Gamma$ . We say that  $f(\tau)$  has a zero (resp., a pole) at  $r$  if  $\mathrm{ord}_r f(\tau) > 0$  (resp.,  $\mathrm{ord}_r f(\tau) < 0$ ).

Cho, Koo and Park [5] gave a nice description on the inequivalent cusps of  $\Gamma_0(N)$  and their widths.

**Lemma 2.1.** *Let  $a, a', c$  and  $c'$  be integers with  $\mathrm{gcd}(a, c) = \mathrm{gcd}(a', c') = 1$ . Let  $S_{\Gamma_0(N)}$  be the set of all inequivalent cusps of  $\Gamma_0(N)$ . We denote  $\pm 1/0$  as  $\infty$ . Then*

- (1)  $a/c$  and  $a'/c'$  are equivalent over  $\Gamma_0(N)$  if and only if there exist an integer  $n$  and an element  $\bar{s} \in (\mathbb{Z}/N\mathbb{Z})^\times$  such that  $(a', c') \equiv (\bar{s}^{-1}a + nc, \bar{s}c) \pmod{N}$ ,
- (2) we have

$$S_{\Gamma_0(N)} = \{a_{c,j}/c \in \mathbb{Q} : 0 < c \mid N, 0 < a_{c,j} \leq N, \mathrm{gcd}(a_{c,j}, N) = 1,$$

$$a_{c,j} = a_{c,j'} \stackrel{\text{def}}{\iff} a_{c,j} \equiv a_{c,j'} \pmod{\mathrm{gcd}(c, N/c)}\},$$

- (3) the width of the cusp  $a/c \in S_{\Gamma_0(N)}$  is  $N/\mathrm{gcd}(c^2, N)$ .

*Proof.* See [5, Corollary 4(1)]. □

Let  $\mathcal{M}^\infty(N)$  be the set of all modular functions on  $\Gamma_0(N)$  that have poles only at infinity. Since  $\Gamma_0(N)$  has genus zero for  $N \in \{12, 16\}$ , we see that  $\mathcal{M}^\infty(N)$  is generated by a single function  $g \in \mathcal{M}^\infty(N)$ . The following result shows that for these values of  $N$ , every element of  $\mathcal{M}^\infty(N)$  can be expressed in terms of  $g$ .

**Lemma 2.2.** *Let  $N \in \{12, 16\}$  and  $h \in \mathcal{M}^\infty(N)$ . Set  $m := \mathrm{ord}_\infty h$ . Then  $h$  can be written as a polynomial in  $g$  of degree  $-m$  with complex coefficients.*

*Proof.* This follows from [11, Theorem 8.3]. □

We next define an  $\eta$ -quotient given by

$$f(\tau) = \prod_{\delta \mid N} \eta(\delta\tau)^{r_\delta}$$

for some indexed set  $\{r_\delta \in \mathbb{Z} : \delta \mid N\}$ , where  $\eta(\tau) = q^{1/24}(q; q)_\infty$  is the Dedekind eta function. The following results give conditions for an  $\eta$ -quotient to be modular on  $\Gamma_0(N)$ .

**Lemma 2.3.** Let  $f(\tau) = \prod_{\delta|N} \eta(\delta\tau)^{r_\delta}$  be an  $\eta$ -quotient with  $k = \frac{1}{2} \sum_{\delta|N} r_\delta \in \mathbb{Z}$  such that

$$\sum_{\delta|N} \delta r_\delta \equiv 0 \pmod{24} \quad \text{and} \quad \sum_{\delta|N} \frac{N}{\delta} r_\delta \equiv 0 \pmod{24}.$$

Then for all  $\begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \Gamma_0(N)$ , we have

$$f\left(\frac{a\tau + b}{c\tau + d}\right) = \left(\frac{(-1)^k \prod_{\delta|N} \delta^{r_\delta}}{d}\right) (c\tau + d)^k f(\tau),$$

where  $\left(\frac{\cdot}{d}\right)$  is the Legendre symbol.

*Proof.* See [10, Theorem 1.64]. □

**Lemma 2.4.** Let  $c, d$  and  $N$  be positive integers with  $d | N$  and  $\gcd(c, d) = 1$  and let  $f(\tau) = \prod_{\delta|N} \eta(\delta\tau)^{r_\delta}$  be an  $\eta$ -quotient satisfying the conditions of Lemma 2.3. Then the order of  $f(\tau)$  at the cusp  $c/d$  is

$$\frac{N}{24d \gcd(d, \frac{N}{d})} \sum_{\delta|N} \gcd(d, \delta)^2 \cdot \frac{r_\delta}{\delta}.$$

*Proof.* See [10, Theorem 1.65]. □

We now define a generalized  $\eta$ -quotient given by

$$f(\tau) = \prod_{1 \leq g \leq \lfloor N/2 \rfloor} \eta_{N,g}(\tau)^{r_g}$$

for some indexed set  $\{r_g \in \mathbb{Z} : 1 \leq g \leq \lfloor N/2 \rfloor\}$ , where

$$\eta_{N,g}(\tau) = q^{NB_2(g/N)/2} (q^g, q^{N-g}; q^N)_\infty$$

is the generalized Dedekind eta function with  $B_2(t) := t^2 - t + 1/6$ . The following results provide the transformation formula of  $\eta_{N,g}$  on  $\Gamma_0(N)$  and conditions for a generalized  $\eta$ -quotient to be modular on  $\Gamma_1(N)$ .

**Lemma 2.5.** The function  $\eta_{N,g}(\tau)$  satisfies  $\eta_{N,g+N}(\tau) = \eta_{N,-g}(\tau) = -\eta_{N,g}(\tau)$ . Moreover, let  $\gamma = \begin{bmatrix} a & b \\ cN & d \end{bmatrix} \in \Gamma_0(N)$  with  $c \neq 0$ . Then we have

$$\eta_{N,g}(\gamma\tau) = \varepsilon(a, bN, c, d) e^{\pi i(g^2 ab/N - gb)} \eta_{N,ag}(\tau),$$

where

$$\varepsilon(a, b, c, d) = \begin{cases} e^{\pi i(bd(1-c^2) + c(a+d-3))/6}, & \text{if } c \equiv 1 \pmod{2}, \\ -ie^{\pi i(ac(1-d^2) + d(b-c+3))/6}, & \text{if } c \equiv 0 \pmod{2}. \end{cases}$$

*Proof.* See [14, Corollary 2]. □

**Lemma 2.6.** Suppose  $f(\tau) = \prod_{1 \leq g \leq \lfloor N/2 \rfloor} \eta_{N,g}(\tau)^{r_g}$  is a generalized  $\eta$ -quotient such that

- (1)  $\sum_{1 \leq g \leq \lfloor N/2 \rfloor} r_g \equiv 0 \pmod{12}$ ,
- (2)  $\sum_{1 \leq g \leq \lfloor N/2 \rfloor} gr_g \equiv 0 \pmod{2}$ , and
- (3)  $\sum_{1 \leq g \leq \lfloor N/2 \rfloor} g^2 r_g \equiv 0 \pmod{2N}$ .

Then  $f(\tau)$  is a modular function on  $\Gamma_1(N)$ .

*Proof.* See [14, Corollary 3]. □

**Lemma 2.7.** Let  $N$  be a positive integer and  $\gamma = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in SL_2(\mathbb{Z})$ . Then the first term of the  $q$ -expansion of  $\eta_{N,g}(\gamma\tau)$  is  $\varepsilon q^\delta$ , where  $|\varepsilon| = 1$  and

$$\delta = \frac{\gcd(c, N)^2}{2N} P_2\left(\frac{ag}{\gcd(c, N)}\right)$$

with  $P_2(t) := B_2(\{t\})$  and  $\{t\}$  is the fractional part of  $t$ .

*Proof.* See [14, Lemma 2]. □

We apply Lemmas 2.1 and 2.7 to find the order of a generalized  $\eta$ -quotient at a cusp of  $\Gamma_0(N)$ .

### 3. PROOFS OF IDENTITY (4) AND THEOREM 1.1

We first derive the level 12 identity (4) and prove Theorem 1.1 using sums of generalized  $\eta$ -quotients on  $\Gamma_0(12)$ .

*Proof of (4).* We replace  $q$  with  $q^{12}$  and set  $b = q^6$  in (10) so that

$$(11) \quad \sum_{n=-\infty}^{\infty} \left[ \frac{aq^{12n}}{(1-aq^{12n})^2} - \frac{q^{12n+6}}{(1-q^{12n+6})^2} \right] = aq^{-3} \Pi_{q^6}^2 \frac{f^2(-aq^6, -\frac{q^6}{a})}{f^2(-a, -\frac{q^{12}}{a})}.$$

Substituting  $a = q, q^3, q^5$  in (11) and adding the resulting identities lead to

$$(12) \quad \frac{1}{\Pi_{q^6}^2} \left( \sum_{n \geq 1} \frac{q^{2n-1}}{(1-q^{2n-1})^2} - 6 \sum_{n \geq 1} \frac{q^{12n-6}}{(1-q^{12n-6})^2} \right) =: h_1(\tau) + 1 + \frac{1}{h_1(\tau)},$$

where

$$h_1(\tau) := \frac{\eta_{12,5}^2(\tau)}{\eta_{12,1}^2(\tau)}$$

is modular on  $\Gamma_1(12)$  by Lemma 2.6 and satisfies  $h_1(\gamma\tau) = 1/h_1(\tau)$  by Lemma 2.5 with  $\gamma := \begin{bmatrix} 5 & 2 \\ 12 & 5 \end{bmatrix}$ . Since  $\Gamma_1(12)$  and  $\gamma$  generate  $\Gamma_0(12)$ , we see that  $h_1(\tau) + 1/h_1(\tau)$  is modular on  $\Gamma_0(12)$ . From Lemma 2.1, we have

$$S_{\Gamma_0(12)} := \{\infty, 0, 1/2, 1/3, 1/4, 1/6\}.$$

We use Lemma 2.7 to obtain the orders of  $h_1(\tau) + 1/h_1(\tau)$  at each cusp  $r \in S_{\Gamma_0(12)}$ , as shown in Table 1 below.

cusp $r$	0	1/2	1/3	1/4	1/6	$\infty$
$\text{ord}_r h_1(\tau)$	0	0	0	0	0	-2
$\text{ord}_r 1/h_1(\tau)$	0	0	0	0	0	2
$\text{ord}_r h_1(\tau) + 1/h_1(\tau)$	$\geq 0$	$\geq 0$	$\geq 0$	$\geq 0$	$\geq 0$	-2

TABLE 1. The orders of  $h_1(\tau) + 1/h_1(\tau)$  at the cusps of  $\Gamma_0(12)$

We see that  $h_1(\tau) + 1/h_1(\tau) \in \mathcal{M}^\infty(12)$  with unique double pole at infinity. We note that the  $\eta$ -quotient

$$h := \frac{\Pi_{q^2}}{\Pi_{q^6}} = \frac{\eta^4(4\tau)\eta^2(6\tau)}{\eta^2(2\tau)\eta^4(12\tau)}$$

generates  $\mathcal{M}^\infty(12)$  with unique simple pole at infinity by Lemmas 2.3 and 2.4. We infer from Lemma 2.2 that  $h_1(\tau) + 1/h_1(\tau)$  is a quadratic polynomial in  $h$ . From the  $q$ -expansion

$$\begin{aligned} h_1(\tau) + \frac{1}{h_1(\tau)} &= q^{-2} + 2q^{-1} + 3 + 4q + 6q^2 + O(q^3), \\ h &= q^{-1} + 2q + q^3 + O(q^4), \end{aligned}$$

we deduce that

$$(13) \quad h_1(\tau) + \frac{1}{h_1(\tau)} = h^2 + 2h - 1.$$

Comparing (12) and (13), we get identity (4). □

We require the following result about a modular function on  $\mathcal{M}^\infty(12)$  involving

$$h_2(\tau) := \frac{\Pi_{q^6}^2}{\Pi_{q^{12}}^2} = \frac{\eta^{12}(12\tau)}{\eta^4(6\tau)\eta^8(24\tau)}.$$

**Lemma 3.1.** *The function*

$$H := H(\tau) := h_2(\tau) + \frac{16}{h_2(\tau)}$$

is modular on  $\Gamma_0(12)$ , and  $hH$  is in  $\mathcal{M}^\infty(12)$  with  $\text{ord}_\infty hH = -4$ .

*Proof.* Observe that by Lemma 2.3,  $h_2(\tau)$  is modular on  $\Gamma_0(24)$ . Since  $\Gamma_0(12)$  is generated by  $\Gamma_0(24)$  and  $\alpha := \begin{bmatrix} 1 & 0 \\ 12 & 1 \end{bmatrix}$ , we know that  $H(\tau) = h_2(\tau) + h_2(\alpha\tau)$  is a modular function on  $\Gamma_0(12)$ . We will show that

$$(14) \quad h_2(\alpha\tau) = \frac{16}{h_2(\tau)}.$$

Consider the function  $H_0(\tau) := h_2(\alpha\tau)h_2(\tau)$ . Then  $H_0(\tau)$  is modular on  $\Gamma_0(24)$  since  $\Gamma_0(24)$  is a normal subgroup of  $\Gamma_0(12)$ . We find the orders of  $H_0(\tau)$  at the cusps of  $\Gamma_0(24)$  given by

$$S_{\Gamma_0(24)} := \{\infty, 0, 1/2, 1/3, 1/4, 1/6, 1/8, 1/12\}$$

from Lemma 2.1 as follows. We first look for the cusps  $r_1, r_2 \in S_{\Gamma_0(24)}$  such that  $\alpha(r_1)$  is equivalent to  $r_2$  over  $\Gamma_0(24)$ , so that by [12, Lemma 2], we have

$$(15) \quad \text{ord}_{r_1} h_2(\alpha\tau) = \text{ord}_{\alpha(r_1)} h_2(\tau) = \text{ord}_{r_2} h_2(\tau).$$

We denote this equivalence by  $\alpha(r_1) \sim r_2$ . Using Lemma 2.1, we find that

$$\begin{aligned} \alpha(0) &\sim 0, & \alpha(1/2) &\sim 1/2, & \alpha(1/3) &\sim 1/3, & \alpha(1/4) &\sim 1/8, \\ \alpha(1/6) &\sim 1/6, & \alpha(1/8) &\sim 1/4, & \alpha(1/12) &\sim \infty, & \alpha(\infty) &\sim 1/12. \end{aligned}$$

We then compute the required orders using Lemma 2.4 and (15), as shown in Table 2 below. Consequently, we see that  $H_0(\tau)$  is holomorphic at every cusp of  $\Gamma_0(24)$ , so  $H_0(\tau)$  is constant by [12, Lemma 5].

cusps $r$	0	1/2	1/3	1/4	1/6	1/8	1/12	$\infty$
$\text{ord}_r h_2(\alpha\tau)$	0	0	0	-1	0	1	-3	3
$\text{ord}_r h_2(\tau)$	0	0	0	1	0	-1	3	-3
$\text{ord}_r H_0(\tau)$	0	0	0	0	0	0	0	0

TABLE 2. The orders of  $H_0(\tau)$  at the cusps of  $\Gamma_0(24)$

We now use the transformation formula [14, Lemma 1]

$$\eta\left(-\frac{1}{\tau}\right) = \sqrt{-i\tau}\eta(\tau)$$

to find the constant  $H_0(0) = h_2^2(0)$ . We compute

$$h_2(0) = \lim_{\tau \rightarrow \infty} \frac{\eta^{12}(-12/\tau)}{\eta^4(-6/\tau)\eta^8(-24/\tau)} = 4 \lim_{\tau \rightarrow \infty} \frac{(q^{1/12}; q^{1/12})_\infty^{12}}{(q^{1/6}; q^{1/6})_\infty^4 (q^{1/24}; q^{1/24})_\infty^8} = 4,$$

so that  $H_0(0) = 16$ . Thus, (14) follows.

We now proceed to finding the orders of  $H(\tau)$  at the cusps of  $\Gamma_0(12)$  as follows. We first write  $h_2(\tau)$  as a generalized  $\eta$ -quotient

$$h_2(\tau) = \frac{\eta_{24,12}^4(\tau)}{\eta_{24,6}^4(\tau)}.$$

For each  $1/c \in S_{\Gamma_0(12)}$ , let  $\alpha_c \in \mathrm{SL}_2(\mathbb{Z})$  be the matrix with  $\alpha_c(\infty) = 1/c$ . By Lemma 2.7, the first term of the  $q$ -expansion of  $h_2(\alpha_c\tau)$  is  $\varepsilon q^{\delta_c}$  for some  $|\varepsilon| = 1$ , where

$$\delta_c := \frac{\mathrm{gcd}(c, 24)^2}{12} \left[ P_2 \left( \frac{12}{\mathrm{gcd}(c, 24)} \right) - P_2 \left( \frac{6}{\mathrm{gcd}(c, 24)} \right) \right].$$

In view of Lemma 2.1, we see that for  $1/c \in S_{\Gamma_0(12)}$ ,

$$(16) \quad \mathrm{ord}_{1/c} h_2(\tau) = \frac{12}{\mathrm{gcd}(c^2, 12)} \delta_c.$$

Using (16), we record the required orders as shown in Table 3 below.

cusps $r$	0	1/2	1/3	1/4	1/6	$\infty$
$\mathrm{ord}_r h_2(\tau)$	0	0	0	1	0	3
$\mathrm{ord}_r 1/h_2(\tau)$	0	0	0	-1	0	-3
$\mathrm{ord}_r H(\tau)$	$\geq 0$	$\geq 0$	$\geq 0$	-1	$\geq 0$	-3

TABLE 3. The orders of  $H(\tau)$  at the cusps of  $\Gamma_0(12)$

Since  $\mathrm{ord}_{1/4} h = 1$  by Lemma 2.4, we surmise that  $hH$  has a unique pole of order  $-4$  at infinity and holomorphic elsewhere, so that  $hH \in \mathcal{M}^\infty(12)$ .  $\square$

*Proof of Theorem 1.1.* Setting  $a = q^2$  and  $a = q^4$  in (11) and adding the resulting identities yield

$$(17) \quad \sum_{n \geq 1} \left[ \frac{q^{2n}}{(1 - q^{2n})^2} - \frac{q^{6n}}{(1 - q^{6n})^2} \right] - 4 \sum_{n \geq 1} \frac{q^{12n-6}}{(1 - q^{12n-6})^2} = \Pi_{q^6}^2 \left( h_3(\tau) + \frac{1}{h_3(\tau)} \right),$$

where

$$h_3(\tau) := \frac{\eta_{12,4}^2(\tau)}{\eta_{12,2}^2(\tau)}$$

is modular on  $\Gamma_1(12)$  by Lemma 2.6 and satisfies  $h_3(\gamma\tau) = 1/h_3(\tau)$  by Lemma 2.5. Thus,  $h_3(\tau) + 1/h_3(\tau)$  is a modular function on  $\Gamma_0(12)$  whose orders at the cusps of  $\Gamma_0(12)$  can be found via Lemma 2.7. Table 4 shows the computed orders. Recalling that  $\mathrm{ord}_{1/4} h = 1$ , we have

cusps $r$	0	1/2	1/3	1/4	1/6	$\infty$
$\mathrm{ord}_r h_3(\tau)$	0	0	0	1	0	-1
$\mathrm{ord}_r 1/h_3(\tau)$	0	0	0	-1	0	1
$\mathrm{ord}_r h_3(\tau) + 1/h_3(\tau)$	$\geq 0$	$\geq 0$	$\geq 0$	-1	$\geq 0$	-1

TABLE 4. The orders of  $h_3(\tau) + 1/h_3(\tau)$  at the cusps of  $\Gamma_0(12)$

that  $h(h_3(\tau) + 1/h_3(\tau)) \in \mathcal{M}^\infty(12)$  with unique double pole at infinity, so that it is a quadratic polynomial in  $h$  according to Lemma 2.2. By looking at the  $q$ -expansion

$$h_3(\tau) + \frac{1}{h_3(\tau)} = q^{-1} + 3q - q^3 + O(q^4),$$

we obtain

$$h \left( h_3(\tau) + \frac{1}{h_3(\tau)} \right) = h^2 + 1,$$

so that the right-hand side of (17) becomes

$$(18) \quad \sum_{n \geq 1} \left[ \frac{q^{2n}}{(1 - q^{2n})^2} - \frac{q^{6n}}{(1 - q^{6n})^2} \right] - 4 \sum_{n \geq 1} \frac{q^{12n-6}}{(1 - q^{12n-6})^2} = \Pi_{q^6}^2 \left( h + \frac{1}{h} \right).$$

Invoking (12), (13), and the identity

$$(19) \quad \sum_{n \geq 1} \left[ \frac{q^n}{(1-q^n)^2} - \frac{q^{2n}}{(1-q^{2n})^2} \right] = \sum_{n \geq 1} \frac{q^{2n-1}}{(1-q^{2n-1})^2},$$

we transform (18) into

$$(20) \quad \frac{1}{\Pi_{q^6}^2} \sum_{n \geq 1} \left[ \frac{q^n}{(1-q^n)^2} - 6 \frac{q^{6n}}{(1-q^{6n})^2} \right] - h^2 - 3h - \frac{1}{h} \\ = \frac{5}{\Pi_{q^6}^2} \sum_{n \geq 1} \left[ \frac{q^{12n-6}}{(1-q^{12n-6})^2} - \frac{q^{12n}}{(1-q^{12n})^2} \right].$$

Replacing  $q$  with  $q^6$  in (1), substituting it on the right-hand side of (20), and then multiplying both sides by 24, we arrive at

$$(21) \quad \frac{1}{\Pi_{q^6}^2} \left[ 24 \left( \sum_{n \geq 1} \frac{q^n}{(1-q^n)^2} - 6 \sum_{n \geq 1} \frac{q^{6n}}{(1-q^{6n})^2} \right) + 5 \right] - 24h^2 - 72h - \frac{24}{h} = 5H.$$

We know from Lemma 3.1 that  $hH \in \mathcal{M}^\infty(12)$  with  $\text{ord}_\infty hH = -4$ . Consequently, by Lemma 2.2,  $hH$  is a quartic polynomial in  $h$ . It follows from the  $q$ -expansion

$$hH = q^{-4} + 2q^{-2} + 1 + 20q^2 + O(q^4)$$

that

$$hH = h^4 - 6h^2 - 3,$$

so we conclude from (21) that

$$\frac{1}{\Pi_{q^6}^2} \left[ 24 \left( \sum_{n \geq 1} \frac{q^n}{(1-q^n)^2} - 6 \sum_{n \geq 1} \frac{q^{6n}}{(1-q^{6n})^2} \right) + 5 \right] = 5h^3 + 24h^2 + 42h + \frac{9}{h},$$

which is precisely identity (7). □

#### 4. PROOF OF THEOREM 1.2

We next prove Theorem 1.2 by employing certain sums of generalized  $\eta$ -quotients on  $\Gamma_0(16)$ .

*Proof of Theorem 1.2.* Replacing  $q$  with  $q^{16}$  and setting  $b = q^8$  in (10) give

$$(22) \quad \sum_{n=-\infty}^{\infty} \left[ \frac{aq^{16n}}{(1-aq^{16n})^2} - \frac{q^{16n+8}}{(1-q^{16n+8})^2} \right] = aq^{-4} \Pi_{q^8}^2 \frac{f^2(-aq^8, -\frac{q^8}{a})}{f^2(-a, -\frac{q^{16}}{a})}.$$

Substituting  $a = q, q^3, q^5, q^7$  in (22) and combining the resulting identities, we have

$$(23) \quad \frac{1}{\Pi_{q^8}^2} \left( \sum_{n \geq 1} \frac{q^{2n-1}}{(1-q^{2n-1})^2} - 8 \sum_{n \geq 1} \frac{q^{16n-8}}{(1-q^{16n-8})^2} \right) =: s_1(\tau) + s_2(\tau) + \frac{1}{s_1(\tau)} + \frac{1}{s_2(\tau)},$$

where

$$s_1(\tau) := \frac{\eta_{16,7}^2(\tau)}{\eta_{16,1}^2(\tau)}, \quad s_2(\tau) := \frac{\eta_{16,5}^2(\tau)}{\eta_{16,3}^2(\tau)}$$

are modular on  $\Gamma_1(16)$  by Lemma 2.6. By Lemma 2.5, we have

$$s_1(\gamma\tau) = s_2(\tau), \quad s_2(\gamma\tau) = \frac{1}{s_1(\tau)},$$

where  $\gamma := \begin{bmatrix} 3 & -1 \\ 16 & -5 \end{bmatrix}$ . As  $\Gamma_1(16)$  and  $\gamma$  generate  $\Gamma_0(16)$ , we deduce that

$$z := s_1(\tau) + s_2(\tau) + \frac{1}{s_1(\tau)} + \frac{1}{s_2(\tau)}$$

is modular on  $\Gamma_0(16)$ . From Lemma 2.1, we compute

$$S_{\Gamma_0(16)} := \{\infty, 0, 1/2, 1/4, 3/4, 1/8\}.$$

Applying Lemma 2.7, we calculate the orders of  $z$  at each element of  $S_{\Gamma_0(16)}$  as shown in Table 5 below.

cusps $r$	0	1/2	1/4, 3/4	1/8	$\infty$
$\text{ord}_r s_1(\tau)$	0	0	0	0	-3
$\text{ord}_r s_2(\tau)$	0	0	0	0	-1
$\text{ord}_r 1/s_1(\tau)$	0	0	0	0	3
$\text{ord}_r 1/s_2(\tau)$	0	0	0	0	1
$\text{ord}_r z$	$\geq 0$	$\geq 0$	$\geq 0$	$\geq 0$	-3

TABLE 5. The orders of  $z$  at the cusps of  $\Gamma_0(16)$

We see that  $z \in \mathcal{M}^\infty(16)$  with unique triple pole at infinity. As the  $\eta$ -quotient

$$s := \frac{\Pi_{q^4}}{\Pi_{q^8}} = \frac{\eta^6(8\tau)}{\eta^2(4\tau)\eta^4(16\tau)}$$

generates  $\mathcal{M}^\infty(16)$  with unique simple pole at infinity by Lemmas 2.3 and 2.4, we see from Lemma 2.2 that  $z$  is a cubic polynomial in  $s$ . We imply from the  $q$ -expansion

$$z = q^{-3} + 2q^{-2} + 4q^{-1} + 4 + 6q + 8q^2 + O(q^3)$$

that

$$(24) \quad z = s^3 + 2s^2 + 4s + 4,$$

and comparing (23) and (24) yields identity (8).

To prove identity (9), we need the following lemma concerning a modular function on  $\Gamma_0(16)$  that involves

$$s_3(\tau) := \frac{\Pi_{q^8}^2}{\Pi_{q^{16}}^2} = \frac{\eta^{12}(16\tau)}{\eta^4(8\tau)\eta^8(32\tau)}.$$

**Lemma 4.1.** *The function*

$$S := S(\tau) = s_3(\tau) + \frac{16}{s_3(\tau)}$$

lies on  $\mathcal{M}^\infty(16)$  with  $\text{ord}_\infty S = -4$ .

*Proof.* We know from Lemma 2.3 that  $s_3(\tau)$  is modular on  $\Gamma_0(32)$ . Since  $\Gamma_0(32)$  and  $\alpha := \begin{bmatrix} 1 & 0 \\ 16 & 1 \end{bmatrix}$  generate  $\Gamma_0(16)$ , we deduce that  $S(\tau) = s_3(\tau) + s_3(\alpha\tau)$  is modular on  $\Gamma_0(16)$ . By similar arguments as in the proof of (14) from Lemma 3.1, we get

$$s_3(\alpha\tau) = \frac{16}{s_3(\tau)}.$$

To find the orders of  $S(\tau)$  at the cusps of  $\Gamma_0(16)$ , we write  $S(\tau)$  as a generalized  $\eta$ -quotient

$$s_3(\tau) = \frac{\eta_{32,16}^4(\tau)}{\eta_{32,8}^4(\tau)}.$$

Following similar arguments as in the proof of (16) from Lemma 3.1 leads us to

$$(25) \quad \text{ord}_{a/c} s_3(\tau) = \frac{\gcd(c, 32)^2}{\gcd(c^2, 16)} \left[ P_2 \left( \frac{16a}{\gcd(c, 32)} \right) - P_2 \left( \frac{8a}{\gcd(c, 32)} \right) \right]$$

for each cusp  $a/c \in S_{\Gamma_0(16)}$ . Using (25), we record the above orders as shown in Table 6 below. We conclude from the table that  $S \in \mathcal{M}^\infty(16)$  with  $\text{ord}_\infty S = -4$ . □

cusps $r$	0	1/2	1/4, 3/4	1/8	$\infty$
$\text{ord}_r s_3(\tau)$	0	0	0	0	4
$\text{ord}_r 1/s_3(\tau)$	0	0	0	0	-4
$\text{ord}_r S(\tau)$	$\geq 0$	$\geq 0$	$\geq 0$	$\geq 0$	-4

TABLE 6. The orders of  $S(\tau)$  at the cusps of  $\Gamma_0(16)$

We are now in a position to prove (9). Setting  $a = q^2, q^4, q^6$  in (22) and combining the resulting identities yield

$$(26) \quad \sum_{n \geq 1} \left[ \frac{q^{2n}}{(1-q^{2n})^2} - \frac{q^{8n}}{(1-q^{8n})^2} \right] - 6 \sum_{n \geq 1} \frac{q^{16n-8}}{(1-q^{16n-8})^2} = \Pi_{q^8}^2 \left( s_4(\tau) + 1 + \frac{1}{s_4(\tau)} \right),$$

where

$$s_4(\tau) := \frac{\eta_{16,6}^2(\tau)}{\eta_{16,2}^2(\tau)}$$

is modular on  $\Gamma_1(16)$  by Lemma 2.6 and satisfies  $s_4(\gamma\tau) = 1/s_4(\tau)$  by Lemma 2.5. Thus,  $s_4(\tau) + 1/s_4(\tau)$  is modular on  $\Gamma_0(16)$ , and its orders at each cusps  $r \in S_{\Gamma_0(16)}$  can be obtained via Lemma 2.7. We tabulate these orders as shown in Table 7 below.

cusps $r$	0	1/2	1/4, 3/4	1/8	$\infty$
$\text{ord}_r s_4(\tau)$	0	0	0	0	-2
$\text{ord}_r 1/s_4(\tau)$	0	0	0	0	2
$\text{ord}_r s_4(\tau) + 1/s_4(\tau)$	$\geq 0$	$\geq 0$	$\geq 0$	$\geq 0$	-2

TABLE 7. The orders of  $s_4(\tau) + 1/s_4(\tau)$  at the cusps of  $\Gamma_0(16)$

We see that  $s_4(\tau) + 1/s_4(\tau) \in \mathcal{M}^\infty(16)$  with unique double pole at infinity, so that it is a quadratic polynomial in  $s$  according to Lemma 2.2. By looking at the  $q$ -expansion

$$s_4(\tau) + \frac{1}{s_4(\tau)} = q^{-2} + 2 + 4q^2 + O(q^6),$$

we infer that

$$s_4(\tau) + \frac{1}{s_4(\tau)} = s^2 + 2$$

and the right-hand side of (26) becomes

$$(27) \quad \sum_{n \geq 1} \left[ \frac{q^{2n}}{(1-q^{2n})^2} - \frac{q^{8n}}{(1-q^{8n})^2} \right] - 6 \sum_{n \geq 1} \frac{q^{16n-8}}{(1-q^{16n-8})^2} = \Pi_{q^8}^2 (s^2 + 3).$$

In view of (19), (23), and (24), we write (27) as

$$(28) \quad \frac{1}{\Pi_{q^8}^2} \sum_{n \geq 1} \left[ \frac{q^n}{(1-q^n)^2} - 8 \frac{q^{8n}}{(1-q^{8n})^2} \right] - s^3 - 3s^2 - 4s - 7 = \frac{7}{\Pi_{q^8}^2} \sum_{n \geq 1} \left[ \frac{q^{16n-8}}{(1-q^{16n-8})^2} - \frac{q^{16n}}{(1-q^{16n})^2} \right].$$

We now replace  $q$  with  $q^8$  in (1), put it on the right-hand side of (28), and then multiply both sides by 24, we get

$$(29) \quad \frac{1}{\Pi_{q^8}^2} \left[ 24 \left( \sum_{n \geq 1} \frac{q^n}{(1-q^n)^2} - 8 \sum_{n \geq 1} \frac{q^{8n}}{(1-q^{8n})^2} \right) + 7 \right] - 24s^3 - 72s^2 - 96s - 168 = 7S.$$

We know from Lemma 4.1 that  $S \in \mathcal{M}^\infty(16)$  with unique pole of order  $-4$  at infinity. Thus,  $S$  is a quartic polynomial in  $s$  by Lemma 2.2. In view of the  $q$ -expansion

$$S = q^{-4} + 20q^4 - 62q^{12} + O(q^{13}),$$

we get

$$S = s^4 - 8,$$

and (29) becomes

$$\frac{1}{\Pi_q^2 s} \left[ 24 \left( \sum_{n \geq 1} \frac{q^n}{(1 - q^n)^2} - 8 \sum_{n \geq 1} \frac{q^{8n}}{(1 - q^{8n})^2} \right) + 7 \right] = 7s^4 + 24s^3 + 72s^2 + 96s + 112,$$

which is identity (9). This completes the proof.  $\square$

#### REFERENCES

- [1] M. El Bachraoui, *On series identities of Gosper and integrals of Ramanujan theta function  $\psi(q)$* , Proc. Amer. Math. Soc. **147** (2019), 4451–4464.
- [2] W. N. Bailey, *Series of hypergeometric type which are infinite in both directions*, Q. J. Math. **os-7** (1936), 105–115.
- [3] ———, *A further note on two of Ramanujan’s formulae*, Q. J. Math. **3** (1952), 158–160.
- [4] B. C. Berndt, *Ramanujan’s Notebooks, Part III*, Springer-Verlag, New York, 1991.
- [5] B. Cho, J. K. Koo, and Y. K. Park, *Arithmetic of the Ramanujan-Göllnitz-Gordon continued fraction*, J. Number Theory **129** (2009), 922–947.
- [6] R. Wm. Gosper, *Experiments and discoveries in  $q$ -trigonometry*, Symbolic Computation, Number Theory, Special Functions, Physics and Combinatorics (F. G. Garvan and M. E. H. Ismail, eds.), Springer, New York, 2001, 79–103.
- [7] R. Guadalupe, *Gosper’s Lambert series identities of level 14*, preprint (2024), <https://doi.org/10.48550/arXiv.2412.18228>.
- [8] B. He, *Proofs for certain  $\Pi_q$  identities of Gosper*, J. Math. Anal. Appl. **492** (2020), paper no. 124486, 14 pp.
- [9] ———, *Proof for a  $\Pi_q$ -identity of Gosper*, Adv. in Appl. Math. **123** (2021), paper no. 102120, 9 pp.
- [10] K. Ono, *The Web of Modularity: Arithmetic of the Coefficients of Modular Forms and  $q$ -Series*, CBMS Regional Conference Series in Mathematics vol. 102, American Mathematical Society, Rhode Island, 2004.
- [11] P. Paule and C.-S. Radu, *Holonomic relations for modular functions and forms: first guess, then prove*, Int. J. Number Theory **17** (2021), 713–759.
- [12] C.-S. Radu, *An algorithmic approach to Ramanujan-Kolberg identities*, J. Symbolic Comput. **68** (2015), 225–253.
- [13] L. Wang, *Modular proofs of Gosper’s identities*, Adv. in Appl. Math. **135** (2022), paper no. 102312, 25 pp.
- [14] Y. Yang, *Transformation formulas for generalized Dedekind eta functions*, Bull. London Math. Soc. **36** (2004), 671–682.
- [15] M. V. Yathirajsharma, K. M. Harshitha, and K. R. Vasuki, *On Gosper’s  $\Pi_q$  and Lambert series identities*, Hiroshima Math. J. **52** (2022), 113–137.

INSTITUTE OF MATHEMATICS, UNIVERSITY OF THE PHILIPPINES, DILIMAN, QUEZON CITY 1101, PHILIPPINES  
*Email address:* rguadalupe@math.upd.edu.ph