Two-Sample Tests for Optimal Lifts, Manifold Stability and Reverse Labeling Reflection Shape

Do Tran Van^{*1}, Susovan Pal^{†2}, Benjamin Eltzner^{‡1}, and Stephan F. Huckemann^{§1}

¹Institute for Mathematical Stochastics, University of Göttingen, Germany ²Mathematics and Data Science Group, Vrije Universiteit Brussel, Belgium

March 25, 2025

Contents

1	Introduction	1
2	Setup	2
3	Manifold Stability and Optimal Lifts	4
4	Smoothness of Optimal Lifts	6
5	The Strong Law for Optimal Lifts	7
6	Almost Everywhere Continuity and Uniqueness of Optimal Lifts	11
7	Two-Sample Tests Using Optimal Lifts	11
8	Reverse Labeling Reflection Shape Spaces	14
9	Simulations and Application	17

Abstract

We consider a quotient of a complete Riemannian manifold modulo an isometrically and properly acting Lie group and lifts of the quotient to the manifolds in optimal position to a reference point on the manifold. With respect to the pushed forward Riemannian volume onto the quotient we derive continuity and uniqueness a.e. and smoothness to large extents also with respect to the reference point. In consequence we derive a general manifold stability theorem: the Fréchet mean lies in the highest dimensional stratum assumed with positive probability, and a strong law for optimal lifts. This allows to define new two-sample tests utilizing individual optimal lifts which outperform existing two-sample tests on simulated data. They also outperform existing tests on a newly derived reverse labeling reflection shape space, that is used to model filament data of microtubules within cells in a biological application.

1 Introduction

Statistical analysis of shape leads to the general task of performing statistics on data in a quotient, often a nonmanifold quotient Q = M/G of a complete Riemannian manifold M (of pre-shapes) modulo an isometric and proper action of a Lie group G (conveying shape equivalence). The very successful and highly popular *Procrustes analysis* introduced by Gower (1975) from Hurley and Cattell (1962) registers pre-shape data to the tangent space of a pre-shape of a Procrustes mean and can then apply a plethora of statistical methods from the toolbox of Euclidean statistics (Dryden

 $^{^*} do. tranvan @uni-goettingen. de$

[†]Susovan.Pal@vub.be

 $^{^{\}ddagger}$ beltzne@uni-goettingen.de

 $[\]label{eq:shuckeman@math.uni-goettingen.de} \$ huckeman@math.uni-goettingen.de$

and Mardia 2016). In the language of geometry this registration procedure amounts to *lifting* the quotient Q to the top manifold M in *optimal position* to some $p \in M$. For less concentrated data, due to the presence of curvature (shape spaces feature infinite positive curvatures near singularities, e.g. Kendall et al. (1999); Huckemann et al. (2010b)) decisive shape information can get lost, however.

In this contribution we aim at filtering out main effects of such curvatures caused by passing to the quotient. To this end, we first show that in general such lifts are measurable (Theorem 3.4) and almost everywhere unique and continuous (Theorem 6.1), even to large extents smooth, also with respect to their basepoints (Corollary 4.4). In passing, this allows to extend the *manifold stability theorem* from Huckemann (2012), now stating that in general a Fréchet mean is assumed on the manifold part of the quotient, whenever it is assumed with positive probability (Theorem 3.1). Hence, in combination with our smoothness Corollary 4.4, in general, for optimal lifts, central limit theorems on manifolds (ranging from Bhattacharya and Patrangenaru (2005) to Eltzner and Huckemann (2019) giving a rather general version including smeariness) can be applied.

From a.e. continuity of optimal lifts in the basepoint we derive a strong law of large numbers for optimal lifts (Theorem 5.1). This is required for our new two-sample tests introduced in Section 7 based on lifting each group separately, optimal with respect to optimally positioned lifts of their individual means, instead of lifting optimally with respect to a lift of the pooled mean (as is, e.g., done in Procrustes analysis). First simulations in Section 9 show that our new tests outperform classical tests.

In view of an application in cell structure biology we introduce in Section 8 a new landmark based shape space modulo similarity and reflection and modulo *reverse labeling* of landmarks, which is appropriate for modeling filament string-like shapes by placing landmarks on them. Thus in Section 9, our new tests can distinguish between two groups of cellular buckle structures which previous tests could not.

We note that the bias introduced by curvature due to passing to the quotient space has been corrected by Miolane et al. (2017).

Since our results build on a fair amount of geometry convolved with statistics, in the following section we recall a considerable body of foundations.

2 Setup

Throughout this contribution we consider the following setup (detailed in standard textbooks, for instance Bredon (1972); Lee (1997); Gray (2003); Bhattacharya and Bhattacharya (2012); Alexandrino and Bettiol (2015); Patrangenaru and Ellingson (2015); Durrett (2019):

1. M is a complete connected finite dimensional Riemannian manifold with countable atlas and intrinsic distance d, tangent space T_pM at $p \in M$, Riemannian inner product $\langle v, w \rangle$ and norm ||v|| for $v, w \in T_pM$ and Riemannian exponential $\exp_p : T_pM \to M$, which has a well defined inverse $\exp_p^{-1} : U \to T_pM$ for some open set $p \in U \subseteq M$, p in M. In particular, with the tangent bundle $TM = \bigcup_{p \in M} (\{p\} \times T_pM)$

$$\exp: TM \to M, \quad (p,v) \mapsto \exp_n v$$

is a smooth map. Here *smooth* means that derivatives of any order exist.

- 2. Notably, for an incomplete Riemannian manifold, the exponential is well defined and smooth in a neighborhood of the zero-section of the tangent bundle.
- 3. For a closed set $A \subset M$ we have the distance $d(p', A) := \min_{p \in A} d(p', p)$ of a point $p' \in M$ from A.
- 4. For a measurable set $A \subseteq M$ we have its (possibly infinite) Riemannian volume vol(A).
- 5. For $p \in M$ we have its *cut locus*

$$\begin{split} \mathrm{Cut}(p) &:= & \{ \exp_p(v) : v \in T_p M \text{ such that} \\ & \quad d(\exp_p(tv), p) = t \|v\| \text{ for all } 0 \leq t \leq 1 \text{ but} \\ & \quad d(\exp_p(tv), p) < t \|v\| \text{ for al } t > 1 \} \,. \end{split}$$

In order to use Theorem 3.6 by Le and Barden (2014) below, we assume that for every $p \in M$ and every $p' \in \operatorname{Cut}(p)$ there are $v_1, v_2 \in T_pM$ with $v_1 \neq v_2$ but $\exp_p(v_1) = p' = \exp_p(v_2)$. Manifolds where this is not the case are rather exotic. For an incomplete, geodesically convex (see below) Riemannian manifold, the cut locus of a point p is defined restricting \exp_p to the largest open set in the tangent space for which \exp_p is defined and smooth (Chavel 1995, p. 117/118).

- 6. G is a Lie group with unit element $e \in G$ acting on M via $G \times M \to M, (g, p) \mapsto g.p$, properly, i.e. preimages of compact sets under $G \times M \to M \times M, (g, p) \mapsto (g.p, p)$ are compact, and isometrically, i.e. d(p, p') = d(g.p, g.p') for all $p, p' \in M$ and $g \in G$.
- 7. Then the quotient $Q := M/G = \{[p] : p \in M\}$ with *orbit* $[p] := \{g.p : g \in M\}$ is Hausdorff when Q is equipped with the canonical quotient topology, i.e. the unique topology making the canonical projection $\pi : M \to Q, p \mapsto [p]$ continuous and open. Further, for every $p \in M$, $[p] \subset M$ is a closed embedded submanifold and

$$d_Q([p], [p']) = \min_{a \in G} d(p, g. p')$$

is the canonical quotients metric. We say that $p, p' \in M$ are in optimal position if $d(p, p') = d_Q([p], [p'])$ (Huckemann et al. 2010b).

For better clarity, from now on for $p \in M$, [p] denotes the corresponding subset of M which, as an element of Q is denoted by q such that $p \in [p] = \pi^{-1}(q)$.

8. $I_p = \{g \in G : g.p = p\}$ is the *isotropy group* at $p \in M$ and we assume that

$$M^* := \{ p \in M : I_p = \{e\} \} \neq \emptyset.$$

Then, the principal orbit theorem teaches that M^* is open and dense in M.

9. For $p \in M$ every tangent space decomposes into a vertical space along the orbit [p] and its orthogonal complement, the horizontal space at $p \in M$:

$$T_p M = T_p[p] \oplus H_p M$$

Further, again by the principal orbit theorem, $Q^* := M^*/G$ carries a canonical Riemannian quotient structure with $T_q Q^* \cong H_p M$ for every $p \in \pi^{-1}(q)$, $q \in Q^*$. For this reason, Q^* is called the *manifold part* of Q and $Q^0 = Q \setminus Q^*$ is the *singular part*, with preimage $M^0 = \pi^{-1}(Q^0)$.

10. In particular, with the Riemann exponential \exp^{Q^*} of Q^* and the identification $T_{\pi(p)}Q^* \cong H_p M$ for $p \in M^*$, we have that

$$\exp_p|_{H_pM} = \exp_{\pi(p)}^{Q^*}$$

whenever the r.h.s. is defined, as Q^* is not complete in general. While only Q is complete, Q^* is, however, geodesically convex, i.e. for any two $q, q' \in Q^*$ there is a shortest curve in Q joining the two and this curve lies fully in Q^* . In fact, with the parallel transport $\theta_{q,q'}: T_q Q^* \to T_{q'} Q^*$ along a length minimizing geodesic from q' to q (which is unique unless $q' \in \operatorname{Cut}(q)$) and $v \in T_q Q^*$ such that $q' = \exp_q^{Q^*} v$, we have

$$q = \exp_{q'}^{Q^*} \left(-\theta_{q,q'}(v)\right).$$

- 11. $X_1, \ldots, X_n \stackrel{\text{i.i.d.}}{\sim} X$ are random variables on M, i.e. they are Borel measurable mappings to M from a silently underlying probability space $(\Omega, \mathbb{P}, \mathcal{A})$ where we denote by $\mathbb{P}^X(B) = \mathbb{P}\{X \in B\}$, for Borel measurable $B \subseteq M$, the *push forward measure* on M. Here $\stackrel{\text{i.i.d.}}{\sim}$ stand for identically and independently distributed, i.e. $\mathbb{P}^{(X_1,\ldots,X_n)}(B_1 \times \ldots \times B_n) = \prod_{j=1}^n \mathbb{P}^{X_j}(B_j)$ for all Borel sets $B_j \in M$, $j = 1, \ldots, n$ and identically distributed, i.e. $\mathbb{P}^{X_j} = \mathbb{P}^X$ for all $j = 1, \ldots, n$.
- 12. These random variables give rise to their *population* and *sample Fréchet functions*, respectively

$$F^X(p) = \mathbb{E}[d(p,X)^2], \quad F_n^X(p) := \frac{1}{n} \sum_{j=1}^n d(p,X_j)^2, \quad p \in M$$

Every minimizer is called a *population* and *sample Fréchet mean*, respectively

$$\mu \in \underset{p \in M}{\operatorname{argmin}} F^X(p), \quad \mu_n \in \underset{p \in M}{\operatorname{argmin}} F^X_n(p).$$

13. The above $X_1, \ldots, X_n \stackrel{\text{i.i.d.}}{\sim} X$ project to random variables $\pi \circ X_1, \ldots, \pi \circ X_n \stackrel{\text{i.i.d.}}{\sim} \pi \circ X$ on Q with the corresponding Fréchet functions

$$F^{\pi \circ X}(a) = \mathbb{E}[d(q, \pi \circ X)^2], \quad F_n^{\pi \circ X}(q) := \frac{1}{n} \sum_{j=1}^n d(q, \pi \circ X_j)^2, \quad q \in Q$$

and Fréchet means

$$\nu \in \operatorname*{argmin}_{q \in Q} F^{\pi \circ X}(q), \quad \nu_n \in \operatorname*{argmin}_{q \in Q} F^{\circ X}_n(q).$$

- 14. Since every random variable Y on Q can be lifted to a random variable $X = \ell_p \circ Y$ as detailed in Theorem 3.4, we can equivalently do our analysis starting with random variables on M, or starting with random variables on Q.
- 15. We assume that $d_Q(q, \pi \circ X)$ is \mathbb{P} -integrable for some $q = q_0 \in Q$, and hence, by the triangle inequality, for any $q \in Q$. Then $\emptyset \neq \operatorname{argmin}_{q \in Q}(F^{\pi \circ X}(q) F^{\pi \circ X}(q_0))$, again by the triangle inequality, i.e., we assume that a population Fréchet mean exists.

3 Manifold Stability and Optimal Lifts

The following main result of this section states that every Fréchet mean lies on the manifold part, if the latter is assumed with positive probability. This strengthens Huckemann (2012, Corollary 1) who assumed that there were only countably many point masses on the singular part. The rest of this section is devoted to its proof and a few more convenient results.

Theorem 3.1 (Manifold Stability). Let $\nu \in \operatorname{argmin}_{q \in Q} F^{\pi \circ X}(q)$ and $\mathbb{P}\{\pi \circ X \in Q^*\} > 0$ then

$$\nu \in Q^*$$

Proof. This is assertion (v) of Lemma 3.7.

Now, we build the necessary machinery. For $p \in M$ let

$$L_p := \{ p' \in M : p' \text{ is in optimal position to } p \},$$

$$L'_p := \{ p' \in L_p : \text{ if } p'' \in [p'] \cap L_p \text{ then } p'' = p' \}.$$

Thus, L'_p denotes all points, uniquely in optimal position to p.

Remark 3.2. Note that for all $p, p' \in M$, due to the proper action, $L_p \cap [p'] \neq \emptyset$. Further, optimal positioning is symmetric, i.e.

$$p' \in L_p \Leftrightarrow p \in L_{p'}$$

Putting uniquely in optimal position, however, is not reflexive:

$$p \in M^*, p' \in L'_p \cap M^0 \Rightarrow p \notin L'_{p'}.$$

On the preimage of the manifold part, however, it is

$$p \in M, p' \in L'_p \cap M^* \Rightarrow p \in L'_{p'}.$$

Example 3.3. Consider on $M = \mathbb{R}^3$ the action of G of rotations about the last axis $N = \{(0,0,z)^T : z \in \mathbb{R}\}$ and the points

$$p_1 = \begin{pmatrix} 0\\0\\1 \end{pmatrix}, \quad p_2 = \begin{pmatrix} 1\\-1\\1 \end{pmatrix}, \quad p_3 = \begin{pmatrix} 2\\-2\\3 \end{pmatrix}, \quad p_4 = \begin{pmatrix} -2\\2\\3 \end{pmatrix}.$$

Then $p_2, p_3, p_4 \in L_{p_1}, p_1, p_3 \in L'_{p_2} \subset L_{p_2}$ but $p_2 \notin L'_{p_1} = N$.

Theorem 3.4 (Existence of optimal lifts). For every $p \in M$ there is a measurable mapping $\ell_p : Q \to M$ such that

- (i) $\pi \circ \ell_p = id_Q$ (lift),
- (ii) $\ell_p(q) \in L_p$ for all $q \in Q$ (optimal).

Proof. Let $p \in M$. By continuity, L_p is closed and hence so is $\emptyset \neq L_p \cap \pi^{-1}(q) \subseteq M$ for all $q \in Q$. Setting

$$\begin{split} \Psi: Q &\to \{ \emptyset \neq A \subset M : A \text{ closed} \} \\ q &\mapsto L_p \cap \pi^{-1}(q) \,, \end{split}$$

apply (Bogachev 2007, Theorem 6.9.3) stating that there exists a measurable map $\xi : Q \to M$ (*M* is called *X* in Bogachev (2007) and *Q* is called Ω there) with $\xi(q) \in \Psi(q)$, if

$$\{q \in Q : \Psi(q) \cap U \neq \emptyset\}$$
 is measurable for all open $U \subseteq M$.

Indeed, this condition is met, since for open $U \subseteq M$, $W := L_p \cap U$ is measurable in M, and hence, so is

$$\{q \in Q : \Psi(q) \cap U \neq \emptyset\} = \pi\{p' \in M : W \cap \pi^{-1}(\pi(p')) \neq \emptyset\} = \pi(W)$$

Thus $\ell_p := \xi$ is a measurable optimal lift.

Definition 3.5. Every such ℓ_p from Theorem 3.4 is called an optimal lift of Q through $p \in M$ (we drop the adjective 'measurable').

The following theorem states that Fréchet means are equilibrium points, i.e. classical means in the inverse exponential chart.

Theorem 3.6 (Le and Barden (2014), Corollaries 2 and 3). If $\mu \in \operatorname{argmin}_{p \in M} F^X(p)$ then

- (i) the domain $0 \in V \subset T_{\mu}M$ of \exp_{μ}^{-1} can be chosen such that $\mathbb{P}\{X \in \exp_{\mu}V\} = 1$ and
- (ii) $\int_M \exp^{-1}(p) d\mathbb{P}^X(p) = 0.$

We are now ready for the toolbox of this section.

Lemma 3.7. The following hold:

(i)
$$F^{\ell_p \circ \pi \circ X}(p) = F^{\pi \circ X}(\pi(p))$$
 for all $p \in M$

Moreover, if $\nu \in \operatorname{argmin}_{q \in Q} F^{\pi \circ X}(q)$, $\mu \in \pi^{-1}(\nu)$ and optimal lift ℓ_{μ} through μ , then

- (*ii*) $\mu \in \operatorname{argmin}_{p \in M} F^{\ell_{\mu} \circ \pi \circ X}(p)$
- (iii) if $p' \in \operatorname{argmin}_{p \in M} F^{\ell_{\mu} \circ \pi \circ X}(p)$ then $\pi(p') \in \operatorname{argmin}_{q \in Q} F^{\pi \circ X}(q)$
- (iv) if ℓ'_{μ} is another optimal lift through μ then $\ell'_{\mu} = \ell_{\mu}$ a.s.
- (v) if $\mathbb{P}\{\pi \circ X \in Q^*\} > 0$ then $\nu \in Q^*$

(vi) if
$$\{\nu\} = \operatorname{argmin}_{q \in Q} F^{\pi \circ X}(q)$$
 and $\mathbb{P}\{X \in L'_{\mu}\} > 0$ then $\{\mu\} = \operatorname{argmin}_{p \in M} F^{\ell_{\mu} \circ \pi \circ X}(p)$

Proof. (i) follows from $d(p, \ell_p(q')) = d_Q(q, q')$ for arbitrary $p \in M$ with $q = \pi(p)$ and from

$$F^{\ell_{\mu}\circ\pi\circ X}(p) = \int_{M} d(p,p')^{2} d\mathbb{P}^{\ell_{\mu}\circ\pi\circ X}(p')$$
$$= \int_{M} d(p,\ell_{p}(q'))^{2} d\mathbb{P}^{\pi\circ X}(q') = F^{\pi\circ X}(\pi(p)).$$

(ii) follows at once from (i).

(iii): if $p' \in \operatorname{argmin}_{p \in M} F^{\ell_{\mu} \circ \pi \circ X}(p)$ then, due to (ii) and twice (i),

$$F^{\ell_{\mu}\circ\pi\circ X}(p') = F^{\ell_{\mu}\circ\pi\circ X}(\mu) = F^{\pi\circ X}(\pi(\mu))$$

$$\leq F^{\pi\circ X}(\pi(p')) = F^{\ell_{p'}\circ\pi\circ X}(p')$$

$$= \int_{Q} d(p'\ell_{p'}(q))^2 d\mathbb{P}^{\pi\circ X}$$

$$\leq \int_{Q} d(p'\ell_{\mu}(q))^2 d\mathbb{P}^{\pi\circ X} = F^{\ell_{\mu}\circ\pi\circ X}(p')$$

Hence, all inequalities are equalities and thus $\pi(p') \in \operatorname{argmin}_{q \in Q} F^{\pi \circ X}(q)$.

(iv): Let ℓ_{μ} and ℓ'_{μ} be two optimal lifts through μ . Then, due to Theorem 3.6 and (ii),

$$\int_{Q} \exp_{\mu}^{-1} \circ \ell_{\mu}(q) \, d\mathbb{P}^{\pi \circ X}(q) = \int_{M} \exp_{\mu}^{-1}(p) \, d\mathbb{P}^{\ell_{\mu} \circ \pi \circ X}(p) = 0$$

$$= \int_{M} \exp_{\mu}^{-1}(p) \, d\mathbb{P}^{\ell'_{\mu} \circ \pi \circ X}(p) = \int_{Q} \exp_{\mu}^{-1} \circ \ell'_{\mu}(q) \, d\mathbb{P}^{\pi \circ X}(q). \quad (3.1)$$

Now, in order to see (iv), it suffices to show that

$$\int_{B} \exp_{\mu}^{-1} \circ \ell_{\mu}(q) \, d\mathbb{P}^{\pi \circ X}(q) = \int_{B} \exp_{\mu}^{-1} \circ \ell_{\mu}'(q) \, d\mathbb{P}^{\pi \circ X}(q) \tag{3.2}$$

for all Borel $B\subseteq Q$ and to this end, fix such a B and define

$$\ell''_{\mu}(q) = \begin{cases} \ell_{\mu}(q) & \text{if } q \in B \\ \ell'_{\mu}(q) & \text{if } q \in Q \setminus B \end{cases}$$

which is yet another optimal lift through μ . Then, due to (3.1),

$$\int_{Q\setminus B} \exp_{\mu}^{-1} \circ \ell_{\mu}(q) \, d\mathbb{P}^{\pi \circ X}(q) + \int_{B} \exp_{\mu}^{-1} \circ \ell_{\mu}(q) \, d\mathbb{P}^{\pi \circ X}(q) = \int_{Q} \exp_{\mu}^{-1} \circ \ell_{\mu}(q) \, d\mathbb{P}^{\pi \circ X}(q)$$
$$= \int_{B} \exp_{\mu}^{-1} \circ \ell_{\mu}'(q) \, d\mathbb{P}^{\pi \circ X}(q)$$
$$\int_{Q\setminus B} \exp_{\mu}^{-1} \circ \ell_{\mu}(q) \, d\mathbb{P}^{\pi \circ X}(q) + \int_{B} \exp_{\mu}^{-1} \circ \ell_{\mu}'(q) \, d\mathbb{P}^{\pi \circ X}(q) \, .$$

Subtracting $\int_{Q \setminus B} \exp_{\mu}^{-1} \circ \ell_{\mu}(q) d\mathbb{P}^{\pi \circ X}(q)$ from both sides yields the desired (3.2).

(v): Assume the contrary, namely that $\nu \in \operatorname{argmin}_{q \in Q} F^{\pi \circ X}(q)$ and $\mathbb{P}\{\pi \circ X \in Q^*\} > 0$ but $\nu \in Q \setminus Q^*$, i.e. $\mu \in M^0$ for any $\mu \in \pi^{-1}(\nu)$ and we show a contradiction. Then, there is $e \neq g \in G$ with $g.\mu = \mu$ but $g.p \neq p$ for all $p \in M^*$. Moreover, then, consider an optimal lift ℓ_{μ} through μ and note that $\ell'_{\mu} = g.\ell_{\mu}$, due to $d(\mu, g.\ell_{\mu}(p')) = d(g^{-1}.\mu, \ell_{\mu}(p')) = d(\mu, \ell_{\mu}(p))$ for all $p \in M$ by virtue of isometric action, is also an optimal lift through μ . Hence, by hypothesis and (iv),

$$\begin{array}{rcl} 0 &<& \mathbb{P}\{\pi \circ X \in Q^*\} &=& \mathbb{P}^X(M^*) \\ &\leq& \mathbb{P}^X\{p \in M : g.p \neq p\} &=& \mathbb{P}^{\pi \circ X}\{q \in Q : \ell_\mu(q) = \ell'_\mu(q)\} &=& 0\,, \end{array}$$

the desired contradiction.

=

(vi): Let $\{\nu\} = \operatorname{argmin}_{q \in Q} F^{\pi \circ X}(q)$ and $\mu, \mu' \in \operatorname{argmin}_{p \in M} F^{\ell_{\mu} \circ \pi \circ X}(p)$. Due to (iii) $\mu, \mu' \in \pi^{-1}(\nu)$, i.e. there is $g \in G$ with $\mu' = g.\mu$. Hence

$$\int_{M} d(\ell_{\mu}(p),\mu)^{2} d\mathbb{P}^{\ell_{\mu}\circ\pi\circ X}(p) = F^{\ell_{\mu}\circ\pi\circ X}(\mu) = F^{\ell_{\mu}\circ\pi\circ X}(g.\mu) = \int_{M} d(\ell_{\mu}(p),g.\mu)^{2} d\mathbb{P}^{\ell_{\mu}\circ\pi\circ X}(p).$$

Since $d(\ell_{\mu}(p),\mu) \leq d(\ell_{\mu}(p),g.\mu)$ for all $p \in M$, in consequence of $a^2 - b^2 = (a+b)(a-b)$ we have

$$\mathbb{P}^{\ell_{\mu} \circ \pi \circ X} \{ p \in \ell_{\mu}(Q) : d(p,\mu) = d(p,g.\mu) \} = 1 \,.$$

Thus, by hypothesis,

$$\mathbb{P}^{\ell_{\mu}\circ\pi\circ X}\left\{p\in\ell_{\mu}(Q)\cap L'_{\mu}:d(p,\mu)=d(p,g.\mu)\right\}>0\,,$$

yielding that there is $p \in \ell_{\mu}(Q) \cap L'_{\mu}$ with $d(p,\mu) = d(g^{-1}.p,\mu)$, i.e. $g^{-1}.p = p$, hence g = e and thus $\mu' = g.\mu = \mu$.

4 Smoothness of Optimal Lifts

Definition 4.1. Let $P \subset M$ be a closed submanifold with normal bundle NP, a subbundle of TM. Then the cut locus of P is defined as

$$\begin{split} \operatorname{Cut}(P) &:= & \{ \exp_p(v) : (p,v) \in NP \text{ such that} \\ & \quad d(\exp_p(tv),P) = t \|v\| \text{ for all } 0 \leq t \leq 1 \text{ but} \\ & \quad d(\exp_p(tv),P) < t \|v\| \text{ for all } t > 1 \} \,. \end{split}$$

Since $\pi^{-1}(q)$ is a closed submanifold of M for every $q \in Q$, with its normal bundle $N\pi^{-1}(q)$, we have a smooth mapping

$$\Phi_q: N\pi^{-1}(q) \to M \quad (p,v) \quad \mapsto \quad \exp_p v \,. \tag{4.1}$$

Note that $(p, v) \in N\pi^{-1}(q)$ if and only if $v \in H_p M$.

Theorem 4.2 (Basu and Prasad (2023), p. 4218). For every $q \in Q$ the inverse map Φ_q^{-1} is well defined and smooth on $M \setminus (\operatorname{Cut}(\pi^{-1}(q)) \cup \pi^{-1}(q))$.

Remark 4.3. Note that Φ_q^{-1} can be continuously extended to any $p \in \pi^{-1}(q)$) via $\Phi_q^{-1}(p) = (p, 0)$, since for $\pi^{-1}(q) \not\supseteq p'_n \to p \in \pi^{-1}(q)$ and $\Phi_q^{-1}(p'_n) = (p_n, v_n)$ with $d(p'_n, p_n) = ||v_n|| \le d(p'_n, p) \to 0$. In fact, this extension is smooth by Gray (2003, Lemma 2.3).

Corollary 4.4. Let $q \in Q$ and $p' \in M \setminus \operatorname{Cut}(\pi^{-1}(q))$ with $\Phi^{-1}(p') = (p, v)$. Then

$$p' \in L_p \text{ and } p \in L'_{p'}$$

Moreover, for every open $U \subseteq Q$ and $p' \in M \setminus C$ where $C := \bigcup_{q \in U} \operatorname{Cut}(\pi^{-1}(q))$, there is an optimal lift through p' of form

$$\ell_{p'}(q) = \exp_p v \text{ for } q \in U, \quad \text{where } \Phi_q^{-1}(p') = (p, v).$$

Further, this lift varies smoothly in $p' \in M \setminus C$ and also in $q \in U$ if $U \subseteq Q^*$ and $p' \in M^* \setminus C$.

Proof. By definition, if $p' = \exp_p v \in M \setminus (\operatorname{Cut}(\pi^{-1}(q)) \cup \pi^{-1}(q))$ with $\Phi^{-1}(p') = (p, v)$ then $d(p', \pi^{-1}(q)) = ||v|| = d(p', p)$ so that $p' \in L_p$. If also $g.p \in L_{p'}$ for $e \neq g \in G$ then there would be two geodesics from $\pi^{-1}(q)$ to p', a contradiction to $p' \notin \operatorname{Cut}(\pi^{-1}(q))$.

The second assertion follows at once from the first assertion by fixing $p' \in M \setminus C$, taking any optimal lift through p' and modifying it on the measurable set U.

For the last assertion, smoothness in $p' \in M \setminus C$ follows from Theorem 4.2 and Remark 4.3.

If $U \subseteq Q^*$, since Q^* is a manifold, due to geodesic convexity of Q^* , there is $v \in T_q M$, varying smoothly in $q \in Q^*$, such that $q' = \pi(p') = \exp_q^{Q^*} v$ and hence

$$q = \exp_{q'}^{Q^*}(-\theta_{q,q'}(v))$$

where $\theta_{q,q'}: T_q Q^* \to T_{q'} Q^*$ is the parallel transport which is smooth. Similarly, for $p \in L'_{p'}$ (see first assertion), $p' = \exp_p v$, where we have identified $H_p M \cong T_q Q^*$, i.e.

$$\ell_{p'}(q) = p = \exp_{p'}(-\theta_{q,q'}(v))$$

which varies smoothly in $q \in U \subseteq Q^*$.

5 The Strong Law for Optimal Lifts

Theorem 5.1 (The Strong Law for Optimal Lifts). Let X be absolutely continuous w.r.t. vol and with unique Fréchet mean

$$\{\nu\} = \underset{q \in Q}{\operatorname{argmin}} F^{\pi \circ X}(q) \text{ and measurable selection } \nu_n \in \underset{q \in Q}{\operatorname{argmin}} F_n^{\pi \circ X}(q)$$

Then, for arbitrary

$$\mu \in \pi^{-1}(\nu)$$
 and measurable selection $\mu_n \in \pi^{-1}(\nu_n) \cap L_{\mu}$

and arbitrary optimal lifts ℓ_{μ} and ℓ_{μ_n} through μ and μ_n , respectively, we have that

$$\ell_{\mu_n} \to \ell_{\mu}$$
 a.s.

in the sense that

$$\int_{Q} d(\ell_{\mu_n}(q), \ell_{\mu}(q)) \, d\mathbb{P}^{\pi \circ X}(q) \to 0 \quad \text{a.s.} \ .$$

Proof. Since \mathbb{P}^X is absolutely continuous w.r.t. to **vol**, we have $0 < \mathbb{P}\{X \in M^*\} \leq \mathbb{P}\{\pi \circ X \in Q^*\}$ and hence by manifold stability (Theorem 3.1) that $\nu \in Q^*$, i.e. $\mu \in M^* \subseteq F$ (*F* is introduced below in Definition 5.2 and only needed to apply Lemma 5.6 below), due to Theorem 5.7, and similarly $\nu_n \in Q^*$ a.s., i.e. $\mu_n \in M^* \subseteq F$ a.s.. Then, by (ii) and (iii) of Lemma 5.6,

$$d(\ell_{\mu_n}(q), \ell_{\mu}(q)) \to 0 \quad \text{for all} \quad q \in Q \quad \mathbb{P}^{\pi \circ X} \text{-a.e.}$$
(5.1)

holds a.s., since by Ziezold's strong law for Fréchet means on metric spaces (Ziezold 1977), $\nu_n \to \nu$ a.s. and hence $d(\mu_n, \mu) = d_Q(\nu_n, \nu) \to 0$ a.s..

Moreover, observe

$$\begin{aligned} d(\ell_{\mu_n}(q), \ell_{\mu}(q)) &\leq d(\ell_{\mu_n}(q), \mu_n) + d(\mu_n, \mu) + d(\mu, \ell_{\mu}(q)) \\ &= d_Q(q, \nu_n) + d(\mu_n, \mu) + d_Q(\nu, q) \\ &\leq 2d(\mu_n, \mu) + 2d_Q(\nu, q) \,. \end{aligned}$$

Since for fixed μ_n , the bottom row as a function of q is $\mathbb{P}^{\pi \circ X}$ integrable (by hypothesis a Fréchet mean exists), by dominated convergence, from (5.1) infer that indeed

$$\int_{Q} d(\ell_{\mu_n}(q), \ell_{\mu}(q)) d\mathbb{P}^{\pi \circ X}(q) \to 0 \quad \text{a.s.} .$$

Definition 5.2. For closed $S \subset M$ define

$$C_S := \{ p \in M : \text{ there are } S \ni s_1 \neq s_2 \in S \text{ such that } d(p, s_1) = d(p, S) = d(p, s_2) \}.$$

Further, we set for $p \in M$,

$$F_{p} := \bigcup_{p' \in L'_{p}} [p'] = \{ p' \in M : \text{ there is a uniuge } s' \in [p'] \text{ with } d(p,s) = d(p,[p']) \}$$

$$F := \{ p \in M : \mathbf{vol}(M \setminus F_{p}) = 0 \}$$

Lemma 5.3. Let $S \subset M$, then $\operatorname{vol}(C_S) = 0$.

Proof. Claim I: $f: M \to [0, \infty), p \mapsto d(p, S)$ is 1-Lipschitz.

Indeed, for $p_1, p_2 \in M$ with $s_1, s_2 \in S$ such that $d(p_i, S) = d(p_i, s_i)$, i = 1, 2, we have by the triangle inequality,

$$\begin{aligned} f(p_1) - f(p_2) &= d(p_1, s_1) - d(p_2, s_2) \leq d(p_1, s_2) - d(p_2, s_2) \leq d(p_1, p_2) \\ - (f(p_1) - f(p_2)) &= d(p_2, s_2) - d(p_1, s_1) \leq d(p_2, s_1) - d(p_1, s_1) \leq d(p_1, p_2) \,, \end{aligned}$$

yielding Claim I.

Claim II: $\operatorname{grad}_p f$ exists vol-a.e. for p with $\|\operatorname{grad}_p f\| \leq 1$ there.

In every local chart (U, \exp_p^{-1}) , the Riemannian volume pushes forward to a measure absolutely continuous with respect to Lebesgue measure in $\exp_p^{-1}(U) \subseteq T_p M$. Making U small enough, the Euclidean distance is bounded by a multiple of the intrinsic distance of preimages. Hence the intrinsic distance in U to $U \cap S$ is Lipschitz in terms of Euclidean distance, and thus by Rademacher's theorem (e.g. Heinonen (2005, Theorem 3.1)) the subset in $\exp_p^{-1}(U)$ where $x \mapsto$ $d(\exp_p x, S)$ is not differentiable, has Lebesgue measure zero. In consequence, $\operatorname{vol}(S_U) = 0$ for the subset S_U in $U \cap S$ where $p' \mapsto d(p', S)$ is not differentiable. Thus the first assertion of Claim II follows since M has a countable atlas.

For the second assertion, consider $p \in M$ where $\operatorname{grad}_p f$ is defined, and arbitrary $v \in T_p M$. Then, due to Claim I,

$$\begin{aligned} |\langle \mathbf{grad}_p f, v \rangle| &= \left| \frac{d}{dt} f(\exp_p(tv)) \right| &= \lim_{t \to 0} \frac{1}{t} \left| f(\exp_p(tv)) - f(p) \right| \\ &\leq \lim_{t \to 0} \frac{1}{t} d(\exp_p(tv), p) &= ||v||, \end{aligned}$$

yielding the second assertion.

Claim III: If $p \in C_S$ then f is not differentiable at p.



Figure 1: Depicting the counterexample from Remark 5.4: All points on the circular segment S are closest to any point in the lower right quadrant which is C_S in the incomplete M which is the plane minus a slit along the positive first axis.

For otherwise, if f was differentiable at $p \in C_S$, there would be $S \ni s_1 \neq s_2 \in S$ with $d(s_i, p) = f(p)$ for i = 1, 2. In particular, since M is complete, there would be two different unit speed geodesics $\gamma_i : t \mapsto \exp_p(tv_i)$ with $T_pM \ni v_1 \neq v_2 \in T_pM$, $||v_1|| = 1 = ||v_2||$ and $s_i = \exp_p(f(p)v_i)$ for i = 1, 2. Thus, in particular,

$$f(\gamma_i(t)) = d(\gamma_i(t), s_i) = f(p) - t$$
, for all $0 \le t \le f(p)$

In consequence

$$-1 = \frac{d}{dt}\Big|_{t=0} f(\gamma_i(t)) = \langle \operatorname{\mathbf{grad}}_p f, v_i \rangle, \text{ for } i = 1, 2,$$

which, in conjunction with the second part of Claim II ($\|\operatorname{\mathbf{grad}}_p f\| \leq 1$), yields that $\operatorname{\mathbf{grad}}_p f = -v_i$ for i = 1, 2, a contradiction.

Hence, due to Claim III, $C_S \subseteq \{p \in M : f \text{ is not differentiable at } p\}$, where the latter has zero volume, due to the first part of Claim II. Thus $\operatorname{vol}(C_S) = 0$ as asserted.

Remark 5.4. Note that the assertion of Lemma 5.3 can be wrong if M ceases to be complete (completeness was essential for proving Claim III above). For instance, with $M = \mathbb{R}^2 \setminus \{(x,0)^T : x > 0\}$ and $S = \{(\cos \phi, \sin \phi)^T : \phi \in [\pi/4, \pi/2]\}$ verify that $C_S = \{(x, y)^T \in M : x \ge 0 \ge y\}$ which has infinite volume, see Figure 1.

Lemma 5.5. The following hold:

- (i) Let $p, p' \in M$, then $p' \in F_p \Leftrightarrow p \in M \setminus C_{[p']}$,
- (ii) $\pi^{-1} \circ \pi(F_p) = F_p$ for all $p \in M$,
- (iii) $\operatorname{vol}(M \setminus F_p) = 0$ for $p \in M$ vol-a.e.,
- (iv) $\operatorname{vol}(M \setminus F) = 0.$

Proof. (i) follows directly from definition.

(i) The inclusion $F_p \subseteq \pi^{-1} \circ \pi(F_p)$. To see the reverse, let $p' \in \pi^{-1} \circ \pi(F_p)$. Then $\pi(p') \in \pi(F_p)$, and hence, there are $g, g' \in G$ with $g'.p' \in g.F_p$, i.e. $g^{-1}g'.p' \in F_p$. Hence, there is a unique $p'' \in [p']$ such that p'' is in optimal position to p, hence $p' \in F_p$, yielding the reverse inclusion $\pi^{-1} \circ \pi(F_p) \subseteq F_p$ and thus equality.

(iii): Since $[p'] \subset M$ is closed, $\operatorname{vol}(C_{[p']}) = 0$ by Lemma 5.3 for all $p' \in M$. Hence,

$$0 = \int_{M} \left(\int_{C_{[p']}} d\operatorname{vol}(p) \right) d\operatorname{vol}(p') = \int_{M} \left(\int_{M \setminus F_{p}} d\operatorname{vol}(p') \right) d\operatorname{vol}(p)$$

by Fubini and (i), yielding $\operatorname{vol}(M \setminus F_p) = 0$ for $p \in M$ vol-a.e. which is (iii).

(iv) follows in consequence of the above, as,

$$0 = \mathbf{vol}\{p \in M : \mathbf{vol}(M \setminus F_p) \neq 0\} = \mathbf{vol}(M \setminus F).$$

	_	

Lemma 5.6. Let $\operatorname{vol}_{\sharp} = \operatorname{vol} \circ \pi^{-1}$ be the pushforward of the Riemannian volume on M to Q. Then

(i) $\operatorname{vol}_{\sharp}(Q \setminus \pi(F_p)) = 0$ for all $p \in F$.

Further assuming that $F \ni p_n \to p \in F$ and setting $\tilde{Q} := \pi(F_p) \cap \bigcap_{n=1}^{\infty} \pi(F_{p_n})$, we have

- (*ii*) $\operatorname{vol}_{\sharp}(Q \setminus \tilde{Q}) = 0$,
- (iii) $\ell_{p_n}(q) \to \ell_p(q)$ for all $q \in \tilde{Q}$; here ℓ_{p_n} and ℓ_p are arbitrary optimal lifts through p_n and p_n respectively.

Proof. (i): From the definition and (ii) of Lemma 5.5 we have

$$vol_{\sharp}(Q \setminus \pi(F_p)) = \mathbf{vol}(M \setminus F_p).$$

where, by definition, the above r.h.s. is zero for all $p \in F$.

(ii): By de-Morgan's law:

$$vol_{\sharp}\left(Q\setminus \tilde{Q}\right) = \mathbf{vol}_{\sharp}\left(\left(Q\setminus \pi(F_p)\right)\cup \bigcup_{n=1}^{\infty}\left(Q\setminus \pi(F_{p_n})\right)\right),$$

whose r.h.s. is zero by (i) since $p, p_n \in F$ for all $n \in \mathbb{N}$.

(iii): Since

$$d(\ell_{p_n}(q), \ell_p(q)) \leq d(\ell_{p_n}(q), p_n) + d(p_n, p) + d(p, \ell_p(q))$$

= $d(\pi^{-1}(q), p_n) + d(p_n, p) + d(p, \pi^{-1}(q))$
 $\rightarrow 2d(p, \pi^{-1}(q)),$

observe that $\ell_{p_n}(q)$ is bounded. Since M is Heine-Borel (by hypothesis), $\ell_{p_n}(q)$ has a cluster point $p' \in \pi^{-1}(q)$. Now suppose that $p'' \in \pi^{-1}(q)$ is another cluster point of $\ell_{p_n}(q)$. Since $d(p_n, \ell_{p_n}(q)) = d(p_n, \pi^{-1}(q)) \to d(p, \pi^{-1}(q))$ we have thus

$$d(p, p') = d(p, \pi^{-1}(q)) = d(p, p'')$$

and from $q \in \tilde{Q}$ we know that $\pi^{-1}(q)$ has a unique point $\ell_p(q) = p' = p''$ in optimal position to p, hence, indeed

$$\ell_{p_n}(q) \to \ell_p(q)$$
.

Theorem 5.7. $F = M^*$.

Proof. In order to see $F \subseteq M^*$ we show $p \in M^0 \Rightarrow p \notin F$. To this end, let $p \in M^0$ and set $q = \pi(p)$. Then there is $e \neq g \in G$ such that g.p = p. Using the notation from Section 4, since M^* is open and dense, there is an open set $U \subset M^* \cap (M \setminus \operatorname{Cut}(\pi^{-1}(q)))$. Then for every $p' \in U$ there is $g' \in G$ such that the first component of $\Phi_q^{-1}(p')$ (Φ_q maps from the tangent bundle into the manifold, cf. (4.1) is $(g')^{-1}.p$, i.e. g'.p' is in optimal position to p. However, also $gg'.p' \neq g'.p'$ (since $p' \in M^*$) is in optimal position to p, yielding $U \cap F_p = \emptyset$. Since $\operatorname{vol}(U) > 0$, thus $p \notin F$.

(since $p' \in M^*$) is in optimal position to p, yielding $U \cap F_p = \emptyset$. Since $\operatorname{vol}(U) > 0$, thus $p \notin F$. $M^* \subseteq F$: Let $p \in M^*$ with $\pi(p) =: q \in Q^*$. If $p' \in M \setminus F_p$ is in optimal position to pthen there is also $p' \neq p'' \in \pi^{-1}(q')$, with $q' := \pi(p') \in Q$, in optimal position to p and hence $v', v'' \in H_p M$ with $v' \neq v''$ such that $p' = \exp_p v'$ and $p'' = \exp_p v''$ as shortest geodesics between points in optimal position are horizontal (Huckemann et al. 2010b, p 10/11). In consequence, since $p \in M^*$, these project to two different geodesics in Q from q to q'. Hence either $q' \in Q^0$, i.e. $p' \in M^0$ which has zero volume (due to Duistermaat and Kolk (2012, Proposition 2.7.1) every tube in M meets only finitely many singular orbits, and M, as a manifold, has a countable atlas) or $q' \in \operatorname{Cut}(q)$ (with respect to the manifold Q^*) which, as $\operatorname{vol}_{\sharp}$ is the Riemannian volume for Q^* , has $\operatorname{vol}_{\sharp}(\operatorname{Cut}(q)) = 0$ (see Chavel (1995, p. 123) and Itoh and Tanaka (2001)). In consequence $p' \in \pi^{-1}(\operatorname{Cut}(q))$ with $\operatorname{vol}(\pi^{-1}(\operatorname{Cut}(q))) = \operatorname{vol}_{\sharp}(\operatorname{Cut}(q)) = 0$. In summary $\operatorname{vol}(M \setminus F_p) = 0$ yielding $p \in F$.

In consequence of the above Theorem, we have the following extension of (vi) of Lemma 3.7.

Corollary 5.8. Let X be absolutely continuous w.r.t. vol and with unique Fréchet mean $\{\nu\} = \operatorname{argmin}_{q \in Q} F^{\pi \circ X}(q)$. If $\mathbb{P}\{X \in Q^*\} > 0$ and if $\mu \in \pi^{-1}(\nu)$ with optimal lift ℓ_{μ} through μ , then

$$\{\mu\} = \operatorname*{argmin}_{p \in M} F^{\ell_{\mu} \circ \pi \circ X}(p) \,.$$

Proof. From $\mathbb{P}\{X \in Q^*\} > 0$ infer with (v) of Lemma 3.7 that $\nu \in Q^*$, i.e $\mu \in M^* = F$ for every $\mu \in \pi^{-1}(\nu)$, due to Theorem 5.7. By definition of F, then $\operatorname{vol}\{p \in M : p \notin L'_{\mu}\} = 0$ and thus $\mathbb{P}\{X \in L'_{\mu}\} > 0$ by absolute continuity of X w.r.t. vol. The assertion now follows from (vi) of Lemma 3.7.

6 Almost Everywhere Continuity and Uniqueness of Optimal Lifts

Theorem 6.1. Let $p \in M^*$. Then there is a Borel measurable $Q_p \subset Q$ such that

- (i) $\operatorname{vol}\left(M \setminus \pi^{-1}(Q_p)\right) = 0,$
- (ii) $\ell_p = \ell'_p$ on Q_p for any two optimal lifts ℓ_p and ℓ'_p through p,
- (iii) $Q_p \to M$, $q \mapsto \ell_p(q)$ is continuous.

Proof. Setting $Q_p = \pi(F_p)$ with $\pi^{-1}(Q_p) = F_p$ (due to (ii) Lemma 5.5), this is a consequence of Lemma 6.2 below, since for $p \in F$, $0 = \operatorname{vol}(M \setminus F_p)$ by definition and $F = M^*$ by Theorem 5.7.

Lemma 6.2. Let $p \in M$ and $\pi(F_p) \ni q'_n \to q' \in \pi(F_p)$ with $p'_n \in L_p \cap \pi^{-1}(q'_n)$ and $p' \in L_p \cap \pi^{-1}(q')$. Then $p'_n \to p'$.

Proof. By definition of F_p , both p'_n and p' are uniquely in optimal position to p. With $q = \pi^{-1}(p)$ we have thus that

$$d(p'_n, p) = d_Q(q'_n, q) \to d_Q(q', q)$$

yielding that p'_n is bounded and hence it has a cluster point $p'' \in M$ which is in $\pi^{-1}(q')$ since the latter is closed. Hence d(p'', p) agrees with the limit $d_Q(q', q)$ of $d(p'_n, p)$, i.e. p'' is in optimal position to p. By uniqueness conclude p'' = p' yielding the assertion.

7 Two-Sample Tests Using Optimal Lifts

Here we are concerned with generalizations of the classical two-sample test for

$$X_1, \ldots, X_n \stackrel{\text{i.i.d.}}{\sim} X \text{ and } Y_1, \ldots, Y_m \stackrel{\text{i.i.d.}}{\sim} Y,$$

jointly independent on $M = \mathbb{R}^d$, $d \in \mathbb{N}$, with population and sample means

$$\bar{X}_n = \frac{1}{n} \sum_{j=1}^n X_j, \quad \mathbb{E}[X], \quad \bar{Y}_m = \frac{1}{m} \sum_{j=1}^m X_j, \quad \mathbb{E}[Y],$$

respectively, covariances

$$\mathbf{cov}_{n}[X] = \frac{1}{n} \sum_{j=1}^{n} (X_{j} - \bar{X}_{n})(X_{j} - \bar{X}_{n})^{T}, \quad \mathbf{cov}[X] = \mathbb{E}[(X - \mathbb{E}[X])(X - \mathbb{E}[X])^{T}],$$
$$\mathbf{cov}_{m}[Y] = \frac{1}{m} \sum_{j=1}^{m} (Y_{j} - \bar{Y}_{m})(Y_{j} - \bar{Y}_{m})^{T}, \quad \mathbf{cov}[Y] = \mathbb{E}[(Y - \mathbb{E}[Y])(Y - \mathbb{E}[Y])^{T}]$$

and pooled covariance

$$\mathbf{cov}_{n,m}[X,Y] = \frac{1}{n+m-2} \left(n \, \mathbf{cov}_n[X] + m \, \mathbf{cov}_m[Y] \right) \,.$$

Then, under $H_0: X \sim Y \sim \mathcal{N}(\mathbb{E}[X], \mathbf{cov}[X])$ (multivariate normal distribution),

$$\hat{T} := \frac{mn}{m+n} (\bar{X}_n - \bar{Y}_m)^T (\mathbf{cov}_{n,m}[X,Y])^{-1} (\bar{X}_n - \bar{Y}_m) \sim T^2_{d,m+n-2}$$
(7.1)

where $T_{d,k}^2$ denotes the well known Hotelling T^2 -distribution with (d,k) degrees of freedom, see, e.g. Mardia et al. (2024), if $\mathbf{cov}_{n,m}[X,Y]$ is of rank d. Here we have utilized the expected value $\mathbb{E}[f(X)] = \int_{\mathbb{R}^d} f(x) d\mathbb{P}^X(x)$ for \mathbb{P}^X -integrable functions. Further in case of arbitrary X, Y which are not necessarily normally distributed but feature second moments $\mathbb{E}[X^T X], \mathbb{E}[Y^T Y] < \infty$ with full rank covariances, robustness under nonnormality holds (Romano and Lehmann 2005, p. 207, 321), i.e.

$$\hat{T} \rightsquigarrow Z$$
 by which we mean $\frac{\hat{T}}{Z} \xrightarrow{\mathcal{D}} 1$

if $Z \sim T_{d,m+n-2}^2$ is independent and $n, m \to \infty$, under

$$H_0: \mathbb{E}[X] = \mathbb{E}[Y] \text{ and } \left(\frac{n}{m} \to 1 \text{ or } \mathbf{cov}[X] = \mathbf{cov}[Y]\right).$$

With the α -quantile $T_{d,m+n-2,\alpha}$ of the Hotelling T^2 -distribution with (d, m+n-2) degrees of freedom defined by $\mathbb{P}\{Z \leq T_{n,m,\alpha}\} = \alpha$ the Hotelling test rejects H_0 at nominal level $\alpha \in [0,1]$ if

$$\tilde{T} > T_{d,m+n-2,1-\alpha} \,.$$

Under the above assumptions, this test has asymptotically the true level α (Romano and Lehmann 2005).

In the following we introduce four different tests for data on a quotient space Q. For shape spaces (Section 8 introduces some classical shape spaces and a new one), the first one gives, among others, the classical two-sample test based on *Procrustes analysis* (Dryden and Mardia 2016) or for *intrinsic MANOVA* (Huckemann et al. 2010a). The other two are new and build on the strong law for optimal lifts (Theorem 5.1) and their local smoothness (Corollary 4.4). For all three tests we have the following setup.

Assumption 7.1. We assume that

$$W_1, \ldots, W_n \overset{\text{i.i.d.}}{\sim} W \text{ and } Z_1, \ldots, Z_m \overset{\text{i.i.d.}}{\sim} Z,$$

are jointly independent on Q, dim $(Q^*) = d$ with population Fréchet means ν^W and ν^Z , respectively, and measurable selections ν_n^W and ν_m^Z of sample Fréchet means, respectively. Further, let $\mu_W \in \pi^{-1}(\nu_W)$ with optimal lift ℓ_{μ_W} through μ_W and $\mu_Z \in \pi^{-1}(\nu_Z)$ with optimal lift ℓ_{μ_Z} through μ_Z . We assume that W, Z are absolutely continuously distributed with respect to $\pi \circ \text{vol}$.

Moreover, we assume that either $n/m \to 1$ or that $\mathbf{cov}[X] = \mathbf{cov}[\theta_{\mu_W,\mu_Z} \circ Y]$ where

$$X := (\exp_{\mu_W})^{-1} \circ \ell_{\mu_W} \circ W,$$

$$Y := (\exp_{\mu_Z})^{-1} \circ \ell_{\mu_Z} \circ Z$$

and θ_{μ_W,μ_Z} is the parallel transport along a length minimizing geodesic from μ_Z to μ_W .

Under these assumptions,

- 1. by Theorem 3.6, X and Y are a.s. well defined,
- 2. $\nu^W, \nu^Z \in Q^*$ by the manifold stability Theorem 3.1,

and in the following, we test for

$$H_0: \nu^W = \nu^Z$$

at a given level $\alpha \in [0, 1]$.

Test 7.2 (Pooled Lifting). Let $\nu_{n,m}^{W,Z}$ be a measurable selection of the pooled sample Fréchet means of $W_1, \ldots, W_n, Z_1, \ldots, Z_m$, let $\mu_{n,m}^{W,Z} \in \pi^{-1}(\nu_{n,m}^{W,Z})$ be a measurable selection and let $\ell_{\mu_{n,m}^{W,Z}}$ be an optimal lift through $\mu_{n,m}^{W,Z}$. Setting

$$\begin{split} X_j &:= (\exp_{\mu_{n,m}^{W,Z}})^{-1} \circ \ell_{\mu_{n,m}^{W,Z}} \circ W_j, \quad 1 \le j \le n, \\ Y_j &:= (\exp_{\mu_{n,m}^{W,Z}})^{-1} \circ \ell_{\mu_{n,m}^{W,Z}} \circ Z_j, \quad 1 \le j \le m \end{split}$$

and $T_{pooled} := \hat{T}$ from (7.1), reject H_0 if $T_{pooled} > T_{n,m,1-\alpha}$.

In this test the sample covariance matrices have been computed using averages in the tangent spaces. Alternatively, these averages can be replaced in an intrinsic fashion by preimages of sample Fréchet means. Test 7.3 (Pooled Lifting Intrinsically). With the notation from Test 7.2, setting

$$\bar{X}_{n}^{(i)} := (\exp_{\mu_{n,m}^{W,Z}})^{-1} \circ \ell_{\mu_{n,m}^{W,Z}} \circ \nu_{n}^{W}, \qquad \bar{Y}_{m}^{(i)} := (\exp_{\mu_{n,m}^{W,Z}})^{-1} \circ \ell_{\mu_{n,m}^{W,Z}} \circ \nu_{m}^{Z}$$

define the intrinsic sample covariances and corresponding intrinsic test statistic

$$\begin{aligned} \mathbf{cov}_{n}^{(i)}[X] &:= \frac{1}{n} \sum_{j=1}^{n} (X_{j} - \bar{X}_{n}^{(i)}) (X_{j} - \bar{X}_{n}^{(i)})^{T}, \\ \mathbf{cov}_{m}^{(i)}[Y] &:= \frac{1}{m} \sum_{j=1}^{m} (Y_{j} - \bar{Y}_{m}^{(i)}) (Y_{j} - \bar{Y}_{m}^{(i)})^{T}, \\ \mathbf{cov}_{n,m}^{(i)}[X,Y] &:= \frac{1}{n+m-2} \left(n \operatorname{cov}_{n}^{(i)}[X] + m \operatorname{cov}_{m}^{(i)}[Y] \right), \\ \hat{T}^{(i)} &:= \frac{mn}{m+n} (\bar{X}_{n}^{(i)} - \bar{Y}_{m}^{(i)})^{T} \left(\operatorname{cov}_{n,m}^{(i)}[X,Y] \right)^{-1} (\bar{X}_{n}^{(i)} - \bar{Y}_{m}^{(i)}) \end{aligned}$$
(7.2)

Then, setting $T_{pooledI} := \hat{T}^{(i)}$ from (7.2), reject H_0 if $T_{pooledI} > T_{d,m+n-2,1-\alpha}$.

Alternatively we can lift each sample separately.

Test 7.4 (Individual Lifting). With the notation from Test 7.2 let $\mu_n^W \in \pi^{-1}(\nu_n^W) \cap L_{\mu_{n,m}^{W,Z}}$, $\mu_n^Z \in \pi^{-1}(\nu_n^Z) \cap L_{\mu_{n,m}^{W,Z}}$, i.e. both are a.s. uniquely (due to Theorems 3.1 and 5.7) in optimal position to $\mu_{n,m}^{W,Z}$ and let $\ell_{\mu_n^W}$ be an optimal lift through μ_n^W , $\ell_{\mu_m^Z}$ be an optimal lift through μ_m^Z . Then setting

$$\bar{X}_{n}^{(i)} := (\exp_{\mu_{n}^{W}})^{-1} \circ \ell_{\mu_{n}^{W}} \circ \nu_{n}^{W}, \qquad \bar{Y}_{m}^{(i)} := (\exp_{\mu_{m}^{Z}})^{-1} \circ \ell_{\mu_{m}^{Z}} \circ \nu_{m}^{Z},$$

as well as,

$$\begin{aligned} X_j &:= (\exp_{\mu_n^W})^{-1} \circ \ell_{\mu_n^W} \circ W_j, \quad 1 \le j \le n, \\ Y_j &:= (\exp_{\mu_m^Z})^{-1} \circ \ell_{\mu_m^Z} \circ Z_j, \quad 1 \le j \le m \end{aligned}$$

and $T_{individual} := \hat{T}^{(i)}$ from (7.2), reject H_0 if $T_{individual} > T_{d,n+m-2,1-\alpha}$.

If it is computationally too expensive to compute the pooled sample mean, in particular in view of the bootstrap tests introduced further below, instead optimal position with respect to the other sample's mean.

Test 7.5 (Individual Asymmetric Lifting). Measurably select $\mu_n^W \in \pi^{-1}(\nu_n^W)$ and $\mu_n^Z \in \pi^{-1}(\nu_n^Z) \cap L_{\mu_n^W}$, i.e. both are a.s. uniquely (due to Theorems 3.1 and 5.7) in optimal position to one another and define all other quantities as in Test 7.4. With those compute $T_{individualA} := \hat{T}^{(i)}$ from (7.1). Then reject H_0 if $T_{individualA} > T_{d,n+m-2,,1-\alpha}$.

Remark 7.6. As detailed above and in Huckemann and Eltzner (2020), if the Hessian of the Fréchet function $q \mapsto F^{\ell_{p'} \circ W}(q)$ in the exponential chart has all eigenvalues equal to 2 (as is the case if Q is Euclidean) under H_0 at $q = \mu^W = \mu^Z$ the various tests have asymptotic level α for the various choices of $p' = \nu^W \in \pi^{-1}(\mu^W)$. The smaller the eigenvalues, the more drastic the deviation (Hundrieser et al. 2024b), and in case of zeroes, a phenomenon called (directional) smeariness kicks in (Eltzner and Huckemann 2019; Tran et al. 2021), making the tests completely inapplicable.

For this reason, Hundrieser et al. (2024a) propose to use bootstrap versions of the above quantile based tests, namely simulating quantiles via suitably resampling, as follows. Here is the generic procedure from Eltzner and Huckemann (2017) applicable to each of the three latter tests. Taking averages in tangent spaces instead of intrinsic sample means, the thus modified bootstrap version is also applicable to the first Test 7.2. In fact, the following test, inspired by the test from Welch (1947), also works well in case of different covariances.

Test 7.7 (Bootstrap Version for Tests 7.3, 7.4 and 7.5). In a first round of bootstrapping from W_1, \ldots, W_n obtain $\mathbf{cov}_n^{*,(i)}[W]$ and of bootstrapping from Z_1, \ldots, Z_m obtain $\mathbf{cov}_m^{*,(i)}[Z]$. Then set

$$C := \mathbf{cov}_n^{*,(i)}[W] + \mathbf{cov}_m^{*,(i)}[Z]$$

In a second independent round of bootstrapping $B \in \mathbb{R}$ (large) times from W_1, \ldots, W_n and from Z_1, \ldots, Z_m obtain

 $d^{*,X,b} := \bar{X}_n^{*,b} - \bar{X}_n^{(i)}, \quad d^{*,Y,b} := \bar{Y}_m^{*,b} - \bar{X}_m^{(i)}, \quad b = 1, \dots, B.$

From these obtain the bootstrap α -quantile $T^*_{m,n,\alpha}$ of the data

$$(d^{*,X,b} - d^{*,Y,b})^T C^{-1} (d^{*,X,b} - d^{*,Y,b})$$
.

Then, reject H_0 if

$$\hat{T}^{(i,*)} := (\bar{X}_n^{(i)} - \bar{Y}_m^{(i)})^T C^{-1} (\bar{X}_n^{(i)} - \bar{Y}_m^{(i)}) > T_{d,m+n-2,\alpha}^* \,.$$

For data on the circle Hundrieser et al. (2024a) have shown that the corresponding tests keep the level α asymptotically. For general (quotient) spaces this is open, but simulations indicate that the levels are asymptotically kept. We illustrate some simulations in Section 9.

We note that Bhattacharya and Patrangenaru (2005) have been the first to introduce intrinsic two-sample tests for manifold data. One of their test is based on quantiles of a χ^2 -distribution and the other is a bootstrap alternative (see also Bhattacharya and Lin (2017)).

8 Reverse Labeling Reflection Shape Spaces

Classical shape spaces and reflection shape spaces (Dryden and Mardia 2016) model, modulo certain group actions, landmark configurations $A = (a_1, \ldots, a_k) \in \mathbb{R}^{m \times k}$, giving the position of $3 \leq k \in \mathbb{N}$ landmark vectors $a_1, \ldots, a_k \in \mathbb{R}^m$ of dimension $m \in \mathbb{N}$, where we assume that $2 \leq m < k$. One basic group action is that of translations and rotations (Euclidean motions), given by

 $SO(m) \rtimes \mathbb{R}^m \text{ acting on } M = \mathbb{R}^{m \times k} \text{ via } A \stackrel{(g,a)}{\to} gA + a\mathbf{1}_k^T \quad (g,a) \in SO(m) \rtimes \mathbb{R}^m \,.$

Here, with the standard unit vectors $e_1, \ldots, e_k \in \mathbb{R}^m$, $1_k := e_1 + \ldots + e_k \in \mathbb{R}^k$ denotes the vector with all entries equal to one and the (semidirect) product of $(g, a), (h, b) \in SO(m) \rtimes \mathbb{R}^m$ is given by (gh, b + ga). Letting $I_k = (e_1, \ldots, e_k) \in \mathbb{R}^{k \times k}$ denote the unit matrix, there are two popular ways to filter out translation:

(i) Centering by subtracting the mean landmark: $\mathbb{R}^{m \times k} \to \mathcal{C}_m^k$, $A \mapsto A(I_k - \frac{1}{k} \mathbf{1}_k \mathbf{1}_k^T)$, here

$$\mathcal{C}_m^k := \{ B \in \mathbb{R}^{m \times k} : B1_k = 0 \}$$

denotes the *centered configurations*.

(ii) Helmertizing: $\mathbb{R}^{m \times k} \to \mathbb{R}^{m \times (k-1)}, A \mapsto AH_k$ where $H_k \in \mathbb{R}^{k \times (k-1)}$ is chosen such that $(H_k|\frac{1}{\sqrt{k}}1_k) \in SO(k)$. Usually H_k is chosen as the *sub Helmert matrix*:

$$H_k = \begin{pmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{6}} & \cdots & \frac{1}{\sqrt{k(k-1)}} \\ -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{6}} & \cdots & \frac{1}{\sqrt{k(k-1)}} \\ 0 & -\frac{2}{\sqrt{6}} & \cdots & \frac{1}{\sqrt{k(k-1)}} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & -\frac{k-1}{\sqrt{k(k-1)}} \end{pmatrix}.$$

As usual, $\mathbb{R}^{m \times k}$ is equipped with the standard Euclidean inner product

$$\langle A, B \rangle := \mathbf{tr}(A^T B), \quad A, B \in \mathbb{R}^{m \times k}$$

yielding the Frobenius norm

$$\|A\| := \sqrt{\mathbf{tr}(A^T A)}$$

where \mathbf{tr} denotes the matrix trace. Since the action of scaling

$$\mathbb{R}_+$$
 acting on $\mathbb{R}^{m \times k}$ via $A \xrightarrow{\lambda} \lambda A, \quad \lambda \in \mathbb{R}_+$

is neither isometric nor proper (the only closed orbit is that of $0 \in \mathbb{R}^{m \times k}$), representatives on the unit hypersphere

$$\mathbb{S}^{m \times k-1} := \{ B \in \mathbb{R}^{m \times l} : \|B\| = 1 \}$$

are chosen. Thus, taking away the problematic diagonal orbit originating from single point configurations,

$$\mathcal{D}_m^k := \left\{ A = (a, \dots, a) \in \mathbb{R}^{m \times k} : a \in \mathbb{R}^m \right\},\$$

filtering out translation and scaling, one arrives at

$$\mathbb{R}^{m \times k} \setminus \mathcal{D}_m^k \to \left\{ \begin{array}{cc} \mathcal{C}_m^k \setminus \{0\} & \text{centering} \\ \mathbb{R}^{m \times (k-1)} \setminus \{0\} & \text{Helmertizing} \end{array} \right\} \xrightarrow{B \mapsto \frac{B}{\|B\|}} \left\{ \begin{array}{cc} \mathbb{S}^{m \times k-1} \cap \mathcal{C}_m^k & \text{centering} \\ \mathbb{S}^{m \times (k-1)-1} & \text{Helmertizing} \end{array} \right\}$$

Usually, $\mathbb{S}^{m \times (k-1)-1}$ is called the *pre-shape sphere* and we also call

$$\mathcal{S}_m^k := \mathbb{S}^{m \times k-1} \cap \mathcal{C}_m^k$$

the centered pre-shape sphere.

In the last step, rotations are filtered out leading to the classical shape space

 Σ_m^k viewed as either $\mathcal{S}_m^k/SO(m)$ or $\mathbb{S}^{m \times (k-1)-1}/SO(m)$,

or orthogonal transformations are filtered out leading to the classical reflection shape space

 $\mathcal{R}\Sigma_m^k$ viewed as either $\mathcal{S}_m^k/O(m)$ or $\mathbb{S}^{m \times (k-1)-1}/O(m)$.

Here SO(m) and O(m) act, as above, via multiplication from the left.

Notably, for $A, B \in \mathbb{S}^{m \times (k-1)-1}$ their spherical distance is $d(A, B) = \arccos \operatorname{tr}(A^T B)$ and hence,

$$\min_{R \in O(m)} d(A, RB) = \arccos\max_{R \in O(m)} \operatorname{tr}(A^T RB) = \arccos\operatorname{tr}\Lambda$$

where $BA^T = U\Lambda V^T$ is a singular value decomposition. In consequence the maximal possible distance in $\mathcal{R}\Sigma_m^k$ is $\pi/2$ and a similar argument yields the same maximal distance in Σ_m^k .

Planar shape and planar reflection shape (see additionally Huckemann and Ziezold (2006); Huckemann and Hotz (2009)) take advantage from identifying $A = (a_1, \ldots, a_k) \in \mathbb{R}^{2 \times k}$ with the complex row vector $z = (z_1, \ldots, z_k) \in \mathbb{C}^k$, where $a_j = (\operatorname{Re}(z_j), \operatorname{Im}(z_j))^T$, $j = 1, \ldots, k$. Rotation by the angle $\phi \in [0, 2\pi)$ corresponds to complex multiplication from the left by $e^{i\phi}$. In consequence, Σ_2^k carries a canonical Riemannian manifold structure of the complex projective space $\mathbb{C}P^{k-2}$ of complex dimension k-2 (real dimension 2k-4).

Reflection then corresponds to joint complex conjugation:

$$R.z = R.(z_1, \ldots, z_k) := (\bar{z}_1, \ldots, \bar{z}_k) = \bar{z}$$

and hence $\mathcal{R}\Sigma_2^k$ also has a singular stratum comprising all shapes

$$[z] = \{e^{i\phi}z : 0 \le \phi < 2\pi\} \text{ where } z \in \{(z_1, \dots, z_k) \in \mathbb{S}^{m \times k-1} \cap \mathcal{C}_m^k : z_j \in \mathbb{R} \text{ for all } 1 \le j \le k\}.$$

In particular Σ_2^3 is a sphere with radius 1/2 and $\mathcal{R}\Sigma_2^3$ can be viewed as its closed upper hemisphere. This is due to the explicit Hopf fibration (the Riemannian isometry is achieved by halving the radius \mathbb{S}^2):

$$\mathbb{S}^{3} \to \mathbb{S}^{3}/\mathbb{S}^{1} \cong \mathbb{S}^{2}, \quad (z_{1}, z_{2}) \mapsto \begin{pmatrix} 2\operatorname{\mathbf{Re}}(z_{1}\bar{z}_{2}) \\ 2\operatorname{\mathbf{Im}}(z_{1}\bar{z}_{2}) \\ |z_{1}|^{2} - |z_{2}|^{2} \end{pmatrix} = \begin{pmatrix} 2\operatorname{\mathbf{Re}}(z_{1}\bar{z}_{2}) \\ 2\operatorname{\mathbf{Im}}(z_{1}\bar{z}_{2}) \\ 2|z_{1}|^{2} - 1 \end{pmatrix}.$$

Further, for $z, w \in \mathbb{S}^{2 \times k-1} \subset \mathbb{C}^k$ with $z \mathbf{1}_k = 0 = w \mathbf{1}_k$ we have

$$d_{\Sigma_2^k}([z]_{\Sigma_2^k}, [w]_{\Sigma_2^k}) = \arccos |z\bar{w}^T|$$

$$d_{\mathcal{R}\Sigma_2^k}([z]_{\mathcal{R}\Sigma_2^k}, [w]_{\mathcal{R}\Sigma_2^k}) = \min \{\arccos |zw^T|, \arccos |z\bar{w}^T|\}.$$

The new reverse labeling reflection shape spaces are required in some applications, for instance when studying the shape of planar of one-dimensional non oriented string structures using landmark configurations, a frequent task in (micro)biology. Then, their shapes are preserved when placing landmarks from one end to the other, also in reversed order. To this end, introduce the *reverse labeling* action

$$\mathbb{R}^{m \times k} \to \mathbb{R}^{m \times k}, \quad A \mapsto AL_k \text{ with } L_k := (e_k, \dots, e_1) = \begin{pmatrix} 0 & \dots & 1 \\ \vdots & \ddots & \vdots \\ 1 & \dots & 0 \end{pmatrix} \in \mathbb{R}^{k \times k}.$$

Notably this action commutes with centering and with the action of O(m). Since $L_k^2 = I_k$, setting $G = \{L_k, I_k\}$, we have the following.

Definition 8.1. The reverse labeling reflection shape space for m-dimensional k-landmark configurations is given by

$$\mathcal{RR}\Sigma_m^k := \mathcal{R}\Sigma_m^k/G := \left(\mathcal{S}_m^k/G\right)/O(m).$$

Reverse labeling is obviously isometric but it does not commute with Helmertizing since $H_k^T L_k H_k \neq L_{k-1}$:

Lemma 8.2. For $H_k \in \mathbb{R}^{k \times k-1}$ with $(H_k | \frac{1}{\sqrt{k}} 1_k) \in SO(k)$, L_k as above and arbitrary $A \in \mathbb{R}^{m \times k}$ we have

$$AL_kH_k = AH_k(H_k^TL_kH_k)$$
, where $H_k^TL_kH_k \in O(k-1)$.

Proof. The first equality follows from $H_k H_k^T L_k = (I_k - \frac{1}{k} \mathbf{1}_k \mathbf{1}_k^T) L_k = L_k - \frac{1}{k} \mathbf{1}_k \mathbf{1}_k^T$ and the fact that $\mathbf{1}_k^T H_k = 0$ by construction. For the second assertion consider

$$(H_k^T L_k H_k)^T H_k^T L_k H_k = H_k^T L_k H_k H_k^T L_k H = H_k^T L_k (L_k - \frac{1}{k} \mathbf{1}_k \mathbf{1}_k^T) H_k = I_{k-1}$$

since $L_k^2 = I_k$ and $H_k^T H_k = I_{k-1}$.

In consequence, for reverse labeling reflection shape we only use the centered pre-shape sphere S_m^k as above, on which G acts from the right and O(m) from the left. Thus, for $A, B \in S_m^k$, we have

$$d_{\mathcal{RR}\Sigma_m^k}([A]_{\mathcal{RR}\Sigma_m^k}, [B]_{\mathcal{RR}\Sigma_m^k}) = \min\{\arccos \operatorname{tr} \Lambda, \arccos \operatorname{tr} \Lambda'\}$$

where $BA^T = U\Lambda V^T$ and $BL_k A^T = U'\Lambda'(V')^T$ are singular value decompositions.

For planar reverse labeling reflection shape in complex representation, for $z = (z_1, \ldots, z_k) \in S_2^k$, we set $\tilde{z} := (z_k, \ldots, z_1) = zL_k$, so that

$$d_{\mathcal{RR}\Sigma_2^k}([z]_{\mathcal{RR}\Sigma_2^k}, [w]_{\mathcal{RR}\Sigma_2^k}) = \min\{\arccos|zw^T|, \arccos|z\bar{w}^T|, \arccos|\tilde{z}w^T|, \arccos|\tilde{z}\bar{w}^T|\}.$$

Lemma 8.3. $\mathcal{RR}\Sigma_2^3$ can be viewed as a closed third of the closed hemisphere $\mathcal{R}\Sigma_2^3$ in the sphere Σ_2^3 . In particular, the maximal distance in $\mathcal{RR}\Sigma_2^3$ is $\pi/6$.

Proof. For $z = (z_1, z_2, z_3) \in \mathcal{S}_2^3$ let $w = (w_1, w_2) = zH_3$. Then, using the Helmert submatrix,

$$(w_1, w_2)H_3^T L_3 H_3 = (w_1, w_2)\frac{1}{2} \begin{pmatrix} 1 & -\sqrt{3} \\ -\sqrt{3} & -1 \end{pmatrix} = -\frac{1}{2} \begin{pmatrix} w_2 \sqrt{3} - w_1, w_2 + w_1 \sqrt{3} \end{pmatrix},$$

and

$$\begin{pmatrix} w_2 \sqrt{3} - w_1 \end{pmatrix} \left(\overline{w_2 + w_1 \sqrt{3}} \right) = -\sqrt{3} (|w_1|^2 - |w_2|^2) - w_1 \overline{w}_2 + 3w_2 \overline{w}_1 = -\sqrt{3} (|w_1|^2 - |w_2|^2) - 2 \operatorname{\mathbf{Re}}(w_1 \overline{w}_2) - i4 \operatorname{\mathbf{Im}}(w_1 \overline{w}_2), \\ w_2 \sqrt{3} - w_1 \Big|^2 - \left| \overline{w_2 + w_1 \sqrt{3}} \right|^2 = 2 (|w_1|^2 - |w_2|^2) - 4\sqrt{3} \operatorname{\mathbf{Re}}(w_1 \overline{w}_2).$$

Hence, under reverse labeling,

$$\begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 2\operatorname{\mathbf{Re}}(w_1\bar{w}_2) \\ 2\operatorname{\mathbf{Im}}(w_1\bar{w}_2) \\ |w_1|^2 - |w_2|^2 \end{pmatrix}$$

is mapped to

$$\begin{pmatrix} \frac{1}{2} \operatorname{\mathbf{Re}} \left(\left(w_2 \sqrt{3} - w_1 \right) \left(\overline{w_2 + w_1 \sqrt{3}} \right) \right) \\ \frac{1}{2} \operatorname{\mathbf{Im}} \left(\left(w_2 \sqrt{3} - w_1 \right) \left(\overline{w_2 + w_1 \sqrt{3}} \right) \right) \\ \frac{1}{4} \left(\left| w_2 \sqrt{3} - w_1 \right|^2 - \left| \overline{w_2 + w_1 \sqrt{3}} \right|^2 \right) \end{pmatrix} = -\frac{1}{2} \begin{pmatrix} x_1 + \sqrt{3} x_3 \\ 2x_2 \\ \sqrt{3} x_1 - x_3 \end{pmatrix},$$

i.e. accounting for a rotation by $\pi/3$ in the (x_1, x_3) -plane. Since reflection is filtered out by assuming $x_2 \ge 0$ and taking into account that in order to achieve an isometry the radius of the unit sphere needs to be halved, the maximal distance in $\mathcal{RR}\Sigma_2^3$ is indeed $\frac{1}{2}\frac{\pi}{3}$.

Remark 8.4. We anticipate that the diameter of $\mathcal{RR}\Sigma_2^k$ increases with k.

9 Simulations and Application

Simulations on Σ_2^5 . Here we consider two different planar configurations of five landmarks, of same shape for H_0 and of shape distance 0.06 for an alternative H_1 , each with random independent identical multinormal isotropic noise on every landmark with standard deviation of 0.2. In Table 1 we compare the various two-samples tests at nominal level $\alpha = 0.05$ for various sample sizes with one another and record the empirical level under H_0 and the power under H_1 ; we also include the bootstrap test T_J proposed by Preston and Wood (2010), originally for three-dimensional shape data. In terms of keeping the level and maximizing power, from inspection of Table 1, the tests based on individual lifting tend to outperform the others tests.

Method	od T		T'		Τ _ο		T ₁		T ₂	
	Null	Power	Null	Power	Null	Power	Null	Power	Null	Power
N1 = N2										
15 vs 15	0.01	0.04	0.005	0.02	0.015	0.055	0.01	0.04	0.015	0.055
30 vs 30	0.04	0.145	0.03	0.125	0.035	0.19	0.04	0.165	0.03	0.21
50 vs 50	0.045	0.335	0.03	0.34	0.05	0.43	0.045	0.35	0.04	0.505
100 vs 100	0.05	0.6	0.045	0.605	0.05	0.67	0.052	0.62	0.045	0.72
N1≠ N2										
15 vs 30	0.03	0.11	0.02	0.095	0.025	0.14	0.035	0.11	0.035	0.135
30 vs 60	0.04	0.315	0.04	0.25	0.04	0.355	0.05	0.36	0.04	0.42
50 vs 100	0.045	0.50	0.052	0.51	0.05	0.55	0.048	0.54	0.048	0.6

Table 1: Comparing true levels (columns labeled Null) at nominal level 0.05 and powers (columns labeled Power) for the bootstrap versions of the various tests from Section 7: T stands for "pooled lifting", T' for T_J from Preston and Wood (2010), T_0 for "pooled lifting intrinsically", T_1 for "individual lifting" and T_2 for "individual asymmetric lifting". The numbers are averages of rejections of H_0 of 1000 tests each. The first column gives the two samples sizes, which are equal for rows 4 - 7 (alleviating asymptotic normality) and different for rows 8 - 10.

In a second simulation, we consider the same configuration data, now in reverse labeling reflection shape space $\mathcal{RR}\Sigma_2^5$. We perform the same tests as above and now record the results in Table 2. While in terms of keeping the level and maximizing power, the tests based on individual lifting still tend to outperform the others tests, this effect is weaker than for shape spaces above. Figure 2 gives more detail by depicting the corresponding power curves. Here, the picture is clearer, the bootstrap tests "pooled lifting" and T_J have least power, the tests "pooled lifting individually" and "individual lifting" have higher power, the test "individual asymmetric lifting" has largest power. Notably their true levels are rather comparable (see Table 2).

Method	Т		T'		T ₀		T ₁		T ₂	
	Null	Power	Null	Power	Null	Power	Null	Power	Null	Power
N1 = N2										
15 vs 15	0.035	0.085	0.015	0.065	0.04	0.12	0.035	0.18	0.03	0.30
30 vs 30	0.04	0.175	0.05	0.155	0.035	0.265	0.35	0.415	0.04	0.475
50 vs 50	0.046	0.31	0.048	0.26	0.046	0.39	0.048	0.60	0.04	0.61
N1≠ N2										
15 vs 30	0.042	0.24	0.03	0.235	0.044	0.305	0.05	0.42	0.04	0.455
30 vs 60	0.046	0.345	0.046	0.355	0.052	0.415	0.048	0.53	0.046	0.58

Table 2: Depicting true levels and powers as in Table 1 for the same tests of the same data, but now for reverse labeling reflection shape in $\mathcal{RR}\Sigma_2^5$. Again the numbers are averages of 1000 tests each.



Figure 2: Power curves for the data and five tests (labeling explained in Table 1. The values at 0.06 correspond to the ones recorded under power in Table 2. Top row for equal sample sizes (rows 4 - 7 in Table 2) and bottom row for different sample sizes (rows 8 - 10 in Table 2). In contrast to Table 2 each test has been repeated 500 times. The horizontal axis records the reverse relabeling reflection shape distance between the corresponding two population Fréchet means and the vertical axis depicts the percentage of rejections.

Application to the interaction of microtubules and intermediate filaments in biological cells. In the analysis of biological cells filament structures, buckles of *microtubules* play an important role, for example Nolting et al. (2014). In particular, intermediate filaments (IF) are important components which provide mechanical stability in cells and tissues. A disruption of the IF or of their connection with other cell structures cause different degenerative diseases of skin, muscle, and neurons, see, for example Lodish et al. (2008). Most notably, even though tumor cells

lose their normal appearance, they retain several expressions of particular IF proteins. Thus, a method to identify the IF proteins in tumor cells could help to develop the most effective treatment to destroy the tumor (Lodish et al. 2008).



Figure 3: Determining 5 landmarks on a typical microtubules filament at mathematically characteristic locations for buckle structure. Landmarks 1 and 5 maximize the ratio between Euclidean distance and curve length between them times the fourth root of the maximal normal distance along the buckle, which gives the 3rd landmark. Landmark 2 is furthest from this normal line (on the side of Landmark 1). Landmark 4 maximizes normal distance from the line connecting Landmarks 5 and 3. Afterwards, all landmarks (2,3,4) have been shifted minimally to make them more equidistant.

With the theory and methods presented in this paper, we were able to shed light on developing a new method to distinguish cells with and without IFs.

Using the *filament sensor* from Eltzner et al. (2015) (see also Hauke et al. (2023)), from cells provided by the Cellular Biophysics Research Group of the Institute for X-ray Physics at the University of Göttingen, we have extracted planar microtubules buckle structures and placed 5 mathematically defined landmarks on them (Figure 3), leading to two groups. The first group contains 65 buckles from cells with intermediate filaments (generating stiffness) and the second group contains 29 buckles from cells without intermediate filaments. Because the buckles' shape are considered to be the same under reflection and traversing in reverse orientation, the two groups of buckles are modeled in the reverse labeling reflection shape space $\mathcal{RR}\Sigma_2^5$. Figure 4 shows typical buckle structures.



Figure 4: Typical microtubules buckle structures discretized by 5 landmarks. Upper row: without intermediate (vimentin) filaments. Lower row: in the presence of intermediate (vimentin) filaments (generating stiffness).

Applying the two-sample test procedures from Section 7 to the data we are able to distinguish the two groups at level $\alpha = 0.05$. In fact, from the *p*-values in Table 3 we can see that only the test "individual lifting" (T_1) and "individual asymmetric lifting" (T_2) detect the difference, while the other two tests "pooled lifting" (T) and "pooled lifting intrinsically" (T_0) do not. Notably, after Bonferroni correction for four tests, only the test "individual asymmetric lifting" (T_1) detects the difference. For convenience, Table 3 also records the *p*-value of test T_J from Preston and Wood (2010).

	Т	T_0	T_1	T_2	T_J
<i>p</i> -value	0.08	0.052	0.011	0.027	0.081

Table 3: Results from different two-sample tests for the two groups (with and without IFs) of reverse relabeling reflection buckles. Test abbreviations as in Table 1.

Acknowledgments

All four authors are very grateful to Alexander Lytchak for very helpful comments in differential geometry. Also, all authors thank Sarah Köster for providing the microtubules data. DTV and SP gratefully acknowledge funding by DFG HU 1575/7. SP acknowledges support by the Research Foundation – Flanders (FWO) via the Odysseus II programme no. G0DBZ23N. BE acknowledges funding by DFG SFB 1456.

References

- Alexandrino, M. M. and R. G. Bettiol (2015). Lie groups and geometric aspects of isometric actions, Volume 82. Springer.
- Basu, S. and S. Prasad (2023). A connection between cut locus, thom space and morse-bott functions. Algebraic & Geometric Topology 23(9), 4185–4233.
- Bhattacharya, A. and R. Bhattacharya (2012). Nonparametric inference on manifolds: with applications to shape spaces, Volume 2. Cambridge University Press.
- Bhattacharya, R. and L. Lin (2017). Omnibus CLTs for Fréchet means and nonparametric inference on non-Euclidean spaces. *Proceedings of the American Mathematical Society* 145(1), 413–428.
- Bhattacharya, R. N. and V. Patrangenaru (2005). Large sample theory of intrinsic and extrinsic sample means on manifolds II. *The Annals of Statistics* 33(3), 1225–1259.
- Bogachev, V. I. (2007). Measure theory, Volume 2. Springer Science & Business Media.
- Bredon, G. E. (1972). Introduction to Compact Transformation Groups, Volume 46 of Pure and Applied Mathematics. New York: Academic Press.
- Chavel, I. (1995). *Riemannian geometry: a modern introduction*. Number 108. Cambridge university press.
- Dryden, I. L. and K. V. Mardia (2016). Statistical Shape Analysis (2nd ed.). Chichester: Wiley.
- Duistermaat, J. J. and J. A. Kolk (2012). Lie groups. Springer Science & Business Media.
- Durrett, R. (2019). Probability: theory and examples, Volume 49. Cambridge university press.
- Eltzner, B. and S. Huckemann (2017). Bootstrapping descriptors for non-Euclidean data. In Geometric Science of Information 2017 proceedings, pp. 12–19. Springer.
- Eltzner, B. and S. F. Huckemann (2019). A smeary central limit theorem for manifolds with application to high-dimensional spheres. *Ann. Statist.* 47(6), 3360–3381.
- Eltzner, B., C. Wollnik, C. Gottschlich, S. Huckemann, and F. Rehfeldt (2015). The filament sensor for near real-time detection of cytoskeletal fiber structures. *PloS one* 10(5), e0126346.
- Gower, J. C. (1975). Generalized Procrustes analysis. *Psychometrika* 40, 33–51.

Gray, A. (2003). Tubes, Volume 221. Springer Science & Business Media.

- Hauke, L., A. Primeßnig, B. Eltzner, J. Radwitz, S. F. Huckemann, and F. Rehfeldt (2023). Filamentsensor 2.0: An open-source modular toolbox for 2d/3d cytoskeletal filament tracking. *Plos one* 18(2), e0279336.
- Heinonen, J. (2005). Lectures on Lipschitz analysis. Number 100. University of Jyväskylä.
- Huckemann, S. (2012). On the meaning of mean shape: Manifold stability, locus and the two sample test. Annals of the Institute of Statistical Mathematics 64(6), 1227–1259.
- Huckemann, S. and T. Hotz (2009). Principal components geodesics for planar shape. Journal of Multivariate Analysis (100), 699–714.
- Huckemann, S., T. Hotz, and A. Munk (2010a). Intrinsic MANOVA for Riemannian manifolds with an application to Kendall's space of planar shapes. *IEEE Transactions on Pattern Analysis* and Machine Intelligence 32(4), 593–603.
- Huckemann, S., T. Hotz, and A. Munk (2010b). Intrinsic shape analysis: Geodesic principal component analysis for Riemannian manifolds modulo Lie group actions (with discussion). *Statistica Sinica* 20(1), 1–100.
- Huckemann, S. and H. Ziezold (2006). Principal component analysis for Riemannian manifolds with an application to triangular shape spaces. *Advances of Applied Probability (SGSA)* 38(2), 299–319.
- Huckemann, S. F. and B. Eltzner (2020). Statistical methods generalizing principal component analysis to non-Euclidean spaces. In *Handbook of Variational Methods for Nonlinear Geometric* Data, pp. 317–388. Springer.
- Hundrieser, S., B. Eltzner, and S. Huckemann (2024a). Finite sample smeariness of Fréchet means with application to climate. *Electron. J. Statist.* 18(2), 3274–3309.
- Hundrieser, S., B. Eltzner, and S. F. Huckemann (2024b). A lower bound for estimating Fréchet means. arXiv preprint arXiv:2402.12290.
- Hurley, J. R. and R. B. Cattell (1962). The procrustes program: Producing direct rotation to test a hypothesized factor structure. *Behavioral Science* 7, 258–262.
- Itoh, J.-i. and M. Tanaka (2001). The lipschitz continuity of the distance function to the cut locus. Transactions of the American Mathematical Society 353(1), 21–40.
- Kendall, D. G., D. Barden, T. K. Carne, and H. Le (1999). Shape and Shape Theory. Chichester: Wiley.
- Le, H. and D. Barden (2014). On the measure of the cut locus of a Fréchet mean. Bulletin of the London Mathematical Society 46(4), 698–708.
- Lee, J. M. (1997). Riemannian manifolds: an introduction to curvature, Volume 176. Springer.
- Lodish, H., A. Berk, C. A. Kaiser, C. Kaiser, M. Krieger, M. P. Scott, A. Bretscher, H. Ploegh, P. Matsudaira, et al. (2008). *Molecular cell biology*. Macmillan.
- Mardia, K. V., J. T. Kent, and C. C. Taylor (2024). Multivariate analysis. John Wiley & Sons.
- Miolane, N., S. Holmes, and X. Pennec (2017). Template shape estimation: correcting an asymptotic bias. SIAM Journal on Imaging Sciences 10(2), 808–844.
- Nolting, J.-F., W. Möbius, and S. Köster (2014). Mechanics of individual keratin bundles in living cells. *Biophysical journal* 107(11), 2693–2699.
- Patrangenaru, V. and L. Ellingson (2015). Nonparametric statistics on manifolds and their applications to object data analysis. CRC Press.
- Preston, S. P. and A. T. Wood (2010). Two-sample bootstrap hypothesis tests for threedimensional labelled landmark data. *Scandinavian Journal of Statistics* 37(4), 568–587.

Romano, J. P. and E. L. Lehmann (2005). Testing statistical hypotheses. Springer, Berlin.

Tran, D., B. Eltzner, and S. F. Huckemann (2021). Smeariness begets finite sample smeariness. In Geometric Science of Information 2021 proceedings, pp. 29–36. Springer.

- Welch, B. L. (1947). The generalization of 'student's'problem when several different population varlances are involved. *Biometrika* 34 (1-2), 28–35.
- Ziezold, H. (1977). Expected figures and a strong law of large numbers for random elements in quasi-metric spaces. Transaction of the 7th Prague Conference on Information Theory, Statistical Decision Function and Random Processes A, 591–602.