

ON CLOSED EMBEDDINGS IN $\mathbb{P}^{\mathbb{N}} \cup \mathbb{Q}^{\mathbb{N}}$

ELŻBIETA POL, ROMAN POL AND MIROŚLAWA REŃSKA

ABSTRACT. Let \mathbb{P} and \mathbb{Q} be the irrationals and the rationals in $[0, 1]$. We prove that if a separable metrizable X is a union of two disjoint 0-dimensional sets E, F , E is absolutely G_{δ} and F is absolutely $F_{\sigma\delta}$, then there is a closed embedding $h : X \rightarrow \mathbb{P}^{\mathbb{N}} \cup \mathbb{Q}^{\mathbb{N}}$ with $E = h^{-1}(\mathbb{P}^{\mathbb{N}})$, $F = h^{-1}(\mathbb{Q}^{\mathbb{N}})$.

We prove also that, for $H = \{x \in I^{\mathbb{N}} : \forall k \exists l x(2^k 3^l) = 0\}$, whenever A is an $F_{\sigma\delta}$ -set in a compact one-dimensional metrizable space X , there is an embedding $h : X \rightarrow (\mathbb{P} \cup \{0\})^{\mathbb{N}} \cup \mathbb{Q}^{\mathbb{N}}$ such that $A = h^{-1}(H)$.

1. INTRODUCTION

We denote by \mathbb{P} and \mathbb{Q} the irrationals and the rationals in $I = [0, 1]$, respectively, and $\mathbb{P}^{\mathbb{N}}, \mathbb{Q}^{\mathbb{N}}$ are the countable products, the subspaces of the Hilbert cube $I^{\mathbb{N}}$.

All our spaces are metrizable separable, and our terminology follows [En] and [Ku] .

In particular, X is absolutely G_{δ} (absolutely $F_{\sigma\delta}$) if for some, equivalently - for any, embedding of X in a completely metrizable space Y , the image of X in Y is a G_{δ} -set ($F_{\sigma\delta}$ -set, respectively); absolutely F_{σ} -sets are exactly the σ -compact sets, cf. [En], 4.5.7.

The space $\mathbb{P}^{\mathbb{N}}$ is absolutely G_{δ} and $\mathbb{Q}^{\mathbb{N}}$ is absolutely $F_{\sigma\delta}$, cf. Comment 4.1.

The following result was proved in [PPR], Theorem 1,1.

Theorem 1.1. *Let $\Gamma = \{x \in \mathbb{Q}^{\mathbb{N}} : x(i) \in \{0, 1\} \text{ for all but finitely many } i\}$. Then for any disjoint zero-dimensional sets G, S in a compact space X , where G is absolutely G_{δ} and S is σ -compact, there is an embedding $h : X \rightarrow I^{\mathbb{N}}$ such that $G = h^{-1}(\mathbb{P}^{\mathbb{N}})$ and $S = h^{-1}(\Gamma)$.*

In this note we shall derive from Theorem 1.1 the following result.

Theorem 1.2. *Let X be a union of two disjoint zero-dimensional sets E, F , where E is absolutely G_{δ} and F is absolutely $F_{\sigma\delta}$. Then there is an*

2010 *Mathematics Subject Classification.* Primary: 54B10, 54C25, 54D40; Secondary: 54F50.

Key words and phrases. one-dimensional spaces, universal spaces, $F_{\sigma\delta}$ -sets, closed embeddings.

embedding $h : X \rightarrow \mathbb{P}^{\mathbb{N}} \cup \mathbb{Q}^{\mathbb{N}}$ onto a closed subspace such that $E = h^{-1}(\mathbb{P}^{\mathbb{N}})$ and $F = h^{-1}(\mathbb{Q}^{\mathbb{N}})$.

In particular, $\mathbb{P}^{\mathbb{N}} \cup \mathbb{Q}^{\mathbb{N}}$, which is a union of a zero-dimensional absolutely G_{δ} -set and a zero-dimensional absolutely $F_{\sigma\delta}$ -set, contains topologically closed copies of every such space.

We shall prove also another result concerning embeddings of one-dimensional absolutely $F_{\sigma\delta}$ -spaces, using some key elements of the reasoning in [PPR], combined with Theorem 1.2.

Before we state this result, let us explain some background. Let

$$(1) H_{kl} = \{x \in I^{\mathbb{N}} : x(2^k 3^l) = 0\}, H = \bigcap_k \bigcup_l H_{kl}.$$

Then H is a $F_{\sigma\delta}$ -set in $I^{\mathbb{N}}$ and, by some standard arguments, one checks that for any $F_{\sigma\delta}$ -set A in a compact space X , there is an embedding $h : X \rightarrow I^{\mathbb{N}}$ with $A = h^{-1}(H)$, cf. [Si].

If X is a compact zero-dimensional space and A is a $F_{\sigma\delta}$ -set in X , there is an embedding $h : X \rightarrow \{0, 1, \frac{1}{2}, \frac{1}{3}, \dots\}^{\mathbb{N}}$ with $A = h^{-1}(H)$, cf. [Ke], sec. 23.A.

We shall obtain a counterpart of these results for one-dimensional spaces.

Theorem 1.3. *For each $F_{\sigma\delta}$ -set A in a compact one-dimensional space X , there is an embedding $h : X \rightarrow (\mathbb{P} \cup \{0\})^{\mathbb{N}} \cup \mathbb{Q}^{\mathbb{N}}$ such that $A = h^{-1}(H)$, cf. (1).*

In particular, $((\mathbb{P} \cup \{0\})^{\mathbb{N}} \cup \mathbb{Q}^{\mathbb{N}}) \cap H$ is a one-dimensional absolutely $F_{\sigma\delta}$ -set containing closed topological copies of any absolutely $F_{\sigma\delta}$ -set of dimension ≤ 1 , cf. Comment 4.1.

2. PROOF OF THEOREM 1.2.

Proof. Let $X = E \cup F$, where E, F are disjoint zero-dimensional sets, E is absolutely G_{δ} and F is absolutely $F_{\sigma\delta}$. Using [Ku], vol. II, §45, VII, Theorem 5, one can extend X to a compact space X^* with $\dim X^* = \dim X \leq 1$ such that F can be extended to a σ -compact zero-dimensional set \tilde{F} in X^* , disjoint from E .

Indeed, F being a F_{σ} -set in X , $F = \bigcup_n F_n$, where F_n is closed in X , and the cited theorem in [Ku] provides a compact extension X^* of X with $\dim X^* = \dim X$ such that for the closure F_n^* of each F_n in X^* , $\dim F_n^* = \dim F_n = 0$.

By absoluteness, E is a G_δ -set in X^* , and hence $\tilde{F} = \bigcup_n F_n^* \setminus E$ has required properties.

Furthermore, we can enlarge the set \tilde{F} to a σ -compact zero-dimensional set C ,

$$(1) \tilde{F} \subset C \subset X^* \setminus E,$$

such that there are G_δ -sets G_1, G_2, \dots in X^* with

$$(2) X^* \setminus C = \bigcup_i G_i, \dim(E \cup G_i) = 0.$$

To get C and G_i , we start from picking a base B_1, B_2, \dots of X^* with boundaries $\text{Fr}B_i$ disjoint from the zero-dimensional set E . For $G_1 = X^* \setminus (\tilde{F} \cup \bigcup_i \text{Fr}B_i)$, we have $E \subset G_1$ and $\dim G_1 = 0$. Since $\dim \text{Fr}B_i \leq 1$, one can choose for each i a zero-dimensional σ -compact set $C_i \subset \text{Fr}B_i$ with $\dim(\text{Fr}B_i \setminus C_i) \leq 0$, and let $C = \tilde{F} \cup \bigcup_i C_i$, $G_{i+1} = \text{Fr}B_i \setminus C$, $i = 1, 2, \dots$

We have then (1), the first part of (2), and since $E \subset G_1$ and G_i is relatively closed in $E \cup G_i$ for $i \geq 2$, we get also the second part of (2).

Having defined C , we shall recall that, by absoluteness, F is a $F_{\sigma\delta}$ -set in C , and therefore, there are σ -compact sets L_k such that

$$(3) L_1 = C \supset L_2 \supset L_3 \supset \dots, F = \bigcap_k L_k,$$

and let

$$(4) L_k = \bigcup_i L_{ki}, L_{ki} \text{ compact.}$$

Let $\phi : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$ be a bijection and, cf. (4), (3),

$$(5) D_{\phi(k,i)} = L_{ki} \setminus L_{k+1}$$

The set L_{ki} being compact and disjoint from E , cf. (1), $D_{\phi(k,i)}$ is relatively closed in $E \cup D_{\phi(k,i)}$, and hence

$$(6) \dim(E \cup D_i) = 0, \text{ for } i = 1, 2, \dots$$

We are now in a position to apply Theorem 1.1 to get embeddings $u_i : X^* \rightarrow I^{\mathbb{N}}$ and $v_i : X^* \rightarrow I^{\mathbb{N}}$, $i = 1, 2, \dots$, such that with $G_0 = \emptyset$, we have, cf. (2), (5) and (6),

$$(7) E \cup G_i = u_i^{-1}(\mathbb{P}^{\mathbb{N}}), C \subset u_i^{-1}(\mathbb{Q}^{\mathbb{N}}),$$

$$(8) E \cup D_i = v_i^{-1}(\mathbb{P}^{\mathbb{N}}), C \setminus D_i \subset v_i^{-1}(\mathbb{Q}^{\mathbb{N}}).$$

We shall show that the embedding

$$(9) h = (u_0, v_1, u_1, v_2, \dots) : X^* \rightarrow (I^{\mathbb{N}})^{\mathbb{N}}$$

satisfies the condition

$$(10) E = h^{-1}((\mathbb{P}^{\mathbb{N}})^{\mathbb{N}}), F = h^{-1}((\mathbb{Q}^{\mathbb{N}})^{\mathbb{N}}).$$

Since $E \cup G_0 = E$, the first part of (10) follows from (7) and (8).

From (3), (5), (7) and (8) we have also $F \subset h^{-1}((\mathbb{Q}^{\mathbb{N}})^{\mathbb{N}})$. Let $x \notin F$. If $x \notin C$, then by (2), $x \in G_i$ for some $i \geq 1$, and by (7), $h(x) \notin (\mathbb{Q}^{\mathbb{N}})^{\mathbb{N}}$. Let $x \in C \setminus F$, and using (3) and (4) pick k, i such that $x \in L_{ki} \setminus L_{k+1}$. Then by (5), for $j = \phi(k, i)$, $x \in D_j$, and by (8) and (9), $h(x) \notin (\mathbb{Q}^{\mathbb{N}})^{\mathbb{N}}$.

This ends the verification of (10).

To complete the proof of the theorem, it is enough to fix a bijection $\sigma : \mathbb{N} \rightarrow \mathbb{N} \times \mathbb{N}$ and to notice that for the homeomorphism $g : (I^{\mathbb{N}})^{\mathbb{N}} = I^{\mathbb{N} \times \mathbb{N}} \rightarrow I^{\mathbb{N}}$, $g(x)(n) = x(\sigma(n))$, we have $g((\mathbb{P}^{\mathbb{N}})^{\mathbb{N}}) = \mathbb{P}^{\mathbb{N}}$, and $g((\mathbb{Q}^{\mathbb{N}})^{\mathbb{N}}) = \mathbb{Q}^{\mathbb{N}}$. □

3. PROOF OF THEOREM 1.3.

A basic ingredient of our reasonings is the following lemma, which we shall derive from Lemma 2.1 in [PPR] and Theorem 1.1.

Lemma 3.1. *For each nonempty closed set F in $\mathbb{P}^{\mathbb{N}} \cup \mathbb{Q}^{\mathbb{N}}$ there is a continuous function $f : \mathbb{P}^{\mathbb{N}} \cup \mathbb{Q}^{\mathbb{N}} \rightarrow I$ such that $f(\mathbb{P}^{\mathbb{N}} \setminus F) \subset \mathbb{P}$, $f(\mathbb{Q}^{\mathbb{N}}) \subset \mathbb{Q}$ and $F = f^{-1}(0)$.*

Proof. Since $\mathbb{Q}^{\mathbb{N}}$ is a zero-dimensional F_{σ} -set in $\mathbb{P}^{\mathbb{N}} \cup \mathbb{Q}^{\mathbb{N}}$, similarly as at the beginning of the proof of Theorem 1.2, one can use [Ku], vol. II, §45, VII, Theorem 5, to get an embedding $u : \mathbb{P}^{\mathbb{N}} \cup \mathbb{Q}^{\mathbb{N}} \rightarrow I^{\mathbb{N}}$ such that $u(\mathbb{Q}^{\mathbb{N}})$ is contained in a zero-dimensional σ -compact set T in $I^{\mathbb{N}} \setminus u(\mathbb{P}^{\mathbb{N}})$.

Let $K = \overline{u(F)}$ be the closure of $u(F)$ in $I^{\mathbb{N}}$ (notice that $K \cap u(\mathbb{P}^{\mathbb{N}} \cup \mathbb{Q}^{\mathbb{N}}) = u(F)$).

Let us split $T \setminus K$ into pairwise disjoint compact sets C_1, C_2, \dots with $\text{diam} C_i \rightarrow 0$ (cf. the beginning of Sec. 4 in [PPR]), and let $q : I^{\mathbb{N}} \rightarrow I^{\mathbb{N}} / \sim$ be the quotient map determined by the equivalence relation \sim , where the equivalence classes are $C_i, i = 1, 2, \dots$ and the singletons $\{x\}, x \notin \bigcup_i C_i = T \setminus K$.

Then $q \circ u : \mathbb{P}^{\mathbb{N}} \cup \mathbb{Q}^{\mathbb{N}} \rightarrow I^{\mathbb{N}} / \sim$ embeds $\mathbb{P}^{\mathbb{N}} \setminus F$ onto a G_{δ} -set G , $S = q \circ u(\mathbb{Q}^{\mathbb{N}} \setminus F)$ is a countable set disjoint from G , and $A = q(K)$ is a compact set in $I^{\mathbb{N}} / \sim$ missing G . Applying Lemma 2.1 from [PPR], we get a continuous map $w : I^{\mathbb{N}} / \sim \rightarrow I$ with $w(G) \subset \mathbb{P}$, $w(S) \subset \mathbb{Q}$ and $A = w^{-1}(0)$. One readily checks that $f = w \circ q \circ u$ has required properties. □

Having proved Lemma 3.1, one can now justify Theorem 1.3 easily.

Since by Theorem 1.1, each one-dimensional compact space embeds in $\mathbb{P}^{\mathbb{N}} \cup \mathbb{Q}^{\mathbb{N}}$, we can assume that

(1) $X \subset \mathbb{P}^{\mathbb{N}} \cup \mathbb{Q}^{\mathbb{N}}$.

The set $A \subset X$ being $F_{\sigma\delta}$ in X ,

(2) $A = \bigcap_k \bigcup_l F_{kl}$, F_{kl} closed in X .

We shall apply to each $F_{kl} \subset \mathbb{P}^{\mathbb{N}} \cup \mathbb{Q}^{\mathbb{N}}$, cf. (1), Lemma 3.1, getting continuous functions $f_{kl} : X \rightarrow I$ such that

(3) $f_{kl}(\mathbb{P}^{\mathbb{N}} \cap (X \setminus F_{kl})) \subset \mathbb{P}$, $f_{kl}(\mathbb{Q}^{\mathbb{N}} \cap X) \subset \mathbb{Q}$, $F_{kl} = f_{kl}^{-1}(0)$.

Let

(4) $\mathbb{M} = \{n \in \mathbb{N} : n \neq 2^k 3^l, k, l = 1, 2, \dots\}$, $\sigma : \mathbb{M} \rightarrow \mathbb{N}$ bijection

and let us define $h : X \rightarrow I^{\mathbb{N}}$ by, cf. (4),

(5) $h(x)(n) = \begin{cases} f_{kl}(x), & \text{if } n = 2^k 3^l, \\ x(\sigma(n)), & \text{if } n \in \mathbb{M}. \end{cases}$

The second part of (5) guarantees that h is an embedding.

If $n = 2^k 3^l$, $h(x)(n) = f_{kl}(x)$, by (5), and hence, by (3), $h(x)(n) \in \mathbb{P}$ for $x \in \mathbb{P}^{\mathbb{N}} \setminus F_{kl}$, $f_{kl}(x) = 0$ for $x \in F_{kl}$, and $f_{kl}(x) \in \mathbb{Q} \setminus \{0\}$ for $x \in \mathbb{Q}^{\mathbb{N}} \setminus F_{kl}$.

It follows that $h(X) \subset (\mathbb{P} \cup \{0\})^{\mathbb{N}} \cup \mathbb{Q}^{\mathbb{N}}$ and moreover, $F_{kl} = f_{kl}^{-1}(0) = h^{-1}(H_{kl})$, cf. (1) in Section 1. By (2), we get $h^{-1}(H) = \bigcap_k \bigcup_l F_{kl} = A$.

4. COMMENTS.

4.1. Concerning $\mathbb{Q}^{\mathbb{N}}$. The space $\mathbb{Q}^{\mathbb{N}}$ (which is zero-dimensional and absolutely $F_{\sigma\delta}$) contains closed topological copies of all zero-dimensional absolutely $F_{\sigma\delta}$ -spaces.

The following elegant reasoning to this effect was kindly communicated to us by Jan van Mill.

Let X be a zero-dimensional absolutely $F_{\sigma\delta}$ -space. We can assume that $X \subset 2^{\mathbb{N}}$ and $X = \bigcap_n F_n$, where $F_1 \supset F_2 \supset \dots$ are σ -compact sets. Since $F_n \times 2^{\mathbb{N}} \times \mathbb{Q}$ is a zero-dimensional σ -compact nowhere countable and nowhere locally compact space, it is homeomorphic to $2^{\mathbb{N}} \times \mathbb{Q}$, by a classical Alexandroff-Urysohn theorem, cf. [vM], Corollary 5.2. In effect, F_n embeds onto a closed subspace of $2^{\mathbb{N}} \times \mathbb{Q}$. It follows that $\prod_n F_n$ embeds onto a closed subspace of $(2^{\mathbb{N}} \times \mathbb{Q})^{\mathbb{N}}$ and so does X , which can be identified with the diagonal of $\prod_n F_n$. Finally, $2^{\mathbb{N}} \subset \mathbb{Q}^{\mathbb{N}}$, so $(2^{\mathbb{N}} \times \mathbb{Q})^{\mathbb{N}}$ embeds onto a closed subspace of $\mathbb{Q}^{\mathbb{N}}$.

The result follows also readily from a difficult theorem of van Engelen, providing a topological characterization of $\mathbb{Q}^{\mathbb{N}}$, cf. [vE], Section 3.

4.2. Concerning Theorem 1.3. The result in Theorem 1.3 can be extended to higher Borel classes $F_{\sigma\delta\sigma}$, $F_{\sigma\delta\sigma\delta}$, \dots

To illustrate this point, let us consider the class of absolutely $F_{\sigma\delta\sigma}$ -sets.

For each triple $(k, l, m) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N}$ let us define, cf. (1) in Section 1 $H_{klm} = \{x \in I^{\mathbb{N}} : x(2^k 3^l 5^m) = 0\}$, and $H' = \bigcup_m \bigcap_k \bigcup_l H_{klm}$.

Then, with minor adjustments of the proof of Theorem 1.3, we get the assertion of this theorem for $F_{\sigma\delta\sigma}$ -sets A in compact one-dimensional spaces X , where H is replaced by H' .

REFERENCES

- [En] R. Engelking, *General Topology*, 2nd edition, Helderman, Berlin 1989.
- [vE] F. van Engelen, *Characterizations of the countable infinite products of rationals and some related problems*, Rend. Circ. Mat. Palermo 11 (1985), 37 - 54.
- [Ke] A. Kechris, *Classical descriptive set theory*, Springer 1995.
- [Ku] K. Kuratowski, *Topology vol. I and II*, Academic Press, New York 1966 and 1968.
- [vM] J. van Mill, *Characterization of some zero-dimensional separable metric spaces*, Trans. Amer. Math. Soc. 264 (1981), 205 - 215.
- [PPR] E. Pol, R. Pol, M. Reńska, *On an extension of the Nöbeling rational universal space*, Fund. Math. (to appear).
- [Si] R. Sikorski, *Some examples of Borel sets*, Coll. Math. 5 (1958), 170 - 171.

INSTITUTE OF MATHEMATICS, UNIVERSITY OF WARSAW, BANACHA 2, 02-097
WARSAWA, POLAND

Email address: e.pol@mimuw.edu.pl

INSTITUTE OF MATHEMATICS, UNIVERSITY OF WARSAW, BANACHA 2, 02-097
WARSAWA, POLAND

Email address: r.pol@mimuw.edu.pl

FACULTY OF MATHEMATICS AND INFORMATION SCIENCE, WARSAW UNIVERSITY
OF TECHNOLOGY, KOSZYKOWA 75, 00-662 WARSAWA, POLAND

Email address: miroszlawa.renska@pw.edu.pl