# ON A FAMILY OF HYPERBOLIC BRUNNIAN LINKS AND THEIR VOLUMES DUŠAN D. REPOVŠ AND ANDREI YU. VESNIN

ABSTRACT. An *n*-component link *L* is said to be *Brunnian* if it is non-trivial but every proper sublink of *L* is trivial. The simplest and best known example of a hyperbolic Brunnian link is the 3-component link known as "Borromean rings". For  $n \ge 2$ , we introduce an infinite family of *n*-component Brunnian links with positive integer parameters  $Br(k_1, \ldots, k_n)$  that generalize examples constructed by Debrunner in 1964. We are interested in hyperbolic invariants of 3-manifolds  $S^3 \setminus Br(k_1, \ldots, k_n)$  and we obtain upper bounds for their volumes. Our approach is based on Dehn fillings on cusped manifolds with volumes related to volumes of ideal right-angled hyperbolic antiprisms.

## 1. INTRODUCTION

If a link L in  $S^3$  is nontrivial, yet every proper sublink of L is trivial, we say that L is *Brunnian* (or that L has the *Brunnian property*). Links with this property were studied by Milnor [18] who called them *almost trivial* links. In 1961 Debrunner [9] renamed them as *Brunnian* links, in honor of Hermann Brunn whose early contributions [8] to knot theory also included examples of such links. In recent decades, Brunnian links have been under investigation from several points of views. In particular, we refer to Bai [5], Bai and Ma [6], Bai and Wang [7], and Kanenobu [14, 15] for hyperbolic and satellite properties of Brunnian links, Lei, Wu and Zhang [17] for intersecting subgroups of Brunnian link groups, and Habiro and Meilhan [13] for finite-type invariants and Milnor invariants of Brunnian links. We would also like to mention molecular Borromean rings considered by Wang and Stoddart [22] and a possible physical realization of higher-order Brunnian structures studied by Baas [4].

The simplest and best known example of a 3-component Brunnian link is the 3-component link  $6_2^3$  in Rolfsen's notations [19] which is also known as the

<sup>2020</sup> Mathematics Subject Classification. 57K10, 57K32, 52B10.

Key words and phrases. Hyperbolic Brunnian link, Adams move, augmented link, ideal right-angled antiprism.

D.R. was supported by the Slovenian Research and Innovation Agency (program P1-0292 and grants J1-4031, J1-4001, N1-0278, N1-0114, and N1-0083). A.V. was supported by the Ministry of Science and Higher Education of Russia under agreement no. 075-02-2024-1437.

"Borromean rings". We denote it by  $\mathcal{B}$ . It is well-known from Thurston [21, Chapter 3] that link  $\mathcal{B}$  is hyperbolic, the complement  $S^3 \setminus \mathcal{B}$  can be decomposed into two copies of an ideal right-angled octahedron, and  $\operatorname{vol}(S^3 \setminus \mathcal{B}) = 7.327724$ , up to six digits.

We recall that a polyhedron in a hyperbolic 3-space  $\mathbb{H}^3$  is said to be *ideal* if all of its vertices belong to  $\partial \mathbb{H}^3$ , and it is called *right-angled* if all of its dihedral angles are equal to  $\pi/2$ . By virtue of the Andreev theorem [3], every ideal rightangled hyperbolic polyhedron is determined by its 1-dimensional skeleton, up to an isometry of  $\mathbb{H}^3$ . An initial list of ideal right-angled hyperbolic polyhedra and their volumes was presented by Egorov and Vesnin [10, 11], and the upper volume bounds depending only of number of vertices were obtained by Alexandrov, Bogachev, Vesnin, and Egorov [2].

In the present chapter, we introduce for every  $n \ge 2$ , an infinite family of *n*-component Brunnian links which generalize examples constructed in 1961 by Debrunner [9]. We are interested in the hyperbolic structure on the complements of these links. By using decompositions of complements of fully augmented links into pairs of ideal right-angled polyhedra by a method from an appendix by Agol and Thurston in Lackenby [16], and the Dehn filling theorem, we provide upper bounds for volumes of such hyperbolic Brunnian links.

The chapter is organized as follows: In Section 2 we introduce a family of (3n+2)-component links  $L_n$ , for every  $n \ge 2$ , and demonstrate that their complement  $S^3 \setminus L_n$  can be decomposed into four ideal right-angled antiprisms  $A_{2n}$ . Then we apply Thurston's formula from [21, Chapter 6] for volumes of antiprisms to obtain a formula for  $vol(S^3 \setminus L_n)$  (see Theorem 2.1). Next, applying Adams moves to 2n vertical components of  $L_n$ , we construct links  $L'_n$  with 3n components such that  $vol(S^3 \setminus L'_n) = vol(S^3 \setminus L_n)$ . In Section 3, we construct by Dehn fillings on 2n components of  $L'_n$ , a family of n-component links  $Br(k_1, \ldots, k_n)$ , depending on the filling parameters  $k_1, \ldots, k_n$  (see Theorems 3.1 and 3.4). In particular, when all  $k_i$  are equal to 1, we get the Brunnian links from Debrunner [9], whose hyperbolicity was established by Bai [5].

### 2. Hyperbolic links and ideal right-angled antiprisms

Recall that a knot or a link  $K \subset S^3$  is said to be *hyperbolic* if its complement  $S^3 \setminus K$  admits a complete metric of constant curvature -1. Equivalently, the 3-manifold  $S^3 \setminus K = \mathbb{H}^3/G$  is hyperbolic, where  $\mathbb{H}^3$  is the hyperbolic 3-space and G is a discrete, torsion-free group of isometries, isomorphic to  $\pi_1(S^3 \setminus K)$ .

For  $n \ge 2$ , let us denote a link with 3n + 2 components by  $L_n$ , where each component is a circle and the components are linked in the same manner as shown in Figure 1 for the case n = 3.

3



FIGURE 1. Link  $L_n$  with 3n + 2 components, case n = 3.

By using the computer program SnapPy [20], one can see that the 8-component link  $L_2$  is hyperbolic and has  $vol(S^3 \setminus L_2) = 24.092184$ , up to six digits. To demonstrate hyperbolicity of  $L_n$  for arbitrary  $n \ge 2$  and find the volume formula for  $S^3 \setminus L_n$ , we shall use the approach from Lackenby [16] (see also Futer and Purcell [12]). In the terminology of Lackenby [16], the link  $L_n$  is augmented, with 2n "vertical" components (colored in red in Figure 1), and  $S^3 \setminus L_n$  admits a decomposition into two ideal polyhedra  $P_n$  and  $P'_n$  with faces identified in pairs. The polyhedra  $P_n$  and  $P'_n$  are identical. By the construction each vertical component of  $L_n$  gives to a pair of triangles in  $P_n$  with a common vertex like a bowtie, as presented in Figure 2 where common vertices are red.



FIGURE 2. Associating bowties to vertical components of  $L_n$ , case n = 3.

At the last step, in order to obtain the polyhedron  $P_n$ , we shall compress black edges which connect two vertices of valence three, to obtain a new vertex of valence four. There are 4n such edges which give us 2n black vertices, as presented in Figure 3 (to complete the construction, it is necessary to identify the vertices A and B, as well as the vertices C and D). As a result,  $P_n$  is an ideal



FIGURE 3. 1-skeleton of the polyhedron  $P_n$ , case n = 3.

right-angled polyhedron with 6n vertices, where 2n red vertices correspond to bowties and 4n black vertices appeared after compressing black edges. Moreover,  $P_n$  has two 2n-gonal faces as its top and bottom, 2n triangles incident to the top, 2n triangles incident to the bottom, and 4n quadrilaterals on the middle level. By cutting  $P_n$  along the middle line passing through quadrilaterals, we shall see that  $P_n$  can be decomposed into two identical ideal right-angled 2ngonal antiprisms  $A_{2n}$ . Recall that  $A_n$  has 2n + 2 faces, where two *n*-gonal faces can be considered as the top and the bottom, and 2n triangular faces on the lateral surface, see Figure 4 for the antprism  $A_4$ .



FIGURE 4. 1-skeleton of antiprism  $A_4$ .

Volumes of right-angled antiprisms  $A_n$ ,  $n \ge 3$ , are given by the following formula obtained by Thurston [21, Example 6.8.7], where an antiprism was named a *drum with triangles*,

(1) 
$$\operatorname{vol}(A_n) = 2n \left[ \Lambda \left( \frac{\pi}{4} + \frac{\pi}{2n} \right) + \Lambda \left( \frac{\pi}{4} - \frac{\pi}{2n} \right) \right].$$

Here,  $\Lambda(\theta)$  is the Lobachevsky function defined in [21, Chapter 7] as

$$\Lambda(\theta) = -\int_0^\theta \log|2\sin(t)|dt.$$

In particular, up to six digits, we have the following volumes of ideal right-angled antiprisms:

$$vol(A_3) = 3.663863,$$
  
 $vol(A_4) = 6.023046,$   
 $vol(A_5) = 8.137885,$   
 $vol(A_6) = 10.149416$ 

**Theorem 2.1.** For every  $n \ge 2$ , the following formula holds:

$$\operatorname{vol}(S^3 \setminus L_n) = 16n \left[ \Lambda \left( \frac{\pi}{4} + \frac{\pi}{4n} \right) + \Lambda \left( \frac{\pi}{4} - \frac{\pi}{4n} \right) \right].$$

*Proof.* Indeed, from the above considerations we obtain

$$\operatorname{vol}(S^3 \setminus L_n) = 2\operatorname{vol}(P_n) = 4\operatorname{vol}(A_{2n}),$$

so the assertion follows by formula (1).

The following was proved by Adams [1, Corollary 5.1].

**Theorem 2.2** (see [1]). Let J be a link in  $S^3$  such that  $S^3 \setminus J$  is hyperbolic and J has a projection for which some part appears as in Figure 5(a). Let J'be the link obtained by replacing that part by the projection of J appearing in Figure 5(a) with the one appearing in Figure 5(b). Then  $S^3 \setminus J'$  is hyperbolic with the same volume as  $S^3 \setminus J$ .



FIGURE 5. Replacing the part of the projection by Theorem 2.2.

We shall call the replacement presented in Figure 5 and its inverse the Adams moves. Let us apply Adams moves to each of 2n parts of related vertical components of the hyperbolic link  $L_n$ . The resulting link  $L'_n$  with 3n components is depicted in Figure 6 for the case n = 3.

Theorems 2.1 and 2.2 now yield the following result.



FIGURE 6. Link  $L'_n$  with 3n components, case n = 3.

**Corollary 2.3.** For every  $n \ge 2$ , the following formula holds:

(2) 
$$\operatorname{vol}(S^3 \setminus L'_n) = 16n \left[ \Lambda \left( \frac{\pi}{4} + \frac{\pi}{4n} \right) + \Lambda \left( \frac{\pi}{4} - \frac{\pi}{4n} \right) \right].$$

# 3. A family of hyperbolic Brunnian links

Suppose that M is a compact orientable 3-manifold with  $\partial M$  a collection of tori, and that the interior  $M \subset \overline{M}$  admits a complete hyperbolic structure. Then M is referred to as a *cusped manifold*. For any  $n \geq 2$ , the complement  $S^3 \setminus L'_n$  is a cusped manifold by construction.

For every  $n \geq 2$  and every positive integers  $k_1, \ldots, k_n$ , we define an *n*component link  $Br(k_1, \ldots, k_n)$  with a diagram having 2n blocks of twist regions as follows: Consider *n* top blocks consisting of  $2k_i + 1$  positive half-twists and *n* bottom blocks consisting of  $2k_i + 1$  negative half-twists for  $i = 1, \ldots, n$ . A diagram of Br(1, 2, 1) is presented in Figure 7. By Rolfsen [19],  $Br(k_1, \ldots, k_n)$ can be considered as a result of Dehn fillings with slopes  $1/k_i$  and  $-1/k_i$  for  $i = 1, \ldots, n$  on 2n cusps corresponding to vertical components of  $S^3 \setminus L'_n$  in Figure 6. Thus, in total we get 2n twist regions with  $2k_i + 1$  positive and negative half-twists,  $i = 1, \ldots, n$ .

If all  $k_i$  are equal to some k, then  $Br(k, k, \ldots, k)$  admits a cyclic rotation symmetry of order n. The links  $Br(1, 1, \ldots, 1)$  appeared already in Debrunner [9, Fig. 2] and were denoted by  $L_F$ . It was demonstrated in [9] that  $L_F$  is unsplittable, but any proper sublink of  $L_F$  is completely splittable. The 5-component link presented in Rolfsen [19, p. 69] is link Br(1, 1, 1, 1, 1) in our notation. By the same argument as in [9], the following result holds.

**Theorem 3.1.** For every n,  $Br(k_1, \ldots, k_n)$  is an n-component Brunnian link.



FIGURE 7. Link Br(1, 2, 1).

*Proof.* It is easy to see that after removing one component of  $Br(k_1, \ldots, k_n)$ , all other components can be transformed to trivial ones by a sequence of simplifying underpassing moves from (a) to (b) shown in Figure 8.



FIGURE 8. The simplifying underpassing move.

A practical method to check the hyperbolicity of Brunnian links was presented by Bai [5] who proved that for  $n \ge 2$  with  $k_i = 1, i = 1, ..., n$ , the link  $Br(k_1, ..., k_n)$  is hyperbolic [5, Theorem 1.3]. The smallest one is a 2component link Br(1, 1) with 12 crossings and  $vol(S^3 \setminus Br(1, 1)) = 12.528922$ , up to six digits. Moreover, it can be recognized by using SnapPy [20] that Br(1, 1) = L12n1180, a non-alternating link with 12 components. At the same time, for  $k_i \ge 3$ , twisted regions in the diagram of  $Br(k_1, \ldots, k_n)$  have more than 6 crossings. Therefore, the following result from Futer and Purcell [12, Theorem 1.7] can be applied.

**Theorem 3.2** (see [12]). Let K be a link in  $S^3$  with a prime, twist-reduced diagram D(K). Suppose that every twist region of D(K) contains at least 6 crossings and that each component of K passes through at least 7 twist regions

7

(counted with their multiplicity). Then every non-trivial Dehn filling of all the components of K is hyperbolic.

The relation between volumes of cusped manifold and hyperbolic manifolds obtained by Dehn filling is due to Gromov and Thurston [21, Theorem 6.5.6].

**Theorem 3.3** (see [21]). Suppose M is a complete hyperbolic manifold of finite volume and that  $N \neq M$  is a complete hyperbolic manifold obtained topologically by replacing certain cusps of M by solid tori. Then vol(N) < vol(M).

**Theorem 3.4.** For hyperbolic links  $Br(k_1, \ldots, k_n)$ , the following upper bound holds:

$$\operatorname{vol}(S^3 \setminus B(k_1, \dots, k_n)) < 16n \left[\Lambda\left(\frac{\pi}{4} + \frac{\pi}{4n}\right) + \Lambda\left(\frac{\pi}{4} - \frac{\pi}{4n}\right)\right]$$

*Proof.* Since  $Br(k_1, \ldots, k_n)$  can be obtained by Dehn filling on 2n cusps of the hyperbolic link  $L'_n$ , the result follows by Corollary 2.3 and Theorem 3.3.

**Corollary 3.5.** For every  $n \ge 2$ , the value  $\beta_n = 16n \left[\Lambda \left(\frac{\pi}{4} + \frac{\pi}{4n}\right) + \Lambda \left(\frac{\pi}{4} - \frac{\pi}{4n}\right)\right]$  is the limit point for volumes of hyperbolic Brunnian links with n components.

We conclude the chapter by the following open problems concerning hyperbolic Brunnian links.

**Problem 3.1.** What is the smallest volume hyperbolic Brunnian link with n components?

It is well known that the link "Borromean rings" is arithmetic [21, Chapter 7].

**Problem 3.2.** Which Brunnian links are arithmetic?

Also, the following interesting problem was formulated by Bai and Ma [6, Problem 7.0.8].

**Problem 3.3** (see [6]). Let B(n) be the number of Brunnian links with n or fewer crossings,  $B_h(n)$  the number of hyperbolic Brunnian links with n or fewer crossings, and denote

$$\limsup_{n \to \infty} \frac{B_h(n)}{B(n)} = a, \qquad \liminf_{n \to \infty} \frac{B_h(n)}{B(n)} = b.$$

Is then a = b, b = 0, or a < 1?

Acknowledgements. We thank the editors and referees for comments and suggestions.

#### References

- C. Adams, *Thrice-punctured spheres in hyperbolic 3-manifolds*, Transactions of the American Mathematical Society 287 (1985), 645-656.
- [2] S. Alexandrov, N. Bogachev, A. Vesnin, A. Egorov, On volumes of hyperbolic right-angled polyhedra, Sbornik: 214:2Mathematics (2023),148 https://doi.org/10.4213/sm9740e Preprint version available 165.is  $\mathbf{at}$ https://arxiv.org/abs/2111.08789
- [3] E. M. Andreev, On convex polyhedra of finite volume in Lobachevskii space, Mathematics of the USSR-Sbornik 12:2 (1970), 255-259.
- [4] N.A. Baas, New states of matter suggested by new topological structures, International Journal of General Systems 42:2 (2013), 137-169. https://doi.org/10.1080/03081079.2012.728403 Preprint version is available at https://arxiv.org/abs/1012.2698.
- [5] S. Bai, *Hyperbolic Brunnian links*, Preprint available at https://arxiv.org/abs/2104.12637.
- [6] S. Bai, J. Ma, Satellite constructions and geometric classification of Brunnian links, Journal of Knot Theory and Its Ramifications 30:10 (2021), art. no. 2140005. https://doi.org/10.1142/S0218216521400058 Preprint version is available at https://arxiv.org/abs/1906.01253.
- S. Bai, W. Wang, New criteria and constructions of Brunnian links, Journal of Knot Theory and Its Ramifications 29:13 (2020), art. no. 2043008. https://doi.org/10.1142/S0218216520430087 Preprint version is available at https://arxiv.org/abs/2006.10290.
- [8] H. Brunn, Über Verkettung, Sitzungsberichte der Bayerischen Akademie der Wissenschaften, Math.-Naturwiss. Klasse 22 (1892), 77-99.
- [9] H. Debrunner, *Links of Brunnian type*, Duke Mathematical Journal **28** (1961), 17-23.
- [10] A. Vesnin, Α. Egorov, Ideal right-angled poluhedra inLobachevsky space. Chebyshevskii Sbornik 21:2 (2020),65 - 83.(in Russian) https://doi.org/10.22405/2226-8383-2020-21-2-65-83 English preprint version is available at https://arxiv.org/abs/1909.11523
- [11] A. Egorov, A. Vesnin, Volume estimates for right-angled hyperbolic polyhedra, Rendiconti dell'Istituto di Matematica dell'Università di Trieste 52 (2020), 565-576. https://doi.org/10.13137/2464-8728/30958 Preprint version is available at https://arxiv.org/abs/2010.11147
- [12] D. Futer, J.S. Purcell, Links with no exceptional surgeries, Commentarii Mathematici Helvetici 82 (2007), 629-664. https://doi.org/10.4171/CMH/105
- [13] K. Habiro, J.-B. Meilhan, Finite type invariants and Milnor invariants for Brunnian links, International Journal of Mathematics 19:6 (2008), 747-766. https://doi.org/10.1142/S0129167X08004820
- [14] T. Kanenobu, Satelite links with Brunnian properties, Archiv der Mathematik 44 (1984), 369-372.
- [15] T. Kanenobu, Hyperbolic links with Brunnian properties, Journal of the Mathematical Society of Japan 38:2 (1986), 295-308.

- [16] M. Lackenby, The volume of hyperbolic alternating link complements. With an appendix by I. Agol and D. Thurston, Proceedings of the London Mathematical Society 88 (2004), 204-224. https://doi.org/10.1112/S0024611503014291
- Lei, [17] F. J. Wu, Υ. Zhang, Onintersecting subgroups ofBrunnian Algebraic and Geometric Topology linkgroups, 16(2016),1043-1061. https://doi.org/10.2140/agt.2016.16.1043
- [18] J. Milnor, *Link groups*, Annals of Mathematics **59** (1954), 177-195.
- [19] D. Rolfsen, Knots and Links, Corrected reprint of the 1976 original, Mathematics Lecture Series Vol. 7, Publish or Perish, Houston, TX, 1990.
- [20] SnapPy, Version 3.1 (May 2023). https://snappy.math.uic.edu
- [21] W. P. Thurston, The Geometry and Topology of Three-Manifolds, Collected Works of William P. Thurston with Commentary, Vol. IV, With a preface by S.P. Kerckhoff, Edited by B. Farb, D. Gabai and S. P. Kerckhoff, American Mathematical Society, Providence, RI, 2022. ISBN: 9781470463915; 9781470451646
- [22] Y. Wang, J. F. Stoddart. Molecular Borromean rings: from controlledconstruction topotentialapplications, Chem 3:1 (2017),17-18. https://doi.org/10.1016/j.chempr.2017.06.009

FACULTY OF EDUCATION, FACULTY OF MATHEMATICS AD PHYSICS, UNIVERSITY OF LJUBLJANA

& Institute of Mathematics, Physics and Mechanics, Ljubljana, 1000, Slovenia https://orcid.org/0000-0002-6643-1271

Email address: dusan.repovs@guest.arnes.si

SOBOLEV INSTITUTE OF MATHEMATICS, RUSSIAN ACADEMY OF SCIENCES, NOVOSIBIRSK, 630090

& Regional Mathematical Center, Tomsk State University, Tomsk, 634050, Russia

HTTPS://ORCID.ORG/0000-0001-7553-1269

Email address: vesnin@math.nsc.ru