Bag Semantics Conjunctive Query Containment. Four Small Steps Towards Undecidability*

Jerzy Marcinkowski

Mateusz Orda

Institute of Computer Science, University of Wrocław; Wrocław, Poland.

Abstract

Query Containment Problem (QCP) is one of the most fundamental decision problems in database query processing and optimization.

Complexity of QCP for conjunctive queries (QCP_{CQ}) has been fully understood since 1970s. But, as Chaudhuri and Vardi noticed in their classical paper [1], this understanding is based on the assumption that query answers are sets of tuples, and it does not transfer to the situation when multi-set (bag) semantics is considered.

Now, 30 years after [1] was written, decidability of QCP_{CQ} for bag semantics remains an open question, one of the most intriguing open questions in database theory.

In this paper we show a series of undecidability results for some generalizations of bag-semantics QCP_{CQ} . We show, for example, that the problem whether, for given two boolean conjunctive queries ϕ_s and ϕ_b , and a linear function \mathbb{F} , the inequality $\mathbb{F}(\phi_s(D)) \leq \phi_b(D)$ holds for each database instance D, is undecidable¹.

1 The context. And our contribution.

1.1 The context

Query Containment Problem (QCP) is one of the most fundamental decision problems in database query processing and optimization. It is formulated as follows:

The instance of QCP are two database queries, Ψ_s and Ψ_b . The question is whether $\Psi_s(D) \subseteq \Psi_b(D)$ holds for each database D.

In the above, by $\Psi(D)$ we denote² the result of applying query Ψ to the database *D*. If the reader wondered what the subscripts *s* and *b* are supposed to mean: *s* stands for ,,small" and *b* stands for ,,big" (and we use this naming convention through this paper, sometimes also using terms *s*-query and *b*-query for Ψ_s and Ψ_b). QCP asks if the answer to the ,,small" query is always contained in the answer to the ,,big" one.

^{*}This research was supported by grant 2022/45/B/ST6/00457 from the Polish National Science Centre (NCN).

¹Notice that if ϕ is a boolean conjunctive query then, under the multiset semantics, $\phi(D)$ is a natural number.

²See Section 2 for more explanation regarding the notations we are using.

As usual in such situations, the problem comes in many variants, depending on two parameters: on the class of queries we allow and on the precise semantics of $\Psi(D)$ (and – in consequence – of the precise semantics of the symbol \subseteq). The classes of queries which have been considered in this context are CQ (conjunctive queries), or UCQ (unions of conjunctive queries) or CQ_{\neq} (conjunctive queries with inequalities), or some subsets of CQ. The possible semantics of $\Psi(D)$ are two: either we can see $\Psi(D)$ as a relation, that is a **set** of tuples, or as a multirelation, that is a **multiset** also known as a **bag** of tuples³. In the first case, the \subseteq in the above statement of QCP is understood to be the set inclusion, in the second case it is the multiset inclusion. We use natural notations to call the variants, for example $QCP_{CQ,\neq}^{bag}$ is QCP for conjunctive queries with inequality, under bag semantics and QCP_{UCQ}^{set} is QCP for unions of CQs under set semantics. QCP^{set} has long been well understood. Already in 1977 Chandra and Merlin [2] realized that QCP_{CQ}^{set} is

 QCP^{set} has long been well understood. Already in 1977 Chandra and Merlin [2] realized that $QCP^{\text{set}}_{\text{cc}}$ is NP-complete. Concerning more general classes of queries, it was shown in [3] that $QCP^{\text{set}}_{\text{ucq}}$ is Π_2^{p} -complete . Then another more general class, conjunctive queries with comparison predicates \neq and \leq , was studied in [4], where it was proven that $QCP^{\text{set}}_{\text{cq},\neq,\leq}$ is also in Π_2^{p} , but no lower bound was established. This gap was finally filled by [5], which proves that $QCP^{\text{set}}_{\text{cq},\neq,\leq}$ is Π_2^{p} -complete.

But a case can be made that in real database systems, where duplicate tuples are not eliminated, queries are usually evaluated under bag semantics, not set semantics.

Unfortunately, as it was realized in the early 1990s, no tools or techniques developed for the analysis of QCP^{set} survive in the context of QCP^{bag} . In the seminal paper [1] the authors observe that the proof of the Chandra-Merlin NP upper bound for QCP^{set}_{CQ} does not survive in the bag-semantics world, and claim a Π_2^P lower bound for QCP^{bag}_{CQ} , deferring the proof however to the full version of the paper.

The same observation was made also in an earlier⁴ paper [6], less well known than [1]. Let us quote [6] here: *The classical theorem by Chandra and Merlin does not hold, because it treats relations as sets and not multisets.* (...). In general, there is almost no theory on the properties of queries and programs that retain duplicates. The development of such a theory is part of our future plans.

But such theory was never really developed, the full version of [1] never appeared, and neither the authors of [1], nor anyone later on, proved any upper bound⁵ for the complexity of QCP_{CQ}^{bag} . So not only nothing is known about the complexity of this fundamental problem but even its decidability has now been an open problem for 30 years. And this is not because people did not try.

When a difficult decision problem is attacked, the action usually takes place in two theaters of operations: on the positive side, where more and more general subcases of the problem are being proven to be decidable, and on the negative side, where undecidability results are shown for some generalizations of the problem.

On the **positive side**, numerous results were produced, which seem to naturally fall into two main lines of attack.

One of these lines includes decidability of QCP^{bag} for projection-free conjunctive queries [7]. In a related paper [8] the authors generalize QCP and then give a partial positive answer for the problem of query containment of (unions of) conjunctive queries over so called annotated databases. This line of research was continued in [9] and [10]. Then, more recently, the decidability result from [7] was extended to the case where Ψ_s is a projection-free CQ and Ψ_b is an arbitrary CQ [11]. The proof is via a reduction to a known decidable class of Diophantine inequalities.

The second line of attack originated from the work of Kopparty and Rossman [12]. They observe that QCP_{CQ}^{bag} is a purely combinatorial (or graph theoretic) phenomenon related to the notion of homomorphism domination exponent. In consequence, they postulate that the existing combinatorial technology could be

³We should probably remark here that, while $\Psi(D)$ may be a multiset, D is always a relational structure in this paper.

⁴It seems that the authors of [1] were not aware of [6].

⁵We of course know that the problem is in co-r.e.

used to approach the problem (and indeed they prove decidability for a case when Ψ_s is series-parallel and Ψ_b is a chordal graph). Their mathematically very attractive toolbox features the information-theoretic notion of entropy. Unfortunately, the paper [13] exhibits the limitations of this attitude, showing that decidability of QCP_{cq}^{bag} , even if restricted to the case where Ψ_b is an acyclic CQ, is already equivalent to an long standing open problem in information theory, decidability of the problem Max-IIP.

On the **negative side**, which is more interesting from the point of view of our paper, the results are so far very few. All we know is that the two most natural extensions of QCP_{cq}^{bag} are undecidable. First [14] proved that QCP_{ucq}^{bag} is undecidable. The proof is quite easy – it is a straightforward encoding of Hilbert's 10th problem. Then, in 2006, [15] have shown that $QCP_{cq,\neq}^{bag}$ is also undecidable. The argument here is much more complicated than the one in [14] and, while "real" conjunctive queries are mentioned in the title of [15], the queries needed for the proof of this negative result require no less than 59¹⁰ inequalities.

No progress has been made in this theater of operations since that time.

1.2 Our contribution

In this paper we present a series of negative results for some generalizations of QCP_{CQ}^{bag} . We also notice that (some of) our results are (in some sense) ultimate: no stronger undecidability result is possible unless the problem QCP_{CQ}^{bag} itself is undecidable.

All our results hold for **boolean conjunctive queries** (with, or without, inequality). For a boolean query, the result of its application to a database is a natural number (see Section 2) and, in consequence, the \subseteq symbol from the QCP statement at the beginning of Section 1.1 turns into \leq .

Call a database *D* **non-trivial** if it contains two **different** constants, σ^* and φ . Our first result is:

Theorem 1. The problem:

Given are boolean conjunctive queries (without inequality) ϕ_s and ϕ_b , and a natural number \mathfrak{c} . Does $\mathfrak{c}\phi_s(D) \leq \phi_b(D)$ hold for each non-trivial database D? is undecidable.

Notice that Theorem 1 would make no sense without the condition that D must be non-trivial, at least not for any c > 1. This is because if D is the "well of positivity" – a structure with a single vertex⁶ such that all atomic formulas are true in D for this vertex then whatever queries ϕ_s and ϕ_b (without inequality) we take, we get $\phi_b(D) = \phi_b(D) = 1$, and thus $c\phi_s(D) > \phi_b(D)$.

As the following theorem says we can however replace the non-triviality condition with an additive constant: it is undecidable, for two conjunctive queries ϕ_s and ϕ_b , without equality, and a linear function \mathbb{F} , whether $\mathbb{F}(\phi_s(D))$ is bounded by $\phi_b(D)$. In other words:

Theorem 2. The problem:

Given are boolean conjunctive queries (without inequality) φ_s and φ_b , and natural numbers \mathfrak{c} and \mathfrak{c}' . Does $\mathfrak{c}\varphi_s(D) \leq \varphi_b(D) + \mathfrak{c}'$ hold for each D? is undecidable.

Our next result does not involve any multiplicative constants:

⁶For a similar reason, the homomorphism domination exponent in [12] is only defined for structures which allow for at least two different homomorphisms, which also rules out single-vertex databases.

Theorem 3. The problem:

Given are boolean conjunctive queries (without inequality) ψ_s and ψ_b (with at most one inequality). Does $\psi_s(D) \leq \psi_b(D)$ hold for each non-trivial database D? is undecidable.

Non-triviality can of course be enforced by replacing ψ_s with $\sigma \neq \varphi \land \psi_s$. So Theorem 3 is an improvement upon the main result from [15]: we get undecidability of $QCP_{co,\neq}^{bag}$ already for queries with one inequality each, instead of 59^{10} .

Notice that again the non-triviality assumption cannot be easily dropped here: by the "well of positivity" argument the query $\rho_b \wedge (x \neq x')$ never contains ρ_s (for queries ρ_s and ρ_b being CQs without equality). The next theorem shows, that this "well of positivity" argument is actually the only reason why we need non-triviality:

Theorem 4. The problem:

Given boolean conjunctive queries ρ_s (without inequality) and ρ_b (with at most one inequality). Does $\rho_s(D) \leq max\{1, \rho_b(D)\}$ hold for each database D? is undecidable.

Theorem 4 says (with one caveat) that the inequality in the *s*-query is not really essential for the negative result from [15]. Finally, we tried to address the question, whether we could get rid of the inequality in the *b*-query rather than the one on the *s*-query and still prove undecidability. The answer is given in Theorem 5: maybe we can, but this can only happen if QCP_{co}^{bag} itself is undecidable:

Theorem 5. The problem:

Given two conjunctive queries ψ_s (with any number of inequalities) and ψ_b (without inequalities), Does $\psi_s(D) \leq \psi_b(D)$ hold for each database D? is decidable if and only if QCP_{co}^{bag} is decidable.

What concerns **our techniques:** all the undecidability results in bag-semantics database theory we are aware of ([14], [15], [16]) use Hilbert's 10th Problem as the source of undecidability (see Section 4.1). In all these papers the database provides a valuation of the numerical variables, and the universal quantification from the Hilbert's Problem is simulated by the universal quantification over databases. The question is how to encode the evaluation of a given polynomial using the syntax under analysis.

This can be easily done if we deal with UCQs, like in [14]: a monomial translates in a very natural way into a CO, and a sum of monomials into a disjunction of COs.

In [15] a trick was found to encode an entire polynomial as one CQ. But this only works well for some "good" databases. So this trick has been married in [15] to an elaborate anti-cheating mechanism, which guarantees that if D is not "good" then (using the language of Theorem 3 above) $\psi_b(D)$ is easily big enough to be greater than $\psi_s(D)$. And it is this anti-cheating mechanism in [15] that requires an astronomical number of inequalities.

The main idea behind our proof of Theorems 1 and 2 is a new polynomial-encoding trick, which superficially looks quite similar to the one from [15], and also only works for "good databases" (which we call correct in Section 4), but is different enough not to require any inequalities in the anti-cheating part (in particular, nothing similar to Lemma 12, which is very important in our proof, seems to be compatible with the trick from [15]). This comes with a cost however, which is the multiplicative constant c.

Theorems 3 and 4 follow from Theorem 1. We use a combinatorial argument to show how to use the single inequality in ψ_h to simulate multiplication by \mathfrak{c} . This part is not related to any previous work we are aware of.

Organization of the paper. In Section 3 we show how Theorem 3 follows from Theorem 1. Theorem 1 itself is proved in Section 4 (due to space limitation proofs of two lemmas are presented in Appendix A and Appendix B). Sections 3 and 4 can be read in any order.

Proofs of Theorems 4 and 2 are deferred to the full paper. They follow the same paths as the proofs of Theorems 1 and 3, but there are some new obstacles there to overcome. This is because the anti-cheating mechanism in the proofs of Theorems 1 and 3 strongly relies on the input database to be non-trivial. Which means that another level of anti-cheating response must be present, in the proofs of Theorems 4 and 2, to make sure that in trivial databases the answer to the *b*-query is always at least big as the answer to the *s*-query.

The (easy) proof of Theorem 5 can be found in Section 5.

We begin the technical part with Section 2 where our notations and other basic concepts are explained.

2 Preliminaries and notations

2.1 Some standard notions and notations

We use the terms **database** and **structure** interchangeably, to denote a finite relational structure over some relational schema (signature). Apart from relations we allow for constants (see Section 2.3) in the signature: *a* and *b* (possibly with subscripts) are used to denote such constants as well as σ^a and φ . We use the letter *D* (possibly with subscripts) to denote structures. If *D* is a structure then by V_D we mean the set of vertices of *D* (or the active domain of *D* if you prefer a more database-theoretical terminology). If *a* is a constant and *D* is a structure then we also use *a* to denote the interpretation of *a* in *D* (instead of the more formally correct a_D).

When we say "query" we always⁷ mean a conjunctive query (CQ). All queries in this paper are boolean. We never explicitly write the existential quantifiers if front of queries, but whenever we write a query it is assumed that all the variables are existentially quantified. To denote queries we always use lower case Greek letters. If ψ is a query then $Var(\psi)$ is the set consisting of all the variables which appear in ψ and V_{ψ} is the set consisting of all the variables **and** all the constants which appear in ψ . This is consistent with V_D being the active domain of a database, since we tacitly identify queries with their **canonical structures**: the active domain of the canonical structure of query ψ is V_{ψ} and the atoms of the canonical structure are the atoms of ψ .

Queries may contain **inequalities**: by inequality we mean an atomic formula of the form $x \neq x'$, where x and x' are variables or constants. We think that \neq is a binary relation symbol which, for each structure D, is interpreted in D as the relation $(V_D \times V_D) \setminus \{[s, s] : s \in V_D\}$.

For two structures D and D' by Hom(D, D') we denote the set of all **homomorphisms** from D to D'. Note that if a is a constant of the language and $h \in Hom(D, D')$ is a homomorphism then h(a) = a. We always use h and g to denote homomorphisms.

If ψ is a query and *D* is a structure then $\psi(D)$ denotes the result of the application of ψ to *D*. Since ψ is boolean, one would intuitively think that the result can only be YES or NO, but since we consider the multiset (bag) semantics in this paper, this YES can be repeated any positive natural number of times, depending on the number of ways ψ can be satisfied in *D*, and the above intuition is formalized as:

$\psi(D) \stackrel{\text{df}}{=} |Hom(\psi, D)|$

⁷With the only exception for Section 1.1, where more general classes of queries are discussed, and where we do not assume that queries are boolean. To distinguish, we use the upper case Greek letters to denote such more general queries.

Notice that $\psi(D)$ is always a natural number, so in particular we can multiply it by a constant, as we do in Theorem 1. We use the notation $D \models \psi$ if the set $Hom(\psi, D)$ is nonempty.

Notice also that the above definition of $\psi(D)$ makes perfect sense for queries with inequalities. This is since if $x \neq x'$ is an inequality in ψ and if $h \in Hom(\psi, D)$ is a homomorphism then:

$$[h(x), h(x')] \in V_D \times V_D \setminus \{[s, s] : s \in V_D\}$$

which means that x and x' are indeed mapped by h to different elements of V_D .

2.2 Disjoint conjunction and query exponentiation

In our proofs we often construct conjunctive queries as conjunctions of smaller conjunctive queries. When, for two conjunctive queries ρ and ρ' , we write $\rho \wedge \rho'$, we think that the conjunction of the quantifier-free parts of ρ and ρ' is taken first, and then the result is existentially quantified.

If we want to treat the variables in ρ and ρ' as local (which amounts to assuming that the existential quantification came first, and conjunction later) we write $\rho \bar{\wedge} \rho'$ instead of $\rho \wedge \rho'$.

Obviously:

Lemma 1. For each D, ρ and ρ' it is: $(\rho \wedge \rho')(D) = \rho(D)\rho'(D)$.

It is equally obvious that the lemma would not be true if, in its statement, we replaced $\overline{\wedge}$ with \wedge . The symbol $\overline{\wedge}$ relates to $\overline{\wedge}$ as \wedge relates to \wedge :

Definition 2. For $k \in \mathbb{N}$ by $\theta \uparrow k$ we denote the query $\overline{\bigwedge}_{1 \leq i \leq k} \theta$.

Clearly, for any database D, query θ , and number $k \in \mathbb{N}$ we have: $(\theta \uparrow k)(D) = (\theta(D))^k$.

2.3 Short remark about the role of constants

As we said in Section 2.1, all the queries we consider in this paper are boolean, but we allow for constants in the language. Can we somehow get rid of the constants, possibly allowing non-boolean queries instead?

Imagine ϕ_s and ϕ_b , boolean queries, with some tuple **a** of constants. And let ϕ'_s and ϕ'_b be syntactically the same queries, but now **a** is understood to be a tuple of variables. Free variables. Then the observation is (and excuse us if it is too obvious) that: ϕ_b contains ϕ_s if and only if ϕ'_b contains ϕ'_s .

This is true for any semantics (set or multiset). And this is also true if **a** are not *all* the constants that occur in ϕ_s and ϕ_b but only some of them.

With the above observation in mind let us see what happens with our results if we ban constants. Since the non-triviality condition is important for us, and to express this condition we need constants, there are two possible versions of such ban: soft one, where all constants *except for* σ^3 and φ are disallowed, and hard, where σ^3 and φ are disallowed too.

In the soft version, Theorems 1 and 3 survive almost intact (after one defines, in the natural way, what $\mathfrak{c} \cdot M$ means, for a natural number \mathfrak{c} and a multiset M). In the hard version, Theorem 3 survives, but with the additional inequality $rightarrow q \mathfrak{p}$ in the *s*-query.

3 How to multiply. From Theorem 1 to Theorem 3.

In this section we show how Theorem 3 follows from Theorem 1 (as stated in Section 1.2). To this end, suppose we have given ϕ_s and ϕ_b , like in Theorem 1, both CQs without inequality, and a natural number c.

Definition 3. For a rational number q > 0 we say that CQs (with or without inequality) ρ_s and ρ_b multiply by q if:

- (=) there exists a non-trivial database D such that $\rho_s(D) = q\rho_b(D) \neq 0$;
- (\leq) for each non-trivial database D it holds that $\rho_s(D) \leq q \rho_b(D)$.

In order to prove Theorem 3 it will now be enough to construct conjunctive queries α_s (without inequalities) and α_b (with at most one inequality) which multiply by c.

Indeed, suppose we have such α_s and α_b , whose schema is disjoint from the schema of ϕ_s and ϕ_b . Define ψ_s as $\alpha_s \bar{\wedge} \phi_s$ and ψ_b as $\alpha_b \bar{\wedge} \phi_b$. Then it is not hard to notice that the following two conditions are equivalent: (i) there exist a non-trivial database D, such that $\mathfrak{c}\phi_s(D) > \phi_b(D)$; (ii) there exist a non-trivial database D, such that $\psi_s(D) > \psi_b(D)$.

Indeed, for the proof that (i) \Rightarrow (ii), let D_1 be such a database, over the schema of ϕ_s and ϕ_b , that $\mathfrak{c}\phi_s(D_1) > \phi_b(D_1)$ and let D_2 be such a database, over the schema of α_s and α_b , that $\alpha_s(D_2) = \mathfrak{c}\alpha_b(D_2)$. Let also $D = D_1 \cup D_2$. Then:

$$\mathfrak{C}\psi_{\mathfrak{s}}(D) = (\mathfrak{C}\phi_{\mathfrak{s}}(D_1))\alpha_{\mathfrak{s}}(D_2) > \phi_{\mathfrak{b}}(D_1)(\mathfrak{C}\alpha_{\mathfrak{b}}(D_2)) = \mathfrak{C}\psi_{\mathfrak{b}}(D)$$

In order to see the $\neg(i) \Rightarrow \neg(ii)$ implication, suppose that for each non-trivial *D* it holds that $\mathbb{c}\phi_s(D) \le \phi_b(D)$. We also know that for each non-trivial *D* the inequality $\alpha_s(D) \le \mathbb{c}\alpha_b(D)$ holds. This implies that for each *D*:

$$(\mathbb{C}\phi_s(D))\alpha_s(D) \le \phi_b(D)(\mathbb{C}\alpha_b(D))$$

which means that $\psi_s(D) \leq \psi_b(D)$.

In the next two subsections we will construct α_s and α_b , as specified above. But before we go there, notice that:

Lemma 4. Suppose ϱ_s and ϱ_b multiply by some \mathfrak{q} and ϱ'_s and ϱ'_b multiply by some \mathfrak{q}' . Assume also that the schema of ϱ_s and ϱ_b is disjoint from the schema of ϱ'_s and ϱ'_b . Then $\varrho_s \wedge \varrho'_s$ and $\varrho_b \wedge \varrho'_b$ multiply by $\mathfrak{q}\mathfrak{q}'$.

3.1 The workhorse: queries β_s and β_b

As we said above, the plan is now to construct conjunctive queries, α_s (with no inequalities) and α_b (with a single inequality) which multiply by c. But is it possible at all⁸ for such queries to multiply by a number greater than 1? And (if so), is it possible for such queries to multiply by an arbitrarily huge number? This is what this subsection is about.

Let *R* be a new relational symbol of some arity $p \ge 3$. Define query *CYCLIQ*(x_1, x_2, \ldots, x_n) as:

$$R(x_1, x_2, \dots, x_p) \wedge R(x_2, \dots, x_p, x_1) \wedge \dots \wedge R(x_p, x_1, \dots, x_{p-1})$$

In this subsection we will frequently need to talk about p-tuples (of variables, or of elements of some structure), and special attention will be paid to the first element of such tuple. For this reason, we will use an overline arrow and overline bar to denote tuples of p - 1 elements. We will write x_1, \vec{x} instead of $x_1, x_2, \ldots x_p$ while by \bar{s} we will mean a tuple [$s, s, \ldots s$] of length p - 1.

Now, we can use these new notations to define β_s as:

$$CYCLIQ(x_1, \vec{x}) \land CYCLIQ(y_1, \vec{y}) \land CYCLIQ(\Diamond, \Diamond) \land CYCLIQ(\varsigma^*, \Diamond)$$

and to define β_b as $CYCLIQ(x_1, \vec{x}) \land CYCLIQ(y_1, \vec{y}) \land x_1 \neq y_1$.

The main lemma of this subsection is:

⁸Let us remark here, that a pair of conjunctive queries without inequality cannot multiply by a number greater that 1.

Lemma 5. β_s and β_b multiply by $\frac{(p+1)^2}{2p}$

The proof of Lemma 5 occupies the rest of this subsection.

To see that the condition (=) of Definition 3 is satisfied for β_s and β_b , take *D* as the canonical structure of the query $CYCLIQ(\varphi, \overline{\varphi}) \wedge CYCLIQ(\sigma^*, \overline{\varphi})$. The active domain of this *D* is of size 2, and it is an easy exercise to verify that in this case $\beta_s(D) = (p + 1)^2$ and $\beta_b(D) = 2p$.

For the proof of condition (\leq) from Definition 3 consider some non-trivial database *D*, which will remain fixed throughout this proof. We need to show that:

$$\beta_b(D) \ge \frac{2\mathrm{p}}{(\mathrm{p}+1)^2} \beta_s(D) \qquad (*)$$

At first, notice that the inequality is trivially satisfied if $\beta_s(D) = 0$. We may thus consider only the case when $\beta_s(D) > 0$. In such situation (*) is equivalent to the inequality:

$$\frac{\beta_b(D)}{\beta_s(D)} \ge \frac{2\mathrm{p}}{(\mathrm{p}+1)^2} \tag{**}$$

Before we can proceed, we need a series of definitions:

Definition 6. • We call a tuple $[s_1, \vec{s}]$ of elements of D a cyclique if $D \models CYCLIQ(s_1, \vec{s})$. • For a cyclique $[s_1, \vec{s}]$, for a natural number $0 \le k < p$, and for a tuple $[t_1, \vec{t}]$ we will say that $[t_1, \vec{t}]$ is a cyclic k-shift of $[s_1, \vec{s}]$ if $\forall_{1 \le i \le n} s_i = t_{((i+k) \mod n)+1}$.

• We write $[s_1, \vec{s}] \approx [t_1, \vec{t}]$ if there exists k such that $[t_1, \vec{t}]$ is a cyclic k-shift of $[s_1, \vec{s}]$.

It is of course easy to see that if $[s_1, \vec{s}] \approx [t_1, \vec{t}]$ then $[t_1, \vec{t}]$ is also a cyclique, and that \approx is an equivalence relation on cycliques. The equivalence class of cyclique *C* with respect to \approx will be denoted as *cyclass*(*C*).

Note that each cyclass has at least one and at most p elements. We are going to consider three kinds of cycliques:

Definition 7. • We say that a cyclique C is homogeneous, if |cyclass(C)| = 1 (note that it holds when $C = [s, \bar{s}]$ for some element s of D).

- For a non-homogeneous cyclique C, we say that C is degenerate if |cyclass(C)| < p.
- A cyclique which is neither homogeneous nor degenerate will be called normal.

Note that the above definition does not depend on the choice of the representative of the equivalence class of \approx , so we can also speak about cyclasses being homogeneous, degenerate or normal. As an example of the new notions, notice also that $[\varphi, \overline{\varphi}]$ is a homogeneous cyclique and $[\sigma, \overline{\varphi}]$ is a normal one.

The proof of the next lemma, which will later be useful, is an easy exercise in elementary group theory:

Lemma 8. If C is a degenerate cyclique then $|cyclass(C)| \le p/2$.

Now we are ready to come back to the proof of Lemma 5.

Inequality (**) begs for a probabilistic interpretation. Consider an experiment in which we draw randomly two cycliques $[s_1, \vec{s}]$ and $[t_1, \vec{t}]$ from D (with repetitions, independently and uniformly). Denote by diff the event that $s_1 \neq t_1$. Then $\mathbb{P}[\text{diff}] = \beta_b(D)/\beta_s(D)$, so to show that (**) is true, it will be sufficient to show that $\mathbb{P}[\text{diff}] \ge 2n/(n+1)^2$. For two sets *X*, *Y* of cycliques by A(X, Y) we will denote the event that one of the two drawn cycliques comes from *X* and another from *Y* (in any order). Denote by *H* the set of all homogeneous cycliques, and let $G = cyclass([\sigma^a, \bar{\varphi}])$. We know that *H* at least contains $[\varphi, \bar{\varphi}]$.

The main milestone in the proof of Lemma 5 is:

Lemma 9. Each of the following conditions (a)-(d) implies that $\mathbb{P}[\text{diff} | A(X, Y)] \ge 2\mathfrak{p}/(\mathfrak{p}+1)^2$:

(a) Y is any cyclass and X is a degenerate cyclass;

(b) $X = Y = G \cup H;$

(c) X and Y are two distinct normal cyclasses;

(**d**) *X* is a normal cyclass other than *G* and $Y = X \cup H$.

Indeed, notice that every possible pair of cycliques falls into exactly one of the events mentioned in Lemma 9 (\mathbf{a})-(\mathbf{d}). So, once Lemma 9 is proved, Lemma 5 will follow by a trivial application of the Law of Total Probability.

Proof of Lemma 9:

(a) Suppose a cyclique $C = [s_1, \vec{s}]$ from Y was picked and now we are about to pick a cyclique C' from X. Since X is a degenerate cyclass (and not a homogeneous one), there is certainly at least one cyclique $[t_1, \vec{t}]$ in X satisfying $t_1 \neq s_1$. And, by Lemma 8, there are at most p/2 cycliques in X. Hence, for each $C \in Y$, the probability that [C, C'] will be in diff is at least 2/p, which is more than $2p/(p + 1)^2$.

(**b**) First consider the case that |H| = 1, which means that $[\varphi, \overline{\varphi}]$ is the only homogeneous cyclique. Then there are $(p + 1)^2$ pairs in A(X, Y) and among them there are 2n pairs in diff. So the inequality from Lemma 9 is true in this case (and, in this case, turns into the equality from condition (=)).

Now let us consider the situation when |H| > 1. Suppose some $[s, \bar{s}]$ for $s \neq \varphi$ was picked, from H, as one element of the pair. And let us now pick a cyclique $C = [t_1, \bar{t}]$ as the other element of this pair. Then, if $C \in H$ then the probability that $t_1 \neq s$ is at least 1/2 (because there are at least 2 elements in H) and if $C \in G$ then this probability is 1 (if $s \neq \sigma^3$) or p - 1/p (if $s = \sigma^3$), in both cases more than $2p/(p + 1)^2$.

(c) Suppose $X = cyclass([t_1, \vec{t}])$ and $Y = cyclass([s_1, \vec{s}])$. Now, if for each t_i there exist at least two elements in the tuple $[s_1, \vec{s}]$ which are different than t_i then $\mathbb{P}[\text{diff} | A(X, Y)] \ge 2/p > 2p/(p + 1)^2$ and we are done. Analogously, if for each s_i there are at least two elements of the tuple $[t_1, \vec{t}]$ which are different than s_i then we are done.

So suppose there is a t_i such that all elements of $[s_1, \vec{s}]$ but one are equal to t_i and that there is s_j such that all elements of $[t_1, \vec{t}]$ but one are equal to s_j . Now, if $t_i \neq s_j$ then among the p^2 possible pairs (we imagine that first we pick an element from X and then from Y) there are at least $(p-1)^2$ pairs in diff, a fraction much greater than $2p/(p+1)^2$. If $t_i = s_j$ then there are at least 2p - 2 pairs in diff, and $2p - 2/p^2 > 2n/(p+1)^2$.

(d) There are p(p + 2|H|) possible pairs in A(X, Y). This number includes p^2 pairs which do not involve a cyclique from H and among them there are at least 2p - 2 pairs in diff (like in the proof of (c)). There are also p|H| pairs which involve a cyclique from H as the first element and p|H| pairs which involve a cyclique from H as the first element and p|H| pairs which involve a cyclique from H as the first element and p|H| pairs which involve a cyclique from H as the second element.

If |H| = 1 then among the 2p pairs in A(X, Y) which involve a cyclique from H there are at least 2 in diff. So, in this case, $\mathbb{P}[\text{diff} | A(X, Y)] \ge 2p/p(p+2)$, which is greater than $2p/(p+1)^2$.

If |H| > 1 then for every cyclique $[t_1, \vec{t}]$ from X there are at least 2|H| pairs in A(X, Y) which involve $[t_1, \vec{t}]$ as one of the elements and a cyclique from H as the other one. At least 2(|H| - 1) of them are in diff. So, in total, there are at least p(2(|H| - 1)) such pairs in diff, and, in this case:

$$\mathbb{P}[\text{diff} \mid A(X, Y)] \ge \frac{2p - 2 + p(2(|H| - 1))}{p(p + 2|H|)} \ge \frac{2p}{(p + 1)^2}$$

This ends the proof of Lemma 9 and Lemma 5

3.2 Constructing α'_s and α_b : fine tuning

The queries β_s and β_b from the previous subsection do a good job multiplying by arbitrarily huge numbers. There is a slight problem however: they only can multiply by numbers of the form $(p + 1)^2/2p$, and there is no way to find, for an arbitrary natural number c, a natural number p such that $(p + 1)^2/2p = c$.

Some fine tuning is needed: we will construct queries γ_s and γ_b which will multiply by something slightly less than 1, and then we will define $\alpha_s = \beta_s \bar{\wedge} \gamma_s$ and $\alpha_b = \beta_b \bar{\wedge} \gamma_b$ and use Lemma 4.

Let us remark here that, for reasons which will become clear in Section 5, we are not able to multiply by anything bigger than 1, without using an inequality in the *b*-query. And we cannot afford having inequality in γ_b , because we only want to have one in α_b and there already was one in β_b .

But, as it turns out, multiplication by a number smaller than 1 does not require inequality in γ_b .

For a unary relational symbol U, and for some fixed new relational symbol P, of arity m define formula $CYCLIQ_U(x_1, x_2, ..., x_m)$ as:

$$P(x_1, x_2, \dots, x_m) \land P(x_2, \dots, x_m, x_1) \land \dots \land P(x_m, x_1, \dots, x_{m-1}) \land U(x_1) \land U(x_2) \land \dots \land U(x_m)$$

Like in Section 3.1 we will be using the notations \bar{s} (to denote a tuple of identical elements, this time of length m - 1), and \vec{x} (for $x_2, x_3, ..., x_m$). Let now A and B be two new unary relation symbols. Define γ_s as $\gamma'_s \wedge \gamma''_s$ where:

$$\gamma'_{s} = CYCLIQ_{A}(\sigma, \bar{\varphi}) \land B(\sigma) \qquad \qquad \gamma''_{s} = CYCLIQ_{B}(x_{1}, \vec{x}) \land A(x_{1})$$

And define γ_b as $\gamma'_b \wedge \gamma''_b$ where:

$$\gamma'_{h} = CYCLIQ_{A}(y_{1}, \vec{y}) \land B(y_{1})$$
 $\gamma''_{h} = CYCLIQ_{B}(x_{1}, \vec{x})$

Lemma 10. The queries γ_s and γ_b multiply by $\frac{m-1}{m}$.

Proof: Let us start from condition (=) of Definition 3. We need to construct a structure D such that:

$$\gamma_s(D) = \frac{m-1}{m} \gamma_b(D) \neq 0$$

So take D as a disjoint union of the canonical structure of γ'_s and of the canonical structure of the query:

$$CYCLIQ_B(x_1, \vec{x}) \land A(x_1) \land A(x_2) \dots \land A(x_{m-1})$$

Now one can easily verify that $\gamma'_s(D) = 1$ and $\gamma'_b(D) = 1$ (notice that the last subscript in the query above is **not** m but m - 1) while $\gamma''_s(D) = m - 1$ and $\gamma''_b(D) = m$.

Proving that condition (\leq) of Definition 3 holds true for γ_s and γ_b is a little bit more complicated. Recall that we need to show that for each non-trivial *D* there is $\gamma_s(D) \leq \frac{m_j-1}{m_j} \gamma_b(D)$. So let now *D* be some fixed non-trivial database, with $\gamma_s(D) \neq 0$.

Analogously to Section 3.1 a tuple satisfying the query $CYCLIQ_U$ will be called a *U-cyclique*. Notice that the concepts we introduced in Section 3.1 survive in this new context: a cyclic *k*-shift of a *U*-cyclique is again a *U*-cyclique, so one can again consider the equivalence relation \approx and cyclasses.

For a unary relation V we will also use the term U-cyclique^V for such a U-cyclique $[s_1, \vec{s}]$ for which $V(s_1)$ also holds.

First observation we can make is that $\gamma''_s(D) \leq \gamma''_b(D)$: this is because $\gamma''_b(D)$ is the cardinality of the set of all *B*-cycliques in *D*, and $\gamma_s''(D)$ is the cardinality of its subset: the set of all *B*-cycliques^A.

Notice also that $\gamma'_s(D) = 1$, because it only mentions constants. So if $\gamma'_b(D) \ge 2$ then $\gamma_s(D) \le \frac{\gamma_b(D)}{2}$ and we are done.

So, from now on we assume that $\gamma'_{b}(D) = 1$. This means that there is exactly one A-cyclique^B in D. And we already know one such A-cyclique: it is $[\sigma, \overline{\varphi}]$.

What remains to be proved is that $\gamma_s''(D) \le \frac{m-1}{m} \gamma_b''(D)$. Now, using the language of probabilities again, let \mathcal{A} be the event that a randomly (uniformly) picked *B*-cyclique is a *B*-cyclique^{*A*}. We need to show that $\mathbb{P}[\mathcal{A}] \leq \frac{n-1}{n_j}$. To this end it will be enough to show, for each *B*-cyclique *C*, that $\mathbb{P}[\mathcal{A} \mid cyclass(C)] \leq \frac{m-1}{m}$ (this is because the cyclasses constitute a partition of the set of all *B*-cycliques). But, for each *B*-cyclique *C* it holds that $|cyclass(C)| \leq m$. So (\leq) will be proven once we show that for each B-cyclique C there exists a B-cyclique C', which is not a B-cyclique^A, and such that $C' \approx C.$

So suppose, towards contradiction, that there is some C such that all B-cycliques in cyclass(C) are B-cycliques^A. This would imply, that all elements of C satisfy A and, in consequence, all cycliques in cyclass(C) would also be A-cycliques^B. Recall that we assumed that there is only one A-cyclique^B in D so cyclass(C) must be a singleton, with $C = [s, \bar{s}]$ for some s. But, because of the same assumption, $[s, \bar{s}]$ must be equal to $[\sigma, \overline{\varphi}]$, which cannot be true in a non-trivial database.

Now take p = 2c - 1 and m = p + 1 and put $\alpha_s = \beta_s \bar{\wedge} \gamma_s$ and $\alpha_b = \beta_b \bar{\wedge} \gamma_b$. Notice that:

$$\frac{(p+1)^2}{2p} \frac{m-1}{m} = \frac{(p+1)^2}{2p} \frac{p}{p+1} = \frac{p+1}{2} = \alpha$$

So, indeed, by Lemma 4, α_s and α_b multiply by c. This ends the proof of Theorem 3 (using Theorem 1).

Proof of Theorem 1 4

In this section we prove Theorem 1, stated in Section 1.2.

The source of undecidability 4.1

As the source of undecidability we are going to use;

Lemma 11. The problem:

Given are a natural number $\mathcal{C} \geq 2$ and two polynomials, P_s and P_b , of numerical⁹ variables x_1, x_2, \ldots, x_m . with natural coefficients, which additionally satisfy the following conditions:

• $P_s = \sum_{m=1}^{m} c_{s,m} \mathbb{T}_m$ and $P_b = \sum_{m=1}^{m} c_{b,m} \mathbb{T}_m$, where \mathbb{T}_m , for $m \in \{1, \dots, m\}$ are monomials, and $c_{s,m}, c_{b,m} \in \mathbb{N}$ are coefficients;

- there exists $d \in \mathbb{N}$ such that each monomial \mathbb{T}_m , for $m \in \{1, \dots, m\}$, is of degree exactly d;
- x_1 occurs as the first variable in each \mathbb{T}_m ;

⁹We will call the variables ranging over \mathbb{N} numerical variables to distinguish them from the first order logic variables that occur in the queries.

• $1 \leq c_{s,m} \leq c_{b,m}$ holds for each $m \in \{1, \dots, m\}$.

Does the inequality $CP_s(\Xi(\vec{x})) \leq \Xi(x_1)^d P_b(\Xi(\vec{x}))$ hold for every valuation $\Xi : \{x_1, x_2, \dots, x_n\} \to \mathbb{N}$? is undecidable.

Proof of Lemma 11 is an easy exercise (assuming, of course, undecidability of the Hilbert's 10th problem), but for the sake of completeness it will be included in the Appendix.

Notice that, in P_s and P_b , we use the *s* and *b* again, and again they stand for "small" and "big". This is how we think of the two polynomials: it is undecidable whether the "small one", multiplied by *c* can ever surpass the "big one" (strictly speaking, the monomial x_1^d is not part of P_b , but we find it convenient to imagine it is).

In order to prove Theorem 1 we will construct, for a natural number c and for two polynomials, P_s and P_b , like in Lemma 11, a tuple $[c, \phi_s, \phi_b]$, consisting of a natural number and two conjunctive queries, such that the two conditions are equivalent:

 \mathscr{X} There exist a valuation $\Xi : \{x_1, x_2, \dots, x_n\} \to \mathbb{N}$ such that: $cP_s(\Xi(\vec{x})) > \Xi(x_1)^d P_b(\Xi(\vec{x})).$

‡ There exists a non-trivial database *D* such that: $c\phi_s(D) > \phi_b(D)$.

Notice that the c in the output tuple $[c, \phi_s, \phi_b]$ of our reduction is not the c in the input tuple $[c, P_s, P_b]$.

4.2 High-level definition of ϕ_s and ϕ_b

In Sections 4.3-4.6 we are going to define queries *Arena*, π_s , π_b , ζ_b and δ_b . Then ϕ_s will be defined as *Arena* $\bar{\wedge} \pi_s$ and ϕ_b will be defined as $\pi_b \bar{\wedge} \zeta_b \bar{\wedge} \delta_b$.

In Section 4.7 we will show that the equivalence $\Re \Leftrightarrow \clubsuit$ holds for such ϕ_s and ϕ_b , and for the number \mathfrak{c} as defined in Section 4.5.

4.3 Introducing the queries π_s and π_b .

We assume that polynomials P_s and P_b are like in Lemma 11, so in particular they have the same monomials $\mathbb{T}_1, \mathbb{T}_2, \ldots, \mathbb{T}_m$, all of them of degree d. Recall that coefficients before $\mathbb{T}_1, \mathbb{T}_2, \ldots, \mathbb{T}_m$ in P_s are (respectively) $c_{s,1}, c_{s,2}, \ldots, c_{s,m}$ and the coefficients before $\mathbb{T}_1, \mathbb{T}_2, \ldots, \mathbb{T}_m$ in P_b are (respectively) $c_{b,1}, c_{b,2}, \ldots, c_{b,m}$.

Recall also that for each $1 \le m \le m$ we have $1 \le c_{s,m} \le c_{b,m}$.

Notice that we use *m* as a natural number ranging from 1 to m. In a similar manner we will always use *n* as a natural number ranging from 1 to n (recall that n is the number of variables in P_s and P_b) and *d* as a natural number ranging from 1 to d. Also, define $\mathbb{I} = m + m + 2$.

Let Σ_0 be the schema comprising a binary relation S_m for each $m \in \{1, ..., m\}$ (so that we have one relation S for each monomial in P_s and P_b), a binary relation R_d for each $d \in \{1, ..., d\}$, and a binary relation E. And let $\Sigma = \Sigma_0 \cup \{X\}$ for some binary relation X.

Now, define the query π_s as:

$$\bigwedge_{m \in \{1,\dots,m\}} S_m(x,x) \wedge S_m(x,x_{c_{s,m}}^m) \wedge \bigwedge_{m \in \{1,\dots,m\}} \bigwedge_{1 \le k < c_{s,m}} S_m(x_{k+1}^m,x_k^m) \wedge \bigwedge_{d \in \{1,\dots,d\}} R_d(x,y_d) \wedge X(y_d,z_d)$$

and let π_b be:

$$\bigwedge_{m \in \{1, \dots, m\}} S_m(x, x) \wedge S_m(x, x_{c_{b,m}}^m) \wedge \bigwedge_{m \in \{1, \dots, m\}} \bigwedge_{1 \le k < c_{b,m}} S_m(x_{k+1}^m, x_k^m) \wedge \bigwedge_{d \in \{1, \dots, d\}} R_d(x, y_d) \wedge X(y_d, z_d) \wedge \bigwedge_{d \in \{1, \dots, d\}} R_1(x, y_d') \wedge X(y_d', z_d')$$

It will be helpful at this point to get some understanding of the structure of π_b : the (canonical structure of the) query is a star, with x as its center. For each monomial \mathbb{T}_m in P_b there is an " S_m -ray" in this star (radiating from the center, as the rays normally do) of length equal to the coefficient before \mathbb{T}_m in P_b . There is also an S_m -loop in x which is very important, as you are soon going to see.

Apart from the " S_m -rays" there are also rays of length two, 2d of them, as many as there are numerical variable occurrences in each term of $x_1^d P_b$. Each such ray begins with R_d (this *d* indicates which numerical variable we have in mind) and then comes the *X* which is supposed (as you are going to see in Section 4.4) to represent the valuation Ξ . The fact that all the "rays" representing x_1^d begin from R_1 is related to the third condition from Lemma 11.

The structure of π_s relates to P_s as π_b relates to P_b .

Lemma 12. For every database D there is $\pi_s(D) \leq \pi_b(D)$.

Proof: Let us start from a simple observation. Suppose for some queries ρ_b and ρ_s there exists an *onto* mapping *h* from the variables of ρ_b to the variables of ρ_s which is a homomorphism of queries. Then for every database *D* there is $\rho_s(D) \le \rho_b(D)$.

Indeed, if *h* is a mapping as in the previous paragraph, then the function *H* (of argument *g*) defined as $H(g) = g \circ h$ is a 1-1 function from $Hom(\rho_s, D)$ to $Hom(\rho_b, D)$. Obviously, if $g \in Hom(\rho_s, D)$ then $g \circ h \in Hom(\rho_b, D)$. To see that *H* is 1-1, take $g, g' \in Hom(\rho_s, D)$ such that $g \neq g'$. Then there must be v_s , a variable in ρ_s , such that $g(v_s) \neq g'(v_s)$. Let v_b be such a variable of ρ_b that $h(v_b) = v_s$ (recall that *h* is onto). Then $gh(v_b) \neq g'h(v_b)$.

What remains to be shown is that there exists an *onto* homomorphism h from π_b to π_s .

To this end, first of all notice that $Var(\pi_s) \subseteq Var(\pi_b)$. Let us first define *h* on the variables which are both in $Var(\pi_s)$ and in $Var(\pi_b)$ as the identity:

$$h \upharpoonright Var(\pi_s) = id_{Var(\pi_s)}$$

At this point we can already be sure that *h* will be *onto*. Now we need to define its values for the variables in $Var(\pi_b) \setminus Var(\pi_s)$ in such a way that the resulting *h* is indeed a homomorphism.

For each variable $x_k^m \in Var(\pi_b) \setminus Var(\pi_s)$ define $h(x_k^m) = x$. See how, thanks to the S_m -loops at x, the S_m -rays in π_b are now homomorphically mapped onto the S_m -rays in π_s (this is the only place in the entire paper where we use the condition, from Lemma 11, that the coefficients in P_b are at least equal to the respective coefficients in P_s).

What still remains to be defined are the values of *h* for the variables (other than *x*) in the subquery $\bigwedge_{d \in \{1,...,d\}} R_1(x, y'_d) \land X(y'_d, z'_d)$. It is not hard to guess that *h* will map all the *y*'s to *y*₁ and it will map all the *z*'s to *z*₁.

4.4 Arena, and how π_s and π_b compute P_s and P_b .

The query $Arena = Arena_{\pi} \wedge Arena_{\delta}$ over Σ_0 will be defined as a conjunction of facts mentioning only constants (and no variables). Because of that, for every database D we will have $Arena(D) \in \{0, 1\}$: the value is 1 if all the atomic formulas in *Arena* are indeed facts in D and it is 0 otherwise.

Definition of the query $Arena_{\pi}$ will follow now and $Arena_{\delta}$ will be defined in Section 4.6, together with the query δ_b .

Let $\mathcal{P} \subseteq \{1, \dots, m\} \times \{1, \dots, m\}$ be the relation saying which numerical variable constitutes which argument in which of the monomials: $\mathcal{P}(n, d, m)$ means that x_n is the *d*-th variable¹⁰ in \mathbb{T}_m .

Now define *Arena*^{π} as the query:

$$\bigwedge_{[n,d,m]\in\mathcal{P}} R_d(a_m,b_n) \wedge \bigwedge_{m,m'\in\{1,\dots,m\}} S_{m'}(a_m,a_m) \wedge \bigwedge_{m\in\{1,\dots,m\}} S_m(a_m,a) \wedge S_m(a,a)$$

Notice that we have one constant (a_m) for each monomial and one constant (b_n) for each numerical variable. As we said, $Arena_{\delta}$ is not going to be defined right now. All you need to know at this point is that the only relation $Arena_{\delta}$ mentions is E, which does not appear in $Arena_{\pi}$.

Let D_{Arena} be the canonical structure of the query Arena.

Definition 13. A database D over Σ , such that $D \models$ Arena, will be called:

• correct if $D \upharpoonright \Sigma_0 = D_{Arena}$, where by $D \upharpoonright \Sigma_0$ we mean the database resulting from D by removing from it all atoms of the relation X;

• slightly incorrect *if it is not correct but* $D \upharpoonright \Sigma_0 \supseteq D_{Arena}$, where \supseteq *is understood to be inclusion of relational structures;*

• seriously incorrect if it is neither correct nor slightly incorrect.

In other words, a correct database is just D_{Arena} with some additional atoms of relation X, a slightly incorrect database is D_{Arena} with some additional atoms of relation X and possibly also of the relations from Σ_0 , and a seriously incorrect database is one which satisfies *Arena* (that is, contains a homomorphic image of D_{Arena}), but which identifies some elements of D_{Arena} (that is, this homomorphism is not 1-1).

As we already mentioned, the relation X will represent a valuation of numerical variables:

Definition 14. For a database D, over the schema Σ , such that $D \models$ Arena let us define a valuation Ξ_D : $\{x_1, x_2, \ldots x_n\} \rightarrow \mathbb{N}$ in the following way:

$$\Xi_D(\mathbf{x}_i) = (\exists x \ X(b_i, x))(D)$$

Translating it to human language, notice that $\exists x \ X(b_i, x)$ is a boolean query, so – when applied to D – it returns a number, which is the number of X-edges that begin at b_i .

Proof of the following lemma can be found in Appendix A:

Lemma 15. If D is a correct database, then $\pi_s(D) = P_s(\Xi_D(\vec{x}))$ and $\pi_b(D) = \Xi_D(x_1)^d P_b(\Xi_D(\vec{x}))$.

Clearly, for every valuation $\Xi : \{x_1, x_2, \dots, x_n\} \to \mathbb{N}$, there exists a correct database *D* such that $\Xi = \Xi_D$. This, together with Lemma 15, implies:

Lemma 16. *The following two conditions are equivalent:*

• there exists a valuation $\Xi : \{x_1, x_2, \dots, x_n\} \to \mathbb{N}$ such that $CP_s(\Xi(\vec{x})) > \Xi(x_1)^d P_b(\Xi(\vec{x}))$

• there exists a correct database D such that $c\pi_s(D) > \pi_b(D)$.

¹⁰If variable x_n occurs in \mathbb{T}_m more than once then $\mathcal{P}(n, d, m)$ will be true for more than one number *d*.

4.5 How to punish for slight incorrectness

Lemma 16 almost looks like the $\Re \Leftrightarrow \Leftrightarrow$ equivalence which we are trying to prove. Almost, because it only works for correct databases. And we have no idea how the values of $\pi_s(D)$ and $\pi_b(D)$ would behave if we were given a slightly incorrect database, or a seriously incorrect one. In Sections 4.5 and 4.6, we create tools for "punishing" such databases. The tools will be then used in Section 4.7.

Let $\Sigma_{RS} = \{S_1, \dots, S_m, R_1, \dots, R_d\}$. For a relation symbol $P \in \Sigma_{RS}$, let j^P denote the number of atoms of the relation *P* in *Arena*.

Let $j = \max(\{j^P : P \in \Sigma_{RS}\})$ and let k be the smallest natural number such that $(\frac{j+1}{j})^k \ge c$. Clearly, we also have $(\frac{j^P+1}{j_m^P})^k \ge c$ for each $P \in \Sigma_{RS}$.

Now, again for a relation symbol $P \in \Sigma_{RS}$, let ζ^P be the query:

 $P(w,v) \uparrow k$

And let ζ_b be defined as $\bar{\bigwedge}_{P \in \Sigma_{RS}} \zeta^P$. Finally, define $\varepsilon_1 = \zeta_b(D_{Arena})$ and $\varepsilon = c \varepsilon_1$ (recall that c comes from our input tuple $[c, P_s, P_b]$, and ε is a part of the output tuple $[\varepsilon, \phi_s, \phi_b]$). Then of course:

Lemma 17. • If D is correct then $\zeta_b(D) = \mathfrak{c}_1$.

• If $D \models Arena then \zeta_b(D) \ge 1$.

The second claim of the above lemma follows since, in order for *Arena* to be satisfied in *D*, there must be at least one atom in *D* of each of the relations in Σ_{RS} . Finally:

Lemma 18. If *D* is slightly incorrect then $\zeta_b(D) \ge \mathbb{C}$

Proof: Clearly, for any database *D* we have:

$$\zeta_b(D) = \prod_{P \in \Sigma_{PS}} \zeta^P(D) \qquad (*)$$

If *D* is slightly incorrect then, by definition of slight incorrectness:

(a) for each $P \in \Sigma_{RS}$ there are at least as many atoms of P in D as in D_{Arena} ;

(b) there is $P_0 \in \Sigma_{RS}$ such that there are more atoms of P_0 in D then in D_{Arena} .

From (a) we get that $\zeta^P(D) \ge \zeta^P(D_{Arena})$ for each $P \in \Sigma_{RS}$. From (b) we get that there are at least $j^{P_0} + 1$ atoms of the relation P_0 in D. It implies that $\zeta^{P_0}(D)/\zeta^{P_0}(D_{Arena}) \ge c$. So from (*), from (a) and from (b) we get that $\zeta_b(D) \ge c\zeta_b(D_{Arena}) = c$.

4.6 How to punish for serious incorrectness

Recall that the query $Arena_{\pi}$ mentions constants $a, a_1, a_2, \dots a_m$ and $b_1, b_2, \dots b_n$ and that $E \in \Sigma_0$ is a binary relation symbol which was not used so far. It is now time to define $Arena_{\delta}$, as the query:

 $E(\mathfrak{S}^{\flat},\mathfrak{S}^{\flat}) \wedge E(\mathfrak{Q},a) \wedge E(a,a_{1}) \wedge E(a_{1},a_{2}) \wedge \ldots \wedge E(a_{\mathfrak{m}-1},a_{\mathfrak{m}}) \wedge E(a_{\mathfrak{m}},b_{1}) \wedge E(b_{1},b_{2}) \wedge \ldots \wedge E(b_{\mathfrak{n}-1},b_{\mathfrak{n}}) \wedge E(b_{\mathfrak{n}},\mathfrak{Q})$

In words, *Arena*_{δ} comprises the self-loop $E(\sigma^2, \sigma^2)$ and an *E*-cycle, of length \mathbb{I} (recall that $\mathbb{I} = n + m + 2$) containing φ and all the constants from *Arena*_{π}.

For a natural number *l* let $\delta_{b,l}$ be the query:

$$E(z_1, z_2) \wedge E(z_2, z_3) \wedge \ldots E(z_{l-1}, z_l) \wedge E(z_l, z_1)$$

Let $L = \{1, 2, \dots, \|-1\} \cup \{\|+1\}$. Define δ_b as $(\overline{\bigwedge}_{l \in L} \delta_{b,l}) \uparrow \mathbb{C}$.

In human language the meaning of $\delta_b(D)$ is as follows: for each $l \in L$ count all (the homomorphic images of) cycles of length l in D, take a product of all the |L| numbers you got and raise this product to the power \mathbb{C} .

Lemma 19. Suppose D is such that $D \models Arena$. Then $\delta_b(D) \ge 1$.

Proof: Recall that if $D \models Arena$ then $E(\mathcal{A}, \mathcal{A})$ must be true in D. Then δ_b can be satisfied in D by mapping all its variables to \mathcal{A} .

Lemma 20. If D is a correct database then $\delta_b(D) = 1$.

Proof: Arena, and hence also any other correct database, contains two cycles (with respect to the relation *E*): one of length 1 (namely, $E(\sigma^2, \sigma^2)$) and one of length 1.

But *L* contains all natural numbers up to $\mathbb{I} + 1$ except for \mathbb{I} . So, for every $l \in L$, there is exactly one (homomorphic image of a) cycle of length l in *Arena* – the loop $E(\mathfrak{G}^n, \mathfrak{G}^n)$. In consequence $\delta_b(D)$ is the product of |L| ones, raised to power \mathbb{I} , and this equals 1.

Lemma 21. Suppose *D* is a seriously incorrect non-trivial database, such that $D \models Arena$. Then $\delta_b(D) \ge 2^{\mathfrak{c}} \ge \mathfrak{c}$.

Proof: Suppose D is a seriously incorrect non-trivial database, such that $D \models Arena$. We need to prove that:

$$(\bigwedge_{l\in L} \delta_{b,l})(D) \geq 2$$

Using the argument from the proof of Lemma 19 we get that for each $l \in L$ there is $\delta_{b,l}(D) \ge 1$. So what remains to be proved is that there exists $l \in L$ such that $\delta_{b,l}(D) \ge 2$. Or, in other words, that there exists $l \in L$ and a homomorphism $h : \delta_{b,l} \to D$ which **does not** map all the variables of $\delta_{b,l}$ to σ^* . Call such homomorphisms *non-trivial*.

There are two cases, depending on which constants of Arena are identified in the (seriously incorrect) D.

Case 1. D identifies σ^a with some other constant of Arena. Then there indeed exists a non-trivial homomorphism $h \in Hom(\delta_{b,l+1}, D)$. To see it first notice that $D \models Arena_{\delta}$, so there must be an *E*-cycle of length \mathbb{I} in *D*. One of the elements of this cycle is identified with σ^a , and $D \models E(\sigma^a, \sigma^a)$, so we have a cycle of length \mathbb{I} whose one vertex has a self-loop, and we can of course homomorphically embed $\delta_{b,l+1}$ (which itself is a cycle of length $\mathbb{I} + 1$) in such a cycle. But why are we sure that this homomorphism is non-trivial? This is because *D* is non-trivial, so we are sure that σ^a and φ are not identified in *D*.

Case 2. D does not identify σ^n with any other constant of Arena but it does identify some other two constants. Then some two vertices of the length- $\mathbb{I} E$ -cycle from $Arena_{\delta}$ are identified in D, which produces a shorter cycle. In consequence, there exists a non-trivial homomorphism $h \in Hom(\delta_{b,l}, D)$ for some $l < \mathbb{I}$. \Box

4.7 Putting all pieces together

What remains to be shown is that the equivalence $\mathfrak{B} \Leftrightarrow \mathfrak{A}$ from Section 11 holds for $\phi_s = Arena \overline{\wedge} \pi_s$, for $\phi_b = \pi_b \overline{\wedge} \zeta_b \overline{\wedge} \delta_b$, and for the \mathfrak{c} as in Section 4.5.

To show that $\mathfrak{B} \Rightarrow \mathfrak{A}$, suppose $\Xi : \{\mathfrak{x}_1, \mathfrak{x}_2, \dots, \mathfrak{x}_n\} \to \mathbb{N}$ is such that $\mathcal{C}P_s(\Xi(\vec{\mathfrak{x}})) > \Xi(\mathfrak{x}_1)^d P_b(\Xi(\vec{\mathfrak{x}}))$. Take a correct database D such that $\Xi_D = \Xi$. By Lemma 15 we get that $\mathcal{C}\pi_s(D) > \pi_b(D)$. Recall that Arena(D) = 1.

From Lemma 20 also $\delta_b(D) = 1$. So $c(Arena \wedge \pi_s)(D) > (\pi_b \wedge \delta_b)(D)$. We also know, from Lemma 17, that $\zeta_b(D) = c_1$. Multiplying both sides by c_1 , we get:

$$\mathcal{C}\mathbb{C}_1(Arena \wedge \pi_s)(D) > (\pi_b \wedge \zeta_b \wedge \delta_b)(D)$$

Now just recall that $c = cc_1$.

For the $\mathscr{X} \leftarrow \mathbf{\dot{x}}$ direction, suppose $CP_s(\Xi(\vec{x})) \leq \Xi(x_1)^d P_b(\Xi(\vec{x}))$ for all valuations Ξ . We want to show that in such case for every non-trivial *D* there is $c\phi_s(D) \leq \phi_b(D)$.

So fix a non-trivial database D. If $D \not\models Arena$ then $D \not\models \phi_s$ and there is nothing to prove. If $D \models Arena$ then there are three cases: D may be correct, or slightly incorrect, or seriously incorrect.

If *D* is correct then just reuse the above argument for $\Re \Rightarrow \Re$. For the remaining two cases, first recall Lemma 12: whatever *D* is, we know for sure that $\pi_s(D) \le \pi_b(D)$. We need to show that $\mathfrak{c} \le (\zeta_b \wedge \delta_b)(D)$.

Now, if *D* is slightly incorrect, then $\delta_b(D) \ge 1$ (by Lemma 19) and $\zeta_b(D) \ge c$ (by Lemma 18).

If *D* is seriously incorrect, then $\delta_b(D) \ge c$ (by Lemma 21) and $\zeta_b(D) \ge 1$ (by the second claim of Lemma 17).

This ends the proof of Theorem 1.

5 Proof of Theorem 5

In this Section we prove Theorem 5. But first we need to introduce the tools.

5.1 Operations on structures

We are going to construct new structures (from some given structures) using two standard operations on graphs (which also apply to any other relational structures): graph product and the the blow-up operation. Let us recall their definitions (see for example [17] Chapter 3.3).

The blow-up operation. For a relational structure *D*, with the set of vertices *V*, and for a natural number \hat{k} , the structure $blowup(D, \hat{k})$ is defined as follows. The set of vertices of $blowup(D, \hat{k})$ is $V_{\hat{k}} = \{[s, i] : s \in V \land i \in \{1, 2, ..., \hat{k}\}$. Then, for each relation¹¹ symbol *R*, and for each $[s, i], [r, j] \in V_{\hat{k}}$, the atom R([s, i], [r, j]) is in $blowup(D, \hat{k})$ if and only if R(s, r) is in *D*.

The product operation. For two structures D_1 and D_2 , with sets of vertices V_1 and V_2 respectively, their product $D_1 \times D_2$ is defined as a structure whose set of vertices is $V_1 \times V_2$, such that R([s, s'], [r, r']) is an atom¹² of $D_1 \times D_2$ if and only if R(s, r) is an atom of D_1 and R(s', r') is an atom of D_2 . For a structure D and natural number k by $D^{\times k}$ we denote the product of k copies of D.

The well-known and important lemma (and also easy to prove) is:

Lemma 22. Suppose D is a structure, ϕ is a CQ, without inequalities, with j variables and \hat{k} is a natural number. Then:

- (i) $\phi(blowup(D, \hat{k})) = \hat{k}^{\dagger}\phi(D)$
- (*ii*) $\phi(D^{\times \hat{k}})) = (\phi(D))^{\hat{k}}$

Notice that it follows directly from Lemma 22 (ii) that if ρ_s and ρ_b (like in Definition 3) are CQs without inequality then they cannot multiply by a number greater than 1.

¹¹In order to keep the notation light (and avoid superscripts) we imagine that R is binary, but the definition is analogous for relations of any arity.

¹²The remark from the previous footnote applies here accordingly.

5.2 **Proof of Theorem 5**

Now, Theorem 5 will follow directly from:

Lemma 23. Let ψ_s and ψ_b be conjunctive queries, ψ_b without inequalities and ψ_s with inequalities. Let ψ'_s be ψ_s with all the inequalities removed. Then the two conditions are equivalent: (a) there exist a structure D, such that $\psi_s(D) > \psi_b(D)$; (b) there exist a structure D_0 , such that $\psi'_s(D_0) > \psi_b(D_0)$;

Proof of Lemma 23: To keep the presentation simple we will show the lemma for ψ_s having exactly one inequality $x \neq x'$. Proof in the general case is similar (we will comment on it at the end of this Section).

Of course, for each *D*, there is $\psi'_s(D) \ge \psi_s(D)$ so (a) implies (b).

Now suppose there exists a structure D_0 , such that $\psi'_s(D_0) > \psi_b(D_0)$. We are going to construct D such that $\psi_s(D) > \psi_b(D)$. One lemma we will need is:

Lemma 24. For every structure D it holds that: $\psi_s(blowup(D,2)) \ge \frac{\psi'_s(blowup(D,2))}{2}$

Proof of Lemma 24: It is clear that:

 $Hom(\psi_s, blowup(D, 2)) \subseteq Hom(\psi'_s, blowup(D, 2))$

In order to prove Lemma 24 it will be enough to show that:

 $|Hom(\psi'_s, blowup(D, 2)) \setminus Hom(\psi_s, blowup(D, 2))| \le |Hom(\psi_s, blowup(D, 2))|$

Translating the above into the human language, we want to show that for each homomorphism $h \in Hom(\psi'_s, blowup(D, 2))$ which **does not** map x and x' to different vertices of blowup(D, 2) there is another such homomorphism which **does**.

To this end it will be enough to construct an injection *F* from the set $Hom(\psi'_s, blowup(D, 2)) \setminus Hom(\psi_s, blowup(D, 2))$ to the set $Hom(\psi_s, blowup(D, 2))$. In other words, it will be enough to construct a 1-1 function *F* which will produce, for each mapping (from the set of variables of ψ_s to blowup(D, 2)) satisfying ψ'_s but not ψ_s a mapping (from the set of variables of ψ_s to blowup(D, 2)) which will satisfy ψ_s .

So take a mapping *h* which is in $Hom(\psi'_s, blowup(D, 2))$ but not in $Hom(\psi_s, blowup(D, 2))$. Clearly, $h(x) = h(x') = [s, \epsilon]$ for some $s \in V_D$ and some $\epsilon \in \{1, 2\}$. And define F(h) as the mapping:

$$(F(h))(y) = \begin{cases} [s, \epsilon] & \text{if } y = x \\ [s, 3 - \epsilon] & \text{if } y = x' \\ h(y) & \text{if } y \neq x \text{ and if } y \neq x' \end{cases}$$

It is now trivial to see that the F we have just defined is indeed as required, which ends the proof of Lemma 24 \Box

Let now j be the number of variables in ψ_b . Since $\psi'_s(D_0) > \psi_b(D_0)$, and by Lemma 22 (ii), there exists k such that:

$$\psi'_{s}(D_{0}^{\times k}) > 2^{j+1}\psi_{b}(D_{0}^{\times k})$$

Now take $D = blowup(D_0^{\times \hat{k}}, 2)$. By Lemma 22 (i) we have:

$$\psi_b(D) = 2^{j} \psi_b(D_0^{\times k})$$

It then follows from the choice of k that:

$$\psi'_{s}(D) > 2\psi_{b}(D)$$

Finally, from Lemma 24 we get that:

$$\psi_s(D) > \psi_b(D)$$

Which ends the proof of Lemma 23 and of Theorem 5

If there were <u>n</u> inequalities in ψ_s rather than one, then we would use 2<u>n</u> rather than 2 in Lemma 24 and we would need to modify the \hat{k} above accordingly.

References

- S. Chaudhuri and M. Y. Vardi, "Optimization of real conjunctive queries," in *Proceedings of the Twelfth* ACM SIGACT-SIGMOD-SIGART Symposium on Principles of Database Systems, PODS '93, (New York, NY, USA), p. 59–70, Association for Computing Machinery, 1993.
- [2] A. K. Chandra and P. M. Merlin, "Optimal implementation of conjunctive queries in relational data bases," in *Proceedings of the 9th Annual ACM Symposium on Theory of Computing, May 4-6, 1977, Boulder, Colorado, USA*, pp. 77–90, ACM, 1977.
- [3] Y. Sagiv and M. Yannakakis, "Equivalences among relational expressions with the union and difference operators," J. ACM, vol. 27, no. 4, pp. 633–655, 1980.
- [4] A. C. Klug, "On conjunctive queries containing inequalities," J. ACM, vol. 35, no. 1, pp. 146–160, 1988.
- [5] R. van der Meyden, "The complexity of querying indefinite data about linearly ordered domains," *Journal of Computer and System Sciences*, vol. 54, no. 1, pp. 113–135, 1997.
- [6] Y. E. Ioannidis and E. Wong, "Towards an algebraic theory of recursion," *Journal of the ACM (JACM)*, vol. 38, no. 2, pp. 329–381, 1991.
- [7] F. N. Afrati, M. Damigos, and M. Gergatsoulis, "Query containment under bag and bag-set semantics," *Information Processing Letters*, vol. 110, no. 10, pp. 360–369, 2010.
- [8] E. V. Kostylev, J. L. Reutter, and A. Z. Salamon, "Classification of annotation semirings over query containment," in Proc. of the 31st ACM SIGMOD-SIGACT-SIGART Symposium on Principles of Database Systems, PODS 2012, Scottsdale, AZ, USA, May 20-24, 2012, pp. 237–248, ACM, 2012.
- [9] S. Cohen, "Equivalence of queries that are sensitive to multiplicities," VLDB J., vol. 18, no. 3, pp. 765– 785, 2009.
- [10] R. Chirkova, "Equivalence and minimization of conjunctive queries under combined semantics," in 15th Int. Conf. on Database Theory, ICDT '12, Berlin, Germany, March 26-29, 2012, pp. 262–273, ACM, 2012.
- [11] G. Konstantinidis and F. Mogavero, "Attacking diophantus: Solving a special case of bag containment," in Proc. of the 38th ACM SIGMOD-SIGACT-SIGAI Symposium on Principles of Database Systems, PODS 2019, Amsterdam, The Netherlands, June 30 - July 5, 2019, pp. 399–413, ACM, 2019.
- [12] S. Kopparty and B. Rossman, "The homomorphism domination exponent," *Eur. J. Comb.*, vol. 32, no. 7, pp. 1097–1114, 2011.
- [13] M. Abo Khamis, P. G. Kolaitis, H. Q. Ngo, and D. Suciu, "Bag query containment and information theory," in *Proceedings of the 39th ACM SIGMOD-SIGACT-SIGAI Symposium on Principles of Database Systems*, PODS'20, (New York, NY, USA), p. 95–112, Association for Computing Machinery, 2020.
- [14] Y. E. Ioannidis and R. Ramakrishnan, "Containment of conjunctive queries: Beyond relations as sets," *ACM Trans. on Database Systems (TODS*, 1995.

- [15] T. S. Jayram, P. G. Kolaitis, and E. Vee, "The containment problem for ibi¿reali/bi¿ conjunctive queries with inequalities," in *Proceedings of the Twenty-Fifth ACM SIGMOD-SIGACT-SIGART Symposium* on *Principles of Database Systems*, PODS '06, (New York, NY, USA), p. 80–89, Association for Computing Machinery, 2006.
- [16] J. Kwiecien, J. Marcinkowski, and P. Ostropolski-Nalewaja, "Determinacy of real conjunctive queries. the boolean case," in *PODS '22: International Conference on Management of Data, Philadelphia, PA, USA, June 12 - 17, 2022* (L. Libkin and P. Barceló, eds.), pp. 347–358, ACM, 2022.
- [17] L. Lovász, *Large Networks and Graph Limits*, vol. 60 of *Colloquium Publications*. American Mathematical Society, 2012.
- [18] M. Davis, "Unsolvable problems," in *Handbook of Mathematical Logic* (J. Barwise, ed.), pp. 567–594, North-Holland, Amsterdam, 1977.

A Proof of Lemma 15

We first deal with the equality for π_s . Suppose *D* is a correct database. In order to count the homomorphisms $h: \pi_s \to D$ let us group them by h(x), that is by the vertex of *D* the variable *x* is mapped to.

Clearly, for *h* to be a homomorphism h(x) must be one of the constants a_m for some $m \in \{1, ..., m\}$ (this is the only way the atoms of the relations R_d from π_x can possibly be satisfied in *D*). So let us fix an *m* such that $h(x) = a_m$.

The Lemma will follow once we are able to prove that:

$$|\{h \in Hom(\pi_s, D) : h(x) = a_m\}| = c_{s,m}t_m(\Xi_D(\vec{x}))$$
 (*)

Let us now split π_s into π'_s and π''_s , where:

$$\pi'_s = \bigwedge_{d \in \{1, \dots, d\}} R_d(x, y_d) \wedge X(y_d, z_d)$$

$$\pi_{s}^{\prime\prime} = \bigwedge_{m \in \{1, \dots, m\}} S_{m}(x, x) \wedge S_{m}(x, x_{c_{s,m}}^{m}) \wedge \bigwedge_{m \in \{1, \dots, m\}} \bigwedge_{1 \le k < c_{s,m}} S_{m}(x_{k+1}^{m}, x_{k}^{m})$$

Since π'_s and π''_s do not share variables (except for *x*), in order to prove (*) it will be enough to show that:

$$|\{h \in Hom(\pi'_{s}, D) : h(x) = a_{m}\}| = t_{m}(\Xi_{D}(\vec{x}))$$
(**)

and that:

$$|\{h \in Hom(\pi''_{s}, D) : h(x) = a_{m}\}| = c_{s,m}$$
(***)

To see (**) notice that for each $d \in \{1, ..., d\}$ there is a unique way for $h(y_d)$ to be defined. But, again for each $d \in \{1, ..., d\}$, the value of $h(z_d)$ can be chosen in $\Xi_D(\mathbf{x}_{m_d})$ ways, where \mathbf{x}_{m_d} is the *d*-th variable of the term t_m . And the choices, for each *d*, are independent, so indeed:

$$|\{h \in Hom(\pi'_{s}, D) : h(x) = a_{m}\}| = \prod_{d \in \{1, \dots, d\}} \Xi_{D}(\mathbf{x}_{m_{d}}) = t_{m}(\Xi_{D}(\vec{\mathbf{x}}))$$

Now, let us concentrate on (***). Recall that π''_s is a star, with x as its center, and (for each $j \in \{1, ..., m\}$) with a "ray" consisting of $c_{s,j} - 1$ edges of relation S_j radiating from this center. On the other hand, in our correct database D, there is a loop $S_j(a_m, a_m)$ for each $j \in \{1, ..., m\}$ (recall that $h(x) = a_m$) and there are no other edges of the form $S_j(a_m, ...)$ for $j \neq m$. This means that if $j \neq m$ then the S_j -ray can only be mapped to D in one way: by mapping all its edges to $S_j(a_m, a_m)$. The situation is different when j = m. Then we still have the loop $S_m(a_m, a_m)$ in D, but there is also $S_m(a_m, a_m)$. But it can also loop in a_m for some time, then have one of its edges mapped to $S_m(a_m, a)$ and then have all the remaining edges mapped to $S_m(a, a)$. The edge to be mapped to $S_m(a_m, a)$ can be chosen in $c_{s,j} - 1$ ways, and hence we get (***).

The argument for π_b is almost analogous. The only difference is that now we need to notice that:

$$\Xi_D(\mathbf{x}_1)^{\mathsf{d}} = (\bigwedge_{d \in \{1, \dots, \mathsf{d}\}} R_1(x, y'_d) \wedge X(y'_d, z'_d) \quad)(D)$$

B Proof of Lemma 11

In this appendix we show how Lemma 11 follows from undecidability of Hilbert's 10th problem.

B.1 Hilbert's 10th problem

Before we start, let us clarify what exactly we mean by Hilbert's 10th problem. In order to do it, we refer to the canonical resource [18]. One of the results presented there, the one we are interested in, can be formulated as:

Theorem 6. (*Hilbert's 10th problem undecidablity*)

The problem:

Given is polynomial Q of numerical variables $\xi_{Q,1}, ..., \xi_{Q,\mathfrak{n}_Q}$ with integer coefficients. Does it hold that $Q(\Xi(\bar{\xi}_Q)) \neq 0$ for every valuation $\Xi : \{\xi_{Q,1}, \xi_{Q,2}, ..., \xi_{Q,\mathfrak{n}_Q}\} \to \mathbb{N}$?

is undecidable.

We are going to show how the above Theorem implies Lemma 11. As a way to do it, for any Q being an instance of 10th Hilbert's problem (understood like in Theorem 6), we will construct a triple $[P_s, P_b, c']$ – the corresponding instance of the problem from Lemma 11. Keeping the notation from Lemma 11 and from Theorem 6, the constructed instance should be such that these are equivalent:

• there exists valuation $\Xi : \{\xi_{Q,1}, ..., \xi_{Q,n_q}\} \to \mathbb{N}$ such that

 $Q(\Xi(\bar{\xi_0})) = 0$

O there exists valuation $\Xi' : \{\xi_1, ..., \xi_n\} \to \mathbb{N}$ such that

$$\mathrm{d}P_s(\Xi'(\bar{\xi})) > \Xi'(\xi_1)^d P_b(\Xi'(\bar{\xi}))$$

B.2 From one polynomial to two

Let Q be any polynomial like in Theorem 6. Rename the variables of Q as $\xi_2, ..., \xi_n$. From this point, we will denote by $\overline{\xi}$ the tuple [$\xi_2, ..., \xi_n$]. Note that we start indexing from 2 on purpose (in particular, $n = n_Q + 1$, if we refer to the notation from Lemma 6). The reason for such choice will become clear soon.

We will now construct two polynomials P_1 and P_2 of the same variables as Q, which both have natural coefficients and such that for any fixed valuation P_1 is greater than P_2 if and only if Q equals zero for the same valuation.

First consider polynomial $Q' = Q^2$ It can of course be written in the form:

$$Q' = \sum_{i=1}^{m_{Q'}} c_i t_i,$$

where $m_{Q'} \in \mathbb{N}$, the values $c_i \in \mathbb{Z}$ are coefficients and t_i are monomials (of any degree each). Let now I_+ and I_- be the sets of indices of positive and negative coefficients, respectively:

$$I_{+} = \{1 \le i \le m_{Q'} : c_i > 0\}$$
$$I_{-} = \{1 \le i \le m_{Q'} : c_i < 0\}$$

This allows us to introduce the polynomials:

$$Q'_+ = \Sigma_{i \in I_+} c_i t_i$$

and

$$Q'_{-} = \sum_{i \in I_{-}} (-c_i) t_i$$

It is easy to see that $Q' = Q'_+ - Q'_-$ and that all the coefficients in Q'_+ and in Q'_- are natural. Let $P_1 = Q_- + 1$ and $P_2 = Q_+$. Observe that:

Lemma 25. For each valuation $\Xi : \{\xi_2, ..., \xi_n\} \to \mathbb{N}$ it holds that $Q(\Xi(\bar{\xi})) = 0$ if and only if $P_1(\Xi(\bar{\xi})) > P_2(\Xi(\bar{\xi}))$.

Proof: For any valuation Ξ like above we have:

$$\begin{array}{l} Q(\Xi(\bar{\xi})) = 0 \Leftrightarrow \\ \Leftrightarrow \ Q^2(\Xi(\bar{\xi})) = Q'(\Xi(\bar{\xi})) < 1 \Leftrightarrow \\ \Leftrightarrow \ Q'_+(\Xi(\bar{\xi})) - Q'_-(\Xi(\bar{\xi})) < 1 \Leftrightarrow \\ \Leftrightarrow \ Q'_+(\Xi(\bar{\xi})) < Q'_-(\Xi(\bar{\xi})) + 1 \Leftrightarrow \\ \Leftrightarrow \ P_2(\Xi(\bar{\xi})) < P_1(\Xi(\bar{\xi})). \end{array}$$

Lemma 25 is the first span in our bridge between Theorem 6 and Lemma 11. But it is not the entire bridge yet, recall that Lemma 11 mentions several conditions which P_s and P_b have to satisfy. And, clearly, there is no reason to think that P_1 and P_2 will always satisfy all of them.

So now we will, step by step, keep adjusting P_1 and P_2 to consecutive conditions from Lemma 11. Finally, we will obtain a pair of polynomials which indeed can be taken as P_s and P_b (the constant c will appear in the meantime).

At the end, we want to refer back to Lemma 25. In order to be able to do this, we need to construct P_s and P_b in such a way that they in some sense inherit the initial inequality between P_1 and P_2 .

B.3 Common monomials

We can express P_1 and P_2 in the form similar to Q', that is:

$$P_1 = \sum_{i=1}^{m_1} c_{1,i} t_{1,i}$$
$$P_2 = \sum_{i=1}^{m_2} c_{2,i} t_{2,i}$$

where $m_1, m_2 \in \mathbb{N}$, $c_{j,i} \in \mathbb{N}$ are again coefficients and $t_{j,i}$ are again monomials.

Denote by T_1 the set of monomials of P_1 , that is $\{t_{1,1}, ..., t_{1,m_1}\}$. Analogously, let T_2 be such set for P_2 . If we wanted to take P_1 and P_2 as P_s and P_b , respectively, then one of the conditions from Lemma 11 can be expressed as $T_1 = T_2$.

However, it may happen that $T_1 \neq T_2$. To deal with this problem, set $T = T_1 \cup T_2$, that is, the set of monomials appearing in P_1 or in P_2 . For further convenience, it is helpful to denote the monomials belonging to T as $t_1, ..., t_m$, where m = |T|. It is also useful to introduce the polynomial $P = \sum_{i=1}^{m} t_i$. Now, instead of P_1 and P_2 , consider:

$$P_1' = P_1 + I$$

 $P'_{2} = P_{2} + P$

and

These polynomials have common set of monomials, namely T. This allows us to express them in the form we are already familiar with:

$$P'_{1} = \sum_{i=1}^{m} c'_{1,i} t_{i}$$
$$P'_{2} = \sum_{i=1}^{m} c'_{2,i} t_{i}$$

Note that the new coefficients c' are different then the ones appearing in P_1 and P_2 , but they still are natural numbers.

B.4 Common degree and the variable ξ_1

The next problem is that we want all $t_i \in T$ to be of the same degree degree. Moreover, we want all t_i to start with the same variable. Let us try to deal with these both constraints simultaneously, introducing the new variable ξ_1 .

Denote by d_i the degree of monomial t_i . Take $d = 1 + max\{d_1, ..., d_m\}$ and define:

$$t_i' = \xi_1^{d-d_i} t_i.$$

Now, define P_1'', P_2'' to be:

$$P_{1}'' = \sum_{i=1}^{m} c_{1,i}' t_{i}'$$
$$P_{2}'' = \sum_{i=1}^{m} c_{2,i}' t_{i}'$$

Notice that the coefficients remained unchanged in P_1'' and P_2'' (compared to P_1' and P_2'), we just modified the monomials a little bit, so that they are all now divisible by ξ_1 and all have degree d.

Once we defined P_1'' and P_2'' , one can notice that:

Lemma 26. For any valuation $\Xi : \{\xi_2, ..., \xi_n\} \to \mathbb{N}$ and $a \in \mathbb{N}$ it holds that:

- $P_1''([1, \Xi(\bar{\xi})]) = P_1'(\Xi(\bar{\xi}))$
- $P_{1}^{\prime\prime}([a, \Xi(\bar{\xi})]) \le a^{d}P_{1}^{\prime}(\Xi(\bar{\xi})) \le a^{d}P_{1}^{\prime\prime}(\Xi(\bar{\xi}))$
- the two above claims hold also for P_2'' and P_2' , respectively

Proof: The first claim is a trivial consequence of the fact that to construct P''_1 from P'_1 , we only multiply each of the terms appearing in P'_1 by some power of ξ_1 .

To make sure the second claim holds, it is convenient to consider each of the inequalities separately. For the first inequality, it is enough to notice that for each $1 \le i \le m$ we have:

$$c'_{1,i}t'_{i}([a, \Xi(\bar{\xi})]) = c'_{1,i}a^{d-d_{i}}t_{i}(\Xi(\bar{\xi})) \le a^{d}c'_{1,i}t_{i}(\Xi(\bar{\xi}))$$

The equality above follows from the definition of t'_i .

The prove the above inequality two cases need to be considered. The simpler one is when a = 0, as the inequality is trivially satisfied then. When a > 0, for each $1 \le i \le m$ we have:

$$a^{d-d_i} \ge 1$$

which implies:

$$c_{1,i}'t_i(\Xi(\bar{\xi})) \leq c_{1,i}'t_i'([a,\Xi(\bar{\xi})])$$

The second second inequality becomes clear after multiplying both sides by a^d .

The last claim can be shown by repeating exactly the same arguments for P_2'' and P_2' .

Now we want to show that the inequality between P_1 and P_2 in some sense survives when we consider P''_1 and P''_2 instead of them. The exact way it happens is presented by the next two lemmas.

Lemma 27. Let Ξ : { $\xi_2, ..., \xi_n$ } $\rightarrow \mathbb{N}$ be any valuation such that $P_1(\Xi(\bar{\xi})) > P_2(\Xi(\bar{\xi}))$. Then $P''_1([1, \Xi(\bar{\xi})]) > P''_2([1, \Xi(\bar{\xi})])$.

Proof:

$$\begin{split} P_1''([1,\Xi[\bar{\xi}]) &= P_1'(\Xi[\bar{\xi}]) = P_1(\Xi[\bar{\xi}]) + P(\Xi[\bar{\xi}]) > \\ &> P_2(\Xi[\bar{\xi}]) + P(\Xi[\bar{\xi}]) = P_2'(\Xi[\bar{\xi}]) = P_2''([1,\Xi[\bar{\xi}]]) \end{split}$$

The first and last equalities come from the first and third claims of Lemma 26.

Lemma 28. Let $\Xi' : \{\xi_1, ..., \xi_n\} \to \mathbb{N}$ be any valuation. Consider the valuation $\Xi : \{\xi_2, ..., \xi_n\} \to \mathbb{N}$ such that $\Xi'([\xi_1, \overline{\xi}]) = [a, \Xi(\overline{\xi})]$ for some $a \in \mathbb{N}$. If additionally $P_1(\Xi(\overline{\xi})) \leq P_2(\Xi(\overline{\xi}))$ holds, then also $P''_1([a, \Xi(\overline{\xi})]) \leq a^d P''_2([a, \Xi(\overline{\xi})])$.

 $\begin{array}{l} Proof: \ P_1''([a, \Xi(\bar{\xi})]) \leq a^d P_1'(\Xi(\bar{\xi})) = a^d(P_1(\Xi(\bar{\xi})) + P(\Xi(\bar{\xi}))) \leq \\ \leq a^d(P_2(\Xi(\bar{\xi})) + P(\Xi(\bar{\xi}))) = a^d P_2'(\Xi(\bar{\xi})) \leq a^d P_2''([a, \Xi(\bar{\xi})]). \end{array}$

Here the first and last inequalities come from the second and third claims of Lemma 26.

B.5 The final construction

 P_1'' and P_2'' are almost what we want to have as P_s and P_b . Almost, because we are still left with the last condition from Lemma 11: for each $1 \le i \le m$ we require $c_{1,i}' \le c_{2,i}'$ to hold.

This is the place where the constant c plays its role: to enforce inequalities mentioned in the previous sentence, we may multiply P_2 by some big number. It is enough to take:

$$c' = max\{c'_{1,1}, c'_{1,2}, ..., c'_{1,m}\}$$

We may thus finally define also:

$$P_s = P_1^{\prime\prime}$$
 and $P_b = c P_2^{\prime\prime}$

The last step is to notice that indeed for each $1 \le i \le m$ we have now:

$$c_{s,i} = c'_{1,i} \le c' \le c' c'_{2,i} = c_{b,i}$$

where $c_{s,i}$ and $c_{b,i}$ are understood like in Lemma 11.

B.6 Proof of the equivalence

If we carefully followed the construction of P_s and P_b , we should be now confident that they (alongside with c) satisfy all the conditions from Lemma 11. Thus, we just need to prove the following lemma, which together with Lemma 25 will tell us that $\bullet \Leftrightarrow O$:

Lemma 29. The following conditions are equivalent:

- there exists valuation $\Xi : \{\xi_2, ..., \xi_n\} \to \mathbb{N}$ such that $P_1(\Xi(\overline{\xi})) > P_2(\Xi(\overline{\xi}));$
- there exists valuation $\Xi': \{\xi_1, \xi_2, ..., \xi_n\} \to \mathbb{N}$ such that $cP_s(\Xi'([\xi_1, \overline{\xi}])) > \Xi'(\xi_1)^d P_b(\Xi'([\xi_1, \overline{\xi}])).$

Proof: Consider first the situation when for some valuation Ξ we have $P_1(\Xi(\bar{\xi})) > P_2(\Xi(\bar{\xi}))$. Take Ξ' such that:

$$\Xi'([\xi_1, \bar{\xi}]) = [1, \Xi(\bar{\xi})].$$

From Lemma 27 we can conclude that:

$$P_1''([1,\Xi(\bar{\xi})]) > P_2''([1,\Xi(\bar{\xi})])$$

and multiplying both sides by c allows us to write:

$$\begin{split} c P_s(\Xi'([\xi_1,\bar{\xi}])) &= c P_s([1,\Xi(\bar{\xi})]) = c P_1''([1,\Xi(\bar{\xi})]) > \\ > c P_2''([1,\Xi(\bar{\xi})]) &= P_b([1,\Xi(\bar{\xi})]) = P_b(\Xi'([\xi_1,\bar{\xi}])) = \Xi'(\xi_1)^d P_b(\Xi'([\xi_1,\bar{\xi}])). \end{split}$$

In the last equality, we were allowed to insert $\Xi'(\xi_1)^d$ before $P_b(\Xi'([\xi_1, \overline{\xi}]))$, as $\Xi'(\xi_1) = 1$.

Now assume that $P_1(\Xi(\bar{\xi})) \leq P_2(\Xi(\bar{\xi}))$ holds for all valuations Ξ . Consider then any valuation Ξ' , we know that there exist $a \in \mathbb{N}$ and valuation Ξ such that $\Xi'([\xi_1, \bar{\xi}]) = [a, \Xi(\bar{\xi})]$. Lemma 28 tells us that we then have $P''_1([a, \Xi(\bar{\xi})]) \leq a^d P''_2([a, \Xi(\bar{\xi})])$. Let us again multiply both sides by c, obtaining in this case:

$$cP_s(\Xi'([\xi_1,\bar{\xi}])) = cP_s([a,\Xi(\bar{\xi})]) = cP''_1([a,\Xi(\bar{\xi})]) \le \le ca^d P''_2([a,\Xi(\bar{\xi})]) = a^d P_b([a,\Xi(\bar{\xi})]) = \Xi'(\xi_1)^d P_b(\Xi'([\xi_1,\bar{\xi}])).$$