

Non-(strong) ergodicity criteria for discrete time Markov chains on general state spaces*

Ling-Di Wang^{a†} Yu Chen^{b*‡} Yu-Hui Zhang^b

March 25, 2025

Abstract

For discrete time Markov chains on general state spaces, we provide criteria for non-ergodicity and non-strong ergodicity. By taking advantage of minimal non-negative solution theory, our criteria are in terms of the existence of solutions to inequalities involving the one step transition semigroup of the chain. Based on Dynkin's formula, Lyapunov-type conditions for non-strong ergodicity are also obtained.

Keywords: Markov chain, non-(strong) ergodicity, test function, general state space.

MSC 2010: 60J05, 37B25, 60J35

1 Introduction

Ergodicity is one of the three fundamental and classical problems (uniqueness, recurrence and ergodicity) for Markov chains([2]). Ordinary ergodicity, strong ergodicity and so on, are usually used to characterize the stability of Markov chains, which play important roles in both theoretical and applied researches([3, 4, 8, 9, 13]). The topic on criteria for various types of ergodicity has been studied by many authors in the past decades, see [2, 5, 7, 10] etc.. Especially, Lyapunov (drift) criteria have always been used to provide sufficient and necessary conditions for ergodicity or strong ergodicity. In principle, these results also provided the conditions for the non-(strong) ergodicity by stating that there does not exist Lyapunov functions, which is usually not practical. Due to this reason, Non-(strong) ergodicity criteria with test functions for Markov chains, which is so-called “inverse problem”, have also been studied specially. We refer the readers to [1, 11, 12]

* Corresponding author: Yu Chen; E-mail: ychen0504@mail.bnu.edu.cn

[†]*a.* School of Mathematics and Statistics, Henan University, Kaifeng 475001, China;

[‡]*b.* School of Mathematics Sciences, Beijing Normal University, Laboratory of Mathematics and Complex Systems, Ministry of Education, Beijing 100875, China.

for the inverse problem of Markov process in denumerable state space and continuous time settings, to [14] in denumerable state space and discrete time settings, to [6] in general state space and continuous time settings. However, to our knowledge, during the existing literature, there are few results on non-ergodicity or non-strong ergodicity for discrete time Markov chain on general state space, which will be studied in the paper, presenting the criteria or sufficient Lyapunov (drift) conditions for its non-(strong) ergodicity.

Consider a homogeneous Markov chain on a general state space \mathcal{X} , which is a locally compact separable metric topological space, endowed with a countably generated σ -field $\mathcal{B}(\mathcal{X})$. Denoted by

$$P^n(x, A) = \mathbb{P}(X_n \in A | X_0 = x), \quad n \geq 0, x \in \mathcal{X}, A \in \mathcal{B}(\mathcal{X}),$$

with $P(x, A) = P^1(x, A)$. The Markov chain $\{X_n\}$ is called ψ -irreducible if for some σ -finite measure ψ on $(\mathcal{X}, \mathcal{B}(\mathcal{X}))$, such that $\sum_{n=1}^{\infty} P^n(x, A) > 0$ for all $x \in \mathcal{X}$ whenever $\psi(A) > 0$. **Throughout the paper, we assume the chain $\{X_n\}$ is ψ -irreducible, where ψ is a maximal irreducibility measure on $(\mathcal{X}, \mathcal{B}(\mathcal{X}))$.** The concepts “aperiodicity”, “Feller” and “ ψ -irreducible” coincide with those in [10] (see Section 5.4.2 and Chapter 6 therein) and we omit them here for simplicity. Write $Pf(x) = \int_E f(y)P(x, dy)$ for some $\mathcal{B}(\mathcal{X})$ measurable function f and

$$\mathcal{B}^+(\mathcal{X}) = \{A \in \mathcal{B}(\mathcal{X}) : \psi(A) > 0\}.$$

Denote by $\|\cdot\|_{TV}$ the total variation norm of some measure. The Markov chain $\{X_n\}$ is called

- 1) **recurrent** if $\sum_{n=1}^{\infty} P^n(x, A) = \infty$, $x \in \mathcal{X}$, $A \in \mathcal{B}^+(\mathcal{X})$, otherwise we call it **transient**;
- 2) **Harris recurrent** if $\mathbb{P}_x(\sum_{n=1}^{\infty} \mathbf{1}_{\{X_n \in A\}} = \infty) = 1$, $x \in \mathcal{X}$ and $A \in \mathcal{B}^+(\mathcal{X})$;
- 3) **positive** if it admits an invariant probability measure;
- 4) **ergodic** if there exists a probability measure π such that

$$\lim_{n \rightarrow \infty} \|P^n(x, \cdot) - \pi\|_{TV} = 0;$$

- 5) **strongly (or uniformly) ergodic** if there exists a probability π such that

$$\lim_{n \rightarrow \infty} \sup_{x \in \mathcal{X}} \|P^n(x, \cdot) - \pi\|_{TV} = 0.$$

Refer to [10, 13] for more details on these notions. In general, for aperiodic Harris recurrent Markov chains, we refer to “positive Harris recurrent chains” as “ergodic chain” ([10, Theorem 13.0.1]). A Harris recurrent chain must be a recurrent chain. Since we are dealing with ergodic properties, we assume that the Markov chains we considered in the following are all Harris recurrent without especially pointed. “Harris recurrent” is short for “recurrent” in the paper.

Petite sets and small sets (see [10, Section 5]) are also needed. Precisely, for some $A \in \mathcal{B}(\mathcal{X})$, if there exists an $m > 0$ and a non-trivial measure ν_m on $\mathcal{B}(\mathcal{X})$ such that for all $x \in A$,

$B \in \mathcal{B}(\mathcal{X})$, $P^m(x, B) \geq \nu_m(B)$, then A is called ν_m -small. $A \in \mathcal{B}(\mathcal{X})$ is called ν_a -petite set if there exists a probability distribution $a = \{a_n\}$ on \mathbb{Z}_+ ,

$$\sum_{n=0}^{\infty} a_n P^n(x, B) \geq \nu_a(B), \quad B \in \mathcal{B}(\mathcal{X})$$

for $x \in A$, where ν_a is a non-trivial measure on $\mathcal{B}(\mathcal{X})$. Under the assumption that $\{X_n\}$ is ψ -irreducible and aperiodic, a set is a petite set if and only if it is small ([10, Theorem 5.5.7]). Petite sets are not rare: the compact sets are petite under some mild conditions ([10, Proposition 6.2.8]). For each $A \in \mathcal{B}^+(\mathcal{X})$, there exists a petite set $B \subseteq A$ and $B \in \mathcal{B}^+(\mathcal{X})$ ([10, Theorem 5.2.2]).

Now, we declare our results.

Assumption 1 Markov chain $\{X_n\}$ is ψ -irreducible, aperiodic, Feller and $\text{supp}\psi$ has nonempty interior.

Assumption 2 The transition kernel $P(x, dy)$ is absolutely continuous with respect to the maximal irreducibility measure ψ .

Theorem 1.1 (Non-ergodicity) Assume that Markov chain $\{X_n\}$ satisfying Assumption 1 and 2 is recurrent. Then $\{X_n\}$ is not ergodic if and only if there exists a set $A \in \mathcal{B}^+(\mathcal{X})$ and a sequence of functions $\{V^{(n)}(x)\}_{n \geq 1}$, $V^{(n)}(x) : \mathcal{X} \rightarrow \mathbb{R}$ satisfies the following conditions:

- 1) $\sup_{x \in A^c} V^{(n)}(x) < \infty$, $n \geq 1$;
- 2) $\int_{A^c} V^{(n)}(y) P(x, dy) \geq V^{(n)}(x) - 1$, $x \in A^c$, $n \geq 1$;
- 3) $\int_A \sup_{n \geq 1} V^{(n)}(y) \psi(dy) = \infty$.

Theorem 1.2 (Non-strong ergodicity) Assume that Markov chain $\{X_n\}$ satisfies Assumption 1. Then $\{X_n\}$ is non-strongly ergodic if and only if there exists a set $A \in \mathcal{B}^+(\mathcal{X})$ and a sequence of functions $\{V^{(n)}(x)\}_{n \geq 1}$, $V^{(n)}(x) : \mathcal{X} \rightarrow \mathbb{R}$ satisfies the following conditions:

- 1) $\sup_{x \in A^c} V^{(n)}(x) < \infty$, $n \geq 1$; $V^{(n)}(x) = 0$, $x \in A$, $n \geq 1$;
- 2) $PV^{(n)}(x) \geq V^{(n)}(x) - 1$, $x \in A^c$, $n \geq 1$;
- 3) $\sup_{x \in A^c, n \geq 1} V^{(n)}(x) = \infty$.

Remark 1.3 We release the assumption of Feller and $\text{supp}\psi$ has nonempty interior in Theorem 1.1 or Theorem 1.2, their sufficiency still holds respectively.

Proposition 1.4 Assume that $\{E_n, n \geq 1\} \subset \mathcal{B}(\mathcal{X})$ is an increasing sequence of bounded sets and $\mathcal{X} = \cup_{m=1}^{\infty} E_m$. There exists a continuous function $V : \mathcal{X} \rightarrow \mathbb{R}$, a set $A \in \mathcal{B}^+(\mathcal{X})$ such that

- 1) $\sup_{x \in A} V(x) < \infty$; $\sup_{x \in A^c} V(x) = \infty$.
- 2) $V(x)$ is locally bounded and $PV(x) \geq V(x) - 1$ for $x \in A^c$;
- 3) $W(x) : \mathcal{X} \rightarrow [0, \infty)$ is continuous, $PW(x) \leq W(x) + d\mathbf{1}_A(x)$ for some constant d ; and
$$\lim_{m \rightarrow \infty} \sup_{x \in E_m^c} \frac{V(x)}{W(x)} = 0.$$

Then $\{X_n\}_{n \geq 0}$ is non-strongly ergodic.

2 Proof of criteria for inverse problems

2.1 Recall minimal solution theory

We first recall some useful results in minimal solution theory from [2].

Define \mathcal{H} a set of mappings from \mathcal{X} to $[0, \infty]$: \mathcal{H} contains the constant 1 and is closed under the nonnegative combination and monotone increasing limit, where the order relation “ \geq ” in \mathcal{H} is defined point-wise. Then \mathcal{H} is a convex cone. $H : \mathcal{H} \rightarrow \mathcal{H}$ is called a cone mapping if $H0 = 0$, and for any nonnegative constants c_1, c_2 , mappings f_1, f_2 in \mathcal{H} ,

$$H(c_1f + c_2f_2) = c_1H(f_1) + c_2H(f_2).$$

Denote by \mathcal{A} the set of all such mappings which also satisfy the following hypothesis: for $f_n \in \mathcal{H}$,

$$f_n \uparrow f \text{ implies } Hf_n \uparrow Hf.$$

Definition 2.1 ([2, Definition 2.1]) Given $H \in \mathcal{A}$ and $g \in \mathcal{H}$. We say the f^* is the minimal nonnegative solution (abbrev. minimal solution) to the equation

$$f(x) = (Hf)(x) + g(x), \quad x \in \mathcal{X}, \quad (2.1)$$

if f^* satisfies (2.1) and for any solution $\tilde{f} \in \mathcal{H}$ of (2.1),

$$\tilde{f}(x) \geq f^*(x), \quad x \in \mathcal{X}.$$

The last property is called the minimum property of f^* .

Lemma 2.2 ([2, Theorem 2.2]) The minimum solution to Equation (2.1) always exists uniquely.

By Lemma 2.2, for each $H \in \mathcal{A}$, we may define a map m_H from \mathcal{H} into itself: $m_H(g) = f^*$. We mention that for $H_1, H_2 \in \mathcal{A}$, $H_1 \geq H_2$ if and only if $H_1f \geq H_2f$ for $f \in \mathcal{H}$.

Lemma 2.3 ([2, Theorem 2.7]) m_H is a cone mapping. Let $\{g_n\}_{n=1}^\infty \subset \mathcal{H}$, $\{H_n\} \subset \mathcal{A}$, $H_n \uparrow H$ and $g_n \uparrow g$. Then $g \in \mathcal{H}$, $H \in \mathcal{A}$ and $m_{H_n}(g_n) \uparrow m_H(g)$.

Lemma 2.4 ([2, Theorem 2.9]) Let $H \in \mathcal{A}$, $\{g_n\}_{n=1}^\infty \subset \mathcal{H}$. Define $\tilde{f}^{(1)} = g_1$, $\tilde{f}^{(n+1)} = H\tilde{f}^{(n)} + g_{n+1}$, $n \geq 1$. If $g_n \uparrow g$ (resp., $\sum_{n=1}^\infty g_n = g$), then $\tilde{f}^{(n)} \uparrow m_Hg$ (resp., $m_Hg = \sum_{n=1}^\infty \tilde{f}^{(n)}$).

2.2 Lower bound for the moments of hitting times and sufficiency

In this section, we prove the sufficiency of Theorem 1.1, Theorem 1.2 and Proposition 1.4 by taking the minimal solution theory and Dynkin's formula.

For any set $A \in \mathcal{B}(\mathcal{X})$, define the first return time τ_A^+ and first hitting time τ_A on A , respectively as follows:

$$\tau_A^+ := \inf\{n \geq 1 : X_n \in A\}; \quad \tau_A := \inf\{n \geq 0 : X_n \in A\}.$$

Denote

$$L(x, A) = \sum_{n=1}^{\infty} \mathbb{P}_x(\tau_A^+ = n) = \mathbb{P}_x(\tau_A^+ < \infty).$$

Before proceeding further, let's briefly describe the main points in our proofs.

Taking non-strong ergodicity for instance, it has other description ([10, Theorem 16.0.2]): for aperiodic chain $\{X_n\}$, it is strongly ergodic if and only if there exists a petite set $C \in \mathcal{B}(\mathcal{X})$ such that $\sup_{x \in \mathcal{X}} \mathbb{E}_x \tau_C^+ < \infty$, and for every set $A \in \mathcal{B}^+(\mathcal{X})$, we have $\sup_{x \in \mathcal{X}} \mathbb{E}_x \tau_A^+ < \infty$. To verify it is non-strongly ergodic, it suffices to prove $\sup_{x \in \mathcal{X}} \mathbb{E}_x \tau_A^+ = \infty$ for some set $A \in \mathcal{B}^+(\mathcal{X})$. We first get a lower bound for the expectation of return time according to minimal solution theory (see Proposition 2.7), then an increasing sequence of lower bounds can imply the desired results. On the other hand, finite approximation method (see Proposition 2.10 and Lemmas 2.13, 2.14) would guarantee the existence of an increasing sequence of lower bounds and therefore the necessity of our conditions.

Besides, one can also get the lower bound of hitting time for the Markov chain according to Dynkin's formula by adding in proper Lyapunov-like type condition. Therefore, we can get some Lyapunov-like type criteria for non-strongly ergodicity (see (??) below for details).

The following assertion, which releases the nonnegative condition, is a modified version of [10, Theorem 8.0.2].

Proposition 2.5 *The ψ -irreducible Markov chain $\{X_n\}$ is transient if and only if there exists such a $C \in \mathcal{B}^+(\mathcal{X})$, and $V(x) : \mathcal{X} \rightarrow \mathbb{R}$, satisfying $\inf_{x \in \mathcal{X}} V(x) > -\infty$,*

$$PV(x) \leq V(x), \quad x \in C^c,$$

and

$$\left\{ x : V(x) < \inf_{y \in C} V(y) \right\} \in \mathcal{B}^+(\mathcal{X}). \quad (2.2)$$

Proof Necessity is obvious by [10, Theorem 8.0.2]. We then prove sufficiency. By [10, Theorem 8.3.6(ii)], it suffices to prove that there exist C and $D \in \mathcal{B}^+(\mathcal{X})$ so that

$$P_x(\tau_C^+ < \infty) < 1, \quad x \in D.$$

Without loss of generality, assume that $V(x)$ is nonnegative (otherwise, replace $V(x)$ by $V(x) - \inf_{x \in \mathcal{X}} V(x)$). For the set $C \in \mathcal{B}^+(\mathcal{X})$, denoted by $a = \inf_{x \in C} V(x)$ and $D = \{x : V(x) < a\}$. By (2.2), $a > 0$, $D \in \mathcal{B}^+(\mathcal{X})$ and $D \subseteq C^c$. For $x \in D$,

$$\begin{aligned} a > V(x) &\geq PV(x) = \int_C V(y)P(x, dy) + \int_{C^c} V(y)P(x, dy) \\ &\geq aP(x, C) + \int_{C^c} PV(y)P(x, dy) \\ &\geq aP(x, C) + a \int_{C^c} P(y, C)P(x, dy) + \int_{C^c} \int_{C^c} V(z_1)P(y, dz_1)P(x, dy) \\ &= a\mathbb{P}_x(\tau_C^+ = 1) + a\mathbb{P}_x(\tau_C^+ = 2) + \int_{C^c} \int_{C^c} V(z_1)P(y, dz_1)P(x, dy). \end{aligned}$$

By induction, we get for $x \in D$,

$$a > V(x) > a \sum_{k=1}^n \mathbb{P}_x(\tau_C^+ = k) + \int_{C^c} \cdots \int_{C^c} V(z_{n-1})P(z_{n-2}, z_{n-1}) \cdots P(y, dz_1)P(x, dy), \quad n \geq 1.$$

Then

$$1 > \mathbb{P}_x(\tau_C^+ < \infty), \quad x \in D$$

by letting $n \rightarrow \infty$. \square

The following minimal solution characterization of the moment of return time is essential for us to exploit minimal solution theory.

Proposition 2.6 *For each $A \in \mathcal{B}(\mathcal{X})$, $\mathbb{E}_x \tau_A^+ \mathbf{1}_{\{\tau_A^+ < \infty\}}$ is the minimal solution to the following equation:*

$$V(x) = \int_{A^c} V(y)P(x, dy) + L(x, A), \quad x \in \mathcal{X}. \quad (2.3)$$

Proof We prove it by the second successive approximation scheme for the minimal solution. Set $V^{(1)}(x) = \mathbb{P}_x(\tau_A^+ = 1)$. Consider the induction equation

$$V^{(n+1)}(x) = \int_{A^c} V^{(n)}(y)P(x, dy) + \mathbb{P}_x(\tau_A^+ = n+1), \quad x \in \mathcal{X}.$$

It has a solution $V^{(n)}(x) = n\mathbb{P}_x(\tau_A^+ = n)$, $n \geq 1$. Note that

$$L(x, A) = \sum_{n=1}^{\infty} \mathbb{P}_x(\tau_A^+ = n).$$

By Lemma 2.4,

$$V^*(x) := \sum_{n=1}^{\infty} V^{(n)}(x) = \mathbb{E}_x \tau_A^+ \mathbf{1}_{\{\tau_A^+ < \infty\}}, \quad x \in \mathcal{X}$$

is the minimal nonnegative solution to (2.3). \square

Notice $\{X_n\}$ is recurrent if and only if $L(x, A) = 1$ for all $x \in \mathcal{X}$ and $A \in \mathcal{B}^+(\mathcal{X})$ ([10, Theorem 8.3.6, Propositions 9.1.1, 9.1.7]). The following assertion provides a “lower bound” of the moment of return time.

Proposition 2.7 *Assume that Markov chain $\{X_n\}$ with one step transition probability kernel $\{P(x, \cdot) : x \in \mathcal{X}\}$ is ψ -irreducible and recurrent. If there exists a set $A \in \mathcal{B}^+(\mathcal{X})$, and a function $V(x) : \mathcal{X} \rightarrow \mathbb{R}$ satisfying $\sup_{x \in A^c} V(x) < \infty$ and*

$$V(x) \leq \int_{A^c} V(y) P(x, dy) + 1, \quad x \in A^c, \quad (2.4)$$

then

$$1) \quad \psi(\{y \in A^c : V(y) > \mathbb{E}_y \tau_A^+\}) = 0.$$

$$2) \quad V(x) \leq \mathbb{E}_x \tau_A^+ + \sup_{x \in A} V(x), \quad x \in \mathcal{X}. \text{ Moreover, if } V(x) = 0 \text{ for } x \in A, \text{ then}$$

$$V(x) \leq \mathbb{E}_x \tau_A^+, \quad x \in \mathcal{X}.$$

Proof 1) Since $\{X_n\}$ is recurrent, $L(x, A) = 1$. By Proposition 2.6,

$$\mathbb{E}_x \tau_A^+ = \int_{A^c} \mathbb{E}_y \tau_A^+ P(x, dy) + 1, \quad x \in \mathcal{X}. \quad (2.5)$$

Let

$$Z(x) = [\mathbb{E}_x \tau_A^+ - V(x)] \mathbf{1}_{A^c}(x).$$

Note that $Z(x)$ is well defined since $\sup_{x \in A^c} V(x) < \infty$, which also implies that $\inf_{x \in \mathcal{X}} Z(x) > -\infty$. For $x \in A^c$, by (2.4) and (2.5),

$$\begin{aligned} PZ(x) &= \int_{A^c} Z(y) P(x, dy) \\ &= \int_{A^c} \mathbb{E}_y \tau_A^+ P(x, dy) - \int_{A^c} V(y) P(x, dy) \\ &\leq \mathbb{E}_x \tau_A^+ - L(x, A) - [V(x) - L(x, A)], \end{aligned}$$

we get $PZ(x) \leq Z(x)$, $x \in A^c$. Since $\{X_n\}$ is recurrent, by Proposition 2.5, we have

$$\psi\left(\left\{x : Z(x) < \inf_{y \in A} Z(y)\right\}\right) = 0.$$

That is $\psi(D) = 0$ with

$$D = \left\{y \in A^c : V(y) > \mathbb{E}_y \tau_A^+ \mathbf{1}_{\{\tau_A^+ < \infty\}}\right\}.$$

2) Since

$$PV(x) = P(V \mathbf{1}_A)(x) + PV \mathbf{1}_{A^c}(x) \geq P(V \mathbf{1}_A)(x),$$

by Dynkin's formula and Markov property (see [10, Theorem 11.3.1]), for $x \in A^c$, $m \geq 1$,

$$\begin{aligned}\mathbb{E}_x V(X_{m \wedge \tau_A}) &= V(x) + \mathbb{E}_x \left[\sum_{i=0}^{m \wedge \tau_A - 1} \left(PV(X_i) - V(X_i) \right) \right] \\ &\geq V(x) - \mathbb{E}_x(m \wedge \tau_A).\end{aligned}\tag{2.6}$$

Since $\{X_n\}$ is Harris recurrent, $\mathbb{P}_x(\tau_A < \infty) = 1$ for $A \in \mathcal{B}^+(\mathcal{X})$, $x \in \mathcal{X}$. Thus,

$$\mathbb{P}_x(\tau_A > m) \rightarrow 0, \quad m \rightarrow \infty, \quad x \in \mathcal{X}.$$

Since

$$V(x) \leq \sup_{x \in A^c} V(x) < \infty,$$

we get

$$\overline{\lim}_{m \rightarrow \infty} \mathbb{E}_x V(X_{m \wedge \tau_A}) \leq \mathbb{E}_x V(X_{\tau_A}) \leq \sup_{x \in A} V(x), \quad x \in \mathcal{X}.$$

Moreover, $\mathbb{E}_x \tau_A^+ + \sup_{x \in A} V(x) \geq V(x)$, $x \in \mathcal{X}$ by (2.6) and the required assertions hold. \square

Proof of sufficiency of Theorem 1.1 We will verify it by finding a contradiction. If $\{X_n\}$ is ergodic, by Condition 1), 2) and Proposition 2.7,

$$\psi(N_n) = 0, \quad N_n := \{y \in A^c : V^{(n)}(y) > \mathbb{E}_y \tau_A^+\}, \quad n \geq 1.$$

Denoted by $N = \cup_{n=1}^{\infty} N_n$. Then $\psi(N) = 0$ and

$$\mathbb{E}_y \tau_A^+ \geq V^{(n)}(y), \quad y \in A^c \setminus N.$$

For $x \in A$, $n \geq 1$, based on Assumption 2 and Condition 2),

$$\begin{aligned}V^{(n)}(x) &\leq \int_{A^c} V^{(n)}(y) P(x, dy) + 1 \\ &\leq \int_{A^c \setminus N} V^{(n)}(y) P(x, dy) + 1 + \int_N V^{(n)}(y) P(x, dy) \\ &\leq \int_{A^c} \mathbb{E}_y \tau_A^+ P(x, dy) + 1 + \int_N \left(V^{(n)}(y) - \mathbb{E}_y \tau_A^+ \right) P(x, dy) \\ &= \mathbb{E}_x \tau_A^+.\end{aligned}$$

Hence,

$$\infty = \int_A \sup_{n \geq 1} V^{(n)}(x) \psi(dx) \leq \int_A \mathbb{E}_x \tau_A^+ \psi(dx),$$

which is a contradiction with ergodicity. As a consequence, the Markov chain is non-ergodic. \square

Proposition 2.8 *Markov chain $\{X_n\}$ is ψ -irreducible. If there exists $A, B \in \mathcal{B}^+(\mathcal{X})$ such that*

$$\{x \in A : \mathbb{E}_x \tau_B^+ = \infty\} \in \mathcal{B}^+(\mathcal{X}),\tag{2.7}$$

then $\{X_n\}$ is non-ergodic.

Proof If $\{X_n\}$ is ergodic, by [10, Theorem 11.1.4], there exists a set S , which has a countable cover $\{S_n\}$, is full(i.e., $\psi(S^c) = 0$), absorbing set(i.e., $P(x, S) = 1, x \in S$) and

$$\sup_{x \in S_n} \mathbb{E}_x \tau_B^+ < \infty, \quad B \in \mathcal{B}^+(\mathcal{X}).$$

Therefore,

$$\psi(\{x \in \mathcal{X} : \mathbb{E}_x \tau_B^+ = \infty\}) = \psi(\{x \in S : \mathbb{E}_x \tau_B^+ = \infty\} \cup S^c) = 0, \quad B \in \mathcal{B}^+(\mathcal{X}),$$

which is a contradiction with (2.7). \square

Corollary 2.9 *Assume that Markov chain $\{X_n\}$ satisfying Assumption 1 and 2 is recurrent. Then $\{X_n\}$ is not ergodic if there exists a set $A, B \in \mathcal{B}^+(\mathcal{X})$ and a sequence of functions $\{V^{(n)}(x)\}_{n \geq 1}, V^{(n)}(x) : \mathcal{X} \rightarrow \mathbb{R}$ satisfies the following conditions:*

- 1) $\sup_{x \in A^c} V^{(n)}(x) < \infty, n \geq 1;$
- 2) $\int_{A^c} V^{(n)}(y) P(x, dy) \geq V^{(n)}(x) - 1, x \in A^c, n \geq 1;$
- 3) $\{x \in B : \sup_{n \geq 1} V^{(n)}(x) = \infty\} \in \mathcal{B}^+(\mathcal{X}).$

Proof This proof differs only in Condition 3) from the sufficiency proof of Theorem 1.1.

Notice that,

$$V^{(n)}(x) \leq \mathbb{E}_x \tau_A^+, \quad x \in \mathcal{X} \setminus N, \quad n \geq 1.$$

For $B \in \mathcal{B}^+(\mathcal{X})$,

$$\{x \in B : \sup_{n \geq 1} V^{(n)}(x) = \infty\} \subset \{x \in B : \mathbb{E}_x \tau_A^+ = \infty\} \in \mathcal{B}^+(\mathcal{X}),$$

by 3). The required assertion holds immediately by Proposition 2.8. \square

Proof of sufficiency of Theorem 1.2 If $\{X_n\}$ is strongly ergodic, then $\mathbb{P}_x(\tau_A^+ < \infty) = 1$ for $A \in \mathcal{B}^+(\mathcal{X}), x \in \mathcal{X}$. By Proposition 2.7 and Conditions 1), 2),

$$\mathbb{E}_x \tau_A^+ \geq V^{(n)}(x), \quad x \in \mathcal{X}.$$

Hence,

$$\sup_{x \in A^c} \mathbb{E}_x \tau_A^+ \geq \sup_{x \in A^c, n \geq 1} V^{(n)}(x) = \infty$$

by 3). This is a contradiction with $\sup_{x \in \mathcal{X}} \mathbb{E}_x \tau_A^+ < \infty$. Therefore, $\{X_n\}$ is non-strong ergodicity. \square

Proof of Proposition 1.4 By a similar argument to the proof of Theorem 1.2, for $x \in E_n \setminus A, m \geq 1$,

$$\mathbb{E}_x V(X_{m \wedge \tau_A \wedge \tau_{E_n^c}}) \geq V(x) - \mathbb{E}_x(m \wedge \tau_A \wedge \tau_{E_n^c}),$$

and

$$\mathbb{E}_x W(X_{m \wedge \tau_A \wedge \tau_{E_n^c}}) \leq W(x). \quad (2.8)$$

If the chain is strongly ergodic, then $\mathbb{P}_x(\tau_A < \infty) = 1$ for $A \in \mathcal{B}^+(\mathcal{X})$. Moreover, $\lim_{m \rightarrow \infty} \mathbb{P}_x(\tau_A > m) = 0$ for $x \in \mathcal{X}$. By (2.8),

$$\mathbb{E}_x W(X_{\tau_A \wedge \tau_{E_n^c}}) \leq \lim_{m \rightarrow \infty} \mathbb{E}_x W(X_{m \wedge \tau_A \wedge \tau_{E_n^c}}) \leq W(x), \quad x \in E_n \setminus A. \quad (2.9)$$

By the local boundedness of V ,

$$|V(X_m)| \mathbf{1}_{\{m < \tau_A \wedge \tau_{E_n^c}\}} \leq \sup_{x \in E_n \setminus A^c} |V(x)| < \infty.$$

Then $\overline{\lim}_{m \rightarrow \infty} \mathbb{E}_x V(X_m) \mathbf{1}_{\{m < \tau_A \wedge \tau_{E_n^c}\}} = 0$, $x \in E_n \setminus A$ and

$$\overline{\lim}_{m \rightarrow \infty} \mathbb{E}_x V(X_{m \wedge \tau_A \wedge \tau_{E_n^c}}) = \overline{\lim}_{m \rightarrow \infty} \mathbb{E}_x V(X_{\tau_A \wedge \tau_{E_n^c}}) \mathbf{1}_{\{m \geq \tau_A \wedge \tau_{E_n^c}\}} \leq \mathbb{E}_x V(X_{\tau_A \wedge \tau_{E_n^c}}).$$

Therefore,

$$\mathbb{E}_x(\tau_A \wedge \tau_{E_n^c}) \geq V(x) - \mathbb{E}_x V(X_{\tau_A \wedge \tau_{E_n^c}}). \quad (2.10)$$

Since W is nonnegative, by (2.9),

$$\mathbb{E}_x \left[W(X_{\tau_{E_n^c}}) \mathbf{1}_{\{\tau_A \geq \tau_{E_n^c}\}} \right] \leq W(x), \quad x \in E_n \setminus A.$$

Furthermore,

$$\begin{aligned} \mathbb{E}_x(\tau_A \wedge \tau_{E_n^c}) &\geq V(x) - \mathbb{E}_x V(X_{\tau_A \wedge \tau_{E_n^c}}) \\ &\geq V(x) - \mathbb{E}_x V(X_{\tau_A}) \mathbf{1}_{\{\tau_A < \tau_{E_n^c}\}} - \left(\sup_{y \in E_n^c} \frac{V(y)}{W(y)} \right) \mathbb{E}_x [W(X_{\tau_{E_n^c}}) \mathbf{1}_{\{\tau_A \geq \tau_{E_n^c}\}}] \\ &\geq V(x) - \mathbb{E}_x V(X_{\tau_A}) \mathbf{1}_{\{\tau_A < \tau_{E_n^c}\}} - \left(\sup_{y \in E_n^c} \frac{V(y)}{W(y)} \right) W(x), \quad x \in E_n \setminus A. \end{aligned}$$

By $\tau_{E_n^c} \uparrow \infty, n \rightarrow \infty$ and 1), we obtain

$$\sup_{x \in A^c} \mathbb{E}_x \tau_A \geq \sup_{x \in A^c} V(x) - \sup_{x \in A} V(x) = \infty,$$

which is contradict with $\sup_{x \in \mathcal{X}} \mathbb{E}_x \tau_A^+ < \infty$. Therefore, $\{X_n\}$ is non-strong ergodic. \square

2.3 Approximation for the moments of hitting times and necessity of Theorem 1.1 and Theorem 1.2

2.3.1 Approximation of Markov chain

Proposition 2.10 *Assume that Markov chain $\{X_n\}$ with Markov semigroup $\{P(x, \cdot)\}$ on locally compact separable metric space $(\mathcal{X}, \mathcal{B}(\mathcal{X}))$ is ψ -irreducible, aperiodic, Feller and $\supp \psi$ has*

nonempty interior. $E \subseteq \mathcal{X}$ is a compact set and $0 < \psi(E) < \infty$. For any $A \in \mathcal{B}^+(\mathcal{X})$, $A \subseteq E$ is a compact set, define $\{\hat{P}(x, \cdot)\}$ on $(E, \mathcal{B}(E))$:

$$\hat{P}(x, B) = \left[P(x, B) + P(x, E^c) \frac{\psi(B \cap A)}{\psi(A)} \right] \mathbf{1}_{E \setminus A}(x) + \frac{\psi(B)}{\psi(E)} \mathbf{1}_A(x), \quad B \in \mathcal{B}(E). \quad (2.11)$$

Then $\{\hat{P}(x, \cdot) : x \in E\}$ is a transition probability and the Markov chain generated by $\{\hat{P}(x, \cdot) : x \in E\}$ is ψ -irreducible, aperiodicity. Moreover, it is strongly ergodic.

Proof It is easy to verify that $\hat{P}(x, E) = 1$ and $\{\hat{P}(x, B), x \in E, B \in \mathcal{B}(E)\}$ is a transition probability on $(E, \mathcal{B}(E))$. Denoted by $\{\hat{X}_n\}$ the Markov chain on state space $(E, \mathcal{B}(E))$ with one step transition probability $\{\hat{P}(x, B), x \in E, B \in \mathcal{B}(E)\}$, denote $\hat{P}_1 = P$, $\hat{P}_n(x, B) = \mathbb{P}_x(\hat{X}_n \in B)$, $n \geq 2$ for $B \in \mathcal{B}(E)$.

We first prove that the chain $\{\hat{X}_n\}$ is ψ -irreducible, i.e., for $x \in E$, and $B \in \mathcal{B}(E)$ with $\psi(B) > 0$, there exists $n \geq 1$, $\hat{P}_n(x, B) > 0$ by [10, Proposition 4.2.1].

1) It is obvious that $\hat{P}(x, B) > 0$ for $x \in A$ and $\psi(B) > 0$, $B \in \mathcal{B}(E)$.

2) We prove that for $\sum_n \hat{P}_n(x, A) > 0$ for $x \in E \cap A^c$. (Equivalently, $x \in E \cap A^c$, there exists $n \geq 1$, $\hat{P}_n(x, A) > 0$.) If it fails, then there exists $x_0 \in E \cap A^c$, $\hat{P}_n(x_0, A) = 0$ for $n \geq 1$. We will find a contradiction in the following. Since

$$\hat{P}_1(x_0, A) = \hat{P}(x_0, A) = P(x_0, A) + P(x_0, E^c) = 0,$$

by (2.11), we have $P(x_0, E^c) = 0$ and $P(x_0, A) = 0$. Moreover, $P(x_0, A^c \cap E) = 1$, $\hat{P}(x_0, A^c \cap E) = 1$ and

$$\hat{P}(x_0, A^c \cap E \cap C) = P(x_0, A^c \cap E \cap C), \quad C \in \mathcal{B}(\mathcal{X}).$$

By Chapman-Kolmogorov equation, for $C \in \mathcal{B}(\mathcal{X})$,

$$\begin{aligned} \hat{P}_2(x_0, A^c \cap E \cap C) &= \int_{E \cap A^c} \hat{P}(x_0, dy) \hat{P}(y, A^c \cap E \cap C) \quad (\text{since } \hat{P}(x_0, A^c \cap E) = 1) \\ &= \int_{E \cap A^c} P(x_0, dy) P(y, A^c \cap E \cap C) \quad (\text{by (2.11)}) \\ &= P_2(x_0, E \cap A^c \cap C) \quad (\text{since } P(x_0, A^c \cap E) = 1). \end{aligned}$$

Similarly, $\hat{P}_2(x, E^c) = 0$ by $P(x_0, E^c) = 0$ and Chapman-Kolmogorov equation. Since $\hat{P}_2(x_0, A) = 0$, we get $\hat{P}_2(x_0, A^c \cap E) = P_2(x_0, E \cap A^c) = 1$. Generally, assume that

$$\hat{P}_n(x_0, A^c \cap E \cap C) = P_n(x_0, A^c \cap E \cap C), \quad C \in \mathcal{B}(\mathcal{X}), \quad (2.12)$$

and $\hat{P}_n(x_0, A^c \cap E) = P_n(x_0, E \cap A^c) = 1$ for $n \leq k$. Then

$$\begin{aligned}\hat{P}_{k+1}(x_0, A^c \cap E \cap C) &= \int_E \hat{P}_k(x_0, dy) \hat{P}(y, A^c \cap E \cap C) \\ &= \int_{E \cap A^c} \hat{P}_k(x_0, dy) \hat{P}(y, A^c \cap E \cap C) \quad (\text{since } \hat{P}_k(x_0, A) = 0) \\ &= \int_{E \cap A^c} P_k(x_0, dy) P(y, A^c \cap E \cap C) \quad (\text{by (2.11) and (2.12)}) \\ &= P_{k+1}(x_0, E \cap A^c \cap C) \quad (\text{since } P_k(x_0, A^c \cap E) = 1)\end{aligned}$$

Hence, if there exists $x_0 \in E \cap A^c$, $\hat{P}_n(x_0, A) = 0$ for $n \geq 1$, then $\hat{P}_n(x_0, A^c \cap E) = P_n(x_0, E \cap A^c) = 1$, $n \geq 1$. Furthermore,

$$\sum_{n=1}^{\infty} P_n(x_0, E^c \cup A) = 0,$$

which is contradiction with the irreducible property of $\{X_n\}$.

3) We prove that $\sum_n \hat{P}_n(x, B) > 0$ for $x \in E \cap A^c$ and $B \in \mathcal{B}(E)$ with $\psi(B) > 0$. Indeed, by 2), there exists $n \geq 1$, $\hat{P}_n(x, A) > 0$. Moreover,

$$\hat{P}_{n+1}(x, B) \geq \int_A \hat{P}_n(x, dy) \hat{P}(y, B) = \frac{\psi(B)}{\psi(E)} \hat{P}_n(x, A) > 0.$$

Therefore, $\{\hat{X}_n\}$ is ψ -irreducible.

Since $A \subseteq E$ is a compact set, $\{\hat{X}_n\}$ is weak Feller by definition obviously. Therefore, it suffices to prove that $\{\hat{X}_n\}$ is aperiodicity according to [10, Theorems 6.2.9, 16.2.5]. Let measure $\hat{\nu}_1(\cdot) = \psi(\cdot)/\psi(E)$. Then $\hat{P}(x, B) = \nu_1(B)$ for $x \in A$, $B \in \mathcal{B}(E)$, we get A is ν_1 -small by definition (see [10, Section 5.2]). Since, $\nu_1(A) > 0$, we get $\{\hat{X}_n\}$ is strongly aperiodic by definition (see [10, Section 5.4.3]). Hence the required assertion holds. \square

Example 2.11 Consider $X_{n+1} = aX_n + W_{n+1}$ on \mathbb{R} with $X_0 = 0$, $|a| < 1$. Then for any $m \geq 1$, $E_m := [-m + \frac{1}{m}, m - \frac{1}{m}] \uparrow \mathbb{R}$, its one step transition kernel is $\{P(x, \cdot) : x \in \mathbb{R}\}$ satisfying $P(x, dy)$ has intensity function $p(x, y) = \frac{1}{\sqrt{2\pi}} \exp\{-\frac{(y-ax)^2}{2}\}$. Let $A = E_1$, $\mu_{Leb}(A) > 0$. Define

$$p^{(m)}(x, y) = \begin{cases} p(x, y) + \frac{P(x, E_m^c)}{\mu_{Leb}(A)}, & x \in E_m \setminus A; \\ \frac{1}{\mu_{Leb}(E_m)}, & x \in A; \end{cases} \quad (2.13)$$

Denoted by $P^{(m)}(x, dy) = p^{(m)}(x, y)dy$. Then $\{P^{(m)}(x, dy)\}$ is a transition kernel on E_m . Let $\{X_n^{(m)}\}$ be Markov chain generated by $\{P^{(m)}(x, \cdot)\}$. Then it is easy to verify that $\{P^{(m)}(x, dy)\}$ is irreducible with respect to Lebesgue measure on E_m . Furthermore, $\{X_n\}$ is strongly ergodic by [10, Theorems 6.2.9, 16.2.5].

Remark 2.12 It is easy to see that $\{p^{(m)}(x, y)\}$ in (2.13) is increasing in m to $P(x, y)$. Therefore, we can verify some properties of $\{X_n\}$ generated by $p(x, \cdot)$ by approximation method according to the description of Markov chain generated by $\{p^{(m)}(x, \cdot)\}$.

2.3.2 Necessity

In this subsection, the locally compact separable metric space $\mathcal{X} = \cup_{m=1}^{\infty} E_m$, where $\{E_m\}_{m \geq 1}$ is a sequence of increasing compact sets and Markov chain $\{X_n\}$ with Markov semigroup $\{P(x, \cdot)\}$ on $(\mathcal{X}, \mathcal{B}(\mathcal{X}))$ always satisfies Assumption 1 (i.e., ψ -irreducible, aperiodic, Feller and $\text{supp}\psi$ has nonempty interior).

The following Lemma holds obviously by Proposition 2.10.

Lemma 2.13 *$A \in \mathcal{B}^+(E_1)$ is a compact set. $\{X_n^{(m)} : n \geq 0\}$ is a sequence of Markov chains on the state space E_m respectively, each of which has one step transition semigroup $P^{(m)}(x, \cdot)$ defined as \hat{P} in (2.11) by replacing E there with E_m . Then Markov chain $\{X_n^{(m)}\}$ on $(E_m, \mathcal{B}(E_m))$ is strongly ergodic.*

For a set $B \in \mathcal{B}(E_m)$, define the Markov chain $\{X_n^{(m)}\}$ as that in Lemma 2.13, set $\tau_B^{(m)+} = \inf\{n \geq 1 : X_n^{(m)} \in B\}$. Notice $\mathbb{P}_x(\tau_B^{(m)+} < \infty) = 1$, $x \in E_m$. Consider the equation

$$V(x) = \int_{B^c \cap E_m} V(y) P^{(m)}(x, dy) + 1, \quad x \in E_m. \quad (2.14)$$

By Proposition 2.6, the minimal solution to (2.14) is

$$V_B^{*(m)}(x) := \mathbb{E}_x \tau_B^{(m)+}, \quad x \in E_m.$$

Denoted by

$$M_m(B) = \sup_{x \in B^c \cap E_m} V_B^{*(m)}(x) = \sup_{x \in B^c \cap E_m} \mathbb{E}_x \tau_B^{(m)+}.$$

By Proposition 2.10, we have the following assertion immediately.

Lemma 2.14 *$A \in \mathcal{B}^+(E_1)$ is a compact set and $\{X_n\}$ is recurrent. Then the following assertions hold:*

- (1) $M_m(A)$ is finite for $m \geq 1$.
- (2) $\mathbb{E}_x \tau_A^+ = \lim_{m \rightarrow \infty} \uparrow V_A^{*(m)}(x) \mathbf{1}_{E_m}(x)$ for $x \in \mathcal{X}$ and $\{M_m(A)\}_{m \geq 1}$ is increasing.
- (3) $\{M_m(A)\}_{m \geq 1}$ is bounded if and only if $\{\mathbb{E}_x \tau_A^+ : x \in A^c\}$ is bounded. That's $\sup_{x \in A^c} \mathbb{E}_x \tau_A^+ < \infty$ if and only if $\sup_{m \geq 1} M_m(A) < \infty$.

Proof (1) Since $\{X_n^{(m)}\}$ is strong ergodicity by Lemma 2.13, $A \in \mathcal{B}^+(E_m)$, we get $L^{(m)}(x, A) = 1$ and $M_m(A) < \infty$.

(2) Since $A \subset E_1$ and $\{X_n^{(m)}\}$ is Harris recurrent, by Proposition 2.6, $V_A^{*(m)}(x)$, $x \in E_m$ is the minimal solution to

$$V(x) = \int_{E_m \cap A^c} V(y) P^{(m)}(x, dy) + 1, \quad x \in E_m.$$

Equivalently, $\tilde{V}_A^{*(m)}(x) := V_A^{*(m)}(x)\mathbf{1}_{E_m}(x)$ is the minimal solution to

$$V(x) = \int_{E_m \cap A^c} V(y)P^{(m)}(x, dy)\mathbf{1}_{E_m}(x) + \mathbf{1}_{E_m}(x), \quad x \in \mathcal{X}.$$

Since $E_m \uparrow \mathcal{X}$,

$$\int_{A^c} V(y)\mathbf{1}_{E_m}(y)P^{(m)}(x, dy)\mathbf{1}_{E_m}(x) \uparrow \int_{A^c} V(y)P(x, dy), \quad m \rightarrow \infty, \quad x \in \mathcal{X}.$$

Notice that $\mathbb{E}_x \tau_A^+$ is the minimal solution to the following equation:

$$V(x) = \int_{A^c} V(y)P(x, dy) + 1, \quad x \in \mathcal{X},$$

we have $\lim_{m \rightarrow \infty} \uparrow \tilde{V}_A^{*(m)}(x) = \mathbb{E}_x \tau_A^+$, $x \in \mathcal{X}$ by Lemma 2.3. Therefore, $V_A^{*(m)}(x)\mathbf{1}_{E_m}(x) \uparrow \mathbb{E}_x \tau_A^+$ for $x \in \mathcal{X}$ and $M_m(A)$ is increasing in m .

(3) By (2), the assertion holds obviously. \square

Proof of necessity of Theorem 1.1 Suppose that $\{X_n\}$ is recurrent but not ergodic, for any petite sets $C \in \mathcal{B}^+(\mathcal{X})$,

$$\sup_{x \in C} \mathbb{E}_x \tau_C^+ = \infty, \quad \int_C \mathbb{E}_x \tau_C^+ \psi(dx) = \infty.$$

For any compact set $A \in \mathcal{B}^+(\mathcal{X})$, without loss of generality, let $A \subset E_1$. Denoted by

$$W^{(m)}(x) := \begin{cases} 0, & x \in E_m^c, \\ \mathbb{E}_x \tau_A^{(m)+}, & x \in E_m \cap A^c, \\ \int_{E_m \cap A^c} \mathbb{E}_y \tau_A^{(m)+} P(x, dy) + 1, & x \in A. \end{cases}$$

If $x \in E_m \cap A^c$,

$$\int_{A^c} W^{(m)}(y)P(x, dy) = \int_{E_m \cap A^c} W^{(m)}(y)P^{(m)}(x, dy) = \mathbb{E}_x \tau_A^{(m)+} - 1,$$

if $x \in A$,

$$\int_{A^c} W^{(m)}(y)P(x, dy) = \int_{E_m \cap A^c} \mathbb{E}_y \tau_A^{(m)+} P(x, dy) = W^{(m)}(x) - 1,$$

and if $x \in E_m^c$, due to the non-negativity of $W^{(m)}$,

$$\int_{A^c} W^{(m)}(y)P(x, dy) \geq 0,$$

hence, Condition 2) holds for $x \in \mathcal{X}$.

From Lemma 2.14,

$$\sup_{x \in A^c} W^{(m)}(x) = \sup_{x \in E_m \cap A^c} \mathbb{E}_x \tau_A^{(m)+} < \infty,$$

which means Condition 1) holds.

Notice that, $\{X_n\}$ is not ergodic,

$$\int_A \sup_{m \geq 1} W^{(m)}(x) \psi(dx) = \int_A \mathbb{E}_x \tau_A^+ \psi(dx) = \infty,$$

that is to say Condition 3) holds.

Based on these, $\{W^{(m)}(x) : m \geq 1\}$ is the sequence of functions that we need for necessity.

□

Proof of necessity of Theorem 1.2 Since the chain $\{X_n\}$ is non-strongly ergodic, there exists $A \in \mathcal{B}^+(\mathcal{X})$, $\sup_{x \in \mathcal{X}} \mathbb{E}_x \tau_A^+ = \infty$. Without loss of generality, we can also assume that A is a compact set (Otherwise, since $\sup_{x \in \mathcal{X}} \mathbb{E}_x \tau_{E_m}^+ = \infty, m \geq 1$ by [10, Theorem 16.0.2, Proposition 6.2.8], one can replace A by the closure of A). Since $\{P^{(m)}(x, \cdot)\}$ on $(E_m, \mathcal{B}(E_m))$ is strong ergodicity, $\sup_{x \in E_m \cap A^c} V_A^{*(m)}(x) < \infty$. Then $W^{(m)}(x) := \mathbf{1}_{E_m \cap A^c}(x) V_A^{*(m)}(x)$, $x \in \mathcal{X}$ is the required sequence. Indeed, it is obvious that $\sup_{x \in A^c} W^{(m)}(x) < \infty$, and

$$\begin{aligned} W^{(m)}(x) &= \int_{A^c \cap E_m} V(y) P^{(m)}(x, dy) + 1 \\ &= \int_{A^c \cap E_m} W^{(m)}(y) P(x, dy) + 1, \quad x \in E_m \cap A^c. \end{aligned}$$

Furthermore, $W^{(m)}(x) \leq P W^{(m)}(x) + 1, x \in A^c$. By Lemma 2.14,

$$\lim_{m \rightarrow \infty} \uparrow W^{(m)}(x) = \mathbb{E}_x(\tau_A^+ \mathbf{1}_{\{\tau_A^+ < \infty\}}), \quad x \in A^c.$$

Hence, $\sup_{x \in A^c, m \geq 1} \uparrow W^{(m)}(x) = \infty$. □

Acknowledgments

This work is supported by the National Natural Science Foundation of China (No. 12101186, No. 12171038) and the National Key R&D Program of China (No. 2020YFA0712900).

References

- [1] B.D. Choi, B. Kim, Non-ergodicity criteria for denumerable continuous time Markov process, Operations Research Letters. 32(6) (2004) 574-580.

- [2] M.-F. Chen, From Markov Chains to Non-Equilibrium Particle Systems. World Scientific, Singapore, 2nd edition, 2004.
- [3] S.F. Jarner, G.O. Roberts, Polynomial convergence rates of markov chains. *Annals of Applied Probability*. 12(1) (2002) 224-247.
- [4] O.S. Lee, Geometric ergodicity and transience for nonlinear autoregressive monels. *Communications of the Korean Mathematical Society*. 10(2) (1995) 409-417.
- [5] Y.-H. Mao, Algebraic convergence for discrete-time ergodic Markov chains, *Sci. China Ser. A*. 46(5) (2003) 621-630.
- [6] Y.-H. Mao, T. Wang, Lyapunov-type Conditions for Non-strong Ergodicity of Markov Processes, *Journal of Applied Probability*. (58) (2021) 238-253.
- [7] Y.-H. Mao, Y.-H. Song, On geometric and algebraic transience for discrete-time Markov chains. *Stochastic Processes and Their Applications* 124(4) (2014) 1648-1678.
- [8] S.P. Meyn, R.L. Tweedie, Stability of Markovian processes I: criteria for discrete-time chains. *Advances in Applied Probability* 24(3) (1992) 542-574.
- [9] S.P. Meyn, R.L. Tweedie, Stability of Markovian Processes III: Foster-Lyapunov Criteria for Continuous-Time Processes. *Advances in Applied Probability*. 25(3) (1993) 518-548.
- [10] S.P. Meyn, R L. Tweedie, *Markov Chains and Stochastic Stability*(Second edition), Cambridge press, 2009.
- [11] L.I. Sennott, H.R.L. Tweedie, Mean drifts and the non-ergodicity of Markov chains. *Operations Research*. 31(4) (1983) 783-789.
- [12] W. Szpankowski, Some sufficient conditions for non-ergodicity of Markov chains. *Journal of Applied Probability*. 22(1) (1985) 138-147.
- [13] D. Tjøstheim, Non-linear time series and Markov chains. *Advances in Applied Probability*, 22 (1990) 587-611.
- [14] Z.F. Wei, Inverse Problems for Ergodicity of Markov Chains, *Journal of Mathematical Analysis and Applicationss*. 505 (2022) 125483.