

# LEIBNIZ CONFORMAL BIALGEBRAS AND THE CLASSICAL LEIBNIZ CONFORMAL YANG-BAXTER EQUATION

ZHONGYIN XU, CHENGMING BAI, AND YANYONG HONG

**ABSTRACT.** We introduce the notion of Leibniz conformal bialgebras, presenting a bialgebra theory for Leibniz conformal algebras as well as the conformal analogues of Leibniz bialgebras. They are equivalently characterized in terms of matched pairs and conformal Manin triples of Leibniz conformal algebras. In the coboundary case, the classical Leibniz conformal Yang-Baxter equation is introduced, whose symmetric solutions give Leibniz conformal bialgebras. Moreover, such solutions are constructed from  $\mathcal{O}$ -operators on Leibniz conformal algebras and Leibniz-dendriform conformal algebras. On the other hand, the notion of Novikov bi-dialgebras is introduced, which correspond to a class of Leibniz conformal bialgebras, lifting the correspondence between Novikov dialgebras and a class of Leibniz conformal algebras to the context of bialgebras. In addition, we introduce the notion of classical duplicate Novikov Yang-Baxter equation whose symmetric solutions produce Novikov bi-dialgebras and thus Leibniz conformal bialgebras.

## CONTENTS

1. Introduction	2
2. Preliminaries on Leibniz conformal algebras	4
2.1. Leibniz conformal algebras and Novikov dialgebras	4
2.2. Matched pairs of Leibniz conformal algebras	6
2.3. Conformal Manin triples of Leibniz conformal algebras	8
3. Leibniz conformal bialgebras	10
4. Coboundary Leibniz conformal bialgebras and classical Leibniz conformal Yang-Baxter equation	13
4.1. Coboundary Leibniz conformal bialgebras	14
4.2. Operator forms of the classical Leibniz conformal Yang-Baxter equation	17
5. A class of Leibniz conformal bialgebras corresponding to Novikov bi-dialgebras	23
5.1. Novikov bi-dialgebras	23
5.2. The classical duplicate Novikov Yang-Baxter equation	25
References	29

---

2010 *Mathematics Subject Classification.* 17A32, 17A60, 17B38, 17B69.

*Key words and phrases.* Leibniz conformal bialgebra, Leibniz-dendriform conformal algebra, classical Leibniz conformal Yang-Baxter equation, Novikov dialgebra, Novikov bi-dialgebra, Rota-Baxter operator.

## 1. INTRODUCTION

The notion of Lie conformal algebras was introduced by Kac in [22, 23] as a formal language describing the algebraic properties of the operator product expansions of chiral fields in two-dimensional conformal field theory. Moreover, Lie conformal algebras are closely related with infinite-dimensional Lie algebras satisfying the locality property [24], local Poisson brackets in the theory of integrable evolution equations [3] and so on. The structure theory [11], representation theory [9, 10] and cohomology theory [2, 13] of finite Lie conformal algebras have been studied and well developed.

Leibniz conformal algebras introduced in [2] are the conformal analogues of Leibniz algebras which were first discovered by Bloh and named D-algebras [6] and then rediscovered by Loday [27, 28]. Leibniz conformal algebras are non-commutative generalizations of Lie conformal algebras and have close connections to field algebras [1]. From the perspective of algebraic structures, a Leibniz conformal algebra is a Leibniz pseudoalgebra [31] which is a Leibniz algebra in the pseudo-tensor category [4]. Thus Leibniz conformal algebras can also be seen as generalizations of Leibniz algebras. Recently, Leibniz conformal algebras have been widely studied, for example, quadratic Leibniz conformal algebras and their central extensions [36], the cohomology theory of Leibniz conformal algebras [35] and so on. For more details, see [16, 12, 21, 33, 32]. It is well known that Lie bialgebras have important applications in both mathematics and mathematical physics [8, 14]. A natural idea is to study “conformal analogues” of bialgebras. Liberati developed the theory of Lie conformal bialgebras in [26] as the conformal analogues of Lie bialgebras, which includes the introduction of the notions of conformal classical Yang-Baxter equation, conformal Manin triples and conformal Drinfeld’s doubles. The operator forms of conformal classical Yang-Baxter equation were investigated in [18]. Similarly, the theories of antisymmetric infinitesimal conformal bialgebras and left-symmetric conformal bialgebras were developed in [17, 20]. However, there is no conformal analogue of “Leibniz bialgebras” as far as we know. Thus the purpose of this paper is to investigate a bialgebra theory for Leibniz conformal algebras, namely, Leibniz conformal bialgebras, giving the conformal analogues of Leibniz bialgebras. That is, the following diagram is commutative.

$$(1) \quad \begin{array}{ccc} \text{Leibniz algebras} & \xrightarrow{\text{conformal analogues}} & \text{Leibniz conformal algebras} \\ \downarrow \text{bialgebras} & & \downarrow \text{bialgebras} \\ \text{Leibniz bialgebras} & \xrightarrow{\text{conformal analogues}} & \text{Leibniz conformal bialgebras} \end{array}$$

It is worth mentioning that [30], [29] and [5] have defined three different kinds of Leibniz bialgebras respectively and our main idea is to extend the study of Leibniz bialgebras given in [30].

In order to characterize Leibniz conformal bialgebras, we introduce the notion of matched pairs of Leibniz conformal algebras and conformal Manin triples of Leibniz conformal algebras. That is, under some conditions, Leibniz conformal bialgebras are equivalent to some matched pairs of Leibniz conformal algebras as well as conformal Manin triples of Leibniz conformal algebras. In the study of a special class of Leibniz conformal bialgebras called coboundary Leibniz conformal bialgebras, we introduce the notion of classical Leibniz conformal Yang-Baxter equation (CLCYBE), as a conformal analogue of the classical Leibniz Yang-Baxter equation.

Notably, its symmetric solutions directly give Leibniz conformal bialgebras. In addition, the CLCYBE is interpreted in terms of its operator forms by  $\mathcal{O}$ -operators on Leibniz conformal algebras. In particular, a symmetric solution of the CLCYBE corresponds to the symmetric part of a conformal linear map  $T$ , where  $T_0 = T_\lambda|_{\lambda=0}$  is an  $\mathcal{O}$ -operator in the conformal sense, which implies that an  $\mathcal{O}$ -operator on a Leibniz conformal algebra gives a symmetric solution of the CLCYBE in a semi-direct product Leibniz conformal algebra. Furthermore, we introduce the notion of Leibniz-dendriform conformal algebras and show that for a Leibniz-dendriform conformal algebra which is finite and free as a  $\mathbf{k}[\partial]$ -module, there is a natural  $\mathcal{O}$ -operator on the associated Leibniz conformal algebra. Therefore there are constructions of symmetric solutions of the CLCYBE and hence Leibniz conformal bialgebras from  $\mathcal{O}$ -operators on Leibniz conformal algebras as well as Leibniz-dendriform conformal algebras. In summary, they are depicted in the diagram



On the other hand, based on the result on [36], we show that there is the following correspondence between Novikov dialgebras and a class of Leibniz conformal algebras.

$$(2) \quad \text{A class of Leibniz conformal algebras} \xleftrightarrow{\text{Proposition 2.8}} \text{Novikov dialgebras}$$

This fact generalizes the known correspondence between Novikov algebras and a class of Lie conformal algebras [34].

$$(3) \quad \text{A class of Lie conformal algebras} \longleftrightarrow \text{Novikov algebras}$$

Note that the notion of Novikov dialgebras was introduced in [25] and perm algebras with derivations can naturally produce Novikov dialgebras [25]. Moreover, it is worth mentioning that the correspondence given in (3) was lifted to the context of bialgebras in [19], which shows that Novikov bialgebras can produce Lie conformal bialgebras. Hence it is natural to introduce the notion of Novikov bi-dialgebras and show that there is a correspondence between Novikov bi-dialgebras and a class of Leibniz conformal bialgebras, lifting the correspondence given in (2) to the context of bialgebras. Moreover, we introduce the notion of classical duplicate Novikov Yang-Baxter equation (CDNYBE), whose symmetric solutions produce Novikov bi-dialgebras and show that symmetric solutions of the CDNYBE correspond to a class of solutions of the CLCYBE.

This paper is organized as follows. In Section 2, we recall the definitions of Leibniz conformal algebras and their representations, and show that there is a correspondence between Novikov dialgebras and a class of Leibniz conformal algebras. Furthermore, we introduce the definitions of matched pairs of Leibniz conformal algebras and conformal Manin triples of Leibniz conformal algebras, and show that under some conditions, a special matched pair of Leibniz conformal algebras is equivalent to a conformal Manin triple of Leibniz conformal algebras. In Section 3, the notion of Leibniz conformal bialgebras is introduced and we show that under some conditions, there is an equivalence between conformal Manin triples of Leibniz conformal algebras and Leibniz conformal bialgebras. In Section 4, the notion of CLCYBE is introduced in the study of a class of Leibniz conformal bialgebras called coboundary Leibniz

conformal bialgebras. In addition, we introduce the definitions of  $\mathcal{O}$ -operators on Leibniz conformal algebras and Leibniz-dendriform conformal algebras to construct (symmetric) solutions of the CLCYBE and hence give rise to Leibniz conformal bialgebras. In Section 5, we define Novikov bi-dialgebras and show that there is a correspondence between Novikov bi-dialgebras and a class of Leibniz conformal bialgebras. Furthermore, we introduce the notion of CDNYBE and show that its symmetric solutions can produce Novikov bi-dialgebras and thus Leibniz conformal bialgebras.

Throughout this paper, let  $\mathbf{k}$  be a field of characteristic zero. All tensors over  $\mathbf{k}$  are denoted by  $\otimes$ . We denote the identity map by  $\text{id}$ . Moreover, if  $A$  is a vector space, then we denote the space of polynomials of  $\lambda$  with coefficients in  $A$  by  $A[\lambda]$ . For a vector space or a  $\mathbf{k}[\partial]$ -module  $A$ , let

$$\tau : A \otimes A \rightarrow A \otimes A, \quad a \otimes b \mapsto b \otimes a \quad \text{for all } a, b \in A,$$

be the flip operator. Then we denote  $\tau_{12} = \tau \otimes \text{id}$ ,  $\tau_{23} = \text{id} \otimes \tau$ ,  $\tau_{13} = (\tau \otimes \text{id})(\text{id} \otimes \tau)(\tau \otimes \text{id})$ .

## 2. PRELIMINARIES ON LEIBNIZ CONFORMAL ALGEBRAS

We recall some facts about Leibniz conformal algebras and show that there is a correspondence between Novikov dialgebras and a class of Leibniz conformal algebras. Moreover, we introduce the notions of dual representations, matched pairs and conformal Manin triples of Leibniz conformal algebras and show that under some conditions, some matched pair of Leibniz conformal algebras is equivalent to a conformal Manin triple of Leibniz conformal algebras.

### 2.1. Leibniz conformal algebras and Novikov dialgebras.

**Definition 2.1.** [1] A **Leibniz conformal algebra**  $Q$  is a  $\mathbf{k}[\partial]$ -module with a  $\lambda$ -bracket  $[\cdot, \cdot]_\lambda$  which is a  $\mathbf{k}$ -bilinear map from  $Q \times Q \rightarrow Q[\lambda]$ , satisfying

$$\begin{aligned} \text{(conformal sesquilinearity)} \quad & [\partial a_\lambda b] = -\lambda[a_\lambda b], & [a_\lambda \partial b] &= (\lambda + \partial)[a_\lambda b], \\ \text{(Jacobi identity)} \quad & [a_\lambda [b_\mu c]] = [[a_\lambda b]_{\lambda+\mu} c] + [b_\mu [a_\lambda c]] \quad \text{for all } a, b, c \in Q. \end{aligned}$$

A Leibniz conformal algebra  $Q$  is called **finite**, if it is finitely generated as a  $\mathbf{k}[\partial]$ -module. Otherwise, we call it **infinite**.

**Remark 2.2.** Recall [22] that a **Lie conformal algebra**  $Q$  is a Leibniz conformal algebra satisfying

$$\text{(skew-symmetry)} \quad [a_\lambda b] = -[b_{-\lambda-\partial} a] \quad \text{for all } a, b \in Q.$$

Thus Lie conformal algebras are Leibniz conformal algebras.

Recall [27] that a **Leibniz algebra**  $(A, [\cdot, \cdot])$  is a vector space  $A$  with a bilinear operation  $[\cdot, \cdot] : A \times A \rightarrow A$ , satisfying

$$[a, [b, c]] = [[a, b], c] + [b, [a, c]] \quad \text{for all } a, b, c \in A.$$

**Example 2.3.** Let  $(A, [\cdot, \cdot])$  be a Leibniz algebra. Then we can naturally define a Leibniz conformal algebra structure on  $\text{Cur } A = \mathbf{k}[\partial] \otimes A$  with the  $\lambda$ -bracket:

$$[a_\lambda b] = [a, b] \quad \text{for all } a, b \in A.$$

$\text{Cur } A$  is called the **current Leibniz conformal algebra** associated with  $(A, [\cdot, \cdot])$ .

**Proposition 2.4.** [21] *Let  $Q = \mathbf{k}[\partial]L$  be a Leibniz conformal algebra which is a free  $\mathbf{k}[\partial]$ -module of rank 1. Then  $Q$  is either abelian or isomorphic to the Virasoro Lie conformal algebra, that is,  $[L_\lambda L] = (\partial + 2\lambda)L$ .*

Next, we recall the definition of Novikov dialgebras.

**Definition 2.5.** [25] *Let  $A$  be a vector space with two binary operations  $\vdash$  and  $\dashv$ . If for all  $a, b, c \in A$ , they satisfy the following equalities*

$$\begin{aligned} (a \vdash b) \dashv c &= (a \vdash c) \vdash b = (a \dashv c) \vdash b, & a \dashv (b \vdash c) &= a \dashv (b \vdash c), \\ (a \dashv b) \dashv c &= (a \dashv c) \dashv b, & (a \dashv b) \dashv c - a \dashv (b \vdash c) &= (b \vdash a) \dashv c - b \vdash (a \dashv c), \\ (a \vdash b) \vdash c - a \vdash (b \vdash c) &= (b \vdash a) \vdash c - b \vdash (a \vdash c), \end{aligned}$$

then  $(A, \dashv, \vdash)$  is called a **(left) Novikov dialgebra**.

**Remark 2.6.** There is also the definition of right Novikov dialgebras. A **right Novikov dialgebra** is a triple  $(A, \dashv, \vdash)$  where  $A$  is a vector space with two binary operations satisfying

$$\begin{aligned} a \vdash (b \vdash c) &= b \vdash (a \vdash c), & a \vdash (b \dashv c) &= b \dashv (a \vdash c), & (a \vdash b - a \dashv b) \dashv c &= 0, \\ a \dashv (b \vdash c - b \dashv c) &= 0, & a \vdash (b \vdash c - c \dashv b) &= (a \vdash b) \vdash c - (a \vdash c) \dashv b, \\ a \dashv (b \vdash c - c \dashv b) &= (a \dashv b) \dashv c - (a \dashv c) \dashv b & \text{ for all } a, b, c \in A. \end{aligned}$$

Note that such algebra structures appear in the study of quadratic Leibniz conformal algebras [36]. Moreover, it is easy to see that  $(A, \dashv, \vdash)$  is a Novikov dialgebra if and only if  $(A, \dashv', \vdash')$  is a right Novikov dialgebra where  $\vdash'$  and  $\dashv'$  are defined by

$$a \dashv' b := b \vdash a, \quad a \vdash' b := b \dashv a \quad \text{for all } a, b \in A.$$

Recall that a **perm algebra** is a pair  $(A, \cdot)$ , where  $A$  is a vector space,  $\cdot$  is a binary operation and the following equalities hold.

$$(4) \quad (a \cdot b) \cdot c = a \cdot (b \cdot c) = b \cdot (a \cdot c) \quad \text{for all } a, b, c \in A.$$

**Lemma 2.7.** [25] *Let  $(A, \cdot)$  be a perm algebra and  $D$  be a derivation, that is,  $D$  satisfies  $D(a \cdot b) = D(a) \cdot b + a \cdot D(b)$  for all  $a, b \in A$ . Define two binary operations  $\vdash$  and  $\dashv$  by*

$$(5) \quad a \vdash b := a \cdot D(b), \quad a \dashv b := D(b) \cdot a \quad \text{for all } a, b \in A.$$

Then  $(A, \dashv, \vdash)$  is a Novikov dialgebra.

Next, we give a correspondence between Novikov dialgebras and a class of Leibniz conformal algebras.

**Proposition 2.8.** *Let  $(A, \dashv, \vdash)$  be a vector space with binary operations  $\vdash$  and  $\dashv$ . Equip the free  $\mathbf{k}[\partial]$ -module  $Q := \mathbf{k}[\partial]A (= \mathbf{k}[\partial] \otimes A)$  with the  $\lambda$ -bracket*

$$(6) \quad [a_\lambda b] := \partial(b \dashv a) + \lambda(a \vdash b + b \dashv a) \quad \text{for all } a, b \in A.$$

Then  $(Q, [\cdot, \cdot]_\lambda)$  is a Leibniz conformal algebra if and only if  $(A, \dashv, \vdash)$  is a Novikov dialgebra. We call  $Q$  the **Leibniz conformal algebra corresponding to  $(A, \dashv, \vdash)$** .

*Proof.* Define

$$(7) \quad [a_\lambda b] := \partial(a \vdash' b) + \lambda(a \vdash' b + b \dashv' a) \quad \text{for all } a, b \in A,$$

where  $\vdash'$  and  $\dashv'$  are two binary operations on  $A$ . By [36, Theorem 3.1],  $(Q, [\cdot, \cdot])$  is a Leibniz conformal algebra if and only if  $(A, \dashv', \vdash')$  is a right Novikov dialgebra. Set

$$a \dashv b := b \vdash' a, \quad a \vdash b := b \dashv' a \quad \text{for all } a, b \in A.$$

By Remark 2.6,  $(A, \dashv', \vdash')$  is a right Novikov dialgebra if and only if  $(A, \dashv, \vdash)$  is a Novikov dialgebra. Then the conclusion follows.  $\square$

**Corollary 2.9.** *Let  $(A, \cdot)$  be a perm algebra and  $D$  be a derivation on  $(A, \cdot)$ . Equip the free  $\mathbf{k}[\partial]$ -module  $Q := \mathbf{k}[\partial]A (= \mathbf{k}[\partial] \otimes A)$  with the  $\lambda$ -bracket*

$$(8) \quad [a_\lambda b] := \partial(D(a) \cdot b) + \lambda(a \cdot D(b) + D(a) \cdot b) \quad \text{for all } a, b \in A.$$

*Then  $(Q, [\cdot, \cdot])$  is a Leibniz conformal algebra.*

*Proof.* Define two binary operations  $\vdash$  and  $\dashv$  by (5). By Lemma 2.7,  $(A, \dashv, \vdash)$  is a Novikov dialgebra. Then it follows directly from Proposition 2.8.  $\square$

## 2.2. Matched pairs of Leibniz conformal algebras.

**Definition 2.10.** [22] Let  $U$  and  $V$  be two  $\mathbf{k}[\partial]$ -modules. A **conformal linear map** from  $U$  to  $V$  is a  $\mathbf{k}$ -linear map  $a_\lambda : U \rightarrow V[\lambda]$  such that  $a_\lambda(\partial u) = (\lambda + \partial)a_\lambda u$  for all  $u \in U$ . The vector space of all conformal linear maps from  $U$  to  $V$ , denoted by  $\text{Chom}(U, V)$ , has a  $\mathbf{k}[\partial]$ -module structure given by

$$(\partial a)_\lambda u = -\lambda a_\lambda u \quad \text{for all } a \in \text{Chom}(U, V) \text{ and } u \in U.$$

In particular, if  $U = V$ , then we denote  $\text{Chom}(U, V)$  by  $\text{Cend}(V)$ . If  $V$  is a finite  $\mathbf{k}[\partial]$ -module, then  $\text{Cend}(V)$  is a Lie conformal algebra defined by

$$[f_\lambda g]_\mu v = f_\lambda(g_{\mu-\lambda} v) - g_{\mu-\lambda}(f_\lambda v) \quad \text{for all } f, g \in \text{Cend}(V) \text{ and } v \in V.$$

We define the conformal dual of a  $\mathbf{k}[\partial]$ -module  $U$  as  $U^{*c} = \text{Chom}(U, \mathbf{k})$ , where  $\mathbf{k}$  is viewed as the trivial  $\mathbf{k}[\partial]$ -module, that is,  $U^{*c} = \{a : U \rightarrow \mathbf{k}[\lambda] \mid \mathbf{k}\text{-linear and } a_\lambda(\partial u) = \lambda a_\lambda u \text{ for all } u \in U\}$ . For convenience in the following, we set  $\langle a, u \rangle_\lambda = a_\lambda u$  for all  $a \in U^{*c}$  and  $u \in U$ .

Let  $U, V$  and  $W$  be  $\mathbf{k}[\partial]$ -modules. A **conformal bilinear map** from  $U \times V$  to  $W$  is a  $\mathbf{k}$ -bilinear map  $f_\lambda : U \times V \rightarrow W[\lambda]$ , such that  $f_\lambda(\partial u, v) = -\lambda f_\lambda(u, v)$  and  $f_\lambda(u, \partial v) = (\lambda + \partial)f_\lambda(u, v)$  for all  $u \in U$  and  $v \in V$ .

**Definition 2.11.** [35] A **representation** of a Leibniz conformal algebra  $(Q, [\cdot, \cdot])$  is a triple  $(M, l, r)$ , where  $M$  is a  $\mathbf{k}[\partial]$ -module and  $l, r : Q \rightarrow \text{Cend}(M)$  are two  $\mathbf{k}[\partial]$ -module homomorphisms satisfying

$$(9) \quad l(a)_\lambda(l(b)_\mu v) = l([a_\lambda b])_{\lambda+\mu} v + l(b)_\mu(l(a)_\lambda v),$$

$$(10) \quad l(a)_\lambda(r(b)_{-\mu-\partial} v) = r(b)_{-\lambda-\mu-\partial}(l(a)_\lambda v) + r([a_\lambda b])_{-\mu-\partial} v,$$

$$(11) \quad r(b)_{-\lambda-\mu-\partial}(r(a)_{-\mu-\partial} v) = -r(b)_{-\lambda-\mu-\partial}(l(a)_\lambda v) \quad \text{for all } a, b \in Q \text{ and } v \in M.$$

A representation  $(M, l, r)$  of  $(Q, [\cdot, \cdot])$  is called **finite**, if  $M$  is a finitely generated  $\mathbf{k}[\partial]$ -module.

**Example 2.12.** Let  $(Q, [\cdot, \cdot])$  be a Leibniz conformal algebra. Define two  $\mathbf{k}[\partial]$ -module homomorphisms  $L$  and  $R$  from  $Q$  to  $\text{Cend}(Q)$  by  $L(a)_\lambda b = [a_\lambda b]$  and  $R(a)_\lambda b = [b_{-\lambda-\partial} a]$  for all  $a, b \in Q$ . Then  $(Q, L, R)$  is a representation of  $(Q, [\cdot, \cdot])$ , which is called the **regular representation** of  $(Q, [\cdot, \cdot])$ . If there is a Leibniz conformal algebra structure on the conformal dual  $Q^{*c}$  of  $Q$ , then we denote the regular representation of  $(Q^{*c}, [\cdot, \cdot])$  by  $(Q^{*c}, \mathcal{L}, \mathcal{R})$ .

**Proposition 2.13.** Let  $Q$  be a Leibniz conformal algebra. Then  $(M, l, r)$  is a representation of  $Q$  if and only if there is a Leibniz conformal algebra structure on the direct sum  $Q \oplus M$  as  $\mathbf{k}[\partial]$ -modules given by

$$(12) \quad [(a+x)_\lambda(b+y)] = [a_\lambda b] + l(a)_\lambda y + r(b)_{-\lambda-\partial} x \quad \text{for all } a, b \in Q \text{ and } x, y \in M.$$

Denote this Leibniz conformal algebra by  $Q \ltimes_{l,r} M$ , which is called a **semi-direct product** of  $Q$  and  $(M, l, r)$ , or a **semi-direct product Leibniz conformal algebra**.

*Proof.* It is straightforward.  $\square$

**Proposition 2.14.** Let  $(M, l, r)$  be a finite representation of a Leibniz conformal algebra  $(Q, [\cdot, \cdot])$ , which is free as a  $\mathbf{k}[\partial]$ -module. Let  $l^*$  and  $r^*$  be two  $\mathbf{k}[\partial]$ -module homomorphisms from  $Q$  to  $\text{Cend}(M^{*c})$  given by

$$(13) \quad \langle l^*(a)_\lambda f, v \rangle_\mu = -\langle f, l(a)_\lambda v \rangle_{\mu-\lambda}, \quad \langle r^*(a)_\lambda f, v \rangle_\mu = -\langle f, r(a)_\lambda v \rangle_{\mu-\lambda},$$

for all  $a \in A$ ,  $f \in M^{*c}$  and  $v \in M$ . Then  $(M^{*c}, l^*, -l^* - r^*)$  is a representation of  $(Q, [\cdot, \cdot])$ , which is called the **dual representation** of  $(M, l, r)$ .

*Proof.* Let  $a, b \in Q$ ,  $f \in M^{*c}$  and  $v \in M$ . We have

$$\begin{aligned} & \langle l^*(a)_\lambda(-l^* + r^*)(b)_{-\mu-\partial} f \rangle + (l^* + r^*)(b)_{-\lambda-\mu-\partial} \langle l^*(a)_\lambda f \rangle + (l^* + r^*)([a_\lambda b])_{-\mu-\partial} \langle f, v \rangle_\theta \\ &= \langle (l^* + r^*)(b)_{-\mu-\partial} f, l(a)_\lambda v \rangle_{\theta-\lambda} - \langle l^*(a)_\lambda f, (l+r)(b)_{-\lambda-\mu+\theta} v \rangle_{\lambda+\mu} - \langle f, (l+r)([a_\lambda b])_{-\mu+\theta} v \rangle_\mu \\ &= \langle f, -(l+r)(b)_{-\lambda-\mu+\theta} (l(a)_\lambda v) + l(a)_\lambda ((l+r)(b)_{-\lambda-\mu+\theta} v) - (l+r)([a_\lambda b])_{-\mu+\theta} v \rangle_\mu \\ &= -\langle f, l(b)_{-\lambda-\mu+\theta} (l(a)_\lambda v) - l(a)_\lambda (l(b)_{-\lambda-\mu+\theta} v) + l([a_\lambda b])_{-\mu+\theta} v \rangle_\mu \\ & \quad - \langle f, r(b)_{\theta-\lambda-\partial} (l(a)_\lambda v) - l(a)_\lambda (r(b)_{\theta-\partial} v) + r([a_\lambda b])_{\theta-\partial} v \rangle_\mu = 0, \end{aligned}$$

which means that (10) holds. Similarly, we can verify that (9) and (11) hold. Therefore,  $(M^{*c}, l^*, -l^* - r^*)$  is a representation of  $(Q, [\cdot, \cdot])$ .  $\square$

**Example 2.15.** Let  $(Q, [\cdot, \cdot])$  be a finite Leibniz conformal algebra which is free as a  $\mathbf{k}[\partial]$ -module. By Proposition 2.14, the dual representation of the regular representation  $(Q, L, R)$  is  $(Q^{*c}, L^*, -L^* - R^*)$ .

**Definition 2.16.** Let  $Q_1$  and  $Q_2$  be two Leibniz conformal algebras. If  $(Q_2, l, r)$  is a representation of  $Q_1$ ,  $(Q_1, \phi, \psi)$  is a representation of  $Q_2$  and the following equations

$$(14) \quad [x_\lambda(r(a)_{-\mu-\partial} y)] + r(\phi(y)_\mu a)_{-\lambda-\partial} x = r(a)_{-\lambda-\mu-\partial} ([x_\lambda y]) + [y_\mu(r(a)_{-\lambda-\partial} x)] + r(\phi(x)_\lambda a)_{-\mu-\partial} y,$$

$$(15) \quad l(a)_\lambda ([x_\mu y]) = [(l(a)_\lambda x)_{\lambda+\mu} y] + l(\psi(x)_{-\lambda-\partial} a)_{\lambda+\mu} y + [x_\mu(l(a)_\lambda y)] + r(\psi(y)_{-\lambda-\partial} a)_{-\mu-\partial} x,$$

$$(16) \quad [x_\lambda(l(a)_\mu y)] + r(\psi(y)_{-\mu-\partial} a)_{-\lambda-\partial} x = l(\phi(x)_\lambda a)_{\lambda+\mu} y + l(a)_\mu ([x_\lambda y]) + [(r(a)_{-\lambda-\partial} x)_{\lambda+\mu} y],$$

$$(17) \quad [a_\lambda(\psi(x)_{-\mu-\partial} b)] + \psi(l(b)_\mu x)_{-\lambda-\partial} a = \psi(x)_{-\lambda-\mu-\partial} [a_\lambda b] + [b_\mu(\psi(x)_{-\lambda-\partial} a)] + \psi(l(a)_\lambda x)_{-\mu-\partial} b,$$

$$(18) \quad [a_\lambda(\phi(x)_\mu b)] + \psi(r(b)_{-\mu-\partial} x)_{-\lambda-\partial} a = [(\psi(x)_{-\lambda-\partial} a)_{\lambda+\mu} b] + \phi(l(a)_\lambda x)_{\lambda+\mu} b + \phi(x)_\mu ([a_\lambda b]),$$

$$(19) \quad [(\phi(x)_\lambda a)_{\lambda+\mu} b] + \phi(r(a)_{-\lambda-\partial} x)_{\lambda+\mu} b + [(\psi(x)_{-\mu-\partial} a)_{\lambda+\mu} b] + \phi(l(a)_\mu x)_{\lambda+\mu} b = 0,$$

hold for all  $a, b \in Q_1$  and  $x, y \in Q_2$ , then  $(Q_1, Q_2; (l, r), (\phi, \psi))$  is called a **matched pair** of Leibniz conformal algebras  $Q_1$  and  $Q_2$ .

**Proposition 2.17.** [21] *Let  $(Q_1, Q_2; (l, r), (\phi, \psi))$  be a matched pair of Leibniz conformal algebras. Then there is a Leibniz conformal algebra structure on the direct sum  $Q_1 \oplus Q_2$  of  $\mathbf{k}[\partial]$ -modules defined by*

$$(20) \quad [(a+x)_\lambda(b+y)] = [a_\lambda b] + \phi(x)_\lambda b + \psi(y)_{-\lambda-\partial} a + [x_\lambda y] + l(a)_\lambda y + r(b)_{-\lambda-\partial} x,$$

for all  $a, b \in Q_1$ ,  $x, y \in Q_2$ . Denote this Leibniz conformal algebra by  $Q_1 \bowtie Q_2$ , which is called a **bicrossed product** of  $Q_1$  and  $Q_2$  associated with the matched pair  $(Q_1, Q_2; (l, r), (\phi, \psi))$  of Leibniz conformal algebras. Moreover, any Leibniz conformal algebra  $E = Q_1 \oplus Q_2$  where the sum is the direct sum of  $\mathbf{k}[\partial]$ -modules such that  $Q_1$  and  $Q_2$  are two Leibniz conformal subalgebras of  $E$ , is isomorphic to  $Q_1 \bowtie Q_2$  associated with some matched pair of Leibniz conformal algebras.

### 2.3. Conformal Manin triples of Leibniz conformal algebras.

**Definition 2.18.** Let  $M$  be a  $\mathbf{k}[\partial]$ -module. A **conformal bilinear form** on  $M$  is a  $\mathbf{k}$ -bilinear map  $B_\lambda(\cdot, \cdot) : M \times M \rightarrow \mathbf{k}[\lambda]$  satisfying

$$B_\lambda(\partial u, v) = -\lambda B_\lambda(u, v), \quad B_\lambda(u, \partial v) = \lambda B_\lambda(u, v) \quad \text{for all } u, v \in M.$$

A conformal bilinear form  $B_\lambda(\cdot, \cdot)$  on  $M$  is called **skew-symmetric** if

$$B_\lambda(u, v) = -B_{-\lambda}(v, u) \quad \text{for all } u, v \in M.$$

**Remark 2.19.** If there is a conformal bilinear form on a  $\mathbf{k}[\partial]$ -module  $M$ , then we have a  $\mathbf{k}[\partial]$ -module homomorphism  $\phi : M \rightarrow M^{*c}$  sending  $v$  to  $\phi_v$  where  $\phi_v$  is defined by

$$(\phi_v)_\lambda w = B_\lambda(v, w) \quad \text{for all } v, w \in M.$$

Moreover, if  $\phi$  is an isomorphism of  $\mathbf{k}[\partial]$ -modules, then this conformal bilinear form is called **non-degenerate**.

**Definition 2.20.** A **conformal Manin triple** of Leibniz conformal algebras is a triple of finite Leibniz conformal algebras  $(Q, Q_0, Q_1)$ , where

- (a) there is a non-degenerate skew-symmetric conformal bilinear form  $B_\lambda(\cdot, \cdot)$  on  $Q$  such that the following invariant condition holds:

$$(21) \quad B_\mu(a, [b_\lambda c]) = B_{-\lambda}([a_\mu c] + [c_{-\mu-\partial} a], b) \quad \text{for all } a, b, c \in Q,$$

- (b)  $Q_0$  and  $Q_1$  are Leibniz conformal subalgebras of  $Q$ , where  $Q = Q_0 \oplus Q_1$  as a direct sum of  $\mathbf{k}[\partial]$ -modules,  
(c)  $Q_0$  and  $Q_1$  are isotropic with respect to  $B_\lambda(\cdot, \cdot)$ , i.e.  $B_\lambda(Q_i, Q_i) = 0$  for  $i = 0, 1$ .

**Example 2.21.** Let  $Q$  be a finite Leibniz conformal algebra which is free as a  $\mathbf{k}[\partial]$ -module. Then  $(Q \bowtie_{L^*, -L^* - R^*} Q^{*c}, Q, Q^{*c})$  is a conformal Manin triple of Leibniz conformal algebras, where the non-degenerate skew-symmetric conformal bilinear form  $B_\lambda(\cdot, \cdot)$  on  $Q \bowtie_{L^*, -L^* - R^*} Q^{*c}$  is given by

$$(22) \quad B_\lambda(a+f, b+g) = \langle f, b \rangle_\lambda - \langle g, a \rangle_{-\lambda} \quad \text{for all } a, b \in Q \text{ and } f, g \in Q^{*c}.$$

In the sequel, for convenience, we set  $\langle a, f \rangle_{-\lambda} := -\langle f, a \rangle_\lambda$  for all  $a \in Q$  and  $f \in Q^{*c}$ .



**Theorem 2.22.** *Let  $Q$  and  $Q^{*c}$  be two finite Leibniz conformal algebras which are free as  $\mathbf{k}[\partial]$ -modules. Then  $(Q \bowtie Q^{*c}, Q, Q^{*c})$  is a conformal Manin triple of Leibniz conformal algebras where the invariant skew-symmetric conformal bilinear form on  $Q \oplus Q^{*c}$  is given by (22) if and only if  $(Q, Q^{*c}; (L^*, -L^* - R^*), (\mathcal{L}^*, -\mathcal{L}^* - \mathcal{R}^*))$  is a matched pair of Leibniz conformal algebras.*

*Proof.* Let  $(Q, Q^{*c}; (L^*, -L^* - R^*), (\mathcal{L}^*, -\mathcal{L}^* - \mathcal{R}^*))$  be a matched pair of Leibniz conformal algebras. Then there is a Leibniz conformal algebra structure on  $Q \oplus Q^{*c}$  with the  $\lambda$ -bracket given by (20). Obviously,  $B_\lambda(\cdot, \cdot)$  is non-degenerate and skew-symmetric. For all  $a, b, c \in Q$  and  $f, g, h \in Q^{*c}$ , we have

$$\begin{aligned} & B_\mu(a + f, [(b + g)_\lambda(c + h)]) \\ &= B_\mu(a + f, [b_\lambda c] + \mathcal{L}^*(g)_\lambda c + (-\mathcal{L}^* - \mathcal{R}^*)(h)_{-\lambda-\partial} b + [g_\lambda z] + L^*(b)_\lambda h + (-L^* - R^*)(c)_{-\lambda-\partial} g) \\ &= \langle f, [b_\lambda c] \rangle_\mu + \langle f, \mathcal{L}^*(g)_\lambda c \rangle_\mu - \langle f, \mathcal{L}^*(h)_{-\lambda-\partial} b + \mathcal{R}^*(h)_{-\lambda-\partial} b \rangle_\mu - \langle [g_\lambda h], a \rangle_{-\mu} \\ &\quad - \langle L^*(b)_\lambda h, a \rangle_{-\mu} + \langle L^*(c)_{-\lambda-\partial} g, a \rangle_{-\mu} + \langle R^*(c)_{-\lambda-\partial} g, a \rangle_{-\mu} \\ &= \langle f, [b_\lambda c] \rangle_\mu + \langle c, [g_\lambda f] \rangle_{-\mu-\lambda} - \langle b, [h_{-\mu-\lambda} f] \rangle_\lambda - \langle b, [f_\mu h] \rangle_\lambda + \langle a, [g_\lambda h] \rangle_\mu \\ &\quad + \langle h, [b_\lambda a] \rangle_{-\lambda-\mu} - \langle g, [c_{-\lambda-\mu} a] \rangle_\lambda - \langle g, [a_\mu c] \rangle_\lambda. \end{aligned}$$

On the other hand, we have

$$\begin{aligned} & B_{-\lambda}([(a + f)_\mu(c + h)] + [(c + h)_{-\mu-\partial}(a + f)], b + g) \\ &= B_{-\lambda}([a_\mu c] + \mathcal{L}^*(f)_\mu c - (\mathcal{L}^* + \mathcal{R}^*)(h)_{-\mu-\partial} a + [f_\mu h] + L^*(a)_\mu h - (L^* + R^*)(c)_{-\mu-\partial} f, b + g) \\ &\quad + B_{-\lambda}([c_{-\mu-\partial} a] + \mathcal{L}^*(h)_{-\mu-\partial} a - (\mathcal{L}^* + \mathcal{R}^*)(f)_\mu c + [h_{-\mu-\partial} f] + L^*(c)_{-\mu-\partial} f - (L^* + R^*)(a)_\mu h, b + g) \\ &= \langle [a_\mu c] + [c_{-\mu-\partial} a], g \rangle_{-\lambda} - \langle c, [f_\mu g] \rangle_{-\lambda-\mu} + \langle a, [h_{-\lambda-\mu} g] \rangle_\mu + \langle a, [g_\lambda h] \rangle_\mu \\ &\quad + \langle c, [g_\lambda f] \rangle_{-\lambda-\mu} + \langle [f_\mu h] + [h_{-\mu-\partial} f], b \rangle_{-\lambda} + \langle f, [b_\lambda c] \rangle_\mu + \langle h, [b_\lambda a] \rangle_{-\lambda-\mu} \\ &= \langle f, [b_\lambda c] \rangle_\mu + \langle c, [g_\lambda f] \rangle_{-\lambda-\mu} - \langle b, [h_{-\lambda-\mu} f] \rangle_\lambda - \langle b, [f_\mu h] \rangle_\lambda + \langle a, [g_\lambda h] \rangle_\mu \\ &\quad + \langle h, [b_\lambda a] \rangle_{-\lambda-\mu} - \langle g, [c_{-\lambda-\mu} a] \rangle_\lambda - \langle g, [a_\mu c] \rangle_\lambda. \end{aligned}$$

Thus  $B_\lambda(\cdot, \cdot)$  satisfies (21). Therefore,  $(Q \bowtie Q^{*c}, Q, Q^{*c})$  is a conformal Manin triple of Leibniz conformal algebras where the invariant skew-symmetric conformal bilinear form  $B_\lambda(\cdot, \cdot)$  on  $Q \oplus Q^{*c}$  is given by (22).

Conversely, let  $(Q \bowtie Q^{*c}, Q, Q^{*c})$  be a conformal Manin triple of Leibniz conformal algebras with the non-degenerate skew-symmetric invariant bilinear form  $B_\lambda(\cdot, \cdot)$  given by (22). Assume that  $Q \bowtie Q^{*c}$  is a bicrossed product associated with a matched pair  $(Q_1, Q_2; (l, r), (\phi, \psi))$  of Leibniz conformal algebras. Therefore, for all  $a \in Q$ ,  $f \in Q^{*c}$ , assume that

$$[a_\lambda f] = l(a)_\lambda f + \psi(f)_{-\lambda-\partial} a, \quad [f_\lambda a] = \phi(f)_\lambda a + r(a)_{-\lambda-\partial} f.$$

Let  $a, b \in Q$  and  $f, g \in Q^{*c}$ . By (21), we get

$$\begin{aligned} \langle g, \psi(f)_{-\lambda-\partial} a \rangle_\mu &= B_\mu(g, [a_\lambda f]) = B_{-\lambda}([g_\mu f] + [f_{-\mu-\partial} g], a) = -\langle a, [g_\mu f] + [f_{-\mu-\lambda} g] \rangle_\lambda \\ &= \langle \mathcal{L}^*(f)_{-\lambda-\partial} a + \mathcal{R}^*(f)_{-\lambda-\partial} a, g \rangle_{-\mu} = -\langle g, (\mathcal{L}^* + \mathcal{R}^*)(f)_{-\lambda-\partial} a \rangle_\mu, \end{aligned}$$

which means that  $\psi = -\mathcal{L}^* - \mathcal{R}^*$ . Moreover, we have

$$\langle l(a)_\lambda f, b \rangle_\mu = B_\mu([a_\lambda f], b) = -B_{-\lambda}([b_{-\mu} f] + [f_{\mu-\partial} b], a) = -\langle f, [a_\lambda b] \rangle_{\mu-\lambda} = \langle L^*(a)_\lambda f, b \rangle_\mu,$$

which means that  $l = L^*$ . Similarly, we can get  $\phi = \mathcal{L}^*$  and  $r = -L^* - R^*$ . Hence,  $(Q, Q^{*c}; (L^*, -L^* - R^*), (\mathcal{L}^*, -\mathcal{L}^* - \mathcal{R}^*))$  is a matched pair of Leibniz conformal algebras.  $\square$

## 3. LEIBNIZ CONFORMAL BIALGEBRAS

We introduce the notion of Leibniz conformal bialgebras and then establish the equivalences among some matched pairs of Leibniz conformal algebras, conformal Manin triples of Leibniz conformal algebras, and Leibniz conformal bialgebras.

First, we introduce the notion of Leibniz conformal coalgebras.

**Definition 3.1.** A **Leibniz conformal coalgebra** is a  $\mathbf{k}[\partial]$ -module  $Q$  endowed with a  $\mathbf{k}[\partial]$ -module homomorphism  $\Delta : Q \rightarrow Q \otimes Q$  such that

$$(23) \quad (\text{id} \otimes \Delta)\Delta(a) = (\Delta \otimes \text{id})\Delta(a) + \tau_{12}(\text{id} \otimes \Delta)\Delta(a) \quad \text{for all } a \in Q,$$

where the module action of  $\mathbf{k}[\partial]$  on  $Q \otimes Q$  is defined as  $\partial(a \otimes b) = (\partial a) \otimes b + a \otimes (\partial b)$  for all  $a, b \in Q$ .

Let  $Q$  be a  $\mathbf{k}[\partial]$ -module. For all  $f, g \in Q^{*c}$  and  $a, b \in Q$ , we define

$$\langle f \otimes g, a \otimes b \rangle_{\lambda, \mu} = \langle f, a \rangle_{\lambda} \langle g, b \rangle_{\mu}.$$

**Proposition 3.2.** (a) *Let  $(Q, \Delta)$  be a finite Leibniz conformal coalgebra. Then  $Q^{*c}$  is a Leibniz conformal algebra with the following  $\lambda$ -bracket:*

$$(24) \quad \langle [f \lambda g], a \rangle_{\mu} = \langle f \otimes g, \Delta(a) \rangle_{\lambda, \mu - \lambda} \quad \text{for all } f, g \in Q^{*c} \text{ and } a \in Q.$$

(b) *Let  $Q$  be a finite Leibniz conformal algebra which is free as a  $\mathbf{k}[\partial]$ -module, that is,  $Q = \sum_{i=1}^n \mathbf{k}[\partial]a^i$ , where  $\{a^1, \dots, a^n\}$  is a  $\mathbf{k}[\partial]$ -basis of  $Q$ . Then  $Q^{*c} = \text{Chom}(A, \mathbf{k}) = \sum_{i=1}^n \mathbf{k}[\partial]a_i$ , where  $\{a_1, \dots, a_n\}$  is the dual  $\mathbf{k}[\partial]$ -basis in the sense that  $(a_i)_{\lambda} a^j = \delta_{ij}$ , is a Leibniz conformal coalgebra with the following co-bracket:*

$$(25) \quad \Delta(f) = \sum_{i,j} f_{\mu}([a_{\lambda}^i a^j])(a_i \otimes a_j)|_{\lambda = \partial \otimes \text{id}, \mu = -\partial \otimes \text{id} - \text{id} \otimes \partial}.$$

*More precisely, if  $[a_{\lambda}^i a^j] = \sum_k P_k^{ij}(\lambda, \partial) a^k$ , where  $P_k^{ij}(\lambda, \partial) \in \mathbf{k}[\lambda, \partial]$ , then the co-bracket is*

$$(26) \quad \Delta(a_k) = \sum_{i,j} Q_k^{ij}(\partial \otimes \text{id}, \text{id} \otimes \partial) a_i \otimes a_j,$$

*where  $Q_k^{ij}(x, y) = P_k^{ij}(x, -x - y)$ .*

*Proof.* It follows directly by the proof of [26, Proposition 2.13]. □

**Corollary 3.3.** *Let  $(Q, \Delta)$  be a Leibniz conformal coalgebra where  $Q$  is free of rank 1 as a  $\mathbf{k}[\partial]$ -module. Then  $(Q, \Delta) \simeq (\mathbf{k}[\partial]a, \Delta')$  with  $\Delta'(a) = 0$  or  $\Delta'(a) = c(\partial a \otimes a - a \otimes \partial a)$ ,  $c \in \mathbf{k} \setminus \{0\}$ .*

*Proof.* It follows directly by Propositions 2.4 and 3.2. □

In the sequel, we always set  $\partial^{\otimes 2} := \partial \otimes \text{id} + \text{id} \otimes \partial$ .

Next, we introduce the definition of Leibniz conformal bialgebras.

**Definition 3.4.** Let  $(Q, [\cdot, \cdot])$  be a Leibniz conformal algebra and  $(Q, \Delta)$  be a Leibniz conformal coalgebra. Then  $(Q, [\cdot, \cdot], \Delta)$  is called a **Leibniz conformal bialgebra** if the following conditions hold:

$$(27) \quad (R(a)_{\lambda-\partial \otimes \text{id}} \otimes \text{id})\Delta(b) = \tau((R(b)_{-\lambda-\partial \otimes \text{id}} \otimes \text{id})\Delta(a)),$$

$$(28) \quad \Delta([a, b]) = (\text{id} \otimes R(b)_{-\lambda-\partial \otimes^2} - L(b)_{-\lambda-\partial \otimes^2} \otimes \text{id} - R(b)_{-\lambda-\partial \otimes^2} \otimes \text{id})(\tau + \text{id})\Delta(a) \\ + (\text{id} \otimes L(a)_\lambda + L(a)_\lambda \otimes \text{id})\Delta(b) \quad \text{for all } a, b \in Q.$$

We give a remark to illustrate the relationship between Leibniz bialgebras and Leibniz conformal algebras.

**Remark 3.5.** Recall [30] that a Leibniz bialgebra is a triple  $(A, [\cdot, \cdot], \delta)$ , where  $(A, [\cdot, \cdot])$  is a Leibniz algebra,  $(A, \delta)$  is a Leibniz coalgebra in the sense that

$$(\text{id} \otimes \delta)\delta(a) = (\delta \otimes \text{id})\delta(a) + \tau_{12}(\text{id} \otimes \delta)\delta(a) \quad \text{for all } a \in A,$$

and they satisfy the following equalities

$$(R_A(a) \otimes \text{id})\delta(b) = \tau((R_A(b) \otimes \text{id})\delta(a)),$$

$$\delta([a, b]) = (\text{id} \otimes R_A(b) - L_A(b) \otimes \text{id} - R_A(b) \otimes \text{id})(\tau + \text{id})\delta(a) + (\text{id} \otimes L_A(a) + L_A(a)_\lambda \otimes \text{id})\delta(b),$$

for all  $a, b \in A$ , where  $L_A(a)(b) = [a, b] = R_A(b)(a)$ .

Let  $(A, [\cdot, \cdot], \delta)$  be a Leibniz bialgebra. Then  $(\text{Cur } A = \mathbf{k}[\partial] \otimes A, [\cdot, \cdot], \Delta)$  is a Leibniz conformal bialgebra, where  $(\text{Cur } A, [\cdot, \cdot])$  is the current Leibniz conformal algebra associated with  $(A, [\cdot, \cdot])$  and  $\Delta$  is a  $\mathbf{k}[\partial]$ -module homomorphism defined by

$$\Delta(a) = \delta(a) \quad \text{for all } a \in A.$$

$(\text{Cur } A = \mathbf{k}[\partial] \otimes A, [\cdot, \cdot], \Delta)$  is called the **current Leibniz conformal bialgebras** associated with  $(A, [\cdot, \cdot], \delta)$ .

Conversely, let  $(Q, [\cdot, \cdot], \Delta)$  be a Leibniz conformal bialgebra and  $\pi : Q \rightarrow Q/\partial Q$  be the natural projection. Set  $[a, b] = \sum_{i=0}^{n_{a,b}} \frac{\lambda^i}{i!} a_{(i)} b$  for all  $a, b \in Q$ . Then it is easy to see that  $(Q/\partial Q, [\cdot, \cdot], \delta)$  is a Leibniz bialgebra, where  $[\cdot, \cdot]$  and  $\delta$  are defined by

$$[a + \partial Q, b + \partial Q] := \pi(a_{(0)} b), \quad \delta(a + \partial Q) := (\pi \otimes \pi)\Delta(a) \quad \text{for all } a, b \in Q.$$

Hence in the above sense, Leibniz conformal bialgebras are regarded as ‘‘conformal analogues’’ of Leibniz bialgebras.

**Proposition 3.6.** *Let  $(Q, [\cdot, \cdot])$  be a finite Leibniz conformal algebra which is free as a  $\mathbf{k}[\partial]$ -module. Suppose that there is a Leibniz conformal algebra structure on  $Q^{*c}$  which is obtained from a  $\mathbf{k}[\partial]$ -module homomorphism  $\Delta : Q \rightarrow Q \otimes Q$ . Then  $(Q, Q^{*c}; (L^*, -L^* - R^*), (\mathcal{L}^*, -\mathcal{L}^* - \mathcal{R}^*))$  is a matched pair of Leibniz conformal algebras if and only if  $(Q, [\cdot, \cdot], \Delta)$  is a Leibniz conformal bialgebra.*

*Proof.* Let  $\{a^1, \dots, a^n\}$  be a  $\mathbf{k}[\partial]$ -basis of  $Q$ ,  $\{a_1, \dots, a_n\}$  be the dual  $\mathbf{k}[\partial]$ -basis of  $Q^{*c}$  and  $I = \{1, \dots, n\}$ . Set

$$[a^i, \lambda a^j] = \sum_k P_k^{ij}(\lambda, \partial) a^k, \quad [a_{i\lambda} a_j] = \sum_k H_k^{ij}(\lambda, \partial) a_k,$$

where  $P_k^{ij}(\lambda, \partial), H_k^{ij}(\lambda, \partial) \in \mathbf{k}[\lambda, \partial]$ . By Proposition 3.2, we get  $\Delta(a^k) = \sum_{i,j} Q_k^{ij}(\partial \otimes \text{id}, \text{id} \otimes \partial) a^i \otimes a^j$ , where  $Q_k^{ij}(x, y) = H_k^{ij}(x, -x - y)$ . Since

$$\langle \mathcal{L}^*(a_i)_\lambda a^j, a_k \rangle_\mu = -\langle a^j, [a_{i\lambda} a_k] \rangle_{\mu-\lambda} = -H_j^{ik}(\lambda, \mu - \lambda),$$

we get  $\mathcal{L}^*(a_i)_\lambda a^j = -\sum_k H_j^{ik}(\lambda, -\lambda - \partial) a^k$ . Similarly, we can get

$$\begin{aligned} \mathcal{R}^*(a_i)_\lambda a^j &= -\sum_k H_j^{ki}(\partial, -\lambda - \partial) a^k, & L^*(a^i)_\lambda a_j &= -\sum_k P_j^{ik}(\lambda, -\lambda - \partial) a_k, \\ R^*(a^i)_\lambda a_j &= -\sum_k P_j^{ki}(\partial, -\lambda - \partial) a_k. \end{aligned}$$

Considering the coefficient of  $a^j \otimes a^h$  in  $(R(a^i)_{\lambda-\partial \otimes \text{id}} \otimes \text{id}) \Delta(a^k) = \tau((R(a^k)_{-\lambda-\partial \otimes \text{id}} \otimes \text{id}) \Delta(a^i))$ , we get

$$(29) \quad \sum_t H_k^{th}(\lambda, -\lambda - \text{id} \otimes \partial) P_j^{ti}(-\lambda, \partial \otimes \text{id}) = \sum_t H_i^{tj}(-\lambda, \lambda - \partial \otimes \text{id}) P_h^{tk}(\lambda, \text{id} \otimes \partial).$$

On the other hand, (19) holds for all  $a, b \in Q$  and  $x \in Q^{*c}$  if and only if

$$-\langle \mathcal{L}^*(R^*(a^i)_{-\lambda-\partial} a_j)_{\lambda+\mu} a^k, a_h \rangle_\theta = \langle [(R^*(a_j)_{-\mu-\partial} a^i)_{\lambda+\mu} a^k], a_h \rangle_\theta \quad \text{for all } i, j, k, h \in I,$$

holds if and only if

$$(30) \quad \sum_t H_k^{th}(\lambda+\mu, -\lambda-\mu+\theta) P_j^{ti}(-\lambda-\mu, \lambda) = \sum_t H_i^{tj}(-\lambda-\mu, \mu) P_h^{tk}(\lambda+\mu, -\theta) \quad \text{for all } i, j, k, h \in I,$$

holds. Obviously, (29) is equivalent to (30) by replacing  $\lambda$  by  $\lambda + \mu$ ,  $\text{id} \otimes \partial$  by  $-\theta$  and  $\partial \otimes \text{id}$  by  $\lambda$  in (29). Then (27) holds if and only if (19) holds.

Note that (18) holds for all  $a, b \in Q$  and  $x \in Q^{*c}$  if and only if for all  $i, j, k, h \in I$ ,

$$\begin{aligned} &\langle [a^i_\lambda (\mathcal{L}^*(a_j)_\mu a^k)] + (\mathcal{L}^* + \mathcal{R}^*)((L^* + R^*)(a^k)_{-\mu-\partial} a_j)_{-\lambda-\partial} a^i, a_h \rangle_\theta \\ &= \langle \mathcal{L}^*(a_j)_\mu [a^i_\lambda a^k] - [(\mathcal{L}^*(a_j)_\mu a^i)_{\lambda+\mu} a^k] + \mathcal{L}^*((L^* + R^*)(a^i)_{-\mu-\partial} a_j)_{\lambda+\mu} a^k, a_h \rangle_\theta \end{aligned}$$

holds, which is equivalent to

$$(31) \quad \begin{aligned} &\sum_t (P_j^{kt}(-\lambda - \mu + \theta, \mu) + P_j^{tk}(\lambda - \theta, \mu))(H_i^{th}(-\lambda + \theta, \lambda) + H_i^{ht}(-\theta, \lambda)) \\ &- \sum_t H_k^{jt}(\mu, -\mu - \lambda + \theta) P_h^{it}(\lambda, -\theta) = -\sum_t P_t^{ik}(\lambda, \mu - \theta) H_t^{jh}(\mu, -\mu + \theta) \\ &+ \sum_t (H_i^{jt}(\mu, \lambda) P_h^{tk}(\lambda + \mu, -\theta) + P_j^{it}(\lambda, \mu) + P_j^{ti}(-\lambda - \mu, \mu)) H_k^{th}(\lambda + \mu, -\lambda - \mu + \theta). \end{aligned}$$

On the other hand, (28) holds if and only if for all  $i, k \in I$ ,

$$(32) \quad \begin{aligned} \Delta([a^i_\lambda a^k]) &= (\text{id} \otimes R(a^k)_{-\lambda-\partial^{\otimes 2}} - L(a^k)_{-\lambda-\partial^{\otimes 2}} \otimes \text{id} - R(a^k)_{-\lambda-\partial^{\otimes 2}} \otimes \text{id})(\tau + \text{id}) \Delta(a^i) \\ &+ (\text{id} \otimes L(a^i)_\lambda + L(a^i)_\lambda \otimes \text{id}) \Delta(a^k) \end{aligned}$$

holds. Considering the coefficient of  $a^j \otimes a^h$  in (32), we get

$$\begin{aligned}
(33) \quad & \sum_t P_t^{ik}(\lambda, \partial^{\otimes 2}) H_t^{jh}(\partial \otimes \text{id}, -\partial^{\otimes 2}) = \sum_t (H_i^{jt}(\partial \otimes \text{id}, \lambda) P_h^{tk}(\lambda + \partial \otimes \text{id}, \text{id} \otimes \partial) \\
& + H_i^{tj}(-\lambda - \partial \otimes \text{id}, \lambda) P_h^{tk}(\lambda + \partial \otimes \text{id}, \text{id} \otimes \partial) - H_i^{th}(-\lambda - \text{id} \otimes \partial, \lambda) P_j^{kt}(-\lambda - \partial^{\otimes 2}, \partial \otimes \text{id}) \\
& - H_i^{ht}(\text{id} \otimes \partial, \lambda) P_j^{kt}(-\lambda - \partial^{\otimes 2}, \partial \otimes \text{id}) - H_i^{th}(-\lambda - \text{id} \otimes \partial, \lambda) P_j^{tk}(\lambda + \text{id} \otimes \partial, \partial \otimes \text{id}) \\
& - H_i^{ht}(\text{id} \otimes \partial, \lambda) P_j^{tk}(\lambda + \text{id} \otimes \partial, \partial \otimes \text{id}) + H_k^{jt}(\partial \otimes \text{id}, -\partial \otimes \text{id} - \lambda - \text{id} \otimes \partial) P_h^{ti}(\lambda, \text{id} \otimes \partial) \\
& + H_k^{th}(\lambda + \partial \otimes \text{id}, -\lambda - \partial^{\otimes 2}) P_j^{ti}(\lambda, \partial \otimes \text{id})).
\end{aligned}$$

By (27), we have

$$\sum_t H_k^{th}(\lambda + \partial \otimes \text{id}, -\lambda - \partial^{\otimes 2}) P_j^{ti}(-\lambda - \partial \otimes \text{id}, \partial \otimes \text{id}) = \sum_t H_i^{tj}(-\lambda - \partial \otimes \text{id}, \lambda) P_h^{tk}(\lambda + \partial \otimes \text{id}, \text{id} \otimes \partial).$$

Then it is clear that (31) is equivalent to (33) by replacing  $\mu$  by  $\partial \otimes \text{id}$  and  $\theta$  by  $-\text{id} \otimes \partial$ . Moreover, (17) can be deduced by (18) and (19).

For all  $a, b \in Q$  and  $x, y \in Q^{*c}$ , by some calculations we have

$$\begin{aligned}
& \langle L^*(a)_\lambda([x_\mu y]) - [(L^*(a)_\lambda x)_{\lambda+\mu} y] + L^*((\mathcal{L}^* + \mathcal{R}^*)(x)_{-\lambda-\partial a})_{\lambda+\mu} y \\
& - [x_\mu (L^*(a)_\lambda y)] - (L^* + R^*)((\mathcal{L}^* + \mathcal{R}^*)(y)_{-\lambda-\partial a})_{-\mu-\partial x}, b \rangle_\theta \\
& = \langle -\mathcal{L}^*(x)_\mu([a_\lambda b]) - \mathcal{L}^*(L^*(a)_\lambda x)_{\lambda+\mu} b + [((\mathcal{L}^* + \mathcal{R}^*)(x)_{-\lambda-\partial a})_{\lambda+\mu} b] \\
& + [a_\lambda (\mathcal{L}^*(x)_\mu b)] + (\mathcal{L}^* + \mathcal{R}^*)((L^* + R^*)(b)_{-\mu-\partial x})_{-\lambda-\partial a}, y \rangle_{\lambda+\mu-\theta},
\end{aligned}$$

which means that (15) is equivalent to (18). Similarly, we can obtain that (16) is equivalent to (19) and (14) is equivalent to (17). Thus,  $(Q, Q^{*c}; (L^*, -L^* - R^*), (\mathcal{L}^*, -\mathcal{L}^* - \mathcal{R}^*))$  is a matched pair of Leibniz conformal algebras if and only if (27) and (28) hold.  $\square$

By Theorem 2.22 and Proposition 3.6, we have the following conclusion.

**Corollary 3.7.** *Let  $(Q, [\cdot, \cdot])$  be a finite Leibniz conformal algebra which is free as a  $\mathbf{k}[\partial]$ -module. Suppose that there is a Leibniz conformal algebra structure on  $Q^{*c}$  which is obtained from a  $\mathbf{k}[\partial]$ -module homomorphism  $\Delta : Q \rightarrow Q \otimes Q$ . Then the following conditions are equivalent:*

- (a)  $(Q, Q^{*c}; (L^*, -L^* - R^*), (\mathcal{L}^*, -\mathcal{L}^* - \mathcal{R}^*))$  is a matched pair of Leibniz conformal algebras;
- (b)  $(Q \oplus Q^{*c}, Q, Q^{*c})$  is a conformal Manin triple of Leibniz conformal algebras  $Q$  and  $Q^{*c}$ , where the invariant skew-symmetric bilinear form  $B_\lambda(\cdot, \cdot)$  on  $Q \oplus Q^{*c}$  is given by (22);
- (c)  $(Q, [\cdot, \cdot], \Delta)$  is a Leibniz conformal bialgebra.

#### 4. COBOUNDARY LEIBNIZ CONFORMAL BIALGEBRAS AND CLASSICAL LEIBNIZ CONFORMAL YANG-BAXTER EQUATION

We define the notion of coboundary Leibniz conformal bialgebras which leads us to introduce the notion of classical Leibniz conformal Yang-Baxter equation (CLCYBE). Moreover, the notions of  $\mathcal{O}$ -operators on Leibniz conformal algebras and Leibniz-dendriform conformal algebras are introduced to construct symmetric solutions of the CLCYBE.

For convenience, set  $\partial_1 := \partial \otimes \text{id} \otimes \text{id}$ ,  $\partial_2 := \text{id} \otimes \partial \otimes \text{id}$ ,  $\partial_3 := \text{id} \otimes \text{id} \otimes \partial$  and  $\partial^{\otimes 3} := \partial_1 + \partial_2 + \partial_3$ .

**4.1. Coboundary Leibniz conformal bialgebras.** Let  $Q$  be a  $\mathbf{k}[\partial]$ -module. Define a linear map  $F : Q \rightarrow \text{End}_{\mathbf{k}}(Q \otimes Q)$  by

$$(34) \quad F(a) = L(a)_\lambda \otimes \text{id} + R(a)_\lambda \otimes \text{id} - \text{id} \otimes R(a)_\lambda|_{\lambda=-\partial^{\otimes 2}} \quad \text{for all } a \in Q.$$

If an element  $r \in Q \otimes Q$  satisfies  $F(a)r = 0$  for all  $a \in Q$ , then we call  $r$  **invariant**.

**Definition 4.1.** For a Leibniz conformal bialgebra  $(Q, [\cdot, \cdot], \Delta)$ , if there exists  $r \in Q \otimes Q$  such that

$$(35) \quad \Delta(a) = \Delta_r(a) = F(a)r \quad \text{for all } a \in Q,$$

where  $F$  is given by (34), then  $(Q, [\cdot, \cdot], \Delta = \Delta_r)$  is called a **coboundary Leibniz conformal bialgebra**.

**Definition 4.2.** Let  $Q$  be a Leibniz conformal algebra and  $r = \sum_i a_i \otimes b_i \in Q \otimes Q$ . Define

$$(36) \quad [[r, r]] = [r_{12\mu}r_{13}]|_{\mu=\partial_2} + [r_{23\mu}r_{13}]|_{\mu=\partial_2} - [r_{12\mu}r_{23}]|_{\mu=\partial_1} - [r_{23\mu}r_{12}]|_{\mu=-\partial_1-\partial_2},$$

where

$$\begin{aligned} [r_{12\mu}r_{13}] &= \sum_{i,j} [a_{i\mu}a_j] \otimes b_i \otimes b_j, & [r_{23\mu}r_{13}] &= \sum_{i,j} a_i \otimes a_j \otimes [b_{j\mu}b_i], \\ [r_{12\mu}r_{23}] &= \sum_{i,j} a_i \otimes [b_{i\mu}a_j] \otimes b_j, & [r_{23\mu}r_{12}] &= \sum_{i,j} a_i \otimes [a_{j\mu}b_i] \otimes b_j. \end{aligned}$$

Then  $[[r, r]] = 0 \pmod{\partial^{\otimes 3}}$  in  $Q \otimes Q \otimes Q$  is called the **classical Leibniz conformal Yang-Baxter equation (CLCYBE)** in  $Q$ .

**Proposition 4.3.** Let  $Q$  be a Leibniz conformal algebra and  $r \in Q \otimes Q$ . Then  $r$  is a solution of the CLCYBE in  $Q$  if and only if  $\tau r$  is a solution of the CLCYBE in  $Q$ .

*Proof.* Set  $r = \sum_i a_i \otimes b_i$ . In  $Q \otimes Q \otimes Q$ , we have

$$\begin{aligned} [[\tau r, \tau r]] &= \sum_{i,j} [b_{i\mu}b_j] \otimes a_i \otimes a_j|_{\mu=\partial_2} + b_i \otimes b_j \otimes [a_{j\mu}a_i]|_{\mu=\partial_2} \\ &\quad - b_i \otimes [a_{i\mu}b_j] \otimes a_j|_{\mu=\partial_1} - b_i \otimes [b_{j\mu}a_i] \otimes a_j|_{\mu=-\partial_1-\partial_2} \pmod{\partial^{\otimes 3}} \\ &= \tau_{13}[[r, r]] \pmod{\partial^{\otimes 3}}. \end{aligned}$$

Thus this conclusion holds.  $\square$

**Proposition 4.4.** Let  $Q$  be a Leibniz conformal algebra and  $r = \sum_i a_i \otimes b_i \in Q \otimes Q$ . Define a co-bracket  $\Delta_r : Q \rightarrow Q \otimes Q$  by (35). Then  $(Q, \Delta_r)$  is a Leibniz conformal coalgebra if and only if for all  $a \in Q$  the following equality holds:

$$\begin{aligned} &(\text{id} \otimes \text{id} \otimes R(a)_{-\partial^{\otimes 3}} - \text{id} \otimes (L + R)(a)_{-\partial^{\otimes 3}} \otimes \text{id})(\tau_{12}[[r, r]] - \sum_j F(a_j)(r - \tau(r)) \otimes b_j) \\ (37) \quad &+ \sum_j ((L + R)(a_j)_{\partial_3} \otimes \text{id} \otimes \text{id})(\tau \circ F(a)(r - \tau(r)) \otimes b_j) \\ &+ ((L + R)(a)_{-\partial^{\otimes 3}} \otimes \text{id} \otimes \text{id})[[r, r]] = 0. \end{aligned}$$

*Proof.* For all  $a \in Q$ , we have

$$(\Delta_r \otimes \text{id})\Delta_r(a) + \tau_{12}(\text{id} \otimes \Delta_r)\Delta_r(a) - (\text{id} \otimes \Delta_r)\Delta_r(a) = \sum_{k=1}^9 A_k,$$

where

$$\begin{aligned} A_1 &= \sum_{i,j} (-[a_{i-\partial^2 \otimes \text{id}} a_j] \otimes b_j \otimes [b_{i\partial^2 \otimes \text{id}} a] - [a_{j\partial_2} a_i] \otimes b_j \otimes [b_{i\partial^2 \otimes \text{id}} a] \\ &\quad + a_j \otimes [b_{j\partial_1} a_i] \otimes [b_{i\partial^2 \otimes \text{id}} a] + a_j \otimes a_i \otimes [b_{j\partial_1} [b_{i\partial_2} a]] - a_i \otimes a_j \otimes [b_{j\partial_2} [b_{i\partial_1} a]]) \\ &= (\text{id} \otimes \text{id} \otimes R(a)_{-\partial^3})(a_j \otimes [b_{j\partial_1} a_i] \otimes b_i + a_i \otimes a_j \otimes [b_{i\partial_1} b_j] \\ &\quad - [a_{i-\partial^2 \otimes \text{id}} a_j] \otimes b_j \otimes b_i - [a_{j\partial_2} a_i] \otimes b_j \otimes b_i) \\ &= (\text{id} \otimes \text{id} \otimes R(a)_{-\partial^3})(\tau_{12}[[r, r]] - \sum_j F(a_j)(r - \tau(r)) \otimes b_j) \\ A_2 &= \sum_{i,j} ([[a_{-\partial^3} a_i]_{-\partial^2 \otimes \text{id}} a_j] \otimes b_j \otimes b_i + [[a_{i\partial_3} a]_{-\partial^2 \otimes \text{id}} a_j] \otimes b_j \otimes b_i) = 0, \\ A_3 &= \sum_{i,j} (-[[b_{i\partial_2} a]_{-\partial_1-\partial_3} a_j] \otimes a_i \otimes b_j - [a_{j\partial_3} [b_{i\partial_2} a]] \otimes a_i \otimes b_j), \\ A_4 &= \sum_{i,j} ([a_{j\partial_2} [a_{-\partial^3} a_i]] \otimes b_j \otimes b_i - [a_{-\partial^3} a_i] \otimes [b_{i-\text{id} \otimes \partial^2} a_j] \otimes b_j - [a_{-\partial^3} a_i] \otimes [a_{j\partial_3} b_i] \otimes b_j \\ &\quad + [a_{-\partial^3} a_i] \otimes a_j \otimes [b_{j\partial_2} b_i]) = (L(a)_{-\partial^3} \otimes \text{id} \otimes \text{id})[[r, r]] + \sum_{i,j} ([[a_{i\partial_2} a]_{-\partial_1-\partial_3}] \otimes b_i \otimes b_j), \\ A_5 &= \sum_{i,j} ([b_{i-\partial_1-\partial_3} a_j] \otimes [a_{-\partial^3} a_i] \otimes b_j + [a_{j\partial_3} b_i] \otimes [a_{-\partial^3} a_i] \otimes b_j - a_j \otimes [a_{-\partial^3} a_i] \otimes [b_{j\partial_1} b_i]) \\ &= -(\text{id} \otimes L(a)_{-\partial^3} \otimes \text{id})\tau_{12}[[r, r]] + \sum_{i,j} (b_i \otimes [a_{-\partial^3} [a_{i\partial_1} a_j]] \otimes b_j), \\ A_6 &= \sum_{i,j} (-a_j \otimes [b_{j\partial_1} [a_{-\partial^3} a_i]] \otimes b_i + a_i \otimes [[b_{i\partial_1} a]_{-\text{id} \otimes \partial^2} a_j] \otimes b_j) = \sum_{i,j} -a_i \otimes [a_{-\partial^3} [b_{i\partial_1} a_j]] \otimes b_j, \\ A_7 &= \sum_{i,j} ([a_{j\partial_2} [a_{i\partial_3} a]] \otimes b_j \otimes b_i - [a_{i\text{id} \otimes \partial^2} a] \otimes [b_{i-\text{id} \otimes \partial^2} a_j] \otimes b_j - [a_{i\text{id} \otimes \partial^2} a] \otimes [a_{j\partial_3} b_i] \otimes b_j \\ &\quad + [a_{i\text{id} \otimes \partial^2} a] \otimes a_j \otimes [b_{j\partial_2} b_i]) = (R(a)_{-\partial^3} \otimes \text{id} \otimes \text{id})[[r, r]] + \sum_{i,j} ([a_{j\partial_3} [a_{i\partial_2} a]] \otimes b_i \otimes b_j), \\ A_8 &= \sum_{i,j} ([b_{i-\partial_1-\partial_3} a_j] \otimes [a_{-\partial^3} a_i] \otimes b_j + [a_{j\partial_3} b_i] \otimes [a_{-\partial^3} a_i] \otimes b_j - a_j \otimes [a_{-\partial^3} a_i] \otimes [b_{j\partial_1} b_i]) \\ &= -(\text{id} \otimes R(a)_{-\partial^3} \otimes \text{id})\tau_{12}[[r, r]] + \sum_{i,j} b_i \otimes [[a_{i\partial_1} a_j]_{\partial_1+\partial_3} a] \otimes b_j, \\ A_9 &= \sum_{i,j} (a_i \otimes [a_{j\partial_3} [b_{i\partial_1} a]] \otimes b_j - a_j \otimes [b_{j\partial_1} [a_{i\partial_3} a]] \otimes b_i) = -\sum_{i,j} a_i \otimes [[b_{i\partial_1} a_j]_{\partial_1+\partial_3} a] \otimes b_j. \end{aligned}$$

Thus we have

$$\begin{aligned}
& \sum_{k=1}^9 A_k - (\text{id} \otimes \text{id} \otimes R(a)_{-\partial^{\otimes 3}})(\tau_{12}[[r, r]] - \sum_j F(a_j)(r - \tau(r)) \otimes b_j) \\
& - ((L + R)(a)_{-\partial^{\otimes 3}} \otimes \text{id} \otimes \text{id})[[r, r]] + (\text{id} \otimes (L + R)(a)_{-\partial^{\otimes 3}} \otimes \text{id})\tau_{12}[[r, r]] \\
& - \sum_j ((L + R)(a_j)_{\partial_3} \otimes \text{id} \otimes \text{id})(\tau \circ (F(a)))(r - \tau(r)) \otimes b_j \\
= & \sum_{i,j} (b_i \otimes [a_{-\partial^{\otimes 3}}[a_{i\partial_1} a_j]] \otimes b_j - a_i \otimes [a_{-\partial^{\otimes 3}}[b_{i\partial_1} a_j]] \otimes b_j + b_i \otimes [[a_{i\partial_1} a_j]_{\partial_1 + \partial_3} a] \otimes b_j \\
& - a_i \otimes [[b_{i\partial_1} a_j]_{\partial_1 + \partial_3} a] \otimes b_j - [b_{i-\partial_1 - \partial_3} a_j] \otimes [a_{-\partial^{\otimes 3}} a_i] \otimes b_j - [b_{i-\partial_1 - \partial_3} a_j] \otimes [a_{i\partial_1 + \partial_3} a] \otimes b_j \\
& + [a_{i-\partial_1 - \partial_3} a_j] \otimes [a_{-\partial^{\otimes 3}} b_i] \otimes b_j + [a_{i-\partial_1 - \partial_3} a_j] \otimes [b_{i\partial_1 + \partial_3} a] \otimes b_j - [a_{j\partial_3} b_i] \otimes [a_{-\partial^{\otimes 3}} a_i] \otimes b_j \\
& - [a_{j\partial_3} b_i] \otimes [a_{i\partial_1 + \partial_3} a] \otimes b_j + [a_{j\partial_3} a_i] \otimes [a_{-\partial^{\otimes 3}} b_i] \otimes b_j + [a_{j\partial_3} a_i] \otimes [b_{i\partial_1 + \partial_3} a] \otimes b_j) \\
= & (\text{id} \otimes (L + R)(a)_{-\partial^{\otimes 3}} \otimes \text{id}) \sum_j F(a_j)(r - \tau(r)) \otimes b_j,
\end{aligned}$$

which means that (37) holds, since

$$\begin{aligned}
& \sum_{i,j} ([a_{i\partial_2} a]_{-\partial_1 - \partial_3}) \otimes b_i \otimes b_j + [a_{j\partial_3} [a_{i\partial_2} a]] \otimes b_i \otimes b_j - [[b_{i\partial_2} a]_{-\partial_1 - \partial_3} a_j] \otimes a_i \otimes b_j \\
& - [a_{j\partial_3} [b_{i\partial_2} a]] \otimes a_i \otimes b_j \\
= & \sum_j ((L + R)(a_j)_{\partial_3} \otimes \text{id} \otimes \text{id})(\tau \circ (F(a)))(r - \tau(r)) \otimes b_j + \sum_{i,j} (-[b_{i-\partial_1 - \partial_3} a_j] \otimes [a_{-\partial^{\otimes 3}} a_i] \otimes b_j \\
& - [b_{i-\partial_1 - \partial_3} a_j] \otimes [a_{i\partial_1 + \partial_3} a] \otimes b_j + [a_{i-\partial_1 - \partial_3} a_j] \otimes [a_{-\partial^{\otimes 3}} b_i] \otimes b_j + [a_{i-\partial_1 - \partial_3} a_j] \otimes [b_{i\partial_1 + \partial_3} a] \otimes b_j \\
& - [a_{j\partial_3} b_i] \otimes [a_{-\partial^{\otimes 3}} a_i] \otimes b_j - [a_{j\partial_3} b_i] \otimes [a_{i\partial_1 + \partial_3} a] \otimes b_j + [a_{j\partial_3} a_i] \otimes [a_{-\partial^{\otimes 3}} b_i] \otimes b_j \\
& + [a_{j\partial_3} a_i] \otimes [b_{i\partial_1 + \partial_3} a] \otimes b_j).
\end{aligned}$$

Therefore, this conclusion holds.  $\square$

**Theorem 4.5.** *Let  $Q$  be a Leibniz conformal algebra and  $r = \sum_i a_i \otimes b_i \in Q \otimes Q$ . The triple  $(Q, [\cdot, \cdot], \Delta_r)$  is a Leibniz conformal bialgebra where  $\Delta_r$  is defined by (35) if and only if (37) and the following conditions hold for all  $a, b \in Q$ :*

$$(38) \quad \tau(R(b)_{-\lambda - \partial^{\otimes \text{id}}} \otimes \text{id})F(a)(r - \tau(r)) = 0,$$

$$(39) \quad ((L + R)(b)_{-\lambda - \partial^{\otimes 2}} \otimes \text{id})\tau(F(a)(r - \tau(r))) - \tau(R(b)_{-\lambda - \partial^{\otimes 2}} \otimes \text{id})(F(a)(r - \tau(r))) = 0.$$

*Proof.* We only need to check that  $\Delta_r$  satisfies (27) and (28) if and only if (38) and (39) hold. By (27), we have

$$\begin{aligned}
& (R(a)_{\lambda - \partial^{\otimes \text{id}}} \otimes \text{id})\Delta_r(b) - \tau((R(b)_{-\lambda - \partial^{\otimes \text{id}}} \otimes \text{id})\Delta_r(a)) \\
= & \sum_i ([b_{-\lambda - \text{id} \otimes \partial} a_i]_{-\lambda} a] \otimes b_i + [[a_{i\text{id} \otimes \partial} b]_{-\lambda} a] \otimes b_i - [a_{i-\lambda} a] \otimes [b_{i\lambda} b] \\
& - b_i \otimes [[a_{\lambda - \partial^{\otimes \text{id}}} a_i]_{\lambda} b] - b_i \otimes [[a_{i\partial^{\otimes \text{id}}} a]_{\lambda} b] + [b_{i-\lambda} a] \otimes [a_{i\lambda} b]) \\
= & \sum_i (-[a_{i-\lambda} a] \otimes [b_{i\lambda} b] + [b_{i-\lambda} a] \otimes [a_{i\lambda} b]) = -\tau(R(b)_{-\lambda - \partial^{\otimes \text{id}}} \otimes \text{id})F(a)(r - \tau(r)),
\end{aligned}$$



which means that (27) holds if and only if (38) holds. By (28), we have

$$\begin{aligned}
& \Delta_r([a_\lambda b]) - (\text{id} \otimes R(b)_{-\lambda-\partial^{\otimes 2}} - L(b)_{-\lambda-\partial^{\otimes 2}} \otimes \text{id} - R(b)_{-\lambda-\partial^{\otimes 2}} \otimes \text{id})(\tau + \text{id})\Delta_r(a) \\
& - (\text{id} \otimes L(a)_\lambda + L(a)_\lambda \otimes \text{id})\Delta_r(b) \\
= & \sum_i ([a_\lambda b]_{-\partial^{\otimes 2}} a_i) \otimes b_i + [a_{i\text{id}\otimes\partial}[a_\lambda b]] \otimes b_i - a_i \otimes [b_{i\partial\text{id}}[a_\lambda b]] - ([b_{-\lambda-\partial^{\otimes 2}} a_i] \otimes [a_\lambda b_i] \\
& + [a_{i\lambda+\text{id}\otimes\partial} b] \otimes [a_\lambda b_i] - a_i \otimes [a_\lambda [b_{i\partial\text{id}} b]] + [a_\lambda [b_{-\lambda-\partial^{\otimes 2}} a_i]] \otimes b_i + [a_\lambda [a_{i\text{id}\otimes\partial} b]] \otimes b_i \\
& - [a_\lambda a_i] \otimes [b_{i\lambda+\partial\text{id}} b]) - ([a_\lambda a_i] \otimes [b_{i\lambda+\partial\text{id}} b] + [a_{i-\lambda-\partial\text{id}} a] \otimes [b_{i\lambda+\partial\text{id}} b] \\
& - a_i \otimes [[b_{i\partial\text{id}} a]_{\lambda+\partial\text{id}} b] + b_i \otimes [[a_\lambda a_i]_{\lambda+\partial\text{id}} b] + b_i \otimes [[a_{i\partial\text{id}} a]_{\lambda+\partial\text{id}} b] - [b_{-\lambda-\partial^{\otimes 2}} [a_\lambda a_i]] \otimes b_i \\
& - [b_{i-\lambda-\partial\text{id}} a] \otimes [a_{i\lambda+\partial\text{id}} b] - [b_{-\lambda-\partial^{\otimes 2}} [a_{i\text{id}\otimes\partial} a]] \otimes b_i + [b_{-\lambda-\partial^{\otimes 2}} a_i] \otimes [b_{i-\lambda-\text{id}\otimes\partial} a] \\
& - [b_{-\lambda-\partial^{\otimes 2}} b_i] \otimes [a_\lambda a_i] - [b_{-\lambda-\partial^{\otimes 2}} b_i] \otimes [a_{i-\lambda-\text{id}\otimes\partial} a] + [b_{-\lambda-\partial^{\otimes 2}} [b_{i\text{id}\otimes\partial} a]] \otimes a_i \\
& - [[a_\lambda a_i]_{\lambda+\text{id}\otimes\partial} b] \otimes b_i - [[a_{i\text{id}\otimes\partial} a]_{\lambda+\text{id}\otimes\partial} b] \otimes b_i + [a_{i\lambda+\text{id}\otimes\partial} b] \otimes [b_{i-\lambda-\text{id}\otimes\partial} a] \\
& - [b_{i\lambda+\text{id}\otimes\partial} b] \otimes [a_\lambda a_i] - [b_{i\lambda+\text{id}\otimes\partial} b] \otimes [a_{i-\lambda-\text{id}\otimes\partial} a] + [[b_{i\text{id}\otimes\partial} a]_{\lambda+\text{id}\otimes\partial} b] \otimes a_i) = \sum_{j=1}^4 B_j,
\end{aligned}$$

where

$$\begin{aligned}
B_1 = & \sum_i ([a_{i\text{id}\otimes\partial}[a_\lambda b]] \otimes b_i - [a_{i\lambda+\text{id}\otimes\partial} b] \otimes [a_\lambda b_i] - [a_\lambda [a_{i\text{id}\otimes\partial} b]] \otimes b_i + [b_{i\lambda+\text{id}\otimes\partial} b] \otimes [a_\lambda a_i] \\
& - [b_{i\text{id}\otimes\partial} a]_{\lambda+\text{id}\otimes\partial} b] \otimes a_i) = (R(b)_{-\lambda-\partial^{\otimes 2}} \otimes \text{id})(\tau - \text{id})(F(a)(r - \tau(r))),
\end{aligned}$$

$$\begin{aligned}
B_2 = & - \sum_i ([b_{-\lambda-\partial^{\otimes 2}} a_i] \otimes [a_\lambda b_i] - [b_{-\lambda-\partial^{\otimes 2}} [a_{i\text{id}\otimes\partial} a]] \otimes b_i + [b_{-\lambda-\partial^{\otimes 2}} a_i] \otimes [b_{i-\lambda-\text{id}\otimes\partial} a] \\
& - [b_{-\lambda-\partial^{\otimes 2}} b_i] \otimes [a_\lambda a_i] - [b_{-\lambda-\partial^{\otimes 2}} b_i] \otimes [a_{i-\lambda-\text{id}\otimes\partial} a] + [b_{-\lambda-\partial^{\otimes 2}} [b_{i\text{id}\otimes\partial} a]] \otimes a_i) \\
= & (L(b)_{-\lambda-\partial^{\otimes 2}} \otimes \text{id})\tau(F(a)(r - \tau(r))),
\end{aligned}$$

$$\begin{aligned}
B_3 = & - \sum_i ([a_{i-\lambda-\partial\text{id}} a] \otimes [b_{i\lambda+\partial\text{id}} b] - [b_{i-\lambda-\partial\text{id}} a] \otimes [a_{i\lambda+\partial\text{id}} b]) \\
= & - \tau(R(b)_{-\lambda-\partial^{\otimes 2}} \otimes \text{id})(F(a)(r - \tau(r))),
\end{aligned}$$

$$B_4 = - \sum_i ([a_{i\lambda+\text{id}\otimes\partial} b] \otimes [b_{i-\lambda-\text{id}\otimes\partial} a] - [b_{i\lambda+\text{id}\otimes\partial} b] \otimes [a_{i-\lambda-\text{id}\otimes\partial} a]) = (R(b)_{-\lambda-\partial^{\otimes 2}} \otimes \text{id})(F(a)(r - \tau(r))).$$

Thus, (28) holds if and only if (39) holds, finishing the proof.  $\square$

By Theorem 4.5, we get the following conclusion.

**Corollary 4.6.** *Let  $(Q, [\cdot, \cdot])$  be a Leibniz conformal algebra and  $r \in Q \otimes Q$  be a solution of the CLCYBE in  $Q$ . Then  $(Q, [\cdot, \cdot], \Delta_r)$  is a Leibniz conformal bialgebra if  $r - \tau(r)$  is invariant, and we call it a **quasi-triangular Leibniz conformal bialgebra**. In particular, if  $r$  is symmetric, we call  $(Q, [\cdot, \cdot], \Delta_r)$  a **triangular Leibniz conformal bialgebra**.*

**4.2. Operator forms of the classical Leibniz conformal Yang-Baxter equation.** First we recall the notion of  $\mathcal{O}$ -operators on Leibniz conformal algebras.

**Definition 4.7.** [15] Let  $Q$  be a Leibniz conformal algebra and  $(M, l, r)$  be a representation of  $Q$ . A  $\mathbf{k}[\partial]$ -module homomorphism  $T : M \rightarrow Q$  is called an  $\mathcal{O}$ -operator on  $Q$  associated to  $(M, l, r)$  if for all  $x, y \in M$ , it satisfies

$$(40) \quad [T(x)_\lambda T(y)] = T(l(T(x))_\lambda y + r(T(y))_{-\lambda-\partial} x).$$

Let  $Q$  be a finite Leibniz conformal algebra which is free as a  $\mathbf{k}[\partial]$ -module. Define a linear map  $\sigma : Q \otimes Q \rightarrow \text{Chom}(Q^{*c}, Q)$  by

$$(41) \quad \langle f, \sigma(a \otimes b)_{\lambda} g \rangle_{\mu} = \langle g \otimes f, a \otimes b \rangle_{-\lambda-\mu, \mu} \quad \text{for all } f, g \in Q^{*c} \text{ and } a, b \in Q.$$

Thus  $\sigma(a \otimes b)_{\lambda} g = \langle g, a \rangle_{-\lambda-\partial} b$ , where  $\partial^Q$  represents the action of  $\partial$  on  $Q$ . Moreover,  $\sigma$  is a  $\mathbf{k}[\partial]$ -module isomorphism [17].

**Lemma 4.8.** *Let  $Q$  be a finite Leibniz conformal algebra which is free as a  $\mathbf{k}[\partial]$ -module and  $r \in Q \otimes Q$ . Define a co-bracket  $\Delta_r : Q \rightarrow Q \otimes Q$  by (35). Then  $Q^{*c}$  is endowed with a Leibniz conformal algebra structure  $(Q^{*c}, [\cdot, \cdot]^*)$ , where the  $\lambda$ -bracket is given by*

$$(42) \quad [f, g]_{\lambda}^* = L^*(\sigma(r)_0 f)_{\lambda} g - (L^* + R^*)(\sigma(\tau r)_0 g)_{-\lambda-\partial} f \quad \text{for all } f, g \in Q^{*c},$$

where  $\sigma(r)_0 f = \sigma(r)_{\lambda} f|_{\lambda=0}$ .

*Proof.* Using the assumptions from the proof of Proposition 3.6, and letting  $r = \sum_{i,j} r_{ij}(\partial \otimes \text{id}, \text{id} \otimes \partial) a^i \otimes a^j$  where  $r_{ij}(\lambda, \mu) \in \mathbf{k}[\lambda, \mu]$ , we have

$$\begin{aligned} \Delta_r(a^k) &= (L(a^k)_{\partial^{\otimes 2}} \otimes \text{id} + R(a^k)_{\partial^{\otimes 2}} \otimes \text{id} - \text{id} \otimes R(a^k)_{\partial^{\otimes 2}}) r \\ &= \sum_{i,j,s} (r_{ij}(-\text{id} \otimes \partial, \text{id} \otimes \partial) P_s^{ki}(-\partial^{\otimes 2}, \partial \otimes \text{id}) a^s \otimes a^j + r_{ij}(-\text{id} \otimes \partial, \text{id} \otimes \partial) P_s^{ik}(\text{id} \otimes \partial, \partial \otimes \text{id}) \\ &\quad a^s \otimes a^j - r_{ij}(\partial \otimes \text{id}, -\partial \otimes \text{id}) P_s^{jk}(\partial \otimes \text{id}, \text{id} \otimes \partial) a^i \otimes a^s), \end{aligned}$$

and  $\sigma(r)_{\lambda}(a_i) = \sum_j r_{ij}(-\lambda - \partial, \partial) a^j$ . Then the facts that

$$\begin{aligned} \langle a_{m\lambda} a_n, a^k \rangle_{\mu} &= \langle a_m \otimes a_n, \Delta_r(a^k) \rangle_{\lambda, \mu-\lambda} \\ &= \sum_i r_{in}(\lambda - \mu, \mu - \lambda) (P_m^{ki}(-\mu, \lambda) + P_m^{ik}(\mu - \lambda, \lambda)) - \sum_j r_{mj}(\lambda, -\lambda) P_n^{jk}(\lambda, \mu - \lambda) \end{aligned}$$

and

$$\begin{aligned} &\langle L^*(\sigma(r)_0 a_m)_{\lambda} a_n - (L^* + R^*)(\sigma(\tau r)_0 a_n)_{-\lambda-\partial} a_m, a^k \rangle_{\mu} \\ &= \langle \sum_j r_{mj} L^*(a^j)_{\lambda} a_n, a^k \rangle_{\mu} + \sum_i r_{in}(\lambda - \mu, \mu - \lambda) \langle (L^* + R^*)(a^i)_{-\lambda+\mu} a_m, a^k \rangle_{\mu} \\ &= - \sum_j r_{mj}(\lambda, -\lambda) P_n^{jk}(\lambda, \mu - \lambda) + \sum_i r_{in}(\lambda - \mu, \mu - \lambda) (P_m^{ki}(-\mu, \lambda) + P_m^{ik}(\mu - \lambda, \lambda)) \end{aligned}$$

imply that the conclusion holds.  $\square$

**Theorem 4.9.** *Let  $Q$  be a finite Leibniz conformal algebra which is free as a  $\mathbf{k}[\partial]$ -module and  $r \in Q \otimes Q$ . Then  $r$  is a solution of the CLCYBE in  $Q$  if and only if  $\sigma(\tau r) \in \text{Chom}(Q^{*c}, Q)$  satisfies*

$$(43) \quad [\sigma(\tau r)_0 x_{\lambda} \sigma(\tau r)_0 y] = \sigma(\tau r)_0 (L^*(\sigma(r)_0 x)_{\lambda} y - (L^* + R^*)(\sigma(\tau r)_0 y)_{-\lambda-\partial} x) \quad \text{for all } x, y \in Q^{*c}.$$

*Moreover, if  $r - \tau r$  is invariant, then  $r$  is a solution of the CLCYBE in  $Q$  if and only if  $\sigma(r)$  satisfies*

$$(44) \quad [\sigma(r)_0 x_{\lambda} \sigma(r)_0 y] = \sigma(r)_0 (L^*(\sigma(r)_0 x)_{\lambda} y - (L^* + R^*)(\sigma(r)_0 y)_{-\lambda-\partial} x) \quad \text{for all } x, y \in Q^{*c}.$$

*Proof.* Set  $r = \sum_i a_i \otimes b_i$ . By (41), we have

$$\sigma(r)_{\lambda} x = \sum_i \langle x, a_i \rangle_{-\lambda-\partial} b_i, \quad \sigma(\tau r)_{\lambda} x = \sum_i \langle x, b_i \rangle_{-\lambda-\partial} a_i \quad \text{for all } x \in Q^{*c}.$$

It is obvious that the following cases are equivalent:

- (a)  $[[r, r]] \bmod (\partial^{\otimes 3}) = 0$ ;
- (b)  $\langle x \otimes y \otimes z, [[r, r]] \bmod (\partial^{\otimes 3}) \rangle_{\lambda, \mu, \xi} = 0$  for all  $x, y, z \in Q^{*c}$ ;
- (c)  $\langle x \otimes y \otimes z, [[r, r]] \rangle_{\lambda, \mu, \xi} = 0 \bmod (\lambda + \mu + \xi)$  for all  $x, y, z \in Q^{*c}$ .

For all  $x, y, z \in Q^{*c}$ , we have

$$\begin{aligned} \sum_{i,j} \langle x \otimes y \otimes z, [a_{i\partial_2} a_j] \otimes b_i \otimes b_j \rangle_{\lambda, \mu, \xi} &= \sum_{i,j} \langle x, [a_{i\mu} a_j] \rangle_{\lambda} \langle y, b_i \rangle_{\mu} \langle z, b_j \rangle_{\xi} \\ &= \langle x, [(\sum_i \langle y, b_i \rangle_{\mu} a_i)_{\mu} \sum_j \langle z, b_j \rangle_{\xi} a_j] \rangle_{\lambda} = \langle x, [\sigma(\tau r)_0(y)_{\mu} \sigma(\tau r)_{-\xi-\mu-\lambda} z] \rangle_{\lambda}, \end{aligned}$$

and

$$\begin{aligned} \sum_{i,j} \langle x \otimes y \otimes z, a_i \otimes a_j \otimes [b_{j\partial_2} b_i] \rangle_{\lambda, \mu, \xi} \\ &= \sum_i \langle x, a_i \rangle_{\lambda} \langle z, [\sigma(r)_0(y)_{\mu} b_i] \rangle_{\xi} = - \sum_i \langle x, a_i \rangle_{\lambda} \langle L^*(\sigma(r)_0(y))_{\mu} z, b_i \rangle_{\mu+\xi} \\ &= - \sum_i \langle x, \langle L^*(\sigma(r)_0(y))_{\mu} z, b_i \rangle_{\mu+\xi} a_i \rangle_{\lambda} = - \langle x, \sigma(\tau r)_{-\lambda-\mu-\xi} (L^*(\sigma(r)_0(y))_{\mu} z) \rangle_{\lambda}. \end{aligned}$$

Similarly, we can get

$$\begin{aligned} \sum_{i,j} \langle x \otimes y \otimes z, a_i \otimes [b_{i\partial_1} a_j] \otimes b_j \rangle_{\lambda, \mu, \xi} &= - \langle x, \sigma(\tau r)_0 (R^*(\sigma(\tau r)_{-\lambda-\mu-\xi}(z))_{-\mu-\partial} y) \rangle_{\lambda}, \\ \sum_{i,j} \langle x \otimes y \otimes z, a_i \otimes [a_{j-\partial^{\otimes 2}} b_i] \otimes b_j \rangle_{\lambda, \mu, \xi} &= - \langle x, \sigma(\tau r)_0 (L^*(\sigma(\tau r)_{-\lambda-\mu-\xi}(z))_{-\mu-\partial} y) \rangle_{\lambda}. \end{aligned}$$

Thus,  $[[r, r]] \bmod (\partial^{\otimes 3}) = 0$  if and only if

$$(45) \quad [\sigma(\tau r)_0(y)_{\mu} \sigma(\tau r)_0(z)] - \sigma(\tau r)_0(L^*(\sigma(r)_0(y))_{\mu} z) + \sigma(\tau r)_0((L^* + R^*)(\sigma(\tau r)_0(z))_{-\mu-\partial} y) = 0$$

for all  $y, z \in Q^{*c}$ . The first conclusion follows by replacing  $\mu$  by  $\lambda$  in (45). On the other hand, we have

$$\begin{aligned} &\langle z, [\sigma(r)_0 x_{\lambda} \sigma(r)_0 y] - \sigma(r)_0(L^*(\sigma(r)_0 x)_{\lambda} y - (L^* + R^*)(\sigma(\tau r)_0 y)_{-\lambda-\partial} x) \rangle_{\mu} \\ &= \sum_{i,j} \langle x \otimes y \otimes z, a_i \otimes a_j \otimes [b_{i\lambda} b_j] + a_j \otimes [b_{j\lambda} a_i] \otimes b_i - [a_{j-\mu-\lambda} a_i] \otimes b_j \otimes b_i - [a_{i\mu} a_j] \otimes b_j \otimes b_i \rangle_{\lambda, -\lambda-\mu, \mu} \\ &= \sum_j \langle x \otimes y \otimes z, \tau_{12}[[r, r]] - F(a_j)(r - \tau r) \rangle_{\lambda, -\lambda-\mu, \mu}, \end{aligned}$$

for all  $x, y, z \in Q^{*c}$ , which means that the second conclusion holds.  $\square$

By Proposition 4.3, Lemma 4.8 and Theorem 4.9, we obtain the following conclusion.

**Corollary 4.10.** *Let  $Q$  be a finite Leibniz conformal algebra which is free as a  $\mathbf{k}[\partial]$ -module and  $r$  be a solution of the CLCYBE in  $Q$ . Then the following conclusions hold:*

- (a)  $\delta(\tau r)_0$  is a Leibniz conformal algebra homomorphism from  $(Q^{*c}, [\cdot, \lambda]^*)$  to  $(Q, [\cdot, \lambda])$ .  
 (b) If  $(Q, [\cdot, \lambda], \Delta_r)$  is a quasi-triangular Leibniz conformal bialgebra, then  $\delta(r)_0$  and  $\delta(\tau r)_0$  are both Leibniz conformal algebra homomorphisms from  $(Q^{*c}, [\cdot, \lambda]^*)$  to  $(Q, [\cdot, \lambda])$ . Particularly, if  $r$  is symmetric, then  $\delta(r)_0$  is an  $\mathcal{O}$ -operator on  $Q$  associated to  $(Q^{*c}, L^*, -L^* - R^*)$ .

Let  $Q$  be a Leibniz conformal algebra and  $(M, l_Q, r_Q)$  be a representation of  $Q$ , where  $Q$  and  $M$  are finite and free as  $\mathbf{k}[\partial]$ -modules. Then  $(M^{*c}, l_Q^*, -l_Q^* - r_Q^*)$  is a representation of  $Q$  by Proposition 2.14. By [7, Proposition 6.1], there is an isomorphism of  $\mathbf{k}[\partial]$ -modules  $\varphi : M^{*c} \otimes Q \rightarrow \text{Chom}(M, Q)$  defined by

$$\varphi(f \otimes a)_\lambda v = \langle v, f \rangle_{-\lambda - \partial} a \quad \text{for all } a \in Q, v \in M, f \in M^{*c}.$$

Therefore, the inverse  $\mathbf{k}[\partial]$ -module isomorphism of  $\varphi$  is  $\eta : \text{Chom}(M, Q) \rightarrow M^{*c} \otimes Q$  given by

$$\langle x \otimes g, \eta(T) \rangle_{\lambda, \mu} = \langle g, T_{-\lambda - \mu}(x) \rangle_\mu \quad \text{for all } x \in M, g \in Q^{*c} \text{ and } T \in \text{Chom}(M, Q).$$

Consequently, for any  $T \in \text{Chom}(M, Q)$ , we have  $\eta(T) \in M^{*c} \otimes Q \subset (Q \times_{l_Q^*, -l_Q^* - r_Q^*} M^{*c}) \otimes (Q \times_{l_Q^*, -l_Q^* - r_Q^*} M^{*c})$ .

**Theorem 4.11.** *Let  $Q$  be a Leibniz conformal algebra,  $(M, l_Q, r_Q)$  be a representation of  $Q$  and both  $Q$  and  $M$  are finite and free as  $\mathbf{k}[\partial]$ -modules. Let  $T \in \text{Chom}(M, Q)$  and  $\eta(T) \in M^{*c} \otimes Q \subset (Q \times_{l_Q^*, -l_Q^* - r_Q^*} M^{*c}) \otimes (Q \times_{l_Q^*, -l_Q^* - r_Q^*} M^{*c})$ . Then  $r = \eta(T) + \tau\eta(T)$  is a symmetric solution of the CLCYBE in the Leibniz conformal algebra  $Q \times_{l_Q^*, -l_Q^* - r_Q^*} M^{*c}$  if and only if  $T_0 = T_{\lambda|_{\lambda=0}}$  is an  $\mathcal{O}$ -operator on  $Q$  associated to  $(M, l_Q, r_Q)$ .*

*Proof.* Let  $\{a^1, \dots, a^n\}$  be a  $\mathbf{k}[\partial]$ -basis of  $Q$ ,  $\{v^1, \dots, v^m\}$  be a  $\mathbf{k}[\partial]$ -basis of  $M$  and  $\{v_1, \dots, v_m\}$  be a  $\mathbf{k}[\partial]$ -basis of  $M^{*c}$ . Assume that for all  $i = 1, \dots, m$ ,

$$T_\lambda(v_i) = \sum_{j=1}^n g_{ij}(\lambda, \partial) a^j, \quad g_{ij}(\lambda, \partial) \in \mathbf{k}[\lambda, \partial].$$

Then we have  $\eta(T) = \sum_i^m \sum_j^n g_{ij}(-\partial^{\otimes 2}, \text{id} \otimes \partial) v_i^* \otimes a^j$ . Thus we obtain

$$r = \eta(T) + \tau\eta(T) = \sum_i^m \sum_j^n (g_{ij}(-\partial^{\otimes 2}, \text{id} \otimes \partial) v_i^* \otimes a^j + g_{ij}(-\partial^{\otimes 2}, \partial \otimes \text{id}) a^j \otimes v_i^*).$$

On the other hand, by (13), we have

$$l_Q^*(a^i)_\lambda v_j^* = - \sum_t \langle v_j^*, l_Q(a^i)_\lambda v_t \rangle_{-\lambda - \partial} v_t^*, \quad r_Q^*(a^i)_\lambda v_j^* = - \sum_t \langle v_j^*, r_Q(a^i)_\lambda v_t \rangle_{-\lambda - \partial} v_t^*.$$

Then we have

$$\begin{aligned}
& [[r, r]] \quad \text{mod } (\partial^{\otimes 3}) \\
&= \sum_{i,t}^m \sum_{s,j}^n (g_{ij}(0, \partial_2) g_{ts}(-\partial^{\otimes 3}, \partial^{\otimes 2} \otimes \text{id}) [v_{i\partial_2}^* a^s] \otimes a^j \otimes v_t^* \\
&\quad + g_{ts}(0, -\partial_2) g_{ij}(-\partial^{\otimes 3}, \partial_3) [a^s_{\partial_2} v_i^*] \otimes v_t^* \otimes a^j + g_{ij}(0, -\partial_2) g_{ts}(-\partial^{\otimes 3}, \partial^{\otimes 2} \otimes \text{id}) [a^j_{\partial_2} a^s] \otimes v_i^* \otimes v_t^* \\
&\quad + g_{ts}(0, -\partial_2) g_{ij}(-\partial^{\otimes 3}, \text{id} \otimes \partial^{\otimes 2}) v_i^* \otimes v_t^* \otimes [a^s_{\partial_2} a^j] + g_{ts}(0, \partial_2) g_{ij}(-\partial^{\otimes 3}, \text{id} \otimes \partial^{\otimes 2}) v_i^* \otimes a^s \otimes [v_{i\partial_2}^* a^j] \\
&\quad + g_{ij}(0, -\partial_2) g_{ts}(-\partial^{\otimes 3}, \partial_1) a^s \otimes v_i^* \otimes [a^j_{\partial_2} v_t^*] - g_{ij}(0, -\partial_1) g_{ts}(-\partial^{\otimes 3}, \partial^{\otimes 2} \otimes \text{id}) v_i^* \otimes [a^j_{\partial_1} a^s] \otimes v_t^* \\
&\quad - g_{ij}(0, -\partial_1) g_{ts}(-\partial^{\otimes 3}, \partial_3) v_i^* \otimes [a^j_{\partial_1} v_t^*] \otimes a^s - g_{ij}(0, \partial_1) g_{ts}(-\partial^{\otimes 3}, \partial^{\otimes 2} \otimes \text{id}) a^j \otimes [v_{i\partial_1}^* a^s] \otimes v_t^* \\
&\quad - g_{ij}(0, -\partial_1) g_{ts}(-\partial^{\otimes 3}, \partial_3) v_i^* \otimes [v_{i-\partial^{\otimes 2}}^* a^j] \otimes a^s - g_{ij}(0, -\partial_1) g_{ts}(-\partial^{\otimes 3}, \partial^{\otimes 2} \otimes \text{id}) v_i^* \otimes [a^s_{-\partial^{\otimes 2}} a^j] \otimes v_t^* \\
&\quad - g_{ij}(0, \partial_1) g_{ts}(-\partial^{\otimes 3}, \partial^{\otimes 2} \otimes \text{id}) a^j \otimes [a^s_{-\partial^{\otimes 2}} v_i^*] \otimes v_t^*) \quad \text{mod } (\partial^{\otimes 3}) \\
&= \sum_{i,t}^m ([v_{i\partial_2}^* T_0(v_t)] \otimes T_0(v_i) \otimes v_t^* + [T_0(v_t)_{\partial_2} v_i^*] \otimes v_t^* \otimes T_0(v_i) + [T_0(v_i)_{\partial_2} T_0(v_t)] \otimes v_i^* \otimes v_t^* \\
&\quad + v_i^* \otimes v_t^* \otimes [T_0(v_t)_{\partial_2} T_0(v_i)] + v_i^* \otimes T_0(v_t) \otimes [v_{i\partial_2}^* T_0(v_i)] + T_0(v_t) \otimes v_i^* \otimes [T_0(v_i)_{\partial_2} v_t^*] \\
&\quad - v_i^* \otimes [T_0(v_i)_{\partial_1} T_0(v_t)] \otimes v_t^* - v_i^* \otimes [T_0(v_i)_{\partial_1} v_t^*] \otimes T_0(v_t) \\
&\quad - T_0(v_i) \otimes [v_{i\partial_1}^* T_0(v_t)] \otimes v_t^* - v_i^* \otimes [v_{i-\partial^{\otimes 2}}^* T_0(v_i)] \otimes T_0(v_t) \\
&\quad - v_i^* \otimes [T_0(v_t)_{-\partial^{\otimes 2}} T_0(v_i)] \otimes v_t^* - T_0(v_i) \otimes [T_0(v_t)_{-\partial^{\otimes 2}} v_i^*] \otimes v_t^*) \quad \text{mod } (\partial^{\otimes 3}).
\end{aligned}$$

Since  $T_0$  is a  $\mathbf{k}[\partial]$ -module homomorphism, we have

$$\begin{aligned}
& \sum_{i,t}^m [v_{i\partial_2}^* T_0(v_t)] \otimes T_0(v_i) \otimes v_t^* \quad \text{mod } (\partial^{\otimes 3}) = - \sum_{i,t}^m (l_Q^* + r_Q^*)(T_0(v_t))_{-\partial^{\otimes 2} \otimes \text{id}} v_i^* \otimes T_0(v_i) \otimes v_t^* \quad \text{mod } (\partial^{\otimes 3}) \\
&= - \sum_{i,t}^m (l_Q^* + r_Q^*)(T_0(v_t))_{\partial_3} v_i^* \otimes T_0(v_i) \otimes v_t^* \quad \text{mod } (\partial^{\otimes 3}) \\
&= \sum_{i,t,j}^m \langle v_i^*, (l_Q + r_Q)(T_0(v_t))_{\partial_3} v_j \rangle_{\partial_2} v_j^* \otimes T_0(v_i) \otimes v_t^* \quad \text{mod } (\partial^{\otimes 3}) \\
&= \sum_{i,t,j}^m v_j^* \otimes T_0(\langle v_i^*, (l_Q + r_Q)(T_0(v_t))_{\partial_3} v_j \rangle_{\partial_2} v_i) \otimes v_t^* \quad \text{mod } (\partial^{\otimes 3}) \\
&= \sum_{t,j}^m v_j^* \otimes T_0((l_Q + r_Q)(T_0(v_t))_{\partial_3} v_j) \otimes v_t^* \quad \text{mod } (\partial^{\otimes 3}).
\end{aligned}$$

Similarly, we get

$$\begin{aligned}
& [[r, r]] \quad \text{mod } (\partial^{\otimes 3}) \\
&= \sum_{i,t}^m (([T_0(v_i)_{\partial_2} T_0(v_t)] - T_0(l_Q(T_0(v_i))_{\partial_2} v_t) - T_0(r_Q(T_0(v_t))_{\partial_3} v_i)) \otimes v_i^* \otimes v_t^* \\
&\quad + v_i^* \otimes (T_0(l_Q(T_0(v_i))_{\partial_1} v_t) + r_Q(T_0(v_t))_{\partial_3} v_i) - [T_0(v_i)_{\partial_1} T_0(v_t)]) \otimes v_t^*
\end{aligned}$$

$$\begin{aligned}
& + v_i^* \otimes (T_0(l_Q(T_0(v_i)))_{\partial_3} v_i + r_Q(T_0(v_i))_{\partial_1} v_i) - [T_0(v_i)_{\partial_3} T_0(v_i)] \otimes v_i^* \\
& + v_i^* \otimes v_i^* \otimes ([T_0(v_i)_{\partial_2} T_0(v_i)] - T_0(l_Q(T_0(v_i)))_{\partial_2} v_i + r_Q(T_0(v_i))_{\partial_1} v_i) \pmod{(\partial^{\otimes 3})}.
\end{aligned}$$

Thus  $r$  is a solution of the CLCYBE in the Leibniz conformal algebra  $Q \ltimes_{l_Q^*, -l_Q^*, -r_Q^*} M^{*c}$  if and only if for all  $i, t \in \{1, \dots, m\}$ ,

$$[T_0(v_i)_\lambda T_0(v_t)] = T_0(l_Q(T_0(v_i)))_\lambda v_t + T_0(r_Q(T_0(v_i)))_{-\lambda-\partial} v_t.$$

Hence this conclusion holds.  $\square$

In what follows, we recall the notion of Leibniz-dendriform conformal algebras.

**Definition 4.12.** [16] A **Leibniz-dendriform conformal algebra (LDCA)** is a triple  $(Q, \triangleleft_\lambda, \triangleright_\lambda)$  consisting of a  $\mathbf{k}[\partial]$ -module  $Q$  and two  $\lambda$ -products  $\triangleleft_\lambda, \triangleright_\lambda : Q \times Q \rightarrow Q[\lambda]$  which satisfy the following conditions:

- (46)  $(\partial a) \triangleleft_\lambda b = -\lambda a \triangleleft_\lambda b, \quad a \triangleleft_\lambda (\partial b) = (\lambda + \partial) a \triangleleft_\lambda b,$
- (47)  $(\partial a) \triangleright_\lambda b = -\lambda a \triangleright_\lambda b, \quad a \triangleright_\lambda (\partial b) = (\lambda + \partial) a \triangleright_\lambda b,$
- (48)  $a \triangleright_\lambda (b \triangleright_\mu c + b \triangleleft_\mu c) = (a \triangleright_\lambda b) \triangleright_{\lambda+\mu} c + b \triangleleft_\mu (a \triangleright_\lambda c),$
- (49)  $a \triangleleft_\lambda (b \triangleright_\mu c) = (a \triangleleft_\lambda b) \triangleright_{\lambda+\mu} c + b \triangleright_\mu (a \triangleleft_\lambda c + a \triangleleft_\lambda c),$
- (50)  $a \triangleleft_\lambda (b \triangleleft_\mu c) = (a \triangleright_\lambda b + a \triangleleft_\lambda b) \triangleleft_{\lambda+\mu} c + b \triangleleft_\mu (a \triangleleft_\lambda c)$  for all  $a, b, c \in Q$ .

**Remark 4.13.** [16] Let  $(Q, \triangleleft_\lambda, \triangleright_\lambda)$  be a LDCA. Define  $[\cdot, \cdot]_{\triangleleft, \triangleright}$  by

$$(51) \quad [a_\lambda b]_{\triangleleft, \triangleright} = a \triangleright_\lambda b + a \triangleleft_\lambda b \quad \text{for all } a, b \in Q.$$

Then  $(Q, [\cdot, \cdot]_{\triangleleft, \triangleright})$  is a Leibniz conformal algebra.

Let  $(Q, \triangleleft_\lambda, \triangleright_\lambda)$  be a LDCA. Define two  $\mathbf{k}[\partial]$ -module homomorphisms  $L_\triangleleft$  and  $R_\triangleright$  from  $Q$  to  $\text{Cend}(Q)$  by  $L_\triangleleft(a)_\lambda b = a \triangleleft_\lambda b, R_\triangleright(a)_{-\lambda-\partial} b = b \triangleright_\lambda a$  for all  $a, b \in Q$ .

**Proposition 4.14.** Let  $(Q, \triangleleft_\lambda, \triangleright_\lambda)$  be a LDCA. Then  $(Q, L_\triangleleft, R_\triangleright)$  is a representation of the Leibniz conformal algebra  $(Q, [\cdot, \cdot]_{\triangleleft, \triangleright})$ , where  $[\cdot, \cdot]_{\triangleleft, \triangleright}$  is defined by (51). Moreover, the identity map  $\text{id} : Q \rightarrow Q$  is an  $\mathcal{O}$ -operator on  $(Q, [\cdot, \cdot]_{\triangleleft, \triangleright})$  associated to  $(Q, L_\triangleleft, R_\triangleright)$ .

*Proof.* It is straightforward.  $\square$

**Proposition 4.15.** [16, Proposition 2.8] Let  $T : M \rightarrow Q$  be an  $\mathcal{O}$ -operator on  $(Q, [\cdot, \cdot]_{\triangleleft, \triangleright})$  associated to  $(M, l_Q, r_Q)$ . Then there is an LDCA structure on  $M$  given by

$$x \triangleleft_\lambda y = l_Q(T(x))_\lambda y, \quad x \triangleright_\lambda y = r_Q(T(y))_{-\lambda-\partial} x \quad \text{for all } x, y \in M.$$

Finally, we give a construction of symmetric solutions of the CLCYBE from LDCAs.

**Theorem 4.16.** Let  $(Q, \triangleleft_\lambda, \triangleright_\lambda)$  be a finite LDCA which is free as a  $\mathbf{k}[\partial]$ -module. Then

$$r = \sum_{i=1}^n (a_i \otimes a^i + a^i \otimes a_i)$$

is a solution of the CLCYBE in the Leibniz conformal algebra  $Q \ltimes_{l_Q^*, -l_Q^*, -r_Q^*} Q^{*c}$ , where  $\{a^i\}_{i=1}^n$  is a  $\mathbf{k}[\partial]$ -basis of  $Q$  and  $\{a_i\}_{i=1}^n$  is the dual  $\mathbf{k}[\partial]$ -basis of  $Q^{*c}$ .

*Proof.* By Proposition 4.14,  $T = \text{id} : Q \rightarrow Q$  is an  $\mathcal{O}$ -operator on  $(Q, [\cdot, \cdot]_{\triangleleft, \triangleright})$  associated to  $(Q, L_\triangleleft, R_\triangleright)$ . Then the conclusion follows from Theorem 4.11.  $\square$

## 5. A CLASS OF LEIBNIZ CONFORMAL BIALGEBRAS CORRESPONDING TO NOVIKOV BI-DIALGEBRAS

We introduce the notion of Novikov bi-dialgebras and show that there is a correspondence between Novikov bi-dialgebras and a class of Leibniz conformal bialgebras. Furthermore, we introduce the notion of classical duplicated Novikov Yang-Baxter equation whose symmetric solutions give Novikov bi-dialgebras. In addition, we show that symmetric solutions of the classical duplicated Novikov Yang-Baxter equation correspond to a class of symmetric solutions of the CLCYBE. In this section, we assume that  $A$  is finite-dimensional as a vector space.

5.1. **Novikov bi-dialgebras.** First, we introduce the notion of Novikov co-dialgebras.

**Definition 5.1.** A **Novikov co-dialgebra** is a triple  $(A, \delta, \Delta)$  where  $\Delta, \delta : A \rightarrow A \otimes A$  are linear maps satisfying

$$(52) \quad (\delta \otimes \text{id})\delta(a) = \tau_{23}(\delta \otimes \text{id})\delta(a),$$

$$(53) \quad (\Delta \otimes \text{id})\delta(a) = \tau_{23}(\Delta \otimes \text{id})\Delta(a) = \tau_{23}(\delta \otimes \text{id})\Delta(a),$$

$$(54) \quad (\text{id} \otimes \delta)\delta(a) = (\text{id} \otimes \Delta)\delta(a),$$

$$(55) \quad (\delta \otimes \text{id} - \text{id} \otimes \delta)\delta(a) = \tau_{12}((\Delta \otimes \text{id})\delta(a) - (\text{id} \otimes \delta)\Delta(a)),$$

$$(56) \quad (\Delta \otimes \text{id} - \text{id} \otimes \Delta)\Delta(a) = \tau_{12}(\Delta \otimes \text{id} - \text{id} \otimes \Delta)\Delta(a) \quad \text{for all } a \in A.$$

It is easy to check that  $(A, \delta, \Delta)$  is a Novikov co-dialgebra if and only if  $(A^*, \dashv, \vdash)$  is a Novikov dialgebra defined by

$$\langle x \dashv y, a \rangle = \langle x \otimes y, \delta(a) \rangle, \quad \langle x \vdash y, a \rangle = \langle x \otimes y, \Delta(a) \rangle \quad \text{for all } x, y \in A^*, a, b \in A.$$

**Proposition 5.2.** Let  $\bar{A} = \mathbf{k}[\partial]A$  be the free  $\mathbf{k}[\partial]$ -module on a vector space  $A$ . Suppose that  $\delta$  and  $\Delta$  are  $\mathbf{k}[\partial]$ -module homomorphisms from  $\bar{A}$  to  $\bar{A} \otimes \bar{A}$  such that  $\delta(A) \subset A \otimes A$  and  $\Delta(A) \subset A \otimes A$ . Define a  $\mathbf{k}[\partial]$ -module homomorphism  $\alpha : \bar{A} \rightarrow \bar{A} \otimes \bar{A}$  by

$$(57) \quad \alpha(a) = (\partial \otimes \text{id})\Delta(a) - \tau(\partial \otimes \text{id})\delta(a) \quad \text{for all } a \in A.$$

Then  $(\bar{A}, \alpha)$  is a Leibniz conformal coalgebra if and only if  $(A, \delta, \Delta)$  is a Novikov co-dialgebra.

*Proof.* Let  $a \in A$ . By the definition of  $\alpha$ , we have

$$\begin{aligned} (\text{id} \otimes \alpha)\alpha(a) &= (\text{id} \otimes \alpha)((\partial \otimes \text{id})\Delta(a) - \tau(\partial \otimes \text{id})\delta(a)) \\ &= (\partial \otimes \partial \otimes \text{id})(\text{id} \otimes \Delta)\Delta(a) - (\partial \otimes \text{id} \otimes \partial)\tau_{23}(\text{id} \otimes \delta)\Delta(a) \\ &\quad - \partial_2(\text{id} \otimes \partial^{\otimes 2})(\text{id} \otimes \Delta)\tau\delta(a) + \partial_3(\text{id} \otimes \partial^{\otimes 2})\tau_{23}(\text{id} \otimes \delta)\tau\delta(a), \\ (\alpha \otimes \text{id})\alpha(a) &= (\alpha \otimes \text{id})((\partial \otimes \text{id})\Delta(a) - \tau(\partial \otimes \text{id})\delta(a)) \\ &= \partial_1(\partial^{\otimes 2} \otimes \text{id})(\Delta \otimes \text{id})\Delta(a) - \partial_2(\partial^{\otimes 2} \otimes \text{id})\tau_{12}(\delta \otimes \text{id})\Delta(a) \\ &\quad - (\partial \otimes \text{id} \otimes \partial)(\Delta \otimes \text{id})\tau\delta(a) + (\text{id} \otimes \partial \otimes \partial)\tau_{12}(\delta \otimes \text{id})\tau\delta(a). \end{aligned}$$

Then we consider

$$(58) \quad (\text{id} \otimes \alpha)\alpha(a) = (\alpha \otimes \text{id})\alpha(a) + \tau_{12}(\text{id} \otimes \alpha)\alpha(a) \quad \text{for all } a \in A.$$

Considering the terms of (58) in  $A \otimes A \otimes \partial^2 A$ , we can get

$$\tau_{13}(\delta \otimes \text{id})\delta(a) = \tau_{13}\tau_{23}(\delta \otimes \text{id})\delta(a) \quad \text{for all } a \in A,$$

which is equivalent to (52). Similarly, by considering the terms of (58) in  $\partial^2 A \otimes A \otimes A$  and  $A \otimes \partial^2 A \otimes A$ , we get that (53) holds. By considering the terms of (58) in  $\partial A \otimes A \otimes \partial A$  and  $A \otimes \partial A \otimes \partial A$ , we obtain (54). Moreover, we have (55) and (56) by considering the terms of (58) in  $A \otimes \partial A \otimes \partial A$  and  $\partial A \otimes \partial A \otimes A$ . Then this conclusion holds.  $\square$

**Definition 5.3.** Let  $(A, \dashv, \vdash)$  be a Novikov dialgebra and  $(A, \delta, \Delta)$  be a Novikov co-dialgebra. If  $\Delta$  and  $\delta$  satisfy the following conditions for all  $a, b \in A$ :

$$(59) \quad (L_-(a) \otimes \text{id})\Delta(b) = -(\text{id} \otimes (L_- + R_+)(b))\delta(a),$$

$$(60) \quad (\text{id} \otimes L_-(b))\delta(a) = (L_-(a) \otimes \text{id})\tau\delta(b),$$

$$(61) \quad (\text{id} \otimes (L_- + R_+)(a))(\delta(b) + \tau\Delta(b)) = (R_+(b) \otimes \text{id})\Delta(a),$$

$$(62) \quad \delta(a \dashv b) = (R_+(b) \otimes \text{id})\delta(a) + (\text{id} \otimes L_+(a))(\text{id} - \tau)(\delta(b) - \Delta(b)),$$

$$(63) \quad \delta(a \vdash b) = (R_+(b) \otimes \text{id})(\delta(a) - \Delta(a)) + (\text{id} \otimes (L_- + R_+)(a))(\delta(b) + \tau\Delta(b)),$$

$$(64) \quad \Delta(a \dashv b) = (\text{id} \otimes (L_- + R_+)(a))(\text{id} - \tau)(\Delta(b) - \delta(b)) + (R_-(b) \otimes \text{id})\Delta(a),$$

$$(65) \quad \Delta(a \vdash b) = ((R_- - R_+)(b) \otimes \text{id})(\delta(a) - \Delta(a)) + (\text{id} \otimes (L_- + R_+)(a))(\tau\delta(b) + \Delta(b)),$$

$$(66) \quad ((L_- + R_+)(b) \otimes \text{id})(\delta(a) - \Delta(a)) + (\text{id} \otimes L_+(b))\tau(\delta(a) - \Delta(a)) \\ - ((L_- + R_+)(a) \otimes \text{id})\delta(b) + (\text{id} \otimes (L_- + R_+)(a))\tau\Delta(b) = 0,$$

$$(67) \quad (\text{id} \otimes R_+(a))(\text{id} - \tau)(\delta(b) - \Delta(b)) - (R_-(b) \otimes \text{id})(\Delta(a) + \tau\delta(a)) \\ = -(\text{id} \otimes R_-(b))(\Delta(a) + \tau\delta(a)) + ((R_- - R_+)(a) \otimes \text{id})(\text{id} - \tau)(\Delta(b) - \delta(b)),$$

then  $(A, \dashv, \vdash, \delta, \Delta)$  is called a **Novikov bi-dialgebra**.

Next, we give a correspondence between Novikov bi-dialgebras and a class of Leibniz conformal bialgebras as follows.

**Theorem 5.4.** Let  $\bar{A} = \mathbf{k}[\partial]A$  be the Leibniz conformal algebra corresponding to a Novikov dialgebra  $(A, \dashv, \vdash)$ . Suppose that  $\delta$  and  $\Delta$  are two  $\mathbf{k}[\partial]$ -module homomorphisms from  $\bar{A}$  to  $\bar{A} \otimes \bar{A}$  such that  $\delta(A) \subset A \otimes A$  and  $\Delta(A) \subset A \otimes A$ . Then  $(\bar{A}, [\cdot, \cdot], \alpha)$  is a Leibniz conformal bialgebra if and only if  $(A, \dashv, \vdash, \delta, \Delta)$  is a Novikov bi-dialgebra, where  $\alpha$  is given by (57). We call  $(\bar{A}, [\cdot, \cdot], \alpha)$  the **Leibniz conformal bialgebra corresponding to a Novikov bi-dialgebra**  $(A, \dashv, \vdash, \delta, \Delta)$ .

*Proof.* Let  $a, b \in A$  and set  $\Delta(a) = \sum a_{(1)} \otimes a_{(2)}$  and  $\delta(a) = \sum a^{(1)} \otimes a^{(2)}$ . Then we have

$$\begin{aligned} & (R(a)_{\lambda-\partial \otimes \text{id}} \otimes \text{id})\alpha(b) - \tau((R(b)_{-\lambda-\partial \otimes \text{id}} \otimes \text{id})\alpha(a)) \\ &= (R(a)_{\lambda-\partial \otimes \text{id}} \otimes \text{id})((\partial \otimes \text{id})\Delta(b) - \tau(\partial \otimes \text{id})\delta(b)) - \tau((R(b)_{-\lambda-\partial \otimes \text{id}} \otimes \text{id})((\partial \otimes \text{id})\Delta(a) - \tau(\partial \otimes \text{id})\delta(a))) \\ &= \sum [\partial b_{(1)-\lambda} a] \otimes b_{(2)} - \sum [b_{-\lambda}^{(2)} a] \otimes \partial b^{(1)} - \tau \left( \sum [\partial a_{(1)\lambda} b] \otimes a_{(2)} - [a_{\lambda}^{(2)} b] \otimes \partial a^{(1)} \right) \\ &= \sum (\lambda(\partial(a \dashv b_{(1)})) - \lambda(a \dashv b_{(1)} + b_{(1)} \vdash a) \otimes b_{(2)}) + \sum (-\partial(a \dashv b^{(2)}) \otimes \partial b^{(1)} \\ &\quad + \lambda(a \dashv b^{(2)} + b^{(2)} \vdash a) \otimes \partial b^{(1)}) + \sum (a_{(2)} \otimes (\lambda\partial(b \dashv a_{(1)}) + \lambda^2(b \dashv a_{(1)} + a_{(1)} \vdash b))) \\ &\quad + \sum (\partial a^{(1)} \otimes (\partial(b \dashv a^{(2)}) + \lambda(a^{(2)} \vdash b + b \dashv a^{(2)}))). \end{aligned}$$

Thus (27) holds if and only if

$$\begin{aligned} (L_-(a) \otimes \text{id})\Delta(b) &= -(\text{id} \otimes (L_- + R_+)(b))\delta(a), & ((L_- + R_+)a \otimes \text{id})\Delta(b) &= (\text{id} \otimes (L_- + R_+)b)\tau\Delta(a), \\ (L_-(a) \otimes \text{id})\tau\delta(b) &= (\text{id} \otimes L_-(b))\delta(a), \end{aligned}$$

by considering the coefficients of  $\lambda(\partial A \otimes A)$ ,  $\lambda^2 A \otimes A$  and  $\partial^2 A \otimes A$ . That is, (27) holds if and only if (59)-(61) hold.



Similarly, we have

$$\begin{aligned}
& (\text{id} \otimes R(b)_{-\lambda-\partial^2} - L(b)_{-\lambda-\partial^2} \otimes \text{id} - R(b)_{-\lambda-\partial^2} \otimes \text{id})(\tau + \text{id})\Delta(a) \\
& + (\text{id} \otimes L(a)_\lambda + L(a)_\lambda \otimes \text{id})\Delta(b) - \alpha([a_\lambda b]) \\
= & (\partial \otimes \partial)((\text{id} \otimes L_+(b))(\text{id} - \tau)(\Delta(a) - \delta(a)) + ((L_- - L_+)(b) \otimes \text{id})(\text{id} - \tau)(\Delta(a) - \delta(a)) \\
& + (\text{id} \otimes R_+(a))\Delta(b) - (R_+(a) \otimes \text{id})\tau\delta(b)) + (\partial^2 \otimes \text{id})(\text{id} \otimes (L_+ + R_+)(b))(\text{id} - \tau)(\Delta(a) - \delta(a)) \\
& + (R_+(a) \otimes \text{id})\Delta(b) + (\text{id} \otimes \partial^2)((L_\star(b) \otimes \text{id})(\text{id} - \tau)(\delta(a) - \Delta(a)) - (\text{id} \otimes R_+(a))\tau\delta(b)) \\
& + \lambda(\partial \otimes \text{id})(\text{id} \otimes (L_+ + R_+)(b))((\text{id} - \tau)(\Delta(a) - \delta(a)) - \tau(\Delta(a) - \delta(a))) \\
& + ((L_- - L_+)(b) \otimes \text{id})(\delta(a) - \Delta(a)) + (\text{id} \otimes (L_+ + R_+)(a))\Delta(b) + (L_- + 2R_- \otimes \text{id})\Delta(b) \\
& + \lambda(\text{id} \otimes \partial)((\text{id} \otimes L_+(b))\tau(\delta(a) - \Delta(a)) + (L_\star(b) \otimes \text{id})(\text{id} - \tau)(\delta(a) - \Delta(a)) \\
& + (L_\star(b) \otimes \text{id})(\delta(a) - \Delta(a)) - (\text{id} \otimes (L_+ + 2R_+)a + (L_+ + R_+)(a) \otimes \text{id})\tau\delta(b)) \\
& + \lambda^2((\text{id} \otimes (L_+ + R_+)(b))\tau(\delta(a) - \Delta(a)) + (L_\star(b) \otimes \text{id})(\delta(a) - \Delta(a)) \\
& - (\text{id} \otimes (L_+ + R_+)(a))\tau\delta(b) + ((L_+ + R_+) \otimes \text{id})\Delta(b)) - (\partial \otimes \partial)(\Delta(b \dashv a) - \tau\delta(b \dashv a)) \\
& - (\partial^2 \otimes \text{id})\Delta(b \dashv a) + (\text{id} \otimes \partial^2)\tau\delta(b \dashv a) - \lambda((\partial \otimes \text{id})\Delta(b \dashv a + a \vdash b) \\
& + (\text{id} \otimes \partial)\tau\delta(b \dashv a + a \vdash b)) = 0.
\end{aligned}$$

Then by considering the coefficients of  $\partial^2 A \otimes A$ ,  $A \otimes \partial^2 A$  and  $\lambda^2 A \otimes A$  in the above equality, we obtain that the three equalities are equivalent to (64), (62) and (66) respectively. Furthermore by considering the coefficients of  $\partial A \otimes \partial A$ ,  $\lambda(\partial A \otimes A)$  and  $\lambda(A \otimes \partial A)$  in the above equality, we get that the three equalities are equivalent to (67), (65) and (63) respectively. Hence the conclusion holds.  $\square$

## 5.2. The classical duplicate Novikov Yang-Baxter equation.

Let  $(A, \dashv, \vdash)$  be a Novikov dialgebra and  $r = \sum_i a_i \otimes b_i \in A \otimes A$ . We define

$$N(r) = \sum_{i,j} a_i \vdash a_j \otimes b_i \otimes b_j + a_i \otimes (b_i \vdash a_j - b_i \dashv a_j) \otimes b_j + a_i \otimes a_j \otimes (b_j \vdash b_i + b_i \dashv b_j).$$

The equation  $N(r) = 0$  is called the **classical duplicate Novikov Yang-Baxter equation (CD-NYBE)** in  $A$ . In addition, we define

$$M(r) = \sum_{i,j} a_i \vdash b_j \otimes b_i \otimes a_j + a_i \otimes (b_i \vdash b_j - b_i \dashv b_j) \otimes a_j + a_i \otimes b_j \otimes (a_j \vdash b_i + b_i \dashv a_j).$$

It is clear that if  $r$  is symmetric, then  $N(r) = M(r)$ .

**Proposition 5.5.** *Let  $(A, \dashv, \vdash)$  be a Novikov dialgebra and  $r = \sum_i a_i \otimes b_i \in A \otimes A$ . Define*

$$(68) \quad \Delta_r : A \rightarrow A \otimes A, \quad \Delta_r(a) = ((L_- - L_+)(a) \otimes \text{id} - \text{id} \otimes (L_+ + R_+)(a))r,$$

$$(69) \quad \delta_r : A \rightarrow A \otimes A, \quad \delta_r(a) = (\text{id} \otimes L_\star(a) + L_+(a) \otimes \text{id})\tau r \quad \text{for all } a \in A.$$

*Then  $(A, \Delta_r, \delta_r)$  is a Novikov co-dialgebra if and only if the following conditions hold for all  $a \in A$  :*

$$(70) \quad (\text{id} - \tau_{23})(\text{id} \otimes \text{id} \otimes (L_\star(a))(\tau_{13}N(r) - N(\tau r)))$$

$$+ (\text{id} - \tau_{23}) \sum_i (\text{id} \otimes \text{id} \otimes L_\star(a_i))(b_i \otimes (\text{id} \otimes L_-(a))(r - \tau r)) = 0,$$

$$(71) \quad -(\text{id} \otimes \text{id} \otimes L_\star(a))\tau_{23}\tau_{12}N(r) + \tau_{12} \sum_i b_i \otimes (\text{id} \otimes (L_+ + R_+)(a \dashv a_i - a \vdash a_i))(r - \tau r)$$

$$- (\text{id} \otimes (L_+ + R_+)(a) \otimes \text{id})\tau_{12}M(\tau r) + (\text{id} \otimes (L_+ + R_+)(b_i) \otimes \text{id})(a_i \otimes (L_-(a) \otimes \text{id}))(r - \tau r)$$

$$- \tau_{23}((L_-(a \dashv a_i - a \vdash a_i) \otimes \text{id})(r - \tau r) \otimes b_i) = 0,$$

$$(72) \quad \sum_i (L_-(a \dashv a_i - a \vdash a_i) \otimes \text{id} + \text{id} \otimes (L_+ + R_+)(a \dashv a_i - a \vdash a_i))(r - \tau r) \otimes b_i$$

$$+ (\text{id} \otimes \text{id} \otimes (L_+ + R_+)(a))(\tau_{23}\tau_{12}M(\tau r) + \tau_{13}M(\tau r) - M(\tau r))$$

$$+ \sum_i (b_i \otimes (\text{id} \otimes L_-(a_i))r - a_i \otimes (\text{id} \otimes L_-(b_i))\tau r) = 0,$$

$$(73) \quad (L_-(a) \otimes \text{id} \otimes \text{id})(M(\tau r) + \tau_{23}M(\tau r) - \tau_{12}\tau_{23}M(\tau r) + \sum_i ((R_+(b_i) \otimes \text{id})r \otimes a_i$$

$$- (R_+(a_i) \otimes \text{id})\tau r \otimes b_i)) + \sum_i b_i \otimes ((\text{id} \otimes (L_+ + R_+)(a \star a_i) - L_-(a \star a_i) \otimes \text{id})(r - \tau r)) = 0$$

$$(74) \quad (L_-(a) \otimes \text{id} \otimes \text{id})(N(\tau r) - \tau_{12}N(\tau r) + \sum_i ((R_+ - R_+)(b_i) \otimes \text{id})(r - \tau r) \otimes a_i)$$

$$+ (\text{id} \otimes (L_- - L_+)(a) \otimes \text{id})(\tau_{12}\tau_{23}N(r) - \tau_{23}N(\tau r) + \sum_i b_i \otimes (L_-(a_i) \otimes \text{id})(r - \tau r))$$

$$+ \sum_i (\text{id} \otimes \text{id} \otimes L_\star(a_i))(b_i \otimes (\text{id} \otimes (L_+ + R_+)(a))(r - \tau r)) = 0,$$

$$(75) \quad ((L_- - L_+)(a) \otimes \text{id} \otimes \text{id})N(r) - \sum_i (R_+(a \dashv a_i - a \vdash a_i) \otimes \text{id})(r - \tau r) \otimes b_i$$

$$+ \sum_i ((R_+ - R_+)(a_i) \otimes \text{id} \otimes \text{id})((R_+(a) \otimes \text{id})(r - \tau r) \otimes b_i) - (\text{id} \otimes (L_- - L_+)(a) \otimes \text{id})\tau_{12}N(r)$$

$$+ \sum_i (\text{id} \otimes R_+(a \dashv a_i - a \vdash a_i) + (\text{id} \otimes (R_+ - R_+)(a_i))(\text{id} \otimes R_+(a)))(r - \tau r) \otimes b_i$$

$$+ \sum_i (\text{id} \otimes \text{id} \otimes (L_+ + R_+)(a))(\text{id} - \tau_{12})(a_i \otimes (\text{id} \otimes L_-(b_i))(r - \tau r)) = 0.$$

*Proof.* We just need to prove that (70)-(75) are equivalent to (52)-(56). For all  $a \in A$ , we have

$$(\delta_r \otimes \text{id})\delta_r(a) - \tau_{23}((\delta_r \otimes \text{id})\delta_r(a))$$

$$= \sum_{i,j} (\text{id} - \tau_{23})(b_j \otimes (b_i \star a_j) \otimes (a \star a_i) + (b_i \dashv b_j) \otimes a_j \otimes (a \star a_i) + b_j \otimes ((a \dashv b_i) \star a_j) \otimes a_i$$

$$+ (a \dashv b_i) \dashv b_j \otimes a_j \otimes a_i) = (\text{id} - \tau_{23})(\text{id} \otimes \text{id} \otimes L_\star(a))(\tau_{13}N(r) - N(\tau r))$$

$$+ (\text{id} - \tau_{23}) \sum_{i,j} (b_i \otimes b_j \otimes a \star (a_i \dashv a_j) + b_j \otimes ((a \dashv b_i) \star a_j) \otimes a_i)$$

$$= (\text{id} - \tau_{23})\{(\text{id} \otimes \text{id} \otimes L_\star(a))(\tau_{13}N(r) - N(\tau r)) + \sum_i (\text{id} \otimes \text{id} \otimes L_\star(a_i))(b_i \otimes (\text{id} \otimes L_-(a))(r - \tau r))\},$$

which means that (52) is equivalent to (70). Similarly, we have

$$(53) \iff (71) \text{ and } (72), \quad (54) \iff (73), \quad (55) \iff (74), \quad (56) \iff (75).$$

Thus this conclusion holds.  $\square$

**Theorem 5.6.** *Let  $(A, \dashv, \vdash)$  be a Novikov dialgebra and  $r \in A \otimes A$ . The maps  $\Delta_r$  and  $\delta_r$  given by (68) and (69) respectively are co-multiplications of a Novikov bi-dialgebra  $A$  if and only if they satisfy (70)-(75) and the following conditions for all  $a, b \in A$  :*

$$(76) \quad (L_{\dashv}(a) \otimes \text{id})(\text{id} \otimes (L_{\dashv} + R_{\vdash})(b))(r - \tau r) = 0,$$

$$(77) \quad (L_{\dashv}(a) \otimes \text{id})(\text{id} \otimes L_{\dashv}(b))(r - \tau r) = 0,$$

$$(78) \quad (R_{\vdash}(b) \otimes \text{id})(\text{id} \otimes (L_{\dashv} + R_{\vdash})(a))(r - \tau r) = 0,$$

$$(79) \quad (\text{id} \otimes L_{\star}(a))(\text{id} \otimes R_{\vdash}(b) + (R_{\vdash} - R_{\dashv})(b) \otimes \text{id})(r - \tau r) = 0,$$

$$(80) \quad ((R_{\vdash} - R_{\dashv})(b) \otimes \text{id})(L_{\dashv}(a) \otimes \text{id} + \text{id} \otimes L_{\star}(a))(r - \tau r) = 0,$$

$$(81) \quad (\text{id} \otimes L_{\star}(b))(\text{id} \otimes L_{\dashv}(a) + L_{\star}(a) \otimes \text{id})(r - \tau r) \\ + ((L_{\dashv} + R_{\vdash})(b) \otimes \text{id})(\text{id} \otimes (L_{\dashv} + R_{\vdash})(a))(r - \tau r) = 0,$$

$$(82) \quad (R_{\vdash}(b) \otimes \text{id})(\text{id} \otimes R_{\vdash}(a))(r - \tau r) + ((R_{\vdash} - R_{\dashv})(a) \otimes \text{id})(\text{id} \otimes (R_{\vdash} - R_{\dashv})(b))(r - \tau r) \\ + (R_{\dashv}(b) \otimes \text{id})(R_{\vdash}(a) \otimes \text{id})(r - \tau r) - (R_{\dashv}(a) \otimes \text{id})(R_{\vdash}(b) \otimes \text{id})(r - \tau r) = 0.$$

*Proof.*  $r = \sum_i a_i \otimes b_i$ . For all  $a, b \in A$ , we have

$$\begin{aligned} & (L_{\dashv}(a) \otimes \text{id})\Delta_r(b) + (\text{id} \otimes (L_{\dashv} + R_{\vdash})(b))\delta_r(a) \\ &= \sum_i ((a \dashv (b \dashv a_i) - a \dashv (b \vdash a_i)) \otimes b_i - (a \dashv a_i) \otimes (b \dashv b_i + b_i \dashv b) \\ & \quad + (a \dashv b_i) \otimes (b \dashv a_i + a_i \vdash b) + b_i \otimes (b \dashv (a \star a_i) + (a \star a_i) \vdash b)) \\ &= - (L_{\dashv}(a) \otimes \text{id})(\text{id} \otimes (L_{\dashv} + R_{\vdash})(b))(r - \tau r), \end{aligned}$$

which means that (59) is equivalent to (76). Similarly, we have

$$(60) \iff (77), \quad (61), (63) \text{ and } (64) \iff (78), \quad (62) \iff (79), \\ (65) \iff (80), \quad (66) \iff (81), \quad (67) \iff (82).$$

Thus this conclusion holds.  $\square$

**Corollary 5.7.** *Let  $(A, \dashv, \vdash)$  be a Novikov dialgebra. If  $r \in A \otimes A$  is a symmetric solution of the CDNYBE in  $A$ , then  $(A, \dashv, \vdash, \delta_r, \Delta_r)$  is a Novikov bi-dialgebra, where  $\Delta_r$  and  $\delta_r$  are defined by (68) and (69) respectively.*

*Proof.* It is straightforward by Theorem 5.6.  $\square$

Next, we give the relationship between the CLCYBE and the CDNYBE as follows.

**Proposition 5.8.** *Let  $(A, \dashv, \vdash)$  be a Novikov dialgebra,  $(\bar{A} = \mathbf{k}[\partial]A, [\cdot, \cdot])$  be the Leibniz conformal algebra corresponding to  $(A, \dashv, \vdash)$  and  $r \in A \otimes A$  be symmetric. Then  $r$  is a solution of the CLCYBE in  $\bar{A}$  if and only if  $r$  is a solution of the CDNYBE in  $A$ .*

*Proof.* Set  $r = \sum_i a_i \otimes b_i$ . Note that

$$\begin{aligned}
[[r, r]] &= \sum_{i,j} (\partial_1(a_i \vdash a_i \otimes b_i \otimes b_j - a_i \otimes (b_i \vdash a_j + a_j \vdash b_i) \otimes b_j) \\
&\quad + \partial_2((a_i \vdash a_j + a_j \vdash a_i) \otimes b_i \otimes b_j + a_i \otimes a_j \otimes (b_j \vdash b_i + b_i \vdash b_j) \\
&\quad - a_i \otimes a_j \vdash (b_i + b_i \vdash a_j) \otimes b_j) + \partial_3(a_i \otimes a_j \otimes b_i + b_j - a_i \otimes (a_j \vdash b_i + b_i \vdash a_j) \otimes b_j)) \\
&= \sum_{i,j} (\partial_1(-a_i \vdash a_j \otimes b_i \otimes b_j - a_i \otimes (b_i \vdash a_j - b_i \vdash a_j) \otimes b_j - a_i \otimes a_j \otimes (b_j \vdash b_i + b_i \vdash b_j)) \\
&\quad + \partial_3(-a_i \otimes a_j \otimes (b_j \vdash b_i) - a_i \otimes (a_j \vdash b_i - a_j \vdash b_i) \otimes b_j \\
&\quad - (a_i \vdash a_j + a_j \vdash a_i) \otimes b_i \otimes b_j)) \pmod{(\partial^{\otimes 3})}.
\end{aligned}$$

Since  $r$  is symmetric, we have  $[[r, r]] = -\partial_1 N(r) - \partial_3 \tau_{13} N(r)$ . Thus this conclusion holds.  $\square$

**Corollary 5.9.** *Let  $(A, \vdash, \dashv)$  be a Novikov dialgebra and  $(\bar{A} = \mathbf{k}[\partial]A, [\cdot, \lambda \cdot])$  be the Leibniz conformal algebra corresponding to  $(A, \vdash, \dashv)$ . Suppose that  $r \in A \otimes A$  is a symmetric solution of the CDNYBE in  $A$ . On the one hand, there is a Novikov bi-dialgebra  $(A, \vdash, \dashv, \delta_r, \Delta_r)$  by Corollary 5.7 where  $\Delta_r$  and  $\delta_r$  are defined by (68) and (69) respectively. By Theorem 5.4, let  $(\bar{A}, [\cdot, \lambda \cdot], \alpha_r)$  be the corresponding Leibniz conformal bialgebra, where  $\alpha_r$  is defined by (57). On the other hand,  $r$  is also the symmetric solution of the CLCYBE in  $\bar{A}$  by Proposition 5.8. By Corollary 5.7, there is a triangular Leibniz conformal bialgebra  $(\bar{A}, [\cdot, \lambda \cdot], \alpha'_r)$ , where  $\alpha'_r$  is defined by (35). Therefore both the Leibniz conformal bialgebras  $(\bar{A}, [\cdot, \lambda \cdot], \alpha_r)$  and  $(\bar{A}, [\cdot, \lambda \cdot], \alpha'_r)$  coincide. That is, there is the following commutative diagram.*

$$(83) \quad \begin{array}{ccc}
\text{symmetric solutions of} & \xrightarrow{\text{Coro.5.7}} & \text{Novikov} \\
\text{the CDNYBE} & & \text{bi-dialgebras} \\
\updownarrow \text{Prop.5.8} & & \updownarrow \text{Thm.5.4} \\
\text{symmetric solutions of} & \xrightarrow{\text{Coro.4.6}} & \text{Leibniz} \\
\text{the CLCYBE} & & \text{conformal bialgebras}
\end{array}$$

*Proof.* Let  $r = \sum_i a_i \otimes b_i$  and  $a \in A$ . By Proposition 2.8, we have

$$\begin{aligned}
\alpha'_r(a) &:= (L(a)_\lambda \otimes \text{id} + R(a)_\lambda \otimes \text{id} - \text{id} \otimes R(a)_\lambda)(r)|_{\lambda=-\partial^{\otimes 2}} \\
&= \sum_i ([a_{-\partial^{\otimes 2} a_i}] \otimes b_i + [a_{\text{id} \otimes \partial a}] \otimes b_i - a_i \otimes [b_{\partial \otimes \text{id} a}]) \\
&= (\partial \otimes \text{id})((-a \vdash a_i \otimes b_i) + (a \vdash a_i \otimes b_i) - (a_i \otimes b_i \vdash a) - (a_i \otimes a \vdash b_i)) \\
&\quad - (\text{id} \otimes \partial)((a_i \vdash a + a \vdash a_i - a_i \vdash a - a \vdash a_i) \otimes b_i + (a_i \otimes a \vdash b_i)) \\
&= (\partial \otimes \text{id})\Delta_r(a) - \tau(\partial \otimes \text{id})\delta_r(a) := \alpha_r(a).
\end{aligned}$$

Hence the conclusion holds.  $\square$

Finally, we present an example to construct Novikov bi-dialgebras by symmetric solutions of the CDNYBE, and hence Leibniz conformal bialgebras.

**Example 5.10.** Let  $A = \mathbf{k}x \oplus \mathbf{k}y$  be a vector space with a basis  $\{x, y\}$  and  $\{x^*, y^*\}$  be the basis of  $A^*$  dual to  $\{x, y\}$ . Then  $(B = A \oplus A^*, \vdash, \dashv)$  is a Novikov dialgebra with the nonzero-products

given by

$$\begin{aligned} y \vdash x &= x \vdash x = -b(x - y), & y \vdash y &= x \vdash y = -c(x - y), & x^* \vdash x &= -y^* \vdash x = b(x^* + y^*), \\ x^* \vdash y &= -y^* \vdash y = c(x^* + y^*), & x \dashv x^* &= -x \dashv y^* = (c - b)y^*, & y \dashv x^* &= -y \dashv y^* = (b - c)x^*, \\ y \vdash x &= x \vdash x^* = -y \vdash y^* = -x \vdash y^* = bx^* + cy^*, \end{aligned}$$

where  $b, c \in \mathbf{k}$ . It is straightforward to check that  $r = (y^* + x^*) \otimes (x - y) + (x - y) \otimes (y^* + x^*)$  is a symmetric solution of the CDNYBE in  $(B, \dashv, \vdash)$ . Thus by Corollary 5.7, there is a Novikov bi-dialgebra  $(B, \dashv, \vdash, \delta, \Delta)$ , where nonzero co-multiplications  $\delta$  and  $\Delta$  are given by

$$-\delta(y) = -\delta(x) = \tau\Delta(y) = \tau\Delta(x) = (c - b)(x - y) \otimes (y^* + x^*).$$

Thus one obtains a Leibniz conformal bialgebra  $(\bar{B} = \mathbf{k}[\partial]B, [\cdot, \cdot], \alpha)$  where the nonzero  $\lambda$ -bracket  $[\cdot, \cdot]_\lambda$  and co-multiplication  $\alpha$  are given by

$$\begin{aligned} [y_\lambda x] &= [x_\lambda x] = -b\lambda(x - y), & [y_\lambda y] &= [x_\lambda y] = -c\lambda(x - y), \\ [x_\lambda^* x] &= -[y_\lambda^* x] = (c - b)\partial y^* + \lambda(bx^* + cy^*), & [x_\lambda^* y] &= -[y_\lambda^* y] = (b - c)\partial x^* + \lambda(bx^* + cy^*), \\ [y_\lambda x^*] &= [x_\lambda x^*] = -[y_\lambda y^*] = -[x_\lambda y^*] = \lambda(bx^* + cy^*), \\ \alpha(y) &= \alpha(x) = (c - b)(\partial(y^* + x^*) \otimes (x - y) + (y^* + x^*) \otimes \partial(x - y)). \end{aligned}$$

**Acknowledgments.** This research is supported by NSFC (12171129, 12271265, 12261131498, W2412041), the Zhejiang Provincial Natural Science Foundation of China (No. Z25A010006) and the Fundamental Research Funds for the Central Universities and Nankai Zhide Foundation.

**Declaration of interests.** The authors have no conflicts of interest to disclose.

**Data availability.** No new data were created or analyzed in this study.

## REFERENCES

- [1] B. Bakalov, V. Kac. Field algebras. *Int. Math. Res. Not.* **3** (2003), 123–159. [2](#), [4](#)
- [2] B. Bakalov, V. Kac, A. Voronov. Cohomology of conformal algebras. *Comm. Math. Phys.* **200** (1999), 561–598. [2](#)
- [3] A. Barakat, A. De Sole, V. Kac. Poisson vertex algebras in the theory of Hamiltonian equations. *Jpn. J. Math.* **4** (2009), 141–252. [2](#)
- [4] B. Bakalov, A. D’Andrea, V. Kac. Theory of finite pseudoalgebras. *Adv. Math.* **162** (2001), 1–140. [2](#)
- [5] E. Barreiro, S. Benayadi. A new approach to Leibniz bialgebras. *Algebr. Represent. Theory.* **19** (2016), 71–101. [2](#)
- [6] A. Blokh. On a generalization of the concept of Lie algebra. *Dokl. Akad. Nauk SSSR.* **165** (1965), 471–473. [2](#)
- [7] C. Boyallian, V. Kac, J. Liberati. On the classification of subalgebras of  $Cend_N$  and  $gc_N$ . *J. Algebra.* **260** (2003), 32–63. [20](#)
- [8] V. Chari, A. Pressley. A guide to quantum groups. Cambridge Univ. Press, Cambridge. (1994). [2](#)
- [9] S. Cheng, V. Kac. Conformal modules. *Asian J. Math.* **1** (1997), 181–193. [2](#)
- [10] S. Cheng, V. Kac, M. Wakimoto. Extensions of conformal modules, in *Topological field theory, primitive forms and related topics (Kyoto, 1996)*, 79–129, *Progr. Math.*, 160, Birkhäuser Boston, Boston, MA. [2](#)
- [11] A. D’Andrea, V. Kac. Structure theory of finite conformal algebras. *Selecta Math. (N.S.)* **4** (1998), 377–418. [2](#)
- [12] A. Das, A. Sahoo. Homotopification and categorification of Leibniz conformal algebras. *Comm. Algebra.* (2025), Doi: 10.1080/00927872.2025.2472255. [2](#)
- [13] A. De Sole, V. Kac. Lie conformal algebra cohomology and the variational complex. *Comm. Math. Phys.* **292** (2009), 667–719. [2](#)

- [14] V. Drinfeld. Hamiltonian structures on Lie groups, Lie bialgebras and the geometric meaning of classical Yang-Baxter equations. Dokl. Akad. Nauk SSSR. **268** (1983), 285–287. [2](#)
- [15] T. Feng, L. Chen. Cohomology and deformations of  $\mathcal{O}$ -operators on Leibniz conformal algebras. (2023), in press. [17](#)
- [16] S. Guo, S. Wang. Twisted relative Rota-Baxter operators on Leibniz conformal algebras. Comm. Algebra. **52** (2024), 3946–3959. [2](#), [22](#)
- [17] Y. Hong, C. Bai. On antisymmetric infinitesimal conformal bialgebras. J. Algebra. **586** (2021), 325–356. [2](#), [18](#)
- [18] Y. Hong, C. Bai. Conformal classical Yang-Baxter equation,  $S$ -equation and  $\mathcal{O}$ -operators. Lett. Math. Phys. **110** (2020), 885–909. [2](#)
- [19] Y. Hong, C. Bai, L. Guo. A bialgebra theory of Gel'fand-Dorfman algebras with applications to Lie conformal bialgebras. ArXiv: 2401.13608. [3](#)
- [20] Y. Hong, F. Li. On left-symmetric conformal bialgebras. J. Algebra Appl. **14** (2015), 1450079, 37 pp. [2](#)
- [21] Y. Hong, L. Yuan. Unified products of Leibniz conformal algebras. Comm. Algebra. **49** (2021), 2074–2090. [2](#), [5](#), [8](#)
- [22] V. Kac. Vertex algebras for beginners. University Lecture Series, 10, Amer. Math. Soc., Providence, RI, (1997). [2](#), [4](#), [6](#)
- [23] V. Kac. Formal distribution algebras and conformal algebras, in: XIIIth International Congress of Mathematical Physics (ICMP '97) (Brisbane), 80–97, Int. Press, Cambridge, MA. [2](#)
- [24] V. Kac. The idea of locality, in: H.-D. Doebner, et al. (Eds.), Physical Applications and Mathematical Aspects of Geometry, Groups and Algebras, World Scientific Publishing, Singapore, (1997), pp. 16–32. [2](#)
- [25] P. Kolesnikov, B. Sartayev. On the embedding of left-symmetric algebras into differential Perm-algebras. Comm. Algebra. **50** (2022), 3246–3260. [3](#), [5](#)
- [26] J. Liberati. On conformal bialgebras. J. Algebra. **319** (2008), 2295–2318. [2](#), [10](#)
- [27] J. Loday. Une version non commutative des algèbres de Lie: les algèbres de Leibniz, in R.C.P. 25, **44** (French) (Strasbourg, 1992), 127–151. [2](#), [4](#)
- [28] J. Loday, T. Pirashvili. Universal enveloping algebras of Leibniz algebras and (co)homology. Math. Ann. **296** (1993), no. 1, 139–158. [2](#)
- [29] A. Rezaei-Aghdam, L. Sedghi-Ghadim, G. Haghighatdoost. Leibniz bialgebras, classical Yang-Baxter equations and dynamical systems. Adv. Appl. Clifford Algebr. **31** (2021), Paper No. 77, 20 pp. [2](#)
- [30] R. Tang, Y. Sheng. Leibniz bialgebras,  $\mathcal{O}$ -operators, and the classical Leibniz Yang-Baxter equation. J. Non-commut. Geom. **16** (2022), 1179–1211. [2](#), [11](#)
- [31] Z. Wu. Leibniz H-pseudoalgebras. J. Algebra. **437** (2015), 1–33. [2](#)
- [32] Z. Wu. Leibniz conformal algebras of rank three, in: Ring theory 2019, 214–231, World Sci. Publ., Hackensack, NJ. [2](#)
- [33] Z. Wu. Leibniz conformal algebras of rank two. Acta Math. Sin. (Engl. Ser.) **36** (2020), 109–120. [2](#)
- [34] X. Xu. Quadratic conformal superalgebras. J. Algebra, **231** (2000), 1–38. [3](#)
- [35] J. Zhang. On the cohomology of Leibniz conformal algebras. J. Math. Phys. **56** (2015), 041703, 14 pp. [2](#), [6](#)
- [36] J. Zhou, Y. Hong. Quadratic Leibniz conformal algebras. J. Algebra Appl. **18** (2019), 1950195, 22 pp. [2](#), [3](#), [5](#), [6](#)

CHERN INSTITUTE OF MATHEMATICS & LPMC, NANKAI UNIVERSITY, TIANJIN, 300071, CHINA  
 Email address: zy\_xu@mail.nankai.edu.cn

CHERN INSTITUTE OF MATHEMATICS & LPMC, NANKAI UNIVERSITY, TIANJIN 300071, PR CHINA  
 Email address: baicm@nankai.edu.cn

SCHOOL OF MATHEMATICS, HANGZHOU NORMAL UNIVERSITY, HANGZHOU, 311121, CHINA  
 Email address: yyhong@hznu.edu.cn