

A SIMPLE WEAK GALERKIN FINITE ELEMENT METHOD FOR A CLASS OF FOURTH-ORDER PROBLEMS IN FLUORESCENCE TOMOGRAPHY

CHUNMEI WANG * AND SHANGYOU ZHANG[†]

Abstract. In this paper, we propose a simple numerical algorithm based on the weak Galerkin (WG) finite element method for a class of fourth-order problems in fluorescence tomography (FT), eliminating the need for stabilizer terms required in traditional WG methods. FT is an emerging, non-invasive 3D imaging technique that reconstructs images of fluorophore-tagged molecule distributions in vivo. By leveraging bubble functions as a key analytical tool, our method extends to both convex and non-convex elements in finite element partitions, representing a significant advancement over existing stabilizer-free WG methods. It overcomes the restrictive conditions of previous approaches, offering substantial advantages. The proposed method preserves a simple, symmetric, and positive definite structure. These advantages are confirmed by optimal-order error estimates in a discrete H^2 norm, demonstrating the effectiveness and accuracy of our approach. Numerical experiments further validate the efficiency and precision of the proposed method.

Key words. weak Galerkin, stabilizer-free, bubble functions, non convex polygonal or polyhedral meshes, fluorescence tomography.

AMS subject classifications. Primary, 65N30, 65N15, 65N12, 74N20; Secondary, 35B45, 35J50, 35J35

1. Introduction. This paper focuses on the development of numerical methods for a class of fourth-order problems with Dirichlet and Neumann boundary conditions. The model problem seeks an unknown function $u = u(x)$ satisfying

$$(1.1) \quad \begin{aligned} (-\nabla \cdot (\kappa \nabla) + \mu)^2 u &= f, & \text{in } \Omega, \\ u &= \xi, & \text{on } \partial\Omega, \\ \kappa \nabla u \cdot \mathbf{n} &= \nu, & \text{on } \partial\Omega, \end{aligned}$$

where Ω is an open, bounded domain in \mathbb{R}^d ($d = 2, 3$) with a Lipschitz continuous boundary $\partial\Omega$. Here, \mathbf{n} denotes the unit outward normal to $\partial\Omega$, κ is a symmetric and positive definite matrix-valued function, and μ is a nonnegative real-valued function. The functions f , ξ , and ν are given in the domain or on its boundary, as appropriate.

For convenience, we denote the second-order elliptic operator $\nabla \cdot (\kappa \nabla)$ by E . To simplify the analysis, and without loss of generality, we assume that κ is piecewise constant matrix and μ is a piecewise constant function.

The fourth order model problem (1.1) is derived from fluorescence tomography (FT) [7, 9, 8, 12, 25, 33, 34, 54], an advanced noninvasive 3D imaging technique used for in vivo applications. Unlike traditional imaging methods that rely on X-rays or strong magnetic fields, FT utilizes highly specific fluorescent probes and non-ionizing near-infrared radiation [10], reducing potential health risks. The primary objective

*Department of Mathematics, University of Florida, Gainesville, FL 32611, USA (chunmei.wang@ufl.edu). The research of Chunmei Wang was partially supported by National Science Foundation Grant DMS-2136380.

[†]Department of Mathematical Sciences, University of Delaware, Newark, DE 19716, USA (szhang@udel.edu).

of FT is to reconstruct the spatial distribution of fluorophores, which are bound to target molecules, based on boundary measurements. Given its capability to provide molecular-level imaging, FT has emerged as a valuable tool for early cancer detection and drug monitoring [1, 11, 35].

We define the function space

$$H_{\kappa}^2(\Omega) = \{v : v \in H^1(\Omega), \kappa \nabla v \in H(\text{div}; \Omega)\},$$

equipped with the norm

$$\|v\|_{\kappa,2} = (\|v\|_1^2 + \|\nabla \cdot (\kappa \nabla v)\|^2)^{\frac{1}{2}}.$$

A variational formulation for the fourth-order model problem (1.1) seeks a function $u \in H_{\kappa}^2(\Omega)$ satisfying the boundary conditions $u|_{\partial\Omega} = \xi$, $\kappa \nabla u \cdot \mathbf{n}|_{\partial\Omega} = \nu$, such that

$$(1.2) \quad (Eu, Ev) + 2\mu(\kappa \nabla u, \nabla v) + \mu^2(u, v) = (f, v), \quad \forall v \in \mathcal{V},$$

where (\cdot, \cdot) denotes the standard inner product in $L^2(\Omega)$. The test space \mathcal{V} is given by

$$\mathcal{V} = \{v \in H_{\kappa}^2(\Omega) : v|_{\partial\Omega} = 0, \kappa \nabla v \cdot \mathbf{n}|_{\partial\Omega} = 0\}.$$

Here, \mathbf{n} represents the outward unit normal to the boundary of Ω .

The weak Galerkin (WG) finite element method represents a significant advancement in numerical techniques for solving partial differential equations (PDEs). It reconstructs or approximates differential operators in a manner analogous to the theory of distributions for piecewise polynomials. Unlike traditional finite element methods, WG relaxes the usual regularity requirements on approximating functions by incorporating carefully designed stabilizers. Extensive research has demonstrated the effectiveness of WG across a variety of model PDEs, as evidenced by a comprehensive set of references [14, 15, 46, 50, 16, 17, 18, 19, 48, 51, 5, 45, 23, 13, 29, 53, 40, 44, 41, 42, 43, 47, 49], underscoring its potential as a powerful tool in scientific computing.

A key distinguishing feature of WG methods is their reliance on weak derivatives and weak continuities to construct numerical schemes based on the weak formulation of PDEs. This structural flexibility enhances their applicability across a broad spectrum of PDEs, ensuring both stability and accuracy in numerical approximations.

An important advancement within the WG framework is the Primal-Dual Weak Galerkin (PDWG) method, which addresses challenges that conventional numerical techniques often encounter [20, 21, 2, 3, 4, 22, 30, 31, 52, 6, 36, 37, 32, 38, 39]. PDWG formulates numerical solutions as constrained minimizations of functionals, where the constraints capture the weak formulation of PDEs through weak derivatives. This leads to an Euler-Lagrange equation that integrates both the primal variable and a dual variable (Lagrange multiplier), resulting in a symmetric numerical scheme.

The variational formulation in (1.2) differs significantly from the standard biharmonic equation. First, conventional H^2 -conforming finite elements designed for the biharmonic problem are not necessarily H_{κ}^2 -conforming and therefore cannot be directly applied to (1.2). Additionally, widely used nonconforming elements, such as the Morley element [26], typically rely on formulations involving the full Hessian. Since it

is unclear whether (1.2) can be reformulated to align with such elements, their direct applicability remains uncertain. In fact, we suspect that these elements may not be suitable for this problem.

This paper introduces a simplified weak Galerkin (WG) finite element formulation that eliminates the need for stabilizers required in [53], significantly streamlining both the numerical scheme and its implementation. Furthermore, our approach extends the applicability of WG methods to both convex and non-convex polytopal meshes, whereas existing methods have been limited to convex cases. A key analytical tool enabling these advancements is the use of bubble functions. Our method preserves the size and global sparsity of the stiffness matrix, reducing programming complexity compared to traditional stabilizer-dependent WG methods. Theoretical analysis establishes that our WG approximations achieve optimal error estimates in the discrete H^2 norm. By introducing a stabilizer-free WG method that maintains accuracy while improving computational efficiency, this paper makes a significant contribution to the development of finite element methods for a class of fourth-order problems on non-convex polytopal meshes.

The paper is structured as follows. Section 2 provides a concise review of weak differential operators and their discrete counterparts. In Section 3, we introduce a simple weak Galerkin algorithm for the fourth-order model problem (1.1), formulated based on the variational approach in (1.2). Section 4 presents local L^2 projection operators and establishes key approximation properties essential for the convergence analysis. Section 5 is dedicated to deriving the error equation for the WG finite element solution. In Section 6, we prove an optimal-order error estimate for the WG finite element approximation in an H^2 -equivalent discrete norm. Finally, Section 7 provides numerical results that validate the theoretical findings presented in the preceding sections.

2. Weak Differential Operators and Their Discrete Counterparts. In this section, we briefly review the definitions of the weak second-order elliptic operator E [53] and the weak gradient operator [49], along with their discrete formulations.

Let \mathcal{T}_h be a partition of the domain Ω into polygons in 2D or polyhedra in 3D. We assume that \mathcal{T}_h satisfies the shape regularity condition as defined in [49]. Denote by \mathcal{E}_h the set of all edges (in 2D) or flat faces (in 3D) within \mathcal{T}_h , and let $\mathcal{E}_h^0 = \mathcal{E}_h \setminus \partial\Omega$ represent the set of all interior edges or faces. A weak function on an element $T \in \mathcal{T}_h$ is defined as a triplet $v = \{v_0, v_b, v_g\}$, where $v_0 \in L^2(T)$ represents the function's value in the interior of T , $v_b \in L^2(\partial T)$ represents the function's value on the boundary of T , and $v_g \in L^2(\partial T)$ approximates the normal flux $\kappa \nabla v \cdot \mathbf{n}$ on ∂T , with \mathbf{n} denoting the outward unit normal. On each interior edge or face $e \in \mathcal{E}_h^0$, shared by two elements T_L and T_R , the function v_g has two values: v_g^L , as seen from element T_L , and v_g^R , as seen from element T_R which satisfy the condition $v_g^L + v_g^R = 0$.

The space of all weak functions on T is given by

$$W(T) = \{v = \{v_0, v_b, v_g\} : v_0 \in L^2(T), v_b \in L^2(\partial T), v_g \in L^2(\partial T)\}.$$

DEFINITION 2.1. [53] *For any $v \in W(T)$, the weak second-order elliptic operator, denoted as $E_w v$, is defined as a linear functional in the dual space of $H^2(T)$. Its action on each $\varphi \in H^2(T)$ is given by*

$$(2.1) \quad (E_w v, \varphi)_T = (v_0, E\varphi)_T - \langle v_b, \kappa \nabla \varphi \cdot \mathbf{n} \rangle_{\partial T} + \langle v_g, \varphi \rangle_{\partial T}.$$

For any non-negative integer $r \geq 0$, let $P_r(T)$ denote the space of polynomials on T with a degree at most r . The discrete weak second-order elliptic operator, denoted by $E_{w,r,T}$, is defined as the unique polynomial $E_{w,r,T}v \in P_r(T)$ satisfying

$$(2.2) \quad (E_{w,r,T}v, \varphi)_T = (v_0, E\varphi)_T - \langle v_b, \kappa \nabla \varphi \cdot \mathbf{n} \rangle_{\partial T} + \langle v_g, \varphi \rangle_{\partial T}, \quad \forall \varphi \in P_r(T).$$

Using integration by parts, this can be rewritten as

$$(2.3) \quad (E_{w,r,T}v, \varphi)_T = (Ev_0, \varphi)_T + \langle v_0 - v_b, \kappa \nabla \varphi \cdot \mathbf{n} \rangle_{\partial T} - \langle \kappa \nabla v_0 \cdot \mathbf{n} - v_g, \varphi \rangle_{\partial T}, \quad \forall \varphi \in P_r(T).$$

DEFINITION 2.2. [49] *The weak gradient operator for any $v \in W(T)$, denoted as $\nabla_w v$, is defined as a linear vector functional in the dual space of $[H^1(T)]^d$. Its action on each $\boldsymbol{\psi} \in [H^1(T)]^d$ is given by*

$$(2.4) \quad (\nabla_w v, \boldsymbol{\psi})_T = -(v_0, \nabla \cdot \boldsymbol{\psi})_T + \langle v_b, \boldsymbol{\psi} \cdot \mathbf{n} \rangle_{\partial T}.$$

The discrete weak gradient operator, denoted by $\nabla_{w,r,T}$, is defined as the unique vector polynomial $\nabla_{w,r,T}v \in [P_r(T)]^d$ satisfying

$$(2.5) \quad (\nabla_{w,r,T}v, \boldsymbol{\psi})_T = -(v_0, \nabla \cdot \boldsymbol{\psi})_T + \langle v_b, \boldsymbol{\psi} \cdot \mathbf{n} \rangle_{\partial T}, \quad \forall \boldsymbol{\psi} \in [P_r(T)]^d.$$

Using integration by parts, this can be reformulated as

$$(2.6) \quad (\nabla_{w,r,T}v, \boldsymbol{\psi})_T = (\nabla v_0, \boldsymbol{\psi})_T - \langle v_0 - v_b, \boldsymbol{\psi} \cdot \mathbf{n} \rangle_{\partial T}, \quad \forall \boldsymbol{\psi} \in [P_r(T)]^d.$$

3. Stabilizer-Free Weak Galerkin Methods. For any given integer $k \geq 1$, let $W_k(T)$ denote the local discrete weak function space defined as

$$(3.1) \quad W_k(T) = \{v = \{v_0, v_b, v_g\} : v_0 \in P_k(T), v_b \in P_k(e), v_g \in P_{k-1}(e), e \subset \partial T\}.$$

By assembling $W_k(T)$ over all the elements $T \in \mathcal{T}_h$ and enforcing continuity across the interior edges \mathcal{E}_h^0 , we define the global weak finite element space V_h as

$$V_h = \{v = \{v_0, v_b, v_g\} : \{v_0, v_b, v_g\}|_T \in W_k(T), \forall T \in \mathcal{T}_h\}.$$

Furthermore, we define the subspace V_h^0 consisting of functions with vanishing traces,

$$V_h^0 = \{v = \{v_0, v_b, v_g\} \in V_h, v_b|_e = 0, v_g|_e = 0, e \subset \partial T \cap \partial \Omega\}.$$

Denote by E_{w,r_1} and ∇_{w,r_2} the discrete weak second-order elliptic operator and the discrete weak gradient operator, respectively, which are computed on each element T for $k \geq 1$ using equations (2.2) and (2.5). Specifically, for $v \in V_h$, we have

$$(E_{w,r_1}v)|_T = E_{w,r_1,T}(v|_T), \quad (\nabla_{w,r_2}v)|_T = \nabla_{w,r_2,T}(v|_T).$$

For simplicity, we omit the subscripts r_1 and r_2 in E_{w,r_1} and ∇_{w,r_2} .

For each element T , let Q_0 be the L^2 projection onto $P_k(T)$, and for each edge or face $e \subset \partial T$, let Q_b and Q_g be the L^2 projections onto $P_k(e)$ and $P_{k-1}(e)$, respectively. Given $u \in H^2(\Omega)$, we define a projection onto the weak finite element space V_h by

$$Q_h u = \{Q_0 u, Q_b u, Q_g(\kappa \nabla u \cdot \mathbf{n})\}.$$

We now present the stabilizer-free weak Galerkin finite element scheme for the fourth-order model problem (1.1), based on the variational formulation (1.2).

WEAK GALERKIN ALGORITHM 1. Find $u_h = \{u_0, u_b, u_g\} \in V_h$ satisfying the boundary conditions $u_b = Q_b \xi$ and $u_g = Q_g \nu$ on $\partial\Omega$, such that

$$(3.2) \quad \sum_{T \in \mathcal{T}_h} (E_w u_h, E_w v)_T + 2\mu(\kappa \nabla_w u_h, \nabla_w v)_T + \mu^2(u_0, v_0)_T = \sum_{T \in \mathcal{T}_h} (f, v_0)_T, \forall v \in V_h^0.$$

For any $v \in V_h$, define an energy norm $\|v\|$ by

$$(3.3) \quad \|v\|^2 = \sum_{T \in \mathcal{T}_h} (E_w v, E_w v)_T + 2\mu(\kappa \nabla_w v, \nabla_w v)_T + \mu^2(v_0, v_0)_T.$$

Additionally, we define the discrete H^2 semi-norm as

$$(3.4) \quad \begin{aligned} \|v\|_{2,h}^2 &= \sum_{T \in \mathcal{T}_h} (E v_0, E v_0)_T + 2\mu(\kappa \nabla v_0, \nabla v_0)_T + \mu^2(v_0, v_0)_T \\ &+ h_T^{-1} \langle \kappa \nabla u_0 \cdot \mathbf{n} - u_g, \kappa \nabla v_0 \cdot \mathbf{n} - v_g \rangle_{\partial T} + h_T^{-3} \langle u_0 - u_b, v_0 - v_b \rangle_{\partial T}. \end{aligned}$$

For any element $T \in \mathcal{T}_h$ and any function $\varphi \in H^1(T)$, the trace inequality (see [49]) asserts that

$$(3.5) \quad \|\varphi\|_{\partial T}^2 \leq C(h_T^{-1} \|\varphi\|_T^2 + h_T \|\nabla \varphi\|_T^2).$$

Moreover, if φ is a polynomial on T , then we have from the inverse inequality (see also [49]) that

$$(3.6) \quad \|\varphi\|_e^2 \leq C h_T^{-1} \|\varphi\|_T^2.$$

Here e is an edge or flat face on the boundary of T .

LEMMA 3.1. For $v = \{v_0, v_b, v_g\} \in V_h$, there exists a constant C such that

$$\|E v_0\|_T \leq C \|E_w v\|_T.$$

Proof. Let $T \in \mathcal{T}_h$ be a polytopal element with N edges or faces, denoted by e_1, \dots, e_N . Note that T may be non-convex. For each edge/face e_i , we define a linear function $l_i(x)$ satisfying $l_i(x) = 0$ on e_i as follows:

$$l_i(x) = \frac{1}{h_T} \overrightarrow{AX} \cdot \mathbf{n}_i,$$

where A is a given point on e_i , X is any point on e_i , \mathbf{n}_i is the unit normal to e_i , and h_T represents the size of T .

The bubble function associated with T is defined as

$$\Phi_B = l_1^2(x) l_2^2(x) \cdots l_N^2(x) \in P_{2N}(T).$$

It is straightforward to verify that $\Phi_B = 0$ on ∂T . Moreover, Φ_B can be normalized so that $\Phi_B(M) = 1$ where M denotes the barycenter of T . Additionally, there exists a sub-domain $\hat{T} \subset T$ where $\Phi_B \geq \rho_0$ for some constant $\rho_0 > 0$.

For $v = \{v_0, v_b, v_g\} \in V_h$, setting $r = 2N + k - 2$ and choosing $\varphi = \Phi_B E v_0 \in P_r(T)$ in (2.3) leads to

$$(3.7) \quad \begin{aligned} & (E_w v, \Phi_B E v_0)_T \\ &= (E v_0, \Phi_B E v_0)_T + \langle v_0 - v_b, \kappa \nabla(\Phi_B E v_0) \cdot \mathbf{n} \rangle_{\partial T} - \langle \kappa \nabla v_0 \cdot \mathbf{n} - v_g, \Phi_B E v_0 \rangle_{\partial T} \\ &= (E v_0, \Phi_B E v_0)_T, \end{aligned}$$

where the boundary condition $\Phi_B = 0$ on ∂T is used.

Using the domain inverse inequality [49], there exists a constant C such that

$$(3.8) \quad (E v_0, \Phi_B E v_0)_T \geq C (E v_0, E v_0)_T.$$

Applying the Cauchy–Schwarz inequality together with (3.7)–(3.8), we obtain

$$(E v_0, E v_0)_T \leq C (E_w v, \Phi_B E v_0)_T \leq C \|E_w v\|_T \|\Phi_B E v_0\|_T \leq C \|E_w v\|_T \|E v_0\|_T.$$

Thus, we conclude

$$\|E v_0\|_T \leq C \|E_w v\|_T.$$

This completes the proof. \square

REMARK 3.1. *When the polytopal element T is convex, the bubble function introduced in Lemma 3.1 can be simplified to*

$$\Phi_B = l_1(x) l_2(x) \cdots l_N(x).$$

This function satisfies the following properties: (1) $\Phi_B = 0$ on ∂T , (2) there exists a subregion $\hat{T} \subset T$ where $\Phi_B \geq \rho_0$ for some constant $\rho_0 > 0$.

Using this simplified bubble function, Lemma 3.1 can be established through the same reasoning as before, with the parameter choice $r = N + k - 2$.

LEMMA 3.2. [27] *For any function $v = \{v_0, v_b, v_g\} \in V_h$, there exists a constant C such that*

$$\|\nabla v_0\|_T \leq C \|\nabla_w v\|_T.$$

To define an edge/face-based bubble function, we introduce

$$\varphi_{e_k} = \prod_{i=1, \dots, N, i \neq k} l_i^2(x).$$

This function satisfies: (1) $\varphi_{e_k} = 0$ on every edge/face e_i for $i \neq k$, (2) there exists a subset $\hat{e}_k \subset e_k$ such that $\varphi_{e_k} \geq \rho_1$ for some constant $\rho_1 > 0$. Now, let

$$\varphi = (v_b - v_0) l_k \varphi_{e_k}.$$

Then, it follows that $\varphi = 0$ on all edges/faces e_i for $i = 1, \dots, N$, and $\nabla \varphi = 0$ on e_i for $i \neq k$. Moreover, the gradient takes the form $\nabla \varphi = (v_0 - v_b)(\nabla l_k) \varphi_{e_k} = \mathcal{O}(\frac{(v_0 - v_b) \varphi_{e_k}}{h_T} \mathbf{C})$ on e_k for some vector constant \mathbf{C} .

LEMMA 3.3. *For $\{v_0, v_b, v_g\} \in V_h$, let $\varphi = (v_b - v_0) l_k \varphi_{e_k}$. Then, the inequality*

$$(3.9) \quad \|\varphi\|_T^2 \leq C h_T \int_{e_k} (v_b - v_0)^2 ds.$$

holds.

Proof. To extend v_b , originally defined on the $(d-1)$ -dimensional edge/face e_k , to the entire d -dimensional element T , we use the extension

$$v_b(X) = v_b(Proj_{e_k}(X)),$$

where X is an arbitrary point in T , $Proj_{e_k}(X)$ represents the orthogonal projection of X onto the hyperplane $H \subset \mathbb{R}^d$ containing e_k . If $Proj_{e_k}(X)$ lies outside e_k , then $v_b(Proj_{e_k}(X))$ is defined as the natural extension of v_b from e_k to H .

We assert that v_b remains a polynomial defined on the element T following the extension.

Consider the hyperplane H that contains the edge/face e_k , which is determined by $d-1$ linearly independent vectors $\beta_1, \dots, \beta_{d-1}$ originating from a point A on e_k . Any point P on e_k can be parametrized as

$$P(t_1, \dots, t_{d-1}) = A + t_1\beta_1 + \dots + t_{d-1}\beta_{d-1},$$

where t_1, \dots, t_{d-1} are parameters.

Since $v_b(P(t_1, \dots, t_{d-1}))$ is a polynomial of degree p defined on e_k , it can be expressed as

$$v_b(P(t_1, \dots, t_{d-1})) = \sum_{|\alpha| \leq p} c_\alpha \mathbf{t}^\alpha,$$

where $\mathbf{t}^\alpha = t_1^{\alpha_1} \dots t_{d-1}^{\alpha_{d-1}}$ and $\alpha = (\alpha_1, \dots, \alpha_{d-1})$ is a multi-index.

For any point X in the element T , the projection of X onto the hyperplane H is the point in H that minimizes the distance to X . This projection, denoted as $Proj_{e_k}(X)$, is an affine transformation given by

$$Proj_{e_k}(X) = A + \sum_{i=1}^{d-1} t_i(X)\beta_i,$$

where A is the origin point on e_k , and $t_i(X)$ are the projection coefficients determined by solving the orthogonality condition

$$(X - Proj_{e_k}(X)) \cdot \beta_j = 0, \quad \forall j = 1, \dots, d-1.$$

This yields a system of linear equations in $t_1(X), \dots, t_{d-1}(X)$, which can be explicitly solved, ensuring that

$$t_i(X) \text{ is a linear function of } X.$$

Consequently, the projection $Proj_{e_k}(X)$ is an affine linear function of X .

To extend v_b from e_k to the entire element T , we define

$$v_b(X) = v_b(Proj_{e_k}(X)) = \sum_{|\alpha| \leq p} c_\alpha \mathbf{t}(X)^\alpha,$$

where $\mathbf{t}(X)^\alpha = t_1(X)^{\alpha_1} \dots t_{d-1}(X)^{\alpha_{d-1}}$. Since each $t_i(X)$ is a linear function of X , it follows that $\mathbf{t}(X)^\alpha$ is a polynomial in $X = (x_1, \dots, x_d)$, thereby confirming that $v_b(X)$ remains a polynomial in d -dimensional coordinates.

Similarly, let v_{trace} denote the trace of v_0 on e_k . We extend v_{trace} to T using the formula

$$v_{trace}(X) = v_{trace}(Proj_{e_k}(X)),$$

For any X in T , if $Proj_{e_k}(X)$ does not lie on e_k , then $v_{trace}(Proj_{e_k}(X))$ is defined as the extension of v_{trace} from e_k to the hyperplane H . By a reasoning analogous to that for v_b , it follows that v_{trace} remains a polynomial after this extension.

Define $\varphi = (v_b - v_0)l_k\varphi_{e_k}$. Then,

$$\begin{aligned} \|\varphi\|_T^2 &= \int_T \varphi^2 dT \leq Ch_T^2 \int_T (\nabla\varphi)^2 dT \\ &\leq Ch_T^2 \int_T (\nabla((v_b - v_{trace})(X)l_k\varphi_{e_k}))^2 dT \\ &\leq Ch_T^3 \int_{e_k} ((v_b - v_{trace})(Proj_{e_k}(X))(\nabla l_k)\varphi_{e_k})^2 ds \\ &\leq Ch_T \int_{e_k} (v_b - v_0)^2 ds, \end{aligned}$$

where we applied Poincare inequality since $\varphi = 0$ on each e_i for $i = 1, \dots, N$, $\nabla\varphi = 0$ on each e_i for $i \neq k$, $\nabla\varphi = (v_0 - v_b)(\nabla l_k)\varphi_{e_k} = \mathcal{O}(\frac{(v_0 - v_b)\varphi_{e_k}}{h_T}\mathbf{C})$ on e_k for some vector constant \mathbf{C} , along with the properties of the projection.

This completes the proof of the lemma.

□

LEMMA 3.4. For any $\{v_0, v_b, v_g\} \in V_h$, define $\varphi = (\kappa\nabla v_0 \cdot \mathbf{n} - v_g)\varphi_{e_k}$. The following inequality holds:

$$(3.10) \quad \|\varphi\|_T^2 \leq Ch_T \int_{e_k} (\kappa\nabla v_0 \cdot \mathbf{n} - v_g)^2 ds.$$

Proof. We begin by extending v_g , originally defined on the $(d-1)$ -dimensional edge/face e_k , to the entire d -dimensional polytopal element T using the formula:

$$v_g(X) = v_g(Proj_{e_k}(X)),$$

where X is any point in T , and $Proj_{e_k}(X)$ denotes the orthogonal projection of the point X onto the hyperplane H containing e_k . If $Proj_{e_k}(X)$ does not lie on e_k , we define $v_g(Proj_{e_k}(X))$ as the extension of v_g from e_k to H .

We assert that after this extension, v_g remains a polynomial on T , which follows from the same reasoning as in Lemma 3.3.

Next, let v_{trace} be the trace of v_0 on e_k . We extend v_{trace} to T using the formula:

$$v_{trace}(X) = v_{trace}(Proj_{e_k}(X)).$$

Again, if $Proj_{e_k}(X)$ is not on e_k , we define $v_{trace}(Proj_{e_k}(X))$ as the extension of v_{trace} from e_k to H . It follows from Lemma 3.3 that v_{trace} remains a polynomial after this extension.

Now, setting $\varphi = (\kappa \nabla v_0 \cdot \mathbf{n} - v_g) \varphi_{e_k}$, we obtain:

$$\begin{aligned} \|\varphi\|_T^2 &= \int_T \varphi^2 dT = \int_T ((\kappa \nabla v_0 \cdot \mathbf{n} - v_g)(X) \varphi_{e_k})^2 dT \\ &\leq Ch_T \int_{e_k} ((\kappa \nabla v_{trace} \cdot \mathbf{n} - v_g)(Proj_{e_k}(X)) \varphi_{e_k})^2 dT \\ &\leq Ch_T \int_{e_k} (\kappa \nabla v_0 \cdot \mathbf{n} - v_g)^2 ds, \end{aligned}$$

where we used the facts that (1) $\varphi_{e_k} = 0$ on each edge/face e_i for $i \neq k$, (2) there exists a subdomain $\hat{e}_k \subset e_k$ where $\varphi_{e_k} \geq \rho_1$ for some constant $\rho_1 > 0$, and applied the properties of the projection.

This completes the proof. \square

LEMMA 3.5. *There exist two positive constants, C_1 and C_2 , such that for every $v = \{v_0, v_b, v_g\} \in V_h$, the following norm equivalence holds:*

$$(3.11) \quad C_1 \|v\|_{2,h} \leq \|v\| \leq C_2 \|v\|_{2,h}.$$

Proof. We begin by recalling that the edge/face-based bubble function is given by

$$\varphi_{e_k} = \prod_{i=1, \dots, N, i \neq k} l_i^2(x).$$

First, we extend the function v_b , originally defined on the edge/face e_k , to the entire element T . Similarly, let v_{trace} denote the trace of v_0 on e_k ; we extend v_{trace} to T as well. For notational convenience, these extensions are still denoted by v_b and v_0 . (Details of these extensions appear in Lemma 3.3.) By substituting

$$\varphi = (v_b - v_0) l_k \varphi_{e_k}$$

into equation (2.3), we obtain

$$\begin{aligned} &(E_w v, \varphi)_T \\ &= (E v_0, \varphi)_T + \langle v_0 - v_b, \kappa \nabla \varphi \cdot \mathbf{n} \rangle_{\partial T} - \langle \kappa \nabla v_0 \cdot \mathbf{n} - v_g, \varphi \rangle_{\partial T} \\ (3.12) \quad &= (E v_0, \varphi)_T + \int_{e_k} |v_b - v_0|^2 \kappa (\nabla l_k) \varphi_{e_k} \cdot \mathbf{n} ds \\ &= (E v_0, \varphi)_T + Ch_T^{-1} \int_{e_k} |v_b - v_0|^2 \kappa \varphi_{e_k} ds, \end{aligned}$$

where we have used that $\varphi = 0$ on each edge/face e_i for $i = 1, \dots, N$, that $\nabla \varphi = 0$ on each edge/face e_i with $i \neq k$ and that on e_k , the gradient satisfies

$$\nabla \varphi = (v_0 - v_b) (\nabla l_k) \varphi_{e_k} = \mathcal{O}\left(\frac{(v_0 - v_b) \varphi_{e_k}}{h_T} \mathbf{C}\right)$$

with \mathbf{C} a constant vector.

Recall additionally that: (1) $\varphi_{e_k} = 0$ on every edge/face e_i for $i \neq k$, (2) there exists a subdomain $\hat{e}_k \subset e_k$ such that $\varphi_{e_k} \geq \rho_1$ for some constant $\rho_1 > 0$. By invoking

the Cauchy–Schwarz inequality and a domain inverse inequality (see [49]), together with (3.12) and the result of Lemma 3.3, we deduce

$$\begin{aligned} \int_{e_k} |v_b - v_0|^2 ds &\leq C \int_{e_k} |v_b - v_0|^2 \kappa \varphi_{e_k} ds \\ &\leq Ch_T (\|E_w v\|_T + \|E v_0\|_T) \|\varphi\|_T \\ &\leq Ch_T^{\frac{3}{2}} (\|E_w v\|_T + \|E v_0\|_T) \left(\int_{e_k} |v_b - v_0|^2 ds \right)^{\frac{1}{2}}, \end{aligned}$$

which, in conjunction with Lemma 3.1, implies that

$$(3.13) \quad h_T^{-3} \int_{e_k} |v_b - v_0|^2 ds \leq C (\|E_w v\|_T^2 + \|E v_0\|_T^2) \leq C \|E_w v\|_T^2.$$

Next, we extend v_g from e_k to T ; for simplicity, the extension is still denoted by v_g (see Lemma 3.4 for details). Letting $\varphi = (\kappa \nabla v_0 \cdot \mathbf{n} - v_g) \varphi_{e_k}$ in (2.3) yields

$$\begin{aligned} &(E_w v, \varphi)_T \\ &= (E v_0, \varphi)_T + \langle v_0 - v_b, \kappa \nabla \varphi \cdot \mathbf{n} \rangle_{\partial T} - \langle \kappa \nabla v_0 \cdot \mathbf{n} - v_g, \varphi \rangle_{\partial T} \\ &= (E v_0, \varphi)_T + \langle v_0 - v_b, \kappa \nabla \varphi \cdot \mathbf{n} \rangle_{\partial T} - \int_{e_k} (\kappa \nabla v_0 \cdot \mathbf{n} - v_g)^2 \varphi_{e_k} ds \end{aligned}$$

where we have used that $\varphi_{e_k} = 0$ on every edge/face e_i with $i \neq k$, and the fact that there exists a subdomain $\hat{e}_k \subset e_k$ for which $\varphi_{e_k} \geq \rho_1 > 0$. By applying the Cauchy–Schwarz inequality, the domain inverse inequality (see [49]), the inverse inequality, and the trace inequality (3.6), together with (3.13) and Lemma 3.4, we obtain

$$\begin{aligned} &\int_{e_k} (\kappa \nabla v_0 \cdot \mathbf{n} - v_g)^2 ds \\ &\leq C \int_{e_k} (\kappa \nabla v_0 \cdot \mathbf{n} - v_g)^2 \varphi_{e_k} ds \\ &\leq C (\|E_w v\|_T + \|E v_0\|_T) \|\varphi\|_T + C \|v_0 - v_b\|_{\partial T} \|\kappa \nabla \varphi \cdot \mathbf{n}\|_{\partial T} \\ &\leq Ch_T^{\frac{1}{2}} (\|E_w v\|_T + \|E v_0\|_T) \left(\int_{e_k} (\kappa \nabla v_0 \cdot \mathbf{n} - v_g)^2 ds \right)^{\frac{1}{2}} \\ &\quad + Ch_T^{\frac{3}{2}} \|E_w v\|_T h_T^{-\frac{1}{2}} h_T^{-1} h_T^{\frac{1}{2}} \left(\int_{e_k} (\kappa \nabla v_0 \cdot \mathbf{n} - v_g)^2 ds \right)^{\frac{1}{2}}. \end{aligned}$$

This, together with Lemma 3.1, gives

$$(3.14) \quad h_T^{-1} \int_{e_k} (\kappa \nabla v_0 \cdot \mathbf{n} - v_g)^2 ds \leq C (\|E_w v\|_T^2 + \|E v_0\|_T^2) \leq C \|E_w v\|_T^2.$$

By combining Lemmas 3.1–3.2 with estimates (3.13) and (3.14), and by employing the norm definitions (3.3) and (3.4), we deduce that

$$C_1 \|v\|_{2,h} \leq \|v\|.$$

For the reverse inequality, we apply Cauchy–Schwarz inequality, the inverse inequality, and the trace inequality (3.6) to (2.3) to obtain

$$\begin{aligned} |(E_w v, \varphi)_T| &\leq \|E v_0\|_T \|\varphi\|_T + \|v_b - v_0\|_{\partial T} \|\kappa \nabla \varphi \cdot \mathbf{n}\|_{\partial T} + \|\kappa \nabla v_0 \cdot \mathbf{n} - v_g\|_{\partial T} \|\varphi\|_{\partial T} \\ &\leq \|E v_0\|_T \|\varphi\|_T + h_T^{-\frac{3}{2}} \|v_b - v_0\|_{\partial T} \|\varphi\|_T + h_T^{-\frac{1}{2}} \|\kappa \nabla v_0 \cdot \mathbf{n} - v_g\|_{\partial T} \|\varphi\|_T. \end{aligned}$$

This implies that

$$\|E_w v\|_T^2 \leq C(\|Ev_0\|_T^2 + h_T^{-3}\|v_b - v_0\|_{\partial T}^2 + h_T^{-1}\|\kappa \nabla v_0 \cdot \mathbf{n} - v_g\|_{\partial T}^2),$$

which, combined with Lemma 3.2, gives

$$\|v\| \leq C_2 \|v\|_{2,h}.$$

The two inequalities together establish the desired norm equivalence, thereby completing the proof. \square

THEOREM 3.6. *The WG Algorithm 3.2 admits a unique solution.*

Proof. Suppose $u_h^{(1)} \in V_h$ and $u_h^{(2)} \in V_h$ are two distinct solutions of the WG Algorithm 3.2. Define their difference as $\eta_h = u_h^{(1)} - u_h^{(2)}$. Then, $\eta_h \in V_h^0$ satisfies

$$\sum_{T \in \mathcal{T}_h} (E_w \eta_h, E_w v)_T + 2\mu(\kappa \nabla_w \eta_h, \nabla_w v)_T + \mu^2(\eta_0, v_0)_T = 0, \quad \forall v \in V_h^0.$$

Choosing $v = \eta_h$ in the above equation yields $\|\eta_h\| = 0$. By the norm equivalence (3.11), we obtain $\|\eta_h\|_{2,h} = 0$, which implies that on each element T ,

$$E\eta_0 = 0, \quad \nabla \eta_0 = 0, \quad \eta_0 = 0,$$

along with the conditions $\eta_0 = \eta_b$ and $\kappa \nabla \eta_0 \cdot \mathbf{n} = \eta_g$ on ∂T . Consequently, η_0 must be a constant within each element T , which, together with the fact that $\eta_0 = \eta_b$ on each ∂T , indicates that η_0 is a constant across the entire domain Ω . Furthermore, since $\eta_0 = \eta_b$ on ∂T and $\eta_b|_{\partial T \cap \partial \Omega} = 0$, we conclude that $\eta_0 = 0$ in Ω . This, together with $\eta_0 = \eta_b$ on ∂T , implies $\eta_b = 0$ in Ω . Similarly, using the boundary condition $\kappa \nabla \eta_0 \cdot \mathbf{n} = \eta_g$ on each ∂T and $\eta_g|_{\partial T \cap \partial \Omega} = 0$, we deduce that $\eta_g = 0$ in Ω . Therefore, $\eta_h = 0$ in Ω , which implies $u_h^{(1)} \equiv u_h^{(2)}$.

This establishes the uniqueness of the solution, completing the proof. \square

4. Technical Results. This section aims to establish essential technical results related to L^2 projections, which play a crucial role in the error analysis of the WG method.

For each element $T \in \mathcal{T}_h$, let Q^{r_1} and Q^{r_2} be the L^2 projection operators onto the finite element spaces consisting of piecewise polynomials of degree at most r_1 and r_2 , respectively.

LEMMA 4.1. *The following properties hold:*

$$(4.1) \quad E_w v = Q^{r_1}(Ev), \quad \forall v \in H_\kappa^2(T),$$

$$(4.2) \quad \nabla_w v = Q^{r_2}(\nabla v), \quad \forall v \in H^1(T).$$

Proof. For any $u \in H_\kappa^2(T)$, applying equation (2.3) yields

$$\begin{aligned} & (E_w v, \varphi)_T \\ &= (Ev, \varphi)_T + \langle v|_T - v|_{\partial T}, \kappa \nabla \varphi \cdot \mathbf{n} \rangle_{\partial T} - \langle \kappa \nabla v|_T \cdot \mathbf{n} - (\kappa \nabla v \cdot \mathbf{n})|_{\partial T}, \varphi \rangle_{\partial T} \\ &= (Ev, \varphi)_T = (Q^{r_1} Ev, \varphi)_T, \end{aligned}$$

for all $\varphi \in P_{r_1}(T)$.

Similarly, for any $u \in H^1(T)$, using equation (2.6), we obtain

$$(\nabla_w v, \boldsymbol{\psi})_T = (\nabla v, \boldsymbol{\psi})_T - \langle v|_T - v|_{\partial T}, \boldsymbol{\psi} \cdot \mathbf{n} \rangle_{\partial T} = (\nabla v, \boldsymbol{\psi})_T = (Q^{r_2} \nabla v, \boldsymbol{\psi})_T$$

for all $\boldsymbol{\psi} \in [P_{r_2}(T)]^d$.

This completes the proof of this lemma. \square

LEMMA 4.2. [53, 49] *Let \mathcal{T}_h be a finite element partition of Ω satisfying the shape regularity assumption as defined in [49]. Then, for any $0 \leq s \leq 2$, $0 \leq m \leq k$, $1 \leq n \leq r_1$, and $0 \leq q \leq r_2$, there exists a constant C such that the following estimates hold:*

$$(4.3) \quad \sum_{T \in \mathcal{T}_h} h_T^{2s} \|u - Q_0 u\|_{s,T}^2 \leq C h^{2(m+1)} \|u\|_{m+1}^2,$$

$$(4.4) \quad \sum_{T \in \mathcal{T}_h} h_T^{2s} \|Eu - Q^{r_1} Eu\|_{s,T}^2 \leq C h^{2(n-1)} \|u\|_{n+1}^2,$$

$$(4.5) \quad \sum_{T \in \mathcal{T}_h} h_T^{2s} \|\kappa \nabla u - Q^{r_2}(\kappa \nabla u)\|_{s,T}^2 \leq C h^{2q} \|u\|_{q+1}^2.$$

LEMMA 4.3. [53] *Let $0 \leq m \leq k$, $1 \leq n \leq r_1$, $0 \leq q \leq r_2$, and $u \in H^{\max\{n+1, 4\}}(\Omega)$. There exists a constant C such that the following estimates hold true:*

$$(4.6) \quad \left(\sum_{T \in \mathcal{T}_h} h_T \|Eu - Q^{r_1}(Eu)\|_{\partial T}^2 \right)^{\frac{1}{2}} \leq C h^{n-1} \|u\|_{n+1},$$

$$(4.7) \quad \left(\sum_{T \in \mathcal{T}_h} h_T^3 \|\kappa \nabla(Eu - Q^{r_1}(Eu)) \cdot \mathbf{n}\|_{\partial T}^2 \right)^{\frac{1}{2}} \leq C h^{n-1} (\|u\|_{n+1} + h \delta_{r_1, 0} \|u\|_4),$$

$$(4.8) \quad \left(\sum_{T \in \mathcal{T}_h} h_T^{-1} \|\kappa \nabla(Q_0 u) \cdot \mathbf{n} - Q_g(\kappa \nabla u \cdot \mathbf{n})\|_{\partial T}^2 \right)^{\frac{1}{2}} \leq C h^{m-1} \|u\|_{m+1},$$

$$(4.9) \quad \left(\sum_{T \in \mathcal{T}_h} h_T^{-3} \|Q_0 u - Q_b u\|_{\partial T}^2 \right)^{\frac{1}{2}} \leq C h^{m-1} \|u\|_{m+1},$$

$$(4.10) \quad \left(\sum_{T \in \mathcal{T}_h} h_T^3 \|(\kappa \nabla u - Q^{r_2}(\kappa \nabla u)) \cdot \mathbf{n}\|_{\partial T}^2 \right)^{\frac{1}{2}} \leq C h^{q+1} \|u\|_{q+1}.$$

Here $\delta_{i,j}$ is the usual Kronecker's delta with value 1 when $i = j$ and value 0 otherwise.

Proof. To prove (4.6), by the trace inequality (3.5) and the estimate (4.4), we get

$$\begin{aligned} & \sum_{T \in \mathcal{T}_h} h_T \|Eu - Q^{r_1}(Eu)\|_{\partial T}^2 \\ & \leq C \sum_{T \in \mathcal{T}_h} \|Eu - Q^{r_1}(Eu)\|_T^2 + h_T^2 \|Eu - Q^{r_1}(Eu)\|_{1,T}^2 \\ & \leq Ch^{2n-2} \|u\|_{n+1}^2. \end{aligned}$$

As to (4.7), by the trace inequality (3.5) and the estimate (4.4), we obtain

$$\begin{aligned} & \sum_{T \in \mathcal{T}_h} h_T^3 \|\kappa \nabla(Eu - Q^{r_1}(Eu)) \cdot \mathbf{n}\|_{\partial T}^2 \\ & \leq C \sum_{T \in \mathcal{T}_h} h_T^2 \|\nabla(Eu - Q^{r_1}(Eu))\|_T^2 + h_T^4 \|\nabla(Eu - Q^{r_1}(Eu))\|_{1,T}^2 \\ & \leq Ch^{2n-2} (\|u\|_{n+1}^2 + h^2 \delta_{r_1,0} \|u\|_4^2). \end{aligned}$$

As to (4.8), by the trace inequality (3.5) and the estimate (4.3), we have

$$\begin{aligned} & \sum_{T \in \mathcal{T}_h} h_T^{-1} \|\kappa \nabla(Q_0 u) \cdot \mathbf{n} - Q_g(\kappa \nabla u \cdot \mathbf{n})\|_{\partial T}^2 \\ & \leq \sum_{T \in \mathcal{T}_h} h_T^{-1} \|\kappa \nabla(Q_0 u) \cdot \mathbf{n} - \kappa \nabla u \cdot \mathbf{n}\|_{\partial T}^2 \\ & \leq C \sum_{T \in \mathcal{T}_h} h_T^{-1} \|\nabla(Q_0 u) - \nabla u\|_{\partial T}^2 \\ & \leq C \sum_{T \in \mathcal{T}_h} h_T^{-2} \|\nabla Q_0 u - \nabla u\|_T^2 + \|\nabla Q_0 u - \nabla u\|_{1,T}^2 \\ & \leq Ch^{2m-2} \|u\|_{m+1}^2. \end{aligned}$$

As to (4.9), by the trace inequality (3.5) and the estimate (4.3), we have

$$\begin{aligned} & \sum_{T \in \mathcal{T}_h} h_T^{-3} \|Q_0 u - Q_b u\|_{\partial T}^2 \\ & \leq \sum_{T \in \mathcal{T}_h} h_T^{-3} \|Q_0 u - u\|_{\partial T}^2 \\ & \leq C \sum_{T \in \mathcal{T}_h} h_T^{-4} \|Q_0 u - u\|_T^2 + h_T^{-2} \|\nabla(Q_0 u - u)\|_T^2 \\ & \leq Ch^{2m-2} \|u\|_{m+1}^2. \end{aligned}$$

Finally, as to (4.10), by the trace inequality (3.5) and the estimate (4.5), we have

$$\begin{aligned} & \sum_{T \in \mathcal{T}_h} h_T^3 \|(\kappa \nabla u - Q^{r_2}(\kappa \nabla u)) \cdot \mathbf{n}\|_{\partial T}^2 \\ & \leq \sum_{T \in \mathcal{T}_h} h_T^2 \|\kappa \nabla u - Q^{r_2}(\kappa \nabla u)\|_T^2 + h_T^4 \|\kappa \nabla u - Q^{r_2}(\kappa \nabla u)\|_{1,T}^2 \\ & \leq Ch^{2q+2} \|u\|_{q+1}^2. \end{aligned}$$

This completes the proof of the lemma. \square

5. An Error Equation. Let u be the exact solution of (1.1), and let $u_h = \{u_0, u_b, u_g\} \in V_h$ be the weak Galerkin finite element approximation satisfying (3.2). Define the error function as

$$(5.1) \quad e_h = u - u_h.$$

The objective of this section is to derive an error equation for e_h .

LEMMA 5.1. *The error function $e_h \in V_h^0$ as defined by (5.1) satisfies the following equation*

$$(5.2) \quad \sum_{T \in \mathcal{T}_h} (E_w e_h, E_w v)_T + 2\mu(\kappa \nabla_w e_h, \nabla_w v)_T + \mu^2(e_0, v_0)_T = \phi_u(v), \quad \forall v \in V_h^0,$$

where

$$(5.3) \quad \begin{aligned} \phi_u(v) = & \sum_{T \in \mathcal{T}_h} \left(-\langle \kappa \nabla(Eu - Q^{r_1}(Eu)) \cdot \mathbf{n}, v_0 - v_b \rangle_{\partial T} \right. \\ & + \langle \kappa \nabla v_0 \cdot \mathbf{n} - v_g, Eu - Q^{r_1}Eu \rangle_{\partial T} \\ & \left. + 2\mu \langle v_0 - v_b, \kappa(\nabla u - Q^{r_2}\nabla u) \cdot \mathbf{n} \rangle_{\partial T} \right). \end{aligned}$$

Proof. Using (2.3) with $\varphi = E_w u$, from (4.1), we obtain

$$\begin{aligned} (E_w v, E_w u)_T &= (E_w v, Q^{r_1}Eu)_T \\ &= (Ev_0, Q^{r_1}Eu)_T + \langle v_0 - v_b, \kappa \nabla(Q^{r_1}Eu) \cdot \mathbf{n} \rangle_{\partial T} \\ &\quad - \langle \kappa \nabla v_0 \cdot \mathbf{n} - v_g, Q^{r_1}Eu \rangle_{\partial T} \\ &= (Ev_0, Eu)_T + \langle v_0 - v_b, \kappa \nabla(Q^{r_1}Eu) \cdot \mathbf{n} \rangle_{\partial T} \\ &\quad - \langle \kappa \nabla v_0 \cdot \mathbf{n} - v_g, Q^{r_1}Eu \rangle_{\partial T}, \end{aligned}$$

which implies that

$$(5.4) \quad \begin{aligned} (Ev_0, Eu)_T &= (E_w u, E_w v)_T - \langle v_0 - v_b, \kappa \nabla(Q^{r_1}Eu) \cdot \mathbf{n} \rangle_{\partial T} \\ &\quad + \langle \kappa \nabla v_0 \cdot \mathbf{n} - v_g, Q^{r_1}Eu \rangle_{\partial T}. \end{aligned}$$

Next, it follows from the integration by parts that

$$(5.5) \quad \sum_{T \in \mathcal{T}_h} (Eu, Ev_0)_T = \sum_{T \in \mathcal{T}_h} (E^2 u, v_0)_T - \langle \kappa \nabla(Eu) \cdot \mathbf{n}, v_0 \rangle_{\partial T} + \langle \kappa \nabla v_0 \cdot \mathbf{n}, Eu \rangle_{\partial T}.$$

Using (2.6) with $\psi = \kappa Q^{r_2}(\nabla u)$, from (4.2) and the integration by parts, we have

$$\begin{aligned}
& \sum_{T \in \mathcal{T}_h} 2\mu(\kappa \nabla_w u, \nabla_w v)_T \\
&= \sum_{T \in \mathcal{T}_h} 2\mu(\kappa Q^{r_2}(\nabla u), \nabla_w v)_T \\
&= \sum_{T \in \mathcal{T}_h} 2\mu(\nabla v_0, \kappa Q^{r_2}(\nabla u))_T - 2\mu\langle v_0 - v_b, \kappa Q^{r_2}(\nabla u) \cdot \mathbf{n} \rangle_{\partial T} \\
(5.6) \quad &= \sum_{T \in \mathcal{T}_h} 2\mu(\nabla v_0, \kappa \nabla u)_T - 2\mu\langle v_0 - v_b, \kappa Q^{r_2}(\nabla u) \cdot \mathbf{n} \rangle_{\partial T} \\
&= \sum_{T \in \mathcal{T}_h} -2\mu(v_0, \nabla \cdot (\kappa \nabla u))_T + 2\mu\langle v_0, \kappa \nabla u \cdot \mathbf{n} \rangle_{\partial T} \\
&\quad - 2\mu\langle v_0 - v_b, \kappa Q^{r_2}(\nabla u) \cdot \mathbf{n} \rangle_{\partial T} \\
&= \sum_{T \in \mathcal{T}_h} -2\mu(v_0, Eu)_T + 2\mu\langle v_0 - v_b, \kappa(\nabla u - Q^{r_2}(\nabla u)) \cdot \mathbf{n} \rangle_{\partial T},
\end{aligned}$$

where we have used the fact that the sum for the terms associated with v_b vanishes (note that v_b vanishes on $\partial T \cap \partial\Omega$).

Testing v_0 on both sides of the first equation of (1.1) gives

$$(5.7) \quad \sum_{T \in \mathcal{T}_h} (E^2 u, v_0)_T - 2\mu(Eu, v_0)_T + \mu^2(u, v_0)_T = \sum_{T \in \mathcal{T}_h} (f, v_0)_T.$$

Adding (5.5)-(5.6) and using (5.7), we have

$$\begin{aligned}
& \sum_{T \in \mathcal{T}_h} (Eu, Ev_0)_T + 2\mu\kappa(\nabla_w u, \nabla_w v)_T + \mu^2(u, v_0)_T \\
&= (f, v_0) + \sum_{T \in \mathcal{T}_h} \left(-\langle \kappa \nabla(Eu) \cdot \mathbf{n}, v_0 - v_b \rangle_{\partial T} + \langle \kappa \nabla v_0 \cdot \mathbf{n} - v_g, Eu \rangle_{\partial T} \right. \\
&\quad \left. + 2\mu\langle v_0 - v_b, \kappa(\nabla u - Q^{r_2} \nabla u) \cdot \mathbf{n} \rangle_{\partial T} \right),
\end{aligned}$$

where we have used the fact that the sum for the terms associated with v_b and v_g vanishes (note that both v_b and v_g vanish on $\partial T \cap \partial\Omega$). Combining the above equation with (5.4) yield

$$\begin{aligned}
& \sum_{T \in \mathcal{T}_h} (E_w u, E_w v)_T + 2\mu\kappa(\nabla_w u, \nabla_w v)_T + \mu^2(u, v_0)_T \\
&= (f, v_0) + \sum_{T \in \mathcal{T}_h} \left(-\langle \kappa \nabla(Eu - Q^{r_1}(Eu)) \cdot \mathbf{n}, v_0 - v_b \rangle_{\partial T} \right. \\
&\quad + \langle \kappa \nabla v_0 \cdot \mathbf{n} - v_g, Eu - Q^{r_1} Eu \rangle_{\partial T} \\
&\quad \left. + 2\mu\langle v_0 - v_b, \kappa(\nabla u - Q^{r_2} \nabla u) \cdot \mathbf{n} \rangle_{\partial T} \right),
\end{aligned}$$

which, subtracting (3.2), completes the proof. \square

6. Error Estimates. This section derives error estimates for the solution obtained using the stabilizer-free weak Galerkin algorithm.

THEOREM 6.1. *Let $u_h \in V_h$ be the weak Galerkin finite element solution resulting from (3.2), using finite elements of order $k \geq 2$. Suppose the exact solution u of (1.1) is sufficiently regular, satisfying $u \in H^{k+1}(\Omega)$. Then, there exists a constant C such that*

$$(6.1) \quad \|u - Q_h u\| \leq Ch^{k-1} \|u\|_{k+1}.$$

Proof. Using (2.3), the trace inequality (3.6), Cauchy-Schwartz inequality, the inverse inequality, the estimates (4.3), (4.8) and (4.9) we derive:

$$\begin{aligned} & \sum_{T \in \mathcal{T}_h} (E_w(u - Q_h u), \phi)_T \\ &= \sum_{T \in \mathcal{T}_h} (E(u - Q_0 u), \phi)_T + \langle Q_b u - Q_0 u, \kappa \nabla \phi \cdot \mathbf{n} \rangle_{\partial T} \\ & \quad - \langle \kappa \nabla(u - Q_0 u) \cdot \mathbf{n} - (u_g - Q_g(\kappa \nabla u \cdot \mathbf{n})), \phi \rangle_{\partial T} \\ &= \sum_{T \in \mathcal{T}_h} (E(u - Q_0 u), \phi)_T + \langle Q_b u - Q_0 u, \kappa \nabla \phi \cdot \mathbf{n} \rangle_{\partial T} \\ & \quad - \langle -\kappa \nabla Q_0 u \cdot \mathbf{n} + Q_g(\kappa \nabla u \cdot \mathbf{n}), \phi \rangle_{\partial T} \\ &\leq \left(\sum_{T \in \mathcal{T}_h} \|E(u - Q_0 u)\|_T^2 \right)^{\frac{1}{2}} \left(\sum_{T \in \mathcal{T}_h} \|\phi\|_T^2 \right)^{\frac{1}{2}} \\ & \quad + \left(\sum_{T \in \mathcal{T}_h} \|Q_b u - Q_0 u\|_{\partial T}^2 \right)^{\frac{1}{2}} \left(\sum_{T \in \mathcal{T}_h} \|\kappa \nabla \phi \cdot \mathbf{n}\|_{\partial T}^2 \right)^{\frac{1}{2}} \\ & \quad + \left(\sum_{T \in \mathcal{T}_h} \|-\kappa \nabla Q_0 u \cdot \mathbf{n} + Q_g(\kappa \nabla u \cdot \mathbf{n})\|_{\partial T}^2 \right)^{\frac{1}{2}} \left(\sum_{T \in \mathcal{T}_h} \|\phi\|_{\partial T}^2 \right)^{\frac{1}{2}} \\ &\leq Ch^{k-1} \|u\|_{k+1} \left(\sum_{T \in \mathcal{T}_h} \|\phi\|_T^2 \right)^{\frac{1}{2}}. \end{aligned}$$

Setting $\phi = E_w(u - Q_h u)$ leads to

$$(6.2) \quad \sum_{T \in \mathcal{T}_h} \|E_w(u - Q_h u)\|_T^2 \leq Ch^{2k-2} \|u\|_{k+1}^2$$

Using (2.6), the trace inequality (3.6), Cauchy-Schwartz inequality, (4.3), (4.9), we have

$$\begin{aligned} & \sum_{T \in \mathcal{T}_h} 2\mu(\kappa \nabla_w(u - Q_h u), \psi)_T \\ &= \sum_{T \in \mathcal{T}_h} 2\mu\kappa((\nabla(u - Q_0 u), \psi)_T - \langle Q_b u - Q_0 u, \psi \cdot \mathbf{n} \rangle_{\partial T}) \\ &\leq \left(\sum_{T \in \mathcal{T}_h} \|\nabla(u - Q_0 u)\|_T^2 \right)^{\frac{1}{2}} \left(\sum_{T \in \mathcal{T}_h} \|\psi\|_T^2 \right)^{\frac{1}{2}} + \left(\sum_{T \in \mathcal{T}_h} \|Q_b u - Q_0 u\|_{\partial T}^2 \right)^{\frac{1}{2}} \left(\sum_{T \in \mathcal{T}_h} \|\psi \cdot \mathbf{n}\|_{\partial T}^2 \right)^{\frac{1}{2}} \\ &\leq Ch^k \|u\|_{k+1} \left(\sum_{T \in \mathcal{T}_h} \|\psi\|_T^2 \right)^{\frac{1}{2}}. \end{aligned}$$

Letting $\psi = \nabla_w(u - Q_h u)$ gives

$$(6.3) \quad \sum_{T \in \mathcal{T}_h} 2\mu(\kappa \nabla_w(u - Q_h u), \nabla_w(u - Q_h u))_T \leq Ch^{2k} \|u\|_{k+1}^2.$$

Using Cauchy-Schwartz inequality, (4.3), we have

$$(6.4) \quad \sum_{T \in \mathcal{T}_h} \mu^2 (u - Q_0 u, u - Q_0 u)_T \leq \left(\sum_{T \in \mathcal{T}_h} \|u - Q_0 u\|_T^2 \right) \leq Ch^{2(k+1)} \|u\|_{k+1}^2.$$

Combining (6.2), (6.3) and (6.4) completes the proof (6.1). \square

THEOREM 6.2. *Let $k \geq 2$. Let $u_h \in V_h$ be the weak Galerkin finite element solution of (3.2). Assume that the exact solution u of (1.1) satisfies $u \in H^{\max\{k+1, 4\}}(\Omega)$. Then, there exists a constant C such that*

$$(6.5) \quad \|u - u_h\| \leq Ch^{k-1} \left(\|u\|_{k+1} + h\delta_{r_1,0} \|u\|_4 \right).$$

Proof. We estimate each term on the right-hand side of (5.2) as follows: For the first term, applying the Cauchy-Schwarz inequality and using estimate (4.7), we obtain

$$\begin{aligned} & \left| \sum_{T \in \mathcal{T}_h} \langle \kappa \nabla (Eu - Q^{r_1} Eu) \cdot \mathbf{n}, v_0 - v_b \rangle_{\partial T} \right| \\ & \leq \left(\sum_{T \in \mathcal{T}_h} h_T^3 \|\kappa \nabla (Eu - Q^{r_1} Eu) \cdot \mathbf{n}\|_{\partial T}^2 \right)^{\frac{1}{2}} \left(\sum_{T \in \mathcal{T}_h} h_T^{-3} \|v_0 - v_b\|_{\partial T}^2 \right)^{\frac{1}{2}} \\ & \leq Ch^{k-1} (\|u\|_{k+1} + h\delta_{r_1,0} \|u\|_4) \|v\|_{2,h}. \end{aligned}$$

For the second term, again applying the Cauchy-Schwarz inequality and using estimate (4.6), we obtain

$$\begin{aligned} & \left| \sum_{T \in \mathcal{T}_h} \langle \kappa \nabla v_0 \cdot \mathbf{n} - v_g, Eu - Q^{r_1} Eu \rangle_{\partial T} \right| \\ & \leq \left(\sum_{T \in \mathcal{T}_h} h_T^{-1} \|\kappa \nabla v_0 \cdot \mathbf{n} - v_g\|_{\partial T}^2 \right)^{\frac{1}{2}} \left(\sum_{T \in \mathcal{T}_h} h_T \|Eu - Q^{r_1} Eu\|_{\partial T}^2 \right)^{\frac{1}{2}} \\ & \leq Ch^{k-1} \|u\|_{k+1} \|v\|_{2,h}. \end{aligned}$$

For the third term, using the Cauchy-Schwarz inequality and estimate (4.10), we derive

$$\begin{aligned} & \sum_{T \in \mathcal{T}_h} 2\mu \langle v_0 - v_b, \kappa (\nabla u - Q^{r_2} \nabla u) \cdot \mathbf{n} \rangle_{\partial T} \\ & \leq \left(\sum_{T \in \mathcal{T}_h} h_T^{-3} \|v_0 - v_b\|_{\partial T}^2 \right)^{\frac{1}{2}} \left(\sum_{T \in \mathcal{T}_h} h_T^3 \|\kappa (\nabla u - Q^{r_2} \nabla u) \cdot \mathbf{n}\|_{\partial T}^2 \right)^{\frac{1}{2}} \\ & \leq Ch^{k+1} \|u\|_{k+1} \|v\|_{2,h}. \end{aligned}$$

Substituting these estimates into (5.2) and using (3.11), we obtain

$$(6.6) \quad \begin{aligned} & (E_w e_h, E_w v) + 2\mu (\kappa \nabla_w e_h, \nabla_w v) + \mu^2 (e_h, v) \\ & \leq Ch^{k-1} (\|u\|_{k+1} + h\delta_{r_1,0} \|u\|_4) \|v\|_{2,h} \\ & \leq Ch^{k-1} (\|u\|_{k+1} + h\delta_{r_1,0} \|u\|_4) \|v\|. \end{aligned}$$

Choosing $v = Q_u - u_h$ in (6.6) and applying the Cauchy-Schwarz inequality and the estimate (6.1), we derive

$$\begin{aligned}
& \|e_h\|^2 \\
&= (E_w e_h, E_w(u - Q_h u)) + 2\mu(\kappa \nabla_w e_h, \nabla_w(u - Q_h u)) + \mu^2(e_h, (u - Q_h u)) \\
&\quad + (E_w e_h, E_w(Q_h u - u_h)) + 2\mu(\kappa \nabla_w e_h, \nabla_w(Q_h u - u_h)) + \mu^2(e_h, (Q_h u - u_h)) \\
&\leq \|e_h\| \|u - Q_h u\| + Ch^{k-1}(\|u\|_{k+1} + h\delta_{r_1,0}\|u\|_4) \|Q_h u - u_h\| \\
&\leq \|e_h\| \|u - Q_h u\| + Ch^{k-1}(\|u\|_{k+1} + h\delta_{r_1,0}\|u\|_4) (\|Q_h u - u\| + \|u - u_h\|) \\
&\leq Ch^{k-1}(\|u\|_{k+1} + h\delta_{r_1,0}\|u\|_4) \|e_h\| + Ch^{k-1}(\|u\|_{k+1} + h\delta_{r_1,0}\|u\|_4) h^{k-1} \|u\|_{k+1}.
\end{aligned}$$

This establishes the desired bound, concluding the proof. \square

7. Numerical test. In the first numerical example, we solve (1.1) on the unit square domain $\Omega = (0, 1) \times (0, 1)$ with the following parameters and the exact solution,

$$\begin{aligned}
(7.1) \quad & \mu = 1, \quad \kappa = \begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix}, \\
& u = (x - x^2)^2 (y - y^2)^2.
\end{aligned}$$




FIG. 7.1. The first three grids for the computation in Tables 7.1–7.2.

We compute the finite element solutions for (7.1) on uniform triangular grids shown in Figure 7.1 by the $P_k/P_k/P_{k-1}$ WG finite elements, defined in (3.1), for $k = 2, 3$ and 4. The results are listed in Table 7.1. The optimal orders of convergence are achieved for all solutions in all norms.

TABLE 7.1

The error and the computed order of convergence for the solution (7.1) on Figure 7.1 triangular meshes.

G_i	$\ Q_0 u - u_0\ _0$	h^r	$\ \nabla_w(Q_h u - u_h)\ _0$	h^r	$\ E_w(Q_h u - u_h)\ _0$	h^r
By the $P_2/P_2/P_1$ WG finite element (3.1).						
4	0.183E-3	1.5	0.196E-2	1.6	0.247E+0	0.9
5	0.502E-4	1.9	0.536E-3	1.9	0.125E+0	1.0
6	0.128E-4	2.0	0.137E-3	2.0	0.629E-1	1.0
By the $P_3/P_3/P_2$ WG finite element (3.1).						
3	0.360E-4	3.2	0.660E-3	2.9	0.835E-1	1.8
4	0.277E-5	3.7	0.808E-4	3.0	0.221E-1	1.9
5	0.184E-6	3.9	0.982E-5	3.0	0.561E-2	2.0
By the $P_4/P_4/P_3$ WG finite element (3.1).						
2	0.243E-4	3.8	0.718E-3	3.2	0.669E-1	2.4
3	0.140E-5	4.1	0.777E-4	3.2	0.126E-1	2.4
4	0.351E-7	5.3	0.528E-5	3.9	0.174E-2	2.9

In the second numerical example, we solve (1.1) on the unit square domain $\Omega = (0, 1) \times (0, 1)$ with the following parameters and the exact solution,

$$(7.2) \quad \begin{aligned} \mu &= 1, \quad \kappa = \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}, \\ u &= (x - x^2)^2 (y - y^2)^2. \end{aligned}$$

We compute the finite element solutions for (7.2) on uniform triangular grids shown in Figure 7.1 by the $P_k/P_k/P_{k-1}$ WG finite elements, defined in (3.1), for $k = 2, 3$ and 4. The results are listed in Table 7.2. The optimal orders of convergence are achieved for all solutions in all norms.

TABLE 7.2

The error and the computed order of convergence for the solution (7.2) on Figure 7.1 triangular meshes.

G_i	$\ Q_0 u - u_0\ _0$	h^r	$\ \nabla_w(Q_h u - u_h)\ _0$	h^r	$\ E_w(Q_h u - u_h)\ _0$	h^r
By the $P_2/P_2/P_1$ WG finite element (3.1).						
4	0.267E-3	1.2	0.258E-2	1.3	0.327E+0	0.9
5	0.858E-4	1.6	0.821E-3	1.7	0.165E+0	1.0
6	0.236E-4	1.9	0.226E-3	1.9	0.830E-1	1.0
By the $P_3/P_3/P_2$ WG finite element (3.1).						
3	0.586E-4	2.9	0.780E-3	2.8	0.106E+0	1.8
4	0.474E-5	3.6	0.910E-4	3.1	0.282E-1	1.9
5	0.313E-6	3.9	0.106E-4	3.1	0.716E-2	2.0
By the $P_4/P_4/P_3$ WG finite element (3.1).						
2	0.291E-4	3.8	0.783E-3	3.2	0.889E-1	2.5
3	0.216E-5	3.8	0.888E-4	3.1	0.162E-1	2.5
4	0.482E-7	5.5	0.578E-5	3.9	0.223E-2	2.9




FIG. 7.2. The first three nonconvex polygonal grids for the computation in Tables 7.3–7.4.

We compute the two numerical examples again on nonconvex polygonal meshes, shown in Figure 7.2. We apply the $P_k/P_k/P_{k-1}$ WG finite elements, defined in (3.1), for $k = 2, 3$ and 4. The results are listed in Tables 7.3–7.4. We obtain the optimal orders of convergence for all cases.

TABLE 7.3

The error and the computed order of convergence for the solution (7.1) on Figure 7.2 polygonal meshes.

G_i	$\ Q_0 u - u_0\ _0$	h^r	$\ \nabla_w(Q_h u - u_h)\ _0$	h^r	$\ E_w(Q_h u - u_h)\ _0$	h^r
By the $P_2/P_2/P_1$ WG finite element (3.1).						
4	0.152E-3	1.6	0.165E-2	1.7	0.199E+0	0.9
5	0.405E-4	1.9	0.435E-3	1.9	0.102E+0	1.0
6	0.102E-4	2.0	0.109E-3	2.0	0.513E-1	1.0
By the $P_3/P_3/P_2$ WG finite element (3.1).						
3	0.229E-4	3.3	0.800E-3	2.5	0.586E-1	2.2
4	0.164E-5	3.8	0.109E-3	2.9	0.156E-1	1.9
5	0.107E-6	3.9	0.138E-4	3.0	0.400E-2	2.0
By the $P_4/P_4/P_3$ WG finite element (3.1).						
2	0.133E-4	5.4	0.957E-3	5.3	0.801E-1	4.8
3	0.484E-6	4.8	0.120E-3	3.0	0.870E-2	3.2
4	0.145E-7	5.1	0.792E-5	3.9	0.118E-2	2.9

TABLE 7.4

The error and the computed order of convergence for the solution (7.2) on Figure 7.2 polygonal meshes.

G_i	$\ Q_0 u - u_0\ _0$	h^r	$\ \nabla_w(Q_h u - u_h)\ _0$	h^r	$\ E_w(Q_h u - u_h)\ _0$	h^r
By the $P_2/P_2/P_1$ WG finite element (3.1).						
4	0.227E-3	1.3	0.225E-2	1.4	0.288E+0	0.9
5	0.676E-4	1.7	0.657E-3	1.8	0.148E+0	1.0
6	0.178E-4	1.9	0.172E-3	1.9	0.743E-1	1.0
By the $P_3/P_3/P_2$ WG finite element (3.1).						
3	0.328E-4	3.3	0.113E-2	2.0	0.770E-1	2.1
4	0.231E-5	3.8	0.158E-3	2.8	0.209E-1	1.9
5	0.148E-6	4.0	0.197E-4	3.0	0.537E-2	2.0
By the $P_4/P_4/P_3$ WG finite element (3.1).						
2	0.157E-4	5.2	0.112E-2	5.2	0.104E+0	4.8
3	0.702E-6	4.5	0.128E-3	3.1	0.112E-1	3.2
4	0.201E-7	5.1	0.919E-5	3.8	0.152E-2	2.9

REFERENCES

- [1] S. BRENNER, V. NTZIACHRISTOS AND R. WEISSLEDER, *Optical-based molecular imaging: contrast agents and potential medical applications*, European Radiology, vol. 13, pp. 231-243, 2003.
- [2] W. CAO, C. WANG AND J. WANG, *An L^p -Primal-Dual Weak Galerkin Method for div-curl Systems*, Journal of Computational and Applied Mathematics, vol. 422, 114881, 2023.
- [3] W. CAO, C. WANG AND J. WANG, *An L^p -Primal-Dual Weak Galerkin Method for Convection-Diffusion Equations*, Journal of Computational and Applied Mathematics, vol. 419, 114698, 2023.
- [4] W. CAO, C. WANG AND J. WANG, *A New Primal-Dual Weak Galerkin Method for Elliptic Interface Problems with Low Regularity Assumptions*, Journal of Computational Physics, vol. 470, 111538, 2022.
- [5] S. CAO, C. WANG AND J. WANG, *A new numerical method for div-curl Systems with Low Regularity Assumptions*, Computers and Mathematics with Applications, vol. 144, pp. 47-59, 2022.
- [6] W. CAO AND C. WANG, *New Primal-Dual Weak Galerkin Finite Element Methods for Convection-Diffusion Problems*, Applied Numerical Mathematics, vol. 162, pp. 171-191, 2021.
- [7] S. CHOW, K. YIN, H. ZHOU AND A. BEHROOZ, *Solving inverse source problems by the orthogonal solution and kernel correction algorithm(OSKCA) with applications in fluorescence tomography*, Inverse Problems and Imaging, vol. 8, pp. 79-102, 2014.
- [8] H. GAO AND H. ZHAO, *Analysis of a numerical solver for radiative transport equation*, Math. Comp., vol. 82, no. 281, pp. 153-172, 2013.
- [9] J. GUERMOND, G. KANSCHAT AND J. RAGUSA, *Discontinuous Galerkin for the radiative transport equation. Recent developments in discontinuous Galerkin finite element methods for partial differential equations*, IMA Vol. Math. Appl., 157, Springer, Cham, pp. 181-193, 2014.
- [10] J. HEBDEN, S. ARRIDGE AND D. DELPY, *Optical imaging in medicine: I. experimental techniques*, Physics in Medicine and Biology, vol. 42, pp. 825-840, 1997.
- [11] D. HAWRYSZ AND E. SEVICK-MURACA, *Developments toward diagnostic breast cancer imaging using near-infrared optical measurements and fluorescent contrast agents*, Neoplasia (New York, NY), vol. 2, pp. 388-417, 2000.
- [12] O. LEHTIKANGAS, T. TARVAINEN, A. KIM AND S. ARRIDGE, *Finite element approximation of the radiative transport equation in a medium with piece-wise constant refractive index*, J. Comput. Phys., vol. 282, pp. 345-359, 2015.
- [13] D. LI, Y. NIE, AND C. WANG, *Superconvergence of Numerical Gradient for Weak Galerkin*

- Finite Element Methods on Nonuniform Cartesian Partitions in Three Dimensions*, Computers and Mathematics with Applications, vol 78(3), pp. 905-928, 2019.
- [14] D. LI, C. WANG AND J. WANG, *An Extension of the Morley Element on General Polytopal Partitions Using Weak Galerkin Methods*, Journal of Scientific Computing, 100, vol 27, 2024.
- [15] D. LI, C. WANG AND S. ZHANG, *Weak Galerkin methods for elliptic interface problems on curved polygonal partitions*, Journal of Computational and Applied Mathematics, pp. 115995, 2024.
- [16] D. LI, C. WANG, J. WANG AND X. YE, *Generalized weak Galerkin finite element methods for second order elliptic problems*, Journal of Computational and Applied Mathematics, vol. 445, pp. 115833, 2024.
- [17] D. LI, C. WANG, J. WANG AND S. ZHANG, *High Order Morley Elements for Biharmonic Equations on Polytopal Partitions*, Journal of Computational and Applied Mathematics, Vol. 443, pp. 115757, 2024.
- [18] D. LI, C. WANG AND J. WANG, *Curved Elements in Weak Galerkin Finite Element Methods*, Computers and Mathematics with Applications, Vol. 153, pp. 20-32, 2024.
- [19] D. LI, C. WANG AND J. WANG, *Generalized Weak Galerkin Finite Element Methods for Biharmonic Equations*, Journal of Computational and Applied Mathematics, vol. 434, 115353, 2023.
- [20] D. LI, C. WANG AND J. WANG, *An L^p -primal-dual finite element method for first-order transport problems*, Journal of Computational and Applied Mathematics, vol. 434, 115345, 2023.
- [21] D. LI AND C. WANG, *A simplified primal-dual weak Galerkin finite element method for Fokker-Planck type equations*, Journal of Numerical Methods for Partial Differential Equations, vol 39, pp. 3942-3963, 2023.
- [22] D. LI, C. WANG AND J. WANG, *Primal-Dual Weak Galerkin Finite Element Methods for Transport Equations in Non-Divergence Form*, Journal of Computational and Applied Mathematics, vol. 412, 114313, 2022.
- [23] D. LI, C. WANG, AND J. WANG, *Superconvergence of the Gradient Approximation for Weak Galerkin Finite Element Methods on Rectangular Partitions*, Applied Numerical Mathematics, vol. 150, pp. 396-417, 2020.
- [24] B. LI, X. XIE AND S. ZHANG, *BPS preconditioners for a weak Galerkin finite element method for 2D diffusion problems with strongly discontinuous coefficients*, Computers & Mathematics with Applications, 76(4), pp.701-724, 2018.
- [25] S. MOHAN, T. TARVAINEN, M. SCHWEIGER, A. PULKKINEN AND S. ARRIDGE, *Variable order spherical harmonic expansion scheme for the radiative transport equation using finite elements*, J. Comput. Phys., vol. 230, no. 19, pp. 7364-7383, 2011.
- [26] L. MORLEY, *The triangular equilibrium element in the solution of plate bending problems*, Aero. Quart., vol. 19, pp. 149-169, 1968.
- [27] C. WANG, *Auto-Stabilized Weak Galerkin Finite Element Methods on Polytopal Meshes without Convexity Constraints*, arXiv:2408.11927.
- [28] C. WANG, *A Preconditioner for the FETI-DP Method for Mortar-Type Crouzeix-Raviart Element Discretization*, Applications of Mathematics, Vol. 59, 6, pp. 653-672, 2014.
- [29] C. WANG, *New Discretization Schemes for Time-Harmonic Maxwell Equations by Weak Galerkin Finite Element Methods*, Journal of Computational and Applied Mathematics, Vol. 341, pp. 127-143, 2018.
- [30] C. WANG, *Low Regularity Primal-Dual Weak Galerkin Finite Element Methods for Ill-Posed Elliptic Cauchy Problems*, Int. J. Numer. Anal. Mod., vol. 19(1), pp. 33-51, 2022.
- [31] C. WANG, *A Modified Primal-Dual Weak Galerkin Finite Element Method for Second Order Elliptic Equations in Non-Divergence Form*, Int. J. Numer. Anal. Mod., vol. 18(4), pp. 500-523, 2021.
- [32] C. WANG, *A New Primal-Dual Weak Galerkin Finite Element Method for Ill-posed Elliptic Cauchy Problems*, Journal of Computational and Applied Mathematics, vol 371, 112629, 2020.
- [33] J. WARSA, M. BENZI, T. WAREING AND J. MOREL, *Two-level preconditioning of a discontinuous Galerkin method for radiation diffusion*, Numerical mathematics and advanced applications, Springer Italia, Milan, pp. 967-977, 2003.
- [34] J. WARSA, M. BENZI, T. WAREING AND J. MOREL, *Preconditioning a mixed discontinuous finite element method for radiation diffusion*, Numer. Linear Algebra Appl., vol. 11, no. 8-9, pp. 795-811, 2004.
- [35] R. WEISSELEDER, C. TUNG, U. MAHMOOD AND A. BOGDANOV, *In vivo imaging of tumors with protease-activated near-infrared fluorescent probes*, Nature Biotechnology, vol. 17, pp. 375-378, 1999.

- [36] C. WANG AND J. WANG, *A Primal-Dual Weak Galerkin Finite Element Method for Fokker-Planck Type Equations*, SIAM Numerical Analysis, vol. 58(5), pp. 2632-2661, 2020.
- [37] C. WANG AND J. WANG, *A Primal-Dual Finite Element Method for First-Order Transport Problems*, Journal of Computational Physics, Vol. 417, 109571, 2020.
- [38] C. WANG AND J. WANG, *Primal-Dual Weak Galerkin Finite Element Methods for Elliptic Cauchy Problems*, Computers and Mathematics with Applications, vol 79(3), pp. 746-763, 2020.
- [39] C. WANG AND J. WANG, *A Primal-Dual Weak Galerkin Finite Element Method for Second Order Elliptic Equations in Non-Divergence form*, Mathematics of Computation, Vol. 87, pp. 515-545, 2018.
- [40] C. WANG AND J. WANG, *Discretization of Div-Curl Systems by Weak Galerkin Finite Element Methods on Polyhedral Partitions*, Journal of Scientific Computing, Vol. 68, pp. 1144-1171, 2016.
- [41] C. WANG AND J. WANG, *A Hybridized Formulation for Weak Galerkin Finite Element Methods for Biharmonic Equation on Polygonal or Polyhedral Meshes*, International Journal of Numerical Analysis and Modeling, Vol. 12, pp. 302-317, 2015.
- [42] J. WANG AND C. WANG, *Weak Galerkin Finite Element Methods for Elliptic PDEs*, Science China, Vol. 45, pp. 1061-1092, 2015.
- [43] C. WANG AND J. WANG, *An Efficient Numerical Scheme for the Biharmonic Equation by Weak Galerkin Finite Element Methods on Polygonal or Polyhedral Meshes*, Journal of Computers and Mathematics with Applications, Vol. 68, 12, pp. 2314-2330, 2014.
- [44] C. WANG, J. WANG, R. WANG AND R. ZHANG, *A Locking-Free Weak Galerkin Finite Element Method for Elasticity Problems in the Primal Formulation*, Journal of Computational and Applied Mathematics, Vol. 307, pp. 346-366, 2016.
- [45] C. WANG, J. WANG, X. YE AND S. ZHANG, *De Rham Complexes for Weak Galerkin Finite Element Spaces*, Journal of Computational and Applied Mathematics, vol. 397, pp. 113645, 2021.
- [46] C. WANG, J. WANG AND S. ZHANG, *Weak Galerkin Finite Element Methods for Optimal Control Problems Governed by Second Order Elliptic Partial Differential Equations*, Journal of Computational and Applied Mathematics, in press, 2024.
- [47] C. WANG, J. WANG AND S. ZHANG, *A parallel iterative procedure for weak Galerkin methods for second order elliptic problems*, International Journal of Numerical Analysis and Modeling, vol. 21(1), pp. 1-19, 2023.
- [48] C. WANG, J. WANG AND S. ZHANG, *Weak Galerkin Finite Element Methods for Quad-Curl Problems*, Journal of Computational and Applied Mathematics, vol. 428, pp. 115186, 2023.
- [49] J. WANG, AND X. YE, *A weak Galerkin mixed finite element method for second-order elliptic problems*, Math. Comp., vol. 83, pp. 2101-2126, 2014.
- [50] C. WANG, X. YE AND S. ZHANG, *A Modified weak Galerkin finite element method for the Maxwell equations on polyhedral meshes*, Journal of Computational and Applied Mathematics, vol. 448, pp. 115918, 2024.
- [51] C. WANG AND S. ZHANG, *A Weak Galerkin Method for Elasticity Interface Problems*, Journal of Computational and Applied Mathematics, vol. 419, 114726, 2023.
- [52] C. WANG AND L. ZIKATANOV, *Low Regularity Primal-Dual Weak Galerkin Finite Element Methods for Convection-Diffusion Equations*, Journal of Computational and Applied Mathematics, vol 394, 113543, 2021.
- [53] C. WANG AND H. ZHOU, *A Weak Galerkin Finite Element Method for a Type of Fourth Order Problem arising from Fluorescence Tomography*, Journal of Scientific Computing, Vol. 71(3), pp. 897-918, 2017.
- [54] K. YIN, *New Algorithms for solving inverse source problems in imaging techniques with applications in fluorescence tomography*, Ph.D. Thesis, Georgia Institute of Technology, 2013.
- [55] Y. XIU AND S. ZHANG, *A stabilizer-free weak Galerkin finite element method on polytopal meshes*, Journal of Computational and Applied Mathematics, vol 371, 112699, 2020.