

New constructions of unbalanced $\{C_4, \theta_{3,t}\}$ -free bipartite graphs

Baran Düzgün ^{*} Ago-Erik Riet [†] Vladislav Taranchuk [‡]

March 25, 2025

Abstract

In 1979, Erdős conjectured that if $m = O(n^{2/3})$, then $ex(n, m, \{C_4, C_6\}) = O(n)$. This conjecture was disproven by several papers and the current best-known bounds for this problem are

$$c_1 n^{1+\frac{1}{15}} \leq ex(n, n^{2/3}, \{C_4, C_6\}) \leq c_2 n^{1+1/9}$$

for some constants c_1, c_2 . A consequence of our work here proves that

$$ex(n, n^{2/3}, \{C_4, \theta_{3,4}\}) = \Theta(n^{1+1/9}).$$

More generally, for each integer $t \geq 2$, we establish that

$$ex(n, n^{\frac{t+2}{2t+1}}, \{C_4, \theta_{3,t}\}) = \Theta(n^{1+\frac{1}{2t+1}})$$

by demonstrating that subsets of points $S \subseteq \text{PG}(n, q)$ for which no $t + 1$ points lie on a line give rise to $\{C_4, \theta_{3,t}\}$ -free graphs, where $\text{PG}(n, q)$ is the projective space of dimension n over the finite field of q elements.

1 Introduction

Let m, n be positive integers and \mathcal{F} be a family of graphs. The bipartite Turán number $ex(m, n, \mathcal{F})$, is the maximum number of edges in a bipartite graph whose part sizes are m and n and such that it contains no graph in \mathcal{F} as a subgraph. The function $ex(m, n, \mathcal{F})$

^{*}Hacettepe University, Graduate School of Science and Engineering, Beytepe 06800 Ankara, Türkiye. E-mail: baranduzgun@hacettepe.edu.tr.

[†]Institute of Mathematics and Statistics, University of Tartu, 51009 Tartu, Estonia. E-mail: ago-erik.riet@ut.ee. This work was supported by the Estonian Research Council grant PRG2531.

[‡]Department of Mathematics: Analysis, Logic and Discrete Mathematics, Ghent University, 9000 Ghent, Belgium. E-mail: vlad.taranchuk@ugent.be.

has been studied extensively for many different sets \mathcal{F} , but many questions remain. See the well-known survey by Füredi and Simonovits [4] for a history of the work done on these types of problems.

One of the most notorious cases of determining $ex(n, m, \mathcal{F})$ is when $\mathcal{F} = C_{2k}$ for some positive integer k . When $m = n$, the order of magnitude of $ex(n, n, \{C_{2k}\})$ is only known for $k = 2, 3, 5$ [4], which coincides with the existence of special finite geometries called generalized polygons. When $m = n^a$ for any $a < 1$, and we allow $n \rightarrow \infty$, even less is known. The best constructions in this aforementioned case which yield C_4 -free graphs come from the point-block incidence graphs of 2-designs.

In 1979, Erdős conjectured that when $m = O(n^{2/3})$, then $ex(m, n, \{C_4, C_6\}) = O(n)$ [3]. This was disproven first by de Caen and Székely [1] who constructed an infinite family of graphs which yielded

$$cn^{1+\frac{1}{57}+o(1)} \leq ex(n, n^{2/3}, \{C_4, C_6\})$$

for a constant c . Later this lower bound was improved by Lazebnik, Ustimenko and Woldar [6] who constructed an infinite family of graphs which yielded

$$cn^{1+\frac{1}{15}} \leq ex(n, n^{2/3}, \{C_4, C_6\}).$$

for some constant c . It is worth mentioning that the class of graphs which is constructed in [6] is very much related to the (q, q^2) -generalized quadrangle. In fact, it seems to have escaped the graph theory community that the existence of such quadrangles also yields the bound obtained in [6] by taking an induced subgraph. The bounds on $ex(n, n^{2/3}, \{C_4, C_6\})$ have not budged in 30 years. In this paper, we establish results which suggest that the upper bound for this problem may be closer to the truth.

A common generalization of the cycle C_{2k} , is the theta graph $\theta_{t,k}$ which is the graph consisting of two vertices joined by t internally vertex-disjoint k -edge paths. While the lower and upper bounds for $ex(n, n, \{C_{2k}\})$ do not have matching orders of magnitude for all $k \neq 2, 3, 5$ [4], it is known that for each k , there exists a (relatively large) constant $t = t(k)$ such that

$$ex(n, n, \theta_{k,t}) = \Theta(n^{1+\frac{1}{k}}).$$

which is the same order of magnitude as the upper bound for $ex(n, n, C_{2k})$ [2]. In [2], Conlon uses a random algebraic method to construct infinite families of graphs not containing $\theta_{k,t}$, where t is fixed, but large relative to k . Therefore, explicit constructions yielding the same bounds and with a smaller t are of great interest.

There has also been some recent progress on determining $ex(n, m, \theta_{t,k})$. Jiang, Ma, and Yepremyan [5] proved that there exists a constant $c = c(k, t)$ such that

$$ex(m, n, \{\theta_{k,t}\}) \leq \begin{cases} c[(mn)^{\frac{k+1}{2k}+m+n}] & \text{if } k \text{ is odd} \\ c[(mn)^{\frac{k+2}{2k}n^{\frac{1}{2}}+m+n}] & \text{if } k \text{ is even} \end{cases}$$

and when $k = 3$, they obtained that

$$ex(m, n, \{\theta_{3,t}\}) \leq 144t^3((mn)^{\frac{2}{3}} + m + n).$$

Theodorakopoulos [8] extended the random algebraic methods used in [2] to prove that for each odd positive integer k and rational number a satisfying $\frac{k-1}{k+1} < a < 1$, there exists a constant $c = c(k)$ such that

$$ex(n, n^a, \{\theta_{k,c_k}\}) = \Theta((n^{1+a})^{\frac{k+1}{2k}})$$

Here we prove that subsets of points S of the projective space $\text{PG}(n, q)$ satisfying the condition that no $t + 1$ points of S lie on a line, produce $\{C_4, \theta_{3,t}\}$ -free graphs via their linear representations. In particular, this implies the following theorem.

Theorem 1.1. *Let q be a prime power and t, n be positive integers. Suppose that S is a subset of points of $\text{PG}(n, q)$ satisfying the condition that no $t + 1$ points of S lie on a common line. Then*

$$|S|q^{n+1} \leq ex(q^{n+1}, |S|q^n, \{C_4, \theta_{3,t}\}).$$

We remark that such sets with many elements are known to exist as shown in Lin and Wolf [7]. Consequently, we obtain our result.

Theorem 1.2. *Let $t \geq 2$ be a positive integer. Then*

$$ex(n, n^{\frac{t+2}{2t+1}}, \{C_4, \theta_{3,t}\}) = \Theta(n^{1+\frac{1}{2t+1}}).$$

Finally, our theorem implies the following corollary which suggests that the true value of $ex(n, n^{2/3}, \{C_4, C_6\})$ may be closer to the best-known upper bound.

Corollary 1.3. *We have*

$$ex(n, n^{2/3}, \{C_4, \theta_{3,4}\}) = \Theta(n^{1+\frac{1}{9}}).$$

2 Linear Representations of Point Sets

Let q be a prime power, let \mathbb{F}_q^{n+1} denote the vector space of dimension $n + 1$ over the finite field \mathbb{F}_q , and let $\text{PG}(n, q)$ be the corresponding projective space.

Definition 2.1. *Let S be a set of points of $\text{PG}(n, q)$ and embed $\text{PG}(n, q)$ as a hyperplane into $\text{PG}(n + 1, q)$. A linear representation of S is the geometry whose points are all the points in $\text{PG}(n + 1, q) \setminus \text{PG}(n, q)$ and the lines are all the lines of $\text{PG}(n + 1, q)$ which intersect $\text{PG}(n, q)$ in precisely one point, namely a point of S .*

Remark: Observe that if two lines in $\text{PG}(n+1, q)$ not contained in $\text{PG}(n, q)$ intersect in a point of S , then they are parallel in the linear representation of S .

From this geometry we may build its point-line incidence graph. This graph is bipartite with bipartition classes given by the points and the lines of the geometry. A point will be adjacent to a line in the graph if they are incident in the geometry, i.e. the point is on the line. Denote this graph by $\Gamma_{S,n,q}$. It can easily be verified that $\Gamma_{S,n,q}$ will have the following properties:

1. There are q^{n+1} point vertices, each of degree $|S|$.
2. There are $|S|q^n$ line vertices, each of degree q .

Theorem 2.2. *Let S be a subset of points of $\text{PG}(n, q)$ such that any line in $\text{PG}(n, q)$ intersects S in at most t points. Then $\Gamma_{S,n,q}$ is $\{C_4, \theta_{3,t}\}$ -free.*

Proof. Note that by construction, the linear representation of S is a geometry in which any two lines intersect in at most one point and any two points lie on at most one line. Thus, the graph $\Gamma_{S,n,q}$ is necessarily C_4 -free.

Suppose that $\Gamma_{S,n,q}$ contains a $\theta_{3,t}$. This implies there exists a point vertex r and line vertex ℓ between which there are t vertex-disjoint paths of length 3. Geometrically, this implies that the linear representation of S contains a configuration consisting of a point r , t lines which contain r , call them m_1, m_2, \dots, m_t , all of which intersect the line ℓ in our geometry, and such that r is not on ℓ . Note that the set of lines $\ell, m_1, m_2, \dots, m_t$ all pairwise intersect, and so no two can be parallel. Thus, in $\text{PG}(n+1, q)$, each of the lines ℓ, m_1, \dots, m_t contains a distinct point in S .

Note that all of the lines of this configuration lie in a common plane, Π , the plane in $\text{PG}(n+1, q)$ spanned by r and ℓ . Since Π is not contained in the hyperplane $\text{PG}(n, q)$, it intersects $\text{PG}(n, q)$ in a line, call it ℓ_∞ . But this implies that there is a set of $t+1$ points in S (one for each line ℓ, m_1, \dots, m_t) which lie on ℓ_∞ . Since we assumed any line in $\text{PG}(n, q)$ intersects S in at most t points, this is a contradiction. Thus $\Gamma_{S,n,q}$ is also $\theta_{3,t}$ -free. \square

In [7], the authors obtained the following result. For completeness, we give an explicit construction of such a set for all integers $t \geq 2$ and prime powers $q > t$.

Theorem 2.3. *Let q be a prime power and $t > q$ be a positive integer. Then there exists a subset S of points of $\text{PG}(t+1, q)$ of size q^t such that no $t+1$ points of S lie on a line.*

It is well-known that the field \mathbb{F}_{q^t} can be viewed as a vector space over \mathbb{F}_q . Fix any basis, $\alpha_1, \dots, \alpha_t$ for \mathbb{F}_{q^t} over \mathbb{F}_q . For each x in \mathbb{F}_{q^t} , denote by $x|_q$ the vector of the field reduced elements of x . That is, $x|_q = (x_1\alpha_1 + \dots + x_t\alpha_t)|_q = (x_1, \dots, x_t)$. Denote by N the norm function from \mathbb{F}_{q^t} to \mathbb{F}_q , i.e. $N(x) = x^{(q^t-1)/(q-1)} = x \cdot x^q \cdot x^{q^2} \cdot \dots \cdot x^{q^{t-1}}$.

Theorem 2.4. *Let $t \geq 2$ be a positive integer and $q > t$ be a prime power. Then the set*

$$S = \{(1, x|_q, N(x)) : x \in \mathbb{F}_{q^t}\}$$

as a subset of points of $\text{PG}(t+1, q)$ contains no $t+1$ points on a line.

Proof. We will omit the notation of field reduction to avoid getting bogged down in notation. Take any two vectors in S , call them $(1, x, N(x))$ and $(1, y, N(y))$. These two points lie on some line in $\text{PG}(t+1, q)$. We will count how many other points $(1, z, N(z))$ in S can lie on this same line. If $(1, z, N(z))$ lies on the same line, it implies that the matrix

$$\begin{bmatrix} 1 & 1 & 1 \\ x & y & z \\ N(x) & N(y) & N(z) \end{bmatrix}$$

has an element in its null space $v = [a, b, c]$ with each $a, b, c \in \mathbb{F}_q^*$. Without loss of generality, we assume $c = -1$. Thus we obtain a linear system

$$\begin{aligned} a + b &= 1, \\ ax + by &= z, \\ aN(x) + bN(y) &= N(z). \end{aligned}$$

This system implies that

$$aN(x) + (1 - a)N(y) = N(ax + (1 - a)y) \tag{1}$$

Since x and y are fixed, that leaves a as the only variable. Since $a \in \mathbb{F}_q$, then $a^q = a$ and so expanding (1), we obtain a polynomial equation in a of degree t . In particular, the largest degree term a^t has coefficient $N(x - y) \neq 0$. Observe that $a = 0$ and $a = 1$ are both solutions to (1), which are not valid choices of a for us (since $a, b \neq 0$). Consequently, there are at most $t - 2$ valid choices of a which solve (1). Each a determines a unique $z = ax + (1 - a)y$, so any line contains at most t points of S . \square

Consequently, the linear representation of this set produces a biregular, unbalanced bipartite graph with bipartition class sizes $m = q^{t+2}$, $n = q^{2t+1}$, and q^{2t+2} edges which is $\{C_4, \theta_{3,t}\}$ -free. By an application of Bertrand's postulate, we obtain that for each t , there exists a constant c_t such that

$$c_t n^{1 + \frac{1}{2t+1}} \leq ex(n, n^{\frac{t+2}{2t+1}}, \{C_4, \theta_{3,t}\}).$$

For each such t , the order of magnitude matches the upper bounds given in [5]. Thus we prove Theorem 1.2

$$ex(n, n^{\frac{t+2}{2t+1}}, \{C_4, \theta_{3,t}\}) = \Theta(n^{1 + \frac{1}{2t+1}}).$$

References

- [1] D. de Caen and L. A. Székely, “The maximum size of 4- and 6-cycle free bipartite graphs on m, n vertices”, in *Graphs, Sets and Numbers*, Proc. of the Conference Dedicated to the 60th Birthdays of A. Hajnal and Vera T. Sós, Budapest, (1991).
- [2] D. Conlon, “Graphs with few paths of prescribed length between any two vertices”, *Bulletin of the London Mathematical Society*, 51(6), (2019), 1015–1021.
- [3] P. Erdős, “Some old and new problems in various branches of combinatorics”, *Proceedings of the Tenth Southeastern Conference on Combinatorics, Graph Theory and Computing*, Boca Raton, Vol. 1, Congressus Numerantium 23, (1979), 19–38.
- [4] Z. Füredi and M. Simonovits, “The History of Degenerate (Bipartite) Extremal Graph Problems”, *Erdős Centennial Bolyai Society Mathematical Studies*, 25, (2013).
- [5] T. Jiang, J. Ma, and L. Yepremyan, “On Turán exponents of bipartite graphs”, *Combinatorics, Probability and Computing*, 31(2), (2022), 333–344.
- [6] F. Lazebnik, V. A. Ustimenko, and A. J. Woldar, “New constructions of bipartite graphs on m, n vertices with many edges and without small cycles”, *Journal of Combinatorial Theory Series B*, 61(1), (1991), 111–117.
- [7] Y. Lin and J. Wolf, “On subsets of \mathbb{F}_q^n containing no k -term progressions”, *European Journal of Combinatorics*, 31(5), (2010), 1391–1403.
- [8] S. Theodorakopoulos, “Some sharp lower bounds for the bipartite Turán number of theta graphs”, , arXiv: 2405.02864, (2024).