A NON-DEGENERACY THEOREM FOR INTERACTING ELECTRONS IN ONE DIMENSION

THIAGO CARVALHO CORSO

ABSTRACT. In this paper, we show that the ground-state of many-body Schrödinger operators for electrons in one dimension is non-degenerate. More precisely, we consider Schrödinger operators of the form

$$H_N(v,w) = -\Delta + \sum_{i\neq j}^N w(x_i,x_j) + \sum_{j=1}^N v(x_i) \quad \text{acting on } \wedge^N \mathrm{L}^2([0,1]),$$

where the external and interaction potentials v and w belong to a large class of distributions. In this setting, we show that the ground-state of the system with Fermi statistics and local boundary conditions is non-degenerate and does not vanish on a set of positive measure. In the case of periodic and anti-periodic (or more general non-local) boundary conditions, we show that the same result holds whenever the number of particles is odd and even, respectively. This non-degeneracy result seems to be new even for regular potentials v and w. As an immediate application of this result, we prove eigenvalue inequalities and the strong unique continuation property for eigenfunctions of the single-particle one-dimensional operators $h(v) = -\Delta + v$. In addition, we prove strict inequalities between the lowest eigenvalues of different self-adjoint realizations of $H_N(v, w)$.

Contents

1.	Introduction	1
2.	Main Results	2
3.	Background on Sobolev Spaces	6
4.	Schrödinger operators with distributional potentials	9
5.	Non-degeneracy of the ground-state	11
6.	Eigenvalue inequalities	17
Ар	pendix A. Quadratic forms and self-adjoint extensions	22
References		24

1. INTRODUCTION

The primary goal of this paper is to study the spectral properties of Schrödinger operators of the form

$$H_N(v,w) = -\Delta + \sum_{i\neq j}^N w(x_i, x_j) + \sum_{j=1}^N v(x_j)$$

acting on the space of antisymmetric wave-functions $\wedge^{N} L^{2}(I)$, where I = (0, 1) (or \mathbb{R}), and v and w belongs to suitable class of distributions. Such operators are used to describe interacting electrons living in one-dimensional space and are therefore ubiquitous in quantum mechanics.

In the mathematical literature, Schrödinger operators with distributional potentials have been extensively studied. Early contributions include [RS72, AGHK84, Bra85, AGHH88, Her89], while more recent works can be found in [AKoo, EGNT15, LR15, EP16, BFK+17]. In particular, special attention has been devoted to the case of one-dimensional Sturm-Liouville operators [Kur97, SS99, AHM08, Dav13, EGNT13a, EGNT13b, MK14, Gul19]. However, most (if not all) of these works focus on the case of what could be called "single-particle electronic systems", i.e., where $H = -\Delta + V$ acts on $L^2(\Omega)$ for some $\Omega \subset \mathbb{R}^d$ instead of the antisymmetric L^2 -space. Of course, the spectral properties of the many-particle non-interacting Hamiltonian

Date: March 25, 2025.

²⁰²⁰ Mathematics Subject Classification. Primary: 81Q10 Secondary: 35J10, 81V74.

Key words and phrases. Spectral theory of Schrödinger operators, eigenvalue inequalities, distributional potentials, many-body quantum systems, electronic systems, Density functional theory, unique continuation.

Funding information: DFG - Project-ID 442047500 - SFB 1481.

^{©2025} by the authors. Faithful reproduction of this article, in its entirety, by any means is permitted for noncommercial purposes.

 $H_N(v, 0)$ acting on $\wedge^N L^2(I)$ are determined be the spectral properties of the single-particle operator $h(v) = -\Delta + v$ acting on $L^2(I)$. Unfortunately, this is no longer true for interacting systems, which is arguably the most interesting case in applications.

In fact, our main motivation for studying the interacting operator $H_N(v, w)$ stems from recent advances [SPR+24, Cor25] towards establishing a mathematically rigorous formulation of Kohn-Sham Density Functional Theory (DFT), a highly successful and widely used approach to study the electronic structure of materials. More precisely, the class of distributional potentials studied here appeared in recent developments [SPR+24] concerning the *v*-representability problem, which, roughly speaking, consists in characterizing the set of all possible single-particle ground state densities of operators of the form $H_N(v, w)$ for a given class of potentials *v*. While a detailed discussion on the foundation of DFT and the *v*-representability problem goes beyond the scope of the current paper, we emphasize that the results obtained here will be used to provide a completely rigorous formulation of Kohn-Sham DFT for electrons in one-dimension in a follow-up article.

In this work, we focus instead on the ground-state properties of $H_N(v, w)$. More precisely, our main contributions can be stated as follows:

- (i) We prove that the ground-state of self-adjoint realizations of $H_N(v, w)$ with local boundary conditions (BCs) is non-degenerate and almost everywhere non-vanishing. In particular, this applies to the Dirichlet and Neumann realizations of $H_N(v, w)$.
- (ii) In the case of non-local boundary conditions such as periodic and anti-periodic BCs, we prove that the ground-state is non-degenerate and almost everywhere non-vanshing whenever the number of particles *N* is odd and even, respectively.
- (iii) As an application of these non-degeneracy results, we obtain eigenvalue inequalities and prove the strong unique continuation property for eigenfunctions of the one-dimensional single-particle operator $h(v) = -\Delta + v$.
- (iv) In addition, we obtain strict inequalities between the lowest eigenvalues (or ground-state energies) of different self-adjoint realizations of $H_N(v, w)$.

2. MAIN RESULTS

In this section, we state our main results precisely. We then outline the key steps in the proofs and how these steps are organized throughout the paper.

2.1. Notation. We start with some notation. Throughout this paper, we let I = (0, 1) be the open unit interval and set $I_N := (0, 1)^N$ for any $N \in \mathbb{N}$.

We denote by $H^1(I)$ the Sobolev space of functions $f \in L^2(I)$ with weak derivative $\partial_x f \in L^2(I)$. More generally, for $1 \le p \le \infty$ and open $\Omega \subset \mathbb{R}^d$, we denote by $W^{1,p}(\Omega)$ the usual Sobolev spaces of functions in $L^p(\Omega)$ with weak gradient in $L^p(\Omega)$, and by $W^{-1,q}(\Omega)$, where 1/q + 1/p = 1, the dual space of $W^{1,p}(\Omega)$. In addition, we denote by $H^{1/2}(\partial\Omega)$ the standard 1/2-Sobolev (or Besov) space along the boundary $\partial\Omega$.

For a given closed subspace $L \subset H^{1/2}(\partial \Omega)$, we denote by $H^1_L(\Omega)$ the Sobolev spaces of functions with trace on L, i.e.,

$$\mathrm{H}^{1}_{L}(\Omega) \coloneqq \{ f \in \mathrm{H}^{1}(\Omega) : \gamma f \in L \},\$$

where $\gamma : H^1(\Omega) \to H^{1/2}(\partial \Omega)$ is the Dirichlet (boundary) trace operator.

Next, we define $\mathcal V$ as the following space of generalized external potentials:

$$\mathcal{V} \coloneqq \{ v \in \mathrm{H}^{-1}(I) : v(\varphi) \in \mathbb{R} \quad \text{for any real-valued } \varphi \in \mathrm{H}^{1}(I) \}$$

Similarly, we define \mathcal{W} as the following space of generalized interaction potentials:

 $\mathcal{W} \coloneqq \{ w \in W^{-1,q}(I_2) : \text{for some } q > 2 \text{ and } w(\varphi) \in \mathbb{R} \text{ for any real-valued } \varphi \in W^{1,p}(I_2) \}.$

For $N \in \mathbb{N}$, we denote by \mathcal{H}_N the usual space of electronic wave-functions, i.e., the antisymmetric *N*-fold tensor product

$$\mathcal{H}_N = \bigwedge^N \mathrm{L}^2(I).$$

For $v \in \mathcal{V}$, $w \in \mathcal{W}$, and $L \subset \mathrm{H}^{1/2}(\partial I_N)$ as above, we denote by $H_N^L(v, w)$ the *N*-particle Hamiltonian

$$H_N^L(v,w) = -\Delta + \sum_{i \neq j}^N w(x_i, x_j) + \sum_{j=1}^N v(x_i) \quad \text{acting on } \mathcal{H}_N \text{ with boundary conditions in } L.$$

More precisely, $H_N^L(v, w)$ is the unique self-adjoint operator associated to the sesquilinear form

$$a_{v,w}(\Psi,\Phi) \coloneqq \int_{I_N} \overline{\nabla \Psi(x_1,...,x_N)} \cdot \nabla \Phi(x_1,...,x_N) dx_1...dx_N + v\left(\rho_{\Psi,\Phi}\right) + w(\rho_{2,\Psi,\Phi})$$
(2.1)

with form domain

$$Q_N^L \coloneqq \mathcal{H}_N \cap \mathrm{H}^1_L(I_N),$$

where $\rho_{\Psi,\Phi}$ is the overlapping single-particle density

$$\rho_{\Psi,\Phi}(x) \coloneqq N \int_{I_{N-1}} \overline{\Psi(x, x_2, ..., x_N)} \Phi(x, x_2, ..., x_N) dx_2 ... dx_N,$$
(2.2)

and $\rho_{2,\Psi,\Phi}$ is the overlapping pair density

$$\rho_{2,\Psi,\Phi}(x,y) \coloneqq N(N-1) \int_{I_{N-2}} \overline{\Psi(x,y,x_3,...,x_N)} \Phi(x,y,x_3,...,x_N) dx_3...dx_N.$$
(2.3)

2.2. Main results. As a first result, we show that the ground-state of self-adjoint realizations of $H_N(v, w)$ with local boundary conditions is non-degenerate. Precisely, let $\Gamma \subset \partial \Omega$ and define

$$H^{1}(I_{N};\Gamma) = \{\Psi \in H^{1}(I_{N}) : (\gamma \Psi)|_{\Gamma} = 0\}.$$

Then the following holds.

Theorem 2.1 (Non-degeneracy theorem with local BCs). Let $v \in \mathcal{V}$, $w \in \mathcal{W}$, $N \in \mathbb{N}$ and $\Gamma \subset \partial \Omega$. Then the operator $H_N(v, w; \Gamma)$ defined as the self-adjoint realization of (2.1) with form domain $H^1(I_N; \Gamma) \cap \mathcal{H}_N$ has a unique ground-state Ψ and $\Psi \neq 0$ almost everywhere.

Remark 2.2 (Unique continuation). The fact that Ψ does not vanish on a set of positive Lebesgue measure is known as the strong unique continuation property or principle (UCP). While the UCP is known to hold for distributional solutions of the Schrödinger equation with a large class of multiplicative potentials, see, e.g. [JK85, Sog90, Wol93, KT01, H0r07, Gar18], the author is not aware of similar results for manybody Schrödinger operators with distributional potentials. In fact, many of the difficulties encountered throughout our proofs can be traced back to this lack of unique continuation results. Interestingly though, Theorem 2.1 allow us to establish a unique continuation result for the single-particle operator $h(v) = -\Delta + v$; see Theorem 2.6.

The next result shows that Theorem 2.1 can be extended to the case of periodic and anti-periodic boundary conditions. In this case, however, we need an additional assumption on the number of particles $N \in \mathbb{N}$. To state this result precisely, let us define, for any $\alpha \in \mathbb{R} \setminus \{0\}$ the space

$$\mathrm{H}^{1}_{\alpha}(I_{N}) \coloneqq \left\{ \Psi \in \mathrm{H}^{1}(I_{N}) : (\gamma \Psi)|_{I_{k-1} \times \{0\} \times I_{N-k}} - \alpha(\gamma \Psi)|_{I_{k-1} \times \{1\} \times I_{N-k}} = 0 \quad \text{for any } 1 \leq k \leq N \right\}.$$

We say that these spaces have non-local boundary conditions as the boundary values along any face of ∂I_N depend on the values along the opposite face¹. Note that $\alpha = 1$ corresponds to periodic BCs and $\alpha = -1$ to antiperiodic BCs. For such spaces, the following result holds.

Theorem 2.3 (Non-degeneracy theorem with non-local BCs). Let $\alpha \in \mathbb{R} \setminus \{0\}$ and suppose that

$$\alpha(-1)^{N-1} > 0. \tag{2.4}$$

Then the self-adjoint realization $H_N^{\alpha}(v, w)$ of (2.1) with form domain $H_{\alpha}^1(I_N) \cap \mathcal{H}_N$ has a unique ground state Ψ and $\Psi \neq 0$ almost everywhere.

Remark 2.4 (On the condition on the number of particles). Theorem 2.3 shows that periodic systems are non-degenerate for an odd number of particles, while anti-periodic systems are non-degenerate for an even number of particles. This result is optimal in the sense that there are explicit examples where the ground-state is non-degenerate if and only if condition (2.4) holds. For instance, the simplest example is the free periodic and anti-periodic Laplacians, where the ground-state is non-degenerate if and only if N is odd respectively even. Moreover, one can also construct examples of periodic systems where the ground-state is non-degenerate for any given number of particles N; thus an improvement of the form "the system is at least two-fold degenerate for even/odd number of particles" is also not possible.

Remark 2.5 (Further extensions). Theorem 2.1 can also be extended in the following directions:

¹We exclude the case $\alpha = 0$ for two reasons. First, this case corresponds to local boundary conditions and are therefore covered by Theorem 2.1. Second, the notation $H_0^1(I_N)$ is reserved to the case of Dirichlet boundary conditions along the whole boundary.

T. CARVALHO CORSO

- (1) (Real line) Theorem 2.1 can be extended to the whole real line $I = \mathbb{R}$ (or half-line). In this case, the existence of a ground-state is not guaranteed and the statement should be changed to "if a ground-state exists" then all of the conclusions of Theorem 2.1 holds.
- (2) (Many-body interactions) We can allow a larger class of distributional interaction potentials w. In fact, Theorem 2.1 holds for *M*-body interactions w_M with arbitrary $M \in \mathbb{N}$, provided that w_M lies in suitable dual Sobolev spaces (see Section 4.3). Notably, the case of two-body 3D Coulomb interactions, which could be seen as a "six-body" interaction of the form

$$w_6(x_1, x_2, x_3, y_1, y_2, y_3) = \left(\sum_{j=1}^3 (x_j - y_j)^2\right)^{-\frac{1}{2}}$$

is also admissible. In particular, one can show that the ground-state of the 3D many-particle system with Coulomb interaction, restricted to maximally antisymmetric wave-functions, i.e., wave-functions that are antisymmetric with respect to the exchange of all possible one-dimensional coordinates, is also non-degenerate.

(3) (Elliptic operators) Theorems 2.1 and 2.3 also hold with the Laplacian replaced by more general elliptic operators

$$\mathcal{L}_{a}\Psi \coloneqq \sum_{i,j=1}^{N} \partial_{x_{i}}\left(a_{ij}(x)\partial_{x_{j}}\Psi\right), \quad \text{with } a_{ij}(x) \in \mathrm{L}^{\infty}(I_{N};\mathbb{R}).$$

The only condition needed for our proof is that weak subsolutions $\mathcal{L}_a u \leq 0$ satisfy the strong maximum principle.

(4) (More general boundary conditions) We can further extend Theorem 2.1 to a larger class of selfadjoint realizations of $H_N(v, w)$. Roughly speaking, the maximal class of spaces for which our proof works is given by all $H^1_L(I_N)$ where the restricted space

$$\{\Psi|_{S_N} : \Psi \in H^1_L(I_N) \cap \mathcal{H}_N\}$$
 where $S_N = \{(x_1, ..., x_N) \in I_N : x_1 < x_2 ... < x_N\}$

is invariant under taking the real and positive part, see Theorem 5.6. In particular, Theorem 2.3 still holds with the following mixed space of non-local and local boundary conditions

$$\mathrm{H}^{1}_{\alpha}(I_{N};\Gamma) = \mathrm{H}^{1}_{\alpha}(I_{N}) \cap \mathrm{H}^{1}(I_{N};\Gamma), \qquad (2.5)$$

for $\alpha \in \mathbb{R} \setminus \{0\}$ satisfying (2.4).

Interestingly, Theorems 2.1 and 2.3 can be applied to obtain information on the whole spectrum of singleparticle operators. More precisely, let $v \in \mathcal{V}$, $L \subset \mathbb{C}^2$ be a (closed) subspace, and denote by $h_L(v) = -\Delta_L + v$ the self-adjoint operator associated to the form

$$a_{v}(\psi,\psi) = \int_{I} |\nabla\psi(x)|^{2} dx + v(|\psi|^{2}) \quad \text{with form domain } H^{1}_{L}(I) = \{\psi \in H^{1}(I) : (\psi(0),\psi(1)) \in L\}.$$
(2.6)

Then by applying Theorems 2.1 and 2.3 to self-adjoint realizations of the non-interacting operator $H_N(v, 0)$, we obtain the following result.

Theorem 2.6 (Single-particle operators). Let $v \in V$ and $L = \{\beta = (\beta_0, \beta_1) \in \mathbb{C}^2 : \beta_0 - \alpha\beta_1 = 0\}$ for some $\alpha \in \mathbb{R} \setminus \{0\}$. Then $h_L(v)$ has discrete spectrum, all eigenfunctions are almost everywhere non-vanishing, and the eigenvalues $\{\lambda_k\}_{k \in \mathbb{N}}$ ordered in non-decreasing order and counting multiplicity satisfy

$$\begin{cases} \lambda_{2k-1} < \lambda_{2k}, & \text{if } \alpha \ge 0, \\ \lambda_{2k} < \lambda_{2k+1}, & \text{if } \alpha \le 0, \end{cases}, \quad \text{for any } k \in \mathbb{N}.$$

$$(2.7)$$

Moreover, in the case where L is one of the following sets

$$\{0\} \times \mathbb{C}, \quad \mathbb{C} \times \{0\}, \quad \{(0,0)\}, \quad or \quad \mathbb{C}^2,$$
 (2.8)

the same holds but all the eigenvalues are simple.

Remark (Self-adjont realizations). We remark that all local self-adjoint realizations of (2.6) on $C_c^{\infty}(I)$ are of the form $h_L(v)$ for some closed subspace $L \subset \mathbb{C}^2$ (see Propostion 3.9). Hence, Theorem 2.6 provides information on the spectrum of all self-adjoint extensions of $h_L(v)$ in $C_c^{\infty}(I)$ whose form domain is closed under complex conjugation.

5

Remark (Sturm-Liouville theory). The result from Theorem 2.6 in the case of local boundary conditions (2.8) is, at least for regular potentials *v*, well-known in the theory of Sturm-Liouville operators, see, e.g., [Zet21, Theorem 1.9.2]. Nevertheless, the approach presented here seems to be completely different from previous proofs and has the important advantage that it works in the interacting case.

Next, we present a monotonicity result for the ground-state energy of $H_N(v, w)$. This result shows that the ground-state energy is strictly monotone with respect to enlarging the Dirichlet set of the form domain of $H_N(v, w)$. Precisely, let us denote by Γ_N the symmetrization of the set $\Gamma \subset \partial I_N$, i.e.,

$$\Gamma_N := \bigcup_{\sigma \in \mathcal{P}_N} \sigma(\Gamma)$$
 where $\sigma(x_1, ..., x_N) = (x_{\sigma(1)}, ..., x_{\sigma(N)}),$

and \mathcal{P}_N is the set of permutations of $\{1, ..., N\}$. Then we have the following result.

Theorem 2.7 (Monotonicity with respect to the Dirichlet set). Let $\Gamma \subset \Gamma' \subset \partial I_N$ be such that $\Gamma'_N \setminus \Gamma_N$ has non-empty (relative) interior in ∂I_N . Let $\lambda_1(\Gamma)$ and $\lambda_1(\Gamma')$ be respectively the lowest eigenvalues of $H_N(v, w; \Gamma)$ and $H_N(v, w; \Gamma')$. Then

$$\lambda_1(\Gamma) < \lambda_1(\Gamma').$$

Moreover, if $\alpha \in \mathbb{R} \setminus \{0\}$ satisfies (2.4) and $\lambda_1(\Gamma)$ and $\lambda_1(\Gamma')$ denote respectively the lowest eigenvalues of the self-adjoint realizations with form domain $H^1_{\alpha}(I_N; \Gamma) \cap \mathcal{H}_N$ and $H^1_{\alpha}(I_N; \Gamma') \cap \mathcal{H}_N$, then the same result holds.

Remark (Strict inequality). The important point in Theorem 2.7 is the strict inequality, since the weaker inequality \leq trivially follows from the variational principle.

Remark (Topological assumption). Our main tool to establish Theorem 2.7 is a weak unique continuation result along the boundary, namely Theorem 6.1. As this result applies only to relatively open subsets of the boundary, the assumption that $\Gamma'_N \setminus \Gamma_N$ has non-empty interior is crucial for our proof. Lifting this assumption seems like a challenging problem.

2.3. **Outline of the proofs**. We now discuss the key steps and main challenges in the proofs of our main results.

The proof of Theorems 2.1 and 2.3 is carried out in Section 5 and relies on two main steps. The first step consists in proving a Perron-Frobenius (PF) theorem for Schrödinger operators of the form $-\Delta + V$ for a large class of distributional potentials V (see Theorem 5.3). In other words, we show that the ground-state of such operators is non-degenerate and strictly positive almost everywhere. This step relies on well-known semigroup techniques and a perturbative argument borrowed from Reed and Simon [RS78][Section XII.12] (see also [GJ70, FS75, Ges84] for similar arguments). Combining this result with a density result for dual Sobolev spaces (Lemma 3.11), we can then establish a PF theorem for Schrödinger operators of the form $H_N(v, w) = -\Delta + \sum_j v_j + \sum_{j \neq i} w_{i,j}$ under a suitable positivity assumption on their form domain. Unfortunately, these results only apply to the operator without Fermi statistics, i.e., without antisymmetry restrictions on the wave-function. Indeed, whenever antisymmetry is present the wave-function must have both positive and negative values and can not be almost everywhere strictly positive. In particular, as warned in [RS78, page 207], PF results cannot be applied to electronic systems.

It turns out, however, that this is not entirely true for the case of systems with antisymmetry with respect to exchange of one-dimensional coordinates. Indeed, in the second and key step of our proof, we show that the operator $H_N(v, w)$, acting on antisymmetric functions in I_N , is unitarily equivalent to a reduced Schrödinger operator acting on functions on the simplex $S_N := \{(x_1, ..., x_N) \in (0, 1)^N : x_1 < x_2 ... < x_N\}$, but without symmetry constraints. This reduction is based on the rather simple but surprisingly useful observation that the box I_N can be tilled by reflections of the simplex S_N . This reduction then allow us to apply the aforementioned PF theorem to establish the non-degeneracy and the strong unique continuation property of the ground-state for the electronic Hamiltonian $H_N(v, w)$. At this step, condition (2.4) can be shown to be equivalent to positivity of the form domain of the reduced operator and is therefore crucial for our proof. This step is carried out in Section 5.2.

In Section 5.4 we present the proof of Theorem 2.6. This result is an immediate application of Theorems 2.1 and 2.3, and a technical lemma that connects the boundary conditions of the non-interacting operator $H_N(v, 0)$ to the boundary conditions of the single-particle operator $h(v) = -\Delta + v$.

The proof of Theorem 2.7 is inspired by ideas from previous works on similar eigenvalue inequalities [Filo5, GM09, BRS18, Roh21]. More precisely, the core idea of the proof is to apply the beautiful domain extension argument of Filonov [Filo5] to establish a unique continuation result along the boundary (cf. Theorem 6.1). Here, however, we face two fundamental problems when trying to apply this argument.

First, it relies on unique continuation results which are not available in the case of distributional potentials. Second, and perhaps more critical, is the fact that the non-local boundary conditions are such that, roughly speaking, the faces of I_N are "glued" to each other, and therefore, no extension argument is possible.

The first problem can be overcome by using the unique continuation property of the ground-state guaranteed by the PF theorem. Note that this property is only proven for the ground-state, which explains why we can only obtain inequalities for the lowest eigenvalue of $H_N(v, w)$. To deal with the second difficulty, we rely on a distributional definition of the Neumann trace (or normal derivative). While this definition is the same as in previous works, e.g., [GMo9] where the Laplacian with non-local (Robin) boundary conditions is considered, we emphasize that the situation here is more delicate for two reasons. First, the non-local boundary conditions here are imposed directly in the form domain of the operator, as opposed to adding a non-local form along the boundary as done in [GMo9]. Second, an eigenfunction Ψ of $H_N(v, w)$ has, in general, no more regularity than H¹. In particular, its Laplacian has no more regularity than H⁻¹, and therefore, the usual definition of the weak Neumann trace, which requires $\Delta \Psi$ to be at least in H^s for s > -1/2, see [GMo9, (3.17)], is not meaningful.

Fortunately, we can still show that, despite this lack of regularity, a meaningful sense can be given to the Neumann trace of eigenfunctions along any open subset of the boundary where they vanish; this result crucially relies on the locality of the operator $H_N(v, w)$, a property that indirectly underlies many of the arguments used here. We then obtain an explicitly formula for this weak Neumann trace that could be new and of independent interest; see Lemma 6.4. By combining this formula with a positivity property of the boundary conditions, which is inherited by the ground-state (see Theorem 5.6), we can show that the Neumann trace of the ground-state vanishes whenever its Dirichlet trace vanishes. This allows us to extend the form domain of $H_N(v, w)$ and carry out the extension argument of Filonov to complete the proof of the unique continuation along the boundary in Theorem 6.1. Theorem 2.7 then follows from standard arguments.

3. BACKGROUND ON SOBOLEV SPACES

In this section we present some well-known results about Sobolev spaces that will be useful throughout our proofs.

3.1. **Definitions and basic properties**. Let us start by recalling the precise definition of Sobolev spaces and setting-up some notation.

Throughout this section $\Omega \subset \mathbb{R}^d$ will be a bounded, open, and connected subset of \mathbb{R}^d with Lipschitz boundary. This means that, for any $x \in \partial\Omega$, there exists a neighborhood $x \in U \subset \mathbb{R}^d$ such that, up to a rigid motion (i.e., translation and rotation), $U \cap \partial\Omega$ is the graph of a Lipschitz function. On such domains, the Sobolev spaces can be defined as follows.

Definition 3.1 (Sobolev spaces). For $1 \le p \le \infty$, we denote by $W^{1,p}(\Omega)$, the space of (complex-valued) functions $f \in L^p(\Omega)$ with weak gradient $\nabla f \in L^p(\Omega; \mathbb{C}^n)$ endowed with the norm

$$\|f\|_{1,p}^{p} \coloneqq \|f\|_{L^{p}}^{p} + \|\nabla f\|_{L^{p}}^{p}.$$
(3.1)

Moreover, we denote by $W_0^{1,p}(\Omega)$ the closure of the space $C_c^{\infty}(\Omega)$ with respect to the W^{1,p}-norm.

Definition 3.2 (Dual Sobolev spaces). For $1 \le p \le \infty$, we denote by $W^{-1,p}(\Omega)$, the dual space of $W^{1,q}(\Omega)$, where *q* is the Hölder conjugate of *p* (i.e., 1/p + 1/q = 1). More precisely,

 $W^{-1,p}(\Omega) \coloneqq \{T : W^{1,q}(\Omega) \to \mathbb{C} \text{ linear and continuous}\}$

endowed with the norm

$$||T||_{-1,p} = \sup_{f \in W^{1,q} \setminus \{0\}} \frac{|T(f)|}{||f||_{1,q}}$$

Similarly, we denote by $W_0^{-1,p}(\Omega)$ respectively $W_L^{-1,p}(\Omega)$ the dual spaces of $W_0^{1,q}(\Omega)$ respectively $W_L^{1,q}(\Omega)$.

Remark 3.3 (Notation). For p = 2, we use the standard notation $H^1(\Omega)$ instead of $W^{1,2}(\Omega)$. Moreover, whenever clear from the context, we shall omit the domain Ω and simply write H^1 , $W^{1,p}$, $W^{1,p}_0$ and so on.

For later reference, let us recall the celebrated Gagliardo-Nirenberg-Sobolev (GNS) inequality. We refer, e.g., to [Leo17, Theorem 12.83] for a proof of the general $1 \le p, q \le \infty$ case, and to [BM18] for precise conditions on the validity of GNS for Sobolev spaces of fractional order.

Lemma 3.4 (GNS interpolation inequality). Let $\Omega \subset \mathbb{R}^d$ be a bounded open and connected domain with Lipschitz boundary. Then for any $f \in W^{1,p}(\Omega)$ with $2 \le p < \infty$ we have

$$||f||_{\mathcal{L}^{p}} \lesssim ||f||_{\mathcal{H}^{1}}^{1-\theta} ||f||_{\mathcal{L}^{2}}^{\theta} \quad where \quad \theta = \frac{d}{2} - \frac{d}{p}.$$
 (3.2)

For d = 1, the case $p = \infty$ is also allowed.

Another fundamental property of Sobolev functions on Lipschitz domains is the existence of a trace along the boundary. Let us recall this fact here as well. For a proof, see [Leo17, Section 18.4].

Theorem 3.5 (Trace of Sobolev functions). For any open, bounded, and Lipschitz $\Omega \subset \mathbb{R}^d$ with $d \ge 2$, there exists a unique continuous trace operator $\gamma : H^1(\Omega) \to H^{\frac{1}{2}}(\partial\Omega)$ satisfying $\gamma f = f|_{\partial\Omega}$ for $f \in H^1(\Omega) \cap C(\overline{\Omega})$. Moreover, the trace operator is surjective and there exists (infinitely many) right inverses, i.e., continuous maps $J : H^{\frac{1}{2}}(\partial\Omega) \to H^1(\Omega)$ such that $\gamma J f = f$ for any $f \in H^{\frac{1}{2}}(\partial\Omega)$.

Remark 3.6 (Fractional Sobolev space on the boundary). Although the precise definition of $H^{\frac{1}{2}}(\partial\Omega)$ will not be relevant to us here, let us state it at least once. We say that a function $f : \partial\Omega \to \mathbb{C}$ belongs to $H^{1/2}(\partial\Omega)$ if f is measurable with respect to the \mathscr{H}^{d-1} Hausdorff measure and satisfies

$$\|f\|_{\mathrm{H}^{1/2}(\partial\Omega)}^2 = \int_{\partial\Omega} |f(x)|^2 \mathcal{H}^{d-1}(\mathrm{d} x) + \int_{\partial\Omega\times\partial\Omega} \frac{|f(x) - f(y)|^2}{|x - y|^d} \mathcal{H}^{d-1}(\mathrm{d} x) \mathcal{H}^{d-1}(\mathrm{d} y) < \infty.$$

Alternatively, $H^{1/2}(\partial \Omega)$ can be defined via localization and the Fourier transform, see [McLoo, Chapter 3].

Remark 3.7 (One dimensional case). For $\Omega = I = (0, 1)$, any function in $H^1(I)$ is continuous up to the boundary. In this case, we set $H^{1/2}(\partial I) = \mathbb{C}^2$ and the trace operator reduces to $(\gamma f) = (f(0), f(1)) \in \mathbb{C}^2$.

Using the trace operator, we can define the following subspaces of $H^1(\Omega)$.

Definition 3.8 (Trace-restricted Sobolev spaces). For any closed subspace $L \subset H^{\frac{1}{2}}(\partial\Omega)$, we denote by $H^{1}_{L}(\Omega)$ the space of Sobolev functions with trace in *L*, i.e.,

$$\mathrm{H}^{1}_{L}(\Omega) = \{ f \in \mathrm{H}^{1}(\Omega) : \gamma f \in L \},\$$

where γ is the trace operator of Theorem 3.5.

From the continuity of the trace operator, it is easy to see that $H^1_L(\Omega)$ is a closed subspace of $H^1(\Omega)$. Moreover, any right inverse $J : H^{\frac{1}{2}}(\partial \Omega) \to H^1(\Omega)$ of the trace operator defines an isomorphism $H^1(\Omega) \cong H^{1/2}_0(\Omega) \oplus H^{1/2}(\partial \Omega)$ via the map

$$G: \mathrm{H}^{1}(\Omega) \to \mathrm{H}^{1}_{0}(\Omega) \oplus \mathrm{H}^{1/2}(\partial\Omega), \quad G(f) = (f - J\gamma(f), \gamma(f)).$$

From this decomposition, one can show that any closed subspace of $H^1(\Omega)$ containing $C_c^{\infty}(\Omega)$ is given by $H^1_t(\Omega)$ for a unique $L \subset H^{\frac{1}{2}}(\partial\Omega)$. More precisely, the following holds.

Proposition 3.9 (Characterization of intermediate spaces of H¹). Let $X \subset H^1(\Omega)$ be a closed subspace such that $C_c^{\infty}(\Omega) \subset X$, then there exists a unique closed subspace $L \subset H^{1/2}(\partial\Omega)$ such that $X = H^1_I(\Omega)$.

3.2. **Representation and density on dual Sobolev spaces**. The following lemma gives a simple but useful representation of the dual Sobolev spaces.

Lemma 3.10 (Representation of dual Sobolev space). Let $1 , then for any <math>T \in W^{-1,p}(\Omega)$, there exists $\alpha \in \mathbb{C}$ and $V \in L^p(\Omega; \mathbb{C}^n)$ such that

$$T(f) = \overline{\alpha} \int_{\Omega} f(x) dx + \int_{\Omega} \overline{V}(x) \cdot \nabla f(x) dx.$$
(3.3)

Proof. First, we note that the map

$$i: \mathbf{W}^{1,q}(\Omega) \to \mathbb{C} \times \mathbf{L}^q(\Omega; \mathbb{C}^n) \quad i(f) = \left(\int_{\Omega} f \mathrm{d}x, \nabla f\right)$$

is a continuous immersion, i.e., i is continuous, injective and has closed range. Indeed, the continuity is immediate from the definition of $W^{1,p}$ and the injectivity follows from the simple fact that

 $\nabla f - \nabla g = 0$ implies f - g = constant for any distributions $f, g \in \mathcal{D}'(\Omega)$,

T. CARVALHO CORSO

To see that *i* has closed range, note that if $\{(\int f_n, \nabla f_n)\}_{n \in \mathbb{N}}$ is a Cauchy sequence in $\mathbb{C} \times L^q(\Omega; \mathbb{C}^d)$, then by Poincare's inequality

$$\lim_{n \to \infty} \|f_n - f_m - \frac{1}{|\Omega|} \int (f_n - f_m)\|_{L^q} = 0.$$

Thus f_n is a Cauchy in $W^{1,q}(\Omega)$; in particular, the limit of this sequence exists and belongs to $f \in W^{1,q}$ by completeness. Hence $i(f) = (\int f, \nabla f) = \lim_{n \to \infty} (\int f_n, \nabla f_n)$.

Now let $T \in W^{-1,p}(\Omega)$. As the inverse of *i* is continuous from ran(*i*) to $W^{1,q}$ by the closed graph theorem, the functional $T \circ i^{-1}$ belongs to the dual of ran(*i*). Since ran(*i*) is a closed subspace of $\mathbb{C} \times L^q(\Omega; \mathbb{C}^d)$, by the Hahn-Banach theorem there exists a continuous extension of $T \circ i^{-1}$ to $\mathbb{C} \times L^q(\Omega; \mathbb{C}^d)$, denoted here by *T'*. Thus, by the Riesz representation theorem in L^q spaces (recall that $1 \le q < \infty$ since 1), $there exists <math>\beta \in \mathbb{C}$ and $V \in L^p(\Omega; \mathbb{C}^d)$ such that

$$T'(\beta, W) = \overline{\alpha}\beta + \int_{\Omega} \overline{V}(x) \cdot W(x) dx, \quad \text{for any } \beta \in \mathbb{C} \text{ and } W \in \mathrm{L}^{q}(\Omega; \mathbb{R}^{d}).$$

In particular, for $(\beta, W) = (\int f, \nabla f)$, equation (3.3) holds and the proof is complete.

An important consequence of the above representation is the following density result.

Lemma 3.11 (Density of smooth functions). Let $T \in W^{-1,p}(\Omega)$ with $1 , then the space of functions <math>\{g + \alpha : g \in C_c^{\infty}(\Omega), \alpha \in \mathbb{C}\} \subset C^{\infty}(\overline{\Omega})$ is dense in $W^{-1,p}(\Omega)$.

Proof. Let $(\alpha, V) \in \mathbb{C} \times L^p(\Omega; \mathbb{C}^d)$ be given by Lemma 3.10. As $C_c^{\infty}(\Omega)$ is dense in $L^p(\Omega)$ for any $1 \le p < \infty$, there exists a sequence of vector fields $V_n \in C_c^{\infty}(\Omega; \mathbb{C}^d)$ such that $||V_n - V||_{L^p} \to 0$ as $n \to \infty$. In particular, the sequence $g_n \coloneqq \alpha - \operatorname{div} V_n \in C_c^{\infty}(\Omega) + \mathbb{C}$ satisfies

$$\begin{aligned} \left| T(f) - \int_{\Omega} \overline{g_n(x)} f(x) dx \right| &= \left| \int_{\Omega} \overline{V(x)} \cdot \nabla f(x) dx + \int_{\Omega} \operatorname{div} \overline{V_n}(x) f(x) dx \right| = \left| \int_{\Omega} \overline{(V - V_n)}(x) \cdot \nabla f(x) dx \right| \\ &\leq \|V - V_n\|_{L^p} \|f\|_{1,q}, \quad \text{for any } f \in W^{1,q}. \end{aligned}$$

Hence, $g_n \to T$ in W^{-1,p}, which completes the proof.

Remark 3.12 (Dual representation on restricted Sobolev spaces). Since $W_L^{1,q}(\Omega)$ is a closed subspace of $W^{1,q}(\Omega)$, any functional in $W_L^{-1,p}(\Omega)$ can be continuously extended to a functional in $W^{-1,p}(\Omega)$ by the Hahn-Banach theorem. In particular, the representation and density results in Lemmas 3.10 and 3.11 hold for $W_L^{-1,p}(\Omega)$ and $W_0^{-1,p}(\Omega)$ as well.

3.3. **Regularity of reduced densities**. We now present a regularity result for reduced densities of wavefunctions with finite kinetic energy. This result is a simple application of the GNS, Hölder's, and Minkowiski integral inequalities.

Lemma 3.13 (Regularity of reduced densities). Let $\Psi, \Phi \in H^1(\Omega \times \Omega')$ where $\Omega \subset \mathbb{R}^d$ and $\Omega' \subset \mathbb{R}^{d'}$ are bounded Lipschitz domains. Let $\rho_{\Psi,\Phi}$ denote the overlapping density

$$\rho_{\Psi,\Phi}(x) \coloneqq \int_{\Omega'} \overline{\Psi(x,y)} \Phi(x,y) \mathrm{d}y.$$
(3.4)

Then we have

$$\|\rho_{\Psi,\Phi}\|_{1,p} \lesssim \|\Psi\|_{H^{1}} \|\Phi\|_{H^{1}}^{1-\frac{d}{p}} \|\Phi\|_{L^{2}}^{\frac{d}{p}} + \|\Phi\|_{H^{1}} \|\Psi\|_{H^{1}}^{1-\frac{d}{p}} \|\Psi\|_{L^{2}}^{\frac{d}{p}}, \quad for \ any \qquad \begin{cases} 1 \le p \le 2, \quad if \ d = 1, \\ 1 \le p < 2, \quad if \ d = 2, \\ 1 \le p \le \frac{d}{d-1}, \quad if \ d \ge 3, \end{cases}$$
(3.5)

where the implicit constant depends on p, Ω , and Ω' , but is independent of Ψ and Φ .

Proof. First, observe that

$$\nabla_{x}\rho_{\Psi,\Phi}(x) = \int_{\Omega'} \overline{\nabla_{x}\Psi(x,y)}\Phi(x,y) + \overline{\Psi(x,y)}\nabla_{x}\Phi(x,y)dy$$

$$\begin{split} \|\nabla_{x}\rho_{\Psi,\Phi}\|_{L^{p}} &= \left(\int_{\Omega} \left|\int_{\Omega'} (\overline{\nabla_{x}\Psi}\Phi + \overline{\Psi}\nabla_{x}\Phi) dy\right|^{p} dx\right)^{\frac{1}{p}} \leq \int_{\Omega'} \left(\int_{\Omega} (|\nabla_{x}\Phi\Psi| + |\Phi\nabla_{x}\Psi|)^{p} dx\right)^{\frac{1}{p}} dy \\ &\leq \int_{\Omega'} \left(\int_{\Omega} |\nabla_{x}\Psi|^{2} dx\right)^{\frac{1}{2}} \left(\int_{\Omega} |\Phi|^{\frac{2p}{2-p}} dx\right)^{\frac{2-p}{2p}} dy + \int_{\Omega'} \left(\int_{\Omega} |\nabla_{x}\Phi|^{2} dx\right)^{\frac{1}{2}} \left(\int_{\Omega} |\Psi|^{\frac{2p}{2-p}} dx\right)^{\frac{2-p}{2p}} dy \\ &\lesssim \int_{\Omega'} \|\nabla_{x}\Psi\|_{L^{2}_{x}} \|\Phi\|_{H^{1}_{x}}^{1-\theta} \|\Phi\|_{L^{2}_{x}}^{\theta} + \|\nabla_{x}\Phi\|_{L^{2}_{x}}^{2} \|\Psi\|_{H^{1}_{x}}^{1-\theta} \|\Psi\|_{L^{2}_{x}}^{\theta} dy \\ &\lesssim \|\Psi\|_{H^{1}} \|\Phi\|_{H^{1}}^{1-\theta} \|\Phi\|_{L^{2}}^{\theta} + \|\Phi\|_{H^{1}} \|\Psi\|_{L^{2}}^{1-\theta} \|\Psi\|_{L^{2}_{x}}^{\theta}, \end{split}$$

where $\theta = d(\frac{1}{2} - \frac{p-2}{2p}) = \frac{d}{p}$. As this holds for any smooth Ψ, Φ , the result follows by an approximation argument. The restriction p < 2 for d = 2 is necessary in (3.5) because the GNS inequality (3.2) is not valid in the end point case $(p, d) = (\infty, 2)$. On the other hand, the case $(p, d) = (\infty, 1)$ is allowed in (3.2) and therefore p = 2 in (3.5) is allowed for d = 1.

An important feature of inequality (3.5) is that only the dimension of the set Ω is relevant for the regularity of the reduced densities. Moreover, in the case where d' = 0, i.e., we consider only $\Psi, \Phi \in H^1(\Omega)$ and $\rho_{\Psi,\Phi} = \overline{\Psi}\Phi$, estimate (3.5) reduces to the standard (Sobolev) regularity expected by the product of two H¹ functions in \mathbb{R}^d . In particular, it implies that the set H¹(Ω) for an interval $\Omega = (a, b) \subset \mathbb{R}$ is an algebra of functions. This property was crucial in [Cor25, SPR+24] but will play no special role here.

4. Schrödinger operators with distributional potentials

In this section, we present the construction of self-adjoint realizations of $H_N(v, w)$ in detail. We also present some important examples of distributional potentials for which this construction holds. For convenience of the reader, we recall the basic results on the theory of quadratic forms and self-adjoint extensions in Appendix A.

4.1. Self-adjoint realizations. The estimate in Lemma 3.13 allows us to define self-adjoint Schrödinger operators with external and interaction potentials in a large class of distributions. Indeed, from estimate (3.5) and Young's inequality, we find that any $v \in \mathcal{V}$ and $w \in \mathcal{W}$ satisfy

$$|v(\rho_{\Psi,\Psi}) + w(\rho_{2,\Psi,\Psi})| \le \epsilon \|\Psi\|_{\mathrm{H}^1}^2 + \frac{C}{\epsilon} \|\Psi\|_{\mathrm{L}^2}^2, \quad \text{for any } \Psi \in \mathrm{H}^1(I_N), \tag{4.1}$$

and any $\epsilon > 0$ for some constant C = C(v, w) > 0. Hence, from the celebrated KLMN theorem (cf. Theorem A.5), for any closed subspace $L \subset H^{1/2}(\partial \Omega)$, we can construct a unique self-adjoint operator associated to the sesquilinear form

$$a_{v,w}(\Psi,\Phi) = \int_{I_N} \overline{\nabla \Psi(x)} \cdot \nabla \Phi(x) dx + v(\rho_{\Psi,\Phi}) + w(\rho_{2,\Psi,\Phi}), \quad \text{for } \Psi, \Phi \in \mathrm{H}^1_L(I_N) \cap \mathcal{H}_N.$$
(4.2)

Precisely, the following holds.

Lemma 4.1 (Schrödinger operators with distributional potentials). Let $N \in \mathbb{N}$, $v \in \mathcal{V}$, and $w \in \mathcal{W}$. Moreover, let $L \subset H^{1/2}(\partial I_N)$ be a closed subspace and $H^1_L(I_N)$ the trace-restricted Sobolev space introduced in Definition 3.8. Then the quadratic form in (4.2) is closed, symmetric and semibounded. In particular, there exists a unique self-adjoint operator $H^L_N(v, w)$ associated to this form. Moreover, $H^L_N(v, w)$ is semibounded and has purely discrete spectrum.

Remark 4.2 (Notation). For the rest of the paper, we shall denote by $H_N(v, w; \Gamma)$ the self-adjoint realization of $a_{v,w}$ with form domain $H^1(I_N; \Gamma) \cap \mathcal{H}_N$ and by $H^{\alpha}_N(v; w; \Gamma)$ the realization with form domain $H^1_{\alpha}(I_N; \Gamma) \cap \mathcal{H}_N$.

4.2. From single-particle to many-particle boundary conditions. Formally, the non-interacting operator $H_N(v, 0)$ can be written as the sum

$$H_N(v,0) = \sum_{j=1}^N 1 \otimes \dots \otimes \overbrace{h(v)}^{j^{th} position} \otimes \dots \otimes 1,$$

where 1 is the identity operator and $h(v) = -\Delta + v$ is the single-particle operator acting on $L^2(I)$. This allow us to reduce the study of non-interacting (electronic) operators to the case of single-particle ones.

However, to make this observation precise, we need to impose appropriate boundary conditions on the form domains of $H_N(v, 0)$ and h(v). In this section, our goal is to clarify this point.

To this end, we start with the following lemma.

Lemma 4.3 (Many-particle form domain). Let $\alpha \in \mathbb{R}$ and $H^1_{\alpha}(I)$ be the space introduced in (3.5). Then for any $N \in \mathbb{N}$, the span of Slater determinants

$$\Phi = \varphi_1 \wedge \dots \wedge \varphi_N, \quad \varphi_j \in \mathrm{H}^1_{\alpha}(I)$$

is dense in $\mathrm{H}^{1}_{\alpha}(I_{N}) \cap \mathcal{H}_{N}$.

Proof. We first observe that the projection on the antisymmetric space, explicitly given by

$$\Phi \mapsto (P_N \Phi)(x) = \frac{1}{N!} \sum_{\sigma \in \mathcal{P}_N} \operatorname{sign}(\sigma) \Phi(\sigma x)$$

is continuous with respect to the H¹ topology and maps *N*-fold tensor products to *N*-fold Slater determinants. Therefore, it suffices to show that the space

$$Y \coloneqq \operatorname{span}\{\varphi_1 \otimes \dots \otimes \varphi_N : \varphi_j \in \operatorname{H}^1_\alpha(I)\}$$

is dense in $H^1_{\alpha}(I_N)$. For this, let $\{\varphi_j\}_{j\in\mathbb{N}}$ be an orthonormal basis of eigenfunctions of the one-dimensional Laplacian in $H^1_{\alpha}(I)$, i.e., φ_j satisfies

$$\int_{I} \overline{\partial_{x} \varphi_{j}(x)} \partial_{x} \psi(x) dx = \lambda_{j} \int_{I} \overline{\varphi_{j}(x)} \psi(x) dx, \quad \text{for any } \psi \in \mathrm{H}^{1}_{\alpha}(I).$$

Then, it is not hard to see that $\varphi_i \in C^{\infty}(\overline{I})$ and

$$\alpha \partial_x \varphi_j(0) - \partial_x \varphi_j(1) = 0$$

Consequently, by integration by parts, for any $\Psi \in H^1_{\alpha}(I_N)$ and $(j_1, ..., j_N) \in \mathbb{N}^N$ we have

$$\begin{split} \langle \Psi, \varphi_{j_1} \otimes \dots \otimes \varphi_{j_N} \rangle_{\mathrm{H}^1} &= \sum_{k=1}^N \langle \partial_{x_k} \Psi, \partial_{x_k} \varphi_{j_1} \otimes \dots \varphi_{j_N} \rangle_{\mathrm{L}^2} + \langle \Psi, \varphi_{j_1} \otimes \dots \varphi_{j_N} \rangle_{\mathrm{L}^2} \\ &= \sum_{k=1}^N \lambda_{j_k} \langle \Psi, \varphi_{j_1} \otimes \dots \otimes \varphi_{j_k} \rangle_{\mathrm{L}^2} + \langle \Psi, \varphi_{j_1} \otimes \dots \otimes \varphi_{j_N} \rangle_{\mathrm{L}^2} \\ &= \left(1 + \sum_{k=1}^N \lambda_{j_k} \right) \langle \Psi, \varphi_{j_1} \otimes \dots \otimes \varphi_{j_N} \rangle. \end{split}$$

Thus, if Ψ is H¹-orthogonal to Y, then Ψ is L²-orthogonal to all $\{\varphi_{j_1} \otimes ... \otimes \varphi_{j_N}\}_{(j_1,...,j_N) \in \mathbb{N}^N}$. As this set is an orthogonal basis of L²(I_N), we must have $\Psi = 0$. Hence, the intersection of H¹_{α}(I_N) with the H¹-orthogonal complement of Y is trivial, which implies that Y is dense in H¹_{α}(I_N).

As a consequence of the previous lemma, we have the following relation between the single-particle operator $h_{\alpha}(v) = -\Delta_{\alpha} + v$ and the *N*-particle operator $H_N^{\alpha}(v, 0)$ associated to the form

$$a_{v}(\Psi,\Psi) = \int_{I_{N}} |\nabla \Psi(x)|^{2} \mathrm{d}x + v(\rho_{\Psi}), \quad \Psi \in \mathrm{H}^{1}_{\alpha}(I_{N}) \cap \mathcal{H}_{N}.$$

Lemma 4.4 (From single-particle to non-interacting many-particle systems). Let $v \in \mathcal{V}$, $\alpha \in \mathbb{C}^2$, and $N \in \mathbb{N}$. Let $\{\psi_j\}_{j\in\mathbb{N}}$ be an orthogonal basis of eigenfunctions of $h_{\alpha}(v) = -\Delta_{\alpha} + v$ with corresponding eigenvalues $\{\lambda_j\}_{j\in\mathbb{N}}$ ordered in non-decreasing order. Then the Slater determinants

$$\{\Psi_J = \psi_{j_1} \land \dots \land \psi_{j_N} : J = \{j_1, \dots, j_N\} \in \mathbb{N}^N \quad \text{such that} \quad 1 \le j_1 < j_2 \dots < j_N\}$$
(4.3)

forms an \mathcal{H}_N -orthogonal basis of eigenfunctions of $H_N^{\alpha}(v, 0)$ with eigenvalues $\lambda_J = \sum_{k=1}^N \lambda_{j_k}$. In particular, the ground-state of $H_N^{\alpha}(v, 0)$ is non-degenerate if and only if λ_N satisfies the closed shell condition $\lambda_N < \lambda_{N+1}$.

Proof. From a long but straightforward calculation we find that

$$a_{v}(\Psi_{J},\varphi_{1}\wedge\ldots\wedge\varphi_{N})=\lambda_{J}\langle\Psi_{J},\varphi_{1}\wedge\ldots\wedge\varphi_{N}\rangle_{L^{2}}, \text{ for any } \{\varphi_{j}\}_{j=1}^{N}\subset \mathrm{H}^{1}_{\alpha}(I),$$

where $\Psi_J = \psi_{j_1} \wedge ... \wedge \psi_{j_N}$ and $\lambda_J = \sum_{k=1}^N \lambda_{j_k}$. Thus by Lemma 4.3 and approximation we have

$$a_v(\Psi_J, \Phi) = \lambda_J \langle \Psi_J, \Phi \rangle$$
 for any $\Phi \in \mathrm{H}^1_\alpha(I_N) \cap \mathcal{H}_N$

Therefore Ψ_J is an eigenfunction of $H_N^{\alpha}(v, 0)$ with eigenvalue λ_J . The fact that the Ψ_J 's form an \mathcal{H}_N orthogonal basis follows from the fact that $\{\psi_i\}_{i\in\mathbb{N}}$ is an L²-orthogonal basis of L²(I).

4.3. **Examples of distributional potentials**. To illustrate how large the class of admissible external and interaction potentials in Lemma 4.1 is, let us present a few important examples.

- (1) The Dirac's delta potential δ_{x_0} with $x_0 \in I$ belongs to \mathcal{V} by the continuous embedding $H^1(I) \subset C(\overline{I})$. Note that $x_0 \in \partial I = \{0, 1\}$ is also allowed.
- (2) The δ -interaction potential $w = \delta(x y)$, defined via

$$w_{\delta}(\rho_{2,\Psi,\Phi}) = \int_{I} \rho_{2,\Psi,\Phi}(x,x) \mathrm{d}x.$$

This follows from the fact that the trace operator $\gamma : W^{1,p}(I_2) \to L^p(D)$ sending ρ_2 to its restriction along the diagonal set $D = \{(x, x) \in I_2 : x \in I\}$ is a continuous operator.

(3) Multiplicative potentials $v \in L^{1}(I)$ and $w \in L^{1}(2I)$, whose actions are defined via

$$v(\rho) = \int_I v(x)\rho(x)dx$$
 and $w(\rho_2) = \int_{I_2} w(x-y)\rho_2(x,y)dxdy$,

are also allowed. The first one is clear from Hölder's inequality. For the second one, note that

$$f(u) \coloneqq \int_{|u|}^{1-|u|} \rho_2\left(\frac{u+v}{2}, \frac{u-v}{2}\right) \mathrm{d}v$$

belongs to $L^{\infty}(I)$ for any $\rho_2 \in W^{1,p}(I_2)$ with $1 \le p \le \infty$. Hence, the map

$$\rho_2 \mapsto \int_{I_2} w(x-y)\rho_2(x,y)\mathrm{d}x\mathrm{d}y = \frac{1}{2}\int_{-1}^1 w(u)f(u)\mathrm{d}u$$

belongs to $W^{-1,q}(I_2)$.

(4) Lemma 4.1 can also be extended to the case of k^{th} -body distributional potentials for $k \ge 3$. To be precise, for any $w_k \in W^{-1,q}(I_k)$ with q > k, one can use Lemma 3.13 and Young's inequality to show that the form $(\Psi, \Phi) \mapsto w_k(\rho_{k,\Psi,\Phi})$, where

$$\rho_{k,\Psi,\Phi}(x_1,..,x_k) = \frac{N!}{(N-k)!k!} \int_{I_{N-k}} \overline{\Psi(x_1,...,x_N)} \Phi(x_1,...,x_N) dx_{k+1}...dx_N,$$

is Δ -bounded with relative bound 0. Therefore, any Schrödinger operator of the form

$$H_N(w_1, w_2, ..., w_N) = -\Delta + \sum_{k=1}^N \sum_{j_1 \neq ... \neq j_k}^N w_k(x_{j_1}, x_{j_2}, ..., x_{j_k}),$$

with real-valued $w_k \in W^{-1,q_k}(I_k)$, where $q_k > k$ for $k \ge 2$ and $w_1 \in \mathcal{V}$, defines a self-adjoint operator with discrete spectrum. In particular, the case of 3D Coulomb interactions

$$w(x_1, ..., x_6) = \left(\sum_{j=1}^3 (x_j - x_{j+3})^2\right)^{-\frac{1}{2}}$$

is also included for $N \ge 6$.

5. Non-degeneracy of the ground-state

Our goal in this section is to prove Theorem 2.1.

5.1. **Perron-Frobenius theorem**. We start with a Perron-Frobenius theorem for the free Laplacian without particle statistics. To properly state this result, we introduce the following definition.

Definition 5.1. (Positivity and reality preserving) Let $L \subset H^{\frac{1}{2}}(\partial \Omega)$ be a closed subspace and $H^{1}_{L}(\Omega)$ the space introduced in Definition 3.8. Then we say that

- (i) $H_L^1(\Omega)$ is reality preserving if $\operatorname{Re} \Psi \in H_L^1(\Omega)$ for any $\Psi \in H_L^1(\Omega)$.
- (ii) $H_{I}^{1}(\Omega)$ is positivity preserving if $\Psi_{+} = \max\{\Psi, 0\} \in H_{I}^{1}(\Omega)$ for any real-valued $\Psi \in H_{I}^{1}(\Omega)$.

Lemma 5.2 (Perron-Frobenius for the free Laplacian). Let $\Omega \subset \mathbb{R}^d$ be open, connected, and bounded subset with Lipschitz boundary, and let $\mathrm{H}^1_L(\Omega)$ be positivity and reality preserving. Then the Laplacian $-\Delta_L$, defined as the unique self-adjoint operator associated to the quadratic form

$$a_0: \mathrm{H}^1_L(\Omega) \times \mathrm{H}^1_L(\Omega) \to \mathbb{C}, \quad a_0(\Psi, \Phi) \coloneqq \int_{\Omega} \overline{\nabla \Psi(x)} \cdot \nabla \Phi(x) \mathrm{d}x$$

has a non-degenerate ground-state and the unique (up to a global phase) normalized ground-state wave-function is strictly positive everywhere in Ω .

Proof. The proof is based on the variational principle and the strong maximum principle. So first, we note that $-\Delta_L$ has discrete spectrum and therefore a ground-state Ψ exists. Next, we claim that the ground-state can be taken real-valued. To see this, note that $\overline{\Phi} \in H^1_L(\Omega)$ for any $\Phi \in H^1_L(\Omega)$ by the reality preserving property. In particular

$$a_0(\overline{\Psi}, \Phi) = \int_{\Omega} \nabla \Psi(x) \cdot \nabla \Phi(x) dx = a_0(\overline{\Phi}, \Psi) = \lambda \langle \overline{\Phi}, \Psi \rangle = \lambda \langle \overline{\Psi}, \Phi \rangle, \quad \text{for any } \Phi \in \mathrm{H}^1_L(\Omega),$$

where $\lambda \ge 0$ is the ground-state energy. Consequently, $\overline{\Psi}$ is also a ground-state of $-\Delta_L$, and therefore $2\text{Re}\Psi = \Psi + \overline{\Psi}$ is a ground-state as well.

Thus, assuming that the ground-state is real-valued, we can define $\Psi_+ = \max\{0, \Psi\}$ and $\Psi_- = (-\Psi)_+$. Since $H^1_L(\Omega)$ is positivity preserving by assumption, both Ψ_+ and Ψ_- belong to $H^1_L(\Omega)$. Moreover, since they have disjoint support and $\Psi = \Psi_+ - \Psi_-$, we have

$$a_{0}(\Psi_{+},\Psi_{+}) = \int_{\Omega} \overline{\nabla \Psi_{+}(x)} \cdot \nabla \Psi_{+}(x) dx = \int_{\Omega} \overline{\nabla \Psi(x)} \cdot \nabla \Psi_{+}(x) dx = \lambda \langle \Psi, \Psi_{+} \rangle = \lambda ||\Psi_{+}||_{L^{2}}^{2}$$

Hence, it follows from the variational principle that

$$\lambda = \min_{\Psi \in H_{1}^{1} \setminus \{0\}} \frac{a_{0}(\Psi, \Psi)}{\|\Psi\|_{12}^{2}} = \frac{a_{0}(\Psi_{+}, \Psi_{+})}{\|\Psi_{+}\|_{12}^{2}}$$

As a consequence, Ψ_+ satisfies the Euler-Lagrange (eigenfunction) equation

$$a_0(\Phi, \Psi_+) = \int_{\Omega} \overline{\nabla \Phi(x)} \cdot \nabla \Psi(x) = \lambda \langle \Phi, \Psi_+ \rangle, \quad \text{for any } \Phi \in C_c^{\infty}(\Omega),$$

or equivalently, $(-\Delta - \lambda)\Psi_+ = 0$ in the distributional sense. From standard elliptic regularity (cf. [Eva10, Section 6.3]), it follows that $\Psi_+ \in C^{\infty}(\Omega)$. As $\Psi_+ \ge 0$ and $\lambda \ge 0$, Ψ_+ is a subsolution of Laplace equation,

$$-\Delta \Psi_+ \ge 0$$

Therefore, we can apply the strong maximum principle [Evalo, Theorem 3 in Section 6.4.2] to conclude that either $\Psi_+(x) > 0$ for any $x \in \Omega$ or $\Psi_+ = 0$. Hence, either $\Psi_+ = 0$ or $\Psi_- = 0$, which implies $\Psi(x)$ has constant sign everywhere in Ω . Thus any real-valued ground-state of $-\Delta_L$ is strictly positive (up to multiplication by a constant). Therefore, the ground-state is non-degenerate as there cannot be two strictly positive functions that are mutually orthogonal.

We can now combine the previous result with the perturbative approach in Reed and Simon [RS78, Section XIII.12] to prove the following Perron-Frobenius theorem for generalized Schrödinger operators. This result will play a key role throughout this paper.

Theorem 5.3 (Perron-Frobenius for Schrödinger operators with distributional potentials). Let $\Omega \subset \mathbb{R}^d$ and $H^1_L(\Omega)$ be as in Lemma 5.2. Let $V : H^1_L(\Omega) \times H^1_L(\Omega) \to \mathbb{C}$ be a sequelinear form and suppose that there exists a sequence $\{V_n\}_{n\in\mathbb{N}} \subset L^{\infty}(\Omega)$ of real-valued functions such that

$$\lim_{n \to \infty} \|V - V_n\|_{H^1 \to H^{-1}} = \lim_{n \to \infty} \sup_{\Psi, \Phi \in H^1_I(\Omega) \setminus \{0\}} \frac{\left| V(\Psi, \Phi) - \int_{\Omega} V_n(x) \overline{\Psi(x)} \Phi(x) dx \right|}{\|\Psi\|_{H^1} \|\Phi\|_{H^1}} = 0.$$

Then there exists a unique self-adjoint operator $H(V) = -\Delta_L + V$ with quadratic form

$$a_{V}(\Psi, \Phi) = \int_{\Omega} \overline{\nabla \Psi(x)} \cdot \nabla \Phi(x) dx + V(\Psi, \Phi), \quad \text{for } \Phi, \Psi \in \mathrm{H}^{1}_{L}(\Omega).$$
(5.1)

Moreover, the ground-state of H(V) is non-degenerate and the unique (up to a global phase) ground-state wave-function can be chosen strictly positive almost everywhere in Ω .

Proof. We first show that V is symmetric and Δ -bounded with relative bound 0. For this, note that, for any $\epsilon > 0$ there exists $n \in \mathbb{N}$ such that $||V - V_n||_{H^1 \to H^{-1}} \leq \epsilon$. Hence,

$$\left| V(\Psi, \Phi) - \int_{\Omega} V_n(x) \overline{\Psi(x)} \Phi(x) \mathrm{d}x \right| \le \epsilon \|\Psi\|_{\mathrm{H}^1} \|\Psi\|_{\mathrm{H}^1} + \|V_n\|_{\mathrm{L}^{\infty}} \|\Psi\|_{\mathrm{L}^2} \|\Phi\|_{\mathrm{L}^2}, \quad \text{for any } \Psi, \Phi \in \mathrm{H}^1_L(\Omega),$$

and

$$\begin{aligned} |V(\Psi, \Phi) - \overline{V(\Phi, \Psi)}| &= \left| \left(V(\Psi, \Phi) - \int_{\Omega} V_n(x) \overline{\Psi}(x) \Phi(x) dx \right) + \left(\overline{V(\Phi, \Psi)} - \int_{\Omega} \overline{V_n(x) \overline{\Phi(x)} \Psi(x)} dx \right) \right| \\ &\leq 2\epsilon \|\Psi\|_{H^1} \|\Phi\|_{H^1}. \end{aligned}$$

As $\epsilon > 0$ can be taken arbitrarily small, we conclude that V is symmetric and Δ -bounded with relative bound 0. Hence, by the KLMN theorem (cf. Theorem A.5), there exists a unique self-adjoint operator with quadratic form (5.1). Moreover, as the quadratic form domain of H(V) is $H_I^1(\Omega)$, which is compactly embedded in $L^2(\Omega)$, the operator H(V) has compact resolvent and therefore purely discrete spectrum. In particular, there exists at least one ground-state.

To show that the ground-state is non-degenerate and strictly positive almost everywhere, we shall use the following lemma, which is a reformulation of results extracted from Reed and Simon [RS78, Theorem XIII.43 and Theorem XIII.45]. For the proof of this lemma, we refer to their book.

Lemma 5.4 (Perron-Frobenius invariance under strong resolvent convergence via multiplicative potentials). Let H and H_0 be self-adjoint operators on $L^2(\Omega)$ and suppose that there exists $\{V_n\}_{n\in\mathbb{N}} \subset L^{\infty}(\Omega)$ such that $H_0 + V_n$ converges in the strong resolvent sense to H. Suppose also that e^{-tH_0} is positivity preserving (i.e., sends non-negative functions to non-negative functions). Then the following holds: if H_0 has a non-degenerate and almost everywhere strictly positive ground-state, then so does H (provided that a ground-state exists).

We now observe that, by Lemma 5.2, the operator $H_0 = -\Delta_L$ has a non-degenerate and strictly positive ground-state. Moreover, e^{-tH_0} is positivity preserving because, from the standard formula

 $\nabla |\Psi| = \nabla \Psi \mathbb{1}_{\{\Psi > 0\}} - \nabla \Psi \mathbb{1}_{\{\Psi < 0\}}, \text{ where } \mathbb{1}_A \text{ is the indicator function on } A,$

for real-valued Ψ , we see that H_0 satisfies the Beurling-Deny criterion (cf. [RS78, Theorem XIII.50]):

$$\langle |\Psi|, H_0|\Psi| \rangle = \int_{\Omega} |\nabla|\Psi|(x)|^2 dx = \int_{\Omega} |\nabla\Psi(x)|^2 dx = \langle \Psi, H\Psi \rangle, \quad \text{for any } \Psi \in \mathrm{H}^1_L(\Omega).$$

To conclude, we now note that, since $V_n \to V$ in $\mathcal{B}(H^1; H^{-1})$, the operators $H_0 + V_n$ converge to H(V) in the strong (in fact norm) resolvent sense. Indeed, this follows from the resolvent formula

$$(z - H(V))^{-1} - (z - H_0 - V_n)^{-1}$$

= $(z - H(V))^{-1/2} \left(I - \left(I + (z - H(V))^{-1/2} (V - V_n) (z - H(V))^{-1/2} \right)^{-1} \right) (z - H(V))^{-1/2}.$

(See [RS80, Theorem VIII.25.(c)] for the detailed argument.) We can thus apply Lemma 5.4 to complete the proof.

5.2. Unitary reduction to the simplex. Throughout this section, we denote by S_N the open simplex

$$S_N \coloneqq \{(x_1, ..., x_N) \in I_N : 0 < x_1 < x_2 < ... < x_N < 1\}.$$

The key observation that allow us to apply Theorem 5.3 to the operator $H_N^L(v, w)$ is that the hypercube I_N can be decomposed in a disjoint union of reflections of the simplex S_N . To make this precise, let us introduce some additional notation.

First, we denote by \mathcal{P}_N the set of all permutations of $\{1, ..., N\}$, i.e.,

• •

$$\mathcal{P}_N = \{ \sigma : \{1, ..., N\} \rightarrow \{1, ..., N\} \text{ bijective} \}.$$

With some abuse of notation, we also denote by σ the linear map of permutation of coordinates

$$\sigma: \mathbb{R}^N \to \mathbb{R}^N, \quad \sigma(x_1, ..., x_N) = (x_{\sigma(1)}, ..., x_{\sigma(N)}).$$
(5.2)

Recall that the sign of the permutation σ is defined as $sgn(\sigma) = \det \sigma \in \{-1, 1\}$. Moreover, we denote by $\{F_j\}_{i=0}^N$ the faces of the boundary of the simplex S_N , i...,

$$F_j := \{ (x_1, ..., x_N) \in \overline{S_N} : x_j = x_{j+1} \},$$
(5.3)



FIGURE 1. Simplex S_N in the case N = 2 (left) and N = 3 (right) with edges of ∂I_N (dashed lines) and interior boundary $\Gamma_{int} \subset \partial S_N$ (in red).

with the convention $x_0 = 0$ and $x_{N+1} = 1$. We can now define the interior part of the boundary of S_N as

 $\Gamma_{\text{int}} \coloneqq \partial S_N \setminus (F_0 \cup F_N) = \{(x_1, ..., x_N) \in \overline{S_N} : x_j = x_{j+1} \text{ for some } j \text{ and } 0 < x_1 \le x_N < 1\}.$ (5.4) For a visual illustration of S_N and the interior boundary Γ_{int} , see Figure 1.

Let us also define the pushforward map $\sigma_{\#}$ as

$$\sigma_{\#}: \mathrm{H}^{1}(S_{N}) \to \mathrm{H}^{1}(\sigma(S_{N})), \quad (\sigma_{\#}\Psi)(x_{1}, ..., x_{N}) = \Psi\left(\sigma^{-1}(x_{1}, ..., x_{N})\right) = \Psi\left(x_{\sigma^{-1}(1)}, ..., x_{\sigma^{-1}(N)}\right).$$

Finally, let us define the reduced trace-restricted space on S_N as

$$\mathrm{H}^{1}_{L}(S_{N}) \coloneqq \{\Psi|_{S_{N}} : \Psi \in \mathrm{H}^{1}_{L}(I_{N}) \cap \mathcal{H}_{N}\}.$$

$$(5.5)$$

We then have the following key decomposition of $\mathcal{H}_N \cap \mathrm{H}^1_I(I_N)$.

Lemma 5.5 (Unitary reduction to the simplex). The map $T : H^1_L(S_N) \to H^1_L(I_N) \cap \mathcal{H}_N$ defined via

$$T(\Psi) = \frac{1}{\sqrt{N!}} \sum_{\sigma \in \mathcal{P}_N} \operatorname{sgn}(\sigma) \sigma_{\#}(\Psi)$$
(5.6)

is an isometric isomorphism.

Proof. The key observations here are the following:

(i) (Tessellating cover) The sets $\sigma(\overline{S_N})$ with $\sigma \in \mathcal{P}_N$ tessellate I_N , i.e.,

$$I_N \subset \bigcup_{\sigma \in \mathcal{P}_N} \sigma(\overline{S_N}) \quad \text{and} \quad \sigma(S_N) \cap \tau(S_N) = \emptyset \quad \text{for any } \sigma \neq \tau \in \mathcal{P}_N.$$
(5.7)

(ii) (Interior boundary) The interior boundary Γ_{int} defined in (5.4) satisfies $\Gamma_{\text{int}} = \partial S_N \cap I_N$.

The fact that $\sigma(\overline{S_N})$ covers I_N is clear. Indeed, for any $(x_1, ..., x_N) \in I_N$ there exists $\sigma \in \mathcal{P}_N$ such that $0 \le x_{\sigma(1)} \le x_{\sigma(2)} ... \le x_{\sigma(N)} \le 1$ and therefore $(x_1, ..., x_N) \in \sigma^{-1}(\overline{S_N})$.

To prove the disjoint property, we first note that, since any $\sigma \in \mathcal{P}_N$ is a homeomorphism in \mathbb{R}^N ,

$$\sigma(S_N) \cap \tau(S_N) = \emptyset$$
 if and only if $(\tau^{-1} \circ \sigma)(S_N) \cap S_N = \emptyset$.

Hence, it suffices to show (5.7) for $\sigma \in \mathcal{P}_N$ and $\tau = I$ (identity map). For this, we define

 $m(\sigma) \coloneqq \max\{j : \sigma^{-1}(k) \le k \text{ for any } k \le j\} \cup \{0\}.$

Then, if $\sigma \neq I$ we must have $0 \leq m(\sigma) < N$. In particular $\sigma^{-1}(m(\sigma) + 1) > m(\sigma) + 1 = \sigma^{-1}(m(\sigma) + k)$ for some $k \geq 2$. Then, on the one hand, any $(x_1, ..., x_N) \in \sigma(S_N)$ satisfies $x_{\sigma^{-1}(j)} < x_{\sigma^{-1}(j+1)}$ for any j and, in particular, $x_{\sigma^{-1}(m(\sigma)+1)} < x_{\sigma^{-1}(m(\sigma)+k)} = x_{m(\sigma)+1}$. On the other hand, any $(x_1, ..., x_N) \in S_N$ satisfies $x_j < x_{j+1}$ for any j. Since $\sigma^{-1}(m(\sigma) + 1) > m(\sigma) + 1$, this implies, in particular, that $x_{m(\sigma)+1} < x_{\sigma^{-1}(m(\sigma)+1)}$. Therefore, we can not have $(x_1, ..., x_N) \in S_N \cap \sigma(S_N)$, which completes the proof of (i).

We now turn to (ii). First, it is clear from the definition (5.3) that $F_0 \cup F_N \subset \partial I_N$ and therefore $(F_0 \cup F_N) \cap I_N = \emptyset$. Hence, $\partial S_N \cap I_N \subset \Gamma_{int}$. On the other hand, if $(x_1, ..., x_N) \in \Gamma_{int} = \partial S_N \setminus (F_0 \cup F_N)$ we must have $0 < x_0 \le x_1 \dots \le x_N < 1$. As every element in the boundary of I_N must have at least one coordinate with value 0 or 1, we must have $(x_1, ..., x_N) \in I_N$ and therefore item (ii) holds.

We now show that T is an isometry from $H^1_t(S_N)$ to $H^1(I_N) \cap \mathcal{H}_N$. To this end, first note that, since Ψ is the restriction of an antisymmetric function, it follows from the definition of Γ_{int} (see (5.4)) that $\gamma \Psi|_{\Gamma_{\text{int}}} = 0$. Consequently, from property (ii), the extension of Ψ to I_N by zero on $I_N \setminus S_N$ belongs to $H^1(I_N)$. Hence $T\Psi \in H^1(I_N)$. Moreover, if we denote the permutation of x_i and x_{i+1} by σ_i , then by construction we have

$$(\sigma_j)_{\#}(T\Psi) = \frac{1}{\sqrt{N!}} \sum_{\sigma \in \mathcal{P}_N} \operatorname{sign}(\sigma)(\sigma_j \circ \sigma)_{\#}(\Psi) = -\frac{1}{\sqrt{N!}} \sum_{\sigma_j \circ \sigma \in \mathcal{P}_N} \operatorname{sgn}(\sigma_j \circ \sigma)(\sigma_j \circ \sigma)_{\#}(\Psi) = -T\Psi.$$

Therefore, $T\Psi \in \mathcal{H}_N \cap H^1(I_N)$. Furthermore, since each $\sigma(S_N)$ is disjoint and σ an isometry, we have

$$\|T\Psi\|_{L^{2}(\Omega)}^{2} + \|\nabla T\Psi\|_{L^{2}(\Omega)}^{2} = \frac{1}{N!} \sum_{\sigma \in \mathcal{P}_{N}} \int_{\sigma(S_{N})} |\Psi(\sigma^{-1}x)|^{2} + |\nabla\Psi(\sigma^{-1}x)|^{2} dx = \|\Psi\|_{H^{1}(S_{N})}^{2},$$

and therefore T is an isometry. Moreover, a similar calculation shows that the restriction map $\Psi \mapsto T^{-1}\Psi =$ $\sqrt{N!}\Psi|_{S_N}$ is an isometry from $H^1(I_N) \cap \mathcal{H}_N$ to $H^1(S_N;\Gamma_{int})$. As T^{-1} maps $H^1_L(I_N) \cap \mathcal{H}_N$ to $H^1_L(S_N)$ by definition, it follows that $T: H^1_I(S_N) \to H^1_I(I_N) \cap \mathcal{H}_N$ is surjective, which completes the proof.

As the operator T in (5.6) is an isometry from $H^1_L(S_N)$ to $H^1_L(I_N)$ and from $L^2(S_N)$ to $L^2(I_N)$, by Lemma A.8, the form

$$\widetilde{a}_{v,w}(\Psi,\Phi) \coloneqq (T^{\#}a_{v,w})(\Psi,\Phi) = a_{v,w}(T\Psi,T\Phi) = \int_{I_N} \overline{\nabla(T\Psi)(x)} \cdot \nabla(T\Phi)(x) dx + v(\rho_{T\Psi,T\Phi}) + w(\rho_{2,T\Phi,T\Phi}),$$

$$= \int_{S_N} \overline{\nabla\Psi(x)} \cdot \nabla\Phi(x) dx + v(\rho_{T\Psi,T\Phi}) + w(\rho_{2,T\Phi,T\Phi}), \quad \text{for } \Psi, \Phi \in \mathrm{H}^1_L(S_N),$$
(5.8)

where the density and pair density are defined as in (2.2) and (2.3), is a semibounded closed form on $L^2(S_N)$. Hence, we can define the operator

$$\widetilde{H}_{N}^{L}(v,w) \coloneqq T^{\#}H_{N}^{L}(v,s)$$
(5.9)

as the unique self-adjoint operator associated to the form $\tilde{a}_{v,w}$ with form domain $H_I^1(S_N)$. Moreover, since T is an isometric isomorphism between $H^1_L(S_N)$ and $H^1_L(I_N) \cap \mathcal{H}_N$, the operator $H^L_N(v, w)$ is unitarily equivalent to $\widetilde{H}_{N}^{L}(v, w)$. In particular, we can now apply Theorem 5.3 to obtain the following result.

Theorem 5.6 (Perron-Frobenius with Fermi statistics). Let $v \in V$, $w \in W$, and $L \subset H^{1/2}(\partial I_N)$ be a closed subspace such that

$$\mathrm{H}^{1}_{L}(S_{N}) = \{\Psi|_{S_{N}} : \Psi \in \mathrm{H}^{1}_{L}(I_{N}) \cap \mathcal{H}_{N}\}$$

is reality and positivity preserving. Then the self-adjoint operator $H_N^L(v,w)$ defined in Lemma 4.1 has a non-degenerate ground-state and the unique (up to a global phase) ground-state wave-function Ψ satisfies $\Psi(x) > 0$ for a.e. $x \in S_N$.

Proof. As T is an isometry, the ground-state of the reduced operator $\widetilde{H}_{N}^{L}(v, w)$ defined in (5.9) is mapped to the ground-state of $H_N^L(v, w)$ via the map $\Psi \mapsto T\Psi$. Hence, it suffices to show that $H_N^L(v, w)$ has a non-degenerate and strictly positive ground-state. In particular, it suffices to show that $H_N^L(v, w)$ satisfy the assumptions from Theorem 5.3.

To this end, we first use Lemma 3.11 to obtain a sequence of bounded (actually smooth) functions $v_n \in L^{\infty}(I)$ and $w_n \in L^{\infty}(I_2)$ such that $\lim_{n\to\infty} ||v_n - v||_{H^{-1}} \to 0$ and $\lim_{n\to\infty} ||w_n - w||_{W^{-1,p}} = 0$. Then, by estimate (3.5) we have

$$\lim_{n \to \infty} \sup_{\Psi, \Phi \in H^{1}(I^{N}) \setminus \{0\}} \frac{|v_{n}(\rho_{\Psi, \Phi}) + w_{n}(\rho_{2, \Psi, \Phi}) - v(\rho_{\Psi, \Phi}) - w(\rho_{2, \Psi, \Phi})|}{\|\Psi\|_{H^{1}} \|\Phi\|_{H^{1}}} \leq \lim_{n \to \infty} \|v_{n} - v\|_{-1, 2} + \|w_{n} - w\|_{-1, p} = 0.$$

As $T: \mathrm{H}^{1}_{L}(S_{N}) \to \mathrm{H}^{1}_{L}(I_{N}) \cap \mathcal{H}_{N}$ is continuous, the forms $T^{\#}v_{n}$ and $T^{\#}w_{n}$, defined as

$$(T^{\#}v_n)(\Psi,\Phi) = v_n(\rho_{T\Psi,T\Phi})$$
 and $(T^{\#}w_n)(\Psi,\Phi) = w_n(\rho_{2,T\Psi,T\Phi})$

also converge to the forms $T^{\#}v$ and $T^{\#}w$ (defined analogously) in the H¹-form norm, i.e.,

$$\lim_{n \to \infty} \sup_{\Psi, \Phi \in H^1_I(S_N) \setminus \{0\}} \frac{|(T^* v_n + T^* w_n - T^* v - T^* w)(\Psi, \Phi)|}{||\Psi||_{H^1} ||\Phi||_{H^1}} = 0.$$
(5.10)

Moreover, it follows from the definition of *T* and the tessellating property (5.7) that, for any $V \in L^{\infty}(I_N)$,

$$\begin{aligned} (T^{\#}V)(\Psi,\Phi) &= V(T\Psi,T\Phi) = \int_{I_N} V(x)\overline{T\Psi(x)}T\Phi(x)dx = \frac{1}{N!}\sum_{\sigma,\tau\in\mathcal{P}_N}\int_{\sigma(S_N)\cap\tau(S_N)} V(x)\overline{\Psi(\sigma^{-1}x)}\Phi(\tau^{-1}x)dx, \\ &= \frac{1}{N!}\sum_{\sigma\in\mathcal{P}_N}\int_{S_N} V(\sigma x)\overline{\Psi(x)}\Phi(x)dx = \int_{S_N}\widetilde{V}(x)\overline{\Psi(x)}\Phi(x)dx. \end{aligned}$$

In particular, this implies that the pull-back of any form on I_N generated by a bounded multiplicative potential is a form in S_N generated by a bounded multiplicative potential. As the forms v_n and w_n are generated by the multiplicative potentials

$$V_n(x_1,...,x_N) = \sum_{j=1}^N v_n(x_j)$$
 and $W_n(x_1,...,x_N) = \sum_{i\neq j} w_n(x_i,x_j),$

the forms $T^{\#}(v_n + w_n)$ are also generated by multiplicative potentials. Thus, by estimate (5.10), we have found a sequence of multiplicative potentials approximating $T^{\#}v + T^{\#}w$. We can therefore apply Theorem 5.3 to complete the proof.

5.3. **Proofs of Theorems 2.1 and 2.3**. We now prove the following result, which together with Theorem 5.6 completes the proof of Theorems 2.1 and 2.3.

Lemma 5.7 (Positivity preserving spaces). Let $N \in \mathbb{N}$ and $\Gamma \subset \partial I_N$, then the space

$$\{\Psi|_{S_N}: \Psi \in \mathrm{H}^1(I_N; \Gamma) \cap \mathcal{H}_N\} \subset \mathrm{H}^1(S_N)$$
(5.11)

is reality and positivity preserving. Moreover, if $\alpha \in \mathbb{R} \setminus \{0\}$ satisfies (2.4), then the space

$$\{\Psi|_{S_N}: \Psi \in \mathrm{H}^1_{\alpha}(I_N; \Gamma) \cap \mathcal{H}_N\} \subset \mathrm{H}^1(S_N)$$
(5.12)

is also reality and positivity preserving.

Proof. That both spaces are reality preserving is clear since $H^1_{\alpha}(I_N; \Gamma)$ with $\alpha \in \mathbb{R}$ and \mathcal{H}_N are both reality preserving. For the positivity preserving part, we first deal with the space in (5.11).

In this case, the proof follows from two observations. First, by antisymmetry, the boundary trace of any function Ψ in $\mathrm{H}^1(I_N;\Gamma) \cap \mathcal{H}_N$ must actually vanish on the (possibly larger) symmetrized set $\Gamma_N = \bigcup_{\sigma \in \mathcal{P}_N} \sigma(\Gamma)$. In particular, the restriction $\Psi|_{S_N}$ vanishes on the intersection $\Gamma_N \cap \partial S_N$. The second observation is that $(\gamma \Psi)_+ = \gamma(\Psi_+)$ because this holds for continuous functions and taking the positive part is (e.g., by [MM79, Theorem 1]) a continuous operation in H^1 . Therefore, $(\Psi|_{S_N})_+ \in \mathrm{H}^1(S_N;\Gamma_N \cap \partial I_N)$. In particular, the boundary trace of the extension $T(\Psi_+|_{S_N})$ vanishes on the symmetrized set Γ_N and therefore on Γ . Thus $T(\Psi_+|_{S_N})$ belongs to $\mathrm{H}^1(I_N;\Gamma) \cap \mathcal{H}_N$ and satisfies $T(\Psi_+|_{S_N})|_{S_N} = \frac{1}{\sqrt{N!}}\Psi_+|_{S_N}$. This shows that Ψ_+ belongs to the space in (5.11) and therefore this space is positivity preserving.

For the second statement, it suffices to show that

$$Y \coloneqq \{\Psi|_{S_N} : \Psi \in \mathrm{H}^1_{\alpha}(I_N) \cap \mathcal{H}_N\}$$

is positivity preserving. Indeed, if *Y* is positivity preserving, then the result follows because the intersection of two positivity preserving spaces is positivity preserving, and

$$\{\Psi|_{S_N}: \Psi \in \mathrm{H}^1_{\alpha}(I_N; \Gamma) \cap \mathcal{H}_N\} = Y \cap \mathrm{H}^1(S_N; \Gamma_N \cap \partial S_N).$$

To prove that *Y* is positivity preserving, we first claim that

$$Y = \{\Psi \in H^1(S_N; \Gamma_{\text{int}}) : \Psi(0, x') - \alpha (-1)^{N-1} \Psi(x', 1) = 0 \text{ for a.e. } x' \in S_{N-1} \}.$$
(5.13)

Assuming this claim for the moment, the result follows from the following observation. Since $\alpha(-1)^{N-1} \ge 0$ and $\Psi(0, x') = (-1)^{N-1} \alpha \Psi(x', 1)$ for (a.e.) $x' \in S_{N-1}$, we have

$$\Psi(0, x')_{+} = \max\{\Psi(0, x'), 0\} = \max\{\alpha(-1)^{N-1}\Psi(x', 1), 0\} = (-1)^{N-1}\alpha\Psi(x', 1)_{+}$$

and therefore *Y* is positivity preserving.

To prove the claim, we first note that, by the definition of $H^1_{\alpha}(I_N)$ and antisymmetry, any $\Psi \in H^1_{\alpha}(I_N) \cap \mathcal{H}_N$ satisfies

$$\Psi(0, x_1, ..., x_{N-1}) = \alpha \Psi(1, x_1, ..., x_{N-1}) = -\alpha \Psi(x_1, 1, x_2, ..., x_{N-1}) = (-1)^{N-1} \alpha \Psi(x_1, ..., x_{N-1}, 1)$$

for any $x' = (x_1, ..., x_{N-1}) \in I_{N-1}$; therefore the inclusion \subset in (5.13) holds. To prove the opposite inclusion, we observe that for any $\Psi \in H^1(S_N; \Gamma_{int})$ satisfying $\Psi(0, x') = (-1)^{N-1} \alpha \Psi(x', 1)$ for a.e. $x' \in S_{N-1}$, we have

$$(T\Psi)(0,\sigma x') = \operatorname{sign}(\sigma)\Psi(0,x') = \operatorname{sign}(\sigma)(-1)^{N-1}\alpha\Psi(x',1) = (-1)^{N-1}\alpha(T\Psi)(\sigma x',1) = \alpha(T\Psi)(1,\sigma x')$$
(5.14)

for any $\sigma \in \mathcal{P}_{N-1}$ and a.e. $x' \in S_{N-1}$. As $\{\sigma(S_{N-1})\}_{\sigma \in \mathcal{P}_{N-1}}$ covers I_{N-1} up to finitely many hyperplanes of dimension N-2, equation (5.14) holds for almost every $x' \in I_{N-1}$. Thus $T\Psi \in H^1_{\alpha}(I_N)$ and satisfies $(T\Psi)|_{S_N} = \frac{1}{\sqrt{N!}}\Psi$, which proves the inclusion \supset in (5.13) and completes the proof.

5.4. Proof of Theorem 2.6. We now prove Theorem 2.6.

Proof of Theorem 2.6. Let $\alpha \in \mathbb{R} \setminus \{0\}$ and $v \in \mathcal{V}$, then by Lemma 4.4, the eigenfunctions of the operator $H_N^{\alpha}(v, 0)$ are given by linear combination of Slater determinants of the eigenfunctions of $h_{\alpha}(v) = -\Delta_{\alpha} + v$. In particular, if we denote by $\{\psi_j\}_{j\in\mathbb{N}}$ and $\{\lambda_j\}_{j\in\mathbb{N}}$ the eigenfunctions and eigenvalues of $h_{\alpha}(v)$ ordered in non-decreasing order, then $H_N(v, 0)$ has a non-degenerate ground-state if and only if $\lambda_N < \lambda_{N+1}$. Thus by Theorem 2.3, we have $\lambda_N < \lambda_{N+1}$ for any $N \in \mathbb{N}$ satisfying (2.4), in particular, for N = 2k - 1 if $\alpha \ge 0$ and N = 2k for $\alpha \le 0$. This proves (2.7). The cases of local (or separable) boundary conditions in (2.8) follows from the same argument.

To see that the eigenfunctions are almost everywhere non-vanishing, let $E := \{\psi_j = 0\}$ and pick N > j satisfying (2.4). Then the unique ground-state of $H_N^{\alpha}(v, 0)$ is given by

$$\Psi(x_1,...,x_N) = \frac{1}{\sqrt{N!}} \sum_{\sigma \in \mathcal{P}} \operatorname{sgn}(\sigma) \psi_1(x_{\sigma(1)}) \dots \psi_j(x_{\sigma(j)}) \dots \psi_N(x_{\sigma(N)}).$$

In particular, for $x \in E^N$ we have $x_{\sigma(j)} \in E$ for any $\sigma \in \mathcal{P}_N$, and therefore Ψ vanishes on the set E^N . By the strong UCP in Theorems 2.1 and 2.3, this implies that $|E^N| = |E|^N = 0$, which completes the proof.

6. EIGENVALUE INEQUALITIES

In this section, our goal is to prove Theorem 2.7.

6.1. Unique continuation along the boundary. The key ingredient in the proof of Theorem 2.7 is the following (weak) unique continuation result along the boundary.

Theorem 6.1 (Weak unique continuation along the boundary). Let $\Gamma \subset \partial I_N$, and let Ψ be the unique ground-state of the self-adjoint realization of $a_{v,w}$ with form domain $\mathrm{H}^1(I_N;\Gamma) \cap \mathcal{H}_N$. Then $\gamma \Psi$ can not vanish identically on a relatively open set

$$U \subset \partial I_N \setminus \Gamma_N$$
, where $\Gamma_N = \bigcup_{\sigma \in \mathcal{P}_N} \sigma(\Gamma)$.

Moreover, if $\alpha \in \mathbb{R} \setminus \{0\}$ satisfies (2.4), then the same holds for the ground-state of the self-adjoint realization with form domain $H^1_{\alpha}(I_N; \Gamma) \cap \mathcal{H}_N$.

Proof of Theorem 6.1 in the case of local BCs. To simplify the notation, we set

$$\Gamma_S \coloneqq \Gamma_{\text{int}} \cup (\Gamma_N \cap \partial S_N)$$

where Γ_{int} is the interior boundary of the simplex S_N (see (5.4)). Then, we recall that the ground-state of $H_N(v, w; \Gamma)$ is given by $T\widetilde{\Psi}$, where T is given by (5.6) and $\widetilde{\Psi}$ is the ground-state of the reduced operator $\widetilde{H}_N(v, w; \Gamma)$ defined via (5.8) with form domain (5.11). Hence, it suffices to show that $\widetilde{\Psi}$ can not vanish on a relatively open subset of $\partial S_N \setminus \Gamma_S$. To prove this, we shall assume that $\widetilde{\Psi}$ vanishes on a relatively open set $U \subset \partial S_N \setminus \Gamma_N$ and argue by contradiction.

First, since $U \subset \partial S_N$ is relatively open, we can find an open ball $B \subset \mathbb{R}^N$ such that $B \cap \partial S_N$ is contained in the intersection of U with the interior of one of the exterior faces of S_N . So without loss of generality, let us assume that $B \cap \partial S_N \subset U \cap F_0$, where we recall that $F_0 = \{(x_1, ..., x_N) \in \partial S_N : x_1 = 0\}$. Next, we define $\Omega := S_N \cup B$. Then note that $S_N \subset \Omega \subset \mathbb{R}^N$, $\Gamma_S \subset \partial \Omega$, and Ω is an open bounded and connected subset of \mathbb{R}^N with Lipschitz boundary (see Figure 2). In particular, the restriction map

$$R: \mathrm{H}^{1}(\Omega; \Gamma_{S}) \to \mathrm{H}^{1}(S_{N}; \Gamma_{S}), \quad \Phi \mapsto R\Phi = \Phi|_{S_{N}},$$

is continuous. We can thus define the extended self-adjoint operator H_{Ω} via the sesquilinear form

$$\widetilde{a}_{v,w,\Omega}(\Psi,\Phi) \coloneqq \int_{\Omega \setminus S_N} \overline{\nabla \Psi(x)} \cdot \nabla \Phi(x) dx + \widetilde{a}_{v,w}(R\Psi, R\Phi) = \int_{\Omega} \overline{\nabla \Psi(x)} \cdot \nabla \Phi(x) dx + (T^{\#}v)(R\Psi, R\Phi) + (T^{\#}w)(R\Psi, R\Phi), \quad \Psi, \Phi \in \mathrm{H}^{1}(\Omega; \Gamma_{S}).$$
(6.1)

Since $(\gamma \widetilde{\Psi})|_{\Omega \cap \partial s_N} = 0$, the extension by zero of $\widetilde{\Psi}$ to Ω , denoted also by $\widetilde{\Psi}$, belongs to $H^1(\Omega; \Gamma_S)$ and satisfies

$$\widetilde{a}_{v,w,\Omega}(\widetilde{\Psi},\Phi) = \int_{\Omega} \overline{\nabla \widetilde{\Psi}} \cdot \nabla \Phi dx + (T^{\#}(v+w))(R\widetilde{\Psi},R\Phi) = \widetilde{a}_{v,w}(\widetilde{\Psi},R\Phi) = \lambda \langle \widetilde{\Psi},R\Phi \rangle = \lambda \langle \widetilde{\Psi},\Phi \rangle,$$

where λ is the ground-state energy of $\widetilde{H}_N(v, w; \Gamma)$. In particular, $\widetilde{\Psi}$ is an eigenfunction of H_{Ω} .

Next, notice that the pullback $R^{\#}V$ of any bounded multiplicative function $V \in L^{\infty}(S_N)$ is simply the extension of V by zero to Ω , so in particular, a bounded function in $L^{\infty}(\Omega)$. Consequently, the pullback via R of any sequence of bounded functions in S_N approximating the quadratic form $T^{\#}(v + w)$ in H¹ (as in (5.10)) yields a sequence of bounded functions approximating $R^{\#}T^{\#}(v + w)$. In particular, the operator H_{Ω} defined via (6.1) satisfies the assumptions of Theorem 5.3, and therefore, has a unique and a.e. strictly positive ground-state Ψ_{Ω} .

This now yields a contradiction for the following reason. As we have shown that $\widetilde{\Psi}$ is an eigenfunction of H_{Ω} , it must be either orthogonal or proportional to Ψ_{Ω} . However, $\widetilde{\Psi}$ cannot be orthogonal to Ψ_{Ω} as they are both non-negative (up to a global phase), and $\widetilde{\Psi}$ cannot be parallel to Ψ_{Ω} as the former vanishes on $\Omega \setminus S_N$ while the later is (a.e.) strictly positive in Ω . This completes the proof.



FIGURE 2. Example of extended set Ω in the case N = 2 (left) and N = 3 (right) with interior boundary Γ_{int} (in red), exterior boundary $\partial \Omega \setminus \Gamma_{int}$ (in blue), and $B \cap \partial S_N \subset F_N$ (in yellow).

In the case of non-local boundary conditions, the proof is more involved. The reason is that the exterior faces F_0 and F_N of the simplex are not independent but "glued" to each other in the sense that the boundary conditions along one exterior face must match (up to a constant) the boundary condition along the other exterior face. Hence, we cannot immediately use the extension argument from the previous proof.

To overcome this issue, the first step is to show that, whenever an eigenfunction vanishes on an open subset of the boundary, a meaningful sense can be given to its normal derivative along this set. For this, let us introduce the following definition.

Definition 6.2 (Weak Neumann trace). Let $\alpha \in \mathbb{R} \setminus \{0\}$ and $\Psi \in H^1_{\alpha}(I_N; \Gamma) \cap \mathcal{H}_N$ be an eigenfunction of $H^{\alpha}_N(v, w; \Gamma)$ with eigenvalue λ . Then for any $f \in H^{1/2}(\partial I_N)$, we define the Neumann trace $\partial_v \Psi$ as

$$(\partial_{\nu}\Psi)(f) = \int_{I_N} \overline{\nabla\Psi(x)} \cdot \nabla F(x) + v(\rho_{\Psi,P_NF}) + w(\rho_{2,\Psi,P_NF}) - \lambda \langle \Psi, F \rangle$$
(6.2)

for any $F \in H^1(I_N)$ satisfying $\gamma F = f$, where γ is the standard (Dirichlet) trace and $P_N : L^2(I_N) \to \mathcal{H}_N$ is the projection on the antisymmetric space,

$$(P_N F)(x) = \frac{1}{N!} \sum_{\sigma \in \mathcal{P}_N} \operatorname{sgn}(\sigma) F(\sigma x).$$

It is not hard to see that $\partial_{\nu}\Psi(f)$ is independent of the extension of f. Indeed, if we have two H¹extensions F and F', then $F - F' \in H_0^1(I_N)$, and therefore $P_N(F - F') \in H_0^1(I_N) \cap \mathcal{H}_N$; in particular, $P_N(F - F')$ belongs to the form domain $H_\alpha^1(I_N; \Gamma) \cap \mathcal{H}_N$, and consequently,

$$\begin{split} &\int_{I_N} \overline{\nabla \Psi(x)} \cdot \nabla (F - F')(x) + v(\rho_{\Psi, P_N(F - F')}) + w(\rho_{2,\Psi, P_N(F - F')}) - \lambda \langle \Psi, F - F' \rangle \\ &= \int_{I_N} \frac{1}{N!} \sum_{\sigma \in \mathcal{P}_N} \operatorname{sgn}(\sigma) \overline{\nabla(\sigma_{\#}\Psi)(x)} \cdot \nabla(F - F')(x) dx + v(\rho_{\Psi, P_N(F - F')}) + w(\rho_{2,\Psi, P_N(F - F')}) - \lambda \langle \Psi, P_N(F - F') \rangle \\ &= a_{v,W}(\Psi, P_N(F - F')) - \lambda \langle \Psi, P_N(F - F') \rangle = 0, \end{split}$$

where the first equality hold because $P_N \Psi = \Psi$ and the second equality because Ψ is an eigenfunction of $H_N^{\alpha}(v, w; \Gamma)$ with eigenvalue λ . Moreover, the Neumann trace $f \in H^{1/2}(\partial I_N) \to (\partial_v \Psi)(f)$ is continuous since we can pick F = Jf for any right inverse $J : H^{1/2}(\partial I_N) \to H^1(I_N)$ of the Dirichlet trace operator γ . However, it should be noted that this is a completely adhoc definition of the Neumann trace as, a priori, there is no connection between $\partial_v \Psi$ and the normal derivative of Ψ along the boundary.

Fortunately, it turns out that this definition is meaningful along the vanishing set of Ψ on the boundary. To make this statement precise, we shall use the following lemma.

Lemma 6.3 (Interior approximation). Let $\Psi \in H^1(I_N)$ satisfy $(\gamma \Psi)|_U = 0$ for some relatively open subset $U \subset \partial I_N$. Let $V \subset \subset U$ and define

$$\eta_{\epsilon}(x) \coloneqq \eta(d_V(x)/\epsilon), \quad \text{where} \quad d_V(x) \coloneqq \inf_{x \in V} ||x - y||_{\epsilon}$$

for some $\eta \in \text{Lip}(\mathbb{R}, \mathbb{R})$ satisfying $0 \le \eta \le 1$, $\eta(x) = 1$ for $|x| \le 1/2$, and $\eta(x) = 0$ for $|x| \ge 1$. Then the function $\Psi_{\epsilon} = \Psi(1 - \eta_{\epsilon}) \in H^1(I_N; U)$ satisfies

$$\lim_{\epsilon \downarrow 0} \|\Psi - \Psi_{\epsilon}\|_{\mathrm{H}^{1}} = \lim_{\epsilon \downarrow 0^{+}} \|\eta_{\epsilon}\Psi\|_{\mathrm{H}^{1}} = 0.$$

Proof. Since $\eta_{\epsilon}(x) \to 0$ as $\epsilon \downarrow 0$ for every $x \in I_N$, by dominated convergence we have

$$\|\Psi - \Psi_{\epsilon}\|_{\mathrm{L}^2}^2 = \int_{I_N} |\Psi(x)|^2 |\eta_{\epsilon}(x)|^2 \mathrm{d}x \to 0 \quad \mathrm{as} \quad \epsilon \downarrow 0.$$

Next, notice that the map $x \mapsto d_V(x)$ is globally Lipschitz with Lipschitz constant smaller than 1. Thus from Rademacher's theorem, the function $d_V(x)$ has a bounded weak gradient, and therefore,

$$\begin{split} \|\nabla(\Psi - \Psi_{\epsilon})\|_{L^{2}}^{2} &\leq 2 \int_{I_{N}} |\nabla\Psi(x)|^{2} |\eta_{\epsilon}(x)|^{2} + |\Psi(x)|^{2} \frac{|\dot{\eta}(d_{V}(x)/\epsilon)|\nabla d_{V}(x)|^{2}}{\epsilon^{2}} \mathrm{d}x \\ &\lesssim \int_{I_{N}} |\nabla\Psi(x)|^{2} |\eta_{\epsilon}(x)|^{2} \mathrm{d}x + \frac{1}{\epsilon^{2}} \int_{V_{\epsilon} \cap I_{N}} |\Psi(x)|^{2} \mathrm{d}x, \end{split}$$

where $V_{\epsilon} := \{x : d_V(x) \le 2\epsilon\}$. The first term converges to zero again by dominated convergence. For the second term, we can assume, via a partition of the unity argument, that $V_{\epsilon} \cap I_N$ is the hypergraph of a Lipschitz function, i.e., up to a rotation and translation, there exists $\kappa : \mathbb{R}^{N-1} \to \mathbb{R}$ Lipschitz such that $\partial I_N \cap V_{\epsilon} = \{(x', \kappa(x')) : x' \in W_{\epsilon}\}$ for some open bounded set $W_{\epsilon} \subset \mathbb{R}^{N-1}$ and $V_{\epsilon} \cap I_N = \{(x', x_N) : x' \in W_{\epsilon}\}$ for some open bounded set $W_{\epsilon} \subset \mathbb{R}^{N-1}$ and $V_{\epsilon} \cap I_N = \{(x', x_N) : x' \in W_{\epsilon}, \kappa(x') < x_N < y(x', \epsilon)\}$, where $y(x', \epsilon) - \kappa(x') \le \epsilon$. Since $V_{\epsilon} \cap \partial I_N \subset U$ for $\epsilon > 0$ small enough, we have $\Psi(x', \kappa(x')) = 0$, and therefore

$$\frac{1}{\epsilon^2} \int_{V_{\epsilon}} |\Psi(x)|^2 \mathrm{d}x = \frac{1}{\epsilon^2} \int_{W_{\epsilon}} \mathrm{d}x' \int_{\kappa(x')}^{y(x',\epsilon)} |\Psi(x',x_N)|^2 \mathrm{d}x_N = \frac{1}{\epsilon^2} \int_{W_{\epsilon}} \int_{\kappa(x')}^{y(x',\epsilon)} \left| \int_{\kappa(x')}^{x_N} \partial_{x_N} \Psi(x',t) \mathrm{d}t \right|^2 \mathrm{d}x.$$

If we now apply Cauchy-Schwarz, use the change of variables $z = (x_N - \kappa(x'))/\epsilon$, and use the estimate $y(x', \epsilon) - \kappa(x') \le C\epsilon$, we find

$$\begin{split} \frac{1}{\epsilon^2} \int_{W_\epsilon} \int_{\kappa(x')}^{y(x',\epsilon)} \left| \int_{\kappa(x')}^{x_N} \partial_{x_N} \Psi(x',t) \mathrm{d}t \right|^2 \mathrm{d}x &\leq \int_{W_\epsilon} \int_{\kappa(x')}^{y(x',\epsilon)} \left(\int_{\kappa(x')}^{x_N} |\partial_{x_N} \Psi(x',t)|^2 \mathrm{d}t \right) \frac{x_N - \kappa(x')}{\epsilon^2} \mathrm{d}x_N \mathrm{d}x' \\ &\lesssim \int_{W_\epsilon} \int_0^C \int_{\kappa(x')}^{\kappa(x') + \epsilon z} |\partial_{x_N} \Psi(x',t)|^2 \mathrm{d}t \mathrm{d}z \mathrm{d}x' \\ &\lesssim \int_{V_\epsilon \cap I_N} |\partial_{x_N} \Psi(x',t)|^2 \mathrm{d}x' \mathrm{d}t. \end{split}$$

Hence, by dominated convergence, the limit $\epsilon \downarrow 0^+$ goes to zero, which completes the proof.

We now show that the Neumann trace is meaningful along any open subset of the boundary where Ψ vanishes. The key to prove this is the locality of the form associated to $H_N(v, w)$.

Lemma 6.4 (Neumann trace formula). Let $U \subset \partial I_N$ be a relatively open subset of I_N and suppose that U is compactly contained in the interior of a face of ∂I_N . Let Ψ be an eigenfunction of $H_N^{\alpha}(v, w; \Gamma)$ such that $(\gamma \Psi)|_U = 0$. Then for any $f \in C_c^{\infty}(U)$, we have

$$(\partial_{\nu}\Psi)(f) = -\lim_{\epsilon \downarrow 0^{+}} \frac{\langle \gamma_{\epsilon}\Psi, f \rangle_{L^{2}(U)}}{\epsilon}, \qquad (6.3)$$

where the above limit exists and $\gamma_{\epsilon}\Psi$ denotes the Dirichlet trace of Ψ along the hyperplane parallel and at distance ϵ of the face containing U.

Proof. Without loss of generality, let us assume that $U \subset C E_0 := \{x = (x_1, x') \in \partial I_N : x_1 = 0, x' \in I_{N-1}\}$. Then, we let $\beta \in C_c^{\infty}(\mathbb{R}; \mathbb{R})$ be a standard mollifier, i.e., $0 \leq \beta \leq 1$, $\operatorname{supp}(\beta) \subset [-1, 1]$ and $\beta = 1$ on a neighborhood of 0, and define the function *F* as

$$F(x_1, x') \coloneqq \beta(x_1) f(x') \quad \text{for } (x_1, x') \in I_N.$$

Clearly, *F* is an H¹-extension of *f*, i.e., $F \in H^1(I_N)$ and $\gamma F = f$.

Next, let $V \subset \partial I_N$ be such that $\operatorname{supp}(f) \subset V \subset U$ and define η_{ϵ} as in Lemma 6.3. Then $F_{\epsilon} := (1 - \eta_{\epsilon})F \in \operatorname{H}_0^1(I_N)$ and therefore

$$0 = a_{v,w}(\Psi, P_N F_{\epsilon}) - \lambda \langle \Psi, P_N F_{\epsilon} \rangle = \int_{I_N} (1 - \eta_{\epsilon}) \overline{\nabla \Psi} \cdot \nabla F dx + v(\rho_{\Psi, P_N F_{\epsilon}}) + w(\rho_{2,\Psi, P_N F_{\epsilon}}) - \lambda \langle \Psi, F_{\epsilon} \rangle$$
(6.4)

$$-\int_{I_N} F(x)\overline{\nabla\Psi(x)} \cdot \nabla\eta_{\epsilon}(x) \mathrm{d}x, \qquad (6.5)$$

where the first equality follows because Ψ is an eigenfunction of $H^{\alpha}_{\mathcal{N}}(v, w; \Gamma)$ with eigenvalue λ .

The key observation now is the following: since

$$\overline{\Psi(x)}(P_N F_{\epsilon})(x) = \frac{1}{N!} \sum_{\sigma \in \mathcal{P}_N} \operatorname{sgn}(\sigma) \overline{\Psi(x) (1 - \eta_{\epsilon}(\sigma x))} F(\sigma x),$$

and $\gamma \Psi$ vanishes on $\sigma^{-1}(U)$ for any $\sigma \in \mathcal{P}_N$ (by antisymmetry), from Lemma 6.3 we have

$$\lim_{\epsilon \downarrow 0^+} \|\Psi(1 - \sigma_{\#}\eta_{\epsilon}) - \Psi\|_{\mathrm{H}^1} = 0 \quad \text{for any } \sigma \in \mathcal{P}_N.$$

Consequently, by Lemma 3.13,

$$\lim_{\epsilon \downarrow 0^+} \rho_{\Psi, P_N F_{\epsilon}} = \rho_{\Psi, P_N F} \quad \text{in } \mathrm{H}^1(I) \quad \text{and} \quad \lim_{\epsilon \downarrow 0^+} \rho_{2, \Psi, P_N F_{\epsilon}} = \rho_{2, \Psi, P_N F} \quad \text{in } \mathrm{W}^{-1, p}(I_2).$$

Using these convergence results and dominated convergence for the kinetic term, we see that equation (6.4) converges to $\partial_{\nu}\Psi(f)$ as $\epsilon \downarrow 0^+$. In particular,

$$\partial_{\nu}\Psi(f) = \lim_{\epsilon \downarrow 0^+} \int_{I_N} F(x) \overline{\nabla \Psi(x)} \cdot \nabla \eta_{\epsilon}(x) \mathrm{d}x,$$

where the limit exists. To compute this limit, let us take

$$\eta(x) = \begin{cases} 1, & \text{for } x \le 1/2, \\ 2 - 2x, & \text{for } \frac{1}{2} \le x \le 1, \\ 0, & \text{otherwise.} \end{cases}$$

Since supp(*F*) \subset [0, 1] × *V*, we have $d_V(x) = x_1$ for any $x = (x_1, x') \in$ supp(*F*). In particular, $\eta_{\epsilon}(x) = \eta(x_1/\epsilon)$ for $x \in$ supp(*F*). Moreover, for $\epsilon > 0$ small enough, we have $\beta(x_1) = 1$ on supp($\eta_{\epsilon}(x)$). Therefore

$$\partial_{\nu}\Psi(f) = \lim_{\epsilon \downarrow 0^{+}} \int_{I_{N}} F(x) \overline{\nabla \Psi(x)} \cdot \nabla \eta_{\epsilon}(x) dx$$

$$= \lim_{\epsilon \downarrow 0^{+}} \int_{[\epsilon/2,\epsilon] \times V} f(x') \overline{\partial_{x_{1}}\Psi(x_{1},x')} \left(\frac{-2}{\epsilon}\right) dx_{1} dx'$$

$$= \lim_{\epsilon \downarrow 0^{+}} \frac{\langle \gamma_{\epsilon/2}\Psi, f \rangle - \langle \gamma_{\epsilon}\Psi, f \rangle}{\epsilon/2}$$
(6.6)

To complete the proof, we now write

$$\langle \gamma_{2^{-n}\epsilon}\Psi, f \rangle - \langle \gamma_{\epsilon}\Psi, f \rangle - (\epsilon - 2^{-n}\epsilon)\partial_{\nu}\Psi(f) = \sum_{k=1}^{n} \left(\langle \gamma_{2^{-k}\epsilon}\Psi, f \rangle - \langle \gamma_{2^{-k+1}\epsilon}f \rangle - 2^{-k}\epsilon\partial_{\nu}\Psi \right)$$

Thus, by (6.6), for any $\delta > 0$ we can take $\epsilon > 0$ so small that the right hand side is controlled by $\delta \epsilon$. Taking the limit $n \to \infty$ and using that $\gamma \Psi = 0$ on *U*, we conclude that

$$|\epsilon \partial_{\nu} \Psi(f) + \langle \gamma_{\epsilon} \Psi, f \rangle| \leq \delta \epsilon.$$

Diving this expression by ϵ and taking the limits $\epsilon \downarrow 0^+$ and $\delta \downarrow 0^+$ completes the proof.

We now show that the ground-state Ψ of $H_N^{\alpha}(v, w; \Gamma)$ satisfies $\partial_v \Psi = 0$ along any open set $U \subset \partial S_N$ such that $\gamma \Psi|_U = 0$. To prove this, the key observation is that Ψ is non-negative in S_N by the Perron-Frobenius theorem.

Lemma 6.5 (Vanishing of normal derivative). Let $\alpha \in \mathbb{R} \setminus \{0\}$ satisfy (2.4) and let Ψ be the groundstate of $H_N^{\alpha}(v, w; \Gamma)$. Suppose that $(\gamma \Psi)|_U = 0$ for some (relatively) open set $U \subset \subset \operatorname{int} F_0 \setminus \Gamma_N$, where $F_0 = \{(x_1, x_2, ..., x_N) \in \partial S_N : x_1 = 0\}$. Then

$$\partial_{\nu}\Psi(f) = 0 \quad \text{for any } f \in \mathrm{H}^{1/2}(\partial\Omega) \text{ with } \mathrm{supp}(f) \subset U.$$
 (6.7)

Proof. Since any function $f \in H^{1/2}(\partial I_N)$ with $\operatorname{supp}(f) \subset U$ can be approximated by functions in $C_c^{\infty}(U)$ it suffices to prove (6.7) for $f \in C_c^{\infty}(U)$. To this end, we first claim that

$$\partial_{\nu}\Psi(f) \le 0.$$
 for any $f \in C_c^{\infty}(U)$ such that $f \ge 0.$ (6.8)

Indeed, since $\Psi \ge 0$ almost everywhere in S_N (by Theorem 5.6), this claim follows from the formula in Lemma 6.4.

Next, we recall that, by the definition of $H^1_{\alpha}(I_N)$ and antisymmetry, Ψ satisfies

$$(\gamma \Psi)(0, x') - (-1)^{N-1} \alpha(\gamma \Psi)(x', 1) = 0$$
 for a.e. $x' \in I_{N-1}$

Therefore $\gamma \Psi$ also vanishes on the set $U' := \{(x', 1) : (0, x') \in U\} \subset F_N$. Consequently, (6.8) also holds for $f \in H^{1/2}(\partial I_N)$ with $supp(f) \subset U'$. Now let $f \in C_c^{\infty}(U)$ and define

$$f'(x) \coloneqq \begin{cases} (-1)^{N-1} \alpha f(0, x') & \text{for } x = (x', 1) \in U', \\ 0, & \text{for } x \in \partial I_N \setminus U'. \end{cases}$$

Let Jf and Jf' be extensions of f and f' in $H^1(I_N)$. Then, recalling the proof of Lemma 5.7, we see that $P_N(Jf + Jf') \in H^1_\alpha(I_N; \Gamma) \cap \mathcal{H}_N$. Therefore, by the definition of the Neumann trace, we have

$$\partial_{\nu}\Psi(f) + \partial_{\nu}\Psi(f') = \partial_{\nu}\Psi(f+f') = a_{v,w}\left(\Psi, P_N(Jf+Jf')\right) - \lambda\langle\Psi, P_N(Jf+Jf')\rangle = 0.$$

Thus if $f \ge 0$, then $f' \ge 0$ by (2.4), and therefore $\partial_{\nu}\Psi(f) = \partial_{\nu}\Psi(f') = 0$ by (6.8). This concludes the proof.

We can now complete the proof of Theorem 6.1.

Proof of Theorem 6.1 in the case of non-local BCs. Let $U \subset \partial I_N \setminus \Gamma_N$ be relatively open. We shall assume that $\Psi|_U = 0$ and obtain a contradiction. To this end, first note that, by the antisymmetry of Ψ , the Dirichlet trace $\gamma \Psi$ vanishes on the symmetrized set $U_N = \bigcup_{\sigma} \sigma(U)$. Moreover, after possibly shrinking U_N , we can assume that $U_N \cap \partial S_N$ is compactly contained on the interior of an exterior face of ∂S_N . So without loss of generality, we assume that $U \subset F_0 = \{(x_1, ..., x_N) \in \partial S_N : x_1 = 0\}$. Hence, if we denote by $\widetilde{\Psi} = T^{-1}\Psi = \sqrt{N!}\Psi|_{S_N}$ the ground-state of the reduced operator $\widetilde{H}_N^{\alpha}(v, w; \Gamma)$, Lemma 6.5 implies that

$$\partial_{\nu}\widetilde{\Psi}(f) \coloneqq \widetilde{a}_{v,w}(\widetilde{\Psi}, Jf) - \lambda \langle \widetilde{\Psi}, Jf \rangle = a_{v,w}(T\widetilde{\Psi}, TJf) - \lambda \langle T\widetilde{\Psi}, TJf \rangle = \sqrt{N!} \partial_{\nu} \Psi(f) = 0$$

for any $f \in H^{1/2}(\partial S_N)$ such that $\operatorname{supp}(f) \subset U$, where $J : H^{1/2}(\partial S_N) \to H^1(S_N)$ is any right inverse of the Dirichlet boundary trace on S_N .

We now claim that this is equivalent to

$$\widetilde{a}_{v,w}(\widetilde{\Psi},\Phi) - \lambda \langle \widetilde{\Psi},\Phi \rangle = 0, \quad \text{for any } \Phi \in Q_{\alpha}(W), \tag{6.9}$$

where

$$Q_{\alpha}(W) := \{ \Phi \in \mathrm{H}^{1}(S_{N}; \Gamma_{N} \cup \Gamma_{\mathrm{int}}) : (\gamma \Phi)(0, x') - (-1)^{N-1} \alpha(\gamma \Phi)(x', 1) \quad \text{for } (0, x') \in F_{0} \setminus W \},$$

T. CARVALHO CORSO

for any neighborhood $W \subset U \subset F_0$. Indeed, if $\Phi \in Q_\alpha(W)$, then Φ can be written as the sum of a function in the form domain of the reduced operator $\widetilde{H}_N^\alpha(v, w; \Gamma)$ and a function with trace supported in $\overline{W} \subset U$. To be precise, for any $\Phi \in Q_\alpha(W)$, we can define

$$f_{\Phi}(x) := \begin{cases} (-1)^{N-1} \frac{1}{\alpha} (\gamma \Phi)(x', 1), & \text{if } x = (0, x') \text{ for } x' \in S_{N-1}, \\ (\gamma \Phi)(x', 1), & \text{if } x = (x', 1) \text{ with } x' \in S_{N-1}, \\ 0, & \text{otherwise,} \end{cases}$$

and note that $f_{\Phi} \in H^{1/2}(\partial S_N)$ because f_{Φ} is locally equal to $\gamma \Phi$ or a rigid motion of $\gamma \Phi$. Moreover, by the definition of $Q_{\alpha}(W)$, we have $\operatorname{supp}(\gamma \Phi - f_{\Phi}) \subset \overline{W}$. Thus, since Jf_{Φ} belongs to the form domain of $\widetilde{H}_N^{\alpha}(v, w; \Gamma)$ (namely (5.13) intersected with $H^1(S_N; \Gamma_N \cap \partial S_N)$), we have the desired decomposition $\Phi = (\Phi - Jf_{\Phi}) + Jf_{\Phi}$, which proves the claim in (6.9).

Next, notice that (6.9) is equivalent to saying that $\widetilde{\Psi}$ is an eigenfunction of the operator associated to the form $\widetilde{a}_{v,w}$ but with the larger form domain $Q_{\alpha}(W)$. The important observation now is that the trace of functions in $Q_{\alpha}(W)$ have no restriction inside W. Therefore, we can repeat the extension argument in the proof of the case of local BCs. More precisely, we extend S_N to Ω by adding an open ball whose intersection with ∂S_N is compactly contained in W. Then $\widetilde{\Psi}$ extended by zero is also an eigenfunction of the extended operator H_{Ω} associated to (6.1) but this time with form domain

$$Q_{\Omega} := \{ \Psi \in \mathrm{H}^{1}(\Omega; \Gamma) : \Psi|_{U \cap \partial \Omega} = 0 \quad \text{and} \quad (\gamma \Phi)(0, x') - (-1)^{N-1} \alpha(\gamma \Phi)(x', 1) \quad \text{for } (0, x') \in F_{0} \setminus W \}$$

Thus, recalling assumption (2.4), one can repeat the arguments in the proof of Lemma 5.7 to show that Q_{Ω} is positivity and reality preserving. In particular, Theorem 5.3 applies to H_{Ω} and therefore its ground-state Ψ_{Ω} is non-degenerate and almost everywhere strictly positive in Ω . This in turn implies that Ψ_{Ω} is either orthogonal or parallel to $\widetilde{\Psi}$. However, neither of these options are possible because both functions are strictly positive in S_N but $\widetilde{\Psi}$ vanishes identically in $\Omega \setminus S_N$ while Ψ_{Ω} does not. This yields the desired contradiction and completes the proof.

6.2. Proof of Theorems 2.7. We can now complete the proof of Theorem 2.7.

Proof of Theorem 2.7. As mentioned in Remark 2.2, the inequality $\lambda_1(\Gamma) \leq \lambda_1(\Gamma')$ is immediate from the variational principle. Hence, it suffices to show that equality cannot hold. For this, note that, if $\lambda_1(\Gamma) = \lambda_1(\Gamma')$, then by the variational principle and the non-degeneracy theorem, the ground-state of $H_N^{\alpha}(v, w; \Gamma)$ and $H_N^{\alpha}(v, w; \Gamma')$ are the same. Hence the ground-state Ψ of $H_N^{\alpha}(v, w; \Gamma)$ vanishes on Γ' . As Ψ is antisymmetric, it must also vanish in the symmetrized set Γ'_N . Hence, Ψ vanishes on $\Gamma'_N \setminus \Gamma_N$. As this set contains a relatively open subset of the boundary by assumption, this is not possible by Theorem 6.1. We thus have a contradiction, which concludes the proof.

Acknowledgements

The author thanks Chokri Manai, Timo Weidl, Robert van Leeuwen, and Markus Penz for fruitful discussions and helpful comments on this article. The author is also grateful to Rafael Antonio Lainez Reyes for pointing out the references on Sobolev spaces.

T.C. Corso acknowledges funding by the *Deutsche Forschungsgemeinschaft* (DFG, German Research Foundation) - Project number 442047500 through the Collaborative Research Center "Sparsity and Singular Structures" (SFB 1481).

Appendix A. Quadratic forms and self-adjoint extensions

In this section we recall the basic theory of self-adjoint extensions via quadratic forms. The results presented here are standard and can be found in several classical references [RS75, RS78, AGHH88, Tes14]. We shall therefore omit the proofs, with an exception for part A.3 for which we have no reference.

A.1. Basic definitions. First, let us recall the definition of sesquilinear forms and self-adjoint operators.

Definition A.1 (Quadratic forms). We say that $a : Q \times Q \to \mathbb{C}$ is a (densely defined) sesquilinear form in a Hilbert space \mathcal{H} if Q is a dense vector subspace of \mathcal{H} and a is antilinear in the first argument and linear in the second one. In this case, we call Q the form domain of a. Moreover, we say that

(i) *a* is symmetric if $a(\Psi, \Phi) = \overline{a(\Phi, \Psi)}$ for any $\Phi, \Psi \in Q$.

(ii) *a* is semi-bounded on \mathcal{H} if $a(\Psi, \Psi) \in \mathbb{R}$ and

$$a(\Psi, \Psi) \ge C \|\Psi\|_{\mathcal{H}}^2$$
, for any $\Psi \in Q$ and some $C \in \mathbb{R}$.

(iii) *a* is closed in \mathcal{H} if for any sequence $\{\Psi_n\}_{n\in\mathbb{N}}\subset Q$ satisfying

$$\lim_{n\to\infty}\Psi_n=\Psi\quad\text{in }\mathcal{H}\text{ and }\quad\lim_{\max\{m,n\}\to\infty}a(\Psi_n-\Psi_m,\Psi_n-\Psi_m)=0,$$

it holds that $\Psi \in Q$ and $\lim_{n\to\infty} a(\Psi - \Psi_n, \Psi - \Psi_n) = 0$.

Remark (Semibounded implies symmetry). By definition any semibounded form satisfies $a(\Psi, \Psi) \in \mathbb{R}$ for $\Psi \in Q$. In particular, from the polarization identity, any semibounded form is automatically symmetric.

Remark (Closed and semibounded). If *a* is semibounded on \mathcal{H} with lower bound *C*, then *a* is closed if and only if *Q* is a (complete) Hilbert space with respect to the norm

$$\|\Psi\|_{a} = a(\Psi, \Psi) + (C+1) \|\Psi\|_{\mathcal{H}}^{2}.$$
(A.1)

Definition A.2 (Self-adjoint). We say that a linear operator $A : D(A) \subset \mathcal{H} \to \mathcal{H}$ is a semibounded self-adjoint operator if it satisfies the following conditions:

- (i) (Densely defined) $D(A) \subset \mathcal{H}$ is dense.
- (ii) (Symmetric) $\langle \Psi, A\Phi \rangle_{\mathcal{H}} = \langle A\Psi, \Phi \rangle_{\mathcal{H}}$, for any $\Psi, \Phi \in D(A)$.
- (iii) (Semibounded) there exists C > 0 such that

$$\langle \Psi, A\Psi \rangle_{\mathcal{H}} \ge C \|\Psi\|_{\mathcal{H}}^2$$
, for any $\Psi \in D(A)$.

(iv) (Self-adjoint) $D(A^*) = D(A)$, where

$$D(A^*) := \{ \Psi \in \mathcal{H} \mid \text{such that the linear map} \mid \Phi \in D(A) \mapsto \langle \Psi, A\Phi \rangle_{\mathcal{H}} \mid \text{is bounded in } \mathcal{H} \}$$

A.2. Quadratic forms and self-adjoint operators. The main result on the theory of self-adjoint extensions via quadratic forms is the following. For a proof, see [RS75].

Theorem A.3 (Equivalence between quadratic forms and self-adjoint operators). Let A be a positive selfadjoint operator on a Hilbert space \mathcal{H} , then the domain $D((A + 1)^{\frac{1}{2}}) \subset \mathcal{H}$ is dense and the sesquilinear form

$$a_A(\Psi, \Phi) = \langle (A+1)^{\frac{1}{2}}\Psi, (A+1)^{\frac{1}{2}}\Phi \rangle - \langle \Psi, \Phi \rangle$$

is a densely defined, symmetric, positive, and closed quadratic form on \mathcal{H} . Conversely, for any densely defined, positive, symmetric and closed quadratic form a, there exists a unique self-adjoint positive operator A such that $a_A = a$.

Remark (Associated quadratic form). We call a_A the associated form of A.

Remark (The shifting trick). The above construction works for any semibounded operator H by first shifting H by a large enough constant such that A = H + C is positive, taking the associated form of A, and then shifting it back to $a_H = a_A - C\langle \cdot, \cdot \rangle_{\mathcal{H}}$. In particular, Theorem A.3 a one-to-one correspondence between semibounded self-adjoint operators and semibounded closed forms.

We now state the celebrated KLMN theorem [RS75, Theorem X.17]. For this, we recall the following definition.

Definition A.4 (Relatively bounded). Let *a* be a closed semibounded form on a Hilbert space \mathcal{H} with form domain *Q*. Then we say that a symmetric sesquilinear form $b : Q \times Q \to \mathbb{C}$ is *a*-bounded if there exists c, C > 0 such that

$$|b(\Psi, \Psi)| \le ca(\Psi, \Psi) + C \|\Psi\|_{\mathcal{H}}^2, \quad \text{for any } \Psi \in Q.$$
(A.2)

The infimum over all c > 0 for which (A.2) holds is called the *a*-bound of *b*. Moreover, if this infimum is zero, then we say that *b* is infinitesimally *a*-bounded.

Theorem A.5 (KLMN). Let a be a closed semibounded form on a Hilbert space \mathcal{H} and suppose that b is a relatively a-bounded with relative bound < 1. Then the form a + b is closed and semibounded, and therefore, there exists a unique self-adjoint operator associated to this form. Moreover, the norm in (A.1) induced by a + b is equivalent to the norm induced by a.

Remark A.6 (Discrete spectrum and form domain). A simple but useful consequence of the KLMN theorem is that, if the form domain of *a* is compactly embedded in \mathcal{H} , then the self-adjoint operator associated to *a*+*b* has purely discrete spectrum. In particular, this operator has an \mathcal{H} -orthonormal basis of eigenfunctions.

A.3. **Pull-back**. We now introduce the notion of the pull-back of a form via a linear transformation. We then collect some elementary properties of the pull-back map.

Definition A.7 (Pull-back). Let $T : Q \to Q'$ be a linear bounded operator between Hilbert spaces Q and Q'. Then for any quadratic form a with form domain Q', we define the pull-back of a via T as the form

$$T^{\#}a: Q \times Q \to \mathbb{C}$$
 $(T^{\#}a)(\Psi, \Phi) = a(T\Psi, T\Phi).$

It is immediate to verify that the pull-back preserves symmetry. The next lemma gives sufficient condition on T for the pull-back to also preserve semiboundeness and closedness.

Lemma A.8 (Pull-back properties). Let $T : \mathcal{H} \to \mathcal{H}'$ be a bounded linear operator such that $T(Q') \subset Q$ for some dense subspaces $Q \subset \mathcal{H}$ and $Q' \subset \mathcal{H}'$. Then the pull-back of any semibounded form in \mathcal{H}' with form domain Q' is a semibounded form in \mathcal{H} with form domain Q. Moreover, if T is bijective and T(Q) = Q', then the pull-back of any closed form with form domain Q' is a closed form with form domain Q.

Proof. For the first part note that if *a* is semibounded in \mathcal{H}' and *T* bounded from \mathcal{H} to \mathcal{H}' , then

$$(T^{\#}a)(\Psi,\Psi) = a(T\Psi,T\Psi) \ge C \|T\Psi\|_{\mathcal{H}'}^2 \ge \min\{0,C\} \|T\|_{\mathcal{H}\to\mathcal{H}'} \|\Psi\|_{\mathcal{H}}^2$$

Thus $T^{\sharp}a$ is semibounded in \mathcal{H} . For the second part, let $\Psi_n \subset Q$ be a sequence such that $\Psi_n \to \Psi$ in \mathcal{H} and

$$\|\Psi_n - \Psi_m\|_{T^{\#}a} = (T^{\#}a)(\Psi_n - \Psi_m, \Psi_n - \Psi_m) \to 0 \quad \text{as} \quad \max\{m, n\} \to \infty.$$

Then, $T\Psi_n \to T\Psi$ in \mathcal{H}' and $a(T\Psi_n - T\Psi_m, T\Psi_n - T\Psi_m) \to 0$ as $m, n \to \infty$. As *a* is closed, this implies that $T\Psi \in Q$ and $a(T\Psi - T\Psi_n, T\Psi - T\Psi_n) \to 0$. Since *T* is bijective and T(Q) = Q', we have $\Psi = T^{-1}T\Psi \in Q'$ and therefore $(T^{\#}a)$ is closed.

References

- [AGHK84] S. ALBEVERIOR, F. GESZTESY, R. HØEGH-KROHN, and W. KIRSCH, On point interactions in one dimension. Journal of Operator Theory 12 (1984), no.1:101–26. MR 757115. Zbl 0561.35023.
- [AGHH88] S. ALBEVERIO, F. GESZTESY, R. HØEGH-KROHN, and H. HOLDEN, Solvable Models in Quantum Mechanics, Springer-Verlag, (1988). MR 958850 Zbl 0679.46057.
- [AHM08] S. ALBEVERIO, A. HRYNIV, and R. MYKYTYUK, On spectra of non-self-adjoint Sturm–Liouville operators. Selecta Mathematica. New Series 13 (2008), no.4, pp. 571–599. MR 2403304. Zbl 1149.34003.
- [AK00] S. ALBEVERIO and P. KURASOV, Singular perturbations of differential operators, Cambridge University Press, (2000). MR 1752110. Zbl 0945.47015.

[BFK+17] J. BEHRNDT, R. L. FRANK, C. KÜHN, V. LOTOREICHIK, and J. ROHLEDER, Spectral theory for Schrödinger operators with δ-interactions supported on curves in R³ Annales Henri Poincaré 18, no.4, pp. 1305–47. MR 3626305. Zbl 1378.81026.

- [BM18] H. BREZIS and P. MIRONESCU, Gagliardo-Nirenberg inequalities and non-inequalities: the full story, Annales de l'Institut Henri Poincaré C. Analyse Non Linéaire 35 (2018), no.5, pp. 1355–1376. MR 3813967. Zbl 1401.46022.
- [Bra85] J. BRASCHE, Perturbation of Schrödinger Hamiltonians by measures—Self-adjointness and lower semiboundedness, Journal of Mathematical Physics 26 (1985), no.4, pp. 621–26. MR 785672. Zbl 0589.35031.
- [BRS18] J. BEHRNDT, J. ROHLEDER, and S. STADLER, Eigenvalue inequalities for Schrödinger operators on unbounded Lipschitz domains, *Journal of Spectral Theory* 8 (2018), no. 2, pp. 493–508. MR 3812805. Zbl 1423.35087.
- [Cor25] T. C. CORSO, v-representability and Hohenberg-Kohn theorem for non-interacting Schrödinger operators with distributional potentials in the one-dimensional torus, *Journal of Physics A: Mathematical and Theoretical* 58 (2025), pp. 125203.

[Dav13] E. B. DAVIES, Singular Schrödinger operators in one dimension, Mathematika. A Journal of Pure and Applied Mathematics 59 (2013), no.1, pp. 141–159. MR 3028176. Zbl 1282.34087.

[EGNT13a] J. ECKHARDT, F. GESZTESY, R. NICHOLS and G. TESCHL, Inverse spectral theory for Sturm-Liouville operators with distributional potentials, *Journal of the London Mathematical Society* 88 (2013), pp. 801–828. MR 3106738. Zbl 1282.34087.
 [EGNT13b] J. ECKHARDT, F. GESZTESY, R. NICHOLS, and G. TESCHL, Weyl-Titchmarsh theory for Sturm-Liouville operators with

distributional potentials, *Opuscula Mathematica* 33 (2013), pp. 467–563. MR 3105767. Zbl 1282.34088. [EGNT15] J. ECKHARDT, F. GESZTESY, R. NICHOLS, and G. TESCHL, Supersymmetry and Schrödinger-type operators with distribu-

tional matrix-Valued potentials, *Journal of Spectral Theory* 4 (2015), no.4, pp. 715–68. MR 3299812. Zbl 1320.34116.

[[]EP16] P. EXNER and J. ROHLEDER, Generalized interactions supported on hypersurfaces, *Journal of Mathematical Physics* 57, no.4, 041507. MR 3490341. Zbl 1339.81038.

[[]Eva10] L. C. EVANS, Partial differential equations, American Mathematical Society (2010). MR 2597943. Zbl 1194.35001.

[[]FS75] W. FARIS and B. SIMON, Degenerate and non-degenerate ground states for Schrödinger operators, Duke Mathematical Journal 42 (1975), no.3, pp. 559–56. MR 381571. Zbl 0339.35033.

[[]Filo5] N. FILONOV, On an inequality between Dirichlet and Neumann eigenvalues for the Laplace operator, St. Petersburg Mathematical Journal 16 (2005), no. 2, pp. 413–416. Zbl 1078.35081.

[[]Gar18] L. GARRIGUE, Unique continuation for many body Schrödinger operators and the Hohenberg-Kohn theorem, Mathematical Physics, Analysis and Geometry 21 (2018), no.3, Paper No. 27, 11. MR 3854406. Zbl 1400.81217.

[[]Ges84] F. GESZTESY, Non-degenerate ground state for Schrödinger operators, *Reports on Mathematical Physics* 20 (1984), no.1, pp. 93–109. Zbl 0567.35075.

[GJ70]	J. GLIMM and A. JAFFE, The $\lambda(\Pi^4)_2$ quantum field theory without cutoffs. II. The field operators and the approximate vacuum, <i>Annals of Mathematics</i> 91 (1970), pp. 362–401. MR 256677. Zbl 0191.27005.
[GM09]	F. GESZTESY and M. MITREA, Nonlocal Robin Laplacians and some remarks on a paper by Filonov on eigenvalue inequalities, <i>Journal of Differential Equations</i> 247 (2009), no. 10, pp. 2871–2896. MR 2568160. Zbl 1181.35155.
[Gul19]	N. J. GULIYEV, Schrödinger operators with distributional potentials and boundary conditions dependent on the eigenvalue parameter, <i>Journal of Mathematical Physics</i> 60 (2019), 063501. MR 3945700. Zbl 1415,34083.
[Her89]	J. HERCZYŃSKI, JAN, On Schrödinger operators with distributional potentials, <i>Journal of Operator Theory</i> 21 (1989), no.2, pp. 273-295. MR 1023316.
[Horo7]	L. HÖRMANDER, The analysis of linear partial differential operators III, Springer (2007). MR 2304165. Zbl 1115.35005.
[JK85]	D. JERISON and C. E. KENIG, Unique continuation and absence of positive eigenvalues for Schrödinger operators, <i>Annals of Mathematics</i> 121 (1985), no.3, pp. 463–494. MR 794370. Zbl 0593.35119.
[KT01]	H. KOCH and D. TARTARU, Carleman estimates and unique continuation for second-order elliptic equations with nonsmooth coefficients, <i>Communications on Pure and Applied Mathematics</i> 54 (2001), no. 3, pp. 339–360. MR 1809741. Zbl 1033.35025.
[Kur97]	K. KURATA, A unique continuation theorem for the Schrödinger equation with singular magnetic field, <i>Proceedings of the American Mathematical Society</i> 125 (1997), no.3, pp. 853–860. MR 1363173. Zbl o887.35026
[Le017]	G. LEONI, A first course in Sobolev spaces, Graduate Studies in Mathematics Volume 181, American Mathematical Society (2017). MR 3726909. Zbl 1382.46001.
[McLoo]	W. MCLEAN, Strongly elliptic systems and boundary integral equations, Cambridge University Press (2000). MR 1742312. Zbl 0948.35001.
[MK14]	K. A. MIRZOEV, N. N. KONECHNAYA, Singular Sturm-Liouville Operators with Distribution Potentials, <i>Journal of Mathematical Sciences</i> 200 (2014), pp. 96–105. MR 3272038. Zbl 1309.34061.
[MM79]	M. MOSHE and V. J. MIZEL, Every superposition operator mapping one Sobolev space into another is continuous, <i>Journal of Functional Analysis</i> 33 (1979), no.2, pp. 217–229. MR 546508, Zbl 0418.46024
[LR15]	V. LOTOREICHIK and J. ROHLEDER, An eigenvalue inequality for Schrödinger operators with δ - and δ' -interactions supported on hypersurfaces, <i>In Operator Algebras and Mathematical Physics</i> , edited by Tirthankar Bhattacharyya and Michael A. Dritschel, 173–84. Cham: Springer International Publishing. MR 3408721. Zbl 1346.35135.
[RS72]	J. F. READING and J. L. SIGEL, Exact solution of the one-dimensional Schrödinger equation with δ -function potentials of arbitrary position and strength <i>Physical Review B</i> 5 (1972), no.2, pp. 556–65. MR 391777.
[Roh21]	J. ROHLEDER, Inequalities between Neumann and Dirichlet eigenvalues of Schrödinger operators, <i>Journal of Spectral Theory</i> 11 (2021), no. 3, pp. 915-933. MR 4322026. Zbl 1484.35303.
[RS ₇₅]	M. REED and B. SIMON, Methods of modern mathematical physics. II. Fourier analysis, self-adjointness, Academic Press (1975). MR 493420.
[RS ₇ 8]	M. REED and B. SIMON, Methods of modern mathematical physics. IV. Analysis of operators, Academic Press (1978). MR 493421.
[RS80]	M. REED and B. SIMON, Methods of modern mathematical physics. I. Functional Analysis, Academic Press (1980). MR 751959
[SS99]	A. M. SAVCHUK, A. A. SHKALIKOV, Sturm-Liouville operators with singular potentials, <i>Mathematical Notes</i> 66 (1999), pp.
[Sog90]	741–753. MR 1746481. Zbl 0962.34067. C. D. SOGGE, Strong uniqueness theorems for second order elliptic differential equations, <i>American Journal of Mathematics</i>
	112 (1990), no. 6, pp. 943–984. MR 1081811. Zbl 0734.35012.
[SPR+24]	S. M. SUTTER, M. PENZ, M. RUGGENTHALER, R. VAN LEUWEEN, and K. J. H. GIESBERTZ, Solution of the <i>v</i> -representability problem on a one-dimensional torus, <i>Journal of Physics A: Mathematical and Theoretical</i> 57 (2024), no 47. MR 4840211.
[Tes14]	G. TESCHL, Mathematical methods in quantum mechanics: with applications to Schrödinger operators, American Mathematical Society, Graduate Studies in Mathematics 157 (2014) MR 2242089. Thi 1242 81002
[Wol93]	T. H. WOLFF, Recent work on sharp estimates in second-order elliptic unique continuation problems, <i>The Journal of Geometric Analysis</i> 3 (1993) no. 6, pp. 621–650, MR 1248088 Zbl 9787 35017
[Zet21]	A. ZETTL, <i>Recent developments in Sturm-Liouville theory</i> De Gruyter Studies in Mathematics, 76 (2021). MR 4510393. Zbl 1479.34004.

(T. Carvalho Corso) Institute of Applied Analysis and Numerical Simulation, University of Stuttgart, Pfaffenwaldring 57, 70569 Stuttgart, Germany

Email address: thiago.carvalho-corso@mathematik.uni-stuttgart.de