

# A LINEAR CONVERGENCE RESULT FOR THE JACOBI-PROXIMAL ALTERNATING DIRECTION METHOD OF MULTIPLIERS

HYELIN CHOI AND WOOCHEOL CHOI

## Abstract

In this paper, we analyze the convergence rate of the Jacobi-Proximal Alternating Direction Method of Multipliers (ADMM), a method initially introduced by Deng et al.[6] for the block-structured optimization problem with linear constraint. We establish the linear convergence of the algorithm when the cost functions are strongly convex and smooth.

**Keywords:** Alternating Direction Method of Multipliers, Jacobi-Proximal ADMM, Linear Convergence, Convex Optimization

## 1. INTRODUCTION

We consider the following convex optimization problem with  $N$ -block variables involving a linear constraint given as follows:

$$\begin{aligned} \min_{x_1, \dots, x_N} \quad & \sum_{i=1}^N \tilde{f}_i(x_i) \\ \text{subject to} \quad & \sum_{i=1}^N A_i x_i = c, \\ & x_i \in \mathcal{X}_i, \quad i = 1, \dots, N. \end{aligned} \tag{1.1}$$

where  $N \geq 2$ ,  $x_i \in \mathbb{R}^{n_i}$ ,  $A_i \in \mathbb{R}^{m \times n_i}$ ,  $c \in \mathbb{R}^m$  and  $\mathcal{X}_i \subset \mathbb{R}^{n_i}$  are closed convex sets and  $\tilde{f}_i : \mathbb{R}^{n_i} \rightarrow (-\infty, +\infty]$  ( $i = 1, 2, \dots, N$ ) are closed proper convex functions. The solution set of (1.1) is assumed to be nonempty. Note that the convex constraints  $x_i \in \mathcal{X}_i$  can be incorporated into the objective  $\tilde{f}_i$  using the indicator function:

$$I_{\mathcal{X}_i}(x_i) = \begin{cases} 0 & \text{if } x_i \in \mathcal{X}_i, \\ +\infty & \text{otherwise.} \end{cases} \tag{1.2}$$

Then the problem (1.1) can be rewritten as

$$\begin{aligned} \min_{x_1, \dots, x_N} \quad & \sum_{i=1}^N f_i(x_i) \\ \text{subject to} \quad & \sum_{i=1}^N A_i x_i = c, \end{aligned} \tag{1.3}$$

where  $f_i(x_i) = \tilde{f}_i(x_i) + I_{\mathcal{X}_i}(x_i)$ . Thus, we consider the equivalent reformulation (1.1) throughout this paper for ease of presentation.

The problem (1.1) appears in various applications that contain the image alignment problem in [33], the robust principal component analysis model with noisy and incomplete data in [36], the latent variable Gaussian graphical model selection in [3, 31] and the quadratic discriminant analysis model in [30].

To solve the problem (1.1), various algorithms have been proposed. A simple distributed algorithm for solving (1.1) is dual decomposition [11], which is essentially a dual ascent method or dual subgradient method [35] as follows. Consider the Lagrangian for problem (1.1):

$$\mathcal{L}(x_1, \dots, x_N, \lambda) = \sum_{i=1}^N f_i(x_i) - \left\langle \lambda, \sum_{i=1}^N A_i x_i - c \right\rangle \quad (1.4)$$

where  $\lambda \in \mathbb{R}^m$  is the Lagrangian multiplier or the dual variable. The dual decomposition updates as follows: for  $k \geq 1$ ,

$$\begin{cases} (x_1^{k+1}, x_2^{k+1}, \dots, x_N^{k+1}) = \arg \min_{\{x_i\}} \mathcal{L}(x_1, \dots, x_N, \lambda^k), \\ \lambda^{k+1} = \lambda^k - \alpha_k \left( \sum_{i=1}^N A_i x_i^{k+1} - c \right), \end{cases}$$

where  $\alpha_k > 0$  is a step-size. Since all the  $x_i$ 's are separable in the Lagrangian function (1.4), the  $x$ -update step reduces to solving  $N$  individual  $x_i$ - subproblems:

$$x_i^{k+1} = \arg \min_{x_i} f_i(x_i) - \langle \lambda^k, A_i x_i \rangle \quad \text{for } i = 1, 2, \dots, N,$$

and thus they can be carried out in parallel. With a suitable choice of  $\alpha_k$  and certain assumptions, dual decomposition is guaranteed to converge to an optimal solution [35]. However, dual decomposition exhibits slow convergence in practice, with a convergence rate of  $\mathcal{O}(1/\sqrt{k})$  for general convex problems.

An effective distributed algorithm for (1.1) can be designed based on the alternating direction method of multipliers (ADMM) [13, 15] using the following augmented Lagrangian:

$$\mathcal{L}_\rho(x_1, \dots, x_N; \lambda) := \sum_{i=1}^N f_i(x_i) - \left\langle \lambda, \sum_{i=1}^N A_i x_i - c \right\rangle + \frac{\rho}{2} \left\| \sum_{i=1}^N A_i x_i - c \right\|^2,$$

where  $\rho > 0$  is a penalty parameter. In the context of ADMM using the augmented Lagrangian, the primal variable  $x_i$  can be updated using different approaches. One common method is the Gauss-Seidel update, in which the primal variable  $x_i$  is updated sequentially as follows:

$$\begin{aligned} x_i^{k+1} &= \arg \min_{x_i} \mathcal{L}_\rho(x_1^{k+1}, \dots, x_{i-1}^{k+1}, x_i, x_{i+1}^k, \dots, x_N^k, \lambda^k) \\ &= \arg \min_{x_i} f_i(x_i) + \frac{\rho}{2} \left\| \sum_{j<i} A_j x_j^{k+1} + A_i x_i + \sum_{j>i} A_j x_j^k - c - \frac{\lambda^k}{\rho} \right\|^2. \end{aligned}$$

Alternatively, the Jacobi-type update provides a parallel update scheme for the primal variable  $x_i$  as follows:

$$\begin{aligned} x_i^{k+1} &= \arg \min_{x_i} \mathcal{L}_\rho(x_1^k, \dots, x_{i-1}^k, x_i, x_{i+1}^k, \dots, x_N^k, \lambda^k) \\ &= \arg \min_{x_i} f_i(x_i) + \frac{\rho}{2} \left\| A_i x_i + \sum_{j \neq i} A_j x_j^k - c - \frac{\lambda^k}{\rho} \right\|_2^2. \end{aligned}$$

Since Gauss-Seidel ADMM updates the primal variables  $x_i$  using the most recent information, the performance of Jacobi ADMM is typically expected to be worse than that of Gauss-Seidel ADMM. For instance, the work [19] found an example for which Jacobi ADMM diverges even when  $N = 2$ . This issue was addressed by Deng et al.[6] that introduced the Jacobi-Proximal ADMM involving a proximal term  $\frac{1}{2} \|x_i - x_i^k\|_{P_i}^2$  for each  $x_i$ -subproblem with positive semi-definite matrix  $P_i$  and a damping parameter  $\gamma > 0$  for the update of  $\lambda$  to guarantee its convergence. The Jacobi-Proximal ADMM is formulated as in Algorithm 1. Here we denote  $\|x_i\|_{P_i}^2 := x_i^T P_i x_i$ . The Jacobi-Proximal

---

**Algorithm 1** Jacobi-Proximal ADMM

---

**Initialize:**  $x_i^0$  ( $i = 1, 2, \dots, N$ ) and  $\lambda^0$ ;

**for**  $k = 0, 1, \dots$  **do**

Update  $x_i$  for  $i = 1, \dots, N$  in parallel by:

$$x_i^{k+1} = \arg \min_{x_i} \left\{ f_i(x_i) + \frac{\rho}{2} \left\| A_i x_i + \sum_{j \neq i} A_j x_j^k - c - \frac{\lambda^k}{\rho} \right\|_2^2 + \frac{1}{2} \|x_i - x_i^k\|_{P_i}^2 \right\}; \quad (1.5)$$

$$\text{Update } \lambda^{k+1} = \lambda^k - \gamma \rho \left( \sum_{i=1}^N A_i x_i^{k+1} - c \right). \quad (1.6)$$

**end for**

---

ADMM has the advantage of being highly parallelizable, as the updates for  $x_i$  for each  $i = 1, 2, \dots, N$  can be computed simultaneously for each  $i$ , offering potential computational efficiency gains. We note that the original  $x_i$ -subproblem (1.5) contains a quadratic term of the form  $\frac{\rho}{2} x_i^T A_i^T A_i x_i$ . If  $A_i^T A_i$  is ill-conditioned or computationally expensive to invert, one approach is to modify  $P_i$  as  $P_i = D_i - \rho A_i^T A_i$ , where  $D_i$  is a well-conditioned and simple matrix (e.g., a diagonal matrix). This modification eliminates the quadratic term  $\frac{\rho}{2} x_i^T A_i^T A_i x_i$ , making the  $x_i$ -subproblem easier to solve. There are various choices for  $P_i$  that can simplify the computation of the  $x_i$ -subproblem. Here are two widely used formulations for  $P_i$  in [7], corresponding to the following methods:

- (1) The *standard proximal method*, where  $P_i$  is taken as  $P_i = \tau_i \mathbf{I}$ , with  $\tau_i > 0$ .
- (2) The *prox-linear method*, where  $P_i$  is taken as  $P_i = \tau_i \mathbf{I} - \rho A_i^T A_i$ , with  $\tau_i > 0$ .

The work of Deng et al.[6] established a sublinear convergence result for the convergence of Jacobi-Proximal ADMM when the proximal matrices  $P_i$  satisfy  $P_i \succ (\frac{N}{2-\gamma} - 1) A_i^T A_i$  for  $i = 1, 2, \dots, N$  with  $\gamma < 2$ . In particular, under specific choices of  $P_i$ , the convergence

conditions reduce to  $\tau_i > \rho(\frac{N}{2-\gamma} - 1)\|A_i\|^2$  for the *standard proximal method* and  $\tau_i > \frac{\rho N}{2-\gamma}\|A_i\|^2$  for the *prox-linear method*.

We then extend the analysis to establish the linear convergence of Jacobi-Proximal ADMM. It is well-known that ADMM converges linearly when the cost functions exhibit additional structure, such as strong convexity. We prove that the Jacobi-Proximal ADMM converges linearly under the assumption that each cost function  $f_j$  is strongly convex. We derive a condition on the parameters  $\rho > 0$  and  $\gamma > 0$  for linear convergence, and we establish a linear convergence rate dependent on these parameters. We refer to Section 2 for the details and we provide a comparison of our convergence result with previous work on the problem (1.1) in Table 1.

**1.1. Related works.** The Alternating Direction Method of Multipliers (ADMM) [13, 15] is an effective approach that simplifies the original problem by decomposing it into multiple easier-to-solve minimization subproblems. Building on the classical Douglas-Rachford operator splitting method [8], Glowinski and Marrocco [16], along with Gabay and Mercier [14], introduced the well-known alternating direction method of multipliers (ADMM) with Gauss-Seidel decomposition for solving (1.1) when  $N = 2$ . It utilizes the augmented Lagrangian for (1.1):

$$\mathcal{L}_\rho(x_1, \dots, x_N; \lambda) := \sum_{i=1}^N f_i(x_i) - \left\langle \lambda, \sum_{i=1}^N A_i x_i - c \right\rangle + \frac{\rho}{2} \left\| \sum_{i=1}^N A_i x_i - c \right\|^2$$

which incorporates a quadratic penalty of the constraints (with a parameter  $\rho > 0$ ) into the Lagrangian. At each iteration, the augmented Lagrangian is sequentially minimized with respect to  $x_1$  and  $x_2$ , followed by an update of the dual variable  $\lambda$ . The ADMM has been extensively investigated for solving two-block convex minimization problems (i.e.,  $N = 2$ ). Its global convergence in the case of  $N = 2$  has been established in [9, 12]. In a more recent work by Deng and Yin [7], a generalized version of ADMM was introduced, where proximal terms were added to the two subproblems in (2.2), demonstrating linear convergence under certain assumptions on the strong convexity of functions  $f_1$  and  $f_2$ , as well as the rank conditions on  $A_1$  and  $A_2$ . Additionally, various studies have explored the convergence rate properties of ADMM when  $N = 2$  (see, e.g., [28, 10, 18, 1, 29, 20, 32, 21]). Although the convergence properties of ADMM for  $N = 2$  are well understood, its behavior for  $N \geq 3$  has remained uncertain for a considerable period. In the work by Chen et al. [4], a counterexample was provided, demonstrating that without additional conditions, the ADMM for  $N \geq 3$  may fail to converge. Building on this, we first discuss studies that establishes the convergence conditions for ADMM using the Gauss-Seidel scheme when  $N \geq 3$ , where the Gauss-Seidel method updates variables sequentially. We refer to the works [2], [23], [5], and [27] that achieved global and sublinear convergence results of the 3-block ADMM. Han and Yuan [17] established the global convergence of ADMM (2.2) under the assumption that  $f_1, \dots, f_N$  are all strongly convex and  $\gamma$  is restricted to a specific range. Both Chen, Shen, and You [5] and Lin, Ma, and Zhang [25] relax the conditions compared to [17] by allowing only  $N - 1$  functions to be strongly convex, with

	Ref.	Convexity	Regularity	Rank Conditions	Rate
Gauss-Seidel ADMM for $N = 2$	[7]	$f_1$ : SC	$f_1$ : $L$ -smooth	$A_1$ : full column rank $A_2$ : full row rank	linear
		$f_1, f_2$ : SC	$f_1$ : $L$ -smooth	$A_2$ : full row rank	
		$f_1$ : SC	$f_1, f_2$ : $L$ -smooth	$A_1$ : full column rank	
		$f_1, f_2$ : SC	$f_1, f_2$ : $L$ -smooth	—	
Gauss-Seidel ADMM for $N \geq 3$	[26]	$f_2, \dots, f_N$ : SC	$f_N$ : $L$ -smooth	$A_N$ : full row rank	linear
		$f_1, \dots, f_N$ : SC	$f_1, \dots, f_N$ : $L$ -smooth	—	
		$f_2, \dots, f_N$ : SC	$f_1, \dots, f_N$ : $L$ -smooth	$A_1$ : full column rank	
Gauss-Seidel ADMM for $N \geq 3$	[22]	$f_i(x_i) = g_i(E_i x_i) + h_i(x_i)$ $g_i$ : strictly C $h_i$ : C	$g_i$ : $L$ -smooth	$E_i$ : full column rank	linear
Jacobi-Proximal ADMM for $N \geq 3$	[6]	$f_1, \dots, f_N$ : C	—	$A_i$ : full column rank	sub-linear
Jacobi-Proximal ADMM for $N \geq 3$	This work	$f_1, \dots, f_N$ : SC	$f_1, \dots, f_N$ : $L$ -smooth	$[A_1^T; A_2^T; \dots; A_N^T]$ : full row rank	linear

TABLE 1. This table summarizes the convergence results for ADMM. In the ‘Convexity’ column, C (resp., SC) means that the total cost function is assumed to be convex (resp., strongly convex). In the ‘Rate’ column, ‘linear’ refers to the cases with linear convergence, while ‘sublinear’ indicates the case with sublinear convergence.  $N$  denotes the number of blocks of objective functions and  $k$  denotes the number of iterations.

with Chen et al. [5] considering the case  $N = 3$  and Lin et al. [25] considering  $N \geq 3$ . Furthermore, Lin, Ma, and Zhang [25] demonstrated that under these conditions, ADMM (2.2) exhibits a sublinear convergence rate in both the ergodic and non-ergodic senses. Lin, Ma, and Zhang [24] further presented sufficient conditions to guarantee a sublinear convergence rate for ADMM when  $N \geq 3$ . Lin, Ma, and Zhang [26] proved the global linear convergence of ADMM under the three scenarios. The global R-linear convergence of ADMM has been established by Hong and Luo [22]. Alternatively, Jacobi updates in ADMM have been extensively studied, offering parallelization advantages for multi-block optimization problems. The linear convergence of Gauss-Seidel ADMM, Proximal ADMM, and modified Jacobi ADMM, as shown in [22], is established under several assumptions. In [6], it was established that the multi-block Jacobi ADMM converges globally at a rate of  $o(1/k)$  in two cases: (i) when all coefficient matrices are nearly mutually orthogonal and possess full column rank, or (ii) when proximal terms are introduced into the subproblems. In [19], the multi-block Jacobi ADMM was combined with a relaxation step, achieving a worst-case convergence rate of  $O(1/k)$  in both ergodic and non-ergodic senses. In a recent work by Xiao, Deng, and Xu [38], the authors introduce the Distributed Jacobi-Proximal ADMM algorithm for solving consensus optimization problems.

**1.2. Contribution.** The main contribution in this paper are as follows. We establish the global linear convergence of Jacobi-Proximal ADMM(Algorithm 1) under the assumption that the objective function  $f_i$  are strictly convex and  $L_i$ -smooth. We establish its convergence at a rate of  $O(\sigma^k)$  for the optimality measure defined in (2.4). Our numerical results on the LCQP model and the optimal resource allocation problem support the linear convergence result of this work.

**1.3. Organization.** We organize the rest of this paper as follows. In Section 2, we present a sufficient condition to ensure the convergence of (2.2). Then based on the analysis in Section 2, we prove the global convergence of the Algorithm 1 in Section 3. In Section 4, we present convergence results by conducting numerical experiments on two examples. Finally, we conclude the paper in Section 5.

## 2. STATEMENT OF THE MAIN RESULT

In this section, we state the main result of this paper concerning the linear convergence of the Jacobi-Proximal ADMM in Algorithm 1. Since both of the objective function and constraints of (1.3) are sums of terms on individual  $x_i$ 's, the problem can be decomposed into  $N$  smaller subproblems, which is solved in a parallel and distributed manner.

Related to the problem (1.3), we first define the following augmented Lagrangian

$$\mathcal{L}_\rho(x_1, \dots, x_N; \lambda) := \sum_{i=1}^N f_i(x_i) - \left\langle \lambda, \sum_{i=1}^N A_i x_i - b \right\rangle + \frac{\rho}{2} \left\| \sum_{i=1}^N A_i x_i - b \right\|^2$$

where  $\lambda \in \mathbb{R}^m$  is the dual variable and parameter  $\rho > 0$ .

We state assumptions on the cost functions which will be used throughout the paper. Here we let  $n = n_1 + \dots + n_N$  and set  $A \in \mathbb{R}^{m \times n}$  by  $A = [A_1 \dots A_N]$ . Also, we set  $x^k \in \mathbb{R}^n$  for  $k \geq 0$  and  $x^* \in \mathbb{R}^n$  as  $x^k = [x_1^k; \dots; x_N^k]$  and  $x^* = [x_1^*; \dots; x_N^*]$ . We then have

$$Ax^k = \sum_{i=1}^N A_i x_i^k.$$

**Assumption 2.1.** *The functions  $f_i : \mathbb{R}^{n_i} \rightarrow (-\infty, +\infty]$  are closed proper convex for  $1 \leq i \leq N$ .*

**Assumption 2.2.** *We assume that  $x^* = (x_1^*, \dots, x_N^*) \in \mathbb{R}^n$  is a minimizer of problem (1.3) accompanied with a multiplier  $\lambda^* \in \mathbb{R}^m$  satisfying the following KKT condition*

$$\nabla f_i(x_i^*) = A_i^T \lambda^* \quad \text{for } i = 1, \dots, N, \quad (2.1)$$

$$Ax^* = \sum_{i=1}^N A_i x_i^* = c. \quad (2.2)$$

**Assumption 2.3.** *There are positive constants  $L_i$  such that for all  $x, y \in \mathbb{R}^{n_i}$ ,*

$$\|\nabla f_i(x) - \nabla f_i(y)\| \leq L_i \|x - y\| \quad \text{for } 1 \leq i \leq N.$$

**Assumption 2.4.** *The function  $f_i : \mathbb{R}^{n_i} \rightarrow (-\infty, +\infty]$  is  $\alpha$ -strongly convex for some  $\alpha > 0$ , i.e.,*

$$f_i(y) \geq f_i(x) + \langle y - x, \nabla f_i(x) \rangle + \alpha \|x - y\|^2$$

for all  $x, y \in \mathbb{R}^{n_i}$ .

**Assumption 2.5.** *We assume that the matrix  $[A_1^T; A_2^T; \dots; A_N^T] \in \mathbb{R}^{n \times m}$  with  $n = n_1 + \dots + n_N$  has full column rank.*

Here  $P_i$  is positive semi-definite for each  $1 \leq i \leq N$ . For  $k \geq 0$ , we denote by  $(x_1^k, \dots, x_N^k; \lambda^k)$  the variable generated by the Jacobi-Proximal ADMM. We state the main result of this paper.

**Theorem 2.6.** *Suppose Assumptions 2.1-2.5 hold. Choose any  $s > 0$  such that*

$$s \left[ \rho^2 D \|A_i\| + \frac{L}{N} \right] < \frac{\alpha}{2N}, \quad \forall 1 \leq i \leq N,$$

and suppose that  $0 < \gamma < 2$ ,  $\rho > 0$  are chosen so that there exist values  $\xi_i > 0$  such that

$$\begin{cases} \rho A_i^T A_i + P_i - 8s(\rho A_i^T A_i + P_i)^T (\rho A_i^T A_i + P_i) - \frac{\rho}{\xi_i} A_i^T A_i \succ 0, \\ \sum_{i=1}^N \xi_i < 2 - \gamma. \end{cases} \quad (2.3)$$

Let  $\sigma = \max \{1 - 2\gamma\rho s c_A^2, \mu_s\}$ . Here  $c_A > 0$  and  $\mu_s \in (0, 1)$  are constants satisfying

$$\begin{cases} c_A^2 I_m \preceq \sum_{i=1}^N A_i A_i^T, \\ (\rho + 4Ns\rho^2 D) A_i^T A_i + P_i \preceq \mu_s (\rho A_i^T A_i + P_i) + 2(\alpha - Ls) I_{n_i}, \end{cases}$$

where the first inequality ensures that  $c_A = \sqrt{\min \text{eig}\left(\sum_{i=1}^N A_i A_i^T\right)} > 0$ . Then we have

$$\phi(x^{k+1}, \lambda^{k+1}) \leq \sigma \phi(x^k, \lambda^k), \quad (2.4)$$

where  $\phi : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}$  is defined as  $\phi(x^k, \lambda^k) = \frac{1}{2\gamma\rho} \|\lambda^k - \lambda^*\|^2 + \frac{1}{2} \sum_{i=1}^N \|x_i^k - x_i^*\|_{\rho A_i^T A_i + P_i + 2(\alpha - 2Ls)I_{n_i}}^2$ . As a result of (2.4), we have the following result

$$\phi(x^k, \lambda^k) = O(\sigma^k).$$

*Remark 2.7.* Choosing  $\xi_i < \frac{2-\gamma}{N}$  leads to a simplification of the convergence condition (2.3), resulting in the following form:

$$\rho A_i^T A_i + P_i - 8s(\rho A_i^T A_i + P_i)^T (\rho A_i^T A_i + P_i) - \frac{\rho N}{2-\gamma} A_i^T A_i \succ 0. \quad (2.5)$$

In the case of  $P_i = \tau_i \mathbf{I}$  (*standard proximal method*), the condition (2.5) further reduces to

$$\rho \|A_i\|^2 + \tau_i - 8s(\rho \|A_i\|^2 + \tau_i)^2 - \frac{\rho N}{2-\gamma} \|A_i\|^2 > 0.$$

Similarly, for  $P_i = \tau_i \mathbf{I} - \rho A_i^T A_i$  (*prox-linear method*), the condition (2.5) simplifies to

$$\tau_i - 8s\tau_i^2 - \frac{\rho N}{2-\gamma} \|A_i\|^2 > 0.$$

### 3. CONVERGENCE

In this section, we prove Theorem 2.6. For this, we prepare the several lemmas.

**Lemma 3.1.** *Suppose Assumption 2.5 holds. Consider matrices  $A_i \in \mathbb{R}^{m \times n_i}$  for  $1 \leq i \leq N$ . Then there exists  $c_A > 0$  such that*

$$\left( \sum_{i=1}^N \|A_i^T \lambda\|^2 \right)^{\frac{1}{2}} \geq c_A \|\lambda\| \quad (3.1)$$

for all  $\lambda \in \mathbb{R}^m$ .

*Proof.* We define  $c_A \geq 0$  by

$$c_A = \min_{\lambda \in \mathbb{R}^m \setminus \{0\}} \frac{\left( \sum_{i=1}^N \|A_i^T \lambda\|^2 \right)^{\frac{1}{2}}}{\|\lambda\|} = \min_{\lambda \in S^{m-1}} \frac{\left( \sum_{i=1}^N \|A_i^T \lambda\|^2 \right)^{\frac{1}{2}}}{\|\lambda\|}.$$

In order to prove that  $c_A > 0$ , we assume  $c_A = 0$  and proceed to derive a contradiction. Since  $S^{m-1}$  is compact, there exists a minimizer  $\lambda_0 \in S^{m-1}$  of (3.1) such that

$$\frac{\left( \sum_{i=1}^N \|A_i^T \lambda_0\|^2 \right)^{\frac{1}{2}}}{\|\lambda_0\|} = 0.$$

Thus we have  $A_i^T \lambda_0 = 0$  for all  $1 \leq i \leq N$ , and this leads to  $\lambda_0 = 0$  since  $[A_1^T; \dots; A_N^T]$  has full column rank. It contradicts to the fact that  $\lambda_0 \in S^{m-1}$ . Therefore, we have  $c_A > 0$ , and the lemma is proved.  $\square$



*Remark 3.2.* The assumption of Lemma 3.1 means that the matrix  $[A_1^T; A_2^T; \dots; A_N^T] \in \mathbb{R}^{n \times m}$  with  $n = n_1 + \dots + n_N$  has only a trivial kernel, i.e.,  $\ker([A_1^T; A_2^T; \dots; A_N^T]) = \{\mathbf{0}\}$ .

By the strongly convexity of the objective function, we have the following lemma.

**Lemma 3.3.** *Suppose Assumptions 2.1, 2.2, and 2.4 hold. Then it holds that*

$$\begin{aligned} & \frac{1}{2\gamma\rho} \|\lambda^k - \lambda^*\|^2 + \frac{1}{2} \sum_{i=1}^N \|x_i^k - x_i^*\|_{\rho A_i^T A_i + P_i}^2 \\ & \geq \alpha \sum_{i=1}^N \|x_i^{k+1} - x_i^*\|^2 + \frac{1}{2\gamma\rho} \|\lambda^{k+1} - \lambda^*\|^2 + \frac{1}{2} \sum_{i=1}^N \|x_i^{k+1} - x_i^*\|_{\rho A_i^T A_i + P_i}^2 + H_k \end{aligned} \quad (3.2)$$

where  $H_k := \frac{1}{2} \sum_{i=1}^N \|x_i^{k+1} - x_i^k\|_{\rho A_i^T A_i + P_i}^2 + \frac{2-\gamma}{2\gamma^2\rho} \|\lambda^{k+1} - \lambda^k\|^2 + \frac{1}{\gamma} \langle \lambda^k - \lambda^{k+1}, A(x^k - x^{k+1}) \rangle$ .

*Proof.* By the optimality of (1.5), we have

$$\nabla f_i(x_i^{k+1}) = A_i^T \lambda^k - \rho A_i^T \left( A_i x_i^{k+1} + \sum_{j \neq i} A_j x_j^k - c \right) + P_i(x_i^k - x_i^{k+1}). \quad (3.3)$$

Combining (3.3) and (2.1), we get

$$\begin{aligned} & A_i^T(\lambda^k - \lambda^*) \\ & = \nabla f_i(x_i^{k+1}) - \nabla f_i(x_i^*) + \rho A_i^T \left( A_i x_i^{k+1} + \sum_{j \neq i} A_j x_j^k - c \right) + P_i(x_i^{k+1} - x_i^k). \end{aligned} \quad (3.4)$$

Since each  $f_i$  is  $\alpha$ -strongly convex, we have

$$\langle x_i^{k+1} - x_i^*, \nabla f_i(x_i^{k+1}) - \nabla f_i(x_i^*) \rangle \geq \alpha \|x_i^{k+1} - x_i^*\|^2. \quad (3.5)$$

Inserting (3.4) into (3.5), we have

$$\begin{aligned} & \left\langle A_i(x_i^{k+1} - x_i^*), \lambda^k - \lambda^* - \rho \left( A_i x_i^{k+1} + \sum_{j \neq i} A_j x_j^k - c \right) \right\rangle \\ & \quad + \left\langle x_i^{k+1} - x_i^*, P_i(x_i^k - x_i^{k+1}) \right\rangle \geq \alpha \|x_i^{k+1} - x_i^*\|^2. \end{aligned}$$

We rewrite above inequality as follows

$$\begin{aligned} & \left\langle A_i(x_i^{k+1} - x_i^*), \lambda^k - \lambda^* - \rho \left( \sum_{i=1}^N A_i x_i^{k+1} + \sum_{j \neq i} A_j (x_j^k - x_j^{k+1}) - c \right) \right\rangle \\ & \quad + \left\langle x_i^{k+1} - x_i^*, P_i(x_i^k - x_i^{k+1}) \right\rangle \geq \alpha \|x_i^{k+1} - x_i^*\|^2. \end{aligned}$$

For simplicity, we let  $\hat{\lambda} = \lambda^k - \rho \left( \sum_{i=1}^N A_i x_i^{k+1} - c \right) = \lambda^k - \frac{1}{\gamma} (\lambda^k - \lambda^{k+1})$ . Then the above inequality reads as follows

$$\begin{aligned} & \left\langle A_i(x_i^{k+1} - x_i^*), \hat{\lambda} - \lambda^* - \rho \left( \sum_{j \neq i} A_j (x_j^k - x_j^{k+1}) \right) \right\rangle \\ & \quad + \left\langle x_i^{k+1} - x_i^*, P_i(x_i^k - x_i^{k+1}) \right\rangle \geq \alpha \|x_i^{k+1} - x_i^*\|^2. \end{aligned}$$

Using that  $\sum_{j \neq i} A_j(x_j^k - x_j^{k+1}) = A(x^k - x^{k+1}) - A_i(x_i^k - x_i^{k+1})$  and summing the above estimate for  $1 \leq i \leq N$  yields

$$\begin{aligned} & \left\langle A(x^{k+1} - x^*), \hat{\lambda} - \lambda^* \right\rangle + \sum_{i=1}^N (x_i^{k+1} - x_i^*)^T (\rho A_i^T A_i + P_i) (x_i^k - x_i^{k+1}) \\ & \geq \alpha \sum_{i=1}^N \|x_i^{k+1} - x_i^*\|^2 + \rho \left\langle A(x^{k+1} - x^*), A(x^k - x^{k+1}) \right\rangle. \end{aligned} \quad (3.6)$$

Combining the update rule (1.6) and (2.2) and using that  $\hat{\lambda} - \lambda^* = (\lambda^{k+1} - \lambda^*) + \frac{\gamma-1}{\gamma} (\lambda^k - \lambda^{k+1})$  in the above inequality, we get

$$\begin{aligned} & \left\langle \frac{1}{\gamma\rho} (\lambda^k - \lambda^{k+1}), \lambda^{k+1} - \lambda^* + \frac{\gamma-1}{\gamma} (\lambda^k - \lambda^{k+1}) \right\rangle \\ & + \sum_{i=1}^N (x_i^{k+1} - x_i^*)^T (\rho A_i^T A_i + P_i) (x_i^k - x_i^{k+1}) \\ & \geq \alpha \sum_{i=1}^N \|x_i^{k+1} - x_i^*\|^2 + \frac{1}{\gamma} \left\langle \lambda^k - \lambda^{k+1}, A(x^k - x^{k+1}) \right\rangle. \end{aligned} \quad (3.7)$$

For the two terms in the left hand side, we have the following identities

$$\begin{aligned} & \left\langle \frac{1}{\gamma\rho} (\lambda^k - \lambda^{k+1}), \lambda^{k+1} - \lambda^* + \frac{\gamma-1}{\gamma} (\lambda^k - \lambda^{k+1}) \right\rangle \\ & = \frac{\gamma-2}{2\gamma^2\rho} \|\lambda^{k+1} - \lambda^k\|^2 - \frac{1}{2\gamma\rho} \|\lambda^{k+1} - \lambda^*\|^2 + \frac{1}{2\gamma\rho} \|\lambda^k - \lambda^*\|^2, \end{aligned}$$

and

$$\begin{aligned} & \sum_{i=1}^N (x_i^{k+1} - x_i^*)^T (\rho A_i^T A_i + P_i) (x_i^k - x_i^{k+1}) \\ & = -\frac{1}{2} \sum_{i=1}^N \left( \|x_i^{k+1} - x_i^*\|_{\rho A_i^T A_i + P_i}^2 + \|x_i^{k+1} - x_i^k\|_{\rho A_i^T A_i + P_i}^2 - \|x_i^k - x_i^*\|_{\rho A_i^T A_i + P_i}^2 \right). \end{aligned}$$

Inserting the above two equalities in (3.7), we obtain

$$\begin{aligned} & \frac{1}{2\gamma\rho} \|\lambda^k - \lambda^*\|^2 + \frac{1}{2} \sum_{i=1}^N \|x_i^k - x_i^*\|_{\rho A_i^T A_i + P_i}^2 \\ & \geq \alpha \sum_{i=1}^N \|x_i^{k+1} - x_i^*\|^2 + \frac{1}{2\gamma\rho} \|\lambda^{k+1} - \lambda^*\|^2 + \frac{1}{2} \sum_{i=1}^N \|x_i^{k+1} - x_i^*\|_{\rho A_i^T A_i + P_i}^2 + H_k \end{aligned}$$

where  $H_k = \frac{1}{2} \sum_{i=1}^N \|x_i^{k+1} - x_i^k\|_{\rho A_i^T A_i + P_i}^2 + \frac{2-\gamma}{2\gamma^2\rho} \|\lambda^{k+1} - \lambda^k\|^2 + \frac{1}{\gamma} \left\langle \lambda^k - \lambda^{k+1}, A(x^k - x^{k+1}) \right\rangle$ . This is the desired inequality.  $\square$

We notice that the term  $\alpha \sum_{i=1}^N \|x_i^{k+1} - x_i^*\|^2$  in the inequality (3.2) is a gain for the strongly convexity of the costs. For the linear convergence, we make it to bound the distance  $\|\lambda^k - \lambda^*\|^2$  between the multipliers  $\lambda^k$  and  $\lambda^*$  in the following lemma.

**Lemma 3.4.** *Suppose Assumption 2.1-2.5 hold. Let  $L = \max_{1 \leq i \leq N} L_i^2$  and  $D = \sum_{i=1}^N \|A_i^T\|^2$ . Then it holds that*

$$\begin{aligned}
& c_A^2 \|\lambda^k - \lambda^*\|^2 \\
& \leq 2L \sum_{i=1}^N \|x_i^{k+1} - x_i^*\|^2 + 4\rho^2 D \left\| \sum_{j=1}^N A_j (x_j^k - x_j^*) \right\|^2 \\
& \quad + 4 \sum_{i=1}^N \left\| (\rho A_i^T A_i + P_i) (x_i^{k+1} - x_i^k) \right\|^2.
\end{aligned} \tag{3.8}$$

*Proof.* Using Lemma 3.1 and (3.4), we get

$$\begin{aligned}
c_A^2 \|\lambda^k - \lambda^*\|^2 & \leq \sum_{i=1}^N \|A_i^T (\lambda^k - \lambda^*)\|^2 \\
& \leq \sum_{i=1}^N \left\| \nabla f_i(x_i^{k+1}) - \nabla f_i(x_i^*) + \rho A_i^T \left( A_i (x_i^{k+1} - x_i^*) + \sum_{j \neq i} A_j (x_j^k - x_j^*) \right) \right. \\
& \quad \left. + P_i (x_i^{k+1} - x_i^k) \right\|^2.
\end{aligned}$$

We apply the Cauchy-Schwartz inequality to deduce

$$\begin{aligned}
& c_A^2 \|\lambda^k - \lambda^*\|^2 \\
& \leq 2 \sum_{i=1}^N \left\| \nabla f_i(x_i^{k+1}) - \nabla f_i(x_i^*) \right\|^2 \\
& \quad + 2 \sum_{i=1}^N \left\| \rho A_i^T \left( A_i (x_i^{k+1} - x_i^*) + \sum_{j \neq i} A_j (x_j^k - x_j^*) \right) + P_i (x_i^{k+1} - x_i^k) \right\|^2 \\
& = 2 \sum_{i=1}^N \left\| \nabla f_i(x_i^{k+1}) - \nabla f_i(x_i^*) \right\|^2 \\
& \quad + 2 \sum_{i=1}^N \left\| \rho A_i^T \left( \sum_{j=1}^N A_j (x_j^k - x_j^*) \right) + (\rho A_i^T A_i + P_i) (x_i^{k+1} - x_i^k) \right\|^2,
\end{aligned}$$

where the latter equality holds by a simple rearrangement. Now we apply Assumption 2.3 and the Cauch-Schwartz inequality to obtain

$$\begin{aligned}
& c_A^2 \|\lambda^k - \lambda^*\|^2 \\
& \leq 2 \sum_{i=1}^N L_i^2 \|x_i^{k+1} - x_i^*\|^2 \\
& \quad + 4\rho^2 \sum_{i=1}^N \left\| A_i^T \left( \sum_{j=1}^N A_j (x_j^k - x_j^*) \right) \right\|^2 + 4 \sum_{i=1}^N \left\| (\rho A_i^T A_i + P_i) (x_i^{k+1} - x_i^k) \right\|^2.
\end{aligned}$$

Applying the Cauchy-Schwartz inequality once more, we obtain

$$\begin{aligned}
& c_A^2 \left\| \lambda^k - \lambda^* \right\|^2 \\
& \leq 2 \left( \max_{1 \leq i \leq N} L_i^2 \right) \sum_{i=1}^N \left\| x_i^{k+1} - x_i^* \right\|^2 \\
& \quad + 4\rho^2 \sum_{i=1}^N \left\| A_i^T \right\|^2 \left\| \sum_{j=1}^N A_j (x_j^k - x_j^*) \right\|^2 + 4 \sum_{i=1}^N \left\| \left( \rho A_i^T A_i + P_i \right) (x_i^{k+1} - x_i^k) \right\|^2 \\
& = 2L \sum_{i=1}^N \left\| x_i^{k+1} - x_i^* \right\|^2 + 4\rho^2 D \left\| \sum_{j=1}^N A_j (x_j^k - x_j^*) \right\|^2 \\
& \quad + 4 \sum_{i=1}^N \left\| \left( \rho A_i^T A_i + P_i \right) (x_i^{k+1} - x_i^k) \right\|^2.
\end{aligned}$$

This proves the lemma.  $\square$

We derive the following result using Lemma 3.3 and Lemma 3.4.

**Proposition 3.5.** *Suppose Assumptions 2.1-2.5 hold. For  $s > 0$ , we have the following inequality*

$$\begin{aligned}
& \left( \frac{1}{2\gamma\rho} - sc_A^2 \right) \left\| \lambda^k - \lambda^* \right\|^2 + \frac{1}{2} \sum_{i=1}^N \left\| x_i^k - x_i^* \right\|_{(\rho+4Ns\rho^2D)A_i^T A_i + P_i}^2 \\
& \geq \frac{1}{2\gamma\rho} \left\| \lambda^{k+1} - \lambda^* \right\|^2 + \frac{1}{2} \sum_{i=1}^N \left\| x_i^{k+1} - x_i^* \right\|_{\rho A_i^T A_i + P_i + 2(\alpha-2Ls)I_{n_i}}^2 + G_k,
\end{aligned} \tag{3.9}$$

where  $H_k := \frac{1}{2} \sum_{i=1}^N \left\| x_i^{k+1} - x_i^k \right\|_{\rho A_i^T A_i + P_i}^2 + \frac{2-\gamma}{2\gamma^2\rho} \left\| \lambda^{k+1} - \lambda^k \right\|^2 + \frac{1}{\gamma} \left\langle \lambda^k - \lambda^{k+1}, A(x^k - x^{k+1}) \right\rangle$

and  $G_k := H_k - 4s \sum_{i=1}^N \left\| \left( \rho A_i^T A_i + P_i \right) (x_i^{k+1} - x_i^k) \right\|^2$ .

*Proof.* We recall from Lemma 3.3 the following inequality

$$\begin{aligned}
& \frac{1}{2\gamma\rho} \left\| \lambda^k - \lambda^* \right\|^2 + \frac{1}{2} \sum_{i=1}^N \left\| x_i^k - x_i^* \right\|_{\rho A_i^T A_i + P_i}^2 \\
& \geq \alpha \sum_{i=1}^N \left\| x_i^{k+1} - x_i^* \right\|^2 + \frac{1}{2\gamma\rho} \left\| \lambda^{k+1} - \lambda^* \right\|^2 + \frac{1}{2} \sum_{i=1}^N \left\| x_i^{k+1} - x_i^* \right\|_{\rho A_i^T A_i + P_i}^2 \\
& \quad + \frac{1}{2} \sum_{i=1}^N \left\| x_i^{k+1} - x_i^k \right\|_{\rho A_i^T A_i + P_i}^2 + \frac{2-\gamma}{2\gamma^2\rho} \left\| \lambda^{k+1} - \lambda^k \right\|^2 + \frac{1}{\gamma} \left\langle \lambda^k - \lambda^{k+1}, A(x^k - x^{k+1}) \right\rangle.
\end{aligned} \tag{3.10}$$

Next we recall the inequality of Lemma 3.4 given as

$$\begin{aligned}
& c_A^2 \left\| \lambda^k - \lambda^* \right\|^2 \\
& \leq 2L \sum_{i=1}^N \left\| x_i^{k+1} - x_i^* \right\|^2 \\
& \quad + 4\rho^2 D \left\| \sum_{j=1}^N A_j (x_j^k - x_j^*) \right\|^2 + 4 \sum_{i=1}^N \left\| \left( \rho A_i^T A_i + P_i \right) (x_i^{k+1} - x_i^k) \right\|^2.
\end{aligned} \tag{3.11}$$

Summing (3.11) multiplied by  $s > 0$  and (3.10), we get

$$\begin{aligned}
& \left( \frac{1}{2\gamma\rho} - s c_A^2 \right) \left\| \lambda^k - \lambda^* \right\|^2 + \frac{1}{2} \sum_{i=1}^N \left\| x_i^k - x_i^* \right\|_{\rho A_i^T A_i + P_i}^2 + 4s\rho^2 D \left\| \sum_{j=1}^N A_j (x_j^k - x_j^*) \right\|^2 \\
& \geq (\alpha - 2Ls) \sum_{i=1}^N \left\| x_i^{k+1} - x_i^* \right\|^2 + \frac{1}{2\gamma\rho} \left\| \lambda^{k+1} - \lambda^* \right\|^2 + \frac{1}{2} \sum_{i=1}^N \left\| x_i^{k+1} - x_i^* \right\|_{\rho A_i^T A_i + P_i}^2 \\
& \quad + \frac{1}{2} \sum_{i=1}^N \left\| x_i^{k+1} - x_i^k \right\|_{\rho A_i^T A_i + P_i}^2 - 4s \sum_{i=1}^N \left\| \left( \rho A_i^T A_i + P_i \right) (x_i^{k+1} - x_i^k) \right\|^2 \\
& \quad + \frac{2-\gamma}{2\gamma^2\rho} \left\| \lambda^{k+1} - \lambda^k \right\|^2 + \frac{1}{\gamma} \left\langle \lambda^k - \lambda^{k+1}, A(x^k - x^{k+1}) \right\rangle.
\end{aligned} \tag{3.12}$$

We use the Cauchy-Schwartz inequality to find

$$4s\rho^2 D \left\| \sum_{j=1}^N A_j (x_j^k - x_j^*) \right\|^2 \leq 4Ns\rho^2 D \sum_{j=1}^N \left\| A_j (x_j^k - x_j^*) \right\|^2.$$

Using this inequality in (3.12) and rearranging, we get

$$\begin{aligned}
& \left( \frac{1}{2\gamma\rho} - s c_A^2 \right) \left\| \lambda^k - \lambda^* \right\|^2 + \frac{1}{2} \sum_{i=1}^N \left\| x_i^k - x_i^* \right\|_{(\rho+4Ns\rho^2 D)A_i^T A_i + P_i}^2 \\
& \geq \frac{1}{2\gamma\rho} \left\| \lambda^{k+1} - \lambda^* \right\|^2 + \frac{1}{2} \sum_{i=1}^N \left\| x_i^{k+1} - x_i^* \right\|_{\rho A_i^T A_i + P_i + 2(\alpha-2Ls)I_{n_i}}^2 \\
& \quad + \frac{1}{2} \sum_{i=1}^N \left\| x_i^{k+1} - x_i^k \right\|_{\rho A_i^T A_i + P_i}^2 - 4s \sum_{i=1}^N \left\| \left( \rho A_i^T A_i + P_i \right) (x_i^{k+1} - x_i^k) \right\|^2 \\
& \quad + \frac{2-\gamma}{2\gamma^2\rho} \left\| \lambda^{k+1} - \lambda^k \right\|^2 + \frac{1}{\gamma} \left\langle \lambda^k - \lambda^{k+1}, A(x^k - x^{k+1}) \right\rangle,
\end{aligned}$$

which is the desired inequality.  $\square$

To show the main convergence result Proposition 2.6, we need the following two lemmas.

**Lemma 3.6.** *If  $s > 0$  satisfies*

$$s \left[ \rho^2 D \|A_i\|^2 + \frac{L}{N} \right] < \frac{\alpha}{2N}, \tag{3.13}$$

there exists  $\mu_s \in (0, 1)$  such that

$$\|x\|_{(\rho+4Ns\rho^2D)A_i^T A_i+P_i}^2 \leq \mu_s \|x\|_{\rho A_i^T A_i+P_i+2(\alpha-2Ls)I_{n_i}}^2 \quad \forall x \in \mathbb{R}^{n_i}. \quad (3.14)$$

*Proof.* If we choose  $s > 0$  that satisfies (3.13), we have

$$(\rho + 4Ns\rho^2D)A_i^T A_i + P_i \prec \rho A_i^T A_i + P_i + 2(\alpha - 2Ls)I_{n_i}.$$

Thus, we obtain the final result for  $\mu_s \in (0, 1)$ , which is sufficiently close to 1.  $\square$

**Lemma 3.7.** *If one chooses some  $\xi_i > 0$  such that  $0 < \gamma < 2$ ,  $\rho > 0$ , and  $s > 0$  satisfies the following conditions:*

$$\begin{cases} \rho A_i^T A_i + P_i - 8s(\rho A_i^T A_i + P_i)^T(\rho A_i^T A_i + P_i) - \frac{\rho}{\xi_i} A_i^T A_i \succ 0, \\ \sum_{i=1}^N \xi_i < 2 - \gamma, \end{cases}$$

then we have

$$\begin{aligned} & \frac{1}{2} \sum_{i=1}^N \|x_i^{k+1} - x_i^k\|_{\rho A_i^T A_i+P_i}^2 - 4s \sum_{i=1}^N \left\| \left( \rho A_i^T A_i + P_i \right) (x_i^{k+1} - x_i^k) \right\|^2 \\ & + \frac{2-\gamma}{2\gamma^2\rho} \|\lambda^{k+1} - \lambda^k\|^2 + \frac{1}{\gamma} \left\langle \lambda^k - \lambda^{k+1}, A(x^k - x^{k+1}) \right\rangle \geq 0. \end{aligned}$$

*Proof.* Using Young's inequality, we have

$$\begin{aligned} \frac{2}{\gamma} \left\langle \lambda^k - \lambda^{k+1}, A(x^k - x^{k+1}) \right\rangle &= \sum_{i=1}^N \frac{2}{\gamma} (\lambda^k - \lambda^{k+1})^T A_i (x_i^k - x_i^{k+1}) \\ &\geq - \sum_{i=1}^N \left( \frac{\xi_i}{\rho\gamma^2} \|\lambda^{k+1} - \lambda^k\|^2 + \frac{\rho}{\xi_i} \|A_i(x_i^k - x_i^{k+1})\|^2 \right) \\ &= - \frac{\sum_{i=1}^N \xi_i}{\rho\gamma^2} \|\lambda^{k+1} - \lambda^k\|^2 - \sum_{i=1}^N \frac{\rho}{\xi_i} \|A_i(x_i^k - x_i^{k+1})\|^2, \end{aligned} \quad (3.15)$$

for  $\xi_j > 0$  ( $1 \leq j \leq N$ ). Adding  $\sum_{i=1}^N \|x_i^{k+1} - x_i^k\|_{\rho A_i^T A_i+P_i}^2 - 8s \sum_{i=1}^N \left\| \left( \rho A_i^T A_i + P_i \right) (x_i^{k+1} - x_i^k) \right\|^2$ , and  $\frac{2-\gamma}{\gamma^2\rho} \|\lambda^{k+1} - \lambda^k\|^2$  to both sides of (3.15), we obtain

$$\begin{aligned} & \sum_{i=1}^N \|x_i^{k+1} - x_i^k\|_{\rho A_i^T A_i+P_i}^2 - 8s \sum_{i=1}^N \left\| \left( \rho A_i^T A_i + P_i \right) (x_i^{k+1} - x_i^k) \right\|^2 \\ & + \frac{2-\gamma}{\gamma^2\rho} \|\lambda^{k+1} - \lambda^k\|^2 + \frac{2}{\gamma} \left\langle \lambda^k - \lambda^{k+1}, A(x^k - x^{k+1}) \right\rangle \\ & \geq \sum_{i=1}^N \|x_i^{k+1} - x_i^k\|_{\rho A_i^T A_i+P_i}^2 - 8s(\rho A_i^T A_i+P_i)^T(\rho A_i^T A_i+P_i) - \frac{\rho}{\xi_i} A_i^T A_i \\ & + \left( \frac{2-\gamma - \sum_{i=1}^N \xi_i}{\gamma^2\rho} \right) \|\lambda^{k+1} - \lambda^k\|^2 \end{aligned}$$

for  $\xi_j > 0$  ( $j = 1, \dots, N$ ). The proof is done.  $\square$

We are now ready to show our main convergence and rate results.

*Proof of Theorem 2.6.* By applying Lemma 3.6 and Lemma 3.7 to the inequality of Proposition 3.5, we get

$$\begin{aligned} & \left( \frac{1}{2\gamma\rho} - sc_A^2 \right) \|\lambda^k - \lambda^*\|^2 + \frac{\mu_s}{2} \sum_{i=1}^N \|x_i^k - x_i^*\|_{\rho A_i^T A_i + P_i + 2(\alpha - 2Ls)I_{n_i}}^2 \\ & \geq \frac{1}{2\gamma\rho} \|\lambda^{k+1} - \lambda^*\|^2 + \frac{1}{2} \sum_{i=1}^N \|x_i^{k+1} - x_i^*\|_{\rho A_i^T A_i + P_i + 2(\alpha - 2Ls)I_{n_i}}^2. \end{aligned}$$

This gives the following inequality

$$\begin{aligned} & \sigma \left( \frac{1}{2\gamma\rho} \|\lambda^k - \lambda^*\|^2 + \frac{1}{2} \sum_{i=1}^N \|x_i^k - x_i^*\|_{\rho A_i^T A_i + P_i + 2(\alpha - 2Ls)I_{n_i}}^2 \right) \\ & \geq \frac{1}{2\gamma\rho} \|\lambda^{k+1} - \lambda^*\|^2 + \frac{1}{2} \sum_{i=1}^N \|x_i^{k+1} - x_i^*\|_{\rho A_i^T A_i + P_i + 2(\alpha - 2Ls)I_{n_i}}^2, \end{aligned}$$

where  $\sigma = \max\{1 - 2\gamma\rho sc_A^2, \mu_s\}$ . The proof is done.  $\square$

#### 4. NUMERICAL EXPERIMENTS

In this section, we present numerical tests of Algorithm 1 supporting the linear convergence property proved in Theorem 2.6. We test the Algorithm 1 for the linearly constrained quadratic programming (LCQP) model [34] and the optimal source allocation problem [37].

**4.1. The LCQP Model.** Here we consider the following LCQP model:

$$\begin{aligned} & \min \sum_{i=1}^N f_i(x_i) \\ & \text{s.t. } \sum_{i=1}^N A_i x_i = c \end{aligned} \tag{4.1}$$

where  $f_i : \mathbb{R}^{n_i} \rightarrow \mathbb{R}$  is given by  $f_i(x_i) = \frac{1}{2}x_i^T H_i x_i + x_i^T q_i$  with  $H_i \in \mathbb{S}_+^{n_i}$ ,  $q_i \in \mathbb{R}^{n_i}$ ,  $A_i \in \mathbb{R}^{m \times n_i}$  and  $c \in \mathbb{R}^m$ . Since  $H_i \in \mathbb{S}_+^{n_i}$ , the quadratic function  $f_i$  is strongly convex for each  $i = 1, \dots, N$ . We take  $n_1 = \dots = n_N := n$  in our experiment. We conduct experiments with the damping parameter  $\gamma \in \{0.5, 1.0, 1.5, 1.9\}$ , and the penalty parameter  $\rho \in \{0.1, 1, 5, 10\}$  in Algorithm 1. The matrices involved in this problem are randomly generated through the following procedure:

- (1) Each  $A_i \in \mathbb{R}^{m \times n_i}$  ( $i = 1, \dots, N$ ) is randomly generated with i.i.d. standard normal entries with satisfying the condition that  $[A_1^T; \dots; A_N^T]$  has full column rank.
- (2) The positive semi-definite matrices  $P_i \in \mathbb{R}^{n_i \times n_i}$  ( $i = 1, \dots, N$ ) are generated by sampling  $B \in \mathbb{R}^{n_i \times n_i}$  with i.i.d. standard normal entries, and then symmetrizing as  $P_i = \frac{1}{2}(B + B^T)$  with checking that the eigenvalues of  $P_i$  are non-negative.

- (3) The positive definite matrices  $H_i \in \mathbb{S}_+^{n_i}$  ( $i = 1, \dots, N$ ) are generated by first sampling  $B \in \mathbb{R}^{n_i \times n_i}$  with i.i.d. standard normal entries, and then symmetrizing as  $H_i = \frac{1}{2}(B + B^T)$  with checking that the eigenvalues of  $H_i$  are positive.
- (4) We first generate the optimal solution  $x_i^* \in \mathbb{R}^{n_i}$  and  $\lambda^* \in \mathbb{R}^m$  as follows:  $x_i^* = \text{randn}(n_i, 1)$  for  $i = 1, \dots, N$ , and  $\lambda^* = \text{randn}(m, 1)$ ;
- (5) Next we define  $q_i \in \mathbb{R}^{n_i}$  ( $i = 1, \dots, N$ ) and  $c \in \mathbb{R}^m$  as follows:  $q_i = -H_i x_i^* + A_i^T \lambda^*$  and  $c = \sum_{i=1}^N A_i x_i^*$  so that  $(x_1^{*T}, \dots, x_N^{*T}, \lambda^{*T})^T$  satisfies the KKT conditions.

We update  $x_i^{k+1}$  in (1.5) using the equation from the optimality given as follows:

$$H_i x_i^{k+1} + q_i + \rho A_i^T \left( A_i x_i^{k+1} + \sum_{j \neq i} A_j x_j^k - c - \frac{\lambda^k}{\rho} \right) + P_i(x_i^{k+1} - x_i^k) = 0.$$

We define the error metric at each  $k$ -th iteration for  $u^k = (x_1^k, \dots, x_N^k; \lambda^k)$  as follows:

$$\text{dis}(u^k) := \max \left\{ \|x_1^k - x_1^*\|, \dots, \|x_N^k - x_N^*\|, \|\lambda^k - \lambda^*\| \right\}. \quad (4.2)$$

The initial values  $u^0 = (x_1^0, \dots, x_N^0; \lambda^0)$  are set to zero. The experiments were conducted for  $N = 3$  with  $m = 100$  and  $n = 40$ , and for  $N = 10$  with  $m = 100$  and  $n = 60$ . The experimental results are plotted in Figures 1 and 2 for the 3-block case, and in Figures 3 and 4 for the 10-block case. We find that the value  $\text{dis}(u^k)$  decreases to zero linearly as expected in Theorem 2.6.

**4.2. The Optimal Resource Allocation Problem.** In this example, we consider the following case:

$$\begin{aligned} \min \quad & \sum_{i=1}^N f_i(x_i) \\ \text{s.t.} \quad & \sum_{i=1}^N x_i = 0 \end{aligned}$$

where  $f_i : \mathbb{R} \rightarrow \mathbb{R}$  is given as  $f_i(x_i) = (1/2)a_i(x_i - c_i)^2 + \log[1 + \exp(b_i(x_i - d_i))]$  with the coefficients  $a_i, b_i, c_i, d_i$  generated randomly with uniform distributions on  $[0, 2]$ ,  $[-2, 2]$ ,  $[-10, 10]$ ,  $[-10, 10]$  respectively. We conduct experiments with the damping parameter  $\gamma \in \{0.5, 1.0, 1.5, 1.9\}$ , and the penalty parameter  $\rho \in \{0.1, 1, 5, 10\}$  in Algorithm 1. We update  $x_i^{k+1}$  in (1.5) using the equation derived from the optimality given as follows:

$$a_i(x_i^{k+1} - c_i) + \frac{b_i \exp[b_i(x_i^{k+1} - d_i)]}{1 + \exp[b_i(x_i^{k+1} - d_i)]} + \rho \left( x_i^{k+1} + \sum_{j \neq i} x_j^k - \frac{\lambda^k}{\rho} \right) + P_i(x_i^{k+1} - x_i^k) = 0.$$

In order to study the convergence behavior of the test algorithms, we test two cases: 6-block and 20-block. The value  $u^* = (x_1^*, \dots, x_N^*; \lambda^*)$  is obtained numerically by running Algorithm 1 5,000 iterations. We employ the same error metric  $\text{dis}(u^k)$  as defined in (4.2) for this experiment. The initial values  $u^0 = (x_1^0, \dots, x_N^0; \lambda^0)$  are set to zero. The experimental results are plotted in Figures 5 and 6 for the 6-block case, and in Figures 7






FIGURE 1. Experimental results on 3-block LCQP model for fixed  $\gamma$  under varying  $\rho$ .




FIGURE 2. Experimental results on 3-block LCQP model for fixed  $\rho$  under varying  $\gamma$ .

and 8 for the 20-block case. We observe that the value  $\text{dis}(u^k)$  converges linearly to zero, as anticipated in Theorem 2.6.




FIGURE 3. Experimental results on 10-block LCQP model for fixed  $\gamma$  under varying  $\rho$ .




FIGURE 4. Experimental results on 10-block LCQP model for fixed  $\rho$  under varying  $\gamma$ .

## 5. CONCLUSION

In this paper, we consider a multi-block Jacobi-Proximal ADMM algorithm for solving linearly coupled optimization problems in a parallel way. We established the linear convergence of the algorithm and derived its convergence rate involving more than two variable blocks when the cost functions are strongly convex and Lipschitz continuous. This is the first linear convergence result for the Jacobi-Proximal ADMM.




FIGURE 5. Experimental results on 6-block problem for the optimal resource allocation model with a fixed  $\gamma$  under varying  $\rho$ .




FIGURE 6. Experimental results on 6-block problem for the optimal resource allocation model with a fixed  $\rho$  under varying  $\gamma$ .

## REFERENCES

- [1] Daniel Boley. Local linear convergence of the alternating direction method of multipliers on quadratic or linear programs. *SIAM Journal on Optimization*, 23(4):2183–2207, 2013.
- [2] Xingju Cai, Deren Han, and Xiaoming Yuan. The direct extension of admm for three-block separable convex minimization models is convergent when one function is strongly convex. *Optimization Online*, 229:230, 2014.




FIGURE 7. Experimental results on 20-block problem for the optimal resource allocation model with a fixed  $\gamma$  under varying  $\rho$ .




FIGURE 8. Experimental results on 20-block problem for the optimal resource allocation model with a fixed  $\rho$  under varying  $\gamma$ .

- [3] Venkat Chandrasekaran, Pablo A. Parrilo, and Alan S. Willsky. Latent variable graphical model selection via convex optimization. *The Annals of Statistics*, 40(4):1935–1967, 2012.
- [4] Caihua Chen, Bingsheng He, Yinyu Ye, and Xiaoming Yuan. The direct extension of admm for multi-block convex minimization problems is not necessarily convergent. *Mathematical Programming*, 155(1):57–79, 2016.

- [5] Caihua Chen, Yuan Shen, and Yanfei You. On the convergence analysis of the alternating direction method of multipliers with three blocks. In *Abstract and Applied Analysis*, volume 2013, page 183961. Wiley Online Library, 2013.
- [6] Wei Deng, Ming-Jun Lai, Zhimin Peng, and Wotao Yin. Parallel multi-block admm with  $o(1/k)$  convergence. *Journal of Scientific Computing*, 71:712–736, 2017.
- [7] Wei Deng and Wotao Yin. On the global and linear convergence of the generalized alternating direction method of multipliers. *Journal of Scientific Computing*, 66:889–916, 2016.
- [8] Jim Douglas and Henry H Rachford. On the numerical solution of heat conduction problems in two and three space variables. *Transactions of the American mathematical Society*, 82(2):421–439, 1956.
- [9] Jonathan Eckstein and Dimitri P Bertsekas. On the douglas—rachford splitting method and the proximal point algorithm for maximal monotone operators. *Mathematical programming*, 55:293–318, 1992.
- [10] Jonathan Eckstein, Dimitri P Bertsekas, et al. An alternating direction method for linear programming, 1990.
- [11] Hugh Everett III. Generalized lagrange multiplier method for solving problems of optimum allocation of resources. *Operations research*, 11(3):399–417, 1963.
- [12] Daniel Gabay. Chapter ix applications of the method of multipliers to variational inequalities. In *Studies in mathematics and its applications*, volume 15, pages 299–331. Elsevier, 1983.
- [13] Daniel Gabay and Bertrand Mercier. A dual algorithm for the solution of nonlinear variational problems via finite element approximation. *Computers & Mathematics with Applications*, 2(1):17–40, 1976.
- [14] Daniel Gabay and Bertrand Mercier. A dual algorithm for the solution of nonlinear variational problems via finite element approximation. *Computers & mathematics with applications*, 2(1):17–40, 1976.
- [15] R. Glowinski and A. Marroco. Sur l’approximation, par éléments finis d’ordre un, et la résolution, par pénalisation-dualité d’une classe de problèmes de Dirichlet non linéaires. *Revue française d’automatique, informatique, recherche opérationnelle. Analyse numérique*, 9(R2):41–76, 1975.
- [16] Roland Glowinski and Americo Marroco. Sur l’approximation, par éléments finis d’ordre un, et la résolution, par pénalisation-dualité d’une classe de problèmes de dirichlet non linéaires. *Revue française d’automatique, informatique, recherche opérationnelle. Analyse numérique*, 9(R2):41–76, 1975.
- [17] Deren Han and Xiaoming Yuan. A note on the alternating direction method of multipliers. *Journal of Optimization Theory and Applications*, 155:227–238, 2012.
- [18] Deren Han and Xiaoming Yuan. Local linear convergence of the alternating direction method of multipliers for quadratic programs. *SIAM Journal on numerical analysis*, 51(6):3446–3457, 2013.
- [19] Bingsheng He, Liusheng Hou, and Xiaoming Yuan. On full jacobian decomposition of the augmented lagrangian method for separable convex programming. *SIAM Journal on Optimization*, 25(4):2274–2312, 2015.
- [20] Bingsheng He and Xiaoming Yuan. On the  $o(1/n)$  convergence rate of the douglas—rachford alternating direction method. *SIAM Journal on Numerical Analysis*, 50(2):700–709, 2012.
- [21] Bingsheng He and Xiaoming Yuan. On non-ergodic convergence rate of douglas—rachford alternating direction method of multipliers. *Numerische Mathematik*, 130(3):567–577, 2015.
- [22] Mingyi Hong and Zhi-Quan Luo. On the linear convergence of the alternating direction method of multipliers. *Mathematical Programming*, 162(1):165–199, 2017.
- [23] Min Li, Defeng Sun, and Kim-Chuan Toh. A convergent 3-block semi-proximal admm for convex minimization problems with one strongly convex block. *Asia-Pacific Journal of Operational Research*, 32(04):1550024, 2015.
- [24] Tian-Yi Lin, Shi-Qian Ma, and Shu-Zhong Zhang. On the sublinear convergence rate of multi-block admm. *Journal of the Operations Research Society of China*, 3:251–274, 2015.
- [25] Tianyi Lin, Shiqian Ma, and Shuzhong Zhang. On the convergence rate of multi-block admm. *arXiv preprint arXiv:1408.4265*, 229, 2014.

- [26] Tianyi Lin, Shiqian Ma, and Shuzhong Zhang. On the global linear convergence of the admm with multiblock variables. *SIAM Journal on Optimization*, 25(3):1478–1497, 2015.
- [27] Tianyi Lin, Shiqian Ma, and Shuzhong Zhang. Global convergence of unmodified 3-block admm for a class of convex minimization problems. *Journal of Scientific Computing*, 76(1):69–88, 2018.
- [28] Pierre-Louis Lions and Bertrand Mercier. Splitting algorithms for the sum of two nonlinear operators. *SIAM Journal on Numerical Analysis*, 16(6):964–979, 1979.
- [29] Zhi-Quan Luo and Paul Tseng. Error bound and convergence analysis of matrix splitting algorithms for the affine variational inequality problem. *SIAM Journal on Optimization*, 2(1):43–54, 1992.
- [30] Geoffrey J McLachlan. *Discriminant Analysis and Statistical Pattern Recognition*. Wiley Series in Probability and Statistics. John Wiley & Sons, Nashville, TN, jul 2004.
- [31] Karthik Mohan, Palma London, Maryam Fazel, Daniela Witten, and Su-In Lee. Node-based learning of multiple gaussian graphical models. *J. Mach. Learn. Res.*, 15(1):445–488, jan 2014.
- [32] Renato DC Monteiro and Benar F Svaiter. Iteration-complexity of block-decomposition algorithms and the alternating direction method of multipliers. *SIAM Journal on Optimization*, 23(1):475–507, 2013.
- [33] Yigang Peng, Arvind Ganesh, John Wright, Wenli Xu, and Yi Ma. Rasl: Robust alignment by sparse and low-rank decomposition for linearly correlated images. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, 34(11):2233–2246, 2012.
- [34] Yuan Shen, Yannian Zuo, and Xiayang Zhang. A faster generalized admm-based algorithm using a sequential updating scheme with relaxed step sizes for multiple-block linearly constrained separable convex programming. *Journal of Computational and Applied Mathematics*, 393:113503, 2021.
- [35] Naum Zuselevich Shor. *Minimization methods for non-differentiable functions*, volume 3. Springer Science & Business Media, 2012.
- [36] Min Tao and Xiaoming Yuan. Recovering low-rank and sparse components of matrices from incomplete and noisy observations. *SIAM Journal on Optimization*, 21(1):57–81, 2011.
- [37] Lin Xiao and Stephen Boyd. Optimal scaling of a gradient method for distributed resource allocation. *Journal of optimization theory and applications*, 129:469–488, 2006.
- [38] Xian-Hong Xiao, Hui Deng, and Yang-Dong Xu. Distributed jacobi-proximal admm for consensus convex optimization, 2024.

SUNGKYUNKWAN UNIVERSITY, SUWON, GYEONGGI, 16419 REPUBLIC OF KOREA  
Email address: hlchoimath@gmail.com

SUNGKYUNKWAN UNIVERSITY, SUWON, GYEONGGI, 16419 REPUBLIC OF KOREA  
Email address: choiwc@skku.edu