

This is a preprint of the following chapter:

Antonio Algaba, M. Cinta Domínguez-Moreno, Manuel Merino, and Alejandro J. Rodríguez-Luis, A degenerate Takens–Bogdanov bifurcation in a normal form of Lorenz’s equations, published in *Advances in Nonlinear Dynamics Proceedings of the Second International Nonlinear Dynamics Conference (NODYCON 2021)*, Volume 1, edited by W. Lacarbonara, B. Balachandran, M.J. Leamy, J. Ma, J.A. Tenreiro Machado, G. Stepan, 2022, Springer, Cham, reproduced with permission of Springer Nature Switzerland AG. The final authenticated version is available online at: http://dx.doi.org/10.1007/978-3-030-81162-4_60.

A degenerate Takens–Bogdanov bifurcation in a normal form of Lorenz’s equations

Antonio Algaba¹, M. Cinta Domínguez-Moreno¹, Manuel Merino¹, and Alejandro J. Rodríguez-Luis²

¹ Departamento de Ciencias Integradas, Centro de Estudios Avanzados en Física, Matemática y Computación, Universidad de Huelva, 21071 Huelva, Spain

`algaba@uhu.es`, `mcinta.dominguez@dmate.uhu.es`, `merino@uhu.es`

² Departamento de Matemática Aplicada II, E.T.S. Ingenieros, Universidad de Sevilla, 41092 Sevilla, Spain

`ajrluis@us.es`

Abstract. In this work we consider an unfolding of a normal form of the Lorenz system near a triple-zero singularity. We are interested in the analysis of double-zero bifurcations emerging from that singularity. Their local study provide partial results that are extended by means of numerical continuation methods. Specifically, a curve of heteroclinic connections is detected. It has a degenerate point from which infinitely many homoclinic connections emerge. In this way, we can partially understand the dynamics near the triple-zero singularity.

Keywords: double-zero, Takens–Bogdanov, Lorenz system, heteroclinic

1 Introduction

In the world of dynamical systems the first known and most famous chaotic one is the Lorenz system [1]

$$\dot{x} = \sigma(y - x), \quad \dot{y} = \rho x - y - xz, \quad \dot{z} = -bz + xy, \quad \sigma, \rho, b \in \mathbb{R}. \quad (1)$$

The study of its riveting and intricate dynamical behavior has been carried out in multitude of works (see the recent papers [2–15] and references therein).

One way to obtain important information on the organizing centers of the dynamics in system (1) is by means of the study of local bifurcations of equilibria. Whereas the Hopf and Takens-Bogdanov bifurcations have been fully studied in Lorenz system [9, 10], the Hopf-pitchfork bifurcation (a pair of imaginary eigenvalues and the third one zero; it occurs when $\sigma = -1$, $b = 0$, $\rho > 1$) and the triple-zero bifurcation (a triple-zero eigenvalue; it arises if $\sigma = -1$, $b = 0$, $\rho = 1$) cannot be analyzed by the standard procedures because non-isolated equilibria appear when $b = 0$.

To avoid this problem, in this paper we consider a three-parameter unfolding, that is close to the normal form of the triple-zero bifurcation exhibited by Lorenz system, given by [16]

$$\dot{x} = y, \quad \dot{y} = \varepsilon_1 x + \varepsilon_2 y + Axz + Byz, \quad \dot{z} = \varepsilon_3 z + Cx^2 + Dz^2, \quad (2)$$

where $\varepsilon_1, \varepsilon_2, \varepsilon_3 \approx 0$ and A, B, C, D are real parameters. System (2) exhibits a triple-zero bifurcation when $\varepsilon_1 = \varepsilon_2 = \varepsilon_3 = 0$. These equations are also invariant under the change $(x, y, z) \rightarrow (-x, -y, z)$. We remark that several systems studied in the literature appear as particular cases of (2) for certain parameter choices [17–22]. Without loss of generality [16], we can take $A = -1, C = 1$

$$\dot{x} = y, \quad \dot{y} = \varepsilon_1 x + \varepsilon_2 y - xz + Byz, \quad \dot{z} = \varepsilon_3 z + x^2 + Dz^2. \quad (3)$$

System (3) can have up to four equilibria, namely $E_1 = (0, 0, 0)$, $E_2 = (0, 0, -\varepsilon_3/D)$ if $D \neq 0$ and $E_{3,4} = (\pm\sqrt{-\varepsilon_1(\varepsilon_3 + D\varepsilon_1)}, 0, \varepsilon_1)$ if $\varepsilon_1(\varepsilon_3 + D\varepsilon_1) < 0$. Note that E_1 and E_2 are placed on the z -axis, which is an invariant set.

The characteristic polynomial of the Jacobian matrix of system (3) at the origin E_1 is given by $P(\lambda) = \lambda^3 + p_1\lambda^2 + p_2\lambda + p_3$, where $p_1 = -(\varepsilon_2 + \varepsilon_3)$, $p_2 = \varepsilon_2\varepsilon_3 - \varepsilon_1$, $p_3 = \varepsilon_1\varepsilon_3$. Therefore, the origin exhibits the following bifurcations: (a) A pitchfork bifurcation when $\varepsilon_1 = 0, \varepsilon_2 \neq 0, \varepsilon_3 \neq 0$. The nontrivial equilibria E_3 and E_4 appear when $-\varepsilon_1(\varepsilon_3 + D\varepsilon_1) > 0$. (b) A transcritical bifurcation (involving also E_2) when $\varepsilon_3 = 0, \varepsilon_1 \neq 0, \varepsilon_2 \neq 0, D \neq 0$. (c) A Hopf bifurcation when $\varepsilon_1 < 0, \varepsilon_2 = 0, \varepsilon_3 \neq 0$. (d) A Takens–Bogdanov bifurcation (nondiagonalizable double-zero eigenvalue) when $\varepsilon_1 = 0, \varepsilon_2 = 0, \varepsilon_3 \neq 0$. It is of homoclinic type when $\varepsilon_3 < 0$ and of heteroclinic type if $\varepsilon_3 > 0$. (e) A Hopf-zero bifurcation when $\varepsilon_2 = 0, \varepsilon_3 = 0, \varepsilon_1 < 0$. (f) A double-zero bifurcation (a diagonalizable double-zero eigenvalue) when $\varepsilon_1 = 0, \varepsilon_3 = 0, \varepsilon_2 \neq 0$ (see [16]). (g) A triple-zero bifurcation when $\varepsilon_1 = \varepsilon_2 = \varepsilon_3 = 0$.

We show now that all the information on the equilibrium E_2 can be obtained from the analysis of E_1 . To study E_2 we translate it to the origin by means of the change $x = \tilde{x}, y = \tilde{y}, z = \tilde{z} - \varepsilon_3/D$, that transforms system (3) into

$$\dot{\tilde{x}} = \tilde{y}, \quad \dot{\tilde{y}} = (\varepsilon_1 + \frac{1}{D}\varepsilon_3)\tilde{x} + (\varepsilon_2 - \frac{B}{D}\varepsilon_3)\tilde{y} - \tilde{x}\tilde{z} + B\tilde{y}\tilde{z}, \quad \dot{\tilde{z}} = -\varepsilon_3\tilde{z} + \tilde{x}^2 + D\tilde{z}^2, \quad (4)$$

with $\varepsilon_3, D \neq 0$.

Since system (3) is symmetric to the change

$$(x, y, z, t, \varepsilon_1, \varepsilon_2, \varepsilon_3, B, D) \rightarrow \left(x, y, z - \frac{\varepsilon_3}{D}, t, \varepsilon_1 + \frac{\varepsilon_3}{D}, \varepsilon_2 - \frac{B\varepsilon_3}{D}, -\varepsilon_3, B, D\right), \quad (5)$$

it is direct to obtain the stability and bifurcations of E_2 from the stability and bifurcations of E_1 . Thus, it is enough to study the bifurcations exhibited by E_1 .

2 Double-zero bifurcations of the origin

The Takens–Bogdanov bifurcation of E_1 occurs when $p_2 = p_3 = 0, p_1 \neq 0$, that is, when $\varepsilon_1 = \varepsilon_2 = 0, \varepsilon_3 \neq 0$. For these values the Jacobian matrix has a nondiagonalizable double-zero and a nonzero eigenvalue. Thus, system (3) reads

$$\dot{x} = y, \quad \dot{y} = -xz + Byz, \quad \dot{z} = \varepsilon_3 z + x^2 + Dz^2. \quad (6)$$

By means of the second-order approximation to the center manifold we obtain the third-order reduced system and compute its normal form

$$\dot{x} = y, \quad \dot{y} = a_3x^3 + b_3x^2y, \quad \text{with} \quad a_3 = 1/\varepsilon_3, b_3 = (2 - \varepsilon_3B)/\varepsilon_3^2. \quad (7)$$

Its unfolding is given by

$$\dot{x} = y, \quad \dot{y} = \varepsilon_1x + \varepsilon_2y + a_3x^3 + b_3x^2y. \quad (8)$$

By means of the rescaling

$$x \rightarrow \frac{|\varepsilon_3|\sqrt{|\varepsilon_3|}}{2 - B\varepsilon_3} \bar{x}, \quad y \rightarrow \frac{\varepsilon_3^2\sqrt{|\varepsilon_3|}}{(2 + B\varepsilon_3)} \bar{y}, \quad t \rightarrow \frac{2 - B\varepsilon_3}{|\varepsilon_3|} \tau, \quad (9)$$

system (8) is transformed into

$$\dot{\bar{x}} = \bar{y}, \quad \dot{\bar{y}} = \frac{\varepsilon_1(2 - B\varepsilon_3)^2}{\varepsilon_3^2} \bar{x} + \frac{\varepsilon_2(2 - B\varepsilon_3)}{|\varepsilon_3|} \bar{y} + \text{sgn}(\varepsilon_3)\bar{x}^3 - \bar{x}^2\bar{y}. \quad (10)$$

Thus, the Takens–Bogdanov bifurcation is of heteroclinic case when $\varepsilon_3 > 0$ and of homoclinic case if $\varepsilon_3 < 0$. Note that if $\varepsilon_3 = 0$ a triple-zero bifurcation is present. Moreover, when $(\varepsilon_1, \varepsilon_2, \varepsilon_3) = (0, 0, 2/B)$, with $B \neq 0$, a degenerate Takens–Bogdanov bifurcation occurs. This case will be analyzed below.

In the homoclinic case, when $\varepsilon_3 < 0$, the equilibria of system (10) are $(0, 0)$ and $(\pm(2 - B\varepsilon_3)\sqrt{-\varepsilon_1}/\varepsilon_3, 0)$, $\varepsilon_3 \neq 2/B$, $\varepsilon_1 > 0$. From the Takens–Bogdanov singularity the curves corresponding to the following bifurcations emerge [23–25]: (a) A pitchfork bifurcation of the origin is present when $\varepsilon_1 = 0$. (b) A subcritical Hopf bifurcation of the origin if $\varepsilon_2 = 0$, $\varepsilon_1 < 0$. (c) A supercritical Hopf bifurcation of the nontrivial equilibria when $\varepsilon_2 \approx (2 - B\varepsilon_3)\varepsilon_1/\varepsilon_3$, for $\varepsilon_1 > 0$. (d) A homoclinic connection to the origin for $\varepsilon_2 \approx 4(2 - B\varepsilon_3)\varepsilon_1/(5\varepsilon_3)$, with $\varepsilon_1 > 0$. Since the third eigenvalue ($\varepsilon_3 < 0$) determines the behavior outside the center manifold, these homoclinic connections are attractive. (e) A saddle-node bifurcation of symmetric periodic orbits when $\varepsilon_2 \approx c\varepsilon_1(2 - B\varepsilon_3)/\varepsilon_3$, where $\varepsilon_1 > 0$, $c \approx 0.752$.

In the heteroclinic case, when $\varepsilon_3 > 0$, the equilibria of system (10) are $(0, 0)$ and $(\pm(2 - B\varepsilon_3)\sqrt{-\varepsilon_1}/\varepsilon_3, 0)$, $\varepsilon_3 \neq 2/B$, $\varepsilon_1 < 0$. The following curves are present [23–25]: (a) A pitchfork bifurcation of the origin for (b) A subcritical Hopf bifurcation of the origin for (c) A heteroclinic connection to nontrivial equilibria if $\varepsilon_2 \approx (2 - B\varepsilon_3)\varepsilon_1/(5\varepsilon_3)$, $\varepsilon_1 < 0$. As the third eigenvalue ($\varepsilon_3 > 0$) determines the behavior outside the center manifold, these heteroclinic connections are repulsive.

2.1 Codimension-three degeneracy

The normal form coefficient b_3 , given in (7), vanishes when $\varepsilon_3 = 2/B$, $B \neq 0$.

Thus, considering the fourth-order approximation to the center manifold we obtain the fifth-order reduced system on the center manifold

$$\dot{x} = y, \quad \dot{y} = a_3x^3 + a_5x^5 + b_5x^4y. \quad (11)$$

Multiplying system (11) by $(1 - (a_5/a_3)x^2)$, the x^5 -term can be eliminated

$$\dot{x} = y + \frac{-a_5}{a_3}x^2y, \quad \dot{y} = a_3x^3 + b_5x^4y, \quad \text{with } a_3 = \frac{1}{\varepsilon_3}, \quad b_5 = \frac{-B^4}{8}(5B + 3D). \quad (12)$$

An unfolding is given by

$$\dot{x} = y, \quad \dot{y} = \mu_1x + \mu_2y + a_3x^3 + \mu_3x^2y + b_5x^4y, \quad (13)$$

with $\mu_1 = \varepsilon_1$, $\mu_2 = \varepsilon_2$, $\mu_3 = (2 - B\varepsilon_3)/\varepsilon_3^2$.

To analyze this degeneracy we use the rescaling

$$x \rightarrow \sqrt[6]{\frac{|a_3|}{b_5^2}}u, \quad y \rightarrow b_5\sqrt[6]{\frac{|a_3|^5}{b_5^{10}}}v, \quad t \rightarrow \sqrt[3]{\frac{b_5}{|a_3|^2}}\tau,$$

and then system (13) is transformed into

$$\dot{u} = v, \quad \dot{v} = \tilde{\mu}_1u + \tilde{\mu}_2v + \text{sgn}(a_3)u^3 + \tilde{\mu}_3u^2v + u^4v, \quad (14)$$

where

$$\tilde{\mu}_1 = \frac{\varepsilon_1}{4}\sqrt[3]{\varepsilon_3^4B^8(5B + 3D)^2}, \quad \tilde{\mu}_2 = \frac{-\varepsilon_2}{2}\sqrt[3]{-B^4\varepsilon_3^2(5B + 3D)}, \quad \tilde{\mu}_3 = \frac{-2(2 - B\varepsilon_3)\sqrt[3]{\varepsilon_3}}{\sqrt[3]{B^4(5B + 3D)}}.$$

The analysis of this three-parameter family can be found in [26, 27]. In the parameter space several codimension-two bifurcation curves emerge from the point $(0, 0, 2/B)$.

In the homoclinic case ($\text{sgn}(a_3) < 0$, i.e., $\varepsilon_3 < 0$):

- A nondegenerate Takens-Bogdanov for $\tilde{\mu}_1 = \tilde{\mu}_2 = 0$, $\tilde{\mu}_3 \neq 0$, i.e., $\varepsilon_1 = \varepsilon_2 = 0$, $\varepsilon_3 \neq 2/B$.
- A degenerate Hopf bifurcation of the origin for $\tilde{\mu}_2 = \tilde{\mu}_3 = 0$, $\tilde{\mu}_1 < 0$, i.e., $\varepsilon_2 = 0$, $\varepsilon_3 = \frac{2}{B}$, $\varepsilon_1 < 0$.
- A degenerate Hopf bifurcation of the nontrivial equilibria when $\tilde{\mu}_2 = -\tilde{\mu}_1^2$, $\tilde{\mu}_3 = 0$, i.e., $\varepsilon_3 = \frac{2}{B}$, $\varepsilon_2 = \frac{-1}{2}\varepsilon_1^2B^2(5B + 3D)$, $\varepsilon_1 > 0$.
- A degenerate (zero-trace) homoclinic connection to the origin when $\tilde{\mu}_2 = 0$, $\tilde{\mu}_3 = \frac{-8\tilde{\mu}_1}{7}$, $\tilde{\mu}_1 > 0$, i.e., $\varepsilon_2 = 0$, $\varepsilon_1 = \frac{7(2 - B\varepsilon_3)}{B^4\varepsilon_3(5B + 3D)} > 0$.
- A cusp of saddle-node bifurcations of periodic orbits when $\tilde{\mu}_2 = c_3\tilde{\mu}_1^2$, $\tilde{\mu}_3 = c_4\tilde{\mu}_1$, $\tilde{\mu}_1 > 0$, with $c_3 \approx 1.5713$, $c_4 \approx -3.3484$. That is, for $\varepsilon_2 = \frac{2c_3(2 - B\varepsilon_3)^2}{c_4^2B^2(5B + 3D)}$, $\varepsilon_1 = \frac{-8(2 - B\varepsilon_3)}{c_4\varepsilon_3B^4(5B + 3D)} > 0$.

In the heteroclinic case ($\text{sgn}(a_3) > 0$, i.e., $\varepsilon_3 > 0$):

- A nondegenerate Takens-Bogadnov bifurcation when $\tilde{\mu}_1 = \tilde{\mu}_2 = 0$, $\tilde{\mu}_3 \neq 0$, i.e., $\varepsilon_1 = \varepsilon_2 = 0$, $\varepsilon_3 \neq 2/B$.
- A degenerate Hopf bifurcation of the origin for $\tilde{\mu}_2 = \tilde{\mu}_3 = 0$, $\tilde{\mu}_1 < 0$, i.e., $\varepsilon_2 = 0$, $\varepsilon_3 = 2/B$, $\varepsilon_1 < 0$.

- A degenerate (zero-trace) heteroclinic connection when $\tilde{\mu}_2 = 0$, $\tilde{\mu}_3 \approx \frac{3}{7}\tilde{\mu}_1$, $\tilde{\mu}_1 < 0$, i.e., $\varepsilon_2 = 0$, $\varepsilon_1 \approx -\frac{7}{3}(2 - B\varepsilon_3)^3$.

We end this section with two useful comments for Sect. 3. On the one hand, the information on the Takens–Bogdanov bifurcation exhibited by E_2 can be easily obtained from the above results by using the symmetry (5): it is of homoclinic-type if $\varepsilon_3 > 0$ and of heteroclinic-type when $\varepsilon_3 < 0$. On the other hand, the (diagonalizable) double-zero bifurcation exhibited by the origin has been analyzed in [16]. There, the following expression is obtained for the curve of heteroclinic connections between E_1 and E_2 (in the nondegenerate case $B \neq 2\Delta - \Delta^3\varepsilon_1$)

$$\varepsilon_1 \approx -\frac{1}{2D}\varepsilon_3 + \frac{a^2(-2\Delta + B)}{4D(3a + 2)(-\Delta^2\varepsilon_1 + 1)}\varepsilon_3^2, \quad \text{where } \Delta = \frac{1}{\varepsilon_2}, \quad a = \frac{\Delta^3\varepsilon_1 - \Delta}{D} > 0.$$

3 Numerical study

With the information provided in Sect. 2, we are going to perform a numerical study of system (3) with the continuation software AUTO [28]. Specifically, we will draw bifurcation sets in the $(\varepsilon_1, \varepsilon_3)$ -parameter plane, in a neighbourhood of the degeneracies DZ (diagonalizable double-zero) and TB (Takens–Bogdanov). We will fix $\varepsilon_2 = -1$, $B = -0.1 < 0$ and $D = 0.01 > 0$ according to the values used in [22].

First we draw, in Figs. 1(a) and 1(b), partial bifurcation sets with the bifurcation curves related to the degeneracies DZ (of the origin) and TB (of E_2) in the fourth and second quadrants, respectively. We can observe the curves **h** (Hopf bifurcation of the equilibria $E_{3,4}$) and **He** (heteroclinic cycle between the equilibria E_1 and E_2). Both curves emerge from the point where the double-zero degeneracy of the origin occurs, DZ = (0, 0) in the $(\varepsilon_1, \varepsilon_3)$ -parameter plane. Moreover, three straight lines intersect at the double-zero point DZ, namely **P**¹ (pitchfork bifurcation of the origin, $\varepsilon_1 = 0$), **P**² (pitchfork bifurcation of E_2 , $\varepsilon_3 = -0.01\varepsilon_1$) and **T** (transcritical bifurcation between E_1 and E_2 , $\varepsilon_3 = 0$). Note that in these pitchfork bifurcations the equilibria $E_{3,4}$ emerge. Fixed a value of ε_3 in a neighborhood of the point DZ, the periodic orbit emerged from **h** is attractive and, increasing the value of ε_1 , it disappears in a heteroclinic cycle **He**.

This cycle **He** is formed by two heteroclinic connections, one is structurally stable (since it goes from E_1 to E_2 on the invariant z -axis) and the other one is more relevant (because it is placed outside the z -axis). To show the differences that exist between the heteroclinic cycles of Figs. 1(a) and 1(b) we draw their projections onto the (x, z) -plane in Figs. 1(c) and 1(d), respectively. Remark that throughout this work, with the aim of simplifying the notation, we will label the heteroclinic cycle (in fact, due to the symmetry, a pair of heteroclinic cycles exist) and the heteroclinic bifurcation with the same symbol, although they are two different objects. Also, when necessary, we will use superscripts to indicate the equilibria that are involved in a certain bifurcation or in its degeneration.

Such heteroclinic cycles are attractive when they emerge from DZ since the saddle quantities $\delta_{E_1} = |\max(\lambda_1, \lambda_3)/\lambda_2|$ and $\delta_{E_2} = |\lambda_2^*/\lambda_3^*|$ satisfy $\delta_{E_1}\delta_{E_2} >$




Fig. 1. For $\varepsilon_2 = -1, B = -0.1, D = 0.01$ partial bifurcation set in a neighborhood of the point DZ: (a) in the fourth quadrant; (b) in the second quadrant. (c) Projection onto the (x, z) -plane of the heteroclinic cycle He of panel (a) that exists when $(\varepsilon_1, \varepsilon_3) \approx (0.2, -0.0043175)$. (d) Projection onto the (x, z) -plane of the heteroclinic cycle He of panel (b) that exists when $(\varepsilon_1, \varepsilon_3) \approx (-0.2, 0.0037414)$.

1. Here we denote the eigenvalues of the Jacobian matrix at the origin E_1 as $\lambda_1, \lambda_3 < 0 < \lambda_2$, where $\lambda_3 = \varepsilon_3$, $\lambda_{2,1} = [\varepsilon_2 \pm \sqrt{\varepsilon_2^2 + 4\varepsilon_1}]/2$, and the eigenvalues of the Jacobian matrix at $E_2 = (0, 0, -100\varepsilon_3)$ as $\lambda_1^* < \lambda_2^* < 0 < \lambda_3^*$, with $\lambda_3^* = -\varepsilon_3$, $\lambda_{2,1}^* = [\varepsilon_2 - \frac{B}{D}\varepsilon_3 \pm \sqrt{(\varepsilon_2 - \frac{B}{D}\varepsilon_3)^2 + 4(\varepsilon_1 + \frac{\varepsilon_3}{D})}]/2$.

As seen in Figs. 1(a) and 1(b), on each of the curves He , there is a point DHe where the heteroclinic cycle is degenerate. In fact, when $\varepsilon_1 > 0$, the equilibrium E_1 is always a real saddle along the curve He but E_2 , that is also a real saddle when it arises from DZ, becomes a saddle-focus from the point $DHe^2 \approx (0.2328879, -0.0050898)$. In the case of the branch located in the fourth quadrant, when $\varepsilon_1 < 0$, the equilibrium E_2 is always a real saddle along the curve He , whereas the equilibrium E_1 , that emerges from DZ as a real saddle, becomes a saddle-focus from $DHe^1 \approx (-0.25, 0.004611)$.

The change in the configuration of E_1 (resp. E_2) on the curve of the heteroclinic connections He in the second (resp. fourth) quadrant implies the appearance of an infinity of bifurcation curves, which arise from the aforementioned point DHe^1 (reps. DHe^2) on the He curve.




Fig. 2. For $\varepsilon_2 = -1, B = -0.1, D = 0.01$ partial bifurcation set: (a) in the second quadrant; (b) zoom of panel (a) in the vicinity of the points DHe^1 (upper panel) and TB (lower panel).

Specifically, in Figs. 2(a) and 2(b), we can see the curve H (of homoclinic connections to the equilibrium E_2) that emerges from the point DHe^1 . In a vicinity of DHe^1 , a saddle periodic orbit emerges from H (since $\delta_{E_2} < 1$ at the points of such a neighborhood). A degenerate point DH appears on H when $(\varepsilon_1, \varepsilon_3) \approx (-8.5339606, 0.0855365)$ because $\delta_{E_2} = 1$. The curve H ends at the point $TB = (-10, 0.1)$ where E_2 undergoes a Takens-Bogdanov bifurcation. This is in agreement with the theoretical results of Sect. 2 that guarantee, for these parameter values, that the Takens-Bogdanov bifurcation of E_2 is of homoclinic type.

In the second quadrant, the Hopf bifurcation h of the nontrivial equilibria $E_{3,4}$ is always supercritical and this curve connects the points DZ and TB (see Fig. 1(b)). In addition to the homoclinic connection curve H , other curves emerge from TB , namely a curve h^2 of Hopf bifurcation of E_2 ($\varepsilon_3 = 0.1$ when $\varepsilon_1 < -10$) and a curve SN_1 of saddle-node bifurcation of periodic orbits. The Hopf bifurcation h^2 is subcritical when it emerges from TB and it becomes supercritical because a degeneracy occurs at $Dh^2 \approx (-10.2487498, 0.1)$ as the first Lyapunov coefficient vanishes. A new curve SN_2 of saddle-node bifurcations of symmetric periodic orbits emerges from Dh^2 . A cusp bifurcation of periodic orbits $CU \approx (-9.7715876, 0.0977398)$ occurs when SN_1 and SN_2 collapse (see lower panel of Fig. 2(b)). The cusp CU is the first one of an infinite sequence of cusps that accumulate to the point DH [24].

Close to the degeneracy DH , other bifurcation curves can be seen in Figs. 3(a) and 3(b). Specifically from the point $TP_p \approx (-8.3738877, 0.0841835)$ (where a T-point heteroclinic loop between E_2 and $E_{3,4}$ exists and whose projection in the (x, z) -plane can be seen in Fig. 3(c)) three curves of global connections arise, namely He^{34} (heteroclinic connection between E_3 and E_4), H^3 (homoclinic connection to $E_{3,4}$) and a curve of homoclinic connections to E_2 (not included in Fig. 3) that ends at DHe^1 (see Figs. 2(a) and 2(b)) [29–31, 6]. The curve H^3 ends




Fig. 3. For $\varepsilon_2 = -1$, $B = -0.1$, $D = 0.01$: (a) partial bifurcation set in the second quadrant. (b) Zoom of panel (a). Projection onto the (x, z) -plane of T-point heteroclinic loops connecting E_2 and the equilibria $E_{3,4}$ (note that because of the symmetry a pair of the corresponding orbits exists): (c) principal T-point for $(\varepsilon_1, \varepsilon_3) \approx (-8.3738877, 0.0841835)$; (d) secondary T-point when $(\varepsilon_1, \varepsilon_3) \approx (-8.4159326, 0.0850368)$.

in a secondary T-point between E_2 and $E_{3,4}$ for $TP_s \approx (-8.4159326, 0.0850368)$ whose projection on the (x, z) -plane appears in Fig. 3(d).

The presence in the second quadrant of the degenerations analyzed (and the bifurcation curves that arise from them) determines the existence of regions where chaotic attractors exist (see Figs. 4(a) and 4(b)). We note that these attractors, obtained for $\varepsilon_3 = 0.085$, are structurally stable when the value of ε_1 is varied. Moreover, as observed in these figures, their size increases when ε_1 augments along the interval $[-8, -6.3]$.

4 Conclusions

In this work we consider an unfolding of a normal form of the Lorenz system near a triple-zero singularity. The combination of analytical and numerical tools allows to obtain partial interesting information on the complicated dynamics exhibited by system (3) related to a Takens–Bogdanov bifurcation and a diagonal-




Fig. 4. For $\varepsilon_2 = -1, \varepsilon_3 = 0.085, B = -0.1, D = 0.01$ chaotic attractor when: (a) $\varepsilon_1 = -8$, with initial conditions $(x_0, y_0, z_0) = (0, 0.1, -8)$. (b) $\varepsilon_1 = -6.3$, with initial conditions $(x_0, y_0, z_0) = (0, 0.1, -7)$.

izable double-zero degeneracy. Specifically, a degenerate heteroclinic connection, among other global connections, gives rise to infinite homoclinic orbits that will lead to the existence of chaos. Completing in the future the analysis of system (3) will shed new light on the behavior of the Lorenz system near its triple-zero singularity.

References

1. Lorenz E.N. (1963) Deterministic non-periodic flows. *J. Atmos. Sci.* **20** 130–141.
2. Doedel E.J., Krauskopf B., Osinga H.M. (2011) Global invariant manifolds in the transition to preturbulence in the Lorenz system. *Indagationes Math.* **22** 222–240.
3. Barrio R., Blesa F., Serrano S. (2011) Global organization of spiral structures in biparameter space of dissipative systems with Shilnikov saddle-foci. *Phys. Rev. E* **84** 035201.
4. Barrio R., Shilnikov A.L., Shilnikov L. (2012) Kneadings, symbolic dynamics and painting Lorenz chaos. *Int. J. Bifurcation Chaos* **22** 1230016.
5. Algaba A., Fernández-Sánchez F., Merino M., Rodríguez-Luis A.J. (2014) Centers on center manifolds in the Lorenz, Chen and Lü systems. *Commun. Nonlinear Sci. Numer. Simul.* **19** 772–775.
6. Algaba A., Fernández-Sánchez F., Merino M., Rodríguez-Luis A.J. (2015) Analysis of the T-point-Hopf bifurcation in the Lorenz system. *Commun. Nonlinear Sci. Numer. Simul.* **22** 676–691.
7. Creaser J.L., Krauskopf B., Osinga H.M. (2015) α -flips and T-points in the Lorenz system. *Nonlinearity* **28** R39–R65.
8. Doedel E.J., Krauskopf B., Osinga H.M. (2015) Global organization of phase space in the transition to chaos in the Lorenz system. *Nonlinearity* **28** R113–R139.
9. Algaba A., Domínguez-Moreno M.C., Merino M., Rodríguez-Luis A.J. (2015) Study of the Hopf bifurcation in the Lorenz, Chen and Lü systems. *Nonlinear Dynam.* **79** 885–902.

10. Algaba A., Domínguez-Moreno M.C., Merino M., Rodríguez-Luis A.J. (2016) Takens–Bogdanov bifurcations of equilibria and periodic orbits in the Lorenz system. *Commun. Nonlinear Sci. Numer. Simul.* **30** 328–343.
11. Algaba A., Gamero E., Merino M., Rodríguez-Luis A.J. (2016) Resonances of periodic orbits in the Lorenz system. *Nonlinear Dynam.* **84** 2111–2136.
12. Algaba A., Merino M., Rodríguez-Luis A.J. (2016) Superluminal periodic orbits in the Lorenz system. *Commun. Nonlinear Sci. Numer. Simul.* **39** 220–232.
13. Creaser J.L., Krauskopf B., Osinga H.M. (2017) Finding first foliation tangencies in the Lorenz system. *SIAM J. Appl. Dyn. Syst.* **16** 2127–2164.
14. Osinga H.M. (2018) Understanding the geometry of dynamics: the stable manifold of the Lorenz system, *J. Roy. Soc. New Zeal.* **48** 203–214.
15. Algaba A., Domínguez-Moreno M.C., Merino M., Rodríguez-Luis A.J. (2018) A Review on Some Bifurcations in the Lorenz System. In: Carmona V., Cuevas-Maraver J., Fernández-Sánchez F., García-Medina E. (eds.) *Nonlinear Systems, Vol. 1. Understanding Complex Systems*. Springer, Cham.
16. Algaba A., Domínguez-Moreno M.C., Merino M., Rodríguez-Luis A.J. (2020) Double-zero degeneracy and heteroclinic cycles in a perturbation of the Lorenz system. Preprint.
17. Shimizu T., Morioka N. (1980) On the bifurcation of a symmetric limit cycle to an asymmetric one in a simple model. *Phys. Lett. A* **76** 201–204.
18. Shil’nikov A.L. (1993) On bifurcations of the Lorenz attractor in the Shimizu-Morioka model. *Physica D* **62** 338–346.
19. Rucklidge A.M. (1993) Chaos in a low-order model of magnetoconvection. *Physica D* **62** 323–337.
20. Liu C., Liu T., Liu L., Liu K. (2004) A new chaotic attractor. *Chaos Soliton Fract.* **22** 1031–1038.
21. Mello L.F., Messias M., Braga D.C. (2008) Bifurcation analysis of a new Lorenz-like chaotic system. *Chaos Soliton Fract.* **37** 1224–1255.
22. Kokubu H., Roussarie R. (2004) Existence of a singularly degenerate heteroclinic cycle in the Lorenz system and its dynamical consequences: Part I. *J. Dyn. Differ. Equ.* **16** 513–557.
23. Guckenheimer, J., Holmes, P.J. (1983) *Nonlinear Oscillations, Dynamical Systems, and Bifurcations of Vector Fields*. Springer, New York.
24. Wiggins S. (2003) *Introduction to Applied Dynamical Systems and Chaos*. Springer, New York.
25. Kuznetsov, Y.A. (2004) *Elements of Applied Bifurcation Theory*. Springer, New York.
26. Li C., Rousseau C. (1990) Codimension 2 symmetric homoclinic bifurcations and application to 1:2 resonance. *Can. J. Math.* **42** 191–212.
27. Rodríguez-Luis A.J., Freire E., Ponce, E. (1991) On a codimension 3 bifurcation arising in an autonomous electronic circuit, in *Bifurcation and Chaos: Analysis, Algorithms, Applications*, R. Seydel et al. (eds.), International Series of Numerical Mathematics, vol. 97, pp. 301–306, Birkhäuser, Basel.
28. Doedel E.J. et al. (2012) AUTO-07P: Continuation and bifurcation software for ordinary differential equations. Technical report, Concordia University.
29. Glendinning P., Sparrow C. (1986) T-points: a codimension two heteroclinic bifurcation, *J. Statist. Phys.* **43** 479–488.
30. Fernández-Sánchez F., Freire E., Rodríguez-Luis A.J. (2002) T-Points in a \mathbb{Z}_2 -symmetric electronic oscillator. *Nonlinear Dynam.* **28** 53–69.
31. Fernández-Sánchez F., Freire E., Rodríguez-Luis A.J. (2008) Analysis of the T-point–Hopf bifurcation. *Physica D* **237** 292–305.