
AN IMPROVED CENTRAL LIMIT THEOREM FOR THE EMPIRICAL SLICED WASSERSTEIN DISTANCE

 **David Rodríguez-Vitores***

Universidad de Valladolid and IMUVa
Valladolid, Spain

 **Eustasio del Barrio Tellado**

Universidad de Valladolid and IMUVa
Valladolid, Spain

 **Jean-Michel Loubes**

Institut de Mathématiques de Toulouse, INRIA
Toulouse, France

ABSTRACT

Optimal transport theory has become a fundamental tool for handling diverse types of data, with growing applications across various fields. However, the Wasserstein distance presents significant computational and statistical challenges in high-dimensional settings. To address these issues, alternative distances such as the sliced Wasserstein distance, which leverages one-dimensional projections, have been introduced. In this work, we establish a novel central limit theorem for the p -sliced Wasserstein distance, for $p > 1$, using the Efron-Stein inequality—a technique that has proven effective in related problems. This approach yields a central limit theorem centered at the expected value of the empirical cost, under mild regularity conditions. Notably, unlike the general Wasserstein distance in arbitrary dimensions, we demonstrate that, under specific assumptions, the centering constants can be replaced by the population cost, which is essential for statistical inference. This generalizes and significantly refines existing results for the one-dimensional case. Consequently, we present the first asymptotically valid inference framework for the sliced Wasserstein distance applicable to measures that are not necessarily compactly supported, for $p > 1$. Finally, we address key practical aspects for inference, including Monte Carlo estimation of the integral and estimation of the asymptotic variance, ensuring applicability in real-world scenarios.

Keywords Optimal transport · Sliced Wasserstein distance · Efron-Stein inequality

1 Introduction

In recent years, the rapid growth in both the quantity and variety of data has driven the development of new methods to tackle emerging challenges. Examples from various fields highlight the need for effective tools to handle distributional data. Optimal transport has proven to be a powerful approach in many applications, as it adapts to the underlying geometric structure of the data. Given P, Q in $\mathcal{P}(\mathbb{R}^d)$, the set of probability measures in \mathbb{R}^d endowed with the Borel σ -field, and a cost function $c : \mathbb{R}^d \times \mathbb{R}^d \rightarrow [0, \infty)$ such that $c(x, y)$ represents the cost of moving mass from the initial point x to the final point y , the optimal transport problem consists in solving

$$\inf_{\pi \in \Pi(P, Q)} \int_{\mathbb{R}^d \times \mathbb{R}^d} c(x, y) d\pi(x, y), \quad (1)$$

where $\Pi(P, Q)$ represents the set of probability measures with fixed marginals P and Q . A minimizer π in (1) is known as an optimal transport plan or, interchangeably, an optimal transport coupling. Intuitively, $\pi(x, y)$ represents the amount of mass transported from x to y in the optimal solution. If there exists a map $T : \mathbb{R}^d \rightarrow \mathbb{R}^d$ satisfying $\pi = (I_d, T)\#P$, where I_d is the identity map and $\#$ denotes the push-forward operator, then T is called an optimal

*Correspondence to: david.rodriguez.vitores@uva.es

transport map. In the particular case where the cost function is given by $c(x, y) = \|x - y\|^p$, with $p \geq 1$, the solution to the optimal transport problem allows the definition of a family of distances by taking the p -th root of the infimum. The resulting distance is known as the p -Wasserstein distance, and it is given by

$$W_p(P, Q) = \inf_{\pi \in \Pi(P, Q)} \left(\int_{\mathbb{R}^d \times \mathbb{R}^d} \|x - y\|^p d\pi(x, y) \right)^{1/p}. \quad (2)$$

Despite its intuitive definition and excellent properties, the Wasserstein distance presents a number of challenges for large-scale problems. The first challenge is the computational complexity. Let P_n and Q_m denote the empirical measures built from n and m independent and identically distributed observations from P and Q , respectively. If $n = m$, computing the Wasserstein distance between one-dimensional measures entails a cost that scales at a rate of $n \log(n)$. This follows from the unique characterization of the Wasserstein distance in terms of quantile functions F^{-1} and G^{-1} of P and Q , respectively, given by

$$W_p(P, Q) = \left(\int_{\mathbb{R}} |F^{-1}(t) - G^{-1}(t)|^p dt \right)^{1/p}. \quad (3)$$

Thus, the optimal transport plan between one-dimensional measures is completely characterized by an increasing rearrangement of the measure. However, such a simple characterization of the solution is not available in higher dimensions, and computing the distance requires solving a linear problem, whose complexity scales as n^3 . This hinders the use of optimal transport in machine learning applications, which are of paramount importance nowadays due to their overwhelming reliance on vast quantities of data.

The second fundamental challenge arises from the significant degradation of the statistical performance of Wasserstein-based statistical analysis in high-dimensional settings. Understanding the convergence rates of empirical estimates of the distance to the theoretical values is a fundamental task. Under suitable moment assumptions, [FG15, WB19] proved that if $P = Q$ and $d > 2p$, then $\mathbb{E}(W_p^p(P_n, Q_n))$ converges to 0 at the rate $n^{-2/d}$. Similarly, if $P \neq Q$ and $d \geq 5$, assuming that both measures are absolutely continuous and sub-Weibull, [MNW24] showed that $\mathbb{E}(W_p^p(P_n, Q_n) - W_p^p(P, Q))$ converges at the rate $n^{-\min(2, p)/d}$, demonstrating that the Wasserstein distance suffers from the curse of dimensionality.

This undesirable asymptotic behavior hinders the statement of central limit theorems for the Wasserstein distance in arbitrary dimension, both for $P = Q$ and $P \neq Q$, provided that P and Q are sufficiently regular. The first result in arbitrary dimension was established in [dBL19], where a central limit theorem for the empirical Wasserstein distance centered at its expected value was given, under mild assumptions, for $p = 2$. This was later extended in [dBGSL24] to a broader class of cost functions, including $c(x, y) = \|x - y\|^p$ for $p > 1$, yielding the central limit theorem

$$\sqrt{n} \left(W_p^p(P_n, Q) - \mathbb{E}(W_p^p(P_n, Q)) \right) \rightsquigarrow N(0, \sigma_{P, Q}^2), \quad (4)$$

where the Gaussian limit is non-degenerate only if $P \neq Q$. The convergence in (4) has limited applicability for inferential purposes, since the centering constant is not $W_p^p(P, Q)$, as one would desire. Moreover, this convergence, along with the previous expectation bounds, ensures that a central limit theorem centered at the population value $W_p^p(P, Q)$ cannot be established in arbitrary dimension if P and Q are sufficiently regular. Therefore, distributional limits are primarily available either in the discrete setting, where they can be derived for both $P = Q$ and $P \neq Q$ (see [SM17, TSM19]), or in the one-dimensional setting. In this case, initial work focused on the case $P = Q$ (see [dBGM99] for $p = 1$, [dBGU05] for $p = 2$), whereas for $P \neq Q$, [dBGL19] provided sufficient assumptions to ensure that the bias induced by replacing the expected value by $W_p^p(P, Q)$ in (4) vanishes as $n \rightarrow \infty$, for $p \geq 2$. Some authors have also explored alternative assumptions. For example, for $P \neq Q$, [HKMS24] establishes a central limit theorem for compactly supported measures in dimension $d \leq 3$.

To address these issues, various alternative distance metrics have been proposed, aiming to retain the favorable properties of the Wasserstein distance while mitigating its computational and statistical challenges. Notably, entropically regularized optimal transport, introduced by [Cut13], adds a convex regularization term to the original problem, resulting in improved computational and statistical properties ([dBSLW23, MNW19]). Numerous approaches have also been developed to leverage the unique aspects of the one-dimensional setting. In this work, we focus in one of the most prominent proposals, the sliced Wasserstein distance ([BRPP15]), which is defined as an average over one-dimensional projections

$$SW_p(P, Q) = \left(\int_{\mathbb{S}^{d-1}} W_p^p(P^\theta, Q^\theta) d\sigma(\theta) \right)^{1/p}, \quad (5)$$

where $\text{Pr}_\theta(x) = \langle \theta, x \rangle$ and $P^\theta = \text{Pr}_\theta \# P$, $Q^\theta = \text{Pr}_\theta \# Q$ are the projected distributions. This distance retains key properties of the Wasserstein distance while mitigating the curse of dimensionality, as demonstrated in prior work. We now review related contributions in this direction.

Related work

The statistical analysis of the sliced Wasserstein distance has evolved alongside other projection-based distances. In particular, the max-sliced distance replaces the integral in (5) by the maximum taken over all directions in \mathbb{S}^{d-1} . The first important results for these projection-based distances were developed specifically for each problem. [NWR22] derived expectation bounds with rates independent of the dimension for the max-sliced version (and a subspace generalization). [MBW22] adapts the previous bounds to the sliced setting, extending their validity to a trimmed version of (5), and demonstrating the first distributional limit for the sliced Wasserstein distance and its trimmed version, for $p > 1$ and $P \neq Q$ (in the sense that the asymptotic distribution vanishes if $P = Q$, as in (4)). Denoting by $F_{\theta,n}$ and F_{θ} the cumulative distribution function of the projected distributions along the direction of θ , their asymptotic result is based on the weak convergence of the empirical process $\mathbb{G}_n(\theta, x) = \sqrt{n}(F_{\theta,n}(x) - F_{\theta}(x))$, along with Hadamard differentiability and the functional delta method. The work primarily focuses on the trimmed version and the assumptions are rather strong for the untrimmed setting, involving

$$SJ_{\infty}(P) = \operatorname{esssup}_{0 < t < 1} \frac{1}{f_{\theta}(F_{\theta}^{-1}(t))} < \infty, \quad (6)$$

where f_{θ} denotes the density associated with F_{θ} . An important step in the derivation of distributional limits for projection-based distances was the unifying approach presented in [XNW22]. Using ideas similar to those in [MBW22], and inspired by the duality-based approach in [HMWM24], [XNW22] investigates the weak convergence of the sliced process $\mathbb{G}_n(\theta) = \sqrt{n}(\operatorname{SW}_p^{\theta}(P_n, Q^{\theta}) - \operatorname{SW}_p^{\theta}(P, Q^{\theta}))$, under the assumptions of compactly supported probability measures such that the support of the projected probabilities is an interval. Weak convergence for various projection-based distance notions, including the sliced Wasserstein distance, follows directly from Hadamard differentiability and the functional delta method. Their main results primarily address the case $p > 1$, $P \neq Q$, although some insights are also provided for $p = 1$.

Independently, [GKRS24] derived similar results using techniques more closely related to optimal transport. In particular, they leverage the dual formulation of the one-dimensional Wasserstein distance with $p = 1$, expressed as a supremum over Lipschitz functions, to establish the weak convergence of the empirical process indexed by functions $\varphi \circ \operatorname{Pr}_{\theta}$, where φ is 1-Lipschitz and $\theta \in \mathbb{S}^{d-1}$. For $p > 1$ and $P \neq Q$, distributional limits are obtained using the extended functional delta method, under the assumption that the probability measures are compactly supported with convex support. Additionally, for $p = 1$, distributional limits under mild assumptions are derived for both $P \neq Q$ and $P = Q$. The assumptions in the latter case were further refined by [XH22].

Recently, [HMWM24] provided a refined version of the results in [XNW22] and [GKRS24], again for compactly supported probabilities but with slightly weaker assumptions on the supports, using similar arguments to those in [XNW22]. In a different vein, employing slightly different proof techniques, this work also explores, with great generality, distributional limits for empirical transport problems where the cost function is also estimated from the data.

Contributions

The aim of this work is to establish a new central limit theorem for the empirical sliced Wasserstein distance, based on the use of the Efron-Stein inequality, for $p > 1$. This technique was initially developed in [dBL19]. The results presented in this article were later enhanced, leading to less restrictive assumptions: first for the one-dimensional setting in [dBGL19], and subsequently for general costs in arbitrary dimensions in [dBGSL24]. The same ideas allowed [MNW19] and [dBSLNW23] to state a similar theory for the entropy-regularized Wasserstein distance. Consequently, a natural question is whether we can extend these ideas to the sliced Wasserstein distance. Section 3 is devoted to answer this question. Theorem 3.3 establishes that, under relatively mild assumptions,

$$\sqrt{n}(\operatorname{SW}_p^p(P_n, Q) - \mathbb{E}(\operatorname{SW}_p^p(P_n, Q))) \rightsquigarrow N(0, v_{P,Q}^2), \quad (7)$$

where $v_{P,Q}^2 < \infty$ is a constant that vanishes if $P = Q$. As a by-product of our analysis, we provide a detailed study of convergence and integrability properties of a particular suitable choice of the optimal potentials of the dual problem across the values of $\theta \in \mathbb{S}^{d-1}$, presented in Section 2, which might be of independent interest. The main contribution of Theorem 3.3 is that it enables the decomposition of the original problem into two distinct components: the analysis of the central limit theorem centered at the expected empirical distance, addressed by (7), and the investigation of the bias

$$\sqrt{n}(\mathbb{E}(\operatorname{SW}_p^p(P_n, Q)) - \operatorname{SW}_p^p(P, Q)). \quad (8)$$

Section 4 is devoted to a detailed study of this subject. Theorem 4.1 provides sufficient conditions to ensure (8) for general $p > 1$, significantly improving upon existing one-dimensional results in [dBGL19]. Combining these results

with Theorem 3.3, we derive a central limit theorem for the sliced Wasserstein distance, centered at the population value, as in [MBW22, XNW22, GKRS24, HMWM24]. A notable contribution of our work is that, for the first time, this asymptotic result applies to measures that are not necessarily compactly supported.

Once we have sufficient assumptions to guarantee our central limit theorem centered at the population value, two important considerations arise when performing inference in real-data scenarios. First, the integral in the definition of $\text{SW}_p^p(P_n, Q)$ is not computationally tractable and is typically approximated by an average over k random directions. Section 5 addresses this computational challenge, and Theorem 5.1 establishes a central limit theorem for this approximation, providing a flexible Gaussian limit that adapts to the projection regime ($\tau = \lim \frac{k}{k+n} \in [0, 1]$). To our knowledge, the existing results in the literature offer only probability bounds on the approximation error [XH22]. Thus, Theorem 5.1 is the first central limit theorem that adaptively accounts for the variability induced by the Monte Carlo approximation of the integral. The second challenge is the consistent estimation of the asymptotic variance, which is addressed in Section 6. Finally, Section 7 presents a brief simulation experiment to highlight the role of the τ in statistical inference. Details on the proofs and auxiliary results are provided in the appendix.

Notation

Throughout this work, $\langle \cdot, \cdot \rangle$ and $\| \cdot \|$ denote the usual inner product and the Euclidean norm of \mathbb{R}^d . Generally, we will denote vectors in \mathbb{R}^d by x, y , while reserving the notation s, t for points in \mathbb{R} . \mathbb{S}^{d-1} denotes the unit sphere, i.e., the set of vectors $\theta \in \mathbb{R}^d$ with $\|\theta\| = 1$. We denote by $\mathcal{P}(\mathbb{R}^d)$ the set of probability measures in \mathbb{R}^d , and by $\mathcal{P}_p(\mathbb{R}^d)$ the subset of probabilities with a finite p -th moment. Given $P \in \mathbb{R}^d \in \mathcal{P}(\mathbb{R}^d)$, $\text{supp}(P)$ denotes the smallest closed subset of \mathbb{R}^d verifying $P(\text{supp}(P)) = 1$. We say that P has negligible boundary if

$$\ell_d(\text{supp}(P) \setminus \text{int}(\text{supp}(P))) = 0,$$

where ℓ_d denotes the Lebesgue measure in \mathbb{R}^d . For each $\theta \in \mathbb{S}^{d-1}$, $P^\theta = \text{Pr}_\theta \# P$ is the projected distribution onto the direction of θ , where $\#$ is the push-forward operator and $\text{Pr}_\theta(x) = \langle \theta, x \rangle$. The distribution function of P^θ will be denoted by F_θ , and the density, if it exists, by f_θ . X_1, \dots, X_n will be i.i.d. (independent and identically distributed) random variables drawn from P , and Y_1, \dots, Y_m i.i.d. random variables from Q , assumed to be independent of the X_i 's. The uniform distribution in \mathbb{S}^{d-1} will be denoted by σ , and $\Theta_1, \dots, \Theta_k$ will represent i.i.d. random variables drawn from σ , which will be assumed to be independent of the X_i 's and Y_i 's. The probability measure induced by a random variable Z will be denoted by $\mathcal{L}(Z)$. The symbols \rightarrow_P and \rightsquigarrow will be used to denote convergence in probability and weak convergence, respectively. Finally, we assume that all the random variables considered are defined in a common probability space $(\Omega, \mathcal{F}, \mathbb{P})$.

2 Sliced optimal transport maps and potentials

This section explains the duality theory that underlies the derivation of our main results. In the first part, we introduce the dual problem and review existing results for optimal transport maps and potentials. The second part focuses on adapting this theory to the sliced setting by considering a particular, carefully chosen set of optimal potentials across the values of $\theta \in \mathbb{S}^{d-1}$.

2.1 Preliminaries

Despite the validity of the duality theory presented here for general cost functions, in this work, our attention will be restricted to the cost $c_p(x, y) = h_p(x - y) = \|x - y\|^p$ for $p > 1$. To simplify the notation, we usually denote the cost by $c = c_p$. With this convention, given $P, Q \in \mathcal{P}_p(\mathbb{R}^d)$, the dual formulation of the p -Wasserstein distance guarantees that

$$\text{W}_p^p(P, Q) = \sup_{\phi \in L^1(P)} \int_{\mathbb{R}^d} \phi(x) dP(x) + \int_{\mathbb{R}^d} \phi^c(y) dQ(y), \quad (9)$$

where ϕ^c is the c -conjugate of ϕ , defined as $\phi^c(y) = \inf_{x \in \mathbb{R}^d} (\|x - y\|^p - \phi(x))$. Therefore, the theory of c -concavity plays a fundamental role, for which a thorough exposition can be found in [GM96] and [Vil03]. In particular, we say that $\phi : \mathbb{R}^d \rightarrow [-\infty, \infty)$ is c -concave if there exists $\psi : \mathbb{R}^d \rightarrow [-\infty, \infty)$ such that $\phi = \psi^c$. Obviously, ϕ^c is c -concave, and it is easy to see that $\phi^{cc} \geq \phi$, where the inequality is an equality if and only if ϕ is c -concave. This implies that we can restrict the supremum in (9) to the subset of c -concave functions. Given a c -concave function $\phi : \mathbb{R}^d \rightarrow [-\infty, \infty)$, the c -superdifferential of ϕ is defined as

$$\partial^c \phi = \left\{ (x, y) \in \mathbb{R}^d \times \mathbb{R}^d : f(z) \leq f(x) + (\|z - y\|^p - \|x - y\|^p) \forall z \in \mathbb{R}^d \right\}. \quad (10)$$

For any $x \in \mathbb{R}^d$, we denote by $\partial^c \phi(x)$ the set of points y such that $(x, y) \in \partial^c \phi$. If $\partial^c \phi(x)$ consists of a single point, we denote $\nabla^c \phi(x) = \partial^c \phi(x)$. It is easy to see that if ϕ is c -concave, $\phi(x) + \phi^c(y) \leq c(x, y)$, and the inequality is an equality if and only if $y \in \partial^c \phi(x)$.

Theorems 2.3 and 2.7 in [GM96] ensure that $\pi \in \Pi(P, Q)$ is an optimal transport plan if and only if $\text{supp}(\pi) \subset \partial^c \phi(x)$ for some c -concave function ϕ , and in this case, ϕ is an optimal transport potential. Similarly, it is easy to see that given any c -concave optimal transport potential, there exists an optimal plan π verifying $\text{supp}(\pi) \subset \partial^c \phi(x)$. Moreover, if P is absolutely continuous, Proposition 3.4 in [GM96] guarantees that ϕ is differentiable at almost every point of $\text{dom}(\phi)$, and Theorem 1.2 in [GM96] ensures that $\nabla^c \phi$ is the P -a.s. unique optimal transport map. In addition, if P has negligible boundary and $\text{int}(\text{supp}(P))$ is connected, Corollary 2.7 in [dBGSL24] ensures the uniqueness, up to constants, of the optimal potentials, and Theorem 3.4 in [dBGSL24] establishes uniform convergence of empirical potentials, up to constants, in the compact sets of $\text{int}(\text{supp}(P))$.

2.2 Adaptation to the sliced setting

Let $P, Q \in \mathcal{P}_p(\mathbb{R}^d)$ be such that P has negligible boundary and $\text{int}(\text{supp}(P))$ is connected, and let P_n be the empirical distribution associated with P . Then, for every $\theta \in \mathbb{S}^{d-1}$, $P^\theta, Q^\theta \in \mathcal{P}_p(\mathbb{R})$, and according to Lemma A.1, P^θ has negligible boundary and $\text{int}(\text{supp}(P^\theta))$ is connected. Therefore, in the set of \mathbb{P} -probability one verifying $P_n \rightsquigarrow P$, Theorem 3.4 in [dBGSL24] applies for every $\theta \in \mathbb{S}^{d-1}$. If we denote by ϕ^θ and ϕ_n^θ any c -concave optimal transport potential from P^θ to Q^θ , and from P_n^θ to Q^θ , respectively, there exist constants $a_n^\theta \in \mathbb{R}$ such that

$$\phi_n^\theta - a_n^\theta \rightarrow \phi^\theta$$

in the sense of uniform convergence in the compact sets of $\text{int}(\text{supp}(P^\theta))$. However, to exploit duality within the sliced framework, we need a suitable choice of constants to obtain regularity properties across the values of $\theta \in \mathbb{S}^{d-1}$.

Inspired by [GKRS24], if we are only interested in the limit potentials ϕ^θ , we can proceed as follows. By the comments after Lemma A.2, given any c -concave optimal transport potential ϕ^θ from P^θ to Q^θ , and any reference point $x_0 \in \text{int}(\text{supp}(P))$, it follows that $\phi^\theta(\langle \theta, x_0 \rangle) \in \mathbb{R}$. Thus, we can define

$$\phi_{x_0}^\theta := \phi^\theta - \phi^\theta(\langle \theta, x_0 \rangle). \quad (11)$$

By Corollary 2.7 in [dBGSL24], $\phi_{x_0}^\theta$ is unique, in the sense that if $\tilde{\phi}_{x_0}^\theta$ is any other c -concave optimal transport potential from P^θ to Q^θ verifying $\tilde{\phi}_{x_0}^\theta(\langle \theta, x_0 \rangle) = 0$, then $\phi_{x_0}^\theta(s) = \tilde{\phi}_{x_0}^\theta(s)$ for every $s \in \text{int}(\text{supp}(P))$. Moreover, Lemma B.2 in [GKRS24] proves continuity of the map $\theta \mapsto \phi_{x_0}^\theta(\langle \theta, x \rangle)$.

However, this definition cannot be extended to empirical potentials ϕ_n^θ . If $\text{supp}(Q^\theta)$ is not lower bounded (resp. upper bounded), and

$$\langle \theta, x_0 \rangle < \min_{i=1, \dots, n} \langle \theta, X_i \rangle \quad (\text{resp. } \langle \theta, x_0 \rangle > \max_{i=1, \dots, n} \langle \theta, X_i \rangle)$$

then every c -concave optimal potential ϕ_n^θ verifies $\phi_n^\theta(\langle \theta, x_0 \rangle) = -\infty$. To check this, consider an optimal transport plan π_n^θ verifying $\text{supp}(\pi_n^\theta) \subset \partial^c \phi_n^\theta$. If Q^θ is not lower bounded, by optimality of π_n^θ , there exists $\{y_n\}_{n=1}^\infty$ such that $y_n \rightarrow -\infty$ and $(\min_{i=1, \dots, n} \langle \theta, X_i \rangle, y_n) \in \text{supp}(\pi_n^\theta) \subset \partial^c \phi_n^\theta$. By definition of $\partial^c \phi_n^\theta$, for every $z < \min_{i=1, \dots, n} \langle \theta, X_i \rangle$,

$$\phi_n^\theta(z) \leq \phi_n^\theta\left(\min_{i=1, \dots, n} \langle \theta, X_i \rangle\right) + \left(|z - y_n|^p - \left|\min_{i=1, \dots, n} \langle \theta, X_i \rangle - y_n\right|^p\right) \xrightarrow{n \rightarrow \infty} -\infty$$

Given any $x_0 \in \text{int}(\text{supp}(P))$, if n is large enough (n depends on $\{X_n\}_{n=1}^\infty$), we can ensure $x_0 \in [\min_{i=1, \dots, n} \langle \theta, X_i \rangle, \max_{i=1, \dots, n} \langle \theta, X_i \rangle]$. Thus, $\phi_{n, x_0}^\theta = \phi_n^\theta - \phi_n^\theta(\langle \theta, x_0 \rangle)$ is well defined for n large enough. Elaborating on these ideas, one could prove convergence $\phi_{n, x_0}^\theta \rightarrow \phi_{x_0}^\theta$. Nevertheless, this approach is not valid to provide integrability guarantees, which are crucial in our analysis.

Throughout this work, we propose an intuitive solution for this problem. Since X_1 is the only point such that its projection belongs to $[\min_{i=1, \dots, n} \langle \theta, X_i \rangle, \max_{i=1, \dots, n} \langle \theta, X_i \rangle]$ for every $\theta \in \mathbb{S}^{d-1}$ and $n \in \mathbb{N}$, we propose to consider X_1 as a reference point. This choice is justified by the following proposition.

Proposition 2.1. *Let $p > 1$, $c = c_p$, and $P, Q \in \mathcal{P}_p(\mathbb{R}^d)$ such that P has negligible boundary and $\text{int}(\text{supp}(P))$ is connected. For each $\theta \in \mathbb{S}^{d-1}$, let $\phi^\theta, \phi_n^\theta, \phi_{n, m}^\theta$ be any c -concave optimal transport potentials between P^θ and Q^θ , P_n^θ and Q^θ , P_n^θ and Q_m^θ , respectively. Then, the following properties hold up to a set of \mathbb{P} -probability zero.*

- *Limit:*

1. For every $\theta \in \mathbb{S}^{d-1}$,

$$\phi_{X_1}^\theta := \phi^\theta - \phi^\theta(\langle \theta, X_1 \rangle) \quad (12)$$

is well defined, and it is a c -concave optimal potential from P^θ to Q^θ verifying $\phi_{X_1}^\theta(\langle \theta, X_1 \rangle) = 0$. Moreover, it is unique in $\text{int}(\text{supp}(P))$.

2. The map $\theta \mapsto \phi_{X_1}^\theta(\langle \theta, x \rangle)$ is continuous for every $x \in \text{int}(\text{supp}(P))$.

- *One-sample:*

3. For every $\theta \in \mathbb{S}^{d-1}$,

$$\phi_{n, X_1}^\theta := \phi_n^\theta - \phi_n^\theta(\langle \theta, X_1 \rangle) \quad (13)$$

is well defined, and it is a c -concave optimal potential from P_n^θ to Q^θ verifying $\phi_{n, X_1}^\theta(\langle \theta, X_1 \rangle) = 0$.

4. For every $\theta \in \mathbb{S}^d$, $\phi_{n, X_1}^\theta \rightarrow \phi_{X_1}^\theta$ in the sense of uniform convergence in the compacts of $\text{int}(\text{supp}(P^\theta))$.

5. For every $\theta \in \mathbb{S}^{d-1}$ and for each compact $K \subset \text{int}(\text{supp}(P^\theta)) \cap \text{dom}(\nabla \phi_{X_1}^\theta)$,

$$\sup_{s \in K} \sup_{t_n \in \partial \phi_{n, X_1}^\theta(s)} |t_n - \nabla \phi_{X_1}^\theta(s)| \rightarrow 0$$

6. Given $x \in \text{int}(\text{supp}(P))$, there exist $n_0 \in \mathbb{N}$ and $M_x < \infty$ such that $|\phi_{n, X_1}^\theta(\langle \theta, x \rangle)| \leq M_x$ for every $n \geq n_0$ and $\theta \in \mathbb{S}^{d-1}$.

- *Two-sample:*

7. Let $m = m(n) \rightarrow \infty$ as $n \rightarrow \infty$. For every $\theta \in \mathbb{S}^{d-1}$,

$$\phi_{n, X_1}^\theta := \phi_{n, m}^\theta - \phi_{n, m}^\theta(\langle \theta, X_1 \rangle) \quad (14)$$

is well defined, and it is a c -concave optimal potential from P_n^θ to Q_m^θ verifying $\phi_{n, X_1}^\theta(\langle \theta, X_1 \rangle) = 0$. Moreover, assertions 4, 5 and 6 are also verified by ϕ_{n, m, X_1}^θ .

Given any family of optimal transport potentials $\{\phi^\theta\}_{\theta \in \mathbb{S}^{d-1}}$, and any probability measure $\mu \in \mathcal{P}(\mathbb{R}^d)$, denote

$$\|\phi^\theta\|_{L^q(\sigma \times \mu)}^q := \int_{\mathbb{S}^{d-1} \times \mathbb{R}^d} |\phi^\theta(\langle \theta, x \rangle)|^q d(\sigma \times \mu)(\theta, x).$$

Our next result establishes expectation bounds on the $L^q(\sigma \times \mu)$ -norm of the c -concave random potentials $\phi_{X_1}^\theta, \phi_{n, X_1}^\theta, \phi_{n, m, X_1}^\theta$ given by (12), (13) and (14), when $\mu = P$ or $\mu = P_n$, and similarly for their c -conjugates. By Proposition 2.1, $\phi_{X_1}^\theta, \phi_{n, X_1}^\theta$ and ϕ_{n, m, X_1}^θ are well defined up to a set of \mathbb{P} -probability zero, which can be ignored, since it does not play any role in the expected values.

Proposition 2.2. *Let $q \geq 1$. Under the same assumptions as in Proposition 2.1, if P, Q have finite moments of order pq , then*

$$\mathbb{E}\left(\|\phi_{X_1}^\theta\|_{L^q(\sigma \times P)}^q\right) \vee \mathbb{E}\left(\|(\phi_{X_1}^\theta)^c\|_{L^q(\sigma \times Q)}^q\right) < \infty, \quad (15)$$

$$\sup_n \mathbb{E}\left(\|\phi_{n, X_1}^\theta\|_{L^q(\sigma \times P_n)}^q\right) \vee \sup_n \mathbb{E}\left(\|(\phi_{n, X_1}^\theta)^c\|_{L^q(\sigma \times Q)}^q\right) < \infty, \quad (16)$$

$$\sup_{n, m} \mathbb{E}\left(\|\phi_{n, m, X_1}^\theta\|_{L^q(\sigma \times P_n)}^q\right) \vee \sup_{n, m} \mathbb{E}\left(\|(\phi_{n, m, X_1}^\theta)^c\|_{L^q(\sigma \times Q_m)}^q\right) < \infty. \quad (17)$$

As an immediate consequence of (15), since $\mathcal{L}(X_1) = P$, we deduce that if we define the map $g(x) = \|\phi_x^\theta\|_{L^q(\sigma \times P)}^q$ or $g(x) = \|(\phi_x^\theta)^c\|_{L^q(\sigma \times Q)}^q$ then $\int g(x) dP(x) < \infty$, which allows us to conclude the following result.

Corollary 2.3. *Under the same assumptions as in Proposition 2.2, for almost every $x_0 \in \text{int}(\text{supp}(P))$,*

$$\|\phi_{x_0}^\theta\|_{L^q(\sigma \times P)}^q \vee \|(\phi_{x_0}^\theta)^c\|_{L^q(\sigma \times Q)}^q < \infty. \quad (18)$$

3 Central limit theorem with centering constant

The aim of this section is to outline the steps leading to Theorem 3.3, which establishes a central limit theorem with a centering constant. The proof hinges on two key results, Propositions 3.1 and 3.2, both relying on the Efron-Stein inequality. The first result leverages the primal representation of the optimal transport problem, while the second is based on its dual formulation, as discussed in Sections 3.1 and 3.2, respectively. Finally, Section 3.3 presents the main result, Theorem 3.3, which follows from Proposition 3.2 using elementary probabilistic arguments.

3.1 Efron-Stein and primal representation

Let $Z_n = f(X_1, \dots, X_n)$, where X_1, \dots, X_n are independent random variables. Given an independent copy (X'_1, \dots, X'_n) of (X_1, \dots, X_n) , denote by $Z_n^i = f(X_1, \dots, X'_i, \dots, X_n)$, for each $i = 1, \dots, n$. The Efron-Stein inequality ensures that

$$\text{Var}(Z_n) \leq \frac{1}{2} \sum_{i=1}^n \mathbb{E} (Z_n - Z_n^i)^2 = \sum_{i=1}^n \mathbb{E} (Z_n - Z_n^i)_+^2.$$

In particular, if X_1, \dots, X_n are i.i.d. and f is a symmetric function of X_1, \dots, X_n , then all the values $\mathbb{E} (Z_n - Z_n^i)_+^2$ are equal and the bound simplifies to

$$\text{Var}(Z_n) \leq n \mathbb{E} (Z_n - Z'_n)_+^2 \quad (19)$$

where $Z'_n = f(X'_1, X_2, \dots, X_n)$. Our first result is based on the application of Efron-Stein to the symmetric variable $Z_n = \text{SW}_p^p(P_n, Q)$, where $P_n = \frac{1}{n} \sum_{i=1}^n \delta_{X_i}$. Thus, we have to compare Z_n with $Z'_n = \text{SW}_p^p(P'_n, Q)$, where $P'_n = \frac{1}{n} (\delta_{X'_1} + \sum_{i=2}^n \delta_{X_i})$. Proposition 3.1 relies on the primary definition of the Wasserstein distance (2) as a minimum between all possible couplings. In particular, for each $\theta \in \mathbb{S}^d$, from the optimal coupling π_n^θ between P_n^θ and Q^θ , we can build a coupling between P'_n^θ and Q^θ , which will be close to the optimal, allowing suitable bounds for the difference $Z_n - Z'_n$. The aforementioned line of reasoning was first presented in Proposition 3.1 in [dBL19], and later generalized to general costs in Lemma 4.1 in [dBGSL24]. Yet, our proof is distinct as it relies on optimal plans rather than optimal maps, offering a simpler method to eliminate smoothness assumptions in Q , without relying on the approximation argument of Corollary 4.3 in [dBGSL24].

Proposition 3.1 (Variance bound). *Let $p > 1$ and $P, Q \in \mathcal{P}_{2p}(\mathbb{R}^d)$. Given independent random variables X_1, X'_1 with probability law P and Y with probability law Q , define the constant*

$$K_p(P, Q) = p^2 \mathbb{E} (\|X - X'\|^{2p})^{\frac{1}{p}} \mathbb{E} (\|X - Y\|^{2p})^{\frac{p-1}{p}}$$

Similarly, define the constant $K_p(Q, P)$. Then,

- (a) $\text{Var}(\text{SW}_p^p(P_n, Q)) \leq \frac{K_p(P, Q)}{n}$.
- (b) $\text{Var}(\text{SW}_p^p(P_n, Q_m)) \leq \frac{K_p(P, Q)}{n} + \frac{K_p(Q, P)}{m}$.

3.2 Efron-Stein and dual representation

In this section, we exploit the dual representation of the Wasserstein distance (9) to obtain an enhanced variance bound. For each $\theta \in \mathbb{S}^{d-1}$ $n \in \mathbb{N}$, let ϕ_n^θ be any c -concave optimal transport potential from P_n^θ to Q^θ . To ensure integrability properties of the limit potentials ϕ^θ from P^θ to Q^θ , we cannot consider any choice across the values of $\theta \in \mathbb{S}^{d-1}$. In particular, we assume that there exists $x_0 \in \text{int}(\text{supp}(P))$ such that $\phi^\theta = \phi_{x_0}^\theta$, defined as in (11). Under suitable regularity assumptions, if we define $\phi_{X_1}^\theta, \phi_{n, X_1}^\theta$ as in (12) and (13),

$$\begin{aligned} \text{W}_p^p(P_n^\theta, Q^\theta) &= \int_{\mathbb{R}} \phi_{n, X_1}^\theta(s) dP_n^\theta(s) + \int_{\mathbb{R}} (\phi_{n, X_1}^\theta)^c(t) dQ^\theta(t) \\ \text{W}_p^p(P'_n{}^\theta, Q^\theta) &\geq \int_{\mathbb{R}} \phi_{n, X_1}^\theta(s) dP'_n{}^\theta(s) + \int_{\mathbb{R}} (\phi_{n, X_1}^\theta)^c(t) dQ^\theta(t) \end{aligned}$$

which allows us to bound the difference $\text{W}_p^p(P_n^\theta, Q^\theta) - \text{W}_p^p(P'_n{}^\theta, Q^\theta)$ by

$$\int_{\mathbb{R}} \phi_{n, X_1}^\theta(s) d(P_n^\theta - P'_n{}^\theta)(s) = \int_{\mathbb{R}} (\phi_{n, X_1}^\theta(s) - \phi_{X_1}^\theta(s)) d(P_n^\theta - P'_n{}^\theta)(s) + \int_{\mathbb{R}} \phi_{X_1}^\theta(s) d(P_n^\theta - P'_n{}^\theta)(s)$$

$$= \int_{\mathbb{R}} (\phi_{n, X_1}^\theta(s) - \phi_{X_1}^\theta(s)) d(P_n^\theta - P_n'^\theta)(s) + \int_{\mathbb{R}} \phi_{x_0}^\theta(s) d(P_n^\theta - P_n'^\theta)(s), \quad (20)$$

where the last equality follows from the definition of $\phi_{X_1}^\theta$. Applying this bound uniformly over \mathbb{S}^{d-1} , we can control the difference $Z_n - Z'_n$. Proposition 3.2 refines Proposition 3.1 by applying the Efron-Stein inequality to a modified version R_n of Z_n , obtained by subtracting the integral over \mathbb{S}^{d-1} of the last term in (20). The proof follows the same approach as Theorem 3.2 in [dBL19] and Theorem 4.6 in [DBGSL24]. The adaptation to the sliced setting is based on the special treatment of optimal potentials described in Section 2.2.

Theorem 3.2. *Let $p > 1$ and $c = c_p$. Assume that $P, Q \in \mathcal{P}_{2p(1+\gamma)}(\mathbb{R}^d)$, for some $\gamma > 0$, and that P is absolutely continuous with negligible boundary and $\text{int}(\text{supp}(P))$ connected. Let $x_0 \in \text{int}(\text{supp}(P))$ verify (18), and $\phi_{x_0}^\theta$ be defined as in (11).*

(a) *If we define*

$$R_n = \text{SW}_p^p(P_n, Q) - \int_{\mathbb{S}^{d-1}} \left(\int_{\mathbb{R}^d} \phi_{x_0}^\theta(\langle \theta, x \rangle) dP_n(x) \right) d\sigma(\theta)$$

then, as $n \rightarrow \infty$, $n\text{Var}(R_n) \rightarrow 0$.

(b) *Let $m = m(n) \rightarrow \infty$ as $n \rightarrow \infty$. If Q is also absolutely continuous with negligible boundary, $\text{int}(\text{supp}(Q))$ is connected, and we define*

$$\begin{aligned} R_{n,m} &= \text{SW}_p^p(P_n, Q_m) - \int_{\mathbb{S}^{d-1}} \left(\int_{\mathbb{R}^d} \phi_{x_0}^\theta(\langle \theta, x \rangle) dP_n(x) \right) d\sigma(\theta) \\ &\quad - \int_{\mathbb{S}^{d-1}} \left(\int_{\mathbb{R}^d} (\phi_{x_0}^\theta)^c(\langle \theta, y \rangle) dQ_m(y) \right) d\sigma(\theta), \end{aligned}$$

then, as $n \rightarrow \infty$, $\frac{nm}{n+m}\text{Var}(R_{n,m}) \rightarrow 0$.

3.3 Central limit theorem

Hereafter, given any c -concave optimal transport potential ϕ^θ and $x \in \mathbb{R}^d$, denote, with a slight abuse of notation, $\phi^\theta(x) = \phi^\theta(\langle \theta, x \rangle)$. Under the assumptions of Proposition 3.2, applying Chebyshev's inequality, $\sqrt{n}(R_n - \mathbb{E}(R_n)) \rightarrow_P 0$. The sequence can be decomposed as

$$\begin{aligned} \sqrt{n}(R_n - \mathbb{E}(R_n)) &= \sqrt{n} \left(\text{SW}_p^p(P_n, Q) - \mathbb{E}(\text{SW}_p^p(P_n, Q)) \right) - \\ &\quad - \sqrt{n} \left(\int_{\mathbb{S}^{d-1}} \int_{\mathbb{R}^d} \phi_{x_0}^\theta(x) dP_n(x) d\sigma(\theta) - \mathbb{E} \left(\int_{\mathbb{S}^{d-1}} \int_{\mathbb{R}^d} \phi_{x_0}^\theta(x) dP_n(x) d\sigma(\theta) \right) \right) \end{aligned}$$

Rewriting the second term,

$$\sqrt{n} \left(\frac{1}{n} \sum_{i=1}^n \int_{\mathbb{S}^{d-1}} \phi_{x_0}^\theta(X_i) d\sigma(\theta) - \mathbb{E} \left(\int_{\mathbb{S}^{d-1}} \phi_{x_0}^\theta(X) d\sigma(\theta) \right) \right) \rightsquigarrow N(0, v_{P,Q}^2),$$

where the convergence follows from the central limit theorem, since the variables have finite second order moment by (18), and the variance is given by

$$v_{P,Q}^2 := \text{Var} \left(\int_{\mathbb{S}^{d-1}} \phi_{x_0}^\theta(X) d\sigma(\theta) \right) = \int_{\mathbb{S}^{d-1}} \int_{\mathbb{S}^{d-1}} \text{Cov}_P(\phi_{x_0}^\theta, \phi_{x_0}^\eta) d\sigma(\theta) d\sigma(\eta).$$

$\text{Cov}_P(\phi_{x_0}^\theta, \phi_{x_0}^\eta)$ does not depend on the particular choice of the constant. Under the assumptions of Theorem 3.2, given any c -concave optimal transport potential ϕ^θ from P^θ to Q^θ , there exists $a^\theta \in \mathbb{R}$ such that $\phi^\theta = \phi_{x_0}^\theta + a^\theta$ in $\text{int}(\text{supp}(P^\theta))$, which implies $\text{Cov}_P(\phi^\theta, \phi^\eta) = \text{Cov}_P(\phi_{x_0}^\theta, \phi_{x_0}^\eta)$. With a similar reasoning for the two sample setting, we conclude:

Theorem 3.3. *Let $p > 1, \gamma > 0$ and $P, Q \in \mathcal{P}_{2p(1+\gamma)}(\mathbb{R}^d)$ be such that P is absolutely continuous with negligible boundary, and $\text{int}(\text{supp}(P))$ is connected. For each $\theta \in \mathbb{S}^{d-1}$, let ϕ^θ be any c -concave optimal transport potential from P^θ to Q^θ . Then,*

(a) Denoting $v_{P,Q}^2 = \int_{\mathbb{S}^{d-1}} \int_{\mathbb{S}^{d-1}} \text{Cov}_P(\phi^\theta, \phi^\eta) d\sigma(\theta) d\sigma(\eta)$,

$$\sqrt{n} \left(\text{SW}_p^p(P_n, Q) - \mathbb{E}(\text{SW}_p^p(P_n, Q)) \right) \rightsquigarrow N(0, v_{P,Q}^2)$$

(b) Let $m = m(n) \rightarrow \infty$ as $n \rightarrow \infty$ be such that $\frac{n}{n+m} \rightarrow \lambda \in [0, 1]$. If Q is also absolutely continuous with negligible boundary, and $\text{int}(\text{supp}(Q))$ is connected, then

$$\sqrt{n} \left(\text{SW}_p^p(P_n, Q_m) - \mathbb{E}(\text{SW}_p^p(P_n, Q_m)) \right) \rightsquigarrow N(0, (1-\lambda)v_{P,Q}^2 + \lambda v_{Q,P}^2)$$

Moreover, in both cases, convergence in stronger W_2 sense is also verified.

4 Removing the centering constant

The aim of this section is to provide sufficient assumptions to ensure

$$0 \leq \sqrt{n} \left(\mathbb{E}(\text{SW}_p^p(P_n, Q)) - \text{SW}_p^p(P, Q) \right) \rightarrow 0. \quad (21)$$

The non-negativity follows from the exact same arguments as in [dBGL19]. Our result relies on the finiteness of the functional $SJ_\alpha(P)$, which was defined in [MBW22] as a generalization of J_α , originally proposed by [BL19]. Given $\alpha \geq 1$ and $P \in \mathcal{P}_p(\mathbb{R}^d)$ such that for every $\theta \in \mathbb{S}^{d-1}$, P^θ has density f_θ , it is defined as

$$SJ_\alpha(P) = \int_{\mathbb{S}^{d-1}} J_\alpha(P^\theta) d\sigma(\theta) = \int_{\mathbb{S}^{d-1}} \left(\int_0^1 \frac{(t(1-t))^{\alpha/2}}{f_\theta(F_\theta^{-1}(t))^\alpha} dt \right) d\sigma(\theta). \quad (22)$$

From the definition, it is easy to see that if $1 \leq \alpha' < \alpha$ and $SJ_\alpha(P) < \infty$, then $SJ_{\alpha'}(P) < \infty$.

Theorem 4.1. Assume $p > 1$ and let $P, Q \in \mathcal{P}_p(\mathbb{R}^d)$ be such that for every $\theta \in \mathbb{S}^{d-1}$, P^θ has distribution function F_θ and density f_θ verifying that $f_\theta(F_\theta^{-1}(t))$ is positive and continuous in $(0, 1)$, and monotone for t sufficiently close to 0 and 1. If there exist $\alpha, \beta > 1$ with $\frac{1}{\alpha} + \frac{1}{\beta} = 1$ and $\gamma > 0$ such that $SJ_{\alpha(1+\gamma)}(P) < \infty$ and the following additional moment assumptions are satisfied

- (i) if $1 < p < 2$, P, Q have finite $\beta(1+\gamma)$ moment,
- (ii) if $p = 2$, Q has finite $\beta(1+\gamma)$ moment,
- (iii) if $p > 2$, P, Q have finite $(p-1)\beta(1+\gamma)$ moment,

then, as $n \rightarrow \infty$ (21) holds. Similarly, if $m = m(n) \rightarrow \infty$ as $n \rightarrow \infty$ and there exist $\alpha', \beta' > 1$ such that all the previous assumptions are also verified exchanging the roles of P and Q , then, as $n \rightarrow \infty$,

$$\sqrt{\frac{nm}{n+m}} \left(\mathbb{E}(\text{SW}_p^p(P_n, Q_m)) - \text{SW}_p^p(P, Q) \right) \rightarrow 0. \quad (23)$$

As a straightforward consequence of Theorems 3.3 and 4.1, we can derive a central limit theorem centered at the population value $\text{SW}_p^p(P, Q)$.

Corollary 4.2. Let $p > 1$, and $P, Q \in \mathbb{R}^d$. If the assumptions of Theorems 3.3 and 4.1 for the one-sample setting are verified, then

$$\sqrt{n} \left(\text{SW}_p^p(P_n, Q) - \text{SW}_p^p(P, Q) \right) \rightsquigarrow N(0, v_{P,Q}^2). \quad (24)$$

Moreover, if the assumptions of the two-sample setting are also verified, then

$$\sqrt{n} \left(\text{SW}_p^p(P_n, Q_m) - \text{SW}_p^p(P, Q) \right) \rightsquigarrow N(0, (1-\lambda)v_{P,Q}^2 + \lambda v_{Q,P}^2) \quad (25)$$

Remark 1. Although Theorems 3.3 and 4.1 require different assumptions, some parts can be related. In particular, if the assumptions of Theorem 3.3 hold, P is absolutely continuous with negligible boundary and $\text{int}(\text{supp}(P))$ is connected. Then, if we further assume that the density of P is continuous and positive in $\text{int}(\text{supp}(P))$, it follows that $f_\theta(F_\theta^{-1}(t))$ is positive and continuous in $(0, 1)$ for every $\theta \in \mathbb{S}^{d-1}$.

Theorem 4.1 and Corollary 4.2 enhance the existing results in the literature. Specifically, Proposition 2.3 in [dBGL19] addresses the same problem in the case $d = 1$. Given $P, Q \in \mathcal{P}(\mathbb{R})$, the assumptions required in [dBGL19] involve twice differentiability of the distribution function F of P ,

$$\sup_{t \in (0,1)} \frac{t(1-t) |f'(F^{-1}(t))|}{f^2(F^{-1}(t))} < \infty, \quad (26)$$

moment assumptions and some additional technical conditions, which are slightly weaker than $J_p(P) < \infty$. Moreover, Proposition 2.3 in [dBGL19] is valid only for $p \geq 2$. First, note that our approach is valid for every $p > 1$. In addition, we do not require the existence of second derivatives, nor (26). Instead, we impose monotonicity of f in the tails. Finally, Theorem 4.1 assumes that there exists $\alpha > 1$ such that $J_\alpha(P) < \infty$, and that P, Q have finite moments of an order that increases as α decreases towards one. Again, this is weaker than $J_p(P) < \infty$. For example, if P and Q are Gaussian, we know that $J_\alpha(P) < \infty$ if and only if $\alpha < 2$, and both P, Q have finite moments of every order. In fact, if P is Gaussian, it is easy to see that $J_\alpha(P) = J_\alpha(N(0, 1))$. Therefore, Theorem 4.1 ensures (21) for every $p > 1$ and every pair of one-dimensional Gaussian distributions P and Q , which was only proven in [dBGL19] for $p \geq 2$.

The same conclusion extends to multivariate Gaussian distributions, which emphasizes the novelty of our results, since it is the first central limit theorem for SW_p^p , $p > 1$, which applies to measures without compact support. Furthermore, in [MBW22], the authors conjectured that a central limit theorem of the form (24) would hold under the assumption $SJ_p(P) < \infty$. In contrast, Corollary 4.2 shows that even $SJ_p(P) < \infty$ is unnecessary. It suffices to assume $SJ_\alpha(P) < \infty$ for some $\alpha > 1$, provided that P and Q have sufficiently many finite moments.

5 Slicing with a finite number of directions

Theorem 4.1 establishes a central limit theorem centered at the population value $\text{SW}_p^p(P, Q)$. However, to carry out inference on this quantity, we must evaluate the integral over \mathbb{S}^{d-1} in the definition of $\text{SW}_p^p(P_n, Q)$. In practice, this integral is typically approximated using the Monte Carlo estimator based on an independent sample $\Theta_1, \dots, \Theta_k$ drawn from σ , given by

$$\text{SW}_{p,k}^p(P_n, Q) := \frac{1}{k} \sum_{i=1}^k \text{W}_p^p(P_n^{\Theta_i}, Q^{\Theta_i}). \quad (27)$$

Thus, an additional source of variation is introduced into the problem, which is controlled by the following proposition.

Proposition 5.1. *Let $p > 1$, and $P, Q \in \mathcal{P}_{2p}(\mathbb{R}^d)$. For each $k \in \mathbb{N}$, let $\Theta_1, \dots, \Theta_k$ be i.i.d. samples drawn from σ . If $k = k(n), m = m(n) \rightarrow \infty$ as $n \rightarrow \infty$, then*

$$\begin{aligned} S_{n,k} &= \sqrt{k} \left(\text{SW}_{p,k}^p(P_n, Q) - \text{SW}_p^p(P_n, Q) \right) \rightsquigarrow N(0, w_{P,Q}^2), \\ S_{n,m,k} &= \sqrt{k} \left(\text{SW}_{p,k}^p(P_n, Q_m) - \text{SW}_p^p(P_n, Q_m) \right) \rightsquigarrow N(0, w_{P,Q}^2), \end{aligned} \quad (28)$$

where $w_{P,Q}^2 = \int_{\mathbb{S}^{d-1}} \text{W}_p^{2p}(P, Q) d\sigma(\theta) - \text{SW}_p^{2p}(P, Q)$. In fact, convergence in the stronger W_2 sense is also verified.

Combining Theorems 3.3 and 4.1, and Proposition 5.1, we obtain:

Corollary 5.2. *If $P, Q \in \mathcal{P}_{2p}(\mathbb{R}^d)$ verify all the assumptions of the one-sample setting in Theorems 3.3 and 4.1, and $k = k(n) \rightarrow \infty$ as $n \rightarrow \infty$, such that $k/(k+n) \rightarrow \tau \in [0, 1]$, then*

$$\sqrt{\frac{kn}{k+n}} \left(\text{SW}_{p,k}^p(P_n, Q) - \text{SW}_p^p(P, Q) \right) \rightsquigarrow N\left(0, (1-\tau)w_{P,Q}^2 + \tau v_{P,Q}^2\right).$$

Moreover, if the assumptions of the two sample-setting are also verified and $k = k(n) \rightarrow \infty$ as $n \rightarrow \infty$ such that $k/(k + \frac{nm}{n+m}) \rightarrow \tau \in [0, 1]$, then

$$\sqrt{\frac{k \frac{nm}{n+m}}{k + \frac{nm}{n+m}}} \left(\text{SW}_{p,k}^p(P_n, Q_m) - \text{SW}_p^p(P, Q) \right) \rightsquigarrow N\left(0, (1-\tau)w_{P,Q}^2 + \tau((1-\lambda)v_{P,Q}^2 + \lambda v_{Q,P}^2)\right). \quad (29)$$

6 Variance estimation

The purpose of this section is to provide consistent estimators of the asymptotic variances $w_{P,Q}^2$, $v_{P,Q}^2$ and $v_{Q,P}^2$. The simplest problem is the estimation of $w_{P,Q}^2$. In this case, given $\Theta_1, \dots, \Theta_k$ i.i.d. from σ , a natural estimator in the one

sample setting is given by

$$\hat{w}_{P_n, Q}^2 := \frac{1}{k} \sum_{i=1}^k \mathbf{W}_p^{2p}(P_n^{\Theta_i}, Q^{\Theta_i}) - \mathbf{SW}_{p,k}^{2p}(P_n, Q).$$

In the two sample setting, we can define \hat{w}_{P_n, Q_m}^2 replacing Q by Q_m .

Proposition 6.1. *Under the assumptions of Proposition 5.1,*

$$\hat{w}_{P_n, Q}^2 \rightarrow_p w_{P, Q}^2, \quad \hat{w}_{P_n, Q_m}^2 \rightarrow_p w_{P, Q}^2.$$

The estimation of $v_{P, Q}^2$ (or $v_{Q, P}^2$) is more complex. Our approach is based on the approximation of the exact formula given in Theorem 3.3. Given a sample X_1, \dots, X_n of i.i.d. random variables drawn from P , we can replace P and ϕ^θ by their empirical counterparts. However, we also need to approximate the integral, considering a finite number of slicing directions $\Theta_1, \dots, \Theta_k$ drawn from σ . Then, our estimator is defined as

$$\hat{v}_{P_n, Q}^2 := \frac{1}{k^2} \sum_{i,j=1}^k \text{Cov}_{P_n}(\phi_n^{\Theta_i}, \phi_n^{\Theta_j}), \quad (30)$$

where ϕ_n^θ denotes any c -concave optimal transport potential from P_n^θ to Q^θ . Note that the definition depends on the particular choice of ϕ_n , which might not be unique (not even up to constants). Proposition 6.2 shows that for any choice of ϕ_n^θ , the estimator (30) is a consistent estimator of $v_{P, Q}^2$. However, (30) does not depend on the particular choice of the constant. If we replace ϕ_n^θ by ϕ_{n, X_1}^θ , defined as in (13), the value of (30) remains unchanged, which will be useful for the proof. The same principles extend to the two-sample setting. Given any c -concave optimal transport potential $\phi_{n, m}^\theta$ from P_n^θ to Q_m^θ , we can define

$$\hat{v}_{P_n, Q_m}^2 := \frac{1}{k^2} \sum_{i,j=1}^k \text{Cov}_{P_n}(\phi_{n, m}^{\Theta_i}, \phi_{n, m}^{\Theta_j}), \quad (31)$$

as an estimator of $v_{P, Q}^2$, and symmetrically \hat{v}_{Q_m, P_n}^2 as an estimator of $v_{Q, P}^2$.

Proposition 6.2. *Under the assumptions of Theorem 3.3 for the one-sample setting, if $\hat{v}_{P_n, Q}^2$ is defined as in (30), then*

$$\hat{v}_{P_n, Q}^2 \xrightarrow{P} v_{P, Q}^2.$$

Moreover, if the assumptions of the two-sample setting are also verified,

$$\hat{v}_{P_n, Q_m}^2 \xrightarrow{P} v_{P, Q}^2 \quad \text{and} \quad \hat{v}_{Q_m, P_n}^2 \xrightarrow{P} v_{Q, P}^2.$$

7 Simulations

This section demonstrates the applicability of our method for statistical inference and examines the role of $\tau = \lim k/(k + \frac{nm}{n+m}) \in [0, 1]$. We focus only on the two-sample setting because of its prominent importance in statistical applications. If the assumptions of the previous sections are verified, then

$$\hat{v}^2 = \left(1 - \frac{k}{k + \frac{nm}{n+m}}\right) \hat{w}_{P_n, Q_m}^2 + \frac{k}{k + \frac{nm}{n+m}} \left(\left(1 - \frac{n}{n+m}\right) \hat{v}_{P_n, Q_m}^2 + \frac{n}{n+m} \hat{v}_{Q_m, P_n}^2 \right)$$

is a consistent estimator of the asymptotic variance in (29). Then, if we define

$$T_{n, m, k}^\Delta := \frac{1}{\hat{v}} \sqrt{\frac{k \frac{nm}{n+m}}{k + \frac{nm}{n+m}}} \left(\mathbf{SW}_{p,k}^p(P_n, Q_m) - \Delta \right), \quad (32)$$

it follows that $T_{n, m, k}^\Delta \rightsquigarrow N(0, 1)$ under the null hypothesis $H_0 : \mathbf{SW}_p^p(P, Q) = \Delta$. This provides an effective tool to perform inference on the value $\mathbf{SW}_p^p(P, Q)$, which is valid for any value of $\tau \in [0, 1]$. However, for inferential purposes, the large number of projections regime ($\tau = 1$), is particularly relevant. Taking a closer look at the proof of Corollary 5.2, we observe that, under the null hypothesis, $T_{n, m, k}^\Delta$ behaves in the limit as $\sqrt{\tau} \times (25) + \sqrt{1 - \tau} \times (28)$. However, under $H_a : \mathbf{SW}_p^p(P, Q) \neq \Delta$,

$$\sqrt{n}(\mathbf{SW}_p^p(P_n, Q_m) - \Delta) \quad (33)$$

does not converge, and $T_{n,m,k}^\Delta$ behaves as $\sqrt{\tau} \times (33) + \sqrt{1-\tau} \times (28)$. While (33) only converges to a Gaussian distribution under H_0 , the convergence in (28) does not depend on the value of Δ . Therefore, to achieve higher statistical power under the alternative hypothesis, more weight should be assigned to (33), i.e., τ should be close to 1. Alternatively, large values of k are required. This behavior is illustrated in Figure 1, for $p = 2$, and P, Q Gaussian distributions in dimension $d = 32$. Using Monte Carlo simulations, the distribution of $T_{n,m,k}^\Delta$ has been approximated under the null hypothesis and different alternatives. In addition, we estimate the exact rejection probability of the test that rejects when $|T_{n,m,k}^\Delta| > q_{0.975}$, where $q_{0.975}$ denotes the 0.975 quantile of a Gaussian distribution. We can conclude as expected. The statistical power of the test under H_a decreases significantly for small values of k . Additional computational details can be found in Appendix F.




Figure 1: Histograms based on 10000 independent samples of (32), for $d = 32$, $p = 2$, $n = 5000$, $m = 3000$, $P = N(0, I_d)$, $Q = Q_h = N((\sqrt{d} + h, 0, \dots, 0), I_d)$ and $\Delta = SW_2^2(P, Q_0) = 1$, for different values of k and h . In each subplot, r denotes the proportion of samples verifying $|T_{n,m,k}^\Delta| > q_{0.975}$. The vertical lines represent the values $\pm q_{0.975}$.

Funding

The first and second authors were supported by MCIN/AEI/10.13039/501100011033 /FEDER under Grant Agreement Number PID2021-128314NB-I00. The third author was supported by the Agence Nationale de la Recherche under grant ANR-23-CE23-0029 Regul-IA. He also acknowledges the support from the AI Cluster ANITI (ANR-19-PI3A-0004).

References

- [BF81] Peter J. Bickel and David A. Freedman. Some asymptotic theory for the bootstrap. *The Annals of Statistics*, 9(6):1196 – 1217, 1981.
- [BL19] Sergey G. Bobkov and Michel Ledoux. One-dimensional empirical measures, order statistics, and Kantorovich transport distances. *Memoirs of the American Mathematical Society*, 261(1259), 2019.
- [BRPP15] Nicolas Bonneel, Julien Rabin, Gabriel Peyré, and Hanspeter Pfister. Sliced and Radon Wasserstein barycenters of measures. *Journal of Mathematical Imaging and Vision*, 51(1):22–45, 2015.
- [CCHM86] Miklos Csörgo, Sandor Csorgo, Lajos Horvath, and David M. Mason. Weighted empirical and quantile processes. *The Annals of Probability*, 14(1):31 – 85, 1986.
- [CH90] Miklós Csörgö and Lajos Horváth. On the distributions of L_p norms of weighted quantile processes. *Annales de l’I.H.P. Probabilités et statistiques*, 26(1):65–85, 1990.
- [Cut13] Marco Cuturi. Sinkhorn Distances: Lightspeed computation of optimal transport. In *Advances in Neural Information Processing Systems*, volume 26. Curran Associates, Inc., 2013.
- [dBGL19] Eustasio del Barrio, Paula Gordaliza, and Jean-Michel Loubes. A central limit theorem for L_p transportation cost on the real line with application to fairness assessment in machine learning. *Information and Inference: A Journal of the IMA*, 8(4):817–849, 2019.
- [dBGM99] Eustasio del Barrio, Evarist Giné, and Carlos Matrán. Central limit theorems for the Wasserstein distance between the empirical and the true distributions. *The Annals of Probability*, 27(2):1009 – 1071, 1999.

- [dBGSL24] Eustasio del Barrio, Alberto González-Sanz, and Jean-Michel Loubes. Central limit theorems for general transportation costs. *Annales de l'Institut Henri Poincaré, Probabilités et Statistiques*, 60(2):847 – 873, 2024.
- [dBGU05] Eustasio del Barrio, Evarist Giné, and Frederic Utzet. Asymptotics for L2 functionals of the empirical quantile process, with applications to tests of fit based on weighted Wasserstein distances. *Bernoulli*, 11(1):131 – 189, 2005.
- [dBL19] Eustasio del Barrio and Jean-Michel Loubes. Central limit theorems for empirical transportation cost in general dimension. *The Annals of Probability*, 47(2):926 – 951, 2019.
- [dBSLNW23] Eustasio del Barrio, Alberto González Sanz, Jean-Michel Loubes, and Jonathan Niles-Weed. An improved central limit theorem and fast convergence rates for entropic transportation costs. *SIAM Journal on Mathematics of Data Science*, 5(3):639–669, 2023.
- [FG15] Nicolas Fournier and Arnaud Guillin. On the rate of convergence in Wasserstein distance of the empirical measure. *Probability Theory and Related Fields*, 162:707–738, 2015.
- [GKRS24] Ziv Goldfeld, Kengo Kato, Gabriel Rioux, and Ritwik Sadhu. Statistical inference with regularized optimal transport. *Information and Inference: A Journal of the IMA*, 13(1):iaad056, 2024.
- [GM96] Wilfrid Gangbo and Robert J. McCann. The geometry of optimal transportation. *Acta Mathematica*, 177(2):113 – 161, 1996.
- [HKMS24] Shayan Hundrieser, Marcel Klatt, Axel Munk, and Thomas Staudt. A unifying approach to distributional limits for empirical optimal transport. *Bernoulli*, 30(4):2846 – 2877, 2024.
- [HMWM24] Shayan Hundrieser, Gilles Mordant, Christoph A. Weitkamp, and Axel Munk. Empirical optimal transport under estimated costs: Distributional limits and statistical applications. *Stochastic Processes and their Applications*, 178:104462, 2024.
- [KB13] Vladimir S Korolyuk and Yu V Borovskich. *Theory of U-statistics*, volume 273. Springer Science & Business Media, 2013.
- [MBW22] Tudor Manole, Sivaraman Balakrishnan, and Larry Wasserman. Minimax confidence intervals for the Sliced Wasserstein distance. *Electronic Journal of Statistics*, 16(1):2252 – 2345, 2022.
- [MNW19] Gonzalo Mena and Jonathan Niles-Weed. Statistical bounds for entropic optimal transport: sample complexity and the central limit theorem. In *Advances in Neural Information Processing Systems*, volume 32. Curran Associates, Inc., 2019.
- [MNW24] Tudor Manole and Jonathan Niles-Weed. Sharp convergence rates for empirical optimal transport with smooth costs. *The Annals of Applied Probability*, 34(1B):1108 – 1135, 2024.
- [NWR22] Jonathan Niles-Weed and Philippe Rigollet. Estimation of Wasserstein distances in the Spiked Transport Model. *Bernoulli*, 28(4):2663 – 2688, 2022.
- [Sen77] Pranab Kumar Sen. Almost sure convergence of generalized U -statistics. *The Annals of Probability*, 5(2):287 – 290, 1977.
- [SM17] Max Sommerfeld and Axel Munk. Inference for empirical Wasserstein distances on finite spaces. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 80(1):219–238, 2017.
- [SW09] Galen R. Shorack and Jon A. Wellner. *Empirical processes with applications to statistics*. Classics in Applied Mathematics. Society for Industrial and Applied Mathematics, 2009.
- [TSM19] Carla Taming, Max Sommerfeld, and Axel Munk. Empirical optimal transport on countable metric spaces: Distributional limits and statistical applications. *The Annals of Applied Probability*, 29(5):2744 – 2781, 2019.
- [Vil03] Cédric Villani. *Topics in optimal transportation*. Graduate studies in mathematics. American Mathematical Society, Providence, R.I, 2003.
- [WB19] Jonathan Weed and Francis Bach. Sharp asymptotic and finite-sample rates of convergence of empirical measures in Wasserstein distance. *Bernoulli*, 25(4 A):2620 – 2648, 2019.
- [XH22] Xianliang Xu and Zhongyi Huang. Central limit theorem for the sliced 1-Wasserstein distance and the max-sliced 1-Wasserstein distance. *arXiv preprint*, 2022.
- [XNW22] Jiaqi Xi and Jonathan Niles-Weed. Distributional convergence of the sliced Wasserstein process. In *Advances in Neural Information Processing Systems*, volume 35, pages 13961–13973. Curran Associates, Inc., 2022.

Appendix

The results presented in this article apply to both the one-sample and two-sample settings, with the proofs being largely similar in most cases. The following sections of the appendix provide detailed proofs for the one-sample setting and highlight the key modifications required to extend these arguments to the two-sample setting.

A Proofs of Section 2

A.1 Preliminary results

Lemma A.1. *Let $P \in \mathcal{P}(\mathbb{R}^d)$ be a probability with negligible boundary and $\text{int}(\text{supp}(P))$ connected. Then, for every $\theta \in \mathbb{S}^{d-1}$, P^θ has negligible boundary, and $\text{int}(\text{supp}(P^\theta))$ is connected.*

Proof. First of all, we prove the set inclusions $\text{Pr}_\theta(\text{int}(\text{supp}(P))) \stackrel{(i)}{\subset} \text{int}(\text{supp}(P^\theta)) \stackrel{(ii)}{\subset} \text{cl}(\text{Pr}_\theta(\text{int}(\text{supp}(P))))$.

- (i) Let $s \in \text{Pr}_\theta(\text{int}(\text{supp}(P)))$. There exist $x \in \text{int}(\text{supp}(P))$ and $\epsilon > 0$ such that $s = \langle \theta, x \rangle$ and $B(x, \epsilon) \subset \text{supp}(P)$. Then, for every $s' = s + t \in B(s, \epsilon)$, $x' = x + \theta t \in B(x, \epsilon)$. Therefore, $x' \in \text{supp}(P)$. Thus, for every $r > 0$,

$$P^\theta(B(s', r)) = P(\{x \in \mathbb{R}^d : \langle \theta, x \rangle \in B(s, r)\}) \geq P(B(x', r)) > 0.$$

Then, $s' \in \text{supp}(P^\theta)$, and it follows that $B(s, \epsilon) \subset \text{supp}(P^\theta)$, which implies that $s \in \text{int}(\text{supp}(P^\theta))$.

- (ii) Let $s \in \text{int}(\text{supp}(P^\theta))$. For each $n \in \mathbb{N}$, since P has negligible boundary,

$$\begin{aligned} P^\theta(B(s, 1/n)) &= P(\{x \in \mathbb{R}^d : \langle \theta, x \rangle \in B(s, 1/n)\}) \\ &= P(\{x \in \text{int}(\text{supp}(P)) : \langle \theta, x \rangle \in B(s, 1/n)\}) > 0 \end{aligned}$$

Then, we can define a sequence of values $x_n \in \text{int}(\text{supp}(P))$ such that $\langle \theta, x_n \rangle \rightarrow s$. Therefore, s belongs to the closure of $\text{Pr}_\theta(\text{int}(\text{supp}(P)))$.

Now, since $\text{int}(\text{supp}(P))$ is connected and Pr_θ is open and continuous, it follows that $\text{Pr}_\theta(\text{int}(\text{supp}(P)))$ is an open interval I^θ , which together with the previous inclusions, allow us to conclude

$$\text{int}(\text{supp}(P^\theta)) = \text{Pr}_\theta(\text{int}(\text{supp}(P))) = I^\theta. \quad (34)$$

Therefore, $\text{int}(\text{supp}(P^\theta))$ is connected. Since P has negligible boundary, $P^\theta(\text{int}(\text{supp}(P^\theta))) = P(\text{int}(\text{supp}(P))) = 1$, which implies that $\text{supp}(P^\theta) = \text{cl}(I^\theta)$. Consequently, P^θ also has negligible boundary. \square

The following lemma summarizes key regularity properties of c -concave optimal transport potentials used in the proof of Proposition 2.1. Assertions (1), (2), and (3) are just an adaptation of Theorem 3.3 and Proposition C.4 in [GM96] for $d = 1$, but these properties hold more generally.

Lemma A.2. *Let $P, Q \in \mathcal{P}_p(\mathbb{R})$, $p > 1$, and $c = c_p$. Denote by ϕ any c -concave optimal transport potential from P to Q . Then,*

1. *There exists a closed interval I such that $\text{int}(I) \subset \text{dom } \phi = \{x \in \mathbb{R} : \phi(x) > -\infty\} \subset I$.*
2. *ϕ is locally Lipschitz in $\text{int}(I)$, and, therefore, continuous.*
3. *For every $s \in \text{int}(I)$, $\partial^c \phi(s) \neq \emptyset$.*
4. *Given $s \in \mathbb{R}$, if $\partial^c \phi(s) \neq \emptyset$, then $\partial^c \phi(s)$ is a single point or a closed interval.*

Proof. The only assertion that requires to be proved is (4). Assume that $\partial^c \phi(s)$ does not consist of a single point. To prove that $\partial^c \phi(s)$ is an interval, we need to see that if $t_1, t_2 \in \partial^c \phi(s)$, and $t \in [t_1, t_2]$, then $t \in \partial^c \phi(s)$. This results from the definition (10) and the inequality

$$\min \left\{ |z - t_1|^p - |x - t_1|^p, |z - t_2|^p - |x - t_2|^p \right\} \leq |z - t|^p - |x - t|^p.$$

which follows from the fact that $t \mapsto |z - t|^p - |x - t|^p$ is monotone increasing if $z \leq x$ and decreasing if $z \geq x$. Thus, $\partial^c \phi(s)$ is an interval. To see that it is closed, let $t \in \text{cl}(\partial^c \phi(s))$, and consider a sequence $\{t_n\}_{n=1}^\infty \subset \partial^c \phi(s)$, such that $t_n \rightarrow t$. Then, for every $z \in \mathbb{R}$, $\phi(z) \leq \phi(s) + (|z - t_n|^p - |x - t_n|^p)$. By continuity, we conclude that $t \in \partial^c \phi(x)$. \square

With the notation of the previous lemma, it follows that $\text{supp}(P) \subset I$. Otherwise, if $s \in \text{supp}(P) \setminus I$, by definition of $\text{supp}(P)$, for every $\epsilon > 0$, $P(B(s, \epsilon)) > 0$. Since I is closed, $\mathbb{R} \setminus I$ is open, and there exists $\epsilon > 0$ such that $B(s, \epsilon) \subset \mathbb{R} \setminus I \subset \mathbb{R} \setminus \text{dom}(\phi)$. This implies that $\int \phi(s) dP(s)$ is not finite, which contradicts the optimality of ϕ .

Therefore, $\text{int}(\text{supp}(P)) \subset \text{int}(I)$. If P is absolutely continuous, $\text{int}(\text{supp}(P))$ defines a set of probability one, and Lemma A.2 allows us to guarantee favorable properties on this set. Now, if P assigns positive probability to $s \in \mathbb{R}$, by the optimality, $\phi(s) \in \text{dom}(f)$. In this case, even if $s \in I \setminus \text{int}(I)$, we can ensure $\partial^c \phi(s) \neq \emptyset$. To check this, let $\pi \in \Pi(P, Q)$ be any optimal transport plan such that $\text{supp}(\pi) \subset \partial^c \phi$. Since $P(x) > 0$, we know that $\pi(\{x\} \times \mathbb{R}) > 0$. Therefore, there exists $A \subset \mathbb{R}$ such that $\{x\} \times A \subset \text{supp}(\pi) \subset \partial^c \phi$, which implies $A \subset \partial^c \phi(x)$.

A.2 Proof of Proposition 2.1

We restrict our attention to the subset of probability one $\Omega_0 \subset \Omega$ verifying that for every $\omega \in \Omega_0$,

- $X_1^\omega \in \text{int}(\text{supp}(P))$
- $P_n^\omega \rightsquigarrow P$

Therefore, it suffices to demonstrate all the assertions for a sequence $\{X_n^\omega\}_{n=1}^\infty$, with $\omega \in \Omega_0$. For the sake of simplicity, we omit the superscript ω . Note that the structure of the proof closely follows that of Lemma B.2 in [GKRS24].

(1) For each $\theta \in \mathbb{S}^{d-1}$, by Lemma A.1, P^θ has density with negligible boundary and $\text{int}(\text{supp}(P^\theta))$ connected. By Corollary 2.7 in [dBGSL24], this guarantees uniqueness of the optimal potentials in $\text{int}(\text{supp}(P))$ up to the addition of a constant. By the comments following Lemma A.2, ϕ^θ is finite in $\text{int}(\text{supp}(P^\theta))$. Since $X_1 \in \text{int}(\text{supp}(P))$, by (34), $\langle \theta, X_1 \rangle \in \text{int}(\text{supp}(P^\theta))$. Therefore, $\phi^\theta(\langle \theta, X_1 \rangle) \in \mathbb{R}$ and $\phi_{X_1}^\theta(s) = \phi^\theta(s) - \phi^\theta(\langle \theta, X_1 \rangle)$ is well defined, it is a c -concave optimal transport potential and verifies $\phi_{X_1}^\theta(\langle \theta, X_1 \rangle) = 0$.

(2) Let $x \in \text{int}(\text{supp}(P))$, and $\theta_n \rightarrow \theta$. First, it is easy to see that $P^{\theta_n} \rightsquigarrow P^\theta$. The first part of Theorem 3.4 in [dBGSL24] implies that there exist constants a_n^θ (which depend on $\omega \in \Omega_0$) such that $\phi_{X_1}^{\theta_n} - a_n^\theta \rightarrow \phi_{X_1}^\theta$ in the sense of uniform convergence in the compacts of $\text{int}(\text{supp}(P^\theta))$. Since $\langle \theta, X_1 \rangle \in \text{int}(\text{supp}(P^\theta))$, and $\phi_{X_1}^{\theta_n}(\langle \theta, X_1 \rangle) = \phi_{X_1}^\theta(\langle \theta, X_1 \rangle) = 0$, then $a_n^\theta \rightarrow 0$, and we can conclude that

$$|\phi_{X_1}^{\theta_n} - \phi_{X_1}^\theta| \leq |(\phi_{X_1}^{\theta_n} - a_n^\theta) - \phi_{X_1}^\theta| + |a_n^\theta| \rightarrow 0$$

uniformly in the compact sets of $\text{int}(\text{supp}(P^\theta))$. Let K be a compact neighborhood of $\langle \theta, x \rangle$. For n large enough, $\langle \theta_n, x \rangle \in K$. By Lemma A.2, ϕ^θ is continuous in $\text{int}(\text{supp}(P^\theta))$. Therefore, we can conclude that

$$\begin{aligned} & |\phi_{X_1}^{\theta_n}(\langle \theta_n, x \rangle) - \phi_{X_1}^\theta(\langle \theta, x \rangle)| \\ & \leq |\phi_{X_1}^{\theta_n}(\langle \theta_n, x \rangle) - \phi_{X_1}^\theta(\langle \theta_n, x \rangle)| + |\phi_{X_1}^\theta(\langle \theta_n, x \rangle) - \phi_{X_1}^\theta(\langle \theta, x \rangle)| \rightarrow 0 \end{aligned}$$

(3) By the discussion after Lemma A.2, $\phi_n^\theta(\langle \theta, X_1 \rangle) < \infty$. Therefore, $\phi_{n, X_1}^\theta(s) = \phi_n^\theta(s) - \phi_n^\theta(\langle \theta, X_1 \rangle)$ is well defined, it is a c -concave optimal transport potential and verifies $\phi_{X_1}^\theta(\langle \theta, X_1 \rangle) = 0$.

(4) For each $\theta \in \mathbb{S}^{d-1}$, the continuous mapping theorem implies $P_n^\theta \rightsquigarrow P^\theta$. Reasoning as in the proof of (2), we can ensure that $\phi_{n, X_1}^\theta(t) \rightarrow \phi_{X_1}^\theta(t)$ uniformly on the compact subsets of $\text{int}(\text{supp}(P^\theta))$.

(5) This follows from $P_n^\theta \rightsquigarrow P^\theta$ and the second part of Theorem 3.4 in [dBGSL24].

(6) Assume that the property is not verified. Then, there exists a subsequence $\{n_k\}_{k=1}^\infty$ and values $\{\theta_{n_k}\}_{k=1}^\infty$ such that $|\phi_{n_k, X_1}^{\theta_{n_k}}(\langle \theta_{n_k}, x \rangle)| \rightarrow \infty$. Since \mathbb{S}^{d-1} is compact, we can extract a further subsequence $\theta_{n_{k_m}} \rightarrow \theta \in \mathbb{S}^{d-1}$. For

this subsequence, $P_{n_{k_m}}^{\theta} \rightsquigarrow P^\theta$ and $Q^{\theta_{n_{k_m}}} \rightsquigarrow Q^\theta$. By the same reasoning as in the proof of (2), we conclude that $\phi_{n_{k_m}, X_1}^{\theta_{n_{k_m}}}(t)$ converges to $\phi_{X_1}^\theta(t)$ uniformly on the compact sets of $\text{int}(\text{supp}(P^\theta))$. Thus,

$$\phi_{n_{k_m}, X_1}^{\theta_{n_{k_m}}}(\langle \theta_{n_{k_m}}, x \rangle) \longrightarrow \phi_{X_1}^\theta(\langle \theta, x \rangle) < \infty$$

which contradicts the hypothesis.

Two-sample setting. For each $m \in \mathbb{N}$, the existence of ϕ_{n, m, X_1} follows from Claim 2 with $Q = Q_m$. Theorem 3.4 in [dBGSL24] also applies to the two-sample setting. Therefore, the same proof of (4), (5) and (6) extends to this case in the set of \mathbb{P} -probability one $\Omega_0 \cap \{\omega : Q_m \rightsquigarrow Q\}$. \square

A.3 Proof of Proposition 2.2

First, we prove (16). By Proposition 2.1, $\phi_{X_1}^\theta$ and ϕ_{n, X_1}^θ are well-defined c -concave optimal potentials with reference point X_1 , up to a set of \mathbb{P} -probability zero. To prove integrability properties, we can assume without loss of generality that it is verified for every point. Therefore, since ϕ_{n, X_1}^θ is c -concave,

$$|s - t|^p = \phi_{n, X_1}^\theta(s) + (\phi_{n, X_1}^\theta)^c(t) \iff t \in \partial^c \phi_{n, X_1}^\theta(s) \iff s \in \partial^c (\phi_{n, X_1}^\theta)^c(t) \quad (35)$$

Let t_{n, X_1}^θ be any element of $\partial^c \phi_{n, X_1}^\theta(\langle \theta, X_1 \rangle)$, which is non-empty by the comments after Lemma A.2. The first part of the argument is valid for any choice of $t_{n, X_1}^\theta \in \partial^c \phi_{n, X_1}^\theta(\langle \theta, X_1 \rangle)$, although to conclude, we will consider a particular choice in (41). By definition of the c -superdifferential, for every $s \in \mathbb{R}$,

$$\phi_{n, X_1}^\theta(s) \leq \phi_{n, X_1}^\theta(\langle \theta, X_1 \rangle) + \left(|s - t_{n, X_1}^\theta|^p - |\langle \theta, X_1 \rangle - t_{n, X_1}^\theta|^p \right). \quad (36)$$

Since $t_{n, X_1}^\theta \in \partial^c \phi_{n, X_1}^\theta(\langle \theta, X_1 \rangle)$, it follows by (35) that $\langle \theta, X_1 \rangle \in \partial^c (\phi_{n, X_1}^\theta)^c(t_{n, X_1}^\theta)$. Therefore, for every $t \in \mathbb{R}$,

$$(\phi_{n, X_1}^\theta)^c(t) \leq (\phi_{n, X_1}^\theta)^c(t_{n, X_1}^\theta) + \left(|t - \langle \theta, X_1 \rangle|^p - |t_{n, X_1}^\theta - \langle \theta, X_1 \rangle|^p \right). \quad (37)$$

By (35) and (37), it follows that for every $(s, t) \in \partial^c \phi_{n, X_1}^\theta$,

$$\phi_{n, X_1}^\theta(s) = |s - t|^p - (\phi_{n, X_1}^\theta)^c(t) \geq \phi_{n, X_1}^\theta(\langle \theta, X_1 \rangle) + |s - t|^p - |t - \langle \theta, X_1 \rangle|^p \quad (38)$$

For every $(s, t) \in \partial^c \phi_{n, X_1}^\theta$, combining (36) and (38) and using inequality (57),

$$|\phi_{n, X_1}^\theta(s) - \phi_{n, X_1}^\theta(\langle \theta, X_1 \rangle)| \leq 2C_p \left(|s|^p + |t|^p + |\langle \theta, X_1 \rangle|^p + |t_{n, X_1}^\theta|^p \right). \quad (39)$$

Similarly, for every $(s, t) \in \partial^c \phi_{n, X_1}^\theta$,

$$|(\phi_{n, X_1}^\theta)^c(t) - (\phi_{n, X_1}^\theta)^c(t_{n, X_1}^\theta)| \leq 2C_p \left(|s|^p + |t|^p + |\langle \theta, X_1 \rangle|^p + |t_{n, X_1}^\theta|^p \right) \quad (40)$$

But optimality implies that there exists an optimal coupling π_n^θ between P_n^θ and Q^θ such that $\text{supp}(\pi_n^\theta) \subset \partial^c \phi_{n, X_1}^\theta$. Therefore, (39) is verified for every $(s, t) \in \text{supp}(\pi_n^\theta)$. Given that $\phi_{n, X_1}^\theta(\langle \theta, X_1 \rangle) = 0$ and $\phi_{n, X_1}^\theta(\langle \theta, X_1 \rangle) + (\phi_{n, X_1}^\theta)^c(t_{n, X_1}^\theta) = |\langle \theta, X_1 \rangle - t_{n, X_1}^\theta|^p$ it follows that both integrals

$$\begin{aligned} \int_{\mathbb{R}^d} |\phi_{n, X_1}^\theta(\langle \theta, x \rangle)|^q dP_n(x) &= \int_{\mathbb{R} \times \mathbb{R}} |\phi_{n, X_1}^\theta(s)|^q d\pi_n^\theta(s, t) \\ \int_{\mathbb{R}^d} |(\phi_{n, X_1}^\theta)^c(\langle \theta, y \rangle)|^q dQ(y) &= \int_{\mathbb{R} \times \mathbb{R}} |(\phi_{n, X_1}^\theta)^c(t)|^q d\pi_n^\theta(s, t) \end{aligned}$$

are bounded, up to a constant, by

$$\int_{\mathbb{R}} |s|^{pq} dP_n^\theta(s) + \int_{\mathbb{R}} |t|^{pq} dQ^\theta(t) + |\langle \theta, X_1 \rangle|^{pq} + |t_{n, X_1}^\theta|^{pq}.$$

Therefore, we only need to uniformly bound the expected value of the integral in \mathbb{S}^{d-1} of each term.

- $\mathbb{E}\left(\int_{\mathbb{S}^{d-1}} \int_{\mathbb{R}} |s|^{pq} dP_n^\theta(s) d\sigma(\theta)\right) \leq \int_{\mathbb{R}} \|x\|^{pq} dP(x)$.
- $\mathbb{E}\left(\int_{\mathbb{S}^{d-1}} \int_{\mathbb{R}} |t|^{pq} dQ^\theta(t) d\sigma(\theta)\right) \leq \int_{\mathbb{R}} \|y\|^{pq} dQ(y)$.
- $\mathbb{E}\left(\int_{\mathbb{S}^{d-1}} |\langle \theta, X_1 \rangle|^{pq} d\sigma(\theta)\right) \leq \int_{\mathbb{R}} \|x\|^{pq} dP(x)$.
- For the last term, we need to consider a particular choice of the elements t_{n, X_1}^θ . Note that the c -superdifferential does not depend on the particular choice of the constant. Thus, $\partial^c \phi_{n, X_1}^\theta(\langle \theta, x \rangle) = \partial^c \phi_n^\theta(\langle \theta, x \rangle)$ and we define for each $i = 1, \dots, n$ and $\theta \in \mathbb{S}^{d-1}$ the random variable

$$t_{n, X_i}^\theta = \operatorname{argmin}_{t \in \partial^c \phi_n^\theta(\langle \theta, X_i \rangle)} |t|, \quad (41)$$

which is well defined since the c -superdifferential of a convex function is a closed (possibly infinite) interval, by Lemma A.2. By definition,

$$t_{n, X_1}^\theta \stackrel{d}{=} \dots \stackrel{d}{=} t_{n, X_n}^\theta \quad (42)$$

Then, given an optimal transport plan π_n^θ verifying $\operatorname{supp}(\pi_n^\theta) \subset \partial^c \phi_n^\theta$,

$$\begin{aligned} \int_{\mathbb{R}^d} \|y\|^{pq} dQ(y) &\geq \int_{\mathbb{R}^d} |\langle \theta, y \rangle|^{pq} dQ(y) = \int_{\mathbb{R} \times \mathbb{R}} |t|^{pq} d\pi_n^\theta(s, t) \\ &= \sum_{i=1}^n \int_{\{s=\langle \theta, X_i \rangle, t \in \partial^c \phi_n^\theta(s)\}} |t|^{pq} d\pi_n^\theta(s, t) \\ &\geq \sum_{i=1}^n \int_{\{s=\langle \theta, X_i \rangle, t \in \partial^c \phi_n^\theta(s)\}} |t_{n, X_i}^\theta|^{pq} d\pi_n^\theta(s, t) \\ &= \frac{1}{n} \sum_{i=1}^n |t_{n, X_i}^\theta|^{pq} \end{aligned} \quad (43)$$

This inequality and (42) allow us to conclude

$$\mathbb{E}\left(|t_{n, X_1}^\theta|^{pq}\right) = \mathbb{E}\left(\frac{1}{n} \sum_{i=1}^n |t_{n, X_i}^\theta|^{pq}\right) \leq \int_{\mathbb{R}^d} \|y\|^{pq} dQ(y).$$

The proof of (15) follows from the same ideas. For each $\theta \in \mathbb{S}^{d-1}$, by Theorem 1.2 in [GM96], $\partial^c \phi_\theta(s) = \{\nabla^c \phi_\theta(s)\}$ P^θ -a.s. and $\nabla^c \phi_\theta$ is an optimal transport map between P^θ and Q^θ . Therefore, $t_{X_1}^\theta := \nabla^c \phi_\theta(\langle \theta, X_1 \rangle)$ is well defined ($\mathbb{P} \times \sigma$)-a.s. and

$$\begin{aligned} \mathbb{E}\left(\int_{\mathbb{S}^{d-1}} |t_{X_1}^\theta|^{pq} d\sigma(\theta)\right) &= \int_{\mathbb{S}^{d-1}} \left(\int_{\mathbb{R}} |\nabla^c \phi_\theta(s)|^{pq} dP^\theta(s)\right) d\sigma(\theta) \\ &= \int_{\mathbb{S}^{d-1}} \left(\int_{\mathbb{R}} |t|^{pq} dQ^\theta(s)\right) d\sigma(\theta) \leq \int_{\mathbb{R}^d} \|y\|^{pq} dQ(y) \end{aligned}$$

Two-sample setting: The proof of (17) follows from the same reasoning, taking into account that $\mathbb{E}\left(\int_{\mathbb{R}} \|y\|^{pq} dQ_m(y)\right) = \int_{\mathbb{R}} \|y\|^{pq} dQ(y)$. \square

B Proofs of Section 3

B.1 Proof of Proposition 3.1

Define the random variable $Z_n = \operatorname{SW}_p^p(P_n, Q)$, where $P_n = \frac{1}{n} \sum_{i=1}^n \delta_{X_i}$, with X_1, \dots, X_n i.i.d. random variables with distribution P . Consider now $Z'_n = \operatorname{SW}_p^p(P'_n, Q)$, where $P'_n = \frac{1}{n} (\delta_{X'_1} + \sum_{i=2}^n \delta_{X_i})$, with X'_1 drawn from P , independent of X_1, \dots, X_n . To simplify the notation, let $X'_j = X_j$ for $j \geq 2$.

For each $\theta \in \mathbb{S}^{d-1}$, consider the following construction. First, define the random map $\tau'_\theta : \{1, \dots, n\} \rightarrow \{1, \dots, n\}$ that maps each index i to the position of $\langle \theta, X'_i \rangle$ in the ordered statistic associated with the variables $\{\langle \theta, X'_i \rangle\}_{i=1}^n$. If there are k equal values in the ordered statistic, then τ' is randomly selected as one of the $k!$ possible permutations. With this construction, $\tau'_\theta(1), \dots, \tau'_\theta(n)$ are identically distributed, and given that X_1 is independent of X'_1, \dots, X'_n ,

$$(X_1, \tau'_\theta(1)) \stackrel{d}{=} \dots \stackrel{d}{=} (X_1, \tau'_\theta(n)). \quad (44)$$

Denote by $F_{\theta,n}, F'_{\theta,n}, G_\theta$ the distribution functions of $P_n^\theta, P_n^{\theta'}$ and Q^θ , respectively. Given a random variable U with uniform distribution in $(0, 1)$, we define for each $j = 1, \dots, n$ the positive measure Q_j^θ on \mathbb{R} by

$$Q_j^\theta(A) = \mathbb{P}\left(G_\theta^{-1}(U) \in A, U \in \left(\frac{j-1}{n}, \frac{j}{n}\right)\right).$$

Note that $Q^\theta = \sum_{j=1}^n Q_j^\theta$, and $Q_j^\theta(\mathbb{R}) = 1/n$. Moreover, since we are working with one-dimensional distributions, $\pi_n^{\theta'} = \mathcal{L}(F'_{\theta,n}^{-1}(U), G_\theta^{-1}(U))$ is an optimal coupling between $P_n^{\theta'}$ and Q^θ . It can be decomposed as $\pi_n^{\theta'} = \sum_{i=1}^n \delta_{\langle \theta, X'_i \rangle} \times Q_{\tau'_\theta(i)}^\theta$. Therefore,

$$\mathbb{W}_p^p(P_n^{\theta'}, Q^\theta) = \sum_{i=1}^n \int_{\mathbb{R}} |\langle \theta, X'_i \rangle - t|^p dQ_{\tau'_\theta(i)}^\theta(t).$$

Now, we can define $\pi_n^\theta = \sum_{i=1}^n \delta_{\langle \theta, X_i \rangle} \times Q_{\tau'_\theta(i)}^\theta$. It is easy to see that π_n^θ is a valid transport plan from P_n^θ to Q^θ . Thus,

$$\mathbb{W}_p^p(P_n^\theta, Q^\theta) \leq \sum_{i=1}^n \int_{\mathbb{R}} |\langle \theta, X_i \rangle - t|^p dQ_{\tau'_\theta(i)}^\theta(t).$$

Therefore, since $X_i = X'_i$ for $i \neq 1$, by convexity of $x \mapsto |\langle \theta, x \rangle - t|^p$,

$$\begin{aligned} \mathbb{W}_p^p(P_n^\theta, Q^\theta) - \mathbb{W}_p^p(P_n^{\theta'}, Q^\theta) &\leq \int_{\mathbb{R}} |\langle \theta, X_1 \rangle - t|^p - |\langle \theta, X'_1 \rangle - t|^p dQ_{\tau'_\theta(1)}^\theta(t) \\ &\leq p |\langle \theta, X_1 - X'_1 \rangle| \int_{\mathbb{R}} |\langle \theta, X_1 \rangle - t|^{p-1} dQ_{\tau'_\theta(1)}^\theta(t) \\ &\leq p \|X_1 - X'_1\| \int_{\mathbb{R}} |\langle \theta, X_1 \rangle - t|^{p-1} dQ_{\tau'_\theta(1)}^\theta(t). \end{aligned}$$

Since this reasoning is valid for every $\theta \in \mathbb{S}^{d-1}$, we can ensure that

$$Z_n - Z'_n \leq p \|X_1 - X'_1\| \int_{\mathbb{S}^{d-1}} \int_{\mathbb{R}} |\langle \theta, X_1 \rangle - t|^{p-1} dQ_{\tau'_\theta(1)}^\theta(y) d\sigma(\theta). \quad (45)$$

Applying Hölder's inequality, with $f = 1, g = |\langle \theta, x \rangle - t|^{p-1}, q_1 = \frac{2p}{p+1}, q_2 = \frac{2p}{p-1}$,

$$\begin{aligned} \int_{\mathbb{R}} |\langle \theta, X_1 \rangle - t|^{p-1} dQ_{\tau'_\theta(1)}^\theta(t) &\leq \left(\int_{\mathbb{R}} dQ_{\tau'_\theta(1)}^\theta(t) \right)^{\frac{p+1}{2p}} \left(\int_{\mathbb{R}} |\langle \theta, X_1 \rangle - t|^{2p} dQ_{\tau'_\theta(1)}^\theta(t) \right)^{\frac{p-1}{2p}} \\ &= \frac{1}{n^{\frac{p+1}{2p}}} \left(\int_{\mathbb{R}} |\langle \theta, X_1 \rangle - t|^{2p} dQ_{\tau'_\theta(1)}^\theta(t) \right)^{\frac{p-1}{2p}} \end{aligned} \quad (46)$$

By exchangeability (44), we know that for every $\theta \in \mathbb{S}^{d-1}$,

$$\int_{\mathbb{R}} |\langle \theta, X_1 \rangle - t|^{2p} dQ_{\tau'_\theta(1)}^\theta(t) \stackrel{d}{=} \dots \stackrel{d}{=} \int_{\mathbb{R}} |\langle \theta, X_1 \rangle - t|^{2p} dQ_{\tau'_\theta(n)}^\theta(t)$$

and therefore

$$\begin{aligned} \mathbb{E} \left(\int_{\mathbb{R}} |\langle \theta, X_1 \rangle - t|^{2p} dQ_{\tau'_\theta(1)}^\theta(t) \right) &= \frac{1}{n} \mathbb{E} \left(\sum_{i=1}^n \int_{\mathbb{R}} |\langle \theta, X_1 \rangle - t|^{2p} dQ_{\tau'_\theta(i)}^\theta(t) \right) \\ &= \frac{1}{n} \int_{\mathbb{R}^d} |\langle \theta, X_1 \rangle - t|^{2p} dQ^\theta(t). \end{aligned} \quad (47)$$

From (45) and a further application of Hölder's inequality with $q_1 = p$, $q_2 = \frac{p}{p-1}$,

$$\begin{aligned} \mathbb{E}\left((Z_n - Z'_n)_+^2\right) &\leq p^2 \mathbb{E}\left(\|X_1 - X'_1\|^2 \left(\int_{\mathbb{S}^{d-1}} \int_{\mathbb{R}} |\langle \theta, X_1 \rangle - t|^{p-1} dQ_{\tau'_\theta(1)}(y) d\sigma(\theta)\right)^2\right) \\ &\leq p^2 \mathbb{E}\left(\|X_1 - X'_1\|^{2p}\right)^{\frac{1}{p}} \mathbb{E}\left(\left(\int_{\mathbb{S}^{d-1}} \int_{\mathbb{R}} |\langle \theta, X_1 \rangle - t|^{p-1} dQ_{\tau'_\theta(1)}(y) d\sigma(\theta)\right)^{\frac{2p}{p-1}}\right)^{\frac{p-1}{p}} \end{aligned} \quad (48)$$

Applying Jensen's inequality, (46) and (47), we obtain the bound

$$\begin{aligned} &\mathbb{E}\left(\left(\int_{\mathbb{S}^{d-1}} \int_{\mathbb{R}} |\langle \theta, X_1 \rangle - t|^{p-1} dQ_{\tau'_\theta(1)}(y) d\sigma(\theta)\right)^{\frac{2p}{p-1}}\right) \\ &\leq \mathbb{E}\left(\int_{\mathbb{S}^{d-1}} \left(\int_{\mathbb{R}} |\langle \theta, X_1 \rangle - t|^{p-1} dQ_{\tau'_\theta(1)}(y)\right)^{\frac{2p}{p-1}} d\sigma(\theta)\right) \\ &\leq \frac{1}{n^{\frac{p+1}{p-1}}} \mathbb{E}\left(\int_{\mathbb{S}^{d-1}} \int_{\mathbb{R}} |\langle \theta, X_1 \rangle - t|^{2p} dQ_{\tau'_\theta(1)}^\theta(t) d\sigma(\theta)\right) \\ &\leq \frac{1}{n^{\frac{2p}{p-1}}} \mathbb{E}\left(\int_{\mathbb{S}^{d-1}} \int_{\mathbb{R}} |\langle \theta, X_1 \rangle - t|^{2p} dQ^\theta(t) d\sigma(\theta)\right) \\ &\leq \frac{1}{n^{\frac{2p}{p-1}}} \mathbb{E}\left(\|X_1 - Y\|^{2p}\right) \end{aligned}$$

where Y represents a random variable independent of X , with $\mathcal{L}(Y) = Q$. Combined with (48),

$$\mathbb{E}\left((Z_n - Z'_n)_+^2\right) \leq \frac{p^2}{n^2} \mathbb{E}\left(\|X_1 - X'_1\|^{2p}\right)^{\frac{1}{p}} \mathbb{E}\left(\|X_1 - Y\|^{2p}\right)^{\frac{p-1}{p}}. \quad (49)$$

Finally, Efron-Stein inequality (19) allows us to conclude (a).

Two-sample setting. Let $Z = \text{SW}_p^p(P_n, Q_m)$, where Q_m is the empirical distribution associated with m i.i.d. random variables Y_1, \dots, Y_m drawn from Q , independent of X_1, \dots, X_n . Z is symmetric with respect to its first n variables and its last m variables. Thus, applying Efron-Stein,

$$\text{Var}(\text{SW}_p^p(P_n, Q_m)) \leq n \mathbb{E}\left((Z_{n,m} - Z'_{n,m})_+^2\right) + m \mathbb{E}\left((Z_{n,m} - Z''_{n,m})_+^2\right)$$

where $Z'_{n,m} = \text{SW}_p^p(P'_n, Q_m)$, $Z''_{n,m} = \text{SW}_p^p(P_n, Q'_m)$, P'_n is defined as in the one-sample case, and Q'_m is the empirical measure associated with Y'_1, Y'_2, \dots, Y'_m , with Y'_1 an independent copy. Denote by \mathbb{E}_X the expectation taken only with respect to X_1, \dots, X_n . Then, (49) applied to the discrete case $Q = Q_m$ ensures that,

$$\mathbb{E}_X\left((Z_{n,m} - Z'_{n,m})_+^2\right) \leq \frac{p^2}{n^2} \mathbb{E}_X\left(\|X_1 - X'_1\|^{2p}\right)^{\frac{1}{p}} \mathbb{E}_X\left(\frac{1}{m} \sum_{j=1}^m \|X_1 - Y_j\|^{2p}\right)^{\frac{p-1}{p}}.$$

Taking the expectation with respect to Y_1, \dots, Y_n and applying Jensen's inequality, it is easy to conclude $\mathbb{E}\left((Z_{n,m} - Z'_{n,m})_+^2\right) \leq \frac{1}{n^2} K_p(P, Q)$. Similarly, exchanging the roles of P and Q , $\mathbb{E}\left((Z_{n,m} - Z''_{n,m})_+^2\right) \leq \frac{1}{m^2} K_p(Q, P)$, which allows us to conclude (b). \square

B.2 Proof of Proposition 3.2

Let P_n, P'_n and Z_n, Z'_n be as in the proof of the previous theorem. Let ϕ_n^θ be any c -concave optimal transport potential between P_n^θ and Q^θ , and define ϕ_{n, X_1}^θ as in (13). In addition, define $\phi_{X_1}^\theta$ as in (12) for $\phi^\theta = \phi_{x_0}^\theta$. By Proposition 2.1, we know that there exists a set of \mathbb{P} -probability one where $\phi_{X_1}^\theta, \phi_{n, X_1}^\theta$ are well-defined c -concave optimal transport potentials verifying

$$\phi_{n, X_1}^\theta(\langle \theta, X_1 \rangle) = \phi_{X_1}^\theta(\langle \theta, X_1 \rangle) = 0 \quad \forall \theta \in \mathbb{S}^{d-1}, n \in \mathbb{N}.$$

Thus, the following equalities and inequalities should be understood in a \mathbb{P} -almost surely sense. If we define R'_n as R_n , replacing P_n by P'_n , by (20),

$$\begin{aligned} R_n - R'_n &\leq \int_{\mathbb{S}^{d-1}} \int_{\mathbb{R}} (\phi_{n, X_1}^\theta(s) - \phi_{X_1}^\theta(s)) d(P_n^\theta - P_n'^\theta)(s) d\sigma(\theta) \\ &= \frac{1}{n} \int_{\mathbb{S}^{d-1}} \left(\phi_{n, X_1}^\theta(\langle \theta, X_1 \rangle) - \phi_{X_1}^\theta(\langle \theta, X_1 \rangle) \right) d\sigma(\theta) \\ &\quad - \frac{1}{n} \int_{\mathbb{S}^{d-1}} \left(\phi_{n, X_1}^\theta(\langle \theta, X'_1 \rangle) - \phi_{X_1}^\theta(\langle \theta, X'_1 \rangle) \right) d\sigma(\theta) \end{aligned}$$

First, we will see that $n(R_n - R'_n)_+ \rightarrow 0$ \mathbb{P} -a.s. For that, we prove that the upper bound converges to 0 \mathbb{P} -a.s., for which it is enough to see that

$$\int_{\mathbb{S}^{d-1}} \phi_{n, X_1}^\theta(\langle \theta, x \rangle) d\sigma(\theta) \longrightarrow \int_{\mathbb{S}^{d-1}} \phi_{X_1}^\theta(\langle \theta, x \rangle) d\sigma(\theta) \quad \forall x \in \text{int}(\text{supp}(P)) \quad \mathbb{P}\text{-a.s.} \quad (50)$$

By Proposition 2.1, we know that \mathbb{P} -a.s., for every $x \in \text{int}(\text{supp}(P))$, $\theta \in \mathbb{S}^{d-1}$, $\phi_{n, X_1}^\theta(\langle \theta, x \rangle) \rightarrow \phi_{X_1}^\theta(\langle \theta, x \rangle)$. Moreover, there exist $n_0 \in \mathbb{N}$ and $M_x > 0$ such that $|\phi_{n, X_1}^\theta(\langle \theta, x \rangle)| \leq M_x$ for every $n \in \mathbb{N}$ and $\theta \in \mathbb{S}^{d-1}$. Therefore, we can conclude (50) from dominated convergence theorem, which ensures $n(R_n - R'_n)_+ \rightarrow 0$ \mathbb{P} -a.s., and consequently $n^2(R_n - R'_n)_+^2 \rightarrow 0$ \mathbb{P} -a.s. Our goal now is to check that $\mathbb{E}(n^2(R_n - R'_n)_+^2) \rightarrow 0$, for which suffices to prove uniform integrability of $n^2(R_n - R'_n)_+^2$. Remember that

$$\begin{aligned} n(R_n - R'_n) &= n(Z_n - Z'_n) - n \int_{\mathbb{S}^{d-1}} \left(\int_{\mathbb{R}^d} \phi_{x_0}^\theta(\langle \theta, x \rangle) d(P_n - P'_n)(x) \right) d\sigma(\theta) \\ &= n(Z_n - Z'_n) - \int_{\mathbb{S}^{d-1}} \phi_{x_0}^\theta(\langle \theta, X_1 \rangle) d\sigma(\theta) + \int_{\mathbb{S}^{d-1}} \phi_{x_0}^\theta(\langle \theta, X'_1 \rangle) d\sigma(\theta) \end{aligned} \quad (51)$$

The last two terms do not depend on n , and their second-order moment is $\|\phi_{x_0}^\theta\|_{L^2(\sigma \times P)}^2 < \infty$, by (18). Thus, it suffices to prove uniform integrability of $n^2(Z_n - Z'_n)_+^2$, for which it suffices to prove uniform integrability of the square of the right-hand side in (45). To do this, we provide a uniform bound on the $1 + \gamma$ moment. Applying Hölder's inequality with $q_1 = p$ and $q_2 = \frac{p}{p-1}$,

$$\begin{aligned} &\mathbb{E} \left(\left(\|X_1 - X'_1\| \int_{\mathbb{S}^{d-1}} \int_{\mathbb{R}} |\langle \theta, X_1 \rangle - t|^{p-1} dQ_{\tau'_\theta(1)}^\theta(y) d\sigma(\theta) \right)^{2(1+\gamma)} \right) \\ &\leq \mathbb{E} \left(\left(\|X_1 - X'_1\| \right)^{2p(1+\gamma)} \right)^{1/p} \\ &\quad \times \mathbb{E} \left(\left(n \int_{\mathbb{S}^{d-1}} \int_{\mathbb{R}} |\langle \theta, X_1 \rangle - t|^{p-1} dQ_{\tau'_\theta(1)}^\theta(y) d\sigma(\theta) \right)^{\frac{2p(1+\gamma)}{p-1}} \right)^{\frac{p-1}{p}} \end{aligned} \quad (52)$$

The first term is finite by the moment assumptions, whereas for the second, we can proceed as in the proof of Theorem 3.1. Applying Hölder's inequality as in (46), with $q_1 = \frac{2p(1+\gamma)}{2p(1+\gamma)-p+1}$ and $q_2 = \frac{2p(1+\gamma)}{p-1}$,

$$\begin{aligned} &\int_{\mathbb{R}} |\langle \theta, X_1 \rangle - t|^{p-1} dQ_{\tau'_\theta(1)}^\theta(t) \\ &\leq \frac{1}{n^{\frac{2p(1+\gamma)-p+1}{2p(1+\gamma)}}} \left(\int_{\mathbb{R}} |\langle \theta, X_1 \rangle - t|^{2p(1+\gamma)} dQ_{\tau'_\theta(1)}^\theta(t) \right)^{\frac{p-1}{2p(1+\gamma)}} \end{aligned} \quad (53)$$

Using the exchangeability as in (47), we deduce that

$$\begin{aligned} &\mathbb{E} \left(\left(\int_{\mathbb{S}^{d-1}} \int_{\mathbb{R}} |\langle \theta, X_1 \rangle - t|^{p-1} dQ_{\tau'_\theta(1)}^\theta(y) d\sigma(\theta) \right)^{\frac{2p(1+\gamma)}{p-1}} \right) \\ &\leq \mathbb{E} \left(\int_{\mathbb{S}^{d-1}} \left(\int_{\mathbb{R}} |\langle \theta, X_1 \rangle - t|^{p-1} dQ_{\tau'_\theta(1)}^\theta(y) \right)^{\frac{2p(1+\gamma)}{p-1}} d\sigma(\theta) \right) \\ &\leq \frac{1}{n^{\frac{2p(1+\gamma)}{p-1}-1}} \mathbb{E} \left(\int_{\mathbb{S}^{d-1}} \int_{\mathbb{R}} |\langle \theta, X_1 \rangle - t|^{2p(1+\gamma)} dQ_{\tau'_\theta(1)}^\theta(t) d\sigma(\theta) \right) \end{aligned}$$

$$\begin{aligned}
 &\leq \frac{1}{n^{\frac{2p(1+\gamma)}{p-1}}} \mathbb{E} \left(\int_{\mathbb{S}^{d-1}} \int_{\mathbb{R}} |\langle \theta, X_1 \rangle - t|^{2p(1+\gamma)} dQ^\theta(t) d\sigma(\theta) \right) \\
 &\leq \frac{1}{n^{\frac{2p(1+\gamma)}{p-1}}} \mathbb{E} \left(\|X_1 - Y\|^{2p(1+\gamma)} \right), \tag{54}
 \end{aligned}$$

which allow us to uniformly bound (52) by

$$n^{2(1+\gamma)} \left(\frac{1}{n^{\frac{2p(1+\gamma)}{p-1}}} \mathbb{E} \left(\|X_1 - Y\|^{2p(1+\gamma)} \right) \right)^{\frac{p-1}{p}} = \mathbb{E} \left(\|X_1 - Y\|^{2p(1+\gamma)} \right)^{\frac{p-1}{p}} < \infty,$$

which proves uniform integrability of $n^2(R_n - R'_n)_+^2$. Therefore, $\mathbb{E}(n^2(R_n - R'_n)_+^2) \rightarrow 0$, and Efron-Stein inequality allows us to conclude.

Two-sample setting. Let P_n, P'_n, Q_m, Q'_m be defined as in the proof of Proposition 3.2. Define $R'_{n,m}$ by replacing P_n by P'_n in the definition of $R_{n,m}$. Similarly, define $R''_{n,m}$ by replacing Q_m by Q'_m . Applying Efron-Stein inequality,

$$\frac{nm}{n+m} \text{Var}(R_{n,m}) \leq n^2 \mathbb{E}((R_{n,m} - R'_{n,m})_+^2) + m^2 \mathbb{E}((R_{n,m} - R''_{n,m})_+^2).$$

The convergence $n^2 \mathbb{E}((R_{n,m} - R'_{n,m})_+^2) \rightarrow 0$, follows from the same argument as in the one-sample setting, considering now any c -concave optimal transport potential $\phi_{n,m}^\theta$ between P_n^θ and Q_m^θ , and defining ϕ_{n,m,X_1}^θ as in (14) and $\phi_{X_1}^\theta$ as before. Property (7) in Proposition 2.1 enables us to extend the previous arguments for the almost sure convergence, whereas the uniform integrability follows from $\|\phi_{x_0}^\theta\|_{L^2(\sigma \times P)}^2 < \infty$, the previous bounds, and the same argument as in the proof of Proposition 3.1 for the two-sample setting.

Similarly, we can prove $m^2 \mathbb{E}(R_{n,m} - R''_{n,m})_+^2 \rightarrow 0$. Again, we need to consider a suitable choice of c -concave optimal transport potentials, but now fixed at Y_1 . Given any c -concave optimal transport potential $\psi_{n,m}^\theta$ between Q_m^θ and P_n^θ (for instance, $\psi_{n,m}^\theta = (\phi_{n,m}^\theta)^c$), define

$$\psi_{n,m,Y_1}^\theta = \psi_{n,m}^\theta - \psi_{n,m}^\theta(\langle \theta, Y_1 \rangle), \quad \psi_{Y_1}^\theta = (\phi_{x_0}^\theta)^c - (\phi_{x_0}^\theta)^c(\langle \theta, Y_1 \rangle).$$

Taking into account the symmetric role of P and Q , property (7) in Proposition 2.1 is verified, exchanging the roles of P and Q . The same argument as before allows us to conclude, now using that, by (18), $\|(\phi_{x_0}^\theta)^c\|_{L^2(\sigma \times Q)}^2 < \infty$. \square

B.3 Proof of Theorem 3.3

Weak convergence follows from the discussion preceding Theorem 3.3. The only thing left to prove is the convergence in the W_2 sense. For each $i = 1 \dots, n$, let $Y_i = \int_{\mathbb{S}^{d-1}} \phi^\theta(X_i) d\sigma(\theta)$ and $\bar{Y}_n = \frac{1}{n} \sum_{i=1}^n Y_i$. Theorem 3.2 ensures that

$$\mathbb{E} \left(\left(\sqrt{n}(\text{SW}_p^p(P_n, Q) - \mathbb{E}\text{SW}_p^p(P_n, Q)) - \sqrt{n}(\bar{Y}_n - \mathbb{E}Y) \right)^2 \right) \rightarrow 0,$$

which implies convergence in the Wasserstein distance W_2 . Since the convergence $\sqrt{n}(\bar{Y}_n - \mathbb{E}Y) \rightsquigarrow N(0, v_{P,Q}^2)$ was obtained from the central limit theorem, we have convergence of second-order moments, and therefore convergence in W_2 (see Lemma 8.3 in [BF81]). Applying the triangle inequality, we deduce

$$W_2 \left(\mathcal{L} \left(\sqrt{n}(\text{SW}_p^p(P_n, Q) - \mathbb{E}(\text{SW}_p^p(P_n, Q))) \right), N(0, v_{P,Q}^2) \right) \rightarrow 0. \quad \square$$

C Proofs of Section 4

C.1 Weak convergence of weighted integrals of the quantile process

Our main result in this section is based on the next lemma, which shows weak convergence of weighted integrals of the empirical quantile process. The proof relies on the argument used for property (2.1) in Theorem 2.1 of [CH90].

Lemma C.1. *Let $P \in \mathcal{P}(\mathbb{R})$, with density f such that $f(F^{-1}(t))$ is positive and continuous in $(0, 1)$, and monotone for t sufficiently close to 0 and 1. Let $h : (0, 1) \rightarrow \mathbb{R}$ be such that there exists $\alpha, \beta > 1$, $\frac{1}{\alpha} + \frac{1}{\beta} = 1$, verifying that $J_\alpha(P) < \infty$ and $\int_0^1 |h(t)|^\beta dt < \infty$. Then*

$$\int_0^1 h(t) \sqrt{n} (F_n^{-1}(t) - t) dt \rightsquigarrow \int_0^1 \frac{h(t)}{f(F^{-1}(t))} \mathbb{B}(t) dt, \quad (55)$$

where \mathbb{B} denotes a Brownian bridge.

Proof. The proof is based on two steps. First, it is easy to show that the tails in (55) do not play any role, in the sense that

$$\begin{aligned} \int_0^{\frac{1}{n+1}} h(t) \sqrt{n} (F_n^{-1}(t) - F^{-1}(t)) dt &\rightarrow_P 0 \\ \int_{\frac{1}{n+1}}^1 h(t) \sqrt{n} (F_n^{-1}(t) - F^{-1}(t)) dt &\rightarrow_P 0 \end{aligned}$$

We prove the first convergence; the second follows by analogy. It follows from Markov's inequality and the bound

$$\begin{aligned} \mathbb{E} \left(\left| \int_0^{\frac{1}{n+1}} h(t) \sqrt{n} (F_n^{-1}(t) - F^{-1}(t)) dt \right| \right) &\leq \left(\int_0^{\frac{1}{n+1}} |h(t)|^\beta dt \right)^{1/\beta} \left(n^{\alpha/2} \mathbb{E}(W_\alpha^\alpha(P_n, P)) \right)^{1/\alpha} \\ &\leq 5\alpha J_\alpha(P)^{1/\alpha} \left(\int_0^{\frac{1}{n+1}} |h(t)|^\beta dt \right)^{1/\beta} \rightarrow 0 \end{aligned}$$

where the last inequality follows from Proposition 5.3 in [BL19], and the convergence in the last line is a consequence of $\int_0^1 |h(t)|^\beta dt < \infty$. Thus, it suffices to prove the weak convergence

$$\int_{\frac{1}{n+1}}^{\frac{n}{n+1}} h(t) \sqrt{n} (F_n^{-1}(t) - F^{-1}(t)) dt \rightsquigarrow \int_0^1 \frac{h(t)}{f(F^{-1}(t))} \mathbb{B}(t) dt$$

where \mathbb{B} is a Brownian bridge. Following Theorem 2.1 in [CH90], let X_1, \dots, X_n be an i.i.d. sample from P , and assume that $X_i = F^{-1}(Z_i)$, where Z_1, \dots, Z_n are i.i.d. from the uniform distribution in $(0, 1)$. If U_n denotes the cumulative distribution function of Z_1, \dots, Z_n , the weighted general quantile process and the uniform quantile process are defined for each $t \in (0, 1)$ as

$$\begin{aligned} \rho_n(t) &:= f(F^{-1}(t)) \sqrt{n} (F_n^{-1}(t) - F^{-1}(t)), \\ u_n(t) &:= \sqrt{n} (U_n^{-1}(t) - t). \end{aligned}$$

If we denote $r(t) = \frac{h(t)}{f(F^{-1}(t))}$, our goal is to show the convergence

$$\int_{\frac{1}{n+1}}^{\frac{n}{n+1}} \rho_n(t) r(t) dt \rightsquigarrow \int_0^1 \mathbb{B}(t) r(t) dt$$

Indeed, we can prove something stronger. By Theorem 2.1 in [CCHM86], we can consider a sequence of Brownian bridges $\{\mathbb{B}_n : n \geq 1\}$ such that, as $n \rightarrow \infty$,

$$n^\nu \sup_{\frac{1}{n+1} \leq t \leq \frac{n}{n+1}} |u_n(t) - \mathbb{B}_n(t)| / (t(1-t))^{(1/2)-\nu} = O_P(1)$$

for every $0 \leq \nu < 1/2$, and $\sup_{0 \leq t \leq 1} |u_n(t) - \mathbb{B}_n(t)| = o_P(1)$. For this particular sequence of Brownian bridges, we will conclude that

$$\int_{\frac{1}{n+1}}^{\frac{n}{n+1}} \rho_n(t) r(t) dt - \int_0^1 \mathbb{B}_n(t) r(t) dt \rightarrow_P 0.$$

For simplicity, we can split the integrals at the middle point $1/2$, to include only one of the tails. By symmetry, it suffices to prove

$$\int_{\frac{1}{n+1}}^{\frac{1}{2}} \rho_n(t) r(t) dt - \int_0^{\frac{1}{2}} \mathbb{B}_n(t) r(t) dt \rightarrow_P 0 \quad (56)$$

In analogy with Theorem 2.1 in [CH90] (with $p = 1$, $w(t) = 1/f(F^{-1}(t))$, $q(t) = 1/r(t)$, here q might be infinite and it is not necessarily a positive function), under our given assumptions, the following properties are verified:

(I) $f(F^{-1}(t))$ is continuous and positive in $(0, 1/2]$.

(II) $\int_0^{1/2} t^{1/2}|r(t)| < \infty$.

$$\begin{aligned} \int_0^{1/2} t^{1/2}|r(t)| &\leq \left(\int_0^{1/2} |h(t)|^\beta dt \right)^{1/\beta} \left(\int_0^{1/2} \frac{t^{\alpha/2}}{f(F^{-1}(t))^\alpha} dt \right)^{1/\alpha} \\ &\leq \sqrt{2} \left(\int_0^{1/2} |h(t)|^\beta dt \right)^{1/\beta} J_\alpha(P)^{1/\alpha} < \infty \end{aligned}$$

(III) $w(t) = 1/f(F^{-1}(t))$ is monotone in $(0, \epsilon]$, for ϵ small enough.

(IV) $\lim_{\epsilon \rightarrow 0} \int_0^\epsilon f(F^{-1}(t))w(t/\lambda)t^{1/2}|r(t)|dt = 0$ for every $\lambda > 1$ if w is non-decreasing, and for every $0 < \lambda < 1$ if w is non-increasing.

Given a fixed value of λ in $(0, 1)$ or $(1, \infty)$, for ϵ small enough, $(0, \epsilon/\lambda) \subset (0, 1/2)$. Thus, for any of such ϵ ,

$$\begin{aligned} \int_0^\epsilon f(F^{-1}(t))w(t/\lambda)t^{1/2}|r(t)|dt &= \int_0^\epsilon \frac{1}{f(F^{-1}(t/\lambda))} t^{1/2}|h(t)|dt = \\ &\leq \left(\int_0^\epsilon \frac{t^{\alpha/2}}{f(F^{-1}(t/\lambda))^\alpha} dt \right)^{1/\alpha} \left(\int_0^\epsilon |h(t)|^\beta \right)^{1/\beta} \\ &= \lambda^{\frac{1}{2} + \frac{1}{\alpha}} \left(\int_0^{\epsilon/\lambda} \frac{s^{\alpha/2}}{f(F^{-1}(s))^\alpha} ds \right)^{1/\alpha} \left(\int_0^\epsilon |h(t)|^\beta \right)^{1/\beta} \rightarrow 0 \end{aligned}$$

where the convergence to 0 follows from the fact that both terms in the last expression converge to 0 as $\epsilon \rightarrow 0$, since $\epsilon/\lambda < 1/2$ and

$$\int_0^{1/2} \frac{s^{\alpha/2}}{f(F^{-1}(s))^\alpha} ds \leq 2^{\alpha/2} J_\alpha(P) < \infty, \quad \int_0^1 |h(t)|^\beta < \infty.$$

Properties (I)-(IV) represent a slight modification of the assumptions (I)-(IV) in Theorem 2.1 in [CH90]. As noted in the last paragraph on page 67 in [CH90], all the proofs remain valid for $p = 1$ without the absolute value. Similarly, it is easy to see that the exact same steps as in the proof of property (2.1) in Theorem 2.1 from [CH90] can be applied to our specific scenario to demonstrate (56), thereby completing the proof. \square

C.2 Additional lemmas

Throughout this section, we repeatedly rely on the following elementary inequality. Given $x, y \in \mathbb{R}^d$ and $q > 0$,

$$\|x + y\|^q \leq C_q (\|x\|^q + \|y\|^q) \quad \text{where} \quad \begin{cases} C_q = 1 & 0 < q \leq 1 \\ C_q = 2^{q-1} & q \geq 1 \end{cases} \quad (57)$$

First, we present a well-known fact about the convergence in mean of the Wasserstein distance (see Theorem 2.14 in [BL19]), for which we include a shorter alternative proof.

Lemma C.2. *Let $p \geq 1$. For any $P \in \mathcal{P}_p(\mathbb{R})$, $\mathbb{E}(\mathbf{W}_p^p(P_n, P)) \rightarrow 0$.*

Proof. The proof is based on studying $|F_n^{-1}(\omega, t) - F^{-1}(t)|^p$ as a random variable in the product space $\Omega \times (0, 1)$, with the product probability $\mathbb{P} \times \ell_1$, where ℓ_1 denotes the Lebesgue measure in \mathbb{R} . The Glivenko-Cantelli theorem ensures that there exists $\Omega_0 \subset \Omega$ with $\mathbb{P}(\Omega_0) = 1$ such that $\sup_{x \in \mathbb{R}} |F_n(\omega, x) - F(x)| \rightarrow 0$, for every $\omega \in \Omega_0$. Thus, for any of such $\omega \in \Omega_0$, $P_n^\omega \rightsquigarrow P$. If $C_{F^{-1}}$ denotes the set of continuity points of F^{-1} , which verifies $\ell_1(C_{F^{-1}}) = 1$, by the Skorohod elementary theorem (e.g. see pp. 9-10 in [SW09]),

$$|F_n^{-1}(\omega, t) - F^{-1}(t)|^p \rightarrow 0 \quad \forall (\omega, t) \in \Omega_0 \times C_{F^{-1}}. \quad (58)$$

Thus, we have almost sure convergence in $\Omega \times (0, 1)$, and the only thing left to prove is uniform integrability of the sequence. From the inequality (57), it suffices to prove uniform integrability of $|F_n^{-1}(\omega, t)|^p$, which follows from

$$\mathbb{P} \times \ell_1 \left(\{(\omega, t) : F_n^{-1}(\omega, t) \leq x\} \right) = \mathbb{P} \times \ell_1 \left(\{(\omega, t) : t \leq F_n(\omega, x)\} \right) = \int_\Omega F_n(\omega, x) d\mathbb{P}(\omega) = F(x). \quad (59)$$

Therefore, as a random variable in $(0, 1) \times \Omega$, $\mathcal{L}(F_n^{-1}) = P$, which does not depend on n and has finite moment of order p . This proves the uniform integrability. \square

Given $p > 1$, the function $h_p(x) = |x|^p$ is differentiable, and its derivative is given by $h'_p(x) = p \operatorname{sgn}(x)|x|^{p-1}$. The following result helps us bound the expected values of integrals involving the difference between the values of h'_p at the empirical quantile function difference and the population value of this difference. The following lemma is presented for general $p > 1$, although improved assumptions could be given for the simpler case $p = 2$. However, the proof of Theorem 4.1 for $p = 2$ relies on a different technique that does not require the use of this lemma. Therefore, for simplicity, we do not pursue this direction further.

Lemma C.3. *Let $p > 1$, $\beta \geq 1$, $\gamma > 0$, and $P, Q \in \mathcal{P}(\mathbb{R})$, with distribution functions F, G , respectively. If P, Q have finite moments of order $\beta(p-1)(1+\gamma)$, then*

$$\mathbb{E} \left(\int_0^1 |h'_p(F_n^{-1}(t) - G^{-1}(t)) - h'_p(F^{-1}(t) - G^{-1}(t))|^\beta dt \right) \rightarrow 0, \quad (60)$$

$$\mathbb{E} \left(\int_0^1 |h'_p(F_n^{-1}(t) - G_m^{-1}(t)) - h'_p(F^{-1}(t) - G^{-1}(t))|^\beta dt \right) \rightarrow 0. \quad (61)$$

Proof. From the inequality (57), it follows that

$$\begin{aligned} & \frac{1}{pC_\beta^\beta} \left| h'_p(F_n^{-1}(t) - G^{-1}(t)) - h'_p(F^{-1}(t) - G^{-1}(t)) \right|^\beta \\ & \leq \left| \operatorname{sgn}(F_n^{-1}(t) - G^{-1}(t)) (|F_n^{-1}(t) - G^{-1}(t)|^{p-1} - |F^{-1}(t) - G^{-1}(t)|^{p-1}) \right|^\beta \end{aligned} \quad (62)$$

$$+ \left| (\operatorname{sgn}(F_n^{-1}(t) - G^{-1}(t)) - \operatorname{sgn}(F^{-1}(t) - G^{-1}(t))) |F^{-1}(t) - G^{-1}(t)|^{p-1} \right|^\beta \quad (63)$$

Applying Hölder's inequality with $q_1 = \frac{1+\gamma}{\gamma}$ and $q_2 = 1 + \gamma$, we can bound the expectation of the integral of (63) by

$$\begin{aligned} & \mathbb{E} \left(\int_{\{t \in (0,1): F^{-1}(t) \neq G^{-1}(t)\}} \left| \operatorname{sgn}(F_n^{-1}(t) - G^{-1}(t)) - \operatorname{sgn}(F^{-1}(t) - G^{-1}(t)) \right|^{\frac{\beta(1+\gamma)}{\gamma}} \right)^{\frac{\gamma}{1+\gamma}} \\ & \quad \times \left(\int_0^1 |F^{-1}(t) - G^{-1}(t)|^{\beta(p-1)(1+\gamma)} dt \right)^{\frac{1}{1+\gamma}} \end{aligned} \quad (64)$$

The second term is finite by the moment assumptions, and we will see that the first term converges to 0. With the same reasoning as in (58), $F_n^{-1}(t) \rightarrow F^{-1}(t)$ for every $t \in C_{F^{-1}}$ in a set of \mathbb{P} -probability one. Therefore,

$$I(F^{-1}(t) \neq G^{-1}(t)) \left| \operatorname{sgn}(F_n^{-1}(t) - G^{-1}(t)) - \operatorname{sgn}(F^{-1}(t) - G^{-1}(t)) \right|^{\frac{\beta(1+\gamma)}{\gamma}} \rightarrow 0$$

almost surely as a random variable in $\Omega \times (0, 1)$. Since it is bounded by $2^{\frac{\beta(1+\gamma)}{\gamma}}$, it is uniformly integrable, and we can conclude that (64) converges to 0. Now, the expected value of the integral of (62) is bounded by

$$2^\beta \mathbb{E} \left(\int_0^1 \left| |F_n^{-1}(t) - G^{-1}(t)|^{p-1} - |F^{-1}(t) - G^{-1}(t)|^{p-1} \right|^\beta dt \right) \quad (65)$$

We distinguish between two different cases:

- If $1 < p \leq 2$, it follows from (57) that $\left| |x|^{p-1} - |y|^{p-1} \right| \leq |x - y|^{p-1}$. Thus, we can bound the (65), up to a constant factor, by

$$\mathbb{E} \left(\int_0^1 |F_n^{-1}(t) - F^{-1}(t)|^{\beta(p-1)} dt \right) = \mathbb{E}(\mathbf{W}_{\beta(p-1)}^{\beta(p-1)}(P_n, P)) \rightarrow 0, \quad (66)$$

where the convergence follows from Lemma C.2.

- If $p > 2$, then $h_{p-1}(x) = |\cdot|^{p-1}$ is convex and differentiable. Thus, we know that for every $x, y \in \mathbb{R}$

$$h_{p-1}(x) - h_{p-1}(y) \leq h'_{p-1}(x)(x - y), \quad (67)$$

which implies

$$|h_{p-1}(x) - h_{p-1}(y)| \leq |h'_{p-1}(x) + h'_{p-1}(y)||x - y|. \quad (68)$$

Using this inequality, we can bound (65), up to a constant factor, by

$$\begin{aligned} & \mathbb{E} \left(\int_0^1 |F_n^{-1}(t) - G^{-1}(t)|^{\beta(p-2)} |F_n^{-1}(t) - F^{-1}(t)|^\beta dt \right) \\ & + \mathbb{E} \left(\int_0^1 |F^{-1}(t) - G^{-1}(t)|^{\beta(p-2)} |F_n^{-1}(t) - F^{-1}(t)|^\beta dt \right) \end{aligned}$$

We prove that the first term converges to 0, for the second term follows similarly. Applying Hölder's inequality with $q_1 = \frac{p-1}{p-2}$, $q_2 = p-1$, the first term is bounded by

$$\mathbb{E} \left(\int_0^1 |F_n^{-1}(t) - G^{-1}(t)|^{\beta(p-1)} dt \right)^{\frac{p-2}{p-1}} \mathbb{E} \left(\int_0^1 |F_n^{-1}(t) - F^{-1}(t)|^{\beta(p-1)} dt \right)^{\frac{1}{p-1}}$$

By the moment assumptions, the first term is bounded, whereas the second term converges to 0 by Lemma C.2.

Two-sample setting: The proof follows from the same steps, and a further application of the inequality (57). \square

C.3 Proof of Theorem 4.1 for $d = 1$

For simplicity, we prove first Theorem 4.1 in the one-dimensional setting. To do this, denote by f and F the density and distribution function of P , which play the role of f_θ, F_θ in the statement of Theorem 4.1, and note that $SJ_{\alpha(1+\gamma)}(P) = J_{\alpha(1+\gamma)}(P)$. The proof in the one-sample setting is divided into two parts. We begin with the simpler proof for the case $p = 2$, and then generalize for $p \neq 2$.

C.3.1 Case $p = 2$

Let F, F_n, G be the distribution function of P, P_n and Q , respectively. From the quantile representation of the Wasserstein distance,

$$\begin{aligned} \mathbb{E}(W_2^2(P_n, Q)) - W_2^2(P, Q) &= \mathbb{E} \left(\int_0^1 (F_n^{-1}(t) - G^{-1}(t))^2 - (F^{-1}(t) - G^{-1}(t))^2 dt \right) \\ &= \mathbb{E} \left(\int_0^1 (F_n^{-1}(t)^2 - F^{-1}(t)^2) dt \right) - 2 \mathbb{E} \left(\int_0^1 G^{-1}(t)(F_n^{-1}(t) - F^{-1}(t)) dt \right) \end{aligned} \quad (69)$$

By (59), we know that $\mathcal{L}(F_n^{-1}) = P$ as a random variable in $\Omega \times (0, 1)$. Therefore, it is easy to see that the first term vanishes. Thus, it suffices to show

$$\mathbb{E} \left(\int_0^1 G^{-1}(t) \sqrt{n} (F_n^{-1}(t) - F^{-1}(t)) dt \right) \rightarrow 0. \quad (70)$$

Under the assumptions of Theorem 4.1, we can apply Lemma C.1 with $h = G^{-1}$ to conclude

$$\int_0^1 G^{-1}(t) \sqrt{n} (F_n^{-1}(t) - F^{-1}(t)) dt \rightsquigarrow \int_0^1 \frac{G^{-1}(t)}{f(F^{-1}(t))} \mathbb{B}(t) dt.$$

The limiting variable is an integral of the Brownian bridge \mathbb{B} , it follows a centered Gaussian distribution. Thus, the only remaining task is to prove moment convergence, which is established using a uniform integrability argument. It suffices to demonstrate boundedness of the sequence

$$\begin{aligned} & \mathbb{E} \left(\left| \int_0^1 G^{-1}(t) \sqrt{n} (F_n^{-1}(t) - F^{-1}(t)) dt \right|^{1+\gamma} \right) \leq \\ & \leq \left(\int_0^1 |G^{-1}(t)|^{\beta(1+\gamma)} dt \right)^{1/\beta} \mathbb{E} \left(\int_0^1 |\sqrt{n} (F_n^{-1}(t) - F^{-1}(t))|^{\alpha(1+\gamma)} dt \right)^{1/\alpha} \\ & = \left(\int |y|^{\beta(1+\gamma)} dQ(y) \right)^{1/\beta} \left(n^{\alpha(1+\gamma)/2} \mathbb{E}(W_{\alpha(1+\gamma)}^{\alpha(1+\gamma)}(P_n, P)) \right)^{1/\alpha} \\ & \leq \mathbb{E}(\|Y\|^{\beta(1+\gamma)})^{1/\beta} (5\alpha(1+\gamma))^{1+\gamma} J_{\alpha(1+\gamma)}^{1/\alpha} \end{aligned} \quad (71)$$

where the last inequality follows from Proposition 5.3. in [BL19], and $\mathcal{L}(Y) = Q$.

Two-sample setting. To prove (23), reasoning as in (69), it suffices to show convergence to 0 of

$$\begin{aligned} & \sqrt{\frac{nm}{n+m}} \mathbb{E} \left(\int_0^1 \left(F_n^{-1}(t) G_m^{-1}(t) - F^{-1}(t) G^{-1}(t) \right) dt \right) \\ &= \sqrt{\frac{m}{n+m}} \mathbb{E} \left(\int_0^1 G^{-1}(t) \sqrt{n} \left(F_n^{-1}(t) - F^{-1}(t) \right) dt \right) \\ &+ \sqrt{\frac{n}{n+m}} \mathbb{E} \left(\int_0^1 F^{-1}(t) \sqrt{m} \left(G_m^{-1}(t) - G^{-1}(t) \right) dt \right) \\ &+ \sqrt{\frac{m}{n+m}} \mathbb{E} \left(\int_0^1 \sqrt{n} \left(F_n^{-1}(t) - F^{-1}(t) \right) \left(G_m^{-1}(t) - G^{-1}(t) \right) dt \right) \end{aligned}$$

The same reasoning as in the one-sample setting allows us to conclude convergence to 0 of the first two terms in the last equality. Applying Hölder's inequality, the absolute value of the last term is bounded by

$$\left(n^{\alpha/2} \mathbb{E} \left(W_\alpha^\alpha(P_n, P) \right) \right)^{1/\alpha} \mathbb{E} \left(W_\beta^\beta(Q_m, Q) \right)^{1/\beta}.$$

Given that $J_\alpha(P) < \infty$, Proposition 5.13 in [BL19] ensures that the expression inside the first parenthesis is bounded, and Lemma C.2 implies $\mathbb{E} \left(W_\beta^\beta(Q_m, Q) \right) \rightarrow 0$.

C.3.2 Case $p \neq 2$

For every $p > 1$, by (67) and the positivity condition in (21), we know that

$$\begin{aligned} 0 &\leq \sqrt{n} \left(\mathbb{E} \left(W_p^p(P_n, Q) \right) - W_p^p(P, Q) \right) = \\ &= \sqrt{n} \mathbb{E} \left(\int_0^1 \left| F_n^{-1}(t) - G^{-1}(t) \right|^p - \left| F^{-1}(t) - G^{-1}(t) \right|^p dt \right) \\ &\leq \mathbb{E} \left(\int_0^1 h'_p \left(F_n^{-1}(t) - G^{-1}(t) \right) \sqrt{n} \left(F_n^{-1}(t) - F^{-1}(t) \right) dt \right) \end{aligned} \quad (72)$$

Thus, we only need to prove that (72) converges to 0. To exploit the same arguments as before, it is convenient to consider the decomposition

$$\begin{aligned} & \int_0^1 h'_p \left(F_n^{-1}(t) - G^{-1}(t) \right) \sqrt{n} \left(F_n^{-1}(t) - F^{-1}(t) \right) dt \\ &= \int_0^1 \left(h'_p \left(F_n^{-1}(t) - G^{-1}(t) \right) - h'_p \left(F^{-1}(t) - G^{-1}(t) \right) \right) \sqrt{n} \left(F_n^{-1}(t) - F^{-1}(t) \right) dt \end{aligned} \quad (73)$$

$$+ \int_0^1 h'_p \left(F_n^{-1}(t) - G^{-1}(t) \right) \sqrt{n} \left(F_n^{-1}(t) - F^{-1}(t) \right) dt \quad (74)$$

and prove independently that the expectation of each term converges to 0. Applying Hölder's inequality, the absolute value of the expectation of (73) is bounded by

$$\begin{aligned} & \mathbb{E} \left(\int_0^1 \left| h'_p \left(F_n^{-1}(t) - G^{-1}(t) \right) - h'_p \left(F^{-1}(t) - G^{-1}(t) \right) \right|^\beta dt \right)^{1/\beta} \\ & \quad \times \left(n^{\alpha/2} \mathbb{E} \left(W_\alpha^\alpha(P_n, P) \right) \right)^{1/\alpha} \rightarrow 0. \end{aligned}$$

The first term converges to 0 by Lemma C.3, whereas the second term is bounded by Proposition 5.3 in [BL19]. Convergence to 0 of the expectation of (74) follows by essentially the same arguments as in the case $p = 2$. Specifically, we apply Lemma C.1 with $h(t) = h'_p \left(F^{-1}(t) - G^{-1}(t) \right)$ to conclude the weak convergence to a centered Gaussian random variable. Uniform integrability follows as in (71).

Two-sample setting: With the same ideas as in (72),

$$0 \leq \sqrt{\frac{nm}{n+m}} \mathbb{E} \left(\int_0^1 \left| F_n^{-1}(t) - G_m^{-1}(t) \right|^p - \left| F^{-1}(t) - G^{-1}(t) \right|^p dt \right)$$

$$\leq \sqrt{\frac{nm}{n+m}} \mathbb{E} \left(\int_0^1 h'_p(F_n^{-1}(t) - G_m^{-1}(t)) (F_n^{-1}(t) - F^{-1}(t) + G^{-1}(t) - G_m^{-1}(t)) dt \right)$$

By the symmetry of the preceding expression, along with the symmetry in the assumptions on P and Q , and the bound $0 \leq \frac{m}{n+m} \leq 1$ it suffices to show

$$\mathbb{E} \left(\int_0^1 h'_p(F_n^{-1}(t) - G_m^{-1}(t)) \sqrt{n} (F_n^{-1}(t) - F^{-1}(t)) dt \right) \rightarrow 0.$$

This can be concluded as in the one-sample case, using now the second part of Lemma C.3. \square

C.4 Proof of Theorem 4.1

We aim to prove (21), which can be rewritten as

$$\int_{\mathbb{S}^{d-1}} \sqrt{n} \left(\mathbb{E} \left(\mathbf{W}_p^p(P_n^\theta, Q^\theta) \right) - \mathbf{W}_p^p(P^\theta, Q^\theta) \right) d\sigma(\theta) \rightarrow 0$$

Given that $SJ_{\alpha(1+\gamma)}(P) < \infty$, then $J_{\alpha(1+\gamma)}(P^\theta) < \infty$ for almost every $\theta \in \mathbb{S}^{d-1}$. In addition, the moment assumptions on P, Q of Theorem 4.1 imply the same moment assumptions for P^θ, Q^θ for each $\theta \in \mathbb{S}^{d-1}$. Thus, from the one-dimensional case, it follows that

$$\sqrt{n} \left(\mathbb{E} \left(\mathbf{W}_p^p(P_n^\theta, Q^\theta) \right) - \mathbf{W}_p^p(P^\theta, Q^\theta) \right) \rightarrow 0,$$

for almost every $\theta \in \mathbb{S}^{d-1}$. We have pointwise convergence towards 0 of the function within the integral. Therefore, we can conclude by the dominated convergence theorem if we bound the function by an integrable expression. Define $h_{\theta,n}(t) = G_\theta^{-1}(t)$ if $p = 2$ and $h_{\theta,n}(t) = |F_{\theta,n}^{-1}(t) - G_\theta^{-1}(t)|^{p-1}$ if $p \neq 2$. For each value of p , reasoning as in (69) or (72),

$$\begin{aligned} 0 &\leq \sqrt{n} \left(\mathbb{E} \left(\mathbf{W}_p^p(P_n^\theta, Q^\theta) \right) - \mathbf{W}_p^p(P^\theta, Q^\theta) \right) \\ &\leq \mathbb{E} \left(\int_0^1 |h_{\theta,n}(t)| \sqrt{n} (F_{\theta,n}^{-1}(t) - F_\theta^{-1}(t)) dt \right) \\ &\leq 5\alpha \mathbb{E} \left(\int |h_{\theta,n}(t)|^\beta dt \right)^{1/\beta} J_\alpha^{1/\alpha}(P^\theta). \end{aligned}$$

In any case, under the given moment assumptions, there exists a constant $M < \infty$ such that $\mathbb{E} \left(\int |h_{\theta,n}(t)|^\beta dt \right)^{1/\beta} \leq M$. Since we are assuming that $\theta \mapsto J_\alpha(P^\theta) \in L_\alpha(\sigma)$, and $\alpha \geq 1$, applying Jensen's inequality, it follows that $\theta \mapsto J_2(P^\theta)^{1/\alpha} \in L_2(\sigma)$, and we can conclude applying dominated convergence theorem.

Two-sample setting: The proof is analogous to the one-sample case. The one-dimensional case ensures that

$$\sqrt{\frac{nm}{n+m}} \left(\mathbb{E} \left(\mathbf{W}_p^p(P_n^\theta, Q_m^\theta) \right) - \mathbf{W}_p^p(P^\theta, Q^\theta) \right) \rightarrow 0.$$

In order to apply dominated convergence theorem, we need to bound this term by an integrable function. We can decompose the problem as follows

$$\begin{aligned} 0 &\leq \sqrt{\frac{nm}{n+m}} \left(\mathbb{E} \left(\mathbf{W}_p^p(P_n^\theta, Q_m^\theta) \right) - \mathbf{W}_p^p(P^\theta, Q^\theta) \right) \\ &\leq \sqrt{m} \left(\mathbb{E} \left(\mathbf{W}_p^p(P_n^\theta, Q_m^\theta) \right) - \mathbf{W}_p^p(P_n^\theta, Q^\theta) \right) \\ &\quad + \sqrt{n} \left(\mathbb{E} \left(\mathbf{W}_p^p(P_n^\theta, Q^\theta) \right) - \mathbf{W}_p^p(P^\theta, Q^\theta) \right) \end{aligned}$$

By the one-sample setting, there exists $C < \infty$ such that the last term is bounded by $CJ_\alpha^{1/\alpha}(P^\theta)$. Similarly,

$$\begin{aligned} \sqrt{n} \left(\mathbb{E} \left(\mathbf{W}_p^p(P_n^\theta, Q_m^\theta) \right) - \mathbf{W}_p^p(P_n^\theta, Q^\theta) \right) &\leq \mathbb{E} \left(\int_0^1 |h_{\theta,n,m}(t)| \sqrt{n} (G_{\theta,m}^{-1}(t) - G_\theta^{-1}(t)) dt \right) \\ &\leq 5\alpha' \mathbb{E} \left(\int |h_{\theta,n,m}(t)|^{\beta'} dt \right)^{1/\beta'} J_{\alpha'}^{1/\alpha'}(Q^\theta) = C' J_{\alpha'}^{1/\alpha'}(Q^\theta), \end{aligned}$$

where $h_{\theta,n,m}(t) = F_{\theta,n}^{-1}(t)$ if $p = 2$ and $h_{\theta,n,m}(t) = |F_{\theta,n}^{-1}(t) - G_{\theta,m}^{-1}(t)|^{p-1}$ if $p \neq 2$, and, therefore, $C' < \infty$. Since $\theta \mapsto CJ_\alpha^{1/\alpha}(P^\theta) + C'J_{\alpha'}^{1/\alpha'}(Q^\theta)$ is integrable under the given assumptions, we can conclude as before. \square

D Proofs of Section 5

D.1 Proof of Proposition 5.1

The proof is inspired by the bootstrap results of [BF81]. First, given a random variable Θ drawn from σ , define $Z_n(\Theta) = \text{SW}_p^p(P_n^\Theta, Q^\Theta)$ and $Z(\Theta) = \text{SW}_p^p(P^\Theta, Q^\Theta)$. Throughout this proof, \mathbb{E}_Θ denotes expectation taken only over $\Theta_1, \dots, \Theta_k$, whereas \mathbb{E} represents expectation over all random variables. Then

$$S_{n,k} = \frac{1}{\sqrt{k}} \sum_{i=1}^k \left(Z_n(\Theta_i) - \mathbb{E}_\Theta(Z_n(\Theta_i)) \right), \quad S_k = \frac{1}{\sqrt{k}} \sum_{i=1}^k \left(Z(\Theta_i) - \mathbb{E}_\Theta(Z(\Theta_i)) \right). \quad (75)$$

By definition, $Z(\Theta)$ is a random variable with mean and variance given by

$$\begin{aligned} \mathbb{E}(Z(\Theta)) &= \int_{\mathbb{S}^{d-1}} \text{W}_p^p(P^\theta, Q^\theta) d\sigma(\theta) = \text{SW}_p^p(P, Q) \\ \text{Var}(Z(\Theta)) &= \int_{\mathbb{S}^{d-1}} \text{W}_p^{2p}(P^\theta, Q^\theta) d\sigma(\theta) - \text{SW}_p^{2p}(P, Q) \end{aligned}$$

which is finite, since $\text{W}_p^{2p}(P^\theta, Q^\theta) \leq \text{W}_p^{2p}(P, Q) \leq \text{W}_{2p}^{2p}(P, Q)$ for every $\theta \in \mathbb{S}^{d-1}$. Therefore, a simple application of the central limit theorem ensures $S_k \rightsquigarrow N(0, w_{P,Q}^2)$ and second-order moment convergence, which implies $W_2(\mathcal{L}(S_k), N(0, w_{P,Q}^2)) \rightarrow 0$, by Lemma 8.3 in [BF81]. Consider now the coupling given by independent variables X_1, \dots, X_n and $\Theta_1, \dots, \Theta_k$.

$$\begin{aligned} \text{W}_2^2(\mathcal{L}(S_{n,k}), \mathcal{L}(S_k)) &\leq \mathbb{E} \left(\left(\frac{1}{\sqrt{k}} \sum_{i=1}^k Z_n(\Theta_i) - \mathbb{E}_\Theta(Z_n(\Theta_i)) - (Z(\Theta_i) - \mathbb{E}_\Theta(Z(\Theta_i))) \right)^2 \right) \\ &= \frac{1}{k} \sum_{i=1}^k \mathbb{E} \left((Z_n(\Theta_i) - Z(\Theta_i) - \mathbb{E}_\Theta(Z_n(\Theta_i) - Z(\Theta_i)))^2 \right) \\ &\leq 2\mathbb{E} \left((Z_n(\Theta) - Z(\Theta))^2 \right) = 2\mathbb{E} \left((\text{W}_p^p(P_n^\Theta, Q^\Theta) - \text{W}_p^p(P^\Theta, Q^\Theta))^2 \right) \end{aligned} \quad (76)$$

From the bound (68) and the triangle inequality, for every $\theta \in \mathbb{S}^{d-1}$,

$$\begin{aligned} |\text{W}_p^p(P_n^\theta, Q^\theta) - \text{W}_p^p(P^\theta, Q^\theta)| &\leq p(\text{W}_p^{p-1}(P_n^\theta, Q^\theta) + \text{W}_p^{p-1}(P^\theta, Q^\theta)) |\text{W}_p(P_n^\theta, Q^\theta) - \text{W}_p(P^\theta, Q^\theta)| \\ &\leq p(\text{W}_p^{p-1}(P_n^\theta, Q^\theta) + \text{W}_p^{p-1}(P^\theta, Q^\theta)) \text{W}_p(P_n^\theta, P^\theta). \end{aligned}$$

By the above inequality, (57) and Hölder's inequality with $q_1 = \frac{p}{p-1}$, $q_2 = p$, (76) is bounded, up to a constant, by

$$\begin{aligned} &\left(\mathbb{E}(\text{W}_p^{2p}(P_n^\Theta, Q^\Theta)) + \mathbb{E}(\text{W}_p^{2p}(P^\Theta, Q^\Theta)) \right)^{\frac{p-1}{p}} \mathbb{E}(\text{W}_p^{2p}(P_n^\Theta, P^\Theta))^{\frac{1}{p}} \\ &\leq \left(\mathbb{E}(\text{W}_{2p}^{2p}(P_n, Q)) + \text{W}_{2p}^{2p}(P, Q) \right)^{\frac{p-1}{p}} \mathbb{E}(\text{W}_{2p}^{2p}(P_n, P))^{\frac{1}{p}} \end{aligned}$$

Since we are assuming that P, Q have finite moment of order $2p$, the first term is bounded, whereas the second term converges to 0 by Lemma C.2. \square

D.2 Proof of Corollary 5.2

The proof is based on the following decomposition.

$$\begin{aligned} &\sqrt{\frac{kn}{k+n}} \left(\text{SW}_{p,k}^p(P_n, Q) - \text{SW}_p^p(P, Q) \right) = \\ &= \sqrt{\frac{n}{k+n}} \sqrt{k} \left(\text{SW}_{p,k}^p(P_n, Q) - \text{SW}_p^p(P_n, Q) \right) \quad \left(\rightsquigarrow \sqrt{1-\tau} \cdot N(0, w_{P,Q}^2) \right) \\ &+ \sqrt{\frac{k}{k+n}} \sqrt{n} \left(\text{SW}_p^p(P_n, Q) - \mathbb{E}(\text{SW}_p^p(P_n, Q)) \right) \quad \left(\rightsquigarrow \sqrt{\tau} \cdot N(0, v_{P,Q}^2) \right) \end{aligned}$$

$$+ \sqrt{\frac{k}{k+n}} \sqrt{n} \left(\mathbb{E}(\text{SW}_p^p(P_n, Q)) - \text{SW}_p^p(P, Q) \right) \quad (\rightarrow \sqrt{\tau} \cdot 0),$$

Under the given assumptions, Theorem 4.1 ensures that the last term converges to 0, while Theorem 5.1 establishes the weak convergence of the first term, and Theorem 3.3 guarantees the weak convergence of the second term. Therefore, the only thing left to prove is that the sum of both sequences weakly converges to the sum of two independent Gaussian random variables with the distribution of the previous weak limits. This convergence is not straightforward since the first and second terms are not independent. Let $S_{n,k}$ and S_k be as in (75), and define

$$R_n = \sqrt{n} \left(\text{SW}_p^p(P_n, Q) - \mathbb{E}(\text{SW}_p^p(P_n, Q)) \right).$$

The variable $S_{n,k}$ depends both on X_1, \dots, X_n and $\Theta_1, \dots, \Theta_k$, whereas S_k only depends on $\Theta_1, \dots, \Theta_k$ and R_n on X_1, \dots, X_k . Therefore,

$$\begin{aligned} & W_2 \left(\mathcal{L} \left(\sqrt{\frac{n}{k+n}} S_{n,k} + \sqrt{\frac{k}{k+n}} R_n \right), \sqrt{1-\tau} \cdot N(0, w_{P,Q}^2) + \sqrt{\tau} \cdot N(0, v_{P,Q}^2) \right) \\ & \leq W_2 \left(\mathcal{L} \left(\sqrt{\frac{n}{k+n}} S_{n,k} + \sqrt{\frac{k}{k+n}} R_n \right), \mathcal{L} \left(\sqrt{\frac{n}{k+n}} S_k + \sqrt{\frac{k}{k+n}} R_n \right) \right) \\ & + W_2 \left(\mathcal{L} \left(\sqrt{\frac{n}{k+n}} S_k + \sqrt{\frac{k}{k+n}} R_n \right), \sqrt{1-\tau} \cdot N(0, w_{P,Q}^2) + \sqrt{\tau} \cdot N(0, v_{P,Q}^2) \right) \end{aligned}$$

The first term is bounded by $W_p^p(\mathcal{L}(S_{n,k}), \mathcal{L}(S_k))$, which converges to 0 as shown in the proof of Proposition 5.1. For the second term, Proposition 5.1 ensures that $W_2(\mathcal{L}(S_k), N(0, w_{P,Q}^2)) \rightarrow 0$, while Theorem 3.3 demonstrates that $W_2(\mathcal{L}(R_n), N(0, v_{P,Q}^2)) \rightarrow 0$. Since S_k and R_n are independent, it is easy to conclude considering the coupling (S_k, U, R_n, V) , where $\mathcal{L}(U) = N(0, w_{P,Q}^2)$, $\mathcal{L}(V) = N(0, v_{P,Q}^2)$, (S_k, U) is the optimal coupling for $\mathcal{L}(S_k), N(0, w_{P,Q}^2)$, (R_n, V) optimal coupling for $\mathcal{L}(S_k), N(0, w_{P,Q}^2)$, and (S_k, U) and (R_n, V) are independent. \square

E Proofs of Section 6

E.1 Proof of Proposition 6.1

Consider the decomposition

$$\hat{w}_{P_n, Q}^2 - w_{P, Q}^2 = \frac{1}{k} \sum_{i=1}^k W_p^{2p}(P_n^{\Theta_i}, Q^{\Theta_i}) - \frac{1}{k} \sum_{i=1}^k W_p^{2p}(P^{\Theta_i}, Q^{\Theta_i}) \quad (77)$$

$$+ \frac{1}{k} \sum_{i=1}^k W_p^{2p}(P^{\Theta_i}, Q^{\Theta_i}) - \int_{\mathbb{S}^{d-1}} W_p^{2p}(P^\theta, Q^\theta) d\sigma(\theta) \quad (78)$$

$$- \left(\frac{1}{k} \sum_{i=1}^k W_p^{2p}(P_n^{\Theta_i}, Q^{\Theta_i}) \right)^2 + \left(\int_{\mathbb{S}^{d-1}} W_p^{2p}(P^\theta, Q^\theta) d\sigma(\theta) \right)^2 \quad (79)$$

We will show convergence in probability to 0 of each term. For (78), this follows from a straightforward application of the strong law of large numbers. For (77), the convergence follows from Markov's inequality and

$$\mathbb{E} \left| \frac{1}{k} \sum_{i=1}^k W_p^{2p}(P_n^{\Theta_i}, Q^{\Theta_i}) - \frac{1}{k} \sum_{i=1}^k W_p^{2p}(P^{\Theta_i}, Q^{\Theta_i}) \right| \leq \mathbb{E} \left(\int_{\mathbb{S}^{d-1}} \left| W_p^{2p}(P_n^\theta, Q^\theta) - W_p^{2p}(P^\theta, Q^\theta) \right| d\sigma(\theta) \right) \rightarrow 0,$$

where the convergence follows from the same arguments as in the proof of Proposition 5.1. Similarly, we can prove

$$\frac{1}{k} \sum_{i=1}^k W_p^{2p}(P_n^{\Theta_i}, Q^{\Theta_i}) - \frac{1}{k} \sum_{i=1}^k W_p^{2p}(P^{\Theta_i}, Q^{\Theta_i}) \rightarrow_p 0,$$

and, by the strong law of large numbers,

$$\frac{1}{k} \sum_{i=1}^k W_p^{2p}(P^{\Theta_i}, Q^{\Theta_i}) \rightarrow_{a.s.} \int_{\mathbb{S}^{d-1}} W_p^{2p}(P^\theta, Q^\theta) d\sigma(\theta)$$

Slutski's lemma and continuous mapping theorem allow us to conclude convergence in probability to 0 of (79). \square

E.2 Proof of Proposition 6.2

For each $\theta \in \mathbb{S}^{d-1}$, given any c -concave optimal potentials $\phi_n^\theta, \phi^\theta$, let $\phi_{n, X_1}^\theta, \phi_{X_1}^\theta$ be the random optimal potentials fixed at X_1 given by (12) and (13), which are well-defined \mathbb{P} -almost surely. As in previous proofs, we can assume without loss of generality that they are well-defined everywhere. In addition, denote by $\phi_{x_0}^\theta$ the optimal potential fixed at $x_0 \in \text{int}(\text{supp}(P))$ defined by (11). By Corollary 2.3, we can assume (18) for $q = 2(1 + \gamma)$. Since the covariance is invariant by the addition of a constant, for each $\theta, \eta \in \mathbb{S}^{d-1}$,

$$\begin{aligned} \text{Cov}_{P_n}(\phi_n^\theta, \phi_n^\eta) &= \text{Cov}_{P_n}(\phi_{n, X_1}^\theta, \phi_{n, X_1}^\eta) \\ \text{Cov}_{P_n}(\phi_{X_1}^\theta, \phi_{X_1}^\eta) &= \text{Cov}_{P_n}(\phi_{x_0}^\theta, \phi_{x_0}^\eta) \\ \text{Cov}_P(\phi_{x_0}^\theta, \phi_{x_0}^\eta) &= \text{Cov}_P(\phi^\theta, \phi^\eta). \end{aligned}$$

To simplify the notation, given any optimal transport potential ϕ^θ , we denote $\phi^\theta(x) = \phi^\theta(\langle \theta, x \rangle)$, with a slight abuse of notation. With this convention,

$$\begin{aligned} \hat{v}_{P_n, Q}^2 - v_{P, Q}^2 &= \frac{1}{k^2} \sum_{i, j=1}^k \text{Cov}_{P_n}(\phi_{n, X_1}^{\Theta_i}, \phi_{n, X_1}^{\Theta_j}) - \frac{1}{k^2} \sum_{i, j=1}^k \text{Cov}_{P_n}(\phi_{X_1}^{\Theta_i}, \phi_{X_1}^{\Theta_j}) + \\ &+ \frac{1}{k^2} \sum_{i, j=1}^k \text{Cov}_{P_n}(\phi_{x_0}^{\Theta_i}, \phi_{x_0}^{\Theta_j}) - \int_{\mathbb{S}^{d-1}} \int_{\mathbb{S}^{d-1}} \text{Cov}_P(\phi_{x_0}^\theta, \phi_{x_0}^\eta) d\sigma(\theta) d\sigma(\eta) \\ &= \frac{1}{k^2 n} \sum_{i, j=1}^k \sum_{l=1}^n (\phi_{n, X_1}^{\Theta_i}(X_l) \phi_{n, X_1}^{\Theta_j}(X_l) - \phi_{X_1}^{\Theta_i}(X_l) \phi_{X_1}^{\Theta_j}(X_l)) \end{aligned} \quad (\text{A})$$

$$- \left(\left(\frac{1}{kn} \sum_{i=1}^k \sum_{l=1}^n \phi_{n, X_1}^{\Theta_i}(X_l) \right)^2 - \left(\frac{1}{kn} \sum_{i=1}^k \sum_{l=1}^n \phi_{X_1}^{\Theta_i}(X_l) \right)^2 \right) \quad (\text{B})$$

$$+ \frac{1}{k^2 n} \sum_{i, j=1}^k \sum_{l=1}^n \phi_{x_0}^{\Theta_i}(X_l) \phi_{x_0}^{\Theta_j}(X_l) - \int_{\mathbb{S}^{d-1}} \int_{\mathbb{S}^{d-1}} \int_{\mathbb{R}^d} (\phi_{x_0}^\theta(x) \phi_{x_0}^\eta(x)) dP(x) d\sigma(\theta) d\sigma(\eta) \quad (\text{C})$$

$$- \left(\left(\frac{1}{kn} \sum_{i=1}^k \sum_{l=1}^n \phi_{x_0}^{\Theta_i}(X_l) \right)^2 - \left(\int_{\mathbb{S}^{d-1}} \int_{\mathbb{R}^d} \phi_{x_0}^\theta(x) dP(x) d\sigma(\theta) \right)^2 \right) \quad (\text{D})$$

We will prove convergence in probability to 0 of each term in the decomposition. First, we start with (D). From the continuous mapping theorem, it suffices to see that

$$\frac{1}{kn} \sum_{i=1}^k \sum_{l=1}^n \phi_{x_0}^{\Theta_i}(X_l) \rightarrow_p \int_{\mathbb{S}^{d-1}} \int_{\mathbb{R}^d} \phi_{x_0}^\theta(x) dP(x) d\sigma(\theta).$$

Consider the function $h(\theta, x) = \phi_{x_0}^\theta(x)$. Since $\Theta_1, \dots, \Theta_k$ and X_1, \dots, X_n are independent sequences, the problem can be reformulated in terms of two sample U-statistics (also known as generalized U-statistics). The above convergence can be rewritten as

$$\frac{1}{kn} \sum_{i=1}^k \sum_{l=1}^n h(\Theta_i, X_l) \rightarrow_p \mathbb{E}(h(\Theta_1, X_1)).$$

In fact, existing versions of the strong law of large numbers for generalized U-statistics allow us to prove convergence almost surely. If n and k are of the same order, i.e. $n/(n+k) \rightarrow \tau \in (0, 1)$, Theorem 3.1.2. in [KB13] proves a.s. convergence provided that $\mathbb{E}(|h(\Theta_1, X_1)|) < \infty$. Assuming only $n, k \rightarrow \infty$, [Sen77] proves a.s. convergence under the stronger assumption $\mathbb{E}(|h(\Theta_1, X_1)| \log_+(|h(\Theta_1, X_1)|)) < \infty$. Both conditions are verified, since

$$\mathbb{E}(|h(\Theta_1, X_1)|^{2(1+\gamma)}) = \|\phi_{x_0}^\theta\|_{L^{2(1+\gamma)}(\sigma \times P)}^{2(1+\gamma)} < \infty. \quad (80)$$

For the term (C), consider the following decomposition

$$\frac{1}{k^2 n} \sum_{i, j=1}^k \sum_{l=1}^n \phi_{x_0}^{\Theta_i}(X_l) \phi_{x_0}^{\Theta_j}(X_l) =$$

$$= \frac{1}{k} \left(\frac{1}{kn} \sum_{i=1}^k \sum_{l=1}^n \phi_{x_0}^{\Theta_i}(X_l)^2 \right) + \frac{2 \binom{k}{2}}{k^2} \left(\frac{1}{\binom{n}{1} \binom{k}{2}} \sum_{1 \leq i < j \leq k} \sum_{l=1}^n \phi_{x_0}^{\Theta_i}(X_l) \phi_{x_0}^{\Theta_j}(X_l) \right)$$

The bound (80), and the same reasoning as before, allow us to conclude that the term inside the first parenthesis converges a.s. to $\mathbb{E}(h(\Theta_1, X_1)^2)$. Since it is multiplied by a factor $1/k \rightarrow 0$, the first term converges a.s. to 0. Now, given that $2 \binom{k}{2} / k^2 = (k-1)/k \rightarrow 1$, if we denote $g(\theta, \theta', x) = \phi_{x_0}^\theta(x) \phi_{x_0}^{\theta'}(x)$, it suffices to prove

$$\frac{1}{\binom{n}{1} \binom{k}{2}} \sum_{1 \leq i < j \leq k} \sum_{l=1}^n g(\Theta_i, \Theta_j, X_l) \rightarrow_p \mathbb{E}(g(\Theta_1, \Theta_2, X_1))$$

to conclude (C). Given that $g(\theta, \theta', x)$ is symmetric with respect to θ, θ' , we can apply again strong law of large numbers for two sample U-statistics to conclude almost sure convergence, since

$$\begin{aligned} \mathbb{E}\left(|g(\Theta_1, \Theta_2, X_1)|^{1+\gamma}\right) &\leq \mathbb{E}\left(|\phi_{x_0}^{\Theta_1}(X_1)|^{2(1+\gamma)}\right)^{1/2} \mathbb{E}\left(|\phi_{x_0}^{\Theta_2}(X_1)|^{2(1+\gamma)}\right)^{1/2} \\ &= \mathbb{E}\left(|\phi_{x_0}^{\Theta_1}(X_1)|^{2(1+\gamma)}\right) = \|\phi_{x_0}^\theta\|_{L^{2(1+\gamma)}(\sigma \times P)}^{2(1+\gamma)} < \infty. \end{aligned}$$

Next, to show the convergence of (B), we will prove the convergence in mean

$$\mathbb{E}\left|\frac{1}{kn} \sum_{i=1}^k \sum_{l=1}^n \phi_{n, X_1}^{\theta_i}(X_l) - \frac{1}{kn} \sum_{i=1}^k \sum_{l=1}^n \phi_{X_1}^{\theta_i}(X_l)\right| \rightarrow 0. \quad (81)$$

By Markov's inequality, this implies convergence in probability

$$\frac{1}{kn} \sum_{i=1}^k \sum_{l=1}^n \phi_{n, X_1}^{\Theta_i}(X_l) - \frac{1}{kn} \sum_{i=1}^k \sum_{l=1}^n \phi_{X_1}^{\Theta_i}(X_l) \xrightarrow{p} 0. \quad (82)$$

Proposition 2.2 guarantees the existence of a constant $M > 0$ such that

$$\begin{aligned} \mathbb{E}\left|\frac{1}{kn} \sum_{i=1}^k \sum_{l=1}^n \phi_{n, X_1}^{\Theta_i}(X_l)\right| &\leq \mathbb{E}\left(\|\phi_{n, X_1}\|_{L_1(\sigma \times P_n)}\right) \leq M \quad \forall n \in \mathbb{N}, \\ \mathbb{E}\left|\frac{1}{kn} \sum_{i=1}^k \sum_{l=1}^n \phi_{X_1}^{\Theta_i}(X_l)\right| &\leq \frac{n-1}{n} \mathbb{E}\left(\|\phi_{X_1}\|_{L_1(\sigma \times P)}\right) \leq M. \end{aligned}$$

Therefore, it follows that the sequence

$$\frac{1}{kn} \sum_{i=1}^k \sum_{l=1}^n \phi_{n, X_1}^{\Theta_i}(X_l) + \frac{1}{kn} \sum_{i=1}^k \sum_{l=1}^n \phi_{X_1}^{\Theta_i}(X_l) \quad (83)$$

is bounded in probability. Multiplying (82) and (83), we can conclude convergence in probability of the sequence in (B) to 0. To prove (81), note that

$$\begin{aligned} \phi_{n, X_1}^\theta(X_1) &= 0 \quad , \quad \phi_{n, X_1}^\theta(X_2) = \dots = \phi_{n, X_1}^\theta(X_n) \\ \phi_{X_1}^\theta(X_1) &= 0 \quad , \quad \phi_{X_1}^\theta(X_2) = \dots = \phi_{X_1}^\theta(X_n) \end{aligned}$$

Therefore, (81) can be bounded by

$$\begin{aligned} \frac{1}{kn} \mathbb{E}\left(\sum_{i=1}^k \sum_{l=1}^n |\phi_{n, X_1}^{\Theta_i}(X_l) - \phi_{X_1}^{\Theta_i}(X_l)|\right) &= \frac{1}{n} \mathbb{E}\left(\sum_{l=1}^n |\phi_{n, X_1}^{\Theta_1}(X_l) - \phi_{X_1}^{\Theta_1}(X_l)|\right) \\ &= \frac{n-1}{n} \mathbb{E}\left(|\phi_{n, X_1}^{\Theta_1}(X_2) - \phi_{X_1}^{\Theta_1}(X_2)|\right) \end{aligned}$$

By property (4) in Proposition 2.1, we know that $|\phi_{n, X_1}^{\Theta_1}(X_2) - \phi_{X_1}^{\Theta_1}(X_2)| \rightarrow 0$ \mathbb{P} -a.s. Uniform integrability follows from Proposition 2.2, since

$$\mathbb{E}\left(|\phi_{n, X_1}^{\Theta_1}(X_2) - \phi_{X_1}^{\Theta_1}(X_2)|^{1+\gamma}\right) = \mathbb{E}\left(\int_{\mathbb{S}^{d-1}} \frac{1}{n-1} \sum_{l=2}^n |\phi_{n, X_1}^\theta(X_l) - \phi_{X_1}^\theta(X_l)|^{1+\gamma} d\sigma(\theta)\right)$$

$$\leq C_{1+\gamma} \left(\frac{n}{n-1} \|\phi_{n,X_1}^\theta\|_{L^{1+\gamma}(\sigma \times P_n)}^{1+\gamma} + \|\phi_{X_1}^\theta\|_{L^{1+\gamma}(\sigma \times P)}^{1+\gamma} \right) < \infty. \quad (84)$$

A similar conclusion is reached for (A). To prove convergence in mean,

$$\begin{aligned} & \mathbb{E} \left| \frac{1}{k^2 n} \sum_{i,j=1}^k \sum_{l=1}^n \left(\phi_{n,X_1}^{\Theta_i}(X_l) \phi_{n,X_1}^{\Theta_j}(X_l) - \phi_{X_1}^{\Theta_i}(X_l) \phi_{X_1}^{\Theta_j}(X_l) \right) \right| \\ & \leq \frac{2}{k^2 n} \sum_{1 \leq i < j \leq k} \sum_{l=1}^n \mathbb{E} \left| \phi_{n,X_1}^{\Theta_i}(X_l) \phi_{n,X_1}^{\Theta_j}(X_l) - \phi_{X_1}^{\Theta_i}(X_l) \phi_{X_1}^{\Theta_j}(X_l) \right| \\ & \quad + \frac{1}{k} \frac{1}{kn} \sum_{i=1}^k \sum_{l=1}^n \mathbb{E} \left| \phi_{n,X_1}^{\Theta_i}(X_l)^2 - \phi_{X_1}^{\Theta_i}(X_l)^2 \right| \\ & = \frac{(k-1)(n-1)}{kn} \mathbb{E} \left| \phi_{n,X_1}^{\Theta_1}(X_2) \phi_{n,X_1}^{\Theta_2}(X_2) - \phi_{X_1}^{\Theta_1}(X_2) \phi_{X_1}^{\Theta_2}(X_2) \right| \\ & \quad + \frac{1}{k} \frac{n-1}{n} \mathbb{E} \left| \phi_{n,X_1}^{\Theta_1}(X_2)^2 - \phi_{X_1}^{\Theta_1}(X_2)^2 \right| \end{aligned}$$

By Proposition 2.1, $\phi_{n,X_1}^{\Theta_1}(X_2) - \phi_{X_1}^{\Theta_1}(X_2)$ \mathbb{P} -a.s. Therefore, we have pointwise convergence of both sequences inside the expectations, and we only need to prove uniform integrability. For the first sequence, it follows from

$$\begin{aligned} & \mathbb{E} \left(\left| \phi_{n,X_1}^{\Theta_1}(X_2) \phi_{n,X_1}^{\Theta_2}(X_2) - \phi_{X_1}^{\Theta_1}(X_2) \phi_{X_1}^{\Theta_2}(X_2) \right|^{1+\gamma} \right) \\ & \leq C_{1+\gamma} \left(\mathbb{E} \left(\left| \phi_{n,X_1}^{\Theta_1}(X_2) \phi_{n,X_1}^{\Theta_2}(X_2) \right|^{1+\gamma} \right) + \mathbb{E} \left(\left| \phi_{X_1}^{\Theta_1}(X_2) \phi_{X_1}^{\Theta_2}(X_2) \right|^{1+\gamma} \right) \right) \\ & \leq C_{1+\gamma} \left(\mathbb{E} \left(\left| \phi_{n,X_1}^{\Theta_1}(X_2) \right|^{2(1+\gamma)} \right) + \mathbb{E} \left(\left| \phi_{X_1}^{\Theta_1}(X_2) \right|^{2(1+\gamma)} \right) \right) \\ & = C_{1+\gamma} \left(\frac{n-1}{n} \|\phi_{n,X_1}^\theta\|_{L^{2(1+\gamma)}(\sigma \times P_n)}^{2(1+\gamma)} + \|\phi_{X_1}^\theta\|_{L^{2(1+\gamma)}(\sigma \times P)}^{2(1+\gamma)} \right) < \infty, \end{aligned}$$

An analogous argument proves uniform integrability of the second sequence, which allow us to finish the proof.

Two-sample setting: By symmetry, it suffices to see $\hat{v}_{P_n, Q_m}^2 \rightarrow_P v_{P, Q}^2$. The proof follows from the same arguments, since the uniform convergence and uniform integrability properties in the above proof also apply to the optimal potentials ϕ_{n,m,X_1}^θ , defined by (14). \square

F Efficient estimation of the asymptotic variance

In this section, we detail the computations required to obtain the value of $SW_{p,k}^p(P_n, Q_m)$ and the variance estimators \hat{w}_{P_n, Q_m}^2 , \hat{v}_{P_n, Q_m}^2 and \hat{v}_{Q_m, P_n}^2 in the two-sample setting. We restrict our experimental setting to the prominent case $p = 2$, where the c -concave optimal transport potentials can be efficiently estimated using the relationship between c -concavity and Legendre duality. For general $p > 1$, different strategies may be adopted to find the c -concave optimal potentials that solve the dual problem. However, it remains unclear whether a straightforward solution exists that can be efficiently computed based on the one-dimensional optimal transport plan, as in the case of $p = 2$.

Let $x_1, \dots, x_n, y_1, \dots, y_m \in \mathbb{R}^d$ and $\theta_1, \dots, \theta_k \in \mathbb{S}^{d-1}$ be the given samples. We begin by computing the projected samples

$$\begin{aligned} s_i^l &= \langle \theta_l, X_i \rangle \quad i = 1, \dots, n, \quad l = 1, \dots, k, \\ t_j^l &= \langle \theta_l, Y_j \rangle \quad j = 1, \dots, m, \quad l = 1, \dots, k. \end{aligned}$$

Let $\{s_{(i)}^l\}_{i=1}^n$ and $\{t_{(j)}^l\}_{j=1}^m$ be the values of the ordered statistic. For each $l = 1, \dots, k$ we need to compute

$$W_2^2(P_n^{\theta_l}, Q_m^{\theta_l}) = \int_0^1 \left(F_{\theta_l, n}^{-1}(u) - G_{\theta_l, m}^{-1}(u) \right)^2 du = \sum_{i=1}^n \sum_{j=1}^m R_{i,j} (s_{(i)}^l - t_{(j)}^l)^2,$$

where $R_{i,j} = m \left(\left(\frac{i-1}{n}, \frac{i}{n} \right] \cap \left(\frac{j-1}{m}, \frac{j}{m} \right] \right)$. The matrix $R = \{R_{i,j}\}_{i,j}$ only depends on n and m , it must be computed just once at the beginning. Once we have the values of $\{W_2^2(P_n^{\theta_l}, Q_m^{\theta_l})\}_{l=1}^k$, we can compute $SW_{2,k}^2(P_n, Q_m)$ as in (27),

and

$$\hat{w}_{P_n, Q_m}^2 = \frac{1}{k} \sum_{l=1}^k \mathbf{W}_2^4(P_n^{\theta_l}, Q_m^{\theta_l}) - \mathbf{SW}_{p,k}^{2p}(P_n, Q_m).$$

To compute \hat{v}_{P_n, Q_m}^2 , we can reuse part of the previous computations. Our approach is based on the characterization of optimal transport plans in terms of lower semicontinuous convex functions for the cost $c = c_2$ (see [Vil03] for details). In this case, the c -concave optimal potentials between $P_n^{\theta_l}$ and $Q_m^{\theta_l}$ can be expressed as $\phi_{n,m}^{\theta_l}(x) = x^2 - 2\varphi_{n,m}^{\theta_l}(x)$, where $\varphi_{n,m}^{\theta_l}$ is convex and lower semicontinuous. Given an optimal transport map $\pi_{n,m}^{\theta_l}$, if $\text{supp}(\pi_{n,m}^{\theta_l}) \subset \partial^c \phi_{n,m}^{\theta_l} = \partial \varphi_{n,m}^{\theta_l}$, where ∂ denotes the subgradient, then $\phi_{n,m}^{\theta_l}$ is optimal. Therefore, we need to build $\varphi_{n,m}^{\theta_l}$, convex and lower semicontinuous verifying $\text{supp}(\pi_{n,m}^{\theta_l}) \subset \partial \varphi_{n,m}^{\theta_l}$. To do this, define

$$r(i) = \max\{j = 1, \dots, m : R_{i,j} > 0\} \quad i = 1, \dots, n.$$

This choice of $\varphi_{n,m}^{\theta_l}$ is not unique. We consider the only piecewise linear continuous function such that $\varphi_{n,m}^{\theta_l}(s_{(1)}^l) = 0$ and

$$(\varphi_{n,m}^{\theta_l})'(s) = \begin{cases} t_{(1)}^l & \text{if } s < s_{(1)}^l \\ t_{(r^l(i))}^l & \text{if } s_{(i)}^l < s < s_{(i+1)}^l \\ t_{(m)}^l & \text{if } s > s_{(n)}^l \end{cases}$$

To compute (31), we are only interested in the values of $\varphi_{n,m}^{\theta_l}$ at the points $\{s_{(i)}^l\}_{i=1}^n$. Since we are imposing $\varphi_{n,m}^{\theta_l}(s_{(1)}^l) = 0$, we can iteratively compute

$$\varphi_{n,m}^{\theta_l}(s_{(i+1)}^l) = \varphi_{n,m}^{\theta_l}(s_{(i)}^l) + t_{(r^l(i))}^l (s_{(i+1)}^l - s_{(i)}^l), \quad i = 1, \dots, n-1.$$

Finally, once we have computed all these values, we can obtain the values $\phi_{n,m}^{\theta_l}(s_{(i)}^l)$, which allows us to compute \hat{v}_{P_n, Q_m}^2 as in (31). Symmetric computations allow us to obtain \hat{v}_{Q_m, P_n}^2 .