

On automatic boundedness of some operators in ordered Banach spaces

March 25, 2025

Eduard Emelyanov¹
¹ Sobolev Institute of Mathematics

Abstract

We study order-to-weak continuous operators from an ordered Banach space to a normed space. It is proved that under rather mild conditions every order-to-weak continuous operator is bounded.

Keywords: ordered Banach space, order-to-norm bounded operator, order-to-norm continuous operator, Lebesgue operator

MSC2020: 46A40, 46B42, 47B65

1 Introduction and preliminaries

Order-to-topology continuous and τ -Lebesgue operators between vector lattices and topological vector spaces were studied recently in [3, 8, 5, 9, 10]. We adopt the definition to the setting of ordered vector spaces in the domain and normed spaces in the range. It is important to find conditions on operators in ordered Banach spaces which provide automatic boundedness (see, for example [1, 2, 4] and the references therein). In the present paper, we continue the search of such conditions following [6, 7].

We abbreviate ordered vector (normed, ordered vector, ordered Banach) spaces by OVSs (NSs, ONSs, OBSs) and vector (normed, Banach) lattices by VLs (NLs, BLs). Throughout the present paper, vector spaces are real, operators are linear, symbol $\mathcal{L}(X, Y)$ stands for the space of operators from a vector space X to a vector space Y , $\mathcal{L}_{ob}(X, Y)$ for the space of order bounded operators from an OVS X to an OVS Y , $L(X, Y)$ for the space of bounded operators from a NS X to a NS Y , $x_\alpha \downarrow 0$ for a decreasing net in an OVS such that $\inf_\alpha x_\alpha = 0$, and B_X for the closed unit ball of a NS X . A net (x_α) in an OVS X

- *order converges* to $x \in X$ (o-converges to x , or $x_\alpha \overset{o}{\rightarrow} x$) whenever there exists $g_\beta \downarrow 0$ in X such that, for each β there is α_β satisfying $x_\alpha - x \leq \pm g_\beta$ for all $\alpha \geq \alpha_\beta$.
- *relative uniform converges* to $x \in X$ (ru-converges to x , or $x_\alpha \overset{ru}{\rightarrow} x$) if, for some $u \in X_+$ there exists an increasing sequence (α_n) of indices with $x_\alpha - x \leq \pm \frac{1}{n}u$ for all $\alpha \geq \alpha_n$.

An operator T from an OVS X to an OVS Y is *order continuous* if $Tx_\alpha \overset{o}{\rightarrow} 0$ in Y whenever $x_\alpha \overset{o}{\rightarrow} 0$ in X (the set of such operators is a vector space which is denoted by $\mathcal{L}_{oc}(X, Y)$).

Definition 1.1. *An operator T from an OVS X to a NS Y is*

- a) *(σ -)Lebesgue if $Tx_\alpha \overset{\|\cdot\|}{\rightarrow} 0$ for every (sequence) net $x_\alpha \downarrow 0$ in X [3, Definition 1.1]. The set of such operators is denoted by $L_{Leb}(X, Y)$ ($L_{Leb}^\sigma(X, Y)$).*

- b) $(\sigma\text{-})w\text{-Lebesgue}$ if $Tx_\alpha \xrightarrow{w} 0$ for every (sequence) net $x_\alpha \downarrow 0$ in X [3, Definition 1.1]. The set of such operators is denoted by $L_{wLeb}(X, Y)$ ($L_{wLeb}^\sigma(X, Y)$).
- c) $(\sigma\text{-})\text{order-to-norm continuous}$ if $Tx_\alpha \xrightarrow{\|\cdot\|} 0$ for every (sequence) net $x_\alpha \xrightarrow{o} 0$ in X [5]. The set of such operators is denoted by $L_{onc}(X, Y)$ ($L_{onc}^\sigma(X, Y)$).
- d) $(\sigma\text{-})\text{order-to-weak continuous}$ if $Tx_\alpha \xrightarrow{w} 0$ for every (sequence) net $x_\alpha \xrightarrow{o} 0$ in X . The set of such operators is denoted by $L_{owc}(X, Y)$ ($L_{owc}^\sigma(X, Y)$) [5].
- e) $\text{order-to-norm bounded}$ if $T[a, b]$ is norm bounded for every order interval $[a, b]$ in X [7, Definition 1.1]. The set of such operators is denoted by $L_{onb}(X, Y)$.

A positive operator T from a VL to a NL is Lebesgue/ σ -Lebesgue iff T is order-to-norm continuous/ σ -order-to-norm continuous by [3, Lemma 2.1] (see, also [9, Proposition 1.5]). The same fact holds under slightly more general assumptions.

Lemma 1.2. *Let X be a OVS, Y a normal ONS, and $T \in \mathcal{L}_+(X, Y)$. Then*

$$T \in L_{Leb}(X, Y) \iff T \in L_{onc}(X, Y) \quad \text{and} \quad T \in L_{Leb}^\sigma(X, Y) \iff T \in L_{onc}^\sigma(X, Y).$$

Proof. As the σ -Lebesgue case is similar, we consider only the first equivalence. It suffices to show $L_{Leb}(X, Y) \subseteq L_{onc}(X, Y)$. Suppose $T \in L_{Leb}(X, Y)$ and take $x_\alpha \xrightarrow{o} 0$ in X . Pick a net $g_\beta \downarrow 0$ in X such that, for each β , there exists α_β with $x_\alpha \leq \pm g_\beta$ for $\alpha \geq \alpha_\beta$. Since $T \geq 0$, we obtain $Tx_\alpha \leq \pm Tg_\beta$ for $\alpha \geq \alpha_\beta$. As T is Lebesgue, $\|Tg_\beta\| \rightarrow 0$. So, since $Tg_\beta \downarrow \geq 0$ then $Tg_\beta \downarrow 0$. It follows from [2, Theorem 2.23] that $\|Tx_\alpha\| \rightarrow 0$. Thus, $T \in L_{onc}(X, Y)$. \square

By [3, Proposition 2.2], a positive order continuous operator T from a VL to a NL is Lebesgue iff T is w -Lebesgue. The next lemma extends this fact to the setting of OVSs.

Lemma 1.3. *Let X be a OVS, Y a normal ONS, and $T \in (\mathcal{L}_{oc})_+(X, Y)$. Then*

$$T \in L_{Leb}(X, Y) \iff T \in L_{wLeb}(X, Y) \quad \text{and} \quad T \in L_{Leb}^\sigma(X, Y) \iff T \in L_{wLeb}^\sigma(X, Y).$$

Proof. We consider only the Lebesgue case. It suffices to show $L_{wLeb}(X, Y) \subseteq L_{Leb}(X, Y)$. Suppose $T \in L_{wLeb}(X, Y)$ and take $x_\alpha \downarrow 0$ in X . Since $T \geq 0$ then $Tx_\alpha \downarrow \geq 0$, and as T is w -Lebesgue, $Tx_\alpha \xrightarrow{w} 0$. By [2, Lemma 2.28], we conclude $\|Tx_\alpha\| \rightarrow 0$. Thus, $T \in L_{Leb}(X, Y)$. \square

The following lemma is a direct consequence of Definition 1.1.

Lemma 1.4. *Let X be an OVS and Y a NS. Then*

$$\begin{aligned} L_{Leb}(X, Y) &\subseteq L_{wLeb}(X, Y), \quad L_{Leb}^\sigma(X, Y) \subseteq L_{wLeb}^\sigma(X, Y), \\ L_{onc}(X, Y) &\subseteq L_{owc}(X, Y), \quad L_{onc}^\sigma(X, Y) \subseteq L_{owc}^\sigma(X, Y), \\ L_{onc}(X, Y) &\subseteq L_{Leb}(X, Y) \cap L_{onc}^\sigma(X, Y) \subseteq L_{Leb}(X, Y) \cup L_{onc}^\sigma(X, Y) \subseteq L_{Leb}^\sigma(X, Y), \quad \text{and} \\ L_{owc}(X, Y) &\subseteq L_{wLeb}(X, Y) \cap L_{owc}^\sigma(X, Y) \subseteq L_{wLeb}(X, Y) \cup L_{owc}^\sigma(X, Y) \subseteq L_{wLeb}^\sigma(X, Y). \end{aligned}$$

Proposition 1.5. *Let X be an OBS with a closed generating normal cone and Y a NS. Then $L_{onb}(X, Y) = L(X, Y)$.*

Proof. The inclusion $L_{onb}(X, Y) \subseteq L(X, Y)$ follows from [7, Corollary 2.9]. The reverse inclusion holds due to normality of X . \square

The present note is organized as follows. Lemma 2.1 states that every σ - w -Lebesgue operator is order-to-norm bounded whenever OBS in the domain has normal cone. Theorem 2.2 states that if the domain has closed generating normal cone then every σ - w -Lebesgue operator is bounded. Theorem 2.6 gives conditions under which $(\sigma\text{-})(w\text{-})$ Lebesgue operators are $(\sigma\text{-})$ order to (weak) norm continuous.

For the terminology and notations that are not explained in the text, we refer to [1, 2].

2 Main results

It is established in [7, Theorem 2.8] that every order-to-norm bounded operator from an OBS with closed generating cone to a NS is bounded (see also [6, Theorem 2.1] for the Banach lattice case). Accordingly to [7, Theorem 2.4], each order bounded operator from an OVS to an OVS with generating cones is ru-continuous. Combining these two results and keeping in mind that order intervals in an OBS with normal cone are bounded, we obtain the following automatic boundedness conditions of order bounded operators:

(*) $\mathcal{L}_{ob}(X, Y) \subseteq L(X, Y)$ whenever X and Y are OBSs with generating cones such that X_+ is closed and Y_+ is normal.

Another case of automatic boundedness of operators concerns the order continuity. Namely, as every order continuous operator from an Archimedean OVS with generating cone to an OVS is order bounded [7, Corollary 2.2], we have:

(**) $\mathcal{L}_{oc}(X, Y) \subseteq L(X, Y)$ whenever X and Y are OBSs with generating cones such that X_+ is closed and Y_+ is normal.

Condition $X_+ - X_+ = X$ is essential in both (*) and (**) since each operator $T : X \rightarrow X$ is order bounded and order continuous whenever $X_+ = \{0\}$. An inspection of [7, Example 2.12 b)] tells us that each infinite dimensional Banach space X has a non-closed generating Archimedean cone so that every operator $T : X \rightarrow X$ is order continuous (and hence order bounded by [7, Corollary 2.2]).

It is worth nothing that the order-to-norm and order-to-weak continuous operators on VLs were defined in [5] without specifying the class of (just linear or continuous linear) operators. Automatic boundedness of such operators was investigated in recent paper [9]. In this section we establish automatic boundedness conditions for order-to-weak continuous operators on OVSs. More precisely, Theorem 2.2 which tells us that every σ -w-Lebesgue operator from an OBS with a closed generating normal cone to a NS is bounded.

We begin with conditions under which σ -w-Lebesgue operators are order-to-norm bounded.

Lemma 2.1. *Let X be a normal OBS and Y a NS. Then $L_{wLeb}^\sigma(X, Y) \subseteq L_{onb}(X, Y)$. In particular, $L_{owc}^\sigma(X, Y) \subseteq L_{onb}(X, Y)$.*

Proof. Let $T : X \rightarrow Y$ be σ -w-Lebesgue. If T is not order-to-norm bounded then $T[0, u]$ is not bounded for some $u \in X_+$. Since X is normal then $[0, u]$ is bounded, say $\sup_{x \in [0, u]} \|x\| \leq M$. Pick

a sequence (u_n) in $[0, u]$ with $\|Tu_n\| \geq n2^n$, and set $y_n := \|\cdot\| - \sum_{k=n}^{\infty} 2^{-k}u_k$ for $n \in \mathbb{N}$. Then $y_n \downarrow \geq 0$. Let $0 \leq y_0 \leq y_n$ for all $n \in \mathbb{N}$. Since $0 \leq y_0 \leq 2^{1-n}u$ and $\|2^{1-n}u\| \leq M2^{1-n} \rightarrow 0$ then $y_0 = 0$ by [2, Theorem 2.23]. Thus, $y_n \downarrow 0$. It follows from $T \in L_{wLeb}^\sigma(X, Y)$ that (Ty_n) is w-null, and hence is norm bounded, that is absurd because $\|Ty_{n+1} - Ty_n\| = \|T(2^{-n}u_n)\| \geq n$ for all $n \in \mathbb{N}$.

The second inclusion follows from Lemma 1.4. □

Combining Lemma 2.1 with Proposition 1.5 and Lemma 1.4, we obtain.

Theorem 2.2. *Let X be an OBS with a closed generating normal cone and Y a NS. Then $L_{wLeb}^\sigma(E, Y) \subseteq L(E, Y)$. In particular, $L_{ows}^\sigma(E, Y) \subseteq L(E, Y)$.*

Corollary 2.3. (cf., [3, Proposition 3.2],[9, Lemma 2.3]) *Each σ -w-Lebesgue (and hence each σ -order-to-weak continuous) operator from a BL to a NS is bounded.*

The next proposition is a straightforward consequence of Lemma 1.4 and Theorem 2.2.

Proposition 2.4. *Let X be an OBS with a closed generating normal cone and Y a NS. Then $L_{Leb}(X, Y)$, $L_{Leb}^\sigma(X, Y)$, $L_{wLeb}(X, Y)$, $L_{wLeb}^\sigma(X, Y)$, $L_{onc}(X, Y)$, $L_{onc}^\sigma(X, Y)$, $L_{owc}(X, Y)$, and $L_{owc}^\sigma(X, Y)$ are norm closed subspaces of $L(X, Y)$.*

Positive order-to-norm continuous order-to-norm bounded operators between NLs are not necessary bounded (see [3, Remark 1.1.c])). From the other hand, even a positive σ -Lebesgue rank one contraction on an AM-space need not to be w-Lebesgue [3, Example 3.1]).

We say that the norm in and ONS X is (σ -)order continuous if $\|x_\alpha\| \rightarrow 0$ for every net (sequence) $x_\alpha \xrightarrow{o} 0$ in X . In certain cases o-convergence is equivalent to ru-convergence.

Lemma 2.5. Let X be an OBS with a closed normal cone.

- i) If the norm in X is order continuous then $x_\alpha \xrightarrow{o} 0 \iff x_\alpha \xrightarrow{ru} 0$ in X .
- ii) If the norm in X is σ -order continuous then $x_n \xrightarrow{o} 0 \iff x_n \xrightarrow{ru} 0$ in X .

Proof. First, we show that if X_+ is normal then $x_\alpha \xrightarrow{ru} 0 \implies x_\alpha \xrightarrow{o} 0$ in X . Suppose $x_\alpha \xrightarrow{ru} 0$. Then, for some $u \in X_+$ there exists an increasing sequence (α_n) , $-\frac{1}{n}u \leq x_\alpha \leq \frac{1}{n}u$ for all $\alpha \geq \alpha_n$. Let $0 \leq y_0 \leq \frac{1}{n}u$ for all $n \in \mathbb{N}$. Since $\|\frac{1}{n}u\| \rightarrow 0$ then $y_0 = 0$ by [2, Theorem 2.23]. It follows $\frac{1}{n}u \downarrow 0$, and hence $x_\alpha \xrightarrow{o} 0$.

i) Let X has order continuous norm and $x_\alpha \xrightarrow{o} 0$ in X . Take a net $g_\beta \downarrow 0$ in X such that, for each β there is α_β satisfying $\pm x_\alpha \leq g_\beta$ for $\alpha \geq \alpha_\beta$. Find an increasing sequence (β_k) such that $\sup_{z \in [-g_{\beta_k}, g_{\beta_k}]} \|z\| \leq k^{-3}$ for all $k \in \mathbb{N}$, and set $u = \|\cdot\| - \sum_{k=1}^{\infty} k g_{\beta_k}$. Then $\pm x_\alpha \leq g_{\beta_k} \leq k^{-1}u$ for $\alpha \geq \alpha_{\beta_k}$. Choose any increasing sequence (α_k) with $\alpha_k \geq \alpha_{\beta_1}, \alpha_{\beta_2}, \dots, \alpha_{\beta_k}$. Then $\pm x_\alpha \leq k^{-1}u$ for $\alpha \geq \alpha_k$, and hence $x_\alpha \xrightarrow{ru} 0$.

ii) The proof is similar. □

Theorem 2.6. *Let X be an OBS with a closed generating normal cone and Y a NS.*

- i) *If X has σ -order continuous norm then*

$$L_{Leb}^\sigma(X, Y) = L_{onc}^\sigma(X, Y) \quad \text{and} \quad L_{wLeb}^\sigma(X, Y) = L_{owc}^\sigma(X, Y).$$

- ii) *If X has order continuous norm then*

$$L_{Leb}(X, Y) = L_{onc}(X, Y) \quad \text{and} \quad L_{wLeb}(X, Y) = L_{owc}(X, Y).$$

Proof. We prove only i) as the proof of ii) is similar. In view of Lemma 1.4, we need to prove $L_{Leb}^\sigma(X, Y) \subseteq L_{onc}^\sigma(X, Y)$ and $L_{wLeb}^\sigma(X, Y) \subseteq L_{owc}^\sigma(X, Y)$. Let $x_n \xrightarrow{o} 0$ in X . Then, $x_n \xrightarrow{ru} 0$ by Lemma 2.5, and hence $\|x_n\| \rightarrow 0$ due to normality of X_+ .

If $T \in L_{Leb}^\sigma(X, Y)$ then $T \in L(X, Y)$ by Theorem 2.2, and hence $\|Tx_n\| \rightarrow 0$. Suppose $T \in L_{wLeb}^\sigma(X, Y)$. Then again $T \in L(X, Y)$. So $\|Tx_n\| \rightarrow 0$, and hence $Tx_n \xrightarrow{w} 0$. □

Corollary 2.7. *Let E be a BL with order continuous norm and Y a NS. Then $L_{Leb}^\sigma(E, Y) = L_{onc}^\sigma(E, Y)$, $L_{Leb}(E, Y) = L_{onc}(E, Y)$, $L_{wLeb}^\sigma(E, Y) = L_{owc}^\sigma(E, Y)$, and $L_{wLeb}(E, Y) = L_{owc}(E, Y)$.*

Corollary 2.7 is a partial extension of results of [5, Lemma 1], [3, Lemma 2.1], [10, Theorem 3], and [9, Proposition 1.5] to the OVS setting.

We give some conditions under which each order bounded operator is order-to-norm continuous.

Proposition 2.8. *Let X be an OBS with a closed normal cone and Y an ONS with a generating normal cone.*

- i) *If X has σ -order continuous norm then $\mathcal{L}_{ob}(X, Y) \subseteq L_{onc}^\sigma(X, Y)$.*
- ii) *If X has order continuous norm then $\mathcal{L}_{ob}(X, Y) \subseteq L_{onc}(X, Y)$.*

Proof. We prove only i) as the proof of ii) is similar. Let $T \in \mathcal{L}_{ob}(X, Y)$ and $x_n \xrightarrow{o} 0$ in X . By Lemma 2.5, $x_n \xrightarrow{ru} 0$. It follows from [7, Corollary 2.5] that $Tx_n \xrightarrow{ru} 0$. The normality of Y provides $\|Tx_n\| \rightarrow 0$. □

References

- [1] Aliprantis, C.D. & Burkinshaw, O. Positive operators, 2nd edition. Springer, Dordrecht, 2006.
- [2] Aliprantis, C.D. & Tourky, R. Cones and Duality. American Mathematical Society, Providence, 2007.
- [3] Alpay, S., Emelyanov, E. & Gorokhova, S. $\sigma\tau$ -Continuous, Lebesgue, KB, and Levi Operators Between Vector Lattices and Topological Vector Spaces. Results Math 77, 117 (2022).
- [4] Arendt, W. & Nittka, R. Equivalent complete norms and positivity. Arch. Math. 92, 414–427 (2009).
- [5] Jalili, S.A., Azar, K.Y. & Moghimi, M.B.F. Order-to-topology continuous operators. Positivity 25, 1313–1322, (2021).
- [6] Emelyanov, E. Collective order convergence and collectively qualified sets of operators. Siberian Electronic Mathematical Reports (to appear) arxiv.org/pdf/2408.03671
- [7] Emelyanov, E., Erkuşun-Özcan, N. & Gorokhova, S. Collective Order Boundedness of Sets of Operators Between Ordered Vector Spaces. Results Math. 80, 70 (2025).
- [8] Gorokhova, S. & Emelyanov, E. On operators dominated by the Kantorovich–Banach and Levi operators in locally solid lattices. Vlad. Math. J. 24(3), 55–61, (2022).
- [9] Keleş, C.S., Turan, B. & Altın, B. Order structure of order-to-topological continuous operator. Turkish Journal Math. 49(2), 3 (2025). <https://doi.org/10.55730/1300-0098.3581>
- [10] Zhang, F., Shen, H. & Chen, Z. Property (h) of Banach Lattice and Order-to-Norm Continuous Operators. Mathematics 11, 2747, (2023).