

Accelerating Langevin Monte Carlo Sampling: A Large Deviations Analysis

Nian Yao¹, Pervez Ali², Xihua Tao³, Lingjiong Zhu⁴

March 26, 2025

Abstract

Langevin algorithms are popular Markov chain Monte Carlo methods that are often used to solve high-dimensional large-scale sampling problems in machine learning. The most classical Langevin Monte Carlo algorithm is based on the overdamped Langevin dynamics. There are many variants of Langevin dynamics that often show superior performance in practice. In this paper, we provide a unified approach to study the acceleration of the variants of the overdamped Langevin dynamics through the lens of large deviations theory. Numerical experiments using both synthetic and real data are provided to illustrate the efficiency of these variants.

1 Introduction

In this paper, we are interested in sampling a distribution μ supported on \mathcal{X} with the probability density function

$$\mu(\theta) \propto \exp(-U(\theta)), \quad \theta \in \mathcal{X}, \quad (1.1)$$

where \mathcal{X} is contained in a high-dimensional space. The sampling problem (1.1) has many applications in machine learning, such as Bayesian learning (inference) where different choices of $U(\cdot)$ in (1.1) correspond to different problems such as Bayesian linear regression [Hof09], Bayesian logistic regression [Hof09], Bayesian deep learning [WY20, PS17] and Bayesian principal component analysis [DRW⁺16].

Langevin algorithms are core Markov chain Monte Carlo (MCMC) methods in statistics that allow one to sample from a given density $\mu(\theta)$ of interest defined in (1.1). A common choice of the space \mathcal{X} in (1.1) is the Euclidean space, e.g. $\mathcal{X} = \mathbb{R}^d$. The classical Langevin Monte Carlo algorithm is based on the discretization of *overdamped (or first-order) Langevin dynamics* [Dal17, DM17, DK19, RRT17]:

$$d\theta_t = -\nabla U(\theta_t)dt + \sqrt{2}dW_t, \quad (1.2)$$

¹College of Mathematics Science, Shenzhen University, 518060 Shenzhen, China; yaonian@szu.edu.cn

²Department of Mathematics, Florida State University, 1017 Academic Way, Tallahassee, FL-32306, United States of America; pa22g@fsu.edu

³College of Mathematics Science, Shenzhen University, 518060 Shenzhen, China; 2247640315@qq.com

⁴Department of Mathematics, Florida State University, 1017 Academic Way, Tallahassee, FL-32306, United States of America; zhu@math.fsu.edu

where $U : \mathbb{R}^d \rightarrow \mathbb{R}$ and W_t is a standard d -dimensional Brownian motion with $\theta_0 \in \mathbb{R}^d$. Under some mild assumptions on $U(\cdot)$, the diffusion (1.2) admits a unique stationary distribution with the density $\mu(\theta) \propto e^{-U(\theta)}$, also known as the *Gibbs distribution* [CHS87, HKS89]. In computing practice, this diffusion is simulated by considering its discretization, and one of the most commonly used discretization scheme is the Euler–Maruyama discretization of (1.2), often known as the unadjusted Langevin algorithm in the literature; see e.g. [DM17]:

$$\theta_{k+1} = \theta_k - \eta \nabla U(\theta_k) + \sqrt{2\eta} \xi_{k+1}, \quad (1.3)$$

where ξ_k are i.i.d. $\mathcal{N}(0, I_d)$ Gaussian vectors.

The first non-asymptotic result of the discretized Langevin dynamics (1.3) is due to [Dal17], which was improved soon after by [DM17] with a particular emphasis on the dependence on the dimension d . Both works consider the total variation as the distance to measure the convergence. Later, [DM19] studied the convergence in the 2-Wasserstein distance, and [DMP18] studied variants of (1.3) when U is not smooth. [CB18] studied the convergence in the Kullback-Leibler (KL) distance. [DK19] studied the convergence when only stochastic gradients are available.

In the literature, there have also been active studies of various variants of the overdamped Langevin dynamics and the discretization schemes. One popular Langevin dynamics is the second-order, also known as kinetic or *underdamped Langevin dynamics*, see e.g. [CCBJ18, CCA⁺18, CLW23, DRD20, GGZ20, MCC⁺21, GGZ22]:

$$\begin{cases} dr_t = -\gamma r_t dt - \nabla U(\theta_t) dt + \sqrt{2\gamma} dW_t, \\ d\theta_t = r_t dt, \end{cases} \quad (1.4)$$

where W_t is a standard d -dimensional Brownian motion with $r_0, \theta_0 \in \mathbb{R}^d$. Under some mild assumptions on U , the diffusion (1.4) admits a unique stationary distribution with the density $\mu(\theta, r) \propto e^{-U(\theta) - \frac{1}{2}|r|^2}$, whose θ -marginal distribution coincides with the stationary distribution of (1.2).

Another popular variant is the *non-reversible Langevin dynamics*, see e.g. [HHMS93, HHMS05, DLP16, DPZ17, FSS20, RBS15, GGZ22, HWG⁺20]:

$$d\theta_t = -(I + J) \nabla U(\theta_t) dt + \sqrt{2} dW_t, \quad (1.5)$$

where W_t is a standard d -dimensional Brownian motion with $\theta_0 \in \mathbb{R}^d$, and J is an *antisymmetric* matrix, i.e. $J^\top = -J$, and under mild conditions, $\mu(\theta) \propto e^{-U(\theta)}$ is the unique stationary distribution of (1.5).

Other popular variants of Langevin dynamics include high-order Langevin dynamics [MMW⁺21], Hessian-free high-resolution dynamics [LZT22], mirror Langevin dynamics [HKRC18, CLGL⁺20, ZPFP20, LTVW22], replica exchange Langevin dynamics [CCD⁺19, DFG⁺20], as well as the Lévy-driven Langevin dynamics such as fractional Langevin

Monte Carlo [Sim17] and fractional underdamped Langevin dynamics [SZTG20], the decentralized Langevin algorithms such as decentralized stochastic gradient Langevin dynamics [GGHZ21] and EXTRA stochastic gradient Langevin dynamics [GIWZ24], and the constrained Langevin algorithms such as projected Langevin Monte Carlo [BEL15, BEL18, Lam21, ZL22], skew-reflected non-reversible Langevin dynamics [DFT+25], proximal Langevin Monte Carlo [BDMP17, SR20], and penalized Langevin algorithms [GHZ24].

In this paper, we consider a *generalized Langevin dynamics*, that is, a Markov process $\{\mathbf{z}_t\}_{t \geq 0}$ evolving in $\mathcal{X} = \mathbb{R}^n$, and satisfy the following stochastic differential equation (SDE):

$$d\mathbf{z}_t = \mathbf{f}(\mathbf{z}_t)dt + \sqrt{2\mathcal{D}(\mathbf{z}_t)}d\mathbf{W}_t, \quad (1.6)$$

where \mathbf{W}_t is an n -dimensional Brownian motion with $\mathbf{z}_0 \in \mathbb{R}^n$ and $\mathcal{D}(\mathbf{z})$ is a positive semidefinite diffusion matrix. We define the Hamiltonian

$$H(\mathbf{z}) = H(\theta, r) = U(\theta) + g(\theta, r), \quad (1.7)$$

where \mathbf{z} could represent θ itself, or an augmented state space such that $\mathbf{z} = (\theta, r)$ and θ is the model parameter in \mathbb{R}^d and r is a set of auxiliary variables in \mathbb{R}^m such that $d + m = n$. We are interested in the case when the choices of $\mathbf{f}(\mathbf{z})$ and $\mathcal{D}(\mathbf{z})$ yield the stationary distribution $\mu \propto \exp(-H(\mathbf{z}))$. We write $\mathbf{f}(\mathbf{z})$ as follows:

$$\mathbf{f}(\mathbf{z}) = -[\mathcal{D}(\mathbf{z}) + \mathcal{Q}(\mathbf{z})]\nabla H(\mathbf{z}) + \Gamma(\mathbf{z}), \quad (1.8)$$

where $\Gamma(\mathbf{z}) = (\Gamma_1(\mathbf{z}), \dots, \Gamma_n(\mathbf{z}))$ with

$$\Gamma_i(\mathbf{z}) := \sum_{j=1}^n \frac{\partial}{\partial \mathbf{z}_j} (\mathcal{D}_{ij}(\mathbf{z}) + \mathcal{Q}_{ij}(\mathbf{z})), \quad (1.9)$$

where $\mathcal{Q}(\mathbf{z})$ is an anti-symmetric curl matrix representing the deterministic traversing effects seen in Hamiltonian Monte Carlo (HMC) procedures. It is known that (see e.g. [MCF15]) $\mu \propto e^{-H(\mathbf{z})}$ is a stationary distribution of the SDE (1.6) under the following condition for $\mathcal{Q}(\mathbf{z})$:

$$\sum_{i=1}^n \sum_{j=1}^n \frac{\partial^2}{\partial \mathbf{z}_i \partial \mathbf{z}_j} (\mathcal{Q}_{ij}(\mathbf{z})e^{-H(\mathbf{z})}) = 0. \quad (1.10)$$

For any Langevin dynamics that converges to a given target distribution, it can be written in the provided framework (1.6) by choosing $\mathcal{D}(\mathbf{z})$ and $\mathcal{Q}(\mathbf{z})$ appropriately.

- (1) In the general framework of (1.6), by choosing $\mathcal{D} = \mathbf{I}$ and $\mathcal{Q} = \mathbf{0}$, and $\mathcal{X} = \mathbb{R}^d$, $\mathbf{z}_t = \theta_t \in \mathbb{R}^d$, and choosing $H(\mathbf{z}_t) = U(\theta_t)$, we recover the overdamped Langevin dynamics (1.2).

- (2) In the general framework of (1.6), supposing that $\mathcal{X} = \mathbb{R}^d \times \mathbb{R}^d$ and setting $\mathbf{z}_t = (\theta_t, p_t) \in \mathbb{R}^d \times \mathbb{R}^d$, and choosing $H(\mathbf{z}_t) = H(\theta_t, p_t) = U(\theta_t) + \frac{1}{2}|p_t|^2$, and

$$\mathcal{D} = \begin{pmatrix} \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \gamma \mathbf{I} \end{pmatrix} \quad \text{and} \quad \mathcal{Q} = \begin{pmatrix} \mathbf{0} & -\mathbf{I} \\ \mathbf{I} & \mathbf{0} \end{pmatrix}.$$

Then this reduces (1.6) to the underdamped Langevin dynamics (1.4).

- (3) In the general framework of (1.6), by choosing $\mathcal{D} = \mathbf{I}$ and $\mathcal{Q} = \mathbf{J}$, a constant $d \times d$ anti-symmetric matrix, i.e. $\mathbf{J}^\top = -\mathbf{J}$, and $\mathcal{X} = \mathbb{R}^d$, $\mathbf{z}_t = \theta_t \in \mathbb{R}^d$, and choosing $H(\mathbf{z}_t) = U(\theta_t)$, we recover the non-reversible Langevin dynamics (1.5).

In addition, the general framework of (1.6) includes the following models. We write ∇ for the gradient operator, ∇^2 for the Hessian matrix, $\nabla \cdot$ for the divergence operator, Δ for the Laplacian operator, and ∇^3 for the third-order tensor of partial derivatives. Please find the more detailed definitions in Appendix A.

- (4) In the general framework of (1.6), by choosing $\mathcal{X} = \mathbb{R}^d$ and

$$H(\mathbf{z}_t) = U(\theta_t), \quad \mathcal{D}(\theta) = [\nabla^2 \phi(\theta)]^{-1}, \quad \mathcal{Q}(\theta) = \begin{pmatrix} 0 & e^{U(\theta)} & \dots & e^{U(\theta)} \\ -e^{U(\theta)} & 0 & \ddots & \vdots \\ \vdots & \vdots & \ddots & e^{U(\theta)} \\ -e^{U(\theta)} & -e^{U(\theta)} & \dots & 0 \end{pmatrix}, \quad (1.11)$$

we reduce (1.6) to the well-known *mirror Langevin dynamics* (see e.g. [HKRC18, ZPFP20, CLGL+20, LTVW22, AC21]), that is,

$$d\theta_t = (\Phi(\theta_t) - [\nabla^2 \phi(\theta_t)]^{-1} \nabla U(\theta_t)) dt + \sqrt{2[\nabla^2 \phi(\theta_t)]^{-1}} dW_t, \quad (1.12)$$

where $\Phi(\theta) := -[\nabla^2 \phi(\theta)]^{-1} \text{Tr}(\nabla^3 \phi(\theta) [\nabla^2 \phi(\theta)]^{-1})$ and W_t is a standard Brownian motions in \mathbb{R}^d with $\phi \in C^3(\mathcal{X})$. For mirror Langevin dynamics (1.12), $h := \nabla \phi : \mathcal{X} \mapsto \mathbb{R}^d$ is known as the *mirror map*, which transforms the coordinates from the primal space to the dual space and $\Phi(\theta) = [\nabla h(\theta)]^{-1} \text{Tr}(\nabla^2 h(\theta) [\nabla h(\theta)]^{-1})$ is a mirror curvature correction that ensures that the Langevin dynamics correctly follows the structure induced by the mirror map h .

- (5) In the general framework of (1.6), by choosing $\mathcal{X} = \mathbb{R}^d \times \mathbb{R}^d \times \mathbb{R}^d$ and setting $\mathbf{z}_t = (\theta_t, p_t, r_t) \in \mathbb{R}^d \times \mathbb{R}^d \times \mathbb{R}^d$, and choosing

$$H(\mathbf{z}_t) = H(\theta_t, p_t, r_t) = U(\theta_t) + \frac{1}{2}|p_t|^2 + \frac{1}{2}|r_t|^2, \quad (1.13)$$

and

$$\mathcal{D} = \begin{pmatrix} \mathbf{0} & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \alpha \mathbf{I} \end{pmatrix} \quad \text{and} \quad \mathcal{Q} = \begin{pmatrix} \mathbf{0} & -\mathbf{I} & \mathbf{0} \\ \mathbf{I} & \mathbf{0} & -\gamma \mathbf{I} \\ \mathbf{0} & \gamma \mathbf{I} & \mathbf{0} \end{pmatrix}. \quad (1.14)$$

Then this reduces (1.6) to the *high-order Langevin dynamics* introduced in [MMW⁺21]; see also [Mon23]:

$$\begin{cases} d\theta_t = p_t dt, \\ dp_t = -\nabla U(\theta_t) dt + \gamma r_t dt, \\ dr_t = -\gamma p_t dt - \alpha r_t dt + \sqrt{2\alpha} dW_t, \end{cases} \quad (1.15)$$

where $\alpha, \gamma > 0$ are friction parameters and W_t is a standard Brownian motions in \mathbb{R}^d .

- (6) In the general framework of (1.6), let $\mathcal{X} = \mathbb{R}^d \times \mathbb{R}^d$ and $\mathbf{z}_t = (\theta_t, r_t) \in \mathbb{R}^d \times \mathbb{R}^d$, and define the function

$$H(\mathbf{z}_t) = H(\theta_t, r_t) = U(\theta_t) + \frac{1}{2}|r_t|^2, \quad (1.16)$$

along with the matrices

$$\mathcal{D} = \begin{pmatrix} \beta \mathbf{I} & \mathbf{0} \\ \mathbf{0} & \alpha \mathbf{I} \end{pmatrix} \quad \text{and} \quad \mathcal{Q} = \begin{pmatrix} \mathbf{0} & -\mathbf{I} \\ \mathbf{I} & \mathbf{0} \end{pmatrix},$$

then we obtain the *Hessian-free high-resolution dynamics* introduced in [LZT22]:

$$\begin{cases} d\theta_t = r_t dt - \beta \nabla U(\theta_t) dt + \sqrt{2\beta} d\bar{W}_t, \\ dr_t = -\alpha r_t dt - \nabla U(\theta_t) dt + \sqrt{2\alpha} d\bar{B}_t, \end{cases} \quad (1.17)$$

where $\beta > 0, \alpha > 0$ are friction parameters, and \bar{W}_t, \bar{B}_t are two independent standard Brownian motions in \mathbb{R}^d .

For any fixed time $t > 0$, the empirical measure of the generalized Langevin dynamics $\{\mathbf{z}_t\}_{t \geq 0}$ defined in (1.6) is given as

$$\pi_t = \frac{1}{t} \int_0^t \delta_{\mathbf{z}_s} ds,$$

where δ_x is the Dirac measure at $x \in \mathcal{X}$, such that $\{\pi_t\}_{t \geq 0}$ is a sequence of random measures, which are random elements of $\mathcal{P}(\mathcal{X})$, that is, the space of probability measures on $\mathcal{X} = \mathbb{R}^n$. We will show that under mild conditions, (1.6) is ergodic and by ergodic theorem,

$$\pi_t \rightarrow \mu \propto e^{-H(\mathbf{z})}, \quad \text{almost surely as } t \rightarrow \infty. \quad (1.18)$$

To understand how fast π_t converges to the stationary distribution μ , we adopt a *large deviations* approach. We present the large deviation principle about the invariant measure of the empirical measure related to the process, and then use the large deviation principle to judge the acceleration effect of the algorithm. The large deviation principle was first formulated in the pioneering work by Varadhan [Var66]. In contrast to the law of large numbers, which studies a typical event, the large deviations study the small probability of rare events [DZ98, Var84, Var08]. In a series of seminal papers, Donsker and Varadhan studied the large deviation principle for the empirical measure π_t for $t \rightarrow \infty$, where the underlying is a Markov process [DV75a, DV75b, DV76, DV83]. Informally, $\mathbb{P}(\pi_t \in \cdot)$ satisfies a large deviation principle with rate function $I(\cdot)$ if for any $\nu \in \mathcal{P}(\mathcal{X})$,

$$\mathbb{P}(\pi_t \approx \nu) = e^{-tI(\nu)+o(t)},$$

as $t \rightarrow \infty$, where $I(\nu) \geq 0$ and $I(\nu) = 0$ if and only if $\nu = \mu$. In other words, when $\nu \neq \mu$, $\mathbb{P}(\pi_t \approx \nu)$ is exponentially small as time $t \rightarrow \infty$, and the rate function $I(\nu)$ characterizes how small this probability is. This suggests the larger the value of the rate function, the smaller the probability the empirical measure deviates away from the Gibbs distribution, and hence one expects faster convergence.

From technical perspective, although Donsker-Varadhan large deviations theory for large-time asymptotics works for the general Markov process, it often requires restrictive assumptions, including for example compact domain (see e.g. [DV75a]). The Langevin dynamics we are interested in lives in the unbounded Euclidean space, and our analysis relies on a more recent large deviations result where the space can be unbounded and the underlying topology is the weighted topology [FS20].

The idea of applying large deviations analysis to study Langevin algorithms and the variants is not new. For example, the large deviations for overdamped and underdamped Langevin dynamics are studied in [FS20]. The large deviations for some variants of Langevin algorithms have been studied in [RBS15, CCD⁺19]. However, these works are mostly restricted to a particular variant of the Langevin algorithm, e.g. the non-reversible Langevin in [RBS15] and replica exchange Langevin in [CCD⁺19]. We take a more unified approach by studying a generalized Langevin dynamics, described in [MCF15], that includes overdamped, underdamped, non-reversible Langevin dynamics as special cases, as well as other variants from the recent literature literature, for example, the mirror Langevin dynamics [HKRC18], the high-order Langevin dynamics [MMW⁺21], and the Hessian-free high-resolution dynamics [LZT22] whose large deviations have never been studied in the literature to the best of our knowledge.

The contributions of our paper can be summarized as follows:

- In Section 2, we first establish large deviations for the empirical measures for the generalized Langevin dynamics (1.6) (Theorem 2.1) assuming Hypocoellipticity (Assumption 2.1), Controllability (Assumption 2.2) and Lyapunov condition (Assumption 2.3). Then, we apply the general framework of the large deviations for the

generalized Langevin dynamics (1.6) to study the large deviations for variants of Langevin dynamics, including mirror Langevin dynamics (1.12) (Theorem 2.2), high-order Langevin dynamics (1.15) (Theorem 2.3), and Hessian-free high-resolution dynamics (1.17) (Theorem 2.4), whose large deviations have never been established in the literature to the best of our knowledge. The technical novelty lies upon a careful analysis to check Hypoellipticity (Assumption 2.1), Controllability (Assumption 2.2) and Lyapunov condition (Assumption 2.3) by constructing novel Lyapunov functions, under mild conditions on the target function $U(\cdot)$, for each variant of Langevin dynamics. To the best of our knowledge, this is the first general and unified framework for the large deviations of Langevin dynamics and a study of most familiar variants of Langevin dynamics of interest in the literature.

- In Section 3, we use the large deviation rate functions obtained in Section 2 as a measure to analyze the speed of convergence to the invariant measure. The larger the rate function, the more concentrated the empirical measure is around the invariant measure, thus indicating faster convergence. Since variants of Langevin dynamics, such as underdamped Langevin dynamics (1.4), high-order Langevin dynamics (1.15), and Hessian-free high-resolution dynamics (1.17), may live in a higher-dimensional space such as \mathbb{R}^{2d} or \mathbb{R}^{3d} than the overdamped Langevin dynamics (1.2) in \mathbb{R}^d , a direct comparison of their rate functions impossible. We introduce a novel method to expand the space of overdamped Langevin dynamics to match the dimensions with those of the variants of Langevin dynamics under study. We then show comparisons of their rate functions for some parameter regimes either on the whole space of probability measures or a subspace of probability measures (Proposition 3.1, Proposition 3.2, Proposition 3.3). In the former case, by applying contraction principle from large deviations, we show the acceleration for the underlying process θ_t in \mathbb{R}^d by establishing the comparison for the LDP rate functions for its empirical measure (Corollary 3.3).
- In Section 4, we conduct numerical experiments for various Langevin algorithms based on the Euler–Maruyama discretization of variants of Langevin dynamics. We compare the numerical results with the unadjusted Langevin algorithm, i.e. the Euler–Maruyama discretization of overdamped Langevin dynamics. These numerical experiments show superior performance or comparable performance using variants of Langevin algorithms.

The rest of the paper is organized as follows. In Section 2, we introduce the main results of our paper, including a large deviation principle for the generalized Langevin dynamics and, under various mild assumptions on the target distribution, the large deviations for mirror Langevin dynamics, high-order Langevin dynamics and Hessian-free high-resolution dynamics. In Section 3, we study the acceleration of variants of Langevin dynamics compared to the overdamped Langevin dynamics by comparing their rate functions from the large deviations theory. Numerical experiments are provided in Section 4. We conclude in

Section 5. The notations used in this paper are summarized in Appendix A, and all the technical proofs are provided in Appendix B.

2 Main Results

2.1 Preliminary

We use $C_c^\infty(\mathcal{X})$ (resp. $C_b(\mathcal{X})$) to denote the space of smooth functions with compact support (resp. continuous and bounded functions), as well as the space of smooth functions growing at most polynomially and whose derivatives also grow at most polynomially:

$$\mathcal{S} = \left\{ \varphi \in C^\infty(\mathcal{X}) \mid \forall \alpha \in \mathbb{N}^d, \exists N > 0, \text{ such that } \sup_{x \in \mathcal{X}} \frac{|\partial^\alpha \varphi(x)|}{(1 + |x|^2)^N} < +\infty \right\}, \quad (2.1)$$

where $\partial^\alpha = \partial_{x_1}^{\alpha_1} \cdots \partial_{x_d}^{\alpha_d}$ and $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_d)$.

Let us recall that given a sequence of empirical measures $\{\pi_t\}_{t \geq 0}$, we say that π_t satisfies a *large deviation principle* (LDP) on $\mathcal{P}(\mathcal{X})$ equipped with the τ^κ topology with the rate function $I : \mathcal{P}(\mathcal{X}) \rightarrow \mathbb{R}$ if I is non-negative, lower semicontinuous and for any τ^κ -measurable set $\Theta \subset \mathcal{P}(\mathcal{X})$,

$$-\inf_{\nu \in \overset{\circ}{\Theta}} I(\nu) \leq \liminf_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{P}(\pi_t \in \Theta) \leq \limsup_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{P}(\pi_t \in \Theta) \leq -\inf_{\nu \in \bar{\Theta}} I(\nu),$$

where $\overset{\circ}{\Theta}$ denotes the interior of Θ and $\bar{\Theta}$ stands for its closure; see e.g. [DZ98, Var84, Var08] for a survey on the theory of large deviations.

By following [FS20], we introduce the following three basic assumptions: hypoellipticity of the generator, controllability (irreducibility of the dynamics), and a Lyapunov condition. First, let us introduce the hypoellipticity assumption.

Assumption 2.1. (*Hypoellipticity*) *The functions \mathbf{f} and $\sqrt{\mathcal{D}}$ in (1.6) belong to \mathcal{S}^n and $\mathcal{S}^{n \times n}$, respectively, and the generator \mathcal{L}_τ defined in (B.1) satisfies the Hypoelliptic Hörmander condition. More precisely, \mathcal{L}_τ can be written as*

$$\mathcal{L}_\tau = \sum_{j=1}^r X_j^* X_j + X_0, \quad (2.2)$$

where X_0, \dots, X_r denote first-order homogeneous differential operators in an open set $\Omega \subset \mathbb{R}^n$ with C^∞ coefficients such that $\{X_j\}_{j=1}^r, \{[X_i, X_j]\}_{i,j=0}^r, \{[[X_i, X_j], X_k]\}_{i,j,k=0}^r, \dots$ span the space Ω at any given point $x \in \mathbb{R}^n$ for a finite number of commutators $n_x \in \mathbb{N}$, where the definition of $[X, Y]$ can be found in Appendix A.

Next, we introduce the following controllability condition about the irreducibility of the dynamics.

Assumption 2.2. (Controllability). For any $x, y \in \mathcal{X}$ and $T > 0$, there exists a control $u \in C^0([0, T], \mathbb{R}^n)$ such that the path $\phi \in C^0([0, T], \mathcal{X})$ satisfying

$$\begin{cases} \phi(0) = x, \\ \dot{\phi}(t) = \mathbf{f}(\phi(t)) + \sqrt{2\mathcal{D}(\phi(t))}u(t), \end{cases} \quad (2.3)$$

and $\phi(T) = y$ is well-defined, where \mathbf{f} and \mathcal{D} are defined in (1.6).

Finally, by following the Lyapunov condition given in Proposition 2.9 in [FS20], we introduce the following assumption.

Assumption 2.3. (Lyapunov condition). There exists a function $W : \mathcal{X} \rightarrow [1, +\infty)$ of class $C^2(\mathcal{X})$ has compact level sets and $|\sqrt{\mathcal{D}}\nabla W|$ has compact level sets. For any $\theta \in (0, 1)$,

$$\mathcal{L}_\tau W - \theta|\sqrt{\mathcal{D}}\nabla W|^2 \sim |\sqrt{\mathcal{D}}\nabla W|^2, \quad (2.4)$$

where for any two functions $g, f : \mathcal{X} \rightarrow \mathbb{R}$, g is said to be equivalent to f (denoted by $g \sim f$) if there exist constants $c, c' > 0$ and $R, R' \in \mathbb{R}$ such that $c'g(x) - R' \leq f(x) \leq cg(x) + R$ for any $x \in \mathcal{X}$.

We first introduce the following technical lemma from [FS20], which will be used in the proof of Theorem 2.2 and in particular showing the Lyapunov condition.

Proposition 2.1 (Proposition 2.9 in [FS20]). Suppose Assumptions 2.1-2.3 hold. Let $\kappa : \mathcal{X} \rightarrow [1, +\infty)$ be a function of class \mathcal{S} that is defined in (2.1). For any $\eta \in (0, 1)$, define

$$W_\eta(\mathbf{z}) = e^{\eta W(\mathbf{z})}.$$

Then,

$$\Psi_\tau := -\frac{\mathcal{L}_\tau W_\eta}{W_\eta} = \eta \left(-\mathcal{L}_\tau W - \eta|\sqrt{\mathcal{D}}\nabla W|^2 \right) \quad (2.5)$$

has compact level sets. It follows that

$$\frac{\Psi_\tau(\mathbf{z})}{\kappa(\mathbf{z})} \xrightarrow{|\mathbf{z}| \rightarrow +\infty} +\infty.$$

Under Assumptions 2.1-2.3, it is known that π_t converges to the invariant distribution μ exponentially fast as $t \rightarrow \infty$. We have the following result from [FS20].

Proposition 2.2 (Proposition 2.10. in [FS20]). Suppose Assumptions 2.1-2.3 hold. There exist $C, c > 0$ such that for any initial measure $\nu \in \mathcal{P}(\mathcal{X})$,

$$d_W(\nu P_t, \mu) \leq Ce^{-ct} d_W(\nu, \mu),$$

where $d_W(\nu, \mu)$ is the associated weighted total variation distance for ν and μ ; see exact definition in Appendix A.

However, Proposition 2.2 does not have an explicit expression for the convergence speed c except that such a positive c exists. As a result, one cannot rely on Proposition 2.2 to compare the performance of the variants of Langevin dynamics. This motivates us to adopt a different approach in our paper, by obtaining large deviations for the variants of Langevin dynamics, and characterizing their rate functions as a measure to compare their convergence to the invariant distributions.

We introduce the carré du champ operator [BGL13] associated with $\mathcal{L} := b \cdot \nabla + \mathcal{S} : \nabla^2$ which is defined as follows. For two regular functions φ and ψ :

$$\mathcal{C}(\varphi, \psi) = \frac{1}{2}(\mathcal{L}(\varphi\psi) - \varphi\mathcal{L}\psi - \psi\mathcal{L}\varphi) = \nabla\varphi \cdot \mathcal{S}\nabla\psi, \quad (2.6)$$

where ∇^2 stands for the Hessian matrix, and for two matrices A, B belonging to $\mathbb{R}^{d \times d}$, we write $A : B = \text{Tr}(A^\top B)$.

For any $\varphi \in C_c^\infty(\mathcal{X})$, we also introduce the seminorms

$$|\varphi|_{\mathcal{H}^1(\nu)}^2 = \int_{\mathcal{X}} \mathcal{C}(\varphi, \varphi) d\nu,$$

and

$$|\varphi|_{\mathcal{H}^{-1}(\nu)}^2 = \sup_{\psi \in C_c^\infty} \left\{ 2 \int_{\mathcal{X}} \varphi\psi d\nu - |\psi|_{\mathcal{H}^1(\nu)}^2 \right\}.$$

Let $\tilde{\nabla}$ denote the adjoint of gradient operator ∇ in $L^2(\nu)$.

In order to provide the exact expression of the rate function of generalized Langevin dynamics (1.6), we decompose the rate function into two parts according to the symmetric and anti-symmetric parts of dynamics. First, for any closed operator T , we denote T^* as its adjoint on $L^2(\mu)$, where μ is the invariant probability measure of the dynamics. Then consider the generator \mathcal{L}_τ of the dynamics (1.6), we can decompose it into symmetric and anti-symmetric parts with respect to μ , reads

$$\mathcal{L}_\tau = \mathcal{L}_S + \mathcal{L}_A, \quad \mathcal{L}_S = \frac{\mathcal{L}_\tau + \mathcal{L}_\tau^*}{2}, \quad \mathcal{L}_A = \frac{\mathcal{L}_\tau - \mathcal{L}_\tau^*}{2}. \quad (2.7)$$

For any $\mu \in \mathcal{P}(\mathcal{X})$, let

$$L^2(\mu) = \left\{ \varphi \text{ measurable} \mid \int_{\mathcal{X}} |\varphi|^2 d\mu < +\infty \right\}.$$

The rate function $I(\nu)$ (often referred to as the Donsker-Varadhan functional in the literature [DV75a, DV75b, DV76, DV83]) takes the form

$$I(\nu) = \sup \left\{ - \int_{\mathcal{X}} \frac{\mathcal{L}u}{u} d\nu, \quad u \in \mathcal{D}^+(\mathcal{L}) \right\}, \quad \text{for any } \nu \in \mathcal{P}(\mathcal{X}), \quad (2.8)$$

where

$$\mathcal{D}^+(\mathcal{L}) = \left\{ u \in \mathcal{D}(\mathcal{L}) \cap \mathcal{C}^0(\mathcal{X}) \mid u > 0, -\frac{\mathcal{L}u}{u} \in B_\kappa^\infty(\mathcal{X}) \right\}.$$

According to the symmetric and anti-symmetric parts of the generator, we can use the decomposition of the operator to obtain the symmetric and anti-symmetric parts of the rate function respectively, as described in the following technical lemma.

Lemma 2.1 (Theorem 3.3 in [FS20]). *Suppose that Assumptions 2.1-2.3 hold. Consider a measure $\nu \in \mathcal{P}_\kappa(\mathcal{X})$ such that $d\nu = e^v d\mu$ with $v \in \mathcal{H}^1(\nu)$ and $\mathcal{L}_A v \in \mathcal{H}^{-1}(\nu)$. Then, the rate function I defined in (2.8) reads:*

$$I(\nu) = I_S(\nu) + I_A(\nu),$$

where

$$I_S(\nu) = \frac{1}{4} |v|_{\mathcal{H}^1(\nu)}^2,$$

and

$$I_A(\nu) = \frac{1}{4} |\mathcal{L}_A(v)|_{\mathcal{H}^{-1}(\nu)}^2.$$

Since $d\nu = e^v d\mu$ is the Radon-Nikodym derivative of ν with respect to μ , we have

$$I(\nu) = \frac{1}{4} \left| \log \frac{d\nu}{d\mu} \right|_{\mathcal{H}^1(\nu)}^2 + \frac{1}{4} \left| \mathcal{L}_A \left(\log \frac{d\nu}{d\mu} \right) \right|_{\mathcal{H}^{-1}(\nu)}^2.$$

2.2 Main Results

Now we introduce our main result, that is, the large deviation for the generalized Langevin dynamics (1.6) under Hypoellipticity (Assumption 2.1), Controllability (Assumption 2.2) and Lyapunov condition (Assumption 2.3), and we obtain an explicit characterization of its rate function.

Theorem 2.1. *Suppose that Assumptions 2.1-2.3 hold, and the function κ as in Proposition 2.1 and for any $\nu \in \mathcal{P}(\mathcal{X})$ of the form $d\nu = e^v d\mu$. Then the empirical measure $\{\pi_t\}_{t \geq 0}$ of the generalized Langevin dynamics defined by (1.6) satisfies a large deviation principle in the τ^κ -topology with the corresponding rate function defined by*

$$I_\tau(\nu) = \frac{1}{4} \int_{\mathcal{X}} \nabla v \cdot \mathcal{D} \nabla v \, d\nu + \frac{1}{4} \int_{\mathcal{X}} \nabla \psi_v \cdot \mathcal{D} \nabla \psi_v \, d\nu, \quad (2.9)$$

where ψ_v is the unique solution in $\mathcal{H}^1(\nu)$ to the Poisson equation

$$\tilde{\nabla}(\mathcal{D} \nabla \psi_v) = \mathcal{L}_A v. \quad (2.10)$$

That is to say for τ^κ -measurable set $\Theta \subset \mathcal{P}(\mathcal{X})$, it holds

$$-\inf_{\nu \in \dot{\Theta}} I_\tau(\nu) \leq \liminf_{t \rightarrow +\infty} \frac{1}{t} \log \mathbb{P}(\pi_t \in \Theta) \leq \limsup_{t \rightarrow +\infty} \frac{1}{t} \log \mathbb{P}(\pi_t \in \Theta) \leq -\inf_{\nu \in \dot{\Theta}} I_\tau(\nu).$$

Next, we are interested in applying the general framework of large deviations for the generalized Langevin dynamics (1.6) to study the large deviations for the variants of Langevin dynamics. For each variant of Langevin dynamics, we carefully verify Hypocoellipticity (Assumption 2.1), Controllability (Assumption 2.2) and Lyapunov condition (Assumption 2.3). According to the form of generalized Langevin dynamics (1.6), we can divide the variants of Langevin dynamics of interest into two cases to study:

- \mathbf{z} represents the model parameter θ itself, that is, $g(\theta, r) = 0$ and $H(\mathbf{z}) = U(\theta)$. This case includes the *overdamped Langevin dynamics* (1.2), the *non-reversible Langevin dynamics* (1.5), and the *mirror Langevin dynamics* (1.12).
- \mathbf{z} represents the extended state space, that is, the case of $g(\theta, r) \neq 0$. This case includes the *underdamped Langevin dynamics* (1.4), the *high-order Langevin dynamics* (1.15) and the *Hessian-free high-resolution dynamics* (1.17).

Before we proceed, let us recall the LDPs for overdamped Langevin dynamics (1.2), underdamped Langevin dynamics (1.4) and non-reversible Langevin dynamics (1.5) have already been obtained in the literature; see e.g. [FS20].

The LDP for the overdamped Langevin dynamics (1.2) is well known, see e.g. [FS20]. Setting $\mathbf{z} = \theta \in \mathbb{R}^d$ and choosing $H(\mathbf{z}) = U(\theta)$, we get the overdamped Langevin dynamics:

$$d\theta_t = -\nabla U(\theta_t)dt + \sqrt{2}d\mathbf{W}_t. \quad (2.11)$$

Lemma 2.2 (Proposition 4.1 in [FS20]). *Suppose that Assumptions 2.1-2.3 holds. Then, the rate function $I(\cdot)$ defined in (2.8) for the large deviations for overdamped Langevin dynamics (2.11) reads*

$$I_o(\nu) = \frac{1}{4} \int_{\mathcal{X}} |\nabla v|^2 d\nu, \quad (2.12)$$

for any measure $\nu \in \mathcal{P}(\mathcal{X})$ of the form $d\nu = e^v d\mu$.

The LDP for the underdamped Langevin dynamics (1.4) has been obtained in e.g. [FS20], which is stated as follows.

Lemma 2.3. (*Theorem 4.6 in [FS20]*) *Assume that $(X_t)_{t \geq 0} = (\theta_t, r_t)_{t \geq 0}$ is the underdamped Langevin in equation (1.4) where U satisfies Assumption 4.4 in [FS20] and consider a smooth function κ with $\kappa(\theta, r) = 1 + |\theta|^\alpha + |r|^\beta$ for $|\theta| + |r| \geq 1$ and $\alpha \in [0, 2)$, $\beta \in [0, 2)$. Then $(X_t)_{t \geq 0}$ is ergodic with respect to the measure μ . Moreover, the empirical measure*

$$\pi_t := \frac{1}{t} \int_0^t \delta_{(\theta_s, r_s)} ds$$

satisfies a LDP in the τ^κ -topology. Finally, for any $\nu \in \mathcal{P}_\kappa(\mathcal{X})$ such that $d\nu = e^v d\mu$ with $v \in \mathcal{H}^1(\nu)$ and $\mathcal{L}_{ham} v \in \mathcal{H}^{-1}(\nu)$, where the generator of the dynamics is

$$\mathcal{L}_\gamma = \mathcal{L}_{ham} + \gamma \mathcal{L}_{FD},$$

with

$$\mathcal{L}_{ham} = r \cdot \nabla_\theta - \nabla U \cdot \nabla_r, \quad \mathcal{L}_{FD} = -r \cdot \nabla_r + \Delta_r.$$

and the rate function reads

$$I_u(\nu) = \frac{\gamma}{4} \int_{\mathcal{X}} |\nabla_r v|^2 d\nu + \frac{1}{4\gamma} \int_{\mathcal{X}} |\nabla_r \psi|^2 d\nu, \quad (2.13)$$

where ψ is the unique solution in $\mathcal{H}^1(\nu)$ to the Poisson problem:

$$-\Delta_r \psi + (r - \nabla_r v) \cdot \nabla_r \psi = \mathcal{L}_{ham} v. \quad (2.14)$$

The LDP for the non-reversible Langevin dynamics (1.5) has been obtained in e.g. [FS20], which can be derived from Proposition 4.3 in [FS20]

Lemma 2.4 (Proposition 4.3 in [FS20]). *Let θ_t follow the non-reversible Langevin dynamics (1.5). Assume that the potential $U \in \mathcal{S}$ has compact level sets, satisfies $e^{-U} \in L^1(\mathcal{X})$ and, for any $\eta \in (0, 1)$, it holds*

$$(1 - \eta) |\nabla U|^2 - \Delta U \xrightarrow{|\theta| \rightarrow +\infty} +\infty.$$

Then, with the notation in (2.7), it holds $\mathcal{L}_S = -\nabla U \cdot \nabla + \Delta$ and $\mathcal{L}_A = -J\nabla U \cdot \nabla$ and $\mathcal{L} = \mathcal{L}_S + \mathcal{L}_A$. Moreover, $\pi_t = \frac{1}{t} \int_0^t \delta_{\theta_s} ds$ satisfies a large deviation principle (LDP) in the τ_κ -topology with compact level sets and such that $\kappa(x) \rightarrow +\infty$ as $|x| \rightarrow \infty$. The associated rate function I_J is given by:

$$I_J(\nu) = \frac{1}{2} \int_{\mathbb{R}^n} (|\nabla \psi_v|^2 + (U - v) \nabla \cdot \nabla \psi_v) d\nu,$$

where ψ_v is the unique $H^1(\nu)$ -solution to the Poisson equation

$$-\Delta \psi + \nabla(U - v) \cdot \nabla \psi_v = (-J\nabla U) \cdot \nabla v.$$

Next, we will derive the LDPs for mirror Langevin dynamics (1.12), high-order Langevin dynamics (1.15) and Hessian-free high-resolution dynamics (1.17), which to the best of our knowledge have never been studied in the previous literature.

2.2.1 Mirror Langevin dynamics

According to the values of \mathcal{D} and \mathcal{Q} set in (1.11) it is a classic case of the model we constructed, through formula (B.8) we have

$$\begin{aligned} \mathcal{L}_{MS} &= \sum_{i=1}^d \sum_{j=1}^d \left[\frac{\partial}{\partial \theta_j} \mathcal{D}_{ij}(\theta) - \mathcal{D}_{ij}(\theta) \frac{\partial}{\partial \theta_j} U(\theta) \right] \frac{\partial}{\partial \theta_i} + \sum_{i=1}^d \sum_{j=1}^d \mathcal{D}_{ij}(\theta) \frac{\partial^2}{\partial \theta_i \partial \theta_j}, \\ \mathcal{L}_{MA} &= \sum_{i=1}^d \sum_{j=1}^d \left[\frac{\partial}{\partial \theta_j} \mathcal{Q}_{ij}(\theta) - \mathcal{Q}_{ij}(\theta) \frac{\partial}{\partial \theta_j} U(\theta) \right] \frac{\partial}{\partial \theta_i} = 0. \end{aligned}$$

We know that $\mathcal{L}_M = \mathcal{L}_{MS} + \mathcal{L}_{MA}$ is the infinite generator of the mirror Langevin dynamics in (1.12). We now give a standard condition on U under which the framework developed in Section 2 and Appendix B applies.

Theorem 2.2. *Assume that the potential $U \in \mathcal{S}$ has compact level sets, $e^{-U} \in L^1(\mathcal{X})$, and for any $\eta \in (0, 1)$, it holds that*

$$(1 - \eta) \left| \sqrt{[\nabla^2 \phi(\theta)]^{-1}} \cdot \nabla U \right|^2 - [\nabla^2 \phi(\theta)]^{-1} \cdot \nabla^2 U - \nabla \cdot [\nabla^2 \phi(\theta)]^{-1} \nabla [\nabla^2 \phi(\theta)]^{-1} \xrightarrow{|\theta| \rightarrow +\infty} +\infty. \quad (2.15)$$

Then the mirror Langevin dynamics (1.12) admits the function

$$W_M(\theta) = e^{\eta U(\theta)}$$

for any $\eta \in (0, 1)$ as a Lyapunov function in the sense of Assumption 2.3. Moreover,

$$\begin{aligned} \Psi_M &:= -\frac{\mathcal{L}_M W_M}{W_M} \\ &= \eta \left((1 - \eta) \left| \sqrt{[\nabla^2 \phi(\theta)]^{-1}} \nabla U \right|^2 - [\nabla^2 \phi(\theta)]^{-1} : \nabla^2 U - \nabla [\nabla^2 \phi(\theta)]^{-1} \cdot \nabla U \right) \end{aligned} \quad (2.16)$$

has compact level sets and, for any $\kappa : \mathcal{X} \rightarrow [1, \infty)$ belonging to \mathcal{S} , bounded or with compact level sets and

$$\frac{\Psi_M(\mathbf{z})}{\kappa(\mathbf{z})} \xrightarrow{|\mathbf{z}| \rightarrow +\infty} +\infty.$$

The empirical measure $\{\pi_t\}_{t \geq 0}$ satisfies a large deviation principle in the τ^κ -topology and the corresponding rate function is defined by

$$I_M(\nu) = \frac{1}{4} \int_{\mathcal{X}} \nabla v \cdot [\nabla^2 \phi(\theta)]^{-1} \nabla v \, d\nu, \quad (2.17)$$

where $d\nu = e^v d\mu$. That is to say for τ^κ -measurable set $\Theta \subset \mathcal{P}(\mathcal{X})$, it holds

$$-\inf_{\nu \in \Theta} I_M(\nu) \leq \liminf_{t \rightarrow +\infty} \frac{1}{t} \log \mathbb{P}(\pi_t \in \Theta) \leq \limsup_{t \rightarrow +\infty} \frac{1}{t} \log \mathbb{P}(\pi_t \in \Theta) \leq -\inf_{\nu \in \Theta} I_M(\nu).$$

2.2.2 High-order Langevin dynamics

For the high-order Langevin dynamics (1.15), according to the values of \mathcal{D} and \mathcal{Q} set in (1.14), through formula (B.8), we can compute that the infinitesimal generator of (1.15) is given by

$$\mathcal{L}_H = \mathcal{L}_{HS} + \mathcal{L}_{HA} = \alpha \Delta_r - (\gamma p + \alpha r) \cdot \nabla_r + p \cdot \nabla_\theta - (\nabla U - \gamma r) \cdot \nabla_p, \quad (2.18)$$

where

$$\begin{aligned}\mathcal{L}_{\text{HS}} &:= -\alpha r \cdot \nabla_r + \alpha \Delta_r, \\ \mathcal{L}_{\text{HA}} &:= p \cdot \nabla_\theta - \nabla U \cdot \nabla_p + \gamma r \cdot \nabla_p - \gamma p \cdot \nabla_r.\end{aligned}$$

In order to obtain the large deviation principle for the high-order Langevin dynamics (1.15), we make the following classical assumptions for the growth of potential function U .

Assumption 2.4. *The potential $U \in \mathcal{S}$ has compact level sets, satisfies $e^{-U} \in L^1(\mathcal{X})$*

(a) *there exist $k > 1$ and $M_U, c_U, m_U > 0$ such that for all $\theta \in \mathbb{R}^d$ with $|\theta| \geq c_U$:*

$$m_U |\theta|^k \leq U(\theta) \leq M_U |\theta|^k \quad \text{and} \quad m_U |\theta|^k \leq \theta \cdot \nabla U(\theta).$$

(b) *$|\nabla U(\theta)| \leq M_U |\theta|^{k-1}$ if $|\theta| \geq c_U$.*

In the next lemma, we show that Hypoellipticity (Assumption 2.1) and Controllability (Assumption 2.2) are satisfied for the high-order Langevin dynamics (1.15).

Lemma 2.5. *(Hypoellipticity and Controllability). The generator \mathcal{L}_H defined in (2.18) satisfies the Hypoelliptic Hörmander and Controllability conditions.*

In the next lemma, we show that Lyapunov condition (Assumption 2.3) is satisfied for the high-order Langevin dynamics (1.15).

Lemma 2.6. *(Witten-Lyapunov condition). Suppose that $(\mathbf{z}_t)_{t \geq 0} = (\theta_t, p_t, r_t)_{t \geq 0}$ in (1.15) where potential U satisfies Assumption 2.4. Then for any $\alpha, \gamma > 0$ and $k \in (1, 2]$, there exists $\delta \in (\frac{2-k}{k}, 1]$ such that*

$$W_\delta(\theta, p, r) = e^{\varphi_{\text{HL}}^\delta(\theta, p, r)} \tag{2.19}$$

is a Lyapunov function, where $\varphi_{\text{HL}}(\theta, p, r) = \varphi_0(\theta, p, r) - \inf_{(\theta, p, r) \in \mathbb{R}^{3d}} \varphi_0(\theta, p, r) + 1$ and $\varphi_0(\theta, p, r) := hH(\theta, p, r) + aL(\theta) \cdot p + ap \cdot r$ for some $h > 0$ with $H(\theta, p, r)$ given in (1.13) and $L(\theta)$ in (B.10). More specifically, for any $\alpha, \gamma > 0$ and $a > 0$ small enough, there exist $h, A, B, C, D > 0$ such that

$$-\frac{\mathcal{L}_H W_\delta}{W_\delta} \geq A|\theta|^{2(k-1)} + B|p|^2 + C|r|^2 - D. \tag{2.20}$$

Now, we are ready to state the following large deviations result for the high-order Langevin dynamics (1.15).

Theorem 2.3. *Assume that $(\mathbf{z}_t)_{t \geq 0} = (\theta_t, p_t, r_t)_{t \geq 0}$ in (1.15) where U satisfies Assumption 2.4, and consider a smooth function κ with $\kappa(\theta, p, r) = 1 + |\theta|^\lambda + |p|^\sigma + |r|^\omega$ with*

$\lambda \in [0, 2), \sigma \in [0, 2), \omega \in [0, 2)$. Then $(\mathbf{z}_t)_{t \geq 0}$ is ergodic with respect to the measure μ , with Lyapunov function defined in (2.19). Moreover, the empirical measure

$$\pi_t := \frac{1}{t} \int_0^t \delta_{(\theta_s, p_s, r_s)} ds$$

satisfies LDP in the τ^κ -topology. Finally, for any $\nu \in \mathcal{P}_\kappa(\mathcal{X})$ such that $d\nu = e^v d\mu$, the rate function reads

$$I_H(\nu) = \frac{\alpha}{4} \int_{\mathcal{X}} |\nabla_r v|^2 d\nu + \frac{1}{4\alpha} \int_{\mathcal{X}} |\nabla_r \psi|^2 d\nu, \quad (2.21)$$

where ψ is the unique solution in $\mathcal{H}^1(\nu)$ to the Poisson equation

$$-\Delta_r \psi - \nabla_r v \cdot \nabla_r \psi = \mathcal{L}_{\text{HA}} v. \quad (2.22)$$

2.2.3 Hessian-free high-resolution dynamics

According to the choices of \mathcal{D} and \mathcal{Q} in (1.17), through formula (B.8), we can compute that the infinitesimal generator of (1.17) is given by

$$\begin{aligned} \mathcal{L}_{\text{RS}} &= -\beta \nabla U(\theta) \cdot \nabla_\theta - \alpha r \cdot \nabla_r + \beta \Delta_\theta + \alpha \Delta_r, \\ \mathcal{L}_{\text{RA}} &= r \cdot \nabla_\theta - \nabla U(\theta) \cdot \nabla_r, \end{aligned}$$

which implies that

$$\mathcal{L}_R = \mathcal{L}_{\text{RS}} + \mathcal{L}_{\text{RA}} = (-\beta \nabla U(\theta) + r) \cdot \nabla_\theta + (-\alpha r - \nabla U(\theta)) \cdot \nabla_r + \beta \Delta_\theta + \alpha \Delta_r. \quad (2.23)$$

In order to obtain the large deviation principle for the Hessian-free high-resolution dynamics (1.17), we make the following classical assumptions for the growth of potential function U .

Assumption 2.5. *The potential $U \in \mathcal{S}$ has compact level sets, satisfies $e^{-U} \in L^1(\mathcal{X})$ and*

- (a) *There exist $c_1 > 0, C_1 \in \mathbb{R}$ such that $\theta \cdot \nabla U(\theta) \geq c_1 |\theta|^2 - C_1$ for all $\theta \in \mathbb{R}^d$.*
- (b) *There exists $m_U > 0$ such that $|\nabla U(\theta_1) - \nabla U(\theta_2)| \leq m_U |\theta_1 - \theta_2|$ for all $\theta_1, \theta_2 \in \mathbb{R}^d$.*

The Hessian-free high-resolution dynamics (1.17) automatically satisfies Hypocoellipticity (Assumptions 2.1) and Controllability (Assumption 2.2) since its infinitesimal generator is an elliptic operator. In the next lemma, we will show that Lyapunov condition (Assumption 2.3) is satisfied for the Hessian-free high-resolution dynamics (1.17).

Lemma 2.7. *(Witten-Lyapunov condition). Suppose that $(\mathbf{z}_t)_{t \geq 0} = (\theta_t, r_t)_{t \geq 0}$ in (1.17) where potential U satisfies Assumption 2.5. Then for any $\alpha, \beta > 0$ and $a \in (0, 1)$, there exists $b > 0$ such that*

$$W_a(\theta, r) = e^{\varphi(\theta, r)} \quad (2.24)$$

is a Lyapunov function, where $\varphi(\theta, r) = aH(\theta, r) + b\theta \cdot r$ with $H(\theta, r)$ given in (1.16). More specifically, for any $\alpha, \beta > 0$ and $a \in (0, 1)$, there exist $A, B, C > 0$ such that

$$-\frac{\mathcal{L}_R W_a}{W_a} \geq A|\theta|^2 + B|r|^2 - C. \quad (2.25)$$

Now, we are ready to state the following large deviations result for the Hessian-free high-resolution dynamics (1.17).

Theorem 2.4. *Assume that $(\mathbf{z}_t)_{t \geq 0} = (\theta_t, r_t)_{t \geq 0}$ in (1.17) where U satisfies Assumption 2.5, and consider a smooth function κ with $\kappa(\theta, r) = 1 + |\theta|^\lambda + |r|^\omega$ and $\lambda \in [0, 2), \omega \in [0, 2)$. Then $(\mathbf{z}_t)_{t \geq 0}$ is ergodic with respect to the measure μ , with Lyapunov function defined in (2.24). Moreover, the empirical measure*

$$\pi_t := \frac{1}{t} \int_0^t \delta_{(\theta_s, r_s)} ds$$

satisfies a LDP in the τ^κ -topology. Finally, for any $\nu \in \mathcal{P}_\kappa(\mathcal{X})$ such that $d\nu = e^v d\mu$, the rate function reads

$$I_R(\nu) = \frac{\beta}{4} \left(\int_{\mathcal{X}} |\nabla_{\theta} v|^2 d\nu + \int_{\mathcal{X}} |\nabla_{\theta} \psi_v|^2 d\nu \right) + \frac{\alpha}{4} \left(\int_{\mathcal{X}} |\nabla_r v|^2 d\nu + \int_{\mathcal{X}} |\nabla_r \psi_v|^2 d\nu \right), \quad (2.26)$$

where ψ is the unique solution in $\mathcal{H}^1(\nu)$ to the Poisson equation

$$-\beta \Delta_{\theta} \psi_v - \alpha \Delta_r \psi_v + \beta \nabla_{\theta} \psi_v \cdot \nabla_{\theta} v + \alpha \nabla_r \psi_v \cdot \nabla_r v = \mathcal{L}_{\text{RA}} v. \quad (2.27)$$

3 Comparisons

In this section, by using the large deviation rate function of empirical measure as a tool to analyze the convergence rate to the invariant measure, and taking the large deviation rate function as a measure of the rate of convergence to equilibrium, we study whether variants of Langevin dynamics have acceleration effect over the overdamped Langevin dynamics. The intuition is that the larger the rate function, the more concentrated the empirical measure is around the invariant measure, thus indicating faster convergence. First, we consider the acceleration of the generalized Langevin dynamics (1.6) We have the following result.

Corollary 3.1. *Suppose that Assumptions 2.1-2.3 hold. For any $\nu \in \mathcal{P}(\mathcal{X} = \mathbb{R}^d)$ of the form $d\nu = e^v d\mu$ and $\nu \neq \mu$. If the matrix $\mathcal{D} - \mathbf{I}$ is a positive semidefinite diffusion matrix, then we have*

$$I_{\tau}(\nu) \geq I_o(\nu), \quad (3.1)$$

where $I_{\tau}(\nu)$ is the rate function of the generalized Langevin dynamics (1.6) and $I_o(\nu)$ is the rate function (2.12) of the overdamped Langevin dynamics (1.2).

Corollary 3.1 shows the acceleration effect of the generalized Langevin dynamics compared to the overdamped Langevin dynamics. In the rest of this section, we show the acceleration of variants of Langevin dynamics over the classical overdamped Langevin dynamics by comparing their rate functions from the large deviations. It is known that the rate function for non-reversible Langevin dynamics (1.5) is greater than that for overdamped Langevin dynamics (1.2), and hence the acceleration [FS20]. In the rest of this section, we will compare the rate functions for mirror Langevin dynamics (1.12), Hessian-free high-resolution dynamics (1.17), underdamped Langevin dynamics (1.4), and high-order Langevin dynamics (1.15) with that of overdamped Langevin dynamics (1.2).

Note that for underdamped Langevin dynamics, high-order Langevin dynamics, and Hessian-free high-resolution dynamics, these processes live in higher dimensions than overdamped Langevin dynamics, which makes a direct comparison of their rate functions impossible. We introduce a novel method to expand the space of overdamped Langevin dynamics to match the dimension with that of the variant of Langevin dynamics under study, e.g. expanding to the space of overdamped Langevin dynamics to \mathbb{R}^{2d} when we compare it with underdamped Langevin dynamics and Hessian-free high-resolution dynamics and to \mathbb{R}^{2d} when we compare it with high-order Langevin dynamics. If the comparison can be made on the whole expanded space, we then apply the contraction principle from the large deviations theory to compare the rate functions for the large deviations of $\frac{1}{t} \int_0^t \delta_{\theta_s} ds$.

First, we compare the rate function of the mirror Langevin dynamics (1.12) with that of the overdamped Langevin dynamics (1.2). The following result can be obtained as a corollary of Corollary 3.1.

Corollary 3.2. *Assume the same condition as Theorem 2.2. For any $\nu \in \mathcal{P}(\mathcal{X} = \mathbb{R}^d)$ of the form $d\nu = e^\nu d\mu$ and $\nu \neq \mu$. Then for the mirror Langevin dynamics defined by (1.12), if the matrix $[\nabla^2 \phi(\mathbf{z})]^{-1} - I$ is a positive semidefinite diffusion matrix, then we have*

$$I_M(\nu) \geq I_o(\nu), \quad (3.2)$$

where $I_o(\nu)$ is the rate function of the overdamped Langevin dynamics (1.2) and $I_M(\nu)$ is the rate function of the mirror Langevin dynamics (2.17).

Corollary 3.2 shows the acceleration effect of the mirror Langevin dynamics (1.12) compared to the overdamped Langevin dynamics (1.2).

Next, we compare the rate functions of Hessian-free high-resolution dynamics (1.17) and underdamped Langevin dynamics (1.4) with overdamped Langevin dynamics (1.2). Note that overdamped Langevin dynamics lives in \mathbb{R}^d , whereas Hessian-free high-resolution dynamics and underdamped Langevin dynamics live in \mathbb{R}^{2d} . Thus, we lift the overdamped Langevin dynamics to \mathbb{R}^{2d} , before we do the comparison. We consider the *expanded second-order overdamped Langevin dynamics*, for $(\theta_t, r_t) \in \mathbb{R}^d \times \mathbb{R}^d$,

$$\begin{cases} d\theta_t = -\nabla U(\theta_t)dt + \sqrt{2}dW_t, \\ dr_t = -r_t dt + \sqrt{2}dB_t, \end{cases} \quad (3.3)$$

where W_t and B_t are independent d -dimensional Brownian motions. We can easily check that its invariant measure is $d\mu = e^{-U(\theta) - \frac{1}{2}|r|^2} d\theta dr$, that is the same as the invariant measure of Hessian-free high-resolution dynamics. By LDP for the overdamped Langevin dynamics (Lemma 2.2), the LDP rate function for the expanded second-order overdamped Langevin dynamics (3.3) is

$$I_{e2o}(\nu) = \frac{1}{4} \int_{\mathcal{X}} |\nabla v|^2 d\nu, \quad (3.4)$$

where $\nu \in \mathcal{P}(\mathcal{X} = \mathbb{R}^{2d})$ of the form $d\nu = e^v d\mu$ and $\nu \neq \mu$.

Proposition 3.1. *Assume the same condition as Theorem 2.4. For any $\nu \in \mathcal{P}(\mathcal{X} = \mathbb{R}^{2d})$ of the form $d\nu = e^v d\mu$ and $\nu \neq \mu$. Then for the Hessian-free high-resolution dynamics defined by (1.17), if $\min(\alpha, \beta) \geq 1$, then we have*

$$I_R(\nu) \geq I_{e2o}(\nu), \quad (3.5)$$

where $I_{e2o}(\nu)$ is in (3.4) the rate function of the expanded second-order overdamped Langevin dynamics (3.3) and $I_R(\nu)$ is the rate function of the Hessian-free high-resolution dynamics (1.17).

Proposition 3.1 show the acceleration effect of the Hessian-free high-resolution dynamics (1.17) compared to the expanded second-order overdamped Langevin dynamics (3.3). Indeed, for the Hessian-free high-resolution dynamics (1.17), the θ -marginal distribution of its stationary distribution is the target distribution $\mu(\theta) \propto e^{-U(\theta)}$ which coincides with the θ -marginal distribution of the stationary distribution of the expanded second-order overdamped Langevin dynamics (3.3), which is indeed the stationary distribution of the overdamped Langevin dynamics (1.2). By contraction principle (see e.g. Theorem 4.2.1. in [DZ98]), for the Hessian-free high-resolution dynamics (1.17), $\frac{1}{t} \int_0^t \delta_{\theta_s} ds$ satisfies a LDP in the τ^κ -topology with the rate function

$$I_{R,\theta}(\nu_\theta) := \inf_{\nu \in \mathcal{P}_\kappa(\mathbb{R}^{2d}): M_\theta(\nu) = \nu_\theta} I_R(\nu), \quad (3.6)$$

where $M_\theta(\nu)$ is the θ -marginal distribution of ν . Similarly, by contraction principle, the rate function (2.12) for overdamped Langevin dynamics (1.2) can be re-written as

$$I_o(\nu_\theta) := \inf_{\nu \in \mathcal{P}_\kappa(\mathbb{R}^{2d}): M_\theta(\nu) = \nu_\theta} I_{e2o}(\nu). \quad (3.7)$$

Therefore, we obtain the following corollary from Proposition 3.1.

Corollary 3.3. *For any $\nu_\theta \in \mathcal{P}(\mathbb{R}^d)$,*

$$I_{R,\theta}(\nu_\theta) \geq I_o(\nu_\theta), \quad (3.8)$$

where $I_{R,\theta}$ and I_o are defined in (3.6) and (2.12).

Corollary 3.3 shows the acceleration effect of the Hessian-free high-resolution dynamics (1.17) compared to the expanded second-order overdamped Langevin dynamics (3.3) in terms of the convergence to the target distribution $\mu(\theta) \propto e^{-U(\theta)}$.

Next, we compare the rate functions of underdamped Langevin dynamics (1.4) and high-order Langevin dynamics (1.15) with overdamped Langevin dynamics (1.2). We will make comparisons on a subspace of the probability measures that is defined as follows. It remains an open problem how to compare the rate functions on the whole space of the probabilities for underdamped Langevin dynamics (1.4) and high-order Langevin dynamics (1.15), which will be left as a future research direction.

Definition 3.1. *The class $\mathcal{P}^H(\mathcal{X})$ consists of $\nu \in \mathcal{P}(\mathcal{X} = \mathbb{R}^d \times \mathbb{R}^d \times \dots \times \mathbb{R}^d)$ of the form $d\nu = e^v d\mu$ and $\nu \neq \mu$, where v is a function of the variable associated the last \mathbb{R}^d component of $\mathcal{X} = \mathbb{R}^d \times \mathbb{R}^d \times \dots \times \mathbb{R}^d$.*

We have the following comparison result for the rate function of underdamped Langevin dynamics (1.4) with overdamped Langevin dynamics (1.2).

Proposition 3.2. *Assume the same condition as Assumption 4.4. in [FS20]. For any $\nu(\theta, r) \in \mathcal{P}^H(\mathcal{X} = \mathbb{R}^{2d})$, ν is of the form $d\nu = e^v d\mu$ and $\nu \neq \mu$ for $v = v(r)$. Then for the underdamped Langevin dynamics defined by (1.4), if $\gamma \geq 1$, then we have*

$$I_u(\nu) \geq I_{e_{2o}}(\nu), \quad (3.9)$$

where $I_{e_{2o}}(\nu)$ is the rate function of the expanded second-order overdamped Langevin dynamics (3.3) and $I_u(\nu)$ is the rate function of the underdamped Langevin dynamics (1.4).

Proposition 3.2 shows the acceleration effect of the underdamped Langevin dynamics (1.4) compared to the expanded second-order overdamped Langevin dynamics (3.3) on a class of probability measures $\mathcal{P}^H(\mathcal{X} = \mathbb{R}^{2d})$.

Finally, we compare the rate functions of high-order Langevin dynamics (1.15) with overdamped Langevin dynamics (1.2). Note that overdamped Langevin dynamics lives in \mathbb{R}^d , whereas high-order Langevin dynamics live in \mathbb{R}^{3d} . Thus, we lift the overdamped Langevin dynamics to \mathbb{R}^{3d} , before we do the comparison. We consider the *expanded third-order overdamped Langevin dynamics*, for $(\theta_t, p_t, r_t) \in \mathbb{R}^d \times \mathbb{R}^d \times \mathbb{R}^d$,

$$\begin{cases} d\theta_t = -\nabla U(\theta_t)dt + \sqrt{2}dW_t, \\ dp_t = -p_t dt + \sqrt{2}dB_t^1, \\ dr_t = -r_t dt + \sqrt{2}dB_t^2, \end{cases} \quad (3.10)$$

where W_t , B_t^1 and B_t^2 are independent d -dimensional Brownian motions. We can easily check that its invariant measure is $d\mu = e^{-U(\theta) - \frac{1}{2}|p|^2 - \frac{1}{2}|r|^2} d\theta dp dr$, that is the same as the invariant measure of high-order Langevin dynamics. By LDP for the overdamped Langevin

dynamics (Lemma 2.2), the LDP rate function for the expanded third-order overdamped Langevin dynamics (3.10) is

$$I_{e_{3o}}(\nu) = \frac{1}{4} \int_{\mathcal{X}} |\nabla v|^2 d\nu, \quad (3.11)$$

where $\nu \in \mathcal{P}(\mathcal{X} = \mathbb{R}^{3d})$ of the form $d\nu = e^v d\mu$ and $\nu \neq \mu$.

Proposition 3.3. *Assume the same condition as Assumption 4.4. in [FS20]. For any $\nu(\theta, p, r) \in \mathcal{P}^H(\mathcal{X} = \mathbb{R}^{3d})$, ν is of the form $d\nu = e^v d\mu$ and $\nu \neq \mu$ for $v = v(r)$. Then for the high-order Langevin dynamics defined by (1.15), if $\alpha \geq 1$, then we have*

$$I_H(\nu) \geq I_{e_{3o}}(\nu), \quad (3.12)$$

where $I_{e_{3o}}(\nu)$ is the rate function of the expanded third-order overdamped Langevin dynamics (3.10) and $I_H(\nu)$ is the rate function of the high-order Langevin dynamics (2.21).

Proposition 3.3 shows the acceleration effect of the underdamped Langevin dynamics (1.15) compared to the expanded third-order overdamped Langevin dynamics (3.10) on a class of probability measures $\mathcal{P}^H(\mathcal{X} = \mathbb{R}^{3d})$.

4 Numerical Experiments

In this section, we conduct numerical experiments for the variants of Langevin algorithms based on the Euler–Maruyama discretization of underdamped Langevin dynamics (1.4), non-reversible Langevin dynamics (1.5), mirror Langevin dynamics (1.12), high-order Langevin dynamics (1.15) and Hessian-free high-resolution dynamics (1.17), and compare our results with the unadjusted Langevin algorithm, i.e. the Euler–Maruyama discretization of overdamped Langevin dynamics (1.2).⁵ We focus on applying variants of Langevin algorithms to Bayesian logistic regression using both synthetic and real data. We first introduce the Bayesian logistic regression framework.

4.1 Bayesian logistic regression framework

Consider a dataset $\mathcal{A} = \{(X_j, y_j)\}_{j=1}^n$, where $X_j \in \mathbb{R}^d$ denotes independent feature vectors and $y_j \in \{0, 1\}$ represents binary labels. The likelihood function follows the logistic model:

$$\mathbb{P}(y_j = 1 | X_j, x) = \sigma(x^\top X_j) = \left(1 + e^{-x^\top X_j}\right)^{-1}, \quad (4.1)$$

where $\sigma(\cdot)$ denotes the sigmoid function and $x \in \mathbb{R}^d$ denotes the regression coefficients.

⁵In the literature, finer discretizations for Langevin algorithms have been used; see e.g. [CCBJ18, DRD20]. To illustrate our theory, which is based on the continuous-time diffusion, instead of showing acceleration due to finer discretizations, we use Euler–Maruyama discretization for all the variants of Langevin dynamics for the sake of fair comparison.

Under the Bayesian paradigm, we impose a Gaussian prior $p(x) = \mathcal{N}(0, \lambda I)$ with $\lambda = 10$ to regularize the parameter space. The posterior distribution $\pi(x) \propto e^{-U(x)}$ combines the likelihood and prior through the potential function:

$$U(x) = - \sum_{j=1}^n \log \mathbb{P}(y_j | X_j, x) - \log p(x) = \sum_{j=1}^n \log \left(1 + e^{-x^\top X_j} \right) + \frac{1}{2\lambda} |x|^2. \quad (4.2)$$

Note that the function $U(x)$ in (4.2) is strongly convex and smooth due to the ℓ_2 -regularization term inherited from the Gaussian prior. This guarantees unique identifiability of the parameters and facilitates efficient sampling.

4.2 Bayesian logistic regression with synthetic data

In this section, we validate our methodology through synthetic data experiments. Let feature vectors $X_j \sim \mathcal{N}(0, 10I_d)$, true parameters $x = [x_1, x_2, \dots, x_d]^\top$ with prior $x \sim \mathcal{N}(0, 10I_d)$ and binary labels for each (X_j, p_j) where $p_j \sim \mathcal{U}[0, 1]$, and $y_j = 1$ if $p_j \leq \sigma(x_{\text{true}}^\top X_j)$ and $y_j = 0$ otherwise. This data generation process ensures the labels y_j adhere to the logistic model while maintaining controlled experimental conditions. The uniform threshold introduces stochasticity consistent with the Bernoulli likelihood structure.

For synthetic data, we generated 5000 data points with d random features and generated binary labels. For synthetic data, for non-reversible Langevin, we generated the anti-symmetric matrix J by first generating a $d \times d$ random matrix A with standard normal random entries and then we set $J = A - A^\top$ such that J is an anti-symmetric matrix i.e $J^\top = -J$; for mirror Langevin, we set the mirror map as $h(x) = \frac{1}{4} \sum_{i=1}^d x_i^4$ for any $x = (x_1, \dots, x_d) \in \mathbb{R}^d$. We chose the stepsize $\eta = 0.0003$ for overdamped, non-reversible, Hessian-free high-resolution and mirror Langevin and the stepsize $\eta = 0.003$ for underdamped and high-order Langevin. We chose $\gamma = 4$ for underdamped Langevin, $\gamma = 20$ and $\alpha = 15$ for high-order Langevin, and $\gamma = 1$ and $\alpha = 30$ for Hessian-free high-resolution.

The accuracy is reported only for test data sets. Our numerical results using synthetic data summarized in Figure 1 show that under for proper choices of hyperparameters (and the anti-symmetric matrix for non-reversible Langevin and the mirror map for mirror Langevin), the variants of Langevin algorithms, including underdamped Langevin (Figure 1b), non-reversible Langevin (Figure 1c), high-order Langevin (Figure 1d), Hessian-free high-resolution (Figure 1e) and mirror Langevin (Figure 1f) can all have faster convergence than overdamped Langevin (Figure 1a). On the other hand, the performance of these variants of Langevin algorithms is sensitive to the choices of hyperparameters (and the anti-symmetric matrix for non-reversible Langevin and the mirror map for mirror Langevin). In another set of experiments reported in Figure 2, with a slight change in the choices of hyperparameters, underdamped Langevin (Figure 2b), high-order Langevin (Figure 2d) and Hessian-free high-resolution (Figure 2e) can outperform overdamped Langevin (Figure 2a), whereas mirror Langevin (Figure 2f) and non-reversible (Figure 2c) cannot, even though

their performance is comparable with overdamped Langevin (Figure 2a). One possible explanation is that the performance of non-reversible Langevin dynamics (1.5) depends on the choice of the anti-symmetric matrix, and the performance of mirror Langevin dynamics (1.12) depends on the choice of the mirror map, such that when the hyperparameters are changed, one has to fine tune the choices of the anti-symmetric matrix and the mirror map simultaneously in order to maintain the good performance.




Figure 1: The plots show the accuracy over the synthetic data with dimension 569×31 , in which all variants of the Langevin algorithms outperform overdamped Langevin algorithm in Figure 1a with an appropriate choice of hyperparameters.

4.3 Bayesian logistic regression with real data

In this section, we validate our methodology through real data experiments. We consider the UCI ML Breast Cancer Wisconsin (Diagnostic) data set [WMSS95]. The data set contains 569 instances with 31 features, where each sample describes characteristics of the cell nuclei present in a digitized image of a fine needle aspirate (FNA) of a breast mass.

For real data, we generated the anti-symmetric matrix for non-reversible Langevin and the mirror map $h(x)$ for mirror Langevin similarly as for synthetic data, and kept the same stepsizes as well. We chose $\gamma = 35$ for underdamped Langevin, $\gamma = 35$ and $\alpha = 35$ for high-order Langevin, and $\gamma = 1$ and $\alpha = 30$ for Hessian-free high-resolution.




Figure 2: With a slight change of hyperparameters, we can see from this figure that underdamped Langevin (Figure 2b), high-order Langevin (Figure 2d) and Hessian-free high-resolution (Figure 2e) can outperform overdamped Langevin (Figure 2a); however, mirror Langevin (Figure 2f) and non-reversible (Figure 2c) cannot, even though their performance is comparable with overdamped Langevin (Figure 2a).

The accuracy is reported only for test data sets. Our numerical results using real data show that for some particular selection of hyperparameters, the variants of Langevin algorithms can all outperform the overdamped Langevin algorithm (Figure 3). These experiments demonstrate the practical applicability of our methods and validate their performance on real-world classification tasks. On the other hand, under a different choice of hyperparameters as in Figure 4, underdamped, high-order Langevin and Hessian-free high-resolution algorithms have faster convergence than overdamped Langevin; see Figures 4b, 4e, 4d and 4a; whereas non-reversible Langevin, and mirror Langevin exhibit similar performance compared with overdamped Langevin; see Figures 4c, and 4a. Non-reversible and mirror Langevin seem to be more sensitive to the choice of hyperparameters, which might be due to the fact that their performance also depends on the choice of the anti-symmetric matrix and the mirror map.




Figure 3: The plots show the accuracy over the real data with dimension 569×31 , in which all variants of the Langevin algorithms outperform overdamped Langevin algorithm in Figure 3a with an appropriate choice of hyperparameters.

5 Concluding Remarks

In this paper, we studied variants of Langevin dynamics through the lens of large deviations theory. We showed the acceleration of convergence of the variants of overdamped Langevin dynamics to the Gibbs distribution, including underdamped Langevin dynamics, non-reversible Langevin dynamics, mirror Langevin dynamics, high-order Langevin dynamics, Hessian-free high-resolution dynamics by comparing the rate functions from the large deviations theory. We provided numerical experiments using both synthetic and real data, based on the Euler–Maruyama discretizations of these variants of Langevin dynamics and demonstrated their efficiency.

Acknowledgements

Nian Yao was supported in part by the Natural Science Foundation of China under Grant 12071361, the Natural Science Foundation of Guangdong Province under Grant 2020A1515010822 and Shenzhen Natural Science Fund (the Stable Support Plan Program 20220810152104001). Lingjiong Zhu is partially supported by the NSF grants DMS-2053454, DMS-2208303 and a Simons Foundation Collaboration Grant. The first author




Figure 4: With a slight change of hyperparameters, we can see from this figure that underdamped Langevin (Figure 4b), high-order Langevin (Figure 4d) and Hessian-free high-resolution (Figure 2e) can outperform overdamped Langevin (Figure 4a); however, mirror Langevin (Figure 4f) and non-reversible (Figure 4c) cannot, even though their performance is comparable with overdamped Langevin (Figure 4a).

would like to thank Prof. Liming Wu, Prof. Fuqing Gao and Prof. Qi Lv for the helpful discussions. The second and fourth authors would like to thank Prof. Mert Gürbüzbalaban and Dr. Yuanhan Hu.

References

- [AC21] Kwangjun Ahn and Sinho Chewi. Efficient constrained sampling via the mirror-Langevin algorithm. In *Advances in Neural Information Processing Systems (NeurIPS)*, volume 34, 2021.
- [BDMP17] Nicolas Brosse, Alain Durmus, Éric Moulines, and Marcelo Pereyra. Sampling from a log-concave distribution with compact support with proximal Langevin Monte Carlo. In *Proceedings of the 2017 Conference on Learning Theory*, volume 65, pages 319–342. PMLR, 2017.
- [Bel06] Luc Rey Bellet. Ergodic properties of Markov processes. In Stéphane Attal, Alain Joye, and Claude-Alain Pillet, editors, *Open Quantum Systems II: The*

- Markovian Approach*, pages 1–39. Springer Berlin Heidelberg, Berlin, Heidelberg, 2006.
- [BEL15] Sebastien Bubeck, Ronen Eldan, and Joseph Lehec. Finite-time analysis of projected Langevin Monte Carlo. In *Advances in Neural Information Processing Systems*, volume 28, 2015.
- [BEL18] Sébastien Bubeck, Ronen Eldan, and Joseph Lehec. Sampling from a log-concave distribution with projected Langevin Monte Carlo. *Discrete & Computational Geometry*, 59(4):757–783, 2018.
- [BGL13] Dominique Bakry, Ivan Gentil, and Michel Ledoux. *Analysis and Geometry of Markov Diffusion Operators*. Springer Science & Business Media, MR-3155209, volume 348 of *grundlehren der mathematischen wissenschaften edition*, 2013.
- [CB18] Xiang Cheng and Peter L. Bartlett. Convergence of Langevin MCMC in KL-divergence. In *Proceedings of the 29th International Conference on Algorithmic Learning Theory (ALT)*, volume 83, pages 186–211. PMLR, 2018.
- [CCA⁺18] Xiang Cheng, Niladri S. Chatterji, Yasin Abbasi-Yadkori, Peter L. Bartlett, and Michael I. Jordan. Sharp Convergence Rates for Langevin Dynamics in the Nonconvex Setting. *arXiv:1805.01648*, 2018.
- [CCBJ18] Xiang Cheng, Niladri S Chatterji, Peter L Bartlett, and Michael I Jordan. Underdamped Langevin MCMC: A non-asymptotic analysis. In *Proceedings of the 31st Conference On Learning Theory*, volume 75, pages 300–323. PMLR, 2018.
- [CCD⁺19] Yi Chen, Jinglin Chen, Jing Dong, Jian Peng, and Zhaoran Wang. Accelerating nonconvex learning via replica exchange Langevin diffusion. In *International Conference on Learning Representations (ICLR)*, 2019.
- [CHS87] Tzoo-Shuh Chiang, Chii-Ruey Hwang, and Shuenn Jyi Sheu. Diffusion for global optimization in \mathbb{R}^n . *SIAM Journal on Control and Optimization*, 25(3):737–753, 1987.
- [CLGL⁺20] Sinho Chewi, Thibaut Le Gouic, Cheng Lu, Tyler Maunu, Philippe Rigollet, and Austin Stromme. Exponential ergodicity of mirror-Langevin diffusions. In *Advances in Neural Information Processing Systems (NeurIPS)*, 2020.
- [CLW23] Yu Cao, Jianfeng Lu, and Lihan Wang. On explicit L^2 -convergence rate estimate for underdamped Langevin dynamics. *Archive for Rational Mechanics and Analysis*, 247(90):1–34, 2023.

- [Dal17] Arnak S Dalalyan. Theoretical guarantees for approximate sampling from smooth and log-concave densities. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 79(3):651–676, 2017.
- [DFG⁺20] Wei Deng, Qi Feng, Liyao Gao, Faming Liang, and Guang Lin. Non-convex learning via replica exchange stochastic gradient MCMC. In *Proceedings of the 37th International Conference on Machine Learning*, volume 119, pages 2474–2483. PMLR, 2020.
- [DFT⁺25] Hengrong Du, Qi Feng, Changwei Tu, Xiaoyu Wang, and Lingjiong Zhu. Non-reversible Langevin algorithms for constrained sampling. *arXiv:2501.11743*, 2025.
- [DK19] Arnak S. Dalalyan and Avetik G. Karagulyan. User-friendly guarantees for the Langevin Monte Carlo with inaccurate gradient. *Stochastic Processes and their Applications*, 129(12):5278–5311, 2019.
- [DLP16] Andrew B. Duncan, Tony Lelièvre, and Grigoris A. Pavliotis. Variance reduction using nonreversible Langevin samplers. *Journal of Statistical Physics*, 163(3):457–491, 2016.
- [DM17] Alain Durmus and Eric Moulines. Non-asymptotic convergence analysis for the Unadjusted Langevin Algorithm. *Annals of Applied Probability*, 27(3):1551–1587, 2017.
- [DM19] Alain Durmus and Eric Moulines. High-dimensional Bayesian inference via the Unadjusted Langevin Algorithm. *Bernoulli*, 25(4A):2854–2882, 2019.
- [DMP18] Alain Durmus, Eric Moulines, and Marcelo Pereyra. Efficient Bayesian computation by proximal Markov Chain Monte Carlo: When Langevin meets Moreau. *SIAM Journal on Imaging Sciences*, 11(1):473–506, 2018.
- [DPZ17] Andrew B. Duncan, Grigoris A. Pavliotis, and Konstantinos C. Zygalakis. Nonreversible Langevin samplers: Splitting schemes, analysis and implementation. *arXiv preprint arXiv:1701.04247*, 2017.
- [DRD20] Arnak S Dalalyan and Lionel Riou-Durand. On sampling from a log-concave density using kinetic Langevin diffusions. *Bernoulli*, 26(3):1956–1988, 2020.
- [DRW⁺16] Kumar Avinava Dubey, Sashank J Reddi, Sinead A Williamson, Barnabas Poczos, Alexander J Smola, and Eric P Xing. Variance reduction in stochastic gradient Langevin dynamics. In *Advances in Neural Information Processing Systems*, pages 1154–1162, 2016.

- [DV75a] Monroe D. Donsker and S. R. Srinivasa Varadhan. Asymptotic evaluation of certain Markov process expectations for large time, I. *Communications on Pure and Applied Mathematics*, 28(1):1–47, 1975.
- [DV75b] Monroe D. Donsker and S. R. Srinivasa Varadhan. Asymptotic evaluation of certain Markov process expectations for large time, II. *Communications on Pure and Applied Mathematics*, 28(2):279–301, 1975.
- [DV76] Monroe D. Donsker and S. R. Srinivasa Varadhan. Asymptotic evaluation of certain Markov process expectations for large time, III. *Communications on Pure and Applied Mathematics*, 29(4):389–461, 1976.
- [DV83] Monroe D. Donsker and S. R. Srinivasa Varadhan. Asymptotic evaluation of certain Markov process expectations for large time, IV. *Communications on Pure and Applied Mathematics*, 36(2):183–212, 1983.
- [DZ98] Amir Dembo and Ofer Zeitouni. *Large Deviations Techniques and Applications*. Springer, New York, 2nd edition, 1998.
- [FS20] Grégoire Ferré and Gabriel Stoltz. Large deviations of empirical measures of diffusions in weighted topologies. *Electronic Journal of Probability*, 25(121):1–52, 2020.
- [FSS20] Futoshi Futami, Iseki Sato, and Masashi Sugiyama. Accelerating the diffusion-based ensemble sampling by non-reversible dynamics. In *Proceedings of the 37th International Conference on Machine Learning*, volume 119, pages 3337–3347. PMLR, 2020.
- [GGHZ21] Mert Gürbüzbalaban, Xuefeng Gao, Yunhan Hu, and Lingjiong Zhu. Decentralized stochastic gradient Langevin dynamics and Hamiltonian Monte Carlo. *Journal of Machine Learning Research*, 22(239):1–69, 2021.
- [GGZ20] Xuefeng Gao, Mert Gürbüzbalaban, and Lingjiong Zhu. Breaking reversibility accelerates Langevin dynamics for global non-convex optimization. In *Advances in Neural Information Processing Systems (NeurIPS)*, volume 33, 2020.
- [GGZ22] Xuefeng Gao, Mert Gürbüzbalaban, and Lingjiong Zhu. Global convergence of Stochastic Gradient Hamiltonian Monte Carlo for non-convex stochastic optimization: Non-asymptotic performance bounds and momentum-based acceleration. *Operations Research*, 70(5):2931–2947, 2022.
- [GHZ24] Mert Gürbüzbalaban, Yuanhan Hu, and Lingjiong Zhu. Penalized overdamped and underdamped Langevin Monte Carlo algorithms for constrained sampling. *Journal of Machine Learning Research*, 25(263):1–67, 2024.

- [GIWZ24] Mert Gürbüzbalaban, Mohammad Rafiqul Islam, Xiaoyu Wang, and Lingjiong Zhu. Generalized EXTRA stochastic gradient Langevin dynamics. *arXiv preprint arXiv:2412.01993*, 2024.
- [GLNW24] Arnaud Guillin, Di Lu, Boris Nectoux, and Liming Wu. Generalized Langevin and Nosé-Hoover processes absorbed at the boundary of a metastable domain. *arXiv:2403.17471*, 2024.
- [HHMS93] Chii-Ruey Hwang, Shu-Yin Hwang-Ma, and Shuenn-Jyi Sheu. Accelerating Gaussian diffusions. *Annals of Applied Probability*, 3:897–913, 1993.
- [HHMS05] Chii-Ruey Hwang, Shu-Yin Hwang-Ma, and Shuenn-Jyi Sheu. Accelerating diffusions. *Annals of Applied Probability*, 15:1433–1444, 2005.
- [HKRC18] Ya-Ping Hsieh, Ali Kavis, Paul Rolland, and Volkan Cevher. Mirrored Langevin dynamics. In *Advances in Neural Information Processing Systems (NeurIPS)*, 2018.
- [HKS89] Richard A Holley, Shigeo Kusuoka, and Daniel W Stroock. Asymptotics of the spectral gap with applications to the theory of simulated annealing. *Journal of Functional Analysis*, 83(2):333–347, 1989.
- [Hof09] Peter D Hoff. *A First Course in Bayesian Statistical Methods*, volume 580. Springer, 2009.
- [HWG⁺20] Yuanhan Hu, Xiaoyu Wang, Xuefeng Gao, Gürbüzbalaban, and Lingjiong Zhu. Non-convex stochastic optimization via non-reversible stochastic gradient Langevin dynamics. *arXiv:2004.02823*, 2020.
- [Lam21] Andrew Lamperski. Projected stochastic gradient Langevin algorithms for constrained sampling and non-convex learning. In *Conference on Learning Theory*, volume 134, pages 2891–2937. PMLR, 2021.
- [LTVW22] Ruilin Li, Molei Tao, Santosh S. Vempala, and Andre Wibisono. The mirror Langevin algorithm converges with vanishing bias. In *Proceedings of The 33rd International Conference on Algorithmic Learning Theory*, volume 167, pages 718–742. PMLR, 2022.
- [LZT22] Ruilin Li, Hongyuan Zha, and Molei Tao. Hessian-free high-resolution Nesterov acceleration for sampling. In *Proceedings of the 39th International Conference on Machine Learning*, volume 162, pages 13125–13162. PMLR, 2022.
- [MCC⁺21] Yi-An Ma, Niladri S. Chatterji, Xiang Cheng, Nicolas Flammarion, Peter L. Bartlett, and Michael I. Jordan. Is there an analog of Nesterov acceleration for gradient-based MCMC? *Bernoulli*, 27(3):1942–1992, 2021.

- [MCF15] Yi-An Ma, Tianqi Chen, and Emily Fox. A complete recipe for stochastic gradient MCMC. In *Advances in Neural Information Processing Systems (NIPS)*, pages 2917–2925, 2015.
- [MMW⁺21] Wenlong Mou, Yi-An Ma, Martin J. Wainwright, Peter L. Bartlett, and Michael I. Jordan. High-order Langevin diffusion yields an accelerated MCMC algorithm. *Journal of Machine Learning Research*, 42:1–41, 2021.
- [Mon23] Pierre Monmarché. Almost sure contraction for diffusions on \mathbb{R}^d . application to generalized Langevin diffusions. *Stochastic Processes and their Applications*, 161:316–349, 2023.
- [PS17] Nicholas G. Polson and Vadim Sokolov. Deep learning: A Bayesian perspective. *Bayesian Analysis*, 12(4):1275–1304, 12 2017.
- [RBS15] Luc Rey-Bellet and Konstantinos Spiliopoulos. Irreversible Langevin samplers and variance reduction: a large deviation approach. *Nonlinearity*, 28:2081–2103, 2015.
- [RRT17] Maxim Raginsky, Alexander Rakhlin, and Matus Telgarsky. Non-convex learning via stochastic gradient Langevin dynamics: a nonasymptotic analysis. In *Proceedings of the 2017 Conference on Learning Theory*, volume 65, pages 1674–1703. PMLR, 2017.
- [Şim17] Umut Şimşekli. Fractional Langevin Monte Carlo: Exploring Lévy driven stochastic differential equations for Markov Chain Monte Carlo. In *International Conference on Machine Learning*, pages 3200–3209. PMLR, 2017.
- [SR20] Adil Salim and Peter Richtárik. Primal dual interpretation of the proximal stochastic gradient Langevin algorithm. In *Advances in Neural Information Processing Systems (NeurIPS)*, volume 33, 2020.
- [ŞZTG20] Umut Şimşekli, Lingjiong Zhu, Yee Whye Teh, and Mert Gürbüzbalaban. Fractional underdamped Langevin dynamics: Retargeting SGD with momentum under heavy-tailed gradient noise. In *International Conference on Machine Learning*, volume 119, pages 8970–8980. PMLR, 2020.
- [Var66] S. R. Srinivasa Varadhan. Asymptotic probabilities and differential equations. *Communications on Pure and Applied Mathematics*, 19(3):261–286, 1966.
- [Var84] S. R. Srinivasa Varadhan. *Large Deviations and Applications*. SIAM, Philadelphia, 1984.
- [Var08] S. R. Srinivasa Varadhan. Large deviations. *Annals of Probability*, 36:397–419, 2008.

- [WMSS95] William Wolberg, Olvi Mangasarian, Nick Street, and W. Street. Breast Cancer Wisconsin (Diagnostic). UCI Machine Learning Repository, 1995. DOI: <https://doi.org/10.24432/C5DW2B>.
- [WY20] Hao Wang and Dit-Yan Yeung. A survey on Bayesian deep learning. *ACM Computing Surveys*, 53(5):1–37, 2020.
- [ZL22] Yuping Zheng and Andrew Lamperski. Constrained Langevin algorithms with L-mixing external random variables. In *Advances in Neural Information Processing Systems (NeurIPS)*, volume 35, 2022.
- [ZPFP20] Kelvin Shuangjian Zhang, Gabriel Peyré, Jalal Fadili, and Marcelo Pereyra. Wasserstein control of mirror Langevin Monte Carlo. In *Proceedings of Thirty Third Conference on Learning Theory*, volume 125, pages 3814–3841. PMLR, 2020.

A Notations

For a function $f : \mathbb{R}^n \rightarrow \mathbb{R}$, we introduce the following notations:

- The gradient of f is defined as $\nabla f := \left(\frac{\partial f}{\partial x_1}, \frac{\partial f}{\partial x_2}, \dots, \frac{\partial f}{\partial x_n} \right)$.
- The Hessian matrix of f is defined as $\nabla^2 f := \left(\frac{\partial^2 f}{\partial x_i \partial x_j} \right)_{i,j=1}^n$.
- The third-order tensor of partial derivatives of f is defined as $\nabla^3 f := \left(\frac{\partial^3 f}{\partial x_i \partial x_j \partial x_k} \right)_{i,j,k=1}^n$.
- The Laplacian of f is given by $\Delta f := \nabla \cdot \nabla f = \sum_{i=1}^n \frac{\partial^2 f}{\partial x_i^2}$.

For a vector field $\mathbf{v} : \mathbb{R}^n \rightarrow \mathbb{R}^n$, its divergence is defined as $\nabla \cdot \mathbf{v} := \sum_{i=1}^n \frac{\partial v_i}{\partial x_i}$.

For an operator $\mathcal{L} := b \cdot \nabla + \mathcal{S} : \nabla^2$, the carré du champ operator Γ associated with \mathcal{L} is defined as follows; see e.g. [BGL13]. For two regular functions φ, ψ :

$$\Gamma(\varphi, \psi) := \frac{1}{2} (\mathcal{L}(\varphi\psi) - \varphi\mathcal{L}\psi - \psi\mathcal{L}\varphi) = \nabla\varphi \cdot \mathcal{S}\nabla\psi.$$

Next, we define various spaces for real-valued functions on \mathcal{X} that are used throughout the paper as follows.

- $C^\infty(\mathcal{X})$ is the space of smooth functions.
- $C_c^\infty(\mathcal{X})$ is the space of smooth functions with compact support.
- $C^k(\mathcal{X})$ is the space of functions are smooth to the k -th derivative.

- $C_b(\mathcal{X})$ is the space of continuous and bounded functions.
- \mathcal{S} is the space of smooth functions growing at most polynomially and whose derivatives also grow at most polynomially:

$$\mathcal{S} = \left\{ \varphi \in C^\infty(\mathcal{X}) \mid \forall \alpha \in \mathbb{N}^d, \exists N > 0 \text{ such that } \sup_{x \in \mathcal{X}} \frac{|\partial^\alpha \varphi(x)|}{(1 + |x|^2)^N} < +\infty \right\},$$

where $\partial^\alpha = \partial_{x_1}^{\alpha_1} \dots \partial_{x_d}^{\alpha_d}$ with $\alpha = (\alpha_1, \dots, \alpha_d)$.

- $B_\infty(\mathcal{X})$ is the space of bounded measurable functions, that is endowed with the norm

$$\|\varphi\|_{B_\infty} := \sup_{x \in \mathcal{X}} |\varphi(x)|.$$

- $B_W^\infty(\mathcal{X})$ is defined as

$$B_W^\infty(\mathcal{X}) = \left\{ \varphi : \mathcal{X} \rightarrow \mathbb{R} \text{ measurable} \mid \|\varphi\|_{B_W^\infty} := \sup_{x \in \mathcal{X}} \frac{|\varphi(x)|}{W(x)} < +\infty \right\},$$

where $W : \mathcal{X} \rightarrow [1, +\infty)$ is any given measurable function.

- $\mathcal{P}_W(\mathcal{X})$ is the associated space of probability measures for duality on measure spaces:

$$\mathcal{P}_W(\mathcal{X}) = \{ \nu \in \mathcal{P}(\mathcal{X}) \mid \nu(W) < +\infty \}.$$

- $C^0([0, T], \mathbb{R}^d)$ refers to the set of continuous functions from the interval $[0, T]$ to \mathbb{R}^d .
- For any probability measure $\mu \in \mathcal{P}(\mathcal{X})$, let

$$L^2(\mu) = \left\{ \varphi \text{ measurable} \mid \int_{\mathcal{X}} |\varphi|^2 d\mu < +\infty \right\}.$$

- For any $\varphi \in C_c^\infty(\mathcal{X})$, we introduce the seminorm

$$|\varphi|_{\mathcal{H}^1(\mu)}^2 = \int_{\mathcal{X}} \mathcal{E}(\varphi, \varphi) d\mu,$$

and the equivalence relation \sim_1 through: $\varphi \sim_1 \psi$ if and only if $|\varphi - \psi|_{\mathcal{H}^1(\mu)} = 0$.

- $\mathcal{H}^1(\mu)$ is the closure of $C_c^\infty(\mathcal{X})$ quotiented by \sim_1 for the norm $|\cdot|_{\mathcal{H}^1(\mu)}$. Note that $\mathcal{H}^1(\mu)$ and $L^2(\mu)$ are not subspaces of each other in general, but $\mathcal{H}^1(\mu) \subset L^2(\mu)$ for instance if μ satisfies a Poincaré inequality and \mathcal{S} is positive definite. The difference between $L^2(\mu)$ and $\mathcal{H}^1(\mu)$ is however important for degenerate dynamics; see the

application in Section 4.2 in [FS20]. We now construct a space dual to $\mathcal{H}^1(\mu)$ with the same density argument by introducing the seminorm: for $\varphi \in C_c^\infty(\mathcal{X})$,

$$|\varphi|_{\mathcal{H}^{-1}(\mu)}^2 = \sup_{\psi \in C_c^\infty(\mathcal{X})} \left\{ 2 \int_{\mathcal{X}} \varphi \psi d\mu - |\psi|_{\mathcal{H}^1(\mu)}^2 \right\}.$$

We define similarly the equivalence relation \sim_{-1} on $C_c^\infty(\mathcal{X})$ by $\varphi \sim_{-1} \psi$ if and only if $|\varphi - \psi|_{\mathcal{H}^{-1}(\mu)} = 0$. The space $\mathcal{H}^{-1}(\mu)$ is then the closure of $C_c^\infty(\mathcal{X})$ quotiented by \sim_{-1} . This is actually the dual space of $\mathcal{H}^1(\mu)$.

Next, we introduce the definitions of various topologies that are used in our paper.

- τ -topology is the weak topology on $\mathcal{P}(\mathcal{X})$ associated with the convergence of measures tested against functions belonging to $B_\infty(\mathcal{X})$ (we may also use the notation $\sigma(\mathcal{P}(\mathcal{X}), B_\infty)$). This means that for a sequence $(\nu_n)_{n \in \mathbb{N}}$ in $\mathcal{P}(\mathcal{X})$, $\nu_n \rightarrow \nu$ in the τ -topology if $\nu_n(\varphi) \rightarrow \nu(\varphi)$ for any $\varphi \in B_\infty(\mathcal{X})$. Recall that the τ -topology is stronger than the usual weak topology $\sigma(\mathcal{P}(\mathcal{X}), C_b(\mathcal{X}))$ on $\mathcal{P}(\mathcal{X})$, which corresponds to the convergence $\nu_n(\varphi) \rightarrow \nu(\varphi)$ for any $\varphi \in C_b(\mathcal{X})$.
- τ_W is the associated topology $\sigma(\mathcal{P}_W(\mathcal{X}), B_W^\infty(\mathcal{X}))$, accounting for the convergence of measures tested against the larger class of functions $\varphi \in B_W^\infty(\mathcal{X})$.
- The weighted total variation distance between two probability measures $\nu, \eta \in \mathcal{P}_W(\mathcal{X})$ is given by

$$d_W(\nu, \eta) = \sup_{\|\varphi\|_{B_W^\infty} \leq 1} \left\{ \int_{\mathcal{X}} \varphi d\nu - \int_{\mathcal{X}} \varphi d\eta \right\} = \int_{\mathcal{X}} W(x) |\nu - \eta|(dx),$$

where $|\nu - \eta|$ is the total variation measure associated with $\nu - \eta$.

Finally, the notation $[\cdot, \cdot]$ refers to the Lie bracket of two vector fields. The Lie bracket is used to describe the symmetry or commutation properties between two vector fields and is defined as follows:

$$[A_i, A_j](f) = A_i(A_j(f)) - A_j(A_i(f)),$$

where f is a smooth function. Intuitively, the Lie bracket measures the non-commutativity of two vector fields, i.e., the difference in their actions when applied in different orders.

B Technical Proofs

B.1 Proof of Theorem 2.1

Proof. Under Assumption 2.1-2.3, the empirical measure π_t of the generalized Langevin dynamics defined in (1.6) satisfies a large deviation principle in the τ^κ -topology, and the

proof is similar as Theorem 2.1 in [FS20] and is omitted here. In the rest of the proof, we will obtain the rate function from the large deviation principle through the decomposition of the infinitesimal generator \mathcal{L}_τ of the dynamics (1.6) which can be easily computed as

$$\begin{aligned} \mathcal{L}_\tau = & \sum_{i=1}^n \sum_{j=1}^n \left[-\mathcal{D}_{ij}(\mathbf{z}) \frac{\partial}{\partial \mathbf{z}_j} H(\mathbf{z}) - \mathcal{Q}_{ij}(\mathbf{z}) \frac{\partial}{\partial \mathbf{z}_j} H(\mathbf{z}) + \frac{\partial}{\partial \mathbf{z}_j} \mathcal{D}_{ij}(\mathbf{z}) + \frac{\partial}{\partial \mathbf{z}_j} \mathcal{Q}_{ij}(\mathbf{z}) \right] \frac{\partial}{\partial \mathbf{z}_i} \\ & + \sum_{i=1}^n \sum_{j=1}^n \mathcal{D}_{ij}(\mathbf{z}) \frac{\partial^2}{\partial \mathbf{z}_i \partial \mathbf{z}_j}. \end{aligned} \quad (\text{B.1})$$

By the definition of adjoint operator, we have

$$\int_{\mathcal{X}} g(\mathbf{z}) \mathcal{L}_\tau h(\mathbf{z}) d\mu = \int_{\mathcal{X}} h(\mathbf{z}) \mathcal{L}_\tau^* g(\mathbf{z}) d\mu, \quad (\text{B.2})$$

where \mathcal{L}_τ is the infinitesimal generator of the generalized Langevin dynamics (1.6) that is given by formula (B.1). By plugging formula (B.1) into the left hand side of (B.2), and taking $\mu(d\mathbf{z}) = e^{-H(\mathbf{z})} d\mathbf{z}$, where $H(\mathbf{z})$ is given in (1.7), we get

$$\begin{aligned} \int_{\mathcal{X}} g \mathcal{L}_\tau h d\mu &= \int_{\mathcal{X}} g(\mathbf{z}) \left\{ \sum_{i=1}^n \sum_{j=1}^n \left[-\mathcal{D}_{ij}(\mathbf{z}) \frac{\partial}{\partial \mathbf{z}_j} H(\mathbf{z}) - \mathcal{Q}_{ij}(\mathbf{z}) \frac{\partial}{\partial \mathbf{z}_j} H(\mathbf{z}) + \frac{\partial}{\partial \mathbf{z}_j} \mathcal{D}_{ij}(\mathbf{z}) \right. \right. \\ &\quad \left. \left. + \frac{\partial}{\partial \mathbf{z}_j} \mathcal{Q}_{ij}(\mathbf{z}) \right] \frac{\partial}{\partial \mathbf{z}_i} h(\mathbf{z}) + \sum_i \sum_j \mathcal{D}_{ij}(\mathbf{z}) \frac{\partial^2}{\partial \mathbf{z}_i \partial \mathbf{z}_j} h(\mathbf{z}) \right\} e^{-H(\mathbf{z})} d\mathbf{z} \\ &= \sum_{i=1}^n \sum_{j=1}^n \int_{\mathcal{X}} g(\mathbf{z}) \left\{ \left[-\mathcal{D}_{ij}(\mathbf{z}) \frac{\partial}{\partial \mathbf{z}_j} H(\mathbf{z}) - \mathcal{Q}_{ij}(\mathbf{z}) \frac{\partial}{\partial \mathbf{z}_j} H(\mathbf{z}) + \frac{\partial}{\partial \mathbf{z}_j} \mathcal{D}_{ij}(\mathbf{z}) \right. \right. \\ &\quad \left. \left. + \frac{\partial}{\partial \mathbf{z}_j} \mathcal{Q}_{ij}(\mathbf{z}) \right] \frac{\partial}{\partial \mathbf{z}_i} h(\mathbf{z}) \right\} \cdot e^{-H(\mathbf{z})} d\mathbf{z} \\ &\quad + \sum_{i=1}^n \sum_{j=1}^n \int_{\mathcal{X}} g(\mathbf{z}) \left(\mathcal{D}_{ij}(\mathbf{z}) \frac{\partial^2}{\partial \mathbf{z}_i \partial \mathbf{z}_j} h(\mathbf{z}) \right) \cdot e^{-H(\mathbf{z})} d\mathbf{z} \\ &= \sum_{i=1}^n \sum_{j=1}^n \int_{\mathcal{X}} -\frac{\partial}{\partial \mathbf{z}_i} \left\{ g(\mathbf{z}) \left[-\mathcal{D}_{ij}(\mathbf{z}) \frac{\partial}{\partial \mathbf{z}_j} H(\mathbf{z}) - \mathcal{Q}_{ij}(\mathbf{z}) \frac{\partial}{\partial \mathbf{z}_j} H(\mathbf{z}) \right. \right. \\ &\quad \left. \left. + \frac{\partial}{\partial \mathbf{z}_j} \mathcal{D}_{ij}(\mathbf{z}) + \frac{\partial}{\partial \mathbf{z}_j} \mathcal{Q}_{ij}(\mathbf{z}) \right] e^{-H(\mathbf{z})} \right\} \cdot h(\mathbf{z}) d\mathbf{z} \\ &\quad + \sum_{i=1}^n \sum_{j=1}^n \int_{\mathcal{X}} \frac{\partial^2}{\partial \mathbf{z}_i \partial \mathbf{z}_j} \left(g(\mathbf{z}) \cdot \mathcal{D}_{ij}(\mathbf{z}) \cdot e^{-H(\mathbf{z})} \right) \cdot h(\mathbf{z}) d\mathbf{z}. \end{aligned} \quad (\text{B.3})$$

By expanding equation (1.10) specifically, we obtain

$$\begin{aligned}
& \sum_{i=1}^n \sum_{j=1}^n \left\{ \frac{\partial^2}{\partial \mathbf{z}_i \partial \mathbf{z}_j} \mathcal{Q}_{ij}(\mathbf{z}) e^{-H(\mathbf{z})} + \frac{\partial}{\partial \mathbf{z}_j} \mathcal{Q}_{ij}(\mathbf{z}) \left(-\frac{\partial}{\partial \mathbf{z}_i} H(\mathbf{z}) \right) e^{-H(\mathbf{z})} \right. \\
& \quad + \frac{\partial}{\partial \mathbf{z}_i} \mathcal{Q}_{ij}(\mathbf{z}) \left(-\frac{\partial}{\partial \mathbf{z}_j} H(\mathbf{z}) \right) e^{-H(\mathbf{z})} + \mathcal{Q}_{ij}(\mathbf{z}) \left(-\frac{\partial^2}{\partial \mathbf{z}_i \partial \mathbf{z}_j} H(\mathbf{z}) \right) e^{-H(\mathbf{z})} \quad (\text{B.4}) \\
& \quad \left. + \mathcal{Q}_{ij}(\mathbf{z}) \left(\frac{\partial}{\partial \mathbf{z}_j} H(\mathbf{z}) \frac{\partial}{\partial \mathbf{z}_i} H(\mathbf{z}) \right) e^{-H(\mathbf{z})} \right\} = 0.
\end{aligned}$$

We can divide equation (B.3) into two parts and simplify it by using the properties of matrices $\mathcal{D}(\mathbf{z})$ and $\mathcal{Q}(\mathbf{z})$. We can compute that

$$\begin{aligned}
& -\frac{\partial}{\partial \mathbf{z}_i} \left\{ g(\mathbf{z}) \left[-\mathcal{D}_{ij}(\mathbf{z}) \frac{\partial}{\partial \mathbf{z}_j} H(\mathbf{z}) - \mathcal{Q}_{ij}(\mathbf{z}) \frac{\partial}{\partial \mathbf{z}_j} H(\mathbf{z}) + \frac{\partial}{\partial \mathbf{z}_j} \mathcal{D}_{ij}(\mathbf{z}) + \frac{\partial}{\partial \mathbf{z}_j} \mathcal{Q}_{ij}(\mathbf{z}) \right] e^{-H(\mathbf{z})} \right\} \\
& = \frac{\partial}{\partial \mathbf{z}_i} \mathcal{D}_{ij}(\mathbf{z}) \frac{\partial}{\partial \mathbf{z}_j} H(\mathbf{z}) \cdot g(\mathbf{z}) \cdot e^{-H(\mathbf{z})} + \frac{\partial}{\partial \mathbf{z}_i} \mathcal{Q}_{ij}(\mathbf{z}) \frac{\partial}{\partial \mathbf{z}_j} H(\mathbf{z}) \cdot g(\mathbf{z}) \cdot e^{-H(\mathbf{z})} \\
& \quad - \frac{\partial^2}{\partial \mathbf{z}_j \partial \mathbf{z}_i} \mathcal{D}_{ij}(\mathbf{z}) \cdot e^{-H(\mathbf{z})} \cdot g(\mathbf{z}) - \frac{\partial^2}{\partial \mathbf{z}_j \partial \mathbf{z}_i} \mathcal{Q}_{ij}(\mathbf{z}) \cdot e^{-H(\mathbf{z})} \cdot g(\mathbf{z}) \\
& \quad + \mathcal{D}_{ij}(\mathbf{z}) \frac{\partial^2}{\partial \mathbf{z}_j \partial \mathbf{z}_i} H(\mathbf{z}) \cdot g(\mathbf{z}) \cdot e^{-H(\mathbf{z})} + \mathcal{Q}_{ij}(\mathbf{z}) \frac{\partial^2}{\partial \mathbf{z}_j \partial \mathbf{z}_i} H(\mathbf{z}) \cdot g(\mathbf{z}) \cdot e^{-H(\mathbf{z})} \\
& \quad + \mathcal{D}_{ij}(\mathbf{z}) \frac{\partial}{\partial \mathbf{z}_j} H(\mathbf{z}) \frac{\partial}{\partial \mathbf{z}_i} g(\mathbf{z}) \cdot e^{-H(\mathbf{z})} + \mathcal{Q}_{ij}(\mathbf{z}) \frac{\partial}{\partial \mathbf{z}_j} H(\mathbf{z}) \frac{\partial}{\partial \mathbf{z}_i} g(\mathbf{z}) \cdot e^{-H(\mathbf{z})} \\
& \quad - \frac{\partial}{\partial \mathbf{z}_j} \mathcal{D}_{ij}(\mathbf{z}) \frac{\partial}{\partial \mathbf{z}_i} g(\mathbf{z}) \cdot e^{-H(\mathbf{z})} - \frac{\partial}{\partial \mathbf{z}_j} \mathcal{Q}_{ij}(\mathbf{z}) \frac{\partial}{\partial \mathbf{z}_i} g(\mathbf{z}) \cdot e^{-H(\mathbf{z})} \\
& \quad - \mathcal{D}_{ij}(\mathbf{z}) \left(\frac{\partial}{\partial \mathbf{z}_j} H(\mathbf{z}) \frac{\partial}{\partial \mathbf{z}_i} H(\mathbf{z}) \right) \cdot g(\mathbf{z}) \cdot e^{-H(\mathbf{z})} - \mathcal{Q}_{ij}(\mathbf{z}) \left(\frac{\partial}{\partial \mathbf{z}_j} H(\mathbf{z}) \frac{\partial}{\partial \mathbf{z}_i} H(\mathbf{z}) \right) g(\mathbf{z}) \cdot \\
& \quad e^{-H(\mathbf{z})} + \frac{\partial}{\partial \mathbf{z}_j} \mathcal{D}_{ij}(\mathbf{z}) \frac{\partial}{\partial \mathbf{z}_i} H(\mathbf{z}) \cdot g(\mathbf{z}) \cdot e^{-H(\mathbf{z})} + \frac{\partial}{\partial \mathbf{z}_j} \mathcal{Q}_{ij}(\mathbf{z}) \frac{\partial}{\partial \mathbf{z}_i} H(\mathbf{z}) g(\mathbf{z}) \cdot e^{-H(\mathbf{z})}, \quad (\text{B.5})
\end{aligned}$$

and

$$\begin{aligned}
& \frac{\partial^2}{\partial \mathbf{z}_i \partial \mathbf{z}_j} \left(g(\mathbf{z}) \cdot \mathcal{D}_{ij}(\mathbf{z}) \cdot e^{-H(\mathbf{z})} \right) \\
&= \frac{\partial^2}{\partial \mathbf{z}_i \partial \mathbf{z}_j} \mathcal{D}_{ij}(\mathbf{z}) \cdot g(\mathbf{z}) \cdot e^{-H(\mathbf{z})} + \frac{\partial}{\partial \mathbf{z}_i} \mathcal{D}_{ij}(\mathbf{z}) \frac{\partial}{\partial \mathbf{z}_j} g(\mathbf{z}) \cdot e^{-H(\mathbf{z})} \\
&+ \frac{\partial}{\partial \mathbf{z}_i} \mathcal{D}_{ij}(\mathbf{z}) \cdot g(\mathbf{z}) \left(-\frac{\partial}{\partial \mathbf{z}_j} H(\mathbf{z}) \right) \cdot e^{-H(\mathbf{z})} + \frac{\partial}{\partial \mathbf{z}_j} \mathcal{D}_{ij}(\mathbf{z}) \frac{\partial}{\partial \mathbf{z}_i} g(\mathbf{z}) \cdot e^{-H(\mathbf{z})} \\
&+ \mathcal{D}_{ij}(\mathbf{z}) \frac{\partial^2}{\partial \mathbf{z}_i \partial \mathbf{z}_j} g(\mathbf{z}) \cdot e^{-H(\mathbf{z})} + \mathcal{D}_{ij}(\mathbf{z}) \frac{\partial}{\partial \mathbf{z}_i} g(\mathbf{z}) \left(-\frac{\partial}{\partial \mathbf{z}_j} H(\mathbf{z}) \right) \cdot e^{-H(\mathbf{z})} \\
&+ \frac{\partial}{\partial \mathbf{z}_j} \mathcal{D}_{ij}(\mathbf{z}) \cdot g(\mathbf{z}) \left(-\frac{\partial}{\partial \mathbf{z}_i} H(\mathbf{z}) \right) \cdot e^{-H(\mathbf{z})} + \mathcal{D}_{ij}(\mathbf{z}) \left(-\frac{\partial}{\partial \mathbf{z}_i} H(\mathbf{z}) \right) \frac{\partial}{\partial \mathbf{z}_j} g(\mathbf{z}) \cdot e^{-H(\mathbf{z})} \\
&+ \mathcal{D}_{ij}(\mathbf{z}) g(\mathbf{z}) \left(-\frac{\partial^2}{\partial \mathbf{z}_i \partial \mathbf{z}_j} H(\mathbf{z}) \right) \cdot e^{-H(\mathbf{z})} + \mathcal{D}_{ij}(\mathbf{z}) g(\mathbf{z}) \left(\frac{\partial}{\partial \mathbf{z}_i} H(\mathbf{z}) \frac{\partial}{\partial \mathbf{z}_j} H(\mathbf{z}) \right) \cdot e^{-H(\mathbf{z})}.
\end{aligned} \tag{B.6}$$

Substituting (B.5) and (B.6) into (B.3), we obtain under condition (B.4)

$$\begin{aligned}
\int_{\mathcal{X}} g \mathcal{L}_\tau h \, d\mu &= \int_{\mathcal{X}} h(\mathbf{z}) \left\{ \sum_{i=1}^n \sum_{j=1}^n \left[\frac{\partial}{\partial \mathbf{z}_i} \mathcal{D}_{ij}(\mathbf{z}) - \mathcal{D}_{ij}(\mathbf{z}) \frac{\partial}{\partial \mathbf{z}_i} H(\mathbf{z}) \right] \frac{\partial}{\partial \mathbf{z}_j} g(\mathbf{z}) \right. \\
&\quad \left. + \sum_{i=1}^n \sum_{j=1}^n \left[\mathcal{Q}_{ij}(\mathbf{z}) \frac{\partial}{\partial \mathbf{z}_j} H(\mathbf{z}) - \frac{\partial}{\partial \mathbf{z}_j} \mathcal{Q}_{ij}(\mathbf{z}) \right] \frac{\partial}{\partial \mathbf{z}_i} g(\mathbf{z}) \right. \\
&\quad \left. + \sum_{i=1}^n \sum_{j=1}^n \mathcal{D}_{ij}(\mathbf{z}) \frac{\partial^2}{\partial \mathbf{z}_i \partial \mathbf{z}_j} g(\mathbf{z}) \right\} d\mu \\
&= \int_{\mathcal{X}} h \mathcal{L}_\tau^* g \, d\mu.
\end{aligned}$$

Hence

$$\begin{aligned}
\mathcal{L}_\tau^* &= \sum_{i=1}^n \sum_{j=1}^n \left[\frac{\partial}{\partial \mathbf{z}_i} \mathcal{D}_{ij}(\mathbf{z}) - \mathcal{D}_{ij}(\mathbf{z}) \frac{\partial}{\partial \mathbf{z}_i} H(\mathbf{z}) \right] \frac{\partial}{\partial \mathbf{z}_j} \\
&\quad - \sum_{i=1}^n \sum_{j=1}^n \left[\frac{\partial}{\partial \mathbf{z}_j} \mathcal{Q}_{ij}(\mathbf{z}) - \mathcal{Q}_{ij}(\mathbf{z}) \frac{\partial}{\partial \mathbf{z}_j} H(\mathbf{z}) \right] \frac{\partial}{\partial \mathbf{z}_i} + \sum_{i=1}^n \sum_{j=1}^n \mathcal{D}_{ij}(\mathbf{z}) \frac{\partial^2}{\partial \mathbf{z}_i \partial \mathbf{z}_j}.
\end{aligned} \tag{B.7}$$

Next, it follow from equation (2.7) that

$$\begin{aligned}
\mathcal{L}_S &= \sum_{i=1}^n \sum_{j=1}^n \left[\frac{\partial}{\partial \mathbf{z}_i} \mathcal{D}_{ij}(\mathbf{z}) - \mathcal{D}_{ij}(\mathbf{z}) \frac{\partial}{\partial \mathbf{z}_i} H(\mathbf{z}) \right] \frac{\partial}{\partial \mathbf{z}_j} + \sum_{i=1}^n \sum_{j=1}^n \mathcal{D}_{ij}(\mathbf{z}) \frac{\partial^2}{\partial \mathbf{z}_i \partial \mathbf{z}_j}, \\
\mathcal{L}_A &= \sum_{i=1}^n \sum_{j=1}^n \left[\frac{\partial}{\partial \mathbf{z}_j} \mathcal{Q}_{ij}(\mathbf{z}) - \mathcal{Q}_{ij}(\mathbf{z}) \frac{\partial}{\partial \mathbf{z}_j} H(\mathbf{z}) \right] \frac{\partial}{\partial \mathbf{z}_i}.
\end{aligned} \tag{B.8}$$

By applying Lemma 2.1, we obtain

$$I_\tau(\nu) = \frac{1}{4} \int_{\mathcal{X}} \nabla v \cdot \mathcal{D} \nabla v \, d\nu + \frac{1}{4} \int_{\mathcal{X}} \nabla \psi_v \cdot \mathcal{D} \nabla \psi_v \, d\nu,$$

where ψ_v is the unique solution in $\mathcal{H}^1(\nu)$ to the Poisson equation

$$\tilde{\nabla}(\mathcal{D} \nabla \psi_v) = \mathcal{L}_A v.$$

This completes the proof of Theorem 2.1. \square

B.2 Proof of Lemma 2.4

Proof. This is a direct consequence of Proposition 4.3 in [FS20]. It follows from Proposition 4.3 in [FS20] that $\pi_t = \frac{1}{t} \int_0^t \delta_{\theta_s} ds$ satisfies a large deviation principle (LDP) in the τ_κ -topology under the assumption that $e^{-U} \in L^1(\mathcal{X})$ and, for any $\eta \in (0, 1)$, it holds

$$(1 - \eta)|\nabla U|^2 - \Delta U \xrightarrow{|\theta| \rightarrow +\infty} +\infty,$$

and $\nabla \cdot (J \Delta U e^{-U}) = 0$ and

$$\frac{J \nabla U \cdot \nabla U}{\Psi} \xrightarrow{|\theta| \rightarrow +\infty} 0,$$

where

$$\Psi = \eta \left((1 - \eta)|\nabla U|^2 - \Delta U \right).$$

Then, with the notation of (2.7), it holds $\mathcal{L}_S = -\nabla U \cdot \nabla + \Delta$ and $\mathcal{L}_A = -J \nabla U \cdot \nabla$ and $\mathcal{L} = \mathcal{L}_S + \mathcal{L}_A$. Moreover, we have the following relation:

$$\begin{aligned} -\frac{(\mathcal{L} - J \nabla U \cdot \nabla) e^{\eta U}}{e^{\eta U}} &= \eta \left((1 - \eta)|\nabla U|^2 - \Delta U + J \nabla U \cdot \nabla U \right) \\ &= \eta \left((1 - \eta)|\nabla U|^2 - \Delta U \right) = \Psi. \end{aligned}$$

Finally, the conclusion follows from $\nabla \cdot (J \Delta U e^{-U}) = 0$ and $J \nabla U \cdot \nabla U = 0$ since J is an anti-symmetric matrix. \square

B.3 Proof of Theorem 2.2

Proof. We know that the generator of the mirrored Langevin dynamics (1.12) is elliptic and therefore it automatically satisfies the hypoellipticity of the generator and the controllability (i.e., irreducibility of the dynamics) properties. By letting $\mathcal{D} = [\nabla^2 \phi(\mathbf{z})]^{-1}$, the Lyapunov condition is satisfied by applying Proposition 2.1. Finally, by applying Theorem 2.1, the conclusion follows. \square

B.4 Proof of Lemma 2.5

Proof. We first show the Hypocoellipticity. We know from equation (2.18) that the generator of high-order Langevin dynamics is

$$\begin{aligned}\mathcal{L}_H &= \alpha\Delta_r - (\gamma p + \alpha r) \cdot \nabla_r + p \cdot \nabla_\theta - (\nabla U - \gamma r) \cdot \nabla_p \\ &= (-\alpha r \cdot \nabla_r + \alpha\Delta_r) + (p \cdot \nabla_\theta - \nabla U \cdot \nabla_p + \gamma r \cdot \nabla_p - \gamma p \cdot \nabla_r) \\ &= -A_1 A_1^* + \mathcal{L}_{HA},\end{aligned}$$

where $A_1 = \sqrt{\alpha}\nabla_r$. We can check that $\frac{1}{2}e^{-\frac{1}{2}|r|^2} dr$ is the invariant measure of \mathcal{L}_{HS} . We can compute that for any $f, g \in L^2(d\mu)$,

$$\begin{aligned}\int_{\mathbb{R}^d} (\sqrt{\alpha}\nabla_r f) g \frac{1}{2}e^{-\frac{1}{2}|r|^2} dr &= - \int_{\mathbb{R}^d} f \sqrt{\alpha}\nabla_r \left(g \frac{1}{2}e^{-\frac{1}{2}|r|^2} \right) dr \\ &= \int_{\mathbb{R}^d} f \sqrt{\alpha}(rg - \nabla_r g) \frac{1}{2}e^{-\frac{1}{2}|r|^2} dr,\end{aligned}$$

that is $A_1^* = \sqrt{\alpha}(r - \nabla_r)$. Now we arrive at that $\mathcal{L}_H = -A_1 A_1^* + \mathcal{L}_{HA}$.

Hörmander's condition requires that the vector field family generated by finite Lie brackets at each point must span the entire tangent space. Next we want to show that these operators $A_1, \mathcal{L}_{HA}, [A_1, \mathcal{L}_{HA}], [[A_1, [A_1, \mathcal{L}_{HA}]], \dots$ span the entire space of \mathcal{X} for a finite number of commutators. We can compute that for any smooth function f in \mathcal{X}

$$\begin{aligned}[A_1, \mathcal{L}_{HA}]f &:= (A_1 \mathcal{L}_{HA} - \mathcal{L}_{HA} A_1)f \\ &= \sqrt{\alpha}\nabla_r(p \cdot \nabla_\theta - \nabla U \cdot \nabla_p + \gamma r \cdot \nabla_p - \gamma p \cdot \nabla_r)f \\ &\quad - (p \cdot \nabla_\theta - \nabla U \cdot \nabla_p + \gamma r \cdot \nabla_p - \gamma p \cdot \nabla_r)\sqrt{\alpha}\nabla_r f \\ &= \gamma\nabla_p f, \\ [[A_1, \mathcal{L}_{HA}], \mathcal{L}_{HA}]f &= (\gamma\nabla_p \mathcal{L}_{HA} - \mathcal{L}_{HA} \gamma\nabla_p)f \\ &= \gamma\nabla_p(p \cdot \nabla_\theta - \nabla U \cdot \nabla_p + \gamma r \cdot \nabla_p - \gamma p \cdot \nabla_r)f \\ &\quad - (p \cdot \nabla_\theta - \nabla U \cdot \nabla_p + \gamma r \cdot \nabla_p - \gamma p \cdot \nabla_r)\gamma\nabla_p f \\ &= \gamma\nabla_p(\nabla_\theta - \gamma\nabla_r)f, \\ [[[A_1, \mathcal{L}_{HA}], \mathcal{L}_{HA}], \mathcal{L}_{HA}]f &= [\gamma\nabla_p(\nabla_\theta - \gamma\nabla_r), p \cdot \nabla_\theta - \nabla U \cdot \nabla_p + \gamma r \cdot \nabla_p - \gamma p \cdot \nabla_r]f \\ &= (\gamma(\nabla_\theta - p \cdot \nabla_p \nabla_\theta) + \gamma^2 r \cdot \nabla_p - \gamma^2 p \cdot \nabla_r + \gamma\nabla_r)f.\end{aligned}$$

Therefore, $[A_1, \mathcal{L}_{HA}], [[A_1, \mathcal{L}_{HA}], \mathcal{L}_{HA}], [[[A_1, \mathcal{L}_{HA}], \mathcal{L}_{HA}], \mathcal{L}_{HA}]$ has rank $3d$ at every point (θ, p, q) because that $\nabla_\theta, \nabla_p, \nabla_r$ are independent in the sense of linear independence. That is the Hypocoellipticity of \mathcal{L}_H such that Assumption 2.1 is satisfied.

Next, we show the Controllability. Given $t > 0$ and two pair of points (θ_0, p_0, r_0) and (θ_t, p_t, r_t) , let $\phi(s)$ be any \mathcal{C}^3 path in \mathbb{R}^d which satisfy $\phi(0) = \theta_0$, $\phi(t) = \theta_t$, $\nabla\phi(0) = p_0$

and $\nabla\phi(t) = p_t$, $\nabla^2\phi(0) = -\nabla U(\theta_0) + \gamma r_0$ and $\nabla^2\phi(t) = -\nabla U(\theta_t) + \gamma r_t$. Consider the control u given by

$$u_t = \frac{1}{\sqrt{2\alpha}} \left(\frac{1}{\gamma} \nabla^3\phi(t) + \frac{\alpha}{\gamma} \nabla^2\phi(t) + \left(\frac{1}{\gamma} \nabla^2 U(\phi(t)) + \gamma \right) \nabla\phi(t) + \frac{\alpha}{\gamma} \nabla U(\phi(t)) \right). \quad (\text{B.9})$$

By definition, $(\phi(t), \nabla\phi(t), \nabla^2\phi(t))$ is a solution of the control system with control u_t . Therefore the Controllability (Assumption 2.2) is satisfied. The proof is complete. \square

Remark B.1. *The u_t drives the system from (θ_0, p_0, r_0) to (θ_t, p_t, r_t) . We denote by $G_t(x)$ the set of accessible points from x in time t . This implies that $G_t(x) = \mathbb{R}^{3d}$ for all $t > 0$ and all $x \in \mathbb{R}^{3d}$. From the support theorem (Theorem 6.1 in [Bel06]) we conclude that $P_t(x, F) > 0$ for all $t > 0$, all $x \in \mathbb{R}^{3d}$, and all open set F .*

B.5 Proof of Lemma 2.6

Proof. Let us construct the Lyapunov functions $W_\delta : \mathbb{R}^{3d} \rightarrow [1, +\infty)$ satisfying Lyapunov condition (2.20). The construction is inspired from Proposition 2.13 in [GLNW24]. We define the vector field L as follows. Let $\chi : \mathbb{R}^d \rightarrow [0, 1]$ be a smooth function such that $\chi(\theta) = 0$ if $|\theta| \leq 1$ and $\chi(\theta) = 1$ if $|\theta| \geq 2$. We define $J(\theta) := \theta|\theta|^{\beta-1}\chi(\theta)$ with $\beta \in [0, 1]$. Note that J is C^1 and the first derivatives of J are bounded over \mathbb{R}^d (because $\beta \leq 1$), say by $C_J := \sup_{\theta \in \mathbb{R}^d} |\text{Jac}(J(\theta))| > 0$ (where $\|M\|_2 := \sup\{|My|, |y| = 1\}$, $M \in \mathcal{M}_d(\mathbb{R})$). One then sets:

$$L := \kappa J, \quad \kappa := \frac{\gamma}{2C_J}, \quad (\text{B.10})$$

so that

$$C_L := \sup_{\theta \in \mathbb{R}^d} |\text{Jac}(L(\theta))| \leq \gamma/2. \quad (\text{B.11})$$

For all $(\theta, p, r) \in \mathbb{R}^{3d}$, $b \geq 0$, and $h, a > 0$, we define:

$$F_0(\theta, p, r) := hH(\theta, p, r) + aL(\theta) \cdot p + bp \cdot r, \quad (\text{B.12})$$

with $H(\theta, p, r)$ given in (1.13). The parameter $\beta > 0$ will be chosen such that

$$\inf_{(\theta, p, r) \in \mathbb{R}^{3d}} F_0(\theta, p, r) \in \mathbb{R}. \quad (\text{B.13})$$

We then set $F_{\text{HL}}(\theta, p, r) := F_0(\theta, p, r) - \inf_{(\theta, p, r) \in \mathbb{R}^{3d}} F_0(\theta, p, r) + 1$ and

$$W_\delta(\theta, p, r) := \exp\left(F_{\text{HL}}^\delta(\theta, p, r)\right), \quad \text{where } 1 - \frac{\beta}{k} < \delta < 1. \quad (\text{B.14})$$

In the following, for ease of notation we simply write F for F_{HL} . Since $F^{1-\delta} \geq 1$, a straightforward computation implies that over \mathbb{R}^{3d} ,

$$\frac{\mathcal{L}_H W_\delta}{W_\delta} \leq \frac{\delta}{F^{1-\delta}} [\mathcal{L}_H F + \delta\alpha |\nabla_p F|^2], \quad (\text{B.15})$$

where

$$\mathcal{L}_H = \alpha \Delta_r - (\gamma p + \alpha r) \cdot \nabla_r + p \cdot \nabla_\theta - (\nabla U - \gamma r) \cdot \nabla_p. \quad (\text{B.16})$$

We also have

$$\begin{aligned} \nabla_\theta F(\theta, p, r) &= h \nabla U(\theta) + a \text{Jac}(L(\theta))p, \\ \nabla_p F(\theta, p, r) &= hp + aL(\theta) + br, \\ \nabla_r F(\theta, p, r) &= hr + bp. \end{aligned}$$

Consequently, one has for all $(\theta, p, r) \in \mathbb{R}^{3d}$,

$$\begin{aligned} \mathcal{L}_H F(\theta, p, r) &= -\gamma b |p|^2 + ap \cdot \text{Jac}(L(\theta)) \cdot p - a \nabla U(\theta) \cdot L(\theta) + (b\gamma - \alpha h) |r|^2 \\ &\quad + a\gamma r \cdot L(\theta) - b \nabla U(\theta) \cdot r - \alpha br \cdot p + hd\alpha, \end{aligned} \quad (\text{B.17})$$

and

$$\delta\alpha |\nabla_{\mathbf{z}} F|^2(\theta, p, r) = \delta\alpha |hr + bp|^2, \quad (\text{B.18})$$

with $\mathbf{z} = (\theta, p, r)$. Set

$$b = a > 0, \quad k \in (1, 2], \quad \text{and} \quad \beta = k - 1. \quad (\text{B.19})$$

Let us first check (B.13). Let $p_1 = k/(k-1) > 1$ and $q_1 = p_1/(p_1 - 1) = k \leq 2$. Using Assumption 2.4, we have for all $(\theta, p, r) \in \mathbb{R}^{3d}$, if $|\theta| \geq c_v$,

$$F_0(\theta, p, r) \geq m_U h |\theta|^k + \frac{h-a}{2} |p|^2 + \frac{h-a}{2} |r|^2 - \frac{a\kappa}{p_1} |\theta|^k - \frac{a\kappa}{k} |p|^k. \quad (\text{B.20})$$

Then, for $h > 0$, choose $a > 0$ small enough such that

$$\frac{a\kappa}{p_1} < m_U h \quad \text{and} \quad \frac{a\kappa}{k} + \frac{a}{2} < \frac{h}{2}. \quad (\text{B.21})$$

Then (B.13) holds. Note also that when (B.19) is satisfied,

$$F_0(\theta, p, r) \leq c'(|\theta|^k + |p|^2 + |r|^2) + C'.$$

Next, let us verify that a Lyapunov function W_δ of the form (B.17) which satisfies the Lyapunov condition (2.20). Recall $b = a$. In the following, $(\theta, p, r) \in \mathbb{R}^{3d}$, $|\theta| \geq$

$\max(2, m_U)$, and $\eta = \sqrt{a}$. Using Assumption 2.4, (B.17), and (B.18), one has:

$$\begin{aligned}
& (\mathcal{L}_H F + \delta\alpha|\nabla_r F|^2)(\theta, p, r) \\
& \leq \alpha h d - \gamma a |p|^2 + a p \cdot \text{Jac}(L(\theta))p - a \nabla U(\theta) \cdot L(\theta) - \alpha h |r|^2 + \gamma a |r|^2 \\
& \quad + \gamma a |r| |L(\theta)| + a |\nabla U(\theta)| |r| + \alpha a |r| |p| + \delta\alpha |hr + ap|^2 \\
& \leq \alpha h d - \gamma a |p|^2 + a C_L |p|^2 - a \kappa_{C_U} |\theta|^{2(k-1)} - \alpha h |r|^2 + \gamma a |r|^2 \\
& \quad + \gamma \kappa a |r| |\theta|^{k-1} + a M_U |\theta|^{k-1} |r| + \alpha a |p| |r| + \delta\alpha |hr + ap|^2 \\
& \leq \alpha h d - \gamma a |p|^2 + a C_L |p|^2 - a \kappa_{C_U} |\theta|^{2(k-1)} - \alpha h |r|^2 + \gamma a |r|^2 + \frac{\gamma \kappa a |r|^2}{2\eta} + \gamma \kappa \eta a |\theta|^{2(k-1)} \\
& \quad + \frac{a M_U |r|^2}{2\eta} + a \eta M_U |\theta|^{2(k-1)} + \frac{\alpha a |r|^2}{2\eta} + \frac{\alpha a \eta |p|^2}{2} + 2\delta\alpha h^2 |r|^2 + 2\delta\alpha a^2 |p|^2 \\
& \leq \alpha h d + |p|^2 \left(-\gamma a + a C_L + 2\delta\alpha a^2 + \alpha a^{\frac{3}{2}}/2 \right) \\
& \quad + |\theta|^{2(k-1)} \left(-\kappa_{C_U} a + \gamma \kappa a^{3/2}/2 + M_U a^{3/2}/2 \right) \\
& \quad + |r|^2 \left(-\alpha h + 2\delta\alpha h^2 + \gamma \kappa \sqrt{a}/2 + M_U \sqrt{a}/2 + \alpha \sqrt{a}/2 + \gamma a \right).
\end{aligned}$$

With (B.14) and (B.19) we know that $\delta > \frac{2-k}{k}$. Let $h > 0$ such that $-\alpha h + 2\delta h^2 < 0$. Using also (B.11), it holds $-\gamma + C_L \leq -\gamma/2$. We then choose $a > 0$ such that (B.21) holds and

$$\begin{aligned}
& -\gamma/2 + 2\delta a^2 + \alpha a^{3/2}/2 < 0, \\
& -\kappa_{C_U} a + \gamma \kappa a^{3/2}/2 + M_U a^{3/2}/2 < 0, \\
& -\alpha h + 2\delta h^2 + \gamma \kappa \sqrt{a}/2 + M_U \sqrt{a}/2 + \alpha \sqrt{a}/2 + \gamma a < 0.
\end{aligned}$$

This concludes the proof. \square

B.6 Proof of Theorem 2.3

Proof. By Lemmas 2.5 and 2.6, we know the infinite generator \mathcal{L}_H satisfies the Hypocoellipticity (Assumption 2.1), Controllability (Assumption 2.2) and Witten-Lyapunov condition (Assumption 2.3). As a special case of the constructed model, by Theorem 2.1, the infinitesimal generator is decomposed such that the rate function is the sum of symmetric and anti-symmetric parts. Through formula (2.23) and (2.6), we have

$$\mathcal{C}(v, v) = \frac{1}{2} (\mathcal{L}_\alpha(v^2) - 2v \mathcal{L}_\alpha v) = \alpha |\nabla_r v|^2.$$

Hence by Lemma 2.1, we obtain

$$I_H(\nu) = \frac{\alpha}{4} \int_{\mathcal{X}} |\nabla_r v|^2 d\nu + \frac{\alpha}{4} \int_{\mathcal{X}} |\nabla_r \psi|^2 d\nu.$$

Write $\psi_v = \gamma\psi$ where ψ_v is the unique solution in $\mathcal{H}^1(\nu)$ to the Poisson equation (2.22) and this completes of the proof of Theorem 2.3. \square

B.7 Proof of Lemma 2.7

Proof. Note that \mathcal{L}_R is an elliptic operator, so Controllability (Assumption 2.2) is satisfied. Now we check the Witten-Lyapunov condition for the Hessian-free high-resolution dynamics (1.17). The form of Lyapunov function we constructed is $W_a(\theta, r) = e^{\varphi(\theta, r)}$, where

$$\varphi(\theta, r) = aH(\theta, r) + b\theta \cdot r = a \left(U(\theta) + \frac{1}{2}|r|^2 \right) + b\theta \cdot r,$$

where $H(\theta, r)$ is given in (1.16) and b is a parameter to be determined. With \mathcal{L}_R given in (2.18), $\frac{\mathcal{L}_R W_a}{W_a}$ can be calculated as:

$$\begin{aligned} \frac{\mathcal{L}_R W_a}{W_a} &= \frac{((-\beta\nabla U(\theta) + r) \cdot \nabla_\theta + (-\alpha r - \nabla U(\theta)) \cdot \nabla_r + \beta\Delta_\theta + \alpha\Delta_r)e^\varphi}{e^\varphi} \\ &= (-\beta\nabla U(\theta) + r) \cdot (a\nabla U(\theta) + br) - (\alpha r + \nabla U(\theta)) \cdot (ar + b\theta) \\ &\quad + \alpha(|ar + b\theta|^2) + \beta(|a\nabla U(\theta) + br|^2) + a\beta\Delta U(\theta) + a\alpha d. \end{aligned} \tag{B.22}$$

Note that Assumption 2.5(b) implies that there exist $M_U > 0$, $k > 1$ such that for all $\theta \in \mathbb{R}^d$ with $|\theta| \geq c_U$: $|\nabla U(\theta)| \leq M_U|\theta|^{k-1}$. Moreover, it follows from Assumption 2.5(b) that $\Delta U(\theta) \leq m_U d$ for any $\theta \in \mathbb{R}^d$.

In order to make $-\frac{\mathcal{L}_R W_a}{W_a}$ satisfy Lyapunov condition (2.25), we just prove when $a \leq \frac{1}{2}$, when $a \geq \frac{1}{2}$ the proof is the similar. We set $\eta = \frac{(1-2a)(\alpha+M_U\beta)}{c_1}$ and $b < \frac{c_1}{2\alpha}$ small enough,

and by Assumption 2.5 such that (B.22) becomes that

$$\begin{aligned}
-\frac{\mathcal{L}_R W_a}{W_a} &= |r|^2 (a\alpha - b - a^2\alpha - b^2\beta) + |\theta|^2 (-\alpha b^2) \\
&\quad + r \cdot \theta (\alpha b - 2ab\alpha) + r \cdot \nabla U(\theta) (b\beta - 2ab\beta) \\
&\quad + b\theta \cdot \nabla U(\theta) + |\nabla U(\theta)|^2 (a\beta - a^2\beta) + \Delta U(\theta) (-a\beta) - a\alpha \\
&\geq |r|^2 (a\alpha - b - a^2\alpha - b^2\beta) - |\theta|^2 (\alpha b^2) - \frac{\alpha b - 2ab\alpha}{2} \eta |r|^2 - \frac{\alpha b - 2ab\alpha}{2\eta} |\theta|^2 \\
&\quad - \frac{M_U (b\beta - 2ab\beta)}{2} \eta |r|^2 - \frac{M_U (b\beta - 2ab\beta)}{2\eta} |\theta|^{2k-2} \\
&\quad + bc_1 |\theta|^2 - bC_1 - a\beta m_U d - a\alpha \\
&= |r|^2 \left(a\alpha - a^2\alpha - b - b^2\beta - \frac{\eta b\alpha(1-2a)}{2} - \frac{M_U \eta b\beta(1-2a)}{2} \right) \\
&\quad + |\theta|^2 \left(bc_1 - \alpha b^2 - \frac{b\alpha(1-2a)}{2\eta} - \frac{bM_U \beta(1-2a)}{2\eta} \right) - a\beta m_U d - a\alpha - bC_1 \\
&= |r|^2 2 \left(\alpha a(1-a) - b \left(1 + b\beta + \frac{(1-2a)^2 (\alpha + M_U \beta)^2}{2c_1} \right) \right) \\
&\quad + |\theta|^2 \left(\frac{c_1}{2} - \alpha b \right) - a\beta m_U d - a\alpha - bC_1.
\end{aligned}$$

Then the function W_a is a Lyapunov function that meets the Lyapunov condition (2.25) and this completes the proof of Lemma 2.6. \square

B.8 Proof of Theorem 2.4

Proof. From formula (2.6), we have

$$\begin{aligned}
\mathcal{E}(v, v) &= \frac{1}{2} (\mathcal{L}_R(v^2) - 2v\mathcal{L}_R v) = \beta |\nabla_\theta v|^2 + \alpha |\nabla_r v|^2, \\
\mathcal{E}(\psi_v, \psi_v) &= \beta |\nabla_\theta \psi_v|^2 + \alpha |\nabla_r \psi_v|^2,
\end{aligned}$$

where \mathcal{L}_R is defined in (2.23). For $\varphi \in C_c^\infty(\mathcal{X})$, we have the seminorm

$$|\varphi|_{\mathcal{H}^1(\nu)}^2 = \int_{\mathcal{X}} \mathcal{E}(\varphi, \varphi) d\nu = \int_{\mathcal{X}} (\beta |\nabla_\theta \varphi|^2 + \alpha |\nabla_r \varphi|^2) d\nu,$$

and

$$|\varphi|_{\mathcal{H}^{-1}(\nu)}^2 = \sup_{\psi \in C_c^\infty} \left\{ 2 \int_{\mathcal{X}} \varphi \psi d\nu - |\psi|_{\mathcal{H}^1(\nu)}^2 \right\}.$$

By Lemma 2.1, we deduce that

$$\begin{aligned}
I_A(\nu) &= \frac{1}{4} |\mathcal{L}_{\text{RA}}(v)|_{\mathcal{H}^{-1}(\nu)}^2 \\
&= \frac{1}{4} \sup_{\psi \in C_c^\infty} \left\{ 2 \int_{\mathcal{X}} \mathcal{L}_{\text{RA}}(v) \psi \, d\nu - |\psi|_{\mathcal{H}^1(\nu)}^2 \right\} \\
&= -\frac{1}{2} \inf_{\psi \in C_c^\infty} \left\{ \int_{\mathcal{X}} \frac{1}{2} \mathcal{C}(\psi, \psi) - \mathcal{L}_{\text{RA}}(v) \psi \, d\nu \right\},
\end{aligned}$$

where \mathcal{L}_{RA} is the anti-symmetric part of \mathcal{L}_R . Now it comes to a variational problem to find the infimum of $\int_{\mathcal{X}} (\frac{1}{2} \mathcal{C}(\psi, \psi) - \mathcal{L}_A(v) \psi) \, d\nu$. Using Euler-Lagrange equation and by Lemma 2.1 we can get that

$$I_R(\nu) = \frac{\beta}{4} \left(\int_{\mathcal{X}} |\nabla_{\theta} v|^2 \, d\nu + \int_{\mathcal{X}} |\nabla_{\theta} \psi_v|^2 \, d\nu \right) + \frac{\alpha}{4} \left(\int_{\mathcal{X}} |\nabla_r v|^2 \, d\nu + \int_{\mathcal{X}} |\nabla_r \psi_v|^2 \, d\nu \right), \tag{B.23}$$

where ψ_v is the unique solution in $\mathcal{H}^1(\nu)$ to the Poisson equation

$$\tilde{\nabla} (\beta \nabla_{\theta} \psi_v + \alpha \nabla_r \psi_v) = \mathcal{L}_{\text{RA}} v, \tag{B.24}$$

where $\tilde{\nabla}$ denoting the adjoint of gradient operator ∇ in $L^2(\nu)$, that is

$$-e^v \nabla (e^{-v} (\beta \nabla_{\theta} \psi_v + \alpha \nabla_r \psi_v)) = \mathcal{L}_{\text{RA}} v,$$

i.e. the Poisson equation (2.27) and this completes of the proof of Theorem 2.4. \square

B.9 Proof of Corollary 3.1

Proof. It follows from Theorem 2.1 that

$$I_{\tau}(\nu) = \frac{1}{4} \int_{\mathcal{X}} \nabla v \cdot \mathcal{D} \nabla v \, d\nu + \frac{1}{4} \int_{\mathcal{X}} \nabla \psi_v \cdot \mathcal{D} \nabla \psi_v \, d\nu.$$

Since \mathcal{D} is a positive semidefinite diffusion matrix, we get

$$\nabla \psi_v \cdot \mathcal{D} \nabla \psi_v \geq 0,$$

that is,

$$\frac{1}{4} \int_{\mathcal{X}} \nabla \psi_v \cdot \mathcal{D} \nabla \psi_v \, d\nu \geq 0.$$

It can be seen from (2.12) that

$$I_o(\nu) = \frac{1}{4} \int_{\mathcal{X}} |\nabla v|^2 \, d\nu.$$

Next, we define

$$J_\tau(\nu) := I_\tau(\nu) - I_0(\nu) = \frac{1}{4} \int_{\mathcal{X}} \nabla v \cdot (\mathcal{D} - \mathbf{I}) \nabla v \, d\nu + \frac{1}{4} \int_{\mathcal{X}} \nabla \psi_v \cdot \mathcal{D} \nabla \psi_v \, d\nu,$$

where \mathbf{I} is the identity matrix. If $\mathcal{D} - \mathbf{I}$ is a positive semidefinite diffusion matrix, by the definition of $J_\tau(\nu)$, it is clear that $J_\tau(\nu) \geq 0$. This completes the proof. \square

B.10 Proof of Corollary 3.2

Proof. In this example just let $\mathcal{D} = [\nabla^2 \phi(\mathbf{z})]^{-1}$, then this corollary is a direct application of Corollary 3.1. \square

B.11 Proof of Proposition 3.1

Proof. Recall that the LDP rate function for Hessian-free high-resolution dynamics is in equation (2.26), that is

$$I_R(\nu) = \frac{\beta}{4} \left(\int_{\mathcal{X}} |\nabla_\theta v|^2 \, d\nu + \int_{\mathcal{X}} |\nabla_\theta \psi_v|^2 \, d\nu \right) + \frac{\alpha}{4} \left(\int_{\mathcal{X}} |\nabla_r v|^2 \, d\nu + \int_{\mathcal{X}} |\nabla_r \psi_v|^2 \, d\nu \right).$$

Since $\nabla v = (\nabla_\theta v, \nabla_r v)$, we have

$$\int_{\mathcal{X}} |\nabla v|^2 \, d\nu = \int_{\mathcal{X}} |\nabla_\theta v|^2 \, d\nu + \int_{\mathcal{X}} |\nabla_r v|^2 \, d\nu.$$

Hence we conclude that if $\min(\alpha, \beta) \geq 1$, then we have $I_R(\nu) \geq I_{eo}(\nu)$. This completes the proof. \square

B.12 Proof of Corollary 3.3

Proof. This corollary follows immediately from Proposition 3.1 and (3.6), (2.12) and (3.7). \square

B.13 Proof of Proposition 3.2

Proof. Recall that the LDP rate function for underdamped Langevin dynamics is in equation (2.13), that is

$$I_u(\nu) = \frac{\gamma}{4} \int_{\mathcal{X}} |\nabla_r v|^2 \, d\nu + \frac{1}{4\gamma} \int_{\mathcal{X}} |\nabla_r \psi|^2 \, d\nu.$$

Similar as in the proof of Proposition 3.1

$$I_{e2o}(\nu) = \frac{1}{4} \int_{\mathcal{X}} |\nabla v|^2 \, d\nu = \frac{1}{4} \int_{\mathcal{X}} |\nabla_\theta v|^2 \, d\nu + \frac{1}{4} \int_{\mathcal{X}} |\nabla_r v|^2 \, d\nu.$$

Thus, when $v = v(r)$, we have $\nabla_\theta v = 0$. If $\gamma \geq 1$, we conclude that $I_u(\nu) \geq I_{eo}(\nu)$ and this completes the proof. \square

B.14 Proof of Proposition 3.3

Proof. Recall the LDP rate function for high-order Langevin dynamics from equation (2.21):

$$I_H(\nu) = \frac{\alpha}{4} \int_{\mathcal{X}} |\nabla_r v|^2 d\nu + \frac{1}{4\alpha} \int_{\mathcal{X}} |\nabla_r \psi|^2 d\nu.$$

Moreover,

$$I_{e_{3o}}(\nu) = \frac{1}{4} \int_{\mathcal{X}} |\nabla v|^2 d\nu = \frac{1}{4} \int_{\mathcal{X}} |\nabla_{\theta} v|^2 d\nu + \frac{1}{4} \int_{\mathcal{X}} |\nabla_p v|^2 d\nu + \frac{1}{4} \int_{\mathcal{X}} |\nabla_r v|^2 d\nu.$$

Thus, when $v = v(r)$, we have $\nabla_{\theta} v = 0$ and $\nabla_p v = 0$. If $\alpha \geq 1$, we conclude that $I_H(\nu) \geq I_{e_{3o}}(\nu)$, and this completes the proof. \square