

Efficient Algorithms for Lipschitz Selections of Set-Valued Mappings in \mathbf{R}^2

By PAVEL SHVARTSMAN

Department of Mathematics, Technion - Israel Institute of Technology,

32000 Haifa, Israel

e-mail: pshv@technion.ac.il

Abstract

Let F be a set-valued mapping from an N -element metric space (\mathcal{M}, ρ) into the family of all closed half-planes in \mathbf{R}^2 . In this paper, we provide an efficient algorithm for a Lipschitz selection of F , i.e., a Lipschitz mapping $f : \mathcal{M} \rightarrow \mathbf{R}^2$ such that $f(x) \in F(x)$ for all $x \in \mathcal{M}$. Given a constant $\lambda > 0$, this algorithm produces the following two outcomes: (1) The algorithm guarantees that there is no Lipschitz selection of F with Lipschitz constant at most λ ; (2) The algorithm returns a Lipschitz selection of F with Lipschitz constant at most 3λ .

The total work and storage required by this selection algorithm are at most CN^2 where C is an absolute constant.

CONTENTS

1. Introduction.	2
1.1 Main results.	2
1.2 The Projection Algorithm: main steps and main properties.	6
2. Notation and preliminaries.	10
2.1 Background notation.	10
2.2 Rectangles and rectangular hulls.	12
2.3 Rectangles: intersections, neighborhoods and selections.	15
3. The key theorem: Lipschitz selections and rectangular hulls.	21
4. Proof of the key theorem: the final step.	25
5. The $\vec{\lambda}$ -Projection Algorithm and its versions.	41
5.1 Main properties of the Projection Algorithm: proofs.	41
5.2 Other versions of the Projection Algorithm.	43
6. From geometric description to analytical: five basic algorithms.	44
6.1 The λ -metric refinement of the set-valued mapping - STEP 1.	46
6.2 The rectangular hulls of metric refinements - STEP 2.	47
6.3 The metric refinements of the rectangular hulls - STEP 3.	48
6.4 The centers of the metric projections onto rectangles - STEP 4.	51
6.5 The final step: metric projections onto the metric refinements - STEP 5.	51
6.6 Lipschitz selections of polygon-set valued mappings.	54
7. Further results and comments.	56
7.1 Generalizations of the PA: sets determined by polynomial inequalities and more.	56
7.2 A polynomial algorithm for the sharp selection problem.	57
7.3 Distance oracle and intrinsic metrics.	59
References	61

Math Subject Classification: 46E35

Key Words and Phrases: Set-valued mapping, Lipschitz selection, metric projection, rectangular hull, efficient algorithm

This research was supported by the ISRAEL SCIENCE FOUNDATION (grant No. 520/22).

1. Introduction.

1.1 Main results.

In this paper we study some aspects of the following selection problem formulated by C. Fefferman [29] (see also [30] and [31, Section 8.7]).

Problem 1.1 *Let (\mathcal{M}, ρ) be an N -point metric space. For each $x \in \mathcal{M}$, let $F(x) \subset \mathbf{R}^D$ be a convex polytope.*

How can one compute a map $f : \mathcal{M} \rightarrow \mathbf{R}^D$ such that $f(x) \in F(x)$ for all $x \in \mathcal{M}$, with Lipschitz norm as small as possible up to a factor $C(D)$?

This is a big ill-conditioned linear programming problem. Can we do better than just applying general-purpose linear programming? How does the work of an optimal algorithm scale with the number of points N ?

For this problem we want an *efficient algorithm* that require a minimal use of computer resources.

In [36] C. Fefferman and B. G. Pegueroles suggested such an algorithm which solves a (general) version of Problem 1.1 related to the so-called “*approximate*” m -smooth selections. In this version the metric space (\mathcal{M}, ρ) is an N -point subset $E \subset \mathbf{R}^n$, and we allow the point $f(x)$, $x \in E$, to belong to a certain enlargement of $F(x)$, the set $(1 + \tau)\diamond F(x)$. Here $\tau > 0$ and $(1 + \tau)\diamond F(x)$ denotes the convex set obtained by dilating $F(x)$ about its center of mass by a factor of $(1 + \tau)$.

In the present paper we consider a special case of this result related to C^1 -selections. Using standard transitions from C^1 -functions to Lipschitz functions and back, we can reformulate the main result of [36] for the case $m = 1$ in the following equivalent form.

Theorem 1.2 *(An Approximate Selection Algorithm [36]) Let n, D, L, N be positive integers and let $M, \tau > 0$. Let $E \subset \mathbf{R}^n$ be an N -point set, and let F be a mapping on E which to every $x \in E$ assigns a compact, convex polytope $F(x) \subset \mathbf{R}^D$. We suppose that each $F(x)$ is defined by at most L linear constraints. Under these conditions, we produce one of the following two outcomes:*

OUTCOME 1 (“No-Go”): *We guarantee that there does not exist a Lipschitz function $f : E \rightarrow \mathbf{R}^D$ with Lipschitz constant at most M such that $f(x) \in F(x)$ for all $x \in E$.*

OUTCOME 2 (“Success”): *We return a Lipschitz function $f : E \rightarrow \mathbf{R}^D$, with Lipschitz constant at most $\gamma(n, D)M$ such that $f(x) \in (1 + \tau)\diamond F(x)$ for all $x \in E$.*

This Selection Algorithm requires at most $CN \log N$ computer operations and at most CN units of computer memory. Here, the constant C depends only on n, D, L and τ .

Our paper is inspired by this striking result. As the authors of [36] point out, the approach developed in that paper does not work for the case $\tau = 0$, i.e., for “exact” (rather than “approximate”) Lipschitz selections.

In our paper, we study the Lipschitz selection Problem 1.1 for the case $\tau = 0$ and $D = 2$, i.e., for the special case of set-valued mappings defined on finite metric spaces and taking values in the family of all *convex polygons* in \mathbf{R}^2 .

In what follows, it will be more convenient for us to consider the problem in a somewhat broader context of *pseudometric* spaces (\mathcal{M}, ρ) , that is, we assume that the “distance function” $\rho : \mathcal{M} \times \mathcal{M} \rightarrow [0, +\infty]$ satisfies

$$\rho(x, x) = 0, \quad \rho(x, y) = \rho(y, x), \quad \text{and} \quad \rho(x, y) \leq \rho(x, z) + \rho(z, y)$$

for all $x, y, z \in \mathcal{M}$. Note that $\rho(x, y) = 0$ may hold with $x \neq y$, and $\rho(x, y)$ may be $+\infty$. We call a pseudometric space (\mathcal{M}, ρ) finite if \mathcal{M} is finite, but we say that the pseudometric ρ is finite if $\rho(x, y)$ is finite for every $x, y \in \mathcal{M}$.

Also note that each polygon is the intersection of a finite number of half-planes. This allows us, instead of Problem 1.1, to study its equivalent version, in which the metric space and the family of convex polygons are replaced by the pseudometric space and the family $\mathcal{HP}(\mathbf{R}^2)$ of *all closed half-planes in \mathbf{R}^2* .

Let us formulate our main result. By $\text{Lip}(\mathcal{M})$ we denote the space of all Lipschitz mappings from \mathcal{M} into \mathbf{R}^2 equipped with Lipschitz seminorm

$$\|f\|_{\text{Lip}(\mathcal{M})} = \inf\{\lambda \geq 0 : \|f(x) - f(y)\| \leq \lambda \rho(x, y) \text{ for all } x, y \in \mathcal{M}\}.$$

Hereafter $\|\cdot\|$ denotes the uniform norm in \mathbf{R}^2 , i.e., $\|x\| = \max\{|x_1|, |x_2|\}$ for $x = (x_1, x_2) \in \mathbf{R}^2$.

Let F be a set-valued mapping which to each element $x \in \mathcal{M}$ assigns a nonempty convex closed set $F(x) \subset \mathbf{R}^2$. A *selection* of F is a map $f : \mathcal{M} \rightarrow \mathbf{R}^2$ such that $f(x) \in F(x)$ for all $x \in \mathcal{M}$. A selection f is said to be Lipschitz if $f \in \text{Lip}(\mathcal{M})$.

We introduce the quantity $|F|_{\mathfrak{M}}$ by letting

$$|F|_{\mathfrak{M}} = \inf\{\|f\|_{\text{Lip}(\mathcal{M})} : f \text{ is a Lipschitz selection of } F\} \quad (1.1)$$

whenever F has a Lipschitz selection, and we set $|F|_{\mathfrak{M}} = +\infty$ otherwise.

Thus, we are given a positive parameter λ , an N -element pseudometric space $\mathfrak{M} = (\mathcal{M}, \rho)$ and a set-valued mapping $F : \mathcal{M} \rightarrow \mathcal{HP}(\mathbf{R}^2)$. We want to either compute a Lipschitz function $f : \mathcal{M} \rightarrow \mathbf{R}^2$ such that $f(x) \in F(x)$ on \mathcal{M} , with $\|f\|_{\text{Lip}(\mathcal{M})} \leq \gamma \lambda$, or show that there is no Lipschitz selection of F with Lipschitz constant at most λ . (Here $\gamma > 0$ is an absolute constant.)

We want to solve the above problem with an algorithm, to be implemented on an (idealized) computer with standard von Neumann architecture, able to work with exact real numbers. We hope our algorithm will be *efficient*, i.e., it requires few computer operations. (By “operation” we mean an arithmetic operation (comparison, multiplication, division, addition and subtraction of two real numbers) or fetching a number from RAM. Also, by “running time” or “work” we mean the number of computer operations.)

In Sections 1.2 and Sections 3-6 we present and study an algorithm which satisfies all these requirements. We call this algorithm the *Projection Algorithm*. This simple five-step geometric algorithm is the main object of our study. Its construction involves such basic geometric operations as intersections of half-planes in \mathbf{R}^2 , rectangular hulls of these intersections, and metric projections onto closed convex sets (in the uniform norm in \mathbf{R}^2). See Section 1.2 and Figures 1-4.

Let us formulate the main result of the paper.

Theorem 1.3 *The Projection Algorithm receives as input a real number $\lambda > 0$, an N -element pseudometric space (\mathcal{M}, ρ) and a half-plane $F(x) \subset \mathbf{R}^2$ for each $x \in \mathcal{M}$.*

Given the above input, we produce one of the two following outcomes:

- **OUTCOME 1 (“No Go”):** *We guarantee that there does not exist $f \in \text{Lip}(\mathcal{M})$ with Lipschitz seminorm $\|f\|_{\text{Lip}(\mathcal{M})} \leq \lambda$ such that $f(x) \in F(x)$ for all $x \in \mathcal{M}$.*

- **OUTCOME 2 (“Success”):** *The algorithm produces a mapping $f : \mathcal{M} \rightarrow \mathbf{R}^2$ with Lipschitz seminorm*

$$\|f\|_{\text{Lip}(\mathcal{M})} \leq 3\lambda \quad (1.2)$$

satisfying $f(x) \in F(x)$ for all $x \in \mathcal{M}$.

This algorithm requires at most CN^2 computer operations and at most CN^2 units of computer memory. Here $C > 0$ is an absolute constant.

Theorem 1.3 implies an analog of Theorem 1.2 (for the case $\tau = 0$ and $D = 2$) for set-valued mappings defined on a pseudometric space and taking values in the family of all convex (not necessarily bounded) polygons in \mathbf{R}^2 . This is Theorem 6.10 which we prove in Section 6.6.

Remark 1.4 (*The efficiency of the Projection Algorithm*) Recall, that for Problem 1.1 we want an *efficient algorithm* that requires minimal use of computer resources, i.e., the number of computer operations and the amount of computer memory. Our input data requires at least $3N$ units of computer memory for the N half-planes $F(x)$, $x \in \mathcal{M}$, and $N(N + 1)/2$ units for the pseudometric $\rho(x, y)$, $x, y \in \mathcal{M}$, defined on the N -point pseudometric space (\mathcal{M}, ρ) . In general, we need to read all this information to produce an optimal Lipschitz selection of F , so any algorithm will require at least $N^2/2$ computer operations and at least $N^2/2$ units of memory.

Theorem 1.3 above states the Projection Algorithm requires *at most* CN^2 computer operations and *at most* CN^2 units of computer memory where $C > 0$ is an absolute constant. From this point of view, the Projection Algorithm being considered on the family of *all N -element pseudometric spaces*, is *maximally efficient*, i.e., it requires minimal (up to some absolute factor) use of computer resources.

This result leads to another interesting problem: Let (\mathcal{M}, ρ) be some special non-trivial N -element (pseudo)metric space. Under what condition on (\mathcal{M}, ρ) can we construct a Lipschitz selection algorithm with significantly fewer computer operations than $O(N^2)$ as in Theorem 1.3? For example, let (\mathcal{M}, ρ) be an N -point subset of \mathbf{R}^n equipped with the Euclidean metric. Can the projection algorithm in Theorem 1.3 be replaced by an algorithm that requires at most $O(N \log N)$ computer operations and at most $O(N)$ storage (as in Theorem 1.2 for the Approximate Selection Algorithm [36])? (This question was formulated by C. Fefferman [30].)

We note that the answer to this question is not clear even in the one dimensional case, i.e., for set-valued mappings taking values in the family of all closed intervals in \mathbf{R} . ◀

Remark 1.5 Recall that in the two dimensional case, Problem 1.1 is the problem of constructing an efficient algorithm (with minimal use of computer resources) which given an N -element pseudometric space $\mathfrak{M} = (\mathcal{M}, \rho)$ and a set-valued mapping $F : \mathcal{M} \rightarrow \mathcal{HP}(\mathbf{R}^2)$, provides a Lipschitz selection f of F with Lipschitz seminorm at most $\gamma|F|_{\mathfrak{M}}$. Here $\gamma > 0$ is an absolute constant.

Recall that $|F|_{\mathfrak{M}}$ is defined by (1.1). This definition allows us to reformulate Theorem 1.3 as follows: given $\lambda > 0$ the Projection Algorithm states that either $|F|_{\mathfrak{M}} \geq \lambda$ (the outcome “No Go”) or the Lipschitz selection \tilde{f} of F produced by this algorithm, has Lipschitz seminorm $\|\tilde{f}\|_{\text{Lip}(\mathcal{M})} \leq 3\lambda$ (the outcome “Success”).

Therefore, in this case the following inequality

$$|F|_{\mathfrak{M}} \leq \|\tilde{f}\|_{\text{Lip}(\mathcal{M})} \leq 3\lambda$$

holds. Thus, Theorem 1.3 provides a certain (rather efficient) tool for obtaining lower (for the outcome “No Go”) and upper (for the outcome “Success”) bounds for $|F|_{\mathfrak{M}}$. However, this result does not provide a complete solution to Problem 1.1.

Nevertheless, if we could compute the value of $|F|_{\mathfrak{M}}$ within some absolute constant γ , and then apply Theorem 1.3 with $\lambda = \gamma|F|_{\mathfrak{M}}$, Problem 1.1 would be completely solved. Indeed, in this case, the Projection Algorithm would provide the outcome “Success” and produce a Lipschitz selection \tilde{f} of F with

$$\|\tilde{f}\|_{\text{Lip}(\mathcal{M})} \leq 3\gamma|F|_{\mathfrak{M}}$$

as required in this problem.

We study the problem of efficient algorithms for computing the quantity $|F|_{\mathfrak{M}}$ and related problems in forthcoming paper [67]. For first results in this direction we refer the reader to [66, Section 6.3]. In particular, in this preprint we present an algorithm which, given an N -element pseudometric space $\mathfrak{M} = (\mathcal{M}, \rho)$ and a set-valued mapping $F : \mathcal{M} \rightarrow \mathcal{HP}(\mathbf{R}^2)$, computes the value of $|F|_{\mathfrak{M}}$ (up to an absolute constant) using work and storage at most $C N^3$ for a universal constant C . ◀

Problem 1.1 is a special case of the general Lipschitz selection problem, which studies the existence and properties of Lipschitz selections of set-valued mappings from (pseudo)metric spaces into various families of convex subsets of Banach spaces. This problem can be viewed as a search for a Lipschitz mapping that approximately agrees with the data.

There is an extensive literature devoted to various aspects of the Lipschitz selection and related smooth selection problems. Among the many results known so far, let us mention those contained in papers and monographs [3, 5, 32, 36, 37, 45–47, 56–58, 61–64]. We refer the reader to all these works and references therein, for numerous results and methods related to this topic..

The Lipschitz selection problem is of great interest in recent years, mainly because of its close connection with the classical *Whitney Extension Problem* [70]: *Given a positive integer m and a function f defined on a closed subset of \mathbf{R}^n , how can one determine whether f extends to a C^m -function on all of \mathbf{R}^n ?*

Over the years (since 1934) this problem has attracted a lot of attention, and there is an extensive literature devoted to this problem and its analogues for various spaces of smooth functions. For a detailed account of the history of extension and restriction problems for m -smooth functions, as well as various references related to this topic, we refer the reader to [7, 8, 25, 26, 28, 31, 33, 34, 44, 64] and references therein.

As an example, let us illustrate the connection between the Lipschitz selection problem and the Whitney problem for the space $C^2(\mathbf{R}^n)$. In [8, 60, 62] we showed that the Whitney problem for the restrictions of C^2 -functions to *finite* subsets of \mathbf{R}^n can be reduced to a certain Lipschitz selection problem for *affine-set valued* mappings. The solution to this special case of the Lipschitz selection problem given in [61–63] led us to an interesting property of the restrictions of C^2 -functions, called by C. Fefferman [26] (for the general case of C^m -spaces) the *Finiteness Principle*.

This principle allows us to reduce the Whitney problem for $C^2(\mathbf{R}^n)$ -restrictions to an *arbitrary finite subsets* of \mathbf{R}^n to a similar problem but for $C^2(\mathbf{R}^n)$ -restrictions to *finite sets consisting of at most $k^\# = 3 \cdot 2^{n-1}$ points*. See [60]. In [25], C. Fefferman showed that the Finiteness Principle holds for the space $C^m(\mathbf{R}^n)$ for arbitrary $m, n \geq 1$ with a certain constant $k^\# = k^\#(m, n)$ depending only on m and n .

Furthermore, in [62] we solved the Lipschitz selection problem for the special case of the *line-set valued* mappings in \mathbf{R}^2 and showed how constructive geometric criteria for Lipschitz selections of such mappings are transformed into purely analytical descriptions of the restrictions of C^2 -functions to finite subsets of the plane.

These close connections between these two problems, the geometric Lipschitz selection problem and the analytic Whitney extension problem, allow us to transform efficient Lipschitz selection algorithms into efficient algorithms for various extension problems for spaces of smooth functions. Note also that efficient algorithms for solving C^m -interpolation problems were developed by C. Fefferman and B. Klartag in [27, 34, 35].

1.2 The Projection Algorithm: main steps and main properties.

In this section we present a geometric description of the general version of the Projection Algorithm for the family $\text{Conv}(\mathbf{R}^2)$ of all convex closed subsets of \mathbf{R}^2 .

The input data of this algorithm are:

- (●1) Two real numbers $\lambda_1, \lambda_2 \geq 0$;
- (●2) An N -point pseudometric space (\mathcal{M}, ρ) ;
- (●3) A set-valued mapping $F : \mathcal{M} \rightarrow \text{Conv}(\mathbf{R}^2)$.

Let us prepare the ingredients that are needed for this geometric description.

Definition 1.6 Given a constant $\lambda \geq 0$ we define a set-valued mapping $F^{[1]}[\cdot : \lambda; \rho]$ on \mathcal{M} by letting

$$F^{[1]}[x : \lambda; \rho] = \bigcap_{y \in \mathcal{M}} [F(y) + \lambda \rho(x, y) \mathcal{Q}_0], \quad x \in \mathcal{M}. \quad (1.3)$$

We refer to the mapping $F^{[1]}[\cdot : \lambda; \rho]$ as *the $(\lambda; \rho)$ -metric refinement of F* . If the pseudometric ρ is clear from the context, we omit ρ in this notation and call the mapping in (1.3) the λ -metric refinement of F .

Given a convex set $S \subset \mathbf{R}^2$, we let $\mathcal{H}[S]$ denote *the smallest closed rectangle (possibly unbounded) with sides parallel to the coordinate axes, containing S* . We refer to $\mathcal{H}[S]$ as *the rectangular hull* of the set S .

By $\text{Pr}(\cdot, S)$ we denote the operator of metric projection onto a closed convex subset $S \subset \mathbf{R}^2$. Finally, we let $\text{center}(\cdot)$ denote the center of a centrally symmetric bounded set in \mathbf{R}^2 .

The Projection Algorithm 1.7 Given a vector $\vec{\lambda} = (\lambda_1, \lambda_2)$ with non-negative coordinates λ_1 and λ_2 , a finite pseudometric space $\mathfrak{M} = (\mathcal{M}, \rho)$, and a set-valued mapping $F : \mathcal{M} \rightarrow \text{Conv}(\mathbf{R}^2)$, the Projection Algorithm which we define below either produces a selection $f_{\vec{\lambda}, F}$ of F (the outcome “**Success**”) or terminates (the outcome “**No go**”). This procedure includes the following five main steps.

STEP 1. At this step we construct the λ_1 -metric refinement of F , i.e., the set-valued mapping

$$F^{[1]}[x : \lambda_1] = \bigcap_{y \in \mathcal{M}} [F(y) + \lambda_1 \rho(x, y) \mathcal{Q}_0]. \quad (1.4)$$

If $F^{[1]}[x : \lambda_1] = \emptyset$ for some $x \in \mathcal{M}$, the algorithm produces the outcome “**No go**” and terminates.

STEP 2. Let us assume that the above condition does not hold, i.e., *for every element $x \in \mathcal{M}$ the λ_1 -metric refinement $F^{[1]}[x : \lambda_1]$ is nonempty*.

In this case, for each $x \in \mathcal{M}$, we construct *the rectangular hull* of $F^{[1]}[x : \lambda_1]$, the set

$$\mathcal{T}_{F, \lambda_1}(x) = \mathcal{H}[F^{[1]}[x : \lambda_1]]. \quad (1.5)$$

Thus, $\mathcal{T}_{F, \lambda_1}$ maps \mathcal{M} into the family $\mathfrak{R}(\mathbf{R}^2)$ of all rectangles in \mathbf{R}^2 .

See Fig. 1.




Fig. 1: The second step of the Projection Algorithm.

STEP 3. For every $x \in \mathcal{M}$, we construct the λ_2 -metric refinement of the mapping $\mathcal{T}_{F, \lambda_1}$, i.e., the rectangle $\mathcal{T}_{F, \lambda_1}^{[1]}[x : \lambda_2]$ defined by

$$\mathcal{T}_{F, \lambda_1}^{[1]}[x : \lambda_2] = \bigcap_{y \in \mathcal{M}} [\mathcal{T}_{F, \lambda_1}(y) + \lambda_2 \rho(x, y) \mathcal{Q}_0]. \quad (1.6)$$

See Fig. 2.




Fig. 2: The third step of the Projection Algorithm.

If

$$\mathcal{T}_{F, \lambda_1}^{[1]}[x : \lambda_2] = \emptyset \quad \text{for some } x \in \mathcal{M},$$

the algorithm produces the outcome “**No go**” and terminates.

STEP 4. At this step, we assume that for each $x \in \mathcal{M}$ the rectangle $\mathcal{T}_{F, \lambda_1}^{[1]}[x : \lambda_2] \neq \emptyset$. Let $O = (0, 0)$ be the origin.

We define a mapping $g_F : \mathcal{M} \rightarrow \mathbf{R}^2$ by letting

$$g_F(x) = \text{center}(\text{Pr}(O, \mathcal{T}_{F, \lambda_1}^{[1]}[x : \lambda_2])), \quad x \in \mathcal{M}. \quad (1.7)$$

See Fig. 3.




Fig. 3: The mapping $g_F : \mathcal{M} \rightarrow \mathbf{R}^2$, a selection of the set-valued mapping $\mathcal{T}_{F, \lambda_1}$.

STEP 5. We define the mapping $f_{\vec{\lambda}; F} : \mathcal{M} \rightarrow \mathbf{R}^2$ by letting

$$f_{\vec{\lambda}; F}(x) = \mathbf{Pr}(g_F(x), F^{[1]}[x : \lambda_1]), \quad x \in \mathcal{M}. \quad (1.8)$$

See Fig. 4.




Fig. 4: The final step of the Projection Algorithm.

At this stage, the algorithm produces the outcome **“Success”** and terminates. ■

To specify the dependence on the parameters λ_1 and λ_2 , we call the above algorithm the $\vec{\lambda}$ -Projection Algorithm ($\vec{\lambda}$ -PA for short) or (λ_1, λ_2) -Projection Algorithm ((λ_1, λ_2) -PA). We refer to (λ, λ) -Projection Algorithm as λ -Projection Algorithm (λ -PA for short).

Remark 1.8 Let us note the following important properties of the (λ_1, λ_2) -Projection Algorithm which are immediate from the construction of this algorithm presented above:

(i) The (λ_1, λ_2) -PA produces the outcome **“No Go”** if and only if

$$\text{there exists } x \in \mathcal{M} \text{ such that either } F^{[1]}[x : \lambda_1] = \emptyset \text{ or } \mathcal{T}_{F, \lambda_1}^{[1]}[x : \lambda_2] = \emptyset; \quad (1.9)$$

(ii) The (λ_1, λ_2) -PA produces the outcome **“Success”** if and only if

$$F^{[1]}[x : \lambda_1] \neq \emptyset \text{ and } \mathcal{T}_{F, \lambda_1}^{[1]}[x : \lambda_2] \neq \emptyset \text{ for every } x \in \mathcal{M}. \quad (1.10)$$

The next theorem describes the main properties of the $\vec{\lambda}$ -Projection Algorithm.

Theorem 1.9 *Let $\lambda_1, \lambda_2 \geq 0$, and let $\vec{\lambda} = (\lambda_1, \lambda_2)$. Let $\mathfrak{M} = (\mathcal{M}, \rho)$ be a finite pseudometric space, and let $F : \mathcal{M} \rightarrow \text{Conv}(\mathbf{R}^2)$ be a set-valued mapping.*

(i) *If $\vec{\lambda}$ -Projection Algorithm produces the outcome “No go”, then we guarantee that there does not exist a Lipschitz selection of F with Lipschitz seminorm at most $\min\{\lambda_1, \lambda_2\}$.*

(ii) *Suppose that the $\vec{\lambda}$ -Projection Algorithm produces the outcome “Success”. In this case this algorithm returns the mapping $f_{\vec{\lambda}, F} : \mathcal{M} \rightarrow \mathbf{R}^2$, see (1.8), with the following properties:*

(★A) *The mapping $f_{\vec{\lambda}, F}$ is well defined. This means the following:*

(a) *For every $x \in \mathcal{M}$, the set $F^{[1]}[x : \lambda_1]$ and the rectangle $\mathcal{T}_{F, \lambda_1}^{[1]}[x : \lambda_2]$ are nonempty;*

(b) *The mapping g_F (see (1.7)) is well defined;*

(c) *The metric projection defined by the right hand side of (1.8) is a singleton.*

(★B) *$f_{\vec{\lambda}, F}$ is a Lipschitz selection of F with Lipschitz seminorm*

$$\|f_{\vec{\lambda}, F}\|_{\text{Lip}(\mathcal{M})} \leq \lambda_1 + 2\lambda_2.$$

In Section 5 we show that these properties of the $\vec{\lambda}$ -Projection Algorithm are immediate from a stronger result, Theorem 1.10, which we formulate below.

Given $\lambda \geq 0$ and $x, x', x'' \in \mathcal{M}$, we let $\mathcal{W}_F[x, x', x'' : \lambda]$ denote a (possibly empty) subset of \mathbf{R}^2 defined by

$$\mathcal{W}_F[x, x', x'' : \lambda] = \mathcal{H}[\{F(x') + \lambda\rho(x', x)Q_0\} \cap \{F(x'') + \lambda\rho(x'', x)Q_0\}].$$

We recall that by $\mathcal{H}[\cdot]$ we denote the rectangular hull of a set.

Theorem 1.10 *Let $\mathfrak{M} = (\mathcal{M}, \rho)$ be a finite pseudometric space, and let $F : \mathcal{M} \rightarrow \text{Conv}(\mathbf{R}^2)$ be a set-valued mapping. Given non-negative constants $\vec{\lambda}$ and λ , let us assume that for every $x, x', x'', y, y', y'' \in \mathcal{M}$ the following condition*

$$\mathcal{W}_F[x, x', x'' : \vec{\lambda}] \cap \{\mathcal{W}_F[y, y', y'' : \vec{\lambda}] + \lambda\rho(x, y)Q_0\} \neq \emptyset \quad (1.11)$$

holds. Then F has a Lipschitz selection with Lipschitz seminorm at most $2\lambda + \vec{\lambda}$.

We refer to Theorem 1.10 as *the key theorem*. This theorem is the most technically difficult part of the present paper. We prove Theorem 1.10 in Sections 3 and 4, and Theorem 1.9 in Section 5.

In Section 5 we also show that these theorems imply the following result.

Theorem 1.11 *Let $\mathfrak{M} = (\mathcal{M}, \rho)$ be a finite pseudometric space, and let $F : \mathcal{M} \rightarrow \text{Conv}(\mathbf{R}^2)$ be a set-valued mapping. Given $\vec{\lambda}, \lambda \geq 0$, let us assume that condition (1.11) holds for every $x, x', x'', y, y', y'' \in \mathcal{M}$. Then the $\vec{\lambda}$ -PA with $\vec{\lambda} = (\vec{\lambda}, \lambda)$ produces the outcome “Success” and returns the mapping $f_{\vec{\lambda}, F} : \mathcal{M} \rightarrow \mathbf{R}^2$ which has the following properties:*

$$f_{\vec{\lambda}, F} \text{ is a Lipschitz selection of } F \text{ with } \|f_{\vec{\lambda}, F}\|_{\text{Lip}(\mathcal{M})} \leq 2\lambda + \vec{\lambda}. \quad (1.12)$$

In other words, Theorem 1.11 states that the mapping $f_{\vec{\lambda}, F}$ provides a selection of the set-valued mapping F mentioned in the statement of Theorem 1.10.

We return to the (λ_1, λ_2) -Projection Algorithm for the family $\mathcal{HP}(\mathbf{R}^2)$ of all closed half-planes in \mathbf{R}^2 . **STEPS 1-5** contain a geometric description of this algorithm. However, this description is

incomplete because we do not specify the algorithms for constructing several geometric objects that occur in **STEPS 2-5**. Given $x \in \mathcal{M}$, such objects include:

STEP 1: The set $F^{[1]}[x : \lambda_1]$, i.e., the λ_1 -metric refinement of F . See (1.4).

STEP 2: The rectangle $\mathcal{T}_{F, \lambda_1}(x)$, see (1.5), i.e., the rectangular hull of the set $F^{[1]}[x : \lambda_1]$.

STEP 3: The rectangle $\mathcal{T}_{F, \lambda_1}^{[1]}[x : \lambda_2]$, see (1.6), i.e., the λ_2 -metric refinement of $\mathcal{T}_{F, \lambda_1}$.

STEP 4: The point $g_F(x)$, the center of the metric projection of the origin onto $\mathcal{T}_{F, \lambda_1}^{[1]}[x : \lambda_2]$. See (1.7).

STEP 5: The point $f_{\tilde{\lambda}; F}(x)$ defined by formula (1.8), i.e., the metric projection of $g_F(x)$ onto the set $F^{[1]}[x : \lambda_1]$.

The corresponding computational algorithms for these objects are presented in Section 6.1 - 6.5. In these sections, we also show that for each $x \in \mathcal{M}$, the work and storage required to execute each of these algorithm are linear in N (with some absolute constants).

Algorithms of this kind are more or less elementary for the objects in **STEPS 3-5**. However, at **STEP 1** and **STEP 2** we encounter classical problems of linear optimization for finite families of half-planes in \mathbf{R}^2 . In particular, at **STEP 2** this is the problem of an efficient algorithm for computing the rectangle hull of a polygon determined by N linear constraints. In this case, to solve these problems, we use classical results on low-dimensional linear programming by N. Megiddo [52] and M. E. Dyer [21], which provide the required efficient algorithms with $O(N)$ running time and $O(N)$ units of computer memory.

Finally, by successively applying all these algorithms to our problem, we obtain the required algorithm for Lipschitz selection satisfying the conditions of Theorem 1.3.

We complete the paper with Section 7 devoted to some generalizations of Theorem 1.3. First of them relates to wider families of convex subsets in \mathbf{R}^2 than the family $\mathcal{HP}(\mathbf{R}^2)$ of all closed half-planes. We also discuss efficiency of a polynomial algorithm for Lipschitz selection which provides the same outcomes as in Theorem 1.3, but with the estimate $\|f\|_{\text{Lip}(\mathcal{M})} \leq \lambda$ instead of inequality (1.2). Our last generalization relates to a variant of Problem 1.1 where the metric ρ is replaced with an arbitrary non-negative function defined on $\mathcal{M} \times \mathcal{M}$.

Acknowledgements. I am very grateful to Charles Fefferman for stimulating discussions and valuable advice.

2. Notation and preliminaries.

2.1 Background notation.

Let A and B be nonempty subsets of \mathbf{R}^2 . We let $A + B = \{a + b : a \in A, b \in B\}$ denote the Minkowski sum of these sets. Given $\lambda \geq 0$, by λA we denote the set $\lambda A = \{\lambda a : a \in A\}$.

We write

$$\text{dist}(A, B) = \inf\{\|a - b\| : a \in A, b \in B\}$$

to denote the distance between A and B . For $x \in \mathbf{R}^2$ we also set $\text{dist}(x, A) = \text{dist}(\{x\}, A)$. We put $\text{dist}(\emptyset, A) = 0$ provided A is an arbitrary (possibly empty) subset of \mathbf{R}^2 .

We let $d_{\text{H}}(A, B)$ denote the Hausdorff distance between A and B in the Banach space $\ell_{\infty}^2 = (\mathbf{R}^2, \|\cdot\|)$:

$$d_{\text{H}}(A, B) = \inf\{r > 0 : A + rQ_0 \supset B, B + rQ_0 \supset A\} \quad (2.1)$$

where $Q_0 = [-1, 1]^2$.

Given $a, b \in \mathbf{R}^2$, by $[a, b]$ we denote the closed interval (or a point if $a = b$) with the ends in a and b . Thus, $[a, b] = \{x \in \mathbf{R}^2 : x = (1 - t)a + tb, 0 \leq t \leq 1\}$.

We write $[x]_+ = \max\{x, 0\}$ for the positive part of the real x . We also use the natural convention that

$$\frac{0}{0} = 0, \quad \frac{a}{0} = +\infty \text{ for } a > 0, \quad a - b = 0 \text{ if } a = b = \pm\infty, \text{ and } (\pm\infty) - (\mp\infty) = \pm\infty. \quad (2.2)$$

If S is a finite set, by $\#S$ we denote the number of elements of S .

We let $\langle \cdot, \cdot \rangle$ denote the standard inner product in \mathbf{R}^2 . By

$$Ox_1 = \{x = (t, 0) : t \in \mathbf{R}\} \quad \text{and} \quad Ox_2 = \{x = (0, t) : t \in \mathbf{R}\}$$

we denote the coordinate axes in \mathbf{R}^2 .

Let $\mathcal{I}(Ox_i)$, $i = 1, 2$, be the family of all nonempty convex closed subsets of the coordinate axis Ox_i . In other words, $\mathcal{I}(Ox_i)$ is the family of all nonempty closed intervals (bounded or unbounded) and points lying on the Ox_i axis. Given $I \in \mathcal{I}(Ox_1)$, let

$$L(I) = \inf\{s \in \mathbf{R} : (s, 0) \in I\} \quad \text{and} \quad R(I) = \sup\{s \in \mathbf{R} : (s, 0) \in I\}. \quad (2.3)$$

We call the numbers $L(I)$ and $R(I)$ *the left end and right ends of I* , respectively.

In the same way we define the left and right ends of an interval $I \in \mathcal{I}(Ox_2)$:

$$L(I) = \inf\{t \in \mathbf{R} : (0, t) \in I\} \quad \text{and} \quad R(I) = \sup\{t \in \mathbf{R} : (0, t) \in I\}. \quad (2.4)$$

Note that the numbers $L(I)$ and $R(I)$ may take the values $\{-\infty\}$ or $\{+\infty\}$ provided I is unbounded.

By PR_i , $i = 1, 2$, we denote the operator of the orthogonal projection onto the axis Ox_i ; thus, given $x = (x_1, x_2) \in \mathbf{R}^2$, we have $\text{PR}_1[x] = (x_1, 0)$ and $\text{PR}_2[x] = (0, x_2)$. Given $i = 1, 2$, and a convex closed $S \subset \mathbf{R}^2$, we let $\text{PR}_i[S]$, denote the orthogonal projection of S onto the axis Ox_i . Thus,

$$\text{PR}_1[S] = \{(s, 0) : (s, t) \in S \text{ for some } t \in \mathbf{R}\} \quad \text{and} \quad \text{PR}_2[S] = \{(0, t) : (s, t) \in S \text{ for some } s \in \mathbf{R}\}.$$

Remark 2.1 Note that the orthogonal projections of a convex closed set $S \subset \mathbf{R}^2$ onto the coordinate axes are determined by the numbers

$$\sigma_1(S) = \inf\{s : (s, t) \in S\}, \quad \sigma_2(S) = \sup\{s : (s, t) \in S\}, \quad (2.5)$$

and

$$\tau_1(S) = \inf\{t : (s, t) \in S\}, \quad \tau_2(S) = \sup\{t : (s, t) \in S\}. \quad (2.6)$$

This means that

$$\sigma_1(S) = L(\text{PR}_1[S]), \quad \sigma_2(S) = R(\text{PR}_1[S])$$

and

$$\tau_1(S) = L(\text{PR}_2[S]) \quad \text{and} \quad \tau_2(S) = R(\text{PR}_2[S]),$$

i.e., $\sigma_i(S)$, $i = 1, 2$, are the left and right ends of $\text{PR}_1[S]$, and $\tau_i(S)$, $i = 1, 2$, are the left and right ends of $\text{PR}_2[S]$. See (2.3) and (2.4). More specifically, $\text{PR}_1[S] = \{(s, 0) : \sigma_1(S) \leq s \leq \sigma_2(S)\}$ provided S is bounded, and

$$\text{PR}_1[S] = \{(s, 0) : -\infty < s \leq \sigma_2(S)\} \quad \text{or} \quad \text{PR}_1[S] = \{(s, 0) : \sigma_1(S) \leq s < +\infty\}$$

provided $\sigma_1(S) = -\infty$, $\sigma_2(S) < \infty$, or $-\infty < \sigma_1(S)$, $\sigma_2(S) = +\infty$, respectively. Finally, $\text{PR}_1[S] = Ox_1$ if $\sigma_1(S) = -\infty$ and $\sigma_2(S) = +\infty$.

In the same fashion the numbers $\tau_1(S)$ and $\tau_2(S)$ determine the set $\text{PR}_2[S]$. ◀

Given sets $A_i \subset Ox_i$, $i = 1, 2$, we let $A_1 \times A_2$ denote a subset of \mathbf{R}^2 defined by

$$A_1 \times A_2 = \{a = (a_1, a_2) \in \mathbf{R}^2 : (a_1, 0) \in A_1, (0, a_2) \in A_2\}. \quad (2.7)$$

Given $a \in \mathbf{R}^2$ and $r > 0$, we let $Q(a, r)$ denote the square with center a and length of side $2r$:

$$Q(a, r) = \{y \in \mathbf{R}^2 : \|y - a\| \leq r\}.$$

In particular, $Q_0 = [-1, 1]^2 = Q(0, 1)$ is the unit ball of the Banach space $\ell_\infty^2 = (\mathbf{R}^2, \|\cdot\|)$.

Let S be a nonempty *convex* closed subset of \mathbf{R}^2 . By $\mathbf{Pr}(\cdot, S)$ we denote the operator of metric projection onto S in ℓ_∞^2 -norm. To each $a \in \mathbf{R}^2$ this operator assigns the set of all points in S that are nearest to a on S in the uniform norm. Thus,

$$\mathbf{Pr}(a, S) = S \cap Q(a, \text{dist}(a, S)). \quad (2.8)$$

Clearly, the set $\mathbf{Pr}(a, S)$ is either a singleton or a line segment in \mathbf{R}^2 parallel to one of the coordinate axes.

If $S \subset \mathbf{R}^2$ is convex bounded and centrally symmetric, by $\text{center}(S)$ we denote the center of S .

Given non-zero vectors $u, v \in \mathbf{R}^2$ we write $u \parallel v$ if u and v are collinear, and we write $u \not\parallel v$ whenever these vectors are non-collinear. We say that the vectors $u, v \in \mathbf{R}^2$ are *co-directed* if $u, v \neq 0$, u and v are collinear and have the same direction, i.e., $v = \alpha u$ for some $\alpha > 0$.

Let \tilde{u} and \tilde{v} be the *unit* vectors in the direction of the vectors u and v respectively. By $\theta(u, v) \in [0, 2\pi)$ we denote the angle of rotation from \tilde{u} to \tilde{v} in *the counterclockwise direction*. (Thus, $\theta(v, u) = 2\pi - \theta(u, v)$.) We refer to $\theta(u, v)$ as the angle between the vectors u and v .

Let ℓ_1 and ℓ_2 be two non-parallel straight lines in \mathbf{R}^2 ; in this case, we write $\ell_1 \not\parallel \ell_2$. Let $V = \ell_1 \cap \ell_2$. These two lines form two angles $\varphi_1, \varphi_2 \in [0, \pi)$, $\varphi_1 + \varphi_2 = \pi$, with the vertex at the point V . Let

$$\varphi(\ell_1, \ell_2) = \min\{\varphi_1, \varphi_2\}; \quad \text{clearly, } \varphi(\ell_1, \ell_2) \in [0, \pi/2].$$

We refer to $\varphi(\ell_1, \ell_2)$ as “*the angle between the straight lines ℓ_1 and ℓ_2* ”. If $\ell_1 \parallel \ell_2$ (i.e., ℓ_1 and ℓ_2 are parallel), we set $\varphi(\ell_1, \ell_2) = 0$.

2.2 Rectangles and rectangular hulls.

Recall that $\mathcal{I}(Ox_i)$, $i = 1, 2$, is the family of all nonempty convex closed subsets of the axis Ox_i . We set

$$\mathfrak{R}(\mathbf{R}^2) = \{\Pi = I_1 \times I_2 : I_1 \in \mathcal{I}(Ox_1), I_2 \in \mathcal{I}(Ox_2)\}. \quad (2.9)$$

We refer to every member of the family $\mathfrak{R}(\mathbf{R}^2)$ as a “*rectangle*”. Furthermore, throughout the paper, the word “rectangle” will mean an element of $\mathfrak{R}(\mathbf{R}^2)$, i.e., a closed rectangle (possibly unbounded) with sides parallel to the coordinate axes.

Clearly, thanks to definition (2.9),

$$\Pi = \text{Pr}_1[\Pi] \times \text{Pr}_2[\Pi] \quad \text{for every rectangle } \Pi \in \mathfrak{R}(\mathbf{R}^2). \quad (2.10)$$

Because Pr_i , $i = 1, 2$, is a continuous operator, for every *convex* closed set $S \subset \mathbf{R}^2$ its orthogonal projection $\text{Pr}_i[S]$ onto the axis Ox_i is a *convex* closed subset of Ox_i , i.e., $\text{Pr}_i[S] \in \mathcal{I}(Ox_i)$. Thus, thanks to this property and (2.7), we have the following:

$$\text{a convex closed set } \Pi \text{ belongs to } \mathfrak{R}(\mathbf{R}^2) \text{ if and only if } \Pi = \text{Pr}_1[\Pi] \times \text{Pr}_2[\Pi]. \quad (2.11)$$

We use the following elementary property of the intersection of rectangles in Section 6.

Claim 2.2 Let $\mathcal{R} \subset \mathfrak{R}(\mathbf{R}^2)$ be a finite collection of rectangles in \mathbf{R}^2 . Then the following equality

$$\bigcap_{\Pi \in \mathcal{R}} \Pi = \left\{ \bigcap_{\Pi \in \mathcal{R}} \text{PR}_1[\Pi] \right\} \times \left\{ \bigcap_{\Pi \in \mathcal{R}} \text{PR}_2[\Pi] \right\}. \quad (2.12)$$

holds. In particular, the intersection of all rectangles of the family \mathcal{R} is nonempty if and only if

$$\text{both } \bigcap_{\Pi \in \mathcal{R}} \text{PR}_1[\Pi] \neq \emptyset \text{ and } \bigcap_{\Pi \in \mathcal{R}} \text{PR}_2[\Pi] \neq \emptyset.$$

The proof of this claim is obvious.

We also note that any closed interval in \mathbf{R}^2 lying on a line parallel to a coordinate axis is a ‘‘rectangle’’. In particular, every closed interval on the axis Ox_1 or Ox_2 belongs to the family $\mathfrak{R}(\mathbf{R}^2)$.

Finally, given a bounded rectangle $\Pi \in \mathfrak{R}(\mathbf{R}^2)$ we let $\text{center}(\Pi)$ denote the center of Π .

We recall that $\text{Conv}(\mathbf{R}^2)$ denotes the family of all nonempty convex closed subsets of \mathbf{R}^n . Given a set $S \in \text{Conv}(\mathbf{R}^2)$, we let $\mathcal{H}[S]$ denote the ‘‘rectangular hull’’ of S , i.e., the smallest (with respect to inclusion) rectangle containing S . Thus,

$$\mathcal{H}[S] = \bigcap \{ \Pi : \Pi \in \mathfrak{R}(\mathbf{R}^2), \Pi \supset S \}. \quad (2.13)$$

Combining this definition with (2.12), we conclude that

$$\mathcal{H}[S] = \text{PR}_1[S] \times \text{PR}_2[S]. \quad (2.14)$$

Thus, given $S \in \text{Conv}(\mathbf{R}^2)$, its rectangular hull $\mathcal{H}[S]$ is the only rectangle Π for which

$$\text{PR}_1[\Pi] = \text{PR}_1[S] \quad \text{and} \quad \text{PR}_2[\Pi] = \text{PR}_2[S]. \quad (2.15)$$

Let us also note the following elementary property of rectangles: for every closed convex set $S \subset \mathbf{R}^2$ and every $r \geq 0$ we have

$$\mathcal{H}[S + rQ_0] = \mathcal{H}[S] + rQ_0. \quad (2.16)$$

In this section we present two important auxiliary results. The first of them is a variant of the classical Helly’s intersection theorem for rectangles.

Lemma 2.3 Let $\mathcal{K} \subset \mathfrak{R}(\mathbf{R}^2)$ be a finite collection of rectangles in \mathbf{R}^2 . Suppose that the intersection of every two rectangles from \mathcal{K} is nonempty. Then there exists a point in \mathbf{R}^2 common to all of the family \mathcal{K} .

Proof. Representation (2.10) reduces the problem to the one dimensional case. In this case the statement of the lemma is a variant of Helly’s theorem in \mathbf{R} . See, e.g. [19]. ■

The second auxiliary result is a Helly-type theorem formulated in terms of the orthogonal projections onto the coordinate axes.

Proposition 2.4 Let \mathcal{C} be a finite family of convex closed subsets in \mathbf{R}^2 such that

$$\text{PR}_1[C_1 \cap C'_1] \cap \text{PR}_1[C_2 \cap C'_2] \neq \emptyset \quad (2.17)$$

for every $C_1, C'_1, C_2, C'_2 \in \mathcal{C}$. Then

$$\bigcap \{ C : C \in \mathcal{C} \} \neq \emptyset. \quad (2.18)$$

Furthermore, in this case

$$\mathcal{H}[\bigcap \{ C : C \in \mathcal{C} \}] = \bigcap \{ \mathcal{H}[C \cap C'] : C, C' \in \mathcal{C} \}. \quad (2.19)$$

Proof. Condition (2.17) tells us that for every $C, C' \in \mathfrak{C}$ the set $C \cap C'$ is a nonempty. Because $C \cap C'$ is a convex subset of \mathbf{R}^2 , its projection onto Ox_1 , the set $\text{PR}_1[C \cap C'] \subset Ox_1$, is convex as well, i.e., this set is a closed interval in Ox_1 .

Let $\mathcal{W} = \{\text{PR}_1[C \cap C'] : C, C' \in \mathfrak{C}\}$. Then \mathcal{W} is a finite family of intervals, and, thanks to (2.17), every two members of this family have a common point. Helly's theorem tells us that in this case there exists a point in Ox_1 common to all of the family \mathcal{W} . See Lemma 2.3. Thus,

$$V = \bigcap_{C, C' \in \mathfrak{C}} \text{PR}_1[C \cap C'] \neq \emptyset. \quad (2.20)$$

Fix a point $v \in V$. Then, thanks to (2.20),

$$v \in \text{PR}_1[C \cap C'] \quad \text{for every } C, C' \in \mathfrak{C}. \quad (2.21)$$

Let

$$L = \{w \in \mathbf{R}^2 : \text{PR}_1[w] = v\} \quad (2.22)$$

be the straight line through v orthogonal to the axis Ox_1 .

Given $C \in \mathfrak{C}$, we set $K(C) = C \cap L$. Thanks to (2.21), $v \in \text{PR}_1[C \cap C] = \text{PR}_1[C]$ so that there exists $u_C \in C$ such that $\text{PR}_1[u_C] = v$. From this and (2.22), we have $u_C \in C \cap L$ proving that $K(C) \neq \emptyset$ for every $C \in \mathfrak{C}$. Clearly, each $K(C)$ is a closed interval lying on the straight line L . Let us show that there exists a point in L common to all these intervals.

Property (2.21) tells us that for every $C, C' \in \mathfrak{C}$ there exists a point $\tilde{u} \in C \cap C'$ such that $\text{PR}_1[\tilde{u}] = v$. Hence, thanks to (2.22), $\tilde{u} \in L$ so that

$$\tilde{u} \in L \cap C \cap C' = (L \cap C) \cap (L \cap C') = K(C) \cap K(C').$$

This proves that any two members of the family $\mathcal{K} = \{K(C) : C \in \mathfrak{C}\}$ have a common point. Furthermore, \mathcal{K} is a *finite* (because \mathfrak{C} is finite) family of intervals lying in L . Helly's theorem tells us that in this case $\bigcap \{K(C) : C \in \mathfrak{C}\} \neq \emptyset$. Thus,

$$\bigcap \{K(C) : C \in \mathfrak{C}\} = \bigcap \{L \cap C : C \in \mathfrak{C}\} = L \cap (\bigcap \{C : C \in \mathfrak{C}\}) \neq \emptyset$$

proving (2.18).

Let us prove (2.19). Clearly, the left hand side of the equality (2.19) is contained in its right hand side. Let us prove the converse statement.

Fix a point

$$u = (u_1, u_2) \in \bigcap \{\mathcal{H}[C \cap C'] : C, C' \in \mathfrak{C}\} \quad (2.23)$$

and prove that $u \in \mathcal{H}[\bigcap \{C : C \in \mathfrak{C}\}]$. Thanks to (2.14), property (2.23) is equivalent to the following one: $u_1 \in \text{PR}_1[\mathcal{T}]$ and $u_2 \in \text{PR}_2[\mathcal{T}]$ where $\mathcal{T} = \bigcap \{C : C \in \mathfrak{C}\}$.

Prove that $u_1 \in \text{PR}_1[\mathcal{T}]$. We let L_1 denote the straight line in \mathbf{R}^2 through the point $u = (u_1, u_2)$ orthogonal to the axis Ox_1 . Thus,

$$L_1 = \{w \in \mathbf{R}^2 : \text{PR}_1[w] = u_1\}. \quad (2.24)$$

Let us show that $L_1 \cap \mathcal{T} \neq \emptyset$. Indeed, thanks to (2.23), $u \in \mathcal{H}[C \cap C']$ provided $C, C' \in \mathfrak{C}$ so that, thanks to (2.14), $u_1 \in \text{PR}_1[C \cap C']$ for every $C, C' \in \mathfrak{C}$. Combining this property with definition

(2.24), we conclude that $L_1 \cap C \cap C' \neq \emptyset$ for all $C, C' \in \mathfrak{C}$. Thus, every two members of the (finite) family $\mathcal{K}_1 = \{L_1 \cap C : C \in \mathfrak{C}\}$ have a common point. Therefore, thanks to Helly's theorem, there exists a point $\tilde{u} \in \mathbf{R}^2$ such that

$$\tilde{u} \in \bigcap \{L_1 \cap C : C \in \mathfrak{C}\} = L_1 \cap \{C : C \in \mathfrak{C}\} = L_1 \cap \mathcal{T}.$$

Thus, $\tilde{u} \in L_1$ so that, thanks to (2.24), $u_1 = \text{Pr}[\tilde{u}]$. Furthermore, $\tilde{u} \in \mathcal{T} = \bigcap \{C : C \in \mathfrak{C}\}$ so that $u_1 \in \text{Pr}_1[\mathcal{T}]$.

In the same way we prove that $u_2 \in \text{Pr}_2[\mathcal{T}]$ completing the proof of the proposition. \blacksquare

We conclude the section with the following claim, which we use in Section 6.4 below.

Claim 2.5 *Let $I_i \in \mathcal{I}(Ox_i)$, be a closed interval on the axis Ox_i , $i = 1, 2$, and let $\mathcal{T} = I_1 \times I_2 \in \mathfrak{R}(\mathbf{R}^2)$ be a rectangle in \mathbf{R}^2 . Let $a_i = L(I_i)$ and $b_i = R(I_i)$, $i = 1, 2$, be the left and right ends of the interval I_i respectively. See (2.3) and (2.4).*

Then the distance $\text{dist}(O, \mathcal{T})$ from the origin $O(0, 0)$ to \mathcal{T} can be calculated as follows:

$$\text{dist}(O, \mathcal{T}) = \max\{|a_1|_+, |-b_1|_+, |a_2|_+, |-b_2|_+\}. \quad (2.25)$$

Furthermore, let $g = (g_1, g_2) = \text{center}(\text{Pr}(O, \mathcal{T}))$ be the center of the metric projection of the origin $O(0, 0)$ onto the rectangle \mathcal{T} , see (2.8). Then

$$g_1 = (L_1 + R_1)/2 \quad \text{and} \quad g_2 = (L_2 + R_2)/2 \quad (2.26)$$

where

$$L_i = \max\{-\text{dist}(O, \mathcal{T}), a_i\} \quad \text{and} \quad R_i = \min\{\text{dist}(O, \mathcal{T}), b_i\}, \quad i = 1, 2. \quad (2.27)$$

We leave the proof of this claim to the reader as an easy exercise.

2.3 Rectangles: intersections, neighborhoods and selections.

In this section we present several criteria and several constructive formulae for the optimal Lipschitz selections of set-valued mappings taking values in the family $\mathfrak{R}(\mathbf{R}^2)$ of all closed rectangles in \mathbf{R}^2 with sides parallel to the coordinate axes. See (2.9).

Let $\mathcal{I}(\mathbf{R})$ be the family of all closed intervals and all points in \mathbf{R} , and let $I_0 = [-1, 1]$. Given $a \in \mathbf{R}$ and $r \geq 0$, we set $rI_0 = [-r, r]$. We also recall that, given a bounded interval $I \in \mathcal{I}(\mathbf{R})$, by $\text{center}(I)$ we denote the center of I .

Lemma 2.6 *Let $\mathcal{K} \subset \mathfrak{R}(\mathbf{R}^2)$ be a family of rectangles in \mathbf{R}^2 with nonempty intersection. Then for every $r \geq 0$ the following equality*

$$\left(\bigcap_{K \in \mathcal{K}} K \right) + rQ_0 = \bigcap_{K \in \mathcal{K}} \{K + rQ_0\} \quad (2.28)$$

holds.

Proof. Obviously, the right hand side of (2.28) contains its left hand side. Let us prove that

$$\left(\bigcap_{K \in \mathcal{K}} K \right) + rQ_0 \supset \bigcap_{K \in \mathcal{K}} \{K + rQ_0\}. \quad (2.29)$$

This inclusion is based on the following simple claim: Let \mathcal{I} be a family of convex closed subsets of \mathbf{R} (intervals) with nonempty intersection. Let $K = [a, b] \subset \mathbf{R}$, be a closed bounded interval such that $K \cap I \neq \emptyset$ for every $I \in \mathcal{I}$. Then there exists a point common to K and all of the members of the family \mathcal{I} . (The proof is immediate from Helly's theorem in \mathbf{R} applied to the family $\mathcal{I} \cup \{K\}$ of closed intervals.

This claim implies the following one dimensional variant of inclusion (2.29): Let \mathcal{I} be a family of intervals in \mathbf{R} with nonempty intersection. Then

$$\left(\bigcap_{I \in \mathcal{I}} I \right) + rI_0 \supset \bigcap_{I \in \mathcal{I}} \{I + rI_0\} \quad \text{where } I_0 = [-1, 1]. \quad (2.30)$$

Indeed, if $u \in \bigcap_{I \in \mathcal{I}} \{I + rI_0\}$ then $[u - r, u + r] \cap I \neq \emptyset$ for every $I \in \mathcal{I}$. Therefore, thanks to the above claim, $[u - r, u + r] \cap (\bigcap_{I \in \mathcal{I}} I) \neq \emptyset$ proving that u belongs to the left hand side of (2.30).

Now, let us prove (2.29) using (2.30) and properties (2.11) and (2.12) of rectangles. For every $i = 1, 2$, we have

$$\text{PR}_i \left[\left(\bigcap_{K \in \mathcal{K}} K \right) + rQ_0 \right] = \left(\bigcap_{K \in \mathcal{K}} \text{PR}_i[K] \right) + r \text{PR}_i[Q_0] = U_i.$$

Furthermore,

$$\text{PR}_i \left[\bigcap_{K \in \mathcal{K}} \{K + rQ_0\} \right] = \bigcap_{K \in \mathcal{K}} \{\text{PR}_i[K] + r \text{PR}_i[Q_0]\} = V_i.$$

Thanks to inclusion (2.30), $U_i \supset V_i$, $i = 1, 2$, proving that the orthogonal projections onto the coordinate axes of the left hand side of (2.29) contain the corresponding projections of its right hand side. Because the left and right hand sides of (2.29) are *rectangles*, inclusion (2.29) holds.

The proof of the lemma is complete. \blacksquare

We will also need the following simple claim.

Claim 2.7 (i) *Let A and B be two closed intervals in \mathbf{R} . Then*

$$d_H(A, B) = \max\{|\inf A - \inf B|, |\sup A - \sup B|\}. \quad (2.31)$$

(See also our convention (2.2) for the cases of $\inf A, \inf B = -\infty$ and $\sup A, \sup B = +\infty$.)

(ii) *Let $\mathcal{A}, \mathcal{B} \in \mathfrak{R}(\mathbf{R}^2)$ be two bounded rectangles in \mathbf{R}^2 . Then*

$$\|\text{center}(\mathcal{A}) - \text{center}(\mathcal{B})\| \leq d_H(\mathcal{A}, \mathcal{B}). \quad (2.32)$$

Proof. (i) Let

$$r = d_H(A, B) \quad \text{and} \quad \delta = \max\{|\inf A - \inf B|, |\sup A - \sup B|\}.$$

Then, thanks to (2.1), $A + rI_0 \supset B$ proving that $\sup A + r \geq \sup B$ and $\inf A - r \leq \inf B$. By interchanging the roles of A and B we obtain also $\sup B + r \geq \sup A$ and $\inf B - r \leq \inf A$ proving that $\delta \leq r$.

Let us prove that $r \leq \delta$. Suppose that both A and B are bounded, i.e., $A = [\inf A, \sup A]$ and $B = [\inf B, \sup B]$. Let $\alpha \in [0, 1]$, and let

$$a_\alpha = \alpha \inf A + (1 - \alpha) \sup A \quad \text{and} \quad b_\alpha = \alpha \inf B + (1 - \alpha) \sup B.$$

Then $a_\alpha \in A$, $b_\alpha \in B$, and $|a_\alpha - b_\alpha| \leq \delta$ proving that $\text{dist}(a, B) \leq r$ for every $a \in A$ and $\text{dist}(b, A) \leq r$ for every $b \in B$. Hence, $A + rI_0 \supset B$ and $B + rI_0 \supset A$, so that, thanks to (2.1), $r \leq \delta$. In a similar way we prove this inequality whenever one of the intervals is unbounded. We leave the details to the interested reader as an easy exercise.

(ii) By orthogonal projecting to the coordinate axes, we can reduce the problem to the one dimensional case. In this case, given bounded intervals $A, B \in \mathcal{I}(\mathbf{R})$, we have

$$\text{center}(A) = (\inf A + \sup A)/2 \quad \text{and} \quad \text{center}(B) = (\inf B + \sup B)/2.$$

This and inequality (2.31) imply the required inequality

$$|\text{center}(A) - \text{center}(B)| \leq d_H(A, B)$$

proving the claim. \blacksquare

Let (\mathcal{M}, ρ) be a finite pseudometric space and let $\mathcal{T} : \mathcal{M} \rightarrow \mathfrak{R}(\mathbf{R}^2)$ be a set-valued mapping.

Given $\eta \geq 0$, let

$$\mathcal{T}^{[1]}[x : \eta] = \bigcap_{z \in \mathcal{M}} [\mathcal{T}(z) + \eta \rho(x, z) Q_0], \quad x \in \mathcal{M}, \quad (2.33)$$

be the η -metric refinement of \mathcal{T} . See Definition 1.6.

Proposition 2.8 (a) *If*

$$\mathcal{T}^{[1]}[x : \eta] \neq \emptyset \quad \text{for every } x \in \mathcal{M}, \quad (2.34)$$

then

$$d_H(\mathcal{T}^{[1]}[x : \eta], \mathcal{T}^{[1]}[y : \eta]) \leq \eta \rho(x, y) \quad \text{for all } x, y \in \mathcal{M}. \quad (2.35)$$

(b) *If*

$$\mathcal{T}(x) \cap \{\mathcal{T}(y) + \eta \rho(x, y) Q_0\} \neq \emptyset \quad \text{for all } x, y \in \mathcal{M}, \quad (2.36)$$

then properties (2.34) and (2.35) hold.

Furthermore, if the set $\mathcal{T}^{[1]}[x : \eta]$ is bounded for every $x \in \mathcal{M}$, then the mapping

$$\tau(x) = \text{center}(\mathcal{T}^{[1]}[x : \eta]), \quad x \in \mathcal{M},$$

is a Lipschitz selection of \mathcal{T} with $\|\tau\|_{\text{Lip}(\mathcal{M})} \leq \eta$.

Proof. (a) We know that $\mathcal{T}^{[1]}[x : \eta] \neq \emptyset$ so that, thanks to (2.34), Lemma 2.6 and definition (2.33), we have

$$\begin{aligned} \mathcal{T}^{[1]}[x : \eta] + \eta \rho(x, y) Q_0 &= \left\{ \bigcap_{z \in \mathcal{M}} [\mathcal{T}(z) + \eta \rho(x, z) Q_0] \right\} + \eta \rho(x, y) Q_0 \\ &= \bigcap_{z \in \mathcal{M}} [\mathcal{T}(z) + (\eta \rho(x, z) + \eta \rho(x, y)) Q_0]. \end{aligned}$$

Hence, thanks to the triangle inequality, we have

$$\mathcal{T}^{[1]}[x : \eta] + \eta \rho(x, y) Q_0 \supset \bigcap_{z \in \mathcal{M}} [\mathcal{T}(z) + \eta \rho(y, z) Q_0] = \mathcal{T}^{[1]}[y : \eta].$$

By interchanging the roles of x and y we also obtain

$$\mathcal{T}^{[1]}[y : \eta] + \eta\rho(x, y) Q_0 \supset \mathcal{T}^{[1]}[x : \eta].$$

These two inclusions imply the required inequality (2.35).

(b) Let us fix $x \in \mathcal{M}$ and prove that

$$\{\mathcal{T}(z) + \eta\rho(x, z) Q_0\} \cap \{\mathcal{T}(z') + \eta\rho(x, z') Q_0\} \neq \emptyset \quad \text{for every } z, z' \in \mathcal{M}. \quad (2.37)$$

Thanks to (2.36), there exist points $g(z) \in \mathcal{T}(z)$ and $g(z') \in \mathcal{T}(z')$ such that $\|g(z) - g(z')\| \leq \eta\rho(z, z')$. From this and the triangle inequality, we have

$$\|g(z) - g(z')\| \leq \eta\rho(z, x) + \eta\rho(x, z').$$

This implies the existence of a point $w \in \mathbf{R}^2$ such that

$$\|g(z) - w\| \leq \eta\rho(z, x) \quad \text{and} \quad \|g(z') - w\| \leq \eta\rho(z', x).$$

But $g(z) \in \mathcal{T}(z)$ and $g(z') \in \mathcal{T}(z')$ so that w belongs to the left hand side of (2.37) proving this property. Then, thanks to (2.37) and Lemma 2.3, the set $\mathcal{T}^{[1]}[x : \eta]$ is nonempty which proves property (2.34). In turn, inequality (2.35) follows from part (a) of the proposition.

Finally, inequalities (2.32) and (2.35) imply the required inequality $\|\tau\|_{\text{Lip}(\mathcal{M})} \leq \eta$ completing the proof of the proposition. ■

Let us give several explicit formulae for Lipschitz selections of set-valued mappings in the one dimensional case. Let $G : \mathcal{M} \rightarrow \mathcal{I}(\mathbf{R})$ be a set-valued mapping. We set

$$a_G(x) = \inf G(x) \quad \text{and} \quad b_G(x) = \sup G(x). \quad (2.38)$$

Thus, a_G and b_G are two functions on \mathcal{M} such that

$$a_G : \mathcal{M} \rightarrow \mathbf{R} \cup \{-\infty\}, \quad b_G : \mathcal{M} \rightarrow \mathbf{R} \cup \{+\infty\} \quad \text{and} \quad a_G(x) \leq b_G(x) \quad \text{for all } x \in \mathcal{M}.$$

Clearly,

$$\text{dist}(G(x), G(y)) = \max\{[a_G(x) - b_G(y)]_+, [a_G(y) - b_G(x)]_+\}. \quad (2.39)$$

(See our convention (2.2) for the case of $a_G(x) = -\infty$, $b_G(x) = +\infty$.)

Given $\eta \geq 0$, we introduce the following functions on \mathcal{M} :

$$a_G^{[1]}[x : \eta] = \sup_{y \in \mathcal{M}} \{a_G(y) - \eta\rho(x, y)\}, \quad b_G^{[1]}[x : \eta] = \inf_{y \in \mathcal{M}} \{b_G(y) + \eta\rho(x, y)\} \quad (2.40)$$

and

$$c_G[x : \eta] = (a_G^{[1]}[x : \eta] + b_G^{[1]}[x : \eta]) / 2. \quad (2.41)$$

Let $G^{[1]}[\cdot : \eta]$ be the η -metric refinement of G , i.e., a set-valued mapping on \mathcal{M} defined by

$$G^{[1]}[x : \eta] = \bigcap_{z \in \mathcal{M}} [G(z) + \eta\rho(x, z) I_0], \quad x \in \mathcal{M}. \quad \text{See (7.1).} \quad (2.42)$$

Comparing this definition with definitions (2.40) and (2.41), we conclude that for every $x \in \mathcal{M}$,

$$a_G^{[1]}[x : \eta] = \inf G^{[1]}[x : \eta], \quad b_G^{[1]}[x : \eta] = \sup G^{[1]}[x : \eta], \quad (2.43)$$

provided $G^{[1]}[x : \eta] \neq \emptyset$ and

$$c_G[x : \eta] = \text{center}(G^{[1]}[x : \eta]) \quad (2.44)$$

provided $G^{[1]}[x : \eta]$ is nonempty and *bounded*.

Remark 2.9 We note that the function $g^+ = b_G^{[1]}[\cdot : \eta]$ maps \mathcal{M} into \mathbf{R} if and only if $g^+ \neq +\infty$, i.e., $g^+(x^+) < \infty$ for some $x^+ \in \mathcal{M}$. Analogously, the mapping $g^- = a_G^{[1]}[\cdot : \eta]$ maps \mathcal{M} into \mathbf{R} if and only if $g^- \neq -\infty$. Finally, the mapping $c_G^{[1]}[\cdot : \eta] = (g^+ + g^-)/2$, see (2.41) and (2.44), is well defined if and only if both $g^+ \neq +\infty$ and $g^- \neq -\infty$. ◀

Proposition 2.10 *Let \mathcal{M} be finite and let $G : \mathcal{M} \rightarrow \mathcal{I}(\mathbf{R})$ be a set-valued mapping.*

Given $x \in \mathcal{M}$ and $\eta \geq 0$, the set $G^{[1]}[x : \eta]$ is nonempty if and only if

$$a_G^{[1]}[x : \eta] \leq b_G^{[1]}[x : \eta]. \quad (2.45)$$

See (2.40). Furthermore, if (2.45) holds, then

$$a_G^{[1]}[x : \eta] = \inf G^{[1]}[x : \eta] \quad \text{and} \quad b_G^{[1]}[x : \eta] = \sup G^{[1]}[x : \eta]. \quad (2.46)$$

Proof. Suppose that $G^{[1]}[x : \eta] \neq \emptyset$. Then for every $y_1, y_2 \in \mathcal{M}$, we have

$$(G(y_1) + \eta\rho(x, y_1)Q_0) \cap (G(y_2) + \eta\rho(x, y_2)Q_0) \neq \emptyset,$$

so that

$$\inf(G(y_1) + \eta\rho(x, y_1)Q_0) \leq \sup(G(y_2) + \eta\rho(x, y_2)Q_0).$$

Therefore, thanks to (2.38),

$$a_G(y_1) - \eta\rho(x, y_1) \leq b_G(y_1) + \eta\rho(x, y_2) \quad \text{for all } y_1, y_2 \in \mathcal{M}, \quad (2.47)$$

so that, thanks to (2.40), inequality (2.45) holds.

Let us now assume that (2.45) holds. Then, for every $y_1, y_2 \in \mathcal{M}$ inequality (2.47) holds as well so that

$$a_G(y_2) - b_G(y_1) \leq \eta\rho(x, y_1) + \eta\rho(x, y_2).$$

By interchanging the roles of y_1 and y_2 , we obtain also

$$a_G(y_1) - b_G(y_2) \leq \eta\rho(x, y_1) + \eta\rho(x, y_2).$$

From these inequalities and formula (2.39), we have

$$\text{dist}(G(y_1), G(y_2)) \leq \eta\rho(x, y_1) + \eta\rho(x, y_2).$$

But because $G(y_1)$ and $G(y_2)$ are two closed intervals in \mathbf{R} , there exist points $u_1 \in G(y_1)$ and $u_2 \in G(y_2)$ such that $\|u_1 - u_2\| \leq \eta\rho(x, y_1) + \eta\rho(x, y_2)$. Therefore, there exists a point $v \in \mathbf{R}$ such that $\|u_1 - v\| \leq \eta\rho(x, y_1)$ and $\|u_2 - v\| \leq \eta\rho(x, y_2)$. Clearly,

$$v \in (G(y_1) + \eta\rho(x, y_1)Q_0) \cap (G(y_2) + \eta\rho(x, y_2)Q_0).$$

This proves that any two closed intervals from the family $\mathcal{G} = \{G(y) + \eta\rho(x, y)Q_0 : y \in \mathcal{M}\}$ have a common point. Therefore, thanks to Helly's theorem in \mathbf{R} , there exists a point $w \in \mathbf{R}$ common to all of the intervals from \mathcal{G} .

Thus, w belongs to the intersection of all these intervals, and therefore, thanks to definition (2.42), $w \in G^{[1]}[x : \eta]$ proving that $G^{[1]}[x : \eta] \neq \emptyset$.

Finally, it remains to note that property (2.46) coincides with (2.43), and the proof of the proposition is complete. ■

Proposition 2.11 (*The Finiteness Principle for Lipschitz selections in \mathbf{R} .*) Let \mathcal{M} be finite and let $G : \mathcal{M} \rightarrow \mathcal{I}(\mathbf{R})$ be a set-valued mapping. Let $\eta \geq 0$.

Suppose that for every $x, y \in \mathcal{M}$ the restriction $G|_{\{x,y\}}$ of G to $\{x, y\}$ has a Lipschitz selection $g_{\{x,y\}}$ with $\|g_{\{x,y\}}\|_{\text{Lip}(\{x,y\}, \mathbf{R})} \leq \eta$. Then G has a Lipschitz selection $g : \mathcal{M} \rightarrow \mathbf{R}$ with Lipschitz seminorm $\|g\|_{\text{Lip}(\mathcal{M}, \mathbf{R})} \leq \eta$.

Furthermore, one can set $g = c_G^{[1]}[\cdot : \eta]$ provided there exist $x^+, x^- \in \mathcal{M}$ such that $\inf G(x^-) > -\infty$ and $\sup G(x^+) < \infty$. Also, one can set $g = b_G^{[1]}[\cdot : \eta]$ if $G(x^+)$ is bounded from above for some $x^+ \in \mathcal{M}$, or $g = a_G^{[1]}[\cdot : \eta]$ if $G(x^-)$ is bounded from below for some $x^- \in \mathcal{M}$.

Proof. The hypothesis of the proposition tells us that for every $x, y \in \mathcal{M}$ there exist points $a \in G(x)$ and $b \in G(y)$ such that $\|a - b\| \leq \eta \rho(x, y)$. Hence,

$$G(x) \cap \{G(y) + \eta \rho(x, y) I_0\} \neq \emptyset \quad \text{for all } x, y \in \mathcal{M}.$$

Here $I_0 = [-1, 1]$. Proposition 2.8 tells us that in this case $G^{[1]}[x : \eta] \neq \emptyset$ for every $x \in \mathcal{M}$, and

$$d_H(G^{[1]}[x : \eta], G^{[1]}[y : \eta]) \leq \eta \rho(x, y) \quad \text{for all } x, y \in \mathcal{M}.$$

From this, part (i) of Claim 2.7, and definitions (2.43) it follows that the inequality

$$\max \left\{ |a_G^{[1]}[x : \eta] - a_G^{[1]}[y : \eta]|, |b_G^{[1]}[x : \eta] - b_G^{[1]}[y : \eta]| \right\} \leq \eta \rho(x, y) \quad (2.48)$$

holds for all $x, y \in \mathcal{M}$.

Clearly, if $G \equiv \mathbf{R}$ then the constant mapping $g \equiv \{0\}$ on \mathcal{M} is a Lipschitz selection of G (with $\|g\|_{\text{Lip}(\mathcal{M}, \mathbf{R})} = 0$). Otherwise, either $g^+ = b_G^{[1]}[\cdot : \eta] \not\equiv +\infty$ or $g^- = a_G^{[1]}[\cdot : \eta] \not\equiv -\infty$. Therefore, thanks to Remark 2.9, either $g^+ : \mathcal{M} \rightarrow \mathbf{R}$ or $g^- : \mathcal{M} \rightarrow \mathbf{R}$.

Let us note that if at least one of the sets $G(x)$ is bounded, then the set $G^{[1]}[x : \eta]$ is bounded as well. In this case, the points

$$a_G^{[1]}[x : \eta], b_G^{[1]}[x : \eta] \in G^{[1]}[x : \eta] \subset G(x).$$

See definition (2.43). Therefore, in this case we can set either $g = g^+$ or $g = g^-$. Then, thanks to (2.48), in both cases the mapping $g : \mathcal{M} \rightarrow \mathbf{R}$ will be a Lipschitz selection of G with $\|g\|_{\text{Lip}(\mathcal{M}, \mathbf{R})} \leq \eta$. Also from this it follows that the mapping $g = c_G[\cdot : \eta] = (g^+ + g^-)/2$, see (2.44), has the same properties.

Note that, if $g^+ = b_G^{[1]}[\cdot : \eta] \not\equiv +\infty$ but $g^- = a_G^{[1]}[\cdot : \eta] \equiv -\infty$, then each interval $G^{[1]}[x : \eta]$ is *unbounded from below*. Because \mathcal{M} is finite, all these intervals have a common point, say A . Then the constant mapping $g \equiv \{A\}$ is a Lipschitz selection of G (with $\|g\|_{\text{Lip}(\mathcal{M}, \mathbf{R})} = 0$). Analogously, if $g^- \not\equiv -\infty$ but $g^+ \equiv +\infty$, there is a constant mapping which provides a Lipschitz selection of G .

Let us suppose that both $g^+ = b_G^{[1]}[\cdot : \eta] \not\equiv +\infty$ and $g^- = a_G^{[1]}[\cdot : \eta] \not\equiv -\infty$. In this case, thanks to Remark 2.9, the mapping $c_G^{[1]}[\cdot : \eta] = (g^+ + g^-)/2$, see (2.41) and (2.44), is well defined, i.e., each set $G^{[1]}[x : \eta]$ is nonempty and bounded. Clearly,

$$g(x) = c_G^{[1]}[x : \eta] \in G^{[1]}[x : \eta] \subset G(x) \quad \text{for every } x \in \mathcal{M},$$

proving that g is a selection of G . Thanks to (2.48), its Lipschitz seminorm $\|g\|_{\text{Lip}(\mathcal{M}, \mathbf{R})} \leq \eta$, and the proof of the proposition is complete. ■

Proposition 2.11 implies the following Finiteness Principle for rectangles in \mathbf{R}^2 .

Proposition 2.12 *Let $\eta \geq 0$. Let (\mathcal{M}, ρ) be a finite pseudometric space, and let $\mathcal{T} : \mathcal{M} \rightarrow \mathfrak{R}(\mathbf{R}^2)$ be a set-valued mapping. Suppose that for every $x, y \in \mathcal{M}$ the restriction $\mathcal{T}|_{\{x,y\}}$ of \mathcal{T} to $\{x, y\}$ has a Lipschitz selection $\tau_{\{x,y\}}$ with $\|\tau_{\{x,y\}}\|_{\text{Lip}(\{x,y\})} \leq \eta$.*

Then \mathcal{T} has a Lipschitz selection $\tau : \mathcal{M} \rightarrow \mathbf{R}^2$ with Lipschitz seminorm $\|\tau\|_{\text{Lip}(\mathcal{M})} \leq \eta$.

Proof. By orthogonal projecting onto the coordinate axes, we reduce the problem to the Finiteness Principle for Lipschitz selections in \mathbf{R} proven in Proposition 2.11. \blacksquare

3. The key theorem: Lipschitz selections and rectangular hulls.

We proceed to the proof of Theorem 1.10. This proof is based on an essential refinement of the approach developed in the author's paper [62, Section 3.1].

Let $\mathfrak{M} = (\mathcal{M}, \rho)$ be a finite pseudometric space, and let $F : \mathcal{M} \rightarrow \text{Conv}(\mathbf{R}^2)$ be a set-valued mapping. We recall that, given $\lambda \geq 0$ and $x, x', x'' \in \mathcal{M}$, by $\mathcal{W}_F[x, x', x'' : \lambda]$ we denote a (possibly empty) subset of \mathbf{R}^2 defined by

$$\mathcal{W}_F[x, x', x'' : \lambda] = \mathcal{H}[\{F(x') + \lambda\rho(x', x)Q_0\} \cap \{F(x'') + \lambda\rho(x'', x)Q_0\}]. \quad (3.1)$$

(Recall also that $\mathcal{H}[\cdot]$ denotes the rectangular hull of a set. See (2.13).)

We begin the proof of the key Theorem 1.10 in this section and conclude it in the next section.

PROOF OF THEOREM 1.10. Suppose that for every $x, x', x'', y, y', y'' \in \mathcal{M}$ condition (1.11) holds. Let us construct a Lipschitz selection $f : \mathcal{M} \rightarrow \mathbf{R}^2$ of F with Lipschitz seminorm $\|f\|_{\text{Lip}(\mathcal{M})} \leq 2\lambda + \tilde{\lambda}$. We will do this in three steps.

THE FIRST STEP. We introduce the $\tilde{\lambda}$ -metric refinement of F , see (1.3), i.e., a set-valued mapping on \mathcal{M} defined by the formula

$$F^{[1]}[x : \tilde{\lambda}] = \bigcap_{y \in \mathcal{M}} [F(y) + \tilde{\lambda}\rho(x, y)Q_0], \quad x \in \mathcal{M}. \quad (3.2)$$

Lemma 3.1 *For each $x \in \mathcal{M}$ the set $F^{[1]}[x : \tilde{\lambda}]$ is a nonempty closed convex subset of \mathbf{R}^2 . Moreover, for every $x \in \mathcal{M}$ the following representation holds:*

$$\mathcal{H}[F^{[1]}[x : \tilde{\lambda}]] = \cap\{\mathcal{W}_F[x, y, y' : \tilde{\lambda}] : y, y' \in \mathcal{M}\}. \quad (3.3)$$

Proof. Let us prove that

$$F^{[1]}[x : \tilde{\lambda}] \neq \emptyset \quad \text{for every } x \in \mathcal{M}. \quad (3.4)$$

Given $x \in \mathcal{M}$, we set

$$\mathfrak{C}_x = \{F(y) + \tilde{\lambda}\rho(x, y)Q_0 : y \in \mathcal{M}\}.$$

Then $F^{[1]}[x : \tilde{\lambda}] = \cap\{C : C \in \mathfrak{C}_x\}$. See (3.2).

Let us prove that for every $y_1, y'_1, y_2, y'_2 \in \mathcal{M}$ the sets

$$C_i = F(y_i) + \tilde{\lambda}\rho(x, y_i)Q_0 \quad \text{and} \quad C'_i = F(y'_i) + \tilde{\lambda}\rho(x, y'_i)Q_0, \quad i = 1, 2, \quad (3.5)$$

satisfy property (2.17).

First, let us note that, thanks to (1.11), the set $\mathcal{W}_F[x, y, z : \tilde{\lambda}] \neq \emptyset$ for all $x, y, z \in \mathcal{M}$.

In particular, from this and definition (3.1), it follows that

$$\{F(y) + \tilde{\lambda}\rho(x, y) Q_0\} \cap \{F(z) + \tilde{\lambda}\rho(x, z) Q_0\} \neq \emptyset \quad \text{for every } y, z \in \mathcal{M}$$

proving that *any two elements of \mathfrak{C}_x have a common point*.

Property (1.11) tells us that

$$\mathcal{W}_F[x, y_1, y'_1 : \tilde{\lambda}] \cap \mathcal{W}_F[x, y_2, y'_2 : \tilde{\lambda}] \neq \emptyset \quad \text{for every } y_1, y'_1, y_2, y'_2 \in \mathcal{M}.$$

Thanks to (3.1) and (3.5),

$$\mathcal{W}_F[x, y_1, y'_1 : \tilde{\lambda}] = \mathcal{H}[C_1 \cap C'_1] \quad \text{and} \quad \mathcal{W}_F[x, y_2, y'_2 : \tilde{\lambda}] = \mathcal{H}[C_2 \cap C'_2]$$

proving that

$$\mathcal{H}[C_1 \cap C'_1] \cap \mathcal{H}[C_2 \cap C'_2] \neq \emptyset \quad \text{for every } C_1, C'_1, C_2, C'_2 \in \mathfrak{C}_x.$$

Hence,

$$\text{Pr}_1[\mathcal{H}[C_1 \cap C'_1]] \cap \text{Pr}_1[\mathcal{H}[C_2 \cap C'_2]] \neq \emptyset$$

so that, thanks to (2.15), $\text{Pr}_1[C_1 \cap C'_1] \cap \text{Pr}_1[C_2 \cap C'_2] \neq \emptyset$ proving (2.17).

Because the set \mathcal{M} is *finite*, the family \mathfrak{C}_x is finite as well. Therefore, thanks to Proposition 2.4, $\cap\{C : C \in \mathfrak{C}_x\} \neq \emptyset$ proving (3.4).

Finally, (2.19) and (3.1) imply formula (3.3) completing the proof of the lemma. \blacksquare

Following **STEP 2** of the Projection Algorithm, given $x \in \mathcal{M}$, we let $\mathcal{T}_{F, \tilde{\lambda}}(x)$ denote the rectangular hull of the set $F^{[1]}[x : \tilde{\lambda}]$. Cf. (1.5). Thus, $\mathcal{T}_{F, \tilde{\lambda}}$ is a set-valued mapping from \mathcal{M} into $\mathfrak{R}(\mathbf{R}^2)$ defined by

$$\mathcal{T}_{F, \tilde{\lambda}}(x) = \mathcal{H}[F^{[1]}[x : \tilde{\lambda}]] = \mathcal{H}\left[\cap\{F(y) + \tilde{\lambda}\rho(x, y) Q_0 : y \in \mathcal{M}\}\right], \quad x \in \mathcal{M}. \quad (3.6)$$

Let us note that formula (3.3) provides the following representation of the mapping $\mathcal{T}_{F, \tilde{\lambda}}$:

$$\mathcal{T}_{F, \tilde{\lambda}}(x) = \cap\{\mathcal{W}_F[x, x', x'' : \tilde{\lambda}] : x', x'' \in \mathcal{M}\}, \quad x \in \mathcal{M}. \quad (3.7)$$

Let us note that, thanks to Lemma 3.1, $F^{[1]}[x : \tilde{\lambda}] \neq \emptyset$ for every $x \in \mathcal{M}$. Therefore, thanks to (3.6),

$$\mathcal{T}_{F, \tilde{\lambda}}(x) \neq \emptyset \quad \text{for all } x \in \mathcal{M}. \quad (3.8)$$

THE SECOND STEP. At this step we prove the existence of a Lipschitz selection of the set-valued mapping $\mathcal{T}_{F, \tilde{\lambda}}$.

First, following **STEP 3** of the Projection Algorithm, we let $\mathcal{T}_{F, \tilde{\lambda}}^{[1]}[\cdot : \lambda]$ denote the λ -metric refinement of the mapping $\mathcal{T}_{F, \tilde{\lambda}}$, see (1.6). Thus,

$$\mathcal{T}_{F, \tilde{\lambda}}^{[1]}[x : \lambda] = \bigcap_{z \in \mathcal{M}} \left[\mathcal{T}_{F, \tilde{\lambda}}(z) + \lambda\rho(x, z) Q_0 \right], \quad x \in \mathcal{M}.$$

Proposition 3.2 (i) *The set-valued mapping $\mathcal{T}_{F, \tilde{\lambda}} : \mathcal{M} \rightarrow \mathfrak{R}(\mathbf{R}^2)$ has a Lipschitz selection $g : \mathcal{M} \rightarrow \mathbf{R}^2$ with Lipschitz seminorm $\|g\|_{\text{Lip}(\mathcal{M})} \leq \lambda$;*

(ii) For every $x \in \mathcal{M}$, the following property

$$\mathcal{T}_{F,\tilde{\lambda}}^{[1]}[x : \lambda] \neq \emptyset \quad (3.9)$$

holds. Furthermore, for every $x, y \in \mathcal{M}$, we have

$$d_H\left(\mathcal{T}_{F,\tilde{\lambda}}^{[1]}[x : \lambda], \mathcal{T}_{F,\tilde{\lambda}}^{[1]}[y : \lambda]\right) \leq \lambda \rho(x, y). \quad (3.10)$$

(Recall that d_H denotes the Hausdorff distance between sets.)

(iii) If each rectangle $\mathcal{T}_{F,\tilde{\lambda}}^{[1]}[x : \lambda]$, $x \in \mathcal{M}$, is bounded, then the mapping

$$g_F(x) = \text{center}\left(\mathcal{T}_{F,\tilde{\lambda}}^{[1]}[x : \lambda]\right), \quad x \in \mathcal{M},$$

is a Lipschitz selection of $\mathcal{T}_{F,\tilde{\lambda}}$ with $\|g_F\|_{\text{Lip}(\mathcal{M})} \leq \lambda$.

Proof. (i) Proposition 2.12 tells us that the required Lipschitz selection g exists provided for every $x, y \in \mathcal{M}$ the restriction $\mathcal{T}_{F,\tilde{\lambda}}|_{\{x,y\}}$ of $\mathcal{T}_{F,\tilde{\lambda}}$ to $\{x, y\}$ has a Lipschitz selection $g_{\{x,y\}}$ with Lipschitz seminorm $\|g_{\{x,y\}}\|_{\text{Lip}(\{x,y\})} \leq \lambda$. Clearly, this requirement is equivalent to the following property:

$$\mathcal{T}_{F,\tilde{\lambda}}(x) \cap \{\mathcal{T}_{F,\tilde{\lambda}}(y) + \lambda \rho(x, y)Q_0\} \neq \emptyset \quad \text{for every } x, y \in \mathcal{M}. \quad (3.11)$$

Let us prove that this property holds. Let $x, y \in \mathcal{M}$ and let

$$\mathfrak{I}_x = \{\mathcal{W}_F[x, x', x'' : \tilde{\lambda}] : x', x'' \in \mathcal{M}\} \quad \text{and} \quad \mathfrak{I}_y = \{\mathcal{W}_F[y, y', y'' : \tilde{\lambda}] : y', y'' \in \mathcal{M}\}.$$

Thanks to (3.7),

$$\mathcal{T}_{F,\tilde{\lambda}}(x) = \cap\{W : W \in \mathfrak{I}_x\} \quad \text{and} \quad \mathcal{T}_{F,\tilde{\lambda}}(y) = \cap\{W : W \in \mathfrak{I}_y\}. \quad (3.12)$$

Since \mathcal{M} is finite, \mathfrak{I}_x and \mathfrak{I}_y are *finite* families of rectangles. Thanks to (3.8), the set $\mathcal{T}_{F,\tilde{\lambda}}(z) \neq \emptyset$ for every $z \in \mathcal{M}$, so that each family has a nonempty intersection.

Let $r = \lambda \rho(x, y)$. Then, thanks to (3.12) and Lemma 2.6,

$$\mathcal{T}_{F,\tilde{\lambda}}(y) + \lambda \rho(x, y)Q_0 = \cap\{W : W \in \mathfrak{I}_y\} + rQ_0 = \cap\{W + rQ_0 : W \in \mathfrak{I}_y\}$$

so that

$$\mathcal{T}_{F,\tilde{\lambda}}(x) \cap \{\mathcal{T}_{F,\tilde{\lambda}}(y) + \lambda \rho(x, y)Q_0\} = [\cap\{W : W \in \mathfrak{I}_x\}] \cap [\cap\{W + rQ_0 : W \in \mathfrak{I}_y\}]. \quad (3.13)$$

Let $\tilde{\mathfrak{I}} = \mathfrak{I}_x \cup \mathfrak{I}_y^+$ where $\mathfrak{I}_y^+ = \{W + rQ_0 : W \in \mathfrak{I}_y\}$. Thanks to (3.13), property (3.11) holds provided the family of rectangles $\tilde{\mathfrak{I}}$ has a common point. Because \mathfrak{I}_x and \mathfrak{I}_y are *finite* families, the family $\tilde{\mathfrak{I}}$ is finite as well. Therefore, thanks to Helly's intersection theorem for rectangles, see Lemma 2.3, there exists a point common to all of the family $\tilde{\mathfrak{I}}$ provided $W' \cap W'' \neq \emptyset$ for every $W', W'' \in \tilde{\mathfrak{I}}$.

Clearly, $W' \cap W'' \neq \emptyset$ if $W', W'' \in \mathfrak{I}_x$ or $W', W'' \in \mathfrak{I}_y^+$ because both \mathfrak{I}_x and \mathfrak{I}_y^+ has a nonempty intersection. Let $W' = \mathcal{W}_F[x, x', x'' : \tilde{\lambda}]$, $x', x'' \in \mathcal{M}$, and let $W'' = \mathcal{W}_F[y, y', y'' : \tilde{\lambda}] + rQ_0$, $y', y'' \in \mathcal{M}$, be two arbitrary members of \mathfrak{I}_x and \mathfrak{I}_y^+ respectively. Then, thanks to assumption (1.11) of Theorem 1.10, $W' \cap W'' \neq \emptyset$.

Thus, the hypothesis of Lemma 2.3 holds for $\tilde{\mathfrak{I}}$. Therefore, this family has a common point proving the required property (3.11).

Let us prove parts (ii) and (iii). We note that, thanks to property (3.11), the mapping $\mathcal{T} = \mathcal{T}_{F, \tilde{\lambda}}$ satisfies the conditions of the hypothesis of part (b) of Proposition 2.8 with $\eta = \lambda$. This proposition tells us that property (3.9) and inequality (3.10) hold proving part (ii). Furthermore, part (b) of Proposition 2.8 proves part (iii).

The proof of Proposition 3.2 is complete. \blacksquare

THE THIRD STEP. At this step we construct a Lipschitz selection f of the set-valued mapping F with Lipschitz seminorm at most $2\lambda + \tilde{\lambda}$.

We recall that the set-valued mapping $F^{[1]}[\cdot : \tilde{\lambda}]$ and its rectangular hull, the set-valued mapping $\mathcal{T}_{F, \tilde{\lambda}} = \mathcal{H}[F^{[1]}[\cdot : \tilde{\lambda}]]$, are defined by formulae (3.2) and (3.6) respectively.

Part (i) of Proposition 3.2 tells us that $\mathcal{T}_{F, \tilde{\lambda}}$ has a Lipschitz selection with Lipschitz seminorm at most λ . In other words, there exists a mapping $g : \mathcal{M} \rightarrow \mathbf{R}^2$ such that

$$g(x) \in \mathcal{T}_{F, \tilde{\lambda}}(x) = \mathcal{H}[F^{[1]}[x : \tilde{\lambda}]] \quad \text{for every } x \in \mathcal{M}, \quad (3.14)$$

and

$$\|g(x) - g(y)\| \leq \lambda \rho(x, y) \quad \text{for all } x, y \in \mathcal{M}. \quad (3.15)$$

Proposition 3.3 *Let $g : \mathcal{M} \rightarrow \mathbf{R}^2$ be an arbitrary Lipschitz selection of the set-valued mapping $\mathcal{T}_{F, \tilde{\lambda}} : \mathcal{M} \rightarrow \mathfrak{R}(\mathbf{R}^2)$ with Lipschitz seminorm at most λ , i.e., a mapping satisfying conditions (3.14) and (3.15).*

We define a mapping $f : \mathcal{M} \rightarrow \mathbf{R}^2$ by letting

$$f(x) = \mathbf{Pr}\left(g(x), F^{[1]}[x : \tilde{\lambda}]\right), \quad x \in \mathcal{M}. \quad (3.16)$$

(Recall that $\mathbf{Pr}(\cdot, S)$ is the operator of metric projection onto a convex closed $S \subset \mathbf{R}^2$. See (2.8).)

Then the following properties hold:

(★1) The mapping f is well defined, i.e., $f(x)$ is a singleton for every $x \in \mathcal{M}$. In this case

$$f(x) = \mathbf{Pr}\left(g(x), F^{[1]}[x : \tilde{\lambda}]\right) \in F^{[1]}[x : \tilde{\lambda}] \subset F(x) \quad \text{for every } x \in \mathcal{M},$$

so that f is a selection of F on \mathcal{M} ;

(★2) The mapping $f : \mathcal{M} \rightarrow \mathbf{R}^2$ is Lipschitz with Lipschitz seminorm $\|f\|_{\text{Lip}(\mathcal{M})} \leq 2\lambda + \tilde{\lambda}$.

The proof of this proposition is based on a number of auxiliary results. The first of these is the following lemma.

Lemma 3.4 *Let $S \subset \mathbf{R}^2$ be a nonempty convex closed set. Then for every point $a \in \mathcal{H}[S]$ the metric projection $\mathbf{Pr}(a, S)$ is a singleton. Furthermore, $\mathbf{Pr}(a, S)$ coincides with a vertex of the square $Q(a, \text{dist}(a, S))$.*

Proof. The proof of this lemma is given in [62, p. 301]. See also [65, p. 68]. For the convenience of the reader, we give it here.

Clearly, if $a \in S$, nothing to prove. Suppose $a \notin S$ so that $r = \text{dist}(a, S) > 0$. Because S is closed, $\mathbf{Pr}(a; S) \neq \emptyset$. Furthermore, $\mathbf{Pr}(a; S) = S \cap Q = S \cap \partial Q$ where $Q = Q(a, r)$.

Because $\mathbf{Pr}(a; S)$ is a nonempty convex set lying on the boundary of Q , it belongs to a certain side of the square Q . In other words, there exist two distinct vertices of Q , say A and B , such that $\mathbf{Pr}(a; S) \subset [A, B]$.

Let us prove that

$$\text{either } \mathbf{Pr}(a; S) = \{A\} \quad \text{or} \quad \mathbf{Pr}(a; S) = \{B\}. \quad (3.17)$$

Indeed, otherwise there exists a point $p \in (A, B) \cap \mathbf{Pr}(a; S)$. Let ℓ be the straight line passing through A and B . Clearly, ℓ is parallel to a coordinate axis. Let H_1, H_2 be the closed half-planes determined by ℓ . (Thus $\ell = H_1 \cap H_2$ and $H_1 \cup H_2 = \mathbf{R}^2$.) Clearly, Q is contained in one of these half-planes, say in H_1 . Because $\text{dist}(a, \ell) = r > 0$, the point $a \in H_1^{\text{int}}$ where H_1^{int} denotes the interior of H_1 .

Prove that in this case $S \subset H_2$, i.e., the straight line ℓ separates (not strictly) the square Q and the set S . Indeed, suppose that there exists a point $b \in S \cap H_1^{\text{int}}$. Then also $(p, b] \subset H_1^{\text{int}}$ because $p \in \partial H_1 = \ell$. But $p \in (A, B)$ so that $(p, b] \cap Q^{\text{int}} \neq \emptyset$. On the other hand, because S is convex and $p \in \partial S$, the interval $(p, b] \subset S$ proving that $S \cap Q^{\text{int}} \neq \emptyset$. But $S \cap Q \subset \partial Q$, a contradiction.

Thus, $S \subset H_2$ and $Q \subset H_1$. But $a \in H_1^{\text{int}}$ so that $a \notin H_2$. Clearly, $H_2 \in \mathfrak{R}(\mathbf{R}^2)$, i.e., H_2 is an (unbounded) rectangle. Therefore $\mathcal{H}[S] \subset H_2$, see definition (2.13). Therefore, thanks to the lemma's hypothesis, $a \in \mathcal{H}[S] \subset H_2$, a contradiction.

This contradiction implies (3.17) completing the proof of the lemma. ■

Clearly, this lemma implies the statement (★1) of Proposition 3.3.

Let us prove the statement (★2) which is equivalent to the inequality

$$\|f(x) - f(y)\| \leq (2\lambda + \tilde{\lambda})\rho(x, y) \quad \text{for every } x, y \in \mathcal{M}. \quad (3.18)$$

The proof of this inequality relies on a number of auxiliary results which we present in the next section.

4. Proof of the key theorem: the final step.

Lemma 4.1 *Let $A, B \subset \mathbf{R}^2$ be nonempty convex closed sets such that $A \subset B$, and let $a \in \mathcal{H}[A]$.*

Then $\mathbf{Pr}(a, A)$ and $\mathbf{Pr}(a, B)$ are singletons having the following properties:

(i) $\mathbf{Pr}(a, B) \in [\mathbf{Pr}(a, A), a]$;

(ii) *The following equality holds:*

$$\|\mathbf{Pr}(a, A) - \mathbf{Pr}(a, B)\| = \text{dist}(a, A) - \text{dist}(a, B).$$

Proof. The proof of the lemma is given in [62, p. 302]. See also [65, p. 69]. For the convenience of the reader, we present it here.

First, we note that if $a \in B$, the statement of the lemma is immediate from Lemma 3.4.

Suppose that $a \notin B$. In this case, Lemma 3.4 tells us that $\mathbf{Pr}(a; A)$ is one of the vertices of the square $Q(a, r)$ with $r = \text{dist}(a, A) > 0$. Because $A \subset B$, the point $a \in \mathcal{H}[B]$ so that, thanks to Lemma 3.4, $\mathbf{Pr}(a; B)$ is a vertex of the square $Q(a, \alpha)$ where $\alpha = \text{dist}(a, B) > 0$.

Using a suitable shift and dilation, without loss of generality, we can assume that

$$a = (0, 0), \quad r = \text{dist}(a, A) = 1, \quad \text{and} \quad \mathbf{Pr}(a; A) = (1, 1).$$

Clearly, in this case $0 < \alpha \leq 1$. Furthermore, under these conditions, the statement of the lemma is equivalent to the property

$$\mathbf{Pr}(a; B) = (\alpha, \alpha). \quad (4.1)$$

Suppose that this property does not hold, i.e., $\Pr(a; B) \in \{(\alpha, -\alpha), (-\alpha, \alpha), (-\alpha, -\alpha)\}$.

In order to get a contradiction, we construct a straight line ℓ_A which passes through $(1, 1)$ and separates (not strictly) the square $Q(a, r) = [-1, 1]^2$ and A . This line determines two closed half-planes, S_A^+ and S_A^- , with the common boundary (i.e., the line ℓ_A) such that $\mathbf{R}^2 = S_A^+ \cup S_A^-$. One of them, say S_A^+ , contains A , so that $S_A^- \supset Q(a, r)$. We know that S_A^+ contains $(1, 1)$ and does not contain intrinsic points of the square $[-1, 1]^2$, so that $Q(a, r) \cap \ell_A = (1, 1)$. Therefore, the half-plane S_A^+ admits the following representation:

$$S_A^+ = \{x = (x_1, x_2) \in \mathbf{R}^2 : (x_1 - 1)h_1 + (x_2 - 1)h_2 \geq 0\} \quad (4.2)$$

where $h_1, h_2 > 0$ are certain numbers.

Let us assume that $\Pr(a; B) = (-\alpha, \alpha)$ and show that this assumption leads to a contradiction. We let ℓ_B denote a straight line which passes through the point $(-\alpha, \alpha)$ and separates the square $Q(a, \text{dist}(a, B)) = [-\alpha, \alpha]^2$ and the set B . Let S_B^+ be the one of the two half-planes determined by ℓ_B which contains B . Then the other half-plane, S_B^- , contains $Q(a, \text{dist}(a, B))$, and $S_B^+ \cap S_B^- = \ell_B$.

We know that S_B^+ contains the point $(-\alpha, \alpha)$ on its boundary and does not contain intrinsic points of the square $[-\alpha, \alpha]^2$. Therefore, this half-plane can be represented in the form

$$S_B^+ = \{(x_1, x_2) \in \mathbf{R}^2 : -(x_1 + \alpha)s_1 + (x_2 - \alpha)s_2 \geq 0\} \quad (4.3)$$

with certain $s_1, s_2 > 0$.

Thus, $A \subset S_A^+$ and $A \subset B \subset S_B^+$, so that $A \subset S_A^+ \cap S_B^+$ proving that for every $x = (x_1, x_2) \in A$ we have

$$(x_1 - 1)h_1 + (x_2 - 1)h_2 \geq 0 \quad \text{and} \quad -(x_1 + \alpha)s_1 + (x_2 - \alpha)s_2 \geq 0. \quad (4.4)$$

See (4.2) and (4.3). Note also that since $S_A^+ \cap S_B^+ \supset A \neq \emptyset$, we have $h_2 + s_2 > 0$.

Let us prove that inequalities (4.4) imply the following inclusion:

$$A \subset \mathcal{H}_\alpha = \{x = (x_1, x_2) \in \mathbf{R}^2 : x_2 \geq \alpha\}. \quad (4.5)$$

Indeed, it is easy to see that from (4.4) we have

$$x_2 - \alpha \geq \frac{s_1((1 + \alpha)h_1 + (1 - \alpha)h_2)}{s_1h_2 + s_2h_1} \geq 0, \quad x = (x_1, x_2) \in A,$$

proving (4.5).

Let us note that $\mathcal{H}_\alpha \in \mathfrak{R}(\mathbf{R}^2)$ so that $\mathcal{H}[A] \subset \mathcal{H}_\alpha$. Therefore, thanks to the lemma's assumption, $a = (0, 0) \in \mathcal{H}_\alpha$. But $\alpha > 0$ so that $a = (0, 0) \notin \mathcal{H}_\alpha$, a contradiction.

In a similar way we get a contradiction provided $\Pr(a; B) = (\alpha, -\alpha)$ or $\Pr(a; B) = (-\alpha, -\alpha)$ proving the required property (4.1) and the lemma. \blacksquare

Lemma 4.2 (i) *Let $u \in \mathcal{M}$, and let $a \in \mathcal{H}[F^{[1]}[u : \tilde{\lambda}]]$. Then*

$$\text{dist}(a, F^{[1]}[u : \tilde{\lambda}]) = \max_{z \in \mathcal{M}} \text{dist}(a, F(z) + \tilde{\lambda}\rho(u, z)Q_0) = \max_{z \in \mathcal{M}} [\text{dist}(a, F(z)) - \tilde{\lambda}\rho(u, z)]_+; \quad (4.6)$$

(ii) *Let $u, v \in \mathcal{M}$, and let $a \in \mathcal{H}[F^{[1]}[u : \tilde{\lambda}]]$ and $b \in \mathcal{H}[F^{[1]}[v : \tilde{\lambda}]]$. Then*

$$|\text{dist}(a, F^{[1]}[u : \tilde{\lambda}]) - \text{dist}(b, F^{[1]}[v : \tilde{\lambda}])| \leq \|a - b\| + \tilde{\lambda}\rho(u, v). \quad (4.7)$$

Proof. (i) Let $A = F^{[1]}[u : \tilde{\lambda}]$ and, given $z \in \mathcal{M}$, let $A_z = F(z) + \tilde{\lambda}\rho(u, z)Q_0$. Then, thanks to (3.2), $A = \cap\{A_z : z \in \mathcal{M}\}$. Our goal is to prove that

$$\text{dist}(a, A) = \max\{\text{dist}(a, A_z) : z \in \mathcal{M}\} \quad \text{provided } a \in \mathcal{H}[A]. \quad (4.8)$$

Lemma 3.1 tells us that A is a nonempty convex and closed subset of \mathbf{R}^2 . Because $A \subset A_z$ for each $z \in \mathcal{M}$, the left hand side of the above equality majorizes its right hand side.

Prove the converse inequality. If $a \in A$, nothing to prove.

Let $a \notin A$, and let $\varepsilon \in (0, \text{dist}(a, A))$ be arbitrary. We know that $a \in \mathcal{H}[A]$ so that, thanks to Lemma 4.1, $\mathbf{Pr}(a, A)$ is a singleton.

We let a_ε denote a point on the interval $(\mathbf{Pr}(a, A), a]$ such that $\|a_\varepsilon - \mathbf{Pr}(a, A)\| < \varepsilon$. Because $a_\varepsilon \notin A$ and $A = \cap\{A_z : z \in \mathcal{M}\}$, there exists an element $\tilde{z} \in \mathcal{M}$ such that $a_\varepsilon \notin A_{\tilde{z}}$. Note that $A \subset A_{\tilde{z}}$. Lemma 4.1 tells us that in this case $\mathbf{Pr}(a, A_{\tilde{z}})$ is a singleton such that $\mathbf{Pr}(a, A_{\tilde{z}}) \in [\mathbf{Pr}(a, A), a]$.

Then $\mathbf{Pr}(a, A_{\tilde{z}}) \in [\mathbf{Pr}(a, A), a_\varepsilon]$; otherwise $a_\varepsilon \in [\mathbf{Pr}(a, A), \mathbf{Pr}(a, A_{\tilde{z}})] \subset A_{\tilde{z}}$, a contradiction.

This proves that $\|\mathbf{Pr}(a, A) - \mathbf{Pr}(a, A_{\tilde{z}})\| < \varepsilon$. Hence,

$$\begin{aligned} \text{dist}(a, A) &= \|a - \mathbf{Pr}(a, A)\| \leq \|a - \mathbf{Pr}(a, A_{\tilde{z}})\| + \|\mathbf{Pr}(a, A_{\tilde{z}}) - \mathbf{Pr}(a, A)\| \\ &\leq \text{dist}(a, A_{\tilde{z}}) + \|a_\varepsilon - \mathbf{Pr}(a, A)\| \leq \text{dist}(a, A_{\tilde{z}}) + \varepsilon. \end{aligned}$$

Since $\varepsilon > 0$ can be chosen as small as desired, this implies the required inequality (4.8) proving part (i) of the lemma.

(ii) Let $A = F^{[1]}[u : \tilde{\lambda}]$ and $B = F^{[1]}[v : \tilde{\lambda}]$. Then, thanks to (4.6),

$$\begin{aligned} |\text{dist}(a, A) - \text{dist}(a, B)| &= \left| \sup_{z \in \mathcal{M}} [\text{dist}(a, F(z)) - \tilde{\lambda}\rho(u, z)]_+ - \sup_{z \in \mathcal{M}} [\text{dist}(a, F(z)) - \tilde{\lambda}\rho(v, z)]_+ \right| \\ &\leq \sup_{z \in \mathcal{M}} |[\text{dist}(a, F(z)) - \tilde{\lambda}\rho(u, z)]_+ - [\text{dist}(a, F(z)) - \tilde{\lambda}\rho(v, z)]_+| \\ &\leq \tilde{\lambda} \sup_{z \in \mathcal{M}} |\rho(u, z) - \rho(v, z)| \end{aligned}$$

so that, thanks to the triangle inequality,

$$|\text{dist}(a, A) - \text{dist}(a, B)| \leq \tilde{\lambda}\rho(u, v). \quad (4.9)$$

Next,

$$|\text{dist}(a, A) - \text{dist}(b, B)| \leq |\text{dist}(a, A) - \text{dist}(a, B)| + |\text{dist}(a, B) - \text{dist}(b, B)|.$$

Because $\text{dist}(\cdot, B)$ is a Lipschitz function, from this and (4.9), we have (4.7) completing the proof of the lemma. ■

Let $\delta \geq 0$, and let

$$H_1 \text{ and } H_2 \text{ be two half-planes with } \text{dist}(H_1, H_2) \leq \delta. \quad (4.10)$$

Let $\ell_i = \partial H_i$ be the boundary of the half-plane H_i , $i = 1, 2$. Let us represent the half-planes H_i , $i = 1, 2$, in the form $H_i = \{u \in \mathbf{R}^2 : \langle \mathbf{h}_i, u \rangle + \alpha_i \leq 0\}$ where \mathbf{h}_i is a unit vector and $\alpha_i \in \mathbf{R}$. (Recall that, given points $a = (a_1, a_2), b = (b_1, b_2) \in \mathbf{R}^2$, by $\langle a, b \rangle = a_1b_1 + a_2b_2$ we denote the standard inner product in \mathbf{R}^2 .)

Thus the vector

$$\mathbf{h}_i \text{ is directed outside of } H_i \text{ and } \mathbf{h}_i \perp \ell_i, \quad i = 1, 2. \quad (4.11)$$

Proposition 4.3 Let a_1 and a_2 be two points in \mathbf{R}^2 such that

$$a_1 \in \mathcal{H}[H_1 \cap (H_2 + \delta Q_0)] \quad \text{and} \quad a_2 \in \mathcal{H}[H_2 \cap (H_1 + \delta Q_0)]. \quad (4.12)$$

Suppose that

$$\mathbf{Pr}(a_1, H_1) \in H_2 + \delta Q_0 \quad \text{and} \quad \mathbf{Pr}(a_2, H_2) \in H_1 + \delta Q_0. \quad (4.13)$$

Then the following inequality

$$\|\mathbf{Pr}(a_1, H_1) - \mathbf{Pr}(a_2, H_2)\| \leq 2\|a_1 - a_2\| + \delta \quad (4.14)$$

holds.

Proof. We will need a number of auxiliary lemmas. Let us formulate the first of them. Let

$$S_1 = H_1 \cap (H_2 + \delta Q_0) \quad \text{and} \quad S_2 = H_2 \cap (H_1 + \delta Q_0). \quad (4.15)$$

We know that $\text{dist}(H_1, H_2) \leq \delta$ so that $S_1 \neq \emptyset$ and $S_2 \neq \emptyset$. Furthermore, thanks to (4.12) and (4.15),

$$a_1 \in \mathcal{H}[S_1] \quad \text{and} \quad a_2 \in \mathcal{H}[S_2]. \quad (4.16)$$

Lemma 4.4 Both $\mathbf{Pr}(a_1, H_1)$ and $\mathbf{Pr}(a_2, H_2)$ are singletons. Furthermore, $\mathbf{Pr}(a_i, H_i) = \mathbf{Pr}(a_i, S_i)$ for every $i = 1, 2$, and the following inequality

$$|\text{dist}(a_1, H_1) - \text{dist}(a_2, H_2)| \leq \delta + \|a_1 - a_2\|$$

holds.

Proof. Thanks to (4.16), the point $a_i \in \mathcal{H}[S_i]$, so that $a_i \in \mathcal{H}[H_i]$ because $S_i \subset H_i$, $i = 1, 2$, see (4.15). Therefore, thanks to Lemma 3.4, $\mathbf{Pr}(a_i, H_i)$ is a singleton for every $i = 1, 2$.

Furthermore, thanks to (4.13), $\mathbf{Pr}(a_i, H_i) \in S_i$. But $S_i \subset H_i$ so that $\mathbf{Pr}(a_i, H_i) = \mathbf{Pr}(a_i, S_i)$, $i = 1, 2$. In particular, $\text{dist}(a_i, H_i) = \text{dist}(a_i, S_i)$, $i = 1, 2$.

Clearly,

$$d_H(S_1, S_2) = d_H(H_1 \cap [H_2 + \delta Q_0], H_2 \cap [H_1 + \delta Q_0]) \leq \delta.$$

See (2.1). Therefore,

$$|\text{dist}(a_1, H_1) - \text{dist}(a_1, H_2)| = |\text{dist}(a_1, S_1) - \text{dist}(a_1, S_2)| \leq d_H(S_1, S_2) \leq \delta.$$

Note also that the function $\text{dist}(\cdot, H_2)$ is Lipschitz. Hence, we have

$$|\text{dist}(a_1, H_1) - \text{dist}(a_2, H_2)| \leq |\text{dist}(a_1, H_1) - \text{dist}(a_1, H_2)| + |\text{dist}(a_1, H_2) - \text{dist}(a_2, H_2)| \leq \delta + \|a_1 - a_2\|$$

proving the lemma. \blacksquare

Lemma 4.5 Inequality (4.14) holds provided either $a_1 \in H_1$ or $a_2 \in H_2$.

Proof. For example, suppose that $a_2 \in H_2$. Then $\mathbf{Pr}(a_2, H_2) = a_2$ and $\text{dist}(a_2, H_2) = 0$. Therefore, thanks to Lemma 4.4, $\text{dist}(a_1, H_1) \leq \delta + \|a_1 - a_2\|$. Hence,

$$\begin{aligned} \|\mathbf{Pr}(a_1, H_1) - \mathbf{Pr}(a_2, H_2)\| &= \|\mathbf{Pr}(a_1, H_1) - a_2\| \leq \|\mathbf{Pr}(a_1, H_1) - a_1\| + \|a_1 - a_2\| \\ &= \text{dist}(a_1, H_1) + \|a_1 - a_2\| \leq \delta + 2\|a_1 - a_2\| \end{aligned}$$

proving the lemma. \blacksquare

Everywhere below, in the proof of inequality (4.14), we will assume that

$$a_1 \notin H_1 \quad \text{and} \quad a_2 \notin H_2. \quad (4.17)$$

Recall that ℓ_i is the boundary of the half-plane H_i , $i = 1, 2$. Let us see that the assumption $a_i \notin H_i$, $i = 1, 2$, implies the following property:

$$\ell_i \nparallel Ox_j \quad \text{for every } i, j = 1, 2. \quad (4.18)$$

Indeed, suppose that this statement is not true, say for $i = 1$, i.e., either $\ell_1 \parallel Ox_1$ or $\ell_1 \parallel Ox_2$. Then $\mathcal{H}[H_1] = H_1$. But $a_1 \in \mathcal{H}[H_1 \cap (H_2 + \delta Q_0)] \subset \mathcal{H}[H_1]$ so that $a_1 \in H_1$ which contradicts our assumption that $a_1 \notin H_1$.

Remark 4.6 Our next result, Lemma 4.5, deals with points a_i and half-planes H_i , $i = 1, 2$, such that the vectors

$$\mathbf{Pr}(a_1, H_1) - a_1 \quad \text{and} \quad \mathbf{Pr}(a_2, H_2) - a_2 \quad \text{are co-directed.} \quad (4.19)$$

Recall that this property means that

$$\mathbf{Pr}(a_2, H_2) - a_2 = \beta (\mathbf{Pr}(a_1, H_1) - a_1) \quad \text{for some } \beta > 0.$$

We also recall the representation of H_1 and H_2 in the form $H_i = \{u \in \mathbf{R}^2 : \langle \mathbf{h}_i, u \rangle \leq \alpha_i\}$, $i = 1, 2$, where each \mathbf{h}_i is a unit vector and $\alpha_i \in \mathbf{R}$. Thanks to (4.11), $\mathbf{h}_i \perp \ell_i (= \partial H_i)$, so that from (4.18) we have $\mathbf{h}_i \nparallel Ox_j$ for every $i, j = 1, 2$. In particular, each \mathbf{h}_i , $i = 1, 2$, has non-zero coordinates.

Finally, let us note the following useful property of metric projections in the space $\ell_\infty^2 = (\mathbf{R}^2, \|\cdot\|)$. Let $\alpha \in \mathbf{R}$ and let $\mathbf{h} = (h_1, h_2)$ with $h_1, h_2 \neq 0$ be a unit vector. Let

$$H = \{u \in \mathbf{R}^2 : \langle \mathbf{h}, u \rangle + \alpha \leq 0\}, \quad \text{and let } a \notin H.$$

Clearly, in this case $\mathbf{Pr}(a, H)$ is a singleton, and $\mathbf{Pr}(a, H) \neq a$. Then the vector

$$a - \mathbf{Pr}(a, H) \quad \text{and the vector } (\text{sign } h_1, \text{sign } h_2) \quad \text{are co-directed.} \quad \blacktriangleleft \quad (4.20)$$

Lemma 4.7 *Inequality (4.14) holds provided condition (4.19) is satisfied.*

Proof. Thanks to the triangle inequality,

$$\|\mathbf{Pr}(a_1, H_1) - \mathbf{Pr}(a_2, H_2)\| \leq \|(\mathbf{Pr}(a_1, H_1) - a_1) - (\mathbf{Pr}(a_2, H_2) - a_2)\| + \|a_1 - a_2\|. \quad (4.21)$$

Because the vectors $\mathbf{Pr}(a_1, H_1) - a_1$ and $\mathbf{Pr}(a_2, H_2) - a_2$ are co-directed,

$$\begin{aligned} \|(\mathbf{Pr}(a_1, H_1) - a_1) - (\mathbf{Pr}(a_2, H_2) - a_2)\| &= \left| \|(\mathbf{Pr}(a_1, H_1) - a_1)\| - \|(\mathbf{Pr}(a_2, H_2) - a_2)\| \right| \\ &= |\text{dist}(a_1, H_1) - \text{dist}(a_2, H_2)|. \end{aligned}$$

Therefore, thanks to Lemma 4.4,

$$\|(\mathbf{Pr}(a_1, H_1) - a_1) - (\mathbf{Pr}(a_2, H_2) - a_2)\| \leq \delta + \|a_1 - a_2\|.$$

Combining this inequality with (4.21), we obtain the required inequality (4.14). \blacksquare

Lemma 4.8 (i) *Inequality (4.14) holds provided*

$$\text{dist}(a_1, H_1) + \text{dist}(a_1, H_2) \leq \delta. \quad (4.22)$$

(ii) *Inequality (4.14) holds if $a_1 \in H_2$.*

Proof. (i) First, let us prove that

$$\|\mathbf{Pr}(a_1, H_2) - \mathbf{Pr}(a_2, H_2)\| \leq 2\|a_1 - a_2\|. \quad (4.23)$$

Indeed, thanks to (4.18), $\ell_2 \nparallel Ox_1$ and $\ell_2 \nparallel Ox_2$ so that $\mathcal{H}[H_2] = \mathbf{R}^2$. Hence, $a_1, a_2 \in \mathcal{H}[H_2]$.

We know that $a_2 \notin H_2$. If $a_1 \in H_2$, then all conditions of Lemma 4.5 are satisfied provided $H_1 = H_2$ and $\delta = 0$. This lemma tells us that in this case inequality (4.23) holds.

Now, suppose that $a_1 \notin H_2$. Then the vectors

$$\mathbf{Pr}(a_1, H_2) - a_1 \quad \text{and} \quad \mathbf{Pr}(a_2, H_2) - a_2 \quad \text{are co-directed.}$$

Therefore, all conditions of Lemma 4.7 are satisfied for the same case, i.e., for $H_1 = H_2$ and $\delta = 0$. This lemma tells us that, in these settings, inequality (4.23) holds.

Thus, we have proved (4.23) for every a_1 and a_2 satisfying (4.12) and (4.17).

From (4.23) and the triangle inequality, we have

$$\begin{aligned} \|\mathbf{Pr}(a_1, H_1) - \mathbf{Pr}(a_2, H_2)\| &\leq \|(\mathbf{Pr}(a_1, H_1) - a_1) - (\mathbf{Pr}(a_1, H_2) - a_1)\| + \|\mathbf{Pr}(a_1, H_2) - \mathbf{Pr}(a_2, H_2)\| \\ &\leq \|\mathbf{Pr}(a_1, H_1) - a_1\| + \|\mathbf{Pr}(a_1, H_2) - a_1\| + 2\|a_1 - a_2\| \\ &= \text{dist}(a_1, H_1) + \text{dist}(a_1, H_2) + 2\|a_1 - a_2\|. \end{aligned}$$

Combining this inequality with (4.22), we obtain inequality (4.14) proving part (i) of the lemma.

(ii) Prove that if $a_1 \in H_2$, then $\|\mathbf{Pr}(a_1, H_1) - \mathbf{Pr}(a_1, H_2)\| \leq \delta$. Indeed, this inequality is immediate from Lemma 4.5 applied to the case $a_1 = a_2$. Now, from this and (4.23), we have

$$\|\mathbf{Pr}(a_1, H_1) - \mathbf{Pr}(a_2, H_2)\| \leq \|\mathbf{Pr}(a_1, H_1) - \mathbf{Pr}(a_1, H_2)\| + \|\mathbf{Pr}(a_1, H_2) - \mathbf{Pr}(a_2, H_2)\| \leq \delta + 2\|a_1 - a_2\|$$

proving (4.14) and the lemma. \blacksquare

Lemma 4.9 *Inequality (4.14) holds provided $\ell_1 \parallel \ell_2$.*

Proof. Because $\ell_1 \parallel \ell_2$ and $\ell_i \nparallel Ox_j$, $i, j = 1, 2$ (see (4.18)), we have

$$\mathcal{H}[H_1 \cap (H_2 + \delta Q_0)] = \mathcal{H}[H_2 \cap (H_1 + \delta Q_0)] = \mathbf{R}^2$$

Let us note that, since $\ell_1 \parallel \ell_2$ and $\mathbf{h}_i \perp \ell_i$ (see (4.11)), the vectors \mathbf{h}_1 and \mathbf{h}_2 are collinear unit vectors. Therefore, either $\mathbf{h}_1 = \mathbf{h}_2$ or $\mathbf{h}_1 = -\mathbf{h}_2$.

If $\mathbf{h}_1 = \mathbf{h}_2$, then H_2 is a shift of H_1 , i.e., $H_2 = H_1 + p$ for some $p \in \mathbf{R}^2$.

Thanks to (4.20), in this case the vectors

$$\mathbf{Pr}(a_1, H_1) - a_1 \quad \text{and} \quad \mathbf{Pr}(a_2, H_2) - a_2 \quad \text{are co-directed.}$$

See Fig. 5-1. Therefore, thanks to Lemma 4.7, inequality (4.14) holds.

Let us prove (4.14) for $\mathbf{h}_2 = -\mathbf{h}_1$. Part (ii) of Lemma 4.8 tells us that (4.14) holds provided $a_1 \in H_2$.

Now, let us suppose that $a_1 \notin H_2$ and prove that (4.14) holds as well. In this case $a_1 \notin H_1 \cup H_2$ (because $a_1 \notin H_1$, see (4.17)) so that $H_1 \cap H_2 = \emptyset$ as it shown on Fig. 5-2.




Fig. 5: Metric projections onto the half-planes H_1 and H_2 with the parallel boundaries.

Let us prove that in this case inequality (4.22) holds. Let T be the closure of the set $\mathbf{R}^2 \setminus (H_1 \cup H_2)$. Clearly, T is the strip between the half-planes H_1 and H_2 , and $\partial T = \ell \cup \ell_2$. Recall that $\ell_1 \parallel \ell_2$ and $\text{dist}(H_1, H_2) \leq \delta$ so that

$$\text{dist}(x, H_2) \leq \delta \text{ for } x \in \ell_1 \text{ and } \text{dist}(x, H_1) \leq \delta \text{ for } x \in \ell_2.$$

We define a function f on T by letting $f(x) = \text{dist}(x, H_1) + \text{dist}(x, H_2)$. Clearly, f is a convex continuous function on T . Therefore, $\sup_T f = \sup_{\partial T} f$. But

$$f(x) = \text{dist}(x, \ell_2) \leq \delta \text{ on } \ell_1 \text{ and } f(x) = \text{dist}(x, \ell_1) \leq \delta \text{ on } \ell_2$$

so that $\sup_{\partial T} f \leq \delta$. Hence, $\sup_T f \leq \delta$ proving (4.22). Therefore, thanks to part (i) of Lemma 4.8, inequality (4.14) holds, and the proof of Lemma 4.9 is complete. ■

Thanks to Lemma 4.5, part (ii) of Lemma 4.8 and Lemma 4.9, it remains to prove that inequality (4.14) holds provided $\ell_1 \not\parallel \ell_2$,

$$a_1 \text{ and } a_2 \text{ satisfy (4.12), } a_1 \notin (H_1 \cup H_2) \text{ and } a_2 \notin H_2. \quad (4.24)$$

Clearly, without loss of generality, we may assume that $\ell_1 \cap \ell_2 = \{0\}$. Then

$$H_i = \{u \in \mathbf{R}^2 : \langle \mathbf{h}_i, u \rangle \leq 0\}, \quad i = 1, 2,$$

where $\mathbf{h}_i, i = 1, 2$, are *non-collinear* unit vectors. In these settings,

$$\ell_i = \{u \in \mathbf{R}^2 : \langle \mathbf{h}_i, u \rangle = 0\}, \quad i = 1, 2. \quad (4.25)$$

Let

$$\mathbf{h}_i = (\cos \varphi_i, \sin \varphi_i) \text{ where the angle } \varphi_i \in [0, 2\pi), \quad i = 1, 2. \quad (4.26)$$

Because the uniform norm on the plane is invariant under reflections with respect to the coordinate axes and with respect to the bisectors of the coordinate angles, we can also assume that the angles φ_1 and φ_2 satisfy the following conditions:

$$\varphi_1 \in (\pi/2, \pi) \quad \text{and} \quad \varphi_2 \in (\varphi_1, \varphi_1 + \pi).$$

We know that $\mathbf{h}_i \perp \ell_i, i = 1, 2$ so that $\mathbf{h}_1 \nparallel \mathbf{h}_2$ (because $\ell_1 \nparallel \ell_2$). Let us also recall that \mathbf{h}_i is directed outside of $H_i, i = 1, 2$. We also note that, in the case under consideration, $H_1 \cap H_2$ is a convex cone with the vertex at 0. Moreover, the sets

$$S_1 = H_1 \cap (H_2 + \delta Q_0) \quad \text{and} \quad S_2 = H_2 \cap (H_1 + \delta Q_0)$$

are convex cones in \mathbf{R}^2 . Let

$$X_1 = (s_1, s_2) \quad \text{and} \quad X_2 = (t_1, t_2) \tag{4.27}$$

be the vertices of the cones S_1 and S_2 respectively.

Thus, X_1 is the point of intersection of the line $\ell_1 = \partial H_1$ and the line $\tilde{\ell}_2 = \partial(H_2 + \delta Q_0)$. In turn,

$$X_2 = \ell_2 \cap \tilde{\ell}_1 \quad \text{where} \quad \ell_2 = \partial H_2 \quad \text{and} \quad \tilde{\ell}_1 = \partial(H_1 + \delta Q_0).$$

Moreover, thanks to (4.15), we have the following representations of the cones S_1 and S_2 :

$$S_1 = H_1 \cap H_2 + X_1, \quad S_2 = H_1 \cap H_2 + X_2. \tag{4.28}$$

Let us give explicit formulae for the points X_1 and X_2 . First, we note that

$$H_i + \delta Q_0 = \{u \in \mathbf{R}^2 : \langle \mathbf{h}_i, u \rangle \leq \delta \|\mathbf{h}_i\|_1\}, \quad i = 1, 2.$$

Here, given $u = (u_1, u_2) \in \mathbf{R}^2$ we let $\|u\|_1 = |u_1| + |u_2|$ denote the ℓ_1 -norm in \mathbf{R}^2 . Hence,

$$\tilde{\ell}_i = \{u \in \mathbf{R}^2 : \langle \mathbf{h}_i, u \rangle = \delta \|\mathbf{h}_i\|_1\}, \quad i = 1, 2. \tag{4.29}$$

Let

$$A = \begin{pmatrix} \cos \varphi_1 & \sin \varphi_1 \\ \cos \varphi_2 & \sin \varphi_2 \end{pmatrix} \quad \text{and let} \quad \Delta = \det A = \sin(\varphi_2 - \varphi_1). \tag{4.30}$$

See (4.26). (Clearly, $\Delta \neq 0$ because $\mathbf{h}_1 \nparallel \mathbf{h}_2$.) We know that

$$X_1 = (s_1, s_2) = \ell_1 \cap \tilde{\ell}_2$$

so that, thanks to (4.25) and (4.29), the vector (s_1, s_2) is the solution of the system of linear equations

$$A \begin{pmatrix} s_1 \\ s_2 \end{pmatrix} = \begin{pmatrix} 0 \\ \delta \|\mathbf{h}_2\|_1 \end{pmatrix}.$$

Therefore,

$$s_1 = \frac{1}{\Delta} \begin{vmatrix} 0 & \sin \varphi_1 \\ \delta \|\mathbf{h}_2\|_1 & \sin \varphi_2 \end{vmatrix} = -\frac{\delta}{\Delta} \|\mathbf{h}_2\|_1 \sin \varphi_1 \quad \text{and} \quad s_2 = \frac{\delta}{\Delta} \|\mathbf{h}_2\|_1 \cos \varphi_1. \tag{4.31}$$

Thus,

$$X_1 = \frac{\delta}{\Delta} \|\mathbf{h}_2\|_1 (-\sin \varphi_1, \cos \varphi_1). \tag{4.32}$$

In the same way we prove that

$$X_2 = \frac{\delta}{\Delta} \|\mathbf{h}_1\|_1 (\sin \varphi_2, -\cos \varphi_2). \tag{4.33}$$

Lemma 4.10 *Inequality (4.14) holds provided $\varphi_1 \in (\pi/2, \pi)$ and $\varphi_2 \in (\varphi_1, \pi)$.*

Proof. We recall that in the case under consideration

$$\mathbf{h}_i = (\cos \varphi_i, \sin \varphi_i) \quad \text{where } \varphi_i \in (\pi/2, \pi), \quad i = 1, 2.$$

Hence, $(\text{sign}(\cos \varphi_i), \text{sign}(\sin \varphi_i)) = (-1, 1)$.

This equality and property (4.20) imply the following: for every $i = 1, 2$, the vector $(1, -1)$ and the vector $\Pr(a_i, H) - a_i$ are co-directed. See Fig. 6.




Fig. 6: The half-planes H_1 and H_2 with the non-parallel boundaries: the first case.

This proves that

$$\Pr(a_1, H) - a_1 \quad \text{and} \quad \Pr(a_2, H) - a_2 \quad \text{are co-directed vectors.}$$

Therefore, thanks to Lemma 4.7, (4.14) holds, and the proof of the lemma is complete. ■

Lemma 4.11 *Inequality (4.14) holds provided $\varphi_1 \in (\pi/2, \pi)$ and $\varphi_2 \in (\pi, \varphi_1 + \pi)$.*

Proof. First, let us prove the following important inequality: Let $T = \Delta(X_1, X_2, O)$ be the triangle with vertices at X_1, X_2 and $O = 0$. Then

$$D(x) \leq \delta \quad \text{where} \quad D(x) = \text{dist}(x, H_1) + \text{dist}(x, H_2), \quad x \in T. \quad (4.34)$$

Indeed, D is a convex continuous function on T so that its maximum is attained on the set of vertices of the triangle T , i.e., at the points O, X_1 and X_2 . But $D(O) = 0$, $D(X_1) = \delta$ (because $X_1 \in \tilde{\ell}_2 = \partial(H_2 + \delta Q_0)$) and $D(X_2) = \delta$ (because $X_2 \in \tilde{\ell}_1 = \partial(H_1 + \delta Q_0)$) proving the required inequality (4.34).

We begin the proof of inequality (4.14) with the case $\varphi_1 \in (\pi/2, \pi)$ and $\varphi_2 \in (\pi, \frac{3}{2}\pi)$. See Fig. 7.




Fig. 7: The half-planes H_1 and H_2 with the non-parallel boundaries: $\varphi_2 \in (\pi, \frac{3}{2}\pi)$.

In this case, the following inequalities hold:

$$\cos \varphi_1 < 0, \quad \sin \varphi_1 > 0, \quad \cos \varphi_2 < 0 \quad \text{and} \quad \sin \varphi_2 < 0. \quad (4.35)$$

Therefore, thanks to (4.20), the vector $\mathbf{Pr}(a_1, H_1) - a_1$ is co-directed with the vector $(1, -1)$, and the vector $\mathbf{Pr}(a_2, H_2) - a_2$ is co-directed with $(1, 1)$. Moreover, in this case the convex cone $H_1 \cap H_2$ (with the vertex in 0) contains the positive semi-axis $Ox_1^+ (= \{(t, 0) : t \geq 0\})$. This implies the following properties of the rectangular hulls of the sets $S_1 = H_1 \cap (H_2 + \delta Q_0)$ and $S_2 = H_2 \cap (H_1 + \delta Q_0)$:

$$\mathcal{H}[S_1] = \{(u_1, u_2) \in \mathbf{R}^2 : u_1 \geq s_1\} \quad \text{and} \quad \mathcal{H}[S_2] = \{(u_1, u_2) \in \mathbf{R}^2 : u_1 \geq t_1\}.$$

We recall that $X_1 = (s_1, s_2)$ and $X_2 = (t_1, t_2)$. See (4.27) and Fig. 7.

Let $\alpha_i \in (0, \pi/2)$, $i = 1, 2$, be the angle between the straight line $\ell_i = \partial H_i$ and the axis Ox_1 . On Fig. 7 we consider the case $\alpha_2 \leq \alpha_1$. We note that no necessity in the additional consideration of the case $\alpha_2 > \alpha_1$ because it can be obtained from the case $\alpha_2 < \alpha_1$ with the help of a suitable reflections with respect to the coordinate axes and the bisectors of the coordinate angles.

Let us prove that if $\alpha_2 \leq \alpha_1$, then

$$t_1 \leq s_1 \leq 0. \quad (4.36)$$

First, let us note that, thanks to (4.30), $\Delta = \sin(\varphi_2 - \varphi_1) > 0$ (because $\varphi_2 - \varphi_1 \in (0, \pi)$.) Also note that, thanks to (4.35), $\sin \varphi_1 > 0$. Hence, thanks to formula (4.32), $s_1 \leq 0$.

Let us see that

$$s_1 - t_1 = \frac{\delta}{\Delta} \sin(\varphi_1 + \varphi_2) = \frac{\delta}{\Delta} \sin(\alpha_1 - \alpha_2). \quad (4.37)$$

Indeed, thanks to (4.35),

$$\|\mathbf{h}_1\|_1 = -\cos \varphi_1 + \sin \varphi_1 \quad \text{and} \quad \|\mathbf{h}_2\|_1 = -\cos \varphi_2 - \sin \varphi_2,$$

so that, thanks to formulae (4.32) and (4.33),

$$\begin{aligned} s_1 - t_1 &= (\delta/\Delta)\{-\|\mathbf{h}_2\|_1 \sin \varphi_1 - \|\mathbf{h}_1\|_1 \sin \varphi_2\} \\ &= (\delta/\Delta)\{-(-\cos \varphi_2 - \sin \varphi_2) \sin \varphi_1 - (-\cos \varphi_1 + \sin \varphi_1) \sin \varphi_2\} \\ &= (\delta/\Delta)\{\cos \varphi_2 \sin \varphi_1 + \cos \varphi_1 \sin \varphi_2\} = (\delta/\Delta) \sin(\varphi_1 + \varphi_2). \end{aligned}$$

We note that $\alpha_1 = \varphi_1 - \pi/2$ and $\alpha_2 = \frac{3}{2}\pi - \varphi_2$. Hence, we have

$$\varphi_1 + \varphi_2 = \alpha_1 - \alpha_2 + 2\pi$$

proving (4.37). It remains to note that $\sin(\alpha_1 - \alpha_2) > 0$ (because $0 < \alpha_2 \leq \alpha_1 < \pi/2$) and $\Delta > 0$ so that, thanks (4.37), $t_1 \leq s_1$ proving (4.36).

In particular, this inequality implies the inclusion $\mathcal{H}[S_1] \subset \mathcal{H}[S_2]$ as it shown on Fig. 7. Furthermore, thanks to (4.24)

$$a_1 \notin H_1 \cup H_2 \quad \text{and} \quad a_1 \in \mathcal{H}[S_1] = \mathcal{H}[H_1 \cap (H_2 + \delta Q_0)]. \quad (4.38)$$

Let Y be the point of intersection of the line ℓ_2 (the boundary of H_2) and the line $\hat{\ell}$ passing through the point $X_1 = (s_1, s_2)$ and parallel to the axis Ox_2 . (Thus, $Y = (s_1, y_2)$ for some $y_2 \in \mathbf{R}$). Recall that the point $X_2 = (t_1, t_2)$ lies on the line ℓ_2 , and, thanks to inequality (4.36),

$$t_1 \leq s_1 \leq 0.$$

In particular, these observations shows that $Y \in [X_2, O]$ where $O = 0$ is the origin.

Conditions (4.38) shows that the point a_1 belongs to the triangle $\tilde{T} = \Delta(X_1, Y, O)$ with the vertices at the points X_1, Y and O . Because $Y \in [X_2, O]$, the triangle \tilde{T} is a subset of the triangle $T = \Delta(X_1, X_2, O)$ with the vertices at X_1, X_2 and $O = 0$. Hence, $a_1 \in T$.

Therefore, thanks to (4.34),

$$D(a_1) = \text{dist}(a_1, H_1) + \text{dist}(a_1, H_2) \leq \delta. \quad (4.39)$$

Thus, condition (4.22) of Lemma 4.8 holds. Thanks to this lemma, inequality (4.14) holds proving Lemma 4.11 for the angles

$$\varphi_1 \in (\pi/2, \pi) \quad \text{and} \quad \varphi_2 \in \left(\pi, \frac{3}{2}\pi\right).$$

Let us prove Lemma 4.11 for

$$\varphi_1 \in (\pi/2, \pi) \quad \text{and} \quad \varphi_2 \in \left(\frac{3}{2}\pi, \varphi_1 + \pi\right). \quad \text{See Fig. 8.} \quad (4.40)$$




Fig. 8: The half-planes H_1 and H_2 with the non-parallel boundaries: $\varphi_2 \in \left(\frac{3}{2}\pi, \varphi_1 + \pi\right)$.

In this case,

$$\cos \varphi_1 < 0, \quad \sin \varphi_1 > 0, \quad \cos \varphi_2 > 0 \quad \text{and} \quad \sin \varphi_2 < 0. \quad (4.41)$$

Therefore, thanks to (4.20), the vector $\mathbf{Pr}(a_1, H_1) - a_1$ is co-directed with $(1, -1)$, and the vector $\mathbf{Pr}(a_2, H_2) - a_2$ is co-directed with $(-1, 1)$. Moreover,

$$\mathcal{H}[S_1] = \{(u_1, u_2) \in \mathbf{R}^2 : u_1 \geq s_1, u_2 \geq s_2\} \quad \text{and} \quad \mathcal{H}[S_2] = \{(u_1, u_2) \in \mathbf{R}^2 : u_1 \geq t_1, u_2 \geq t_2\}.$$

Let us prove that

$$X_1 - X_2 = \delta(1, -1). \quad (4.42)$$

(Recall that $X_1 = (s_1, s_2)$ and $X_2 = (t_1, t_2)$ are the points defined by (4.27).) For explicit formulae for $s_i, t_i, i = 1, 2$, see (4.32) and (4.33).

Thanks to (4.41), we have

$$\|\mathbf{h}_1\|_1 = |\cos \varphi_1| + |\sin \varphi_1| = -\cos \varphi_1 + \sin \varphi_1, \quad \|\mathbf{h}_2\|_1 = |\cos \varphi_2| + |\sin \varphi_2| = \cos \varphi_2 - \sin \varphi_2.$$

Therefore, thanks to (4.32) and (4.33),

$$X_1 = \frac{\delta}{\Delta} (\cos \varphi_2 - \sin \varphi_2) (-\sin \varphi_1, \cos \varphi_1) \quad \text{and} \quad X_2 = \frac{\delta}{\Delta} (-\cos \varphi_1 + \sin \varphi_1) (\sin \varphi_2, -\cos \varphi_2).$$

Hence,

$$\begin{aligned} X_1 - X_2 &= \frac{\delta}{\Delta} ((\cos \varphi_2 - \sin \varphi_2) (-\sin \varphi_1) - (-\cos \varphi_1 + \sin \varphi_1) \sin \varphi_2, \\ &\quad (\cos \varphi_2 - \sin \varphi_2) \cos \varphi_1 - (-\cos \varphi_1 + \sin \varphi_1) (-\cos \varphi_2)) \\ &= \frac{\delta}{\Delta} (-\cos \varphi_2 \sin \varphi_1 + \cos \varphi_1 \sin \varphi_2, -\sin \varphi_2 \cos \varphi_1 + \sin \varphi_1 \cos \varphi_2) \\ &= \frac{\delta}{\Delta} \sin(\varphi_2 - \varphi_1)(1, -1). \end{aligned}$$

Thanks to (4.30), $\Delta = \sin(\varphi_2 - \varphi_1)$, and the proof of (4.42) is complete.

Thanks to this equality, $t_1 = s_1 - \delta$ and $t_2 = s_2 + \delta$. Furthermore, (4.42) and (4.28) imply the following:

$$S_2 = S_1 + \delta(-1, 1) \quad \text{and} \quad \mathcal{H}[S_2] = \mathcal{H}[S_1] + \delta(-1, 1).$$

Let us prove that

$$t_1 \leq s_1 \leq 0 \quad \text{and} \quad s_2 \leq t_2 \leq 0. \quad (4.43)$$

In fact, we know that

$$\varphi_1 \in (\pi/2, \pi) \quad \text{and} \quad \varphi_2 \in ((3/2)\pi, \varphi_1 + \pi).$$

Hence, $0 < \varphi_2 - \varphi_1 < \pi$ proving that $\Delta = \sin(\varphi_2 - \varphi_1) > 0$. We also know that $\sin \varphi_1 > 0$ so that, thanks to (4.31), $s_1 \leq 0$. In addition, $t_1 = s_1 - \delta \leq s_1$ proving the first inequality in (4.43).

Next, thanks to (4.33),

$$t_2 = -(\delta/\Delta) \|\mathbf{h}_1\|_1 \cos \varphi_2.$$

But $\Delta > 0$ and $\cos \varphi_2 > 0$ so that $t_2 \leq 0$. Moreover, $s_2 = t_2 - \delta$, so that $s_2 \leq t_2$, and the proof of (4.43) is complete.

We recall that the point a_1 satisfies conditions (4.38). Therefore,

$$a_1 \in T = \Delta(X_1, X_2, O).$$

See Fig. 8. From this and (4.39) it follows that condition (4.22) of Lemma 4.8 is satisfied. This lemma tells us that inequality (4.14) holds proving Lemma 4.11 for the case (4.40).

The proof of Lemma 4.11 is complete. ■

Finally, the results of Lemmas 4.4 – 4.11 imply the required inequality (4.14) completing the proof of Proposition 4.3. ■

We are in a position to complete the proof of Proposition 3.3.

PROOF OF INEQUALITY (3.18). Let us fix elements $x, y \in \mathcal{M}$. We set

$$A_1 = F^{[1]}[x : \tilde{\lambda}], \quad A_2 = F^{[1]}[y : \tilde{\lambda}]$$

and $a_1 = g(x)$, $a_2 = g(y)$. Recall that $g : \mathcal{M} \rightarrow \mathbf{R}$ is the mapping satisfying (3.14) and (3.15).

We also recall that the mapping $F^{[1]}[\cdot : \tilde{\lambda}]$ is defined by (3.2). Thus,

$$A_i = \cap \{A_i^{[u]} : u \in \mathcal{M}\}, \quad i = 1, 2, \quad (4.44)$$

where given $u \in \mathcal{M}$, we set

$$A_1^{[u]} = F(u) + \tilde{\lambda} \rho(u, x) Q_0 \quad \text{and} \quad A_2^{[u]} = F(u) + \tilde{\lambda} \rho(u, y) Q_0. \quad (4.45)$$

Lemma 3.1 tells us that each A_i , $i = 1, 2$, is a nonempty closed convex subset of \mathbf{R}^2 .

Thanks to inequality (3.15), we have

$$\|a_1 - a_2\| \leq \lambda \rho(x, y), \quad (4.46)$$

and, thanks to (3.14),

$$a_i \in \mathcal{H}[A_i], \quad i = 1, 2. \quad (4.47)$$

Furthermore, formula (3.16) tells us that

$$f(x) = \mathbf{Pr}(a_1, A_1) \quad \text{and} \quad f(y) = \mathbf{Pr}(a_2, A_2).$$

(We also recall that, thanks to Lemma 3.4, the metric projection $\mathbf{Pr}(a_i, A_i)$ is well defined, i.e., $\mathbf{Pr}(a_i, A_i)$ is a singleton.)

In these settings, the required inequality (3.18) reads as follows:

$$\|\mathbf{Pr}(a_1, A_1) - \mathbf{Pr}(a_2, A_2)\| \leq (2\lambda + \tilde{\lambda})\rho(x, y). \quad (4.48)$$

Let us note that this inequality is immediate from (4.46) provided $a_i \in A_i$, $i = 1, 2$, (because in this case $\mathbf{Pr}(a_i, A_i) = a_i$).

Suppose that either $a_1 \notin A_1$ or $a_2 \notin A_2$. Without loss of generality, we may assume that $a_1 \notin A_1$. Fix $\varepsilon > 0$ and prove that there exists a half-plane $H_1 \in \mathcal{HP}(\mathbf{R}^2)$ such that

$$H_1 \supset A_1, \quad H_1 + \tilde{\lambda}\rho(x, y)Q_0 \supset A_2, \quad (4.49)$$

and

$$\|\mathbf{Pr}(a_1, A_1) - \mathbf{Pr}(a_1, H_1)\| < \varepsilon. \quad (4.50)$$

We construct the half-plane H_1 as follows: Because $a_1 \notin A_1$, we have $\mathbf{Pr}(a_1, A_1) \neq a_1$ so that $(\mathbf{Pr}(a_1, A_1), a_1]$ is a nonempty semi-open interval in \mathbf{R}^2 . Let us pick a point

$$a^{(\varepsilon)} \in (\mathbf{Pr}(a_1, A_1), a_1]$$

such that

$$\|a^{(\varepsilon)} - \mathbf{Pr}(a_1, A_1)\| < \varepsilon. \quad (4.51)$$

Because $\mathbf{Pr}(a_1, A_1)$ is the nearest to a_1 point on A_1 , we have

$$(\mathbf{Pr}(a_1, A_1), a_1] \cap A_1 = \emptyset.$$

Therefore,

$$a^{(\varepsilon)} \notin A_1 = \bigcap \{A_1^{[u]} : u \in \mathcal{M}\}.$$

See (4.44). This implies the existence of an element $u \in \mathcal{M}$ such that $a^{(\varepsilon)} \notin A_1^{[u]}$. We let B denote the set $A_1^{[u]}$. Thus,

$$a^{(\varepsilon)} \notin B = A_1^{[u]} = F(u) + \tilde{\lambda}\rho(u, x)Q_0. \quad (4.52)$$

See (4.45). Thanks to (4.47) and (4.44),

$$a_1 \in \mathcal{H}[A_1] \quad \text{and} \quad A_1 \subset B. \quad (4.53)$$

Therefore, thanks to Lemma 4.1, the metric projections $\mathbf{Pr}(a_1, A_1)$ and $\mathbf{Pr}(a_1, B)$ are singletons such that

$$\mathbf{Pr}(a_1, B) \in [\mathbf{Pr}(a_1, A_1), a_1].$$

See Fig. 9.




Fig. 9: Metric projections of a_1 onto A_1 and B .

We note that $\mathbf{Pr}(a_1, B) \in [\mathbf{Pr}(a_1, A_1), a^{(\epsilon)}]$; indeed, otherwise $a^{(\epsilon)} \in [\mathbf{Pr}(a_1, A_1), \mathbf{Pr}(a_1, B)] \subset B$, a contradiction. See (4.52). Hence, thanks to (4.51),

$$\|\mathbf{Pr}(a_1, A_1) - \mathbf{Pr}(a_1, B)\| \leq \|a^{(\epsilon)} - \mathbf{Pr}(a_1, A_1)\| < \varepsilon. \quad (4.54)$$

Let $\tilde{Q} = Q(a_1, r)$ where $r = \text{dist}(a_1, B)$. Thus, $\tilde{Q} \cap B = \{\mathbf{Pr}(a_1, B)\}$. Therefore, thanks to the separation theorem, there exists a half-plane $H_1 \in \mathcal{HP}(\mathbf{R}^2)$ which contains B and separates (not strictly) \tilde{Q} and B . Thus, $B \subset H_1$ and $\tilde{Q} \cap H_1 = \mathbf{Pr}(a_1, B)$ as it shown on Fig. 9. In particular, these properties imply the equality $\mathbf{Pr}(a_1, H_1) = \mathbf{Pr}(a_1, B)$.

Let us see that inclusions (4.49) and inequality (4.50) hold for the half-plane H_1 . In fact, (4.50) is immediate from (4.54) and the last equality.

Prove (4.49). We know that $A_1 \subset B$, see (4.53), so that $A_1 \subset B \subset H_1$. We also recall that $B = F(u) + \tilde{\lambda}\rho(u, x)Q_0$ (see (4.52)). Therefore,

$$H_1 + \tilde{\lambda}\rho(x, y)Q_0 \supset B + \tilde{\lambda}\rho(x, y)Q_0 = F(u) + \tilde{\lambda}\rho(u, x)Q_0 + \tilde{\lambda}\rho(x, y)Q_0 = F(u) + \tilde{\lambda}(\rho(u, x) + \rho(x, y))Q_0.$$

Therefore, thanks to the triangle inequality, (4.44) and (4.45), we have

$$H_1 + \tilde{\lambda}\rho(x, y)Q_0 \supset F(u) + \tilde{\lambda}\rho(u, y)Q_0 = A_2^{[u]} \supset A_2$$

proving (4.49).

Next, let us construct a half-plane $H_2 \in \mathcal{HP}(\mathbf{R}^2)$ having the following properties:

$$H_2 \supset A_2, \quad H_2 + \tilde{\lambda}\rho(x, y)Q_0 \supset A_1, \quad (4.55)$$

and

$$\|\mathbf{Pr}(a_2, A_2) - \mathbf{Pr}(a_2, H_2)\| < \varepsilon. \quad (4.56)$$

If $a_2 \notin A_2$, we define H_2 in the same way as we have defined H_1 for a_1 . In this case, properties (4.55), (4.56) are a complete analog of properties (4.49) and (4.50) obtained for the point a_1 .

If $a_2 \in A_2$, we set

$$H_2 = H_1 + \tilde{\lambda}\rho(x, y)Q_0.$$

Clearly, H_2 is a half-plane. Let us see that inclusions (4.55) and inequality (4.56) hold for this choice of H_2 . Indeed, thanks to the second inclusion in (4.49), we have $H_2 \supset A_2$. In turn, thanks to the first inclusion,

$$H_2 + \tilde{\lambda}\rho(x, y)Q_0 = (H_1 + \tilde{\lambda}\rho(x, y)Q_0) + \tilde{\lambda}\rho(x, y)Q_0 \supset H_1 \supset A_1,$$

proving (4.55). Finally, inequality (4.56) is trivial because $\mathbf{Pr}(a_2, A_2) = \mathbf{Pr}(a_2, H_2)(= a_2)$. (Recall that $a_2 \in A_2 \subset H_2$.)

Now, we set

$$\delta = \tilde{\lambda}\rho(x, y) + \varepsilon. \quad (4.57)$$

Let us prove that the points a_1, a_2 and the half-planes H_1 and H_2 satisfy conditions (4.10), (4.12) and (4.13).

Thanks to (4.49), $H_1 \supset A_1$, and, thanks to (4.55), $H_2 + \tilde{\lambda}\rho(x, y)Q_0 \supset A_1$. Hence,

$$H_1 \cap (H_2 + \tilde{\lambda}\rho(x, y)Q_0) \supset A_1.$$

Note that $\tilde{\lambda}\rho(x, y) < \delta$; see (4.57). Therefore,

$$H_1 \cap (H_2 + \delta Q_0) \supset A_1. \quad (4.58)$$

We also know that $A_1 \neq \emptyset$ so that $H_1 \cap (H_2 + \delta Q_0) \neq \emptyset$ as well. This proves that $\text{dist}(H_1, H_2) \leq \delta$ so that the condition (4.10) is satisfied.

Let us prove that the points a_1 and a_2 satisfy condition (4.12). Indeed, inclusion (4.58) tells us that

$$\mathcal{H}[A_1] \subset \mathcal{H}[H_1 \cap (H_2 + \delta Q_0)].$$

But, thanks to (4.47), $a_1 \in \mathcal{H}[A_1]$ proving that $a_1 \in \mathcal{H}[H_1 \cap (H_2 + \delta Q_0)]$.

In the same fashion we show that $a_2 \in \mathcal{H}[H_2 \cap (H_1 + \delta Q_0)]$ proving that condition (4.12) holds.

Let us show that condition (4.13) is satisfied as well. Thanks to (4.55),

$$\mathbf{Pr}(a_1, A_1) \in A_1 \subset H_2 + \tilde{\lambda}\rho(x, y)Q_0,$$

and, thanks to (4.50),

$$\mathbf{Pr}(a_1, H_1) \in \mathbf{Pr}(a_1, A_1) + \varepsilon Q_0 \subset A_1 + \varepsilon Q_0.$$

Therefore, thanks to (4.57),

$$\mathbf{Pr}(a_1, H_1) \in A_1 + \varepsilon Q_0 \subset (H_2 + \tilde{\lambda}\rho(x, y)Q_0) + \varepsilon Q_0 = H_2 + \delta Q_0.$$

In the same way we show that $\mathbf{Pr}(a_2, H_2) \in H_1 + \delta Q_0$ completing the proof of (4.13).

Therefore, thanks to Proposition 4.3, inequality (4.14) holds. This inequality together with (4.46) and (4.57) imply the following:

$$\|\mathbf{Pr}(a_1, H_1) - \mathbf{Pr}(a_2, H_2)\| \leq 2\|a_1 - a_2\| + \delta \leq 2\lambda\rho(x, y) + \tilde{\lambda}\rho(x, y) + \varepsilon.$$

From this, (4.50) and (4.56), we have

$$\begin{aligned} \|\Pr(a_1, A_1) - \Pr(a_2, A_2)\| &\leq \|\Pr(a_1, A_1) - \Pr(a_1, H_1)\| + \|\Pr(a_1, H_1) - \Pr(a_2, H_2)\| \\ &\quad + \|\Pr(a_2, H_2) - \Pr(a_2, A_2)\| \leq (2\lambda + \tilde{\lambda})\rho(x, y) + 3\varepsilon. \end{aligned}$$

Since $\varepsilon > 0$ is arbitrary, this implies (4.48) proving the required inequality (3.18) and completing the proof of Proposition 3.3. ■

Finally, combining part (i) of Proposition 3.2 with Proposition 3.3, we conclude that the mapping f defined by formula (3.16), is a Lipschitz selection of the set-valued mapping F with Lipschitz seminorm at most $2\lambda + \tilde{\lambda}$.

The proof of Theorem 1.10 is complete. ■

5. The $\vec{\lambda}$ -Projection Algorithm and its versions.

5.1 Main properties of the Projection Algorithm: proofs.

PROOF OF THEOREM 1.9. (i) Let us assume that the $\vec{\lambda}$ -PA produces the outcome “**No go**” and prove that there does not exist a Lipschitz selection of F with Lipschitz seminorm at most $\lambda^{\min} = \min\{\lambda_1, \lambda_2\}$. We recall that, thanks to (1.9), in this case there exists a point $\tilde{x} \in \mathcal{M}$ such that

$$\text{either } F^{[1]}[\tilde{x} : \lambda_1] = \emptyset \text{ or } \mathcal{T}_{F, \lambda_1}^{[1]}[\tilde{x} : \lambda_2] = \emptyset.$$

Let us suppose that F has a Lipschitz selection $f : \mathcal{M} \rightarrow \mathbf{R}^2$ with $\|f\|_{\text{Lip}(\mathcal{M})} \leq \lambda^{\min}$, and prove that $F^{[1]}[\tilde{x} : \lambda_1] \neq \emptyset$ and $\mathcal{T}_{F, \lambda_1}^{[1]}[\tilde{x} : \lambda_2] \neq \emptyset$.

Indeed, we know that $f(x) \in F(x)$ and

$$\|f(x) - f(y)\| \leq \lambda^{\min} \rho(x, y) \leq \lambda_1 \rho(x, y) \text{ for all } x, y \in \mathcal{M}.$$

Therefore, $f(x) \in F(y) + \lambda_1 \rho(x, y)Q_0$ for all $x, y \in \mathcal{M}$ so that $f(x) \in F^{[1]}[x : \lambda_1]$ for every $x \in \mathcal{M}$. See definition (1.3). In particular, $f(\tilde{x}) \in F^{[1]}[\tilde{x} : \lambda_1]$ proving that $F^{[1]}[\tilde{x} : \lambda_1] \neq \emptyset$.

Let us show that $\mathcal{T}_{F, \lambda_1}^{[1]}[\tilde{x} : \lambda_2] \neq \emptyset$ as well. Indeed,

$$f(x) \in F^{[1]}[x : \lambda_1] \subset \mathcal{H}[F^{[1]}[x : \lambda_1]] = \mathcal{T}_{F, \lambda_1}(x) \text{ for every } x \in \mathcal{M}.$$

See (1.5). Therefore, $f(\tilde{x}) \in \mathcal{T}_{F, \lambda_1}(\tilde{x})$, $f(y) \in \mathcal{T}_{F, \lambda_1}(y)$ and

$$\|f(\tilde{x}) - f(y)\| \leq \lambda^{\min} \rho(\tilde{x}, y) \leq \lambda_2 \rho(\tilde{x}, y) \text{ for every } y \in \mathcal{M}.$$

Hence,

$$f(\tilde{x}) \in \mathcal{T}_{F, \lambda_1}(y) + \lambda_2 \rho(\tilde{x}, y)Q_0 \text{ for every } y \in \mathcal{M},$$

so that, thanks to (1.6), $f(\tilde{x}) \in \mathcal{T}_{F, \lambda_1}^{[1]}[\tilde{x} : \lambda_2]$. Hence, $\mathcal{T}_{F, \lambda_1}^{[1]}[\tilde{x} : \lambda_2] \neq \emptyset$, a contradiction.

(ii) Let us assume that the (λ_1, λ_2) -PA produces the outcome “**Success**”. Then, thanks to (1.10), for every $x \in \mathcal{M}$, we have

$$F^{[1]}[x : \lambda_1] \neq \emptyset \text{ and } \mathcal{T}_{F, \lambda_1}^{[1]}[x : \lambda_2] \neq \emptyset. \quad (5.1)$$

Let us prove property (★A). Clearly, the part (a) of this property is immediate from (5.1). Prove part (b). Let us note that for every $x \in \mathcal{M}$, the metric projection of the origin O onto the rectangle

$\mathcal{T}_{F,\lambda_1}^{[1]}[x : \lambda_2]$, is either a singleton or a compact line segment parallel to one of the coordinate axes. Therefore, the point $g_F(x)$, the center of the line segment $\mathbf{Pr}(O, \mathcal{T}_{F,\lambda_1}^{[1]}[x : \lambda_2])$, see (1.7), is well defined.

Let us prove part (c). Thanks to (1.5), (1.6) and (1.7),

$$g_F(x) \in \mathcal{T}_{F,\lambda_1}^{[1]}[x : \lambda_2] \subset \mathcal{T}_{F,\lambda_1}(x) = \mathcal{H}[F^{[1]}[x : \lambda_1]]. \quad (5.2)$$

Therefore, thanks to Lemma 3.4 and definition (1.7), $f_{\tilde{\lambda},F}(x)$, the metric projection of $g_F(x)$ onto $F^{[1]}[x : \lambda_1]$, see (1.8), is a singleton, and the proof of property ($\star A$) is complete.

Let us prove property ($\star B$) of the theorem. We will follow the scheme of the proof of the key Theorem 1.10 for the special case

$$\tilde{\lambda} = \lambda_1 \quad \text{and} \quad \lambda = \lambda_2. \quad (5.3)$$

Thanks to (5.1), $\mathcal{T}_{F,\tilde{\lambda}}^{[1]}[x : \lambda] \neq \emptyset$ for every $x \in \mathcal{M}$. Part (a) of Proposition 2.8 tells us that in this case the mapping $\mathcal{T}_{F,\tilde{\lambda}}^{[1]}[\cdot : \lambda]$ is Lipschitz with respect to the Hausdorff distance. See (2.35). Thus,

$$d_H(\mathcal{T}_{F,\tilde{\lambda}}^{[1]}[x : \lambda], \mathcal{T}_{F,\tilde{\lambda}}^{[1]}[y : \lambda]) \leq \lambda \rho(x, y) \quad \text{for all } x, y \in \mathcal{M}. \quad (5.4)$$

Let $\delta(x) = \text{dist}(O, \mathcal{T}_{F,\tilde{\lambda}}^{[1]}[x : \lambda])$. (Recall that $O = (0, 0)$ is the origin.) It is known that the distance from a fixed point to a set is a Lipschitz function of the set (with respect to the Hausdorff distance). Therefore,

$$|\delta(x) - \delta(y)| = |\text{dist}(O, \mathcal{T}_{F,\tilde{\lambda}}^{[1]}[x : \lambda]) - \text{dist}(O, \mathcal{T}_{F,\tilde{\lambda}}^{[1]}[y : \lambda])| \leq d_H(\mathcal{T}_{F,\tilde{\lambda}}^{[1]}[x : \lambda], \mathcal{T}_{F,\tilde{\lambda}}^{[1]}[y : \lambda])$$

so that, thanks to (5.4),

$$|\delta(x) - \delta(y)| \leq \lambda \rho(x, y) \quad \text{for every } x, y \in \mathcal{M}. \quad (5.5)$$

We note that

$$\mathbf{Pr}(O, \mathcal{T}_{F,\tilde{\lambda}}^{[1]}[x : \lambda]) = Q(O, \delta(x)) \cap \mathcal{T}_{F,\tilde{\lambda}}^{[1]}[x : \lambda].$$

From this and Lemma 2.6, we have

$$\begin{aligned} \mathbf{Pr}(O, \mathcal{T}_{F,\tilde{\lambda}}^{[1]}[x : \lambda]) + \lambda \rho(x, y) Q_0 &= \{Q(O, \delta(x)) + \lambda \rho(x, y) Q_0\} \cap \{\mathcal{T}_{F,\tilde{\lambda}}^{[1]}[x : \lambda] + \lambda \rho(x, y) Q_0\} \\ &= Q(O, \delta(x) + \lambda \rho(x, y)) \cap \{\mathcal{T}_{F,\tilde{\lambda}}^{[1]}[x : \lambda] + \lambda \rho(x, y) Q_0\}. \end{aligned}$$

Note that, thanks to (5.5), $\delta(y) \leq \delta(x) + \lambda \rho(x, y)$, and, thanks to (5.4),

$$\mathcal{T}_{F,\tilde{\lambda}}^{[1]}[y : \lambda] \subset \mathcal{T}_{F,\tilde{\lambda}}^{[1]}[x : \lambda] + \lambda \rho(x, y) Q_0.$$

Hence,

$$\mathbf{Pr}(O, \mathcal{T}_{F,\tilde{\lambda}}^{[1]}[x : \lambda]) + \lambda \rho(x, y) Q_0 \supset Q(O, \delta(y)) \cap \mathcal{T}_{F,\tilde{\lambda}}^{[1]}[y : \lambda] = \mathbf{Pr}(O, \mathcal{T}_{F,\tilde{\lambda}}^{[1]}[y : \lambda]).$$

By interchanging the roles of x and y , we obtain also

$$\mathbf{Pr}(O, \mathcal{T}_{F,\tilde{\lambda}}^{[1]}[y : \lambda]) + \lambda \rho(x, y) Q_0 \supset \mathbf{Pr}(O, \mathcal{T}_{F,\tilde{\lambda}}^{[1]}[x : \lambda]).$$

These two inclusions imply the following inequality:

$$d_H(\mathbf{Pr}(O, \mathcal{T}_{F,\tilde{\lambda}}^{[1]}[x : \lambda]), \mathbf{Pr}(O, \mathcal{T}_{F,\tilde{\lambda}}^{[1]}[y : \lambda])) \leq \lambda \rho(x, y), \quad x, y \in \mathcal{M}. \quad (5.6)$$

As we have noted above, the set $\mathbf{Pr}(O, \mathcal{T}_{F, \tilde{\lambda}}^{[1]}[x : \lambda])$ is either a singleton or a compact line segment. Thus $\mathbf{Pr}(O, \mathcal{T}_{F, \tilde{\lambda}}^{[1]}[x : \lambda])$ is a *bounded rectangle*, so that, thanks to part (ii) of Claim 2.7 and (5.6), the mapping g_F defined by formula (1.7) is Lipschitz with Lipschitz seminorm at most λ .

Let us also note that, thanks to (5.2) and (5.3), for every $x \in \mathcal{M}$, we have

$$g_F(x) \in \mathcal{T}_{F, \tilde{\lambda}}(x) = \mathcal{H}[F^{[1]}[x : \tilde{\lambda}]].$$

These properties of g_F show that the mapping $g = g_F$ satisfies conditions (3.14) and (3.15).

Furthermore, comparing (3.16) with (1.8), we conclude that $f = f_{\tilde{\lambda}, F}$ where f is the mapping defined by the formula (3.16). Proposition 3.3 tells us that this mapping is a selection of F with Lipschitz seminorm

$$\|f\|_{\text{Lip}(\mathcal{M})} = \|f_{\tilde{\lambda}, F}\|_{\text{Lip}(\mathcal{M})} \leq \tilde{\lambda} + 2\lambda = \lambda_1 + 2\lambda_2.$$

The proof of Theorem 1.9 is complete. \blacksquare

PROOF OF THEOREM 1.11. Let $\mathfrak{M} = (\mathcal{M}, \rho)$ be a finite pseudometric space, and let $F : \mathcal{M} \rightarrow \text{Conv}(\mathbf{R}^2)$ be a set-valued mapping. We know that, given $\tilde{\lambda}, \lambda \geq 0$, condition (1.11) holds for every $x, x', x'', y, y', y'' \in \mathcal{M}$. Lemma 3.1 tells us that, in this case the set $F^{[1]}[x : \tilde{\lambda}]$ is nonempty for every $x \in \mathcal{M}$. Furthermore, part (ii) of Proposition 3.2 tells us that, for every $x \in \mathcal{M}$, the set $\mathcal{T}_{F, \tilde{\lambda}}^{[1]}[x : \lambda]$ is nonempty as well. See (3.9).

Therefore, thanks to (1.10), the $\vec{\lambda}$ -PA with $\vec{\lambda} = (\tilde{\lambda}, \lambda)$ produces the outcome “**Success**”. Furthermore, part (ii) of Theorem 1.9 tells us that, in this case the $\vec{\lambda}$ -PA returns the mapping $f_{\tilde{\lambda}, F} : \mathcal{M} \rightarrow \mathbf{R}^2$ having the following properties: $f_{\tilde{\lambda}, F}$ is a Lipschitz selection of F with

$$\|f_{\tilde{\lambda}, F}\|_{\text{Lip}(\mathcal{M})} \leq 2\lambda + \tilde{\lambda}.$$

Thus, (1.12) holds and the proof of Theorem 1.11 is complete. \blacksquare

5.2 Other versions of the Projection Algorithm.

Theorems 1.9 and 1.11 hold for various versions of the $\vec{\lambda}$ -Projection Algorithm relating to the particular choice of the mapping g_F in **STEP 4** of the algorithm. In particular, the proof of Theorem 1.9 presented above shows that the only requirement for g_F is that it be a Lipschitz selection of the set-valued mapping

$$\mathcal{T}_{F, \lambda_1} = \mathcal{H}[F^{[1]}[\cdot : \lambda_1]] \quad \text{with} \quad \|g_F\|_{\text{Lip}(\mathcal{M})} \leq \lambda_2.$$

In other words, g_F have to satisfy (3.14) and inequality (3.15) with constants $\tilde{\lambda} = \lambda_1$ and $\lambda = \lambda_2$.

Let us indicate some of these versions.

(★1) We can define g_F by formula (1.7) with replacing the origin O by an *arbitrary* point in \mathbf{R}^2 ;

(★2) Suppose that

$$\text{for every } x \in \mathcal{M} \text{ the set } F^{[1]}[x : \lambda_1] \text{ is bounded.} \quad (5.7)$$

Then the rectangle $\mathcal{T}_{F, \lambda_1}^{[1]}[x : \lambda_2]$ (see (1.6)) defined at **STEP 3** of the algorithm is also bounded for all $x \in \mathcal{M}$. In this case, we can define g_F by the formula

$$g_F(x) = \text{center} \left(\mathcal{T}_{F, \lambda_1}^{[1]}[x : \lambda_2] \right), \quad x \in \mathcal{M}. \quad (5.8)$$

See Fig. 10.




Fig. 10: **STEP 4** of the Projection Algorithm: the case of bounded sets $F^{[1]}[x : \lambda_1]$.

Then, thanks to part (iii) of Proposition 3.2, g_F is Lipschitz with $\|g_F\|_{\text{Lip}(\mathcal{M})} \leq \lambda_2$. Thus, for this choice of g_F both property (3.14) and inequality (3.15) hold.

We note that, thanks to Lemma 3.1, property (5.7) holds provided the set \mathcal{M} is *infinite*, so that in this case we can define g_F by formula (5.8).

Of course, in general we cannot guarantee that property (5.7) holds. However, in this case, the following property of the rectangles $\{\mathcal{T}_{F,\lambda_1}^{[1]}[x : \lambda_2] : x \in \mathcal{M}\}$ maybe useful: if one of the rectangles of this family is a *bounded set*, then *all rectangles* from this family are bounded as well, i.e., (5.7) holds. This property is immediate from the fact that at **STEP 4** the mapping $\mathcal{T}_{F,\lambda_1}^{[1]}[\cdot : \lambda_2]$ is Lipschitz with respect to the Hausdorff distance. See inequality (5.4). (Recall that $\vec{\lambda} = \lambda_1$ and $\lambda = \lambda_2$.)

(★3) Because \mathcal{M} is finite, there exists $R > 0$ such that,

$$\mathcal{T}_{F,\lambda_1}^{[1]}[x : \lambda_2] \cap Q(O, R) \neq \emptyset \quad \text{for every } x \in \mathcal{M}.$$

This enables us to define g_F by the formula

$$g_F(x) = \text{center} \left(\mathcal{T}_{F,\lambda_1}^{[1]}[x : \lambda_2] \cap Q(O, R) \right), \quad x \in \mathcal{M}.$$

Clearly,

$$g_F(x) \in \mathcal{T}_{F,\lambda_1}^{[1]}[x : \lambda_2] \subset \mathcal{T}_{F,\lambda_1}(x) \quad \text{on } \mathcal{M}$$

so that property (3.14) holds. The proof of (3.15) in this case follows the same scheme as the proof of this inequality for g_F defined by formula (1.7). We leave the details to the interested reader.

6. From geometric description to analytical: five basic algorithms.

Let us proceed to a detailed analytical description of the $\vec{\lambda}$ -Projection Algorithm.

First, let us describe *the input data*.

Statement 6.1 *The inputs of the $\vec{\lambda}$ -PA are the following objects:*

- (●1) A positive integer N ;
- (●2) A vector $\vec{\lambda} = (\lambda_1, \lambda_2)$ with non-negative coordinates λ_1 and λ_2 ;
- (●3) An $N \times N$ distance matrix $\mathfrak{D} = \{r_{ij} : i, j = 1, \dots, N\}$ with real entries $r_{ij} \geq 0$ satisfying the standard (pseudo)metric axioms:
 - (i) $r_{ii} = 0$ for all $1 \leq i \leq N$;
 - (ii) $r_{ij} = r_{ji}$ for all $1 \leq i, j \leq N$, i.e., \mathfrak{D} is a symmetric matrix;

(iii) $r_{ij} \leq r_{ik} + r_{kj}$ for all $i, j, k = 1, \dots, N$ (the triangle inequality).

(●4) N two dimensional non-zero vectors $h_i = (a_i, b_i) \in \mathbf{R}^2$, and N real numbers $\alpha_i, i = 1, \dots, N$.

These inputs generate a finite pseudometric space

$$\mathfrak{M} = (\mathcal{M}, \rho) \quad \text{with elements} \quad x_i = i, \quad i = 1, \dots, N, \quad (6.1)$$

a pseudometric

$$\rho(x_i, x_j) = r_{ij}, \quad i, j = 1, \dots, N, \quad (6.2)$$

and two mappings, $h : \mathcal{M} \rightarrow \mathbf{R}^2 \setminus \{0\}$ and $\alpha : \mathcal{M} \rightarrow \mathbf{R}$ defined by the formulae

$$h(x) = h_i \quad \text{and} \quad \alpha(x) = \alpha_i \quad \text{for} \quad x = i \in \mathcal{M}, \quad i = 1, \dots, N. \quad (6.3)$$

We also define a set-valued mapping $F : \mathcal{M} \rightarrow \mathcal{HP}(\mathbf{R}^2)$ by letting

$$F(x) = \{u \in \mathbf{R}^2 : \langle h(x), u \rangle + \alpha(x) \leq 0\}, \quad x \in \mathcal{M}. \quad (6.4)$$

Thus, each $F(x)$ is a half-plane in \mathbf{R}^2 and $h(x)$ is a vector directed outside of $F(x)$ and orthogonal its boundary, the straight line $\ell(x) = \{u \in \mathbf{R}^2 : \langle h(x), u \rangle + \alpha(x) = 0\}$.

Given these inputs, the $\vec{\lambda}$ -Projection Algorithm produces the outcomes described in Theorem 1.9. This procedure follows the approach outlined in Section 1.2 and includes five main algorithms, which we present in this section.

As we noted in Section 1.2, our approach to the corresponding algorithms at **STEPS 1-2** of the $\vec{\lambda}$ -PA relies on a classical result on low-dimensional linear programming which we cite below.

Consider a linear program in \mathbf{R}^d expressed in the inequality form

$$\text{maximize} \quad z = \langle c, x \rangle \quad \text{subject to} \quad Ax \leq b \quad (6.5)$$

where $c \in \mathbf{R}^d$, $b \in \mathbf{R}^m$ and the matrix $A \in \mathbf{R}^{m \times d}$ are the input data, and $x \in \mathbf{R}^d$. (The vector inequality $Ax \leq b$ in (6.5) is with respect to the componentwise partial order in \mathbf{R}^m .)

M. E. Dyer [21] and N. Megiddo [52] found, independently, an efficient algorithm for the linear programming problem (6.5) in the cases $d = 2$.

Theorem 6.2 *The algorithms presented in [21] and [52] give a solution (or say it is unbounded or unfeasible) to the linear program (6.5) in two variables and m constraints.*

The total work and storage required by this algorithm are at most Cm where C is an absolute constant.

Remark 6.3 As usual, the notion of “feasibility” that we use in the statement of Theorem 6.2 means the following: in $O(m)$ running time and $O(m)$ storage, the above algorithm determines *whether the feasible set of a linear program* (i.e., the set of points $x \in \mathbf{R}^d$ satisfying the constraints $Ax \leq b$) *is empty or not.* ◀

Remark 6.4 Let us recall something of the history of algorithms for the linear programming problem, where the running time is analyzed as a function of the number of constraints m and the number of variables d . See [13] for more details.

First, we note that it is also shown in [52] that Theorem 6.2 holds in the three dimensional case.

N. Megiddo [53] generalized Theorem 6.2 to an arbitrary d , proving that a linear program in d variables and m constraints can be solved in $2^{O(2^d)m}$ time. This result has become a cornerstone of computational geometry, with many applications in low-dimensional settings. See also [24].

It was improved by K. L. Clarkson [17] and M. E. Dyer [22] to $O((5/9)^d 3^{d^2} m)$ and $O(3^{(d+2)^2} m)$ respectively. Further improvements were obtained by P. K. Agarwal, M. Sharir and S. Toledo [1] to $O(d)^{10d} (\log d)^{2d} m$, by B. Chazelle and J. Matoušek [15] to $O(d)^{7d} (\log d)^d m$ and H. Brönnimann, B. Chazelle and J. Matoušek [6] to $O(d)^{7d} (\log d)^d m$. The fastest algorithm known to the moment is due to T. M. Chan [13] with the running time $O(d)^{d/2} (\log d)^{3d} m$. ◀

6.1 The λ -metric refinement of the set-valued mapping - STEP 1.

The goal of this section is to provide an efficient algorithm for recognizing whether, for each $x \in \mathcal{M}$, the set $F^{[1]}[x : \lambda_1]$ defined by formula (1.4) (the λ_1 -metric refinement of F) is nonempty or not.

Algorithm 6.5 *There exists an algorithm with the inputs given in Statement 6.1, which produces one of the following outcomes:*

OUTCOME 1: *We guarantee that there exists $x \in \mathcal{M}$ such that the set $F^{[1]}[x : \lambda_1]$ is empty.*

OUTCOME 2: *We guarantee that for each $x \in \mathcal{M}$, the set $F^{[1]}[x : \lambda_1] \neq \emptyset$.*

The total work and storage required by this algorithm are at most CN^2 where C is an absolute constant.

Explanation: We recall that, for every $x \in \mathcal{M}$,

$$F^{[1]}[x : \lambda_1] = \bigcap_{y \in \mathcal{M}} H(x; y) \quad (6.6)$$

where

$$H(x; y) = F(y) + \lambda_1 \rho(x, y) Q_0, \quad y \in \mathcal{M}.$$

See (1.3). Thanks to (6.4),

$$H(x; y) = \{u \in \mathbf{R}^2 : \langle h(y), u \rangle + \alpha(y) \leq 0\} + \lambda_1 \rho(x, y) Q_0.$$

Thus, $H(x, y)$ is also a half-plane having the following representation:

$$H(x; y) = \{u \in \mathbf{R}^2 : \langle h(y), u \rangle + \alpha(y) \leq \lambda_1 \rho(x, y) \|h(y)\|_1\}, \quad y \in \mathcal{M}. \quad (6.7)$$

Here $\|h\|_1 = |h_1| + |h_2|$ for $h = (h_1, h_2) \in \mathbf{R}^2$ is the ℓ_1^2 -norm in \mathbf{R}^2 .

Fix $x \in \mathcal{M}$ and consider a system of linear inequalities in two variables $(u_1, u_2) = u$:

$$\langle h(y), u \rangle + \alpha(y) \leq \lambda_1 \rho(x, y) \|h(y)\|_1, \quad y \in \mathcal{M}. \quad (6.8)$$

Thanks to (6.6) and (6.7), this system has a solution if and only if the set $F^{[1]}[x : \lambda_1] \neq \emptyset$.

We apply Theorem 6.2 to the system (6.8) of linear inequalities with respect to two variables. This theorem tells us that in $O(N)$ running time with $O(N)$ storage we can check the feasibility of this system, i.e., to say whether it has a solution or not.

We apply this procedure to each $x = i \in \mathcal{M}$, $i = 1, \dots, N$, starting from $x = 1$. If for a certain element $\hat{x} = i_0$, $1 \leq i_0 \leq N$, Theorem 6.2 will tell us that the system has no a feasible solution, then $F^{[1]}[\hat{x} : \lambda_1] = \emptyset$. In this case we produce OUTCOME 1.

Recall that in this case the $\vec{\lambda}$ -Projection Algorithm also produces the outcome “No Go” (see Theorem 1.3) and terminates.

If Theorem 6.2 tells us that for every $x \in \mathcal{M}$ the system (6.8) has a solution, we produce OUTCOME 2. In this case, the set $F^{[1]}[x : \lambda_1] \neq \emptyset$ for every $x \in \mathcal{M}$ which corresponds to the outcome “Success” in Theorem 1.3.

Since we apply the algorithm of Theorem 6.2 at most N times, and each time this algorithm produces an answer in $O(N)$ running time with $O(N)$ storage, the work and storage required to perform all steps of this procedure are $O(N^2)$. ■

6.2 The rectangular hulls of metric refinements - STEP 2.

In this section we present an efficient algorithm for computing the rectangle $\mathcal{T}_{F,\lambda_1}(x)$, see (1.5), i.e., the rectangular hull of the set $F^{[1]}[x : \lambda_1]$, $x \in \mathcal{M}$.

Algorithm 6.6 *There exists an algorithm which for any inputs from Statement 6.1 such that the set $F^{[1]}[x : \lambda_1] \neq \emptyset$ for every $x \in \mathcal{M}$, produces $4N$ numbers*

$$s_1(x), s_2(x), t_1(x), t_2(x) \in \mathbf{R} \cup \{\pm\infty\}, \quad s_1(x) \leq s_2(x), \quad t_1(x) \leq t_2(x), \quad x \in \mathcal{M},$$

which determine the rectangle $\mathcal{T}_{F,\lambda_1}(x) = \mathcal{H}[F^{[1]}[x : \lambda_1]]$ for every $x \in \mathcal{M}$.

This means the following: $\mathcal{T}_{F,\lambda_1}(x) = I_1(x) \times I_2(x)$ where $I_1(x) \in \mathcal{I}(Ox_1)$ and $I_2(x) \in \mathcal{I}(Ox_2)$ are nonempty closed intervals on the axes Ox_1 and Ox_2 . These intervals have the following properties: $s_1(x)$ and $s_2(x)$ are the left and right ends of $I_1(x)$, see (2.3), and $t_1(x)$ and $t_2(x)$ are the left and right ends of $I_2(x)$, see (2.4).

The work and storage required by this algorithm are at most CN^2 where C is an absolute constant.

Explanation: Note that

$$I_i(x) = \text{PR}_i[\mathcal{T}_{F,\lambda_1}(x)] = \text{PR}_i[F^{[1]}[x : \lambda_1]], \quad i = 1, 2.$$

Following Remark 2.1 and formulae (2.5) and (2.6), we set $S = F^{[1]}[x : \lambda_1]$,

$$s_1(x) = \sigma_1(S), \quad s_2(x) = \sigma_2(S) \quad \text{and} \quad t_1(x) = \tau_1(S), \quad t_2(x) = \tau_2(S), \quad x \in \mathcal{M}.$$

Remark 2.1 tells us that in this case the numbers $s_i(x), t_i(x), i = 1, 2$, determine the rectangle $\mathcal{T}_{F,\lambda_1}(x)$ in the sense of Algorithm 6.6. Thus,

$$s_1(x) = \inf\{\langle \xi_1, u \rangle : u \in S\}, \quad s_2(x) = \sup\{\langle \xi_1, u \rangle : u \in S\}, \quad (6.9)$$

and

$$t_1(x) = \inf\{\langle \xi_2, u \rangle : u \in S\}, \quad t_2(x) = \sup\{\langle \xi_2, u \rangle : u \in S\} \quad (6.10)$$

provided $S = F^{[1]}[x : \lambda_1]$, $\xi_1 = (1, 0)$ and $\xi_2 = (0, 1)$.

We recall, that, thanks to (6.6) and (6.7), given $x \in \mathcal{M}$, the set $S = F^{[1]}[x : \lambda_1]$ is a polygon determined by N linear constraints given in (6.8).

Given $\xi \in \mathbf{R}^2$, let us consider a linear program in \mathbf{R}^2

$$\text{maximize} \quad z = \langle \xi, u \rangle \quad \text{subject to} \quad u \in S. \quad (6.11)$$

Then, thanks to (6.9) and (6.10), the numbers $s_2(x)$ and $t_2(x)$ are solutions to this problem where ξ equals to $(1, 0)$ or $(0, 1)$ respectively. In turn, the numbers $s_1(x)$ and $t_1(x)$ are solutions to the linear program

$$\text{minimize } z = \langle \xi, u \rangle \quad \text{subject to } u \in S, \quad (6.12)$$

with the same vectors $\xi = (1, 0)$ or $\xi = (0, 1)$.

Theorem 6.2 tells us that each of these four linear programs (i.e., the linear programs (6.11) and (6.12) for computing the numbers $s_1(x), s_2(x), t_1(x)$ and $t_2(x)$) can be solved in $O(N)$ running time with $O(N)$ storage.

Then, we apply this procedure to every $x_i \in \mathcal{M}, i = 1, \dots, N$, starting from $i = 1$. After N steps, we will determine all N rectangles $\mathcal{T}_{F,\lambda_1}(x_i), i = 1, \dots, N$, so that the total work and storage required to proceed all steps of this procedure are $O(N^2)$. ■

6.3 The metric refinements of the rectangular hulls - STEP 3.

At this step of the $\vec{\lambda}$ -Projection Algorithm our inputs are $4N$ numbers

$$s_1(x), s_2(x), t_1(x), t_2(x) \in \mathbf{R} \cup \{\pm\infty\}, \quad s_1(x) \leq s_2(x), \quad t_1(x) \leq t_2(x), \quad x \in \mathcal{M},$$

obtained in Algorithm 6.6. These numbers determine the rectangles $\{\mathcal{T}_{F,\lambda_1}(x) : x \in \mathcal{M}\}$, which means that for each $x \in \mathcal{M}$ the numbers $s_1(x)$ and $s_2(x)$ are the left and right ends of $\text{PR}_1[\mathcal{T}_{F,\lambda_1}(x)]$, and $t_1(x)$ and $t_2(x)$ are the left and right ends of $\text{PR}_2[\mathcal{T}_{F,\lambda_1}(x)]$. Let us also note the following properties of these numbers:

$$s_1(x), t_1(x) < +\infty \quad \text{and} \quad s_2(x), t_2(x) > -\infty.$$

Our aim is to give an efficient algorithm to recognize whether for every $x \in \mathcal{M}$ the λ_2 -metric refinement of the rectangle $\mathcal{T}_{F,\lambda_1}$, i.e., the set

$$\mathcal{T}_{F,\lambda_1}^{[1]}[x : \lambda_2] = \bigcap_{y \in \mathcal{M}} [\mathcal{T}_{F,\lambda_1}(y) + \lambda_2 \rho(x, y) \mathcal{Q}_0] \quad (6.13)$$

is nonempty or not. Recall that, according to **STEP 3** of the $\vec{\lambda}$ -Projection Algorithm, if this set is empty for some $x \in \mathcal{M}$, this algorithm produces the outcome “**No Go**” and terminates. If the set $\mathcal{T}_{F,\lambda_1}^{[1]}[x : \lambda_2] \neq \emptyset$ for all $x \in \mathcal{M}$, we produce the ends of the projections of this set onto the coordinate axes and turn to **STEP 4** of the $\vec{\lambda}$ -PA.

Algorithm 6.7 *There exists an algorithm with the inputs $\{s_1(x), s_2(x), t_1(x), t_2(x) : x \in \mathcal{M}\}$ which produces one of the following outcomes:*

OUTCOME 1: *We guarantee that there exists $x \in \mathcal{M}$ such that the set $\mathcal{T}_{F,\lambda_1}^{[1]}[x : \lambda_2]$ is empty.*

OUTCOME 2: *We guarantee that, for every $x \in \mathcal{M}$, the set $\mathcal{T}_{F,\lambda_1}^{[1]}[x : \lambda_2] \neq \emptyset$.*

Furthermore, at this stage the algorithm produces $4N$ numbers

$$s_1^{[1]}(x), s_2^{[1]}(x), t_1^{[1]}(x), t_2^{[1]}(x) \in \mathbf{R} \cup \{\pm\infty\}, \quad s_1^{[1]}(x) \leq s_2^{[1]}(x), \quad t_1^{[1]}(x) \leq t_2^{[1]}(x), \quad x \in \mathcal{M}, \quad (6.14)$$

which for each $x \in \mathcal{M}$ determine the rectangle $\mathcal{T}_{F,\lambda_1}^{[1]}[x : \lambda_2]$.

The total work and storage required by this algorithm are at most CN^2 where C is a universal constant.

Explanation: Let us note that the numbers $s_i^{[1]}(x)$ and $t_i^{[1]}(x)$, $i = 1, 2$, determine the rectangle $\mathcal{T}_{F,\lambda_1}^{[1]}[x : \lambda_2]$ in the same sense as in formulation of Algorithm 6.6: the rectangle

$$\mathcal{T}_{F,\lambda_1}^{[1]}[x : \lambda_2] = \widetilde{I}_1(x) \times \widetilde{I}_2(x) \quad \text{where} \quad \widetilde{I}_i(x) \in \mathcal{I}(Ox_i), \quad i = 1, 2,$$

are nonempty closed intervals on the axes Ox_1 and Ox_2 . Thus,

$$\widetilde{I}_i(x) = \text{PR}_i[\mathcal{T}_{F,\lambda_1}^{[1]}[x : \lambda_2]], \quad i = 1, 2.$$

These intervals have the following properties: $s_1^{[1]}(x)$ and $t_2^{[1]}(x)$ are the left and right ends of $\widetilde{I}_1(x)$, see (2.3), and $t_1^{[1]}(x)$ and $s_2^{[1]}(x)$ are the left and right ends of $\widetilde{I}_2(x)$, see (2.4).

We define the numbers from (6.14) as follows: we set

$$s_1^{[1]}(x) = \max\{s_1(y) - \lambda_2 \rho(x, y) : y \in \mathcal{M}\}, \quad s_2^{[1]}(x) = \min\{s_2(y) + \lambda_2 \rho(x, y) : y \in \mathcal{M}\},$$

and

$$t_1^{[1]}(x) = \max\{t_1(y) - \lambda_2 \rho(x, y) : y \in \mathcal{M}\}, \quad t_2^{[1]}(x) = \min\{t_2(y) + \lambda_2 \rho(x, y) : y \in \mathcal{M}\}.$$

We turn to the description of the promised algorithm. It includes the following two steps:

(★1) If there exist $x \in \mathcal{M}$ such that

$$\text{either } s_1^{[1]}(x) > s_2^{[1]}(x) \quad \text{or} \quad t_1^{[1]}(x) > t_2^{[1]}(x) \tag{6.15}$$

then we produce OUTCOME 1. We state that in this case the set $\mathcal{T}_{F,\lambda_1}^{[1]}[x : \lambda_2] = \emptyset$.

(★2) If for every $x \in \mathcal{M}$, we have

$$s_1^{[1]}(x) \leq s_2^{[1]}(x) \quad \text{and} \quad t_1^{[1]}(x) \leq t_2^{[1]}(x), \tag{6.16}$$

we produce OUTCOME 2. In this case we guarantee that $\mathcal{T}_{F,\lambda_1}^{[1]}[x : \lambda_2] \neq \emptyset$ for every $x \in \mathcal{M}$. Furthermore, we state that the rectangle $\mathcal{T}_{F,\lambda_1}^{[1]}[x : \lambda_2]$ is determined by the numbers $s_i^{[1]}(x), t_i^{[1]}(x)$, $i = 1, 2$.

Let us explain part (★1) of this algorithm. We have to prove that the set $\mathcal{T}_{F,\lambda_1}^{[1]}[x : \lambda_2] \neq \emptyset$ provided (6.15) holds. Let us suppose, for instance, that

$$s_1^{[1]}(x) > s_2^{[1]}(x). \tag{6.17}$$

Let

$$\mathcal{R} = \{\Pi = \mathcal{T}_{F,\lambda_1}(y) + \lambda_2 \rho(x, y) Q_0 : y \in \mathcal{M}\}.$$

Then, thanks to (6.13),

$$\mathcal{T}_{F,\lambda_1}^{[1]}[x : \lambda_2] = \bigcap_{\Pi \in \mathcal{R}} \Pi. \tag{6.18}$$

Let us show that

$$\bigcap_{\Pi \in \mathcal{R}} \text{PR}_1[\Pi] = \emptyset \tag{6.19}$$

provided (6.17) holds. Indeed, thanks to (2.16),

$$\text{PR}_1[\Pi] = \text{PR}_1[\mathcal{T}_{F,\lambda_1}(y) + \lambda_2 \rho(x, y) Q_0] = \text{PR}_1[\mathcal{T}_{F,\lambda_1}(y)] + \lambda_2 \rho(x, y) I_1 \tag{6.20}$$

where $I_1 = \text{Pr}_1[Q_0] = [(-1, 0), (1, 0)]$.

We know that the closed interval $I(y) = \text{Pr}_1[\mathcal{T}_{F,\lambda_1}(y)]$ has the ends at $s_1(y)$ and $s_2(y)$, i.e., $L(I(y)) = s_1(y)$ and $R(I(y)) = s_2(y)$, see (2.3).

Let us identify the axis Ox_1 with \mathbf{R} and every interval $I(y)$, $y \in \mathcal{M}$, on $O(x_1)$ with a closed interval in \mathbf{R} with the ends in $s_1(y)$ and $s_2(y)$. Let us denote this interval by $G(y)$. Thus, we have a set-valued mapping $G : \mathcal{M} \rightarrow \mathcal{I}(\mathbf{R})$ such that $\inf G(y) = s_1(y)$ and $\sup G(y) = s_2(y)$ for every $y \in \mathcal{M}$.

This identification implies the following: thanks to (6.20), property (6.19) holds provided the set

$$G^{[1]}[x : \lambda_2] = \bigcap_{y \in \mathcal{M}} [G(y) + \lambda_2 \rho(x, y) I_0] \neq \emptyset. \quad (6.21)$$

Let us apply Proposition 2.10 to the mapping G . But first, let us align our settings with the settings of this proposition. We note that, thanks to (2.38), we have

$$a_G(x) = \inf G(x) = s_1(x) \quad \text{and} \quad b_G(x) = \sup G(x) = s_2(x). \quad (6.22)$$

Furthermore, thanks to (2.40),

$$a_G^{[1]}[x : \lambda_2] = s_1^{[1]}(x) \quad \text{and} \quad b_G^{[1]}[x : \lambda_2] = s_2^{[1]}(x). \quad (6.23)$$

Therefore, thanks to assumption (6.17), $a_G^{[1]}[x : \lambda_2] > b_G^{[1]}[x : \lambda_2]$.

Proposition 2.10 and the criterion (2.45) tell us that in this case the set $G^{[1]}[x : \lambda_2] = \emptyset$, i.e., (6.21) holds. This proves (6.19) and guarantees that the set $\mathcal{T}_{F,\lambda_1}^{[1]}[x : \lambda_2] = \emptyset$ justifying OUTCOME 1.

We turn to the part (★2) of the algorithm. In the same fashion, using Proposition 2.10, we show that if both inequalities in (6.16) hold, then $\mathcal{T}_{F,\lambda_1}^{[1]}[x : \lambda_2] \neq \emptyset$.

Indeed, suppose that $s_1^{[1]}(x) \leq s_2^{[1]}(x)$ on \mathcal{M} . In this case,

$$a_G^{[1]}[x : \lambda_2] \leq b_G^{[1]}[x : \lambda_2] \quad \text{for every } x \in \mathcal{M}.$$

Therefore, thanks to (2.45) and Proposition 2.10, we have $\bigcap \{\text{Pr}_1[\Pi] : \Pi \in \mathcal{R}\} \neq \emptyset$. In the same way we prove that

$$\bigcap_{\Pi \in \mathcal{R}} \text{Pr}_2[\Pi] \neq \emptyset \quad \text{provided} \quad t_1^{[1]}(x) \leq t_2^{[1]}(x) \quad \text{for all } x \in \mathcal{M}.$$

Therefore, thanks to (6.18) and Claim 2.2,

$$\mathcal{T}_{F,\lambda_1}^{[1]}[x : \lambda_2] = \bigcap_{\Pi \in \mathcal{R}} \Pi \neq \emptyset \quad \text{for every } x \in \mathcal{M}.$$

Furthermore, equalities (6.23) and (2.46) tell us that $s_1^{[1]}(x)$ and $s_2^{[1]}(x)$ are the ends of the set $G^{[1]}[x : \lambda_2]$ defined by (6.21). Because the projection $\text{Pr}_1[\mathcal{T}_{F,\lambda_1}^{[1]}[x : \lambda_2]]$ can be identified with the interval $G^{[1]}[x : \lambda_2]$, we conclude that $\text{Pr}_1[\mathcal{T}_{F,\lambda_1}^{[1]}[x : \lambda_2]]$ has the ends at $s_1^{[1]}(x)$ and $s_2^{[1]}(x)$.

In the same way we prove that $\text{Pr}_2[\mathcal{T}_{F,\lambda_1}^{[1]}[x : \lambda_2]]$ has the ends at $t_1^{[1]}(x)$ and $t_2^{[1]}(x)$.

Let us estimate the running time of the algorithm. Clearly, given $x \in \mathcal{M}$, thanks to (6.22), we need N arithmetic operation to compute $s_1^{[1]}(x)$. The same is true for the numbers $s_2^{[1]}(x)$, $t_1^{[1]}(x)$ and $t_2^{[1]}(x)$. This shows that for each $x \in \mathcal{M}$, the algorithm requires $O(N)$ running time to compute all numbers $s_i^{[1]}(x)$, $t_i^{[1]}(x)$, $i = 1, 2$. Therefore, the total running time for computing these numbers for all N -element set is $O(N^2)$.

Furthermore, if the numbers $s_i^{[1]}(x)$, $t_i^{[1]}(x)$, $i = 1, 2$, $x \in \mathcal{M}$, are computed, we need $O(N)$ operations of comparison to check (6.15) and (6.16). Therefore, the total work of the algorithm suggested

in (★1) and (★2) is $O(N^2)$. It is also clear this algorithm requires at most CN^2 units of the compute memory where C is a universal constant. ■

6.4 The centers of the metric projections onto rectangles - STEP 4.

At this stage of the Projection Algorithm our inputs are $4N$ numbers

$$s_1^{[1]}(x), s_2^{[1]}(x), t_1^{[1]}(x), t_2^{[1]}(x) \in \mathbf{R} \cup \{\pm\infty\}, \quad s_1^{[1]}(x) \leq s_2^{[1]}(x), \quad t_1^{[1]}(x) \leq t_2^{[1]}(x), \quad x \in \mathcal{M}, \quad (6.24)$$

obtained with the help of Algorithm 6.7. These numbers determine the rectangles $\mathcal{T}_{F,\lambda_1}^{[1]}[x : \lambda_2]$, $x \in \mathcal{M}$. Our goal is to present an algorithm that, based on these inputs, constructs the mapping $g_F : \mathcal{M} \rightarrow \mathbf{R}^2$ defined by formula (1.7).

Algorithm 6.8 *There exists an algorithm which for the inputs (6.24) produces N points $g_F(x) \in \mathbf{R}^2$ defined by the formula*

$$g_F(x) = \text{center} \left(\mathbf{Pr} \left(O, \mathcal{T}_{F,\lambda_1}^{[1]}[x : \lambda_2] \right) \right), \quad x \in \mathcal{M}.$$

The work and storage required by this algorithm are at most CN where C is an absolute constant.

Explanation: We compute the coordinates of the points $g_F(x) = (g_1(x), g_2(x)) \in \mathbf{R}^2$, $x \in \mathcal{M}$, using the explicit formulae for these coordinates given in Claim 2.5. Indeed, we know that for each $x \in \mathcal{M}$, the rectangle

$$\mathcal{T}_{F,\lambda_1}^{[1]}[x : \lambda_2] = I_1(x) \times I_2(x)$$

where $I_i(x) \in \mathcal{I}(Ox_i)$ are closed intervals on the axes Ox_i determined by the numbers $s_i^{[1]}(x)$, $t_i^{[1]}(x)$, $i = 1, 2$, from the inputs (6.24). More specifically, $s_1^{[1]}(x)$ and $s_2^{[1]}(x)$ are the left and right ends of $I_1(x)$ while $t_1^{[1]}(x)$ and $t_2^{[1]}(x)$ are the left and right ends of $I_2(x)$.

Claim 2.5 tells us that we can compute the required coordinates $g_1(x)$ and $g_2(x)$ by letting $g_1(x) = g_1$ and $g_2(x) = g_2$ where the numbers g_1 and g_2 are determined by formulae (2.25), (2.26) and (2.27) with the initial parameters

$$a_1 = s_1^{[1]}(x), \quad b_1 = s_2^{[1]}(x), \quad a_2 = t_1^{[1]}(x) \quad \text{and} \quad b_2 = t_2^{[1]}(x).$$

Clearly, for every $x \in \mathcal{M}$, we need at most C arithmetic operations and C units of computer memory to calculate the coordinates of the point $g_F(x)$. Here C is a universal constant.

Therefore, the total work and storage required by this algorithm for computing the mapping g_F on \mathcal{M} are at most CN . ■

6.5 The final step: metric projections onto the metric refinements - STEP 5.

We move on to the final part of the Projection Algorithm. At this stage our input data are the inputs given in Statement 6.1 and the mapping g_F constructed in Algorithm 6.8. Below we give an efficient algorithm which produces the mapping $f_{\vec{\lambda};F} : \mathcal{M} \rightarrow \mathbf{R}^2$ defined by formula (1.8), i.e., the metric projection of $g_F(x)$ onto the set $F^{[1]}[x : \lambda_1]$, $x \in \mathcal{M}$. The mapping $f_{\vec{\lambda};F}$ is the finite goal of the $\vec{\lambda}$ -Projection Algorithm. Theorem 1.9 tells us that $f_{\vec{\lambda};F}$ is a Lipschitz selection of F with Lipschitz seminorm at most $\lambda_1 + 2\lambda_2$.

Algorithm 6.9 *There exists an algorithm that, for the mapping g_F and the inputs given in Statement 6.1, produces the mapping $f_{\vec{\lambda};F}$ defined by the formula*

$$f_{\vec{\lambda};F}(x) = \mathbf{Pr} \left(g_F(x), F^{[1]}[x : \lambda_1] \right), \quad x \in \mathcal{M}. \quad (6.25)$$

The work and storage required by this algorithm are at most CN^2 where C is a universal constant.

Explanation: First, we present several simple formulas related to calculating the distance from a point $g \in \mathbf{R}^2$ to a half-plane H , as well as the closest point to g on H in the ℓ_∞^2 -norm.

Let $h = (h_1, h_2)$ be a point in \mathbf{R}^2 with coordinates $h_1 \neq 0$ and $h_2 \neq 0$, and let $\alpha \in \mathbf{R}$. Let $H \in \mathcal{HP}(\mathbf{R}^2)$ be a half-plane determined by h and α :

$$H = \{u \in \mathbf{R}^2 : \langle h, u \rangle + \alpha \leq 0\}.$$

Let $g = (g_1, g_2) \in \mathbf{R}^2$. The following formula is well known:

$$\text{dist}(g, H) = |\langle h, g \rangle + \alpha|_+ / \|h\|_1. \quad (6.26)$$

(Recall that $\|h\|_1 = |h_1| + |h_2|$.)

Let $f = (f_1, f_2) \in \mathbf{R}^2$ be a point in H nearest to g (in $\|\cdot\|$ -norm). (Since $h_i \neq 0$, $i = 1, 2$, such a point is well defined and unique.) Clearly,

$$f = g \quad \text{provided} \quad \langle h, g \rangle + \alpha \leq 0 \quad (6.27)$$

(because in this case $g \in H$).

Suppose that $\langle h, g \rangle + \alpha > 0$. Then

$$f_1 = \frac{h_2(g_1 - g_2) - \alpha}{h_1 + h_2} \quad \text{and} \quad f_2 = \frac{h_1(g_2 - g_1) - \alpha}{h_1 + h_2} \quad \text{if} \quad h_1 \cdot h_2 > 0, \quad (6.28)$$

and

$$f_1 = \frac{h_2(g_1 + g_2) - \alpha}{h_2 - h_1} \quad \text{and} \quad f_2 = \frac{h_1(g_2 + g_1) - \alpha}{h_1 - h_2} \quad \text{if} \quad h_1 \cdot h_2 < 0. \quad (6.29)$$

The proof of these formulae we leave the reader as an easy exercise.

Let us also recall that, for every $x \in \mathcal{M}$, the set $F^{[1]}[x : \lambda_1]$ is given by the formula

$$F^{[1]}[x : \lambda_1] = \bigcap_{y \in \mathcal{M}} H(x; y) \quad (6.30)$$

where

$$H(x; y) = \{u \in \mathbf{R}^2 : \langle h(y), u \rangle + \alpha(y) - \lambda_1 \rho(x, y) \|h(y)\|_1 \leq 0\}, \quad y \in \mathcal{M}. \quad (6.31)$$

See (6.6) and (6.7).

Given $x \in \mathcal{M}$, we construct the point $f(x) = f_{x, F}(x)$, defined by the formula (6.25), in two steps.

Let $\mathcal{H} = \{H(x; y) : y \in \mathcal{M}\}$. Clearly, thanks to representation (6.30),

$$\text{dist}(g_F(x), F^{[1]}[x : \lambda_1]) = \max \{\text{dist}(g_F(x), H(x; y)) : y \in \mathcal{M}\}. \quad (6.32)$$

At the first step of our algorithm, we fix a half-plane $H_0 = H(x; y_0) \in \mathcal{H}$ for which the maximum in the right hand side of the equality (6.32) is attained. We can find y_0 in N -steps because, thanks to (6.26) and (6.31), for each $y \in \mathcal{M}$, we have an explicit formula for the distance from $g_F(x)$ to $H(x; y)$:

$$\text{dist}(g_F, H(x; y)) = |\langle h(y), g_F(x) \rangle + \alpha(y) - \lambda_1 \rho(x, y) \|h(y)\|_1|_+ / \|h(y)\|_1.$$

Thus, in N steps we calculate the maximum in the right hand side of (6.32) using at most C operation at each step. This enables us in at most CN running time to determine the half-plane $H_0 = H(x; y_0)$ with the property

$$\text{dist}(g_F(x), F^{[1]}[x : \lambda_1]) = \text{dist}(g_F(x), H(x; y_0)).$$

We know that the point $g_F(x)$ belongs to the rectangular hull of the set $F^{[1]}[x : \lambda_1]$, i.e., to the set $\mathcal{H}[F^{[1]}[x : \lambda_1]]$. Because $F^{[1]}[x : \lambda_1] \subset H_0$, we have $g_F(x) \in \mathcal{H}[H_0]$, so that, thanks to Lemma 3.4, the metric projection $\mathbf{Pr}(g_F(x), H_0)$ of $g_F(x)$ onto H_0 is a *singleton*. This proves that

$$\mathbf{Pr}(g_F(x), F^{[1]}[x : \lambda_1]) = \mathbf{Pr}(g_F(x), H_0).$$

Hence, thanks to (6.25),

$$f(x) = f_{\vec{\lambda}; F}(x) = \mathbf{Pr}(g_F(x), H_0).$$

It remains to calculate the coordinates of the point $f(x) = (f_1(x), f_2(x))$ as the nearest point to $g_F(x)$ on the half-plane $H_0 = H(x; y_0)$. Recall that, thanks to (6.31),

$$H(x; y_0) = \{u \in \mathbf{R}^2 : \langle h(y_0), u \rangle + \alpha(y_0) - \lambda_1 \rho(x, y_0) \|h(y_0)\|_1 \leq 0\}.$$

This enables us to make use of the formulae (6.27), (6.28) and (6.29). We set in these formulas

$$g = g_F(x), \quad h = h(y_0) \quad \text{and} \quad \alpha = \alpha(y_0) - \lambda_1 \rho(x, y_0) \|h(y_0)\|_1,$$

and get the required point $f(x) = f_{\vec{\lambda}; F}(x)$. Clearly, these formulas require at most C computer operations to calculate $f(x) = f_{\vec{\lambda}; F}(x)$. Here C is an absolute constant.

Summarizing, we conclude that, following the above procedure, for every $x \in \mathcal{M}$ we need $O(N)$ running time and $O(N)$ storage to compute the coordinates of the point $f_{\vec{\lambda}; F}(x)$.

Therefore, the total work and storage required by this algorithm for computing the desired mapping $f_{\vec{\lambda}; F}$ on \mathcal{M} are at most CN^2 . ■

We are in a position to prove the main Theorem 1.3.

PROOF OF THEOREM 1.3. Let

$$\lambda_1 = \lambda_2 = \lambda \quad \text{and let} \quad \vec{\lambda} = (\lambda_1, \lambda_2) = (\lambda, \lambda).$$

Without loss of generality we can assume that our pseudometric space (\mathcal{M}, ρ) and the set-valued mapping F are generated by the inputs given in Statement 6.1 and formulas (6.1) - (6.4).

We successively apply Algorithms 6.5 - 6.9 to these initial data. These algorithms corresponds to **STEP 1 - STEP 5** of the $\vec{\lambda}$ -Projection Algorithm. They produces the outcome (“**No Go**”) and the outcome (“**Success**”) described in Theorem 1.9. This theorem tells us that in the case of the outcome “**No go**” we guarantee that there does not exist a Lipschitz selection of F with Lipschitz seminorm at most

$$\min\{\lambda_1, \lambda_2\} = \min\{\lambda, \lambda\} = \lambda.$$

Thus, this outcome coincides with **OUTCOME 1 (“No Go”)** of Theorem 1.3.

Theorem 1.9 also tells us that in the case of the outcome “**Success**”, the $\vec{\lambda}$ -Projection Algorithm returns the mapping $f_{\vec{\lambda}; F} : \mathcal{M} \rightarrow \mathbf{R}^2$, which is a Lipschitz selection of F on \mathcal{M} with Lipschitz seminorm

$$\|f_{\vec{\lambda}; F}\|_{\text{Lip}(\mathcal{M})} \leq \lambda_1 + 2\lambda_2 = \lambda + 2\lambda = 3\lambda.$$

Thus, this outcome coincides with **OUTCOME 1 (“Success”)** of Theorem 1.3.

Let us also note that the work and memory space needed for each of Algorithms 6.5 - 6.9 are at most $O(N^2)$. Therefore, the total work and storage used by $\vec{\lambda}$ -Projection Algorithm are at most CN^2 , where $C > 0$ is a universal constant.

The proof of Theorem 1.3 is complete. ■

6.6 Lipschitz selections of polygon-set valued mappings.

As promised in Section 1.1, in this section we give an analogue of Theorem 1.2 for the case $\tau = 0$ and $D = 2$.

Let L be a positive integer, and let $\mathcal{PG}_L(\mathbf{R}^2)$ be the family of all polygons in \mathbf{R}^2 (not necessarily bounded), each of which is defined by at most L linear constraints.

Theorem 6.10 *There exists an algorithm which, given an N -element pseudometric space (\mathcal{M}, ρ) , a set-valued mapping $F : \mathcal{M} \rightarrow \mathcal{PG}_L(\mathbf{R}^2)$, and a constant $\lambda > 0$, produces the following two outcomes:*

OUTCOME 1 (“No-Go”): *The algorithm guarantees that there does not exist a Lipschitz mapping $f : \mathcal{M} \rightarrow \mathbf{R}^2$ with Lipschitz constant at most λ such that $f(x) \in F(x)$ for all $x \in \mathcal{M}$.*

OUTCOME 2 (“Success”): *The algorithm returns a Lipschitz mapping $f : \mathcal{M} \rightarrow \mathbf{R}^2$, with Lipschitz constant at most 3λ such that $f(x) \in F(x)$ for all $x \in \mathcal{M}$.*

This Selection Algorithm requires at most $CL^2 N^2$ computer operations and at most $CL^2 N^2$ units of computer memory. Here $C > 0$ is a universal constant.

Proof. We know that for every $x \in \mathcal{M}$ the polygon $F(x)$ can be represented as an intersection of at most L half-planes. We denote this family of half-planes by $\mathcal{H}_F(x)$. Thus, $\#\mathcal{H}_F(x) \leq L$ and

$$F(x) = \cap\{H : H \in \mathcal{H}_F(x)\} \quad \text{for every } x \in \mathcal{M}. \quad (6.33)$$

Let us introduce a new pseudometric space $\widetilde{\mathfrak{M}} = (\widetilde{\mathcal{M}}, \tilde{\rho})$ whose elements are all couples $u = (x, H)$ where $x \in \mathcal{M}$ and $H \in \mathcal{H}_F(x)$. We define a pseudometric $\tilde{\rho}$ on $\widetilde{\mathcal{M}}$ by letting

$$\tilde{\rho}(u, u') = \rho(x, x') \quad \text{for every } u = (x, H), u' = (x', H') \in \widetilde{\mathcal{M}}. \quad (6.34)$$

Finally, we define a new set-valued mapping $\widetilde{F} : \widetilde{\mathcal{M}} \rightarrow \mathcal{HP}(\mathbf{R}^2)$ by letting

$$\widetilde{F}((x, H)) = H \quad \text{provided } x \in \mathcal{M} \text{ and } H \in \mathcal{H}_F(x). \quad (6.35)$$

In particular,

$$\tilde{\rho}((x, H), (x, H')) = 0 \quad \text{for every } x \in \mathcal{M} \text{ and } H, H' \in \mathcal{H}_F(x).$$

It is also clear that

$$\#\widetilde{\mathcal{M}} = \sum_{x \in \mathcal{M}} \#\mathcal{H}_F(x) \leq L \cdot N.$$

Let us apply Theorem 1.3 to the constant λ , the pseudometric space $\widetilde{\mathfrak{M}} = (\widetilde{\mathcal{M}}, \tilde{\rho})$ and the set-valued mapping $\widetilde{F} : \widetilde{\mathcal{M}} \rightarrow \mathcal{HP}(\mathbf{R}^2)$. Theorem 1.3 tells us that the Projection Algorithm with these inputs produces one of the two following outcomes:

(• $\tilde{1}$) OUTCOME (“No Go”): We guarantee that there does not exist $\tilde{f} \in \text{Lip}(\widetilde{\mathcal{M}}, \tilde{\rho})$ with Lipschitz seminorm $\|\tilde{f}\|_{\text{Lip}(\widetilde{\mathcal{M}}, \tilde{\rho})} \leq \lambda$ such that $\tilde{f}(u) \in \widetilde{F}(u)$ for all $u \in \widetilde{\mathcal{M}}$.

(• $\tilde{2}$) OUTCOME (“Success”): The algorithm produces a mapping $\tilde{f} : \widetilde{\mathcal{M}} \rightarrow \mathbf{R}^2$ with Lipschitz seminorm $\|\tilde{f}\|_{\text{Lip}(\widetilde{\mathcal{M}}, \tilde{\rho})} \leq 3\lambda$ such that $\tilde{f}(u) \in \widetilde{F}(u)$ for all $u \in \widetilde{\mathcal{M}}$.

Furthermore, this algorithm requires at most $C(\#\widetilde{\mathcal{M}})^2$ computer operations and at most $C(\#\widetilde{\mathcal{M}})^2$ units of computer memory. Here $C > 0$ is an absolute constant.

Our algorithm produces OUTCOME 1 (“No-Go”) if *the outcome (• $\tilde{1}$) holds*, and it produces OUTCOME 2 (“Success”) provided *the outcome (• $\tilde{2}$) holds*.

Prove that these outcomes satisfy the requirements of Theorem 6.10. Indeed, let us show that in the case of the outcome $(\bullet\tilde{1})$ the algorithm guarantees that there does not exist a Lipschitz mapping $f : \mathcal{M} \rightarrow \mathbf{R}^2$ with Lipschitz constant at most λ such that $f(x) \in F(x)$ for all $x \in \mathcal{M}$.

Suppose that it is not true, i.e., there exist a mapping $f \in \text{Lip}(\mathcal{M})$ with

$$\|f\|_{\text{Lip}(\mathcal{M})} \leq \lambda \quad (6.36)$$

such that $f(x) \in F(x)$ for all $x \in \mathcal{M}$. This mapping generates a new mapping $\tilde{f} : \widetilde{\mathcal{M}} \rightarrow \mathbf{R}^2$ defined as follows:

$$\tilde{f}((x, H)) = f(x) \quad \text{for every } x \in \mathcal{M} \text{ and } H \in \mathcal{H}_F(x).$$

Then, thanks to (6.33), (6.34) and (6.35), for every $x \in \mathcal{M}$ and $H \in \mathcal{H}_F(x)$, we have

$$\tilde{f}((x, H)) = f(x) \in F(x) = \cap\{\mathcal{H} : \mathcal{H} \in \mathcal{H}_F(x)\} \subset H = \widetilde{F}((x, H))$$

which proves that $\tilde{f}(u) \in \widetilde{F}(u)$ for every $u \in \widetilde{\mathcal{M}}$.

Furthermore, let $u = (x, H), u' = (x', H') \in \widetilde{\mathcal{M}}$, i.e., $H \in \mathcal{H}_F(x)$ and $H' \in \mathcal{H}_F(x')$. Then, thanks to (6.36),

$$\|\tilde{f}(u) - \tilde{f}(u')\| = \|\tilde{f}((x, H)) - \tilde{f}((x, H'))\| = \|f(x) - f(x')\| \leq \lambda \rho(x, x') = \lambda \tilde{\rho}(u, u').$$

Hence, $\|\tilde{f}\|_{\text{Lip}(\widetilde{\mathcal{M}}, \tilde{\rho})} \leq \lambda$.

The existence of the mapping \tilde{f} with these properties contradicts the requirement of the outcome $(\bullet\tilde{1})$ proving that the statement of OUTCOME 1 (“No-Go”) of Theorem 6.10 holds.

Let us see that if the outcome $(\bullet\tilde{2})$ holds, then OUTCOME 2 (“Success”) of Theorem 6.10 holds as well. In fact, let us define a mapping $f : \mathcal{M} \rightarrow \mathbf{R}^2$ as follows: given $x \in \mathcal{M}$ let us fix some $H \in \mathcal{H}_F(x)$ and set

$$f(x) = \tilde{f}((x, H)). \quad (6.37)$$

Prove that f is well defined on \mathcal{M} , i.e., the value of $f(x)$ does not depend on the choice of the half-plane $H \in \mathcal{H}_F(x)$ in definition (6.37).

Indeed, thanks to (6.34), for every $H, H' \in \mathcal{H}_F(x)$, we have

$$\|\tilde{f}((x, H)) - \tilde{f}((x, H'))\| \leq 3\lambda \tilde{\rho}(u, u') = 3\lambda \rho(x, x) = 0.$$

Thus, $\tilde{f}((x, H)) = \tilde{f}((x, H'))$ for every $H, H' \in \mathcal{H}_F(x)$ proving that the value of $f(x)$ in (6.37) is well defined.

Furthermore, given $x, x' \in \mathcal{M}$ and $H \in \mathcal{H}_F(x)$ and $H' \in \mathcal{H}_F(x')$, we have

$$\|f(x) - f(x')\| = \|\tilde{f}((x, H)) - \tilde{f}((x, H'))\| \leq 3\lambda \tilde{\rho}((x, H), (x, H')) = 3\lambda \rho(x, x')$$

proving that $\|f\|_{\text{Lip}(\mathcal{M})} \leq 3\lambda$.

Finally, let $x \in \mathcal{M}$ and $H \in \mathcal{H}_F(x)$. Then,

$$f(x) = \tilde{f}((x, H)) \in \widetilde{F}((x, H)) = H.$$

From this and (6.33), we have

$$f(x) \in \cap\{H : H \in \mathcal{H}_F(x)\} = F(x)$$

proving that f is a selection of F .

Thus, f satisfies all the requirements of **OUTCOME 2 (“Success”)** of Theorem 6.10 proving that this outcome holds.

It remains to note that the Projection Algorithm produces the outcomes $(\bullet\tilde{1})$ and $(\bullet\tilde{2})$ using the work and storage at most $C(\#\mathcal{M})^2$ where $C > 0$ is an absolute constant. Therefore, the Selection Algorithm proposed in this proof produces **OUTCOME 1 (“No-Go”)** and **OUTCOME 2 (“Success”)** using the work and storage at most $CL^2 N^2$.

The proof of the theorem is complete. ■

Let $\mathcal{PG}(\mathbf{R}^2)$ be the family of all (not necessarily bounded) polygons in \mathbf{R}^2 . Each of these polygons is determined by a finite number of linear constraints. Given a polygon $P \in \mathcal{PG}(\mathbf{R}^2)$ we let $L(P)$ denote the number of linear constraints determining this polygon.

The method of proving Theorem 6.10 allows us to show that the following, slightly more general version of this theorem holds.

Theorem 6.11 *There exists an algorithm which, given a finite pseudometric space (\mathcal{M}, ρ) , a set-valued mapping $F : \mathcal{M} \rightarrow \mathcal{PG}(\mathbf{R}^2)$, and a constant $\lambda > 0$, produces **OUTCOME 1 (“No-Go”)** and **OUTCOME 2 (“Success”)** described in Theorem 6.10.*

This Selection Algorithm requires at most CM^2 work and storage where $C > 0$ is a universal constant and

$$M = \sum_{x \in \mathcal{M}} L(F(x))$$

is the total number of linear constraints determining the polygons $\{F(x) : x \in \mathcal{M}\}$.

7. Further results and comments

7.1 Generalizations of the PA: sets determined by polynomial inequalities and more.

Theorem 1.3 can be generalized to wider families of convex subsets in \mathbf{R}^2 than the family $\mathcal{HP}(\mathbf{R}^2)$ of all closed half-planes. (We study this problem in the forthcoming paper [67].) Let us discuss some of these generalizations.

Let \mathfrak{T} be a family of convex closed subsets of \mathbf{R}^2 . Analyzing the projection algorithm for family \mathfrak{T} , we conclude that **STEP 1** and **STEP 2** are the most complicated parts of this algorithm.

For instance, at **STEP 2** we construct the rectangular hull $\mathcal{H}[F^{[1]}[x : \lambda]]$ of the set $F^{[1]}[x : \lambda]$ defined by formula (1.4). The problem of computing of this rectangular is equivalent to the following problem of convex optimization:

$$\text{maximize } z = u_1 \quad \text{subject to } u = (u_1, u_2) \in \bigcap_{y \in \mathcal{M}} [F(y) + \lambda \rho(x, y) Q_0]. \quad (7.1)$$

Here $\lambda > 0$, $x \in \mathcal{M}$ and the set-valued mapping $F : \mathcal{M} \rightarrow \mathfrak{T}$ defined on the pseudometric space (\mathcal{M}, ρ) are the input data.

As we have noted in Section 6.2, if $\mathfrak{T} = \mathcal{HP}(\mathbf{R}^2)$ is the family of all closed half-planes in \mathbf{R}^2 , then, thanks to the results of N. Megiddo [52] and M. E. Dyer [21], the program (7.1) can be solved in $O(m)$ time provided $\#\mathcal{M} = m$. See Theorem 6.2. This allows us, for each $x \in \mathcal{M}$, to construct the set $\mathcal{H}[F^{[1]}[x : \lambda]]$ in $O(m)$ time. Applying this algorithm to all elements $x \in \mathcal{M}$, we build the family of rectangles $\{\mathcal{H}[F^{[1]}[x : \lambda]] : x \in \mathcal{M}\}$ using at most $O(m^2)$ computer operations.

Thus, if we want to have at **STEP 2** similar estimates for a set \mathfrak{T} of convex closed subsets in \mathbf{R}^2 , we need an analog of Theorem 6.2 for the convex program (7.1).

A family of sets \mathfrak{T} with this property was described by J. Puerto, A. M. Rodríguez-Chía and A. Tamir in [59]. Let k be a positive integer. Let us consider the following convex programming problem, defined on \mathbf{R}^2 :

$$\text{maximize } z = u_1 \quad \text{subject to } u = (u_1, u_2) \in \bigcap_{i=1}^m S_i. \quad (7.2)$$

Here

$$S_i = \{x \in \mathbf{R}^2 : P_i(x) \leq 0\}, \quad i = 1, \dots, m,$$

where each $P_i = P_i(x_1, x_2)$, $i = 1, \dots, m$, is a *convex* polynomial on \mathbf{R}^2 of degree at most k . It is shown in [59, Section 3] that this problem can be solved in $O(m)$ time. The proof of this statement relies on the results of the works [2] and [15].

Basing on this property, we can show that the convex program (7.1) also can be solved in $O(m)$ time provided each set $S \in \mathfrak{T}$ can be represented in the form

$$S = \{x \in \mathbf{R}^2 : P(x) \leq 0\}$$

where P is a *convex* polynomial on \mathbf{R}^2 of degree at most k .

We let $\mathcal{PC}_k(\mathbf{R}^2)$ denote the family of all such sets S . For instance, if $k = 2$, as a family \mathfrak{T} we can take the family of all disks or all ellipses in \mathbf{R}^2 .

Another approach to the problem (7.2) with the family of convex sets even wider than $\mathcal{PC}_k(\mathbf{R}^2)$ was suggested by M. E. Dyer in [23].

T. M. Chan [9] proposed an approach that solves the problem (7.2) for a very large family $\mathcal{S} = \{S_1, \dots, S_m\}$ of convex closed subsets of \mathbf{R}^2 , but with somewhat more computer operations.

Let us introduce two operations with the sets from \mathcal{S} :

(D1) Given $S, S' \in \mathcal{S}$ find the ends the points $v_{min}, v_{max} \in S \cap S'$ such that the projections of the line segment $[v_{min}, v_{max}]$ and the intersection $S \cap S'$ onto the axis Ox_1 coincide;

(D2) Given $S \in \mathcal{S}$ and a straight line $\ell \subset \mathbf{R}^2$, compute the line segment $S \cap \ell$. We refer to (D1) and (D2) as *primitive operations*.

The main result of [9] states that *the convex program (7.2) in \mathbf{R}^2 with m constraints can be solved using $O(m \log m)$ primitive operations.*

Applying this result to the optimization problem (7.1) we can essentially extend the class of sets in the Projection Algorithm. But in this version, this algorithm will require at most $CN^2 \log N$ computer operations (and CN^2 units of computer memory). Here $C > 0$ is an absolute constant.

7.2 A polynomial algorithm for the sharp selection problem.

Let $\lambda > 0$, an N -point pseudometric space (M, ρ) , and a set-valued mapping $F : M \rightarrow \mathcal{HP}(\mathbf{R}^2)$ be the same as in Theorem 1.3. We ask the following question: *Does there exist a polynomial algorithm which provides the same outcomes as in this theorem, but with the estimate $\|f\|_{\text{Lip}(M)} \leq \lambda$ instead of inequality (1.2)?*

We refer to this problem as *the sharp selection problem*. The next theorem gives an affirmative answer to this problem.

Theorem 7.1 *There exists a polynomial algorithm which receives as input a real number $\lambda > 0$, a finite pseudometric space (\mathcal{M}, ρ) with $\#\mathcal{M} = N$, and a set-valued mapping $F : \mathcal{M} \rightarrow \mathcal{HP}(\mathbf{R}^2)$, and produces the same outcomes as in Theorem 1.3 (i.e., (“Success”) and (“No Go”)) with the following refinement of the outcome (“Success”): in this case the algorithm produces a Lipschitz selection f of F with Lipschitz seminorm $\|f\|_{\text{Lip}(\mathcal{M})} \leq \lambda$.*

This algorithm requires at most $CN^4 \log N$ computer operations and at most CN^3 units of computer memory. Here $C > 0$ is an absolute constant.

Proof. This result is immediate from polynomial algorithms for solving systems of linear inequalities of order $m \times n$ with at most two variables per inequality. N. Megiddo [51] was the first to discover such an algorithm requiring $O(mn^3 \log m)$ computer operations. Then E. Cohen and N. Megiddo [16] obtained a new $O(mn^2(\log m + \log^2 n))$ running time algorithm. The fastest algorithm known so far for this problem is by D. S. Hochbaum and J. Naor [43] with $O(mn^2 \log m)$ running time. Note also that all these algorithms require $O(mn)$ units of computer memory.

Let us apply these algorithms to our problem. This problem can be formulated as follows: construct an algorithm which, in polynomial time, given $\lambda > 0$, (\mathcal{M}, ρ) with $\#\mathcal{M} = N$ and $F : \mathcal{M} \rightarrow \mathcal{HP}(\mathbf{R}^2)$, either tells us that no exists a Lipschitz selection of F with Lipschitz seminorm at most λ (outcome (“No Go”)) or produces such a selection (outcome (“Success”)).

In turn, this problem is equivalent the feasibility of the following linear programming problem: Let $\mathcal{M} = \{x_1, \dots, x_N\}$ and let

$$F(x_i) = \{(u, v) \in \mathbf{R}^2 : a_i u + b_i v \leq c_i\}, \quad i = 1, \dots, N.$$

Let us consider the system of linear inequalities with respect to the $2N$ variables $u_1, v_1, \dots, u_N, v_N$ defined as follows:

$$\begin{aligned} \text{Linear Program:} \quad & a_i u_i + b_i v_i \leq c_i, \quad i = 1, \dots, N, & (7.3) \\ & u_i - u_j \leq \lambda \rho(x_i, x_j), \quad v_i - v_j \leq \lambda \rho(x_i, x_j), \\ & u_j - u_i \leq \lambda \rho(x_i, x_j), \quad v_j - v_i \leq \lambda \rho(x_i, x_j), \\ & i, j = 1, \dots, N. \end{aligned}$$

This is a system of $m = N + N(N + 1) = N(N + 2)$ linear inequalities with respect to $n = 2N$ variables $u_1, v_1, \dots, u_N, v_N$ where each inequality involves at most two variables.

Our goal is either find a point that satisfies the all inequalities (the feasible solution of the system) or conclude that no such point exists (i.e., Linear Program (7.3) is infeasible).

Applying to this Linear Program the algorithm suggested by D. S. Hochbaum and J. Naor in [43], we conclude that this algorithm solves this problem in

$$O(mn^2 \log m) = O(N(N + 2)(2N)^2 \log N(N + 2)) = O(N^4 \log N)$$

running time. Also, this algorithm requires

$$O(mn) = O(N(N + 2)(2N)) = O(N^3)$$

units of computer memory.

The proof of the theorem is complete. ■

Remark 7.2 Let us note that applying to Linear Program (7.3) general-purpose linear programming we obtain algorithms with running time depending on N exponentially.

For instance, as we noted in Remark 6.4, the fastest of them known to the moment is the algorithm presented in T. M. Chan [13]. For solving linear programs with m constraints and n variables this algorithm requires $O(n)^{n/2}(\log n)^{3n}m$ running time. In case of Linear Program (7.3), we have $n = 2N$ and $m = N(N + 2)$ which leads to an algorithm with the exponential growth on N of the number of computer operations. ◀

7.3 Distance oracle and intrinsic metrics.

In this section we give some remarks related to the following version of Problem 1.1 formulated by C. Fefferman [31, Chapter 8.7].

Problem 7.3 Let D, L be positive integers. Suppose that (\mathcal{M}, ρ) is an N -point metric space. Given $x, y \in \mathcal{M}$, we suppose that a *distance oracle* tells us $\rho(x, y)$ (perhaps only up to a universal constant factor). For each $x \in \mathcal{M}$, suppose we are given a convex polytope $F(x) \subset \mathbf{R}^D$, defined by at most L linear constraints.

We would like to compute a function $f : \mathcal{M} \rightarrow \mathbf{R}^D$ satisfying $f(x) \in F(x)$ for all $x \in \mathcal{M}$, with Lipschitz seminorm as small as possible up to a factor $C(D)$. How many computer operations does it take?

Our remarks concern the concept of *the distance oracle* mentioned in Problem 7.3. In many cases, in applications, we do not know the true value of the distance $\rho(x, y)$ between any two elements $x, y \in \mathcal{M}$ (even up to an absolute constant factor). Instead of this exact value of $\rho(x, y)$, we often know only a certain upper bound for this distance. In other words, our input is a non-negative weight function $W : \mathcal{M} \times \mathcal{M} \rightarrow [0, +\infty]$ defined on pairs of elements of \mathcal{M} . We assume that

$$W(x, x) = 0 \quad \text{and} \quad W(x, y) = W(y, x) \quad \text{for all } x, y \in \mathcal{M}. \quad (7.4)$$

Note that, in general, W does not necessarily satisfy the triangle inequality. We refer to the function W as a “pre-metric” on \mathcal{M} .

The following selection problem arises in various applications.

Problem 7.4 Let W be non-negative weight function on \mathcal{M} satisfying conditions (7.4). For each $x \in \mathcal{M}$, suppose we are given a convex closed set $F(x) \subset \mathbf{R}^D$. Find an efficient algorithm for constructing a nearly-optimal selection f of F which is “Lipschitz” with respect to W , i.e., $f(x) \in F(x)$ on \mathcal{M} , and for some constant $\lambda \geq 0$ we have

$$\|f(x) - f(y)\| \leq \lambda W(x, y) \quad \text{for all } x, y \in \mathcal{M}. \quad (7.5)$$

Here “nearly-optimal” means “with λ as small as possible” (up to a universal constant factor).

Let us note that Problem 7.4 easily reduces to an equivalent problem where the pre-metric W is replaced with a certain *pseudometric* ρ_W generated by W . The pseudometric ρ_W is the classical object of the graph theory known in the literature as *the intrinsic pseudometric* induced by the pre-metric W . We recall that ρ_W is defined as follows: given $x, y \in \mathcal{M}$, we set

$$\rho_W(x, y) = \inf_{\{u_0, \dots, u_k\}} \sum_{i=0}^{k-1} W(u_i, u_{i+1}), \quad (7.6)$$

where the infimum is taken over all subsets $\{u_0, \dots, u_k\} \subset \mathcal{M}$ such that $u_0 = x$ and $u_k = y$.

Clearly, ρ_W is a pseudometric. It is also clear that a function $f : \mathcal{M} \rightarrow \mathbf{R}^D$ satisfies inequality (7.5) if and only if f is λ -Lipschitz with respect to ρ_W , i.e.,

$$\|f(x) - f(y)\| \leq \lambda \rho_W(x, y) \quad \text{for every } x, y \in \mathcal{M}.$$

This observation shows that the problems of constructing a nearly optimal Lipschitz selection with respect to W and with respect to ρ_W are equivalent. Therefore, at the preparatory stage of our algorithm we can construct ρ_W , and then, at the following stages of the algorithm, deal with the pseudometric ρ_W rather than the pre-metric W .

The problem of computing the intrinsic metric ρ_W is one of the classic problems in the algorithmic graph theory. Let us recall something of the history of this problem.

Let $G = (V, E)$ be a graph with the sets of the nodes V and the sets of the edges E . Let $n = \#V$ and $m = \#E$ be the number of nodes and the number of edges respectively. Finally, let $W : E \rightarrow [0, \infty)$ be a length function. (In particular, in these settings, the intrinsic pseudometric $\rho_W(x, y)$ induced by W (see (7.6)) is the infimum of the lengths of all paths from x to y in the complete graph with the nodes in \mathcal{M} .)

Let us note several results related to two problems known in the literature as the *Single Source Shortest Path problem*, SSSP in short, and the *All Pairs Shortest Path problem*, APSP in short.

The SSSP problem is the problem of finding the length of the shortest path from a single-source node $x \in V$ to all other nodes $y \in V$. The APSP is to find, for each pair of nodes, $x, y \in V$, the length of the shortest path from x to y . (A length of a path is defined as the sum of the length of its edges.) We refer the reader to the papers [4, 10–12, 14, 18, 20, 38, 40–42, 48, 50, 68, 71] and references therein for numerous results on the SSSP and APSP algorithms.

Among the multitude of results devoted to the SSSP we mention the classical Dijkstra's algorithm [20] which solves the SSSP in $O(n^2)$ time; see also Bellman's algorithm [4] with running time $O(mn)$. Let us note various improvements of these result: $O(m)$ [68] ($W(x)$, $x \in V$, are non-negative integers), $O((m+n) \log n)$ [48], $O(m+n \log n)$ [40] and $O(m+n \log n / \log \log n)$ [41].

There is also an extensive literature devoted to different aspects of the APSP problem. We refer the reader to the papers [10, 71] for the detailed reviews of the results obtained in this directions. Here we mention the algorithms with the following running time: $O(n^3)$ [38, 69] (the classical Floyd - Warshall algorithm), $O(n^3(\log \log n / \log n)^{1/3})$ [39], $O(n^3 / \log n)$ [10], $O(n^3(\log \log n)^3 / \log^2 n)$ [11], $O(n^3 \log \log n / \log^2 n)$ [42]. Finally, the algorithm suggested by T. M. Chan and R. Williams [14] improves the running time to $O(n^3 / 2^{30(\log n)^{1/2}})$.

For the case of the *sparse* graphs (i.e., the graphs in which the number of edges is much less than the maximal number of edges) we mention the algorithms of the papers [48, 49] with the running time $O(mn + n^2 \log n)$, [54] with $O(mn \log \alpha(m, n))$ (here α is the inverse-Ackermann function), and [55] with the running time $O(mn + n^2 \log \log n)$. See also [12] for a faster algorithm which provides running time $o(mn)$.

Remark 7.5 In Problem 7.4, sparse graphs appear in a natural way for a weight W that takes the value $+\infty$. Indeed, we introduce a graph structure on \mathcal{M} , allowing $x, y \in \mathcal{M}$ to be joined by an edge if and only if $W(x, y) < +\infty$. Clearly, in definition (7.6) of the intrinsic pseudometric ρ_W it suffices to consider only paths $\{u_0, \dots, u_k\}$ from x to y such that u_i and u_{i+1} are joined by an edge for all $i = 0, \dots, k-1$. In other words, ρ_W is the intrinsic pseudometric induced by the graph $G = (\mathcal{M}, E)$ and the weight function $W|_E$ where $E = \{(x, y) \in \mathcal{M} \times \mathcal{M} : W(x, y) < +\infty\}$.

It may happen that G is a sparse graph, i.e., $\#E \ll n(n+1)/2$.

This will allow us to use the above-mentioned fast APSP algorithms for sparse graphs when constructing the pseudometric ρ_W . ◀

References

- [1] P. K. Agarwal, M. Sharir, S. Toledo, An Efficient Multi-Dimensional Searching Technique and Its Applications. Technical Report CS-1993-20. Department of Computer Science, Duke University, 1993.
- [2] N. Amenta, Helly-type theorems and generalized linear programming. ACM Symposium on Computational Geometry (San Diego, CA, 1993). *Discrete Comput. Geom.* 12 (1994), no. 3, 241–261.
- [3] J.-P. Aubin, H. Frankowska, Set-valued analysis, *Systems & Control: Foundations & Applications*, 2. Birkhauser Boston, 1990.
- [4] R. Bellman, On a routing problem. *Quart. Appl. Math.* 16 (1958), 87–90.
- [5] Y. Benyamini, J. Lindenstrauss, Geometric nonlinear functional analysis, Vol. 1, in: *American Mathematical Society Colloquium Publications*, 48. American Mathematical Society, Providence, RI, 2000. xii+488 pp.
- [6] H. Brönnimann, B. Chazelle, J. Matoušek, Product range spaces, sensitive sampling, and derandomization. *SIAM J. Comput.* 28 (1999), no. 5, 1552–1575.
- [7] Yu. Brudnyi, P. Shvartsman, Generalizations of Whitney’s extension theorem. *Internat. Math. Res. Notices* (1994), no. 3, 129–139.
- [8] Yu. Brudnyi, P. Shvartsman, Whitney Extension Problem for Multivariate $C^{1,\omega}$ -functions. *Trans. Amer. Math. Soc.* 353 (2001), no. 6, 2487–2512.
- [9] T. M. Chan, Deterministic algorithms for 2-d convex programming and 3-d online linear programming. *J. Algorithms* 27 (1998), no. 1, 147–166.
- [10] T. M. Chan, All-pairs shortest paths with real weights in $O(n^3 / \log n)$ time. *Algorithmica* 50 (2008), no. 2, 236–243.
- [11] T. M. Chan, More algorithms for all-pairs shortest paths in weighted graphs. *SIAM J. Comput.* 39 (2010), no. 5, 2075–2089.
- [12] T. M. Chan, All-pairs shortest paths for unweighted undirected graphs in $o(mn)$ time. *ACM Trans. Algorithms* 8 (2012), no. 4, Art. 34, 17 pp.
- [13] T. M. Chan, Improved deterministic algorithms for linear programming in low dimensions. *ACM Trans. Algorithms* 14 (2018), no. 3, Art. 30, 10 pp.
- [14] T. M. Chan, R. Williams, Deterministic APSP, orthogonal vectors, and more: quickly derandomizing Razborov-Smolensky. *ACM Trans. Algorithms* 17 (2021), no. 1, Art. 2, 14 pp.
- [15] B. Chazelle, J. Matoušek, On linear-time deterministic algorithms for optimization problems in fixed dimension. *J. Algorithms* 21 (1996), no. 3, 579–597.
- [16] E. Cohen, N. Megiddo, Improved algorithms for linear inequalities with two variables per inequality. *SIAM J. Comput.* 23 (1994), no. 6, 1313–1347.
- [17] K. L. Clarkson, Linear programming in $O(n \times 3^{d^2})$ time. *Inform. Process. Lett.* 22 (1986), no. 1, 21–24.

- [18] T. H. Cormen, C. E. Leiserson, R. L. Ronald, C. Stein, Introduction to algorithms. Third edition. MIT Press, Cambridge, MA, 2009.
- [19] L. Danzer, B. Grünbaum, V. Klee, Helly's Theorem and its relatives. in: AMS Symposium on Convexity, Seattle, Proceedings of Symposium on Pure Mathematics, Vol. 7, Amer. Math. Soc., Providence, RI, 1963, pp. 101–180.
- [20] E. W. Dijkstra, A note on two problems in connexion with graphs. Numer. Math. 1 (1959), 269–271.
- [21] M. E. Dyer, Linear time algorithms for two- and three-variable linear programs. SIAM J. Comput. 13 (1984), no. 1, 31–45.
- [22] M. E. Dyer, On a multidimensional search problem and its application to the Euclidean one-centre problem, SIAM J. Comput. 15 (1986), no. 3, 725–738.
- [23] M. E. Dyer, A Class of Convex Programs with Applications to Computational Geometry. 8th Annual Computational Geometry, 6/92, Berlin, Germany, 9–15.
- [24] M. Dyer, B. Gärtner, N. Megiddo, E. Welzl, Linear programming. Handbook of discrete and computational geometry, 1291–1309, CRC Press, Boca Raton, FL, 2018.
- [25] C. Fefferman, A sharp form of Whitney's extension theorem. Ann. of Math. (2) 161 (2005), no. 1, 509–577.
- [26] C. Fefferman, Whitney extension problems and interpolation of data. Bull. Amer. Math. Soc. 46 (2009), no. 2, 207–220.
- [27] C. Fefferman, Fitting a C^m -smooth function to data. III. Ann. of Math. (2) 170 (2009), no. 1, 427–441.
- [28] C. Fefferman, Smooth interpolation of data by efficient algorithms. Excursions in harmonic analysis. Vol. 1, 71–84, Appl. Numer. Harmon. Anal., Birkhäuser/Springer, New York, 2013.
- [29] C. Fefferman, Unsolved Problems, 11th Whitney Extension Problems Workshop, Trinity College Dublin, August 13-17, 2019, <https://cms-math.net.technion.ac.il/open-problems-whitney/>
- [30] C. Fefferman, Introduction to Whitney problems. 14th Whitney Problems Online Workshop, 2021, <https://vimeo.com/showcase/8488319/video/552948900>, password: Whitney2021
- [31] C. Fefferman, A. Israel, Fitting Smooth Functions to Data. CBMS Regional Conference Series in Mathematics, 135. American Mathematical Society, Providence, RI, 2020. xi+160 pp.
- [32] C. Fefferman, A. Israel, G. K. Luli, Finiteness principles for smooth selection. Geom. Funct. Anal. 26 (2016), no. 2, 422–477.
- [33] C. Fefferman, F. Jiang, G. K. Luli, C^2 -interpolation with range restriction. Rev. Mat. Iberoam. 39 (2023), no. 2, 649–710.
- [34] C. Fefferman, B. Klartag, Fitting a C^m -smooth function to data. I., Ann. of Math. (2) 169 (2009), no. 1, 315–346.
- [35] C. Fefferman, B. Klartag, Fitting a C^m -smooth function to data. II., Rev. Mat. Iberoam. 25 (2009), no. 1, 49–273.

- [36] C. Fefferman, B. Pegueroles, Efficient Algorithms for Approximate Smooth Selection. *J. Geom. Anal.* 31 (2021), no. 7, 6530–6600.
- [37] C. Fefferman, P. Shvartsman, Sharp finiteness principles for Lipschitz selections. *Geom. Funct. Anal.* 28 (2018), no. 6, 1641–1705.
- [38] R. W. Floyd, Algorithm 97: Shortest path. *Comm. ACM*, 5 (1962), 345.
- [39] M. L. Fredman, New bounds on the complexity of the shortest path problem. *SIAM J. Comput.* 5 (1976), no. 1, 83–89.
- [40] M. L. Fredman, R. E. Tarjan, Fibonacci heaps and their uses in improved network optimization algorithms. *J. Assoc. Comput. Mach.* 34 (1987), no. 3, 596–615.
- [41] M. L. Fredman, D. E. Willard, Surpassing the information-theoretic bound with fusion trees. *Proceedings of the 22nd Annual ACM Symposium on Theory of Computing (Baltimore, MD, 1990)*. *J. Comput. System Sci.* 47 (1993), no. 3, 424–436.
- [42] Y. Han, T. Takaoka, An $O(n^3 \log \log n / \log^2 n)$ time algorithm for all pairs shortest paths. *J. Discrete Algorithms* 38/41 (2016), 9–19.
- [43] D. S. Hochbaum, J. Naor, Simple and fast algorithms for linear and integer programs with two variables per inequality. *SIAM J. Comput.* 23 (1994), no. 6, 1179–1192.
- [44] F. Jiang, G. K. Luli, Algorithms for nonnegative $C^2(\mathbb{R}^2)$ interpolation. *Adv. Math.* 385 (2021), Paper No. 107756, 43 pp.
- [45] F. Jiang, G. K. Luli, K. O’Neill, Smooth selection for infinite sets. *Adv. Math.* 407 (2022), Paper No. 108566, 62 pp.
- [46] F. Jiang, G. K. Luli, K. O’Neill, On the Shape Fields Finiteness Principle. *IMRN* 2022 (2022), no. 23, 18895–18918.
- [47] F. Jiang, C. Lian, Y. Liang, G. K. Luli, Univariate range-restricted C^2 interpolation algorithms. *J. Comput. Appl. Math.* 425 (2023), Paper No. 115040, 19 pp.
- [48] D. B. Johnson, Efficient algorithms for shortest paths in sparse networks. *J. Assoc. Comput. Mach.* 24 (1977), no. 1, 1–13.
- [49] D. R. Karger, D. Koller, S. J. Phillips, Finding the hidden path: time bounds for all-pairs shortest paths. *SIAM J. Comput.* 22 (1993), no. 6, 1199–1217.
- [50] M. J. Kashyop, T. Nagayama, K. Sadakane, Faster algorithms for shortest path and network flow based on graph decomposition. *J. Graph Algorithms Appl.* 23 (2019), no. 5, 781–813.
- [51] N. Megiddo, Towards a genuinely polynomial algorithm for linear programming. *SIAM J. Comput.* 12 (1983), no. 2, 347–353.
- [52] N. Megiddo, Linear-time algorithms for linear programming in \mathbf{R}^3 and related problems. *SIAM J. Comput.* 12 (1983), no. 4, 759–776.
- [53] N. Megiddo, Linear programming in linear time when dimension is fixed. *J. Assoc. Comput. Mach.* 31 (1984), no. 1, 114–127.
- [54] S. Pettie, A new approach to all-pairs shortest paths on real-weighted graphs. *Automata, languages and programming. Theoret. Comput. Sci.* 312 (2004), no. 1, 47–74.

- [55] S. Pettie, V. A. Ramachandran, A shortest path algorithm for real-weighted undirected graphs. *SIAM J. Comput.* 34 (2005), no. 6, 1398–1431.
- [56] K. Przesławski, L. E. Rybinski, Concepts of lower semicontinuity and continuous selections for convex valued multifunctions. *J. Approx. Theory* 68 (1992), no. 3, 262–282.
- [57] K. Przesławski, D. Yost, Continuity properties of selectors and Michael’s theorem. *Mich. Math. J.* 36 (1989), no. 1, 113–134.
- [58] K. Przesławski, D. Yost, Lipschitz Retracts, Selectors and Extensions. *Mich. Math. J.* 42 (1995), no. 3, 555–571.
- [59] J. Puerto, A. M. Rodríguez-Chía, A. Tamir, On the planar piecewise quadratic 1-center problem. *Algorithmica* 57 (2010), no. 2, 252–283.
- [60] P. A. Shvartsman, Traces of functions of Zygmund class. (Russian) *Sibirsk. Mat. Zh.* 28 (1987), no. 5, 203–215; English transl. in: *Sib. Math. J.* 28 (1987), 853–863.
- [61] P. Shvartsman, On Lipschitz selections of affine-set valued mappings. *Geom. Funct. Anal.* 11 (2001), no. 4, 840–868.
- [62] P. Shvartsman, Lipschitz selections of set-valued mappings and Helly’s theorem. *J. Geom. Anal.* 12 (2002), no. 2, 289–324.
- [63] P. Shvartsman, Barycentric selectors and a Steiner-type point of a convex body in a Banach space. *J. Funct. Anal.* 210 (2004), no. 1, 1–42.
- [64] P. Shvartsman, The Whitney extension problem and Lipschitz selections of set-valued mappings in jet-spaces. *Trans. Amer. Math. Soc.* 360 (2008), no. 10, 5529–5550.
- [65] P. Shvartsman, The Core of a 2-Dimensional Set-Valued Mapping. Existence Criteria and Efficient Algorithms for Lipschitz Selections of Low Dimensional Set-Valued Mappings. arXiv: 2010.04540v2
- [66] P. Shvartsman, Existence Criteria for Lipschitz Selections of Set-Valued Mappings in \mathbf{R}^2 . arXiv:2306.14042v1
- [67] P. Shvartsman, Efficient Algorithms for Lipschitz Selections of Low Dimensional Set-Valued Mappings. (to appear)
- [68] M. Thorup, Undirected single-source shortest paths with positive integer weights in linear time. *J. ACM* 46 (1999), no. 3, 362–394.
- [69] S. Warshall, A theorem on boolean matrices. *J. Assoc. Comput. Mach.* 9 (1962), 11–12.
- [70] H. Whitney, Analytic extension of differentiable functions defined in closed sets. *Trans. Amer. Math. Soc.* 36 (1934), no. 1, 63–89.
- [71] R. Williams, From circuit complexity to faster all-pairs shortest paths. *SIAM Rev.* 63 (2021), no. 3, 559–582.