

Solvability of the Dirichlet problem using a weaker Carleson condition in the upper half plane

Martin Ulmer

April 3, 2025

Abstract

We study an elliptic operator $L := \operatorname{div}(A\nabla\cdot)$ on the upper half plane \mathbb{R}_+^2 . There are several conditions on the behavior of the matrix A in the transversal t -direction that yield $\omega \in A_\infty(\sigma)$. These include the t -independence condition, a mixed $L^1 - L^\infty$ condition on $\partial_t A$, and Dini-type conditions. We introduce an L^1 Carleson condition on $\partial_t A(x, t)$ that extends the class of elliptic operators for which we have $\omega \in A_\infty(\sigma)$, i.e. solvability of the L^p Dirichlet problem for some $1 < p < \infty$.

Contents

1	Introduction	2
1.1	Idea and sketch of the proof	4
2	Notations and Setting	5
3	Consequences of the Kato conjecture and properties of the heat semigroup	7
4	Hodge decomposition, Approximation function ρ and difference function θ	9
4.1	Hodge decomposition	9
4.2	The different parts of the derivatives of θ	11
5	Proof of <i>Theorem 1.5</i>	14
6	Square function bounds on the parts of the derivative of θ	19
6.1	The partial derivative parts $\partial_s \phi^s$ and $w_s^{(1)}$	19
6.2	The partial derivative part $w_s^{(2)}$	21
6.3	The partial derivative part w_t	25
7	L^2 estimates for square functions	31

1 Introduction

In this work let $\Omega := \mathbb{R}_+^2 := \mathbb{R} \times (0, \infty)$ be the upper half space and $L := \operatorname{div}(A\nabla \cdot)$ an uniformly elliptic operator, i.e. there exists $\lambda_0 > 0$ such that

$$\lambda_0 |\xi|^2 \leq \xi^T A(x, t) \xi \leq \lambda_0^{-1} |\xi|^2 \quad \text{for all } \xi \in \mathbb{R}^2, \quad (1.1)$$

and a.e. $(x, t) \in \mathbb{R}_+^2$. Furthermore assume that the coefficients of A are bounded and merely measurable. We are interested in the solvability of the L^p Dirichlet boundary value problem

$$\begin{cases} Lu = \operatorname{div}(A\nabla u) = 0 & \text{in } \Omega, \\ u = f & \text{on } \partial\Omega, \end{cases}$$

where $f \in L^p(\partial\Omega)$ (see *Definition 2.1*). It is well known that solvability for some $1 < p < \infty$ is equivalent to the elliptic measure ω_L lying in the Muckenhoupt space $A_\infty(\sigma)$. This Muckenhoupt space yields a version of scale-invariant absolute continuity between the elliptic measure ω and the surface measure σ . Due to the counterexamples in [CFK81] and [MM80] we do not always have absolute continuity between ω and σ , even if we assume that the coefficients of A are continuous. In fact, these examples show that some regularity on the coefficients in transversal direction is necessary to obtain absolute continuity. This observation gave rise to the study of so called *t-independent* elliptic operators L , i.e. operators where $A(x, t) = A(x)$ is independent of the transversal direction. The first breakthrough in this direction came in [JK81], where Jerison and Kenig used a "Rellich" identity to show that if A is symmetric and t -independent with bounded and measurable coefficients on the unit ball then $\omega \in B_2(\sigma) \subset A_\infty(\sigma)$. Although this "Rellich" identity could not be transferred to the case of nonsymmetric matrices A , the work [KKPT00] was able to establish $\omega \in A_\infty(\sigma)$ for nonsymmetric operators in the upper half plane \mathbb{R}_+^2 after about 20 years. Additionally, they give a counterexample why $\omega \in B_2(\sigma)$ cannot be expected for nonsymmetric matrices and the space $A_\infty(\sigma)$ is sharp in that case.

For dimension n however, it took until the Kato conjecture was resolved in [AHL⁺02] after being open for 50 years, before the authors of [HKMP14] could extend this result to matrices that are not necessarily symmetric and have merely bounded and measurable coefficients. Later, this work got streamlined in [HLMP22a], where the authors also extend the result to the case of matrices whose antisymmetric part might be unbounded and instead has a uniformly bounded BMO norm of its antisymmetric part. The t -independence condition could also be adapted for the parabolic setting as a sufficient condition for solvability of the L^p Dirichlet problem (see [AEN18]) and was also established for the elliptic regularity problem as sufficient condition for solvability (see [HKMP15a]).

A little while after the breakthrough [JK81] the same authors together with Fabes published [FJK84], where they showed that t -independence can be relaxed if we use continuous coefficients. More precisely, they show that if a symmetric

A has continuous coefficients, Ω is a bounded C^1 -domain, and the modulus of continuity

$$\eta(s) = \sup_{P \in \partial\Omega, 0 < r < s} |A_{ij}(P - rV(P)) - A_{ij}(P)|$$

with outer normal vector field V satisfies the Dini-type condition

$$\int_0^\infty \frac{\eta(s)^2}{s} ds < \infty, \quad (1.2)$$

then $\omega \in B_2(\sigma) \subset A_\infty(\sigma)$. Together with the counterexample in [JK81], where for a given η with $\int_0^\infty \frac{\eta(s)^2}{s} ds = +\infty$ the authors construct completely singular measures ω with respect to the surface measure, the Dini type condition (1.2) turns out to be sufficient for $\omega \in A_\infty(d\sigma)$ and kind of necessary, if A is symmetric with continuous bounded coefficients. Necessity here does not mean that every elliptic operator whose matrix does not satisfy the Dini-type condition (1.2) cannot have an elliptic measure in $A_\infty(\sigma)$, and hence this is not a classification of solvability. A little bit later, [Dah86] extended this result to symmetric matrices with merely bounded and measurable coefficients by considering perturbations from continuous matrices.

The third condition that leads to an elliptic measure in the Muckenhoupt class $A_\infty(\sigma)$ is an $L^1 - L^\infty$ condition that appears in [Ulm24]. This condition states that if $|\partial_t A(x, t)| < \frac{c}{t}$ for a.e. $(x, t) \in \mathbb{R}_+^{n+1}$ and

$$\int_0^\infty \|\partial_t A(\cdot, t)\|_{L^\infty(\mathbb{R}^n)} dt < \infty, \quad (1.3)$$

then $\omega \in A_\infty(\sigma)$. This condition generalizes the t -independence condition and hence extends the class of elliptic operators with $\omega \in A_\infty(\sigma)$. Interestingly, this condition is also different from the Dini-type condition (1.2) with examples that satisfy (1.3) but not (1.2) and vice versa (cf. [Ulm24]). We would like to point out that the Dini-type condition (1.2) has not yet been shown on the unbounded domain that we are considering here or for non symmetric matrices, and this comparison illustrates that (1.3) covers indeed a different class of elliptic operators.

However, the question of full classification of matrices for which we obtain $\omega \in A_\infty(\sigma)$ remains open, even for smooth coefficients and symmetric matrices. In trying to make progress towards this question we only look at the case of $n = 1$, i.e. the upper half plane $\Omega := \mathbb{R}_+^2$. We assume matrices with bounded and merely measurable coefficients which might be non symmetric and introduce the L^1 Carleson condition on $\partial_t A(x, t)$ as

$$\sup_{\Delta \subset \partial\Omega} \sup_{\text{boundary ball}} \frac{1}{\sigma(\Delta)} \int_{T(\Delta)} \sup_{(y, s) \in B(x, t, t/2)} |\partial_s A(y, s)| dx dt \leq C < \infty. \quad (1.4)$$

Now we can state the main theorem of this work.

Theorem 1.5. *Let $L := \operatorname{div}(A\nabla\cdot)$ be an elliptic operator satisfying (1.1) on $\Omega = \mathbb{R}_+^2$. If condition (1.4) is satisfied and $|\partial_t A|t \leq C < \infty$, then $\omega \in A_\infty(d\sigma)$, i.e. the L^p Dirichlet boundary value problem for L is solvable for some $1 < p < \infty$.*

It is clear that on the upper half plane the L^1 Carleson condition on $\partial_t A$ (1.4) is weaker than the mixed $L^1 - L^\infty$ condition (1.3). Hence, this new condition gives rise to a larger class of elliptic operators with $\omega \in A_\infty(\sigma)$ that is not covered by any other previously mentioned condition.

The L^1 Carleson condition on $\partial_t A$ can also be put into context from a different point of view. In fact, Carleson conditions are a widely applied tool in the area of solvability of boundary value problems in the elliptic and parabolic setting. The famous DKP condition can be stated as

$$\sup_{\Delta \subset \partial\Omega} \sup_{\text{boundary ball}} \frac{1}{\sigma(\Delta)} \int_{T(\Delta)} |\nabla A(y, s)|^2 t dx dt \leq C < \infty, \quad (1.6)$$

if we assume $|\nabla A(x, t)| \leq \frac{C}{t}$, or with $\operatorname{osc}_{(y,s) \in B(x,t,t/2)} |A(y, s)|$ instead of the term $|\nabla A(y, s)|$, and originates from [KP01]. The DKP condition and versions thereof have been studied for elliptic and parabolic boundary value problems and yield solvability of the corresponding boundary value problem in many cases. The elliptic Dirichlet boundary value problem was studied in [KP01], [DPP07], [HMM⁺21], and [DLM23], and also for elliptic operators with complex coefficients (cf. [DP19]) or elliptic systems (cf. [DHM21]). Furthermore, the DKP condition was also successfully studied for the elliptic regularity problem in [DPR17], [MPT22], [DHP23], and [Fen23]. A helpful survey article of the elliptic setting is [DP23] which contains further references. Lastly, we also have positive results for the parabolic Dirichlet and regularity boundary value problem in [DDH20] and [DLP24].

In contrast to the typical DKP condition (1.6), the new Carleson condition in this work (1.4) does not contain any derivative in any other direction than the transversal t -direction and hence it is easy to see that (1.4) applies to a different class of operators.

1.1 Idea and sketch of the proof

According to [DKP11] it is enough to show the local square function bound

$$\sup_{\Delta \subset \partial\Omega} \sigma(\Delta)^{-1} \int_{T(\Delta)} |\nabla u|^2 \delta dX \lesssim \|f\|_{L^\infty(\partial\Omega)}^2 \quad (1.7)$$

to get $\omega \in A_\infty(d\sigma)$ for small cubes Δ . The first part of the proof follows the outline from [HLMP22a] or [Ulm24] in which we go back to the idea of a Hodge decomposition of the component c of A which originated in [HKMP15b]. However, since c does depend on t we have to deal with a family of Hodge decompositions

$$c(x, t)\chi_{2\Delta}(x) = A_{||}(x, t)\nabla\phi_t(x) + h_t(x) \quad \text{with } x\text{-divergence free } h_t.$$

and have to control this family uniformly. Since we are only in $n = 1$, we get pointwise bounds of $\partial_x \phi^s$ and $\partial_s \partial_x \phi^s$ which we do not find in [Ulm24]. These allow us to find square function bounds of the partial derivatives of the approximating function

$$\rho(x, t) := e^{-t^2 L^t} \phi_t(x),$$

where $L^t := \operatorname{div}_x(A(x, t)\nabla_x \cdot)$ is a family of one-dimensional differential operators. This approximating function ρ plays an important role in the proof as it does in [HLMP22a] and [Ulm24] and bounding the necessary square function terms of the partial derivatives of ρ requires the observation in [Ulm24] to decouple the different dependencies of t by considering

$$w(x, t, s) := e^{-t^2 L^s} \phi_s(x)$$

with three variables. This will allow us to obtain a new explicit representation of ρ . The slight improvement of regularity of ϕ^s that we obtain through the restriction to $n = 1$ allows us to eventually replace the mixed $L^1 - L^\infty$ condition (1.3) from [Ulm24] by something weaker, namely (1.4). The local nature of the Carleson condition require much more delicate handling of each term. In addition to the tools needed in [Ulm24] like the Kato conjecture and heat semigroup properties, we also make use of off-diagonal estimates for the heat semigroup in delicate ways.

As in [Ulm24] the majority of the work lies in establishing the involved square function bounds in Lemma 7.1 and properties of the family $(\nabla \phi^s)_s$, which have to be done differently for our weaker condition (1.4).

Acknowledgements

The author expresses gratitude to Jill Pipher and Martin Dindoš for many insightful discussions in this topic that helped with refining the applied methods.

2 Notations and Setting

Throughout this work let $\Omega = \mathbb{R}_+^2 = \mathbb{R} \times (0, \infty)$ and $\Delta = \Delta(P, r) := B(P, r) \cap \partial\Omega$ denotes a boundary ball centered in point $P \in \partial\Omega$ with radius $l(\Delta) = r > 0$ and

$$T(\Delta) := \{(x, t) \in \Omega; x \in \Delta, 0 < t < l(\Delta)\}$$

its Carleson region. The cone with vertex in $P \in \partial\Omega$ and aperture α is denoted by

$$\Gamma_\alpha(P) := \{(x, t) \in \Omega; |x - P| < \alpha t\}.$$

Furthermore, we set $\mathcal{D}_k^\eta(\Delta)$ as the collection of certain boundary balls with radii $\eta 2^{-k}$ in Δ such that their union covers Δ but they have finite overlap.

We define the nontangential maximal function as

$$N_\alpha(u)(P) := \sup_{(x,t) \in \Gamma_\alpha(P)} |u(x,t)| \quad \text{for } P \in \mathbb{R}^n,$$

and the mean-valued nontangential maximal function as

$$\tilde{N}_\alpha(u)(P) := \sup_{(x,t) \in \Gamma_\alpha(P)} \left(\int_{\{(y,s); |y-x| < \frac{t}{2}\}} |u(y,s)|^2 dy ds \right)^{1/2} \quad \text{for } P \in \mathbb{R}^n.$$

We can now consider the L^p -Dirichlet boundary value problem for the elliptic operator $Lu(x,t) := \operatorname{div}(A(x,t)\nabla u(x,t))$, i.e. for an operator satisfying (1.1).

Definition 2.1. We say the L^p -Dirichlet boundary value problem is solvable for L if for all boundary data $f \in C_c(\partial\Omega) \cap L^p(\Omega)$ the unique existent solution $u \in W^{1,2}(\Omega)$ of

$$\begin{cases} Lu = 0 & \text{in } \Omega, \\ u = f & \text{on } \partial\Omega. \end{cases}$$

satisfies

$$\|N(u)\|_{L^p(\partial\Omega)} \leq C \|f\|_{L^p(\partial\Omega)},$$

where the constant C is independent of u and f .

As usual, we denote the elliptic measure by ω , the usual n -dimensional Hausdorff surface measure by σ and the Muckenhoupt and Reverse Hölder spaces by $A_\infty(d\sigma)$ and $B_q(d\sigma)$ respectively. The solvability of the L^p Dirichlet boundary value problem is equivalent to $\omega \in B_{p'}(d\sigma)$, which means that solvability for some p is equivalent to $\omega \in A_\infty(d\sigma)$.

In our setting we would like to introduce a bit more notation and specification. In the following we are considering the matrix

$$A(x,t) = \begin{pmatrix} A_{\parallel}(x,t) & b(x,t) \\ c(x,t) & d(x,t) \end{pmatrix} \quad (2.2)$$

for $(x,t) \in \Omega = \mathbb{R}_+^{n+1}$, where A_{\parallel}, b, c, d are functions from Ω to \mathbb{R} . We will also be discussing the family of operators

$$L^s v(x) := \operatorname{div}_x(A_{\parallel}(x,s)\nabla v(x)) \quad \text{for } v \in W^{1,2}(\partial\Omega) = W^{1,2}(\mathbb{R}^n), s > 0.$$

At last, we notice that from $\frac{|\partial_t A|}{t} \leq C$ and the L^1 Carleson condition (1.4), which is

$$\frac{1}{\sigma(\Delta)} \int_{T(\Delta)} \sup_{(y,s) \in B(x,t,t/2)} |\partial_s A(y,s)| dx dt \leq C < \infty,$$

we have as a consequence the weaker L^2 Carleson type condition

$$\frac{1}{\sigma(\Delta)} \int_{T(\Delta)} \sup_{(y,s) \in B(x,t,t/2)} |\partial_s A(y,s)|^2 s dx dt \leq C < \infty. \quad (2.3)$$

Since it is sufficient to check (1.7) on only small cubes for $\omega_L \in A_\infty(\sigma)$, we restrict to boundary cubes of size $l(\Delta) \leq R$ for some fixed $R > 0$.

3 Consequences of the Kato conjecture and properties of the heat semigroup

The following two results are the solution to the Kato conjecture which was fully resolved in [AHL⁺02] for $p = 2$. The L^p theory for other p was first fully established in [Aus07] with partial L^p theory results appearing earlier (please refer to the introduction of [Aus07] for the full history). Note that we define $\dot{W}^{1,p}(\mathbb{R}^n)$ as the closure of $C_0^\infty(\mathbb{R}^n)$ under the seminorm given by $\|f\|_{\dot{W}^{1,p}} := \|\nabla f\|_{L^p(\mathbb{R}^n)}$. The Kato L^p theory for the most general elliptic operator is shown in Thm 4.77 in [HLMP22b].

Proposition 3.1. *For a function $f \in \dot{W}^{1,p}(\mathbb{R}^n)$ we have*

$$\|L^{1/2}f\|_{L^p(\mathbb{R}^n)} \leq C(n, \lambda_0, \Lambda_0)\|\nabla f\|_{L^p(\mathbb{R}^n)}$$

for all $1 < p < \infty$. Furthermore, there exists $\varepsilon_1 > 0$ such that if $1 < p < 2 + \varepsilon_1$ then

$$\|\nabla f\|_{L^p(\mathbb{R}^n)} \leq C(n, \lambda_0, \Lambda_0)\|L^{1/2}f\|_{L^p(\mathbb{R}^n)}.$$

The classical Kato solution is the L^2 case.

Proposition 3.2. *For a function $f \in L^2(\mathbb{R}^n)$ we have*

$$\|\nabla L^{-1/2}f\|_{L^2(\mathbb{R}^n)} \approx \|f\|_{L^2(\mathbb{R}^n)}$$

and if f is a vector valued function

$$\|L^{-1/2} \operatorname{div}(f)\|_{L^2(\mathbb{R}^n)} \approx \|f\|_{L^2(\mathbb{R}^n)}.$$

The following result assumes a uniform elliptic t -independent operator L , i.e. $Lu(x) = \operatorname{div}_x(A_{\parallel}(x)\nabla u(x))$ for $x \in \mathbb{R}^n$.

Proposition 3.3 (Proposition 4.3 from [HLMP22a]). *If L is t -independent then the following norm estimates hold for $f \in L^2(\mathbb{R}^n)$, $l \in \mathbb{N}_0$, $t > 0$ and constants $c_l, C > 0$:*

- $\|\partial_t^l e^{-tL}f\|_{L^2(\mathbb{R}^n)} \leq c_l t^{-l}\|f\|_{L^2(\mathbb{R}^n)}$,
- $\|\nabla e^{-tL}f\|_{L^2(\mathbb{R}^n)} \leq C t^{-1/2}\|f\|_{L^2(\mathbb{R}^n)}$.

We have the two following important results

Proposition 3.4. [Proposition 11 in [HLMP22a]] *Let $\eta > 0, \alpha > 0$ and L be t -independent. Then we have for all $(y, t) \in \Gamma_{\eta\alpha}(x)$*

$$\eta^{-1}\partial_t e^{-(\eta t)^2 L}f(y) \lesssim M(\nabla f)(x),$$

and hence

$$\|\eta^{-1}N_{\eta\alpha}[\partial_t e^{-(\eta t)^2 L}f]\|_{L^p} \lesssim \|\nabla f\|_{L^p}$$

for all $p > 1$ and $f \in W^{1,p}(\mathbb{R}^n)$.

Proposition 3.5 (Proposition 12 in [HLMP22a]). *Let $\eta > 0$ and L be t -independent. Then we have*

$$\|\tilde{N}_\eta[\nabla e^{-(\eta t)^2 L} f]\|_{L^p} \lesssim \|\nabla f\|_{L^p}$$

for all $p > 2$ and $f \in W^{1,p}(\mathbb{R}^n)$.

We will also apply the following very basic lemma which can be proved with the above introduced statements and ideas.

Lemma 3.6. *Furthermore for t -independent L , if $f \in W_0^{1,2}(\mathbb{R}^n)$ then*

$$\|\nabla e^{-tL} f\|_{L^2(\mathbb{R}^n)} \leq C \|\nabla f\|_{L^2(\mathbb{R}^n)},$$

and

$$\|\nabla \partial_t e^{-t^2 L} f\|_{L^2(\mathbb{R}^n)} \leq C t^{-1} \|\nabla f\|_{L^2(\mathbb{R}^n)}$$

Proof. The first inequality is an easy corollary from the Kato conjecture, i.e. we have with [Proposition 3.1](#) and [Proposition 3.3](#)

$$\begin{aligned} \|\nabla e^{-tL} f\|_{L^2(\mathbb{R}^n)} &\lesssim \|L^{1/2} e^{-tL} f\|_{L^2(\mathbb{R}^n)} \lesssim \|e^{-tL} L^{1/2} f\|_{L^2(\mathbb{R}^n)} \\ &\lesssim \|L^{1/2} f\|_{L^2(\mathbb{R}^n)} \lesssim \|\nabla f\|_{L^2(\mathbb{R}^n)}. \end{aligned}$$

For the second one we have

$$\begin{aligned} \|\nabla \partial_t e^{-t^2 L} f\|_{L^2(\mathbb{R}^n)}^2 &\lesssim \int_{\mathbb{R}^n} A \nabla \partial_t e^{-t^2 L} f \cdot \nabla \partial_t e^{-t^2 L} f dx \\ &\lesssim \int_{\mathbb{R}^n} \frac{\partial_{tt} e^{-t^2 L} f}{2t} \partial_t e^{-t^2 L} f dx \\ &\lesssim \frac{1}{t} \|\partial_{tt} e^{-t^2 L} f\|_{L^2(\mathbb{R}^n)} \|\partial_t e^{-t^2 L} f\|_{L^2(\mathbb{R}^n)} \end{aligned}$$

Again, we can observe in the proof of in [Proposition 3.4](#) that the only properties of $\partial_t e^{-t^2 L}$ that were used are bounds of its kernel which can be found in Proposition 4.3 of [HLMP22b]. However, the same Proposition gives the same bounds for $t \partial_{tt} e^{-t^2 L}$ and hence we get

$$\|\partial_{tt} e^{-t^2 L} f\|_{L^2(\mathbb{R}^n)} \lesssim \frac{1}{t} \|\nabla f\|_{L^2(\mathbb{R}^n)} \quad \text{and} \quad \|\partial_t e^{-t^2 L} f\|_{L^2(\mathbb{R}^n)} \lesssim \|\nabla f\|_{L^2(\mathbb{R}^n)}.$$

So in total we have

$$\|\nabla \partial_t e^{-t^2 L} f\|_{L^2(\mathbb{R}^n)}^2 \lesssim \frac{1}{t^2} \|\nabla f\|_{L^2(\mathbb{R}^n)}^2.$$

□

Furthermore, we have off-diagonal estimates for operators involving the heat semi group. We quote the following special case of [HLMP22b]:

Proposition 3.7. *We say an operator family $T = (T_t)_{t>0}$ satisfies $L^2 - L^2$ off-diagonal estimates, if there exists $C, \alpha > 0$ such that for all closed sets E and F and all $t > 0$*

$$\|T_t(h)\|_{L^2(F)} \leq C e^{-\alpha \frac{d(E,F)^2}{t}} \|h\|_{L^2(E)},$$

where $\text{supp}(h) = E$ and $d(E, F)$ is the semi-distance induced on sets by Euclidean distance.

Then the families $(e^{-tL})_{t>0}$, $(t\partial_t e^{-tL})_{t>0}$, and $(\sqrt{t}\nabla e^{-tL})_{t>0}$ satisfy $L^2 - L^2$ off-diagonal estimates.

4 Hodge decomposition, Approximation function ρ and difference function θ

4.1 Hodge decomposition

For each $s > 0$ we can find a Hodge decomposition consisting of $\phi^s \in W_0^{1,2}(3\Delta)$ and $h^s \in L^2(3\Delta)$, where h^s is divergence free and

$$c(x, s)\chi_{3\Delta}(x) = A_{\parallel}(x, s)\partial_x \phi^s(x) + h^s(x).$$

Since this is a PDE in one dimension, if we write $3\Delta = (a, a + 3l(\Delta))$, the divergence free function is the constant

$$h^s = \frac{\int_a^{a+3l(\Delta)} \frac{c(x, s)}{A_{\parallel}(x, s)} dx}{\int_a^{a+3l(\Delta)} \frac{1}{A_{\parallel}(x, s)} dx},$$

and

$$\phi^s(y) = \int_a^y \frac{c(x, s) - h^s}{A_{\parallel}(x, s)} dx \quad \text{for } x \in 3\Delta.$$

Hence we obtain the following uniform bounds on $\partial_x \phi^s$ and $\partial_s \partial_x \phi^s$:

Lemma 4.1. *There exists $\kappa_0 > 0$ such that*

$$M[|\partial_x \phi^s|](x) \leq C(\lambda, \Lambda_0)l(\Delta) \leq \kappa_0,$$

and hence

$$\|M[\partial_x \phi^s]\|_{L^2} \leq C(\lambda, \Lambda_0)l(\Delta) \leq \kappa_0|\Delta|^{1/2}.$$

Furthermore, we have for every $x \in 3\Delta$

$$|\partial_s \partial_x \phi^s(x)| \lesssim |\partial_s A(x, s)| + \int_{3\Delta} |\partial_s A(y, s)| dy.$$

Let $\eta > 0$ be a parameter to be determined later. We define

$$w(x, t, s) := e^{-tL^s} \phi^s(x)$$

as the solution to the ("t-independent") heat equation

$$\begin{cases} \partial_t w(x, t, s) - L^s w(x, t, s) = 0 & , (x, t) \in \Omega \\ w(x, 0, s) = \phi^s(x) & , x \in \partial\Omega \end{cases}$$

using the heat semigroup.

We define the by η scaled *approximation function* with ellipticized homogeneity

$$\rho_\eta(x, s) := w(x, \eta^2 s^2, s),$$

and the *difference function*

$$\theta_\eta(x, s) := \phi^s(x) - \rho_\eta(x, s).$$

As a consequence of [Lemma 4.1](#) we have the following bound on the difference function.

Lemma 4.2. *For all $(x, s) \in T(3\Delta)$*

$$\theta_\eta(x, s) \leq C(\lambda_0, \Lambda_0) \kappa_0 \eta s.$$

Furthermore, we have any aperture $\alpha > 0$ and

$$\int_{(x-\eta\alpha s, x+\eta\alpha s)} |\partial_x \rho_\eta(y, s)|^2 dy \lesssim_{\alpha, \eta} M[\partial_x \phi^s]^2(x) \lesssim \kappa_0^2$$

Proof. The first statement follows from the maximum principle for parabolic operators. For the second statement we have with a smooth cut-off function $\psi \in C^\infty$ with $\text{supp}(\psi) \subset (x - 2\eta\alpha s, x + 2\eta\alpha s)$ and $\psi \equiv 1$ on $(x - \eta\alpha s, x + \eta\alpha s)$, while $|\partial_t \psi| + |\partial_x \psi| \lesssim \frac{1}{\varepsilon \eta \alpha s}$. We choose the mean value $m := \int_{(x-2\eta\alpha s, x+2\eta\alpha s)} \phi^s(x) dx = (\phi^s(\cdot))_{B_x(x, 2\eta\alpha s)}$. Then we get

$$\begin{aligned} \int_{(x-2\eta\alpha s, x+2\eta\alpha s)} |\partial_x \rho_\eta|^2 \psi^2 dy &\leq \int_{(x-2\eta\alpha s, x+2\eta\alpha s)} A_{\parallel} \partial_x \rho_\eta \cdot \partial_x \rho_\eta \psi^2 dy \\ &= \int_{(x-2\eta\alpha s, x+2\eta\alpha s)} A_{\parallel} \partial_x \rho_\eta \cdot \partial_x ((\rho_\eta - m) \psi^2) dy \\ &\quad + \int_{(x-2\eta\alpha s, x+2\eta\alpha s)} A_{\parallel} \partial_x \rho_\eta \cdot (\rho_\eta - m) \psi \partial_x \psi dy =: I + J. \end{aligned}$$

Since $sL^s \rho_\eta(x, s) = w_t(x, s)$ we have by [Corollary 4.3](#)

$$I = \int_{(x-2\eta\alpha s, x+2\eta\alpha s)} \partial_t e^{-\eta^2 t^2 L^s} \phi^s|_{t=s} \frac{(\rho_\eta(y, s) - m) \psi^2}{\eta \alpha s} dy \lesssim M[\partial_x \phi^s](P)^2$$

for every $(P, s) \in \Gamma_{\eta\alpha}(x)$. Choosing $P \in F$ leads with [Lemma 4.1](#) to $I \lesssim \kappa_0^2$.

For J we can calculate for $0 < \sigma < 1$

$$\begin{aligned} J &\lesssim \left(\int_{(1+\varepsilon)B_x(x, \eta\alpha s)} |\partial_x \rho_\eta|^2 \psi^2 dy \right)^{1/2} \left(\int_{(1+\varepsilon)B_x(x, \alpha\eta s)} \frac{|\rho_\eta - m|^2}{s^2} dy \right)^{1/2} \\ &\lesssim \sigma \int_{(1+\varepsilon)B_x(x, \alpha\eta s)} |\partial_x \rho_\eta|^2 \psi^2 dy + C_\sigma M [\partial_x \phi^s](P)^2. \end{aligned}$$

Hiding the first term on the left side yields

$$\int_{B_x(x, \alpha\eta s)} |\partial_x \rho_\eta|^2 dy \lesssim \kappa_0^2.$$

□

The previous proof used the following result, which is a corollary of [Proposition 3.4](#). We introduce the function w_t in more detail in the next section when discussing all partial derivatives of θ .

Corollary 4.3. *Let $\eta > 0, \alpha > 0$. Then we have for all $(y, t) \in \Gamma_{\eta\alpha}(x)$*

$$\eta^{-1} w_t(y, s) = \eta^{-1} \partial_t e^{-(\eta t)^2 L^s} \phi^s \Big|_{t=s}(y) \lesssim M(\partial_x \phi^s)(x), \quad (4.4)$$

and

$$\frac{\rho_\eta(y, s)}{\eta s} = \frac{e^{-(\eta s)^2 L^s} (\phi^s - m)(y)}{s} \lesssim M(\partial_x \phi^s)(x), \quad (4.5)$$

where

$$m(s) := \int_{B_x(x, 2\eta\alpha s)} \phi^s(x) dx = (\phi^s(\cdot))_{B_x(x, 2\eta\alpha s)}.$$

Proof. The first inequality (4.4) follows directly from [Proposition 3.4](#). For (4.5) we can observe in the proof of [Proposition 3.4](#) that the only properties of $\partial_t e^{-t^2 L}$ that were used are bounds of its kernel which can be found in [Proposition 4.3](#) of [\[HLMP22b\]](#). However, the same [Proposition](#) gives the same bounds for $\frac{e^{-t^2 L}}{t}$ and hence we get with a similar proof to that of (4.4) also (4.5). □

4.2 The different parts of the derivatives of θ

We will have to take the partial derivative of θ_η in the transversal t -direction. Thereby, we see that

$$\partial_s \theta_\eta(x, s) = \partial_s \phi^s - \partial_s w(x, \eta^2 s^2, s) = \partial_s \phi^s - \partial_t w(x, t, s)|_{t=\eta^2 s^2} - \partial_s w(x, t, s)|_{t=\eta^2 s^2}.$$

We can calculate the second and third term more explicitly. For the first of them we get

$$\partial_t w(x, t, s)|_{t=\eta^2 s^2} = 2\eta^2 s L^s w(x, \eta^2 s^2, s).$$

For the second one we need to work a bit more. To start with we can observe that

$$\begin{aligned}\partial_s \partial_t w(x, t, s) &= \partial_s \partial_x (A_{\parallel}(x, s) \partial_x w(x, t, s)) \\ &= \partial_x (\partial_s A_{\parallel}(x, s) \partial_x w(x, t, s)) + L^s \partial_s w(x, t, s).\end{aligned}$$

We can now set $v_1(x, t)$ as the solution to

$$\begin{cases} \partial_t v_1(x, t) = L^s v_1(x, t) & \text{for } (x, t) \in \Omega, \\ v_1(x, 0) = \partial_s \phi^s(x) & \text{for } (x, 0) \in \partial\Omega. \end{cases}$$

and $v_2(x, t)$ as the solution to

$$\begin{cases} \partial_t v_2(x, t) = L^s v_2(x, t) + \partial_x (\partial_s A_{\parallel}(x, s) \partial_x w(x, t, s)) & \text{for } (x, t) \in \Omega, \\ v_2(x, 0) = 0 & \text{for } (x, 0) \in \partial\Omega, \end{cases}$$

Since $\partial_s w(x, 0, s) = \partial_s \phi^s$ we note that $\partial_s w$ and $v_1 + v_2$ satisfy the same linear PDE with same boundary data and hence must be equal. Next, we can give explicit representations for v_2 by applying Duhamel's principle and for v_1 by the heat semigroup. These are

$$v_1(x, t, s) = e^{-tL^s} \partial_s \phi^s(x),$$

and

$$v_2(x, t, s) = \int_0^t e^{-(t-\tau)L^s} \partial_x (\partial_s A_{\parallel}(x, s) \partial_x w(x, \tau, s)) d\tau.$$

Together we get for the derivative of θ in t -direction

$$\begin{aligned}\partial_s \theta(x, s) &= \partial_s \phi^s - 2\eta^2 s L^s w(x, \eta^2 s^2, s) - e^{-\eta^2 s^2 L^s} \partial_s \phi^s(x) \\ &\quad - \int_0^{\eta^2 s^2} e^{-(\eta^2 s^2 - \tau)L^s} \partial_x (\partial_s A_{\parallel}(x, s) \partial_x w(x, \tau, s)) d\tau \\ &= \partial_s \phi^s - 2\eta^2 s L^s w(x, \eta^2 s^2, s) - e^{-\eta^2 s^2 L^s} \partial_s \phi^s(x) \\ &\quad - \int_0^s 2\eta^2 \tau e^{-\eta^2 (s^2 - \tau^2)L^s} \partial_x (\partial_s A_{\parallel}(x, s) \partial_x w(x, \eta^2 \tau^2, s)) d\tau \\ &=: \partial_s \phi^s(x) - w_t(x, s) - w_s^{(1)}(x, s) - w_s^{(2)}(x, s).\end{aligned}$$

Lemma 4.6. *Let $\tilde{Q} \subset \Omega$ be a cube and assume also $4\tilde{Q} \subset \Omega$ and $s \approx l(\tilde{Q})$. If we set*

$$v(x, t, s) := \partial_t w(x, t^2, s) = \partial_t e^{-t^2 L^s} \phi_s,$$

we have the Caccioppoli type inequality

$$\int_{\tilde{Q}} |\partial_x v|^2 dx dt \lesssim \frac{1}{s^2} \int_{2\tilde{Q}} |v|^2 dx dt.$$

The proof works analogously to the proof of the other two Caccioppoli type inequalities in [Lemma 4.7](#).

Proof. We observe that v satisfies the PDE

$$\begin{aligned}\partial_t v(x, t, s) &= \partial_t \partial_t w(x, t^2, s) = \partial_t (-2tL^s w(x, t^2, s)) \\ &= -2L^s w(x, t^2, s) - 2tL^s v(x, t, s) = \frac{v(x, t, s)}{t} - 2tL^s v(x, t^2, s).\end{aligned}$$

The rest of the proof follows standard arguments for Caccioppoli inequalities. \square

To proof [Lemma 6.4](#) we need two more Caccioppoli type inequalities which are the following

Lemma 4.7. *Let $2\hat{Q} \subset \Omega$ be a Whitney cube, i.e. $\text{dist}(\hat{Q}, \partial\Omega) \approx l(\hat{Q}) \approx s$. If we set*

$$v(x, t, s) := v_2(x, t^2, s) = \int_0^t 2\tau e^{-(t^2-\tau^2)L^s} \text{div}(\partial_s A(x, s) \partial_x e^{-\tau^2 L^s} \phi_s(x)) d\tau,$$

then we have

$$\int_{\hat{Q}} |\partial_x v(x, t, s)|^2 dx dt \lesssim \frac{1}{s^2} \int_{2\hat{Q}} |v(x, t, s)|^2 dx dt + \int_{2\hat{Q}} |\partial A_{\parallel}(x, s) \partial_x e^{-t^2 L^s} \phi_s|^2 dx dt,$$

and

$$\begin{aligned}\int_{\hat{Q}} |\partial_x \partial_t v(x, t, s)|^2 dx dt &\lesssim \frac{1}{s^2} \int_{2\hat{Q}} |\partial_t v(x, t, s)|^2 dx dt \\ &\quad + \frac{1}{s^2} \int_{2\hat{Q}} |\partial_t A_{\parallel}(x, s) \partial_x e^{-t^2 L^s} \phi_s|^2 dx dt \\ &\quad + \int_{2Q} |\partial_t A_{\parallel}(x, s) \partial_x \partial_t e^{-t^2 L^s} \phi_s|^2 dx dt.\end{aligned}$$

Proof. First we denote $\tilde{v} := \partial_t v$ to shorten notation. We observe that v and \tilde{v} satisfy the following PDEs

$$\begin{aligned}\partial_t v &= 2t \text{div}(\partial_s A_{\parallel} \partial_x e^{-t^2 L^s} \phi_s) - \int_0^t 4t\tau L^s e^{-(t^2-\tau^2)L^s} \text{div}(\partial_s A_{\parallel}(x, s) \partial_x e^{-\tau^2 L^s} \phi_s(x)) d\tau \\ &= 2t \text{div}(\partial_s A_{\parallel} \partial_x e^{-t^2 L^s} \phi_s) - 2tL^s v,\end{aligned}$$

and

$$\partial_t \tilde{v} = \text{div}(\partial_s A_{\parallel} \partial_x e^{-t^2 L^s} \phi_s) + 2t \text{div}(\partial_s A_{\parallel} \partial_x \partial_t e^{-t^2 L^s} \phi_s) - 2L^s v - 2tL^s \tilde{v}.$$

The rest of the proof is standard for Caccioppoli inequalities of inhomogeneous parabolic PDEs and we omit it here. \square

5 Proof of *Theorem 1.5*

To show that $\omega \in A_\infty(d\sigma)$ we would like to show (1.7). It is well known that it is enough to show the Carleson measure estimate for only all small boundary balls, i.e. boundary balls Δ with $l(\Delta) \leq R$ for some fixed $R > 0$. First, we fix a boundary cube $\Delta \subset \partial\Omega$ and a small $\eta > 0$ that we determine later and assume without loss of generality that $\|f\|_{L^\infty(\partial\Omega)} \leq 1$. Note that the maximum principle implies $\|u\|_\infty \leq 1$. To start with we introduce a smooth cut-off of $T(\Delta)$. Let $\psi \in C^\infty(\mathbb{R}_+^{n+1})$ with

$$\psi \equiv 1 \text{ on } T(\Delta),$$

and

$$\psi \equiv 0 \text{ on } \Omega \setminus (T(3\Delta)).$$

We also have

$$|\nabla\psi(x, t)| \lesssim \frac{1}{l(\Delta)} \quad \text{for all } (x, t) \in \Omega.$$

We can start to estimate the left side of (1.7). Since $u\Psi^2t \in W_0^{1,2}(\Omega)$ we have

$$\int_{\Omega} A\nabla u \cdot \nabla(u\psi^2t) dxdt = 0,$$

and hence

$$\begin{aligned} \int_{T(\Delta)} |\nabla u|^2 t dxdt &\leq \int_{\Omega} |\nabla u|^2 \psi^2 t dxdt \\ &\lesssim \int_{\Omega} A\nabla u \cdot \nabla u \psi^2 t dxdt \\ &= \int_{\Omega} A\nabla u \cdot \nabla(u\psi^2t) dxdt + \int_{\Omega} A\nabla u \cdot \nabla\psi\psi ut dxdt \\ &\quad + \int_{\Omega} A\nabla u \cdot \vec{e}_2 u \psi^2 dxdt \\ &= \int_{\Omega} A\nabla u \cdot \nabla\psi\psi ut dxdt + \int_{\Omega} A\nabla u \cdot \vec{e}_2 u \Psi^2 dxdt \\ &=: J_1 + J_2 \end{aligned}$$

For the first term J_1 we have with boundedness of A that

$$\begin{aligned} J_1 &\leq \left(\int_{\Omega} |\nabla u|^2 \psi^2 t dxdt \right)^{1/2} \left(\int_{\Omega} |\nabla\psi|^2 u^2 t dxdt \right)^{1/2} \\ &\leq \sigma \int_{\Omega} |\nabla u|^2 \psi^2 t dxdt + C_\sigma |\Delta|, \end{aligned}$$

and for sufficiently small $\sigma > 0$ we can hide the first term on the left side.

For J_2 to continue to break up the integral to

$$\int_{\Omega} A\nabla u \cdot \vec{e}_2 u \psi^2 dxdt = \int_{\Omega} c \cdot \partial_x u u \psi^2 dxdt + \int_{\Omega} d \partial_t u u \psi^2 dxdt =: J_{21} + J_{22}.$$

We have for J_{22}

$$\begin{aligned} J_{22} &= \int_{\Omega} \partial_t (du^2 \Psi^2) dxdt + \int_{\Omega} du^2 \partial_t \Psi^2 dxdt + \int_{\Omega} \partial_t du^2 \Psi^2 \\ &\lesssim \int_{3\Delta} |du^2 \Psi^2| dxdt + |\Delta| + \int_{\Omega} |\partial_t d| \\ &\lesssim |\Delta|. \end{aligned}$$

Thus it remains to bound J_{21} .

We proceed with

$$J_{21} = \int_{\Omega} c \cdot \partial_x (u^2 \psi^2) dxdt + \int_{\Omega} c \cdot \partial_x \psi u^2 \psi dxdt := J_{211} + |\Delta|.$$

For the term J_{211} we apply the Hodge decomposition of c and get

$$\begin{aligned} J_{211} &= \int_{\Omega} A_{\parallel} \partial_x \phi^s \cdot \partial_x (u^2 \psi^2) dxdt \\ &= \int_{\Omega} A_{\parallel} \partial_x \theta_{\eta} \cdot \partial_x (u^2 \psi^2) dxdt - \int_{\Omega} A_{\parallel} \partial_x \rho \cdot \partial_x (u^2 \psi^2) dxdt \\ &= \int_{\Omega} A_{\parallel} \partial_x \theta_{\eta} \cdot \partial_x (u^2 \psi^2) dxdt - \int_{\Omega} A_{\parallel} \partial_x \partial_t \rho \cdot \partial_x (u^2 \psi^2) t dxdt \\ &\quad - \int_{\Omega} A_{\parallel} \partial_x \rho \cdot \partial_x \partial_t (u^2 \psi^2) t dxdt - \int_{\Omega} \partial_t A_{\parallel} \partial_x \rho \cdot \partial_x (u^2 \psi^2) t dxdt \\ &=: J_{2111} + I_1 + I_2 + I_3. \end{aligned}$$

First we deal with I_1, I_2 and I_3 . By [Lemma 7.1](#) we have

$$I_1 \lesssim \left(\int_{T(\Delta)} |\partial_x \partial_t \rho|^2 t dxdt \right)^{1/2} \left(\int_{T(\Delta)} |\partial_x (u^2 \psi^2)|^2 t dxdt \right)^{1/2} \lesssim |\Delta|^{1/2} J_1^{1/2} \lesssim |\Delta|.$$

Next, for I_2 we have by [Lemma 7.1](#)

$$I_2 \lesssim \left(\int_{T(\Delta)} |\operatorname{div}(A_{\parallel} \partial_x \rho)|^2 t dxdt \right)^{1/2} \left(\int_{T(\Delta)} |\partial_t (u^2 \psi^2)|^2 t dxdt \right)^{1/2} \lesssim |\Delta|^{1/2} J_1^{1/2} \lesssim |\Delta|.$$

At last, for I_3 and by [Lemma 7.1](#) we get

$$I_3 \lesssim \left(\int_{T(\Delta)} |\partial_t A_{\parallel}|^2 |\partial_x \rho|^2 t dxdt \right)^{1/2} \left(\int_{T(\Delta)} |\partial_x (u^2 \psi^2)|^2 t dxdt \right)^{1/2} \lesssim |\Delta|^{1/2} J_1^{1/2} \lesssim |\Delta|.$$

We continue with J_{2111} and have

$$\int_{\Omega} A_{\parallel} \partial_x \theta_{\eta} \cdot \partial_x (u^2 \psi^2) dxdt + \int_{\Omega} A_{\parallel} \partial_x \theta_{\eta} \cdot u^2 \partial_x (\psi^2) dxdt =: II_1 + II_2.$$

For II_2 we observe that

$$II_2 \lesssim \left(\int_{T(3\Delta)} \frac{|\partial_x \theta_{\eta}|^2}{l(\Delta)^2} dxdt \right)^{1/2} |\Delta|^{1/2} \lesssim |\Delta|.$$

For II_1 we get

$$\begin{aligned} II_1 &= \int_{\Omega} A_{\parallel} \partial_x u \cdot \partial_x (\theta_{\eta} u \psi^2) dx dt + \int_{\Omega} A_{\parallel} \partial_x u \cdot \partial_x u \theta_{\eta} \psi^2 dx dt + \int_{\Omega} A_{\parallel} \partial_x u^2 \cdot \partial_x \psi^2 \theta_{\eta} dx dt \\ &=: II_{11} + II_{12} + II_{13}. \end{aligned}$$

Due to [Lemma 4.2](#) we have that II_{13} can be handled like J_1 and

$$II_{12} = \eta \kappa_0 \int_{T(\Delta)} |\partial_x u|^2 \psi^2 t dx dt,$$

and choosing η sufficiently small let us hide this term on the left side. To deal with II_{11} we use that u is a weak solution and $\theta_{\eta} u \psi^2 \in W_0^{1,2}(\Omega)$ to conclude

$$\begin{aligned} II_{11} &= \int_{\Omega} A_{\parallel} \partial_x u \cdot \partial_x (\theta_{\eta} u \psi^2) dx dt \\ &= \int_{\Omega} b \partial_t u \cdot \partial_x (\theta_{\eta} u \psi^2) dx dt + \int_{\Omega} c \cdot \partial_x u \partial_t (\theta_{\eta} u \psi^2) dx dt + \int_{\Omega} d \partial_t u \partial_t (\theta_{\eta} u \psi^2) dx dt \\ &=: II_{111} + II_{112} + II_{113}. \end{aligned}$$

We have with [Lemma 4.2](#)

$$\begin{aligned} II_{113} &= \int_{\Omega} d \partial_t u \partial_t \theta_{\eta} u \psi^2 dx dt + \int_{\Omega} d \partial_t u \theta_{\eta} \partial_t u \psi^2 dx dt + \int_{\Omega} d \partial_t u \theta_{\eta} u \partial_t \psi^2 dx dt \\ &\lesssim \left(\int_{T(3\Delta)} \frac{|\partial_t \theta_{\eta}|^2}{t} dx dt \right)^{1/2} \left(\int_{T(\Delta)} |\partial_t u|^2 \psi^2 t dx dt \right)^{1/2} + \int_{T(\Delta)} |\partial_t u|^2 \psi^2 \theta_{\eta} dx dt \\ &\quad + \int_{\Omega} |\partial_t u \theta_{\eta} u \partial_t \psi^2| dx dt \\ &\lesssim C_{\sigma} \int_{T(3\Delta)} \frac{|\partial_t \theta_{\eta}|^2}{t} dx dt + \sigma \int_{T(\Delta)} |\partial_t u|^2 \psi^2 t dx dt \\ &\quad + \eta \int_{T(\Delta)} |\partial_t u|^2 \psi^2 t dx dt + \int_{\Omega} |\partial_t u^2 \partial_t \psi^2| t dx dt. \end{aligned}$$

Hence, for a sufficiently small choice of η and σ we can hide the second and third term on the left side, while the fourth term is bounded like J_1 and the first one with [Lemma 7.1](#).

Analogously, we get for II_{112}

$$\begin{aligned} II_{112} &= \int_{\Omega} c \cdot \partial_x u \partial_t \theta_{\eta} u \psi^2 dx dt + \int_{\Omega} c \cdot \partial_x u \theta_{\eta} \partial_t u \psi^2 dx dt + \int_{\Omega} c \cdot \partial_x u \theta_{\eta} u \partial_t \psi^2 dx dt \\ &\lesssim C_{\sigma} \int_{T(3\Delta)} \frac{|\partial_t \theta_{\eta}|^2}{t} dx dt + \sigma \int_{T(\Delta)} |\partial_x u|^2 \psi^2 t dx dt \\ &\quad + \eta \int_{T(\Delta)} |\partial_x u|^2 \psi^2 t dx dt + \int_{\Omega} |\partial_x u^2 \partial_t \psi^2| t dx dt, \end{aligned}$$

and hence $II_{113}, II_{112} \lesssim |\Delta|$. For II_{111} however, we write

$$\begin{aligned} II_{111} &= \int_{\Omega} b \cdot \partial_x \theta_{\eta} \partial_t u^2 \psi^2 dxdt + \int_{\Omega} b \cdot \partial_x u \partial_t u^2 \psi^2 \theta_{\eta} dxdt + \int_{\Omega} b \cdot \partial_x \psi u \partial_t u \psi \theta_{\eta} dxdt \\ &=: II_{1111} + II_{1112} + II_{1113}. \end{aligned}$$

Since

$$II_{1112} \lesssim \eta \int_{T(\Delta)} |\partial_x u|^2 \psi^2 t dxdt,$$

and

$$II_{1113} \lesssim \left(\int_{T(3\Delta)} \frac{|u|^2}{l(\Delta)^2} t dxdt \right)^{1/2} \left(\int_{T(3\Delta)} |\partial_t u|^2 \psi^2 t dxdt \right)^{1/2}$$

we get boundedness of these two terms like before.

Next, we recall that $\partial_t \theta_{\eta} = \partial_t \phi^t - (w_t + w_s^{(1)} + w_s^{(2)})$ and get by integration by parts

$$\begin{aligned} II_{1111} &= \int_{\Omega} b \cdot \partial_x \partial_t \theta_{\eta} u^2 \psi^2 dxdt + \int_{\Omega} b \cdot \partial_x \theta_{\eta} u^2 \psi \partial_t \psi dxdt + \int_{\Omega} \partial_t b \cdot \partial_x \theta_{\eta} u^2 \psi^2 dxdt \\ &\lesssim \int_{\Omega} b \cdot \partial_x (w_t u^2 \psi^2) dxdt + \int_{\Omega} b \cdot \partial_x (\partial_t \phi^t - w_s^{(1)} + w_s^{(2)}) u^2 \psi^2 dxdt \\ &\quad + \int_{\Omega} b \cdot \partial_x (u^2 \psi^2) w_t dxdt + \int_{\Omega} b \cdot \partial_x \theta_{\eta} u^2 \psi \partial_t \psi dxdt + \int_{\Omega} \partial_t b \cdot \partial_x \theta_{\eta} u^2 \psi^2 dxdt \\ &=: III_1 + III_2 + III_3 + III_4 + III_5. \end{aligned}$$

We can bound III_4 like II_2 , while with Cauchy-Schwarz, [Lemma 6.9](#) and with hiding of terms on the left side we have

$$\begin{aligned} III_3 &\lesssim \int_{T(3\Delta)} |w_t \partial_x u u \psi^2| + |w_t u^2 \partial_x \psi \psi| dxdt \\ &\lesssim \left(\int_{T(3\Delta)} \frac{|w_t|^2}{t} dxdt \right)^{1/2} \left(\int_{T(3\Delta)} |\partial_x u|^2 \psi^2 t dxdt \right)^{1/2} \\ &\quad + \left(\int_{T(3\Delta)} \frac{|w_t|^2}{t} dxdt \right)^{1/2} \left(\int_{T(3\Delta)} |u \partial_x \psi|^2 t dxdt \right)^{1/2}. \end{aligned}$$

We obtain boundedness by $|\Delta|$ for III_3 .

For III_2 we observe that due to (6.7) in [Lemma 6.4](#), and (iii) in [Lemma 6.1](#) we have

$$III_2 \lesssim \int_{T(3\Delta)} |\partial_x \partial_t \phi^t| + |\partial_x w_s^{(1)}| + |\partial_x w_s^{(2)}| dxdt \lesssim |\Delta|$$

For III_5 we get with [Lemma 4.1](#) and [Lemma 4.2](#)

$$\int_{\Omega} |\partial_t b \cdot \partial_x \theta_{\eta} u^2 \psi^2| dxdt \lesssim \int_{T(3\Delta)} \sup_{B(x,t,t/2)} |\partial_s b(y,s)| dxdt \lesssim |\Delta|.$$

For III_1 however, we need the Hodge decomposition of b . Since $\partial\theta_\eta\psi^2u^2 \in W_0^{1,2}(T(3\Delta))$ we have

$$III_1 = \int_{\Omega} A_{\parallel} \partial_x \tilde{\theta} \cdot \partial_x (w_t u^2 \psi^2) dxdt + \int_{\Omega} A_{\parallel} \partial_x \tilde{\rho} \cdot \partial_x (w_t u^2 \psi^2) dxdt =: III_{11} + III_{12}.$$

With integration by parts the second term becomes

$$III_{12} \lesssim \left(\int_{T(3\Delta)} \frac{|w_t|^2}{t} dxdt \right)^{1/2} \left(\int_{T(3\Delta)} |L^t \tilde{\rho}|^2 t dxdt \right)^{1/2} \lesssim |\Delta|,$$

where the last inequality follows from [Lemma 7.1](#) and [Lemma 6.9](#).

For III_{11} we obtain

$$\begin{aligned} III_{11} &= \int_{\Omega} A_{\parallel} \partial_x \tilde{\theta} \cdot \partial_x u^2 w_t \psi^2 dxdt + \int_{\Omega} A_{\parallel} \partial_x \tilde{\theta} \cdot \partial_x \psi^2 w_t u^2 dxdt + \int_{\Omega} A_{\parallel} \partial_x \tilde{\theta} \cdot \partial_x w_t u^2 \psi^2 dxdt \\ &=: III_{111} + III_{112} + III_{113}. \end{aligned}$$

First, we see that

$$III_{112} = \left(\int_{T(3\Delta)} \frac{|w_t|^2}{t} dxdt \right)^{1/2} \left(\int_{T(3\Delta)} |\partial_x \tilde{\theta}|^2 |\partial_x \psi|^2 t dxdt \right)^{1/2}$$

can be dealt with by II_2 and [Lemma 6.9](#).

Next, for III_{113} we obtain by integration by parts

$$III_{113} = \int_{\Omega} L^t w_t u^2 \psi^2 \tilde{\theta} dxdt + \int_{\Omega} A_{\parallel} \partial_x w_t \partial_x (u^2 \psi^2) \tilde{\theta} dxdt,$$

where the first term can be bounded by

$$\left(\int_{T(3\Delta)} \frac{|\tilde{\theta}|^2}{t^3} dxdt \right)^{1/2} \left(\int_{T(3\Delta)} |L^t w_t|^2 t^3 dxdt \right)^{1/2} \lesssim |\Delta|$$

due to [Lemma 7.1](#) and [Lemma 6.9](#), while the second term is bounded by

$$\begin{aligned} &\int_{T(3\Delta)} |\partial_x w_t \partial_x u w \psi^2| t + |\partial_x w_t u^2 \partial_x \psi \psi| t dxdt \\ &\lesssim \left(\int_{T(3\Delta)} |\partial_x w_t|^2 t dxdt \right)^{1/2} \left(\int_{T(3\Delta)} |\partial_x u|^2 \psi^2 t dxdt \right)^{1/2} \\ &\quad + \left(\int_{T(3\Delta)} |\partial_x w_t|^2 t dxdt \right)^{1/2} \left(\int_{T(3\Delta)} |u \partial_x \psi|^2 t dxdt \right)^{1/2}, \end{aligned}$$

and hence by hiding terms on the left side and [Lemma 6.9](#).

At last, it remains to bound III_{111} . For that we have with [Lemma 4.2](#)

$$III_{111} \lesssim \sigma \int_{T(3\Delta)} |\partial_x u|^2 \psi^2 t dxdt + C_{\sigma} \int_{T(3\Delta)} \frac{|w_t|^2}{t} dxdt.$$

The first term can be hidden on the left side with sufficiently small choice of σ . For the second one we write

$$\begin{aligned} \int_{T(3\Delta)} \frac{|\partial_x \tilde{\theta}|^2 |w_t|^2}{t} dxdt &= \sum_{k \leq k_0} \sum_{Q' \in \mathcal{D}_k^n(5\Delta)} \int_{Q'} \int_{2^{-k}}^{2^{-k+1}} \frac{|\partial_x \tilde{\theta}|^2 |w_t|^2}{t} dxdt \\ &= \sum_{k \leq k_0} \sum_{Q' \in \mathcal{D}_k^n(5\Delta)} \left(\int_{Q'} \int_{2^{-k}}^{2^{-k+1}} \frac{|\partial_x \tilde{\theta}|^2}{t} dxdt \right) \left(|Q| \sup_{Q \times (2^{-k}, 2^{-k+1}]} |w_t| \right) \end{aligned}$$

Using [Lemma 4.1](#) and [Lemma 4.1](#) we see that

$$\int_{Q'} \int_{2^{-k}}^{2^{-k+1}} |\partial_x \tilde{\theta}|^2 dxdt = \int_{Q'} \int_{2^{-k}}^{2^{-k+1}} |\partial_x \tilde{\rho}|^2 dxdt + \int_{Q'} \int_{2^{-k}}^{2^{-k+1}} |\partial_x \tilde{\phi}^t|^2 dxdt \lesssim \kappa_0,$$

and hence with [Lemma 6.10](#)

$$\int_{T(3\Delta)} \frac{|\partial_x \tilde{\theta}|^2 |w_t|^2}{t} dxdt \lesssim \sum_{k \leq k_0} \sum_{Q' \in \mathcal{D}_k^n(5\Delta)} \int_{Q'} \int_{2^{-k}}^{2^{-k+1}} \frac{|w_t|^2}{t} dxdt \lesssim |\Delta|.$$

The last inequality follows again from [Lemma 7.1](#). In total we verified the Carleson measure estimate (1.7).

6 Square function bounds on the parts of the derivative of θ

We investigate square function type expression involving of $\partial_s \phi^s, w_t, w_s^{(1)}$ and $w_s^{(2)}$. For all of the results in this chapter we assume the L^1 Carleson condition (1.4).

6.1 The partial derivative parts $\partial_s \phi^s$ and $w_s^{(1)}$

Instead of investigating $\partial_s \phi^s$ and $w_s^{(1)}$ separately, we are also going to use that their difference

$$\partial_s \phi^s - w_s^{(1)} = \partial_s \phi^s - e^{-(\eta s)^2 L^s} \partial_s \phi^s$$

gives us better properties.

Lemma 6.1. *The following square function bounds hold*

(i)

$$\int_{T(3\Delta)} \frac{|\partial_s \phi^s - w_s^{(1)}|^2}{s} dxds \lesssim |\Delta|,$$

(ii)

$$\int_{T(3\Delta)} \frac{|s \partial_x \partial_s \phi^s|^2}{s} dxds + \int_{T(3\Delta)} \frac{|s \partial_x w_s^{(1)}|^2}{s} dxds \lesssim |\Delta|.$$

(iii)

$$\int_{T(3\Delta)} |\partial_x \partial_s \phi^s| dx ds + \int_{T(3\Delta)} |\partial_x w_s^{(1)}| dx ds \lesssim |\Delta|.$$

Proof. For (i) we use an analogous way to the proof of [Lemma 4.2](#), where we use [Proposition 3.4](#) to get

$$\begin{aligned} (\partial_s \phi^s - e^{-(\eta s)^2 L^s} \partial_s \phi^s)(x) &= \int_0^s \partial_t e^{-(\eta t)^2 L^s} \partial_s \phi^s dt \lesssim \eta \int_0^s M[\partial_x \partial_s \phi^s](x) dt \\ &\lesssim M[\partial_x \partial_s \phi^s](x) \eta s. \end{aligned}$$

Hence by [Lemma 4.1](#),

$$\begin{aligned} \int_{T(3\Delta)} \frac{|\partial_s \phi^s - w_s^{(1)}|^2}{s} dx ds &\lesssim \int_0^{l(3\Delta)} \int_{3\Delta} |M[\partial_x \partial_s \phi^s](x)|^2 s dx ds \\ &\lesssim \int_0^{l(3\Delta)} \int_{3\Delta} |\partial_x \partial_s \phi^s(x)|^2 s dx ds \\ &\lesssim \int_0^{l(3\Delta)} \int_{3\Delta} |\partial_s A|^2 s dx ds \\ &\lesssim |\Delta|, \end{aligned}$$

and observation (2.3) finishes the proof of (i).

For (ii) we just observe that $\|\partial_x w_s^{(1)}\|_{L^2(\mathbb{R}^n)} \lesssim \|\partial_x \partial_s \phi^s\|_{L^2(\mathbb{R}^n)}$, which implies

$$\begin{aligned} \int_{T(3\Delta)} \frac{|s \partial_x \partial_s \phi^s|^2}{s} dx ds, \int_{T(3\Delta)} \frac{|s \partial_x w_s^{(1)}|^2}{s} dx ds \\ \lesssim \int_0^{l(3\Delta)} \int_{3\Delta} |\partial_x \partial_s \phi^s(x)|^2 s dx ds. \end{aligned}$$

Following the argument for (i) from here finishes the proof for (ii).

Lastly, we deal with (iii). By [Lemma 4.1](#) and [Lemma 4.2](#) we have

$$\int_{T(3\Delta)} |\partial_x \partial_s \phi^s| dx ds \lesssim \int_{T(3\Delta)} |\partial_s A_{||}| dx ds \lesssim |\Delta|,$$

and

$$\begin{aligned} \int_{T(3\Delta)} |\partial_x w_s^{(1)}| dx ds &= \int_0^{3l(\Delta)} \sum_{\substack{Q \subset 5\Delta, \\ l(Q) \approx s, Q \text{ finite overlap, } \cup Q \supset 3\Delta}} \left(\int_Q |\partial_x e^{-\eta^2 s^2 L^s} \partial_s \phi^s|^2 dx \right)^{1/2} |Q| \\ &\lesssim \int_0^{3l(\Delta)} \sum_{\substack{Q \subset 5\Delta, \\ l(Q) \approx s, Q \text{ finite overlap, } \cup Q \supset 3\Delta}} \sup_{B(x, s, s/2)} |\partial_s A_{||}| |Q| \\ &\lesssim \int_{T(5\Delta)} \sup_{B(x, s, s/2)} |\partial_s A_{||}| \lesssim |\Delta|. \end{aligned}$$

□

6.2 The partial derivative part $w_s^{(2)}$

Lemma 6.2. *It holds that*

$$\int_{T(3\Delta)} \frac{|w_s^{(2)}|^2}{s} dx ds \lesssim |\Delta|. \quad (6.3)$$

Proof. We use Minkowski inequality and [Proposition 3.1](#) to obtain

$$\begin{aligned} \int_{T(3\Delta)} \frac{|w_s^{(2)}|^2}{s} dx ds &= \int_0^{3l(\Delta)} \int_{3\Delta} \frac{1}{s} \left| \int_0^s \eta^2 \tau e^{-\eta^2(s^2-\tau^2)L^s} \partial_x(\partial_s A_{\parallel}(x, s)) \partial_x w(x, \eta^2 \tau^2, s) d\tau \right|^2 dx ds \\ &\lesssim \eta^2 \int_0^{3l(\Delta)} \frac{1}{s} \left(\int_0^s \tau \|(L^s)^{\frac{1}{2}} e^{-\eta^2(s^2-\tau^2)L^s} (L^s)^{-\frac{1}{2}} \partial_x(\partial_s A_{\parallel}(x, s)) \partial_x w(x, \eta^2 \tau^2, s)\|_{L^2(\mathbb{R}^n)} d\tau \right)^2 ds \\ &\lesssim \eta^2 \int_0^{3l(\Delta)} \frac{1}{s} \left(\int_0^s \tau \|\partial_x e^{-\eta^2(s^2-\tau^2)L^s} (L^s)^{-\frac{1}{2}} \partial_x(\partial_s A_{\parallel}(x, s)) \partial_x w(x, \eta^2 \tau^2, s)\|_{L^2(\mathbb{R}^n)} d\tau \right)^2 ds \\ &\lesssim \eta^2 \int_0^{3l(\Delta)} \frac{1}{s} \left(\int_0^s \frac{\tau}{\sqrt{s^2-\tau^2}} \|(L^s)^{-\frac{1}{2}} \partial_x(\partial_s A_{\parallel}(x, s)) \partial_x w(x, \eta^2 \tau^2, s)\|_{L^2(\mathbb{R}^n)} d\tau \right)^2 ds \\ &\lesssim \eta^2 \int_0^{3l(\Delta)} \frac{1}{s} \left(\int_0^s \frac{\tau}{\sqrt{s^2-\tau^2}} \left(\int_{\mathbb{R}^n} |\partial_s A(x, s) \partial_x w(x, \eta^2 \tau^2, s)|^2 dx \right)^{1/2} d\tau \right)^2 ds \end{aligned}$$

Let us now use [Proposition 3.7](#) for $w(x, \eta^2 t^2, s) = e^{-\eta^2 t^2 L^s} \phi^s$. We have

$$\begin{aligned} &\left(\int_{\mathbb{R}^n} |\partial_s A(x, s) \partial_x e^{-\eta^2 \tau^2 L^s} \phi^s(x)|^2 dx \right)^{1/2} \\ &= \sum_{k=1}^{\infty} \left(\int_{2^k 3\Delta \setminus 2^{k-1} 3\Delta} |\partial_s A(x, s) \partial_x e^{-\eta^2 \tau^2 L^s} \phi^s(x)|^2 dx \right)^{1/2} + \left(\int_{3\Delta} |\partial_s A(x, s) \partial_x e^{-\eta^2 \tau^2 L^s} \phi^s(x)|^2 dx \right)^{1/2} \\ &= \sum_{k=1}^{\infty} \frac{1}{s} e^{-c \frac{2^k l(\Delta)^2}{\tau^2}} \|\phi^s\|_{L^2} + \left(\sum_{\substack{W \in \mathcal{D}(5\Delta) \\ l(W) \approx \tau}} \sup_{x \in W} |\partial_s A(x, s)|^2 \int_W |\partial_x e^{-\eta^2 \tau^2 L^s} \phi^s(x)|^2 dx \right)^{1/2}, \end{aligned}$$

where we made use of the pointwise bound of $|\partial_s A| \lesssim \frac{1}{s}$. By Poincaré inequality, we obtain $\|\phi^s\|_{L^2} \lesssim l(\Delta) \|\partial_x \phi^s\|_{L^2} \lesssim l(\Delta) |\Delta|^{1/2}$, whence

$$\sum_{k=1}^{\infty} \frac{l(\Delta)}{s} e^{-c \frac{2^k l(\Delta)^2}{\tau^2}} |\Delta|^{1/2} \leq \sum_{k=1}^{\infty} \frac{l(\Delta)}{\tau} e^{-c \frac{2^k l(\Delta)^2}{\tau^2}} |\Delta|^{1/2} \lesssim \frac{\tau}{l(\Delta)} |\Delta|^{1/2} \lesssim |\Delta|^{1/2}.$$

For the second term we apply [Lemma 4.2](#) and obtain

$$\begin{aligned}
& \left(\sum_{\substack{W \in \mathcal{D}(5\Delta) \\ l(W) \approx \tau}} \sup_{x \in W} |\partial_s A(x, s)|^2 \int_W |\partial_x e^{-\eta^2 \tau^2 L^s} \phi^s(x)|^2 dx \right)^{1/2} \\
& \lesssim \left(\sum_{\substack{W \in \mathcal{D}(5\Delta) \\ l(W) \approx \tau}} \sup_{x \in W} |\partial_s A(x, s)|^2 \int_W |M[\partial_x \phi^s]|^2 dx \right)^{1/2} \\
& \lesssim \left(\int_{5\Delta} \sup_{x \in B(x, s, s/2)} |\partial_s A(x, s)|^2 dx \right)^{1/2}.
\end{aligned}$$

Hence in total we get

$$\begin{aligned}
& \int_{T(3\Delta)} \frac{|w_s^{(2)}|^2}{s} dx ds \\
& \lesssim \eta^2 \int_0^{3l(\Delta)} \frac{1}{s} \left(\int_0^s \frac{\tau}{\sqrt{s^2 - \tau^2}} \left(|\Delta|^{1/2} + \left(\int_{5\Delta} \sup_{x \in B(x, s, s/2)} |\partial_s A(x, s)|^2 dx \right)^{1/2} \right) d\tau \right)^2 ds \\
& \lesssim \eta^2 \int_0^{3l(\Delta)} \frac{1}{s} \left(|\Delta| + \int_{5\Delta} \sup_{x \in B(x, s, s/2)} |\partial_s A(x, s)|^2 dx \right) \left(\int_0^s \frac{\tau}{\sqrt{s^2 - \tau^2}} d\tau \right)^2 ds \\
& \lesssim \eta^2 \int_0^{3l(\Delta)} s \left(|\Delta| + \int_{5\Delta} \sup_{x \in B(x, s, s/2)} |\partial_s A(x, s)|^2 dx \right) ds \\
& \lesssim \eta^2 l(\Delta)^2 |\Delta| + \int_{T(5\Delta)} \sup_{x \in B(x, s, s/2)} |\partial_s A(x, s)|^2 s dx ds \lesssim |\Delta|.
\end{aligned}$$

□

Lemma 6.4. *Let $Q \subset \partial\Omega$ be a boundary cube of size s , i.e. $l(Q) \approx s$, $(t_j)_{j \in \mathbb{Z}} = ((c^2)^j)_{j \in \mathbb{Z}}$ be a partition of $(0, \infty)$ for a constant $\frac{1}{2} < c < 1$, and the index i such that $s \in (t_i, t_{i+1})$. We have the following local bound involving $w_s^{(2)}$*

$$\int_Q |\partial_x w_s^{(2)}(x, s)|^2 dx \lesssim \frac{1}{s^2} \int_{t_{i-1}}^{t_{i+2}} \int_{2Q} |v(x, k, s)|^2 dk dx + \|\partial_s A\|_{L^\infty(2Q \times [t_{i-3}, t_{i+4}])}^2 |Q| \kappa_0^2. \tag{6.5}$$

As a consequence we have the square function bound

$$\int_{T(\Delta)} |\partial_x w_s^{(2)}|^2 s ds \lesssim |\Delta| \tag{6.6}$$

and

$$\int_{T(\Delta)} |\partial_x w_s^{(2)}| ds \lesssim |\Delta|. \tag{6.7}$$

Proof. To start with note that from [Ulm24] we already know that

$$\begin{aligned}
\int_Q |\partial_x w_s^{(2)}(x, s)|^2 dx &\lesssim \frac{1}{s^2} \int_{t_{l-1}}^{t_{l+2}} \int_{2Q} |v(x, k, s)|^2 dk dx \\
&+ \|\partial_s A(\cdot, s)\|_{L^\infty(2Q \times (t_{l-3}, t_{l+4}))}^2 \int_{t_{l-3}}^{t_{l+4}} \int_{2Q} |\partial_x e^{-t^2 L^s} \phi_s|^2 dx dt \\
&+ \|\partial_s A(\cdot, s)\|_{L^\infty(2Q \times (t_{l-3}, t_{l+4}))}^2 s^2 \int_{t_{l-3}}^{t_{l+4}} \int_{2Q} |\partial_x \partial_k e^{-k^2 L^s} \phi_s|^2 dx dk.
\end{aligned} \tag{6.8}$$

By [Lemma 4.2](#) we obtain that

$$\int_{t_{l-3}}^{t_{l+4}} \int_{2Q} |\partial_x e^{-k^2 L^s} \phi_s|^2 + s^2 |\partial_x \partial_k e^{-k^2 L^s} \phi_s|^2 dx dk \lesssim |Q| \inf_{x \in Q} M[\Phi](x) \lesssim \kappa_0 |Q|,$$

whence [\(6.5\)](#) follows.

To show [\(6.6\)](#) we note that

$$\begin{aligned}
\int_{3\Delta} |v(x, k, s)|^2 dx &\leq \left(\int_0^k 2\tau \|e^{-(k^2 - \tau^2)L^s} \partial_x (\partial_s A_{\parallel}) \partial_x e^{-\tau^2 L^s} \phi^s\|_{L^2} d\tau \right)^2 \\
&\leq \left(\int_0^k \frac{2\tau}{\sqrt{k^2 - \tau^2}} \|\partial_s A_{\parallel} \partial_x e^{-\tau^2 L^s} \phi^s\|_{L^2} d\tau \right)^2.
\end{aligned}$$

Observing that [Lemma 4.2](#), $L^2 - L^2$ off-diagonal estimates in [Proposition 3.7](#), and Poincaré yield

$$\begin{aligned}
\|\partial_s A_{\parallel} \partial_x e^{-\tau^2 L^s} \phi^s\|_{L^2} &\lesssim \left(\sum_{\substack{Q \subset 5\Delta \\ Q \text{ finite overlap, } l(Q) \approx s \approx k}} \sup_Q |\partial_s A_{\parallel}|^2 |Q| \kappa_0^2 \right)^{1/2} \\
&\quad + \frac{1}{s} \|\partial_x e^{-\tau^2 L^s} \phi^s\|_{L^2(\mathbb{R}^n \setminus 5\Delta)} \\
&\lesssim \left(\int_{5\Delta} \sup_{B(x, s, s/2)} |\partial_s A_{\parallel}|^2 dx \right)^{1/2} + \frac{1}{s} \frac{1}{\tau} e^{-c \frac{l(\Delta)^2}{\tau^2}} \|\phi^s\|_{L^2(\mathbb{R}^n \setminus 5\Delta)} \\
&\lesssim \left(\int_{5\Delta} \sup_{B(x, s, s/2)} |\partial_s A_{\parallel}|^2 dx \right)^{1/2} + \frac{1}{l(\Delta)^2} \|\phi^s\|_{L^2(\mathbb{R}^n \setminus 5\Delta)} \\
&\lesssim \left(\int_{5\Delta} \sup_{B(x, s, s/2)} |\partial_s A_{\parallel}|^2 dx \right)^{1/2} + \frac{1}{\sqrt{l(\Delta)}},
\end{aligned}$$

gives

$$\int_{3\Delta} |v(x, k, s)|^2 dx \leq s^2 \int_{5\Delta} \sup_{B(x, s, s/2)} |\partial_s A_{\parallel}|^2 dx + \frac{s^4}{l(\Delta)}.$$

Hence in total, we obtain

$$\begin{aligned} \int_{T(3\Delta)} |\partial_x w_s^{(2)}(x, s)|^2 s dx ds &\lesssim \int_{T(3\Delta)} 2 \sup_{B(x, s, s/2)} |\partial_s A_{\parallel}|^2 s dx ds + \int_0^{l(\Delta)} \frac{s^3}{l(\Delta)} ds \\ &\lesssim |\Delta|, \end{aligned}$$

which completes the proof of (6.6).

Lastly for (6.7), we first fix $s > 0$ and let $Q_j = Q_j^s \subset 5\Delta$ such that $l(Q_j) \approx s$, the Q_j and Q_{j+1} have nontrivial intersection but all the other pairs Q_j and Q_i do not intersect, and the union of the Q_j covers 3Δ . Then we observe for a dualising functions $h_i \in L^2(Q_i)$ and $Q_i \subset 3\Delta$

$$\begin{aligned} \left(\int_{Q_i} |v(x, k, s)|^2 dx \right)^{1/2} &\leq \int_0^k 2\tau \|e^{-(k^2-\tau^2)L^s} \partial_x (\partial_s A_{\parallel} \partial_x e^{-\tau^2 L^s} \phi^s)\|_{L^2(Q)} d\tau \\ &\leq \int_0^k \tau \int_{\mathbb{R}^n} \partial_s A_{\parallel} \partial_x e^{-\tau^2 L^s} \phi^s \partial_x e^{-(k^2-\tau^2)L^s} h_i dx d\tau \\ &\leq \int_0^k \tau \left(\int_{4\Delta} \partial_s A_{\parallel} \partial_x e^{-\tau^2 L^s} \phi^s \partial_x e^{-(k^2-\tau^2)L^s} h_i dx \right. \\ &\quad \left. + \frac{1}{s} \sum_{j \geq 2} \int_{2^{j+1}\Delta \setminus 2^j\Delta} \partial_s A_{\parallel} \partial_x e^{-\tau^2 L^s} \phi^s \partial_x e^{-(k^2-\tau^2)L^s} h_i dx \right) d\tau. \end{aligned}$$

For the first term we have by [Lemma 4.2](#) and Cauchy Schwarz inequality and $L^2 - L^2$ off diagonal estimates ([Proposition 3.7](#))

$$\begin{aligned} &\int_{4\Delta} \partial_s A_{\parallel} \partial_x e^{-\tau^2 L^s} \phi^s \partial_x e^{-(k^2-\tau^2)L^s} h_i dx \\ &\leq \sum_j \int_{Q_j} \partial_s A_{\parallel} \partial_x e^{-\tau^2 L^s} \phi^s \partial_x e^{-(k^2-\tau^2)L^s} h_i dx \\ &\lesssim \sum_j \sup_{Q_j} |\partial_s A_{\parallel}| \kappa_0 \sqrt{|Q_j|} \frac{1}{\sqrt{k^2-\tau^2}} e^{-c\frac{|i-j|^2 s^2}{k^2-\tau^2}} \|h_i\|_{L^2} \\ &\lesssim \sum_j \sup_{Q_j} |\partial_s A_{\parallel}| \kappa_0 \frac{1}{\sqrt{s}} e^{-c|i-j|^2} \|h_i\|_{L^2}. \end{aligned}$$

We can also notice that summing over i yields

$$\begin{aligned} &\sum_i \int_{4\Delta} \partial_s A_{\parallel} \partial_x e^{-\tau^2 L^s} \phi^s \partial_x e^{-(k^2-\tau^2)L^s} h_i dx \\ &\lesssim \sum_j \sup_{Q_j} |\partial_s A_{\parallel}| \kappa_0 \frac{1}{\sqrt{s}} \sum_i e^{-c|i-j|^2} \|h_i\|_{L^2} \\ &\lesssim \sum_j \sup_{Q_j} |\partial_s A_{\parallel}| \kappa_0 \frac{1}{\sqrt{s}}. \end{aligned}$$

For the second term we have by $L^2 - L^2$ off diagonal estimates

$$\begin{aligned}
& \int_{2^{j+1}\Delta \setminus 2^j\Delta} \partial_s A_{\parallel} \partial_x e^{-\tau^2 L^s} \phi^s \partial_x e^{-(k^2 - \tau^2)L^s} h_i dx \\
& \lesssim \frac{1}{s} \|\partial_x e^{-\tau^2 L^s} \phi^s\|_{L^2(2^{j+1}\Delta \setminus 2^j\Delta)} \|\partial_x e^{-(k^2 - \tau^2)L^s} h_i\|_{L^2(2^{j+1}\Delta \setminus 2^j\Delta)} \\
& \lesssim \frac{1}{s} 2^{(j-1)/2} l(\Delta)^{1/2} \kappa_0 \frac{1}{\sqrt{k^2 - \tau^2}} e^{-c \frac{2^{2j} l(\Delta)^2}{k^2 - \tau^2}} \|h_i\|_{L^2(Q)}.
\end{aligned}$$

Summing over j yields

$$\begin{aligned}
& \sum_j \frac{1}{s} 2^{(j-1)/2} l(\Delta)^{1/2} \kappa_0 \frac{1}{\sqrt{k^2 - \tau^2}} e^{-c \frac{2^{2j} l(\Delta)^2}{k^2 - \tau^2}} \|h_i\|_{L^2(Q)} \\
& \lesssim \frac{1}{s} l(\Delta)^{1/2} \kappa_0 \frac{1}{\sqrt{k^2 - \tau^2}} \frac{(k^2 - \tau^2)^{3/4}}{l(\Delta)^{3/2}} \\
& \lesssim \frac{1}{\sqrt{s} l(\Delta)}.
\end{aligned}$$

In total this concludes the proof of (6.7) by

$$\begin{aligned}
\int_{T(\Delta)} |\partial_x w_s^{(2)}| dx ds & \lesssim \int_0^{l(\Delta)} \sum_{Q_i=Q_i^s} \left(\int_{Q_i} |\partial_x w_s^{(2)}|^2 dx \right)^{\frac{1}{2}} |Q_i|^{\frac{1}{2}} ds \\
& \lesssim \int_0^{l(\Delta)} \sum_{Q_i=Q_i^s} \left(\frac{1}{s} \int_{t_i}^{t_{i+1}} \int_{Q_i} |v|^2 dx dk + \sup_{B(x,s,s/2)} |\partial_s A_{\parallel}| |Q_i|^{1/2} \right) |Q_i|^{\frac{1}{2}} ds \\
& \lesssim \int_0^{l(\Delta)} \frac{1}{s} \int_{t_i}^{t_{i+1}} \int_0^k \tau \sum_{Q_j=Q_j^s} \left(\sup_{Q_j} |\partial_s A_{\parallel}| \kappa_0 + \frac{1}{l(\Delta)} \right) d\tau ds \\
& \quad + \int_0^{l(\Delta)} \sum_{Q_i=Q_i^s} \sup_{B(x,s,s/2)} |\partial_s A_{\parallel}| |Q_i| ds \\
& \lesssim \int_0^{l(\Delta)} \sum_{Q_j=Q_j^s} \sup_{Q_j} |\partial_s A_{\parallel}| s ds + \int_0^{l(\Delta)} \frac{l(\Delta)}{s} \frac{s}{l(\Delta)} ds \\
& \quad + \int_{T(3\Delta)} \sup_{B(x,s,s/2)} |\partial_s A_{\parallel}| dx ds \\
& \lesssim \int_{T(3\Delta)} \sup_{B(x,s,s/2)} |\partial_s A_{\parallel}| dx ds + l(\Delta).
\end{aligned}$$

□

6.3 The partial derivative part w_t

Lemma 6.9. *The following square function bounds involving w_t hold:*

$$(i) \quad \int_{T(\Delta)} \frac{|w_t|^2}{s} dx ds = \int_{T(\Delta)} \frac{|sL^s e^{-\eta^2 s^2 L^s} \phi^s|^2}{s} dx ds \lesssim |\Delta|;$$

$$(ii) \quad \int_{T(\Delta)} \frac{|s\partial_x w_t|^2}{s} dx ds \lesssim |\Delta|;$$

$$(iii) \quad \int_{T(\Delta)} \frac{|s^2 L^s w_t|^2}{s} dx ds \lesssim |\Delta|.$$

Proof. We start with proving (i). First, note that we can write $A_{\parallel} = B^2$ for $B > 0$. And hence we obtain

$$\begin{aligned} \int_{T(\Delta)} \frac{|sL^s e^{-\eta^2 s^2 L^s} \phi^s|^2}{s} dx ds &\leq \int_{\mathbb{R}^n \times (0, l(\Delta))} \frac{|sL^s e^{-\eta^2 s^2 L^s} \phi^s|^2}{s} dx ds \\ &= \int_{\mathbb{R}^n \times (0, l(\Delta))} B\partial_x L^s e^{-\eta^2 t L^s} \phi^s \cdot B\partial_x e^{-\eta^2 s^2 L^s} \phi^s dx dt \\ &= \int_{\mathbb{R}^n \times (0, l(\Delta))} B\partial_s L^s e^{-\eta^2 s^2 L^s} \phi^s \cdot B\partial_x e^{-\eta^2 s^2 L^s} \phi^s dx ds \\ &\quad - \int_{\mathbb{R}^n \times (0, l(\Delta))} B\partial_x w_s^{(1)} \cdot B\partial_x e^{-\eta^2 s^2 L^s} \phi^s dx ds \\ &\quad - \int_{\mathbb{R}^n \times (0, l(\Delta))} B\partial_x w_s^{(2)} \cdot B\partial_x e^{-\eta^2 s^2 L^s} \phi^s dx ds \\ &= \int_0^{l(\Delta)} \partial_s \|B\partial_x e^{-\eta^2 s^2 L^s} \phi^s\|_{L^2(\mathbb{R}^n)}^2 dx ds \\ &\quad + \int_{\mathbb{R}^n \times (0, l(\Delta))} \partial_s A_{\parallel} \partial_x e^{-\eta^2 s^2 L^s} \phi^s \cdot \partial_x e^{-\eta^2 s^2 L^s} \phi^s dx ds \\ &\quad - \int_{\mathbb{R}^n \times (0, l(\Delta))} B\partial_x w_s^{(1)} \cdot B\partial_x e^{-\eta^2 s^2 L^s} \phi^s dx ds \\ &\quad - \int_{\mathbb{R}^n \times (0, l(\Delta))} B\partial_x w_s^{(2)} \cdot B\partial_x e^{-\eta^2 s^2 L^s} \phi^s dx ds \\ &=: I + II + III + IV. \end{aligned}$$

Looking at the integrals separately, first we have

$$I = \|B\partial_x e^{-\eta^2 (3l(\Delta))^2 L^s} \phi^{3l(\Delta)}\|_{L^2(\mathbb{R}^n)}^2 - \|B\partial_x \phi^0\|_{L^2(\mathbb{R}^n)}^2 \lesssim |\Delta|,$$

and

$$II \lesssim \|\sup |\partial_s A_{\parallel}|\|_c \int_{\mathbb{R}^n} \tilde{N}_2(|\partial_x e^{-\eta^2 s^2 L^s} \phi^s|)^2 dx.$$

From [Lemma 4.1](#) we have

$$\int_{5\Delta} \tilde{N}_2(|\partial_x e^{-\eta^2 s^2 L^s} \phi^s|)^2 dx \lesssim |\Delta|,$$

and using $L^2 - L^2$ off diagonal estimates and Poincaré's inequality, we also have

$$\begin{aligned} \int_{\mathbb{R}^n \setminus 4\Delta} \tilde{N}_2(|\partial_x e^{-\eta^2 s^2 L^s} \phi^s|)^2 dx &\lesssim \sum_{j \geq 3} \int_{2^j \Delta \setminus 2^{j-1} \Delta} \tilde{N}_2(|\partial_x e^{-\eta^2 s^2 L^s} \phi^s|)^2 dx \\ &\lesssim \sum_{j \geq 3} \int_{2^j \Delta \setminus 2^{j-1} \Delta} \frac{1}{2^{2j}} \frac{1}{l(\Delta)^2} \sup_s \|\phi^s\|_{L^2}^2 dx \\ &\lesssim \sum_{j \geq 3} \int_{2^j \Delta \setminus 2^{j-1} \Delta} \frac{1}{2^{2j}} dx \\ &\lesssim |\Delta|. \end{aligned}$$

Next, we have by [Lemma 4.2](#)

$$\begin{aligned} III &= \int_0^{l(\Delta)} \int_{\mathbb{R}^n} w_s^{(1)}(x, s) L^s e^{-\eta^2 s^2 L^s} \phi^s dx ds \\ &= \int_0^{l(\Delta)} \int_{\mathbb{R}^n} A \partial_x \partial_s \phi^s(x, s) \partial_x e^{-2\eta^2 s^2 L^s} \phi^s dx ds \\ &\leq \int_0^{l(\Delta)} \sum_{\substack{Q \subset 5\Delta \\ l(Q) \approx s, Q \text{ finite overlap}}} \sup_{y \in Q} (|\partial_s A|(y, s)| + \int_{3\Delta} |\partial_s A| dx) \left(\int_Q |\partial_x e^{-2\eta^2 s^2 L^s} \phi^s|^2 \right)^{1/2} ds \\ &\leq \int_{T(5\Delta)} \sup_{(y,t) \in B(x,s,s/2)} |\partial_s A(y, t)| dx ds \lesssim |\Delta|. \end{aligned}$$

At last, we have

$$\begin{aligned} |IV| &= \int_0^{l(\Delta)} \int_0^s \int_{\mathbb{R}^n} |2\tau \partial_s A| |\partial_x e^{-\eta^2 \tau^2 L^s} \phi^s \cdot \partial_x e^{-\eta^2 (s^2 - \tau^2) L^s} L^s e^{-\eta^2 s^2 L^s} \phi^s| dx dt ds \\ &\leq \|\sup \partial_s A\|_C \int_{\mathbb{R}^n} \tilde{N} \left(\int_0^s \tau |\partial_x e^{-\eta^2 \tau^2 L^s} \phi^s| |\partial_x e^{-\eta^2 (s^2 - \tau^2) L^s} L^s e^{-\eta^2 s^2 L^s} \phi^s| d\tau \right) (x) dx \end{aligned}$$

We note that the appearing nontangential maximal function is pointwise bounded.

To see this we take $x \in 5\Delta, t > 0$ and have

$$\begin{aligned} &\int_{B(x,t,t/2)} \int_0^s \tau |\partial_x e^{-\eta^2 \tau^2 L^s} \phi^s| |\partial_x e^{-\eta^2 (s^2 - \tau^2) L^s} L^s e^{-\eta^2 s^2 L^s} \phi^s| d\tau dx ds \\ &\lesssim \int_{t/2}^{3t/2} \int_0^s \tau \left(\int_{B(x,t,t/2)} |\partial_x e^{-\eta^2 \tau^2 L^s} \phi^s|^2 dx \right)^{1/2} \left(\int_{B(x,t,t/2)} |\partial_x e^{-\eta^2 (s^2 - \tau^2) L^s} L^s e^{-\eta^2 s^2 L^s} \phi^s|^2 dx \right)^{1/2} d\tau ds \end{aligned}$$

By $L^2 - L^2$ -off diagonal estimates ([Proposition 3.7](#)), [Lemma 4.2](#), and [Corollary 4.3](#) we obtain

$$\left(\int_{B(x,t,t/2)} |\partial_x e^{-\eta^2 \tau^2 L^s} \phi^s|^2 dx \right)^{1/2} \lesssim \frac{1}{t} \frac{t}{\tau} \tau \kappa_0 \lesssim \kappa_0,$$

and

$$\begin{aligned}
\left(\int_{B(x,t/2)} |\partial_x e^{-\eta^2(s^2-\tau^2)L^s} L^s e^{-\eta^2 s^2 L^s} \phi^s|^2 dx \right)^{1/2} &\lesssim \frac{1}{\sqrt{s^2-\tau^2}} M[|L^s e^{-\eta^2 s^2 L^s} \phi^s|^2]^{1/2} \\
&\lesssim \frac{1}{\sqrt{s^2-\tau^2}} \frac{1}{s} M[|w_t(\cdot, s)|^2]^{1/2} \\
&\lesssim \frac{1}{\sqrt{s^2-\tau^2}} \frac{1}{s} M[M[|\partial_x \phi^s|^2]]^{1/2}.
\end{aligned}$$

Hence in total with [Lemma 4.1](#)

$$\begin{aligned}
&\int_{B(x,t,t/2)} \int_0^s \tau |\partial_x e^{-\eta^2 \tau^2 L^s} \phi^s| |\partial_x e^{-\eta^2(s^2-\tau^2)L^s} L^s e^{-\eta^2 s^2 L^s} \phi^s| d\tau dx ds \\
&\lesssim \int_{t/2}^{3t/2} \int_0^s \tau \frac{1}{\sqrt{s^2-\tau^2}} \frac{1}{s} d\tau ds \leq C.
\end{aligned}$$

and

$$\begin{aligned}
&\int_{5\Delta} \tilde{N} \left(\int_0^s \tau |\partial_x e^{-\eta^2 \tau^2 L^s} \phi^s| |\partial_x e^{-\eta^2(s^2-\tau^2)L^s} L^s e^{-\eta^2 s^2 L^s} \phi^s| d\tau \right) (x) dx \\
&\lesssim |\Delta|
\end{aligned}$$

For $x \in 2^j \Delta \setminus 2^{j-1} \Delta \mathbb{R}^n \setminus 4\Delta$, $3l(\Delta) > t > 0$ and $j \geq 3$ however, we obtain with $L^2 - L^2$ off diagonal estimates

$$\begin{aligned}
\left(\int_{B(x,t/2)} |\partial_x e^{-\eta^2 \tau^2 L^s} \phi^s|^2 dx \right)^{1/2} &\lesssim \frac{1}{\sqrt{t}} \frac{1}{\tau} e^{-c \frac{2^{2j} l(\Delta)^2}{\tau^2}} \|\phi^s\|_{L^2(3\Delta)} \lesssim \frac{1}{\sqrt{t}} \frac{1}{2^j l(\Delta)} \|\phi^s\|_{L^2(3\Delta)} \\
&\lesssim \frac{1}{2^j \sqrt{t}} \|\partial_x \phi^s\|_{L^2(3\Delta)} \lesssim \frac{1}{2^j \sqrt{t}} \sqrt{|\Delta|},
\end{aligned}$$

and

$$\begin{aligned}
\left(\int_{B(x,t/2)} |\partial_x e^{-\eta^2(s^2-\tau^2)L^s} L^s e^{-\eta^2 s^2 L^s} \phi^s|^2 dx \right)^{1/2} &\lesssim \frac{1}{\sqrt{t}} \frac{1}{\sqrt{s^2-\tau^2}} e^{-c \frac{2^{2j} l(\Delta)^2}{s^2-\tau^2}} \|L^s e^{-\eta^2 s^2 L^s} \phi^s\|_{L^2} \\
&\lesssim \frac{1}{\sqrt{t}} \frac{1}{2^j l(\Delta)} \frac{1}{s} \|M[\partial_x \phi^s]\|_{L^2(3\Delta)} \\
&\lesssim \frac{1}{2^j \sqrt{t}} \frac{1}{s} \frac{1}{\sqrt{|\Delta|}} \kappa_0
\end{aligned}$$

Hence we obtain for $x \in 2^j \Delta \setminus 2^{j-1} \Delta$

$$\begin{aligned}
&\int_{B(x,t,t/2)} \int_0^s \tau |\partial_x e^{-\eta^2 \tau^2 L^s} \phi^s| |\partial_x e^{-\eta^2(s^2-\tau^2)L^s} L^s e^{-\eta^2 s^2 L^s} \phi^s| d\tau dx ds \\
&\lesssim \int_{t/2}^{3t/2} \int_0^s \tau \frac{1}{2^j \sqrt{t}} \sqrt{|\Delta|} \frac{1}{2^j \sqrt{t}} \frac{1}{s} \frac{1}{\sqrt{|\Delta|}} d\tau ds \leq \frac{1}{2^{2j}},
\end{aligned}$$

and

$$\begin{aligned}
& \int_{\mathbb{R}^n \setminus 5\Delta} \tilde{N} \left(\int_0^s \tau |\partial_x e^{-\eta^2 \tau^2 L^s} \phi^s| |\partial_x e^{-\eta^2 (s^2 - \tau^2) L^s} L^s e^{-\eta^2 s^2 L^s} \phi^s| d\tau \right) (x) dx \\
& \lesssim \sum_{j \geq 3} \int_{2^j \Delta \setminus 2^{j-1} \Delta} \tilde{N} \left(\int_0^s \tau |\partial_x e^{-\eta^2 \tau^2 L^s} \phi^s| |\partial_x e^{-\eta^2 (s^2 - \tau^2) L^s} L^s e^{-\eta^2 s^2 L^s} \phi^s| d\tau \right) (x) dx \\
& \lesssim \sum_{j \geq 3} \int_{2^j \Delta \setminus 2^{j-1} \Delta} \frac{1}{2^{2j}} dx \lesssim |\Delta|,
\end{aligned}$$

whence the integral $|IV| \lesssim |\Delta|$.

The proofs of (ii) and (iii) rely on (i) and [Proposition 3.3](#). For (ii) we see that

$$\begin{aligned}
& \int_0^{l(\Delta)} \int_{\Delta} \frac{|s^2 \partial_x L^s e^{-\eta^2 s^2 L^s} \phi^s|^2}{s} dx ds \\
& \leq \int_0^{l(\Delta)} \int_{\mathbb{R}^n} \frac{|s^2 \partial_x L^s e^{-\eta^2 s^2 L^s} \phi^s|^2}{s} dx ds \\
& = \int_0^{l(\Delta)} \int_{\mathbb{R}^n} \frac{|s \partial_x e^{-\frac{\eta^2 s^2}{2} L^s} (s L^s e^{-\frac{\eta^2 s^2}{2} L^s} \phi^s)|^2}{s} dx ds \\
& \lesssim \int_0^{l(\Delta)} \int_{\mathbb{R}^n} \frac{|s L^s e^{-\frac{\eta^2 s^2}{2} L^s} \phi^s|^2}{s} dx ds,
\end{aligned}$$

where the last integral is bounded by (i) after a change of variable argument. Similar we observe for (iii)

$$\begin{aligned}
& \int_{T(\Delta)} \frac{|s^2 L^s L^s e^{-\eta^2 s^2 L^s} \phi^s|^2}{s} dx ds \\
& = \int_0^{l(\Delta)} \int_{\mathbb{R}^n} \frac{|s^2 L^s L^s e^{-\eta^2 s^2 L^s} \phi^s|^2}{s} dx ds \\
& = \int_0^{l(\Delta)} \int_{\mathbb{R}^n} \frac{|s L^s e^{-\frac{\eta^2 s^2}{2} L^s} (s L^s e^{-\frac{\eta^2 s^2}{2} L^s} \phi^s)|^2}{s} dx ds \\
& \lesssim \int_0^{l(\Delta)} \int_{\mathbb{R}^n} \frac{|s L^s e^{-\frac{\eta^2 s^2}{2} L^s} \phi^s|^2}{s} dx ds.
\end{aligned}$$

□

In contrast to $\partial_s \phi^s - w_s^{(1)}$ and $w_s^{(2)}$ we have a certain local Harnack-type property for w_t . First we have

Lemma 6.10. *Let $Q \subset \mathbb{R}^n$ be a cube with side length $l(Q) \approx s \approx R_0$ and $\tilde{Q} := (1 + \varepsilon)Q$ be an enlarged cube for some fixed $\frac{1}{2} > \varepsilon > 0$. Then*

$$\begin{aligned} \sup_{(x,s) \in Q \times (R_0, 2R_0]} w_t(x, s) &\lesssim \int_{(1-\varepsilon)R_0}^{2(1+\varepsilon)R_0} \int_{\tilde{Q} \times ((1-\varepsilon)R_0, 2(1+\varepsilon)R_0)} \frac{|w_t(x, s)|^2}{s} dx ds \\ &\quad + \int_{(1-\varepsilon)R_0}^{2(1+\varepsilon)R_0} \int_{\tilde{Q} \times ((1-\varepsilon)R_0, 2(1+\varepsilon)R_0)} |\partial_x v_1(x, t^2, s)|^2 s dx dt ds \\ &\quad + \int_{(1-\varepsilon)R_0}^{2(1+\varepsilon)R_0} \int_{\tilde{Q} \times ((1-\varepsilon)R_0, 2(1+\varepsilon)R_0)} |\partial_x v_2(x, t^2, s)|^2 s dx dt ds. \end{aligned}$$

As a consequence, we have

$$\sum_{k \leq k_0} \sum_{Q \in \mathcal{D}_k(\Delta)} |Q| \sup_{Q \times (2^{-k}, 2^{-k+1}]} |w_t|^2 \lesssim |\Delta|.$$

The consequence in above lemma resembles a version of (i) in [Lemma 6.9](#) where the supremum is taken on each Whitney cube. Hence it is not surprising that it holds, but the proof is technical.

Proof. First, we would like to refer to the proof of Lemma 7.9 in [\[Ulm24\]](#), where an intermediary step proves

$$\begin{aligned} \sup_{(x,s) \in Q \times (R_0, 2R_0]} w_t(x, s) &\lesssim \int_{(1-\varepsilon)R_0}^{2(1+\varepsilon)R_0} \int_{\tilde{Q} \times ((1-\varepsilon)R_0, 2(1+\varepsilon)R_0)} \frac{|w_t(x, s)|^2}{s} dx ds \\ &\quad + \int_{(1-\varepsilon)R_0}^{2(1+\varepsilon)R_0} \int_{\tilde{Q} \times ((1-\varepsilon)R_0, 2(1+\varepsilon)R_0)} |\partial_x v_1(x, t^2, s)|^2 s dx dt ds \\ &\quad + \int_{(1-\varepsilon)R_0}^{2(1+\varepsilon)R_0} \int_{\tilde{Q} \times ((1-\varepsilon)R_0, 2(1+\varepsilon)R_0)} |\partial_x v_2(x, t^2, s)|^2 s dx dt ds. \end{aligned}$$

By [Lemma 4.1](#) and [Lemma 4.2](#) we obtain

$$\begin{aligned} &\int_{(1-\varepsilon)R_0}^{2(1+\varepsilon)R_0} \int_{\tilde{Q} \times ((1-\varepsilon)R_0, 2(1+\varepsilon)R_0)} |\partial_x v_1(x, t^2, s)|^2 s dx dt ds \\ &\lesssim \int_{(1-\varepsilon)R_0}^{2(1+\varepsilon)R_0} \int_{\tilde{Q} \times ((1-\varepsilon)R_0, 2(1+\varepsilon)R_0)} |M[\partial_s \partial_x \phi^s]|^2 s dx dt ds \\ &\lesssim \int_{\tilde{Q} \times ((1-\varepsilon)R_0, 2(1+\varepsilon)R_0)} \sup_{\tilde{Q} \times ((1-\varepsilon)R_0, 2(1+\varepsilon)R_0)} |\partial_s A_{\parallel}|^2 s dx ds \\ &\quad + \int_{\tilde{Q} \times ((1-\varepsilon)R_0, 2(1+\varepsilon)R_0)} \int_{\Delta} |\partial_s A_{\parallel}|^2 s dx, \end{aligned}$$

whence

$$\begin{aligned} & \sum_{k \leq k_0} \sum_{Q \in \mathcal{D}_k(\Delta)} \int_{\hat{Q} \times ((1-\varepsilon)R_0, 2(1+\varepsilon)R_0)} \int_{(1-\varepsilon)2^{-k}}^{(1+\varepsilon)2^{-k+1}} |\partial_x v_1(x, t^2, s)|^2 s dt dx ds \\ & \lesssim \int_{5\Delta} \sup_{B(x, s, s/2)} |\partial_s A|^2 s dx ds \lesssim |\Delta|. \end{aligned}$$

Furthermore, we can refer to the proof of (6.6) in [Lemma 6.4](#) to see that

$$\sum_{k \leq k_0} \sum_{Q \in \mathcal{D}_k(\Delta)} \int_{\hat{Q} \times ((1-\varepsilon)R_0, 2(1+\varepsilon)R_0)} \int_{(1-\varepsilon)2^{-k}}^{(1+\varepsilon)2^{-k+1}} |\partial_x v_2(x, t^2, s)|^2 s dt dx ds \lesssim |\Delta|.$$

Hence

$$\sum_{k \leq k_0} \sum_{Q \in \mathcal{D}_k(\Delta)} |Q| \sup_{Q \times (2^{-k}, 2^{-k+1})} |w_t|^2 \lesssim \int_{3\Delta} \frac{|w_t(x, s)|^2}{s} dx ds + |\Delta| \lesssim |\Delta|.$$

□

7 L^2 estimates for square functions

Lemma 7.1. *Assuming the L^1 Carleson condition (1.4) we have the following estimates:*

$$(i) \quad \int_{T(\Delta)} |\nabla \partial_s \rho(x, s)|^2 s dx ds \lesssim |\Delta|;$$

$$(ii) \quad \int_{T(\Delta)} |L^s \rho(x, s)|^2 s dx ds \lesssim |\Delta|;$$

$$(iii) \quad \int_{T(\Delta)} |\partial_s A(x, s) \nabla \rho(x, s)|^2 s dx ds \lesssim |\Delta|;$$

$$(iv) \quad \int_{T(\Delta)} \frac{|\partial_s \theta(x, s)|^2}{s} dx ds \lesssim |\Delta|;$$

$$(v) \quad \int_{T(\Delta)} \frac{|\theta(x, s)|^2}{s^3} dx ds \lesssim |\Delta|.$$

Proof. (i) We split

$$\begin{aligned} \int_{T(\Delta)} |\nabla \partial_s \rho(x, s)|^2 s dx ds &\leq \int_{T(\Delta)} |\nabla w_t(x, s)|^2 s dx ds + \int_{T(\Delta)} |\nabla w_s^{(1)}(x, s)|^2 s dx ds \\ &\quad + \int_{T(\Delta)} |\nabla w_s^{(2)}(x, s)|^2 s dx ds, \end{aligned}$$

and apply [Lemma 6.9](#), [Lemma 6.1](#) and [Lemma 6.4](#) to get the required bound.

(ii) We can observe that

$$\int_{T(\Delta)} |L^s \rho(x, s)|^2 s dx ds = \int_{T(\Delta)} \frac{|w_t(x, s)|^2}{s} dx ds \lesssim |\Delta|$$

by [Lemma 6.9](#).

(iii) We can calculate

$$\begin{aligned} \int_{T(\Delta)} |\partial_s A(x, s) \nabla \rho(x, s)|^2 s dx ds &\leq \int_0^{l(\Delta)} \|\partial_s A(\cdot, s)\|_\infty \left(\int_\Delta |\nabla \phi^s|^2 + |\nabla e^{-\eta^2 s^2 L^s} \phi^s|^2 dx \right) s ds \\ &\lesssim \int_0^{l(\Delta)} \|\partial_s A(\cdot, s)\|_\infty |\Delta| s ds \lesssim |\Delta| \end{aligned}$$

by [\(2.3\)](#).

(iv) We split

$$\begin{aligned} \int_{T(\Delta)} \frac{|\partial_s \theta(x, s)|^2}{s} dx ds &\leq \int_{T(\Delta)} \frac{|(\partial_s \phi^s - w_s^{(1)})(x, s)|^2}{s} dx ds + \int_{T(\Delta)} \frac{|w_s^{(2)}(x, s)|^2}{s} dx ds \\ &\quad + \int_{T(\Delta)} \frac{|w_t(x, s)|^2}{s} dx ds \end{aligned}$$

and the bound follows from [Lemma 6.1](#), [Lemma 6.4](#), and [Lemma 6.9](#).

(v) The proof works analogously to the proof of [Lemma 9](#) in [\[HLMP22a\]](#). \square

References

- [AEN18] Pascal Auscher, Moritz Egert, and Kaj Nyström. The Dirichlet problem for second order parabolic operators in divergence form. *Journal de l'École polytechnique — Mathématiques*, 5:407–441, 2018.
- [AHL⁺02] Pascal Auscher, Steve Hofmann, Michael Lacey, Alan McIntosh, and Ph. Tchamitchian. The solution of the kato square root problem for second order elliptic operators on rn. *Annals of Mathematics*, 156(2):633–654, 2002.

- [Aus07] Pascal Auscher. On necessary and sufficient conditions for L^p -estimates of Riesz transforms associated to elliptic operators on \mathbb{R}^n and related estimates, 2007. ISBN: 9780821839416 9780821866122 9781470404758 ISSN: 0065-9266, 1947-6221 Issue: 871 Publisher: American Mathematical Society Series: Memoirs of the American Mathematical Society Volume: 186.
- [CFK81] Luis A. Caffarelli, Eugene B. Fabes, and Carlos E. Kenig. Completely Singular Elliptic-Harmonic Measures. *Indiana University Mathematics Journal*, 30(6):917–924, 1981. Publisher: Indiana University Mathematics Department.
- [Dah86] Björn E. J. Dahlberg. On the Absolute Continuity of Elliptic Measures. *American Journal of Mathematics*, 108(5):1119–1138, 1986. Publisher: Johns Hopkins University Press.
- [DDH20] Martin Dindoš, Luke Dyer, and Sukjung Hwang. Parabolic L^p Dirichlet boundary value problem and VMO-type time-varying domains. *Analysis & PDE*, 13(4):1221–1268, June 2020. Publisher: Mathematical Sciences Publishers.
- [DHM21] Martin Dindoš, Sukjung Hwang, and Marius Mitrea. The L^p Dirichlet boundary problem for second order elliptic Systems with rough coefficients. *Transactions of the American Mathematical Society*, 374(5):3659–3701, May 2021.
- [DHP23] Martin Dindoš, Steve Hofmann, and Jill Pipher. Regularity and Neumann problems for operators with real coefficients satisfying Carleson conditions. *Journal of Functional Analysis*, 285(6):110024, 2023.
- [DKP11] Martin Dindos, Carlos Kenig, and Jill Pipher. BMO Solvability and the A^∞ Condition for Elliptic Operators. *Journal of Geometric Analysis*, 21(1):78–95, January 2011.
- [DLM23] Guy David, Linhan Li, and Svitlana Mayboroda. Small A_∞ results for Dahlberg-Kenig-Pipher operators in sets with uniformly rectifiable boundaries. *Transactions of the American Mathematical Society*, 376(11):7857–7909, November 2023.
- [DLP24] Martin Dindoš, Linhan Li, and Jill Pipher. The L^p regularity problem for parabolic operators, November 2024. arXiv:2410.23801.
- [DP19] Martin Dindoš and Jill Pipher. Regularity theory for solutions to second order elliptic operators with complex coefficients and the L^p Dirichlet problem. *Advances in Mathematics*, 341:255–298, 2019.

- [DP23] Martin Dindoš and Jill Pipher. Boundary Value Problems for Elliptic Operators Satisfying Carleson Condition. *Vietnam Journal of Mathematics*, October 2023.
- [DPP07] Martin Dindos, Stefanie Petermichl, and Jill Pipher. The L^p Dirichlet problem for second order elliptic operators and a p -adapted square function. *Journal of Functional Analysis*, 249(2):372–392, August 2007.
- [DPR17] Martin Dindoš, Jill Pipher, and David Rule. Boundary Value Problems for Second-Order Elliptic Operators Satisfying a Carleson Condition. *Communications on Pure and Applied Mathematics*, 70(7):1316–1365, 2017. eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1002/cpa.21649>.
- [Fen23] Joseph Feneuil. An alternative proof of the L^p -Regularity problem for Dahlberg-Kenig-Pipher operators on \mathbb{R}_+^n , October 2023. arXiv:2310.00645 [math].
- [FJK84] Eugene B. Fabes, David S. Jerison, and Carlos E. Kenig. Necessary and Sufficient Conditions for Absolute Continuity of Elliptic-Harmonic Measure. *Annals of Mathematics*, 119(1):121–141, 1984. Publisher: Annals of Mathematics.
- [HKMP14] Steve Hofmann, Carlos Kenig, Svitlana Mayboroda, and Jill Pipher. Square function/non-tangential maximal function estimates and the Dirichlet problem for non-symmetric elliptic operators. *Journal of the American Mathematical Society*, 28(2):483–529, May 2014.
- [HKMP15a] Steve Hofmann, Carlos Kenig, Svitlana Mayboroda, and Jill Pipher. The regularity problem for second order elliptic operators with complex-valued bounded measurable coefficients. *Mathematische Annalen*, 361(3):863–907, April 2015.
- [HKMP15b] Steve Hofmann, Carlos Kenig, Svitlana Mayboroda, and Jill Pipher. Square function/non-tangential maximal function estimates and the dirichlet problem for non-symmetric elliptic operators. *Journal of the American Mathematical Society*, 28(2):483–529, 2015.
- [HLMP22a] Steve Hofmann, Linhan Li, Svitlana Mayboroda, and Jill Pipher. The Dirichlet problem for elliptic operators having a BMO anti-symmetric part. *Mathematische Annalen*, 382(1):103–168, February 2022.
- [HLMP22b] Steve Hofmann, Linhan Li, Svitlana Mayboroda, and Jill Pipher. L^p theory for the square roots and square functions of elliptic operators having a BMO anti-symmetric part. *Mathematische Zeitschrift*, 301(1):935–976, May 2022. arXiv:1908.01030 [math].

- [HMM⁺21] Steve Hofmann, José María Martell, Svitlana Mayboroda, Tatiana Toro, and Zihui Zhao. Uniform Rectifiability and Elliptic Operators Satisfying a Carleson Measure Condition. *Geometric and Functional Analysis*, 31(2):325–401, April 2021.
- [JK81] David S. Jerison and Carlos E. Kenig. The Dirichlet Problem in Non-Smooth Domains. *Annals of Mathematics*, 113(2):367–382, 1981. Publisher: Annals of Mathematics.
- [KKPT00] C. Kenig, H. Koch, J. Pipher, and T. Toro. A New Approach to Absolute Continuity of Elliptic Measure, with Applications to Non-symmetric Equations. *Advances in Mathematics*, 153(2):231–298, August 2000.
- [KP01] Carlos E. Kenig and Jill Pipher. THE DIRICHLET PROBLEM FOR ELLIPTIC EQUATIONS WITH DRIFT TERMS. *Publicacions Matemàtiques*, 45(1):199–217, 2001. Publisher: Universitat Autònoma de Barcelona.
- [MM80] Luciano Modica and Stefano Mortola. Construction of a singular elliptic-harmonic measure. *Manuscripta Mathematica*, 33(1):81–98, March 1980.
- [MPT22] Mihalis Mourgoglou, Bruno Poggi, and Xavier Tolsa. L^p -solvability of the Poisson-Dirichlet problem and its applications to the Regularity problem, July 2022. arXiv:2207.10554 [math].
- [Ulm24] Martin Ulmer. On a mixed $L^1 - L^\infty$ type Carleson condition on the nontangential derivative of A for an elliptic operator, January 2024. arXiv:2311.00614 [math].