

EMBEDDED CONSTANT MEAN CURVATURE HYPERTORI IN THE $2n$ -SPHERE

JUNQI LAI AND GUOXIN WEI

ABSTRACT. Brendle [3] proved Lawson [12] conjecture about minimal embedded torus in the round three-dimensional sphere. Carlotto and Schulz [6] constructed a minimal embedded three-dimensional hypertorus in the round four-dimensional sphere and conjectured that their hypertorus is a unique minimal embedded three-dimensional hypertorus in the round four-dimensional sphere. In this paper, we construct two different constant mean curvature embedded $(2n - 1)$ -dimensional hypertori (that is, topological type $\mathbb{S}^{n-1} \times \mathbb{S}^{n-1} \times \mathbb{S}^1$) which have the same negative mean curvature H in the round $2n$ -dimensional sphere $\mathbb{S}^{2n}(1)$.

1. INTRODUCTION

A hypersurface in the unit sphere $\mathbb{S}^{n+1}(1)$ is said to have constant mean curvature (CMC) if its mean curvature H , defined as the trace of the second fundamental form, is constant. CMC hypersurfaces generalize the concept of minimal hypersurfaces (where mean curvature $H = 0$). The simplest examples of CMC hypersurfaces in the unit sphere $\mathbb{S}^{n+1}(1)$ are spheres $\mathbb{S}^n(r)$ and Clifford tori $\mathbb{S}^k(r) \times \mathbb{S}^{n-k}(\sqrt{1-r^2})$, where $0 < r < 1$.

For $n = 2$, Lawson [11] proved in 1970 that any closed surface of orientable topological type can be minimally embedded into the three-dimensional unit sphere, while for non-orientable types, there is always a minimal immersion except $\mathbb{R}\mathbb{P}^2$ (real projection plane). In the same year, Lawson [12] conjectured that the Clifford torus $\mathbb{S}^1\left(\frac{\sqrt{2}}{2}\right) \times \mathbb{S}^1\left(\frac{\sqrt{2}}{2}\right)$ is the only compact embedded minimal surface in $\mathbb{S}^3(1)$ of topological type $\mathbb{S}^1 \times \mathbb{S}^1$. By applying the maximum principle technique to a two-point function defined on the space of pairs of points, Brendle [3] proved this conjecture affirmatively in 2013. This technique turned out to be quite useful, see [4] for an overview of two-point functions and [1, 8, 10, 13] for further interesting applications.

For $n = 3$, Carlotto and Schulz [6] were able to construct a minimal embedding of three-dimensional hypertorus (that is, topological type $\mathbb{S}^1 \times \mathbb{S}^1 \times \mathbb{S}^1$) in $\mathbb{S}^4(1)$. In fact, they proved:

Theorem 1.1 (Carlotto and Schulz). *For any $2 \leq n \in \mathbb{N}$, there exists a minimal embedding of $\mathbb{S}^{n-1} \times \mathbb{S}^{n-1} \times \mathbb{S}^1$ in $\mathbb{S}^{2n}(1)$.*

In the same paper, Carlotto and Schulz [6] conjectured the following:

Conjecture (Carlotto and Schulz). *There exists, up to ambient isometry, a unique minimal embedding of the three-dimensional hypertorus in the round four-dimensional sphere $\mathbb{S}^4(1)$.*

This conjecture can be regarded as a higher-dimensional counterpart of Lawson's conjecture. For $n = 4$, they also constructed a minimal embedding of four-dimensional torus $\mathbb{S}^1 \times \mathbb{S}^1 \times \mathbb{S}^1 \times \mathbb{S}^1$ in $\mathbb{S}^5(1)$. However, for higher dimensions, their method fails to construct such embeddings.

Bobenko [2] gave explicit formulas for CMC tori and obtained a "Weierstrass representation" for CMC surfaces of finite-gap type in $\mathbb{S}^3(1)$. Pinkall and Sterling [18] conjectured that embedded CMC tori are surfaces of revolution. Andrews and Li [1] confirmed the conjecture and proved that any constant mean curvature embedded torus in $\mathbb{S}^3(1)$ is axially symmetric. We note that the embeddedness assumption is crucial in the results of Brendle [3] and Andrews-Li [1]: Otsuki [14] constructed a lot of rotationally symmetric immersed minimal tori in $\mathbb{S}^3(1)$; Bobenko [2] constructed an infinite family of non-rotationally symmetric immersed CMC tori in $\mathbb{S}^3(1)$ (see also Brito-Leite [5], Cheng-Lai-Wei [7], Wei-Cheng-Li [19], Perdomo [15], [16]).

For CMC hypertorus (that is, topological type $\mathbb{S}^{n-1} \times \mathbb{S}^{n-1} \times \mathbb{S}^1$), Huang and Wei [9] extended the work of Carlotto and Schulz [6] about minimal cases to CMC hypersurfaces for all mean curvature $H \geq 0$. That is, Huang and Wei [9] constructed compact immersed or embedded CMC $(2n - 1)$ -dimensional hypertori. In this paper, we consider CMC hypersurfaces for negative mean curvature and obtain the following result.

Theorem 1.2. *For any $2 \leq n \in \mathbb{N}$, there exists $\varepsilon = \varepsilon(n) > 0$ such that for any $-\varepsilon < H < 0$, there exist two CMC embeddings of $\mathbb{S}^{n-1} \times \mathbb{S}^{n-1} \times \mathbb{S}^1$ into $\mathbb{S}^{2n}(1)$ which have same mean curvature H and are not isometric to each other.*

Remark 1.1. *For $H \geq 0$, one can construct a embedding of the three-dimensional CMC H hypertorus in $\mathbb{S}^4(1)$. But for $-\varepsilon < H < 0$, from our result, we can construct two different embedding of the three-dimensional CMC H hypertori in $\mathbb{S}^4(1)$.*

Remark 1.2. *Perdomo [17] told us that he can give some numerical construction of the embedded CMC hypersurfaces.*

2. PROOF OF THEOREM 1.2

Let $\gamma(s) = (x(s), y(s), z(s))$ be a arc-length curve in $\{(x, y, z) \in \mathbb{S}^2(1) \mid x > 0, y > 0\}$, then one can construct an immersion

$$\begin{aligned} F : \mathbb{S}^{n-1}(1) \times \mathbb{S}^{n-1}(1) \times I &\rightarrow \mathbb{S}^{2n}(1), \\ (p_1, p_2, s) &\mapsto (x(s)p_1, y(s)p_2, z(s)), \end{aligned}$$

where I is some interval. Let $(x, y, z) = (\sin r \cos \theta, \sin r \sin \theta, \cos r)$ and $\alpha(s)$ denote the angle between the tangent vector of $\gamma(s)$ and $\frac{\partial}{\partial r}$, where $(r, \theta) \in (0, \pi) \times (0, \frac{\pi}{2})$. Huang and Wei [9] proved that the immersion F has constant mean curvature H if and only if $r(s), \theta(s), \alpha(s)$ satisfy the following autonomous system of differential equations:

$$\begin{aligned} (2.1) \quad & \begin{cases} \dot{r} = \cos \alpha, \\ \dot{\theta} = \frac{\sin \alpha}{\sin r}, \\ \dot{\alpha} = \frac{(2n-2) \cot 2\theta}{\sin r} \cos \alpha - (2n-1) \cot r \sin \alpha + H. \end{cases} \\ (2.2) \quad & \\ (2.3) \quad & \end{aligned}$$

Let us begin with a lemma on the symmetry of the solutions to this system. One can check it by a direct computation, so we omit the proof.

Lemma 2.1. *Let $(r(s), \theta(s), \alpha(s))$ be a solution of (2.1), (2.2), (2.3), then*

$$(\tilde{r}(s), \tilde{\theta}(s), \tilde{\alpha}(s)) := (\pi - r(-s), \theta(-s), -\alpha(-s)),$$

$$(\hat{r}(s), \hat{\theta}(s), \hat{\alpha}(s)) := (r(-s), \frac{\pi}{2} - \theta(-s), \pi - \alpha(-s))$$

and

$$(\check{r}(s), \check{\theta}(s), \check{\alpha}(s)) := (r(s), \theta(s), \alpha(s) + 2k\pi), \quad \forall k \in \mathbb{Z},$$

are also solutions of that system.

Lemma 2.2. *Let $H < 0$, then for any $r_0 \in (0, \operatorname{arccot} \frac{-H}{2n-1})$, there exists $0 < s_* < \infty$ so that the system (2.1), (2.2), (2.3) has a unique right maximal solution:*

$$(r, \theta, \alpha) : (0, s_*) \rightarrow B := (0, \frac{\pi}{2}) \times (0, \frac{\pi}{4}) \times (-\frac{\pi}{2}, 0)$$

with initial data $(r, \theta, \alpha)(0) = (r_0, \frac{\pi}{4}, -\frac{\pi}{2})$ and $\dot{\alpha}(s) > 0$ in $(0, s_*)$. Moreover, $r(s_*) = \frac{\pi}{2}$ or $\alpha(s_*) = 0$ or $\dot{\alpha}(s_*) = 0$; in either case $\theta(s_*) > 0$.

Proof. We first prove that $\theta(s_*) > 0$. Note that $\dot{r}(s) > 0$, $\dot{\theta}(s) < 0$ and $\dot{\alpha}(s) > 0$ for all $s \in (0, s_*)$. Suppose $\theta(s_*) = \lim_{s \rightarrow s_*} \theta(s) = 0$, then

$$\lim_{s \rightarrow s_*} \frac{(n-1) \cot 2\theta(s) \cos \alpha(s)}{\sin r(s)} = \infty.$$

Hence, there exists $s_0 \in (0, s_*)$ such that

$$\frac{(n-1) \cot 2\theta(s) \cos \alpha(s)}{\sin r(s)} \geq -H$$

for $s \in [s_0, s_*)$. It follows that

$$\dot{\alpha} \geq \frac{(n-1) \cot 2\theta \cos \alpha}{\sin r} = (n-1) \dot{\theta} \cot 2\theta \cot \alpha,$$

namely,

$$\dot{\alpha} \tan \alpha \leq (n-1) \dot{\theta} \cot 2\theta$$

for $s \in [s_0, s_*)$. Integrating this inequality from s_0 to s gives

$$\log \frac{\cos \alpha(s_0)}{\cos \alpha(s)} \leq \frac{1}{2} (n-1) \log \frac{\sin 2\theta(s)}{\sin 2\theta(s_0)}$$

which is impossible as s tends to s_* , since the right hand side tends to $-\infty$ as s tends to s_* . Now

$$r(s_*) = \lim_{s \rightarrow s_*} r(s) \in (r_0, \frac{\pi}{2}],$$

$$\theta(s_*) = \lim_{s \rightarrow s_*} \theta(s) \in (0, \frac{\pi}{4}),$$

$$\alpha(s_*) = \lim_{s \rightarrow s_*} \alpha(s) \in (-\frac{\pi}{2}, 0],$$

which follows that $s_* < \infty$ since system (2.1), (2.2), (2.3) has no equilibrium. Furthermore, $s_* < S$, where S is the right end point of the right maximal solution of (2.1), (2.2), (2.3) with initial data $r(0) = r_0$, $\theta(0) = \frac{\pi}{4}$, $\alpha(0) = -\frac{\pi}{2}$. By the definition of s_* we have either $r(s_*) = \frac{\pi}{2}$, $\alpha(s_*) = 0$ or $\dot{\alpha}(s_*) = 0$, otherwise $(0, s_*)$ is not right maximally extended with given property. \square

Lemma 2.3. *Let $H < 0$ and $(r, \theta, \alpha) : (0, s_*) \rightarrow B$ be a solution as in lemma 2.2. If $(2n-2) \cot 2\theta \cos \alpha|_{s=s_0} + H \geq 0$ for some $s_0 \in (0, s_*)$, then $\dot{\alpha}(s_*) > 0$.*

Proof. Since $\dot{\alpha}(s) > 0$ for $s \in (0, s_*)$ and $(2n-2) \cot 2\theta \cos \alpha|_{s=s_0} + H \geq 0$, one gets that

$$(2n-2) \cot 2\theta \cos \alpha + H > 0$$

for $s \in (s_0, s_*]$. In particular

$$(2n-2) \cot 2\theta \cos \alpha|_{s=s_*} + H > 0,$$

which follows that

$$\begin{aligned} \dot{\alpha}(s_*) &= \frac{(2n-2) \cot 2\theta \cos \alpha}{\sin r} \Big|_{s=s_*} - (2n-1) \cot r \sin \alpha|_{s=s_*} + H \\ &\geq (2n-2) \cot 2\theta \cos \alpha|_{s=s_*} + H - (2n-1) \cot r \sin \alpha|_{s=s_*} \\ &> 0. \end{aligned}$$

□

Lemma 2.4. *Let $H < 0$, $(r, \theta, \alpha) : (0, s_*) \rightarrow B$ be a solution as in lemma 2.2 and $\dot{\alpha}(s_*) = 0$. Then $(2n-2) \cot 2\theta \cos \alpha + H < 0$ for $s \in s_*$. Moreover, $\alpha(s) + \frac{\pi}{2} < (2n-1) \left(\frac{\pi}{4} - \theta(s) \right) + H \left(1 - \frac{1}{\sin r_0} \right) s$ for $s \in (0, s_*]$.*

Proof. It follows from lemma 2.3 that $(2n-2) \cot 2\theta \cos \alpha + H < 0$ for $s \in (0, s_*)$. Hence

$$\begin{aligned} \dot{\alpha} &= \frac{(2n-2) \cot 2\theta \cos \alpha}{\sin r} - (2n-1) \cot r \sin \alpha + H \\ &< \frac{(2n-2) \cot 2\theta \cos \alpha}{\sin r_0} - (2n-1) \cot r \sin \alpha + \frac{H}{\sin r_0} + \left(1 - \frac{1}{\sin r_0} \right) H \\ &< -(2n-1) \dot{\theta} \cos r + H \left(1 - \frac{1}{\sin r_0} \right) \\ &< -(2n-1) \dot{\theta} + H \left(1 - \frac{1}{\sin r_0} \right) \end{aligned}$$

for $s \in (0, s_*)$. Integrating this inequality from 0 to s gives

$$\alpha + \frac{\pi}{2} < (2n-1) \left(\frac{\pi}{4} - \theta \right) + H \left(1 - \frac{1}{\sin r_0} \right) s$$

for $s \in (0, s_*]$. □

Lemma 2.5. *Let $H < 0$ and $(r, \theta, \alpha) : (0, s_*) \rightarrow B$ be a solution as in lemma 2.2. Then $s_* < \frac{3\sqrt{2}\pi}{4}$.*

Proof. Let s_0 in $(0, s_*)$ be the point, if any, at which $\alpha(s_0) = -\frac{\pi}{4}$, then

$$\dot{\theta} = \frac{\sin \alpha}{\sin r} < \sin \alpha < -\frac{\sqrt{2}}{2}$$

for $s \in (0, s_0)$. Integrating from 0 to s_0 gives

$$\theta(s_0) - \frac{\pi}{4} < -\frac{\sqrt{2}}{2} s_0,$$

which implies that

$$(2.4) \quad s_0 < \frac{\sqrt{2}\pi}{4}.$$

On the other hand, $\dot{r} = \cos \alpha > \frac{\sqrt{2}}{2}$ for $s \in (s_0, s_*)$. Integrating from s_0 to s_* gives that

$$r(s_*) - r(s_0) > \frac{\sqrt{2}}{2}(s_* - s_0),$$

which implies that

$$(2.5) \quad s_* - s_0 < \sqrt{2}(r(s_*) - r(s_0)) < \sqrt{2} \times \frac{\pi}{2} = \frac{\sqrt{2}\pi}{2}.$$

Combining (2.4) and (2.5) gives that

$$s_* < \frac{3\sqrt{2}\pi}{4}.$$

□

Lemma 2.6. *Let $H < 0$, $(r, \theta, \alpha) : (0, s_*) \rightarrow B$ be a solution as in lemma 2.2 and $\dot{\alpha}(s_*) = 0$. If $3\sqrt{2}H \left(1 - \frac{1}{\sin r_0}\right) < 1$ and*

$$\sqrt{2}(n-1) \tan \left(\frac{2}{2n-1} \left(\frac{\pi}{4} - H \left(1 - \frac{1}{\sin r_0}\right) \frac{3\sqrt{2}\pi}{4} \right) \right) + H \geq 0,$$

then $\alpha(s) < -\frac{\pi}{4}$ for $s \in (0, s_*]$.

Proof. Suppose $\alpha(s_0) = -\frac{\pi}{4}$ for some $s_0 \in (0, s_*]$, then we have

$$(2.6) \quad \frac{\pi}{4} < (2n-1) \left(\frac{\pi}{4} - \theta(s_0) \right) + H \left(1 - \frac{1}{\sin r_0}\right) s_0$$

by lemma 2.4. By lemma 2.5 we have

$$(2.7) \quad s_0 \leq s_* < \frac{3\sqrt{2}\pi}{4}.$$

Combining (2.6) and (2.7) gives that

$$2\theta(s_0) < \frac{\pi}{2} - \frac{2}{2n-1} \left(\frac{\pi}{4} - H \left(1 - \frac{1}{\sin r_0}\right) \frac{3\sqrt{2}\pi}{4} \right).$$

Hence

$$\begin{aligned} & (2n-2) \cot 2\theta \cos \alpha|_{s=s_0} + H \\ & > \sqrt{2}(n-1) \tan \left(\frac{2}{2n-1} \left(\frac{\pi}{4} - H \left(1 - \frac{1}{\sin r_0}\right) \frac{3\sqrt{2}\pi}{4} \right) \right) + H \\ & \geq 0. \end{aligned}$$

But this is impossible by lemma 2.3. □

Lemma 2.7. *Let $H < 0$, $(r, \theta, \alpha) : (0, s_*) \rightarrow B$ be a solution as in lemma 2.2. If $H \left(1 - \frac{1}{\sin r_0}\right) 3\sqrt{2} < 1$, $H \geq -\frac{\sqrt{2}}{2}(2n-3)$ and*

$$\sqrt{2}(n-1) \tan \left(\frac{2}{2n-1} \left(\frac{\pi}{4} - H \left(1 - \frac{1}{\sin r_0}\right) \frac{3\sqrt{2}\pi}{4} \right) \right) + H \geq 0,$$

then $\dot{\alpha}(s_*) \neq 0$.

Proof. Suppose $\dot{\alpha}(s_*) = 0$, then $\alpha(s_*) \leq -\frac{\pi}{4}$ and

$$(2n-2) \cot 2\theta(s_*) \cos \alpha(s_*) \leq -H$$

by lemma 2.6 and lemma 2.3. Since $\dot{\alpha}(s) > 0$ for $s \in (0, s_*)$, we have $\ddot{\alpha}(s_*) \leq 0$. On the other hand, we have

$$\begin{aligned} \ddot{\alpha} &= \frac{-(4n-4) \sin \alpha \cos \alpha}{\sin^2 r \sin^2 2\theta} - \frac{(2n-2) \cos r \cos^2 \alpha}{\sin^2 r \tan 2\theta} + (2n-1) \csc^2 r \cos \alpha \sin \alpha \\ &= \frac{\cos \alpha}{\sin^2 r} \left[\sin \alpha \left(-(4n-4) \csc^2 2\theta + 2n-1 \right) - (2n-1) \cot 2\theta \cos \alpha \cos r \right] \\ &\geq \frac{\cos \alpha}{\sin^2 r} \left[\sin \alpha \left(-(4n-4) \csc^2 2\theta + 2n-1 \right) + H \cos r \right] \\ &> \frac{\cos \alpha}{\sin^2 r} \left[\sin \alpha \left(-(4n-4) \csc^2 2\theta + 2n-1 \right) + H \right] \\ &\geq \frac{\cos \alpha}{\sin^2 r} \left[-\frac{\sqrt{2}}{2} \left(-(4n-4) \csc^2 2\theta + 2n-1 + 2n-3 \right) \right] \\ &> 0 \end{aligned}$$

at s_* . We arrive at a contradiction, hence $\dot{\alpha}(s_*) \neq 0$. \square

Lemma 2.8. *Let $H < 0, (r, \theta, \alpha) : (0, s_*) \rightarrow B$ be a solution as in lemma 2.2. Then*

$$\lim_{r_0 \rightarrow \operatorname{arccot} \frac{-H}{2n-1}} r(s_*) = \operatorname{arccot} \frac{-H}{2n-1}.$$

Proof. For simplicity, we denote $\operatorname{arccot} \frac{-H}{2n-1}$ by $a(H)$. Let us first note that $\liminf_{r_0 \rightarrow a(H)} \geq a(H)$, so it suffice to prove that $\limsup_{r_0 \rightarrow a(H)} \leq a(H)$. We adopt a contradiction argument. Suppose $\limsup_{r_0 \rightarrow a(H)} r(s_*) > a(H)$ and let R be in the interval

$$(a(H), \limsup_{r_0 \rightarrow a(H)} r(s_*)),$$

then there exists a sequence $\{\rho_n\}_{n=1}^\infty$ increasingly converging to $a(H)$ so that $r_n(s_*) > R$. We use the notation $(r_n(s), \theta_n(s), \alpha_n(s))$ to denote the solution of system (2.1),(2.2),(2.3) with initial data $(r_n(0), \theta_n(0), \alpha_n(0)) = (\rho_n, \frac{\pi}{4}, -\frac{\pi}{2})$. Now for each n , there are $s_{1,n}, s_{2,n}$ in $(0, s_*)$ such that

$$r_n(s_{1,n}) = \frac{2a(H) + R}{3}, \quad r_n(s_{2,n}) = \frac{a(H) + 2R}{3}.$$

Since $(r_n(s), \theta_n(s), \alpha_n(s))$ converges to $(a(H), -\frac{1}{\sin a(H)}s + \frac{\pi}{4}, -\frac{\pi}{2})$ on compacta as $n \rightarrow \infty$, we see that

$$(2.8) \quad \lim_{n \rightarrow \infty} \theta_n(s_{1,n}) = 0, \quad \lim_{n \rightarrow \infty} \theta_n(s_{2,n}) = 0.$$

For each n , there is a $\xi_n \in (s_{1,n}, s_{2,n})$ such that

$$\frac{\theta_n(s_{2,n}) - \theta_n(s_{1,n})}{r_n(s_{2,n}) - r_n(s_{1,n})} = \frac{\sin \alpha_n(\xi_n)}{\cos \alpha_n(\xi_n) \sin r_n(\xi_n)}.$$

by Cauchy intermediate theorem. Since $\lim_{n \rightarrow \infty} \theta_n(s_{i,n}) = 0$, $i = 1, 2$, we obtain that

$$(2.9) \quad \lim_{n \rightarrow \infty} \alpha_n(\xi_n) = 0.$$

Combining (2.8), (2.9) with that $\dot{\theta}_n < 0$, $\dot{\alpha}_n > 0$ for every n and $s \in (0, s_*)$, we obtain that

$$\frac{(n-1) \cot 2\theta_n \cos \alpha_n}{\sin r_n} + H > 0$$

and

$$\frac{(n-1) \cot 2\theta_n}{\sin r_n} > \frac{3\pi}{2(R-a(H))}$$

for large n and $s \in [s_{2,n}, s_*)$. It follows that

$$\dot{\alpha}_n > \frac{3\pi}{2(R-a(H))} \dot{r}$$

for large n and $s \in [s_{2,n}, s_*)$. Integrating from $s_{2,n}$ to s_* gives that

$$\begin{aligned} \alpha_n(s_*) - \alpha_n(s_{2,n}) &> \frac{3\pi}{2(R-a(H))} (r_n(s_*) - r_n(s_{2,n})) \\ &> \frac{3\pi}{2(R-a(H))} \left(R - \frac{a(H) + 2R}{3} \right) \\ &= \frac{\pi}{2}, \end{aligned}$$

which is impossible. Hence $\liminf_{r_0 \rightarrow a(H)} r(s_*) = \limsup_{r_0 \rightarrow a(H)} r(s_*) = a(H)$, namely, $\lim_{r_0 \rightarrow a(H)} r(s_*) = a(H)$. \square

Lemma 2.9. *Let $(r, \theta, \alpha) : (0, s_*) \rightarrow B$ be a solution as in lemma 2.2. There exists $0 < r'_0 < r''_0 < \frac{\pi}{2}$ and $0 < \delta < (2n-1) \cot r''_0$ such that for $-\delta \leq H \leq 0$, one has $\alpha(s_*) = 0$ with $r_0 = r'_0$ and $r(s_*) = \frac{\pi}{2}$ with $r_0 = r''_0$.*

Proof. For $H = 0$, Carlotto and Schulz [6] proved that there exists $0 < r'_0 < r''_0 < \frac{\pi}{2}$ such that $\alpha(s_*) = 0$ with $r_0 = r'_0$ and $r(s_*) = \frac{\pi}{2}$ with $r_0 = r''_0$. The lemma follows by the parameter dependence of the system (2.1), (2.2), (2.3) on the parameter H . \square

Proposition 2.1. *Let r'_0, r''_0, δ be as in lemma 2.9,*

$$-\min \left\{ \frac{\sqrt{2} \sin \frac{r'_0}{2}}{12 \left(1 - \sin \frac{r'_0}{2} \right)}, \delta, \sqrt{2}(n-1) \tan \frac{\pi}{4(2n-1)} \right\} \leq H < 0$$

and $(r, \theta, \alpha) : (0, s_*) \rightarrow B$ be a solution as in lemma 2.2. Then there exists $r'_0 < \varrho_1 < r''_0$ and $r''_0 < \varrho_2 < \arccot \frac{-H}{2n-1}$ such that $\alpha(s_*) = 0$, $r(s_*) = \frac{\pi}{2}$ with initial value $r_0 = \varrho_i$, $i = 1, 2$.

Proof. Lemma 2.7 shows that $\dot{\alpha}(s_*) \neq 0$ for any $r_0 \in (\frac{r'_0}{2}, \arccot \frac{-H}{2n-1})$. Hence $\dot{\alpha}(s) > 0$, $s \in [0, s_*)$ for any $r_0 \in (\frac{r'_0}{2}, \arccot \frac{-H}{2n-1})$. Note that $r'_0, r''_0 \in (\frac{r'_0}{2}, \arccot \frac{-H}{2n-1})$. Lemma 2.9 shows that $r(s_*) < \frac{\pi}{2}$, $\alpha(s_*) = 0$ for $r_0 = r'_0$ and $r(s_*) = \frac{\pi}{2}$, $\alpha(s_*) < 0$ for $r_0 = r''_0$. Moreover, lemma 2.8 shows that $r(s_*) < \frac{\pi}{2}$, $\alpha(s_*) = 0$ for r_0 close to $\arccot \frac{-H}{2n-1}$. Now the lemma follows by the initial value dependence of the system (2.1), (2.2), (2.3), also see the proof of theorem 1.1 in [6]. \square

Proof of theorem 1.2. Let r'_0, δ be as in lemma 2.9 and

$$\varepsilon := \min \left\{ \frac{\sqrt{2} \sin \frac{r'_0}{2}}{12 \left(1 - \sin \frac{r'_0}{2} \right)}, \delta, \sqrt{2}(n-1) \tan \frac{\pi}{4(2n-1)} \right\}.$$

Given $-\varepsilon < H < 0$, let ϱ_1, ϱ_2 be as in proposition 2.1 and $(r_i(s), \theta_i(s), \alpha_i(s))$ be the solutions of (2.1), (2.2), (2.3) with initial value $(r_i(0), \theta_i(0), \alpha_i(0)) = (\varrho_i, \frac{\pi}{4}, -\frac{\pi}{2})$, $i = 1, 2$. By lemma 2.1, one sees that the curves $(r_i(s), \theta_i(s))$ are smooth simple closed curves.

Hence they generate two CMC embeddings of $\mathbb{S}^{n-1} \times \mathbb{S}^{n-1} \times \mathbb{S}^1$ in $\mathbb{S}^{2n}(1)$ which have same negative mean curvature H . \square

Figure 2.1 shows two distinct simple closed profile curves (red and green) of two 3-dimensional hypertori sharing the same mean curvature $H = -0.2$.




Figure 2.1 Profile curves with $n = 2$ and $H = -0.2$, where the initial value r_0 of the red curve is 1.29691 and that of green curve is 1.44086.

REFERENCES

- [1] B. Andrews and H. Li, *Embedded constant mean curvature tori in the three-sphere*, J. Differential Geom. **99** (2015), no. 2, 169–189; MR3302037
- [2] A. I. Bobenko, *All constant mean curvature tori in \mathbb{R}^3 , S^3 , H^3 in terms of theta-functions*, Math. Ann. **290** (1991), no. 2, 209–245; MR1109632
- [3] S. Brendle, *Embedded minimal tori in S^3 and the Lawson conjecture*, Acta Math. **211** (2013), no. 2, 177–190; MR3143888
- [4] S. Brendle, *Two-point functions and their applications in geometry*, Bull. Amer. Math. Soc. (N.S.) **51** (2014), no. 4, 581–596; MR3237760
- [5] F. Brito and M. L. Leite, *A remark on rotational hypersurfaces of S^n* , Bull. Soc. Math. Belg. **42** (1990), 303–318; MR1081607
- [6] A. Carlotto and M. B. Schulz, *Minimal hypertori in the four-dimensional sphere*, Ars Inven. Anal. **2023**, Paper No. 8, 33 pp.; MR4645955
- [7] Q.-M. Cheng, J. Lai and G. Wei, *Examples of compact embedded convex λ -hypersurfaces*, J. Funct. Anal. **286** (2024), Paper No. 110211, 15 pp.; MR4661640
- [8] G. De Philippis and M. Goldman, *A two-point function approach to connectedness of drops in convex potentials*, Comm. Anal. Geom. **30** (2022), no. 4, 815–841; MR4545852
- [9] C. Huang and G. Wei, *New examples of constant mean curvature hypersurfaces in the sphere*, arXiv:2209.13236, 2022.
- [10] H. Kröner, *Non-collapsing in homogeneity greater than one via a two-point method for a special case*, Proc. Roy. Soc. Edinburgh Sect. A **149** (2019), no. 6, 1627–1635; MR4034585
- [11] H. B. Lawson Jr., *Complete minimal surfaces in S^3* , Ann. of Math. (2) **92** (1970), 335–374; MR0270280
- [12] H. B. Lawson Jr., *The unknottedness of minimal embeddings*, Invent. Math. **11** (1970), 183–187; MR0287447
- [13] P. J. McGrath and E. Meekins, *Two-point functions and constant mean curvature surfaces in \mathbb{R}^3* , Involve **16** (2023), no. 3, 467–482; MR4625886
- [14] T. Otsuki, *Minimal hypersurfaces in a Riemannian manifold of constant curvature*, Amer. J. Math. **92** (1970), 145–173; MR0264565

- [15] O. M. Perdomo, *Embedded constant mean curvature hypersurfaces on spheres*, Asian J. Math. **14** (2010), 73–108; MR2726595
- [16] O. M. Perdomo, *Rotational surfaces in S^3 with constant mean curvature*, J. Geom. Anal. **26** (2016), 2155–2168; MR3511472
- [17] O. M. Perdomo, *Navigating the space of compact CMC hypersurfaces in spheres, Part I*, arXiv:2503.13823, 2025
- [18] U. Pinkall and I. Sterling, *On the classification of constant mean curvature tori*, Ann. of Math. (2) **130** (1989), 407–451; MR1014929
- [19] G. Wei, Q. -M. Cheng and H. Li, *Embedded hypersurfaces with constant m th mean curvature in a unit sphere*, Commun. Contemp. Math. **12** (2010), 997–1013; MR2748282

JUNQI LAI, SCHOOL OF MATHEMATICAL SCIENCES, SOUTH CHINA NORMAL UNIVERSITY, 510631, GUANGZHOU, CHINA, 2019021668@M.SCNU.EDU.CN

GUOXIN WEI, SCHOOL OF MATHEMATICAL SCIENCES, SOUTH CHINA NORMAL UNIVERSITY, 510631, GUANGZHOU, CHINA, WEIGUOXIN@TSINGHUA.ORG.CN