

Some New Congruences For Overpartition Function With ℓ -Regular Non-Overlined Parts

Nipen Saikia* and Adam Paksok

*Department of Mathematics, Rajiv Gandhi University,
Rono Hills, Doimukh, Arunachal Pradesh, India-791112.
E. Mail(s): nipen.saikia@rgu.ac.in; adam.paksok@rgu.ac.in*

**Corresponding author*

Abstract: Alanzi et al. (2022) investigated overpartition of a positive integer n with ℓ -regular non-overlined parts denoted by $\overline{R}_\ell^*(n)$, and proved some results for the case $\ell = 3$. As extension to the results of Alanzi et al., Sellers (2024) proved some new congruences for $\overline{R}_3^*(n)$. In this paper, we prove some new infinite families and particular congruences for $\overline{R}_\ell^*(n)$ for $\ell = 4, 5k, 6$, and 8 , where k is any positive integer. We also offer some congruences connecting $\overline{R}_\ell^*(n)$ with some other partition functions.

Keywords and Phrases: overpartition; ℓ -regular partition with non-overlined parts; congruence; q -series.

Mathematics Subject Classifications: 11P83; 05A17.

1. INTRODUCTION

A partition of a positive integer n is a finite non-increasing sequence of positive integers s_1, s_2, \dots, s_k satisfying $s_1 + s_2 + \dots + s_k = n$, where the integers s_1, s_2, \dots, s_k are called as parts or summands of the partition. For example, the partitions of $n = 3$ are given by $3, 2 + 1, 1 + 1 + 1$. The number of partition of a non-negative integer n is usually denoted by $p(n)$ and its generating function is given by

$$\sum_{n=0}^{\infty} p(n)q^n = \frac{1}{(q; q)_{\infty}}, \quad p(0) = 1, \quad (1.1)$$

where for any complex number a and $|q| < 1$,

$$(a; q)_{\infty} = \prod_{n=0}^{\infty} (1 - aq^n). \quad (1.2)$$

For brevity, we will use the notation $f_h := (q^h; q^h)_{\infty}$ for any positive integer h . Ramanujan [20, 21](also see [6]) offered following beautiful congruences for the partition function $p(n)$:

$$p(5n + 4) \equiv 0 \pmod{5}, \quad p(7n + 5) \equiv 0 \pmod{7}, \quad p(11n + 6) \equiv 0 \pmod{11}. \quad (1.3)$$

For any positive integer ℓ , an ℓ -regular partition of a positive integer n is a partition of n such that none of its part is divisible by ℓ . If $b_\ell(n)$ denotes the number of ℓ -regular

partitions of n , then the generating function of $b_\ell(n)$ is given by

$$\sum_{n=0}^{\infty} b_\ell(n)q^n = \frac{(q^\ell; q^\ell)_\infty}{(q; q)_\infty} = \frac{f_\ell}{f_1}. \quad (1.4)$$

The arithmetic properties of ℓ -regular partition have been studied by different authors, for example see [3, 4, 8, 10, 11, 15, 17, 19, 25, 27].

An overpartition of a positive integer n is a partition of n wherein first occurrence of any part may be overlined. The number of overpartitions of a positive integer n is usually denoted by $\bar{p}(n)$ which is first studied by Corteel and Lovejoy [9]. The generating function of $\bar{p}(n)$ [9] is given by

$$\sum_{n \geq 0} \bar{p}(n)q^n = \frac{(q^2; q^2)_\infty}{(q; q)_\infty^2} = \frac{f_2}{f_1^2}. \quad (1.5)$$

For instance, $\bar{p}(3) = 8$ with overpartitions of 3 given by $3, \bar{3}, 2 + 1, \bar{2} + 1, 2 + \bar{1}, \bar{2} + \bar{1}, 1 + 1 + 1, \bar{1} + 1 + 1$. One may see [14, 13, 16, 18, 22, 26] and references therein for congruence and other arithmetic properties of overpartition function $\bar{p}(n)$.

As an extension to the overpartition function $\bar{p}(n)$, overpartition with ℓ -regular overlined parts denoted by $\bar{p}_\ell(n)$ was defined. For example, $\bar{p}_2(3) = 6$ with the relevant partitions given by $3, \bar{3}, 2 + 1, 2 + \bar{1}, 1 + 1 + 1, \bar{1} + 1 + 1$. The generating function for $\bar{p}_\ell(n)$ [2] is given by

$$\sum_{n=0}^{\infty} \bar{p}_\ell(n)q^n = \frac{(q^2; q^2)_\infty (q^\ell; q^\ell)_\infty}{(q; q)_\infty^2 (q^{2\ell}; q^{2\ell})_\infty} = \frac{f_2 f_\ell}{f_1^2 f_{2\ell}}. \quad (1.6)$$

Ballantine and Merca [2] investigated the connections between $\bar{p}_\ell(n)$ and other restricted overpartition functions, and also established recurrence relations for $\bar{p}_3(n)$. Shivanna and Chandrappa [24] proved congruences modulo 5, 7, 40 and powers of 2 for $\bar{p}_\ell(n)$ with $\ell \in \{4, 8\}$. Buragohain and Saikia [7] proved some congruences for $\bar{p}_\ell(n)$ with $\ell \in \{3, 5, 9, 12, 16, 18, 24\}$ using q -series manipulations.

Recently, Alanzi et al. [1] considered overpartition of positive integer n with ℓ -regular non-overlined parts denoted by $\bar{R}_\ell^*(n)$. For instance, $\bar{R}_2^*(3) = 6$ with the relevant partitions given by $3, \bar{3}, \bar{2} + 1, \bar{2} + \bar{1}, 1 + 1 + 1, \bar{1} + 1 + 1$. Alanzi et al. [1] proved some general parity results and specific congruences for the case $\ell = 3$ of $\bar{R}_\ell^*(n)$. The generating function of $\bar{R}_\ell^*(n)$ was given by Sellers [23] as

$$\sum_{n=0}^{\infty} \bar{R}_\ell^*(n)q^n = \frac{(q^\ell; q^\ell)_\infty (q^2; q^2)_\infty}{(q; q)_\infty^2} = \frac{f_2 f_\ell}{f_1^2}. \quad (1.7)$$

As an extension to the work of Alanzi et al. [1], Sellers [23] found some infinite families and some specific congruences modulo 3 and 4 for $\bar{R}_3^*(n)$. In closing remarks of [23], further congruences for the partition function $\bar{R}_\ell^*(n)$ are desired.

In this paper, we will prove some infinite families and particular congruences for $\overline{R}_\ell^*(n)$ for $\ell = 4, 5k, 6, 8$, where k is any positive integer. We also offer some congruence relations connecting $\overline{R}_\ell^*(n)$ with some other partition functions.

2. PRELIMINARIES

Ramanujan's general theta-function $F(x, y)$ is defined by

$$F(x, y) = \sum_{n=-\infty}^{\infty} x^{n(n+1)/2} y^{n(n-1)/2}, \quad |xy| < 1. \quad (2.1)$$

Three important special cases of $F(x, y)$ [5, p. 36, Entry 22 (i), (ii)] are the theta-functions $\phi(q)$, $\psi(q)$ and $f(-q)$ given by

$$\phi(q) := F(q, q) = \sum_{\nu=0}^{\infty} q^{\nu^2} = (-q; q^2)_{\infty}^2 (q^2; q^2)_{\infty} = \frac{f_2^5}{f_1^2 f_4^2}, \quad (2.2)$$

$$\psi(q) := F(q, q^3) = \sum_{\nu \geq 0} q^{\nu(\nu+1)/2} = \frac{(q^2; q^2)_{\infty}}{(q; q^2)_{\infty}} = \frac{f_2^2}{f_1}, \quad (2.3)$$

and

$$f(-q) := F(-q, -q^2) = \sum_{\nu=-\infty}^{\infty} (-1)^{\nu} q^{\nu(3\nu+1)/2} = f_1. \quad (2.4)$$

Employing elementary q -operations, it is easily seen that

$$\phi(-q) = \frac{f_1^2}{f_2}. \quad (2.5)$$

Lemma 2.1. [10, Theorem 2.1] *For any odd prime, we have*

$$\psi(q) = \sum_{j=0}^{(p-3)/2} q^{(j^2+j)/2} F\left(q^{(p^2+(2j+1)p)/2}, q^{(p^2-(2j+1)p)/2}\right) + q^{(p^2-1)/8} \psi(q^{p^2}). \quad (2.6)$$

Furthermore, $(j^2 + j)/2 \not\equiv (p^2 - 1)/8 \pmod{p}$ for $0 \leq j \leq (p - 3)/2$.

Lemma 2.2. [10, Theorem 2.2] *For any prime $p \geq 5$, we have*

$$\begin{aligned} f_1 = & \sum_{\substack{k=-(p-1)/2 \\ k \neq (\pm p-1)/6}}^{(p-1)/2} (-1)^k q^{(3k^2+k)/2} F\left(-q^{(3p^2+(6k+1)p)/2}, -q^{(3p^2-(6k+1)p)/2}\right) \\ & + (-1)^{(\pm p-1)/6} q^{(p^2-1)/24} f_{p^2}, \end{aligned} \quad (2.7)$$

where

$$\frac{\pm p - 1}{6} = \begin{cases} \frac{(p-1)}{6}, & \text{if } p \equiv 1 \pmod{6} \\ \frac{(-p-1)}{6}, & \text{if } p \equiv -1 \pmod{6}. \end{cases}$$

Furthermore, if $\frac{-(p-1)}{2} \leq k \leq \frac{(p-1)}{2}$ and $k \neq \frac{(\pm p-1)}{6}$ then

$$\frac{3k^2 + k}{2} \not\equiv \frac{p^2 - 1}{24} \pmod{p}.$$

Lemma 2.3. [12] *We have*

$$f_1^2 = \frac{f_2 f_8^5}{f_4^2 f_{16}^2} - 2q \frac{f_2 f_{16}^2}{f_8}, \quad (2.8)$$

$$\frac{1}{f_1^2} = \frac{f_8^5}{f_2^5 f_{16}^2} + 2q \frac{f_4^2 f_{16}^2}{f_2^5 f_8}, \quad (2.9)$$

$$\frac{1}{f_1^4} = \frac{f_4^{14}}{f_2^{14} f_8^4} + 4q \frac{f_4^2 f_8^4}{f_2^{10}} \quad (2.10)$$

and

$$f_1^4 = \frac{f_4^{10}}{f_2^2 f_8^4} - 4q \frac{f_2^2 f_8^4}{f_4^2}. \quad (2.11)$$

Lemma 2.4. [14] *We have*

$$\frac{1}{\phi(-q)} = \frac{1}{\phi(-q^4)^4} \left(\phi(q^4)^3 + 2q\phi(q^4)^2\psi(q^8) + 4q^2\phi(q^4)\psi(q^8)^2 + 8q^3\psi(q^8)^3 \right). \quad (2.12)$$

Lemma 2.5. [14] *We have*

$$\begin{aligned} \frac{1}{\phi(q)} = \frac{\phi(q^{25})}{\phi(q^5)^6} & \left[\phi(q^{25})^4 - 2q\phi(q^{25})^3 X(q^5) + 4q^2\phi(q^{25})^2 X(q^5)^2 - 8q^3\phi(q^{25}) X(q^5)^3 \right. \\ & + q^4 (16X(q^5)^4 - 2\phi(q^{25})^3 Y(q^5)) - 12q^5\phi(q^{25})^2 X(q^5) Y(q^5) + 16q^6\phi(q^{25}) X(q^5)^2 Y(q^5) \\ & - 16q^7 X(q^5)^3 Y(q^5) + 4q^8\phi(q^{25})^2 Y(q^5)^2 + 16q^9\phi(q^{25}) X(q^5) Y(q^5)^2 \\ & \left. + 16q^{10} X(q^5)^2 Y(q^5)^2 - 8q^{12}\phi(q^{25}) Y(q^5)^3 - 16q^{13} X(q^5) Y(q^5)^3 + 16q^{16} Y(q^5)^4 \right], \end{aligned}$$

where

$$X(q) = \sum_{r=-\infty}^{\infty} q^{5r^2+2r} \quad \text{and} \quad Y(q) = \sum_{r=-\infty}^{\infty} q^{5r^2+4r}.$$

Lemma 2.6. [5, p. 49] *We have*

$$\psi(q) = F(q^3, q^6) + q\psi(q^9).$$

We will also employ following congruences which can be easily proved by using the binomial theorem: For any prime p ,

$$f_p \equiv f_1^p \pmod{p}, \quad (2.13)$$

and

$$f_1^{p^2} \equiv f_p^p \pmod{p^2}. \quad (2.14)$$

3. CONGRUENCES FOR $\overline{R}_\ell^*(n)$

Theorem 3.1. *We have*

(i) $\overline{R}_4^*(4n + \xi) \equiv 0 \pmod{4}; \quad \xi = 2, 3,$

(ii) *Let $p \geq 13$ be a prime with $\left(\frac{-6}{p}\right) = -1$ and $1 \leq r \leq p - 1$. Then for any integers $\alpha \geq 0$ and $n \geq 0$, we have*

$$\sum_{n=0}^{\infty} \overline{R}_4^* \left(4 \cdot p^{2\alpha} n + \frac{7 \cdot p^{2\alpha} - 1}{6} \right) \equiv 2f_1 \psi(q^2) \pmod{4} \quad (3.1)$$

and

$$\overline{R}_4^* \left(4 \cdot p^{2\alpha+1} (pn + r) + \frac{7 \cdot p^{2\alpha+2} - 1}{6} \right) \equiv 0 \pmod{4}. \quad (3.2)$$

Proof. Setting $\ell = 4$ in (1.7) and simplifying, we obtain

$$\sum_{n=0}^{\infty} \overline{R}_4^*(n) q^n = \frac{f_4 f_2 f_1^2}{f_1^4}. \quad (3.3)$$

Employing (2.14) with $p = 2$ in (3.3) and then simplifying using (2.5), we obtain

$$\sum_{n=0}^{\infty} \overline{R}_4^*(n) q^n \equiv f_4 \phi(-q) \pmod{4}. \quad (3.4)$$

From [5, p.49], we note that

$$\phi(q) = \phi(q^{n^2}) + \sum_{r=1}^{n-1} q^{r^2} F(q^{n(n-2r)}, q^{n(n+2r)}). \quad (3.5)$$

Setting $n = 2$ in (3.5) and simplifying, we obtain

$$\phi(q) = \phi(q^4) + q\psi(q^8). \quad (3.6)$$

Replacing q by $-q$ in (3.6) and then employing in (3.4), we obtain

$$\sum_{n=0}^{\infty} \overline{R}_4^*(n) q^n \equiv f_4 (\phi(q^4) - 2q\psi(q^8)) \pmod{4}. \quad (3.7)$$

Extracting the terms involving $q^{4n+\xi}$ for $\xi = 2, 3$ from (3.7), we arrive at (i).

To prove (ii), we employ the principle of mathematical induction. Extracting the terms involving q^{4n+1} from (3.7), dividing by q and replacing q^4 by q , we obtain

$$\sum_{n=0}^{\infty} \overline{R}_4^*(4n + 1) q^n \equiv 2f_1 \psi(q^2) \pmod{4}, \quad (3.8)$$

which is $\alpha = 0$ case of (ii). Assume that (ii) is true for some integer $\alpha \geq 0$. Using (2.6) and (2.7) in (ii), we obtain

$$\begin{aligned} & \sum_{n=0}^{\infty} \overline{R}_4^* \left(4 \cdot p^{2\alpha} n + \frac{7 \cdot p^{2\alpha} - 1}{6} \right) \\ & \equiv 2 \left[\sum_{\substack{k=-(p-1)/2 \\ k \neq (\pm p-1)/6}}^{(p-1)/2} (-1)^k q^{\frac{3k^2+k}{2}} F \left(-q^{(3p^2+(6k+1)p)/2}, -q^{(3p^2-(6k+1)p)/2} \right) + (-1)^{(\pm p-1)/6} q^{(p^2-1)/24} f_{p^2} \right] \\ & \quad \times \left[\sum_{j=0}^{(p-3)/2} q^{2(j^2+j)/2} F \left(q^{2(p^2+(2j+1)p)/2}, q^{2(p^2-(2j+1)p)/2} \right) + q^{2(p^2-1)/8} \psi(q^{2p^2}) \right] \pmod{4}. \end{aligned} \quad (3.9)$$

Consider the congruence

$$\left(\frac{3k^2 + k}{2} \right) + 2 \left(\frac{j^2 + j}{2} \right) \equiv \frac{7(p^2 - 1)}{24} \pmod{p},$$

which is similar to

$$(6k + 1)^2 + 6(2j + 1)^2 \equiv 0 \pmod{p}.$$

Since $\left(\frac{-6}{p} \right) = -1$, the above congruence has only solution $k = (\pm p - 1)/6$ and $j = (\pm p - 1)/2$. So, extracting the terms involving $q^{pn+7(p^2-1)/24}$ from both sides of (3.9), dividing by $q^{7(p^2-1)/24}$ and then replacing q^p by q , we obtain

$$\sum_{n=0}^{\infty} \overline{R}_4^* \left(4p^{2\alpha+1} n + \frac{7p^{2\alpha+2} - 1}{6} \right) q^n \equiv 2f_p \psi(q^{2p}) \pmod{4}. \quad (3.10)$$

Extracting the terms involving q^{pn} from (3.10) and replacing q^p by q , we obtain

$$\sum_{n=0}^{\infty} \overline{R}_4^* \left(4p^{2(\alpha+1)} n + \frac{7p^{2\alpha+2} - 1}{6} \right) q^n \equiv 2f_1 \psi(q^2) \pmod{4}, \quad (3.11)$$

which is $\alpha + 1$ case of (3.1). Extracting the coefficients of the terms involving q^{pn+r} , for $1 \leq r \leq p - 1$, from both sides (3.10), we arrive (3.2). \square

Theorem 3.2. *For any positive integer k , we have*

- (i) $\overline{R}_{5k}^*(5n + \xi) \equiv 0 \pmod{4}$; $\xi = 2, 3$,
- (ii) $\overline{R}_{5k}^*(5n + 1) \equiv 0 \pmod{2}$.

Proof. Setting $\ell = 5k$ in (1.7) and employing (2.5), we obtain

$$\sum_{n=0}^{\infty} \overline{R}_{5k}^*(n) q^n = \frac{f_{5k} f_2}{f_1^2} = \frac{f_{5k}}{\phi(-q)}. \quad (3.12)$$

Employing Lemma 2.5 in (3.12) and simplifying using (2.14) with $p = 2$, we obtain

$$\sum_{n=0}^{\infty} \overline{R}_{5k}^*(n)q^n \equiv \frac{f_{5k} \phi(-q^{25})}{\phi(-q^5)^6} \left[\phi(-q^{25})^4 + 2q\phi(-q^{25})^3 X(q^5) - 2q^4 \phi(-q^{25})^3 Y(-q^5) \right] \pmod{4}. \quad (3.13)$$

Extracting the terms involving $q^{5n+\xi}$ for $\xi = 2, 3$ from (3.13), we arrive at (i). Similarly, (ii) follows from (3.12) by employing (2.13) with $p = 2$ and then extracting the terms involving q^{5n+1} . \square

Theorem 3.3. *For any integer $\alpha \geq 0$, we have*

- (i) $\overline{R}_6^* \left(9^\alpha n + \frac{9^\alpha - 1}{4} \right) \equiv \overline{R}_6^*(n) \pmod{3}$,
- (ii) $\overline{R}_6^* \left(9^{\alpha+1} n + \frac{21 \cdot 9^\alpha - 1}{4} \right) \equiv 0 \pmod{3}$,
- (iii) $\overline{R}_6^* \left(9^{\alpha+1} n + \frac{33 \cdot 9^\alpha - 1}{4} \right) \equiv 0 \pmod{3}$.

Proof. Setting $\ell = 6$ in (1.7), we obtain

$$\sum_{n=0}^{\infty} \overline{R}_6^*(n)q^n = \frac{f_6 f_2}{f_1^2}. \quad (3.14)$$

Simplifying (3.14) by employing (2.13) with $p = 3$ and, we obtain

$$\sum_{n=0}^{\infty} \overline{R}_6^*(n)q^n \equiv \frac{f_2^4}{f_1^2} \pmod{3}. \quad (3.15)$$

Employing (2.3) in (3.15), we obtain

$$\sum_{n=0}^{\infty} \overline{R}_6^*(n)q^n \equiv \psi^2(q) \pmod{3}. \quad (3.16)$$

Employing Lemma 2.6 in (3.16), we obtain

$$\sum_{n=0}^{\infty} \overline{R}_6^*(n)q^n \equiv F^2(q^3, q^6) + q^2 \psi^2(q^9) + 2qF(q^3, q^6) \psi(q) \pmod{3}. \quad (3.17)$$

Extracting the terms involving q^{3n+2} from (3.17), dividing by q^2 and replacing q^3 by q , we obtain

$$\sum_{n=0}^{\infty} \overline{R}_6^*(3n+2)q^n \equiv \psi^2(q^3) \pmod{3}. \quad (3.18)$$

Extracting the terms involving q^{3n} from (3.18) and replacing q^3 by q , we obtain

$$\sum_{n=0}^{\infty} \overline{R}_6^*(9n+2)q^n \equiv \psi^2(q) \pmod{3}. \quad (3.19)$$

Employing (3.16) in (3.19), we obtain

$$\sum_{n=0}^{\infty} \overline{R}_6^*(9n+2)q^n \equiv \sum_{n=0}^{\infty} \overline{R}_6^*(n)q^n \pmod{3}. \quad (3.20)$$

Iterating (3.20) with n replaced by $9n+2$ and simplifying, we obtain

$$\sum_{n=0}^{\infty} \overline{R}_6^* \left(9^\alpha n + \frac{9^\alpha - 1}{2} \right) q^n \equiv \sum_{n=0}^{\infty} \overline{R}_6^*(n)q^n \pmod{3}, \quad (3.21)$$

where $\alpha \geq 0$ is any integer. Equating the coefficients of q^n on both sides of (3.21), we arrive at (i).

Combining (3.17) and (3.21), we obtain

$$\sum_{n=0}^{\infty} \overline{R}_6^* \left(9^\alpha n + \frac{9^\alpha - 1}{4} \right) q^n \equiv F^2(q^3, q^6) + q^2 \psi^2(q^9) + 2qF(q^3, q^6)\psi(q) \pmod{3}, \quad (3.22)$$

$\alpha \geq 0$ is any integer.

Extracting the terms involving q^{3n+2} from (3.22), dividing by q^2 and replacing q^3 by q , we obtain

$$\sum_{n=0}^{\infty} \overline{R}_6^* \left(9^\alpha (3n+2) + \frac{9^\alpha - 1}{4} \right) q^n \equiv \psi^2(q^3) \pmod{3}, \quad (3.23)$$

Extracting the terms involving q^{3n+1} and q^{3n+2} from (3.23), we complete the proof of (ii) and (iii), respectively. \square

Theorem 3.4. *Let $p \geq 3$ be a prime with $\left(\frac{-1}{p}\right) = -1$ and $1 \leq r \leq p-1$. Then for any integers $\alpha \geq 0$ and $n \geq 0$, we have*

$$(i) \quad \sum_{n=0}^{\infty} \overline{R}_6^* \left(2 \cdot p^{2\alpha} n + \frac{5 \cdot p^{2\alpha} - 1}{4} \right) \equiv 2\psi(q)\psi(q^4) \pmod{3},$$

$$(ii) \quad \overline{R}_6^* \left(2 \cdot p^{2\alpha+1}(pn+r) + \frac{5 \cdot p^{2\alpha+2} - 1}{4} \right) \equiv 0 \pmod{3}.$$

Proof. Employing (2.9) in (3.15), we obtain

$$\sum_{n=0}^{\infty} \overline{R}_6^*(n)q^n \equiv \frac{f_8^5}{f_2 f_{16}^2} + 2q \frac{f_4^2 f_{16}^2}{f_2 f_8} \pmod{3}. \quad (3.24)$$

Extracting the terms involving q^{2n+1} , dividing by q and replacing q^2 by q from (3.24), we obtain

$$\sum_{n=0}^{\infty} \overline{R}_6^*(2n+1)q^n \equiv 2 \frac{f_2^2}{f_1} \cdot \frac{f_8^2}{f_4} \pmod{3}. \quad (3.25)$$

Employing (2.3) in (3.25), we obtain

$$\sum_{n=0}^{\infty} \overline{R}_6^*(2n+1)q^n \equiv 2\psi(q)\psi(q^4) \pmod{3}. \quad (3.26)$$

which is $\alpha = 0$ case of (i). Assume that (i) is true for some integer $\alpha \geq 0$. Using (2.6) in (i), we obtain

$$\begin{aligned} & \sum_{n=0}^{\infty} \overline{R}_6^* \left(2 \cdot p^{2\alpha} n + \frac{5 \cdot p^{2\alpha} - 1}{4} \right) \\ & \equiv 2 \left[\sum_{j=0}^{(p-3)/2} q^{(j^2+j)/2} F \left(q^{(p^2+(2j+1)p)/2}, q^{(p^2-(2j+1)p)/2} \right) + q^{(p^2-1)/8} \psi(q^{p^2}) \right] \\ & \quad \times \left[\sum_{t=0}^{(p-3)/2} q^{4(t^2+t)/2} F \left(q^{4(p^2+(2t+1)p)/2}, q^{4(p^2-(2t+1)p)/2} \right) + q^{4(p^2-1)/8} \psi(q^{4p^2}) \right] \pmod{3}. \end{aligned} \quad (3.27)$$

Consider the congruence

$$\binom{j^2+j}{2} + 4 \binom{t^2+t}{2} \equiv \frac{5(p^2-1)}{8} \pmod{p},$$

which is equivalent to

$$(2j+1)^2 + (4t+2)^2 \equiv 0 \pmod{p}.$$

Since $\left(\frac{-1}{p}\right) = -1$, the above congruence has only solution $j = t = (\pm p - 1)/2$.

Therefore, extracting the terms involving $q^{pn+5(p^2-1)/8}$ from both sides of (3.27), dividing throughout by $q^{5(p^2-1)/8}$ and then replacing q^p by q , we obtain

$$\sum_{n=0}^{\infty} \overline{R}_6^* \left(2 \cdot p^{2\alpha+1} n + \frac{5 \cdot p^{2\alpha+2} - 1}{4} \right) \equiv 2\psi(q^p)\psi(q^{4p}) \pmod{3}. \quad (3.28)$$

Extracting the terms involving q^{pn} from (3.28) and replacing q^p by q , we obtain

$$\sum_{n=0}^{\infty} \overline{R}_6^* \left(2 \cdot p^{2(\alpha+1)} n + \frac{5 \cdot p^{2(\alpha+1)} - 1}{4} \right) \equiv 2\psi(q)\psi(q^4) \pmod{3} \quad (3.29)$$

which is $\alpha + 1$ case of (i). Thus, by induction, we complete the proof of (i). Extracting the terms involving q^{pn+r} , for $1 \leq r \leq p-1$, from both sides (3.28), we complete the proof (ii). \square

Theorem 3.5. *We have*

- (i) $\overline{R}_8^*(4n + \xi) \equiv 0 \pmod{4}; \quad \xi = 2, 3,$
- (ii) $\overline{R}_8^*(16n + 4\xi + 1) \equiv 0 \pmod{4}, \quad \xi = 1, 2, 3,$

Proof. Setting $\ell = 8$ in (1.7), we obtain

$$\sum_{n=0}^{\infty} \overline{R}_8^*(n)q^n = \frac{f_2 f_8}{f_1^2}. \quad (3.30)$$

Employing (2.9) in (3.30), we obtain

$$\sum_{n=0}^{\infty} \overline{R}_8^*(n)q^n = \frac{f_8^6}{f_2^4 f_{16}^2} + 2q \frac{f_4^2 f_{16}^2}{f_2^4}. \quad (3.31)$$

Employing (2.14) with $p = 2$ in (3.31), we obtain

$$\sum_{n=0}^{\infty} \overline{R}_8^*(n)q^n \equiv \frac{f_8^2}{f_4^2} + 2q f_{16}^2 \pmod{4}. \quad (3.32)$$

Extracting the terms involving $q^{4n+\xi}$ for $\xi = 2, 3$ from (3.32), we arrive at (i).

Again, extracting the terms q^{4n+1} from (3.32), dividing by q and replacing q^4 by q , we obtain

$$\sum_{n=0}^{\infty} \overline{R}_8^*(4n+1)q^n \equiv 2f_4^2 \pmod{4}. \quad (3.33)$$

Extracting the terms involving $q^{4n+\xi}$ for $\xi = 1, 2, 3$ from (3.33), we arrive at (ii). □

Theorem 3.6. *We have*

- (i) $\overline{R}_8^*(4n+3) \equiv 0 \pmod{8}$,
- (ii) $\overline{R}_8^*(8n+2\xi+1) \equiv 0 \pmod{8}$, $\xi = 1, 2, 3$.

Proof. Extracting the terms involving q^{2n+1} from (3.31), dividing by q and replacing q^2 by q , we obtain

$$\sum_{n=0}^{\infty} \overline{R}_8^*(2n+1)q^n = 2 \frac{f_2^2 f_8^2}{f_1^4}. \quad (3.34)$$

Employing (2.10) in (3.34), we obtain

$$\sum_{n=0}^{\infty} \overline{R}_8^*(2n+1)q^n = 2 \frac{f_4^{14}}{f_2^{12} f_8^2} + 8q \frac{f_4^2 f_8^6}{f_2^8}. \quad (3.35)$$

Employing (2.14) with $p = 2$ in (3.35), we obtain

$$\sum_{n=0}^{\infty} \overline{R}_8^*(2n+1)q^n \equiv 2f_8^2 \pmod{8} \quad (3.36)$$

Extracting the terms involving q^{2n+1} on both sides of (3.36), we arrive at (i).

Similarly, (ii) follows from (3.36) by extracting the terms involving $q^{4n+\xi}$ for $\xi = 1, 2, 3$. □

Theorem 3.7. Let $p \geq 5$ be a prime with $\left(\frac{-3}{p}\right) = -1$ and $1 \leq r \leq p-1$. Then for any integers $\alpha \geq 0$ and $n \geq 0$, we have

$$(i) \quad \sum_{n=0}^{\infty} \overline{R}_8^* \left(8p^{2\alpha}n + \frac{4 \cdot p^{2\alpha} - 1}{3} \right) q^n \equiv 2f_1\psi(q) \pmod{8},$$

$$(ii) \quad \overline{R}_8^* \left(8p^{2\alpha+1}(pn+r) + \frac{4 \cdot p^{2\alpha+2} - 1}{3} \right) \equiv 0 \pmod{8},$$

$$(iii) \quad \frac{1}{2} \overline{R}_8^*(16n+1) \equiv p(n) + \sum_{\nu=1}^{\infty} p \left(n - \frac{\nu(\nu+1)}{2} \right) \pmod{4}.$$

Proof. Extracting the terms involving q^{4n} from (3.36) and simplifying, we obtain

$$\sum_{n=0}^{\infty} \overline{R}_8^*(8n+1)q^n \equiv 2 \frac{f_1 f_2^2}{f_1} \pmod{8}. \quad (3.37)$$

Employing (2.3) in (3.37), we obtain

$$\sum_{n=0}^{\infty} \overline{R}_8^*(8n+1)q^n \equiv 2f_1\psi(q) \pmod{8}. \quad (3.38)$$

which is the $\alpha = 0$ case of (i). Assume that (i) is true for some integer $\alpha \geq 0$. Using (2.6) in (2.7), we find that

$$\begin{aligned} & \sum_{n=0}^{\infty} \overline{R}_8^* \left(8p^{2\alpha}n + \frac{4 \cdot p^{2\alpha} - 1}{3} \right) q^n \\ & \equiv 2 \left[\sum_{\substack{k=-(p-1)/2 \\ k \neq (\pm p-1)/6}}^{(p-1)/2} (-1)^k q^{\frac{3k^2+k}{2}} F \left(-q^{(3p^2+(6k+1)p)/2}, -q^{(3p^2-(6k+1)p)/2} \right) + (-1)^{(\pm p-1)/6} q^{(p^2-1)/24} f_{p^2} \right] \\ & \quad \times \left[\sum_{j=0}^{(p-3)/2} q^{(j^2+j)/2} F \left(q^{(p^2+(2j+1)p)/2}, q^{(p^2-(2j+1)p)/2} \right) + q^{(p^2-1)/8} \psi(q^{p^2}) \right] \pmod{8}. \end{aligned} \quad (3.39)$$

Consider the congruence

$$\frac{(3k^2+k)}{2} + \frac{(j^2+j)}{2} \equiv \frac{4(p^2-1)}{24} \pmod{p},$$

which is equivalent to

$$(6k+1)^2 + 3(2j+1)^2 \equiv 0 \pmod{p}.$$

Since $\left(\frac{-3}{p}\right) = -1$, the above congruence has only solution $k = (\pm p - 1)/6$ and $j = (\pm p - 1)/2$. Therefore, extracting the terms involving $q^{pn+(p^2-1)/6}$ from both sides of

(3.39), dividing throughout by $q^{(p^2-1)/6}$ and then replacing q^p by q , we obtain

$$\sum_{n=0}^{\infty} \overline{R}_8^* \left(8p^{2\alpha+1}n + \frac{4p^{2\alpha+2} - 1}{3} \right) q^n \equiv 2f_p \psi(q^p) \pmod{8}. \quad (3.40)$$

Extracting the terms involving q^{pn} from (3.40) and replacing q^p by q , we obtain

$$\sum_{n=0}^{\infty} \overline{R}_8^* \left(8p^{2(\alpha+1)}n + \frac{4p^{2\alpha+2} - 1}{3} \right) q^n \equiv 2f_1 \psi(q) \pmod{8}, \quad (3.41)$$

which is $\alpha + 1$ case of (i). Thus, by induction the proof of (i) complete.

Extracting the terms involving q^{pn+r} , for $1 \leq r \leq p - 1$, from both sides (3.40), we arrive at (ii).

Extracting the terms involving q^{16n+1} from (3.32), dividing by q and replacing q^4 by q , we obtain

$$\sum_{n=0}^{\infty} \overline{R}_8^*(16n+1)q^n \equiv 2f_1^2 \equiv 2\frac{f_2^2}{f_1^2} \pmod{4}, \quad (3.42)$$

where we used (2.14) with $p = 2$.

Employing (1.1) and (2.3) in (3.42), we obtain

$$\frac{1}{2} \sum_{n=0}^{\infty} \overline{R}_8^*(16n+1)q^n \equiv \sum_{n=0}^{\infty} p(n)q^n + \sum_{\nu=0}^{\infty} p(n - \nu(\nu+1)/2)q^n \pmod{4}. \quad (3.43)$$

Equating the coefficient of q^n on both sides of (3.43), we arrived at (iii). \square

Theorem 3.8. *We have*

$$(i) \quad \overline{R}_\ell^*(n) + \sum_{\nu=1}^{\infty} \overline{R}_\ell^*(n - \ell\nu)p(\nu) = \overline{p}(n),$$

$$(ii) \quad \overline{R}_2^*(n) = D_2(n).$$

where $D_2(n)$ is the number of partitions of n into distinct parts such that each part has two copies.

Proof. From (1.7), we have

$$\sum_{n=0}^{\infty} \overline{R}_\ell^*(n)q^n = \frac{f_2 f_\ell}{f_1^2} \quad (3.44)$$

Using (1.5) in (3.44), we obtain

$$\sum_{n=0}^{\infty} \overline{R}_\ell^*(n)q^n \cdot \frac{1}{f_\ell} = \sum_{n=0}^{\infty} \overline{p}(n)q^n. \quad (3.45)$$

Employing (1.1) in (3.45)

$$\sum_{n=0}^{\infty} \overline{R}_\ell^*(n)q^n \left(1 + \sum_{\nu=1}^{\infty} p(\nu)q^{\ell\nu} \right) = \sum_{n=0}^{\infty} \overline{p}(n)q^n \quad (3.46)$$

Extracting the coefficient of q^n from both sides of (3.46), we arrive at (i).

Setting $\ell = 2$ in (3.44) and simplifying, we obtain

$$\sum_{n=0}^{\infty} \bar{R}_{\ell}^*(n) = \frac{f_2^2}{f_1^2} = (-q; q)_{\infty}^2. \quad (3.47)$$

Now (ii) follows immediately from (3.47). \square

Acknowledgement

The second author thanks University Grants Commission (UGC) of India for supporting her research work through CSIR-UGC Junior Research Fellowship (JRF) vide NTA-Ref. No.: 221610074648, Dated 29/11/2022.

Declarations

Author Contributions. Both authors contributed equally in the article.

Conflict of Interest. The authors declare that there is no conflict of interest regarding the publication of this article.

Human and animal rights. The authors declare that there is no research involving human participants or animals in the contained of this paper.

Data availability statements. Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

REFERENCES

- [1] Alanazi A. M., Alenazi B. M., W. J. Keith and Munagi A. O.: Refining overpartitions by properties of non overlined parts, *Contrib. Discrete Math.* **17**(2) (2022), 96-111.
- [2] Ballantine, C. and Merca, M.: Almost 3-regular overpartitions, *Ramanujan J.* **58** (2022), 957-971.
- [3] Baruah, N. D. and Ahmed, Z.: New congruences for ℓ -regular partitions for $\ell \in \{5, 6, 7, 49\}$, *The Ramanujan J.* **40** (2016), 649-668.
- [4] Baruah, N. D. and Das, K.: Parity results for 7-regular and 23-regular partitions, *Int. J. Number Theory* **11** (2015), 2221-2238.
- [5] Berndt, B. C.: *Ramanujan's Notebooks, Part III*. Springer-Verlag, New York, 1991.
- [6] Berndt, B. C. and Rankin, R. A.: *Ramanujan: Letters and Commentary*, Amer. Math. Soc. (1995).
- [7] Buragohain, P. and Saikia, N.: New congruences for overpartitions with ℓ -regular overlined parts. *J. Anal.* **31** (2023), 1819-1837.
- [8] Carlson, R. and Webb, J. J.: Infinite families of congruences for k -regular partitions, *Ramanujan J.* **33** (2014), 329-337.
- [9] Corteel, S. and Lovejoy, J.: Overpartitions, *T. Am. Math. Soc. Comp.* **356**, (2004), 1623-1635.
- [10] Cui, S. P. and Gu, N. S. S.: Arithmetic properties of l -regular partitions, *Adv. Appl. Math.* **51** (2013) 507-523.
- [11] Dandurand, B. and Penniston, D.: ℓ -divisibility of ℓ -regular partition functions, *Ramanujan J.* **19** (2009), 63-70.

- [12] Hirschhorn, M. D.: *The Power of q* , A Personal Journey, Developments in Mathematics, **49**. Springer, Cham (2017).
- [13] Hirschhorn, M. D. and Sellers, J. A.: An infinite family of overpartition congruences modulo 12, *Integers* **5** (2005), A20 1-4.
- [14] Hirschhorn, M. D. and Sellers, J. A.: Arithmetic relations for overpartitions, *J. Combin. Math. Combin. Comput.* **53** (2005), 65-73.
- [15] Hirschhorn, M. D. and Sellers, J. A.: Elementary proofs of parity results for 5-regular partitions, *Bull. Aust. Math. Soc.* **81** (2010), 58-63.
- [16] Kim, B.: A short note on the overpartition function, *Discrete Math.* **309** (2009), 2528-2532.
- [17] Lin, B. L. S.: An infinite family of congruences modulo 3 for 13-regular bipartitions, *Ramanujan J.* **39**(1) (2016), 169-178.
- [18] Mahlburg, K.: The overpartition function modulo small powers of 2, *Discrete Math.* **286** (2004), 263-267.
- [19] Penniston, D.: Arithmetic of ℓ -regular partition functions, *Int. J. Number Theory* **4** (2008), 295-302.
- [20] Ramanujan, S.: Congruence properties of partitions, *Math. Z.* **9** (1921), 147-153.
- [21] Ramanujan, S. and Hardy, G. H.: *Collected papers*, Chelsea, New York, (1962).
- [22] Saikia, N. and Boruah, C.: Congruences for ℓ -regular overpartition for $\ell \in \{5, 6, 8\}$. *Indian J. Pure Appl. Math.* **48**(2) (2017), 295-308.
- [23] Sellers, J. A.: Extending congruences for overpartitions with ℓ -regular nonoverlined parts, *Bull. Aust. Math. Soc.* (2024), doi:10.1017/S0004972724001023.
- [24] Shivanna, G. D. and Chandrappa, S.: Congruences for overpartitions with l -regular overlined parts, *J. Anal.* **31** (2023), 1819-1837
- [25] Webb, J. J.: Arithmetic of the 13-regular partition function modulo 3, *Ramanujan J.* **25** (2011), 49-56.
- [26] Xia, E. X. W.: Congruences modulo 9 and 27 for overpartitions, *Ramanujan J.* **42**(2) (2017), 301-323.
- [27] Xia, E. X. W. and Yao, O. X. M.: A proof of Keith's conjecture for 9-regular partitions modulo 3, *Int. J. Number Theory* **10** (2014), 669-674.