

ARITHMETIC PROPERTIES OF GENERALIZED CUBIC AND OVERCUBIC PARTITIONS

HIRAKJYOTI DAS , SAIKAT MAITY, AND MANJIL P. SAIKIA 

ABSTRACT. We prove several congruences satisfied by the generalized cubic and generalized overcubic partition functions, recently introduced by Amdeberhan, Sellers, and Singh. We also prove infinite families of congruences modulo powers of 2 and modulo 12 satisfied by the generalized overcubic partitions, as well as some density results that they satisfy. We use both elementary q -series techniques as well as the theory of modular forms to prove our results.

1. INTRODUCTION

An (integer) *partition* of a natural number n is a non-increasing sequence of integers $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_k$, such that $\sum_{i=1}^k \lambda_i = n$. We denote by $p(n)$, the number of partitions of n . For instance $p(5) = 7$ and the seven partitions of 5 are

$$(5), (4, 1), (3, 2), (3, 1, 1), (2, 2, 1), (2, 1, 1, 1), (1, 1, 1, 1, 1).$$

It is well-known that their generating function is given by

$$\sum_{n \geq 0} p(n)q^n = \prod_{n \geq 1} \frac{1}{(1 - q^n)} = \frac{1}{(q; q)_\infty} = \frac{1}{f_1},$$

where we have used the shorthand notations

$$(a; q)_\infty := \prod_{n \geq 0} (1 - aq^n), \quad f_k := (q^k; q^k)_\infty.$$

The arithmetic properties of partitions have been a fruitful source of mathematical discovery since their inception. Ramanujan gave three intriguing congruences for the partition function, namely he proved that, for all $n \geq 0$, we have

$$p(5n + 4) \equiv 0 \pmod{5}, \quad p(7n + 5) \equiv 0 \pmod{7}, \quad p(11n + 6) \equiv 0 \pmod{11}.$$

Since then, mathematicians have been on the lookout for such Ramanujan congruences for other restricted partition functions as well. This has led to a proliferation of research in the theory of partitions by giving rise to numerous generalizations of partitions.

One such generalization was recently studied by Amdeberhan, Sellers, and Singh [ASS25]. They studied generalized cubic partitions and generalized cubic overpartitions. A *generalized cubic partition* of weight n is a partition of $n \geq 1$ wherein each part may appear in $c \geq 1$ different colors. We denote by $a_c(n)$ the number of such generalized cubic partitions of n , and set $a_c(0) := 1$ for all $c \geq 1$. The generating function for $a_c(n)$ is given by

$$(1.1) \quad \sum_{n \geq 0} a_c(n)q^n = \frac{1}{f_1 f_2^{c-1}}.$$

2010 *Mathematics Subject Classification.* Primary 05A17, 11F03, 11P83.

Key words and phrases. cubic partitions, overcubic partitions, Ramanujan-type congruences, modular forms, Radu's algorithm.

The last two authors are supported by a Start-Up Grant from Ahmedabad University, India (Reference No. URBASASI24A5).

If we set $c = 1$ then we obtain the ordinary integer partitions, and if we set $c = 2$ then we obtain the *cubic partitions* of n . The case $c = 2$ was first studied by Chan [Cha10a]. Following Chan's work, there has been a lot of follow-up work studying cubic partitions, in particular mathematicians such as Chan [Cha10b], Xiong [Xio11], Sellers [Sel14], Gireesh [Gir17], Yao [Yao19], Hirschhorn [Hir20], Merca [Mer22], Baruah & Das [BD22], and Buragohain & Saikia [BS24] have studied arithmetic properties of cubic partitions over the years.

Amdeberhan, Sellers, and Singh [ASS25] also considered *generalized overcubic partitions*. These are defined in the same way as generalized cubic partitions and denoted by $\bar{a}_c(n)$. The generating function for $\bar{a}_c(n)$ is given by

$$(1.2) \quad \sum_{n \geq 0} \bar{a}_c(n) q^n = \frac{f_4^{c-1}}{f_1^2 f_2^{2c-3}}.$$

The case $c = 1$ are the ordinary overpartitions first studied by Corteel and Lovejoy [CL04], while the case $c = 2$ was first studied by Kim [Kim10]. Following the work of Kim, overcubic partitions and their generalizations have received a lot of attention. Mathematicians such as, Hirschhorn [Hir12], Lin [Lin14, Lin17], Sellers [Sel14], Lin–Shen–Wang [LSW17], Ray & Barman [RB20], Nayaka–Dharmendra–Mahesh Kumar [SNDMK24], and Saikia & Sarma [SS25] have studied their arithmetic properties, and it is an area of active research.

Amdeberhan, Sellers, and Singh [ASS25] obtained several congruences for the $a_c(n)$ and $\bar{a}_c(n)$ functions using both elementary and modular forms techniques. Subsequently, Guadalupe [Gua25] obtained a few more congruences for the $a_c(n)$ function. The purpose of this paper is to extend their lists and prove several more congruences for both $a_c(n)$ and $\bar{a}_c(n)$.

First, we prove the following isolated congruences for $a_c(n)$.

Theorem 1.1. *For all $n \geq 0$, we have*

$$(1.3) \quad a_{37}(43n + 12) \equiv 0 \pmod{43},$$

$$(1.4) \quad a_{41}(47n + 21) \equiv 0 \pmod{47},$$

$$(1.5) \quad a_{53}(59n + 56) \equiv 0 \pmod{59},$$

$$(1.6) \quad a_{61}(67n + 19) \equiv 0 \pmod{67},$$

$$(1.7) \quad a_{65}(71n + 32) \equiv 0 \pmod{71},$$

$$(1.8) \quad a_{73}(79n + 62) \equiv 0 \pmod{79},$$

$$(1.9) \quad a_{77}(83n + 79) \equiv 0 \pmod{83}.$$

Remark 1.2. *After we finished the proof of Theorem 1.1, Guadalupe [Gua25] updated his preprint on arXiv, which actually generalizes the above result. However, he uses elementary techniques to achieve this, so we keep our proof in this version.*

Theorem 1.3. *For all $n \geq 0$, we have*

$$(1.10) \quad a_3(7^2n + 39) \equiv 0 \pmod{7^2}.$$

Theorems 1.1 and 1.3 are proved using the theory of modular forms in Section 3.

We also give a complete characterization of the $\bar{a}_c(n)$ function modulo 4 and 8 in the following results.

Theorem 1.4. *For all $n \geq 1$, we have*

$$\bar{a}_c(n) \equiv \begin{cases} 2 \pmod{4} & \text{if } n = k^2, k \in \mathbb{Z}, \\ 2(c+1) \pmod{4} & \text{if } n = 2k^2, k \in \mathbb{Z}, \\ 0 \pmod{4} & \text{otherwise.} \end{cases}$$

Remark 1.5. *Theorem 1.4 generalizes a result of Sellers [Sel14, Theorem 2.5].*

Theorem 1.6. *For all $n \geq 1$, we have*

$$\bar{a}_c(n) \equiv \begin{cases} 2 \pmod{8} & \text{if } n = k^2, k \in \mathbb{Z}, \\ 6(c+1) \pmod{8} & \text{if } n = 2(2k)^2, k \in \mathbb{Z}, \\ 2(c+1) \pmod{8} & \text{if } n = 2(2k-1)^2, k \in \mathbb{Z}, \\ 2(c+1)(c+2) \pmod{8} & \text{if } n = 4k^2, k \in \mathbb{Z}, \\ 4(c+1) \pmod{8} & \text{if } n = k^2 + 2\ell^2, k, \ell \in \mathbb{Z}, \\ 0 \pmod{8} & \text{otherwise.} \end{cases}$$

Remark 1.7. *Theorem 1.6 generalizes a result of Sellers [Sel14, Theorem 2.7].*

Theorems 1.4 and 1.6 are proved in Section 4 using elementary techniques.

We now move on to prove a few infinite family of congruences.

Theorem 1.8. *For all $n \geq 0$ and $i \geq 1$, we have*

$$(1.11) \quad \bar{a}_{2i}(4n+3) \equiv 0 \pmod{4},$$

$$(1.12) \quad \bar{a}_{2i}(8n+6) \equiv 0 \pmod{4},$$

$$(1.13) \quad \bar{a}_{2i}(8n+5) \equiv 0 \pmod{4},$$

$$(1.14) \quad \bar{a}_{2i}(8n+7) \equiv 0 \pmod{8}.$$

Theorem 1.9. *For all $n \geq 0$, $i \geq 1$, and $\alpha \geq 0$, we have*

$$\bar{a}_{2i}(2^\alpha(4n+3)) \equiv 0 \pmod{4}.$$

Theorem 1.10. *For all $n \geq 0$, $i \geq 1$ and $k \geq 3$, we have*

$$(1.15) \quad \bar{a}_{2^{k_i-2^{k-1}-2}}(8n+5) \equiv 0 \pmod{2^{k+1}},$$

$$(1.16) \quad \bar{a}_{2^{k_i-2^{k-1}-2}}(8n+7) \equiv 0 \pmod{2^{k+2}}.$$

Theorem 1.11. *For all $n \geq 0$, $k \geq 1$ and $i \geq 0$, we have*

$$(1.17) \quad \bar{a}_{2^{k_i+\frac{2k-6}{2}}}(8n+7) \equiv 0 \pmod{2^{k+3}},$$

$$(1.18) \quad \bar{a}_{2^{k_i+\frac{2k-2}{2}}}(8n+7) \equiv 0 \pmod{2^{k+3}}.$$

We also prove a family of congruences modulo 12.

Theorem 1.12. *For all $n \geq 0$ and $i \geq 1$, we have*

$$(1.19) \quad \bar{a}_{9i}(9n+6) \equiv 0 \pmod{12},$$

$$(1.20) \quad \bar{a}_{9i+4}(9n+6) \equiv 0 \pmod{12},$$

$$(1.21) \quad \bar{a}_{9i+8}(9n+6) \equiv 0 \pmod{12}.$$

Theorems 1.8, 1.9, 1.10, 1.11, and 1.12 are proved in Section 5 using elementary techniques.

Along with the study of Ramanujan-type congruences, the study of distribution of the coefficients of a formal power series modulo M is also an interesting problem to explore. Given an integral power series $A(q) := \sum_{n=0}^{\infty} a(n)q^n$ and $0 \leq r \leq M$, the arithmetic density $\delta_r(A, M; X)$ is defined as

$$\delta_r(A, M; X) = \frac{\#\{n \leq X : a(n) \equiv r \pmod{M}\}}{X}.$$

An integral power series A is called *lacunary modulo M* if

$$\lim_{X \rightarrow \infty} \delta_0(A, M; X) = 1,$$

which means that almost all the coefficients of A are divisible by M . It turns out that the partition function $\bar{a}_c(n)$ also satisfies such results. Specifically, we prove the following results in this section.

Theorem 1.13. *Let $k \geq 1$, $\alpha \geq 1$ and $m \geq 1$ be integers with $\gcd(m, 2) = 1$. Then the set*

$$\{n \in \mathbb{N} : \bar{a}_{2^\alpha m} \equiv 0 \pmod{2^k}\}$$

has arithmetic density 1, namely,

$$\lim_{X \rightarrow \infty} \frac{\#\{0 \leq n \leq X : \bar{a}_{2^\alpha m} \equiv 0 \pmod{2^k}\}}{X} = 1.$$

Theorem 1.14. *Let $k \geq 2$, $\alpha \geq 1$ and $m \geq 1$ be integers with $\gcd(m, 2) = 1$ such that $3^{k-2} \geq 2^{\alpha-3}m$, then the set*

$$\{n \in \mathbb{N} : \bar{a}_{2^\alpha m} \equiv 0 \pmod{3^k}\}$$

has arithmetic density 1, namely,

$$\lim_{X \rightarrow \infty} \frac{\#\{0 \leq n \leq X : \bar{a}_{2^\alpha m} \equiv 0 \pmod{3^k}\}}{X} = 1.$$

Theorem 1.15. *Let $k \geq 1$, a_1, a_2, \dots, a_m be positive integers for some positive integer m , for $t = p_1^{a_1} \cdots p_m^{a_m}$ where p_i 's are prime numbers greater than or equal to 5 with $2p_i^{k+a_i} \geq 5t$, the set*

$$\{n \in \mathbb{N} : \bar{a}_t \equiv 0 \pmod{p_i^k}\}$$

has arithmetic density 1, namely,

$$\lim_{X \rightarrow \infty} \frac{\#\{0 \leq n \leq X : \bar{a}_t \equiv 0 \pmod{p_i^k}\}}{X} = 1.$$

The fact that the action of Hecke algebras on the spaces of modular forms of level 1 modulo 2 is locally nilpotent was first observed by Serre [Ser75a, Ser75b] and proved by Tate [Tat94]. This was further generalized by Ono and Taguchi [OT05] to higher levels. We can use Ono and Taguchi's result to prove the following theorem.

Theorem 1.16. *Let n be a non-negative integer, $\alpha \geq 1$, $m \geq 1$ are integers with $\gcd(m, 2) = 1$, and N is a positive integer such that $N = 1, 3, 5, 15$ or 17 . Then there exists an integer $i \geq 0$ such that for every $j \geq 1$ and q_1, q_2, \dots, q_{i+j} are odd primes not dividing N , we have*

$$\bar{a}_{2^\alpha m} \left(\frac{q_1 \cdots q_{i+j} \cdot n}{24} \right) \equiv 0 \pmod{3^j}.$$

Theorems 1.13, 1.14, 1.15, and 1.16 are proved in Section 6.

This paper is arranged as follows: Section 2 contains some preliminary theory required in our later proofs, Sections 3 – 6 contains the proofs of the above results, and finally we close the paper with some concluding remarks in Section 7.

2. PRELIMINARIES

2.1. Various results on modular forms. Let \mathbb{H} be the complex upper half-plane. We define the following matrix groups

$$\Gamma := \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} : a, b, c, d \in \mathbb{Z}, ad - bc = 1 \right\},$$

$$\Gamma_\infty := \left\{ \begin{bmatrix} 1 & n \\ 0 & 1 \end{bmatrix} \in \Gamma : n \in \mathbb{Z} \right\}.$$

For a positive integer N , let

$$\Gamma_0(N) := \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \Gamma : c \equiv 0 \pmod{N} \right\},$$

$$\Gamma_1(N) := \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \Gamma_0(N) : a \equiv d \equiv 1 \pmod{N} \right\}.$$

We denote by $M_\ell(\Gamma_1(N))$ for a positive integer l , the complex vector space of modular forms of weight ℓ with respect to $\Gamma_1(N)$. The index of $\Gamma_0(N)$ in Γ is

$$[\Gamma : \Gamma_0(N)] = N \prod_{\ell|N} \left(1 + \frac{1}{\ell}\right),$$

where ℓ is a prime divisor of N .

Definition 2.1. *If χ is a Dirichlet character modulo N , then a form $f(z) \in M_l(\Gamma_1(N))$ has Nebentypus character χ if*

$$f\left(\frac{az+b}{cz+d}\right) = \chi(d)(cz+d)^l f(z),$$

for all $z \in \mathbb{H}$ and all $\begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \Gamma_0(N)$. The space of such modular forms is denoted by $M_\ell(\Gamma_0(N), \chi)$.

Recall that the Dedekind's eta-function $\eta(z)$ is defined by

$$\eta(z) := q^{1/24}(q; q)_\infty = q^{1/24} \prod_{n=1}^{\infty} (1 - q^n),$$

where $q := e^{2\pi iz}$ and $z \in \mathbb{H}$. A function $f(z)$ is called an eta-quotient if it is of the form

$$f(z) = \prod_{\delta|N} \eta(\delta z)^{r_\delta},$$

where N is a positive integer and r_δ is an integer. We now recall two theorems from [Ono04, p. 18] which will be used to prove some of our results.

Theorem 2.2. [Ono04, Theorems 1.64 and 1.65] *If $f(z) = \prod_{\delta|N} \eta(\delta z)^{r_\delta}$ is an eta-quotient with $\ell = \frac{1}{2} \sum_{\delta|N} r_\delta \in \mathbb{Z}$, with*

$$\sum_{\delta|N} \delta r_\delta \equiv 0 \pmod{24} \quad \text{and} \quad \sum_{\delta|N} \frac{N}{\delta} r_\delta \equiv 0 \pmod{24},$$

then $f(z)$ satisfies

$$f\left(\frac{az+b}{cz+d}\right) = \chi(d)(cz+d)^\ell f(z),$$

for every $\begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \Gamma_0(N)$. Here the character χ is defined by $\chi(d) := \left(\frac{(-1)^\ell \prod_{\delta|N} \delta^{r_\delta}}{d}\right)$. In addition, if c , d , and N are positive integers with $d|N$ and $\gcd(c, d) = 1$, then the order of vanishing of $f(z)$ at the cusp $\frac{c}{d}$ is $\frac{N}{24} \sum_{\delta|N} \frac{\gcd(d, \delta)^2 r_\delta}{\gcd(d, \frac{N}{d}) d \delta}$.

We also recall the following result of Sturm [Stu87] which will be used in the next section.

Theorem 2.3. *Let k be an integer and $g(z) = \sum_{n=0}^{\infty} a(n)q^n$ a modular form of weight k for $\Gamma_0(N)$. For any given positive integer m , if $a(n) \equiv 0 \pmod{m}$ holds for all $n \leq \frac{kN}{12} \prod_{p \text{ prime}, p|N} \left(1 + \frac{1}{p}\right)$ then $a(n) \equiv 0 \pmod{m}$ holds for any $n \geq 0$.*

We now recall the definition of Hecke operators. Let m be a positive integer and $f(z) = \sum_{n=0}^{\infty} a(n)q^n \in M_\ell(\Gamma_0(N), \chi)$. Then the action of Hecke operator T_m on $f(z)$ is defined by

$$f(z) | T_m := \sum_{n=0}^{\infty} \left(\sum_{d|\gcd(n,m)} \chi(d) d^{\ell-1} a\left(\frac{nm}{d^2}\right) \right) q^n.$$

In particular, if $m = p$ is prime, we have

$$(2.1) \quad f(z) | T_p := \sum_{n=0}^{\infty} \left(a(pn) + \chi(p) p^{\ell-1} a\left(\frac{n}{p}\right) \right) q^n.$$

We take by convention that $a(n/p) = 0$ whenever $p \nmid n$. The next result follows directly from (2.1).

Theorem 2.4. *Let p be a prime, $g(z) \in \mathbb{Z}[[q]]$, $h(z) \in \mathbb{Z}[[q^p]]$, and $k > 1$. Then, we have*

$$(g(z)h(z)) | T_p \equiv (g(z) | T_p \cdot h(z/p)) \pmod{p}.$$

We also need the following result in Section 6.

Theorem 2.5. [Ono04, pp. 43] *Let $g(z) \in M_\ell(\Gamma_0(N), \chi)$ has a Fourier expansion*

$$g(z) = \sum_{n=0}^{\infty} b(n)q^n \in \mathbb{Z}[[q]],$$

Then for a positive integer r , there is a constant $\alpha > 0$ such that

$$\#\{0 < n \leq X : b(n) \not\equiv 0 \pmod{r}\} = \mathcal{O}\left(\frac{X}{(\log X)^\alpha}\right).$$

Equivalently

$$\lim_{X \rightarrow \infty} \frac{\#\{0 < n \leq X : b(n) \not\equiv 0 \pmod{r}\}}{X} = 0.$$

2.2. Radu's Algorithm. We also need some results which describe an algorithmic approach to proving partition concurrences, developed by Radu [Rad09]. For integers $M \geq 1$, suppose that $R(M)$ be the set of all the integer sequences

$$(r_\delta) := (r_{\delta_1}, r_{\delta_2}, r_{\delta_3}, \dots, r_{\delta_k})$$

indexed by all the positive divisors δ of M , where $1 = \delta_1 < \delta_2 < \dots < \delta_k = M$. For positive integers m , and any integer s , let $[s]_m$ denote the residue class of s in $\mathbb{Z}_m := \mathbb{Z}/m\mathbb{Z}$. Let \mathbb{Z}_m^* be the set of all invertible elements in \mathbb{Z}_m . Let $\mathbb{S}_m \subseteq \mathbb{Z}_m$ be the set of all squares in \mathbb{Z}_m^* . For $(r_\delta) \in R(M)$ and $t \in \{0, 1, \dots, m-1\}$, we define a subset $P(t) \subseteq \{0, 1, \dots, m-1\}$ by

$$(2.2) \quad P(t) := \left\{ t' \in \{0, 1, \dots, m-1\} : t' \equiv ts + \frac{s-1}{24} \sum_{\delta|M} \delta r_\delta \pmod{m}, \text{ for some } [s]_{24m} \in \mathbb{S}_{24m} \right\}.$$

Let m, M, N be positive integers. For $\gamma = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \Gamma$, $(r_\delta) \in R(M)$ and $(r'_\delta) \in R(N)$, we also define

$$p(\gamma) := \min_{\lambda \in \{0, 1, \dots, m-1\}} \frac{1}{24} \sum_{\delta|M} r_\delta \frac{\gcd(\delta(a + k\lambda c), mc)^2}{\delta m}$$

and

$$p'(\gamma) := \frac{1}{24} \sum_{\delta|N} r'_\delta \frac{\gcd(\delta, c)^2}{\delta}.$$

Suppose m, M and N are positive integers, $t \in \{0, 1, \dots, m-1\}$ and $(r_\delta) \in R(M)$. Let $k := \gcd(m^2 - 1, 24)$ and define the set Δ^* consists of all tuples $(m, M, N, t, (r_\delta))$ such that all of the following conditions are satisfied:

- (1) Prime divisors of m are also prime divisors of N .
- (2) If $\delta|M$, then $\delta|mN$ for every $\delta \geq 1$ with $r_\delta \neq 0$.
- (3) $24|kN \sum_{\delta|M} \frac{r_\delta m N}{\delta}$.
- (4) $8|kN \sum_{\delta|M} r_\delta$.
- (5) $\frac{24m}{\gcd(-24kt - k \sum_{\delta|M} \delta r_\delta, 24m)} | N$.
- (6) If $2|m$, then either $4|kN$ and $8|sN$ or $2|s$ and $8|(1-j)N$, where $\prod_{\delta|M} \delta^{|r_\delta|} = 2^s \cdot j$, here s and j are nonnegative integers with j odd.

We have the following two results useful for us in the next section.

Lemma 2.6. [Rad09, Lemma 4.5] *Suppose that $(m, M, N, t, (r_\delta)) \in \Delta^*$, $(r'_\delta) := (r'_\delta)_{\delta|N} \in R(N)$, $\{\gamma_1, \gamma_2, \dots, \gamma_n\} \subseteq \Gamma$ is a complete set of representatives of the double cosets of $\Gamma_0(N) \backslash \Gamma / \Gamma_\infty$, and $t_{min} = \min_{t' \in P(t)} t'$,*

$$\nu := \frac{1}{24} \left\{ \left(\sum_{\delta|M} r_\delta + \sum_{\delta|N} r'_\delta \right) [\Gamma : \Gamma_0(N)] - \sum_{\delta|N} \delta r'_\delta - \frac{1}{m} \sum_{\delta|M} \delta r_\delta \right\} - \frac{t_{min}}{m},$$

$p(\gamma_j) + p'(\gamma_j) \geq 0$ for all $1 \leq j \leq n$, and $\sum_{n=0}^{\infty} A(n)q^n := \prod_{\delta|M} f_\delta^{r_\delta}$. If for some integers $u \geq 1$, all $t' \in P(t)$, and $0 \leq n \leq \lfloor \nu \rfloor$, $A(mn + t') \equiv 0 \pmod{u}$ is true, then for integers $n \geq 0$ and all $t' \in P(t)$, we have $A(mn + t') \equiv 0 \pmod{u}$.

Lemma 2.7. [RS11, Lemma 2.6] *If N or $\frac{1}{2}N$ is a square-free integer, then*

$$\bigcup_{\delta|N} \Gamma_0(N) \begin{bmatrix} 1 & 0 \\ \delta & 1 \end{bmatrix} \Gamma_\infty = \Gamma.$$

3. PROOF OF THEOREMS 1.1 AND 1.3

Proof of Theorem 1.1. Since the proofs of the congruences are similar, we only prove (1.3) in detail. We consider the η -quotient

$$G_1(z) := \frac{\eta(z)^{816}}{\eta(2z)^{36}}.$$

Using Theorem 2.2, we see that $G_1(z)$ is a modular form of weight 390, and level 4, so $G_1(z) \in \mathcal{M}_{390}(\Gamma_0(4))$. From the definition of $a_c(n)$, we have

$$G_1(z) = \frac{q^{31}}{f_1 f_2^{36}} \cdot f_1^{817} = \left(\sum_{n=0}^{\infty} a_{37}(n) q^{n+31} \right) \cdot f_1^{817}.$$

Applying the Hecke operator T_{43} on $G_1(z)$, we calculate that

$$G_1(z) | T_{43} \equiv \left(\sum_{n=0}^{\infty} a_{37}(43n + 12) q^{n+1} \right) \cdot f_1^{817} \pmod{43}.$$

We know that $G_1(z) | T_{43} \in \mathcal{M}_{390}(\Gamma_0(4))$, By Theorem 2.3, the Sturm bound for this space of forms is 195. Using Mathematica, we verify that the coefficients of q -expansion of $G_1(z) | T_{43}$ up to the desired bound are congruent to 0 (mod 43). Thus, $G_1(z) | T_{43} \equiv 0 \pmod{43}$, which proves (1.3).

For the other congruences in Theorem 1.1, we mention the relevant data required to prove them in the table below.

Congruences	η -quotient	Level	Weight	Sturm bound
(1.4)	$\frac{\eta(z)^{704}}{\eta(2z)^{40}}$	2	332	83
(1.5)	$\frac{\eta(z)^{176}}{\eta(2z)^{52}}$	4	62	31
(1.6)	$\frac{\eta(z)^{1272}}{\eta(2z)^{60}}$	4	606	303
(1.7)	$\frac{\eta(z)^{1064}}{\eta(2z)^{64}}$	2	500	125
(1.8)	$\frac{\eta(z)^{552}}{\eta(2z)^{72}}$	2	240	60
(1.9)	$\frac{\eta(z)^{248}}{\eta(2z)^{76}}$	4	86	43

□

Proof of Theorem 1.3. We notice

$$\sum_{n \geq 0} a_3(n)q^n = \frac{1}{f_1 f_2^2} \equiv \frac{f_1^{48}}{f_2^2 f_7^7} \pmod{49}.$$

So, we choose $(m, M, N, t, (r_\delta)) = (49, 14, 14, 39, (48, -2, -7, 0))$. It is easy to verify that this choice of $(m, M, N, t, (r_\delta))$ satisfies the Δ^* conditions and from equation (2.2) we see that $P(t) = \{39\}$.

By Lemma 2.7, we know that $\left\{ \begin{bmatrix} 1 & 0 \\ \delta & 1 \end{bmatrix} : \delta | 14 \right\}$ forms a complete set of double coset representatives of

$\Gamma_0(N) \backslash \Gamma / \Gamma_\infty$. Let $\gamma_\delta = \begin{bmatrix} 1 & 0 \\ \delta & 1 \end{bmatrix}$, and for the choice of $(r'_\delta) = (18, 0, 0, 0)$ we see that

$$p(\gamma_\delta) + p'(\gamma_\delta) \geq 0, \quad \text{for all } \delta | N.$$

By Lemma 2.6, we compute $\lfloor \nu \rfloor = 57$. Using Mathematica we verify (1.10) for $n \leq 57$, which proves the result. □

4. PROOF OF THEOREMS 1.4 AND 1.6

Let

$$\bar{F}_c(q) := \sum_{n \geq 0} \bar{a}_c(n)q^n.$$

From (1.2), we can easily prove that

$$(4.1) \quad \bar{F}_c(q) = \varphi(q)\varphi(q^2)^{c-1}\bar{F}_c(q^2)^2,$$

where

$$\varphi(q) := \sum_{k=-\infty}^{\infty} q^{k^2} = \frac{f_2^5}{f_1^2 f_4^2}$$

is Ramanujan's φ -function. We can then iterate (4.1) to prove the next theorem.

Lemma 4.1. *Let $c \geq 3$ be an integer. Then*

$$\bar{F}_{c-1}(q) = \varphi(q) \prod_{i \geq 1} \varphi(q^{2^i})^{c \cdot 2^{i-1}}.$$

Proof. We have

$$\begin{aligned}
\bar{F}_{c-1}(q) &= \frac{f_4^{c-2}}{f_1^2 f_2^{2c-5}} = \varphi(q)\varphi(q^2)^{c-2}\bar{F}_{c-1}(q^2)^2 \\
&= \varphi(q)\varphi(q^2)^{c-2}(\varphi(q^2)\varphi(q^4)^{c-2}\bar{F}_{c-1}(q^4)^2)^2 \\
&= \varphi(q)\varphi(q^2)^c\varphi(q^4)^{2(c-2)}\bar{F}_{c-1}(q^4)^4 \\
&= \varphi(q)\varphi(q^2)^c\varphi(q^4)^{2(c-2)}(\varphi(q^4)\varphi(q^8)^{c-2}\bar{F}_{c-1}(q^8)^2)^4 \\
&= \varphi(q)\varphi(q^2)^c\varphi(q^4)^{2c}\varphi(q^8)^{4(c-2)}\bar{F}_{c-1}(q^8)^8 \\
&\vdots
\end{aligned}$$

The result follows by continuing to iterate (4.1) indefinitely. \square

Using Lemma 4.1, we now prove Theorems 1.4 and 1.6.

Proof of Theorem 1.4. From the Binomial theorem, it is clear that $\varphi(q^{2i})^{t \cdot 2^{i-1}} \equiv 1 \pmod{4}$ for each $i \geq 2$. Thus,

$$\begin{aligned}
\sum_{n \geq 0} \bar{a}_c(n)q^n &= \varphi(q) \prod_{i \geq 1} \varphi(q^{2^i})^{(c+1)2^{i-1}} \\
&\equiv \varphi(q)\varphi(q^2)^{c+1} \pmod{4} \\
&= \left(1 + 2 \sum_{n \geq 1} q^{n^2}\right) \left(1 + 2 \sum_{n \geq 1} q^{2n^2}\right)^{c+1} \\
&= \left(1 + 2 \sum_{n \geq 1} q^{n^2}\right) \left(\sum_{k=0}^{c+1} \binom{c+1}{k} 2^k \left(\sum_{n \geq 1} q^{2n^2}\right)^k\right) \\
&\equiv \left(1 + 2 \sum_{n \geq 1} q^{n^2}\right) \left(1 + 2(c+1) \sum_{n \geq 1} q^{2n^2}\right) \pmod{4} \\
&\equiv 1 + 2 \sum_{n \geq 1} q^{n^2} + 2(c+1) \sum_{n \geq 1} q^{2n^2} \pmod{4}.
\end{aligned}$$

Hence, the result follows from the above identity. \square

Proof of Theorem 1.6. From the Binomial theorem, it is clear that $\varphi(q^{2i})^{t \cdot 2^{i-1}} \equiv 1 \pmod{8}$ for each $i \geq 3$. Thus,

$$\begin{aligned}
\sum_{n \geq 0} \bar{a}_c(n)q^n &= \varphi(q) \prod_{i \geq 1} \varphi(q^{2^i})^{(c+1)2^{i-1}} \\
&\equiv \varphi(q)\varphi(q^2)^{c+1}\varphi(q^4)^{2(c+1)} \pmod{8} \\
&= \left(1 + 2 \sum_{n \geq 1} q^{n^2}\right) \left(1 + 2 \sum_{n \geq 1} q^{2n^2}\right)^{c+1} \left(1 + 2 \sum_{n \geq 1} q^{4n^2}\right)^{2(c+1)} \\
&= \left(1 + 2 \sum_{n \geq 1} q^{n^2}\right) \left(\sum_{k=0}^{c+1} \binom{c+1}{k} 2^k \left(\sum_{n \geq 1} q^{2n^2}\right)^k\right) \left(\sum_{k=0}^{2(c+1)} \binom{2(c+1)}{k} 2^k \left(\sum_{n \geq 1} q^{4n^2}\right)^k\right) \\
&\equiv \left(1 + 2 \sum_{n \geq 1} q^{n^2}\right) \left(1 + 2(c+1) \sum_{n \geq 1} q^{2n^2} + 2c(c+1) \sum_{n \geq 1} q^{4n^2}\right) \\
&\quad \left(1 + 4(c+1) \sum_{n \geq 1} q^{4n^2} + 4(c+1) \sum_{n \geq 1} q^{8n^2}\right) \pmod{8} \\
&\equiv 1 + 2 \sum_{n \geq 1} q^{n^2} + 2(c+1) \sum_{n \geq 1} q^{2(2n-1)^2} + 6(c+1) \sum_{n \geq 1} q^{2(2n)^2}
\end{aligned}$$

$$+ 2(c+1)(c+2) \sum_{n \geq 1} q^{4n^2} + 4(c+1) \sum_{m, n \geq 1} q^{n^2+2m^2} \pmod{8}.$$

Hence, the result follows from the above identity. \square

5. PROOFS OF THEOREMS 1.8, 1.9, 1.10, 1.11, AND 1.12

We need the following 2- and 3-dissection formulas.

Lemma 5.1. *We have*

$$(5.1) \quad \varphi(q) = \varphi(q^4) + 2q\psi(q^8),$$

$$(5.2) \quad f_1^2 = \frac{f_2 f_8^5}{f_4^2 f_{16}^2} - 2q \frac{f_2 f_{16}^2}{f_8},$$

$$(5.3) \quad \frac{1}{f_1^2} = \frac{f_8^5}{f_2^5 f_{16}^2} + 2q \frac{f_4^2 f_{16}^2}{f_2^5 f_8},$$

$$(5.4) \quad \varphi(q) = \varphi(q^9) + 2q\Omega(q^3),$$

$$(5.5) \quad \frac{f_2^2}{f_1} = \frac{f_6 f_9^2}{f_3 f_{18}} + q \frac{f_{18}^2}{f_9},$$

$$(5.6) \quad \frac{f_1}{f_4} = \frac{f_6 f_9 f_{18}}{f_{12}^3} - q \frac{f_3 f_{18}^4}{f_9^2 f_{12}^3} - q^2 \frac{f_6^2 f_9 f_{36}^3}{f_{12}^4 f_{18}^2},$$

where $\psi(q) := \frac{f_2^2}{f_1}$ and $\Omega(q) := \frac{f_2^2 f_3 f_{12}}{f_1 f_4 f_6}$.

Proof. (5.1) and (5.3) are the same identity in [Hir17, Eq. (1.9.4)], (5.2) can be obtained by replacing q by $-q$ in (5.1), while (5.4) and (5.5) are [Hir17, Eq. (26.1.1), (26.1.2)]. We prove (5.6) below. We need the following identities

$$(5.7) \quad \frac{f_6 f_{12}^2 f_{18}^2 f_{36}^2}{f_3} = \frac{f_{18}^9}{f_9^3} + q^3 \frac{f_6^3 f_{36}^6}{f_3 f_{12}^2},$$

$$(5.8) \quad f_9^3 f_{12} = \frac{f_3 f_{12}^4 f_{18}^2}{f_6^2 f_{36}} + q^3 f_3 f_{36}^3,$$

$$(5.9) \quad \frac{f_1}{f_2^2} = \frac{f_3^2 f_9^3}{f_6^6} - q \frac{f_3^3 f_{18}^3}{f_6^7} + q^2 \frac{f_3^4 f_{18}^6}{f_6^8 f_9^3},$$

$$(5.10) \quad \frac{f_1 f_4}{f_2} = \frac{f_3 f_{12} f_{18}^5}{f_6^2 f_9^2 f_{36}^2} - q \frac{f_9 f_{36}}{f_{18}}.$$

(5.8) and (5.9) are [Hir17, Eqs. (22.7.5) and (37.2.3)] in disguise, respectively. (5.7) and (5.10) can be obtained by replacing q by $-q$ in (5.8) and (5.5), respectively. With the help of (5.9) and (5.10), we have

$$\begin{aligned} \frac{f_1}{f_4} &= \frac{f_2}{f_4^2} \cdot \frac{f_1 f_4}{f_2} \\ &= \frac{f_3 f_9}{f_{12}^5 f_{18} f_{36}^2} \left(\frac{f_{18}^9}{f_9^3} + q^3 \frac{f_6^3 f_{36}^6}{f_3 f_{12}^2} \right) - q \frac{f_6^2 f_{18}^2 f_{36}}{f_9^2 f_{12}^2} (f_9^3 f_{12} - f_3 f_{36}^3 q^3) \\ &\quad - q^2 \frac{f_3 f_6 f_9 f_{36}}{f_{12} f_{18}^4} \left(\frac{f_{18}^9}{f_9^3} + q^3 \frac{f_6^3 f_{36}^6}{f_3 f_{12}^2} \right). \end{aligned}$$

Hence, (5.6) is evident after employing (5.7) and (5.8) in the above identity. \square

We also need the following easily proved generating functions.

Lemma 5.2. For all $c \geq 1$, we have

$$\sum_{n \geq 0} \bar{a}_c(2n)q^n = \frac{f_2^{c-1} f_4^5}{f_1^{2c+2} f_8^2},$$

$$\sum_{n \geq 0} \bar{a}_c(2n+1)q^n = 2 \frac{f_8^2}{f_4} \left(\frac{f_2}{f_1^2} \right)^{c+1}.$$

Proof. We have

$$\sum_{n \geq 0} \bar{a}_c(n)q^n = \frac{f_4^{c-1}}{f_1^2 f_2^{2c-3}} = \frac{f_4^{c-1}}{f_2^{2c-3}} \left(\frac{f_8^5}{f_2^5 f_2^2} + 2q \frac{f_4^2 f_2^2}{f_2^5 f_8} \right) \text{ thanks to Lemma 5.1 .}$$

Now, extracting the terms involving even and odd powers of q respectively from the above and then setting $q^2 \rightarrow q$, we obtain the given generating functions in the lemma. \square

Now, we are ready to prove Theorem 1.8.

Proof of Theorem 1.8. Using (5.3) and Lemma 5.2, we have

$$\begin{aligned} \sum_{n \geq 0} \bar{a}_c(2n+1)q^n &= 2 \frac{f_8^2}{f_4} \left(\frac{f_2}{f_1^2} \right)^{c+1} = 2 \frac{f_8^2 f_2^{c+1}}{f_4} \left(\frac{f_8^5}{f_2^5 f_2^2} + 2q \frac{f_4^2 f_2^2}{f_2^5 f_8} \right)^{c+1} \\ &= 2 \frac{f_8^2 f_2^{c+1}}{f_4} \sum_{k=0}^{c+1} \binom{c+1}{k} 2^k q^k \left(\frac{f_4^2 f_2^2}{f_2^5 f_8} \right)^k \left(\frac{f_8^5}{f_2^5 f_2^2} \right)^{c+1-k} \\ (5.11) \quad &= 2 \frac{f_8^{5c+7}}{f_2^{4c+4} f_4^{2c+2} f_8} \sum_{k=0}^{c+1} 2^k q^k \binom{c+1}{k} \frac{f_4^{2k} f_8^{4k}}{f_8^{6k}}. \end{aligned}$$

From the above identity, extracting the terms with even and odd exponents of q and then letting $q^2 \rightarrow q$ and $c = 2i$ for $i \geq 1$, we arrive at

$$(5.12) \quad \sum_{n \geq 0} \bar{a}_{2i}(4n+1)q^n = 2 \frac{f_4^{10i+7}}{f_1^{8i+4} f_2^{4i+2} f_8} \sum_{k=0}^i \binom{2i+1}{2k} 2^{2k} q^k \frac{f_2^{4k} f_8^{8k}}{f_4^{12k}},$$

$$(5.13) \quad \sum_{n \geq 0} \bar{a}_{2i}(4n+3)q^n = 4 \frac{f_2 f_4^{10i+1}}{f_1^{8i+4} f_8^{4i-2}} \sum_{k=0}^i \binom{2i+1}{2k+1} 2^{2k} q^k \frac{f_2^{4k} f_8^{8k}}{f_4^{12k}}.$$

Equation (5.13) proves the congruence (1.11).

By a similar process, starting from the generating function for $\bar{a}_c(2n)$ from Lemma 5.2, we can arrive at

$$(5.14) \quad \sum_{n \geq 0} \bar{a}_{2i}(4n+2)q^n = 2 \frac{f_2^7 f_4^{10i+1}}{f_1^{8i+6} f_8^{4i-2}} \sum_{k=0}^i 2^{2k} q^k \binom{2i+1}{2k+1} \frac{f_2^{4k} f_8^{8k}}{f_4^{12k}}.$$

Using (5.14) and (5.3), and then extracting the coefficients of odd powers of q , we obtain

$$\sum_{n \geq 0} \bar{a}_{2i}(8n+6)q^n \equiv 0 \pmod{4}.$$

This clearly proves (1.12).

Similarly, using (5.12), (5.13) and (5.3), and then extracting the coefficients of odd powers of q , by a similar process, we prove that

$$\sum_{n \geq 0} \bar{a}_{2i}(8n+5)q^n \equiv 0 \pmod{4},$$

$$\sum_{n \geq 0} \bar{a}_{2i}(8n+7)q^n \equiv 0 \pmod{8}.$$

These above prove (1.13) and (1.14). \square

Proof of Theorem 1.9. This follows easily by observing from (1.2) and Lemma 5.2 that

$$\bar{a}_{2i}(2n) \equiv \bar{a}_{2i}(n) \pmod{4}.$$

Together with (1.11), (1.12), the above gives the result. \square

Proof of Theorem 1.10. Using Lemma 5.2, we have

$$\sum_{n \geq 0} \bar{a}_{2^k i - 2^{k-1} - 2}(2n+1)q^n = 2 \frac{f_2^{2^k i - 2^{k-1} - 1} f_8^2}{f_1^{2^{k+1} i - 2^{k-2}} f_4} \equiv 2 \frac{f_1^{2^k+2} f_8^2}{f_2^{2^{k-1}+1} f_4} \pmod{2^{k+2}},$$

which, with the help of the 2-dissection (5.2) of f_1^2 , gives the following dissections

$$(5.15) \quad \sum_{n \geq 0} \bar{a}_{2^k i - 2^{k-1} - 2}(4n+1)q^n \equiv 2 \sum_{r=0}^{2^k-2} 2^{2r} \binom{2^{k-1}+1}{2r} q^r \frac{f_4^{5 \times 2^{k-1} - 12r + 7}}{f_2^{2^k - 4r + 3} f_8^{2^k - 8r + 2}} \pmod{2^{k+2}},$$

$$(5.16) \quad \sum_{n \geq 0} \bar{a}_{2^k i - 2^{k-1} - 2}(4n+3)q^n \equiv -4 \sum_{r=0}^{2^k-2} 2^{2r} \binom{2^{k-1}+1}{2r+1} q^r \frac{f_4^{5 \times 2^{k-1} - 12r + 1}}{f_2^{2^k - 4r + 1} f_8^{2^k - 8r - 2}} \pmod{2^{k+2}}.$$

By the properties of binomial coefficients, for $r \geq 1$, we have

$$2^{2r} \binom{2^{k-1}+1}{2r} \equiv 0 \pmod{2^k}, \quad 2^{2r} \binom{2^{k-1}+1}{2r+1} \equiv 0 \pmod{2^k}.$$

Therefore, (5.15) and (5.16) reduce to

$$\sum_{n \geq 0} \bar{a}_{2^k i - 2^{k-1} - 2}(4n+1)q^n \equiv 2 \frac{f_4^{5 \times 2^{k-1} + 7}}{f_2^{2^k + 3} f_8^{2^k + 2}} \pmod{2^{k+1}},$$

$$\sum_{n \geq 0} \bar{a}_{2^k i - 2^{k-1} - 2}(4n+3)q^n \equiv -4(2^{k-1}+1) \frac{f_4^{5 \times 2^{k-1} + 1}}{f_2^{2^k + 1} f_8^{2^k - 2}} \pmod{2^{k+2}}.$$

Extracting the coefficients of the odd powers of q from the above clearly proves Theorem 1.10. \square

Proof of Theorem 1.11. Using Lemma 5.2, we have

(5.17)

$$\begin{aligned} \sum_{n \geq 0} \bar{a}_{2^k i + 2^{k-1} - 3}(2n+1)q^n &\equiv 2 \frac{f_2^{2^k i + 2^{k-1} - 2} f_8^2}{f_1^{2^{k+1} i + 2^k - 4} f_4} = 2 \frac{\psi(q^4)}{\varphi(-q)^{2^k i + 2^{k-1} - 2}} \pmod{2^{k+3}} \\ (5.18) \quad &\equiv 2 \frac{\varphi(-q)^{2^{k-1}(6i+7)+2} \psi(q^4)}{\varphi(-q)^{2^{k+2}(i+1)}} \equiv 2 \frac{\varphi(-q)^{2^{k-1}(6i+7)+2} \psi(q^4)}{\varphi(-q^2)^{2^{k+1}(i+1)}} \pmod{2^{k+3}}. \end{aligned}$$

Employing the 2-dissection of $\varphi(q)$ (5.1) and then extracting the terms with odd powers of q , we obtain

$$\begin{aligned} \sum_{n \geq 0} \bar{a}_{2^k i + 2^{k-1} - 3}(4n+3)q^n &\equiv -2^2 \frac{\psi(q^2)}{\varphi(-q)^{2^{k+1}(i+1)}} \sum_{r=0}^{2^{k-2}(6i+7)} 2^{2r} \binom{2^{k-1}(6i+7)+2}{2r+1} q^r \\ (5.19) \quad &\times \varphi(q^2)^{2^{k-1}(6i+7)-2r+1} \psi(q^4)^{2r+1} \pmod{2^{k+3}}. \end{aligned}$$

By the properties of binomial coefficients, for $r \geq 1, i \geq 0$, we have

$$2^{2r} \binom{2^{k-1}(6i+7)+2}{2r+1} \equiv 0 \pmod{2^{k+1}}.$$

Therefore, (5.19) reduces to

$$\sum_{n \geq 0} \bar{a}_{2^k i + 2^{k-1} - 3}(4n+3)q^n \equiv -2^2(2^{k-1}(6i+7)+2) \frac{\varphi(q^2)^{2^{k-1}(6i+7)+1} \psi(q^2) \psi(q^4)}{\varphi(-q^2)^{2^k(i+1)}} \pmod{2^{k+3}}.$$

From the above identity, we evidently arrive at (1.17) by extracting out the odd powers of q . The proof of (1.18) is similar to that of (1.17), so we skip it. \square

Proof of Theorem 1.12. The proofs of the congruences in the theorem are similar. So, we prove only the first one. Under modulo 3, we have

$$\sum_{n=0}^{\infty} \bar{a}_{9i}(n)q^n = \frac{f_4^{9i-1}}{f_1^2 f_2^{18i-3}} \equiv \frac{f_{12}^{3i}}{f_3 f_6^{6i-1}} \cdot \frac{f_1}{f_4} \pmod{3}$$

Invoking (5.6) and then extracting the terms involving q^{3n} , we obtain

$$\sum_{n=0}^{\infty} \bar{a}_{9i}(3n)q^n \equiv \frac{f_3 f_6 f_2^{2-6i} f_4^{3i-3}}{f_1} \equiv \frac{f_3 f_6 f_{12}^{i-1}}{f_6^{2i}} \cdot \frac{f_2^2}{f_1} \pmod{3}.$$

Since there is no term involving q^{3n+2} in the 3-dissection of f_2^2/f_1 , for all $n \geq 0$, the above identity gives

$$(5.20) \quad \bar{a}_{9i}(9n+6) \equiv 0 \pmod{3}.$$

Again, we have

$$\begin{aligned} \sum_{n=0}^{\infty} \bar{a}_{9i}(n)q^n &= \frac{f_4^{9i-1}}{f_1^2 f_2^{18i-3}} = \frac{1}{\varphi(-q) \varphi(-q^2)^{9i-1}} \\ &= \frac{\varphi(-\omega q) \varphi(-\omega^2 q)}{\varphi(-q) \varphi(-\omega q) \varphi(-\omega^2 q)} \left(\frac{\varphi(-\omega q^2) \varphi(-\omega^2 q^2)}{\varphi(-q^2) \varphi(-\omega q^2) \varphi(-\omega^2 q^2)} \right)^{9i-1}, \end{aligned}$$

where $\omega \neq 1$ is a cube root of unity. Due to (5.4), the above identity is equivalent to

$$\begin{aligned} \sum_{n=0}^{\infty} \bar{a}_{9i}(n)q^n &= \frac{\varphi(-q^9) \varphi(-q^{18})^{9i-1}}{\varphi(-q^3)^4 \varphi(-q^6)^{36i-4}} \left((\varphi(-q^9) - 2\omega q \Omega(-q^3)) \times (\varphi(-q^9) - 2\omega^2 q \Omega(-q^3)) \right. \\ &\quad \left. \times (\varphi(-q^{18}) - 2\omega q^2 \Omega(-q^6))^{9i-1} \times (\varphi(-q^{18}) - 2\omega^2 q^2 \Omega(-q^6))^{9i-1} \right). \end{aligned}$$

Under modulo 4, we have

$$\begin{aligned} \sum_{n=0}^{\infty} \bar{a}_{9i}(n)q^n &\equiv \frac{\varphi(-q^9) \varphi(-q^{18})^{9i-1}}{\varphi(-q^3)^4 \varphi(-q^6)^{36i-4}} \left((\varphi(-q^9)^2 + 2q \varphi(-q^9) \Omega(-q^3)) \right. \\ &\quad \left. \times (\varphi(-q^{18})^2 + 2q^2 \varphi(-q^{18}) \Omega(-q^6))^{9i-1} \right) \pmod{4} \\ &\equiv \frac{\varphi(-q^9) \varphi(-q^{18})^{9i-1}}{\varphi(-q^3)^4 \varphi(-q^6)^{36i-4}} \left((\varphi(-q^9)^2 + 2q \varphi(-q^9) \Omega(-q^3)) \right. \\ &\quad \left. \times (\varphi(-q^{18})^{18i-2} + 2(9i-1)q^2 \varphi(-q^{18})^{18i-3} \Omega(-q^6)) \right) \pmod{4}. \end{aligned}$$

Extracting the terms involving q^{3n} from the above identity, we find that

$$\sum_{n=0}^{\infty} \bar{a}_{9i}(3n)q^n \equiv \frac{\varphi(-q^3)^3 \varphi(-q^6)^{27i-3}}{\varphi(-q)^4 \varphi(-q^2)^{36i-4}} \pmod{4}.$$

Repeating the above process, we obtain

$$\sum_{n=0}^{\infty} \bar{a}_{9i}(3n)q^n \equiv \frac{\varphi(-q^9)^{12} \varphi(-q^{18})^{108i-12}}{\varphi(-q^3)^{16} \varphi(-q^6)^{144i-16}} \pmod{4},$$

from which, it is clear that for all $n \geq 0$,

$$(5.21) \quad \bar{a}_{9i}(9n+6) \equiv 0 \pmod{4}.$$

The congruences (5.20) and (5.21) together imply the first congruence of the theorem. \square

6. PROOFS OF THEOREM 1.13, 1.14, 1.15, AND 1.16

The proof of Theorem 1.13 is immediate from a result of Cotroneo *et. al* [CMSZ20]. We define

$$(6.1) \quad G(\tau) := \frac{\eta(\delta_1\tau)^{r_1}\eta(\delta_2\tau)^{r_2}\cdots\eta(\delta_u\tau)^{r_u}}{\eta(\gamma_1\tau)^{s_1}\eta(\gamma_2\tau)^{s_2}\cdots\eta(\gamma_t\tau)^{s_t}},$$

where r_i, s_i, δ_i , and γ_i are positive integers with $\delta_1, \dots, \delta_u, \gamma_1, \dots, \gamma_t$ distinct, $u, t \geq 0$. The weight of $G(\tau)$ is given by

$$\frac{1}{2} \left(\sum_{i=1}^u r_i - \sum_{i=1}^t s_i \right).$$

Also define $\mathcal{D}_G := \gcd(\delta_1, \delta_2, \dots, \delta_u)$. Then, we have the following result.

Theorem 6.1. [CMSZ20, Theorem 1.1] *Suppose $G(\tau)$ is an eta-quotient of the form (6.1) with integer weight. If p is a prime such that p^a divides \mathcal{D}_G and*

$$(6.2) \quad p^a \geq \sqrt{\frac{\sum_{i=1}^t \gamma_i s_i}{\sum_{i=1}^u \frac{r_i}{\delta_i}}},$$

then $G(\tau)$ is lacunary modulo p^j for any positive integer j . Moreover, there exists a positive constant α , depending on p and j , such that the number of integers $n \leq X$ with p^j not dividing $b(n)$ is $O\left(\frac{X}{\log^\alpha X}\right)$.

Proof of Theorem 1.13. We recall

$$\sum_{n \geq 0} \bar{a}_c(n) q^n = \frac{f_4^{c-1}}{f_1^2 f_2^{2c-3}} = \frac{\eta^{c-1}(4z)}{\eta^2(z) \eta^{2c-3}(2z)}.$$

Following the notations used in (6.1) and the paragraph succeeding it, we have

$$\delta_1 = 4, \quad r_1 = c-1, \quad \gamma_1 = 1, \quad \gamma_2 = 2, \quad s_1 = 2, \quad \text{and} \quad s_2 = 2c-3.$$

Also $\mathcal{D}_G = 4$ and the weight is $-c/2 \in \mathbb{Z}$ iff $c = 2^\alpha m$ with $\alpha \geq 1$ and $\gcd(2, m) = 1$. Next, we see that $2^2 | 4$ and

$$2^2 \geq \sqrt{\frac{2 + (4c-6)}{\frac{c-1}{4}}} = \sqrt{16} = 4.$$

Choosing $p = 2$ and $a = 2$ in Theorem 6.1, we complete the proof. \square

The proofs of Theorems 1.14 and 1.15 do not follow from Theorem 6.1. We present them below.

Proof of Theorem 1.14. Putting $c = 2^\alpha m$ in (1.2), we have

$$(6.3) \quad \sum_{n \geq 0} \bar{a}_{2^\alpha m}(n) q^n = \frac{f_4^{2^\alpha m-1}}{f_1^2 f_2^{2^\alpha m-3}}.$$

We define

$$A_{\alpha, m}(z) := \frac{\eta^3(24z)}{\eta(72z)}.$$

By the binomial theorem, for any positive integers m, k , and prime p we have

$$f_m^k \equiv f_{mp}^{k-1} \pmod{p^k}.$$

Therefore,

$$(6.4) \quad A_{\alpha,m}^{3^k}(z) := \frac{\eta^{3^{k+1}}(24z)}{\eta^{3^k}(72z)} \equiv 1 \pmod{3^{k+1}}.$$

Next we define,

$$B_{\alpha,m,k}(z) = \frac{\eta^{2^\alpha m-1}(96z)}{\eta^2(24z)\eta^{2^{\alpha+1}m-3}(48z)} A_{\alpha,m}^{3^k}(z) = \frac{\eta^{3^{k+1}-2}(24z)\eta^{2^\alpha m-1}(96z)}{\eta^{2^{\alpha+1}m-3}(48z)\eta^{3^k}(72z)}.$$

From (6.3) and (6.4), we have

$$(6.5) \quad B_{\alpha,m,k}(z) = \frac{\eta^{2^\alpha m-1}(96z)}{\eta^2(24z)\eta^{2^{\alpha+1}m-3}(48z)} \equiv \frac{f_{96}^{2^\alpha m-1}}{f_{24}^2 f_{48}^{2^{\alpha+1}m-3}} \equiv \sum_{n=0}^{\infty} \bar{a}_{2^\alpha m}(n) q^{24n} \pmod{3^{k+1}}.$$

Now, we will prove that $B_{\alpha,m,k}(z)$ is a modular form. From Theorem 2.2, we find that the level of $B_{\alpha,m,k}(z)$ is $N = 96M$, where M is the smallest positive integer such that

$$96M \left(\frac{3^{k+1}-2}{24} + \frac{3-2^{\alpha+1}m}{48} - \frac{3^k}{72} + \frac{2^\alpha m-1}{96} \right) \equiv 0 \pmod{24}$$

which implies

$$3M \cdot (4(3^k - 3^{k-2}) + 2^\alpha m - 1) \equiv 0 \pmod{24}.$$

Therefore, $M = 2^3$ and $N = 2^{83}$. The cusps of $\Gamma_0(2^{83})$ are given by fractions c/d where $d|2^{83}$ and $\gcd(c, d) = 1$. By Theorem 2.2, $B_{\alpha,m,k}(z)$ is holomorphic at a cusp c/d if and only if

$$(3^{k+1}-2) \frac{\gcd(d, 24)^2}{24} - (2^{\alpha+1}m-3) \frac{\gcd(d, 48)^2}{48} - 3^k \frac{\gcd(d, 72)^2}{72} + (2^\alpha m-1) \frac{\gcd(d, 96)^2}{96} \geq 0.$$

Equivalently, $B_{\alpha,m,k}(z)$ is holomorphic at a cusp c/d if and only if the following inequalities are satisfied for the different values of d as mentioned in the table below (note that we have simplified the inequalities as far as possible).

Value(s) of d	Inequality
1, 2, 4, 8, 24	$32 \cdot 3^k - 9 \cdot 2^\alpha m - 9 \geq 0$
3, 6, 12	$3^k - \frac{9}{32}(2^\alpha m + 1) \geq 0$
16, 48	$8 \cdot 3^k - 9 \cdot 2^\alpha m + 9 \geq 0$
32, 64, 128, 256	$3^{k-2} \geq 0$
96, 192, 384, 768	$3^k \geq 0$

Since $k \geq 2$, the last two inequalities in the above list are already satisfied. Also we note that if the third inequality is satisfied then the first two are automatically satisfied. So, we only need to verify if

$$8 \cdot 3^k - 9 \cdot 2^\alpha m + 9 \geq 0.$$

But this is true because we have $3^{k-2} \geq 2^{\alpha-3}m$ in our hypothesis. Therefore, $B_{\alpha,m,k}(z)$ is holomorphic at a cusp c/d . The weight of $B_{\alpha,m,k}(z)$ is $\ell = 3^k - 2^{\alpha-1}m$, which is a positive integer, and the associated character is given by

$$\chi(\bullet) = \left(\frac{(-1)^\ell 2^{2 \cdot 3^{k+1} - 3 \cdot 2^\alpha m + 1} 3^{2 \cdot 3^k - 2^\alpha m}}{\bullet} \right).$$

Thus, $B_{\alpha,m,k}(z) \in M_{3^k - 2^{\alpha-1}m}(\Gamma_0(2^{83}), \chi(\bullet))$.

Therefore by Theorem 2.5, the Fourier coefficients of $B_{\alpha,m,k}(z)$ are almost divisible by 3^k . Due to (6.5), this holds for $\bar{a}_{2\alpha_m}$ also. This completes the proof. \square

Proof of Theorem 1.15. Consider $t = p_1^{a_1} \cdots p_m^{a_m}$ where p_i 's are prime numbers greater than or equal to 5. Then, we have

$$(6.6) \quad \bar{a}_t(n)q^n = \frac{f_4^{t-1}}{f_1^2 f_2^{2t-3}}.$$

We define

$$G_i(z) := \frac{\eta^{p_i^{a_i}}(24z)}{\eta(24p_i^{a_i}z)}.$$

From the binomial theorem, for any integer $k \geq 1$, and prime p we have

$$(6.7) \quad G_i^{p_i^k}(z) := \frac{\eta^{p_i^{a_i+k}}(24z)}{\eta^{p_i^k}(24p_i^{a_i}z)} \equiv 1 \pmod{p_i^{k+1}}.$$

Next we define,

$$H_{i,k,t}(z) = \frac{\eta^{t-1}(96z)}{\eta^2(24z)\eta^{2t-3}(48z)} A_i^{p_i^k}(z) = \frac{\eta^{p_i^{a_i+k}-2}(24z)\eta^{t-1}(96z)}{\eta^{2t-3}(48z)\eta^{p_i^k}(24p_i^{a_i}z)}.$$

From (6.6) and (6.7), we have

$$(6.8) \quad H_{i,k,t}(z) = \frac{\eta^{t-1}(96z)}{\eta^2(24z)\eta^{2t-3}(48z)} \equiv \frac{f_{96}^{t-1}}{f_{24}^2 f_{48}^{2t-3}} \equiv \sum_{n=0}^{\infty} \bar{a}_t(n)q^{24n} \pmod{p_i^{k+1}}.$$

Now, we will prove that $H_{i,k,t}(z)$ is a modular form. From Theorem 2.2, we find that the level of $H_{i,k,t}(z)$ is $N = 96M$, where M is the smallest positive integer such that

$$96M \left(\frac{p_i^{a_i+k} - 2}{24} - \frac{2t-3}{48} + \frac{t-1}{96} - \frac{p_i^k}{24p_i^{a_i}} \right) \equiv 0 \pmod{24}$$

which implies

$$M \cdot \left(4p_i^k \left(p_i^{a_i} - \frac{1}{p_i^{a_i}} \right) - 3(t+1) \right) \equiv 0 \pmod{24}.$$

Therefore, $M = 2^3 3$ and $N = 2^8 3^2$. The cusps of $\Gamma_0(2^8 3^2)$ are given by fractions c/d where $d|2^8 3^2$ and $\gcd(c, d) = 1$. By Theorem 2.2, $H_{i,k,t}(z)$ is holomorphic at a cusp c/d if and only if

$$(p_i^{a_i+k} - 2) \frac{\gcd(d, 24)^2}{24} - (2t-3) \frac{\gcd(d, 48)^2}{48} + (t-1) \frac{\gcd(d, 96)^2}{96} - p_i^k \frac{\gcd(d, 24p_i^{a_i})^2}{24p_i^{a_i}} \geq 0.$$

Equivalently, $H_{i,k,t}(z)$ is holomorphic at a cusp c/d if and only if the following inequalities are satisfied for the different values of d as mentioned in the table below (note that we have simplified the inequalities as far as possible).

Value(s) of d	Inequality
1, 2, 3, 4, 6, 8, 9, 12, 18, 24, 36, 72	$8p_i^{k+a_i} - (1+5t) \geq 0$
16, 48, 144	$2p_i^{k+a_i} + 5(1-t) \geq 0$
32, 64, 96, 128, 192, 256, 288, 384, 576, 768, 1152, 2304	$2p_i^{k+a_i} + 8(1-t) \geq 0$

Due to our hypothesis, all the above inequalities in the list are already satisfied.

Therefore, $H_{i,k,t}(z)$ is holomorphic at a cusp c/d . The weight of $H_{i,k,t}(z)$ is $\ell = \frac{1}{2}(p_i^k(p_i^{a_i} - 1) - t)$, which is a positive integer, and the associated character is given by

$$\chi(\bullet) = \left(\frac{(-1)^\ell 2^{\ell} 3^{p_i^{a_i} - 3t + 1} 3^{p_i^{a_i} - t} (p_i^{a_i})^{-p_i^k}}{\bullet} \right).$$

Thus, $H_{i,k,t}(z) \in M_{\frac{1}{2}(p_i^k(p_i^{a_i} - 1) - t)}(\Gamma_0(2^8 3^2), \chi(\bullet))$. Thus by Theorem 2.5, the Fourier coefficients of $H_{i,k,t}(z)$ are almost divisible by $r = p_i^k$. Due to (6.8), this holds for \bar{a}_t also. This completes the proof. \square

We recall the following theorem of Ono and Taguchi [OT05] before proving Theorem 1.16.

Theorem 6.2. [OT05, Theorem 1.3 (1)] *Let n be a non-negative integer and k be a positive integer. Suppose that χ is a Dirichlet character with conductor $2^3 \cdot N$, where $N = 1, 3, 5, 15, 17$. Then there is an integer $c \geq 0$ such that for every $f(z) \in M_k(\Gamma_0(2^a N), \chi) \cap \mathbb{Z}[[q]]$ and every $t \geq 1$,*

$$f(z)|T_{p_1}|T_{p_2}|\cdots T_{p_{c+t}} \equiv 0 \pmod{3^t}$$

whenever p_1, p_2, \dots, p_{c+t} are odd primes not dividing N .

Proof of Theorem 1.16. From (6.5), we have

$$B_{\alpha,m,k} \equiv \sum_{n=0}^{\infty} \bar{a}_{2^{\alpha}m}(n) q^{24n} \pmod{3^{k+1}}$$

which yields

$$(6.9) \quad B_{\alpha,m,k} := \sum_{n=0}^{\infty} \mathcal{B}_{\alpha,m,k}(n) q^n \equiv \sum_{n=0}^{\infty} \bar{a}_{2^{\alpha}m}\left(\frac{n}{24}\right) q^n \pmod{3^{k+1}}.$$

Now, $B_{\alpha,m,k}(z) \in M_{3^k - 2^{\alpha-1}m}(\Gamma_0(2^8 3), \chi(\bullet))$, where $\chi(\bullet)$ is the associated character. From the Theorem 6.2, we get that there is an integer $i \geq 0$ such that for any $j \geq 1$,

$$B_{\alpha,m,k}(z)|T_{q_1}|T_{q_2}|\cdots T_{q_{i+j}} \equiv 0 \pmod{3^j}$$

whenever q_1, q_2, \dots, q_{i+j} are odd primes not dividing N . From the definition of Hecke operators, we have that if q_1, q_2, \dots, q_{i+j} are distinct primes and if n is coprime to $q_1 \cdots q_{i+j}$, then

$$(6.10) \quad \mathcal{B}_{\alpha,m,k}(q_1 \cdots q_{i+j} \cdot n) \equiv 0 \pmod{3^j}$$

Combining (6.9) and (6.10), we complete the proof. \square

7. CONCLUDING REMARKS

(1) We have the following conjecture.

Conjecture 7.1. *For all $n \geq 0$, we have*

$$(7.1) \quad a_{25}(31^2 n + 644) \equiv 0 \pmod{31^2},$$

$$(7.2) \quad a_{41}(47^2 n + 256) \equiv 0 \pmod{47^2},$$

$$(7.3) \quad a_{61}(67^2 n + 555) \equiv 0 \pmod{67^2}.$$

If we use a procedure similar to the proof of Theorem 1.3, it would yield us the following data.

Congruences	$(m, M, N, t, (r_\delta))$ and (r'_δ)	$P(t)$	$\lfloor \nu \rfloor$
(7.1)	$(961, 62, 62, 644, (960, -24, -31, 0))$ and $(49, 0, 0, 0)$	$\{644\}$	3813
(7.2)	$(2209, 94, 94, 256, (2208, -40, -47, 0))$ and $(81, 0, 0, 0)$	$\{256\}$	13208
(7.3)	$(4489, 134, 134, 555, (4488, -60, -67, 0))$ and $(121, 0, 0, 0)$	$\{555\}$	38091

Since the bound ν is large for the computation to take a significant amount of time in a modest computer, we leave the above as an open problem.

- (2) We also have the following conjecture, which cannot be proved similarly like the proof of Theorem 1.3.

Conjecture 7.2. *For all $n \geq 0$, we have*

$$\begin{aligned} a_{25}(31^2n + r) &\equiv 0 \pmod{31^2} \quad r \in \{41, 81, 585\}, \\ a_{37}(43^2n + r) &\equiv 0 \pmod{43^2} \quad r \in \{716, 987\}, \\ a_{53}(59^2n + 100) &\equiv 0 \pmod{59^2}, \\ a_{65}(71^2n + r) &\equiv 0 \pmod{71^2} \quad r \in \{41, 47\}, \\ a_{77}(83^2n + 157) &\equiv 0 \pmod{83^2}. \end{aligned}$$

- (3) It is desirable to have elementary proofs for Theorem 1.3 and the above conjectures.
(4) It can be noted that the congruences in Theorem 1.8 are valid for any arbitrary subscript c , not necessarily for $2i$. We restrict to the given case for ease in notations while writing the proof.
(5) Theorem 1.8 can be further strengthened to the following: for all $n \geq 0$ and $i \geq 1$, we have

$$\begin{aligned} \bar{a}_{2i}(8n + 5) &\equiv 0 \pmod{8}, \\ \bar{a}_{2i}(8n + 7) &\equiv 0 \pmod{16}. \end{aligned}$$

We leave the relevant details for the reader.

- (6) In this paper we have not explored moduli other than powers of 2 in detail. It would be natural to explore what happens if we change the modulus to another prime. To this end, we make the following conjecture, whose proof should be similar to the proof of Theorem 1.12.

Conjecture 7.3. *For all $n \geq 0$ and $i \geq 1$, we have*

$$\begin{aligned} \bar{a}_{3i+2}(9n + 2) &\equiv 0 \pmod{6}, \\ \bar{a}_{9i+5}(9n + 3) &\equiv 0 \pmod{12}, \\ \bar{a}_{9i+8}(9n + 3) &\equiv 0 \pmod{12}, \\ \bar{a}_{3i+2}(9n + 5) &\equiv 0 \pmod{6}, \\ \bar{a}_{3i+2}(9n + 8) &\equiv 0 \pmod{6}. \end{aligned}$$

REFERENCES

- [ASS25] Tewodros Amdeberhan, James A. Sellers, and Ajit Singh. Arithmetic properties for generalized cubic partitions and overpartitions modulo a prime. *Aequationes Math.*, to appear, 2025.
[BD22] Nayandeep Deka Baruah and Hirakjyoti Das. On 3^k -regular cubic partitions. *J. Korean Math. Soc.*, 59(4):685–697, 2022.

- [BS24] Pujashree Buragohain and Nipen Saikia. Some new congruences for ℓ -regular cubic partitions with odd parts overlined. *Integers*, 24:Paper No. A101, 21, 2024.
- [Cha10a] Hei-Chi Chan. Ramanujan’s cubic continued fraction and an analog of his ”most beautiful identity”. *Int. J. Number Theory*, 6(3):673–680, 2010.
- [Cha10b] Hei-Chi Chan. Ramanujan’s cubic continued fraction and Ramanujan type congruences for a certain partition function. *Int. J. Number Theory*, 6(4):819–834, 2010.
- [CL04] Sylvie Corteel and Jeremy Lovejoy. Overpartitions. *Trans. Am. Math. Soc.*, 356(4):1623–1635, 2004.
- [CMSZ20] Tessa Cotron, Anya Michaelson, Emily Stamm, and Weitao Zhu. Lacunary eta-quotients modulo powers of primes. *Ramanujan J.*, 53(2):269–284, 2020.
- [Gir17] D. S. Gireesh. Formulas for cubic partition with 3-cores. *J. Math. Anal. Appl.*, 453(1):20–31, 2017.
- [Gua25] Russelle Guadalupe. A note on congruences for generalized cubic partitions modulo primes. *Integers*, 25:paper a20, 5, 2025.
- [Hir12] Michael D. Hirschhorn. A note on overcubic partitions. *New Zealand J. Math.*, 42:229–234, 2012.
- [Hir17] Michael D. Hirschhorn. *The power of q* , volume 49 of *Developments in Mathematics*. Springer, Cham, 2017. A personal journey, With a foreword by George E. Andrews.
- [Hir20] Michael D. Hirschhorn. Cubic partitions modulo powers of 5. *Ramanujan J.*, 51(1):67–84, 2020.
- [Kim10] Byungchan Kim. The overcubic partition function mod 3. In *Ramanujan rediscovered. Proceedings of a conference on elliptic functions, partitions, and q -series in memory of K. Venkatachaliengar, Bangalore, India, June 1–5, 2009*, pages 157–163. Mysore: Ramanujan Mathematical Society, 2010.
- [Lin14] Bernard L. S. Lin. Arithmetic properties of overcubic partition pairs. *Electron. J. Combin.*, 21(3):Paper 3.35, 12, 2014.
- [Lin17] Bernard L. S. Lin. Congruences modulo 27 for cubic partition pairs. *J. Number Theory*, 171:31–42, 2017.
- [LSW17] Bernard Lishuang Lin, Erin Yiyang Shen, and Andrew Yezhou Wang. Two infinite families of congruences modulo 9 for overcubic partition pairs. *Colloq. Math.*, 147(1):115–123, 2017.
- [Mer22] Mircea Merca. A further look at cubic partitions. *Ramanujan J.*, 59(1):253–277, 2022.
- [Ono04] K. Ono. *The Web of Modularity: Arithmetic of the Coefficients of Modular Forms and q -series: Arithmetic of the Coefficients of Modular Forms and Q -series*. Regional conference series in mathematics. American Mathematical Society, 2004.
- [OT05] Ken Ono and Yuichiro Taguchi. 2-adic properties of certain modular forms and their applications to arithmetic functions. *International Journal of Number Theory*, 01(01):75–101, 2005.
- [Rad09] Silviu Radu. An algorithmic approach to Ramanujan congruences. *Ramanujan Journal*, 20(2):215–251, 2009.
- [RB20] Chiranjit Ray and Rupam Barman. Arithmetic properties of cubic and overcubic partition pairs. *Ramanujan J.*, 52(2):243–252, 2020.
- [RS11] Silviu Radu and James Sellers. Congruence Properties Modulo 5 and 7 for the pod Function. *International Journal of Number Theory*, 8:2249–2259, 2011.
- [Sel14] James A. Sellers. Elementary proofs of congruences for the cubic and overcubic partition functions. *Australas. J. Comb.*, 60:191–197, 2014.
- [Ser75a] Jean-Pierre Serre. Divisibilité de certaines fonctions arithmétiques. In *Séminaire Delange-Pisot-Poitou (16e année: 1974/75), Théorie des nombres, Fasc. 1*, pages Exp. No. 20, 28. Secrétariat Math., Paris, 1975.
- [Ser75b] Jean-Pierre Serre. Valeurs propres des opérateurs de Hecke modulo l . In *Journées Arithmétiques de Bordeaux (Conf., Univ. Bordeaux, Bordeaux, 1974)*, Astérisque, No. 24-25, pages 109–117. Soc. Math. France, Paris, 1975.
- [SNDMK24] S. Shivaprasada Nayaka, B. N. Dharmendra, and M. C. Mahesh Kumar. Divisibility properties for overcubic partition triples. *Integers*, 24:Paper No. A80, 9, 2024.
- [SS25] Manjil P. Saikia and Abhishek Sarma. Further arithmetic properties of overcubic partition triples. *Bull. Aust. Math. Soc.*, to appear, 2025.
- [Stu87] Jacob Sturm. On the congruence of modular forms. In *Number theory (New York, 1984–1985)*, volume 1240 of *Lecture Notes in Math.*, pages 275–280. Springer, Berlin, 1987.
- [Tat94] John Tate. The non-existence of certain Galois extensions of \mathbf{Q} unramified outside 2. In *Arithmetic geometry (Tempe, AZ, 1993)*, volume 174 of *Contemp. Math.*, pages 153–156. Amer. Math. Soc., Providence, RI, 1994.
- [Xio11] Xinhua Xiong. Two identities involving the cubic partition function. *Int. J. Number Theory*, 7(6):1615–1626, 2011.

- [Yao19] Olivia X. M. Yao. Infinite families of congruences modulo 5 and 7 for the cubic partition function. *Proc. Roy. Soc. Edinburgh Sect. A*, 149(5):1189–1205, 2019.

(H Das) DEPARTMENT OF MATHEMATICS, B. BOROOAH COLLEGE, GUWAHATI 781007, ASSAM, INDIA

Email address: hirak@bboroahcollege.ac.in

(S. Maity & M. P. Saikia) MATHEMATICAL AND PHYSICAL SCIENCES DIVISION, SCHOOL OF ARTS AND SCIENCES, AHMEDABAD UNIVERSITY, AHMEDABAD 380009, GUJARAT, INDIA

Email address: saikat.maity@ahduni.edu.in & manjil@saikia.in