HAUSDORFF STABILITY OF THE CUT LOCUS UNDER C²-PERTURBATIONS OF THE METRIC

ARITRA BHOWMICK

Department of Mathematics, Indian Institute of Science, Bengaluru, India

JIN-ICHI ITOH

School of Education, Sugiyama Jogakuen University, Nagoya, Japan

SACHCHIDANAND PRASAD

School of Mathematics, Jilin University, China

Faculty of Mathematics and Computer Science, Göttingen University, Germany

ABSTRACT. In this article, we prove the stability with respect to the Hausdorff metric d_H of the cut locus $\operatorname{Cu}(p, \mathfrak{g})$ of a point p in a compact Riemannian manifold (M, \mathfrak{g}) under C^2 perturbation of the metric. Specifically, given a sequence of metrics \mathfrak{g}_i on M, converging to \mathfrak{g} in the C^2 topology, and a sequence of points p_i in M, converging to p, we show that $\lim_i d_H (\operatorname{Cu}(p_i, \mathfrak{g}_i), \operatorname{Cu}(p, \mathfrak{g})) = 0$. Along the way, we also prove the continuous dependence of the cut time map on the metric.

1. INTRODUCTION

The principal objects of study in Riemannian geometry are *geodesics*, which are *locally* distance minimizing curves. If (M, \mathfrak{g}) is a connected, complete Riemannian manifold, then between any two points, say, $p, q \in M$ there exists a geodesic which is *globally* distance minimizing. We call such geodesics *minimizers*. The cut locus $\operatorname{Cu}(p, \mathfrak{g})$ of p consists of those points $q \in M$ beyond which a minimizer from p to q fails to be distance minimizing from p. Originally introduced by Poincarè [Poi05], the notion of cut locus has been extensively studied in the literature [Kob67, Buc77a, Wol79, Sak96]. A closely related set is $\operatorname{Se}(p, \mathfrak{g})$, which consists of points q admitting at least two distinct minimizers from p to q. It is well known that $\operatorname{Cu}(p, \mathfrak{g})$ is the closure of $\operatorname{Se}(p, \mathfrak{g})$, i.e., $\operatorname{Cu}(p, \mathfrak{g}) = \overline{\operatorname{Se}(p, \mathfrak{g})}$, and consequently, $\operatorname{Cu}(p, \mathfrak{g})$ is a closed set of M. The concept of cut locus can be easily generalized to a closed submanifold (or even a closed set) of M, see [BP23] for a survey of results.

E-mail addresses: aritrab@iisc.ac.in, j-itoh@sugiyama-u.ac.jp, sachchidanand.prasad1729@gmail.com. *Date*: March 26, 2025.

²⁰²⁰ Mathematics Subject Classification. Primary: 53C22, 35D40; Secondary: 35F21, 49J52.

Key words and phrases. cut locus, eikonal equation, viscosity solution, Hausdorff distance, injectivity radius.

The stability question of the cut locus has been studied in the literature as well. In [Buc77b], Buchner considered the notion of stability from the point of view of singularity theory [GG73]. Given \mathcal{G} as the space of smooth Riemannian metrics on a compact manifold M, equipped with the Whitney C^{∞} topology, Buchner proved that $\operatorname{Cu}(p, \mathfrak{g})$ is stable for \mathfrak{g} in an open dense subset $\mathcal{O} \subset \mathcal{G}$, provided dim $M \leq 6$. On the other hand, in [Alb16], Albano considered the stability in the sense of Hausdorff distance. Given the collection \mathcal{A} of bounded open domains $\Omega \subset \mathbb{R}^n$ with C^2 -smooth boundary, and the collection \mathcal{G}' of C^2 -smooth Riemannian metrics on \mathbb{R}^n , Albano considered the cut locus $\operatorname{Cu}^{\operatorname{in}}(\partial\Omega,\mathfrak{g}) \coloneqq \operatorname{Cu}(\partial\Omega,\mathfrak{g}) \cap \Omega$ of the boundary $\partial\Omega$ inside Ω , with respect to a metric $\mathfrak{g} \in \mathcal{G}'$. Using the viscosity solutions of the eikonal equations, it was proved that given a C^2 -convergent sequence $(\Omega_j,\mathfrak{g}_j) \to (\Omega,\mathfrak{g})$, the cut locus $\operatorname{Cu}(\Omega_i,\mathfrak{g}_i)$ converges to $\operatorname{Cu}(\Omega,\mathfrak{g})$ in the Hausdorff metric. We refer to [Alb16] for the notion of C^2 convergence in \mathcal{A} . In yet another direction, the topological stability of the cut locus is studied in [EGGHT21] from the point of view of metric measure theory.

In this article, we follow the approach of [Alb16], and extend it to compact Riemannian manifolds. Suppose M is a compact manifold, and let \mathcal{G} be the collection of smooth Riemannian metrics on M with the Whitney C^2 topology. Then, for any $(p, \mathfrak{g}) \in M \times \mathcal{G}$, the cut locus $\operatorname{Cu}(p, \mathfrak{g})$, being a closed subset, is compact. In other words, we have a map $\operatorname{Cut} : M \times \mathcal{G} \to \mathcal{K}$ (Equation 12), where \mathcal{K} is the collection of compact sets in M. We topologize \mathcal{K} with the Hausdorff metric. The main goal of this article is to prove the following.

Theorem (Theorem 3.1). *The map* Cut *is continuous*.

As a corollary to a lemma (Lemma 3.3) used in the proof of the above, we also show the continuous dependence of the cut time map on the C^2 perturbation of the metric (Corollary 3.4).

Conventions. All manifolds are connected, and without boundary. Boldface letters, e.g., $\mathbf{u}, \mathbf{v}, \mathbf{n}$, etc. will always denote tangent vectors, whereas letters in Fraktur font, e.g., $\mathfrak{g}, \mathfrak{h}$ etc. are reserved for Riemannian metrics. For any metric \mathfrak{g} , the induced distance, length, exponential map etc. are explicitly denoted with the subscript (or superscript) \mathfrak{g} . For a tangent vector \mathbf{v} , the unique geodesic with respect to \mathfrak{g} with initial velocity \mathbf{v} is denoted as $\gamma_{\mathbf{v}}^{\mathfrak{g}}(t)$.

Organization of the paper. In section 2, we recall some basic definitions and results on cut locus, viscosity solutions of the eikonal equation, and convergence of metrics. Then, in section 3, we prove the main theorem of this article.

2. Preliminaries

In this section we collect some preliminary results on cut locus in a Riemannian manifold from the viewpoint of viscosity solutions of the Eikonal equation. We also recall the notion of Hausdorff convergence of compact sets, and the C^2 convergence of Riemannian metrics, along with some immediate consequences.

2.1. **Cut Locus.** We recall the definition of the cut locus of a point and more generally of a submanifold, while deferring to [Sak96] for details.

Let us fix a complete Riemannian manifold (M, \mathfrak{g}) . Given a C^1 curve $\gamma : [a, b] \to M$, the length is defined as

$$L_{\mathfrak{g}}(\gamma) \coloneqq \int_{a}^{b} \left\| \dot{\gamma} \right\|_{\mathfrak{g}} dt.$$

Given $x, y \in M$, the induced distance is denoted as $d_{\mathfrak{g}}(x, y) = \inf L_{\mathfrak{g}}(\gamma)$, where the infimum is taken over the family of (piecewise) C^1 curves joining x to y.

Given a vector $\mathbf{v} \in T_p M$, let us denote the unique g-geodesic with initial velocity \mathbf{v} as $\gamma_{\mathbf{v}}^{\mathfrak{g}}$: $[0,\infty) \to M$. This lets us define the exponential map

$$\exp^{\mathfrak{g}}: T_p M \to M$$
$$\mathbf{v} \mapsto \gamma^{\mathfrak{g}}_{\mathbf{v}}(1)$$

By the Hopf-Rinow theorem, for any $q \in M$, we have a unit-speed g-geodesic, called a *minimizer*, say, $\gamma : [0, L] \to M$ such that $\gamma(0) = p, \gamma(L) = q$ and $L = L_{\mathfrak{g}}(\gamma) = d_{\mathfrak{g}}(p, q)$. The *cut locus* $\operatorname{Cu}(p, \mathfrak{g})$ consists of those points $q \in M$ beyond which a minimizer joining p to q fails to be distance minimizing from p.

Given a closed submanifold $N \subset M$, the normal bundle with respect to \mathfrak{g} is denoted as

$$TN^{\perp_{\mathfrak{g}}} \coloneqq \bigcup_{p \in N} \left\{ \mathbf{n} \in T_pM \mid \mathfrak{g}(\mathbf{n}, \mathbf{v}) = 0, \ \forall \, \mathbf{v} \in T_pN \right\}.$$

An *N*-geodesic with respect to \mathfrak{g} joining *N* to *q* is a \mathfrak{g} -geodesic joining a point in *N* to *q*, with initial velocity in $TN^{\perp_{\mathfrak{g}}}$. A unit speed *N*-geodesic $\gamma : [0, L] \to M$ is said to be an *N*-minimizer (with respect to \mathfrak{g}) joining *N* to *q* if $\gamma(0) \in N, \dot{\gamma}(0) \in TN^{\perp_{\mathfrak{g}}}$, and $\gamma(L) = q$, where $L = L_{\mathfrak{g}}(\gamma) = d_{\mathfrak{g}}(N, q)$. By the first variational principle, any unit-speed \mathfrak{g} -geodesic that minimizes the distance from *N* to a point is an *N*-minimizer. Since *N* is closed, the completeness of \mathfrak{g} implies that given any $q \in M$ there exists an *N*-minimizer joining *N* to *q*. We then define the cut locus $\operatorname{Cu}(N, \mathfrak{g})$ as the set of points $q \in M$, beyond which a minimizer joining *N* to *q* fails to be distance minimizing from *N*. The set of separating points for *N* is the set

 $Se(N, \mathfrak{g}) \coloneqq \{q \in M \mid \text{there are at least two distance minimizing geodesics joining } N \text{ to } q\}.$

It is well-known that $\operatorname{Se}(N, \mathfrak{g}) \subset \operatorname{Cu}(N, \mathfrak{g})$, and moreover

$$\operatorname{Cu}(N,\mathfrak{g}) \coloneqq \overline{\operatorname{Se}(N,\mathfrak{g})}.$$
 (1)

Given a unit vector \mathbf{n} in the normal bundle $TN^{\perp_{\mathfrak{g}}}$ of N, the *cut time* is defined as

$$\rho_{\mathfrak{g}}(N,\mathbf{n}) \coloneqq \sup\left\{t \mid d_{\mathfrak{g}}(N,\gamma_{\mathbf{n}}^{\mathfrak{g}}(t)) = t\right\}.$$
(2)

As \mathfrak{g} is complete, it follows that $\operatorname{Cu}(N,\mathfrak{g}) = \Big\{ \gamma_{\mathbf{n}}^{\mathfrak{g}}(\rho_{\mathfrak{g}}(N,\mathbf{n})) \mid \mathbf{n} \in TN^{\perp_{\mathfrak{g}}}, \|\mathbf{n}\|_{\mathfrak{g}} = 1 \Big\}.$

The *injectivity radius* of (M, \mathfrak{g}) is defined as

$$\operatorname{Inj}(\mathfrak{g}) \coloneqq \inf_{p \in M} \operatorname{Inj}(p, \mathfrak{g}), \tag{3}$$

where

$$\operatorname{Inj}(p, \mathfrak{g}) \coloneqq \inf \left\{ \rho_{\mathfrak{g}}(p, \mathbf{v}) \mid \mathbf{v} \in T_p M, \|\mathbf{v}\|_{\mathfrak{g}} = 1 \right\}$$

If M is compact, from the Whitehead convexity theorem [Whi32] it follows that $\text{Inj}(\mathfrak{g}) > 0$. For any $0 < \delta < \text{Inj}(\mathfrak{g})$, the set $S = S_{\mathfrak{g}}(p, \delta) \coloneqq \{q \mid d_{\mathfrak{g}}(p, q) = \delta\}$ is diffeomorphic to a sphere of dimension $\dim M - 1$. It is easy to see that

$$\operatorname{Cu}(S) = \{p\} \cup \operatorname{Cu}(p, \mathfrak{g}). \tag{4}$$

The *diameter* of (M, \mathfrak{g}) is defined as the supremum

$$\operatorname{Diam}(\mathfrak{g}) \coloneqq \sup_{p,q \in M} d_{\mathfrak{g}}(p,q), \tag{5}$$

which is a maximum whenever M is compact.

2.2. Convergence of Compact Sets. Let (M, d) be a metric space. For arbitrary subsets $A, B \subset M$, the Hausdorff distance between them is defined as

$$d_H(A,B) \coloneqq \max\left\{\sup_{a \in A} d(a,B), \sup_{b \in B} d(A,b)\right\},\tag{6}$$

where $d(a, B) := \inf_{b \in B} d(a, b)$, and $d(A, b) := \inf_{a \in A} d(a, b)$. We shall need the following characterization for convergence with respect to the Hausdorff distance.

Lemma 2.1. Suppose M is compact, and $X_j, X \subset M$ are closed (hence compact) subsets. Then, $\lim_i d_H(X_i, X) = 0$ if and only if the following holds:

- (1) for any $x \in X$, there exists a sequence $x_i \in X_i$ with $\lim_i x_i = x$, and
- (2) for any convergent sequence $x_{i_j} \in X_{i_j}$, we have $\lim_j x_{i_j} \in X$

Proof. Suppose, $\lim_{j \to 0} d_H(X_j, X) = 0$. Let us prove (1) first. Pick $x \in X$. We have,

$$0 \le d(X_j, x) \le \sup_{b \in X} d(X_j, b) \le d_H(X_j, X)$$

Then, for each $n \ge 1$ we can get a strictly increasing sequence N_n such that

$$d(X_j, x) < \frac{1}{n}, \quad \forall j \ge N_n.$$

Now, for each $N_n \leq j < N_{n+1}$, we have $d(X_j, x) = \inf_{a \in X_j} d(a, x) < \frac{1}{n}$, and hence, we can pick $x_j \in X_j$ so that $d(x_j, x) < \frac{1}{n}$. For $1 \leq j < N_1$, pick arbitrary $x_j \in X_j$. Then, we have $\lim_j x_j = x$, as required, showing (1). Next, we prove (2). Assume, $x_{i_j} \in X_{i_j}$ is a sequence converging to $y \coloneqq \lim_j x_{i_j}$. Now, for $\epsilon > 0$, there exists $N \geq 1$ such that

$$d(x_{i_j}, y) < \frac{\epsilon}{2}$$
, and $d_H(X_{i_j}, X) < \frac{\epsilon}{2}$, $\forall j \ge N$.

For $j \geq N$, we then have

$$\sup_{a \in X_{i_j}} d(a, X) \le d_H(X_{i_j}, X) < \frac{\epsilon}{2} \Rightarrow d(x_{i_j}, X) < \frac{\epsilon}{2} \Rightarrow \exists y_j \in X, \text{ such that } d(x_{i_j}, y_j) < \frac{\epsilon}{2}.$$

Thus, for $j \geq N$, we get

$$d(y_j, y) \le d(y_j, x_{i_j}) + d(x_{i_j}, y) < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon.$$

In other words, $y \in \overline{X} = X$, as X is assumed to be closed. This proves (2)

Let us now assume (1) and (2). Fix some $\epsilon > 0$. By compactness of X, we may assume that there are $y_1, \ldots, y_K \in X$ such that

$$X = \bigcup_{i=1}^{K} B\left(y_i, \frac{\epsilon}{2}\right) = \bigcup_{i=1}^{K} \left\{ x \mid d(y_i, x) < \frac{\epsilon}{2} \right\}.$$

Then, from (1), there exists $x_j^i \in X_j$ such that $\lim_j x_j^i = y_i$ for each $1 \le i \le K$. Consequently, we have $N \ge 1$ such that

$$d(X_j, y_i) < \frac{\epsilon}{2}, \quad \forall j \ge N, \ \forall 1 \le i \le K.$$

Now, for any $y \in X$, we have some $1 \le i_0 \le K$ such that $d(y_{i_0}, y) < \frac{\epsilon}{2}$, and hence

$$d(X_j, y) \le d(X_j, y_{i_0}) + d(y_{i_0}, y) < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon, \quad \forall j \ge N.$$

In other words, $\sup_{y\in X} d(X_j,y) \leq \epsilon$ for $j\geq N.$ As $\epsilon>0$ is arbitrary, we get

$$\lim_{j} \sup_{b \in X} d(X_j, b) = 0.$$

Thus, we now need to show that $\lim_{j \to a \in X_j} d(a, X) = 0$. Let us assume the contrary. Then, for some $\epsilon > 0$, there exists an increasing sequence i_j such that

$$\sup_{a \in X_{i_j}} d(a, X) > \epsilon$$

We can pick $x_{i_j} \in X_{i_j}$ such that $d(x_{i_j}, X) > \epsilon$. Now, since M is compact, a subsequence of x_{i_j} , say, $x_{i_{j_k}}$ converges to some y. By (2), we then have $y \in X$, which contradicts $d(x_{i_{j_k}}, y) \ge d(x_{i_{j_k}}, X) > \epsilon$. Hence, $\lim_j \sup_{a \in X_j} d(a, X) = 0$ as well. But then we have $\lim_j d_H(X_j, X) = 0$, concluding the proof.

2.3. Convergence of Riemannian Metrics. Let us briefly recall the Whitney C^r -topology on function spaces, we refer to [GG73, Mic80] for details.

Given two topological spaces X, Y, on the collection of continuous maps $\mathcal{C}(X, Y)$ one can consider two natural topologies. The *strong* topology has the basic open sets $\{g \in \mathcal{C}(X, Y) \mid g(X) \subset U\}$ for $U \subset X$ open. On the other hand, the *weak* topology is the usual compact-open topology, where the basic open sets are of the form $\{g \in \mathcal{C}(X, Y) \mid g(K) \subset U\}$ for $K \subset X$ compact and $U \subset Y$ open. Note that the strong topology is not even Hausdorff, as it fails to separate any two surjective maps.

Now, suppose (Y, d) is a metric space. Then, the weak topology on $\mathcal{C}(X, Y)$ is also known as the compact convergence topology. In particular, if $f_n \to f$ in the weak topology, then $f_n|_K \Rightarrow f|_K$ uniformly for all compact set $K \subset X$. On the other hand, further assuming that X is paracompact, a neighborhood basis of some $f \in \mathcal{C}(X, Y)$ in the strong topology is given as

$$\{g \in \mathcal{C}(X, Y) \mid d(f(x), f(y)) < \epsilon(x), \ x \in X\},\$$

where $\epsilon : X \to (0, \infty)$ is a continuous map. Consequently, $f_n \to f$ in the strong topology implies that there exists a compact set $K \subset X$, such that $f_n|_{X \setminus K} = f|_{X \setminus K}$ for n large, and $f_n|_K \Rightarrow f|_K$ uniformly.

Now, assume that M, N are smooth manifolds. Then, the collection $J^r(M, N)$ of r-jets of maps $M \to N$ is itself a manifold. We have a natural injection

$$j^r: C^{\infty}(M, N) \hookrightarrow \mathcal{C}(M, J^r(M, N))$$

The induced topology on $C^{\infty}(M, N)$ given by the strong (resp. weak) topology on $C(M, J^{r}(M, N))$ is called the strong (resp. weak) Whitney C^{r} topology. If M is compact, the two topologies coincide.

Given a smooth vector bundle $E \to M$, one can construct the *r*-jet space of sections of *E*, and denote it as $E^{(r)} \subset J^r(M, E)$. Then, we have the injection $j^r : \Gamma(E) \hookrightarrow E^{(r)}$, which induces the strong (or weak) Whitney C^r topology on the space ΓE of smooth sections of *E*. Again, if *M* is compact the two topologies are the same, and it is then a metric topology.

Let us now consider $E = T^*M \odot T^*M$, the symmetric tensor product of the cotangent bundles on a smooth manifold M of dimension n. One can then identify the collection of smooth Riemannian metrics on M as a subset $\mathcal{G} \subset \Gamma E$. Let us explain a C^r -neighborhood of a metric $\mathfrak{g} \in \mathcal{G}$ in the weak topology. Fix locally finite, closure compact, coordinate charts $\{U_\alpha, \phi_\alpha\}$ covering M, and write \mathfrak{g} as $\mathfrak{g}^{\alpha}_{ij} dx^i_{\alpha} \odot dx^j_{\alpha}$ on each U_{α} . Set $\tilde{\mathfrak{g}}^{\alpha}_{ij} := \mathfrak{g}^{\alpha}_{ij} \circ \phi^{-1}_{\alpha} : \phi_{\alpha}(U_{\alpha}) \subset \mathbb{R}^n \to \mathbb{R}$. Then, for a multi-index β with order $|\beta| \leq r$, we have the map $\partial_{\beta} \tilde{\mathfrak{g}}_{ij}^{\alpha} : \mathbb{R}^n \to \mathbb{R}$. Fix arbitrary positive numbers $\epsilon = \{\epsilon_{\alpha}\}$. Then, another metric $\mathfrak{h} \in \mathcal{G}$ is ϵ close to \mathfrak{g} in the C^r topology if for each α and for each i, j we have

$$\left|\partial_{\beta}\tilde{\mathfrak{g}}_{ij}^{\alpha}(x) - \partial_{\beta}\tilde{\mathfrak{h}}_{ij}^{\alpha}(x)\right| < \epsilon_{\alpha}, \quad x \in \phi_{\alpha}(U_{\alpha}), \ |\beta| \le r.$$

In particular, $\mathfrak{g}_i \to \mathfrak{g}$ in the weak C^r topology can be understood as the uniform C^r convergence of the local coefficients (with some fixed coordinate system).

We observe that the metric convergence gives rise to the convergence in the norm and the length of a smooth curve.

Lemma 2.2. Let \mathfrak{g}_i be a sequence of Riemannian metric on M, converging in the weak C^0 -topology to the metric \mathfrak{g} . Then, the induced norm $\|\cdot\|_{\mathfrak{g}_i}$ converges to $\|\cdot\|_{\mathfrak{g}}$ in the weak C^0 -sense. Consequently, for any C^1 curve $\gamma : [0,1] \to M$, the length $L_{\mathfrak{g}_i}(\gamma) \coloneqq \int_0^1 \|\dot{\gamma}\|_{\mathfrak{g}_i} dt$ converges to $L_{\mathfrak{g}}(\gamma) \coloneqq \int_0^1 \|\dot{\gamma}\|_{\mathfrak{g}} dt$.

Proof. Let $U \subset M$ be a coordinate chart on M with compact closure, denote the coordinate functions as x^1, \ldots, x^n . Fix some compact set $\mathcal{K} \subset TU$. For any $\mathbf{v} = \sum_i v^i \partial_{x^i} \in TU$, we have a continuous function $f(\mathbf{v}) = \sum_{i,j} |v_i v_j|$, which is bounded from above by some C > 0 on \mathcal{K} . Fix $\epsilon > 0$. Let us write $\mathfrak{g}_i \coloneqq \mathfrak{g}_i^{jk} dx^j \odot dx^k, \mathfrak{g} \coloneqq \mathfrak{g}_i^{jk} dx^j \odot dx^k$ on U. Since $\mathfrak{g}_i \to \mathfrak{g}$, we then have some $N \ge 1$ such that

$$\sum_{ij} \left| \mathfrak{g}_i^{jk}(x) - \mathfrak{g}^{jk}(x) \right| < \frac{\epsilon}{C}, \quad 1 \le j, k \le n, \ x \in U, \ i \ge N.$$

Then, for any $\mathbf{v} \in \mathcal{K}$, we have

$$\left|\mathfrak{g}_{i}(\mathbf{v},\mathbf{v}) - \mathfrak{g}(\mathbf{v},\mathbf{v})\right| \leq \sum_{i,j} \left|\mathfrak{g}_{i}^{jk} - \mathfrak{g}^{jk}\right| \left|v_{i}v_{j}\right| \leq C \sum_{i,j} \left|\mathfrak{g}_{i}^{jk} - \mathfrak{g}^{jk}\right| < \epsilon,$$

for $i \geq N$. In other words, $\|\cdot\|_{\mathfrak{g}_i}^2 \to \|\cdot\|_{\mathfrak{g}}^2$ uniformly on $\mathcal{K} \subset TU$. For an arbitrary compact set $\mathcal{K} \subset TM$, we can consider finitely many charts, covering the compact set $\pi(\mathcal{K}) \subset M$, where $\pi: TM \to M$ is the projection. Finally, as $x \mapsto \sqrt{x}$ is uniformly continuous on $[0, \infty)$, the proof follows immediately.

Remark 2.3. In [Ehr74, Sak83], the authors have used the notation $\mathfrak{g} \leq \mathfrak{h}$ for two given metrics on M to mean that $\mathfrak{g}(\mathbf{v}, \mathbf{v}) \leq \mathfrak{h}(\mathbf{v}, \mathbf{v})$ for all $\mathbf{v} \in TM$. This leads to a weaker formulation of C^0 -convergence when M is compact. In particular, if $\mathfrak{g}_i \to \mathfrak{g}$ in the C^0 -sense, then one can show that

$$(1 - \epsilon_i)\mathfrak{g} \le \mathfrak{g}_i \le (1 + \epsilon_i)\mathfrak{g},\tag{7}$$

for some real numbers $\epsilon_i \to 0$. Note that if we prove the inequality for some $\mathbf{v} \neq 0$, we can deduce it for all $\lambda \mathbf{v}$ with $\lambda \geq 0$. Since the unit bundle $\{\mathbf{v} \mid \|\mathbf{v}\|_{\mathfrak{g}} = 1\}$ is compact, using Lemma 2.2, inductively we have $N_1 < N_2 < \ldots$ such that

$$|\mathfrak{g}_i(\mathbf{v},\mathbf{v}) - \mathfrak{g}(\mathbf{v},\mathbf{v})| < \frac{1}{k}, \quad \forall \mathbf{v} \in TM, \ \|\mathbf{v}\|_{\mathfrak{g}} = 1, \ \forall i \ge N_k.$$

Set

$$\epsilon_i \coloneqq \begin{cases} 2 \max_{\|\mathbf{v}\|_{\mathfrak{g}}=1} |\mathfrak{g}_i(\mathbf{v}, \mathbf{v}) - \mathfrak{g}(\mathbf{v}, \mathbf{v})|, & 1 \le i < N_1, \\ \frac{1}{j}, & N_j \le i < N_{j+1}, \end{cases}$$

so that $\epsilon_i \to 0$. Then, for any v, with $\mathfrak{g}(\mathbf{v}, \mathbf{v}) = 1$, we have

$$\begin{aligned} |\mathfrak{g}_i(\mathbf{v},\mathbf{v}) - \mathfrak{g}(\mathbf{v},\mathbf{v})| &< \epsilon_i \Rightarrow 1 - \epsilon_i < \mathfrak{g}_i(\mathbf{v},\mathbf{v}) < 1 + \epsilon_i \\ \Rightarrow (1 - \epsilon_i)\mathfrak{g}(\mathbf{v},\mathbf{v}) < \mathfrak{g}_i(\mathbf{v},\mathbf{v}) < (1 + \epsilon_i)\mathfrak{g}(\mathbf{v},\mathbf{v}) \end{aligned}$$

Thus, Equation 7 follows.

As an application, we get the convergence of the induced distance on a compact manifold.

Lemma 2.4. Let \mathfrak{g}_i be sequence of Riemannian metrics on a compact manifold M, converging to the metric \mathfrak{g} in the C^0 -sense. Suppose, $p_i, q_i \in M$ are sequences of points, with $p = \lim p_i, q = \lim q_i$. Then, $\lim d_{\mathfrak{g}_i}(p_i, q_i) = d_{\mathfrak{g}}(p, q)$.

Proof. Suppose γ is a g-minimizer joining p to q so that $d_{\mathfrak{g}}(p,q) = L_{\mathfrak{g}}(\gamma)$. Fix some $\epsilon > 0$. By Lemma 2.2, we have $\lim L_{\mathfrak{g}_i}(\gamma) = L_{\mathfrak{g}}(\gamma)$. Thus, we have some $N_{\epsilon}^1 \ge 1$ so that

$$L_{\mathfrak{g}_i}(\gamma) \le L_{\mathfrak{g}}(\gamma) + \epsilon = d_{\mathfrak{g}}(p,q) + \epsilon, \quad i \ge N_{\epsilon}^1.$$

We also have some $N_{\epsilon}^2 \geq 1$ so that

$$d_{\mathfrak{g}}(p_i, p) < \epsilon, \quad d_{\mathfrak{g}}(q_i, q) < \epsilon, \quad i \ge N_{\epsilon}^2.$$

Lastly, by Lemma 2.2, we have some N_{ϵ}^3 so that

$$\|\mathbf{v}\|_{\mathfrak{a}_i} \le (1+\epsilon) \|\mathbf{v}\|, \quad \mathbf{v} \in TM, \quad i \ge N_{\epsilon}^3.$$

Set $N_{\epsilon} \coloneqq \max\{N_{\epsilon}^1, N_{\epsilon}^2, N_{\epsilon}^3\}$. Let $\eta_i : [0, d_{\mathfrak{g}}(p, p_i)] \to M$ be a \mathfrak{g} -minimizer joining p_i to p_i and $\zeta_i : [0, d_{\mathfrak{g}}(q, q_i)] \to M$ be a \mathfrak{g} -minimizer joining q to q_i . Note that, for $i \ge N_{\epsilon}$ we have,

$$L_{\mathfrak{g}_i}(\eta_i) = \int_0^{d_\mathfrak{g}(p,p_i)} \|\dot{\eta}_i\|_{\mathfrak{g}_i} \, dt \le (1+\epsilon) \int_0^{d_\mathfrak{g}(p,p_i)} \underbrace{\|\dot{\eta}_i\|_\mathfrak{g}}_1 \, dt = (1+\epsilon) d_\mathfrak{g}(p,p_i) < (1+\epsilon)\epsilon,$$

and similarly, $L_{\mathfrak{g}_i}(\zeta_i) < (1+\epsilon)\epsilon$ as well. Then, for all $i \ge N_\epsilon$ we get $d_{\mathfrak{g}_i}(p_i, q_i) \le L_{\mathfrak{g}_i}(\eta_i) + L_{\mathfrak{g}_i}(\gamma) + L_{\mathfrak{g}_i}(\zeta_i) < (1+\epsilon)\epsilon + d_{\mathfrak{g}}(p,q) + \epsilon + (1+\epsilon)\epsilon = d_{\mathfrak{g}}(p,q) + (3+2\epsilon)\epsilon$ which implies, $\limsup d_{\mathfrak{g}_i}(p_i, q_i) \le d_{\mathfrak{g}}(p,q) + (3+2\epsilon)\epsilon$. As $\epsilon > 0$ is arbitrary, we get

$$\limsup d_{\mathfrak{g}_i}(p_i, q_i) \le d_{\mathfrak{g}}(p, q). \tag{*1}$$

Next, let us consider \mathfrak{g}_i -minimizers $\gamma_i : [0, L_i] \to M$ joining p_i to q_i , where $L_i \coloneqq d_{\mathfrak{g}_i}(p_i, q_i)$. Since $\limsup L_i \leq d_{\mathfrak{g}}(p, q) < \infty$, we have $L \coloneqq \sup L_i < \infty$. Extend γ_i by constant to [0, L], i.e., $\gamma_i(t) = q_i$ for $L_i \leq t \leq L$. Clearly $\{\gamma_i\}$ is a uniformly bounded family, since M is compact. As $\|\cdot\|_{\mathfrak{g}_i} \to \|\cdot\|_{\mathfrak{g}}$ uniformly, we have some $N_0 \geq 1$ so that

$$\frac{1}{2} \|\mathbf{v}\|_{\mathfrak{g}} \le \|\mathbf{v}\|_{\mathfrak{g}_i} \le \frac{3}{2} \|\mathbf{v}\|_{\mathfrak{g}}, \quad \mathbf{v} \in TM$$

holds for $i \ge N_0$. Now, for $0 \le t_1 \le t_2 \le L_i$ we have,

$$d_{\mathfrak{g}}(\gamma_{i}(t_{1}),\gamma_{i}(t_{2})) \leq \int_{t_{1}}^{t_{2}} \|\dot{\gamma}(t)\|_{\mathfrak{g}} dt \leq 2 \int_{t_{1}}^{t_{2}} \underbrace{\|\dot{\gamma}(t)\|_{\mathfrak{g}_{i}}}_{1} dt = 2(t_{2}-t_{1}).$$

Since γ_i is constant for $t \geq L_i$, we see that

$$d_{\mathfrak{g}}(\gamma_i(t_1))d_{\mathfrak{g}}(\gamma_i(t_1),\gamma_i(t_2)) \le 2|t_2-t_1|, \quad 0 \le t_1 \le t_2 \le L, \ \forall i \ge N.$$

Thus, the family $\{\gamma_i\}_{i\geq N_0}$ is uniformly Lipschitz. By the Arzelà-Ascoli theorem, passing to a subsequence, we see that $\gamma_i \to \eta$ uniformly, for some curve $\eta : [0, L] \to M$. As the convergence is uniform, and the family $\{\gamma_i\}_{i\geq N_0}$ is uniformly Lipschitz, it follows that η is a Lipschitz (and hence, rectifiable) curve. In particular, the g-length of η is defined as

$$L_{\mathfrak{g}}(\eta) \coloneqq \sup_{0=t_0 < \cdots < t_m = L} \sum_{i=1}^m d_{\mathfrak{g}}(\eta(t_{i-1}), \eta(t_i)).$$

Clearly, $\eta(0) = \lim \gamma_i(0) = \lim p_i = p$ and $\eta(L) = \lim \gamma_i(L) = \lim q_i = q$, i.e., η is a Lipschitz curve joining p to q. We claim that $L_{\mathfrak{g}}(\eta) \leq \liminf L_i = \liminf L_{\mathfrak{g}_i}(\gamma_i)$. Fix some $\epsilon > 0$. Then, we have some partition of [0, L] so that $L_{\mathfrak{g}}(\eta) - \epsilon < \sum_{j=1}^m d_{\mathfrak{g}}(\eta(t_{j-1}), \eta(t_j))$. As $\gamma_i \to \eta$ pointwise, we have some $N_{\epsilon} \geq 1$ such that

$$d_{\mathfrak{g}}(\eta(t_j), \gamma_i(t_j)) < \frac{\epsilon}{2m}, \quad 0 \le j \le m, \ i \ge N_{\epsilon}.$$

We also assume, from Lemma 2.2 that

$$\|\mathbf{v}\|_{\mathfrak{g}} \le (1+\epsilon) \|\mathbf{v}\|_{\mathfrak{g}_i}, \quad \forall \mathbf{v} \in TM, \ i \ge N_{\epsilon}.$$

In particular, for $i \geq N_{\epsilon}$ we have

$$\sum_{j=1}^{m} d_{\mathfrak{g}}(\gamma_{i}(t_{j-1}), \gamma_{i}(t_{j})) \leq \sum_{j=1}^{m} \int_{t_{j-1}}^{t_{j}} \|\dot{\gamma}_{i}(t)\|_{\mathfrak{g}} dt = \int_{0}^{L} \|\dot{\gamma}_{i}(t)\|_{\mathfrak{g}} dt$$
$$\leq (1+\epsilon) \int_{0}^{L} \|\dot{\gamma}_{i}(t)\|_{\mathfrak{g}_{i}} dt = (1+\epsilon) d_{\mathfrak{g}_{i}}(p_{i}, q_{i})$$

Then, for all $i \geq N_{\epsilon}$, by the triangle inequality of $d_{\mathfrak{g}}$ we get

$$L_{\mathfrak{g}}(\eta) - \epsilon < \sum_{j=1}^{m} d_{\mathfrak{g}}\left(\gamma_i(t_{j-1}), \gamma_i(t_j)\right) + \epsilon \le (1+\epsilon)d_{\mathfrak{g}_i}(p_i, q_i) + \epsilon,$$

which implies $L_{\mathfrak{g}}(\eta) \leq \liminf(1+\epsilon)d_{\mathfrak{g}_i}(p_i,q_i) + 2\epsilon$. Since $\epsilon > 0$ is arbitrary, we have

$$d_{\mathfrak{g}}(p,q) \le L_{\mathfrak{g}}(\eta) \le \liminf d_{\mathfrak{g}_i}(p_i,q_i). \tag{*2}$$

But then from $(*_1)$ and $(*_2)$, we get $d_{\mathfrak{g}}(p,q) = \lim_i d_{\mathfrak{g}_i}(p_i,q_i)$, concluding the proof.

We shall need the following.

Theorem 2.5. [Ehr74, Sak83] Suppose \mathfrak{g}_i is a sequence of Riemannian metric on a compact manifold, converging to the metric \mathfrak{g} in the C^0 -topology. Then, $\lim_i \operatorname{Diam}(\mathfrak{g}_i) = \operatorname{Diam}(\mathfrak{g})$. Furthermore, if $\mathfrak{g}_i \to \mathfrak{g}$ in C^2 -topology, then $\lim_i \operatorname{Inj}(\mathfrak{g}_i) = \operatorname{Inj}(\mathfrak{g})$.

We then have a stronger version of Lemma 2.4.

Lemma 2.6. Suppose, \mathfrak{g}_i is a sequence of Riemannian metric on a compact manifold M, converging to the metric \mathfrak{g} in C^0 topology, and $p_i \in M$ is a sequence of points converging to p. Set, $u_i := d_{\mathfrak{g}_i}(p_i, _)$ and $u := d_{\mathfrak{g}}(p, _)$. Then, $u_i \Rightarrow u$ uniformly.

Proof. It is clear from Lemma 2.4 that $u_i \rightarrow u$ pointwise. By Lemma 2.2, we have some $N \ge 1$ such that

$$\|\mathbf{v}\|_{\mathfrak{g}_i} \le 2 \|\mathbf{v}\|_{\mathfrak{g}}, \quad \mathbf{v} \in TM, \ i \ge N.$$

Now, for $x, y \in M$, consider a g-minimizer $\gamma : [0, L] \to M$ joining x to y, where $L = d_{\mathfrak{g}}(x, y)$. Then, for $i \geq N$ we have

$$|u_i(x) - u_i(y)| \le d_{\mathfrak{g}_i}(x, y) \le L_{\mathfrak{g}_i}(\gamma) = \int_0^L \|\dot{\gamma}\|_{\mathfrak{g}_i} \le 2\int_0^L \|\dot{\gamma}\|_{\mathfrak{g}} = 2L_{\mathfrak{g}}(\gamma) = 2d_{\mathfrak{g}}(x, y).$$

Thus, the family $\{u_i\}_{i \ge N}$ is uniformly Lipschitz, with respect to the fixed metric $d_{\mathfrak{g}}$. Since $\lim \operatorname{Diam}(\mathfrak{g}_i) = \operatorname{Diam}(\mathfrak{g}) < \infty$ by Theorem 2.5, by taking N larger, we may assume that $\operatorname{Diam}(\mathfrak{g}_i) < K := \operatorname{Diam}(\mathfrak{g}) + 1$ for all $i \ge N$. Consequently, the family $\{u_i\}_{i \ge N}$ is uniformly bounded as well, as $u_i(x) = d_{\mathfrak{g}_i}(p_i, x) \le \operatorname{Diam}(\mathfrak{g}_i) < K$.

Now, if possible, suppose $u_i \not\Rightarrow u$ uniformly. Then, there exists an $\epsilon > 0$ and a subfamily, $\{u_{i_j}\}$ of $\{u_i\}_{i \ge N}$ such that $\sup_{x \in M} |u_{i_j}(x) - u(x)| \ge \epsilon$. But the family $\{u_{i_j}\}$ is both uniformly Lipschitz and uniformly bounded. Hence, by the Arzelà-Ascoli theorem, we have a uniformly convergent subsequence $u_{i_{j_k}}$, which necessarily converges to u. This is a contradiction. Hence, $u_i \Rightarrow u$ uniformly. \Box

2.4. Viscosity Solution. In this section, we recall the notion of viscosity solutions of the eikonal equation, we refer to [CIL92] for details. Let (M, \mathfrak{g}) be a Riemannian manifold. Given an open set $\Omega \subset M$, consider the eikonal equation on Ω

$$\|\nabla^{\mathfrak{g}}u\|_{\mathfrak{g}} = 1$$
 (or equivalently, $\|du\|_{\mathfrak{g}} = 1$), (8)

where $\nabla^{\mathfrak{g}}$ is the gradient operator associated to \mathfrak{g} .

Definition 2.7. A continuous map $u : \Omega \to \mathbb{R}$ is said to be a *viscosity* solution of Equation 8 at $x_0 \in \Omega$ if it satisfies the following.

- u is a viscosity sub-solution at x_0 : for any C^1 function φ near x_0 , such that $u \varphi$ attains a local maximum at x_0 (or equivalently, $u \varphi \leq 0$ and $u(x_0) = \varphi(x_0)$), we have $\|\nabla^{\mathfrak{g}}\varphi\|_{\mathfrak{g}} \leq 1$ at x_0 .
- u is a viscosity super-solution at x₀: for any C¹ function ψ near x₀, such that u ψ attains a local minimum at x₀ (or equivalently, u ψ ≥ 0 and u(x₀) = ψ(x₀)), we have ||∇^gψ||_g ≥ 1 at x₀.

Remark 2.8. Note that without loss of generality, we can ask for a *strict* local maximum at x_0 in the definition above. Indeed, suppose $u - \phi$ attains a local maximum at x_0 for some C^1 map ϕ defined near x_0 . Define $\varphi(x) \coloneqq \phi(x) + (d_{\mathfrak{g}}(x_0, x))^2$. Clearly, φ is C^1 in a small neighborhood of U, and furthermore, $d_{x_0}\varphi = d_{x_0}\phi$. Now, for $x \neq x_0$ we have,

$$(u - \varphi)(x) = (u - \phi)(x) - d_{\mathfrak{g}}(x_0, x)^2 \le u(x_0) - \underbrace{d_{\mathfrak{g}}(x_0, x)^2}_{\ge 0} < u(x_0),$$

and $(u - \varphi)(x_0) = (u - \phi)(x_0)$. Thus, $u - \varphi$ attains a strict local maximum at x_0 , while having $\|\nabla^{\mathfrak{g}}\phi\| = \|\nabla^{\mathfrak{g}}\varphi\|$ at x_0 . Similarly, if $u - \psi$ attains a local minimum at x_0 , it follows that $u - \Psi$ attains a *strict* local minimum at x_0 , with $d_{x_0}\psi = d_{x_0}\Psi$, where $\Psi(x) \coloneqq \psi(x) - (d_{\mathfrak{g}}(x_0, x))^2$.

Next, let us consider the Dirichlet problem

$$\|du\|_{\mathfrak{a}} = 1, \quad u|_{\partial\Omega} = 0. \tag{9}$$

We have the following existence and uniqueness result for the viscosity solution of the above.

Theorem 2.9. [MM03, Theorem 3.1] Suppose Ω is bounded (which is the case since M is compact). Then, the distance function $u := d_{\mathfrak{g}}(\partial\Omega, _)$ on Ω is the unique viscosity solution (among the positive functions) for the Dirichlet problem Equation 9. The only other solution being -u.

Let us denote the sets

$$\operatorname{Cu}^{\operatorname{in}}(\Omega,\mathfrak{g}) \coloneqq \operatorname{Cu}(\partial\Omega,\mathfrak{g}) \cap \Omega, \quad \operatorname{Se}^{\operatorname{in}}(\Omega,\mathfrak{g}) \coloneqq \operatorname{Se}(\partial\Omega,\mathfrak{g}) \cap \Omega.$$
(10)

Then, the distance function $u \coloneqq d_{\mathfrak{g}}(\partial\Omega, _)$ is not C^1 on points of $\operatorname{Se}^{\operatorname{in}}(\Omega, \mathfrak{g})$, and it is smooth on $\Omega \setminus \operatorname{Cu}^{\operatorname{in}}(\Omega, \mathfrak{g})$. It follows from Equation 1, that the C^1 singular support of u inside Ω is precisely the set $\operatorname{Cu}^{\operatorname{in}}(\Omega, \mathfrak{g})$. We shall need the following characterization. We provide a proof, though it may already be known to experts, while referring to [MM03] for the terminology used within.

Proposition 2.10. $x \in \text{Se}^{\text{in}}(\Omega, \mathfrak{g})$ if and only if there exists C^1 function φ near x such that $u - \varphi$ attains a local maximum at x and $\|d\varphi\|_{\mathfrak{g}} < 1$ at x (or equivalently, $\|\nabla_x^{\mathfrak{g}}\varphi\|_{\mathfrak{g}} < 1$), where $u := d_{\mathfrak{g}}(\partial\Omega, \underline{\})$.

Proof. Consider the Hamiltonian $H : TM \to \mathbb{R}$ given by $H(x, \mathbf{v}) = \|\mathbf{v}\|_{\mathfrak{g}} - 1 = \mathfrak{g}_x(\mathbf{v}, \mathbf{v}) - 1$. Observe that for any $\mathbf{u} \neq \mathbf{v} \in T_x M$ with $\|\mathbf{u}\|_{\mathfrak{g}} = 1 = \|\mathbf{v}\|_{\mathfrak{g}}$ we have the *strict* inequality

$$H(x, \lambda \mathbf{u} + (1 - \lambda)\mathbf{v}) < \lambda H(x, \mathbf{u}) + (1 - \lambda)H(x, \mathbf{v}), \quad \lambda \in (0, 1).$$

Indeed, it follows from the triangle inequality, which is strict since \mathbf{u} and \mathbf{v} being distinct unit vectors, cannot be collinear. Next, consider the set of *supergradients*

$$D^+u(x) \coloneqq \left\{ \nabla^{\mathfrak{g}}_x \varphi \mid \varphi \text{ is } C^1 \text{ near } x, \text{ and } u - \varphi \text{ attains a local maximum at } x. \right\} \subset T_x M.$$

If u is differentiable at x, then $D^+u(x) = \{\nabla_x^{\mathfrak{g}}u\}$, whence $\|\nabla_x^{\mathfrak{g}}u\| = 1$. Moreover, as the distance function u is *locally semiconcave* in Ω [MM03, Proposition 3.4], it follows that $D^+u(x) \neq \emptyset$ for all $x \in \Omega$ [MM03, Proposition 2.8]. It is easy to see that $D^+u(x)$ is convex: if for some C^1 functions φ, Ψ near x, both $u - \varphi$ and $u - \Psi$ attains a local maximum at x, then so does the function $\zeta_{\lambda} \coloneqq \lambda \varphi + (1 - \lambda) \Psi$ for $\lambda \in [0, 1]$, and hence, $\nabla_x^{\mathfrak{g}}\zeta_{\lambda} = \lambda \nabla_x^{\mathfrak{g}}\varphi + (1 - \lambda) \nabla_x^{\mathfrak{g}}\Psi \in D^+u(x)$. Thus, it follows from the strict convexity of H that $D^+u(x)$ contains at least two vectors if and only if there exists some φ such that $u - \varphi$ attains a local maximum at x and $\|\nabla_x^{\mathfrak{g}}\varphi\|_{\mathfrak{g}} < 1$. Since u is nondifferentiable precisely in $\mathrm{Se}^{\mathrm{in}}(\Omega, \mathfrak{g})$, the claim follows immediately.

We also have the following stability theorem.

Theorem 2.11. [CIL92] Let \mathfrak{g}_i be a sequence of Riemannian metric on a compact manifold M, converging to \mathfrak{g} in the C^0 sense. Let u_i be a viscosity solution defined on $\Omega \subset M$ to the eikonal equation $\|df\|_{\mathfrak{g}_i} = 1$. Suppose $u_i \Rightarrow u$ uniformly on compact sets. Then, u is a viscosity solution on Ω to the eikonal equation $\|df\|_{\mathfrak{g}} = 1$.

Proof. Since $\mathfrak{g}_i \Rightarrow \mathfrak{g}$ in the C^0 sense, we have $\|\cdot\|_{\mathfrak{g}_i} \Rightarrow \|\cdot\|_{\mathfrak{g}}$ uniformly on TM (Lemma 2.2). Fix some $x_0 \in \Omega$. Suppose, φ is a smooth function defined on a neighborhood $x_0 \in U \subset \Omega$ such that $u - \varphi$ attains a local maximum at x_0 . As observed in Remark 2.8, we assume that $u - \varphi$ attains a strict local maximum at x_0 . Then, for $\epsilon > 0$ small, we have some $\delta_{\epsilon} > 0$ such that

$$(u-\varphi)(x) + \delta_{\epsilon} < (u-\varphi)(x_0), \quad \forall d_{\mathfrak{g}}(x_0,x) = \epsilon.$$

Since $u_i \Rightarrow u$ uniformly on compacts, we have some $N_{\epsilon} \ge 1$ such that

$$u(x) - \frac{\delta_{\epsilon}}{2} < u_i(x) < u(x) + \frac{\delta_{\epsilon}}{2}, \quad d_{\mathfrak{g}}(x_0, x) \le \epsilon, \ i \ge N_{\epsilon}.$$

Hence, for any $i \geq N_{\epsilon}$ and for $d_{\mathfrak{g}}(x_0, x) = \epsilon$, we have

$$(u_i - \varphi)(x) = (u_i - u)(x) + (u - \varphi)(x) < \frac{\delta_{\epsilon}}{2} + (u - \varphi)(x_0) - \delta_{\epsilon}$$
$$= (u - \varphi)(x_0) - \frac{\delta_{\epsilon}}{2} < (u_i - \varphi)(x_0).$$

But then, $(u_i - \varphi)$ attains a local maximum at some x_i with $d_{\mathfrak{g}}(x_0, x_i) < \epsilon$, whenever $i \ge N_{\epsilon}$. Taking $\epsilon \to 0$, we can then have a subsequence $x_i \in U$, with $x_i \to x_0$, so that $(u_i - \varphi)$ attains a local maximum at x_i . Now, since u_i is a viscosity subsolution of $\|df\|_{\mathfrak{g}_i} = 1$, we have

$$\left\| d_{x_i} \varphi \right\|_{\mathfrak{g}_i} \le 1$$

As $\|\cdot\|_{\mathfrak{g}_i} \Rightarrow \|\cdot\|_{\mathfrak{g}}$ uniformly, taking $i \to \infty$, we have $\|d_{x_0}\varphi\|_{\mathfrak{g}} \le 1$. But then, u is a subsolution. Arguing similarly, we can show that u is a supersolution as well. Hence, u is a viscosity solution to the eikonal equation $\|df\|_{\mathfrak{g}} = 1$.

3. CONTINUITY OF THE CUT LOCUS MAP

Let M be a compact manifold, without boundary. Fix an auxiliary Riemannian metric \mathfrak{h} on M. We consider the following collections:

- $\mathcal{G} = \mathcal{G}(M) := \{ \mathfrak{g} \in \Gamma T^*M \odot T^*M \mid \mathfrak{g} \text{ is a } C^2 \text{ Riemannian metric} \}, \text{ and }$
- $\mathcal{K} = \mathcal{K}(M) \coloneqq \{K \subset M \mid K \text{ is compact}\}.$

Note that as M is compact, every Riemannian metric \mathfrak{g} on M is complete. We consider the C^2 Whitney topology on \mathcal{G} . For $A, B \in \mathcal{K}$, define the *Hausdorff metric* as

$$d_H(A,B) \coloneqq \max\left\{\sup_{a\in A} d_{\mathfrak{h}}(a,B), \sup_{b\in B} d_{\mathfrak{h}}(A,b)\right\} = \max\left\{\sup_{a\in A} \inf_{b\in B} d_{\mathfrak{h}}(a,b), \sup_{b\in B} \inf_{a\in A} d_{\mathfrak{h}}(a,b)\right\}.$$
(11)

The topology on \mathcal{K} induced by d_H is independent of the choice of the metric \mathfrak{h} . For $(p, \mathfrak{g}) \in M \times \mathcal{G}$, the cut locus $\operatorname{Cu}(p, \mathfrak{g})$ of p with respect to the metric \mathfrak{g} is a closed subset of M, and hence it is compact. In other words, we have a map

$$\operatorname{Cut}: M \times \mathcal{G} \to \mathcal{K}$$
$$(p, \mathfrak{g}) \mapsto \operatorname{Cu}(p, \mathfrak{g}) \tag{12}$$

The main goal of this section is to prove the following.

Theorem 3.1. The cut locus map $Cut : M \times \mathcal{G} \to \mathcal{K}$ is continuous.

Since we are dealing with metric spaces, it is enough to prove the sequential continuity. Suppose $(p_i, \mathfrak{g}_i) \in M \times \mathcal{G}$ converges to (p, \mathfrak{g}) . We shall require the following two lemmas.

Lemma 3.2. Given any $x \in Cu(p, \mathfrak{g})$, there exists a sequence $x_i \in Cu(p_i, \mathfrak{g}_i)$ with $\lim_i x_i = x$.

Proof. Since $\operatorname{Cu}(p, \mathfrak{g}) = \overline{\operatorname{Se}(p, \mathfrak{g})}$, without loss of generality we may assume that $x \in \operatorname{Se}(p, \mathfrak{g})$. As $\lim_i \operatorname{Inj}(\mathfrak{g}_i) = \operatorname{Inj}(\mathfrak{g})$ (Theorem 2.5), we can fix some $\delta > 0$ such that $\delta < \operatorname{Inj}(\mathfrak{g}_i)$ for all i and $\delta < \operatorname{Inj}(\mathfrak{g})$. Without loss of generality, we assume $d_{\mathfrak{g}}(p, p_i) < \delta$ for $i \geq 1$ as well. Denote the sets

$$\Omega_i \coloneqq \left\{ x \in M \mid d_{\mathfrak{g}_i}(p_i, x) > \delta \right\}, \quad \Omega \coloneqq \left\{ x \in M \mid d_{\mathfrak{g}}(p, x) > \delta \right\},$$

and the distance functions

$$u_i \coloneqq d_{\mathfrak{g}_i}(\partial \Omega_i, _) = d_{\mathfrak{g}_i}(p_i, _) - \delta, \quad u \coloneqq d_{\mathfrak{g}}(\partial \Omega, _) = d_{\mathfrak{g}}(p, _) - \delta.$$

It follows from Lemma 2.6 that $u_i \Rightarrow u$ uniformly. Also, note that by Equation 4 we have

$$\operatorname{Cu}^{\operatorname{in}}(\Omega_i, \mathfrak{g}_i) = \operatorname{Cu}(p_i, \mathfrak{g}_i), \quad \operatorname{Cu}^{\operatorname{in}}(\Omega, \mathfrak{g}) = \operatorname{Cu}(p, \mathfrak{g}),$$

and similar relations hold for $Se(\cdot)$ as well. Then, by Proposition 2.10, we have a C^1 function φ defined on a neighborhood $x \in U$ such that, $u - \varphi$ attains a local maximum at x and

$$\left\| d_x \varphi \right\|_{\mathfrak{g}} < 1.$$

By Remark 2.8, we assume that $u - \varphi$ attains a *strict* local maximum at x_0 . Now, $u_i \Rightarrow u$ uniformly. Hence, as in the proof of Theorem 2.11, we have a sequence $x_i \in U$ such that $\lim x_i = x$ and $u_i - \varphi$ attains a local maximum at x_i . By Theorem 2.9, u_i is a viscosity sub-solution for the eikonal equation $\|df\|_{\mathfrak{g}_i} = 1$ in Ω_i . Hence, we have $\|d\varphi\|_{\mathfrak{g}_i} \leq 1$ at x_i . As φ is C^1 , from the convergence of the norm, we have

$$\lim_{i} \left\| d_{x_i} \varphi \right\|_{\mathfrak{g}_i} = \left\| d_x \varphi \right\|_{\mathfrak{g}} < 1.$$

Hence, for *i* sufficiently large, we must have $\|d_{x_i}\varphi\|_{\mathfrak{g}_i} < 1$. But then again by Proposition 2.10, we have $x_i \in \text{Se}(p_i, \mathfrak{g}_i)$. Hence, the claim follows.

Lemma 3.3. Suppose $x_{i_j} \in Cu(p_{i_j}, \mathfrak{g}_{i_j})$ is a convergent sequence with $x := \lim_j x_{i_j}$. Then, $x \in Cu(p, \mathfrak{g})$.

Proof. Firstly, to keep the notation light, we assume $x_i \in Cu(p_i, \mathfrak{g}_i)$. Next, in view of Equation 1, without loss of generality, we may assume that $x_i \in Se(p_i, \mathfrak{g}_i)$ so that $x := \lim x_i$. If possible, suppose, $x \notin Cu(p, \mathfrak{g})$.

Set $L_i = d_{\mathfrak{g}_i}(p_i, x_i)$ and $L = d_{\mathfrak{g}}(p, x)$. From Lemma 2.4, we have $L = \lim L_i$. Let $\gamma_{\mathbf{v}}^{\mathfrak{g}} : [0, L] \to M$ be the *unique* \mathfrak{g} -minimizer joining p to x, where $\mathbf{v} \in T_p M$ satisfies $\|\mathbf{v}\|_{\mathfrak{g}} = 1$. Since $x_i \in \operatorname{Se}(p_i, \mathfrak{g}_i)$, we have $\mathbf{v}_i^1 \neq \mathbf{v}_i^2 \in T_{p_i} M$ with $\|\mathbf{v}_i^1\|_{\mathfrak{g}_i} = 1 = \|\mathbf{v}_i^2\|_{\mathfrak{g}_i}$, such that for j = 1, 2, the \mathfrak{g}_i -geodesics

$$\gamma_i^j \coloneqq \gamma_{\mathbf{v}_i^j}^{\mathfrak{g}_i} : [0, L_i] \to M$$

are distinct \mathfrak{g}_i -minimizers joining p_i to x_i . By a standard Arzelà-Ascoli type argument (see e.g. [Bus55, Theorem 5.16]), passing to a subsequence, we have $\gamma_i^1 \Rightarrow \eta$ uniformly, where $\eta : [0, L] \to M$ is a Lipschitz curve joining p to x. But $L_{\mathfrak{g}}(\eta) = \lim L_{\mathfrak{g}_i}(\gamma_i^1) = \lim L_i = L$. Hence, $\eta = \gamma_v^{\mathfrak{g}}$ and $\lim \mathbf{v}_i^1 = \mathbf{v}$. Note that since $x \notin \operatorname{Cu}(p, \mathfrak{g})$, a sufficiently small extension of η is still the unique minimizer joining p to x.

By Theorem 2.5, we have $\lim \operatorname{Inj}(\mathfrak{g}_i) = \operatorname{Inj}(\mathfrak{g}) > 0$. Hence, we may fix $\delta > 0$ small such that the following holds for *i* sufficiently large:

$$\delta < \frac{1}{2} \operatorname{Inj}(\mathfrak{g}_i), \quad \delta < \frac{1}{2} \operatorname{Inj}(\mathfrak{g}), \quad d_{\mathfrak{g}}(p, p_i) < \frac{\delta}{2}$$

In particular, the sets $S_i := \{z \in M \mid d_{\mathfrak{g}_i}(p_i, z) = \delta\}$ and $S := \{z \in M \mid d_{\mathfrak{g}}(p, z) = \delta\}$ are embedded submanifolds, diffeomorphic to spheres of dimension $\dim M - 1$. Set $q = \eta(\delta) = \gamma^{\mathfrak{g}}_{\mathbf{v}}(\delta) \in S$. We have neighborhoods $q \in U \subset M$ and $\delta \mathbf{v} \in \tilde{U} \subset T_p M$, such that

$$\exp_n^{\mathfrak{g}}: \tilde{U} \to U$$

is a diffeomorphism. We assume that $\overline{U} \subset \{z \in M \mid d_{\mathfrak{g}}(q, z) < \frac{\delta}{2}\}$, and so, \overline{U} is compact. Set $\widetilde{V} := \{\frac{L}{\delta}\mathbf{u} \mid \mathbf{u} \in \widetilde{U}\}$, which is a neighborhood of $L\mathbf{v}$. Shrinking \widetilde{V} (and hence \widetilde{U} and U as necessary), we have $\exp_p^{\mathfrak{g}}|_{\widetilde{V}} : \widetilde{V} \to V$ is a diffeomorphism, where V is a neighborhood of x. This is possible because $x \notin \operatorname{Cu}(p, \mathfrak{g})$. Thus, we have defined a diffeomorphism $\Phi : U \to V$ as the composition of three diffeomorphisms

$$U \xrightarrow{\left(\exp_{p}^{\mathfrak{g}}|_{\tilde{U}}\right)^{-1}} \tilde{U} \xrightarrow{\mathbf{u} \mapsto \frac{L}{\delta}\mathbf{u}} \tilde{V} \xrightarrow{\exp_{p}^{\mathfrak{g}}|_{\tilde{V}}} V,$$

which maps a neighborhood of q to a neighborhood of x. Next, set $q_i \coloneqq \gamma_i^1(\delta)$. Since $\lim q_i = \lim \gamma_i^1(\delta) = q$ and $x = \lim x_i$, we have $x_i \in V$ and $q_i \in U$ for i large. As $q_i \notin \operatorname{Cu}(p_i, \mathfrak{g}_i)$, we can define the compositions $\Phi_i : U \to M$ as follows

$$U \xrightarrow{\left(\exp_{p_{i}}^{\mathfrak{g}_{i}}|_{\tilde{U}_{i}}\right)^{-1}} \tilde{U}_{i} \xrightarrow{\mathbf{u} \mapsto \frac{L_{i}}{\delta}\mathbf{u}} \tilde{V}_{i} \xrightarrow{\exp_{p}^{\mathfrak{g}_{i}}|_{\tilde{V}_{i}}} M,$$

where the first two maps are diffeomorphisms onto the image. Clearly, $\mathbf{v}_i^1 \in U_i$ and $\Phi_i(q_i) = x_i$. Let us fix neighborhood $q \in W \subset U$ with $\overline{W} \subset U$, such that $q_i \in W$. Note that \overline{W} is compact, as \overline{U} is so.

Firstly, we show that for *i* sufficiently large, Φ_i maps W in to V (not-necessarily diffeomorphically). Suppose this is not the case. Then, we have a sequence, $x_{i_j} \in W$, $\mathbf{u}_{i_j} \in \tilde{U}_{i_j}$, with $\exp_{p_{i_j}}^{\mathfrak{g}_{i_j}}(\mathbf{u}_{i_j}) = x_{i_j}$, such that $y_{i_j} \coloneqq \Phi_{i_j}(x_{i_j}) \notin V$. Passing to a farther subsequence, $x_{i_j} \to x \in \overline{W} \subset U$. Let $\mathbf{u} \in \tilde{U}$ so that $\exp_p^{\mathfrak{g}}(\mathbf{u}) = x$. It follows that the minimizers $\gamma_{\mathbf{u}_{i_j}}^{\mathfrak{g}_{i_j}}$ joining p_{i_j} to x_{i_j} must converge, possibly passing to a subsequence, to the unique minimizer $\gamma_{\mathbf{u}}^{\mathfrak{g}}$ joining p to x. Consequently, $\mathbf{u} = \lim \mathbf{u}_{i_j}$. Now, we have

$$y_{i_j} = \Phi_{i_j}(x_{i_j}) = \exp_{p_{i_j}}^{\mathfrak{g}_{i_j}} \left(\frac{L_i}{\delta} \mathbf{u}_{i_j}\right) = \gamma_{\mathbf{u}_{i_j}}^{\mathfrak{g}_{i_j}} \left(\frac{L_i}{\delta} \mathbf{u}_{i_j}\right) \to \gamma_{\mathbf{u}}^{\mathfrak{g}} \left(\frac{L}{\delta}\right) = \exp_p^{\mathfrak{g}} \left(\frac{L}{\delta} \mathbf{u}\right) = \Phi(x)$$

which contradicts $y_{i_j} \notin V$, as $\lim_j y_{i_j} = \Phi(x) \in V$. Hence, for *i* sufficiently large, we get the maps $\Phi_i : W \to V$.

Next, we claim that for i even larger, $\Phi_i: W \to V$ is an embedding. Consider the maps

$$\begin{aligned} \mathbf{h}_i &: \tilde{U}_i \to V & \mathbf{h} &: \tilde{U} \to V \\ \mathbf{u} &\mapsto \exp_{p_i}^{\mathfrak{g}_i} \left(\frac{L_i}{\delta} \mathbf{u} \right), & \mathbf{u} &\mapsto \exp_p^{\mathfrak{g}} \left(\frac{L}{\delta} \mathbf{u} \right) \end{aligned}$$

so that $\Phi_i = h_i \circ \left(\exp_{p_i}^{\mathfrak{g}_i}|_{\tilde{U}_i}\right)^{-1}$ and $\Phi = h \circ \left(\exp_p^{\mathfrak{g}}|_{\tilde{U}}\right)^{-1}$. Recall that h is a diffeomorphism. If possible, suppose, for infinitely many i, we have $a_i^1, a_i^2 \in W$ such that $\Phi_i(a_i^1) = \Phi_i(a_i^2)$. We have $\mathbf{u}_i^1 \neq \mathbf{u}_i^2 \in \tilde{U}_i$ so that $\exp_{p_i}^{\mathfrak{g}_i}(\mathbf{u}_i^j) = a_i^j$ for j = 1, 2. Passing to a subsequence, j = 1, 2, we have $a_i^j \rightarrow a^j \in \bar{W} \subset U$, and hence, $\mathbf{u}^j \in \tilde{U}$ so that $\exp_p^{\mathfrak{g}}(\mathbf{u}^j) = a^j$. Arguing as in the previous paragraph, we see that, possibly for a subsequence, $\mathbf{u}_i^j \rightarrow \mathbf{u}^j$ for j = 1, 2. Note that, since $\gamma_{\mathbf{u}_i^1}^{\mathfrak{g}_i}\left(\frac{L_i}{\delta}\right) = \Phi_i(a_i^2) = \gamma_{\mathbf{u}_i^2}^{\mathfrak{g}_i}\left(\frac{L_i}{\delta}\right)$ for all i, we have

$$\mathsf{h}\left(\frac{L}{\delta}\mathbf{u}^{1}\right) = \gamma_{\mathbf{u}^{1}}^{\mathfrak{g}}\left(\frac{L}{\delta}\right) = \lim_{i} \gamma_{\mathbf{u}_{i}^{1}}^{\mathfrak{g}_{i}}\left(\frac{L_{i}}{\delta}\right) = \lim_{i} \gamma_{\mathbf{u}_{i}^{2}}^{\mathfrak{g}_{i}}\left(\frac{L_{i}}{\delta}\right) = \gamma_{\mathbf{u}^{2}}^{\mathfrak{g}}\left(\frac{L}{\delta}\right) = \mathsf{h}\left(\frac{L}{\delta}\mathbf{u}^{2}\right)$$

As h is a diffeomorphism, we have $\mathbf{u}^1 = \mathbf{u}^2 = \mathbf{u} \in \tilde{U}$. Set $\mathbf{e}_i \coloneqq \frac{\mathbf{u}_i^2 - \mathbf{u}_i^1}{\|\mathbf{u}_i^2 - \mathbf{u}_i^1\|_{\mathfrak{g}_i}}$, so that $\|\mathbf{e}_i\|_{\mathfrak{g}_i} = 1$. Passing to a subsequence, we have $\mathbf{e}_i \to \mathbf{e} \in T_p M$. In particular, $\|\mathbf{e}\|_{\mathfrak{g}} = 1$, and so $\mathbf{e} \neq 0$. Without loss of generality, we can assume a priori that V is a coordinate chart, with $\varphi: V \to \mathbb{R}^{n=\dim M}$ a coordinate map. Then, by the Taylor's theorem, we have

$$d(\varphi \circ \mathsf{h}_{i})|_{\mathbf{u}_{i}^{1}}(\mathbf{u}_{i}^{2}-\mathbf{u}_{i}^{1}) = -\int_{0}^{1} (1-t) \left(d(\varphi \circ \mathsf{h}_{i})|_{t\mathbf{u}_{i}^{2}+(1-t)\mathbf{u}_{i}^{1}} - d(\varphi \circ \mathsf{h}_{i})|_{\mathbf{u}_{i}^{1}} \right) (\mathbf{u}_{i}^{2}-\mathbf{u}_{i}^{1}) dt.$$

Dividing both sides by $\left\|\mathbf{u}_{i}^{2}-\mathbf{u}_{i}^{2}
ight\|_{\mathfrak{g}_{i}}$ gives,

$$d(\varphi \circ \mathsf{h}_i)|_{\mathbf{u}_i^1}(\mathbf{e}_i) = -\int_0^1 (1-t) \left(d(\varphi \circ \mathsf{h}_i)|_{t\mathbf{u}_i^2 + (1-t)\mathbf{u}_i^1} - d(\varphi \circ \mathsf{h}_i)|_{\mathbf{u}_i^1} \right) (\mathbf{e}_i) dt.$$
(13)

If we denote the geodesic flows associated to $\mathfrak{g}_i, \mathfrak{g}$ respectively as

$$\mathfrak{G}_i^t: TM \to TM, \quad \mathfrak{G}^t: TM \to TM,$$

it is easy to see that

$$\mathbf{h}_{i} = \pi \circ \mathfrak{G}_{i}^{\frac{L_{i}}{\delta}}|_{\tilde{U}_{i}}, \quad \mathbf{h} = \pi \circ \mathfrak{G}^{\frac{L}{\delta}}|_{\tilde{U}},$$

where $\pi: TM \to M$ is the projection. Now, it follows from [Sak83, Lemma 1.6] that $\mathfrak{G}_i^{\frac{L_i}{\delta}} \Rightarrow \mathfrak{G}^{\frac{L}{\delta}}$ in the C^1 topology. This implies that the integrand in Equation 13 converges uniformly to the limit, as we can find C^1 bound for π and φ . Hence, taking $i \to \infty$ we get,

$$d(\varphi \circ \mathsf{h})|_{\mathbf{u}}(\mathbf{e}) = -\int_0^1 (1-t) \left(d(\varphi \circ \mathsf{h})|_{\mathbf{u}} - d(\varphi \circ \mathsf{h})|_{\mathbf{u}} \right) (\mathbf{e}) dt = 0.$$

But this contradicts the fact h is a diffeomorphism. Hence, for *i* large enough, we must have $\Phi_i: W \to V$ is injective. A similar argument shows that $\Phi_i: W \to V$ is an immersion for *i* large.

Now, consider the points $q'_i \coloneqq \gamma_i^2(\delta)$. Clearly $q'_i \neq q_i$, as the geodesics γ_i^1, γ_i^2 intersects at time L_i for the first time, and $L_i \ge \text{Inj}(\mathfrak{g}_i) > \delta$. If $q'_i \in U$, then $q'_i \notin \text{Cu}(p_i, \mathfrak{g}_i)$ and clearly, $\left(\exp_{p_i}^{\mathfrak{g}_i}|_{\tilde{U}_i}\right)^{-1}(q'_i) = \delta \mathbf{v}_i^2$. Consequently, $\Phi_i(q'_i) = \exp_{p_i}^{\mathfrak{g}_i}(L_i\mathbf{v}_i^2) = \gamma_i^2(L_i) = x_i = \Phi_i(q_i)$, contradicting the injectivity of Φ_i . Thus, $q'_i \notin U$ for *i* large. On the other hand, by a Busemann type argument as above, we must have a subsequence of γ_i^2 converging to a curve joining *p* to *x*, which is necessarily the unique minimizer $\eta = \gamma_{\mathbf{v}}^{\mathfrak{g}}$. This is a contradiction as $\gamma_i^2(\delta)$ cannot converge to $\eta(\delta)$.

Hence, $x \in Cu(p, \mathfrak{g})$, proving the claim.

Proof of Theorem 3.1. For any sequence $(p_i, \mathfrak{g}_i) \in M \times \mathcal{G}$ converging to (p, \mathfrak{g}) , let us denote $\mathcal{X}_i \coloneqq \operatorname{Cu}(p_i, \mathfrak{g}_i)$ and $\mathcal{X} \coloneqq \operatorname{Cu}(p, \mathfrak{g})$. We apply Lemma 2.1 to show that $\lim d_H(\mathcal{X}_i, \mathcal{X}) = 0$. Clearly, Lemma 3.2 justifies Lemma 2.1 (1), whereas Lemma 2.1 (2) follows from Lemma 3.3. This concludes the proof.

We finish by observing an immediate corollary to Lemma 3.3. Recall that for a fixed point $p \in M$, the continuity of the cut time map $\mathbf{v} \mapsto \rho_{\mathfrak{g}}(p, \mathbf{v})$ is well known [Sak96, Proposition 4.1]. More generally, for a closed submanifold $N \subset M$, the cut time map $\mathbf{n} \mapsto \rho_{\mathfrak{g}}(N, \mathbf{n})$ is also continuous, where \mathbf{n} varies in the normal bundle of N [BP23, BP24]. We have the following, which improves up on the result from [Sak96] for M compact.

Corollary 3.4. Suppose \mathfrak{g}_i is a sequence of Riemannian metric on a compact manifold M, converging in the C^2 norm to the metric \mathfrak{g} . Let $\mathbf{v}_i \in T_{p_i}M$ be vectors with $\|\mathbf{v}_i\|_{\mathfrak{g}_i} = 1$, converging to $\mathbf{v} \in T_pM$ (necessarily with $\|\mathbf{v}\|_{\mathfrak{g}} = 1$), where $p = \lim p_i$. Then, $\lim \rho_{\mathfrak{g}_i}(p_i, \mathbf{v}_i) = \rho_{\mathfrak{g}}(p, \mathbf{v})$.

Proof. It follows from Theorem 2.5 that $\lim \operatorname{Diam}(\mathfrak{g}_i) = \operatorname{Diam}(\mathfrak{g}) < \infty$. Hence, without loss of generality, we have $\operatorname{Diam}(\mathfrak{g}_i) < K \coloneqq \operatorname{Diam}(\mathfrak{g}) + 1$ for all *i*. But then $R_i \coloneqq \rho_{\mathfrak{g}_i}(p_i, \mathbf{v}_i) < K$ as well. Also, $R_i \ge \operatorname{Inj}(\mathfrak{g}_i) > 0$. In other words, $\{R_i\} \subset [0, K]$ is a bounded sequence in \mathbb{R} . Suppose, $\{R_{i_j}\}$ is a convergent subsequence, with $R \coloneqq \lim_j R_{i_j}$. Set $x_{i_j} \coloneqq \gamma_{\mathbf{v}_{i_j}}^{\mathfrak{g}_{i_j}}(R_{i_j})$ and $x \coloneqq \gamma_{\mathbf{v}}^{\mathfrak{g}}(R)$. Clearly, $x_{i_j} \in \operatorname{Cu}(p_{i_j}, \mathfrak{g}_{i_j})$ and also, $x = \lim x_{i_j}$. But then by Lemma 3.3, we have $x \in \operatorname{Cu}(p, \mathfrak{g})$. Consequently, $R = \rho_{\mathfrak{g}}(p, \mathbf{v})$. The claim then follows.

ACKNOWLEDGMENTS

The first and third authors would like to express their gratitude to Prof. P. Albano for clarifying certain points from the article [Alb16] through email correspondence. The first author was supported by the NBHM grant no. 0204/1(5)/2022/R&D-II/5649. The second author was partially supported by Grant-in Aid for Scientific Research (C) (No. 21K03238), Japan Science for the Promotion of Science. The third author was supported by Jilin University.

STABILITY OF THE CUT LOCUS

References

- [Alb16] Paolo Albano. On the stability of the cut locus. Nonlinear Anal., 136:51-61, 2016. doi:10.1016/j.na.2016.02.008.
- [BP23] Somnath Basu and Sachchidanand Prasad. A connection between cut locus, Thom space and Morse-Bott functions. *Algebr. Geom. Topol.*, 23(9):4185–4233, 2023. doi:10.2140/agt.2023.23.4185.
- [BP24] Aritra Bhowmick and Sachchidanand Prasad. On the cut locus of submanifolds of a Finsler manifold. J. Geom. Anal., 34(10):Paper No. 308, 38, 2024. doi:10.1007/s12220-024-01751-1.
- [Buc77a] Michael A. Buchner. Simplicial structure of the real analytic cut locus. *Proc. Amer. Math. Soc.*, 64(1):118–121, 1977. doi:10.2307/2040994.
- [Buc77b] Michael A. Buchner. Stability of the cut locus in dimensions less than or equal to 6. *Invent. Math.*, 43(3):199–231, 1977. doi:10.1007/BF01390080.
- [Bus55] Herbert Busemann. The geometry of geodesics. Academic Press Inc., New York, N. Y., 1955.
- [CIL92] Michael G. Crandall, Hitoshi Ishii, and Pierre-Louis Lions. User's guide to viscosity solutions of second order partial differential equations. Bull. Amer. Math. Soc. (N.S.), 27(1):1–67, 1992. doi:10.1090/S0273-0979-1992-00266-5.
- [EGGHT21] Benjamin Eltzner, Fernando Galaz-García, Stephan Huckemann, and Wilderich Tuschmann. Stability of the cut locus and a central limit theorem for Fréchet means of Riemannian manifolds. Proc. Amer. Math. Soc., 149(9):3947–3963, 2021. doi:10.1090/proc/15429.
- [Ehr74] Paul E. Ehrlich. Continuity properties of the injectivity radius function. *Compositio Math.*, 29:151–178, 1974.
- [GG73] M. Golubitsky and V. Guillemin. Stable mappings and their singularities, volume Vol. 14 of Graduate Texts in Mathematics. Springer-Verlag, New York-Heidelberg, 1973.
- [Kob67] Shoshichi Kobayashi. On conjugate and cut loci. In Studies in Global Geometry and Analysis, pages 96–122. Math. Assoc. Amer. (distributed by Prentice-Hall, Englewood Cliffs, N.J.), 1967.
- [Mic80] Peter W. Michor. *Manifolds of differentiable mappings*, volume 3 of *Shiva Mathematics Series*. Shiva Publishing Ltd., Nantwich, 1980.
- [MM03] Carlo Mantegazza and Andrea Carlo Mennucci. Hamilton-Jacobi equations and distance functions on Riemannian manifolds. *Appl. Math. Optim.*, 47(1):1–25, 2003. doi:10.1007/s00245-002-0736-4.
- [Poi05] Henri Poincaré. Sur les lignes géodésiques des surfaces convexes. *Trans. Amer. Math. Soc.*, 6(3):237–274, 1905. doi:10.2307/1986219.
- [Sak83] Takashi Sakai. On continuity of injectivity radius function. Math. J. Okayama Univ., 25(1):91–97, 1983.
- [Sak96] Takashi Sakai. Riemannian geometry, volume 149 of Translations of Mathematical Monographs. American Mathematical Society, Providence, RI, 1996. Translated from the 1992 Japanese original by the author. doi:10.1090/mmono/149.
- C. [Whi32] Η. Whitehead. J. Convex Regions in the Geometry of Paths. The Quarterly Journal of Mathematics, os-3(1):33-42, 01 1932. URL: https://academic.oup.com/qjmath/article-pdf/os-3/1/33/4486996/os-3-1-33.pdf, doi:10.1093/qmath/os-3.1.33.
- [Wol79] Franz-Erich Wolter. Distance function and cut loci on a complete Riemannian manifold. Arch. Math. (Basel), 32(1):92–96, 1979. doi:10.1007/BF01238473.