

Bayesian Outlier Detection for Matrix–variate Models

Monica Billio¹ Roberto Casarin^{1,2} Fausto Corradin¹ Antonio Peruzzi¹

March 26, 2025

Abstract

Bayes Factor (BF) is one of the tools used in Bayesian analysis for model selection. The predictive BF finds application in detecting outliers, which are relevant sources of estimation and forecast errors. An efficient framework for outlier detection is provided and purposely designed for large multidimensional datasets. Online detection and analytical tractability guarantee the procedure’s efficiency. The proposed sequential Bayesian monitoring extends the univariate setup to a matrix–variate one. Prior perturbation based on power discounting is applied to obtain tractable predictive BFs. This way, computationally intensive procedures used in Bayesian Analysis are not required. The conditions leading to inconclusive responses in outlier identification are derived, and some robust approaches are proposed that exploit the predictive BF’s variability to improve the standard discounting method. The effectiveness of the procedure is studied using simulated data. An illustration is provided through applications to relevant benchmark datasets from macroeconomics and finance.

Keywords: Bayesian Modelling, Bayes Factor, Matrix–variate, Sequential Model Assessment, Outliers.

1 Introduction

Robust methods are commonly used in data analysis to identify outliers and then to conduct inference after removing the outliers, defined as observations that differ significantly from other data points in the sample (e.g., see Tsukurimichi et al., 2022). An outlier may be due to variability in the measurement or significant experimental errors; the latter are sometimes excluded from the data set. Outlier detection is a statistical problem that has received considerable attention from the frequentist and Bayesian perspectives. As argued by Bayarri and Berger (2004), both frameworks should be considered in applications since they offer distinct perspectives. In addition, real-time or sequential monitoring for high dimensional datasets requires easy and effective tools, which naturally call for tractable iterative procedures for assessing model fitting and prediction. This paper aims to contribute to the Bayesian literature by extending monitoring procedures for univariate models to matrix variate models. We follow a Bayesian approach since it naturally allows for sequential updating of the estimators involved in the monitoring and derive analytical results to avoid computationally intensive approximation procedures.

Within a Bayesian framework, a common approach assumes that (possible) outliers are generated by contaminating models different from the one generating the rest of the data. Previous Bayesian works along these lines can be found in Box and Tiao (1968); Guttman (1973);

¹Ca’ Foscari University of Venice

²Corresponding author address: r.casarin@unive.it. Other contacts: Monica Billio (billio@unive.it), Fausto Corradin (fausto.corradin@unive.it), Antonio Peruzzi (antonio.peruzzi@unive.it)

Abraham and Box (1979); Guttman et al. (1978); Pettit and Smith (1983, 1985); Pettit (1988, 1992, 1990); Verdinelli and Wasserman (1991); Bayarri and Berger (1994); Hoeting et al. (1996). We follow a Bayes Factor (BF) approach, which is widely used in Bayesian analysis (Penny et al., 2004; Schrider et al., 2016; Ly et al., 2016; Chen et al., 2018; Li et al., 2021; Stefan et al., 2019) including selective inference (Yekutieli, 2012; Zhao et al., 2022; Rasines and Young, 2022) and outliers detection (Bayarri and Morales, 2003; Guan and Tibshirani, 2022). The BF is also connected to Schwartz’s criterion or Bayesian Information Criterion (Ghosh et al., 2006), which is another approach commonly used in the outlier detection literature (e.g., see She and Owen, 2011; Riani et al., 2024). See Lahiri (2001) for a review of BF calculation methods and their use in model selection.

Due to the extreme sensitivity (De Santis and Spezzaferri, 1997) of the BF to the choice of the alternative distribution, robust methodologies have been developed (Kosheleva et al., 2021; Li et al., 2021; Schad et al., 2023). In this paper, we follow the power discounting approach of West (1986) and assume the alternative distribution is proportional to the principal distribution raised to a power. We consider the Gaussian family since it is a standard assumption in many fields (e.g., see Billio et al., 2021, 2023; Guhaniyogi et al., 2017) and extend the existing procedure for univariate Gaussian models (West, 1986) to the matrix-variate case, thus providing an original contribution to the expanding literature in this area (e.g., see Landim and Gamerman, 2000; Triantafyllopoulos, 2008; Wang and West, 2009; Carvalho and West, 2007; Carvalho et al., 2007; Viroli, 2011; Leng and Tang, 2012; Viroli and Anderlucci, 2013; Zhu et al., 2017; Chen et al., 2019; Thompson et al., 2020; Chen et al., 2021; Yu et al., 2022; Billio et al., 2021, 2023; Tomarchio et al., 2022). This paper presents not only new results for detecting outliers in the matrix case but also uncovers novel results for the univariate Gaussian model. Furthermore, some pre-existing results for the univariate model are recovered as special cases.

We also prove analytically that the outcome of the BF procedure for detecting outliers heavily depends on the choice of the discounting factor. Thus, we propose robust BFs to alleviate the problem while maintaining a certain degree of tractability. The first solution is a conservative approach based on the minimum BF, a strategy commonly used in the literature to reduce the dependence on the prior choice. A second solution is the integrated BF, which arises from averaging BFs across various discounting factor values, assuming the factor is random and follows a beta distribution. The distribution parameters are selected to return a well-defined integrated BF. The outcomes based on these solutions are sensitive to the choice of the hyper-parameters. Thus, we propose a new BF calibration procedure for the matrix Gaussian model. We derive the analytical distribution of the predictive BF and use it to select the calibrated discounting factor and to derive the inconclusiveness region for the calibrated BF. Through some simulation experiments, we investigate the properties of a testing procedure based on the calibrated BF under various outlier generation settings.

Since outliers can have a large impact on the outcome of empirical analysis of economics data, after the 2009 crisis and the COVID-19 pandemic event, the treatment of outliers attracted the attention of both researchers and the institutes of official statistics, which provided some guidelines on monitoring the effects of outliers when using their data. Thus, this paper illustrates our sequential outlier detection procedure on three relevant economic datasets: a multiple-unit dataset of macroeconomic variables, a network-valued dataset of international trade and a network-valued financial dataset. For those datasets, linear Gaussian models with constant parameters (e.g., see Canova and Ciccarelli, 2004, 2009; Feldkircher et al., 2022; Billio et al., 2023) have been extended to account for various forms of instabilities (Billio et al., 2016, 2022). Thus, the approach developed in this paper can serve as a preliminary procedure for detecting breaks and outliers before a suitable modelling framework is chosen.

The paper is organized as follows. Section 2 introduces the monitoring procedure based on BF and presents the main results. Section 3 provides a simulation study of the monitoring procedures. Section 4 presents some real-data illustrations. Section 5 concludes.

2 Bayesian Model Monitoring

2.1 Bayes Factor

Consider a sequence of observations Y_t , $t = 1, 2, \dots$ with $Y_t \in \mathcal{Y}$ where \mathcal{Y} is a possibly multidimensional sample space and t is a time index. We denote with y_t a realization of the random variable (r.v.) Y_t and assume the information available at time t is given by the collection of past observations $\mathcal{D}_t = \{y_1, \dots, y_t\}$. Given the parameter and past observations, we assume the conditional sampling distribution does not depend on \mathcal{D}_{t-1} and belongs to a parametric family with unknown parameter $\theta \in \Theta$. The parameter space Θ is endowed with a prior distribution $p(\theta)$.

Since our outlier detection procedure applies sequentially over time, we assume at time $t - 1$ a posterior distribution for θ is formed given \mathcal{D}_{t-1} with density given by $p(\theta_t|\mathcal{D}_{t-1}) \propto p(\theta)p(y_1|\theta) \cdot \dots \cdot p(y_{t-1}|\theta)$. We denote with θ_t the random variable $\theta|\mathcal{D}_{t-1}$ and assume its prior distribution is $p(\theta_t|\mathcal{D}_{t-1})$. The marginal predictive distribution for Y_t has density:

$$p(y|\mathcal{D}_{t-1}) = \int_{\Theta} g(y|\theta_t)p(\theta_t|\mathcal{D}_{t-1})d\theta_t, \quad y \in \mathcal{Y}. \quad (1)$$

The distribution $p(y|\mathcal{D}_{t-1})$ naturally provides a measure of the predictive ability of the model $g(y|\theta)$. In the following, we assume the predictive distribution belongs to the same distribution family as the sampling distribution, that is, $g(y|\theta) = p(y|\theta)$. In hypothesis testing, a model for the alternative hypothesis \mathcal{H}_1 is assumed, $p_A(y|\mathcal{D}_{t-1})$ and the prior predictive distribution

$$p_A(y|\mathcal{D}_{t-1}) = \int_{\Theta} p(y|\theta_t)p_A(\theta_t|\mathcal{D}_{t-1})d\theta_t, \quad y \in \mathcal{Y} \quad (2)$$

is obtained as the marginal distribution with respect to the prior distribution at time t under the alternative hypothesis. The Bayes Factor is defined as the ratio of the two marginal distributions:

$$H_t = \frac{p(y|\mathcal{D}_{t-1})}{p_A(y|\mathcal{D}_{t-1})} = \frac{\int_{\Theta} p(y|\theta_t)p(\theta_t|\mathcal{D}_{t-1})d\theta_t}{\int_{\Theta} p(y|\theta_t)p_A(\theta_t|\mathcal{D}_{t-1})d\theta_t} \quad (3)$$

and results as an optimal decision of an inference problem under a 0-1 loss function (e.g., see Robert, 2007, chs. 5,7). If $H_t > 1$, the null hypothesis \mathcal{H}_0 is not rejected, then we conclude there is evidence from the observation y in favour of the null hypothesis. This paper considers the BF as a testing procedure for model failure, which includes sensitivity to structural changes and outliers. To define the distribution under the alternative, we follow an approach based on a parametric perturbation of the posterior predictive distribution

$$p_A(\theta_t|\mathcal{D}_{t-1}) = G(p(\theta_t|\mathcal{D}_{t-1}), \alpha_t), \quad (4)$$

where α_t is a possibly time-varying perturbation parameter and $G(\cdot, \cdot)$ is a perturbation function from $[0, 1] \times [0, 1]$ onto $[0, 1]$. Examples of perturbation functions can be derived from the approaches based on mixtures of distributions, $p_A(\theta_t|\mathcal{D}_{t-1}) = \alpha_t p(\theta_t) + (1 - \alpha_t)p(\theta_t|\mathcal{D}_{t-1})$ (Robert, 2022), and on distortion of probability measures, $p_A(\theta_t|\mathcal{D}_{t-1}) = g(\int_{-\infty}^{\theta_t} p(u|\mathcal{D}_{t-1})du, \alpha_t)p(\theta_t|\mathcal{D}_{t-1})$ where g is the partial derivative of G with respect to its first argument.

In this paper, we assume the following perturbation function $G(p, \alpha_t) = p^{\alpha_t} C(\alpha_t)$ where

$$C(\alpha_t) = \left(\int_{\Theta} p(\theta_t | \mathcal{D}_{t-1})^{\alpha_t} d\theta_t \right)^{-1} \quad (5)$$

is the inverse normalising constant of the density under the alternative hypothesis. The constant satisfies $C(\alpha_t) \rightarrow 1$ as $\alpha_t \rightarrow 1^-$ and $C(\alpha_t) \rightarrow \lambda(\Theta)^{-1}$ as $\alpha_t \rightarrow 0^+$, where $\lambda(\Theta) < \infty$ denotes the Lebesgue measure of Θ . If $\lambda(\Theta)$ is unbounded, then to prevent $C(\alpha_t) \rightarrow 0$ as $\alpha_t \rightarrow 0^+$, some restrictions can be introduced such as $\alpha_t \in (\underline{\alpha}, \bar{\alpha}) \subset (0, 1)$, $\underline{\alpha} > 0$. These limits are relevant in the analysis of the BF, denoted as $H_t(\alpha_t)$ in what follows. This choice of the calibration function returns the power discounting approach proposed in the seminal paper West (1986) for eliciting the prior under the alternative hypothesis

$$p_A(\theta_t | \mathcal{D}_{t-1}) \propto (p(\theta_t | \mathcal{D}_{t-1}))^{\alpha_t}, \quad (6)$$

where the discounting parameter α_t takes values in the interval $(0, 1)$. An advantage of the power discounting approach is that it returns the popular variance contamination model for outlier detection (e.g., see Pettit, 1992; Weiss, 1997; Bayarri and Morales, 2003; Page and Dunson, 2011; Tomarchio et al., 2022) and does not require the estimation of the contamination parameter, which would be difficult, since very little information is available about the outlier generating process. See also van der Merwe et al. (2021) for a comparison between Bayesian contamination procedures and alternative frequentist and Bayesian procedures.

We assume Gaussian likelihood and conjugate prior distribution to preserve some analytical tractability and extend the univariate discounting approach to a matrix-variate setting. We will prove that the BF is generally well-defined and bounded by a function of α_t . However, the results of the hypothesis testing can be significantly influenced by the value of α_t , and the BF can become unbounded in the limit for $\alpha_t \rightarrow 0^+$. Consequently, the discounting parameter should be chosen carefully. The new results for the matrix setting readily apply to the previous univariate and multivariate approaches.

2.2 Robust Bayes Factors

As one can expect, the outcome of a decision based on the BF may depend on the value of α_t . See also the discussion in West (1986). We prove that for a given level H^* of the threshold (which is assumed equal to one in the following), for some values of α_t , the evidence is against the null hypothesis, and for some others, it is against the alternative. In the application to outlier detection, one can face a situation in which the new observation y_t is considered an outlier or not depending on the value of α_t , also in a very sensitive way. The following result sheds some light on the indeterminacy of the outcome of the BF procedure.

Proposition 1. *Assume the likelihood satisfies*

$$\int_{\Theta} p(y_t | \theta_t) (\lambda(\Theta) p(\theta_t | \mathcal{D}_{t-1}) - 1) \lambda(d\theta_t) > 0 \quad (7)$$

with respect to the reference measure λ , and there exists a unique α_t^ such that $q_1(\alpha_t) - q_2(\alpha_t) < 0$ for $\alpha_t < \alpha_t^*$ and $q_1(\alpha_t) - q_2(\alpha_t) > 0$ for $\alpha_t > \alpha_t^*$ and $q_1(\alpha_t^*) - q_2(\alpha_t^*) = 0$, where*

$$q_1(\alpha_t) = \int_{\Theta} p(y_t | \theta_t) p(\theta_t | \mathcal{D}_{t-1})^{\alpha_t} \lambda(d\theta_t) \int_{\Theta} p(\theta_t | \mathcal{D}_{t-1})^{\alpha_t} \log p(\theta_t | \mathcal{D}_{t-1}) \lambda(d\theta_t) \quad (8)$$

$$q_2(\alpha_t) = \int_{\Theta} p(\theta_t|\mathcal{D}_{t-1})^{\alpha_t} \lambda(d\theta_t) \int_{\Theta} p(y_t|\theta_t) p(\theta_t|\mathcal{D}_{t-1})^{\alpha_t} \log p(\theta_t|\mathcal{D}_{t-1}) \lambda(d\theta_t). \quad (9)$$

Then there exists a partition (U_1, U_2) of the unit interval such that $H_t(\alpha_t) > 1$ for $\alpha_t \in U_1$ and $H_t(\alpha_t) < 1$ for $\alpha_t \in U_2$.

Proposition 1 establishes the conditions such that different values of α_t may cause the BF $H_t(\alpha_t)$ to uniquely cross the threshold value $H_t(\alpha_t^*) = 1$ at α_t^* . It can be seen from the proof that for $\lambda(\Theta)$ unbounded, the evidence against the presence of an outlier is negative for $\alpha_t \rightarrow 0^+$, consistently with the Jeffreys–Lindley Paradox (e.g., see Robert, 2014; Bernardo and Smith, 2009, ch. 6). See also Robert et al. (2009) and Wagenmakers and Ly (2023) for a review with a historical perspective. In addition, we shall notice that, in the Bayesian current practice, the BF is usually interpreted following Jeffrey’s scale of evidence against the null hypothesis (e.g., see Kass and Raftery, 1995), which suggests the evidence is negative for $1/H_t < 1$, not worth more than a bare mention for $1 < 1/H_t < 10^{1/2}$, substantial for $10^{1/2} < 1/H_t < 10$, strong for $10 < 1/H_t < 10^{3/2}$, very strong for $10^{3/2} < 1/H_t < 10^2$ and decisive for $1/H_t > 10^2$. The following example illustrates the result given in the previous proposition and shows that an observation can be regarded as an outlier depending on the value of α_t . The illustration is general since a stepwise uniform prior is assumed, and the differentiability of the prior is not required to find the threshold value of α_t .

Remark 1 (Univariate Gaussian model). Assume a Gaussian likelihood with location θ and variance σ^2 , that is $Y_t \sim \mathcal{N}(\theta, \sigma^2)$ i.i.d. $t = 1, \dots, T$. Let us assume σ^2 is known and θ follows a prior distribution $p(\theta_t|\mathcal{D}_{t-1})$ which is absolutely continuous with respect to the Lebesgue measure λ , and has density function

$$p(\theta_t|\mathcal{D}_{t-1}) = \sum_{j=1}^n g_j \mathbb{I}(\theta_t \in \Theta_j), \quad (10)$$

where $\{\Theta_j, j = 1, \dots, n\}$ and $\Theta = \Theta_1 \cup \dots \cup \Theta_n$ with $n > 1$ and such that $g_1 \lambda(\Theta_1) + \dots + g_n \lambda(\Theta_n) = 1$. In Table 1, we report three cases in which an observation y_t can be regarded as an outlier depending on the value of α_t . The table reports together with y_t , the value α_t^* such that $H_t(\alpha_t^*) = 1$, the step-function prior specification with possibly disconnected support given by the union of $\Theta_j = (\underline{\theta}_j, \bar{\theta}_j)$, and density g_j for $j \in 1, \dots, 3$. The table reports also the likelihood values $p_j = p(y_t|\theta_j)$ with $\theta_j \in \Theta_j$.

| α_t^* | y_t | σ | $\underline{\theta}_1$ | $\bar{\theta}_1$ | $\underline{\theta}_2$ | $\bar{\theta}_2$ | $\underline{\theta}_3$ | $\bar{\theta}_3$ | g_1 | g_2 | g_3 | p_1 | p_2 | p_3 |
|--------------|--------|----------|------------------------|------------------|------------------------|------------------|------------------------|------------------|-------|-------|--------|-------|-------|-------|
| 0.054 | -4.422 | 1.12 | 0.471 | 0.565 | 1.487 | 1.507 | 2.863 | 5.863 | 0.44 | 50 | 0.001 | 0.18 | 0.001 | 0.001 |
| 0.076 | -5.69 | 1.054 | -1.112 | 1.689 | 2.422 | 2.588 | 4.78 | 4.8 | 0.002 | 0.492 | 45.645 | 0.002 | 0.153 | 0.002 |
| 0.412 | -8.723 | 1.939 | 0.996 | 1.409 | 2.628 | 4.209 | 5.498 | 5.549 | 0.934 | 0.023 | 11.51 | 0.146 | 0.019 | 0.041 |

Table 1: Prior hyper-parameters $\underline{\theta}_j$, $\bar{\theta}_j$ and g_j and the likelihood values $p_j = p(y_t|\theta_j)$, with $j = 1, \dots, 3$.

The marginal likelihood under the alternative hypothesis is derived as follows. By the mean-value theorem for integrals on a bounded domain, there exists $\theta_j \in \Theta_j$ such that

$$\int_{\Theta} p(y_t|\theta) p_A(\theta|\mathcal{D}_{t-1}) d\theta = C(\alpha_t) \sum_{j=1}^n \int_{\Theta_j} p(y_t|\theta) g_j^{\alpha_t} d\theta = C(\alpha_t) \sum_{j=1}^n g_j^{\alpha_t} p_j \lambda(\Theta_j), \quad (11)$$

where $p_j = p(y_t|\theta_j)$ and $C(\alpha_t) = g_1^{\alpha_t} + g_2^{\alpha_t} + g_3^{\alpha_t}$ is the normalizing constant. This expression can be used to show that the conditions in Prop. 1 are satisfied for the cases provided in Table 1. The




Figure 1: Value of the Bayes Factor $H_t(\alpha_t)$ (—) as a function of α_t (horizontal axis) for the parameter settings given in Tab. 1. Each plot's thresholds are at 1 and $10^{-1/2}$ (---).

values of p_j show that the prior predictive assigns a non-negligible probability to the observations considered in the example. Figure 1 shows that the observation y_t is an outlier for values of $\alpha_t < \alpha_t^*$ ($H_t(\alpha_t) > 1$), while y_t is considered an outlier for $\alpha_t > \alpha_t^*$ ($H_t(\alpha_t) < 1$). Also, for some observations and settings, comparing the BF with the first threshold of Jeffrey's scale, $10^{1/2}$, returns substantial evidence against the null (first and second panel) depending on α_t . The third example shows that the farther the observed value from Θ , e.g. $y_t = -8.723$, the larger α_t needs to be in order to make the presence of an outlier a convincing hypothesis, i.e. $\alpha_t^* = 0.412$.

Since the outcome of the testing procedure strongly depends on the choice of the discounting parameter α_t and since different thresholds for H_t can be chosen, we propose three alternative Bayesian decision rules to conduct outlier detection. These rules are robust and exploit the variability of the BF due to the discounting parameter and the sampling distribution.

A first decision rule is the Minimum BF (MBF), i.e. the smallest possible BF within the class of alternative distributions specified in Eq. 6, that is

$$H_t^{(MBF)} = \inf_{\alpha_t \in (\underline{\alpha}, \bar{\alpha}) \subset (0,1)} H_t(\alpha_t).$$

The MBF has been used in other testing problems such as unit root testing (Berger and Yang, 1994), correlation testing (Fang Chen and Wang, 2021) and reverse-Bayes procedures (Pawel and Held, 2022). See also Held and Ott (2018) for a review of minimum BFs and a comparison with standard BF. The rationale of this decision rule is that the evidence for the null is at least the MBF, (e.g., see Kosheleva et al., 2021). In the example of Remark 1, the three settings return an MBF below one. However, the MBF is not too far below the threshold in the first setting compared to the other two scenarios. Moreover, we can have cases in which $H_t < 1$ for a coarse set of α_t values. These considerations motivate the need for alternative decision rules.

We consider the Integrated BF (IBF) as a second decision rule. With this rule, one assumes a prior distribution for the discounting parameter α_t and averages the BF over all possible discounting values, that is

$$H_t^{(IBF)} = \int_{\underline{\alpha}}^{\bar{\alpha}} H_t(\alpha_t) \pi_t(\alpha_t) d\alpha_t - 1,$$

where $\pi_t(\alpha)$ is a suitable probability density function for α_t with support $(\underline{\alpha}, \bar{\alpha}) \subset (0, 1)$. The choice of the measure π_t would be crucial to achieve a well-defined IBF in the matrix variate case.

Since the BF can be bounded from above under mild regularity conditions, the IBF can be modified to account for the relative magnitude of the evidence in favour of the null. We thus

introduce the Normalized IBF (NIBF) defined as

$$H_t^{(NIBF)} = \left(\int_{\underline{\alpha}}^{\bar{\alpha}} H_t(\alpha_t) \pi_t(\alpha_t) d\alpha_t - 1 \right) \left(\int_{\underline{\alpha}}^{\bar{\alpha}} \kappa_t(\alpha_t) \pi_t(\alpha_t) d\alpha_t - 1 \right)^{-1},$$

where $\kappa_t(\alpha_t) < \infty$ is an upper bound for $H_t(\alpha_t)$. The upper bound will be used later in this paper to show the BF integrability with respect to the discounting parameter.

IBF and NIBF do not account for the sampling variability in hypothesis testing. For this reason, we propose a predictive BF approach to incorporate such variability. As y_t is not observed at time $t - 1$, the predictive BF is a random variable whose marginal distribution can be used to derive a calibrated value for the discounting parameter α_t . With our approach, we account for the sampling variability of the predictive BF and find the analytical distributions $F_{j,t}$ of the random BF, $H_t(\alpha_t)$, under the null hypothesis \mathcal{H}_0 of the absence of outliers ($j = 0$) and the alternative hypothesis \mathcal{H}_1 of the presence of outliers ($j = 1$). The calibrated predictive BF is derived through the following steps.

1. Derive the threshold function $\underline{h}(\alpha_t)$, $\alpha_t \in (\underline{\alpha}, \bar{\alpha})$ for rejecting the null \mathcal{H}_0 , such that $\mathbb{P}(\{H_t < \underline{h}(\alpha_t)\}) = \tau$ under the null for a given test size τ . The probability is evaluated using the $F_{0,t}$ distribution.
2. Determine the discounting parameter $\alpha_t^* \in (\underline{\alpha}, \bar{\alpha})$ such that for a given power level β $\mathbb{P}(\{H_t > \bar{h}(\alpha_t^*)\}) = 1 - \beta$ under the alternative, that is:

$$\beta = 1 - F_{1,t}(\bar{h}(\alpha_t^*)), \quad \bar{h}(\alpha_t^*) = 2 - \underline{h}(\alpha_t^*),$$

where the constraint comes from the symmetric assumption of the thresholds.

3. Define the inconclusive interval $C_H = (\underline{h}(\alpha_t^*), \bar{h}(\alpha_t^*))$ where the BF is statistically equal to one and define the calibrated BF as $H_t^* = H_t(\alpha_t^*)$.

The conclusion of the test procedure is rejecting the null hypothesis \mathcal{H}_0 when $H_t < \underline{h}$, accepting it when $H_t > \bar{h}$ and randomizing when $H_t \in C_H$. As randomization is usually not appealing in some applications, an alternative procedure can be used where only one threshold $\bar{h}(\alpha_t^*) = \underline{h}(\alpha_t^*) < 1$ is considered to define a critical region $(0, \underline{h}(\alpha_t^*))$. Our procedure for the calibrated value of α_t is similar in spirit to the one proposed in Weiss (1997), and also presented in Pawel and Held (2025), for determining the optimal sample size.

3 Matrix Normal Monitoring

In this section, we review the notation for the matrix normal model, present some preliminary results, and state the main findings on the BF properties for the normal matrix model. Proofs of the results are given in Appendix A.

3.1 A Bayesian Matrix Normal Model

We adapt to our notation the definition of matrix normal distribution provided in Definition 2.2.1, p. 55 of Gupta and Nagar (1999). In addition, we provide some results on the Bayesian inference for the location matrix of the matrix normal distribution under a conjugate prior assumption.

Definition 1 (Matrix normal). *The random matrix X ($p \times n$) is said to have a matrix variate normal distribution with mean matrix M ($p \times n$) and covariance matrix $\Sigma \otimes V$, where Σ ($p \times p$) > 0 and V ($n \times n$) > 0 , if $\text{vec}(X') \sim \mathcal{N}_{p,n}(\text{vec}(M'), \Sigma \otimes V)$, where \otimes denotes the Kronecker product and $\text{vec}(\cdot)$ the vectorization operator which stacks matrix columns vertically. We shall use the notation $X \sim \mathcal{N}_{p,n}(M, \Sigma, V)$ and the pdf of X is given in Theorem 2.2.1, p. 55 of Gupta and Nagar (1999):*

$$f(X|M, \Psi, \Sigma) = \frac{\exp\left\{-\frac{1}{2}\text{tr}\left[\Sigma^{-1}(X-M)\Psi^{-1}(X-M)'\right]\right\}}{(2\pi)^{\frac{np}{2}}|\Sigma|^{\frac{n}{2}}|\Psi|^{\frac{p}{2}}}. \quad (12)$$

We shall notice that the parameters Σ and Ψ are not identifiable since the likelihood is invariant to the rescaling of the two parameters. We refer the reader to Anderlucci and Viroli (2015); Gallagher and McNicholas (2018) for a treatment of this identification issue. In this paper, to achieve analytical tractability, we assume Σ and Ψ are given, as it is done for the univariate setting in West (1986). Nevertheless, some of our analytical results can be extended to cases where the covariance matrix is estimated following a Bayesian procedure. A standard prior assumption for a covariance matrix is the Inverse Wishart distribution, which is defined using the following parametrization.

Definition 2 (Inverse Wishart). *The random matrix V ($n \times n$) is said to have an inverse Wishart distribution with location parameter Ψ ($n \times n$) and degrees of freedom parameter m , where Ψ ($n \times n$) > 0 and $m > 2n$, if the pdf of V is:*

$$g(V|\Psi, m) = \frac{|\Psi|^{\frac{m-n-1}{2}} \exp\left\{-\frac{1}{2}\text{tr}(\Psi V^{-1})\right\}}{2^{\frac{(m-n-1)n}{2}} \Gamma_n\left(\frac{m-n-1}{2}\right) |V|^{\frac{m}{2}}}, \quad (13)$$

where $\Gamma_n(a)$ denotes the multivariate gamma function $\Gamma_n(a) = \pi^{n(n-1)/4} \prod_{j=1}^n \Gamma(a + (1-j)/2)$. We shall use the notation $V \sim \mathcal{IW}_n(\Psi, m)$. See Definition 3.4.1, p. 111 of Gupta and Nagar (1999).

Since the matrix normal distribution belongs to the exponential family (e.g., see Gupta and Nagar, 1999), it is possible to show that the matrix normal prior distribution for the location parameter of a matrix normal likelihood is conjugate.

Proposition 2. *Let $\mathbf{Y} = \{Y_j$ ($p \times n$), $j = 1, \dots, t-1\}$, be a collection of i.i.d. random matrices from a matrix normal distribution with unknown mean B ($p \times n$) and known covariance $\Sigma_L \otimes V$, that is $Y_j \sim \mathcal{MN}_{p,n}(B, \Sigma_L, V)$ i.i.d., with Σ_L ($p \times p$) > 0 and V ($n \times n$) > 0 . Assume a conjugate matrix variate normal distribution for B , that is $B \sim \mathcal{N}_{p,n}(M, \Sigma_P, V)$. The posterior distribution of B at time t conditioned to the information available at time $t-1$ is the matrix variate normal $B|\mathbf{Y} \sim \mathcal{N}_{p,n}(M_*, \Sigma_*, V)$ with:*

$$M_* = M + \Sigma_P (\boldsymbol{\nu}_T \otimes I_p)' \left[I_T \otimes \Sigma_L + (\boldsymbol{\nu}_T \otimes I_p) \Sigma_P (\boldsymbol{\nu}_T \otimes I_p)' \right]^{-1} (\mathbf{Y} - \boldsymbol{\nu}_T \otimes M) \quad (14)$$

$$\Sigma_* = \Sigma_P - \Sigma_P (\boldsymbol{\nu}_T \otimes I_p)' \left[I_T \otimes \Sigma_L + (\boldsymbol{\nu}_T \otimes I_p) \Sigma_P (\boldsymbol{\nu}_T \otimes I_p)' \right]^{-1} (\boldsymbol{\nu}_T \otimes I_p) \Sigma_P. \quad (15)$$

The parameters Σ_P and M describe the nature and quality of prior information about B . In the following we assume $\Sigma_P = \Sigma_L/\varphi$, with $\varphi > 0$ as commonly done for Gaussian models (e.g., see Zellner, 1986). We assume that the information set at time $T = t-1$ is given by the sigma-algebra \mathcal{D}_{t-1} generated by the elements of \mathbf{Y} and the matrices Σ_P and Σ_L/φ are \mathcal{D}_{t-1} -measurable. We denote with B_t the random variable $B|\mathbf{Y}$. Following the notation in the previous section, B_t corresponds to θ_t and the distribution of B_t corresponds to the posterior distribution $p(\theta_t|\mathcal{D}_{t-1})$. When V is unknown, the Normal-Inverse Wishart prior for B and V is conjugate, as shown below.

Proposition 3. Let $\mathbf{Y} = \{Y_j (p \times n), j = 1, \dots, t-1\}$, be a collection of i.i.d. random matrices from a matrix normal distribution with unknown mean $B (p \times n)$ and covariance $V (n \times n)$ and known covariance $\Sigma_L (p \times p)$. Assume a conjugate matrix normal inverse Wishart distribution for B and V , that is $B|V \sim \mathcal{N}_{p,n}(M, \Sigma_L/\varphi, V/\rho)$ and $V \sim \mathcal{IW}_n(\Psi, m)$. The joint posterior distribution of B and V at time t given the information available at time $t-1$ is the matrix normal inverse Wishart $B|V, \mathbf{Y} \sim \mathcal{N}_{p,n}(M_*, \Sigma_*, V)$ and $V|\mathbf{Y} \sim \mathcal{IW}(\Psi_*, m_*)$ with:

$$\begin{aligned} M_* &= \frac{kM + T\bar{Y}}{k+T}, \quad \Sigma_* = \frac{1}{k_*} \Sigma_L, \quad \Psi_* = \Psi + kT \frac{(M - \bar{Y})' \Sigma_L^{-1} (M - \bar{Y})}{(k+T)} + TS \\ k_* &= k + T, \quad k = \rho\varphi, \quad m_* = m + Tp \end{aligned} \quad (16)$$

where, $\bar{Y} = \frac{1}{T} \sum_{i=1}^T Y_i$ and $S = \frac{1}{T} \sum_{i=1}^T (Y_i - \bar{Y})' \Sigma_L^{-1} (Y_i - \bar{Y})$.

For the unknown V case, the pair B_t and V_t corresponds to θ_t and their joint distribution corresponds to the posterior $p(\theta_t|\mathcal{D}_{t-1})$. For the two conjugate prior assumptions in Prop. 2 and Prop. 3, it is possible to derive the BF.

Proposition 4 (Bayes Factor). *i) Under the assumptions in Prop. 2, we have that $Y_t|\mathcal{D}_{t-1} \sim \mathcal{N}_{p,n}(M_*, \Sigma_d, V)$, and $Y_t|\mathcal{D}_{t-1} \sim \mathcal{N}_{p,n}(M_*, \Sigma_{A,d}, V)$ under the alternative hypothesis and the Bayes Factor H_t is given by*

$$H_t(\alpha_t) = \frac{|\Sigma_{A,d}|^{\frac{n}{2}}}{|\Sigma_d|^{\frac{n}{2}}} \exp \left\{ -\frac{1}{2} \text{tr} \left[\left(\Sigma_d^{-1} - \Sigma_{A,d}^{-1} \right) (Y_t - M_*) V^{-1} (Y_t - M_*)' \right] \right\}, \quad (17)$$

where $\Sigma_d = \Sigma_L + \Sigma_*$, $\Sigma_{A,d} = \Sigma_L + \Sigma_{A,*}$, $\Sigma_{A,*} = \Sigma_*/\alpha_t$ and the posterior parameters M_* and Σ_* are given in Eq. 14 and 15.

ii) Under the assumptions in Prop. 3, we have that $Y_t|\mathcal{D}_{t-1} \sim \mathcal{T}_{p,n}(m_ - 2n, M_*, \Sigma_L, L_*)$ and $Y_t|\mathcal{D}_{t-1} \sim \mathcal{T}_{p,n}(m_{A,*} - 2n, M_*, \Sigma_L, L_{A,*})$ under the alternative hypothesis with $\mathcal{T}_{p,n}(\nu, M, \Sigma, \Omega)$ denoting the matrix Student- t distributions with degrees of freedom parameter $\nu > 0$, location parameter M , and scale parameters $\Sigma > 0$ and $\Omega > 0$, $L_* = \Psi_* k_d/k_*$, and $L_{A,*} = \Psi_{A,*} k_d/k_{A,*}$ (Gupta and Nagar, 1999, def. 4.2.1). The Bayes Factor H_t is given by*

$$H_t(\alpha_t) = G \frac{\left| \Psi_{A,*} + \frac{\alpha_t k_*}{(\alpha_t k_* + 1)} (M_* - Y_t)' \Sigma_L^{-1} (M_* - Y_t) \right|^{\frac{m_{A,d} - n - 1}{2}}}{\left| \Psi_* + k_* \frac{(M_* - Y_t)' \Sigma_L^{-1} (M_* - Y_t)}{(k_* + 1)} \right|^{\frac{m_d - n - 1}{2}}}, \quad (18)$$

where

$$G = \frac{|\Psi_*|^{\frac{m_* - n - 1}{2}} k_*^{\frac{np}{2}} \Gamma_n \left(\frac{m_d - n - 1}{2} \right) k_{A,d}^{\frac{np}{2}} \Gamma_n \left(\frac{m_{A,*} - n - 1}{2} \right)}{k_d^{\frac{np}{2}} \Gamma_n \left(\frac{m_* - n - 1}{2} \right) |\Psi_{A,*}|^{\frac{m_{A,*} - n - 1}{2}} k_{A,*}^{\frac{np}{2}} \Gamma_n \left(\frac{m_{A,d} - n - 1}{2} \right)}$$

and $k_d = k_* + 1$, $k_{A,*} = \alpha_t k_*$, $k_{A,d} = \alpha_t k_* + 1$, $m_d = m_* + p$, $m_{A,*} = \alpha_t (m_* + p) - p$, $m_{A,d} = m_{A,*} + p$, and m_*, M_*, Ψ_* as in Eq. 16.

Remark 2 (Interpretability). In the case V is known, it can be easily shown that the discounting approach provides a tractable prior distribution under the alternative. The approach is equivalent to assuming, under the alternative hypothesis, the following hierarchical prior distribution $B|\tilde{M}, \mathbf{Y} \sim \mathcal{N}_{p,n}(\tilde{M}, \Sigma_*, V)$ with $\tilde{M}|\mathbf{Y} \sim \mathcal{N}_{p,n}(M_*, \Sigma_*(1/\alpha_t - 1), V)$, which is well-defined since $\alpha_t \in (0, 1)$. This can be interpreted as a random perturbation of the posterior distribution, which inflates the variance of the location parameter distribution.

The properties of the normalizing constant of the alternative distribution can be derived from Prop. 4 and are given in the following.

Corollary 1. 1. Under the assumptions in Prop. 2 the inverse normalizing constant is

$$C(\alpha_t) = \alpha_t^{\frac{np}{2}} (2\pi)^{\frac{(\alpha_t-1)np}{2}} |\Sigma_*|^{\frac{(\alpha_t-1)n}{2}} |V|^{\frac{(\alpha_t-1)p}{2}} \quad (19)$$

and satisfies $C(\alpha_t) \rightarrow 0$ as $\alpha_t \rightarrow 0^+$ and $C(\alpha_t) \rightarrow 1$ as $\alpha_t \rightarrow 1^-$.

2. Under the assumptions in Prop. 3 the inverse normalizing constant is

$$C(\alpha_t) = \frac{|\Psi_{A,*}|^{\frac{m_{A,*}-n-1}{2}} (2\pi)^{\frac{\alpha_t np}{2}} |\Sigma_*|^{\frac{\alpha_t n}{2}} 2^{\frac{\alpha_t(m_*-n-1)n}{2}} \Gamma_n\left(\frac{m_*-n-1}{2}\right)^{\alpha_t}}{(2\pi)^{\frac{np}{2}} |\Sigma_{A,*}|^{\frac{n}{2}} 2^{\frac{(m_{A,*}-n-1)n}{2}} \Gamma_n\left(\frac{m_{A,*}-n-1}{2}\right) |\Psi_*|^{\frac{\alpha_t(m_*-n-1)}{2}}} \quad (20)$$

and satisfies $C(\alpha_t) \rightarrow 0$ as $\alpha \rightarrow \underline{\alpha}^+$ and $C(\alpha_t) \rightarrow 1$ as $\alpha_t \rightarrow 1^-$, where $\underline{\alpha} = (p+2n)/m_d$.

Remark 3 (Univariate Gaussian model). Let $Y_j \sim \mathcal{N}(\theta, \sigma^2)$ i.i.d. for $j = 1, \dots, t-1$ and assume a conjugate prior for θ , that is $\theta \sim \mathcal{N}(m, \sigma^2/\varphi)$ with $\varphi > 0$. It is easy to show that the predictive under the alternative is given by $p_A(y_t|\mathcal{D}_{t-1}) = (2\pi\sigma_*^2/\alpha_t)^{1/2} \exp\{-(\theta - m_*)^2\alpha_t/2\sigma_*^2\}$ with $m_* = \varphi m/(\varphi + t - 1) + (t-1)/(\varphi + t - 1)\bar{y}$ with $\bar{y} = (t-1)^{-1} \sum_{j=1}^{t-1} y_j$, $\sigma_*^2 = \sigma^2/(\varphi + t - 1)$ and the normalising constant is $C(\alpha_t) = (2\pi\sigma_*^2)^{(\alpha_t-1)/2} \alpha_t^{1/2}$. It follows that the BF is

$$H_t(\alpha_t) = \kappa_t(\alpha_t) \exp\{(y_t - m_*)^2 A_t^{-1}\}, \quad (21)$$

where

$$A_t = \frac{2\sigma_*^2(\varphi + t)(\alpha_t(\varphi + t - 1) + 1)}{(\alpha_t - 1)(\varphi + t - 1)} \quad \text{and} \quad \kappa_t(\alpha_t) = \left(\frac{\alpha_t(\varphi + t - 1) + 1}{\alpha_t(\varphi + t)}\right)^{1/2} \quad (22)$$

naturally represents an upper bound of $H_t(\alpha_t)$. Intuitively for small values of α_t the null hypothesis is not rejected, that is $H_t(\alpha_t) \rightarrow +\infty$ as $\alpha_t \rightarrow 0^+$, whereas for large values, the BF gets closer to one, that is $H_t(\alpha_t) \rightarrow 1$ as $\alpha_t \rightarrow 1^-$. Its derivative $H_t'(\alpha_t) = 0$ for α_t is equal to

$$\alpha_{t,0} = ((y_t - m_*)^2/\sigma_*^2 - 1)^{-1}(\varphi + t - 1)^{-1} \quad (23)$$

provided $(y_t - m_*)^2/\sigma^2 - 1 > (\varphi + t - 1)^{-1}$. The conditions in Prop. 1 are thus satisfied, and the rejection of the null hypothesis depends on the values of α_t .

3.2 Bayes Factor Properties

Proposition 5 (Bayes Factor properties, known V). Under the assumptions in Prop. 2, the BF $H_t(\alpha_t)$ satisfies the following properties.

- i) There exists a function $\kappa_t(\alpha_t) = |\Sigma_{A,d}|^{\frac{n}{2}} |\Sigma_d|^{-\frac{n}{2}} \geq 1$ which does not depend on Y_t such that $H_t(\alpha_t) \leq \kappa_t(\alpha_t)$, $\kappa_t \rightarrow 1$ as $\alpha_t \rightarrow 1^-$, $\kappa_t \rightarrow \infty$ as $\alpha_t \rightarrow 0^+$ and κ_t is decreasing in α_t ;
- ii) $H_t(\alpha_t) \rightarrow 1$ as $\alpha_t \rightarrow 1^-$ and $H_t(\alpha_t) \rightarrow \infty$ as $\alpha_t \rightarrow 0^+$;
- iii) $\partial_{\alpha_t} H_t(\alpha_t) = H_t(\alpha_t) (2\alpha_t)^{-1} \text{tr} \left(-n\tilde{B}^{-1}\Sigma_* + \alpha_t\Upsilon(\alpha_t)\tilde{A} + \alpha_t(1 - \alpha_t)\Upsilon(\alpha_t)\Sigma_L\tilde{B}^{-1}\tilde{A} \right)$ where we defined $\tilde{A} = (Y_t - M_*)V^{-1}(Y_t - M_*)'$, $\tilde{B} = (\alpha_t\Sigma_L + \Sigma_*)$ and $\Upsilon(\alpha_t) = (\Sigma_L + \Sigma_*)^{-1}\Sigma_*\tilde{B}^{-1}$.

The derivative of the BF, given in *iii*) of the proposition, is not tractable; thus, it is not easy to find analytically a stationary point. Nevertheless, we shall notice that the second and third elements within the trace are positive. In contrast, the first one is negative and, for each Y_t , $\partial_{\alpha_t} H_t(\alpha_t) \rightarrow -\infty$ as $\alpha_t \rightarrow 0^+$ and $\partial_{\alpha_t} H_t(\alpha_t) \rightarrow (-n \text{tr}((\Sigma_L + \Sigma_*)^{-1}) + \text{tr}(\Upsilon(1)\tilde{A})) / 2$ as $\alpha_t \rightarrow 1^-$. Thus there is a change of the sign, provided $\text{tr}((\Sigma_L + \Sigma_*)^{-1} \Sigma_* ((\Sigma_L + \Sigma_*)^{-1} \tilde{A} - nI)) > 0$, and the BF has at least one stationary point.

Proposition 6 (Bayes Factor properties, unknown V). *Under the assumptions in Prop. 3, the BF $H_t(\alpha_t)$ satisfies the following properties.*

- i)* There exists a function $\kappa_t(\alpha_t) = (k_* k_{A,d} (k_d k_{A,*}))^{-np/2} \Gamma_n((m_d - n - 1)/2) \Gamma_n((m_{A,*} - n - 1)/2) (\Gamma_n((m_* - n - 1)/2) \Gamma_n((m_{A,d} - n - 1)/2))^{-1}$ which does not depend on Y_t such that $H_t(\alpha_t) \leq \kappa_t(\alpha_t)$, κ_t is decreasing in α_t , $\kappa_t \rightarrow 1$ as $\alpha_t \rightarrow 1^-$, $\kappa_t \rightarrow +\infty$ as $\alpha_t \rightarrow \underline{\alpha}^+$ where $\underline{\alpha} = (2n + p)/m_d$;
- ii)* $H_t(\alpha_t) \rightarrow 1$ as $\alpha_t \rightarrow 1^-$ and $H_t(\alpha_t) \rightarrow \infty$ as $\alpha_t \rightarrow \underline{\alpha}^+$;
- iii)* $\partial_{\alpha_t} H_t(\alpha_t) = L (|\Psi_{A,*}|^{(m_{A,*} - n - 1)/2} k_{A,*}^{np/2} \Gamma_n((m_{A,d} - n - 1)/2))^{-2} (B_1 \partial_{\alpha_t} A_1 - A_1 \partial_{\alpha_t} B_1)$, where $L = |\Psi|_*^{(m_* - n - 1)/2} k_*^{np/2} \Gamma_n((m_d - n - 1)/2) (k_d^{np/2} |\Psi_d|^{(m_d - n - 1)/2} \Gamma_n((m_* - n - 1)/2))^{-1}$, $A_1 = a_1 a_2 a_3$, $B_1 = b_1 b_2 b_3$ with $a_1 = k_{A,d}^{np/2}$, $a_2 = \Gamma_n((m_{A,*} - n - 1)/2)$, $a_3 = |\Psi_{A,d}|^{(m_{A,d} - n - 1)/2}$, $b_1 = k_{A,*}^{np/2}$, $b_2 = \Gamma_n((m_{A,d} - n - 1)/2)$, $b_3 = |\Psi_{A,*}|^{(m_{A,*} - n - 1)/2}$.

In the derivative $\partial_{\alpha_t} H_t(\alpha_t)$, the terms $a_i, b_i, i = 1, \dots, 3$ are positive, and the terms $\partial_{\alpha_t} A_1$ and $\partial_{\alpha_t} B_1$ can take both positive and negative values since they are sums of digamma functions. Thus, for some values of α_t , the $\partial_{\alpha_t} H_t(\alpha_t)$ can go to zero, and the BF has at least one stationary point.

The following remark discusses the relationship with the univariate case, whereas the empirical section will provide further illustrations for the matrix case.

Remark 4 (Univariate Gaussian model). The univariate monitoring BF and its upper bound $\kappa_t(\alpha_t)$ given in Remark 3 can be easily obtained setting $n = p = 1$ and $\Sigma_P = \sigma^2/\varphi$ in Eq. 17, in Prop. 4 and in *i*) of Prop. 5. Since A_t in Remark 3 is strictly negative, then the upper bound satisfies the properties in Prop. 5. The derivative of the BF is

$$\partial_{\alpha_t} H_t(\alpha_t) = \frac{H_t(\alpha_t)}{2} \left(\frac{(\varphi + t - 1)(y_t - m_*)^2}{\sigma^2(\alpha_t(\varphi + t - 1) + 1)^2} - \kappa_t(\alpha_t) \frac{1}{\alpha_t^2(\varphi + t)} \right) \quad (24)$$

which goes to $-\infty$ as $\alpha_t \rightarrow 0^+$ and is strictly positive for $\alpha_t > 1/((\varphi + t - 1)((y_t - m_*)^2/\sigma^2 - 1))$ provided that $(y_t - m_*)^2/\sigma^2 > (\varphi + t)/(\varphi + t - 1)$. This implies there exists a stationary point and that $H_t < 1$ for some values of α_t provided the conditions in Prop. 1 are satisfied.

Let us now find the subset of the sample space such that for a given threshold $0 < h_0(\alpha_t) < \kappa_t(\alpha_t)$, the BF leads us to accept the null hypothesis, that is, $H_t \geq h_0(\alpha_t)$. Define $\mathbf{y}_t = \text{vec}(Y_t)$ and $\mathbf{m}_* = \text{vec}(M_*)$ and let $\kappa_t(\alpha_t)$ be the upper bound given in Prop. 5. From the expression of the BF given in Prop. 4 the condition $H_t \geq h_0(\alpha_t)$ is satisfied for \mathbf{y}_t in the ellipsoid:

$$(\mathbf{y}_t - \mathbf{m}_*)' (\Sigma_H \otimes V)^{-1} (\mathbf{y}_t - \mathbf{m}_*) \leq 2 \log \left(\frac{\kappa_t(\alpha_t)}{h_0(\alpha_t)} \right) \quad (25)$$

centered in \mathbf{m}_* , with axis in the direction of the eigenvector ν_k of $\Sigma_H \otimes V$, where $\Sigma_H = (1 - \alpha_t)^{-1} (\alpha_t \Sigma_L + \Sigma_*) \Sigma_*^{-1} (\Sigma_L + \Sigma_*)$. The length of the ellipsoid axes along the ν_k eigenvector's direction is:

$$\ell_k = 2 \sqrt{2 \log \left(\frac{\kappa_t(\alpha_t)}{h_0(\alpha_t)} \right)} \xi_k, \quad (26)$$




Figure 2: Bayes Factor Critical regions. Left: increasing α_t from 0.2 (— $h_0 = 0.8$, - - - $h_0 = 1.1$) to 0.6 (— $h_0 = 0.8$, - - - $h_0 = 1.1$). Right: increasing the threshold h_0 from 0.8 (—) to 1.1 (—).

where ξ_k is the corresponding eigenvalue of $\Sigma_H \otimes V$. Let γ_i and ζ_i be the eigenvalues and the corresponding eigenvectors of Σ_H and τ_j and δ_j the eigenvalues and eigenvectors of V . Then $\Sigma_H \otimes V$ has eigenvalues $\xi_k = \gamma_i \tau_j$ with corresponding eigenvectors $\nu_k = \zeta_i \otimes \delta_j$. If we consider $h_0 = 1$, then Y_t is not considered an outlier for $H_t \geq 1$. Figure 2 provides a numerical illustration for $p = 2$ and $n = 1$. The left plot shows that increasing the discounting parameter value reduces the evidence in favor of the null for $h_0 > 1$ (dashed lines) and increases it for $h_0 < 1$ (solid lines). The right plot shows that for increasing value of h_0 , the evidence against the null becomes stronger. From Eq. 25, we obtain the following result for the univariate monitoring.

Remark 5 (Univariate Gaussian model). In the univariate case, the null hypothesis is accepted, i.e. $H_t > h_0$, for values of y_t in the interval (y_{1t}, y_{2t}) with

$$y_{jt} = m_* + (-1)^j \sqrt{\log \left(\frac{h_0(\alpha_t)}{\kappa_t(\alpha_t)} \right) \frac{2\sigma^2(\varphi + t)(\alpha_t(\varphi + t - 1) + 1)}{(\alpha_t - 1)(\varphi + t - 1)}}. \quad (27)$$

The BF and the outcome of the testing procedure depend on the choice of the discounting parameter α_t . In this paper, we propose the integrated BF and its normalized version as a solution, which requires the integral with respect to a probability measure $\pi_t(\alpha_t)$ is bounded. In the following, we provide sufficient conditions for the existence of the IBF and NIBF.

Proposition 7 (Integrability). *Assume a general beta distribution $\mathcal{B}e(a, b)$ truncated on the interval $(\underline{\alpha}, \bar{\alpha})$. The IBF, $H_t = \int_0^1 H_t(\alpha) \pi_t(\alpha) d\alpha$, satisfies the following properties.*

1. Under the assumptions in Prop. 2, for $a > np/2$, $b > 0$, $\underline{\alpha} = 0$ and $\bar{\alpha} = 1$

$$H_t < \sum_{\ell=0}^{\infty} \sum_{\underline{k} \in \mathcal{K}_\ell} d_{\ell, \underline{k}} \bar{c}_{\underline{k}} \frac{\Gamma(n/2 + 1) B(a - np/2 + w_{\underline{k}}, b)}{\Gamma(\ell + 1) \Gamma(n/2 - \ell) B(a, b)} < \infty \quad (28)$$

with $\underline{k} = (k_0, \dots, k_p)$, $\mathcal{K}_\ell = \{\underline{k} | k_0 + \dots, k_p = \ell\}$, $\bar{c}_0 = \bar{c}_0 - 1$, $\bar{c}_j = \bar{c}_j$, $j = 1, \dots, p$, and

$$d_{\ell, \underline{k}} = \binom{\ell}{k_0, k_1, k_2, \dots, k_p}, \quad \bar{c}_{\underline{k}} = \prod_{j=0}^p \bar{c}_j^{k_j},$$




Figure 3: Bayes Factor distribution, $F_{j,t}(h|\alpha_t)$, under the null hypothesis ($j = 0$) (—) and the alternative hypothesis ($j = 1$) (---), for different values of α_t .

where $d_{\ell, \underline{k}}$ denotes the multinomial coefficient, \tilde{c}_j $j = 0, \dots, p$ are the coefficients of the characteristic polynomial of $\Sigma_L^{-1} \Sigma_*$.

2. Under the assumption in Prop. 3, for $a > 1$ and $b > 0$, $\underline{\alpha} = (2n + p)/m_d$ and $\bar{\alpha} = 1$.

$$H_t < (\underline{\alpha} k_* / (\underline{\alpha} (k_* + 1)))^{np/2} \Gamma\left(\frac{p}{2}\right)^{-n} \left(\frac{4}{p}\right)^n \frac{B(1 - \underline{\alpha}; ra - r + 1, qb - q + 1)}{B(1 - \underline{\alpha}; a, b)} \xi(r, q) \quad (29)$$

where $B(c, a, b) = \int_0^c x^{a-1} (1-x)^{b-1} dx$ denotes the incomplete beta function, $\xi(r, q) = (((m_d(1 - \underline{\alpha}) + p)^{1-qn} - p^{1-qn}) / (m_d(1 - qn)))^{1/q}$ and $q = r / (r - 1)$ with $r > 1$.

In the empirical illustration, we will also show that the choice of the hyper-parameters, e.g. b , can still affect the outcome of the hypothesis testing, which calls for calibrated BFs. As stated in the following proposition, the integral of the upper bound for the univariate case can be derived analytically and is well-defined under the assumption of standard uniform distribution for the discounting parameter.

Corollary 2 (IBF in the univariate model). *The integral of the BF H_t given in Remark 3 is bounded and satisfies:*

$$\int_0^1 H_t(\alpha_t) d\alpha_t - 1 \leq \frac{1}{2\sqrt{(\varphi + t - 1)(\varphi + t)}} \log \left(\frac{\sqrt{(\varphi + t)} + \sqrt{(\varphi + t - 1)}}{\sqrt{(\varphi + t)} - \sqrt{(\varphi + t - 1)}} \right). \quad (30)$$

One can assume the variable \mathbf{y}_t in the predictive BF is not observed at time t . Thus, the predictive BF is random, and we prove that its distribution is a mixture of gamma distributions, as stated in the following. Controlling for the test's size and power and using the BF distribution allow us to derive a calibrated BF and a suitable critical region for the test.

Proposition 8 (Distribution of the BF, known V). *Assume $Y_t | \mathcal{D}_{t-1} \sim \mathcal{N}_{p,n}(\tilde{M}, \tilde{\Sigma}, V)$, let $\lambda_1, \dots, \lambda_p$ be the eigenvalues of the covariance matrix $(\Sigma_H^{-1/2})' \tilde{\Sigma} \Sigma_H^{-1/2}$ with $\Sigma_H = (1 - \alpha_t)^{-1} (\alpha_t \tilde{\Sigma} + \Sigma_*) \Sigma_*^{-1} (\tilde{\Sigma} + \Sigma_*)$ and Q the associated eigenvector matrix. Under the assumptions in Prop. 2, the distribution of the BF H_t defined in Eq. 17 has pdf and cdf:*

$$f_{H_t}(h|\alpha_t) = \frac{2}{h} \sum_{k=0}^{\infty} c_k g(-2 \log(h/\kappa_t); \frac{np}{2} + k, 2\lambda), \quad (31)$$

$$F_{H_t}(h|\alpha_t) = \sum_{k=0}^{\infty} c_k (1 - G(-2 \log(h/\kappa_t); \frac{np}{2} + k, 2\lambda)), \quad (32)$$

respectively, with support $0 < h \leq \kappa_t$, where $\kappa_t(\alpha_t)$ is defined in Prop. 5, the parameter $0 < \lambda < \infty$ is arbitrarily chosen, $g(x; a, b)$ and $G(x; a, b)$ are the pdf and cdf of a gamma distribution with shape and scale parameters $a > 0$ and $b > 0$, respectively, and the coefficients c_k , d_k and f_k satisfy

$$c_k = \exp\left(-\sum_{j=1}^p U_{jj}\right) \prod_{j=1}^p (\lambda_j/\lambda)^{-n/2} f_k, \quad d_k = \frac{n}{2k} \sum_{j=1}^p (1 - \lambda/\lambda_j)^k + \lambda \sum_{j=1}^p \frac{U_{jj}}{\lambda_j} (1 - \lambda/\lambda_j)^{k-1},$$

$$f_{k+1} = \frac{1}{k+1} \sum_{j=1}^{k+1} j d_j f_{k+1-j}, \quad k = 0, 1, 2, \dots, \quad \text{with } f_0 = 1,$$

where U_{jj} are the (j, j) -th entry of $U = Q'(\Sigma_H^{1/2} \tilde{\Sigma}^{-1}(\tilde{M} - M_*)V^{-1})(\tilde{M} - M_*)'\Sigma_H^{-1/2}Q$.

If in the previous proposition, we set $\tilde{M} = M_*$ and $\tilde{\Sigma} = \Sigma_d$, that is the location and scale of the marginal likelihood under the null (from i) in Prop. 4) we denote the distribution with $F_{0,t}(h)$. In contrast, if $\tilde{M} = M_*$ and $\tilde{\Sigma} = \Sigma_{A,d}$, that is the location and scale of the marginal likelihood under the alternative (from i) in Prop. 4), we denote the distribution with $F_{1,t}(h)$. Figure 3 shows the sensitivity of the BF distribution to the perturbation parameter α_t , under the null and the alternative hypothesis (dashed and solid line). Increasing α_t from 0.05 to 0.67, the dispersion of the two distributions decreases, and their distance reduces.

From the properties of the gamma distribution, $f_{H_t}(h|\alpha_t)$ is continuous at 0 and κ_t . As stated in the following, the result of Prop. 8 provides the BF distribution for the univariate Gaussian model given in previous studies (e.g., see Weiss, 1997; De Santis, 2004; Pawel and Held, 2025).

Corollary 3 (Univariate Gaussian model). *In the univariate case, i.e. $n = p = 1$, following the notation in Remark 3, we set $\Sigma = \sigma^2$, $V = 1$ and $B = \theta$. Assuming $\tilde{M} = B$ and $\tilde{\Sigma} = \Sigma_L$ in Prop. 8 the distribution of the BF is*

$$F_{H_t}(h|\alpha_t) = 1 - \Phi(\sqrt{-2 \log(h/\kappa_t)\sigma_H^2/\sigma_L^2} - \sqrt{\gamma}) + \Phi(-\sqrt{-2 \log(h/\kappa_t)\sigma_H^2/\sigma_L^2} - \sqrt{\gamma}), \quad (33)$$

where $\gamma = (\theta - \tilde{m}_*)^2/\sigma^2$, the variances $\sigma_H^2 = \Sigma_H$ and $\sigma_L^2 = \Sigma_L$ are given in Prop. 4, the bound κ_t is given in Remark 3 and $\Phi(\cdot)$ is the cdf of a standard normal distribution.

Our perturbation framework for matrix-variate observations is a form of global regularization affecting all entries within the matrix. Later in this paper, we will explore the sensitivity of the testing procedure in response to variations in the patterns, proportion and magnitude of outliers. As an extension, local contamination frameworks can be devised to detect patterns within the outliers and potential dependencies among outlying observations. In the context of matrices, a multiplicative perturbation can be employed to define $p_A(Y_t|\mathcal{D}_t)$, assuming, for instance, the perturbed normal distribution $\mathcal{N}_{p,n}(M_*, A_1 \Sigma_d A_1, A_2 V A_2)$, where A_1 and A_2 are two diagonal matrices scaling along the rows and columns with varying levels. An additive perturbation would lead to a distribution like $\mathcal{N}_{p,n}(M_*, \Sigma_d + A_1, V + A_2)$, with A_1 and A_2 shifting the posterior's covariance matrix diagonal. On the one hand, local contamination allows for patterns among outliers; however, it requires the identification of the patterns, which calls for inference on the contamination parameters. Therefore, alternative approaches, such as multiple-outlier models presented in Page and Dunson (2011); Tomarchio et al. (2022), might be preferable. This typically involves introducing appropriate prior specifications and has the drawback of requiring computationally intensive procedures for posterior approximation. We will postpone this extension to future research.




Figure 4: Simulated dataset. Top: the BFs $H_t(\alpha_t)$ and their upper bounds $\kappa_t(\alpha_t)$ as a function of α_t for different samples (left, —, —); the BF lower and upper thresholds $\underline{h}(\alpha_t)$ and $\bar{h}(\alpha_t)$ (middle, —) as a function of α_t and the BFs for the calibrated discount factor (right, •) together with the calibrated lower and upper bounds $\underline{h}(\alpha^*)$ and $\bar{h}(\alpha^*)$ (---). Bottom: Minimum BF (left), Integrated BF (middle) and Normalized Integrated BF (right) for different beta distribution settings (—, - - -). In all plots, the reference lines at 1 (---) and $10^{-1/2}$ (- - -).

3.3 Simulation Study

In the following simulation exercise, we generate data with outliers and study some finite-sample properties of the decision procedures based on standard BF, MBF, IBF and NIBF, and the calibrated predictive BF approach we introduced. All analyses were implemented in Matlab 2023a and carried out on a 12-core computing system with 128GB RAM. The parameter values have been randomly generated as follows: $M \sim \mathcal{N}_{p,n}(O_{p \times n}, I_p, I_n)$, $\Sigma = SS'$, $S \sim \mathcal{N}_{p,p}(O_{p \times p}, I_p, I_p)$, $\Psi = GG'$, $G \sim \mathcal{N}_{n,n}(O_{n \times n}, I_n, I_n)$. We consider two scenarios: the case of moderate-size observation matrix, denoted as $Case_1$, with $p = 30$ and $n = 10$, and the case of a large-size observation matrix, $Case_2$, with $p = 50$ and $n = 50$. We generate the synthetic data with an outlier as follows. First a noise sequence is generated, $E_t \sim \mathcal{N}_{p,n}(O_{p \times p}, \Sigma, \Psi)$ i.i.d. $t = 1, \dots, 100$. Secondly, the observable sequence is defined as $X_t = M + E_t$ for $t \neq 80$ and $X_t = M + uR_t + E_t$ for $t = 80$, where R_t is a binary matrix encoding the position of the outliers and u is 0.5, 1, 1.5, 3, 5 and 15 that are 1/30, 1/15, 1/10, 1/5, 1/3 and 1 average standard deviation of the matrix normal.

Our simulation study considered various settings obtained by varying the configurations of the matrix R_t and the magnitude of the outliers. For each setting J independent datasets have been generated $X_t^{(j)}$, $t = 1, \dots, T$ for $j = 1, \dots, J$, with $T = 100$ and the BFs $H_t^{(j)}$ computed. The probabilities $p_I = P(H_t > \bar{h})$, $p_{II} = P(H_t < \underline{h})$ and $p_{III} = P(\bar{h} < H_t < \underline{h})$ have been estimated as follows

$$p_I^J = \frac{1}{J} \sum_{j=1}^J \mathbb{I}(H_t^{(n)} > \bar{h}), \quad p_{II}^J = \frac{1}{J} \sum_{j=1}^J \mathbb{I}(\underline{h} < H_t^{(n)} < \bar{h}), \quad p_{III}^J = \frac{1}{J} \sum_{j=1}^J \mathbb{I}(H_t^{(n)} < \underline{h})$$

under the null hypothesis of the absence of outliers at $t \neq 80$ and under the alternative hypothesis of a certain number of outlying observations in the observation matrix at $t = 80$. The results for $Case_1$ are given in Tab. 2, whereas the ones for $Case_2$ are given in Tab. 3. The results are obtained with $J = 100$ experiments for the two outlier settings. The two settings differ for the positions of the outliers within the matrix R_t : i) row and column patterns in the positions, panel (a); and ii) completely random positions, panel (b).

Figure 4 illustrates the result of the BF procedure on one of the simulated datasets for $u = 0.5$ (top left panel a in Tab. 2). In the BF procedure the calibrated value of α_t for $\tau = 0.01$ and $\beta = 0.8$ is $\alpha^* = 0.750$ for all t and the inconclusive interval C_H has lower and upper bounds $\underline{h}(\alpha^*) = 0.839$ $\bar{h}(\alpha^*) = 1.161$, respectively. Starting from the left in the first row, we notice that the observations with a change in the convexity of $H_t(\alpha_t)$ are recognized as outliers based on the value of α_t . Notably, for the outlier observation introduced in the simulation, the BF is consistently convex and remains far below 1. The second plot illustrates the behaviour of the thresholds $\underline{h}(\alpha_t)$ and $\bar{h}(\alpha_t)$ defined at the end of Section 2 for given power and size values. The length of the randomized decision interval C_h (vertical axis) reduces as α_t goes to 1. The third plot presents the BF and the inconclusive region for $\alpha^* = 0.750$. The BF of the outlying observation is located outside the inconclusive interval, whereas about 16 observations exhibit a BF below 1 within the inconclusive interval. Moving to the second row, the Minimum BF suggests positive evidence against the null hypothesis of the absence of outliers ($MBF < 10^{-1/2}$ following Jeffrey's scale of evidence) for three observations. The IBF and NIBF (solid line in the second and third plot at the bottom) provide strong evidence against the null ($IBF, NIBF < 10^{-1/2} - 1 \approx -0.687$) only for the 80th observation and barely worth mentioning evidence ($IBF, NIBF > 10^{-1/2} - 1 \approx -0.687$) for the other observations. Nevertheless, comparing the solid and dashed lines shows that the outcome of the IBF and NIBF procedures strongly depends on the choice of the hyperparameters of the beta distribution. Thus, the third robust method should be applied, exploiting the BF's sampling variability to find the calibrated α_t and the reference thresholds.

In all the experiments, when data are generated under the null, the type I error probability $P(H_t < \underline{h} | \mathcal{H}_0)$ is about 2%, whereas when data are generated under the alternative hypothesis (first column), the power $P(H_t < \underline{h} | \mathcal{H}_1)$ of the test gets close to one, increasing the number of outliers (e.g., see columns of the panels (b) in Tab. 2). As we can expect, the convergence is faster for larger number of observations (see panels (b) in Tab. 3). Also, the effective size and power may depend on the position of outliers in the rows or columns of the matrix R_t . The presence of patterns in the position of the outliers reduces the power compared to the case of a completely random position within the matrix R_t (compare columns 20×10 and 200 in panels (a) and (b), respectively). The power decreases below 80% for small outlier amplitude (e.g. 1/10 standard deviation) and a small number of outlying observations (e.g. 10 out of 300 elements) within the matrix. When all entries are outliers, the power of the test converges to 1, increasing the outliers' magnitude (different rows in the last column in Tab. 2).

| | \mathcal{H}_0 | \mathcal{H}_1 | | | | | | | | | | | | | |
|------------------------------------|-----------------|--|---------------|--------------|--------------|---------------|---------------|----------------|----------------|--|------|------|------|------|------|
| | | (a) Patterns in the binary matrix $R_t \in \mathbb{R}^{30 \times 10}$ (number of random rows \times number of random columns) | | | | | | | | (b) Random Positions within $R_t \in \mathbb{R}^{30 \times 10}$ (number of randomly selected entries) | | | | | |
| | | 3×2 | 1×10 | 7×5 | 9×7 | 11×9 | 15×9 | 20×10 | 25×10 | 50 | 100 | 150 | 200 | 250 | 300 |
| | | One thirtieth (1/30) of the standard deviation, i.e. $u = 0.5$ in $X_t = M + uR_t + E_t$ for $t = 80$ | | | | | | | | | | | | | |
| $P(H_t > \bar{h})$ | 0.05 | 0.06 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| $P(\underline{h} < H_t < \bar{h})$ | 0.93 | 0.72 | 0.76 | 0.6 | 0.58 | 0.52 | 0.3 | 0.3 | 0.3 | 0.44 | 0.32 | 0.32 | 0.3 | 0.24 | 0.22 |
| $P(H_t < \underline{h})$ | 0.02 | 0.22 | 0.24 | 0.4 | 0.42 | 0.48 | 0.7 | 0.7 | 0.7 | 0.56 | 0.68 | 0.68 | 0.7 | 0.76 | 0.78 |
| | | One fifteenth (1/15) of the standard deviation, i.e. $u = 1$ in $X_t = M + uR_t + E_t$ for $t = 80$ | | | | | | | | | | | | | |
| $P(H_t > \bar{h})$ | 0.05 | 0.01 | 0 | 0 | 0.01 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| $P(\underline{h} < H_t < \bar{h})$ | 0.93 | 0.55 | 0.35 | 0.24 | 0.18 | 0.09 | 0.1 | 0.04 | 0.04 | 0.14 | 0.02 | 0.01 | 0.01 | 0.02 | 0.02 |
| $P(H_t < \underline{h})$ | 0.02 | 0.44 | 0.65 | 0.76 | 0.81 | 0.91 | 0.9 | 0.96 | 0.96 | 0.86 | 0.98 | 0.99 | 0.99 | 0.98 | 0.98 |
| | | One tenth (1/10) of the standard deviation, i.e. $u = 1.5$ in $X_t = M + uR_t + E_t$ for $t = 80$ | | | | | | | | | | | | | |
| $P(H_t > \bar{h})$ | 0.05 | 0.06 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| $P(\underline{h} < H_t < \bar{h})$ | 0.93 | 0.36 | 0.26 | 0.09 | 0.07 | 0.09 | 0.05 | 0.03 | 0.02 | 0.04 | 0.01 | 0 | 0 | 0 | 0 |
| $P(H_t < \underline{h})$ | 0.02 | 0.63 | 0.74 | 0.91 | 0.93 | 0.91 | 0.95 | 0.97 | 0.98 | 0.96 | 0.99 | 1 | 1 | 1 | 1 |
| | | One fifth (1/5) of the standard deviation, i.e. $u = 3$ in $X_t = M + uR_t + E_t$ for $t = 80$ | | | | | | | | | | | | | |
| $P(H_t > \bar{h})$ | 0.05 | 0.06 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| $P(\underline{h} < H_t < \bar{h})$ | 0.93 | 0.22 | 0.1 | 0.05 | 0.02 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| $P(H_t < \underline{h})$ | 0.02 | 0.78 | 0.9 | 0.95 | 0.98 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| | | One tenth (1/3) of the standard deviation, i.e. $u = 5$ in $X_t = M + uR_t + E_t$ for $t = 80$ | | | | | | | | | | | | | |
| $P(H_t > \bar{h})$ | 0.05 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| $P(\underline{h} < H_t < \bar{h})$ | 0.93 | 0.08 | 0.02 | 0.02 | 0 | 0.01 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| $P(H_t < \underline{h})$ | 0.02 | 0.92 | 0.98 | 0.98 | 1 | 0.99 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| | | One (1) standard deviation, i.e. $u = 15$ in $X_t = M + uR_t + E_t$ for $t = 80$ | | | | | | | | | | | | | |
| $P(H_t > \bar{h})$ | 0.05 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| $P(\underline{h} < H_t < \bar{h})$ | 0.93 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| $P(H_t < \underline{h})$ | 0.02 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |

Table 2: **Moderate-size observation matrix.** The observable sequence is defined as $X_t = M + E_t$ for $t \neq 80$ and $X_t = M + uR_t + E_t$ for $t = 80$, with $X_t, M, R_t, E_t \in \mathbb{R}^{30 \times 10}$, where u is the magnitude of the outlier and R_t is a random binary matrix. Note: the position of the outliers in entries of R_t can follow row and column patterns ($r \times c$ in (a) with r and c the given number of rows and columns, respectively, selected randomly) or can be completely random (r in (b) with r the given total number of outlying entries selected randomly).

| | \mathcal{H}_0 | \mathcal{H}_1 | | | | | | | | | | | | | |
|------------------------------------|-----------------|--|---------------|--------------|--------------|---------------|---------------|----------------|----------------|--|------|------|------|------|-----|
| | | (a) Patterns in the binary matrix $R_t \in \mathbb{R}^{50 \times 50}$ (number of random rows \times number of random columns) | | | | | | | | (b) Random Positions within $R_t \in \mathbb{R}^{50 \times 50}$ (number of randomly selected entries) | | | | | |
| | | 3×2 | 1×10 | 7×5 | 9×7 | 11×9 | 15×9 | 20×10 | 25×10 | 50 | 100 | 150 | 200 | 250 | 300 |
| | | One thirtieth (1/30) of the standard deviation, i.e. $u = 0.5$ in $X_t = M + uR_t + E_t$ for $t = 80$ | | | | | | | | | | | | | |
| $P(H_t > h)$ | 0.05 | 0.03 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| $P(\underline{h} < H_t < \bar{h})$ | 0.93 | 0.57 | 0.45 | 0.35 | 0.15 | 0.12 | 0.09 | 0.06 | 0.06 | 0.09 | 0.02 | 0.04 | 0.01 | 0.01 | 0 |
| $P(H_t < \underline{h})$ | 0.02 | 0.4 | 0.55 | 0.65 | 0.85 | 0.88 | 0.91 | 0.94 | 0.94 | 0.91 | 0.98 | 0.96 | 0.99 | 0.99 | 1 |
| | | One fifteenth (1/15) of the standard deviation, i.e. $u = 1$ in $X_t = M + uR_t + E_t$ for $t = 80$ | | | | | | | | | | | | | |
| $P(H_t > h)$ | 0.05 | 0.02 | 0.01 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| $P(\underline{h} < H_t < \bar{h})$ | 0.93 | 0.33 | 0.23 | 0.11 | 0.06 | 0.03 | 0.01 | 0 | 0 | 0.01 | 0 | 0 | 0 | 0 | 0 |
| $P(H_t < \underline{h})$ | 0.02 | 0.65 | 0.76 | 0.89 | 0.94 | 0.97 | 0.99 | 1 | 1 | 0.99 | 1 | 1 | 1 | 1 | 1 |
| | | One tenth (1/10) of the standard deviation, i.e. $u = 1.5$ in $X_t = M + uR_t + E_t$ for $t = 80$ | | | | | | | | | | | | | |
| $P(H_t > h)$ | 0.05 | 0 | 0.01 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| $P(\underline{h} < H_t < \bar{h})$ | 0.93 | 0.23 | 0.16 | 0.04 | 0.01 | 0.02 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| $P(H_t < \underline{h})$ | 0.02 | 0.77 | 0.83 | 0.96 | 0.99 | 0.98 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| | | One fifth (1/5) of the standard deviation, i.e. $u = 3$ in $X_t = M + uR_t + E_t$ for $t = 80$ | | | | | | | | | | | | | |
| $P(H_t > h)$ | 0.05 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| $P(\underline{h} < H_t < \bar{h})$ | 0.93 | 0.08 | 0.04 | 0.01 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| $P(H_t < \underline{h})$ | 0.02 | 0.92 | 0.96 | 0.99 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| | | One tenth (1/3) of the standard deviation, i.e. $u = 5$ in $X_t = M + uR_t + E_t$ for $t = 80$ | | | | | | | | | | | | | |
| $P(H_t > h)$ | 0.05 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| $P(\underline{h} < H_t < \bar{h})$ | 0.93 | 0.02 | 0.01 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| $P(H_t < \underline{h})$ | 0.02 | 0.98 | 0.99 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| | | One (1) standard deviation, i.e. $u = 15$ in $X_t = M + uR_t + E_t$ for $t = 80$ | | | | | | | | | | | | | |
| $P(H_t > h)$ | 0.05 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| $P(\underline{h} < H_t < \bar{h})$ | 0.93 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| $P(H_t < \underline{h})$ | 0.02 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |

Table 3: **Large-size observation matrix.** The observable sequence is defined as $X_t = M + E_t$ for $t \neq 80$ and $X_t = M + uR_t + E_t$ for $t = 80$, with $X_t, M, R_t, E_t \in \mathbb{R}^{50 \times 50}$, where u is the magnitude of the outlier, and R_t is a random binary matrix. Note: the position of the outliers in entries of R_t can follow row and column patterns ($r \times c$ in (a) with r and c the given number of rows and columns, respectively, selected randomly) or can be completely random (r in (b) with r the given total number of outlying entries selected randomly).

4 Empirical Illustration

We illustrate our sequential matrix monitoring on three relevant benchmark datasets: a multi-country dataset of macroeconomic variables (see Canova and Ciccarelli, 2009), an international trade network dataset (see Rose, 2004) and a financial network dataset (Billio et al., 2023).

4.1 Inflation and Unemployment Dataset

We consider three relevant variables for macroeconomic research: the Industrial Production Index, the Price Index and the Unemployment Rate, available from the Federal Reserve Economic Data and the EUROSTAT databases. We selected observations from 11 countries of the EU (Austria, Belgium, Estonia, Finland, France, Germany, Greece, Ireland, Italy, The Netherlands, Portugal, and Spain). Series are seasonally and working day adjusted, and when necessary, they have been differentiated to get stationarity. Variables are sampled at a monthly frequency from January 2002 to October 2022. Thus, our dataset consists of a sequence of $T = 250$ $p \times n$ matrices with $p = 11$ and $n = 3$. The total number of observations is 8,250, representing an example of big data in this field.

The top plots in Figure 5 display the BF $H_t(\alpha_t)$ (solid grey) and the upper bound of the BF $\kappa_t(\alpha_t)$ (solid red) as functions of α_t for various dates (each represented by a different line). The findings in the top-left plot suggest that it is crucial to compare the BF to the upper bound. The BF tends to hover around one for higher values of α_t , with some exceptions, and it significantly exceeds one only for small values of α_t (illustrated by the dark grey lines). It frequently intersects the threshold at lower values of α_t (light grey) as well as for higher values. Additionally, there are instances where it crosses the threshold intermittently.

The left-bottom plot offers a different illustration of the effect of α_t on the outcome of the sequential monitoring for the entire sample from September 2018 to October 2022. In all settings, an outlier was detected in March 2020 (i.e. at the pandemic outbreak). For $\alpha_t = 0.054$ (solid line), a sequence of outliers is detected before February 2020 with BF far from 1, whereas, for α_t equal to 0.402, 0.801 and 0.851 (dashed, dotted and dashed-dotted, respectively), the BF is close to one before March. Qualitatively speaking, after the outbreak, for $\alpha_t = 0.402$, the BF is close to one after December 2021, whereas larger α_t values return BF close to just one after March 2020.

For illustrative purposes, we present in detail the results for some relevant dates (Figure 5 in the Appendix). For the observations in February 2020, during the pandemic outbreak, the null hypothesis of the absence of an outlier is not rejected for any choice of α_t . Nevertheless, the BF is close to one for large values of α_t . In contrast, in March 2020, there is strong evidence of an outlier. On the other dates in the figure, the alternative hypothesis is accepted for some values of α_t . The dashed vertical lines indicate the stationary point. The stationary point is not near zero on some dates, such as February 2022. On other dates, the stationary point is near zero, and the BF is far below one for a large part of the α_t values (e.g., March 2020, March 2022 and October 2022). The top-left plot in Figure 6 shows the results of the alternative procedures.

The MBF and IBFs provide clearer identification of outlying observations, enhancing the standard BF and supporting the outcome of the testing procedure endowed with the inconclusive interval given in the first line of Figure 7. Dashed lines indicate the lower and upper bounds of the inconclusive region, the dots show the value of the BF. For $\alpha_t = 0.75$ and a test size of 1%, the power is approximately 96%. Outlying observations are detected in periods related to three main events: i) the COVID-19 pandemic (from March 2020 to August 2020 and in November 2020); ii) the Ukrainian conflict (March and April 2022); iii) and the rise of inflation (September and October 2022). Our procedure results indicate that the pandemic outbreak's main effects are in March




Figure 5: BF for the three datasets: Inflation and Unemployment (left), International Trade (middle), and Volatility Network (right). Top: the BF $H_t(\alpha_t)$ and its upper bound $\kappa_t(\alpha_t)$ as functions of α_t at different dates (— and —, respectively). Bottom: the BF over time for $\alpha_t = 0.75, 0.80, 0.85$ and 0.90 (—, ---, and ----, respectively). In all plots, the reference lines at 1 (---) and $10^{-1/2}$ (-.-).

2020 since the BF of February 2020 falls within the inconclusive interval. Also, the effects of the Ukrainian conflict on the global economy began in March 2020, and the beginning of inflation rise was detected in September 2020. Another outlying observation was detected in October 2021 and is related to significant changes in the unemployment series. The outcome of the sequential test is in line with those of classical frequentist tests for outliers such as the Grubb’s (G) test (Grubbs, 1950) and the Generalized Extreme Studentized Deviate (GESD) test Rosner (1983). The second row of Figure 7 reports the number of outliers detected at the 1% level by applying the two tests element-wise to each entry of observation matrices.

4.2 Trade Network Dataset

We consider an international trade dataset made available by the International Monetary Fund (IMF). This dataset is now a reference in international trade studies (e.g., see Rose, 2004; Alcalá and Ciccone, 2004; Brancaccio et al., 2020) and combines the country reports with other sources such as United Nations COMTRADE and EUROSTAT. The data frequency is annual. The dataset includes 159 countries from 1995 to 2017. The trade network corresponds to the value of all goods imported (CIF) between pairs of countries collected by the IMF in U.S. dollars, which are transformed into constant terms using the US GDP PPP deflator from the same source with the base year 2010. All series have been differentiated to remove unit roots, and the units sub-selected to include larger GDP countries and to filter out series with more than 8% of zeroes. The final sample includes a sequence of 22 networks between 27 countries, i.e. $n = p = 27$, with the following




Figure 6: Bayes Factors for the three datasets: Inflation and Unemployment (left), International Trade (middle) and Volatility Network (right). The discount factor distribution is $\alpha_t \sim \mathcal{B}e(a, b)$ with the following parameter settings: $a = 16.5001$ and $b = 7.6429$ (left, —) and $b = 37.1667$ (left, ---); $a = 364.5001$ and $b = 156.7857$ (mid, —) and $b = 849.1667$ (mid, ---); $a = 1250.5001$ and $b = 536.2857$ (right, —) and $b = 2915.3000$ (right, ---). In all plots, the reference lines at 1 (---) and $10^{-1/2}$ (-.-).

ISO 3166-1 alpha-3 codes: AUS, AUT, BRA, CAN, CHE, CHN, DEU, DNK, ESP, FIN, FRA, GBR, GRC, IDN, IND, IRL, ITA, JPN, KOR, NLD, NOR, NZL, POL, PRT, SWE, TUR, and USA.

Following the results in the top-middle plot of Figure 5, we find evidence of outlier observations on all dates except for 2016. Nevertheless, the BF is close to one in some cases, such as 2015 and




Figure 7: Calibrated robust BF (top) and Grubb's test (bottom) for the three datasets: Inflation and Unemployment (left), International Trade (middle) and Volatility Network (right). Panel (a): Calibrated BF (—) with thresholds (---). The discounting factor is set at $\alpha_t = 0.75$, and the size is less or equal to 1% and the standard reference lines are at 1 (---) and $10^{-1/2}$ (-.-). Panel (b): Grubb's (—) and GESD (-.-) tests at the 1% significance level

2017. From the sensitivity analysis in Figure B.2, one can see that the test outcome depends crucially on the choice of α_t . The top-middle plot in Figure 6 indicates that minimum and integrated BF's provide evidence of the absence of an outlier in 2016 and the presence of outlying observations in 2015 and 2017, supporting the conclusion of the calibrated BF procedure with inconclusive intervals presented in the Panel (a) middle plot of Figure 7. In the year 2016, the classical G and GESD tests for outliers, applied entry-wise to the observation matrix, detected one outlying series out of 729 series (mid plot in Panel (b)).

4.3 Volatility Network Dataset

We investigate the presence of outliers in a volatility network among European firms with the largest market capitalization. The analyzed firms (22 German, 24 French, 4 Italian) belong to 11 GICS sectors: Financials (7 firms), Communication Services (4 firms), Consumer Discretionary (11 firms), Consumer Staples (5 firms), Health Care (6 firms), Energy (2 firms), Industrials (5 firms), Information Technology (3 firms), Materials (2 firms), Real Estate (1 firm), Utilities (3 firms), and not classified in a specific GICS sector (1 firm). The temporal networks have been extracted by pairwise Granger causality approach (e.g., see Billio et al., 2021) at a weekly frequency (Friday-Friday). The sample period ranges from the 4th of January 2016 to the 30th of September 2020, and periods before and after the outbreak of COVID-19 are included in the time interval. The dataset consists of 145 temporal networks between 50 firms, i.e. $T = 145$ and $n = p = 50$, for 362,500

observations. In the sequential monitoring, we used a rolling window of 90 observations. The plots in the top-right and bottom-right sections of Figure 5 indicate that the rejection of the hypothesis is independent of the chosen discounting coefficient. However, BF exhibits greater sensitivity to discounting towards the end of 2019. The sensitivity analysis of the BF to α_t is presented for selected dates in Figure B.3 in the Appendix.

The top-right plot in Figure 6 shows the results of the minimum BF, which support the main findings of the calibrated BF procedure (Figure 7, Panel(a), right plot). The identified outliers in 2019 are associated with transitions between different volatility levels: i) from “low” to “very low” volatility on 11-15 November; ii) from “low” to “moderate” on 2-6 December; iii) from “moderate” to “low” on the week of 16-20 December. In 2020, the following transitions were detected: i) from “low” to “high” during the week of 17-21 February; ii) a progressive transition from “high” to “moderate” volatility between 7 and 24 April; iii) from “moderate” to “low” volatility during the week of 3-7 August; iv) a sequence of outliers corresponding to a period of switches between “low” and “moderate” between 1 September and 2 October. The BFs of the transitory periods mentioned belong to the inconclusive interval: 25-28 February 2020 (“low” to “high”), 27-30 April (“high” to “moderate”), and 21-25 September (“low” to “moderate”). When the BF is far above one, the classical G and GESD tests for outliers, applied entry-wise to the observation matrix, detect a reduction in the number of outliers. In summary, the rapid changes in volatility and the persistence of volatility regimes call for nonlinear models, such as switching or threshold models, which account for structural breaks and recurrent regimes.

5 Conclusion

The assessment of the model performance is relevant in many applications and becomes crucial in forecasting. This paper proposes sequential monitoring for the matrix normal model. The hypothesis testing procedure extends the predictive Bayes Factor (BF) with the power discounting to matrix models. The proposed approach relies on normality, which is now a default assumption in many modelling applications and serves as a preliminary test for outliers in sequences of matrix-valued data. Some solutions are proposed to mitigate the test outcome’s dependence on the discounting coefficient value. First, the minimum and the integrated BFs are proposed as alternatives to the standard BF. Second, the finite-sample distribution of the predictive BF is derived, and a testing procedure is proposed based on inconclusive region and calibrated discounting. Simulation experiments are conducted to study the properties of our Bayesian test. Numerical illustrations with simulated and some relevant economic and financial datasets are given. They include a comparison with classical tests for outliers and a validation based on major global event dates (e.g., the pandemic, the Ukrainian conflict, and the rise in inflation), which significantly impacted the economic system.

Acknowledgement

The authors’ research used the HPC multiprocessor systems at the Ca’ Foscari University of Venice. This work was partly funded by the MUR – PRIN project ‘*Discrete random structures for Bayesian learning and prediction*’ under g.a. n. 2022CLTYP4 and the Next Generation EU – ‘*GRINS – Growing Resilient, INclusive and Sustainable*’ project (PE0000018), National Recovery and Resilience Plan (NRRP) – PE9. The views and opinions expressed are only those of the authors and do not necessarily reflect those of the European Union or the European Commission. Neither the European Union nor the European Commission can be held responsible for them.

A Proofs of the Results

Proof of Proposition 1.

Proof. For ease of notation in the following, we drop the subscript t from α_t . Since $H_t(1) = 1$, we need to show that under the proposition assumptions, there exists $\alpha_g \in (0, 1)$ such that $H_t(\alpha) > 1$ for $0 < \alpha < \alpha_g$ and $H_t(\alpha) < 1$ for $\alpha_g < \alpha < 1$. The limit of the normalizing constant $C(\alpha) = \left(\int_{\Theta} p(\theta_t|\mathcal{D}_{t-1})^\alpha \lambda(d\theta_t)\right)^{-1}$ for $\alpha \rightarrow 0^+$ is $\lambda(\Theta)^{-1}$ where $\lambda(\Theta)$ is the Lebesgue measure of Θ . It follows that

$$\lim_{\alpha \rightarrow 0^+} H_t(\alpha) = \lim_{\alpha \rightarrow 0^+} \frac{1}{C(\alpha)} \frac{\int_{\Theta} p(y_t|\theta_t)p(\theta_t|\mathcal{D}_{t-1})\lambda(d\theta)}{\int_{\Theta} p(y_t|\theta_t)p(\theta_t|\mathcal{D}_{t-1})^\alpha \lambda(d\theta)} = \frac{\lambda(\Theta) \int_{\Theta} p(y_t|\theta_t)p(\theta_t|\mathcal{D}_{t-1})\lambda(d\theta)}{\int_{\Theta} p(y_t|\theta_t)\lambda(d\theta)} \geq 1 \quad (\text{A.1})$$

by the assumption. The inequality $H_t(\alpha) < 1$ can be equivalently written as

$$\int_{\Theta} p(y_t|\theta_t)(p(\theta_t|\mathcal{D}_{t-1}) - C(\alpha)p(\theta_t|\mathcal{D}_{t-1})^\alpha)\lambda(d\theta_t) < 0. \quad (\text{A.2})$$

and is satisfied if $H_t(\alpha)$ has a unique minimum value for some α . Let us assume the distribution $p(\theta_t|\mathcal{D}_t)$ is absolutely continuous with respect to the Lebesgue measure λ , and define

$$m_0 = \int_{\Theta} p(y_t|\theta_t)p(\theta_t|\mathcal{D}_{t-1})\lambda(d\theta_t), \quad m_1(\alpha) = \int_{\Theta} p(y_t|\theta_t)p(\theta_t|\mathcal{D}_{t-1})^\alpha \lambda(d\theta_t).$$

Finding a necessary condition on α for H_t to admit a minimum is possible. The first derivative of the BF is

$$\begin{aligned} \partial_\alpha H(\alpha) &= \frac{\partial}{\partial \alpha} \frac{m_0}{C(\alpha)m_1(\alpha)} = \frac{m_0}{C(\alpha)m_1(\alpha)^2} \left(C(\alpha)m_1(\alpha) \int_{\Theta} p(\theta_t|\mathcal{D}_{t-1})^\alpha \log p(\theta_t|\mathcal{D}_{t-1})\lambda(d\theta_t) - \right. \\ &\quad \left. \int_{\Theta} p(y_t|\theta_t)p(\theta_t|\mathcal{D}_{t-1})^\alpha \log p(\theta_t|\mathcal{D}_{t-1})\lambda(d\theta_t) \right) \\ &= \frac{m_0}{m_1(\alpha)^2} \left(m_1(\alpha) \int_{\Theta} p(\theta_t|\mathcal{D}_{t-1})^\alpha \log p(\theta_t|\mathcal{D}_{t-1})\lambda(d\theta_t) - \right. \\ &\quad \left. \int_{\Theta} p(\theta_t|\mathcal{D}_{t-1})^\alpha \lambda(d\theta_t) \int_{\Theta} p(y_t|\theta_t)p(\theta_t|\mathcal{D}_{t-1})^\alpha \log p(\theta_t|\mathcal{D}_{t-1})\lambda(d\theta_t) \right), \end{aligned} \quad (\text{A.3})$$

where we used

$$\partial_\alpha C(\alpha) = -C(\alpha)^2 \int_{\Theta} p(\theta_t|\mathcal{D}_{t-1})^\alpha \log p(\theta_t|\mathcal{D}_{t-1})\lambda(d\theta_t),$$

which is well-defined since $x^\alpha \log x \rightarrow 0$ as $x \rightarrow 0^+$ for all $\alpha > 0$. If α_0 satisfies the proposition's assumptions, then the first derivative changes sign only once, and the BF H_t takes values above and below one. \square

Proof of Proposition 2. Let us recall the following standard properties for matrix variate normal variables.

Proposition A.1 (Gupta, 1992, Theorem 2.2). *Let X and Y be two random matrices of dimension $p \times n$ and $q \times n$, respectively. Assume that $Y|X \sim \mathcal{N}_{q,n}(C + DX, \Sigma_2, \Phi)$ and $X \sim \mathcal{N}_{p,n}(F, \Sigma_1, \Phi)$ where C is $q \times n$, D $q \times p$, Σ_2 $q \times q$, F $p \times n$, Φ $n \times n$, Σ_1 $p \times p$, $\Sigma_2 > 0$, $\Sigma_1 > 0$, $\Phi > 0$. Then:*

$$Z = \begin{pmatrix} X \\ Y \end{pmatrix} \sim \mathcal{N}_{q+p,n} \left[\begin{pmatrix} F \\ DF + C \end{pmatrix}, \begin{pmatrix} \Sigma_1 & \Sigma_1 D' \\ D \Sigma_1 & \Sigma_2 + D \Sigma_1 D' \end{pmatrix}, \Phi \right]. \quad (\text{A.4})$$

Proposition A.2 (Gupta, 1999, Theorem 2.3.12, pg. 65). *Let $Z \sim \mathcal{N}_{q,n}(H, \Sigma, Q)$ and partitions:*

$$Z = \begin{pmatrix} Z_{1r} \\ Z_{2r} \end{pmatrix} \begin{matrix} m \\ q-m \end{matrix}; \quad H = \begin{pmatrix} H_{1r} \\ H_{2r} \end{pmatrix} \begin{matrix} m \\ q-m \end{matrix}; \quad \Sigma = \begin{pmatrix} \Sigma_{11} & \Sigma_{12} \\ \Sigma_{21} & \Sigma_{22} \end{pmatrix} \begin{matrix} m \\ q-m \end{matrix}. \quad (\text{A.5})$$

Then:

$$Z_{1r} \sim \mathcal{N}_{m,n}(H_{1r}, \Sigma_{11}, Q), \quad Z_{2r}|Z_{1r} \sim \mathcal{N}_{q-m,n}(H_{2r} + \Sigma_{21}\Sigma_{11}^{-1}(Z_{1r} - H_{1r}), \Sigma_{22.1}, Q), \quad (\text{A.6})$$

where $\Sigma_{22.1} = \Sigma_{22} - \Sigma_{21}\Sigma_{11}^{-1}\Sigma_{12}$. It follows that:

$$Z_{2r} \sim \mathcal{N}_{q-m,n}(H_{2r}, \Sigma_{22}, Q), \quad Z_{1r}|Z_{2r} \sim \mathcal{N}_{m,n}(H_{1r} + \Sigma_{12}\Sigma_{22}^{-1}(Z_{2r} - H_{2r}), \Sigma_{11.2}, Q), \quad (\text{A.7})$$

where $\Sigma_{11.2} = \Sigma_{11} - \Sigma_{12}\Sigma_{22}^{-1}\Sigma_{21}$.

Proof. Let $Y_i (p \times n)$ $i = 1, \dots, t-1$ be a random matrix i.i.d. sequence from a matrix-variate normal distribution with mean matrix $B (p \times n)$ and covariance matrices $\Sigma_L (p \times p) > 0$ and $V (n \times n) > 0$, then $\text{vec}(Y'_i) \sim \mathcal{N}_{pn}(\text{vec}(B'), \Sigma_L \otimes V)$ and the pdf is:

$$f(Y_i|B, V, \Sigma_L) = (2\pi)^{-\frac{np}{2}} |\Sigma_L|^{-\frac{n}{2}} |V|^{-\frac{p}{2}} \exp \left\{ -\frac{1}{2} \text{tr} \left[\Sigma_L^{-1} (Y_i - B) V^{-1} (Y_i - B)' \right] \right\}. \quad (\text{A.8})$$

Define $T = t - 1$, then the likelihood function is:

$$\begin{aligned} f(Y_1, Y_2, \dots, Y_T|B, V, \Sigma_L) &= \prod_{i=1}^T \frac{\exp \left\{ -\frac{1}{2} \text{tr} \left[\Sigma_L^{-1} (Y_i - B) V^{-1} (Y_i - B)' \right] \right\}}{(2\pi)^{\frac{np}{2}} |\Sigma_L|^{\frac{n}{2}} |V|^{\frac{p}{2}}} \\ &= \frac{\exp \left\{ -\frac{1}{2} \text{tr} \left[\sum_{i=1}^T \Sigma_L^{-1} (Y_i - B) V^{-1} (Y_i - B)' \right] \right\}}{(2\pi)^{\frac{Tnp}{2}} |\Sigma_L|^{T\frac{n}{2}} |V|^{T\frac{p}{2}}} \\ &= \frac{\exp \left\{ -\frac{1}{2} \text{tr} \left[(I_T \otimes \Sigma_L^{-1}) (\mathbf{Y} - \boldsymbol{\iota}_T \otimes B) V^{-1} (\mathbf{Y} - \boldsymbol{\iota}_T \otimes B)' \right] \right\}}{(2\pi)^{\frac{Tnp}{2}} |\Sigma_L|^{T\frac{n}{2}} |V|^{\frac{Tp}{2}}}, \end{aligned} \quad (\text{A.9})$$

where $\mathbf{Y}' = [Y'_1, Y'_2, \dots, Y'_T]$ is a matrix ($Tp \times n$) obtained by stacking vertically the T matrices Y_i , $i = 1, \dots, T$ and $I_T \otimes \Sigma_L$ is ($Tp \times Tp$). We can say that the previous expression is the pdf of the random matrix $\mathbf{Y} (Tp \times n)$ with mean $\boldsymbol{\iota}_T \otimes B (Tp \times n)$ and a covariance matrix $(I_T \otimes \Sigma_L) \otimes V$ where I_T and $\boldsymbol{\iota}_T$ are the T -dimensional identity matrix and unit column vector, respectively. The matrix $I_T \otimes \Sigma_L (Tp \times Tp)$ is positive definite, which follows from $\Sigma_L > 0$, $I_T > 0$ and $V > 0$.

Since $|(I_T \otimes \Sigma_L) \otimes V| = |(I_T \otimes \Sigma_L)|^n |V|^{Tp} = |I_T|^{np} |\Sigma_L|^{nT} |V|^{Tp} = |\Sigma_L|^{Tn} |V|^{Tp}$ the pdf of \mathbf{Y} is:

$$\begin{aligned} f(\mathbf{Y}|B, V, \Sigma_L) &= \frac{\exp \left\{ -\frac{1}{2} \text{tr} \left[(I_T \otimes \Sigma_L)^{-1} (\mathbf{Y} - \boldsymbol{\iota}_T \otimes B) V^{-1} (\mathbf{Y} - \boldsymbol{\iota}_T \otimes B)' \right] \right\}}{(2\pi)^{\frac{Tnp}{2}} |I_T \otimes \Sigma_L|^{\frac{n}{2}} |V|^{\frac{Tp}{2}}} \\ &= \frac{\exp \left\{ -\frac{1}{2} \text{tr} \left[(I_T \otimes \Sigma_L)^{-1} (\mathbf{Y} - \boldsymbol{\iota}_T \otimes B) V^{-1} (\mathbf{Y} - \boldsymbol{\iota}_T \otimes B)' \right] \right\}}{(2\pi)^{\frac{Tnp}{2}} |\Sigma_L|^{\frac{Tn}{2}} |V|^{\frac{Tp}{2}}} \end{aligned} \quad (\text{A.10})$$

which is equal to the pdf in Equation (A.9). In the following, we assume the $p \times n$ matrix B is random, and the matrices Σ_P and V are known. Assume a matrix normal prior distribution for B , i.e. $B \sim \mathcal{N}_{p,n}(M, \Sigma_P, V)$, with pdf

$$f(B|M, V, \Sigma_P) = \frac{\exp \left\{ -\frac{1}{2} \text{tr} \left[\Sigma_P^{-1} (B - M) V^{-1} (B - M)' \right] \right\}}{(2\pi)^{\frac{np}{2}} |\Sigma_P|^{\frac{n}{2}} |V|^{\frac{p}{2}}}. \quad (\text{A.11})$$

Note that $\boldsymbol{\nu}_T \otimes B = (\boldsymbol{\nu}_T \otimes I_p) B$ and apply Prop. A.2 to (A.10) and (A.11), replacing $X, Y|X, C, D, F, \Sigma_1, \Sigma_2, V$ with $B_t, \mathbf{Y}|B_t, 0, \boldsymbol{\nu}_T \otimes I_p, M, \Sigma_P, \Sigma_L, \Phi$, where I_p is the identity matrix of order p , we get:

$$Z \sim \mathcal{N}_{T_{p+p}, n} \left[\begin{pmatrix} M \\ (\boldsymbol{\nu}_T, I_p) M + 0 \end{pmatrix}, \begin{pmatrix} \Sigma_P & \Sigma_P (\boldsymbol{\nu}_T, I_p)' \\ (\boldsymbol{\nu}_T, I_p) \Sigma_P & (I_T \otimes \Sigma_L) + (\boldsymbol{\nu}_T \otimes I_p) \Sigma_P (\boldsymbol{\nu}_T \otimes I_p)' \end{pmatrix} \otimes V \right], \quad (\text{A.12})$$

where $Z' = (B_t', \mathbf{Y}')$. By applying Prop. A.2 to Z and defining:

$$M_* = M + \Sigma_P (\boldsymbol{\nu}_T \otimes I_p)' \left[I_T \otimes \Sigma_L + (\boldsymbol{\nu}_T \otimes I_p) \Sigma_P (\boldsymbol{\nu}_T \otimes I_p)' \right]^{-1} (\mathbf{Y} - \boldsymbol{\nu}_T \otimes M) \quad (\text{A.13})$$

$$\Sigma_* = \Sigma_P - \Sigma_P (\boldsymbol{\nu}_T \otimes I_p)' \left[I_T \otimes \Sigma_L + (\boldsymbol{\nu}_T \otimes I_p) \Sigma_P (\boldsymbol{\nu}_T \otimes I_p)' \right]^{-1} (\boldsymbol{\nu}_T \otimes I_p) \Sigma_P, \quad (\text{A.14})$$

we obtain the posterior distribution of B at time $t - 1$ with pdf

$$f(B_t | \mathbf{Y}, M, V, \Sigma_*) = \frac{\exp \left\{ -\frac{1}{2} \text{tr} \left[\Sigma_*^{-1} (B_t - M_*) V^{-1} (B_t - M_*)' \right] \right\}}{(2\pi)^{\frac{np}{2}} |\Sigma_*|^{\frac{n}{2}} |V|^{\frac{p}{2}}} \quad (\text{A.15})$$

that is B follows a matrix normal $B_t | \mathbf{Y} \sim \mathcal{N}_{p, n} (M_*, \Sigma_*, V)$. \square

Proof of Proposition 3. Let $f(B_t, V_t | \mathbf{Y}, M, \Sigma_L, \Psi, m)$ be the density of the Matrix Normal Inverse Wishart prior, the joint posterior density of B_t and V_t is:

$$\begin{aligned} f(B_t, V_t | \mathbf{Y}, M, \Sigma_L, \Psi, m) &= \frac{h(B_t, V_t | M, \Sigma_L, \Psi, m) f(\mathbf{Y} | B_t, V_t, \Sigma_L)}{\int_{V_t > 0} \int_{B_t} h(B_t, V_t | M, \Sigma_L, \Psi, m) f(\mathbf{Y} | B_t, V_t, \Sigma_L) dB_t dV_t} \\ &= \frac{|V|^{-\frac{m+p}{2}} |V|^{-\frac{Tp}{2}} \exp \left\{ -\frac{1}{2} \text{tr} [\mathcal{A}] \right\}}{\int_{V > 0} \int_B |V|^{-\frac{m+p}{2}} |V|^{-\frac{Tp}{2}} \exp \left\{ -\frac{1}{2} \text{tr} [\mathcal{A}] \right\} dB dV}, \end{aligned}$$

where we defined $\mathcal{A} = (B_t - M)' k \Sigma_L^{-1} (B_t - M) V^{-1} + \Psi V_t^{-1} + \sum_{s=1}^T (Y_s - B_t)' \Sigma_L^{-1} (Y_s - B_t) V_t^{-1}$ with $k = \rho\varphi$ and $\bar{Y} = T^{-1} \sum_{s=1}^T Y_s$.

Regarding the numerator, since $S V_t^{-1} = \sum_{s=1}^T (Y_s - B_t)' \Sigma_L^{-1} (Y_s - B_t) V_t^{-1} - T (\bar{Y} - B_t)' \Sigma_L^{-1} (\bar{Y} - B_t) V_t^{-1}$ with $S = \sum_{s=1}^T (Y_s - \bar{Y})' \Sigma_L^{-1} (Y_s - \bar{Y}) / T$ and

$$\begin{aligned} &k(B_t' \Sigma_L^{-1} B_t - M' \Sigma_L^{-1} B_t - B_t' \Sigma_L^{-1} M + M' \Sigma_L^{-1} M) + T(\bar{Y}' \Sigma_L^{-1} \bar{Y} - B_t' \Sigma_L^{-1} \bar{Y} - \bar{Y}' \Sigma_L^{-1} B_t + B_t' \Sigma_L^{-1} B_t) \\ &+ \frac{(kM' + T\bar{Y}') \Sigma_L^{-1} (kM + T\bar{Y})}{(k+T)} - \frac{(kM' + T\bar{Y}') \Sigma_L^{-1} (kM + T\bar{Y})}{(k+T)} \\ &= (k+T) \left(B_t - \frac{kM + T\bar{Y}}{(k+T)} \right)' \Sigma_L^{-1} \left(B_t - \frac{kM + T\bar{Y}}{(k+T)} \right) + kT \frac{(M - \bar{Y})' \Sigma_L^{-1} (M - \bar{Y})}{(k+T)} \end{aligned}$$

the quantity \mathcal{A} in the exponential term can be written as

$$\begin{aligned} \mathcal{A} &= (k+T) \left(B_t - \frac{kM + T\bar{Y}}{(k+T)} \right)' \Sigma_L^{-1} \left(B_t - \frac{kM + T\bar{Y}}{(k+T)} \right) V_t^{-1} + kT \frac{(M - \bar{Y})' \Sigma_L^{-1} (M - \bar{Y})}{(k+T)} V_t^{-1} \\ &+ \sum_{i=1}^T (Y_i - \bar{Y})' \Sigma_L^{-1} (Y_i - \bar{Y}) V_t^{-1} + \Psi V_t^{-1} = [k_* (B_t - M_*)' \Sigma_L^{-1} (B_t - M_*) + \Psi_*] V_t^{-1}, \end{aligned}$$

where $k_* = k + T$, $m_* = m + Tp$, $M_* = \frac{kM + T\bar{Y}}{k+T}$, and $\Psi_* = \Psi + kT \frac{(M - \bar{Y})' \Sigma_L^{-1} (M - \bar{Y})}{(k+T)} + TS$.

The integral at the denominator is

$$\begin{aligned}
& \int_{V_t > 0} \int_{B_t} |V|^{-\frac{m+p}{2}} |V_t|^{-\frac{Tp}{2}} \exp \left\{ -\frac{1}{2} \text{tr} [\mathcal{A}] \right\} dB_t dV_t = \\
& = \int_{V_t > 0} \int_{B_t} |V|^{-\frac{m+Tp}{2}} |V_t|^{-\frac{p}{2}} \exp \left\{ -\frac{1}{2} \text{tr} \left[k_* (B_t - M_*)' \Sigma_L^{-1} (B_t - M_*) + \Psi_* \right] V_t^{-1} \right\} dB_t dV_t \\
& = \int_{V_t > 0} \int_{B_t} |V_t|^{-\frac{p}{2}} \exp \left\{ -\frac{1}{2} \text{tr} \left[k_* (B_t - M_*)' \Sigma_L^{-1} (B_t - M_*) \right] V_t^{-1} \right\} dB_t dV_t \\
& = (2\pi)^{\frac{np}{2}} |\Sigma_L / k_*|^{\frac{n}{2}} \int_{V_t > 0} |V_t|^{-\frac{m_*}{2}} \exp \left\{ -\frac{1}{2} \text{tr} [\Psi_*] V_t^{-1} \right\} dV_t \\
& = (2\pi)^{\frac{np}{2}} |\Sigma_L / k_*|^{\frac{n}{2}} 2^{\frac{(m_* - n - 1)n}{2}} \Gamma_n \left(\frac{m_* - n - 1}{2} \right) |\Psi_*|^{-\frac{m_* - n - 1}{2}}. \tag{A.16}
\end{aligned}$$

Thus the ratio

$$\begin{aligned}
& \frac{|V_t|^{-\frac{m+Tp}{2}} |V_t|^{-\frac{p}{2}} \exp \left\{ -\frac{1}{2} \text{tr} \left[k_* (B - M_*)' \Sigma_L^{-1} (B - M_*) + \Psi_* \right] V_t^{-1} \right\}}{(2\pi)^{\frac{np}{2}} |\Sigma_L / k_*|^{\frac{n}{2}} 2^{\frac{(m_* - n - 1)n}{2}} \Gamma_n \left(\frac{m_* - n - 1}{2} \right) |\Psi_*|^{-\frac{m_* - n - 1}{2}}} \\
& = \frac{|\Psi_*|^{\frac{m_* - n - 1}{2}} \exp \left\{ -\frac{1}{2} \text{tr} \left[(B_t - M_*)' \Sigma_*^{-1} (B_t - M_*) + \Psi_* \right] V_t^{-1} \right\}}{(2\pi)^{\frac{np}{2}} |\Sigma_*|^{\frac{n}{2}} 2^{\frac{(m_* - n - 1)n}{2}} \Gamma_n \left(\frac{m_* - n - 1}{2} \right) |V_t|^{\frac{m_* + p}{2}}} \tag{A.17}
\end{aligned}$$

returns the density of a $\mathcal{MNITW}(B_t, V_t | M_*, \Sigma_*, \Psi_*, k_*, m_*)$, where $\Sigma_* = \Sigma_L / k_*$. \square

Proof of Proposition 4. i) Under the assumption in Prop. 2, apply Prop. A.1 with $X = B | \mathbf{Y}$, $Y = Y_t$, $C = 0$, $D = I_p$, $F = M_*$, $\Sigma_1 = \Sigma_*$ and $\Sigma_2 = \Sigma_L$, where Y_t is the sample at time t and find:

$$Z = \begin{pmatrix} B | \mathbf{Y} \\ Y_t \end{pmatrix} \sim N_{p+p,n} \left[\begin{pmatrix} M_* \\ M_* \end{pmatrix}, \begin{pmatrix} \Sigma_* & \Sigma_* \\ \Sigma_* & \Sigma_d \end{pmatrix} \otimes V \right], \tag{A.18}$$

where $\Sigma_d = \Sigma_L + \Sigma_*$. The posterior predictive distribution returns the numerator of the BF:

$$p(Y_t | \mathcal{D}_{t-1}) = \int p(Y_t | \theta_t) p(\theta_t | \mathcal{D}_{t-1}) d\theta_t \tag{A.19}$$

$$= \int f(Y_t | B_t, V, \Sigma_L) f(B_t | \mathbf{Y}, M, V, \Sigma_P) dB_t \tag{A.20}$$

$$= \frac{\exp \left\{ -\frac{1}{2} \text{tr} \left[\Sigma_d^{-1} (Y_t - M_*) V^{-1} (Y_t - M_*)' \right] \right\}}{(2\pi)^{\frac{np}{2}} |\Sigma_d|^{\frac{n}{2}} |V|^{\frac{p}{2}}} \tag{A.21}$$

that is $Y_t | \mathcal{D}_{t-1} \sim \mathcal{N}_{p,n}(M_*, \Sigma_d, V)$. The denominator of the BF is derived from the alternative distribution of the parameter

$$p_A(\theta_t | \mathcal{D}_{t-1}) = C(\alpha_t) p(B_t | \mathbf{Y}, M, V, \Sigma_P)^{\alpha_t} = \frac{\exp \left\{ -\frac{1}{2} \text{tr} \left[\Sigma_{A,*}^{-1} (B_t - M_*) V^{-1} (B_t - M_*)' \right] \right\}}{(2\pi)^{\frac{np}{2}} |\Sigma_{A,*}|^{\frac{n}{2}} |V|^{\frac{p}{2}}} \tag{A.22}$$

with $\Sigma_{A,*} = \Sigma_* / \alpha_t$ and the constant

$$C(\alpha_t) = \frac{(2\pi)^{\frac{\alpha_t np}{2}} \alpha_t^{\frac{\alpha_t np}{2}} \left| \frac{\Sigma_*}{\alpha_t} \right|^{\frac{\alpha_t n}{2}} |V|^{\frac{\alpha_t p}{2}}}{(2\pi)^{\frac{np}{2}} \left| \frac{\Sigma_*}{\alpha_t} \right|^{\frac{n}{2}} |V|^{\frac{p}{2}}} = \frac{\alpha_t^{\frac{p}{2}} (2\pi)^{\frac{\alpha_t np}{2}} \left| \Sigma_* \right|^{\frac{\alpha_t n}{2}} |V|^{\frac{\alpha_t p}{2}}}{(2\pi)^{\frac{np}{2}} \left| \Sigma_* \right|^{\frac{n}{2}} |V|^{\frac{p}{2}}}. \quad (\text{A.23})$$

We apply Prop. A.1 for (A3.1) and (A2.2) with $i = t$:

$$Z = \begin{bmatrix} (B|\mathbf{Y})_A \\ Y_t \end{bmatrix} \sim \mathcal{N}_{p+p,n} \left[\begin{pmatrix} M_* \\ M_* \end{pmatrix}, \begin{pmatrix} \Sigma_{A,*} & \Sigma_{A,*} \\ \Sigma_{A,*} & \Sigma_{A,d} \end{pmatrix}, V \right], \quad (\text{A.24})$$

where $(B|\mathbf{Y})_A$ means that we consider the alternative distribution as is in (A.22) and $\Sigma_{A,d} = \Sigma_L + \Sigma_{A,*}$. We get:

$$\begin{aligned} p_A(Y_t|\mathcal{D}_{t-1}) &= \int p(Y_t|\theta_t) p_A(\theta_t|\mathcal{D}_{t-1}) d\theta_t = \int f(Y_t|B_t, V, \Sigma_L) f_A(B_t|\mathbf{Y}, M, V, \Sigma_P) dB_t \\ &= \frac{\exp \left\{ -\frac{1}{2} \text{tr} \left[\Sigma_{A,d}^{-1} (Y_t - M_*) V^{-1} (Y_t - M_*)' \right] \right\}}{(2\pi)^{\frac{np}{2}} \left| \Sigma_{A,d} \right|^{\frac{n}{2}} |V|^{\frac{p}{2}}} \end{aligned} \quad (\text{A.25})$$

that is the pdf of $\mathcal{N}_{p,n}(M_*, \Sigma_{A,d}, V)$. The BF follows from its definition

$$H_t = \frac{p(Y_t|\mathcal{D}_{t-1})}{p_A(Y_t|\mathcal{D}_{t-1})} = \frac{\left| \Sigma_{A,d} \right|^{\frac{n}{2}}}{\left| \Sigma_d \right|^{\frac{n}{2}}} \exp \left\{ -\frac{1}{2} \text{tr} \left[\left(\Sigma_d^{-1} - \Sigma_{A,d}^{-1} \right) (Y_t - M_*) V^{-1} (Y_t - M_*)' \right] \right\}. \quad (\text{A.26})$$

ii) Under the assumptions in Prop. 3, since the observation Y_t follows a matrix normal with density $f(Y_t|B_t, V_t, \Sigma)$, the posterior predictive at the numerator of the BF is:

$$p(Y_t|\mathcal{D}_{t-1}) = \int_{V_t > 0} \int_{B_t} f(Y_t|B_t, V_t, \Sigma_L) f(B_t, V_t|\mathbf{Y}, M_*, \Sigma_*, \Psi_*, m_*) dB_t dV_t, \quad (\text{A.27})$$

where the argument of the integral is:

$$\frac{|\Psi_*|^{\frac{m_*-n-1}{2}} \exp \left\{ -\frac{1}{2} \text{tr} \left[\left(k_* (B_t - M_*)' \Sigma_L^{-1} (B_t - M_*) + \Psi_* + (Y_t - B_t)' \Sigma_L^{-1} (Y_t - B_t) \right) V_t^{-1} \right] \right\}}{(2\pi)^{\frac{np}{2}} \left| \Sigma_L / k_* \right|^{\frac{n}{2}} 2^{\frac{(m_*-n-1)n}{2}} \Gamma_n \left(\frac{m_*-n-1}{2} \right) |V_t|^{\frac{m_*+p}{2}} (2\pi)^{\frac{np}{2}} \left| \Sigma_L \right|^{\frac{n}{2}} |V_t|^{\frac{p}{2}}}$$

The matrix premultiplying V_t^{-1} in the argument of the tr is:

$$\begin{aligned} k_* (B_t - M_*)' \Sigma_L^{-1} (B_t - M_*) + \Psi_* + (Y_t - B_t)' \Sigma_L^{-1} (Y_t - B_t) &= \\ k_d (B_t - M_d)' \Sigma_L^{-1} (B_t - M_d) + \Psi_d &= (B_t - M_d)' \Sigma_d^{-1} (B_t - M_d) + \Psi_d, \end{aligned} \quad (\text{A.28})$$

where the equality follows from the fact that S in Ψ_d is equal to zero and $\bar{Y} = Y_t$ for $T = 1$, and where we defined $k_d = k_* + 1$, $m_d = m_* + p$, and

$$M_d = \frac{k_* M_* + Y_t}{k_* + 1}, \quad \Sigma_d = \Sigma_L / k_d, \quad \Psi_d = \Psi_* + k_* \frac{(M_* - Y_t)' \Sigma_L^{-1} (M_* - Y_t)}{(k_* + 1)}. \quad (\text{A.29})$$

The integral becomes:

$$\int_{V_t > 0} \frac{|\Psi_*|^{\frac{m_*-n-1}{2}} k_*^{\frac{np}{2}}}{(2\pi)^{\frac{np}{2}} k_d^{\frac{np}{2}} 2^{\frac{(m_*-n-1)n}{2}} \Gamma_n \left(\frac{m_*-n-1}{2} \right) |V_t|^{\frac{m_d}{2}} \left| \Sigma_L \right|^{\frac{n}{2}}}$$

$$\begin{aligned}
& \int_{B_t} \frac{\exp \left\{ -\frac{1}{2} \text{tr} \left[\left((B_t - M_d)' \Sigma_d^{-1} (B_t - M_d) + \Psi_d \right) V_t^{-1} \right] \right\}}{|\Sigma_d|^{\frac{n}{2}} |V_t|^{\frac{np}{2}} (2\pi)^{\frac{np}{2}}} dB_t dV_t \\
&= \int_{V_t > 0} \frac{|\Psi_*|^{\frac{m_* - n - 1}{2}} k_*^{\frac{np}{2}} \exp \left\{ -\frac{1}{2} \text{tr} \left[\Psi_d V_t^{-1} \right] \right\}}{(2\pi)^{\frac{np}{2}} k_d^{\frac{np}{2}} 2^{\frac{(m_* - n - 1)n}{2}} \Gamma_n \left(\frac{m_* - n - 1}{2} \right) |V_t|^{\frac{m_d}{2}} |\Sigma_L|^{\frac{n}{2}}} dV_t \\
&= \int_{V_t > 0} \frac{|\Psi_*|^{\frac{m_* - n - 1}{2}} |\Psi_d|^{\frac{m_d - n - 1}{2}} k_*^{\frac{np}{2}} \Gamma_n \left(\frac{m_d - n - 1}{2} \right) \exp \left\{ -\frac{1}{2} \text{tr} \left[\Psi_d V_t^{-1} \right] \right\}}{(2\pi)^{\frac{np}{2}} k_d^{\frac{np}{2}} |\Psi_d|^{\frac{m_d - n - 1}{2}} 2^{\frac{(m_d - n - 1)n}{2}} 2^{-\frac{np}{2}} \Gamma_n \left(\frac{m_* - n - 1}{2} \right) \Gamma_n \left(\frac{m_d - n - 1}{2} \right) |V_t|^{\frac{m_d}{2}} |\Sigma_L|^{\frac{n}{2}}} dV_t \\
&= \frac{|\Psi_*|^{\frac{m_* - n - 1}{2}} k_*^{\frac{np}{2}} \Gamma_n \left(\frac{m_d - n - 1}{2} \right)}{\pi^{\frac{np}{2}} k_d^{\frac{np}{2}} |\Psi_d|^{\frac{m_d - n - 1}{2}} \Gamma_n \left(\frac{m_* - n - 1}{2} \right) |\Sigma_L|^{\frac{n}{2}}} \tag{A.30}
\end{aligned}$$

which is the density of a matrix Student-t $\mathcal{T}_{p,n}(m_* - 2n, M_*, \Sigma_L, L_*)$ following the definition in Gupta and Nagar (1999) p. 134, given:

$$\begin{aligned}
|\Psi_d|^{\frac{m_d - n - 1}{2}} &= \left| \Psi_* + k_* \frac{(M_* - Y_t)' \Sigma_L^{-1} (M_* - Y_t)}{(k_* + 1)} \right|^{\frac{m_d - n - 1}{2}} \\
&= \left| \Psi_* \left[I_n + L_*^{-1} (M_{1,*} - Y_{1,t}) \Sigma_L^{-1} (M_{1,*} - Y_{1,t})' \right] \right|^{\frac{m_d - n - 1}{2}} \\
&= |\Psi_*|^{\frac{m_* + p - n - 1}{2}} \left| I_n + L_*^{-1} (M_{1,*} - Y_{1,t}) \Sigma_L^{-1} (M_{1,*} - Y_{1,t})' \right|^{\frac{m_* + p - n - 1}{2}} \\
&\quad (\pi)^{-\frac{np}{2}} |L_*|^{-\frac{p}{2}} |\Sigma_L|^{-\frac{n}{2}} \Gamma_n \left(\frac{m_1 + p + n - 1}{2} \right) \\
&= \frac{(\pi)^{-\frac{np}{2}} |L_*|^{-\frac{p}{2}} |\Sigma_L|^{-\frac{n}{2}} \Gamma_n \left(\frac{m_1 + p + n - 1}{2} \right)}{\left| I_n + L_*^{-1} (M_{1,*} - Y_{1,t}) \Sigma_L^{-1} (M_{1,*} - Y_{1,t})' \right|^{\frac{m_1 + p + n - 1}{2}} \Gamma_n \left(\frac{m_1 + n - 1}{2} \right)}, \tag{A.31}
\end{aligned}$$

is the probability density function of a $\mathcal{T}_{n,p}(m_* - 2n, M_{1,*}, L_*, \Sigma_L)$ for the quantity $Y_{1,t}$, where we have defined $L_* = \Psi_* k_d / k_*$, $M_{1,*} = M_*'$, $Y_{1,t} = Y_t'$, $m_d = m_* + p$, and $m_1 = m_* - 2n = m + Tp - 2n$ (assuming $m > 2n - Tp$) and given the Theorem 4.4.4 in Gupta and Nagar (1999) about the probability density function of the transposed of a matrix Student-t distribution.

For what concerns the denominator of the BF, after some algebraic manipulations, one obtains:

$$\begin{aligned}
p_A(\theta_t | \mathcal{D}_{t-1}) &= C(\alpha_t) p(B_t, V_t | \mathbf{Y}, M_*, \Sigma_*, \Psi_*, m_*)^{\alpha_t} \\
&= \frac{|\alpha_t \Psi_*|^{\frac{m_{A,*} - n - 1}{2}} \exp \left\{ -\frac{1}{2} \text{tr} \left[\alpha_t k_* (B_t - M_*)' \Sigma_L^{-1} (B_t - M_*) + \alpha_t \Psi_* \right] V^{-1} \right\}}{(2\pi)^{\frac{np}{2}} |\Sigma_L / \alpha_t k_*|^{\frac{n}{2}} 2^{\frac{(m_{A,*} - n - 1)n}{2}} \Gamma_n \left(\frac{m_{A,*} - n - 1}{2} \right) |V|^{\frac{m_{A,*} + p}{2}}} \\
&= \frac{|\Psi_{A,*}|^{\frac{m_{A,*} - n - 1}{2}} \exp \left\{ -\frac{1}{2} \text{tr} \left[(B_t - M_*)' \Sigma_{A,*}^{-1} (B_t - M_*) + \Psi_{A,*} \right] V_t^{-1} \right\}}{(2\pi)^{\frac{np}{2}} |\Sigma_{A,*}|^{\frac{n}{2}} 2^{\frac{(m_{A,*} - n - 1)n}{2}} \Gamma_n \left(\frac{m_{A,*} - n - 1}{2} \right) |V_t|^{\frac{m_{A,*} + p}{2}}}, \tag{A.32}
\end{aligned}$$

where we defined $m_{A,*} = \alpha_t (m_* + p) = \alpha_t m_d$, $\Sigma_{A,*} = \Sigma_* / \alpha_t$, $\Psi_{A,*} = \alpha_t \Psi_*$, and the constant

$$C(\alpha_t) = \frac{|\Psi_{A,*}|^{\frac{m_{A,*} - n - 1}{2}} (2\pi)^{\frac{\alpha_t np}{2}} |\Sigma_*|^{\frac{\alpha_t n}{2}} 2^{\frac{\alpha_t (m_* - n - 1)n}{2}} \Gamma_n \left(\frac{m_* - n - 1}{2} \right)^{\alpha_t}}{(2\pi)^{\frac{np}{2}} |\Sigma_{A,*}|^{\frac{n}{2}} 2^{\frac{(m_{A,*} - n - 1)n}{2}} \Gamma_n \left(\frac{m_{A,*} - n - 1}{2} \right) |\Psi_*|^{\frac{\alpha_t (m_* - n - 1)}{2}}}. \tag{A.33}$$

The posterior predictive under the alternative is

$$p_A(Y_t|\mathcal{D}_{t-1}) = \frac{|\Psi_{A,*}|^{\frac{m_{A,*}-n-1}{2}} k_{A,*}^{\frac{np}{2}} \Gamma_n\left(\frac{m_{A,d}-n-1}{2}\right)}{(\pi)^{\frac{np}{2}} k_{A,d}^{\frac{np}{2}} |\Psi_{A,d}|^{\frac{m_{A,d}-n-1}{2}} \Gamma_n\left(\frac{m_{A,*}-n-1}{2}\right) |\Sigma_L|^{\frac{n}{2}}} \quad (\text{A.34})$$

which is a Student-t distribution $\mathcal{T}(m_{A,*} - 2n, M_*, \Sigma_L, L_{A,*})$ where $L_{A,*} = \Psi_{A,*} k_{A,d} / k_{A,*}$, where $k_{A,*} = \alpha_t k_*$, $k_{A,d} = \alpha_t k_* + 1$, $m_{A,d} = m_{A,*} + p$, and $\Psi_{A,d} = \Psi_{A,*} + \alpha_t k_* (M_* - Y_t)' \Sigma_L^{-1} (M_* - Y_t) / ((\alpha_t k_* + 1))$ following an argument analogous to the one used for the numerator. We conclude with the BF, which is the ratio of $p(Y_t|\mathcal{D}_{t-1})$ given in Eq. A.32 and $p_A(Y_t|\mathcal{D}_{t-1})$ given in Eq. A.34:

$$H_t(\alpha_t) = G |\Psi_{A,d}|^{\frac{m_{A,d}-n-1}{2}} |\Psi_d|^{-\frac{m_d-n-1}{2}} \quad (\text{A.35})$$

where

$$G = \frac{|\Psi_*|^{\frac{m_*-n-1}{2}} k_*^{\frac{np}{2}} \Gamma_n\left(\frac{m_d-n-1}{2}\right) k_{A,d}^{\frac{np}{2}} \Gamma_n\left(\frac{m_{A,*}-n-1}{2}\right)}{k_d^{\frac{np}{2}} \Gamma_n\left(\frac{m_*-n-1}{2}\right) |\Psi_{A,*}|^{\frac{m_{A,*}-n-1}{2}} k_{A,*}^{\frac{np}{2}} \Gamma_n\left(\frac{m_{A,d}-n-1}{2}\right)}. \quad (\text{A.36})$$

$H_t(\alpha_t)$ goes to 1 when $\alpha_t \rightarrow 1^-$, $H_t(\alpha_t) \rightarrow \infty$ as $\alpha_t \rightarrow \underline{\alpha}^+$, due to the fact that $G \rightarrow \infty$ as $\alpha_t \rightarrow \underline{\alpha}^+$ because $\Gamma_n\left(\frac{m_{A,*}-n-1}{2}\right) \rightarrow \infty$. □

Proof of Corollary 1. i) In the first case

$$C(\alpha_t) = \frac{\alpha_t^{\frac{p}{2}} (2\pi)^{\frac{\alpha_t np}{2}} |\Sigma_*|^{\frac{\alpha_t n}{2}} |V|^{\frac{\alpha_t p}{2}}}{(2\pi)^{\frac{np}{2}} |\Sigma_*|^{\frac{n}{2}} |V|^{\frac{p}{2}}} \quad (\text{A.37})$$

which goes to zero for $\alpha_t \rightarrow 0^+$ and goes to one for $\alpha_t \rightarrow 1^-$.

ii) In the second case, from the proof of Prop. 4 the constant is

$$C(\alpha_t) = \frac{|\Psi_{A,*}|^{\frac{m_{A,*}-n-1}{2}} (2\pi)^{\frac{\alpha_t np}{2}} |\Sigma_*|^{\frac{\alpha_t n}{2}} 2^{\frac{\alpha_t(m_*-n-1)n}{2}} \Gamma_n\left(\frac{m_*-n-1}{2}\right)^{\alpha_t}}{(2\pi)^{\frac{np}{2}} |\Sigma_{A,*}|^{\frac{n}{2}} 2^{\frac{(m_{A,*}-n-1)n}{2}} \Gamma_n\left(\frac{m_{A,*}-n-1}{2}\right) |\Psi_*|^{\frac{\alpha_t(m_*-n-1)}{2}}}. \quad (\text{A.38})$$

Note that $m_{A,*} = \alpha_t m_d \rightarrow 0$ for $\alpha_t \rightarrow 0^+$ and the argument of the multivariate Gamma function $\Gamma_n((m_{A,*} - n - 1)/2)$ becomes negative. To guarantee the positivity of the constant, the condition $\Gamma_n((m_{A,*} - n - 1)/2) > 0$ is needed noticing that $m_{A,*} \leq m_* \leq m_d \leq m_{A,d}$. A sufficient condition is that the n -th element of the multivariate gamma product $\Gamma((m_{A,*} - n - 1)/2)$ is strictly positive, that is $(m_{A,*} - n - 1)/2 + (1 - n)/2 = (\alpha_t(m + Tp + p) - p - n - 1 + 1 - n)/2 > 0$, or equivalently $\alpha_t > \underline{\alpha}$ where $\underline{\alpha} = (p + 2n)/m_d$. Under this condition $C(\alpha_t) \rightarrow 0^+$ for $\alpha_t \rightarrow \underline{\alpha}^+$. Moreover, $C(\alpha_t) \rightarrow 1$ for $\alpha_t \rightarrow 1^-$. □

Proof of Proposition 5. i) Define:

$$\Sigma_H = \left[\left(\frac{1}{\alpha_t} - 1 \right) (\Sigma_L + \Sigma_*)^{-1} \Sigma_* \left(\Sigma_L + \frac{\Sigma_*}{\alpha_t} \right)^{-1} \right]^{-1} \quad (\text{A.39})$$

$$= \left(\frac{1}{1 - \alpha_t} \right) (\alpha_t \Sigma_L + \Sigma_*) \Sigma_*^{-1} (\Sigma_L + \Sigma_*) \quad (\text{A.40})$$

and $\kappa_t(\alpha_t) = \left| \Sigma_L + \frac{\Sigma_*}{\alpha_t} \right|^{\frac{n}{2}} |\Sigma_L + \Sigma_*|^{-\frac{n}{2}} = |\Sigma_{A,d}|^{\frac{n}{2}} |\Sigma_d|^{-\frac{n}{2}}$. Since: $(\Sigma_L + \Sigma_*)^{-1} - (\Sigma_L + \Sigma_*/\alpha_t)^{-1} = (\Sigma_L + \Sigma_*)^{-1} [\Sigma_*/\alpha_t - \Sigma_*] (\Sigma_L + \Sigma_*/\alpha_t)^{-1} = (1/(\alpha_t - 1)) (\Sigma_L + \Sigma_*)^{-1} \Sigma_* (\Sigma_L + \Sigma_*/\alpha_t)^{-1}$, then the BF can be written as

$$H_t = \kappa_t(\alpha_t) \exp \left\{ -\frac{1}{2} \text{tr} \left[\left[(\Sigma_L + \Sigma_*)^{-1} - \left(\Sigma_L + \frac{\Sigma_*}{\alpha_t} \right)^{-1} \right] (Y_t - M_*) V^{-1} (Y_t - M_*)' \right] \right\} \quad (\text{A.41})$$

$$= \kappa_t(\alpha_t) \exp \left\{ -\frac{1}{2} \text{tr} \left[\Sigma_H^{-1} (Y_t - M_*) V^{-1} (Y_t - M_*)' \right] \right\}. \quad (\text{A.42})$$

The assumptions $(1/\alpha_t - 1) > 0$, $\Sigma_* > 0$, $\Sigma_L > 0$, $\Sigma_L + \Sigma_* > 0$ imply the eigenvalues of $\Sigma_L + \Sigma_*$ and of $\Sigma_L + \Sigma_*/\alpha_t$ are greater than zero, and the product of the three matrices $(\Sigma_L + \Sigma_*)^{-1}$, Σ_* , $(\Sigma_L + \Sigma_*/\alpha_t)^{-1}$ has positive eigenvalues. Moreover the assumption Σ_L , Σ_* are symmetric and positive definite implies $((1 - \alpha_t)^{-1}) (\Sigma_L + \Sigma_*)^{-1} \Sigma_* (\Sigma_L + \Sigma_*/\alpha_t)^{-1}$ is symmetric and has positive eigenvalues, it is positive definite. Since the argument of the exponential function is negative, we get an upper bound for H_t :

$$H_t \leq \kappa_t(\alpha_t) \quad (\text{A.43})$$

such that $\lim_{\alpha_t \rightarrow 1^-} \kappa_t = 1$. From Th. 7.7 in Zhang (2011) it follows:

$$|\Sigma_L + \Sigma_*| \geq |\Sigma_L| + |\Sigma_*| \Leftrightarrow \left| \Sigma_L + \frac{\Sigma_*}{\alpha_t} \right| \geq |\Sigma_L| + \left| \frac{\Sigma_*}{\alpha_t} \right| = |\Sigma_L| + \frac{|\Sigma_*|}{\alpha_t^p}, \quad (\text{A.44})$$

or equivalently

$$\left| \Sigma_L + \frac{\Sigma_*}{\alpha_t} \right| - |\Sigma_L + \Sigma_*| \geq \frac{|\Sigma_*|}{\alpha_t^p} - |\Sigma_*| = \left(\frac{1}{\alpha_t^p} - 1 \right) |\Sigma_*| \geq 0 \quad (\text{A.45})$$

and $\kappa_t \geq 1$. By Jacobi's formula on the derivative of the matrix determinant (Magnus and Neudecker, 1999), we obtain

$$\begin{aligned} \partial_{\alpha_t} \kappa_t(\alpha_t) &= \frac{n}{2} \frac{\left| \Sigma_L + \frac{\Sigma_*}{\alpha_t} \right|^{\frac{n}{2}-1}}{|\Sigma_L + \Sigma_*|^{\frac{n}{2}}} \frac{d}{d\alpha_t} \left| \Sigma_L + \frac{\Sigma_*}{\alpha_t} \right| \\ &= \frac{n}{2} \kappa_t(\alpha_t) \text{tr} \left[\left(\Sigma_L + \frac{\Sigma_*}{\alpha_t} \right)^{-1} \left(-\frac{\Sigma_*}{\alpha_t^2} \right) \right] = -\frac{n}{2\alpha_t^2} \kappa_t(\alpha_t) \text{tr} \left[\left(\Sigma_L + \frac{\Sigma_*}{\alpha_t} \right)^{-1} \Sigma_* \right]. \end{aligned} \quad (\text{A.46})$$

Note that $\left(\Sigma_L + \frac{\Sigma_*}{\alpha_t} \right)^{-1} > 0$ and $\Sigma_* > 0$, thus the trace of the product of positive matrices is positive and $\partial_{\alpha_t} \kappa_t(\alpha_t) < 0$ for every α_t . In conclusion, we showed $\kappa_t(\alpha_t)$ is a decreasing function that goes to infinite when $\alpha_t \rightarrow 0^+$ and goes to 1 when $\alpha_t \rightarrow 1^-$.

ii) From its definition it trivially follows $\lim_{\alpha_t \rightarrow 1^-} H_t(\alpha_t) = 1$. Consider now the term inside the trace and write:

$$\left(\Sigma_L + \frac{\Sigma_*}{\alpha_t} \right)^{-1} = \Sigma_L^{-1} - \Sigma_L^{-1} \left[\Sigma_L^{-1} + \left(\frac{\Sigma_*}{\alpha_t} \right)^{-1} \right]^{-1} \Sigma_L^{-1} = \Sigma_L^{-1} - \Sigma_L^{-1} [\Sigma_L^{-1} + \alpha_t \Sigma_*^{-1}]^{-1} \Sigma_L^{-1}. \quad (\text{A.47})$$

This term goes to zero as $\alpha_t \rightarrow 0^+$, and the exponential in (A.42) has a bounded limit since the other terms don't depend by α_t . Furthermore, due to the:

$$\left| \Sigma_L + \frac{\Sigma_*}{\alpha_t} \right| = \left| \frac{\alpha_t \Sigma_L + \Sigma_*}{\alpha_t} \right| = \frac{1}{\alpha_t^p} |\alpha_t \Sigma_L + \Sigma_*|. \quad (\text{A.48})$$

We have:

$$\lim_{\alpha_t \rightarrow 0^+} \left| \Sigma_L + \frac{\Sigma_*}{\alpha_t} \right|^{\frac{n}{2}} = \lim_{\alpha_t \rightarrow 0^+} \frac{1}{\alpha_t^{\frac{np}{2}}} |\alpha_t \Sigma_L + \Sigma_*|^{\frac{n}{2}} = \infty. \quad (\text{A.49})$$

From (A.42) and (A.49) it follows that $\lim_{\alpha_t \rightarrow 0^+} H_t(\alpha_t) = \infty$.

iii) Let us define $\tilde{A} = (Y_t - M_*) V^{-1} (Y_t - M_*)'$ and $\kappa_t(\alpha_t) = |\Sigma_L + \Sigma_*/\alpha_t|^{\frac{n}{2}} |\Sigma_L + \Sigma_*|^{-\frac{n}{2}}$ then the derivative of the BF is

$$\partial_{\alpha_t} H_t(\alpha_t) = \exp \left\{ -\frac{1}{2} \text{tr} \left[\Sigma_H^{-1} \tilde{A} \right] \right\} \frac{d}{d\alpha_t} \kappa_t(\alpha_t) + \kappa_t(\alpha_t) \frac{d}{d\alpha_t} \exp \left\{ -\frac{1}{2} \text{tr} \left[\Sigma_H^{-1} \tilde{A} \right] \right\}. \quad (\text{A.50})$$

From i) we know:

$$\partial_{\alpha_t} \kappa_t(\alpha_t) = -\frac{n \left| \Sigma_L + \frac{\Sigma_*}{\alpha_t} \right|^{\frac{n}{2}}}{2\alpha_t^2 \left| \Sigma_L + \Sigma_* \right|^{\frac{n}{2}}} \text{tr} \left[\left(\Sigma_L + \frac{\Sigma_*}{\alpha_t} \right)^{-1} \Sigma_* \right] \quad (\text{A.51})$$

and since $\Sigma_H^{-1} = (1/\alpha_t - 1)(\Sigma_L + \Sigma_*)^{-1} \Sigma_* (\Sigma_L + \Sigma_*/\alpha_t)^{-1} = (1 - \alpha_t) \Upsilon$, where we define $\Upsilon = (\Sigma_L + \Sigma_*)^{-1} \Sigma_* (\alpha_t \Sigma_L + \Sigma_*)^{-1}$, then we obtain

$$\begin{aligned} \frac{d}{d\alpha_t} \exp \left\{ -\frac{1}{2} \text{tr} \left[(1 - \alpha_t) \Upsilon \tilde{A} \right] \right\} &= \exp \left\{ -\frac{1}{2} \text{tr} \left[\Sigma_H^{-1} \tilde{A}(\alpha_t) \right] \right\} \left\{ \frac{1}{2} \text{tr} \left[\Upsilon \tilde{A} \right] + \frac{(\alpha_t - 1)}{2} \frac{d}{d\alpha_t} \text{tr} \left[\Upsilon \tilde{A} \right] \right\} \\ &= \exp \left\{ -\frac{1}{2} \text{tr} \left[\Sigma_H^{-1} \tilde{A}(\alpha_t) \right] \right\} \left\{ \frac{1}{2} \text{tr} \left[\Upsilon \tilde{A} \right] + \frac{(\alpha_t - 1)}{2} \text{tr} \left[(\Sigma_L + \Sigma_*)^{-1} \Sigma_* \frac{d}{d\alpha_t} (\alpha_t \Sigma_L + \Sigma_*)^{-1} \tilde{A} \right] \right\}. \end{aligned}$$

The derivative in the last line is:

$$\frac{d}{d\alpha_t} (\alpha_t \Sigma_L + \Sigma_*)^{-1} = -(\alpha_t \Sigma_L + \Sigma_*)^{-1} \Sigma_L (\alpha_t \Sigma_L + \Sigma_*)^{-1}. \quad (\text{A.52})$$

See Abadir and Magnus (2005) and Magnus and Neudecker (1999). Thus, we get the result:

$$\begin{aligned} \partial_{\alpha_t} H_t(\alpha_t) &= \kappa_t(\alpha_t) \exp \left\{ -\frac{1}{2} \text{tr} \left[\Sigma_H^{-1} \tilde{A} \right] \right\} \left\{ -\frac{n}{2\alpha_t^2} \text{tr} \left[\left(\Sigma_L + \frac{\Sigma_*}{\alpha_t} \right)^{-1} \Sigma_* \right] + \right. \\ &\quad \left. \frac{1}{2} \text{tr} \left[\Upsilon \tilde{A} \right] + \frac{(1 - \alpha_t)}{2} \text{tr} \left[\Upsilon \Sigma_L (\alpha_t \Sigma_L + \Sigma_*)^{-1} \tilde{A} \right] \right\} = \\ &= \frac{H_t(\alpha_t)}{2} \text{tr} \left\{ -\frac{n}{\alpha_t^2} \left(\Sigma_L + \frac{\Sigma_*}{\alpha_t} \right)^{-1} \Sigma_* + \Upsilon \tilde{A} + (1 - \alpha_t) \Upsilon \Sigma_L (\alpha_t \Sigma_L + \Sigma_*)^{-1} \tilde{A} \right\}. \end{aligned}$$

□

Proof of Proposition 6. Note that from Prop. 4 the BF can be written as $H_t(\alpha_t) = \Psi(\alpha_t) \kappa_{1t}(\alpha_t) \kappa_{2t}(\alpha_t)$ where the first term depend on Y_t through \tilde{Y}

$$\Phi(\alpha_t) = \frac{|\Psi_{A,d}|^{\frac{m_{A,d}-n-1}{2}} |\Psi_*|^{\frac{m_*-n-1}{2}}}{|\Psi_d|^{\frac{m_d-n-1}{2}} |\Psi_{A,*}|^{\frac{m_{A,*}-n-1}{2}}} = \frac{|\Psi_{A,d}|^{\frac{\alpha_t m_d - n - 1}{2}} |\Psi_*|^{\frac{m_* - n - 1}{2}}}{|\Psi_d|^{\frac{m_d - n - 1}{2}} |\Psi_{A,*}|^{\frac{\alpha_t m_d - p - n - 1}{2}}} \quad (\text{A.53})$$

and the remaining two terms depend only on α_t

$$\kappa_{1t}(\alpha_t) = \left(\frac{k_* k_{A,d}}{k_d k_{A,*}} \right)^{\frac{np}{2}}, \quad \kappa_{2t}(\alpha_t) = \frac{\Gamma_n \left(\frac{m_d - n - 1}{2} \right) \Gamma_n \left(\frac{m_{A,*} - n - 1}{2} \right)}{\Gamma_n \left(\frac{m_* - n - 1}{2} \right) \Gamma_n \left(\frac{m_{A,d} - n - 1}{2} \right)}. \quad (\text{A.54})$$

First, we prove that $\Phi(\alpha_t) < 1$ to find the upper bound to the BF and then study the properties of the remaining terms to find the limits of the BF.

i) Define the positive definite matrix $E = (M_* - Y_t)' \Sigma^{-1} (M_* - Y_t)$ and write

$$\Psi_{A,d} = \alpha_t \Psi_* + \frac{\alpha_t k_*}{\alpha_t k_* + 1} E = \alpha_t \Psi_* \left(I_n + \frac{\alpha_t k_*}{\alpha_t k_* + 1} \Psi_*^{-1} E \right) \quad (\text{A.55})$$

$$\Psi_d = \Psi_* + \frac{k_*}{k_* + 1} E = \Psi_* \left(I_n + \frac{k_*}{k_* + 1} \Psi_*^{-1} E \right) \quad (\text{A.56})$$

and $\Psi_{A,*} = \alpha_t \Psi_*$. Substituting in the expression of $\Phi_t(\alpha_t)$ their determinants

$$|\Psi_{A,d}|^{\frac{\alpha_t m_d - n - 1}{2}} = \alpha_t^{\frac{n(\alpha_t m_d - n - 1)}{2}} |\Psi_*|^{\frac{\alpha_t m_d - n - 1}{2}} \left| I_n + \frac{\alpha_t k_*}{\alpha_t k_* + 1} \Psi_*^{-1} E \right|^{\frac{\alpha_t m_d - n - 1}{2}} \quad (\text{A.57})$$

$$|\Psi_d|^{\frac{m_d - n - 1}{2}} = |\Psi_*|^{\frac{m_d - n - 1}{2}} \left| I_n + \frac{k_*}{k_* + 1} \Psi_*^{-1} E \right|^{\frac{m_d - n - 1}{2}} \quad (\text{A.58})$$

$$|\Psi_{A,*}|^{\frac{\alpha_t m_d - p - n - 1}{2}} = \alpha_t^{\frac{n(\alpha_t m_d - p - n - 1)}{2}} |\Psi_*|^{\frac{\alpha_t m_d - p - n - 1}{2}} \quad (\text{A.59})$$

and using $m_d = m_* + p$ yield

$$\begin{aligned} \Psi(\alpha_t) &= \frac{\alpha_t^{\frac{np}{2}} |\Psi_*|^{\frac{\alpha_t m_d + m_* - m_d - \alpha_t m_d + p}{2}} \left| I_n + \frac{\alpha_t k_*}{\alpha_t k_* + 1} \Psi_*^{-1} E \right|^{\frac{\alpha_t m_d - n - 1}{2}}}{\left| I_n + \frac{k_*}{k_* + 1} \Psi_*^{-1} E \right|^{\frac{m_d - n - 1}{2}}} \\ &= \alpha_t^{\frac{np}{2}} \left| I_n + \frac{\alpha_t k_*}{\alpha_t k_* + 1} \Psi_*^{-1} E \right|^{\frac{\alpha_t m_d - n - 1}{2}} \left| I_n + \frac{k_*}{k_* + 1} \Psi_*^{-1} E \right|^{-\frac{m_d - n - 1}{2}}. \end{aligned} \quad (\text{A.60})$$

In the sense of positive definite matrices, the following inequality is satisfied:

$$I_n + \frac{\alpha_t k_*}{\alpha_t k_* + 1} \Psi_*^{-1} E < I_n + \frac{k_*}{k_* + 1} \Psi_*^{-1} E \quad (\text{A.61})$$

since $\alpha_t < 1$, and $\alpha_t k_*/(\alpha_t k_* + 1) - k_*/(k_* + 1) = k_*(\alpha_t - 1)/((\alpha_t k_* + 1)(k_* + 1)) < 0$. From Th. 7.8 in Zhang (2011) on the relationship between determinants of positive definite matrices we obtain: $|I_n + \alpha_t k_*/(\alpha_t k_* + 1) \Psi_*^{-1} E| < |I_n + k_*/(k_* + 1) \Psi_*^{-1} E|$ which implies

$$\left| I_n + \frac{\alpha_t k_*}{\alpha_t k_* + 1} \Psi_*^{-1} E \right|^{\frac{\alpha_t m_d - n - 1}{2}} < \left| I_n + \frac{k_*}{k_* + 1} \Psi_*^{-1} E \right|^{\frac{m_d - n - 1}{2}} \quad (\text{A.62})$$

since the arguments of the power functions are larger than one. For the first argument, this can be proved using Th. 7.7 and 7.8 in Zhang (2011) and the positive definiteness:

$$\left| I_n + \frac{\alpha_t k_*}{\alpha_t k_* + 1} \Psi_*^{-1} E \right| > |I_n| + \left| \frac{\alpha_t k_*}{\alpha_t k_* + 1} \Psi_*^{-1} E \right| > 1. \quad (\text{A.63})$$

For the second argument, the same inequality for determinants can be used together with $\alpha_t m_d + n - 1 > 0$ and the assumption $\alpha_t > \underline{\alpha} = (2n + p)/m_d$. It follows that $\Phi(\alpha_t) < 1$ and $H_t(\alpha_t) < \kappa_{1t}(\alpha_t) \kappa_{2t}(\alpha_t)$ and the upper bound $\kappa_t(\alpha_t) = \kappa_{1t}(\alpha_t) \kappa_{2t}(\alpha_t)$ does not depend on Y_t and \bar{Y} .

We show now that $\kappa_t(\alpha_t)$ is monotone decreasing since $\kappa_{1t}(\alpha_t)$ and $\kappa_{2t}(\alpha_t)$ are monotone decreasing. Regarding the first derivative of $\kappa_{1t}(\alpha_t)$

$$\partial_{\alpha_t} \left(\frac{\alpha_t k_* + 1}{\alpha_t (k_* + 1)} \right)^{\frac{np}{2}} = \frac{np}{2} \left(\frac{\alpha_t k_* + 1}{\alpha_t (k_* + 1)} \right)^{\frac{np}{2}-1} (-\alpha_t^{-2} (k_* + 1)^{-1}) < 0. \quad (\text{A.64})$$

Regarding the first derivative of $\kappa_2(\alpha_t)$ consider the following preliminary results, which can be easily proved using the properties of the multivariate function of order n (e.g., see Gupta and Nagar, 1999)

$$\partial_{\alpha_t} \Gamma_n \left(\frac{m_{A,*} - n - 1}{2} \right) = \frac{m_* + p}{2} \Gamma_n \left(\frac{m_{A,*} - n - 1}{2} \right) \sum_{i=1}^n \gamma_n \left(\frac{m_{A,*} - n - 1}{2} + \frac{1-i}{2} \right) \quad (\text{A.65})$$

$$\partial_{\alpha_t} \Gamma_n \left(\frac{m_{A,d} - n - 1}{2} \right) = \frac{m_* + p}{2} \Gamma_n \left(\frac{m_{A,d} - n - 1}{2} \right) \sum_{i=1}^n \gamma_n \left(\frac{m_{A,d} - n - 1}{2} + \frac{1-i}{2} \right) \quad (\text{A.66})$$

where $\gamma_n(a)$ is the multivariate digamma function of order n . Note that since $m_{A,d} \geq m_{A,*}$ the smallest argument of γ_n in Eq. A.65-A.66 is $(\alpha_t(m_* + p) - p - n - 1 + 1 - n)/2$ which is positive for $\alpha_t \geq \underline{\alpha} = (p + 2n)/(m_* + p)$. The first derivative of $\kappa_{2t}(\alpha_t)$ becomes:

$$\begin{aligned} \partial_{\alpha_t} \kappa_2(\alpha_t) &= \frac{\Gamma_n \left(\frac{m_d - n - 1}{2} \right)}{\Gamma_n \left(\frac{m_* - n - 1}{2} \right)} \partial_{\alpha_t} \frac{\Gamma_n \left(\frac{m_{A,*} - n - 1}{2} \right)}{\Gamma_n \left(\frac{m_{A,d} - n - 1}{2} \right)} \\ &= \frac{\Gamma_n \left(\frac{m_d - n - 1}{2} \right)}{\Gamma_n \left(\frac{m_* - n - 1}{2} \right)} \frac{(m_* + p)}{2} \Gamma_n \left(\frac{m_{A,*} - n - 1}{2} \right) \Gamma_n \left(\frac{m_{A,d} - n - 1}{2} \right) \\ &\quad \times \frac{\sum_{i=1}^n \left[\gamma_n \left(\frac{m_{A,*} - n - 1}{2} + \frac{i-1}{2} \right) - \gamma_n \left(\frac{m_{A,d} - n - 1}{2} + \frac{i-1}{2} \right) \right]}{\Gamma_n \left(\frac{m_{A,d} - n - 1}{2} \right)^2}. \end{aligned} \quad (\text{A.67})$$

Since $m_{A,d} \geq m_{A,*}$ and the $\gamma(a)$ function is an increasing function for $a \geq 0$ the differences of gamma functions at the numerator are negative, provided $\alpha_t \geq \underline{\alpha}$. This means that $\partial_{\alpha_t} \kappa_2(\alpha_t) \leq 0$.

Finally, the limits of $\kappa_{1t}(\alpha_t)$ as $\alpha_t \rightarrow 1^-$ is 1, since $\kappa_{1t}(\alpha_t)$ is decreasing and the lower bound is

$$\kappa_{1t}(\alpha_t) = \left(\frac{k_* k_{A,d}}{k_d k_{A,*}} \right)^{\frac{np}{2}} = \left(\frac{k_* (\alpha_t k_* + 1)}{(k_* + 1) \alpha_t k_*} \right)^{\frac{np}{2}} = \left(\frac{\alpha_t k_* + 1}{\alpha_t (k_* + 1)} \right)^{\frac{np}{2}} \geq 1 \quad (\text{A.68})$$

whereas $\kappa_{1t}(\alpha_t)$ converges to $(\underline{\alpha} k_* / (\underline{\alpha} (k_* + 1)))^{np/2}$ that is larger than 1 if $\alpha_t \rightarrow \underline{\alpha}$. We also get $\kappa_{2t}(\alpha_t) \rightarrow 1$ as $\alpha_t \rightarrow 1^-$, since $m_{A,d} = m_d$ and $m_{A,*} = m_*$. For $\alpha_t \rightarrow \underline{\alpha}$ we get $\kappa_2(\alpha_t) \rightarrow +\infty$ because $\Gamma_n[(m_{A,*} - n - 1)/2] \rightarrow +\infty$ for $\alpha_t \rightarrow \underline{\alpha}$ and $m_{A,*} \leq m_* \leq m_d \leq m_{A,d}$ for $\alpha_t \leq 1$.

In conclusion we get that $\kappa(\alpha_t) = \kappa_{1t}(\alpha_t) \kappa_{2t}(\alpha_t)$ is the product of two functions monotone decreasing, thus $\kappa(\alpha_t)$ is monotone decreasing and $\kappa(\alpha_t) \rightarrow \infty$ as $\alpha \rightarrow \underline{\alpha}_t^+$ and $\kappa(\alpha_t) \rightarrow 1$ as $\alpha \rightarrow 1^-$.

ii) From its definition it trivially follows $\lim_{\alpha_t \rightarrow 1^-} H_t(\alpha_t) = 1$. Regarding the second limit, note that all terms at the numerator and denominator of $H_t(\alpha_t)$ have bounded limit for $\alpha_t \rightarrow \underline{\alpha}$, expect for $\Gamma_n \left(\frac{m_{A,*} - n - 1}{2} \right)$ which is unbounded since $m_{A,*} - n - 1 = \alpha_t(m_* + p) - p - n - 1 \rightarrow 0$. Thus $H_t(\alpha_t) \rightarrow \infty$ for $\alpha_t \rightarrow \underline{\alpha}$.

iii) We factorize out the terms that depend on α_t and write H_t as $H_t(\alpha_t) = L(a_1 a_2 a_3) / (b_1 b_2 b_3)$, where $L = |\Psi|_*^{\frac{m_* - n - 1}{2}} k_*^{\frac{np}{2}} \Gamma_n \left(\frac{m_d - n - 1}{2} \right) k_d^{\frac{-np}{2}} |\Psi_d|^{-\frac{m_d - n - 1}{2}} \Gamma_n \left(\frac{m_* - n - 1}{2} \right)^{-1}$ and

$$a_1 = k_{A,d}^{\frac{np}{2}}; a_2 = \Gamma_n \left(\frac{m_{A,*} - n - 1}{2} \right); a_3 = |\Psi_{A,d}|^{\frac{m_{A,d} - n - 1}{2}};$$

$$b_1 = k_{A,*}^{\frac{np}{2}}; b_2 = \Gamma_n \left(\frac{m_{A,d} - n - 1}{2} \right); b_3 = |\Psi_{A,*}|^{\frac{m_{A,*} - n - 1}{2}} \quad (\text{A.69})$$

The first derivative of $H(\alpha_t)$ is: $\partial_{\alpha_t} H(\alpha_t) = L(b_1 b_2 b_3)^{-2} \left((a'_1 a_2 a_3 + a_1 a'_2 a_3 + a_1 a_2 a'_3) b_1 b_2 b_3 - (b'_1 b_2 b_3 + b_1 b'_2 b_3 + b_1 b_2 b'_3) a_1 a_2 a_3 \right)$ where

$$a'_1 = \partial_{\alpha_t} k_{A,d}^{\frac{np}{2}} = \partial_{\alpha_t} (\alpha_t k_* + 1)^{\frac{np}{2}} = \frac{np}{2} k_* (\alpha_t k_* + 1)^{\frac{np-2}{2}} \quad (\text{A.70})$$

$$\begin{aligned} a'_2 &= \partial_{\alpha_t} \Gamma_n \left(\frac{m_{A,*} - n - 1}{2} \right) = \partial_{\alpha_t} \Gamma_n \left[\frac{\alpha_t (m_* + p) - p - n - 1}{2} \right] \\ &= (m_* + p) \Gamma_n \left(\frac{m_{A,*} - n - 1}{2} \right) \sum_{i=1}^n \psi \left(\frac{m_{A,*} - n - i}{2} \right) \end{aligned}$$

$$\begin{aligned} a'_3 &= \partial_{\alpha_t} |\Psi_{A,d}|^{\frac{m_{A,d} - n - 1}{2}} \\ &= |\Psi_{A,d}|^{\frac{m_{A,d} - n - 1}{2}} \left[\partial_{\alpha_t} \left(\frac{m_{A,d} - n - 1}{2} \right) \log |\Psi_{A,d}| + \frac{m_{A,d} - n - 1}{2} \frac{\partial_{\alpha_t} |\Psi_{A,d}|}{|\Psi_{A,d}|} \right] \end{aligned}$$

with $\psi(a)$ the digamma function and

$$\begin{aligned} \partial_{\alpha_t} \frac{m_{A,d} - n - 1}{2} &= \frac{m_* + p}{2} \\ \partial_{\alpha_t} |\Psi_{A,d}| &= |\Psi_{A,d}| \operatorname{tr} \left[\Psi_{A,d}^{-1} \left(\Psi_* + \frac{k_* (M_* - Y_t)' \Sigma^{-1} (M_* - Y_t)}{(\alpha_t k_* + 1)^2} \right) \right]. \end{aligned}$$

The remaining three terms have derivatives: b'_1 :

$$b'_1 = \partial_{\alpha_t} k_{A,*}^{\frac{np}{2}} = \partial_{\alpha_t} \alpha_t k_*^{\frac{np}{2}} = \frac{np}{2} k_* k_{A,*}^{\frac{np-2}{2}} = \frac{np}{2} k_* (\alpha_t k_*)^{\frac{np-2}{2}} \quad (\text{A.71})$$

$$\begin{aligned} b'_2 &= \partial_{\alpha_t} \Gamma_n \left(\frac{m_{A,d} - n - 1}{2} \right) = \partial_{\alpha_t} \Gamma_n \left[\frac{\alpha_t (m_* + p) - p + p - n - 1}{2} \right] \\ &= (m_* + p) \Gamma_n \left(\frac{m_{A,d} - n - 1}{2} \right) \sum_{i=1}^n \psi \left(\frac{m_{A,d} - n - i}{2} \right) \end{aligned} \quad (\text{A.72})$$

$$\begin{aligned} b'_3 &= \partial_{\alpha_t} |\Psi_{A,*}|^{\frac{m_{A,*} - n - 1}{2}} \\ &= |\Psi_{A,*}|^{\frac{m_{A,*} - n - 1}{2}} \left[\partial_{\alpha_t} \left(\frac{m_{A,*} - n - 1}{2} \right) \log |\Psi_{A,*}| + \frac{m_{A,*} - n - 1}{2} \frac{\partial_{\alpha_t} |\Psi_{A,*}|}{|\Psi_{A,*}|} \right], \end{aligned} \quad (\text{A.73})$$

where $\partial_{\alpha_t} (m_{A,*} - n - 1)/2 = (m_* + p)/2$ and $\partial_{\alpha_t} |\Psi_{A,*}| = |\Psi_{A,*}| \operatorname{tr} (\Psi_{A,*}^{-1} \Psi_*) = |\Psi_{A,*}| \alpha_t^{-1}$ since $\Psi_{A,*} = \alpha_t \Psi_*$. In conclusion, the derivative of $H_t(\alpha_t)$ is:

$$\partial_{\alpha_t} H_t(\alpha_t) = L \left[|\Psi_{A,*}|^{\frac{m_{A,*} - n - 1}{2}} k_{A,*}^{\frac{np}{2}} \Gamma_n \left(\frac{m_{A,d} - n - 1}{2} \right) \right]^{-2} (A'_1 B_1 - A_1 B'_1) \quad (\text{A.74})$$

where $A_1 = a_1 a_2 a_3$, $B_h = b_1 b_2 b_3$ and their derivatives A_1' , B_1' are defined as:

$$A_1 = k_{A,d}^{\frac{np}{2}} \Gamma_n \left(\frac{m_{A,*} - n - 1}{2} \right) |\Psi_{A,d}|^{\frac{m_{A,d} - n - 1}{2}} \quad (\text{A.75})$$

$$B_1 = k_{A,*}^{\frac{np}{2}} \Gamma_n \left(\frac{m_{A,d} - n - 1}{2} \right) |\Psi_{A,*}|^{\frac{m_{A,*} - n - 1}{2}} \quad (\text{A.76})$$

$$\begin{aligned} A'_1 &= a'_1 a_2 a_3 + a_1 a'_2 a_3 + a_1 a_2 a'_3 \\ &= \frac{np}{2} k_* k_{A,d}^{\frac{np-2}{2}} \Gamma_n \left(\frac{m_{A,*} - n - 1}{2} \right) |\Psi_{A,d}|^{\frac{m_{A,d} - n - 1}{2}} + \\ &\quad k_{A,d}^{\frac{np}{2}} (m_* + p) \Gamma_n \left(\frac{m_{A,*} - n - 1}{2} \right) \sum_{i=1}^n \psi \left(\frac{m_{A,*} - n - i}{2} \right) |\Psi_{A,d}|^{\frac{m_{A,d} - n - 1}{2}} + \\ &\quad k_{A,d}^{\frac{np}{2}} \Gamma_n \left(\frac{m_{A,*} - n - 1}{2} \right) |\Psi_{A,d}|^{\frac{m_{A,d} - n - 1}{2}}. \\ &\quad \left\{ \frac{m_* + p}{2} \log |\Psi_{A,d}| + \frac{m_{A,d} - n - 1}{2} \text{tr} \left[\Psi_{A,d}^{-1} \left(\Psi_* + \frac{k_* (M_* - Y_t)' \Sigma^{-1} (M_* - Y_t)}{(\alpha_t k_* + 1)^2} \right) \right] \right\} \quad (\text{A.77}) \\ B'_1 &= b'_1 b_2 b_3 + b_1 b'_2 b_3 + b_1 b_2 b'_3 \\ &= \frac{np}{2} k_* k_{A,*}^{\frac{np-2}{2}} \Gamma_n \left(\frac{m_{A,d} - n - 1}{2} \right) |\Psi_{A,*}|^{\frac{m_{A,*} - n - 1}{2}} + \\ &\quad k_{A,*}^{\frac{np}{2}} (m_* + p) \Gamma_n \left(\frac{m_{A,d} - n - 1}{2} \right) \sum_{i=1}^n \psi \left(\frac{m_{A,d} - n - i}{2} \right) |\Psi_{A,*}|^{\frac{m_{A,*} - n - 1}{2}} + \\ &\quad k_{A,*}^{\frac{np}{2}} \Gamma_n \left(\frac{m_{A,d} - n - 1}{2} \right) |\Psi_{A,*}|^{\frac{m_{A,*} - n - 1}{2}} \left[\left(\frac{m_* + p}{2} \right) \log |\Psi_{A,*}| + \frac{m_{A,*} - n - 1}{2\alpha_t} \right]. \quad (\text{A.78}) \end{aligned}$$

□

Proof of Corollary 4. Follows from some long but straightforward algebra from *iv)* of *Prop. 5* □

Proof of Remark 5. From the BF in Eq. 17, assuming $n = p = 1$, $\Sigma_P = \sigma^2/\varphi$, $\Sigma_L = \sigma^2$, $V = 1$, $M = m$ and $\mathbf{Y} = \mathbf{y} = (y_1, \dots, y_{t-1})$ one obtains $H_t(\alpha_t) > H_0$ for

$$\begin{aligned} \exp \left\{ \frac{(\alpha_t - 1)(\varphi + T)(y_t - m_*)^2}{2\sigma^2(\varphi + T + 1)(\alpha_t \varphi + \alpha_t T + 1)} \right\} &> H_0 \frac{\sqrt{\alpha_t(\varphi + T + 1)}}{\sqrt{\alpha_t(\varphi + T) + 1}} \iff \\ (y_t - m_*)^2 &\leq \frac{2\sigma^2(\varphi + T + 1)(\alpha_t \varphi + \alpha_t T + 1)}{(\alpha_t - 1)(\varphi + T)} \log \left\{ H_0 \frac{\sqrt{\alpha_t(\varphi + T + 1)}}{\sqrt{\alpha_t(\varphi + T) + 1}} \right\}, \quad (\text{A.79}) \end{aligned}$$

where $m_* = \varphi/(\varphi + T)m + T/(\varphi + T)\bar{y}$, $\bar{y} = (y_1 + \dots + y_T)/T$ and $T = t - 1$. The inequality is satisfied for $y_{1t} \leq y_t \leq y_{2t}$ there y_{jt} are the solutions of the second order equation associated with the above inequality. □

Proof of Proposition 7. i) Since $H_t(\alpha_t) < \kappa_t(\alpha_t)$ it is sufficient to find a , b , $\underline{\alpha}$ and $\bar{\alpha}$ such that $\int_0^1 \kappa_t(\alpha_t) \pi_t(\alpha_t) d\alpha_t < \infty$. By multinomial theorem, it follows

$$\begin{aligned} \frac{|\Sigma_L + \Sigma_*/\alpha_t|^{n/2}}{|\Sigma_L + \Sigma_*|^{n/2}} &= |\Sigma_L + \Sigma_*|^{-n/2} \left(\sum_{j=0}^p c_j \alpha_t^{-j} \right)^{n/2} = |\Sigma_L + \Sigma_*|^{-n/2} \left(\sum_{\underline{k} \in \mathcal{K}_n} d_{n,\underline{k}} \prod_{j=0}^p c_j^{k_j} \alpha_t^{-jk_j} \right)^{1/2} \\ &= |\Sigma_L + \Sigma_*|^{-n/2} \left(\sum_{\underline{k} \in \mathcal{K}} d_{n,\underline{k}} \alpha_t^{-\sum_{j=0}^p jk_j} c_{\underline{k}} \right)^{1/2} < c \alpha_t^{-pn/2} \quad (\text{A.80}) \end{aligned}$$

with c_j the j -th coefficient of the characteristic polynomial of $\Sigma_L + \Sigma_*$, $\underline{k} = (k_0, \dots, k_p)$ a multi-index with values in $\mathcal{K}_n = \{\underline{k} | k_0 + \dots + k_p = n\}$ and

$$c = |\Sigma_L + \Sigma_*|^{-n/2} \left(\sum_{\underline{k} \in \mathcal{K}_n} d_{\underline{k}} c_{\underline{k}} \right)^{1/2}, \quad d_{n, \underline{k}} = \binom{n}{k_0, k_1, k_2, \dots, k_p}, \quad c_{\underline{k}} = \prod_{j=0}^p c_j^{k_j}.$$

We used the relationship $\sum_{j=1}^p j k_j = \sum_{\ell=1}^p \sum_{j=\ell}^p k_j < \sum_{\ell=1}^p \sum_{j=0}^p k_j = pn$, along with the condition $\alpha_t < 1$, to derive the inequality here above. Thus we conclude

$$\int_0^1 \kappa_t(\alpha) \pi_t(\alpha) d\alpha_t < c \int_0^1 \frac{1}{B(a, b)} \alpha_t^{-pn/2} \alpha_t^{a-1} (1 - \alpha_t)^{b-1} d\alpha_t < \infty \quad (\text{A.81})$$

for $a - pn/2 - 1 > -1$, that is $a > pn/2$, and $b > 0$. Under the integrability conditions given above, the integral of the upper bound is

$$\begin{aligned} & \int_0^1 \frac{|\Sigma_L + \Sigma_*/\alpha_t|^{n/2}}{|\Sigma_L + \Sigma_*|^{n/2}} \frac{1}{B(a, b)} \alpha_t^{a-1} (1 - \alpha_t)^{b-1} d\alpha_t = \\ & = \int_0^1 \alpha^{-np/2} \left(1 + \sum_{j=0}^p \tilde{c}_j \alpha_t^j - 1 \right)^{n/2} \frac{1}{B(a, b)} \alpha_t^{a-1} (1 - \alpha_t)^{b-1} d\alpha_t \\ & = \sum_{\ell=0}^{\infty} \frac{\Gamma(n/2 + 1)}{\Gamma(\ell + 1) \Gamma(n/2 - \ell + 1)} \int_0^1 \alpha_t^{-np/2} \left(\sum_{j=0}^p \tilde{c}_j \alpha_t^j + 1 \right)^{\ell} \frac{1}{B(a, b)} \alpha_t^{a-1} (1 - \alpha_t)^{b-1} d\alpha_t \quad (\text{A.82}) \end{aligned}$$

with \tilde{c}_j the j -th coefficient of the characteristic polynomial of $\Sigma_L^{-1} \Sigma_*$, $\bar{c}_0 = \tilde{c}_0 - 1$, $\bar{c}_j = \tilde{c}_j$, $j = 1, \dots, p$, and where we used $0 < |\alpha_t I_p + A| / |I_p + A| < |\alpha_t I_p + A| / (|I_p \alpha| + |I_p(1 - \alpha_t) + A|) < 1$ with $A = \Sigma_L^{-1} \Sigma_*$, and the generalized binomial formula. The above expression becomes

$$\begin{aligned} & \sum_{\ell=0}^{\infty} \sum_{\underline{k} \in \mathcal{K}_{\ell}} \frac{\Gamma(n/2 + 1)}{\Gamma(\ell + 1) \Gamma(n/2 - \ell + 1)} d_{\ell, \underline{k}} \bar{c}_{\underline{k}} \int_0^1 \frac{1}{B(a, b)} \alpha_t^{a-np/2+w_{\underline{k}}-1} (1 - \alpha_t)^{b-1} d\alpha \\ & = \sum_{\ell=0}^{\infty} \sum_{\underline{k} \in \mathcal{K}_{\ell+1}} d_{\ell, \underline{k}} \bar{c}_{\underline{k}} \frac{\Gamma(n/2 + 1) B(a - np/2 + w_{\underline{k}}, b)}{\Gamma(\ell + 1) \Gamma(n/2 - \ell + 1) B(a, b)} \quad (\text{A.83}) \end{aligned}$$

by multinomial theorem, with

$$\bar{c}_{\underline{k}} = \prod_{j=0}^p c_j^{k_j}.$$

ii) Since from Corollary 1 the normalizing constant is well defined for $\alpha_t > (p + 2n)/m_d$ it is natural to assume in the general beta distribution $\underline{\alpha} = (p + 2n)/m_d$. It follows from Prop. 6 the BF $H_t(\alpha_t) < \Psi(\alpha_t) \kappa_{1t}(\alpha_t) \kappa_{2t}(\alpha_t)$, with $\Psi(\alpha_t) < 1$, $\kappa_{1t}(\alpha_t) < (\underline{\alpha} k_*/(\underline{\alpha}(k_* + 1)))^{np/2}$. To show the integrability it is sufficient to prove that $\int \kappa_{2t}(\alpha) \pi_t(\alpha) d\alpha < \infty$. We need to show that

$$\frac{\Gamma_n((m_d - n - 1)/2)}{\Gamma_n((m_* - n - 1)/2)} \int_{\underline{\alpha}} \frac{\Gamma_n\left(\frac{m_{A,*} - n - 1}{2}\right)}{\Gamma_n\left(\frac{m_{A,d} - n - 1}{2}\right)} \pi(\alpha_t) d\alpha_t < \infty. \quad (\text{A.84})$$

Note that we can write:

$$\int_{\underline{\alpha}} \frac{\Gamma_n\left(\frac{m_{A,*} - n - 1}{2}\right)}{\Gamma_n\left(\frac{m_{A,d} - n - 1}{2}\right)} \pi(\alpha_t) d\alpha_t = \int_{\underline{\alpha}} \frac{\pi^{\frac{n(n-1)}{4}} \prod_{j=1}^n \Gamma\left(\frac{m_{A,*} - n - 1}{2} + \frac{1-j}{2}\right)}{\pi^{\frac{n(n-1)}{4}} \prod_{j=1}^n \Gamma\left(\frac{m_{A,d} - n - 1}{2} + \frac{1-j}{2}\right)} \pi(\alpha_t) d\alpha_t$$

$$= \int_{\underline{\alpha}}^1 \frac{\prod_{j=1}^n \Gamma\left(\frac{\alpha_t m_d - p - n - j}{2}\right)}{\prod_{j=1}^n \Gamma\left(\frac{\alpha_t m_d - n - j}{2}\right)} \pi(\alpha_t) d\alpha_t = \int_{\underline{\alpha}}^1 \prod_{j=1}^n \frac{B\left((\alpha_t m_d - n - j)/2, \frac{p}{2}\right)}{\Gamma\left(\frac{p}{2}\right)} \pi(\alpha_t) d\alpha_t, \quad (\text{A.85})$$

where the last line follows from the property: $\Gamma(c-d)\Gamma(d) = B(c,d)\Gamma(c)$. Given the upper bound for the beta function, $B(x,y) \leq \frac{1}{xy}$ (e.g., see From and Ratnasingam, 2022, Th. 1.2)), we can write

$$\begin{aligned} B\left(\frac{\alpha_t m_d - n - j}{2}, \frac{p}{2}\right) &= \int_0^1 t^{\frac{m_d(\alpha_t - \underline{\alpha}) + m_d \underline{\alpha} - n - j}{2} - 1} (1-t)^{\frac{p}{2} - 1} dt \\ &= \int_0^1 t^{\frac{m_d}{2}(\alpha_t - \underline{\alpha}) + \frac{m_d \underline{\alpha} - n - j}{2} - 1} (1-t)^{\frac{p}{2} - 1} dt \leq \frac{2}{p \left[\frac{m_d}{2}(\alpha_t - \underline{\alpha}) + \frac{m_d \underline{\alpha} - n - j}{2} \right]}. \end{aligned}$$

From the above inequality and using $j \leq n$ and by Hölder's inequality

$$\begin{aligned} &\int_{\underline{\alpha}}^1 \frac{\Gamma_n\left(\frac{m_{A,*} - n - 1}{2}\right)}{\Gamma_n\left(\frac{m_{A,d} - n - 1}{2}\right)} \pi(\alpha_t) d\alpha_t \leq \Gamma\left(\frac{p}{2}\right)^{-n} \int_{\underline{\alpha}}^1 \prod_{j=1}^n \frac{4}{p [m_d(\alpha_t - \underline{\alpha}) + (m_d \underline{\alpha} - n - j)]} \pi(\alpha_t) d\alpha_t \\ &\leq \Gamma\left(\frac{p}{2}\right)^{-n} \int_{\underline{\alpha}}^1 \prod_{j=1}^n \frac{4}{p [m_d(\alpha_t - \underline{\alpha}) + (m_d \underline{\alpha} - n - n)]} \pi(\alpha_t) d\alpha_t \\ &= \Gamma\left(\frac{p}{2}\right)^{-n} \left(\frac{4}{p}\right)^n \int_0^{1-\underline{\alpha}} \frac{x^{(a-1)} (1-x)^{(b-1)}}{B(1-\underline{\alpha}; a, b) (m_d x + p)^n} dx \\ &\leq \Gamma\left(\frac{p}{2}\right)^{-n} \left(\frac{4}{p}\right)^n B(1-\underline{\alpha}; a, b)^{-1} \left[\int_0^{1-\underline{\alpha}} x^{r(a-1)} (1-x)^{r(b-1)} dx \right]^{1/r} \left[\int_0^{1-\underline{\alpha}} (m_d x + p)^{-qn} dx \right]^{1/q} \\ &= \Gamma\left(\frac{p}{2}\right)^{-n} \left(\frac{4}{p}\right)^n \frac{B(1-\underline{\alpha}; ra - r + 1, qb - q + 1)^{1/r}}{B(1-\underline{\alpha}; a, b)} \left[\int_0^{1-\underline{\alpha}} (m_d x + p)^{-qn} dx \right]^{1/q} \\ &= \Gamma\left(\frac{p}{2}\right)^{-n} \left(\frac{4}{p}\right)^n \frac{B(1-\underline{\alpha}; ra - r + 1, qb - q + 1)}{B(1-\underline{\alpha}; a, b)} \left[\frac{(m_d(1-\underline{\alpha}) + p)^{-qn+1} - p^{-qn+1}}{m_d(-qn+1)} \right]^{1/q} < \infty. \end{aligned}$$

Note that, since r and q are such that $r^{-1} + q^{-1} = 1$, the upper bound in the last line be minimize in r after choosing $q = r/(1-r)$. \square

Proof of Corollary 2. From Remark 3 an upper bound of the BF is $A_t(\alpha_t) = (\alpha_t(\varphi + t - 1) + 1)/(\alpha_t(\varphi + t))^{1/2}$ and its integral is bounded since $\int_0^1 A_t(\alpha_t) d\alpha_t \leq \int_0^1 \alpha_t^{-1/2} d\alpha_t = 2$ even if $\alpha_t = 0$ is not a continuity point for A_t . By the change of variable $A_t(\alpha_t) = z$ we obtain

$$\begin{aligned} &\int_0^1 A_t(\alpha_t) d\alpha_t = \int_{-\infty}^1 \frac{-2(\varphi + t)z^2}{((\varphi + t)z^2 - (\varphi + t - 1))^2} dz = 1 - \int_{-\infty}^1 \frac{1}{(\varphi + t)z^2 - (\varphi + t - 1)} dz \\ &= 1 - \frac{1}{(\varphi + t - 1)} \int_{-\infty}^{\sqrt{\frac{\varphi+t-1}{\varphi+t-1}}} \sqrt{\frac{\varphi+t-1}{\varphi+t}} \frac{1}{w^2 - 1} dw \\ &= 1 + \frac{1}{2\sqrt{(\varphi+t-1)(\varphi+t)}} \log \left(\frac{\sqrt{(\varphi+t)} + \sqrt{(\varphi+t-1)}}{\sqrt{(\varphi+t)} - \sqrt{(\varphi+t-1)}} \right), \end{aligned}$$

where the second line follows from integration by parts and the last line from the change of variable $w = z(\varphi + t)^{1/2}/(\varphi + t - 1)^{1/2}$. \square

Proof of Proposition 8. Let us define $E_t = Y_t - M_*$, then following the expression of BF given in i) Prop. 5, the cdf of the BF is

$$\begin{aligned}
F_{H_t}(h|\alpha_t) &= \mathbb{P}(H_t < h|\alpha_t) = \mathbb{P}\left(\frac{|\Sigma_{A,d}|^{\frac{n}{2}}}{|\Sigma_d|^{\frac{n}{2}}} \exp\left\{-\frac{1}{2}\text{tr}\left[\Sigma_H^{-1}E_t V^{-1}E_t'\right]\right\} < h|\alpha_t\right) \\
&= \mathbb{P}\left(\exp\left\{-\frac{1}{2}\text{tr}\left[\Sigma_H^{-1}E_t V^{-1}E_t'\right]\right\} < \frac{h}{\kappa_t}|\alpha_t\right) \\
&= \mathbb{P}\left(\text{tr}\left[(\Sigma_H^{-1/2})E_t V^{-1/2}(V^{-1/2})'E_t'(\Sigma_H^{-1/2})'\right] > -2\log(h/\kappa_t)|\alpha_t\right), \tag{A.86}
\end{aligned}$$

where $-\log(h/\kappa_t) > 0$ since $h_t < \kappa_t$, and $A^{1/2}$ denotes the Cholesky decomposition of A and $A^{-1/2}$ denotes its inverse. From Th. 2.3.10 in Gupta and Nagar (1999), the random matrix $W = (\Sigma_H^{-1/2})E_t V^{-1/2}$ follows the matrix normal $\mathcal{N}_{p,n}(M, \Sigma, I_n)$ with $M = (\Sigma_H^{-1/2})(\tilde{M} - M_*)V^{-1/2}$ and $\Sigma = (\Sigma_H^{-1/2})\tilde{\Sigma}\Sigma_H^{-1/2}$. The product WW' follows a non-central Wishart distribution $\mathcal{W}_{p,n}(\Sigma, \Sigma^{-1}MM')$ and following the results in Mathai (1980) and Pham-Gia et al. (2015) its trace $r = \text{tr}(WW')$ is the weighted sum of non-central chi-square variables $r = \lambda_1 r_1 + \dots + \lambda_p r_p$, where r_j follows a non-central chi-square $f_{n,U_{jj}}$ with non-centrality parameter U_{jj} , with U_{jj} being the element j -th diagonal element of $U = Q'\Sigma^{-1}MM'Q = Q'(\Sigma_H^{1/2}\tilde{\Sigma}^{-1}(\tilde{M} - M_*)V^{-1})(\tilde{M} - M_*)'\Sigma_H^{-1/2}Q$, and Q an orthogonal matrix such that $Q'\Sigma Q = \text{diag}(\lambda_1, \dots, \lambda_n)$. Then $F_{H_t}(h|\alpha_t) = 1 - F_r(-2\log(h/\kappa_t))$ and from Kourouklis and Moschopoulos (1985) the density of H_t is

$$f_{H_t}(h|\alpha_t) = \frac{2}{h} \sum_{k=0}^{\infty} c_k g(-2\log(h/\kappa_t); \frac{np}{2} + k, 2\lambda) \tag{A.87}$$

with support $0 < h < \kappa_t$, where $0 < \lambda < \infty$ is arbitrarily chosen, $g(x; a, b)$ is the density of a gamma distribution with shape and scale parameters $a > 0$ and $b > 0$, respectively, and the coefficients c_k , d_k and f_k satisfy

$$\begin{aligned}
c_k &= \exp\left(-\sum_{j=1}^p U_{jj}\right) \prod_{j=1}^p (\lambda_j/\lambda)^{-n/2} f_k, \quad d_k = \frac{n}{2k} \sum_{j=1}^p (1 - \lambda/\lambda_j)^k + \lambda \sum_{j=1}^p \frac{U_{jj}}{\lambda_j} (1 - \lambda/\lambda_j)^{k-1}, \\
f_{k+1} &= \frac{1}{k+1} \sum_{j=1}^{k+1} j d_j f_{k+1-j}, \quad k = 0, 1, 2, \dots \quad \text{with } f_0 = 1.
\end{aligned}$$

The cdf can be obtained by a change of variable argument as

$$\begin{aligned}
F_{H_t}(h|\alpha_t) &= \int_0^h f_{H_t}(u|\alpha_t) du = \sum_{k=0}^{\infty} c_k \int_{-2\log(h/\kappa_t)}^{\infty} g(u; \frac{np}{2} + k, 2\lambda) du \\
&= \sum_{k=0}^{\infty} c_k (1 - G(-2\log(h/\kappa_t); \frac{np}{2} + k, 2\lambda)), \tag{A.88}
\end{aligned}$$

where $G(x; a, b)$ is the cdf of the gamma density $g(x; a, b)$. □

Proof of Corollary 3. In the univariate case, following the notation in Example 3, we set $n = p = 1$, $\tilde{\Sigma} = \sigma_L^2$, $V = 1$ and $\tilde{M} = \theta$. From Prop. 8 one obtains $r = \lambda_1 r_1$ where $\lambda_1 = \sigma_L^2/\sigma_H^2$ and $U_{11} = (\theta - m_*)^2/\sigma_L^2$ and r_1 follows a non-central chi-square $f_{n,\gamma}(z)$ with $\gamma = U_{11}$. Then it follows $F_{H_t}(h|\alpha_t) = 1 - F_r(-2\log(h/\kappa_t)) = 1 - F_{n,U_{11}}(-2\log(h/\kappa_t) \sigma_H^2/\sigma_L^2)$ with density

$$f_{H_t}(h|\alpha_t) = 2 \frac{\sigma_H^2}{h\sigma_L^2} f_{n,U_{11}}(-2\log(h/\kappa_t) \sigma_H^2/\sigma_L^2). \tag{A.89}$$

From Th. 1.3 in András and Baricz (2008) the distribution of a noncentral chi-square with 1 degree of freedom can be written as $f_{n,\gamma}(z) = \exp(-(z + \gamma)/2)(2\pi z)^{-1/2} \cosh(\sqrt{\gamma z})$ for $n = 1$, and the cdf of H_t is

$$\begin{aligned}
F_{H_t}(h|\alpha_t) &= \int_0^h \frac{\sigma_H^2}{u\sigma_L^2} \exp(-(-2 \log(u/\kappa_t)\sigma_H^2/\sigma_L^2 + \gamma)/2) \frac{\cosh(\sqrt{-2\gamma \log(u/\kappa_t)\sigma_H^2/\sigma_L^2})}{\sqrt{-4\pi \log(u/\kappa_t)\sigma_H^2/\sigma_L^2}} du \\
&= \int_{\sqrt{-2 \log(h/\kappa_t)\sigma_H^2/\sigma_L^2}}^{+\infty} \exp(-(u^2 + \gamma)/2) \frac{1}{\sqrt{2\pi}} \exp(\sqrt{\gamma}u) du + \\
&\quad \int_{\sqrt{-2 \log(h/\kappa_t)\sigma_H^2/\sigma_L^2}}^{+\infty} \exp(-(u^2 + \gamma)/2) \frac{1}{\sqrt{2\pi}} \exp(-\sqrt{\gamma}u) du \\
&= 1 - \Phi(\sqrt{-2 \log(h/\kappa_t)\sigma_H^2/\sigma_L^2} - \sqrt{\gamma}) + \Phi(-\sqrt{-2 \log(h/\kappa_t)\sigma_H^2/\sigma_L^2} - \sqrt{\gamma}), \tag{A.90}
\end{aligned}$$

where $\gamma = (\theta - m_*)^2/\sigma^2$ and $\Phi(\cdot)$ is the cdf of a standard normal distribution. \square

B Further Empirical Results




Figure B.1: Macroeconomic dataset. Value of the BF $H_t(\alpha_t)$ (solid black), the upper bound $\kappa_t(\alpha_t)$ (solid red) at different dates. In all plots, the reference lines at 1 (---) and $10^{-1/2}$ (-.-).




Figure B.2: International trade dataset. Value of the BF $H_t(\alpha_t)$ (solid black), the upper bound $\kappa_t(\alpha_t)$ (solid red) at different dates. In all plots, the reference lines at 1 (---) and $10^{-1/2}$ (-.-).




Figure B.3: Realized volatility network dataset. Value of the BF $H_t(\alpha_t)$ (solid black), the upper bound $\kappa_t(\alpha_t)$ (solid red) at different dates. In all plots, the reference lines at 1 (---) and $10^{-1/2}$ (-.-.).

References

- Abadir, K. M. and Magnus, J. R. (2005). *Matrix Algebra*. Econometric Exercises. Cambridge University Press.
- Abraham, B. and Box, G. E. P. (1979). Bayesian Analysis of Some Outlier Problems in Time Series. *Biometrika*, 66(2):229.
- Alcalá, F. and Ciccone, A. (2004). Trade and Productivity. *The Quarterly Journal of Economics*, 119(2):613–646.
- Anderlucci, L. and Viroli, C. (2015). Covariance Pattern Mixture Models for the Analysis of Multivariate Heterogeneous Longitudinal Data. *Annals of Applied Statistics*, 9(2):777–800.
- András, S. and Baricz, A. (2008). Properties of the Probability Density Function of the Non-central Chi-squared Distribution. *Journal of Mathematical Analysis and Applications*, 346(2):395–402.
- Bayarri, M. and Berger, J. O. (1994). Robust Bayesian Bounds for Outlier Detection. *Recent Advances in Statistics and Probability (Vilaplana, JP and Puri, ML, eds.)*, pages 175–190.
- Bayarri, M. and Morales, J. (2003). Bayesian Measures of Surprise for Outlier Detection. *Journal of Statistical Planning and Inference*, 111(1):3–22.
- Bayarri, M. J. and Berger, J. O. (2004). The Interplay of Bayesian and Frequentist Analysis. *Statistical Science*, 19(1):58 – 80.
- Berger, J. O. and Yang, R.-Y. (1994). Noninformative Priors and Bayesian Testing for the AR(1) Model. *Econometric Theory*, 10(3-4):461–482.
- Bernardo, J. M. and Smith, A. F. (2009). *Bayesian Theory*, volume 405. John Wiley & Sons.
- Billio, M., Casarin, R., Costola, M., and Iacopini, M. (2021). A Matrix-Variate t Model for Networks. *Frontiers in Artificial Intelligence*, 4:1–49.
- Billio, M., Casarin, R., Costola, M., and Iacopini, M. (2022). Matrix-variate Smooth Transition Models for Temporal Networks. In Bekker, A., Ferreira, J. T., Arashi, M., and Chen, D.-G., editors, *Innovations in Multivariate Statistical Modeling: Navigating Theoretical and Multidisciplinary Domains*, pages 137–167. Springer, Cham.
- Billio, M., Casarin, R., Kaufmann, S., and Iacopini, M. (2023). Bayesian Dynamic Tensor Regression. *Journal of Business and Economic Statistics*, 41(2):429–439.
- Billio, M., Casarin, R., Ravazzolo, F., and Van Dijk, H. K. (2016). Interconnections between Eurozone and US booms and busts using a Bayesian panel Markov-switching VAR model. *Journal of Applied Econometrics*, 31(7):1352–1370.
- Box, G. E. P. and Tiao, G. C. (1968). A Bayesian Approach to Some Outlier Problems. *Biometrika*, 55(1):119.
- Brancaccio, G., Kalouptsi, M., and Papageorgiou, T. (2020). Geography, Transportation, and Endogenous Trade Costs. *Econometrica*, 88(2):657–691.
- Canova, F. and Ciccarelli, M. (2004). Forecasting and turning point predictions in a Bayesian panel VAR model. *Journal of Econometrics*, 120(2):327–359.

- Canova, F. and Ciccarelli, M. (2009). Estimating Multicountry VAR models. *International Economic Review*, 50(3):929–959.
- Carvalho, C. M., Massam, H., and West, M. (2007). Simulation of Hyper-inverse Wishart Distributions in graphical models. *Biometrika*, 94(3):647–659.
- Carvalho, C. M. and West, M. (2007). Dynamic Matrix-Variate Graphical Models. *Bayesian Analysis*, 2(1):69–97.
- Chen, E. Y., Tsay, R. S., and Chen, R. (2019). Constrained Factor Models for High-dimensional Matrix-variate Time Series. *Journal of the American Statistical Association*, 115(530):775–793.
- Chen, R., Xiao, H., and Yang, D. (2021). Autoregressive Models for Matrix-valued Time Series. *Journal of Econometrics*, 222(1):539–560.
- Chen, X., Irie, K., Banks, D., Haslinger, R., Thomas, J., and West, M. (2018). Scalable Bayesian Modeling, Monitoring, and Analysis of Dynamic Network Flow Data. *Journal of the American Statistical Association*, 113(522):519–533.
- De Santis, F. (2004). Statistical Evidence and Sample Size Determination for Bayesian Hypothesis testing. *Journal of Statistical Planning and Inference*, 124(1):121–144.
- De Santis, F. and Spezzaferri, F. (1997). Alternative Bayes Factors for Model Selection. *Canadian Journal of Statistics*, 25(4):503–515.
- Fang Chen, K. Y. and Wang, M. (2021). The minimum Bayes Factor Hypothesis Test for Correlations and Partial Correlations. *Communications in Statistics - Theory and Methods*, 50(11):2467–2480.
- Feldkircher, M., Huber, F., Koop, G., and Pfarrhofer, M. (2022). Approximate Bayesian Inference and Forecasting in Huge-dimensional Multicountry VARs. *International Economic Review*, 63(4):1625–1658.
- From, S. G. and Ratnasingam, S. (2022). Some New Inequalities for the Beta Function and Certain Ratios of Beta Functions. *Results in Applied Mathematics*, 15:100302.
- Gallaughan, M. P. and McNicholas, P. D. (2018). Finite Mixtures of Skewed Matrix Variate Distributions. *Pattern Recognition*, 80:83–93.
- Ghosh, J. K., Delampady, M., and Samanta, T. (2006). *An Introduction to Bayesian Analysis: Theory and Methods*, volume 725. Springer.
- Grubbs, F. E. (1950). Sample Criteria for Testing Outlying Observations. *The Annals of Mathematical Statistics*, 21(1):27 – 58.
- Guan, L. and Tibshirani, R. (2022). Prediction and Outlier Detection in Classification Problems. *Journal of the Royal Statistical Society, Series B*, 84(2):524–546.
- Guhaniyogi, R., Qamar, S., and Dunson, D. B. (2017). Bayesian Tensor Regression. *The Journal of Machine Learning Research*, 18(1):2733–2763.
- Gupta, A. K. and Nagar, D. K. (1999). *Matrix Variate Distributions*. Chapman & Hall, CRC Press.

- Guttman, I. (1973). Care and Handling of Univariate or Multivariate Outliers in Detecting Spuriousity—A Bayesian Approach. *Technometrics*, 15(4):723–738.
- Guttman, I., Dutter, R., and Freeman, P. R. (1978). Care and Handling of Univariate Outliers in the General Linear Model to Detect Spuriousity: A Bayesian Approach. *Technometrics*, 20(2):187.
- Held, L. and Ott, M. (2018). On p-Values and Bayes Factors. *Annual Review of Statistics and Its Application*, 5:393–419.
- Hoeting, J., Raftery, A. E., and Madigan, D. (1996). A Method for Simultaneous Variable Selection and Outlier Identification in Linear Regression. *Computational Statistics and Data Analysis*, 22(3):251–270.
- Kass, R. E. and Raftery, A. E. (1995). Bayes Factors. *Journal of the American Statistical Association*, 90(430):773–795.
- Kosheleva, O., Kreinovich, V., Trung, N. D., and Autcharyapanitkul, K. (2021). How to Make a Decision Based on the Minimum Bayes Factor (MBF): Explanation of the Jeffreys scale. In *Data Science for Financial Econometrics*, Studies in computational intelligence, pages 115–120. Springer, Cham.
- Kourouklis, S. and Moschopoulos, P. (1985). On the Distribution of the Trace of a Noncentral Wishart Matrix. *Metron*, 43(2):85–92.
- Lahiri, P. (2001). *Model selection, Volume 38*. IMS.
- Landim, F. and Gamerman, D. (2000). Dynamic hierarchical models: An extension to matrix-variate observations. *Computational Statistics & Data Analysis*, 35(1):11–42.
- Leng, C. and Tang, C. Y. (2012). Sparse matrix graphical models. *Journal of the American Statistical Association*, 107(499):1187–1200.
- Li, H., Sheng, X. S., and Yang, J. (2021). Monitoring Recessions: A Bayesian Sequential Quickest Detection Method. *International Journal of Forecasting*, 37(2):500–510.
- Ly, A., Verhagen, J., and Wagenmakers, E.-J. (2016). Harold Jeffreys’s Default Bayes Factor Hypothesis Tests: Explanation, Extension, and Application in Psychology. *Journal of Mathematical Psychology*, 72:19–32.
- Magnus, J. R. and Neudecker, H. (1999). *Matrix Differential Calculus with Applications in Statistics and Econometrics, 2nd Ed.* Wiley.
- Mathai, A. (1980). Moments of the Trace of a Noncentral Wishart Matrix. *Communications in Statistics, Theory and Methods*, 9(8):795–801.
- Page, G. L. and Dunson, D. B. (2011). Bayesian Local Contamination Models for Multivariate Outliers. *Technometrics*, 53(2):152–162.
- Pawel, S. and Held, L. (2022). The Sceptical Bayes Factor for the Assessment of Replication Success. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 84(3):879–911.
- Pawel, S. and Held, L. (2025). Closed-form Power and Sample Size Calculations for Bayes Factors. *The American Statistician*, pages 1–34.

- Penny, W. D., Stephan, K. E., Mechelli, A., and Friston, K. J. (2004). Comparing Dynamic Causal Models. *Neuroimage*, 22(3):1157–1172.
- Pettit, L. and Smith, A. (1983). Bayesian Model Comparison in the Presence of Outliers. *Bulletin of the International Statistical Institute*, 50:292–309.
- Pettit, L. and Smith, A. (1985). Outliers and Influential Observations in Linear Models. *Bayesian Statistics*, 2(1):473–94.
- Pettit, L. I. (1988). Bayes Methods for Outliers in Exponential Samples. *Journal of the Royal Statistical Society, Series B*, 50(3):371–380.
- Pettit, L. I. (1990). The Conditional Predictive Ordinate for the Normal Distribution. *Journal of the Royal Statistical Society, Series B*, 52(1):175–184.
- Pettit, L. I. (1992). Bayes Factors for Outlier Models using the Device of Imaginary Observations. *Journal of the American Statistical Association*, 87(418):541–545.
- Pham-Gia, T., Thanh, D. N., Phong, D. T., et al. (2015). Trace of the Wishart matrix and applications. *Open Journal of Statistics*, 5(03):173.
- Rasines, D. G. and Young, G. A. (2022). Empirical Bayes and Selective Inference. *J Indian Inst Sci*, 102(4):1205–1217.
- Riani, M., Atkinson, A. C., Corbellini, A., Farcomeni, A., and Laurini, F. (2024). Information Criteria for Outlier Detection Avoiding Arbitrary Significance Levels. *Econometrics and Statistics*, 29:189–205.
- Robert, C. P. (2007). *The Bayesian choice: from decision-theoretic foundations to computational implementation*, volume 2. Springer.
- Robert, C. P. (2014). On the Jeffreys-Lindley paradox. *Philosophy of Science*, 81(2):216–232.
- Robert, C. P. (2022). 50 shades of Bayesian Testing of Hypotheses. In Young, A. G., Srinivasa Rao, A. S., and Rao, C. R., editors, *Advancements in Bayesian Methods and Implementations*, chapter 5, pages 103–120. Academic Press.
- Robert, C. P., Chopin, N., and Rousseau, J. (2009). Harold Jeffreys’s Theory of Probability Revisited. *Statistical Science*, pages 141–172.
- Rose, A. (2004). Do We Really Know that the WTO Increases Trade. *American Economic Review*, 94(1):98–114.
- Rosner, B. (1983). Percentage Points for a Generalized ESD Many-outlier Procedure. *Technometrics*, 25(2):165–172.
- Schad, D. J., Nicenboim, B., Bürkner, P.-C., Betancourt, M., and Vasishth, S. (2023). Workflow Techniques for the Robust use of Bayes Factors. *Psychological Methods*, 28(6):1404–1426.
- Schrider, D. R., Shanku, A. G., and Kern, A. D. (2016). Effects of Linked Selective Sweeps on Demographic Inference and Model Selection. *Genetics*, 204(3):1207–1223.
- She, Y. and Owen, A. B. (2011). Outlier Detection Using Nonconvex Penalized Regression. *Journal of the American Statistical Association*, 106(494):626–639.

- Stefan, A. M., Gronau, Q. F., Schönbrodt, F. D., and Wagenmakers, E.-J. (2019). A Tutorial on Bayes Factor Design Analysis using an Informed Prior. *Behavior Research Methods*, 51(3):1042–1058.
- Thompson, G. Z., Maitra, R., Meeker, W. Q., and Bastawros, A. F. (2020). Classification with the Matrix-variate-t Distribution. *Journal of Computational and Graphical Statistics*, 29(3):668–674.
- Tomarchio, S. D., Gallagher, M. P., Punzo, A., and McNicholas, P. D. (2022). Mixtures of Matrix-variate Contaminated Normal Distributions. *Journal of Computational and Graphical Statistics*, 31(2):413–421.
- Triantafyllopoulos, K. (2008). Missing Observation Analysis for Matrix-variate Time Series Data. *Statistics & Probability Letters*, 78(16):2647–2653.
- Tsukurimichi, T., Inatsu, Y., Duy, V. N. L., and Takeuchi, I. (2022). Conditional Selective Inference for Robust Regression and Outlier Detection using Piecewise-Linear Homotopy Continuation. *Annals of the Institute of Statistical Mathematics*, 74(6):1197–1228.
- van der Merwe, A. J., Groenewald, P. C. N., Sjölander, M. R., and Meyer, J. H. (2021). A Bayesian-Frequentists Approach for Detecting Outliers in a One-way Variance Components Model. *Communications in Statistics-Theory and Methods*, 50(23):5652–5677.
- Verdinelli, I. and Wasserman, L. (1991). Bayesian Analysis of Outlier Problems using the Gibbs Sampler. *Statistics and Computing*, 1(2):105–117.
- Viroli, C. (2011). Finite Mixtures of Matrix Normal Distributions for Classifying Three-way Data. *Statistics and Computing*, 21(4):511–522.
- Viroli, C. and Anderlucci, L. (2013). Modelling Longitudinal Data through Matrix-variate Normal Mixtures. In *Advances in Latent Variables - Methods, Models and Applications*.
- Wagenmakers, E.-J. and Ly, A. (2023). History and Nature of the Jeffreys–Lindley Paradox. *Archive for History of Exact Sciences*, 77(1):25–72.
- Wang, H. and West, M. (2009). Bayesian analysis of matrix normal graphical models. *Biometrika*, 96(4):821–834.
- Weiss, R. (1997). Bayesian Sample Size Calculations for Hypothesis Testing. *Journal of the Royal Statistical Society: Series D (The Statistician)*, 46(2):185–191.
- West, M. (1986). Bayesian Model Monitoring. *Journal of the Royal Statistical Society: Series B (Methodological)*, 48(1):70–78.
- Yekutieli, D. (2012). Adjusted Bayesian Inference for Selected Parameters. *Journal of the Royal Statistical Society, Series B*, 74(3):515–541.
- Yu, L., He, Y., Kong, X., and Zhang, X. (2022). Projected Estimation for Large-dimensional Matrix Factor Models. *Journal of Econometrics*, 229(1):201–217.
- Zellner, A. (1986). On Assessing Prior Distributions and Bayesian Regression Analysis with g-prior Distributions. *Bayesian inference and decision techniques*.
- Zhang, F. (2011). *Matrix Theory: Basic Results and Techniques*. Springer.

- Zhao, Q., Small, D. S., and Ertefaie, A. (2022). Selective Inference for Effect Modification via the Lasso. *Journal of the Royal Statistical Society, Series B*, 84(2):382–413.
- Zhu, X., Pan, R., Li, G., Liu, Y., and Wang, H. (2017). Network Vector Autoregression. *The Annals of Statistics*, 45(3):1096–1123.