

NON-FACTORIZABLE RIBBON HOPF ALGEBRAS

Quentin Faes and Maksymilian Manko

March 2025

Abstract

Building on the work of Nenciu we provide examples of non-factorizable ribbon Hopf algebras, and introduce a stronger notion of non-factorizability. These algebras are designed to provide invariants of 4-dimensional 2-handlebodies up to 2-deformations. We prove that some of the invariants derived from these examples are invariants dependent only on the boundary or on the presentation of the fundamental group of the 2-handlebody.

Contents

1	Introduction	1
1.1	Main results	2
1.2	Structure of the paper	3
2	Preliminaries	3
2.1	Notations for Hopf algebras	3
2.2	Integrals and unimodularity	4
2.3	Quasitriangular and ribbon structures	5
2.4	Algebra generators and tuple notation	6
3	The Nenciu construction	7
3.1	Hopf algebra structure	7
3.2	Examples	8
3.3	Unimodular structure	11
3.4	Quasitriangular structure	11
3.5	Ribbon structure	14
3.6	Examples revisited	14
4	Non-factorizable biproducts with $u_q\mathfrak{sl}_2$	17
4.1	The small quantum group $u_q\mathfrak{sl}_2$	17
4.2	The semi-direct biproduct structure	18
4.3	Non-factorizable biproducts with $u_q\mathfrak{sl}_2$	19
4.4	Examples	21
5	Topological applications	25
5.1	Handlebodies	25
5.2	Invariants of handlebodies up to 2-equivalence	25
5.3	Cases of degeneration of the invariant	26
5.4	Degeneration of the invariant associated to the Nenciu algebras	27
A	Proofs	29
A.1	Proof of Theorem 4.8	29
A.2	Proof of Theorem 4.13	30
A.3	Proofs of Propositions 4.21 and 4.25	33
B	Summary of examples	34
B.1	Nenciu type examples	34
B.2	Non-factorizable extensions of $u_q\mathfrak{sl}_2$ examples	37

1 Introduction

Unimodular, ribbon categories introduced by Turaev in [Tur94] constitute an important topic in quantum algebra, chiefly for their many applications in quantum topology. Since the 1990s, multiple constructions of 3-manifold and 4-manifold invariants have been constructed using these essential ingredients [Tur94], [KL01], [CY93]. Ribbon Hopf algebras are a natural source of such categories through their representation theory. A ribbon Hopf algebra H is a quasitriangular Hopf algebra (i.e. it comes with the extra data of an invertible element R of $H \otimes H$ called the R-matrix) together with a central element of H called the ribbon element. Both R and v are required to satisfy some equations, see Section 2.

In particular, one may define a 3-manifold invariant from any so-called *factorizable* unimodular ribbon Hopf algebra. Factorizability is a non-degeneracy condition related to the *monodromy matrix* $M := R_{21}R$ of the algebra, where R_{21} is the R-matrix with permuted entries (see Prop. 2.10). Recall also that a Hopf algebra H is unimodular if it admits an integral $\lambda \in H^*$ and a cointegral $\Lambda \in H$ satisfying some properties, as detailed in Def. 2.5.

There is, similarly, a strong interest in *triangular* Hopf algebras. By the work of Bobtcheva [Bob23] these provide invariants of 2-dimensional CW-complexes, and are closely related to a well-known problem in combinatorial group theory, the *Andrews-Curtis conjecture* [AC65]. Finite-dimensional triangular Hopf algebras over \mathbb{C} were fully classified in [AEG02].

More recently there has been a considerable interest in the intermediate case, the unimodular, *non-factorizable* ribbon Hopf algebras, as ingredients of non-semisimple 4-manifold invariants. In [BdR23] Beliakova and De Renzi introduced Kerler-Lyubashenko-type invariants of 4-dimensional 2-handlebodies up to 2-deformations [BdR23] based on the work of Bobtcheva and Piergallini [BP06], and presented in a more streamlined way in [BBDRP23]. These invariants are the main motivation for this paper. They are meant to provide an angle on a conjecture of Gompf, see Conjecture 5.4, that asserts any diffeomorphic 4-dimensional 2-handlebodies are related by a sequence of Kirby moves involving handles of index at most 2, and is expected to be false. A TQFT construction producing invariants of 4-manifolds up to diffeomorphism introduced in [CGHPM23] similarly relies on representation categories of non-semisimple, unimodular, non-factorizable ribbon Hopf algebras.

At the topological level, non-factorizability is directly related to the ability of the invariants from [BdR23] to differentiate between 1-handles and 2-handles, which in turn is related to the ability of the invariant to see beyond the boundary of the 4-dimensional 2-handlebody. Non (co)semi-simplicity is related to the sensitivity of the invariant to a stabilization operation on handlebodies. This is detailed in Section 5.

Several examples of non-factorizable unimodular ribbon Hopf algebras are known and include the small quantum group $u_q\mathfrak{sl}_2$ at even root of unity q . It was shown in [BDR22] that the scalar Kerler-Lyubashenko type invariant of [BdR23] applied to a connected 4-dimensional 2-handlebody only sees its 3-dimensional boundary and homological information. Another example is the Hopf algebra of symplectic fermions SF_{2n} , which admits a continuous family of non-factorizable ribbon structures, including a triangular one. The generic case, for $n = 1$, was shown by Kerler in [Ker03] to give essentially invariants of the 3-dimensional boundary with this approach. In both cases, the Hopf algebras are, in some sense, “close to factorizable”. Indeed, in the unimodular case, factorizability is equivalent to requiring that applying the integral to one side of the monodromy matrix gives the cointegral. In these non-factorizable cases we mentioned, this equation will still be verified up to an invertible element of the algebra.

1.1 Main results

Hence, in an attempt to avoid such “degeneration to the boundary”, in this paper, we introduce the stricter notion of *strong non-factorizability*. We demand that the result of applying the integral to either side of the monodromy is zero, and hence cannot be related to the cointegral by any invertible element of the algebra:

$$(\lambda \otimes \text{Id})M = 0.$$

Non-semisimple triangular Hopf algebras provide examples of this property, but yield handlebody invariants only related to the presentation of the fundamental group. Here, we present two constructions capable of producing ribbon Hopf algebras which admit only strongly non-factorizable quasitriangular structures, and are generically not triangular.

First we discuss a family of Hopf algebras that was introduced in [BDG00] and analysed from the angle of quasitriangularity and ribbonness in [Nen04]. We will call these Hopf algebras *Nenciu algebras*. Let $\mathbf{m} \in \mathbb{Z}^s$ be a row tuple of integers of length $s \in \mathbb{Z}$, $t \in \mathbb{Z}$ and \mathbf{d}, \mathbf{u} be $t \times s$ matrices with integer entries. A Nenciu algebra is generated by grouplike generators K_a , $a = 1, \dots, s$ such that $K_a^{m_a} = \mathbf{1}$, commuting with each other, as well as skew-primitive nilpotent generators X_k , $k = 1, \dots, t$ such that $X_k^2 = 0$. The relations between the generators are prescribed by the matrices \mathbf{d}, \mathbf{u} and are *diagonal*, that is commuting any two elements results only in a constant in front of the swapped product. We denote the resulting algebra by $H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$, and a full definition is provided in Definition 3.1. In [Nen04], an array of previously existing quasitriangular Hopf algebra constructions, including the family SF_{2n} , were unified, and sufficient conditions for such an algebra to be ribbon were determined, but no new ones were given. In particular, the questions of unimodularity and factorizability were not addressed.

In this paper, we exhibit new examples. Moreover we determine sufficient conditions for unimodularity, and for a number of instances we establish when non-factorizable and strongly non-factorizable ribbon structures are admitted, relying in particular on a theorem of Radford [Rad94], reproduced below as Theorem 2.9. We also study another property: the algebra is called *anomalous* if $\lambda(v) \neq 1$ and anomaly-free otherwise (see Definition 2.20). Topologically, as explained in [BdR23], when the algebra is both factorizable and anomaly-free, the 4-dimensional 2-handlebody invariant associated to the algebra is actually an invariant of the boundary of the handlebody. For completeness, we check that anomaly-freeness is not satisfied in the examples where the corresponding quasitriangular structure is strongly non-factorizable. Combining the results of [Nen04] with our own, we have the following.

Main Theorem A. *Under an appropriate choice of parameters a Nenciu type Hopf algebra $H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$ can be simultaneously*

1. *quasitriangular* (Theorem 2.9 and [Nen04, Theorem 3.4]),
2. *ribbon* (Theorem 3.21 and [Nen04, Proposition 4.9]),
3. *unimodular* (Proposition 3.15),
4. *strongly non-factorizable* (Proposition 3.28),
5. *and anomalous* (Corollary 3.29).

However, we show in Theorem 5.8 that the related Kerler-Lyubashenko invariants of 4-dimensional 2-handlebodies degenerate to invariants of the 2-dimensional *spine*, the 2-dimensional CW-complex resulting from collapsing the cocores of the handles attached or equivalently to invariants of the presentation of the fundamental group of the handlebody (see the discussion above the aforementioned proposition).

In an attempt to remedy this, and inspired by the work of Majid [Maj00], we present a second family of non-factorizable Hopf algebras where we form a sort of semi-direct product of $u_q\mathfrak{sl}_2$ where q is a root of unity of order divisible by 4, and a Nenciu-type Hopf algebra $H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$, where the former (co)acts on the latter on the right. While on the Hopf algebraic level, the construction utilises standard techniques, the fine-tuning required to retrieve the desired properties does not seem to have appeared in the literature before. The action and coaction involve only the K generator of $u_q\mathfrak{sl}_2$ and are designed so that the result is unimodular and carries a ribbon structure being a slightly modified product of the R-matrices and ribbon elements of the two respective factors. It turns out this Hopf algebra has different properties than a trivial tensor product Hopf algebra $u_q\mathfrak{sl}_2 \otimes H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$, or standard smash and cosmash products affecting only the algebra or coalgebra (and the antipode), respectively, presented in [Mol77]. Thus, we dub it the *semidirect biproduct* as both algebra and coalgebra structures interact, and denote $u_q\mathfrak{sl}_2 \ltimes H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$ to emphasize that it is the $u_q\mathfrak{sl}_2$ factor that (co)acts, see Definition 4.4. Its key property is that if the Nenciu factor carries a strongly non-factorizable, anomalous ribbon structure, so does the semidirect biproduct with $u_q\mathfrak{sl}_2$. We summarise our results as follows.

Main Theorem B. *Under an appropriate choice of parameters a semidirect biproduct $u_q\mathfrak{sl}_2 \ltimes H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$ can be simultaneously*

1. *quasitriangular (Theorem 4.13)*
2. *ribbon (Theorem 4.13),*
3. *unimodular (Theorem 4.13),*
4. *strongly non-factorizable (Proposition 4.22),*
5. *and anomalous (Corollary 3.29).*

However, we again observed that no true 4-dimensional information seems to be carried by the 4-dimensional 2-handlebody invariants given by this family of Hopf algebras. More precisely, computations for multiple examples suggest that the invariant might be given by the product of the 2-dimensional invariant of the spine of the handlebody obtained from $H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$ and 3-dimensional invariant of its boundary associated to $u_q\mathfrak{sl}_2$. Thus, we leave it as Conjecture 5.11.

1.2 Structure of the paper

The paper is organised as follows: in Section 2 we introduce all the relevant definitions pertaining to unimodular ribbon Hopf algebras and the notation used to describe the Nenciu algebras. In Section 3 the Nenciu algebras are introduced together with the conditions for a given one to admit unimodular, ribbon, and strongly non-factorizable structures. In Section 4, we introduce the second family of examples and once again give conditions for strong non-factorizability. In Section 5 we recall some topological background, including the Kerler-Lyubashenko invariants of 4-dimensional 2-handlebodies, and discuss the efficacy of the invariants provided by our examples. Finally in Appendix A we list all the examples of Hopf algebras found throughout the paper, and in Appendix B we collect some proofs that are too long to be presented in the main body.

Acknowledgements: The authors would like to thank Ivelina Bobtcheva, Marco de Renzi, Azat Gainutdinov and Cris Negron for early comments on the paper and fruitful discussions, as well as Riccardo Piervallini for making his Mathematica computer program available, and Anna Beliakova for her help and introducing them to the topic. Both authors were supported by Simons Collaboration on New Structures in Low-Dimensional Topology and Grant 200020_207374 of the Swiss National Science Foundation.

2 Preliminaries

Let us recall some definitions and results regarding Hopf algebras (in the category of complex vector spaces), and fix the notation used in the rest of the paper.

2.1 Notations for Hopf algebras

In this paper, we consider Hopf algebras over the field of complex numbers \mathbb{C} .

Definition 2.1. A Hopf algebra $(H, \mu, \eta, \Delta, \epsilon, S)$ is the data of a vector space (over \mathbb{C}) H together with

- a *unit* $\mathbf{1} : \mathbb{C} \rightarrow H$ and a *product* $\mu : H \otimes H \rightarrow H$,
- a *counit* $\epsilon : H \rightarrow \mathbb{C}$, and a *coproduct* $\Delta : H \rightarrow H \otimes H$
- an invertible *antipode* $S : H \rightarrow H$,

that are \mathbb{C} -linear maps satisfying the axioms

$$\begin{aligned}
\mu \circ (\mu \otimes \text{id}) &= \mu \circ (\text{id} \otimes \mu), \\
\mu \circ (\eta \otimes \text{id}) &= \text{id} = \mu \circ (\text{id} \otimes \eta), \\
(\Delta \otimes \text{id}) \circ \Delta &= (\text{id} \otimes \Delta) \circ \Delta, \\
(\varepsilon \otimes \text{id}) \circ \Delta &= \text{id} = (\text{id} \otimes \varepsilon) \circ \Delta, \\
(\mu \otimes \mu) \circ (\text{id} \otimes \tau \otimes \text{id}) \circ (\Delta \otimes \Delta) &= \Delta \circ \mu, \\
\varepsilon \circ \mu &= \varepsilon \otimes \varepsilon, \\
\Delta \circ \eta &= \eta \otimes \eta, \\
\varepsilon \circ \eta &= 1, \\
\mu \circ (S \otimes \text{id}) \circ \Delta &= \eta \circ \varepsilon = \mu \circ (\text{id} \otimes S) \circ \Delta,
\end{aligned}$$

where $\tau : H \otimes H \rightarrow H \otimes H$ is the *flip map* exchanging the factors in the tensor product.

We will often abusively denote the Hopf algebra by H , and write gh instead of $\mu(g, h)$ for $g, h \in H$. We also note that $H \otimes H$ can be endowed with an algebra structure, using the product

$$(a \otimes b)(c \otimes d) := ac \otimes bd.$$

where $a, b, c, d \in H$.

Remark 2.2. Hopf algebras may be defined in any braided monoidal category \mathcal{C} . The algebra structure of $H \otimes H$ is then defined using the braiding.

Notation. We use the Sweedler notation for the coproduct: if

$$\Delta(h) = \sum_{i=1}^n h_{(1)i} \otimes h_{(2)i}$$

then we write

$$\Delta(h) = h_{(1)} \otimes h_{(2)},$$

for some $n \in \mathbb{Z}_{>0}$. By $\Delta^{cop} := \tau \circ \Delta$ we denote the *coopposite coproduct*, with $\tau : H \otimes H \rightarrow H \otimes H$ the flip map. Recall that if H is a Hopf algebra, then $H^{cop} := (H, \mu, \eta, \Delta^{cop}, \varepsilon, S)$ is also a Hopf algebra.

Recall that if H is a Hopf algebra, the dual vector space

$$H^* := \text{Hom}_{\mathbb{C}}(H, \mathbb{C}).$$

can also be provided with a Hopf algebra structure.

Definition 2.3. An element $g \in H$ is called *grouplike* if $\Delta(g) = g \otimes g$. The subset $G(H)$ of all such elements carries a group structure, and generates a Hopf subalgebra which is isomorphic to the group algebra of $G(H)$. An element $X \in H$ is called *skew-primitive with respect to* $L \in G(H)$ if $\Delta(X) = \mathbf{1} \otimes X + X \otimes L$, and *primitive* if $L = \mathbf{1}$.

Definition 2.4. The *left adjoint action* of H is the left H -module structure on H given by

$$\triangleright : H \otimes H \rightarrow H, \quad a \triangleright b := a_{(1)} b S(a_{(2)})$$

in the Sweedler notation.

2.2 Integrals and unimodularity

Let H be a finite dimensional Hopf algebra over an algebraically closed field. Then it admits a left (resp. right) integral $\lambda_L \in H^*$ (resp. λ_R) and cointegral $\Lambda_L \in H$ (resp. Λ_R), see [Rad12, Chapter 10.1].

Definition 2.5. An element $\lambda_L \in H^*$ is a *left* (resp. $\lambda_R \in H^*$ a *right*) *integral* of H if for all $h \in H$,

$$h_{(1)} \otimes \lambda_L(h_{(2)}) = \lambda_L(h) \quad (\text{resp.} \quad \lambda_R(h_{(1)}) \otimes h_{(1)} = \lambda_R(h)).$$

An element $\Lambda_L \in H$ is a *left* (resp. $\Lambda_R \in H$ a *right*) *cointegral* if for all $h \in H$,

$$h \Lambda_L = \varepsilon(h) \Lambda_L \quad (\text{resp.} \quad \Lambda_R h = \Lambda_R \varepsilon(h)).$$

An integral (resp. cointegral) is called *two-sided* if its both a left and right integral (resp. cointegral), or equivalently, if

$$\lambda \circ S = \lambda \quad (\text{resp.} \quad S(\Lambda) = \Lambda).$$

It is also known that Λ_L, Λ_R generate 1-dimensional subalgebras of H and λ_L, λ_R generate 1-dimensional subalgebras of H^* , thus different choices differ up to a constant in the ground field.

Definition 2.6 ([Rad12] Definition 10.2.3). We will call H *unimodular* if Λ is two-sided.

2.3 Quasitriangular and ribbon structures

Let us recall the definition of the quasitriangular Hopf algebra. We need the following notation.

Notation. For an element $R \in H \otimes H$ such as the R -matrix of a quasitriangular Hopf algebra of Definition 2.7 below, we write

$$R = R' \otimes R'',$$

where the summation is again implicit. Sometimes, we will need many copies of R , in which case they will be labeled with numbers. We also denote $R_{21} := \tau(R)$.

Definition 2.7. Let H be a Hopf algebra and R be an element of $H \otimes H$. The pair (H, R) is quasitriangular if the following axioms are satisfied.

$$(QT1) \quad \Delta(R') \otimes R'' = R'_1 \otimes R'_2 \otimes R'_1 R''_2$$

$$(QT2) \quad \epsilon(R')R'' = \mathbf{1}$$

$$(QT3) \quad R' \otimes \Delta^{cop}(R'') = R'_1 R'_2 \otimes R''_1 \otimes R''_2$$

$$(QT4) \quad \epsilon(R'')R' = \mathbf{1}$$

$$(QT5) \quad \Delta^{cop}(h)R = R\Delta(h), \forall h \in H$$

Then R is called a *universal R -matrix* (where we usually drop the “universal”), and it follows from (QT2) and (QT4) that it is invertible in the algebra $H \otimes H$, with the inverse denoted R^{-1} .

Definition 2.8. The *monodromy matrix* is defined as

$$M := R_{21}R = R''_1 R'_2 \otimes R'_1 R''_2.$$

If $M = \mathbf{1} \otimes \mathbf{1}$, the quasitriangular structure is called *triangular*.

In order to exhibit quasitriangular structures on Hopf algebras, we will strongly rely on the following result.

Theorem 2.9 ([Rad94], Section 2.1). *Let H be a Hopf algebra. Any element $R \in H \otimes H$ satisfying (QT1)-(QT4) induces a Hopf algebra map*

$$f_R : H^* \rightarrow H^{cop} \\ h^* \mapsto h^*(R')R'',$$

where H^{cop} is the coopposite Hopf algebra of H , defined above.

By the contrapositive of Theorem 2.9, if there are no bialgebra maps between H^* and H^{cop} , then there can be no quasitriangular structures on H .

Proposition 2.10. [[Shi16] Theorem 1.1, [EGNO16], Exercise 8.6.4] For a finite-dimensional quasitriangular Hopf algebra (H, R) the following two conditions are equivalent

1. the monodromy matrix M induces a non-degenerate pairing on $H^* \otimes H^* \rightarrow \mathbb{C}$ given by $h_1^* \otimes h_2^* \mapsto h_1^*(M')h_2^*(M'')$, or
2. the Drinfeld-Reshetikhin map

$$H^* \rightarrow H \\ h^* \mapsto h^*(M')M''$$

is an isomorphism of vector spaces.

If either item is fulfilled (H, R) is called *factorizable*, and otherwise *non-factorizable*.

For the purpose of this discussion we introduce a stronger notion.

Definition 2.11. A unimodular, quasitriangular Hopf algebra (H, R) with a two-sided integral $\lambda \in H^*$ is called *strongly non-factorizable* if any of the two equivalent conditions is fulfilled

$$\lambda(M')M'' = 0,$$

or

$$M'\lambda(M'') = 0.$$

It is easy to see from Definition 2.10 that any strongly non-factorizable Hopf algebra is non-factorizable.

Remark 2.12. Note that if (H, R) is non-cosemisimple, triangular, and has a two-sided integral λ , then it is strongly non-factorizable. Indeed $\lambda(\mathbf{1}) = 0$ is equivalent to the non-cosemisimplicity condition for a unimodular Hopf algebra (see [Rad12], Theorem 10.3.2).

We now turn to the ribbon structure.

Definition 2.13. Let (H, R) be a finite dimensional, quasitriangular Hopf algebra with monodromy matrix M . A *ribbon element* $v \in Z(H)$ is a central element of H such that

- (R1) $S(v) = v$
(R2) $\epsilon(v) = 1$
(R3) $M\Delta(v) = v \otimes v.$

The triple (H, R, v) is called a *ribbon Hopf algebra*.

Definition 2.14. The *Drinfeld element* of a quasitriangular Hopf algebra (H, R) is

$$u := S(R'')R'.$$

It is invertible with inverse given by

$$u^{-1} = S^{-1}((R^{-1})'')(R^{-1})'.$$

Proposition 2.15 ([Kas12], VIII.4.1). Let (H, R) be a quasitriangular Hopf algebra and u be the corresponding Drinfeld element. For any $h \in H$

$$S^2(h) = uhu^{-1}.$$

Definition 2.16 ([Rad94]). Let H be a Hopf algebra. A *pivotal element* is a grouplike element $g \in G(H)$ such that for any $h \in H$.

$$S^2(h) = ghg^{-1}.$$

As explained in [Rad94, Prop. 2] in a ribbon Hopf algebra, a ribbon element $v = g^{-1}u$ is the product of the Drinfeld element u and the inverse of a pivotal element. However, not every pivotal element can be used to define a ribbon element.

Definition 2.17. Let H be Hopf algebra admitting a left integral $\lambda_L \in H^*$. The *distinguished grouplike element* $a \in G(H)$ is the unique element such that $\lambda_L \circ S(h) = \lambda_L(ah)$.

Lemma 2.18 ([Rad94] Proposition 2.(b)). Let (H, R) be a finite-dimensional quasitriangular Hopf algebra with a two-sided cointegral and denote the Drinfeld element by u . Let $g \in G(H)$ be a pivotal element and a be the distinguished grouplike element. Then $v := g^{-1}u$ is a ribbon element if and only if

$$g^2 = a^{-1}.$$

Lemma 2.19. Let (H, R) be a finite-dimensional quasitriangular Hopf algebra with a two-sided integral λ . If $g \in G(H)$ is a pivotal element, then $g^2 = \mathbf{1}$.

Proof. If λ is two-sided, then the distinguished grouplike element a of Lemma 2.18 satisfies $a = \mathbf{1}$. Thus, $g^2 = \mathbf{1}$. \square

We introduce one more property of the ribbon structure that bears significance in many topological constructions.

Definition 2.20. Let H be a Hopf algebra carrying a (left) integral λ and a ribbon element v . We call H *anomaly-free* if

$$\lambda(v) = 1.$$

Otherwise we will refer to it as *anomalous*.

2.4 Algebra generators and tuple notation

All the constructions considered in this paper produce Hopf algebras generated by grouplike and skew-primitive generators.

Notation. We denote by K_a , $a = 1, \dots, s$ the grouplike generators that commute with one another and are of respective orders m_1, \dots, m_s . By X_k, X_k^\pm, Z_l^\pm for $k = 1, \dots, t_1$ and $l = 1, \dots, t_2$ we denote skew-primitive generators such that $(X_k)^\pm = (X_k^\pm)^2 = (Z_l^\pm)^2 = 0$.

We will repeatedly appeal to the following notion.

Definition 2.21. Let $g, h \in H$. We say g, h have a diagonal relation if there exists $\gamma \in \mathbb{C}$ such that

$$gh = \gamma hg.$$

Notation. We will use boldface for tuples of numbers and algebra generators, for instance, let $\mathbf{0} = (0, \dots, 0)$, and let

$$\boldsymbol{\xi} = (\xi_1, \dots, \xi_s) := (\exp^{2\pi/m_1}, \dots, \exp^{2\pi/m_s}) \in \mathbb{C}^s$$

be the tuple of primitive roots of unity of orders $\mathbf{m} = (m_1, \dots, m_s) \in \mathbb{Z}_{>0}^s$. We similarly express the lists of algebra generators as $\mathbf{K} = (K_1, \dots, K_s)$, or $\mathbf{X} = (X_1, \dots, X_k)$.

We will often use the following algebraic structures.

Notation. Let $\mathbb{Z}_{\mathbf{m}} := \mathbb{Z}_{m_1} \times \dots \times \mathbb{Z}_{m_s}$, where $\mathbb{Z}_{m_a} := \mathbb{Z}/m_a\mathbb{Z}$, and let $\mathbb{C}[\mathbf{K}] := \mathbb{C}[K_1, \dots, K_{m_s}] \subset H$ be the group subalgebra generated by K_1, \dots, K_s . It is isomorphic to $\mathbb{C}[\mathbb{Z}_{\mathbf{m}}]$ as a Hopf algebra.

We also introduce some operations on tuples.

Notation. For two tuples of the same shape, for instance $\mathbf{w}, \mathbf{v} \in \mathbb{Z}_{\mathbf{m}}$ we define the element-wise sum

$$\mathbf{w} + \mathbf{v} := (w_1 + v_1, \dots, w_s + v_s),$$

and the element-wise product

$$\mathbf{w} \cdot \mathbf{v} := (w_1 v_1, \dots, w_s v_s).$$

By a power of a tuple by a tuple, for instance for $\mathbf{v} \in \mathbb{Z}_{\mathbf{m}}$, $\boldsymbol{\xi} \in \mathbb{C}^s$ and \mathbf{K} , we mean

$$\boldsymbol{\xi}^{\mathbf{v}} = (\xi_1)^{v_1} (\xi_1)^{v_2} \dots (\xi_s)^{v_s},$$

and

$$\mathbf{K}^{\mathbf{v}} = (K_1)^{v_1} (K_2)^{v_2} \dots (K_s)^{v_s}.$$

Notation. We will also encounter *matrices*, denoted with boldface sans-serif font, for instance

$$\boldsymbol{\alpha} = (\alpha_{kl})_{1 \leq k, l \leq t} \in \text{Mat}_{t \times t}(\mathbb{C}) \quad \mathbf{d} = (d_{ka})_{\substack{1 \leq a \leq s \\ 1 \leq k \leq t}} \in \text{Mat}_{t \times s}(\mathbb{Z}_{\mathbf{m}}).$$

The rows $\mathbf{d}_k = (d_{ka})_{0 \leq a \leq s}$, $\mathbf{u}_k = (u_{ka})_{0 \leq a \leq s}$ are considered as tuples (and conform to the boldface notation). Products with no operation symbols mean the usual matrix multiplication, provided the participating matrices have appropriate dimensions and entries in the same ring. For $\mathbf{w}, \mathbf{v} \in \mathbb{Z}_{\mathbf{m}}$,

$$\mathbf{w}\mathbf{v}^T = w_1 v_1 + \dots + w_s v_s,$$

is the inner product of row tuples, valued in $\mathbb{Z}_{\text{lcm}(m_1, \dots, m_s)}$, where $\text{lcm}(m_1, \dots, m_s)$ is the least common multiple of m_1, \dots, m_s . For a matrix $\mathbf{z} \in \text{Mat}_{s \times s}(\mathbb{Z}_{\mathbf{m}})$, the expression $\mathbf{v}\mathbf{z}$ is a row tuple again, as is $(\mathbf{w}\mathbf{z}) \cdot \mathbf{v}$, but $\mathbf{w}\mathbf{z}\mathbf{v}^T$ is a scalar in $\mathbb{Z}_{\text{lcm}(m_1, \dots, m_s)}$.

3 The Nenciu construction

The Nenciu construction of [Nen04] produces finite-dimensional ribbon Hopf algebras generated by grouplike generators collected in a tuple $\mathbf{K} = (K_1, \dots, K_s)$ and nilpotent generators collected in a tuple $\mathbf{X} = (X_1, \dots, X_t)$, such that the group subalgebra $\mathbb{C}[\mathbf{K}]$ is abelian, the nilpotent generators are skew-primitive with respect to $\mathbb{C}[\mathbf{K}]$ and all commutation relations between generators are diagonal. The choice of the generators and relations is strongly constrained, and sufficient conditions for the result to be a ribbon Hopf algebra are determined in [Nen04]. We will see that strongly non-factorizable examples of this construction exist.

3.1 Hopf algebra structure

We start by recalling Nenciu's construction: the fact that Definition 3.1 indeed defines a Hopf algebra is a Theorem of [Nen04].

Definition 3.1. Let $\mathbf{m} \in \mathbb{Z}_{>0}^s$ be a tuple of positive integers of length s , and $t \in \mathbb{Z}_{>0}$ and \mathbf{d}, \mathbf{u} be dimension $t \times s$, \mathbb{Z} -valued matrices, such that

$$\boldsymbol{\xi}^{\mathbf{d}_k \cdot \mathbf{u}_l} \boldsymbol{\xi}^{\mathbf{d}_l \cdot \mathbf{u}_k} = 1, \text{ and } \boldsymbol{\xi}^{\mathbf{d}_k \cdot \mathbf{u}_k} = -1, \quad (1)$$

where $\boldsymbol{\xi} := (\exp^{2\pi/m_1}, \dots, \exp^{2\pi/m_s})$ is the tuple of primitive roots of unity of orders given by the tuple $\mathbf{m} \in \mathbb{Z}^s$. Define $H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$ to be the Hopf algebra generated by grouplike generators $\mathbf{K} = (K_1, \dots, K_s)$, and skew-primitive generators $\mathbf{X} = (X_1, \dots, X_t)$, and with the relations

$$\begin{aligned} K_a^{m_a} &= \mathbf{1} & K_a K_b &= K_b K_a & K_a X_k &:= \xi_a^{d_{ka}} X_k K_a \\ X_k^2 &= 0 & X_l X_k &:= \boldsymbol{\xi}^{\mathbf{d}_k \cdot \mathbf{u}_l} X_k X_l, \end{aligned}$$

coalgebra structure determined by

$$\begin{aligned} \epsilon(K_a) &:= 1 & \Delta(K_a) &:= K_a \otimes K_a \\ \epsilon(X_k) &:= 0 & \Delta(X_k) &:= \mathbf{1} \otimes X_k + X_k \otimes \mathbf{K}^{\mathbf{u}_k}, \end{aligned}$$

and antipode determined by

$$S(K_a) = K_a^{-1} := K_a^{m_a-1} \quad S(X_k) := -X_k \mathbf{K}^{-\mathbf{u}_k},$$

for $a, b = 1, \dots, s$, $s := |\mathbf{m}|$ and $k, l = 1, \dots, t$. Here \mathbf{m} is the tuple of orders of the grouplike elements, \mathbf{d} encode the diagonal relations for the grouplike elements, and \mathbf{u} the relations for the skew-primitive elements.

Remark 3.2. Note that in Definition 3.1 the relations $X_k^2 = 0$, for $k = 1, \dots, t$, result from the requirement that

$$\boldsymbol{\xi}^{\mathbf{d}_k \cdot \mathbf{u}_k} = -1.$$

holds, rather than being imposed independently. We include them in the definition for clarity.

Sometimes, we will choose several variables X_k to have the same commutation properties. We refer to this as *type*, and in examples we will indicate each type with a different letter X, Y, Z etc. if necessary.

Definition 3.3. By the *type* of X_k we mean the prescription of \mathbf{d}_k and \mathbf{u}_k , that is X_l is of the same type as X_k if $\mathbf{d}_k = \mathbf{d}_l$ and $\mathbf{u}_k = \mathbf{u}_l$. Since all relations involving X_k and X_l are diagonal, we will call the two generators of *opposite type* if for $a = 1, \dots, s$

$$d_{ka} \equiv -d_{la} \pmod{m_a}$$

and

$$u_{ka} \equiv -u_{la} \pmod{m_a}.$$

This essentially means that generators of opposite type commute with all other generators over reciprocal constants. We will sometimes indicate variables of opposite type using superscripts, for instance X_k^+, X_k^- .

Proposition 3.4. Let X_k^+, X_k^- be nilpotent generators of opposite type, then $\{X_k^+, X_k^-\} = 0$.

Proof. We have directly

$$X_k^+ X_k^- = \xi^{\mathbf{d}_k \cdot \mathbf{u}_k} X_k^- X_k^+ = (-1)^{-1} X_k^- X_k^+ = -X_k^- X_k^+.$$

□

Remark 3.5. When making general statements about $H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$ that do not invoke the types we will use the notation X_k for any nilpotent generator, with no superscripts. In such sense, two nilpotent generators with distinct indices, say X_k, X_l , can be of different, opposite or same type.

We can define a *monomial basis* for $H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$.

Proposition 3.6. The Hopf algebra $H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$ has a monomial basis

$$\{\mathbf{K}^{\mathbf{v}} \mathbf{X}^{\mathbf{r}} \mid \mathbf{v} \in \mathbb{Z}_{\mathbf{m}}, \mathbf{r} \in \mathbb{Z}_2^t\}.$$

Proof. The family $\{\mathbf{K}^{\mathbf{v}} \mathbf{X}^{\mathbf{r}} \mid \mathbf{v} \in \mathbb{Z}_{\mathbf{m}}, \mathbf{r} \in \mathbb{Z}_2^t\}$ is free since different elements involve different sets of generators, and the relations are diagonal. Furthermore, any element in the Hopf algebra is, by definition, a linear combination of products of elements of \mathbf{K} and \mathbf{X} . Since the relations are diagonal, up to a scalar, each term can be reordered so that the grouplike generators come first with a fixed order for all generators. Finally, the term is 0 if a given skew-primitive generator appears twice. Indeed, the relations are diagonal and the nilpotent generators square to 0.

□

Definition 3.7. We call $T := \mathbf{X}^{\mathbf{r}}$ for $\mathbf{r} = (1, \dots, 1)$ the *top element*.

It will play an important role in the unimodular structure we will sometimes define on $H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$. We will also need to use the adjoint action of generators on one another (see Definition 2.4). By direct computation, we get:

Proposition 3.8. The adjoint action of the generators of H is given by the following formulas. A grouplike $\mathbf{K}^{\mathbf{w}}$, $\mathbf{w} \in \mathbb{Z}_{\mathbf{m}}$ acts on the skew-primitive generator X_k as

$$\mathbf{K}^{\mathbf{w}} \triangleright X_k = \mathbf{K}^{\mathbf{w}} X_k (\mathbf{K}^{\mathbf{w}})^{-1} = \xi^{\mathbf{w} \cdot \mathbf{d}_k} X_k,$$

a skew-primitive X_k acts on a grouplike $\mathbf{K}^{\mathbf{w}}$ as

$$X_k \triangleright \mathbf{K}^{\mathbf{w}} = \mathbf{K}^{\mathbf{w}} S(X_k) + X_k \mathbf{K}^{\mathbf{w}} \mathbf{K}^{-\mathbf{u}_k} = (1 - \xi^{\mathbf{w} \cdot \mathbf{d}_k}) X_k \mathbf{K}^{-\mathbf{u}_k} \mathbf{K}^{\mathbf{w}},$$

and a skew-primitive X_k acts on a skew-primitive X_l as

$$X_k \triangleright X_l = X_l S(X_k) + X_k X_l \mathbf{K}^{-\mathbf{u}_k} = (\xi^{\mathbf{u}_k \cdot \mathbf{d}_l} - 1) X_k \mathbf{K}^{-\mathbf{u}_k} X_l.$$

3.2 Examples

In this subsection we study explicit examples of the Nenciu construction. We start with a (not strongly) non-factorizable Hopf algebra, which can be retrieved from Nenciu's construction, and appears in [Nen04] as $E(n)$, first studied in [PO99]. Here we will refer to it as *symplectic fermions*, after [GR17], also appearing in [FGS24]. The notation using the variables L, Z_l^{\pm} is introduced to be used in further examples.

Definition 3.9. The Hopf algebra of symplectic fermions, SF_{2n} is the Hopf algebra generated by $K_1 = L$ and, $X_{2l-1} = Z_l^+$, $X_{2l} = Z_l^-$, for $k, l = 1, \dots, n$, subject to the relations

$$L^2 = \mathbf{1}, \quad LZ_l^{\pm} = -Z_l^{\pm}L, \quad \{Z_l^{\pm}, Z_k^{\pm}\} = \{Z_l^{\pm}, Z_k^{\mp}\} = 0.$$

The Hopf structure is defined by

$$\begin{aligned} \epsilon(L) &= 1, \quad \epsilon(Z_l^{\pm}) = 0, \\ \Delta(L) &= L \otimes L, \quad \Delta(Z_l^{\pm}) = \mathbf{1} \otimes Z_l^{\pm} + Z_l^{\pm} \otimes L, \\ S(L) &= L^{-1} = L, \quad S(Z_l^{\pm}) = -Z_l^{\pm}L. \end{aligned}$$

The dimension of the algebra is 2^{2n+1} . The corresponding Nenciu data is

- $\mathbf{m} = (2)$, so $s = 1$
- $t = 2n$

- $\mathbf{d} = \mathbf{u} = (1, \dots, 1)^T$ are column vectors wherein each row is $d_k = 1$ and $u_k = 1$, $k = 1, \dots, t$

$$\mathbf{d} = \begin{pmatrix} 1 \\ \vdots \\ 1 \end{pmatrix}, \quad \mathbf{u} = \begin{pmatrix} 1 \\ \vdots \\ 1 \end{pmatrix}.$$

We now increase the number of grouplike and nilpotent generators, to produce previously unknown examples that will all turn out to admit only strongly non-factorizable quasitriangular structures.

Example 3.10. Let \mathbb{N}_1 be the Hopf algebra generated by K_a, X_k, Z_l^\pm for $a \in \{1, 2\}$, $j, k \in \{1, \dots, t_1\}$; $t_1 \in 4\mathbb{N}$ and $l, m \in \{1, \dots, t_2\}$; $t_2 \in \mathbb{N}$, subject to the following relations

$$K_a^4 = \mathbf{1}, \quad K_a X_j = i X_j K_a, \quad K_1 Z_l^\pm = \pm i Z_l^\pm K_1, \quad K_2 Z_l^\pm = Z_l^\pm K_2,$$

$$Z_l^\pm X_k = \mp i X_k Z_l^\pm, \quad \{X_j, X_k\} = \{Z_l^\pm, Z_m^\pm\} = \{Z_l^\pm, Z_m^\mp\} = 0,$$

where $i^2 = -1$. The Hopf structure is defined by

$$\epsilon(K_a) = 1, \quad \epsilon(X_k) = \epsilon(Z_l^\pm) = 0,$$

$$\Delta(K_a) = K_a \otimes K_a, \quad \Delta(X_k) = \mathbf{1} \otimes X_k + X_k \otimes K_1 K_2, \quad \Delta(Z_l^\pm) = \mathbf{1} \otimes Z_l^\pm + Z_l^\pm \otimes K_1^{\pm 2} K_2^{\pm 1},$$

$$S(K_a) = K_a^{-1}, \quad S(X_k) = -X_k (K_1 K_2)^{-1}, \quad S(Z_l^\pm) = -Z_l^\pm K_1^{\mp 2} K_2^{\mp 1}.$$

The Nenciu data for this Hopf algebra is

- $\mathbf{m} = (4, 4)$, with $s = 2$
- $t = t_1 + 2t_2$
- \mathbf{d} and \mathbf{u} are $t \times 2$ matrices, such that for $k = 1, \dots, t_1$; $\mathbf{d}_k = (1, 1)$, and $\mathbf{u}_k = (1, 1)$ corresponding to X_k . For $l = t_1 + 1, \dots, t_2$ the rows are $\mathbf{d}_{2l-1} = (1, 0)$, $\mathbf{u}_{2l-1} = (2, 1)$ corresponding to Z_l^+ and $\mathbf{d}_{2l} = (-1, 0)$, $\mathbf{u}_{2l} = (-2, -1)$ corresponding to Z_l^- .

$$\mathbf{d} = \begin{pmatrix} 1 & 1 \\ \vdots & \vdots \\ 1 & 1 \\ 1 & 0 \\ -1 & 0 \\ \vdots & \vdots \\ 1 & 0 \\ -1 & 0 \end{pmatrix}, \quad \mathbf{u} = \begin{pmatrix} 1 & 1 \\ \vdots & \vdots \\ 1 & 1 \\ 2 & 1 \\ -2 & -1 \\ \vdots & \vdots \\ 2 & 1 \\ -2 & -1 \end{pmatrix}.$$

The instance of Example 3.10 with the smallest dimension occurs for $t_1 = 4$, $t_2 = 1$, and its dimension is $4 \times 4 \times 2^6 = 2^{10} = 1024$. It will be shown later this Hopf algebra admits only triangular ribbon structures. The following examples were designed to admit strongly non-factorizable quasitriangular structures which are not triangular, as will be shown later.

Example 3.11. Let \mathbb{N}_2 be the Hopf algebra generated by K_a, X_k^\pm and Z_l^\pm , for $a = 1, 2, 3$, $j, k = 1, \dots, t_1$; $t_1 \in \mathbb{N}$ and $l, m = 1, \dots, t_2$, $t_2 \in \mathbb{N}$, subject to the following relations

$$K_a^4 = \mathbf{1}, \quad K_a X_k^\pm = \pm i X_k^\pm K_a$$

$$K_1 Z_l^\pm = Z_l^\pm K_1, \quad K_2 Z_l^\pm = -Z_l^\pm K_2, \quad K_3 Z_l^\pm = \pm i Z_l^\pm K_3$$

$$\{X_j^\pm, X_k^\pm\} = \{X_j^\pm, X_k^\mp\} = \{Z_l^\pm, X_k^\pm\} = \{Z_l^\pm, X_k^\mp\} = \{Z_l^\pm, Z_m^\pm\} = \{Z_l^\pm, Z_m^\mp\} = 0.$$

Let also $L := K_3^2$ as a shorthand, note that $L^2 = \mathbf{1}$. The Hopf structure is defined by

$$\epsilon(K_a) = 1, \quad \epsilon(X_k^\pm) = \epsilon(Z_l^\pm) = 0,$$

$$\Delta(K_a) = K_a \otimes K_a, \quad \Delta(X_k^\pm) = \mathbf{1} \otimes X_k^\pm + X_k^\pm \otimes (K_1 K_2)^{\pm 1}, \quad \Delta(Z_l^\pm) = \mathbf{1} \otimes Z_l^\pm + Z_l^\pm \otimes L$$

$$S(K_a) = K_a^{-1}, \quad S(X_k^\pm) = -X_k^\pm (K_1 K_2)^{\mp 1}, \quad S(Z_l^\pm) = -Z_l^\pm L.$$

The Nenciu data for this Hopf algebra is

- $\mathbf{m} = (4, 4, 4)$, so $s = 3$
- $t = 2t_1 + 2t_2$
- Then \mathbf{d} and \mathbf{u} are $t \times 3$ matrices, such that for $j = 1, \dots, t_1$ $\mathbf{d}_{2j-1} = (1, 1, 1)$, and $\mathbf{u}_{2j-1} = (1, 1, 0)$ corresponding to X_j^+ and $\mathbf{d}_{2j} = (-1, -1, -1)$, and $\mathbf{u}_{2j} = (-1, -1, 0)$ corresponding to X_j^- .

For $l = 2t_1 + 1, \dots, t_2$ the rows are $\mathbf{d}_{2l-1} = (0, 2, 1)$, $\mathbf{u}_{2l-1} = (0, 0, 2)$ corresponding to Z_l^+ and $\mathbf{d}_{2l} = (0, -2, -1)$, $\mathbf{u}_{2l} = (0, 0, -2)$ corresponding to Z_l^- .

$$\mathbf{d} = \begin{pmatrix} 1 & 1 & 1 \\ -1 & -1 & -1 \\ & \vdots & \\ 1 & 1 & 1 \\ -1 & -1 & -1 \\ 0 & 2 & 1 \\ 0 & -2 & -1 \\ & \vdots & \\ 0 & 2 & 1 \\ 0 & -2 & -1 \end{pmatrix}, \quad \mathbf{u} = \begin{pmatrix} 1 & 1 & 0 \\ -1 & -1 & 0 \\ & \vdots & \\ 1 & 1 & 0 \\ -1 & -1 & 0 \\ 0 & 0 & 2 \\ 0 & 0 & -2 \\ & \vdots & \\ 0 & 0 & 2 \\ 0 & 0 & -2 \end{pmatrix}.$$

The instance of Example 3.11 with the smallest dimension occurs for $t_1 = 1$, $t_2 = 1$, and its dimension is $4 \times 4 \times 4 \times 2^4 = 2^{10} = 1024$.

Note that in Example 3.11 all the relations between skew-primitive generators are anti-commutations. By introducing new skew-primitive generators Y_k^\pm , we can produce commutations involving constants of higher order, while retaining the strongly non-factorizable, but not triangular, quasitriangular structure.

Example 3.12. Let \mathbb{N}_3 be the Hopf algebra generated by K_a , X_k^\pm , Y_k^\pm and Z_l^\pm , for $a = 1, 2, 3$, $j, k = 1, \dots, t_1$; $t_1 \in \mathbb{N}$ and $l, m = 1, \dots, t_2$; $t_2 \in \mathbb{N}$, subject to the following relations

$$\begin{aligned} K_a^4 &= \mathbf{1}, & K_a X_k^\pm &= \pm i X_k^\pm K_a, \\ K_1 Y_k^\pm &= \pm i Y_k^\pm K_1, & K_2 Y_k^\pm &= Y_k^\pm K_2, & K_3 Y_k^\pm &= \pm i Y_l^\pm K_3 \\ K_1 Z_l^\pm &= Z_l^\pm K_1, & K_2 Z_l^\pm &= -Z_l^\pm K_2, & K_3 Z_l^\pm &= \pm i Z_l^\pm K_3 \\ \{X_j^\pm, X_k^\pm\} &= \{X_j^\pm, X_k^\mp\} = \{Y_j^\pm, Y_k^\pm\} = \{Y_j^\pm, Y_k^\mp\}, & X_j Y_k &= i Y_k X_j \\ \{Z_l^\pm, X_k^\pm\} &= \{Z_l^\pm, X_k^\mp\} = \{Z_l^\pm, Y_k^\pm\} = \{Z_l^\pm, Y_k^\mp\} = \{Z_l^\pm, Z_m^\pm\} = \{Z_l^\pm, Z_m^\mp\} = 0. \end{aligned}$$

Let also $L := K_3^2$ as a shorthand, note that $L^2 = \mathbf{1}$. The Hopf structure is defined by

$$\begin{aligned} \epsilon(K_a) &= 1, & \epsilon(X_k^\pm) &= \epsilon(Y_k^\pm) = \epsilon(Z_l^\pm) = 0, \\ \Delta(K_a) &= K_a \otimes K_a, & \Delta(X_k^\pm) &= \mathbf{1} \otimes X_k^\pm + X_k^\pm \otimes (K_1 K_2)^{\pm 1}, \\ \Delta(Y_k^\pm) &= \mathbf{1} \otimes Y_k^\pm + Y_k^\pm \otimes (K_1 K_2^2)^{\pm 1}, & \Delta(Z_l^\pm) &= \mathbf{1} \otimes Z_l^\pm + Z_l^\pm \otimes L \\ S(K_a) &= K_a^{-1} & S(X_k^\pm) &= -X_k^\pm (K_1 K_2)^{\mp 1}, & S(Y_k^\pm) &= -Y_k^\pm (K_1^2 K_2)^{\mp 1} & S(Z_l^\pm) &= -Z_l^\pm L. \end{aligned}$$

The Nenciu data for this Hopf algebra is

- $\mathbf{m} = (4, 4, 4)$, so $s = 3$
- $t = 4t_1 + 2t_2$
- Then \mathbf{d} and \mathbf{u} are $t \times 3$ matrices, such that for $j = 1, \dots, t_1$ $\mathbf{d}_{2j-1} = (1, 1, 1)$, and $\mathbf{u}_{2j-1} = (1, 1, 0)$ corresponding to X_j^+ and $\mathbf{d}_{2j} = (-1, -1, -1)$, and $\mathbf{u}_{2j} = (-1, -1, 0)$ corresponding to X_j^- . For $k = t_1 + 1, \dots, 2t_1$, $\mathbf{d}_{2k-1} = (1, 0, 1)$, and $\mathbf{u}_{2j-1} = (2, 1, 0)$ corresponding to Y_k^+ and $\mathbf{d}_{2k} = (-1, -0, -1)$, and $\mathbf{u}_{2k} = (-2, -1, 0)$ corresponding to Y_k^- . For $l = 2t_1 + 1, \dots, t_2$ the rows are $\mathbf{d}_{2l-1} = (0, 2, 1)$, $\mathbf{u}_{2l-1} = (0, 0, 2)$ corresponding to Z_l^+ and $\mathbf{d}_{2l} = (0, -2, -1)$, $\mathbf{u}_{2l} = (0, 0, -2)$ corresponding to Z_l^- .

$$\mathbf{d} = \begin{pmatrix} 1 & 1 & 1 \\ -1 & -1 & -1 \\ & \vdots & \\ 1 & 1 & 1 \\ -1 & -1 & -1 \\ 1 & 0 & 1 \\ -1 & 0 & -1 \\ & \vdots & \\ 1 & 0 & 1 \\ -1 & 0 & -1 \\ 0 & 2 & 1 \\ 0 & -2 & -1 \\ & \vdots & \\ 0 & 2 & 1 \\ 0 & -2 & -1 \end{pmatrix}, \quad \mathbf{u} = \begin{pmatrix} 1 & 1 & 0 \\ -1 & -1 & 0 \\ & \vdots & \\ 1 & 1 & 0 \\ -1 & -1 & 0 \\ 2 & 1 & 0 \\ -2 & -1 & 0 \\ & \vdots & \\ 2 & 1 & 0 \\ -2 & -1 & 0 \\ 0 & 0 & 2 \\ 0 & 0 & -2 \\ & \vdots & \\ 0 & 0 & 2 \\ 0 & 0 & -2 \end{pmatrix}.$$

The instance of Example 3.12 with the smallest dimension occurs for $t_1 = 1$, $t_2 = 1$, and its dimension is $4 \times 4 \times 4 \times 2^6 = 2^{12} = 4096$.

3.3 Unimodular structure

In this short section we construct integrals and cointegrals on $H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$ and determine the sufficient condition for unimodularity.

Definition 3.13 ([Nen04], Proposition 4.7). Let $H = H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$ be a Hopf algebra such as in Definition 3.1. Then H admits a left cointegral

$$\Lambda_L := \prod_{a=1}^s \left(\sum_{b=0}^{m_a-1} K_a^b \right) \prod_{k=1}^t X_k,$$

and a right cointegral

$$\Lambda_R := \prod_{k=1}^t X_k \prod_{a=1}^s \left(\sum_{b=0}^{m_a-1} K_a^b \right).$$

Let us denote the top element by $T := \prod_{k=1}^t X_k = \mathbf{X}^{(1, \dots, 1)}$ then a two-sided integral λ is defined on the monomial basis by the formula

$$\lambda(\mathbf{K}^{\mathbf{v}} \mathbf{X}^{\mathbf{r}}) := \begin{cases} 1 & \text{if } \mathbf{K}^{\mathbf{v}} \mathbf{X}^{\mathbf{r}} = T \\ 0 & \text{otherwise.} \end{cases}$$

Remark 3.14. Note that $\prod_{a=1}^s \left(\sum_{b=0}^{m_a-1} K_a^b \right)$ is simply the sum of all monomials in the Hopf subalgebra $\mathbb{C}[\mathbf{K}]$.

Proposition 3.15. We have $\Lambda_L = \Lambda_R$ if T is central, which happens if and only if for all $a = 1, \dots, s$

$$\sum_{k=1}^t d_{ka} \equiv 0 \pmod{m_a}.$$

In particular Λ_L is then a two-sided cointegral.

Proof. Firstly, using the monomial basis, and the fact that relations are diagonal, we see that $\Lambda_L = \Lambda_R$ if and only if $\mathbf{K}^{\mathbf{v}}$ commutes with T for every \mathbf{v} . This is tantamount to T being central, since T always commutes with any X_j . Indeed $TX_j = X_jT = 0$, as all skew-primitive generators square to 0 and relations are diagonal. Secondly, this happens if and only if T commutes with K_a for every a . But we can compute directly that

$$K_a T = \prod_{k=1}^t \xi_a^{d_{ka}} T K_a.$$

Thus, T is central if and only if $\sum_{k=1}^t d_{ka} \equiv 0$ in \mathbb{Z}_{m_a} . \square

3.4 Quasitriangular structure

In [Nen04] a sufficient list of conditions is given for $H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$ to admit a ribbon structure. In this section, we recall these results, and we fix an imprecision concerning the existence of the pivotal element in the unimodular case, which we discuss in further sections. These rely on Theorem 2.9. We then revisit some examples of Nenciu and prove they admit unimodular and strongly non-factorizable structures. We end the section by computing the monodromy matrix associated to these examples.

As we have recalled for any Hopf algebra H its dual H^* is again a Hopf algebra. In our setting we have the following result:

Proposition 3.16 ([Nen04], Proposition 3.1). Let $H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$ be as in Definition 3.1, then $H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})^*$ is isomorphic to $H(\mathbf{m}, t, \mathbf{u}, \mathbf{d})$ as a Hopf algebra, and the duals of the generators K_a and X_k , denoted \mathcal{K}_a and \mathcal{X}_k , correspond to the generators of $H(\mathbf{m}, t, \mathbf{u}, \mathbf{d})$ through this isomorphism.

Notation. In the discussion that follows we will distinguish the algebra $H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$ and its dual, so we introduce a notation where $H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})^* \cong H(\mathbf{m}, t, \mathbf{u}, \mathbf{d})$ is generated by grouplike generators \mathcal{K}_a dual to K_a , for $a = 1, \dots, s$ and skew-primitive generators \mathcal{X}_k dual to X_k for $k = 1, \dots, t$ in the caligraphic font, satisfying the relations

$$\begin{aligned} \mathcal{K}_a^{m_a} &= \mathbf{1}, \quad \mathcal{K}_a \mathcal{K}_b = \mathcal{K}_b \mathcal{K}_a, \quad \mathcal{K}_a \mathcal{X}_k := \xi^{u_{ka}} \mathcal{X}_k \mathcal{K}_a, \quad \mathcal{X}_l \mathcal{X}_k = \xi^{\mathbf{u}^k \cdot \mathbf{d}_l} \mathcal{X}_k \mathcal{X}_l, \\ \epsilon(\mathcal{K}_a) &:= 1, \quad \Delta(\mathcal{K}_a) := \mathcal{K}_a \otimes \mathcal{K}_a, \quad \epsilon(\mathcal{X}_k) := 0, \quad \Delta(\mathcal{X}_k) := \mathbf{1} \otimes \mathcal{X}_k + \mathcal{X}_k \otimes \mathcal{K}^{\mathbf{d}_k} \end{aligned}$$

and antipode

$$S(\mathcal{K}_a) := \mathcal{K}_a^{-1} = \mathcal{K}_a^{m_a-1} \qquad S(\mathcal{X}_k) := -\mathcal{X}_k \mathcal{K}^{-\mathbf{d}_k}, \qquad (2)$$

for $a, b = 1, \dots, s$, and $k, l = 1, \dots, t$. We also have the the monomial basis of the dual $H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$ denoted by

$$\{\mathcal{K}^{\mathbf{v}} \mathcal{X}^{\mathbf{r}} \mid \mathbf{v} \in \mathbb{Z}_{\mathbf{m}}, \mathbf{r} \in \mathbb{Z}_2^t\},$$

with the duality pairing for any $\mathbf{w}, \mathbf{v} \in \mathbb{Z}_{\mathbf{m}}$ and $\mathbf{p}, \mathbf{r} \in \mathbb{Z}_2^t$ we have

$$\mathcal{K}^{\mathbf{w}} \mathcal{X}^{\mathbf{p}}(\mathcal{K}^{\mathbf{v}} \mathcal{X}^{\mathbf{r}}) = \prod_{a=1}^s \delta_{w_a, v_a} \prod_{k=1}^t \delta_{p_k, r_k}.$$

The following is a Corollary of Prop. 3.16.

Corollary 3.17 ([Nen04], Section 3). Any Hopf algebra map $f : H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})^* \rightarrow H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})^{cop}$ is parametrized by a \mathbb{Z} -valued $s \times s$ matrix \mathbf{z} and a \mathbb{C} -valued $t \times t$ matrix $\boldsymbol{\alpha}$ as follows:

$$\mathcal{K}_a \mapsto \mathbf{K}^{\mathbf{z}^a},$$

where $\mathbf{z}_a \in \mathbb{Z}^s$ is the a -th row of \mathbf{z} for $a = 1, \dots, s$, and

$$\mathcal{X}_k \mapsto \sum_{l=1}^t \alpha_{kl} \mathbf{K}^{-\mathbf{u}_l} X_l$$

for $k, l = 1, \dots, t$. With these definitions the image of an arbitrary element $\mathbf{K}^{\mathbf{v}} \mathcal{X}^{\mathbf{p}}$ for $\mathbf{v} \in \mathbb{Z}_{\mathbf{m}}$ and $\mathbf{p} \in \mathbb{Z}_2^t$, is

$$f(\mathbf{K}^{\mathbf{v}} \mathcal{X}^{\mathbf{p}}) = \sum_{\mathbf{q} \in \mathbb{Z}_2^t} \left(\prod_{j,k=1}^t \alpha_{jk} p_j q_k \right) \mathbf{K}^{\mathbf{v}\mathbf{z}} (\mathbf{K}^{-\mathbf{u}_1} X_1, \dots, \mathbf{K}^{-\mathbf{u}_t} X_t)^{\mathbf{q}}.$$

Thus, we have a general parametrization for any Hopf map $f : H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})^* \rightarrow H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})^{cop}$, where $H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})^{cop}$ is the Hopf algebra $H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$ with the coproduct modified to Δ^{cop} . We denote the elements of $H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})^{cop}$ with the same symbols as those of $H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$, as the algebra structure remains unchanged.

The \mathbf{p} and \mathbf{q} are row tuples with values in $\{0, 1\}$ and they select a subset of skew-primitive generators present. In particular, $(\mathbf{K}^{-\mathbf{u}_1} X_1, \dots, \mathbf{K}^{-\mathbf{u}_t} X_t)^{\mathbf{q}}$ is a product of terms of the form $\mathbf{K}^{-\mathbf{u}_k} X_k$ according to the selection by \mathbf{q}

$$(\mathbf{K}^{-\mathbf{u}_1} X_1, \dots, \mathbf{K}^{-\mathbf{u}_t} X_t)^{\mathbf{q}} = \prod_{k=1}^t q_k \mathbf{K}^{-\mathbf{u}_k} X_k.$$

Remark 3.18. Not any choice of matrices \mathbf{z} and $\boldsymbol{\alpha}$ gives a valid bialgebra map, and we will list the necessary criteria below, in Theorem 3.19.

This construction provides a weak converse to Theorem 2.9, by reconstructing the R-matrix from the bialgebra map in the chosen parametrization. In essence, every term in the R-matrix expansion is a tensor product of a monomial in $H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$ and the image of its dual under f .

Theorem 3.19 ([Nen04], Theorem 4.1). Let $R \in H(\mathbf{m}, t, \mathbf{d}, \mathbf{u}) \otimes H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$ be the element defined by

$$R := R_{\mathbf{z}} R_{\boldsymbol{\alpha}},$$

for

$$R_{\mathbf{z}} = \frac{1}{\prod_{a=1}^s m_a} \sum_{\mathbf{v}, \mathbf{w} \in \mathbb{Z}_{\mathbf{m}}} \xi^{-\mathbf{v} \cdot \mathbf{w}} \mathbf{K}^{\mathbf{w}} \otimes \mathbf{K}^{\mathbf{v}\mathbf{z}},$$

and

$$R_{\boldsymbol{\alpha}} = \exp \left(\sum_{k,l=1}^t \alpha_{kl} X_k \otimes \mathbf{K}^{-\mathbf{u}_l} X_l \right),$$

where $\mathbf{z} \in \text{Mat}_{s \times s}(\mathbb{Z}_{\mathbf{m}})$ and $\boldsymbol{\alpha} \in \text{Mat}_{t \times t}(\mathbb{C})$, are matrices as in Corollary 3.17. Then R is an R-matrix, if and only if the following constraints are satisfied.

(A1) for any $a = 1, \dots, s$, $b = 1, \dots, s$

$$m_a z_{ab} \equiv 0 \pmod{m_b}$$

(A2) for any $a = 1, \dots, s$ and $k, l = 1, \dots, t$

$$\alpha_{kl} \boldsymbol{\xi}^{\mathbf{d}_l \cdot \mathbf{z}_a} = \alpha_{kl} \boldsymbol{\xi}_a^{u_{ka}}$$

(A3) for any $i, j, k, l = 1, \dots, t$

$$(\alpha_{jl} \alpha_{ik} \boldsymbol{\xi}^{\mathbf{d}_k \cdot \mathbf{u}_l} + \alpha_{jk} \alpha_{il}) \boldsymbol{\xi}^{\mathbf{d}_k \cdot \mathbf{u}_l} = (\alpha_{jl} \alpha_{ik} \boldsymbol{\xi}^{\mathbf{d}_k \cdot \mathbf{u}_l} + \alpha_{jk} \alpha_{il}) \boldsymbol{\xi}^{\mathbf{d}_j \cdot \mathbf{u}_i}$$

(B) for any $k = 1, \dots, t$

$$\sum_{l=1}^t \alpha_{kl} \left(\mathbf{1} \otimes \mathbf{K}^{-\mathbf{u}_l} X_l + \mathbf{K}^{-\mathbf{u}_l} X_l \otimes \mathbf{K}^{\mathbf{d}_k \mathbf{z}} \right) = \sum_{l=1}^t \alpha_{kl} \left(\mathbf{1} \otimes \mathbf{K}^{-\mathbf{u}_l} X_l + \mathbf{K}^{-\mathbf{u}_l} X_l \otimes \mathbf{K}^{-\mathbf{u}_l} \right),$$

(C1) for all $k = 1, \dots, t$

$$\mathbf{d}_k \mathbf{z} = -\mathbf{u}_k$$

(C2) for all $k = 1, \dots, t$, and $\mathbf{v} \in \mathbb{Z}_{\mathbf{m}}$

$$\boldsymbol{\xi}^{\mathbf{v} \cdot \mathbf{u}_k} = \boldsymbol{\xi}^{\mathbf{d}_k \cdot \mathbf{v}\mathbf{z}}$$

(C3) whenever $(\mathbf{p}\boldsymbol{\alpha})^{\mathbf{q}} \neq 0$, for some $\mathbf{p}, \mathbf{q} \in \mathbb{Z}_2^t$ and for all $a = 1, \dots, s$

$$\sum_l^t (p_l + q_l) d_{la} \equiv 0 \pmod{m_a}$$

(C4) whenever $(\mathbf{p}\boldsymbol{\alpha})^{\mathbf{q}} \neq 0$, for some $\mathbf{p}, \mathbf{q} \in \mathbb{Z}_2^t$, then for $k = 1, \dots, t$

$$\boldsymbol{\xi}^{\mathbf{d}_k \cdot (\mathbf{p}\mathbf{u})} \boldsymbol{\xi}^{\mathbf{d}_k \cdot (\mathbf{q}\mathbf{u})} = 1.$$

In (B) we have in particular that for any k, l , either $\alpha_{kl} = 0$ or $\mathbf{d}_k \mathbf{z} = -\mathbf{u}_l$. Note also that the latter two items (C3), (C4) are required if $R_{\boldsymbol{\alpha}} \neq \mathbf{1} \otimes \mathbf{1}$, otherwise they are void. They are involved in realizing (QT5) for the grouplike K_a and nilpotent X_k generators for $a = 1, \dots, s$ and $k = 1, \dots, t$ respectively.

Proof. Let $f : H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})^* \rightarrow H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})^{cop}$ be the map induced by R as in Theorem 2.9. In particular, (A1)-(A3) together with (B) are equivalent to (QT1)-(QT4) and (C1)-(C4) are equivalent to (QT5). We first of all note that (A1), (C1)-(C4) are a reformulation of [Nen04] Theorem 4.1 in our notation. We discuss the remaining conditions that, while implied in [Nen04], were not collected in Theorem 4.1 there.

Conditions (A1)-(A3) imply that f is an algebra map. Towards (A2) let $\mathcal{K}_a, \mathcal{X}_k$ be the images of K_a, X_k , respectively, under the duality using the notation we introduced. Item (A2) is the expression of the fact that the commutation of $\mathcal{K}_a, \mathcal{X}_k$ has to be respected by the map f . Explicitly, if $\mathcal{K}_a \mathcal{X}_k = \boldsymbol{\xi}^{\mathbf{d}_a \cdot \mathbf{u}_k} \mathcal{X}_k \mathcal{K}_a$, then

$$f(\mathcal{K}_a \mathcal{X}_k) = \mathbf{K}^{\mathbf{z}_a} \sum_{l=1}^t \alpha_{kl} \mathbf{K}^{-\mathbf{u}_l} X_l = \sum_{l=1}^t \alpha_{kl} \boldsymbol{\xi}^{\mathbf{z}_a \cdot \mathbf{d}_l} \mathbf{K}^{-\mathbf{u}_l} X_l \mathbf{K}^{\mathbf{z}_a}.$$

On the other hand

$$f(\mathcal{X}_k \mathcal{K}_a) = \xi_a^{u_{ka}} f(\mathcal{K}_a \mathcal{X}_k) = \sum_{l=1}^t \alpha_{kl} \xi_a^{u_{ka}} \mathbf{K}^{-\mathbf{u}_l} X_l.$$

Comparing coefficients elementwise yields the required condition. This is enough as we defined the map on the monomial basis and all the relations are diagonal.

Similarly, (A3) provides this for the commutation of \mathcal{X}_l with \mathcal{X}_k . The calculation is analogous to the case (A2).

Condition (B) expresses the fact that f is a coalgebra map for \mathcal{X}_k , for \mathcal{K}_a this already follows from (A1). We have that

$$\begin{aligned} \Delta^{cop}(f(\mathcal{X}_k)) &= \Delta^{cop} \left(\sum_{l=1}^t \alpha_{kl} \mathbf{K}^{-\mathbf{u}_l} X_l \right) \\ &= \sum_{l=1}^t \alpha_{kl} (\mathbf{K}^{-\mathbf{u}_l} X_l \otimes \mathbf{K}^{-\mathbf{u}_l} + \mathbf{1} \otimes \mathbf{K}^{-\mathbf{u}_l} X_l), \end{aligned}$$

but

$$\begin{aligned} f(\Delta(\mathcal{X}_k)) &= f(\mathbf{1}) \otimes f(\mathcal{X}_k) + f(\mathcal{X}_k) \otimes f(\mathbf{K}^{\mathbf{d}_k}) \\ &= \mathbf{1} \otimes \sum_{l=1}^t \alpha_{kl} \mathbf{K}^{-\mathbf{u}_l} X_l + \sum_{l=1}^t \alpha_{kl} \mathbf{K}^{-\mathbf{u}_l} X_l \otimes \mathbf{K}^{-\mathbf{u}_k} \end{aligned}$$

since $\mathbf{d}_k \mathbf{z} = -\mathbf{u}_k$. In the latter expression combining the two sums yields

$$\sum_{l=1}^t \alpha_{kl} (\mathbf{1} \otimes \mathbf{K}^{-\mathbf{u}_l} X_l + \mathbf{K}^{-\mathbf{u}_l} X_l \otimes \mathbf{K}^{\mathbf{d}_k \mathbf{z}}).$$

Elementwise comparison of the two expressions retrieves (B), as above.

The exponential in the piece $R_{\boldsymbol{\alpha}}$ expands to the summation of all partial products between the elements of the sum in the exponent. In more detail, it can be expanded as

$$R_{\boldsymbol{\alpha}} = \exp \left(\sum_{k,l=1}^t \alpha_{kl} X_k \otimes \mathbf{K}^{-\mathbf{u}_l} X_l \right) = \sum_{j \in \text{Mat}_{t \times t}(\mathbb{Z}_2)} \prod_{k,l=1}^t \alpha_{kl} (X_k)^{j_{kl}} \otimes (\mathbf{K}^{-\mathbf{u}_l} X_l)^{j_{kl}}.$$

Each piece of the rightmost sum corresponds in the former tensor factor to a monomial in X_k generators, $\mathbf{X}^{\mathbf{r}}$, and in the latter tensor factor its image under the composition of the duality (which, recall, is only a linear map), and the bialgebra map so the element $f((\mathbf{X}^{\mathbf{r}})^*) \in H^{cop}$. The parameterization coefficients α_{kl} control whether a term appears in the sum, or equivalently in the image of f , or not, and with which coefficient.

To complete the argument, we can verify that $f_R : H^* \rightarrow H$ of Theorem 2.9 retrieved from $R_{\mathbf{z}} R_{\boldsymbol{\alpha}}$ in fact coincides with f . This is because for any $\mathbf{u}, \mathbf{o} \in \mathbb{Z}_{\mathbf{m}}$ and $\mathbf{p}, \mathbf{r} \in \mathbb{Z}_2^t$ we have

$$\mathcal{K}^{\mathbf{u}} \mathcal{X}^{\mathbf{p}} (\mathbf{K}^{\mathbf{o}} \mathbf{X}^{\mathbf{r}}) = \prod_{a=1}^s \delta_{u_a, o_a} \prod_{k=1}^t \delta_{p_k, r_k}.$$

Therefore

$$\begin{aligned}
& f_R(\mathcal{K}^{\mathbf{u}}\mathcal{X}^{\mathbf{p}}) \\
&= \sum_{\mathbf{w}, \mathbf{v} \in \mathbb{Z}_{\mathbf{m}}} \sum_{j \in \text{Mat}_{t \times t}(\mathbb{Z}_2)} \prod_{k,l=1}^t \alpha_{kl} \xi^{-\mathbf{w} \cdot \mathbf{v}} (\mathcal{K}^{\mathbf{u}}(\mathcal{X}_k)^{p_k}) (\mathbf{K}^{\mathbf{w}}(X_k)^{j_{kl}}) \otimes \mathbf{K}^{\mathbf{vz}}(\mathbf{K}^{-\mathbf{u}_l} X_l)^{j_{kl}} \\
&= \sum_{\mathbf{w}, \mathbf{v} \in \mathbb{Z}_{\mathbf{m}}} \sum_{j \in \text{Mat}_{t \times t}(\mathbb{Z}_2)} \prod_{k,l=1}^t \alpha_{kl} \xi^{-\mathbf{w} \cdot \mathbf{v}} \prod_{a=1}^s \delta_{u_a, w_a} \delta_{p_k, j_{kl}} \mathbf{K}^{\mathbf{vz}}(\mathbf{K}^{-\mathbf{u}_l} X_l)^{j_{kl}}
\end{aligned}$$

which is exactly $f(\mathcal{K}^{\mathbf{u}}\mathcal{X}^{\mathbf{p}})$, as expected. □

3.5 Ribbon structure

We established in Lemma 2.19 that a unimodular quasitriangular Hopf algebra with two-sided integral and the Drinfeld element u can be endowed with a ribbon structure if we can find a pivotal element $g \in G(H)$ such that $g^2 = \mathbf{1}$.

Lemma 3.20. Let K_a and X_k be generators of $H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$ and R be an R-matrix. Denote u the Drinfeld element as in Definition 2.14. Then

$$[u, K_a] = \{u, X_k\} = 0.$$

Proof. By Proposition 2.15

$$S^2(K_a) = K_a = uK_a u^{-1},$$

and

$$S^2(X_k) = -X_k = uX_k u^{-1},$$

the conclusion is immediate. □

Hence we have the following result.

Theorem 3.21. Assume $H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$ admits a two-sided integral, a two-sided cointegral, an R-matrix R , and a pivotal element g . Then $g^2 = \mathbf{1}$ and $v = g^{-1}u = gu$ is a ribbon element.

Proof. This follows from Lemmas 2.18, 2.19 and 3.20. □

3.6 Examples revisited

We revisit examples of the Nenciu construction, and show they all admit unimodular, strongly non-factorizable ribbon structures.

Notation. In this section the expression R_{α} will recur in the same form given by

$$R_{\alpha} := \exp \left(\sum_{l=1}^{t_2} \alpha_l (Z_l^+ \otimes LZ_l^- - Z_l^- \otimes LZ_l^+) \right).$$

Here we stick to the notation for the nilpotent generators from the previous examples emphasizing the types. In particular, let Z_l^+ and Z_l^- be nilpotent generators in $H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$ of opposite type, with neighbouring indices, that is $Z_l^+ = X_k$ and $Z_l^- = X_{k+1}$ for some k . Let moreover $\{Z_l^{\pm}, X_k\} = 0$ for all X_k , including themselves. We changed the notation in the expression above and we denoted $\alpha_{k,k+1} = -\alpha_{k+1,k} := \alpha_l$. We collect α_l in the tuple α .

We again start the list of examples with SF_{2n} .

Proposition 3.22. The algebra SF_{2n} of Definition 3.9 is

1. unimodular, with a two-sided cointegral

$$\Lambda := (\mathbf{1} + L) \prod_{l=1}^n Z_l^+ \prod_{l=1}^n Z_l^-,$$

and a two-sided integral defined on the monomial basis by

$$\lambda(L^v \mathbf{X}^{\mathbf{r}}) := \begin{cases} 1 & \text{if } v = 0, \mathbf{r} = (1, 1, \dots, 1) \\ 0 & \text{otherwise,} \end{cases}$$

2. quasitriangular, with the R-matrix defined for the grouplike generators as

$$R_{\mathbf{z}} := \frac{1}{2} \sum_{v,w \in \mathbb{Z}_2} (-1)^{-vw} L^w \otimes L^{vz} = \frac{1}{2} (\mathbf{1} \otimes \mathbf{1} + \mathbf{1} \otimes L + L \otimes \mathbf{1} - L \otimes L),$$

where $z = 1$. for some $\alpha = (\alpha_1, \dots, \alpha_{t_2}) \in \mathbb{C}^{2t_2}$, so that

$$R = \frac{1}{2} (\mathbf{1} \otimes \mathbf{1} + \mathbf{1} \otimes L + L \otimes \mathbf{1} - L \otimes L) \exp \left(\sum_{l=1}^n \alpha_l (Z_l^+ \otimes LZ_l^- - Z_l^- \otimes LZ_l^+) \right),$$

3. ribbon, with the ribbon element

$$v := \exp \left(-2 \sum_{k=l}^n \alpha_l Z_l^+ Z_l^- \right),$$

corresponding to the pivotal element $g = L$, which in this example is the only such element available.

Proof. See [PO99], Propositions 8, 2 and 11, respectively. \square

Remark 3.23. Note that Theorem 2.9 allows multiple quasitriangular structures for $H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$. Indeed, it is known from [GR17] that, for instance, SF_2 admits a quasitriangular structure where $R := R_z \tilde{R}_\alpha$, with R_z as in Proposition 3.22, but

$$\tilde{R}_\alpha = \exp(\alpha^+ Z^+ \otimes LZ^+ + \alpha^- Z^- \otimes LZ^-)$$

where $\alpha = (\alpha^+, \alpha^-) \in \mathbb{C}^2$. It admits only a trivial ribbon element $v = \mathbf{1}$.

We now move on to more complicated examples, that did not appear in [Nen04], starting with one that is in fact *triangular*.

Proposition 3.24. The algebra N_1 of Example 3.10 is

1. unimodular, with a two-sided cointegral

$$\Lambda := \left(\sum_{a,b=0}^3 K_1^a K_2^b \right) \prod_{k=1}^{t_1} X_k \prod_{l=1}^{t_2} Z_l^+ \prod_{l=1}^{t_2} Z_l^-,$$

and a two-sided integral defined on the monomial basis by

$$\lambda(\mathbf{K}^{\mathbf{v}} \mathbf{X}^{\mathbf{r}}) := \begin{cases} 1 & \text{if } \mathbf{v} = (0, 0), \mathbf{r} = (1, 1, \dots, 1) \\ 0 & \text{otherwise.} \end{cases}$$

2. triangular, with R-matrix defined for the grouplike elements as

$$R_{\mathbf{z}} := \frac{1}{16} \sum_{\mathbf{v}, \mathbf{w} \in \mathbb{Z}_4^2} i^{-\mathbf{v}\mathbf{w}^T} \mathbf{K}^{\mathbf{w}} \otimes \mathbf{K}^{\mathbf{vz}},$$

where $\mathbf{K} = (K_1, K_2)$, $\boldsymbol{\xi} = (i, i)$, and $\mathbf{z} = \begin{pmatrix} 2 & 3 \\ 1 & 0 \end{pmatrix}$.

3. and ribbon, with ribbon element $v = \mathbf{1}$, corresponding to the unique pivotal element $g = K_1^2$,

Remark 3.25. The Hopf algebra of Example 3.10 admits no non-triangular structures. The behaviour of Z_l^\pm generators suggests that a non-triangular R-matrix could be built of terms of the shape $Z_l^\pm \otimes K_1^{\pm 2} K_2^{\pm 1} Z_l^\mp$, as the two generators are of opposite type. However, any Hopf algebra map f_R , sending $Z_l^\pm \mapsto K_1^{-2} K_2^{-1} (Z_l^\mp)$, satisfies all constraints apart from (B1), which is equivalent to $(K_1^2 K_2)^2 = \mathbf{1}$, which is false in this case. The element X_k similarly fails (B1) for any such Hopf algebra map.

With this observation, we can build a non-triangular example, at the cost of introducing a K_3 , another grouplike element of order 4. We can choose $L = K_3^2$, which implies that the corresponding Z_l^\pm will at most anti-commute with all other nilpotent generators, according to the notation we established.

Proposition 3.26. The algebra N_2 of Example 3.11 is

1. unimodular, with a two-sided cointegral

$$\Lambda := \left(\sum_{a,b,c=0}^3 K_1^a K_2^b K_3^c \right) \prod_{k=1}^{t_1} X_k^+ \prod_{k=1}^{t_1} X_k^- \prod_{l=1}^{t_2} Z_l^+ \prod_{l=1}^{t_2} Z_l^-,$$

and a two-sided integral defined on the monomial basis as

$$\lambda(\mathbf{K}^{\mathbf{v}} \mathbf{X}^{\mathbf{r}}) := \begin{cases} 1 & \text{if } \mathbf{v} = (0, 0, 0), \mathbf{r} = (1, 1, \dots, 1) \\ 0 & \text{otherwise,} \end{cases}$$

2. quasitriangular, with the R-matrix defined for the grouplike generators as

$$R_{\mathbf{z}} := \frac{1}{64} \sum_{\mathbf{v}, \mathbf{w} \in \mathbb{Z}_4^3} i^{-\mathbf{v}\mathbf{w}^T} \mathbf{K}^{\mathbf{w}} \otimes \mathbf{K}^{\mathbf{vz}},$$

where $\mathbf{K} = (K_1, K_2, K_3)$ and $\mathbf{z} = \begin{pmatrix} 0 & 3 & 2 \\ 1 & 0 & 0 \\ 2 & 0 & 2 \end{pmatrix}$.

The part containing skew-primitive generators is

$$R_\alpha := \exp \left(\sum_{l=1}^{t_2} \alpha_l (Z_l^+ \otimes LZ_l^- - Z_l^- \otimes LZ_l^+) \right)$$

for some $\alpha = (\alpha_1, \dots, \alpha_{t_2}) \in \mathbb{C}^{2t_2}$, so that

$$R_z R_\alpha := \frac{1}{64} \sum_{\mathbf{v}, \mathbf{w} \in \mathbb{Z}_4^3} i^{-\mathbf{v}\mathbf{w}^T} \mathbf{K}^{\mathbf{w}} \otimes \mathbf{K}^{\mathbf{vz}} \exp \left(\sum_{l=1}^{t_2} \alpha_l (Z_l^+ \otimes LZ_l^- - Z_l^- \otimes LZ_l^+) \right),$$

3. ribbon, with the ribbon element

$$v := \exp \left(-2 \sum_{l=0}^{t_2} \alpha_l Z_l^+ Z_l^- \right).$$

corresponding to the pivotal element $g = L = K_3^2$, which in this example is the only such element available.

We can introduce more nilpotent generators while retaining the same quasitriangular structure.

Proposition 3.27. The algebra N_3 of Example 3.12 is

1. unimodular, with a two-sided cointegral

$$\Lambda := \left(\sum_{a,b,c=0}^3 K_1^a K_2^b K_3^c \right) \prod_{k=1}^{t_1} X_k^+ \prod_{k=1}^{t_1} X_k^- \prod_{k=1}^{t_1} Y_k^+ \prod_{k=1}^{t_1} Y_k^- \prod_{l=1}^{t_2} Z_l^+ \prod_{l=1}^{t_2} Z_l^-,$$

and a two-sided integral defined on the monomial basis by

$$\lambda(\mathbf{K}^{\mathbf{v}} \mathbf{X}^{\mathbf{r}}) := \begin{cases} 1 & \text{if } \mathbf{v} = (0, 0, 0), \mathbf{r} = (1, 1, \dots, 1) \\ 0 & \text{otherwise.} \end{cases}$$

2. quasitriangular, with the R-matrix defined for the grouplike generators as

$$R_{\mathbf{z}} := \frac{1}{64} \sum_{\mathbf{v}, \mathbf{w} \in \mathbb{Z}_4^3} i^{-\mathbf{v}\mathbf{w}^T} \mathbf{K}^{\mathbf{w}} \otimes \mathbf{K}^{\mathbf{vz}},$$

where $\mathbf{K} = (K_1, K_2, K_3)$ and $\mathbf{z} = \begin{pmatrix} 0 & 3 & 2 \\ 1 & 0 & 0 \\ 2 & 0 & 2 \end{pmatrix}$.

The skew-primitive part is

$$R_{\boldsymbol{\alpha}} := \exp \left(\sum_{l=1}^{t_2} \alpha_l (Z_l^+ \otimes LZ_l^- - Z_l^- \otimes LZ_l^+) \right)$$

for some $\boldsymbol{\alpha} = (\alpha_1, \dots, \alpha_{t_2}) \in \mathbb{C}^{2t_2}$, so that

$$R_{\mathbf{z}} R_{\boldsymbol{\alpha}} := \frac{1}{64} \sum_{\mathbf{v}, \mathbf{w} \in \mathbb{Z}_4^3} i^{-\mathbf{v}\mathbf{w}^T} \mathbf{K}^{\mathbf{w}} \otimes \mathbf{K}^{\mathbf{vz}} \exp \left(\sum_{l=1}^{t_2} \alpha_l (Z_l^+ \otimes LZ_l^- - Z_l^- \otimes LZ_l^+) \right),$$

3. ribbon, with ribbon element

$$v := \exp \left(-2 \sum_{k=1}^{t_2} \alpha_k Z_k^+ Z_k^- \right),$$

corresponding to the unique pivotal element $g = L = K_3^2$.

Proof of Propositions 3.24, 3.26, 3.27. The propositions can be verified by a direct check of unimodularity, quasitriangularity and ribbon axioms, or equivalently using Theorem 3.15 for the cointegral, 2.9 for the R-matrix and 3.21 for the ribbon element. \square

Proposition 3.28. The Hopf algebras of examples 3.10, 3.11, 3.12 admit only strongly non-factorizable quasitriangular structures.

Proof. The theorem is verified by essentially checking all choices of bialgebra maps $f : H^* \rightarrow H^{cop}$. Indeed, since the relations are diagonal, the grouplike generators K_b have to be mapped to grouplike generators K_a and nilpotent generators $\mathcal{X}_j, \mathcal{X}_j^\pm, \mathcal{Y}_j^\pm, \mathcal{Z}_l^\pm$ to nilpotent generators $X_k, X_k^\pm, Y_k^\pm, Z_m^\pm$. We highlight the key points.

First, we explain why the examples do not admit any quasitriangular structures such that generators X_k^\pm , as well as Y_k^\pm in Example 3.12 appear in $R_{\boldsymbol{\alpha}}$. One can simply check all possible assignment and find that the only way to satisfy (C1) is to map

$$f(\mathcal{X}_k^\pm) = (K_1 K_2)^{\mp 1} X_k^\pm, \text{ resp. } f(\mathcal{Y}_k^\pm) = (K_1^2 K_2)^{\mp 1} Y_k^\pm,$$

but it can be verified it fails (C4) for all K_a , $a = 1, \dots, s$. The mapping

$$f(\mathcal{X}_k^\pm) = (K_1 K_2)^{\pm 1} X_k^\mp, \text{ resp. } f(\mathcal{Y}_k^\pm) = (K_1^2 K_2)^{\pm 1} Y_k^\mp,$$

analogous to the terms involving Z_l^\pm , satisfies (B1) only if $(K_1 K_2) = (K_1 K_2)^{-1}$, which is not the case for our choice of relations. If the image of \mathcal{X}_k^\pm or \mathcal{Y}_k^\pm involves Z_l^\pm , (B1) is not satisfied, which can be checked directly.

Finally, from the definitions of the integral and the R-matrix, we observe that all nilpotent generators except Z_l^\pm (or all in Example 3.10) are absent from the respective R-matrices. There is no relation available that introduces them, as the commutations are necessarily diagonal, thus they are absent from the monodromy matrix as well. Since the integral is non-zero only on the linear combinations of monomials containing the top element $T = \mathbf{X}^{(1, \dots, 1)}$, and since T contains all the nilpotent generators, we retrieve the result. \square

We can also deduce the following about the ribbon structure.

Corollary 3.29. The Hopf algebras of examples 3.10, 3.11, 3.12 admit only anomalous ribbon structures.

Proof. In the proof of Proposition 3.28, we showed that there is a subset of nilpotent generators that cannot appear in the R-matrix, R . It is a straightforward consequence of the diagonality of all relations that these generators are also necessarily absent from the corresponding Drinfeld element $u = S(R'')R'$ and so is the ribbon element v , as in all cases the pivotal element $g \in \mathbb{C}[\mathbf{K}]$ and $v := g^{-1}u$. Hence by the same reasoning $\lambda(v) = 0$ and so from Definition 2.20 only anomalous ribbon structures appear. \square

We will later need formulas for the monodromy matrices corresponding to the class of R-matrices $R_{\mathbf{z}}R_{\alpha}$ that appeared extensively in the examples.

Proposition 3.30. Let $H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$ be a Hopf algebra containing nilpotent generators Z_l^{\pm} , for $l = 1, \dots, t_2$. Moreover let Z_l^+ and Z_l^- be of opposite type and $\{Z_{i_1}^{\pm}, Z_{i_2}^{\pm}\} = \{Z_{i_1}^{\pm}, Z_{i_2}^{\mp}\} = 0$. Denote $L_l := \mathbf{K}^{\mathbf{u}_l}$. Let $R := R_{\mathbf{z}}R_{\alpha}$ be an R-matrix of the form as in Theorem 3.19. Then the monodromy matrix is given by

$$M := R_{21}R = \exp\left(2 \sum_{l=1}^{t_2} \alpha_l (Z_l^+ \otimes L_l Z_l^- - Z_l^- \otimes L_l Z_l^+)\right).$$

Proof. We first consider the behaviour of the terms $L_l Z_l^{\mp} \otimes Z_l^{\pm}$, that appear in $R_{\alpha, 21}$, when they commute past $R_{\mathbf{z}}$. We have

$$\begin{aligned} (L Z_l^{\mp} \otimes Z_l^{\pm}) R_{\mathbf{z}} &= (L_l Z_l^{\mp} \otimes Z_l^{\pm}) \frac{1}{\prod_{a=1}^s m_a} \sum_{\mathbf{v}, \mathbf{w} \in \mathbb{Z}_{\mathbf{m}}} \xi^{-\mathbf{v} \cdot \mathbf{w}} \mathbf{K}^{\mathbf{w}} \otimes \mathbf{K}^{\mathbf{vz}} \\ &= \frac{1}{\prod_{a=1}^s m_a} \sum_{\mathbf{v}, \mathbf{w} \in \mathbb{Z}_{\mathbf{m}}} \xi^{-\mathbf{v} \cdot \mathbf{w} + \mathbf{w} \cdot \mathbf{d}_l - (\mathbf{vz}) \cdot \mathbf{d}_l} \mathbf{K}^{\mathbf{w} + \mathbf{u}_l} \otimes \mathbf{K}^{\mathbf{vz} - \mathbf{u}_l} (Z_l^{\mp} \otimes L_l Z_l^{\pm}), \end{aligned}$$

where we commuted Z_l^{\pm} past $\mathbf{K}^{\mathbf{w}}$ and $\mathbf{K}^{\mathbf{vz}}$, absorbing the L_l in $\mathbf{K}^{\mathbf{w}}$ and factoring it out of $\mathbf{K}^{\mathbf{vz}}$ in every element of the sum. Note that this is allowed because $L^2 = \mathbf{1}$. Now using (C1), (C2) of Theorem 3.19, we note that $(\mathbf{vz}) \cdot \mathbf{d}_l = \mathbf{v} \cdot \mathbf{u}_l$. Now we introduce the shifts in the sum $\mathbf{w} \mapsto \mathbf{w} - \mathbf{u}_l$ and $\mathbf{vz} \mapsto \mathbf{vz} + \mathbf{u}_l$, that is $\mathbf{v} \mapsto \mathbf{v} + \mathbf{d}_l$. Now the coefficient in the sum becomes

$$\xi^{-(\mathbf{v} + \mathbf{d}_l) \cdot (\mathbf{w} - \mathbf{u}_l) + (\mathbf{w} - \mathbf{u}_l) \cdot \mathbf{d}_l - (\mathbf{vz} + \mathbf{u}_l) \cdot \mathbf{d}_l}$$

which again by (C1), (C2) of Theorem 3.19 simplifies to

$$\xi^{-\mathbf{v} \cdot \mathbf{w} - \mathbf{d}_l \cdot \mathbf{u}_l} = \xi^{-\mathbf{v} \cdot \mathbf{w}} (-1),$$

since by Definition 3.1, $\xi^{\mathbf{d}_l \cdot \mathbf{u}_l} = -1$. Thus, the commutation stands

$$\frac{1}{\prod_{a=1}^s m_a} \sum_{\mathbf{v}, \mathbf{w} \in \mathbb{Z}_{\mathbf{m}}} \xi^{-\mathbf{v} \cdot \mathbf{w}} \mathbf{K}^{\mathbf{w}} \otimes \mathbf{K}^{\mathbf{vz}} (-Z_l^{\mp} \otimes L_l Z_l^{\pm}).$$

Now we can consider the exponential term in full to notice that

$$\exp\left(\sum_{l=1}^{t_2} \alpha_l (L_l Z_l^- \otimes Z_l^+ - L_l Z_l^+ \otimes Z_l^-)\right) R_{\mathbf{z}} = R_{\mathbf{z}} \exp\left(\sum_{l=1}^{t_2} \alpha_l (Z_l^+ \otimes L_l Z_l^- - Z_l^- \otimes L_l Z_l^+)\right) = R_{\mathbf{z}} R_{\alpha}.$$

So we retrieve

$$R_{\mathbf{z}, 21} R_{\alpha, 21} R_{\mathbf{z}} R_{\alpha} = R_{\mathbf{z}, 21} R_{\mathbf{z}} R_{\alpha} R_{\alpha} = \exp\left(2 \sum_{l=1}^{t_2} \alpha_l (Z_l^+ \otimes L_l Z_l^- - Z_l^- \otimes L_l Z_l^+)\right).$$

\square

4 Non-factorizable biproducts with $u_q \mathfrak{sl}_2$

In this section we construct a family of Hopf algebras, which augment $u_q \mathfrak{sl}_2$, for q a primitive r -th root of unity and $r \equiv 0 \pmod{4}$ by a Hopf algebra of Nenciu type $H = H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$ so that both pieces interact. The main property of the construction is that the result is ribbon, with the new R-matrix and ribbon element constructed from those carried by U and H . The idea of extending $u_q \mathfrak{sl}_2$ using nilpotent generators is inspired by the work of Majid, see for instance [Maj00] Example 10.2.13, but the strategy used to find the non-factorizable examples did not appear in the literature to our best knowledge.

4.1 The small quantum group $u_q \mathfrak{sl}_2$

We use this section to set the conventions for the small quantum group. We follow mostly the conventions of [CP95][Section 9.1] and [EGNO16][Section 5.7], used also in [BdR23]. Let $r' = r/2$ and $r'' = r/4$ throughout.

Definition 4.1. The Hopf algebra $u_q\mathfrak{sl}_2$ where q , is a primitive r -th root of unity, $r \equiv 0 \pmod{4}$, is the algebra generated by elements K, E , and F , satisfying the relations

$$K^{r'} = \mathbf{1}, \quad E^{r'} = F^{r'} = 0$$

$$KEK^{-1} = q^2E, \quad KFK^{-1} = q^{-2}F, \quad [E, F] = \frac{K - K^{-1}}{q - q^{-1}}.$$

The Hopf structure is given by

$$\epsilon(K) = 1, \quad \epsilon(E) = \epsilon(F) = 0,$$

$$\Delta(K) = K \otimes K, \quad \Delta(E) = \mathbf{1} \otimes E + E \otimes K, \quad \Delta(F) = F \otimes \mathbf{1} + K^{-1} \otimes F$$

and

$$S(K) = K^{-1} = K^{r'-1}, \quad S(E) = -EK^{-1}, \quad S(F) = -KF.$$

Recall also the notation for quantum integers, for each $k \in \mathbb{Z}_{r'}$:

$$\{k\} := q^k - q^{-k}, \quad [k] := \frac{\{k\}}{\{1\}}, \quad [k]! := [k][k-1] \dots [1].$$

Proposition 4.2. The Hopf algebra $u_q\mathfrak{sl}_2$ is

1. unimodular with a two-sided cointegral

$$\Lambda := \frac{\{1\}^{r'-1}}{\sqrt{r''}[r'-1]!} \sum_{a=0}^{r'-1} E^{r'-1} F^{r'-1} K^a,$$

and a left integral

$$\lambda(E^b F^c K^a) := \frac{\sqrt{r''}[r'-1]!}{\{1\}^{r'-1}} \delta_{a,r'-1} \delta_{b,r'-1} \delta_{c,r'-1},$$

2. quasitriangular with $R := D\Theta$, where

$$D := \frac{1}{r'} \sum_{b,c=0}^{r'-1} q^{-2bc} K^b \otimes K^c,$$

and

$$\Theta := \sum_{a=0}^{r'-1} \frac{\{1\}^a}{[a]!} q^{\frac{a(a-1)}{2}} E^a \otimes F^a.$$

3. ribbon, with the ribbon element for $r \equiv 4 \pmod{8}$

$$v = \frac{i^{\frac{r''-1}{2}}}{\sqrt{r''}} \sum_{a=0}^{r'-1} \sum_{b=0}^{r''-1} \frac{\{-1\}^a}{[a]!} q^{-\frac{(a+3)a}{2} + \frac{(r''+1)^3}{2}(2b-1)^2} E^a F^a K^{-a-2b},$$

and for $r \equiv 0 \pmod{8}$

$$v = \frac{1-i}{\sqrt{r'}} \sum_{a=0}^{r'-1} \sum_{b=0}^{r''-1} \frac{\{-1\}^a}{[a]!} q^{-\frac{(a+3)a}{2} + 2b^2} E^a F^a K^{-a-2b-1}.$$

corresponding to the pivotal element $g = K$.

Remark 4.3. To make a connection with the notation of Theorem 2.9 for D , note that $q^{2r'} = 1$, but $K^{r'} = \mathbf{1}$. Thus, we can write

$$D = \frac{1}{r'} \sum_{v,w \in \mathbb{Z}_{r'}} (q^2)^{-wv} K^w \otimes K^{zv}$$

which corresponds to a $R_{\mathbf{z}}$ for $\xi = q^2$ and $\mathbf{z} = 1$.

4.2 The semi-direct biproduct structure

Towards the goal of constructing non-factorizable extensions of $u_q\mathfrak{sl}_2$ we introduce versions of *semi-direct* (or *smash*) product and coproduct. While these constructions are well-known in the study of Hopf algebras, see for instance [Mol77], [AS06], we use slightly non-standard conventions to achieve the properties studied in the examples below.

Notation. The Sweedler notation will often appear in this section when considering a semi-direct biproduct $U \times H$ of two Hopf algebras U and H . Then it refers to the corresponding coproducts Δ_U and Δ_H of the respective tensor components, rather than the new coproduct Δ of $U \times H$.

Definition 4.4. Let U, H be a Hopf algebras. Let

$$\triangleleft : H \otimes U \rightarrow H$$

be a right algebra action of U on H (see for instance [Kas12], Chapter III.5). We define the *right semi-direct product* of U and H to be the algebra $U \otimes H$ with the unit $\mathbf{1}_U \otimes \mathbf{1}_H$ and the product μ given by

$$\mu(g \otimes h, g' \otimes h') = (g \otimes h)(g' \otimes h') := gg'_{(1)} \otimes (h \triangleleft g'_{(2)})h'.$$

Let also

$$\delta : H \rightarrow H \otimes U$$

be a right coaction of U on H (see for instance [Kas12] Chapter III.6). We adopt the Sweedler notation

$$\delta : h \mapsto h^{[0]} \otimes h^{[1]}.$$

We define the *right semi-direct coproduct* of U and H as the coalgebra $U \otimes H$ with the counit $\epsilon_U \otimes \epsilon_H$, and the coproduct

$$\Delta(g \otimes h) := (g_{(1)} \otimes h^{[0]}_{(1)}) \otimes (g_{(2)} h^{[1]}_{(1)} \otimes h_{(2)}).$$

Let $S : U \otimes H \rightarrow U \otimes H$ be the invertible map given by

$$S(g \otimes h) \mapsto S_U(gh^{[1]}) \otimes S_H(h^{[0]}).$$

If $(U \otimes H, \mu, \mathbf{1}_U \otimes \mathbf{1}_H, \Delta, \epsilon_U \otimes \epsilon_H, S)$ is a Hopf algebra then we will call it a *semi-direct biproduct* of U and H and denote $U \rtimes H$ see for instance [Maj00, Section 6].

Remark 4.5. We emphasize that all tensor products between U and H in Definition 4.4 are taken in the category $\mathbf{Vect}_{\mathbb{C}}$, which carries the trivial braiding. In particular, this is a different setting than the *Radford biproduct* which is taken in a braided category of Yetter-Drinfeld modules over the acting algebra, see [AS06].

Remark 4.6. We do not attempt to determine the conditions under which the semi-direct product and semi-direct coproduct in Definition 4.4 give rise to the semi-direct biproduct. We merely present in the sequel a construction where the requirements are indeed satisfied.

4.3 Non-factorizable biproducts with $u_q \mathfrak{sl}_2$

In this section we build biproducts of $u_q \mathfrak{sl}_2$ and $H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$, and we show they can have the properties of unimodularity, quasitriangularity and ribbonness, and can be (strongly) non-factorizable. For clarity of discussion and the applications we keep in mind, we restrict the order of q, r to be a multiple of 8.

Definition 4.7. Let $U = u_q \mathfrak{sl}_2$ and $H = H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$, where q , is a primitive r -th root of unity, $r \equiv 0 \pmod{8}$ and $r' = r/2$, $r'' = r/4$. We define $U \rtimes H$, where

- the right action $\triangleleft : H(\mathbf{m}, t, \mathbf{d}, \mathbf{u}) \otimes u_q \mathfrak{sl}_2 \rightarrow H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$ for $K_a \in H$, $a = 1, \dots, s$ by

$$K_a \triangleleft K = K_a, \quad K_a \triangleleft E = K_a \triangleleft F = 0$$

and for $X_k \in H$ by

$$X_k \triangleleft K = -X_k, \quad X_k \triangleleft E = X_k \triangleleft F = 0,$$

- the right coaction $\delta : H(\mathbf{m}, t, \mathbf{d}, \mathbf{u}) \rightarrow H(\mathbf{m}, t, \mathbf{d}, \mathbf{u}) \otimes u_q \mathfrak{sl}_2$ for $K_a \in H$ by

$$K_a \mapsto K_a \otimes \mathbf{1}_U,$$

for $X_k \in H$ by

$$X_k \mapsto X_k \otimes K^{r''},$$

and on an arbitrary monomial $\mathbf{K}^{\mathbf{v}} \mathbf{X}^{\mathbf{r}} \in H$ for $\mathbf{v} \in \mathbb{Z}_{\mathbf{m}}$ and $\mathbf{r} \in \mathbb{Z}_2^t$ by

$$\mathbf{K}^{\mathbf{v}} \mathbf{X}^{\mathbf{r}} \mapsto \mathbf{K}^{\mathbf{v}} \mathbf{X}^{\mathbf{r}} \otimes K^{|\mathbf{r}|r''}.$$

Theorem 4.8. Let $u_q \mathfrak{sl}_2 \rtimes H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$ be as in Definition 4.7. Then it is a Hopf algebra with the following semi-direct biproduct structure:

- the new unit $\mathbf{1} = \mathbf{1}_U \otimes \mathbf{1}_H$ and counit $\epsilon := \epsilon_U \otimes \epsilon_H$,
- the algebra structure where we suppress the tensor product structure

$$K := K \otimes \mathbf{1}_H \quad E := E \otimes \mathbf{1}_H \quad F := F \otimes \mathbf{1}_H$$

$$K_a := \mathbf{1}_U \otimes K_a \quad X_k := \mathbf{1}_U \otimes X_k,$$

so that for any $a, b, c, \in \mathbb{Z}_{r'}$, $\mathbf{v} \in \mathbb{Z}_{\mathbf{m}}$ and $\mathbf{r} \in \mathbb{Z}_2^t$

$$E^a F^b K^c \mathbf{X}^{\mathbf{r}} \mathbf{K}^{\mathbf{v}} := E^a F^b K^c \otimes \mathbf{X}^{\mathbf{r}} \mathbf{K}^{\mathbf{v}}$$

and find the new algebra relations between the the elements of U and H to be

$$[K, K_a] = [E, K_a] = [F, K_a] = 0, \quad \{K, X_k\} = \{E, X_k\} = [F, X_k] = 0.$$

- new Hopf structure for X_k give by

$$\Delta(X_k) = \mathbf{1} \otimes X_k + X_k \otimes K^{r''} \mathbf{K}^{\mathbf{u}_k} \quad S(X_k) = -X_k K^{r''} \mathbf{K}^{-\mathbf{u}_k},$$

while for the remaining generators E, F, K and K_a in this notation the coproduct Δ and antipode S are unchanged.

Proof. See Section A.1 in Appendix A. □

Proposition 4.9. The Hopf algebra $u_q \mathfrak{sl}_2 \rtimes H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$ has a monomial basis

$$\{E^e F^f K^k \mathbf{K}^{\mathbf{w}} \mathbf{X}^{\mathbf{r}} | e, f, k = 1, \dots, r', \mathbf{w} \in \mathbb{Z}_{\mathbf{m}}, \mathbf{r} \in \mathbb{Z}_2^t\}.$$

Proof. This follows directly from the fact that both U and H have monomial bases, by the Poincaré-Birkhoff-Witt theorem (see for instance [Kas12]) and Proposition 3.6, respectively. Indeed, all newly introduced relations between the two bases are diagonal by Theorem 4.7. Hence by the same argument as in the proof of Proposition 3.6, we can establish the existence of the monomial basis. □

Remark 4.10. Let U, H and $U \rtimes H$ be as in Definition 4.7, in particular let the pivotal element of H be g_H . Then we claim the following maps are Hopf algebra maps, which do *not* give a short exact sequence:

$$U \hookrightarrow U \rtimes H \rightarrow H.$$

The map $\iota : U \hookrightarrow U \rtimes H$ is given by $u \mapsto u \otimes \mathbf{1}_H$ for $u \in U$, and has a left inverse $j : U \rtimes H \rightarrow U$ given by

$$j(K) = K, \quad j(E) = E, \quad j(F) = F, \quad j(K_a) = \mathbf{1}_U, \quad j(X_k) = 0.$$

The map $p : U \rtimes H \rightarrow H$, is given by

$$p(K) = g_H, \quad p(E) = p(F) = 0, \quad p(K_a) = K_a, \quad p(X_k) = X_k.$$

Remark 4.11. The map p in Remark 4.10 has a right inverse as an algebra map, given by the map

$$q : H \hookrightarrow H \rtimes U, \quad q(h) = \mathbf{1}_U \otimes h,$$

but it is *not* a Hopf algebra map. For instance $S(X_k) = K^{r''} \mathbf{K}^{\mathbf{u}_k} X_k$, so the image of H under q is not closed under taking antipodes (as it is not under taking coproducts). Nevertheless, it makes sense to speak, for instance, of elements $u, h \in U \rtimes H$ as belonging to $u \in U$ and $h \in H$, and we will often slightly abuse the notation in this way.

Remark 4.12. Let U, H and $U \rtimes H$ be as in Theorem 4.8. Then in light of Remark 4.10 $U \rtimes H$ is not isomorphic to $U \otimes H$ as Hopf algebra in the usual sense. In particular, $U \otimes H$ has no non-trivial commutation relations between the elements belonging to U and H . Similarly, $U \rtimes H$ is not isomorphic to $U \oplus H$, that is the short exact sequence is not fully split, as per Remark 4.11.

Now we state the main theorem of the section.

Theorem 4.13. Let $U = u_q \mathfrak{sl}_2$ where q is a primitive r -th root of unity, $r \equiv 0 \pmod{8}$ as in Definition 4.1 carrying the integral λ_U , the cointegral Λ_U , the R -matrix $D\Theta$ and ribbon element v_U corresponding to a pivotal element g_U , as in Proposition 4.2. Let also $H = H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$ with

1. a two-sided cointegral

$$\Lambda_H := \prod_{a=1}^s \left(\sum_{b=0}^{m_a-1} K_a^b \right) \prod_{k=1}^t X_k,$$

2. a two-sided integral

$$\lambda_H(\mathbf{K}^{\mathbf{v}} \mathbf{X}^{\mathbf{r}}) := \begin{cases} 1 & \text{if } \mathbf{K}^{\mathbf{v}} \mathbf{X}^{\mathbf{r}} = T \\ 0 & \text{otherwise,} \end{cases}$$

3. an R -matrix $R_H := R_{\mathbf{z}} R_{\alpha}$ where

$$R_{\mathbf{z}} = \frac{1}{\prod_{a=1}^s m_a} \sum_{\mathbf{v}, \mathbf{w} \in \mathbb{Z}_{\mathbf{m}}} \xi^{-\mathbf{v} \cdot \mathbf{w}} \mathbf{K}^{\mathbf{w}} \otimes \mathbf{K}^{\mathbf{vz}}$$

$$R_{\alpha} = \exp \left(\sum_{l=0}^{t_2} \alpha_l (Z_l^+ \otimes L_l Z_l^- - Z_l^- \otimes L_l Z_l^+) \right),$$

where $L_l := \mathbf{K}^{\mathbf{u}_l}$

4. a ribbon element

$$v_H := \exp \left(-2 \sum_{k=1}^{t_2} \alpha_k Z_k^+ Z_k^- \right)$$

and the corresponding pivotal element $g = L$ and the Drinfeld element

$$u_H := L \exp \left(-2 \sum_{k=1}^{t_2} \alpha_k Z_k^+ Z_k^- \right).$$

Then, $U \rtimes H$, for the choices of action and coaction as in Definition 4.7, carries the following structures

1. a two-sided cointegral

$$\Lambda := \Lambda_U \Lambda_H = \frac{\{1\}^{r'-1}}{\sqrt{r''} [r'-1]!} \sum_{a=0}^{r'-1} E^{r'-1} F^{r'-1} K^a \prod_{a=1}^s \left(\sum_{b=0}^{m_a-1} K_a^b \right) \prod_{k=1}^t X_k,$$

2. a left integral $\lambda \in (U \rtimes H)^*$, $\lambda := \lambda_U \otimes \lambda_H$, equivalently defined on the monomial basis by

$$\lambda(E^b F^c K^a \mathbf{K}^{\mathbf{v}} \mathbf{X}^{\mathbf{r}}) := \begin{cases} \frac{\sqrt{r''} [r'-1]!}{\{1\}^{r'-1}} & \text{if } \mathbf{v} = (0, \dots, 0), \mathbf{r} = (1, \dots, 1), a = b = c = r' - 1 \\ 0 & \text{otherwise,} \end{cases}$$

3. an R -matrix $R := DR_{\mathbf{z}} \Theta \bar{R}_{\alpha}$, where

$$\bar{R}_{\alpha} := \exp \left(\sum_{l=0}^{t_2} \alpha_l (Z^+ \otimes \bar{L}_l Z_l^- - Z_l^- \otimes \bar{L}_l Z_l^+) \right)$$

for $\bar{L}_l = K^{r''} L_l$,

4. a ribbon element

$$v := v_U v_H =$$

$$\frac{1-i}{\sqrt{r'}} \sum_{a=0}^{r'-1} \sum_{b=0}^{r''-1} \frac{\{-1\}^a}{[a]!} q^{-\frac{(a+3)a}{2} + 2b^2} E^a F^a K^{-a-2b-1} \prod_{l=1}^{t_2} L_l \exp \left(-2 \sum_{k=1}^{t_2} \alpha_k Z_l^+ Z_l^- \right)$$

with the pivotal element $g = K$, and the Drinfeld element

$$u := u_U u_H =$$

$$\frac{1-i}{\sqrt{r'}} \sum_{a=0}^{r'-1} \sum_{b=0}^{r''-1} \frac{\{-1\}^a}{[a]!} q^{-\frac{(a+3)a}{2} + 2b^2} E^a F^a K^{-a-2b} \prod_{l=1}^{t_2} L_l \exp \left(-2 \sum_{k=1}^{t_2} \alpha_k Z_l^+ Z_l^- \right).$$

Proof. The proof is rather technical and lengthy, so we postpone it to Section A.2 in Appendix A. \square

Proposition 4.14. Let $u_q \mathfrak{sl}_2 \rtimes H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$ be as above. If $H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$ carries a strongly non-factorizable ribbon structure, then it induces a strongly non-factorizable ribbon structure on $u_q \mathfrak{sl}_2 \rtimes H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$.

Proof. By the reasoning of Proposition 3.28, if $H = H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$ is strongly non-factorizable, the top element $\mathbf{X}^{(1, \dots, 1)}$ is missing from the monodromy matrix $M_H \in H \otimes H$. Since all new relations are also diagonal, it will be missing from the new monodromy matrix $M \in (U \rtimes H) \otimes (U \rtimes H)$, so by the same argument the result follows. \square

4.4 Examples

As in the preceding section, we restrict to $r \equiv 0 \pmod{8}$, $r \geq 8$ so we retain a non-trivial relation of $[E, F]$ in the $u_q \mathfrak{sl}_2$. We start with an example related to the Hopf algebra SF_2 (the version involving SF_{2n} is analogous), but one which generically admits non-strongly non-factorizable quastriangular structures.

Example 4.15. Let $u_q \mathfrak{sl}_2 \rtimes H$, where $H = \text{SF}_2$ be a Hopf algebra generated by K, E, F with relations and morphisms of Definition 4.1, as well as L, Z^{\pm} with the following relations

$$\begin{aligned} L^2 &= \mathbf{1}, \quad (Z^{\pm})^2 = 0, \\ KZ^{\pm}K^{-1} &= q^{r'} Z^{\pm} = -Z^{\pm}, \quad LZ^{\pm}K_2^{-1} = -Z^{\pm}, \\ [L, K] &= [L, E] = [L, F] = 0 \\ EZ^{\pm} &= q^{r'} Z^{\pm} E = -Z^{\pm} E, \quad [Z^{\pm}, F] = 0. \end{aligned}$$

Let $\bar{L} := K^{r''} L$ for convenience. The Hopf structure is

$$\begin{aligned} \epsilon(L) &= 1, \quad \epsilon(Z^{\pm}) = 0, \\ \Delta(L) &= L \otimes L, \quad \Delta(Z^{\pm}) = \mathbf{1} \otimes Z^{\pm} + Z^{\pm} \otimes \bar{L}^{\pm 1}. \end{aligned}$$

and

$$S(L) = L^{-1}, \quad S(Z^{\pm}) = -Z^{\pm} \bar{L}^{\mp 1}.$$

The smallest instance with a non-trivial commutation relation $[E, F]$ occurs at $r = 8$. The dimension is then $4^3 \times 2^3 = 2^8 = 512$.

Proposition 4.16. The algebra $u_q \mathfrak{sl}_2 \rtimes \text{SF}_2$ is

1. unimodular with a two-sided cointegral

$$\Lambda := \frac{\{1\}^{r'-1}}{8\sqrt{r''[r'-1]!}} \sum_{a=0}^{r'-1} \sum_b^2 E^{r'-1} F^{r'-1} K^a L^b Z^+ Z^-,$$

and a left integral expressed on the monomial basis by

$$\lambda(E^e F^f K^a L^b (Z^+)^g (Z^-)^h) := \frac{\sqrt{r''[r'-1]!}}{\{1\}^{r'-1}} \delta_{a,r'-1} \delta_{b,0} \delta_{e,r'-1} \delta_{f,r'-1} \delta_{g,1} \delta_{h,1},$$

2. quasitriangular, with the R-matrix $R := R_{\mathbf{z}} D \Theta \bar{R}_{\alpha}$, where D, Θ were defined in Proposition 4.2, $R_{\mathbf{z}}$ in Proposition 3.22, and

$$\bar{R}_{\alpha} := \exp(\alpha(Z^+ \otimes \bar{L}Z^- - Z^- \otimes \bar{L}Z^+))$$

for $\bar{L} = K^{r''}L$,

3. ribbon, with the ribbon element

$$v = \frac{1-i}{\sqrt{r'}} \sum_{a=0}^{r'-1} \sum_{b=0}^{r''-1} \frac{\{-1\}^a}{[a]!} q^{-\frac{(a+3)a}{2}+2b^2} E^a F^a K^{-a-2b-1} L \exp(-2\alpha Z^+ Z^-)$$

corresponding to the pivotal element $g = K$.

The presence of R_{α} can lead to $\lambda(M') \otimes M'' \neq 0$ being non-zero. In particular, the resulting expression is dependent on α and turns out to be non-zero as long as $\alpha \neq 0$. Thus, $u_q \mathfrak{sl}_2 \rtimes \mathbf{SF}_2$ is not generically strongly non-factorizable. We now construct an example admitting only strongly non-factorizable quasitriangular structures. To do it, we introduce one more Nenciu-type algebra, supressing the Nenciu data for brevity.

Example 4.17. Let $N_4 := H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$ be a Nenciu type algebra generated by grouplike K_1, K_2 and nilpotent X^{\pm} generators, and the following relations

$$K_1^4 = K_2^4 = \mathbf{1}, \quad (X^{\pm})^2 = 0,$$

$$K_1 X^{\pm} K_2^{-1} = K_2 X^{\pm} K_2^{-1} = \pm i X^{\pm}.$$

Let $L := K_1 K_2$ for convenience. The Hopf structure is

$$\epsilon(K_1) = \epsilon(K_2) = 1, \quad \epsilon(X^{\pm}) = 0,$$

$$\Delta(K_1) = K_1 \otimes K_1, \quad \Delta(K_2) = K_2 \otimes K_2, \quad \Delta(X^{\pm}) = \mathbf{1} \otimes X^{\pm} + X^{\pm} \otimes L^{\pm 1}.$$

and

$$S(K_1) = K_1^{-1}, \quad S(K_2) = K_2^{-1}, \quad S(X^{\pm}) = -X^{\pm} L^{\mp 1}.$$

Proposition 4.18. The Hopf algebra N_4 of Example 4.17 is

1. unimodular, with two-sided cointegral

$$\Lambda := \left(\sum_{a,b=0}^3 K_1^a K_2^b \right) X^+ X^-$$

and a two-sided integral expressed on the monomial basis by

$$\lambda(\mathbf{K}^{\mathbf{v}} \mathbf{X}^{\mathbf{r}}) := \begin{cases} 1 & \text{if } \mathbf{v} = (0, 0), \mathbf{r} = (1, 1) \\ 0 & \text{otherwise.} \end{cases}$$

2. triangular, with the R-matrix defined for the grouplike elements as

$$R_{\mathbf{z}} := \frac{1}{16} \sum_{\mathbf{v}, \mathbf{w} \in \mathbb{Z}_4^2} i^{-\mathbf{v}\mathbf{w}^T} \mathbf{K}^{\mathbf{w}} \otimes \mathbf{K}^{\mathbf{v}\mathbf{z}},$$

where $\mathbf{K} = (K_1, K_2)$, $\boldsymbol{\xi} = (i, i)$, and $\mathbf{z} = \begin{pmatrix} 2 & 3 \\ 1 & 0 \end{pmatrix}$.

3. and trivially ribbon, with the ribbon element $\mathbf{1}$, corresponding to the unique pivotal element $g = K_1^2$.

Remark 4.19. Note $N_4 \subset N_1$ of Example 3.10. In analogy with N_1 it can be checked directly from Theorem 2.9 that the Hopf algebra N_4 admits only triangular quasitriangular structures.

Example 4.20. Let $u_q \mathfrak{sl}_2 \rtimes N_4$, be the Hopf algebra generated by K, E, F with relations and morphisms of Definition 4.1, as well as K_1, K_2, X^{\pm} with the following relations

$$K_1^4 = K_2^4 = \mathbf{1}, \quad (X^{\pm})^2 = 0,$$

$$K X^{\pm} K^{-1} = q^{r'} X^{\pm} = -X^{\pm}, \quad K_1 X^{\pm} K_2^{-1} = K_2 X^{\pm} K_3^{-1} = \pm i X^{\pm}.$$

$$[K_1, K_2] = [K_1, K] = [K_1, E] = [K_1, F] = [K_2, K] = [K_2, E] = [K_2, F] = 0$$

$$EX^\pm = q^{r'}X^\pm E = -X^\pm E, \quad [X^\pm, F] = 0.$$

Let $\bar{L} := K^{r''}K_1K_2$ for convenience. The Hopf structure is

$$\epsilon(K_1) = \epsilon(K_2) = 1, \quad \epsilon(X^\pm) = 0,$$

$$\Delta(K_1) = K_1 \otimes K_1, \quad \Delta(K_2) = K_2 \otimes K_2, \quad \Delta(X^\pm) = \mathbf{1} \otimes X^\pm + X^\pm \otimes \bar{L}^{\pm 1}.$$

and

$$S(K_1) = K_1^{-1}, \quad S(K_2) = K_2^{-1}, \quad S(X^\pm) = -X^\pm \bar{L}^{\mp 1}.$$

The smallest instance with a non-trivial commutation relation $[E, F]$ occurs at $r = 8$. The dimension is then $4^3 \times 2^4 \times 2^2 = 2^{12} = 4096$. Also, X^\pm share many features with X_a of Example 3.10. We have to modify the unimodular and ribbon structure accordingly.

Proposition 4.21. The algebra $u_q\mathfrak{sl}_2 \rtimes \mathbb{N}_4$ is

1. unimodular with a two-sided cointegral

$$\Lambda := \frac{\{1\}^{r'-1}}{\sqrt{r''}[r'-1]!} \sum_{a=0}^{r'-1} \sum_{b,c=0}^4 E^{r'-1} F^{r'-1} K^a K_1^b K_2^c X^+ X^-,$$

and a left integral expressed on the monomial basis by

$$\lambda(E^e F^f K^a K_1^b K_2^c (X^+)^g (X^-)^h) = \frac{\sqrt{r''}[r'-1]!}{\{1\}^{r'-1}} \delta_{a,r'-1} \delta_{b,0} \delta_{c,0} \delta_{e,r'-1} \delta_{f,r'-1} \delta_{g,1} \delta_{h,1},$$

2. quasitriangular, with the R-matrix $R := R_{\mathbf{z}} D \Theta$, where D, Θ were defined in Proposition 4.2, and

$$R_{\mathbf{z}} := \frac{1}{16} \sum_{\mathbf{v}, \mathbf{w} \in \mathbb{Z}_4^2} i^{-\mathbf{v}\mathbf{w}^T} (K_1, K_2)^{\mathbf{w}} \otimes (K_1, K_2)^{\mathbf{v}\mathbf{z}},$$

where $\mathbf{z} = \begin{pmatrix} 2 & 3 \\ 1 & 0 \end{pmatrix}$,

3. ribbon, with the ribbon element

$$v = \frac{1-i}{\sqrt{r'}} \sum_{a=0}^{r'-1} \sum_{b=0}^{r''-1} \frac{\{-1\}^a}{[a]!} q^{-\frac{(a+3)a}{2} + 2b^2} E^a F^a K^{-a-2b-1} K_2^2.$$

corresponding to the pivotal element $g = K$.

Proof. See Section A.3 in Appendix A. □

Proposition 4.22. The Hopf algebra $u_q\mathfrak{sl}_2 \rtimes H$ carries a strongly non-factorizable ribbon Hopf algebra, and does not admit any non-strongly non-factorizable quasitriangular structures.

Proof. As pointed out in Remark 4.19, applying the argument of Proposition 3.28 to \mathbb{N}_4 , we see can easily see it admits only strongly non-factorizable (in fact triangular) quasitriangular structures. Then by Proposition 4.14 we find that the quasitriangular structures induced on $u_q\mathfrak{sl}_2 \rtimes H$ are necessarily strongly non-factorizable.

Thus, we aim to use Theorem 2.9 to rule out quasitriangular structures not induced in the above way, that is to show there are no bialgebra maps $f : (u_q\mathfrak{sl}_2 \rtimes H)^* \rightarrow (u_q\mathfrak{sl}_2 \rtimes H)^{cop}$ that assign non-zero values to \mathcal{X}^\pm , the image of X^\pm under duality as given by Proposition 3.16 (in fact \mathbb{N}_4 is found to be self-dual so we drop the calligraphic notation in this proof), as well as ones containing X^\pm in their image. We first understand which map produces the $D\Theta$ part of the R-matrix, and as we already established that the H piece contributes only triangular structures, it suffices to rule out any potential cross-terms. It is enough to verify the claims on the relevant generators, as any bialgebra map needs to, in particular, respect the monomial basis and all Hopf morphisms.

1. It is a well-known fact (see for instance [Kas12]) that the map in Theorem 2.9 producing the R-matrix of $u_q\mathfrak{sl}_2$ in Proposition 2.9 is the Cartan involution $f : (u_q\mathfrak{sl}_2)^* \rightarrow (u_q\mathfrak{sl}_2)^{cop}$

$$f(E) = F, \quad f(F) = E, \quad f(K) = K^{-1}.$$

Crucially, $u_q\mathfrak{sl}_2$ is self-dual, as explained in [BdR23] and we do not introduce any new notation for the duals of E and F . The co-opposite algebra is given by the image of this map, the E and F generators are exchanged and K inverted. Since by Proposition 3.16 \mathbb{N}_4 is also self-dual, we readily verify $(U \rtimes H)^* \cong U \rtimes H$, as Hopf algebras.

2. Firstly, we need to exclude assignments of the form $X^\pm \mapsto bE + cF$, for some constants $b, c \in \mathbb{C}$. Since f needs to be an algebra map we seek $\Delta(f(X^\pm)^2) = 0$. We find

$$f(X^\pm)^2 = (bE + cF)^2 = b^2 E^2 + c^2 F^2 + bc(EF + FE) \neq 0,$$

unless $b = c = 0$. Hence, necessarily $f(X^\pm) = 0$.

3. Secondly, we need to exclude assignments of the form $E \mapsto F + aL^{\mp 1}X^{\pm}$ or $F \mapsto E + a'L^{\mp 1}X^{\pm}$ for $a, a' \in \mathbb{C}$. Consider

$$\Delta(E) = \mathbf{1} \otimes E + E \otimes K.$$

against

$$\Delta^{\text{cop}}(F) + \Delta^{\text{cop}}(L^{\mp 1}X^{\pm}) = \mathbf{1} \otimes (F + L^{\mp 1}X^{\pm}) + F \otimes K^{-1} + aL^{\mp 1}X^{\pm} \otimes L^{\mp 1}.$$

We need f to be a coalgebra map so

$$f(\Delta(E)) = f(\mathbf{1}) \otimes f(E) + f(k) \otimes f(E),$$

which requires,

$$F \otimes K^{-1} + aL^{\mp 1}X^{\pm} \otimes L^{\mp 1} = E \otimes f(K)$$

so $a = 0$ and $f(K) = K^{-1}$ or $a = 1$ and $f(K) = K^{-1} + L^{\pm 1}$. But in the latter case f is not a coalgebra map as the image of the grouplike K is the non-grouplike $K^{-1} + L^{\pm 1}$. Thus, $a = 1$, and $f(K) = K^{-1}$ and $f(E) = F$. The assignment of $f(F) = E$ is fixed by the same argument and we retrieve the assignments of the Cartan involution.

Thus we see the top monomial picked up by the integral of $U \times H$ is absent from the monodromy matrix, as in Proposition 3.28. We conclude the proof with the remark that we need not ask about the fate of generators K_a under the map f , as this does not affect the strong non-factorizability condition. \square

Corollary 4.23. The Hopf algebra of example 4.20 admits only anomalous ribbon structures.

Proof. Analogous to the proof of Corollary 3.29. \square

It is also possible to include the exponential piece of the R-matrix as in Example 3.11, while retaining strong non-factorizability.

Example 4.24. Let $u_q\mathfrak{sl}_2 \times \mathbb{N}_2$, be the Hopf algebra generated by K, E, F with relations and morphisms of def. 4.1, as well as $K_1, K_2, K_3, X^{\pm}, Z^{\pm}$ with the following relations, for $a = 1, 2, 3$

$$\begin{aligned} [K_a, K] &= [K_a, E] = [K_a, F] = 0, \quad KX^{\pm}K^{-1} = -X^{\pm}, \quad K_a^4 = \mathbf{1}, \quad K_aX^{\pm} = \pm iX^{\pm}K_a \\ KZ^{\pm}K^{-1} &= -Z^{\pm} \quad K_1Z^{\pm}K_2^{-1} = Z^{\pm}, \quad K_2Z^{\pm}K_3^{-1} = -Z^{\pm}, \quad K_3Z^{\pm}K_4^{-1} = \pm iZ^{\pm} \\ \{X^{\pm}, X^{\pm}\} &= \{X^{\pm}, X^{\mp}\} = \{Z^{\pm}, X^{\pm}\} = \{Z^{\pm}, X^{\mp}\} = \{Z^{\pm}, Z^{\pm}\} = \{Z^{\pm}, Z^{\mp}\} = 0 \\ EX^{\pm} &= -X^{\pm}E, \quad EZ^{\pm} = -Z^{\pm}E, \quad [X^{\pm}, F] = [Z^{\pm}, F] = 0. \end{aligned}$$

Let also $\bar{L} := K^{r''}K_3^2$ as a shorthand. The Hopf structure is defined by

$$\begin{aligned} \epsilon(K_a) &= 1, \quad \epsilon(X^{\pm}) = \epsilon(Z^{\pm}) = 0, \\ \Delta(K_a) &= K_a \otimes K_a, \quad \Delta(X^{\pm}) = \mathbf{1} \otimes X^{\pm} + X^{\pm} \otimes (K^{r''}K_1K_2)^{\pm 1}, \quad \Delta(Z^{\pm}) = \mathbf{1} \otimes Z^{\pm} + Z^{\pm} \otimes \bar{L} \\ S(K_a) &= K_a^{-1} \quad S(X^{\pm}) = -X^{\pm}(K^{r''}K_1K_2)^{\mp 1} \quad S(Z^{\pm}) = -Z^{\pm}\bar{L}. \end{aligned}$$

At $r = 8$ the dimension is $4^4 \times 2^4 \times 2^2 \times 2^2 = 2^{16} = 65536$.

Proposition 4.25. The algebra $u_q\mathfrak{sl}_2 \times \mathbb{N}_2$ is

1. unimodular with a two-sided cointegral

$$\Lambda := \frac{\{1\}^{r'-1}}{\sqrt{r''}[r'-1]!} \sum_{a=0}^{r'-1} \sum_{b,c,d=0}^4 E^{r'-1} F^{r'-1} K^a K_1^b K_2^c K_3^d X^+ X^- Z^+ Z^-,$$

and a left integral expressed on the monomial basis by

$$\begin{aligned} &\lambda(E^e F^f K^a K_1^b K_2^c K_4^d (X^+)^g (X^-)^h (Z^+)^i (Z^-)^j) \\ &= \frac{\sqrt{r''}[r'-1]!}{\{1\}^{r'-1}} \delta_{a,r'-1} \delta_{b,0} \delta_{c,0} \delta_{d,0} \delta_{e,r'-1} \delta_{f,r'-1} \delta_{g,1} \delta_{h,1} \delta_{a,1} \delta_{j,1}. \end{aligned}$$

2. quasitriangular, with the R-matrix $R := R_{\mathbf{z}} D \Theta R_{\alpha}$, where D, Θ were defined in Proposition 4.2, and

$$R_{\mathbf{z}} := \frac{1}{64} \sum_{\mathbf{v}, \mathbf{w} \in \mathbb{Z}_4^3} i^{-\mathbf{v}\mathbf{w}^T} (K_1, K_2, K_3)^{\mathbf{w}} \otimes (K_1, K_2, K_3)^{\mathbf{v}\mathbf{z}},$$

$$\text{for } \mathbf{z} = \begin{pmatrix} 0 & 3 & 2 \\ 1 & 0 & 0 \\ 2 & 0 & 2 \end{pmatrix}, \text{ and}$$

$$R_{\alpha} := \exp(\alpha(Z^+ \otimes \bar{L}Z^- - Z^- \otimes \bar{L}Z^+)).$$

3. ribbon, with the ribbon element

$$v = \frac{1-i}{\sqrt{r'}} \sum_{a=0}^{r'-1} \sum_{b=0}^{r''-1} \frac{\{-1\}^a}{[a]!} q^{-\frac{(a+3)a}{2} + 2b^2} K^{-a-2b-1} K_3^2 E^a F^a \exp(-2\alpha Z^+ Z^-).$$

corresponding to the pivotal element $g = K$.

Proof. See A.3 in Appendix A. \square

5 Topological applications

As mentioned in the introduction, the main motivation to study non-factorizable ribbon Hopf algebras is that they produce invariants of 4-dimensional 2-handlebodies as described by Beliakova and De Renzi in [BdR23]. More details can be found in [BBDRP23], which relies partly on [BP06]. We first introduce some definitions, and then recall some of the results we mentioned. For a precise introduction to the theory of handles and handlebodies, and more precise definitions, we refer to [GS99, Sections 4.1 and 4.2].

5.1 Handlebodies

Recall that a n -dimensional k -handle is a copy of the 4-ball $D^k \times D^{n-k}$ destined to be attached to another manifold along $(\partial D^k) \times D^{n-k}$. The *core* of the handle is a k -cell $D^k \times \{0\}$, and the handle is a thickening of it.

Definition 5.1. A 4-dimensional 2-handlebody is a smooth manifold W together with a filtration

$$W_0 \subset W_1 \subset W_2,$$

where W_0 is a finite disjoint union of 4-balls, W_1 is obtained from W_0 by gluing 1-handles and W_2 is obtained from W_1 by gluing 2-handles.

Remark 5.2. Up to 1-deformations (defined below), if W is connected, we can always assume that W_0 is connected, i.e. that there is a single 4-ball. Hence from now on we shall consider only this case.

Note that such an object can be encoded in a Kirby diagram as explained in [GS99, Section 4.4]. For us, a regular Kirby diagram is a diagram of a link in S^3 that splits as a disjoint union of a trivial unlink diagram whose components are dotted, and a framed link. Such a datum is meant to represent the attaching maps of the handles in $S^3 = \partial D^4$. The framed knots represent embedded tori where the 2-handles are attached, while the dotted unlink represents 1-handles as explained in [GS99, Section 5.4]. Roughly speaking, instead of adding 1-handles, one can equivalently push a disk bounded by an undotted component inside D^4 and excise a neighborhood of it.

A famous theorem of Cerf implies that two diffeomorphic 2-handlebodies are related by a sequence of handle slides, creation/removal of cancelling pairs of handles, and isotopies respecting the filtration. This motivates the following definition.

Definition 5.3. Let $1 \leq l \leq 4$. An l -deformation of a 4-dimensional 2-handlebody is a finite sequence of

- isotopies of attaching maps of i -handles in ∂W_{i-1} , for $1 \leq i \leq l$,
- handle slides of i -handles over other i -handles for $1 \leq i \leq l$
- creations/removals of cancelling pairs of $(i/i-1)$ -handles for $1 \leq i \leq l$.

Two handlebodies related by an l -deformation are called l -equivalent.

It is known (see [BP06, Section 1.2]) that two diffeomorphic handlebodies are 3-equivalent, but it is not known if they are always 2-equivalent:

Conjecture 5.4 (Generalized Gompf conjecture). Two diffeomorphic 4-dimensional 2-handlebodies are 2-equivalent.

There exist potential counterexamples to this conjecture, that is pairs of 3-equivalent handlebodies that have not been proven not to be 2-equivalent. Hence it is interesting to be provided with invariants of handlebodies up to 2-equivalence.

5.2 Invariants of handlebodies up to 2-equivalence

In [BBDRP23], a category 4HB whose objects are 3-dimensional 1-handlebodies and whose morphisms are 4-dimensional 2-handlebodies up to 2-equivalence is defined, and proven to be equivalent as a braided category to the strict braided monoidal category freely generated by a single so-called BP-Hopf algebra H . Here BP is short for Bobtcheva-Piergallini, and the definition of a BP-Hopf algebra object can be found at [BBDRP23, Definition 2.4.1]. We shall not need this result in its full extent, but we observe that any BP-Hopf algebra in a braided category \mathcal{C} yields a functor from 4HB to \mathcal{C} . In particular, the objects of 4HB are in bijection with the natural numbers, and the morphisms from 0 to 0 are precisely the set of 4-dimensional 2-handlebodies up to 2-equivalence. Thus any BP-Hopf algebra gives an invariant of 4-dimensional 2-handlebodies up to 2-equivalence.

A very important feature of a unimodular ribbon Hopf algebra H is that it can be "transmuted" into a BP-Hopf algebra \underline{H} in the category $H\text{-mod}$ (as a special case of [BdR23, Prop. 7.3]), where \underline{H} is H as an H -module for the left adjoint action from Proposition 3.8. Summarizing, a unimodular ribbon Hopf algebra yields an invariant J_H of 2-handlebodies through a rather complicated process. Fortunately, we do not have to go through the details since a direct algorithm computing the invariant is given in [BdR23, Section 8]. Let us sketch the algorithm, since it will have an importance in the proofs below.

Algorithm 5.5 (Bead Algorithm, [BdR23]). Let W be a 4-dimensional 2-handlebody and L be an associated regular Kirby diagram, then the invariant $J_H(W)$ can be computed in the following way. From the initial diagram, we get another diagram with labeled beads by

1. deleting crossings and placing beads labeled with the components of the R-matrix,

$$\begin{array}{ccc}
\begin{array}{c} \diagup \\ \diagdown \\ \diagup \\ \diagdown \end{array} & \mapsto & \begin{array}{c} \diagup \\ \diagdown \\ \circ R' \\ \diagup \\ \diagdown \\ \circ R'' \end{array} & \text{and} & \begin{array}{c} \diagup \\ \diagdown \\ \diagup \\ \diagdown \end{array} & \mapsto & \begin{array}{c} \diagup \\ \diagdown \\ \circ R'' \\ \diagup \\ \diagdown \\ \circ S(R') \end{array}
\end{array}$$

2. deleting 1-handles and placing beads corresponding to an iterated coproduct of the cointegral,

$$\begin{array}{ccc}
\begin{array}{c} | \\ | \\ \cdots \\ | \\ | \end{array} & \mapsto & \begin{array}{c} | \\ | \\ \Lambda_{(1)} \circ \cdots \circ \Lambda_{(m)} \\ | \\ | \end{array}
\end{array}$$

3. deleting framings and placing beads labeled with copies of the pivotal element g

$$\begin{array}{ccc}
\begin{array}{c} | \\ \curvearrowright \\ | \end{array} & \mapsto & \begin{array}{c} | \\ \circ g \\ | \end{array} & \text{and} & \begin{array}{c} | \\ \curvearrowleft \\ | \end{array} & \mapsto & \begin{array}{c} | \\ \circ g^{-1} \\ | \end{array}
\end{array}$$

We collect the beads together using the rules

$$\begin{array}{ccc}
\begin{array}{c} \circ x \\ | \\ \cup \\ | \end{array} & = & \begin{array}{c} | \\ \cup \\ | \\ \circ S(x) \end{array}, & S(x) \begin{array}{c} | \\ \cup \\ | \end{array} & = & \begin{array}{c} | \\ \cup \\ | \\ \circ x \end{array} \\
\text{and} & & \begin{array}{c} \circ x \\ | \\ \circ y \\ | \end{array} & = & \begin{array}{c} | \\ \circ xy \\ | \end{array}
\end{array}$$

and isotope the components until we get a trivial diagram of an unlink with a single labeled bead on every component:

$$\begin{array}{ccc}
\begin{array}{c} \circ z_1 \\ \bigcirc \end{array} & \begin{array}{c} \circ z_2 \\ \bigcirc \end{array} & \cdots & \begin{array}{c} \circ z_k \\ \bigcirc \end{array}
\end{array}$$

Then

$$J_H(W) = \prod_{j=1}^k \lambda(z_j),$$

where the value is computed using the Sweedler notation. That is, here, $J_H(W)$ is computed as a sum of products, that is a linear combination of terms in $\underline{H}^{\otimes k}$, and not a product.

5.3 Cases of degeneration of the invariant

When the unimodular ribbon Hopf algebra is factorizable Beliakova and De Renzi proved that the invariant we obtained is actually an invariant of the boundary of the handlebody. We call this situation *degeneration to dimension 3*.

Proposition 5.6 ([BdR23], Theorem 1.2 and Section 8.2). If the unimodular ribbon category H is factorizable, then the invariant J_H degenerates to dimension 3.

Furthermore, it is known that two diffeomorphic 4-dimensional 2-handlebodies become 2-equivalent after connected summing a certain number of times with $S^2 \times D^2$. Since the invariant J_4 is additive with respect to the connected summing, this implies that in order to tackle Conjecture 5.4, we would like the invariant $J_4(S^2 \times D^2)$ to be trivial. This is tantamount to H not being cosemisimple.

Proposition 5.7 ([BdR23], Appendix C). If the unimodular ribbon category H is cosemisimple, and W_1 and W_2 are diffeomorphic handlebodies, then $J_H(W_1) = J_H(W_2)$.

Finally, roughly speaking, the invariant can degenerate to dimension 2. Indeed any 4-dimensional 2-handlebody W deformation-retracts onto its *spine*: the 2CW-complex of dimension 2 determined by the cores of its handles. This complex, in turn, yields a presentation of the fundamental group of the 4-dimensional 2-handlebody. A 2-deformation of the handlebody is reflected as an Andrews-Curtis (or 2-equivalence) move for the group presentation. We refer to [Bob23] for more details. If an invariant of 4-dimensional 2-handlebody cannot distinguish between handlebodies with 2-equivalent group presentations of their fundamental groups, then we say it *degenerates to their spines*. The following is a consequence of [Bob23].

Proposition 5.8. If the unimodular ribbon Hopf algebra H is triangular (i.e. has trivial ribbon element), then the invariant J_H degenerates to its spine.

The generalized Andrews-Curtis conjecture states that any balanced presentation of the trivial group can be reduced to the empty presentation through a sequence of Andrews-Curtis moves. Hence an invariant that degenerates to the spines is still a useful invariant for tackling the Andrews-Curtis conjecture.

We have justified the interest in searching for non-factorizable non-cosemisimple unimodular ribbon Hopf algebras. Nevertheless we show that, for most of the non-factorizable examples H we gave in the previous sections, the invariant J_H is still not rich enough for the Gompf conjecture.

5.4 Degeneration of the invariant associated to the Nenciu algebras

In this section, we show that for any of the strongly non-factorizable Hopf algebra

$$H = H(\boldsymbol{\alpha}) := (H(\mathbf{m}, \mathbf{t}, \mathbf{d}, \mathbf{u}), R(\boldsymbol{\alpha}))$$

constructed using the approach of the previous sections, the invariant J_H degenerates to the spines. For the purpose of exposition, as in Example 3.11, we will call Z_l^\pm , $l = 1, \dots, t_2$ nilpotent generators participating in the R-matrix, and X_k , $k = 1, \dots, t_1$ the ones appearing only in the cointegral. In the notation, we emphasized the tuple $\boldsymbol{\alpha} \in \mathbb{C}^l$ that parametrizes the R-matrix as in in Example 3.11.

Theorem 5.9. *Consider a Kirby diagram for W , with k_1 1-handles and k_2 2-handles. Then the invariant J_H obtained using an strongly non-factorizable Hopf algebra $H(\boldsymbol{\alpha})$ constructed as in previous sections, such as the one in Example 3.11 behaves as follows.*

- If $k_1 = k_2$, then $J_H(W) = J_{H(0)}(W)$, where the latter invariant degenerates to the spine.
- If $k_1 \neq k_2$, the invariant vanishes: $J_H(W) = 0$.




Figure 1: A Kirby diagram with k_1 1-handles isotoped to the bottom, k_2 (possibly knotted) 2-handles and m crossings, and the image under the bead algorithm, where each 2-handle is linked to at least one 1-handle. The beads have been colored arbitrarily and not all labeled for a better readability.

Proof. To compute the invariant, we isotope the 1-handles to the bottom of the diagram, as in Figure 1, and we use Algorithm 5.5. Recall that in H , the generators X_k do not appear in the R-matrix nor the ribbon element, and that the integral λ will vanish on a product of generators unless it contains precisely once each skew-primitive generator. First of all, we exclude some simple cases, where the invariant vanishes regardless of k_1 and k_2 .

- If the diagram contains an isolated 1-handle (not attached to any 2-handles), then there is a contribution of $\epsilon(\Lambda) = 0$ by non-semisimplicity of the algebra $H(\boldsymbol{\alpha})$.
- If there is an isolated (unlinked from 1-handles and 2-handles) 2-handle in the diagram, carrying a bead with a label P , this gives a contribution of $\lambda(P) = 0$, as P is a sum of products where none of the X_k generators appear.
- Similarly, if there is a 2-handle linked only to another 2-handle and no 1-handles, carrying an bead with a label P , there is a contribution $\lambda(P) = 0$, as P is obtained from copies of the R-matrix and the ribbon element, corresponding to crossings and framings.

With these cases excluded, we can ensure that if $k_1 > 0$ and every 2-handle links to at least one 1-handle, as in the top of Fig. 1. We apply Algorithm 5.5 and find the bottom picture of Fig. 1. Using the commutation relations, we arrive at the following value of the invariant

$$J_H(W) = \prod_{j=1}^{k_2} C \lambda_j (\Lambda_{(1_{1_j})}^{1_j} \cdots \Lambda_{(i_{1_j})}^{1_j} \cdots \Lambda_{(1_{m_j})}^{m_j} \Lambda_{(i_{m_j})}^{m_j} R'_a \cdots S(R''_b) \cdots),$$

where the Sweedler notation hides that it is a sum over all indices. Here $C \in \mathbb{C}$ is a constant resulting from the commutations between the generators, and depends both on j and the choice of the term in the sum. The j -th 2-handle links to m_j 1-handles, indexed with $1_j, \dots, m_j$, and it links precisely i_{l_j} times with the l -th 1-handle. Every linking carries a piece of $\Delta(\Lambda)$ in the Sweedler notation. This is followed by a product of copies of R', R'' and their antipodes. Once again, recall that R is generated by grouplike generators \mathbf{K} and nilpotent generators Z_l^\pm , $l = 1, \dots, t_2$, while Λ contains also X_k , $k = 1, \dots, t_1$, and λ will vanish on any product of generators that does not contain precisely once each of the variable X_k . When taking coproducts of the integral element, the t_1 variables X_k are distributed among the factors of the tensor product. Then, for fixed indices the term

$$\prod_{j=1}^{k_2} C \lambda_j (\Lambda_{(1_{1_j})}^{1_j} \cdots \Lambda_{(i_{1_j})}^{1_j} \cdots \Lambda_{(1_{m_j})}^{m_j} \Lambda_{(i_{m_j})}^{m_j} R'_a \cdots S(R''_b) \cdots)$$

contains exactly $k_1 t_1$ of the variables X_k . But if we want this term to be non-trivial, then there should be also t_1 variables X_k for each j , in order to “feed” the integral λ . Thus we should have $k_1 t_1 = k_2 t_1$. Then we should have $k_1 = k_2$. As a heuristic summary, we have the following three cases.

Case 1: “underfed” If $k_1 < k_2$, then the invariant vanishes, as in every term, there will be at least one j such λ_j does not have enough X_k generators to “eat” in order to be non-zero.

Case 2: “overfed” If $k_1 > k_2$, then in every term, there is a j with too many of the X_k variables. Thus, the invariant vanishes again.

Case 3: “balanced” If $k_1 = k_2$, then, at least in some terms, there is exactly enough copies of each X_k^\pm to feed all integrals. Since $\Delta(\Lambda)$ produces all shuffles of T (with grouplike prefactors), for at least one configuration the obstacles of cases 1 and 2 can be avoided, since we assumed $m_j > 0$ for all j if $k_1 > 0$. But now the cointegrals introduced also the correct number of copies of Z_l^\pm . But

$$R = 1 \otimes 1 + \alpha_1 (Z_1^+ \otimes L_1 Z_1^- - Z_1^- \otimes L_1 Z_1^+) + \dots,$$

so that every term involving the variable α_l will have too many copies of variables Z . Hence, setting the parameter α_l to 0 does not change the final value of the invariant.

In conclusion, the invariant can be non-vanishing only in the balanced case, and then is tantamount to the invariant obtained from the triangular Hopf algebra $H(\boldsymbol{\alpha} = \mathbf{0})$. \square

Remark 5.10. Here is a series of facts clarifying where the algebras $H(\boldsymbol{\alpha})$ lie regarding the Gompf conjecture.

1. Notice that the Euler characteristic of the handlebody is given by $1 - k_1 + k_2$. Hence if the Euler characteristic of W is different from 1, then $J_H(W) = 0$.
2. Since two diffeomorphic handlebodies have the same Euler characteristic, tackling the Gompf conjecture with the invariant J_H would be possible only for handlebodies of Euler characteristic 1, corresponding to the balanced case, and then we could alternatively work with the invariant $J_H(\boldsymbol{\alpha} = \mathbf{0})$ associated to a triangular algebra. Hence, we would be looking at the Andrews-Curtis conjecture.
3. The algebras are applicable indeed to the Andrews-Curtis case, but by the classification of triangular Hopf algebras of [AEG02] and their correspondence to Hopf supergroups, the cases where $\boldsymbol{\alpha} = \mathbf{0}$ are incarnations of the latter.
4. The above reasoning for the balanced case breaks down if we introduce non-diagonal relations involving some of the Z_l^\pm which do not preserve the number of them in the expression, but produces a non-zero expression nevertheless. An example is $[E, F]$ in $u_q \mathfrak{sl}_2$. Then in the balanced case the generators coming from R (possibly in various powers) can in principle survive up to the integrals. This motivated the construction of the semi-direct biproducts $U \ltimes H$.

Let $U = u_q \mathfrak{sl}_2$ and $H = H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$ and let $U \ltimes H$ be their semidirect biproduct as in Definition 4.22. The authors observed through direct computation on many examples where W is a 0 to 0 morphism in the category **4HB**, that the invariant $J_{U \ltimes H}$ factorizes: it satisfies $J_{U \ltimes H}(W) = J_U(W) J_H(W)$. In particular, it does not seem to carry any truly 4-dimensional information, thus we leave this as a conjecture

Conjecture 5.11. Let $U = u_q \mathfrak{sl}_2$ and $H = H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$ and let $U \ltimes H$ be their semidirect biproduct as in Definition 4.22. Let also W be a 0 to 0 morphism in the category **4HB**. Then the invariant given by Algorithm 5.5 satisfies

$$J_{U \ltimes H}(W) = J_U(W) J_H(W).$$

It has to be noted, however, that for higher-rank, say 1 to 0 or 1 to 1, morphisms of **4HB**, the interaction of U and H pieces becomes manifest, so there is a possibility the invariant can detect 4-dimensional information there.

A Proofs

In this appendix, in a series of propositions, we prove Theorem 4.8, Theorem 4.13 as well as Propositions 4.21 and 4.25.

A.1 Proof of Theorem 4.8

Proof of Theorem 4.8. Let $X_k \in H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$, for $k = 1, \dots, t_t$. Let also $L_k := \mathbf{K}^{\mathbf{u}^k}$ for convenience. Since there is no non-trivial action on the elements $K_a \in H$, we concentrate on the generators $X_k \in H$. From the formulas of Definition 4.4 we first fix for any $u \in U$, $h \in H$

$$u \otimes h := (u \otimes \mathbf{1}_H)(\mathbf{1}_U \otimes h)$$

and obtain the following algebra relations on generators

$$\begin{aligned} (\mathbf{1}_U \otimes K_a)(K \otimes \mathbf{1}_H) &= (K \otimes \mathbf{1}_H)(\mathbf{1}_U \otimes K_a) = (K \otimes K_a) \\ (\mathbf{1}_U \otimes K_a)(E \otimes \mathbf{1}_H) &= (E \otimes \mathbf{1}_H)(\mathbf{1}_U \otimes K_a) = (E \otimes K_a) \\ (\mathbf{1}_U \otimes K_a)(F \otimes \mathbf{1}_H) &= (F \otimes \mathbf{1}_H)(\mathbf{1}_U \otimes K_a) = (F \otimes K_a). \\ (\mathbf{1}_U \otimes X_k)(K \otimes \mathbf{1}_H) &= -(K \otimes \mathbf{1}_H)(\mathbf{1}_U \otimes X_k) = -(K \otimes X_k) \\ (\mathbf{1}_U \otimes X_k)(E \otimes \mathbf{1}_H) &= -(E \otimes \mathbf{1}_H)(\mathbf{1}_U \otimes X_k) = -(E \otimes X_k) \\ (\mathbf{1}_U \otimes X_k)(F \otimes \mathbf{1}_H) &= (F \otimes \mathbf{1}_H)(\mathbf{1}_U \otimes X_k) = (F \otimes X_k). \end{aligned}$$

We can easily extend these relations to arbitrary monomials $E^a F^b K^c \in U$ and $\mathbf{K}^{\mathbf{v}} \mathbf{X}^{\mathbf{r}} \in H$ for $a, b, c \in \mathbb{Z}_{r'}$, $\mathbf{v} \in \mathbb{Z}_{\mathbf{m}}$ and $\mathbf{r} \in \mathbb{Z}_2^t$, resulting in

$$(E^a F^b K^c \otimes \mathbf{1}_H)(\mathbf{1}_U \otimes \mathbf{K}^{\mathbf{v}} \mathbf{X}^{\mathbf{r}}) = E^a F^b K^c \otimes \mathbf{K}^{\mathbf{v}} \mathbf{X}^{\mathbf{r}},$$

and

$$(\mathbf{1}_U \otimes \mathbf{K}^{\mathbf{v}} \mathbf{X}^{\mathbf{r}})(E^a F^b K^c \otimes \mathbf{1}_H) = (-1)^{|\mathbf{r}|(a+c)} E^a F^b K^c \otimes \mathbf{K}^{\mathbf{v}} \mathbf{X}^{\mathbf{r}}. \quad (3)$$

Here we used the fact the the map i of Remark 4.10 is an algebra map which can be checked directly. The list of the relations between the generators is an exhaustive one, as the choices of action and coaction force them to be diagonal. Moreover, with the operations defined for arbitrary monomials as above, we see that the new relations respect the monomial bases of both pieces. Since both U and H are fully defined by the relations on generators and their extension over the PBW- and monomial bases respectively, and this fact is respected by the new operations, we can conclude there are no relations between the pieces U and H not resulting from the relation of Equation 3. That is, we obtain the exhaustive list of relations of $U \times H$ by deriving them from the relations between the generators and applying the fact they respect the new operations of the Hopf algebra.

Using the algebra structure that is now fixed, it is proved in Proposition 4.9 that $U \times H$ has a monomial basis consisting of terms of the form $E^a F^b K^c \otimes \mathbf{K}^{\mathbf{v}} \mathbf{X}^{\mathbf{r}}$, so we can suppress the tensor product by defining

$$E^a F^b K^c \mathbf{K}^{\mathbf{v}} \mathbf{X}^{\mathbf{r}} := E^a F^b K^c \otimes \mathbf{K}^{\mathbf{v}} \mathbf{X}^{\mathbf{r}}$$

for the entire monomial basis.

We now compute the coproduct of any such monomial in $U \times H$ and show that Δ is an algebra map. Define the "parity" of monomials $|E^a F^b K^c| := a + c$ and of $|\mathbf{K}^{\mathbf{v}} \mathbf{X}^{\mathbf{r}}| := |\mathbf{r}|$. Firstly, it is clear from Definition 4.4 that

$$\Delta(E^a F^b K^c \otimes \mathbf{1}_H) = ((E^a F^b K^c)_{(1)} \otimes \mathbf{1}_H) \otimes ((E^a F^b K^c)_{(2)} \otimes \mathbf{1}_H),$$

so it is an algebra map for any product of monomials purely in U . Then, we also have

$$\Delta(\mathbf{1}_U \otimes \mathbf{K}^{\mathbf{v}} \mathbf{X}^{\mathbf{r}}) = (\mathbf{1}_U \otimes (\mathbf{K}^{\mathbf{v}} \mathbf{X}^{\mathbf{r}})_{(1)}) \otimes \left(K^{|\mathbf{K}^{\mathbf{v}} \mathbf{X}^{\mathbf{r}}|_{(1)}|r''} \otimes (\mathbf{K}^{\mathbf{v}} \mathbf{X}^{\mathbf{r}})_{(2)} \right)$$

now let $\mathbf{1}_U \otimes \mathbf{K}^{\mathbf{v}_1} \mathbf{X}^{\mathbf{r}_1}$ and $\mathbf{1}_U \otimes \mathbf{K}^{\mathbf{v}_2} \mathbf{X}^{\mathbf{r}_2}$ be two such monomials. Then

$$\begin{aligned} &\Delta(\mathbf{1}_U \otimes \mathbf{K}^{\mathbf{v}_1} \mathbf{X}^{\mathbf{r}_1}) \Delta(\mathbf{1}_U \otimes \mathbf{K}^{\mathbf{v}_2} \mathbf{X}^{\mathbf{r}_2}) \\ &= (\mathbf{1}_U \otimes (\mathbf{K}^{\mathbf{v}_1} \mathbf{X}^{\mathbf{r}_1})_{(1)}) \otimes \left(K^{|\mathbf{K}^{\mathbf{v}_1} \mathbf{X}^{\mathbf{r}_1}|_{(1)}|r''} \otimes (\mathbf{K}^{\mathbf{v}_1} \mathbf{X}^{\mathbf{r}_1})_{(2)} \right) \\ &\quad (\mathbf{1}_U \otimes (\mathbf{K}^{\mathbf{v}_2} \mathbf{X}^{\mathbf{r}_2})_{(1)}) \otimes \left(K^{|\mathbf{K}^{\mathbf{v}_2} \mathbf{X}^{\mathbf{r}_2}|_{(1)}|r''} \otimes (\mathbf{K}^{\mathbf{v}_2} \mathbf{X}^{\mathbf{r}_2})_{(2)} \right). \end{aligned}$$

Now $K^{r''}$ commutes with all K_a and X_k , so using the fact Δ_H is an algebra map on H , the expression rearranges to

$$\begin{aligned} &(\mathbf{1}_U \otimes (\mathbf{K}^{\mathbf{v}_1} \mathbf{X}^{\mathbf{r}_1} \mathbf{K}^{\mathbf{v}_2} \mathbf{X}^{\mathbf{r}_2})_{(1)}) \otimes \left(K^{|\mathbf{K}^{\mathbf{v}_1} \mathbf{X}^{\mathbf{r}_1} \mathbf{K}^{\mathbf{v}_2} \mathbf{X}^{\mathbf{r}_2}|_{(1)}|r''} \otimes (\mathbf{K}^{\mathbf{v}_1} \mathbf{X}^{\mathbf{r}_1} \mathbf{K}^{\mathbf{v}_2} \mathbf{X}^{\mathbf{r}_2})_{(2)} \right) \\ &= \Delta(\mathbf{1}_U \otimes \mathbf{K}^{\mathbf{v}_1} \mathbf{X}^{\mathbf{r}_1} \mathbf{K}^{\mathbf{v}_2} \mathbf{X}^{\mathbf{r}_2}). \end{aligned}$$

Thus we checked that Δ behaves as an algebra map on products of monomials coming from H . It is left to check the cross-terms. First consider

$$\begin{aligned}
& \Delta(E^a F^b K^c \otimes \mathbf{1}_H) \Delta(\mathbf{1}_U \otimes \mathbf{K}^\mathbf{v} \mathbf{X}^\mathbf{r}) \\
&= (((E^a F^b K^c)_{(1)} \otimes \mathbf{1}_H) \otimes ((E^a F^b K^c)_{(2)} \otimes \mathbf{1}_H)) \\
&\quad \left((\mathbf{1}_U \otimes (\mathbf{K}^\mathbf{v} \mathbf{X}^\mathbf{r})_{(1)}) \otimes K^{|\mathbf{K}^\mathbf{v} \mathbf{X}^\mathbf{r}|_{(1)}|r''} \otimes (\mathbf{K}^\mathbf{v} \mathbf{X}^\mathbf{r})_{(2)} \right) \\
&= ((E^a F^b K^c)_{(1)}) \otimes (\mathbf{K}^\mathbf{v} \mathbf{X}^\mathbf{r})_{(1)} \otimes \left((E^a F^b K^c)_{(2)} K^{|\mathbf{K}^\mathbf{v} \mathbf{X}^\mathbf{r}|_{(2)}|r''} \otimes (\mathbf{K}^\mathbf{v} \mathbf{X}^\mathbf{r})_{(2)} \right) \\
&= \Delta(E^a F^b K^c \otimes \mathbf{K}^\mathbf{v} \mathbf{X}^\mathbf{r}).
\end{aligned}$$

Then for $d, e, f \in \mathbb{Z}_{r'}$ using the above

$$\begin{aligned}
& \Delta(E^a F^b K^c \otimes \mathbf{1}_H) \Delta(E^e F^f K^d \otimes \mathbf{K}^\mathbf{v} \mathbf{X}^\mathbf{r}) \\
&= \Delta(E^a F^b K^c \otimes \mathbf{1}_H) \Delta(E^e F^f K^d \otimes \mathbf{1}_H) \Delta(\mathbf{1}_U \otimes \mathbf{K}^\mathbf{v} \mathbf{X}^\mathbf{r}) \\
&= \Delta(E^a F^b K^c E^e F^f K^d \otimes \mathbf{1}_H) \Delta(\mathbf{1}_U \otimes \mathbf{K}^\mathbf{v} \mathbf{X}^\mathbf{r}) \\
&= \Delta(E^a F^b K^c E^e F^f K^d \otimes \mathbf{K}^\mathbf{v} \mathbf{X}^\mathbf{r})
\end{aligned}$$

Similarly,

$$\begin{aligned}
& \Delta(E^a F^b K^c \otimes \mathbf{K}^{\mathbf{v}_1} \mathbf{X}^{\mathbf{r}_1}) \Delta(\mathbf{1}_U \otimes \mathbf{K}^{\mathbf{v}_2} \mathbf{X}^{\mathbf{r}_2}) \\
&= \Delta(E^a F^b K^c \otimes \mathbf{1}_H) \Delta(\mathbf{1}_U \otimes \mathbf{K}^{\mathbf{v}_1} \mathbf{X}^{\mathbf{r}_1}) \Delta(\mathbf{1}_U \otimes \mathbf{K}^{\mathbf{v}_2} \mathbf{X}^{\mathbf{r}_2}) \\
&= \Delta(E^a F^b K^c \otimes \mathbf{1}_H) \Delta(\mathbf{1}_U \otimes \mathbf{K}^{\mathbf{v}_1} \mathbf{X}^{\mathbf{r}_1} \mathbf{K}^{\mathbf{v}_2} \mathbf{X}^{\mathbf{r}_2}) \\
&= \Delta(E^a F^b K^c \otimes \mathbf{K}^{\mathbf{v}_1} \mathbf{X}^{\mathbf{r}_1} \mathbf{K}^{\mathbf{v}_2} \mathbf{X}^{\mathbf{r}_2})
\end{aligned}$$

Thus, it is now enough to establish

$$\begin{aligned}
& \Delta(\mathbf{1}_U \otimes \mathbf{K}^\mathbf{v} \mathbf{X}^\mathbf{r}) \Delta(E^a F^b K^c \otimes \mathbf{1}_H) \\
&= \left((\mathbf{1}_U \otimes (\mathbf{K}^\mathbf{v} \mathbf{X}^\mathbf{r})_{(1)}) \otimes (K^{|\mathbf{K}^\mathbf{v} \mathbf{X}^\mathbf{r}|_{(1)}|r''} \otimes (\mathbf{K}^\mathbf{v} \mathbf{X}^\mathbf{r})_{(2)}) \right) \\
&\quad \left(((E^a F^b K^c)_{(1)} \otimes \mathbf{1}_H) \otimes ((E^a F^b K^c)_{(2)} \otimes \mathbf{1}_H) \right) \\
&= (-1)^{|(E^a F^b K^c)_{(1)}| |(\mathbf{K}^\mathbf{v} \mathbf{X}^\mathbf{r})_{(1)}| + |(E^a F^b K^c)_{(2)}| |(\mathbf{K}^\mathbf{v} \mathbf{X}^\mathbf{r})_{(2)}|} \\
&\quad \left((E^a F^b K^c)_{(1)} \otimes (\mathbf{K}^\mathbf{v} \mathbf{X}^\mathbf{r})_{(1)} \right) \otimes \left((E^a F^b K^c)_{(2)} K^{|\mathbf{K}^\mathbf{v} \mathbf{X}^\mathbf{r}|_{(2)}|r''} \otimes (\mathbf{K}^\mathbf{v} \mathbf{X}^\mathbf{r})_{(2)} \right).
\end{aligned}$$

But after considering the sums hidden in the Sweedler notation we have the sign factor

$$(-1)^{|(E^a F^b K^c)_{(1)}| |(\mathbf{K}^\mathbf{v} \mathbf{X}^\mathbf{r})_{(1)}| + |(E^a F^b K^c)_{(2)}| |(\mathbf{K}^\mathbf{v} \mathbf{X}^\mathbf{r})_{(2)}|} = (-1)^{|E^a F^b K^c| |\mathbf{K}^\mathbf{v} \mathbf{X}^\mathbf{r}|}$$

so the result is

$$(-1)^{|E^a F^b K^c| |\mathbf{K}^\mathbf{v} \mathbf{X}^\mathbf{r}|} \Delta(E^a F^b K^c \otimes \mathbf{K}^\mathbf{v} \mathbf{X}^\mathbf{r}).$$

The sign factor is exactly $(-1)^{|\mathbf{r}|(a+c)}$ as above. Then we have the general formula

$$\begin{aligned}
& \Delta(E^a F^b K^c \otimes \mathbf{K}^{\mathbf{v}_1} \mathbf{X}^{\mathbf{r}_1}) \Delta(E^e F^f K^d \otimes \mathbf{K}^{\mathbf{v}_2} \mathbf{X}^{\mathbf{r}_2}) \\
&= \Delta(E^a F^b K^c \otimes \mathbf{1}_H) \Delta(\mathbf{1}_U \otimes \mathbf{K}^{\mathbf{v}_1} \mathbf{X}^{\mathbf{r}_1}) \Delta(E^e F^f K^d \otimes \mathbf{1}_H) \Delta(\mathbf{1}_U \otimes \mathbf{K}^{\mathbf{v}_2} \mathbf{X}^{\mathbf{r}_2}) \\
&= (-1)^{|E^e F^f K^g| |\mathbf{K}^{\mathbf{v}_1} \mathbf{X}^{\mathbf{r}_1}|} \Delta(E^a F^b K^c E^e F^f K^d \otimes \mathbf{K}^{\mathbf{v}_1} \mathbf{X}^{\mathbf{r}_1} \mathbf{K}^{\mathbf{v}_2} \mathbf{X}^{\mathbf{r}_2}).
\end{aligned}$$

Thus, Δ is an algebra map, and $U \times H$ is a bialgebra. Now, the antipode is unique if it exists, and a similar check shows it is given by the formula in Definition 4.4. Thus, we can consistently suppress the internal tensor product, and we proved $U \times H$ is indeed a Hopf algebra as required. \square

A.2 Proof of Theorem 4.13

We prove Theorem 4.13 in a series of propositions.

Proposition A.1. Let U carry a left integral λ_U and a two-sided cointegral Λ_U , and H carry a two-sided integral λ_H and a two-sided cointegral Λ_H as in the statement of Theorem 4.13. Then $U \times H$ admits the following structures

1. a two-sided cointegral $\Lambda := \Lambda_U \Lambda_H$

$$\Lambda := \frac{\{1\}^{r'-1}}{\sqrt{r''} [r'-1]!} \sum_{a=0}^{r'-1} E^{r'-1} F^{r'-1} K^a \prod_{a=1}^s \left(\sum_{b=0}^{m_a-1} K_a^b \right) \prod_{k=1}^t X_k,$$

2. a left integral $\lambda \in (U \times H)^*$ defined on the monomial basis of Proposition

$$\begin{aligned}
\lambda(E^b F^c K^a \mathbf{K}^\mathbf{v} \mathbf{X}^\mathbf{r}) := & \begin{cases} \frac{\sqrt{r''} [r'-1]!}{\{1\}^{r'-1}} & \text{if } \mathbf{v} = (0, \dots, 0), \mathbf{r} = (1, \dots, 1), a = b = c = r' - 1 \\ 0 & \text{otherwise.} \end{cases}
\end{aligned}$$

Proof. We need to verify the properties of the integrals according to the Definition 2.5, except for the two-sidedness of the integral which is already not satisfied by the $u_q\mathfrak{sl}_2$ piece as remarked in [BdR23], since the distinguished grouplike element is $K^2 \neq \mathbf{1}$. By Theorem 4.8, it suffices to do it on the monomial basis.

1. $\underline{h\Lambda = \Lambda h = \epsilon(h)\Lambda}$:

Consider Λ multiplying any other element of the monomial basis $h \in H$. If $h = E^b F^c \mathbf{X}^{\mathbf{r}}$ with $b, c \in \{1, \dots, r' - 1\}$ and $\mathbf{r} \in \mathbb{Z}_2^t$, that is it contains nilpotent generators, we clearly get $h\Lambda = \Lambda h = \epsilon(h)\Lambda = 0$, because for the generators E, F the cointegral of Proposition 4.2 is two-sided, and T is central (recall $T := \mathbf{X}^{(1, \dots, 1)}$ is the top element of H). Moreover, $E^{r'-1} F^{r'-1} T$ commutes with all grouplike generators, so since $K \sum_{i=0}^{r'-1} K^i = \sum_{i=0}^{r'-1} K^i$ and $K_a \sum_{i=0}^{m_a-1} K_a^i = \sum_{i=0}^{m_a-1} K_a^i$, we find $K\Lambda = \Lambda K = \epsilon(K)\Lambda = \Lambda$ and $K_a\Lambda = \Lambda K_a = \epsilon(K_a)\Lambda = \Lambda$ for $a = 1, 2, 3$. Thus, Λ is a two-sided cointegral for H .

2. $\underline{(\text{Id} \otimes \lambda)\Delta(h) = \mathbf{1}\lambda(h)}$:

We know that only the monomial $E^{r'-1} F^{r'-1} K^{r'-1} T$ is picked up by the integral, and it is vanishing for any other element of the monomial basis. We need to verify only those monomial basis elements where all nilpotent generators appear, so of the form $E^{r'-1} F^{r'-1} K^a \mathbf{K}^{\mathbf{w}} T$ for some $a \in \{0, \dots, r'\}$ and $\mathbf{w} \in \mathbb{Z}_{\mathbf{m}}$. Then the only term in the coproduct carrying the top piece in the right factor is

$$\Delta(E^{r'-1} F^{r'-1} \mathbf{K}^{\mathbf{w}} T) = K^{1-r'} \mathbf{K}^{\mathbf{w}} \otimes E^{r'-1} F^{r'-1} K^a \mathbf{K}^{\mathbf{w}} T + \dots$$

If we now act with $\text{Id} \otimes \lambda$, we see that $K_a \mathbf{K}^{\mathbf{w}} = K^{r'-1}$, so $a = r' - 1$, $\mathbf{w} = (0, \dots, 0)$ is required for the top term to survive, all other vanish. But then we have

$$\Delta(E^{r'-1} F^{r'-1} K_1^{r'-1} T) = \mathbf{1} \otimes E^{r'-1} F^{r'-1} K^{r'-1} T + \dots$$

so the left factor is proportional to $\mathbf{1}$ as required. Thus, after acting by $\text{Id} \otimes \lambda$, we retrieve the axiom.

3. $\underline{\lambda(\Lambda)=1}$:

Immediately satisfied by construction. □

Proposition A.2. Let $U = u_q\mathfrak{sl}_2$ carry the R-matrix $R_U = D\Theta$ and $H = H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$ carry the R-matrix $R_H = R_{\mathbf{z}} R_{\alpha}$ respectively, such that

$$R_{\alpha} = \exp \left(\sum_{l=0}^{t_2} \alpha_l (Z^+ \otimes LZ^- - Z^- \otimes LZ^+) \right).$$

Then $u_q\mathfrak{sl}_2 \ltimes H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$ admits the R-matrix $R := DR_{\mathbf{z}} \Theta \bar{R}_{\alpha}$, where

$$\bar{R}_{\alpha} := \exp \left(\sum_{l=0}^{t_2} \alpha_l (Z^+ \otimes \bar{L}Z^- - Z^- \otimes \bar{L}Z^+) \right)$$

for $\bar{L} = K^{r''} L$.

Proof. Firstly, we check whether $DR_{\mathbf{z}} \bar{R}_{\alpha}$ fulfills (QT1)-(QT4) of Definition 2.9. But the only change is the appearance of the $K^{r''}$ factors which commute with the entire $H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$, and $(K^{r''})^2 = \mathbf{1}$. Furthermore D and $R_{\mathbf{z}}$ can be collected to one R-matrix containing all grouplike generators, by including K and \mathbf{K} in the tuple $\bar{\mathbf{K}} := (K, K_1, \dots, K_s)$, as well as q^2 and ξ in $\bar{\xi} := (q^2, \xi_1, \dots, \xi_s)$, and extending \mathbf{z} into the block matrix

$$\bar{\mathbf{z}} = \begin{pmatrix} \mathbf{z} & 0 \\ 0 & 1 \end{pmatrix},$$

so that

$$R_{\bar{\mathbf{z}}} = \sum_{\mathbf{v}, \mathbf{w} \in \mathbb{Z}_{r'} \oplus \mathbb{Z}_{\mathbf{m}}} \bar{\xi}^{-\mathbf{v} \cdot \mathbf{w}} \bar{\mathbf{K}}^{\mathbf{w}} \otimes \bar{\mathbf{K}}^{\mathbf{v} \mathbf{z}} = DR_{\mathbf{z}}.$$

This is easily seen to satisfy (QT1)-(QT4) on its own, directly or by the relevant parts of Theorem 2.9. Similarly, it satisfies (QT5) for all $X_k \in H$ as well. Secondly, it is an easy check that

$$[E \otimes F, Z^{\pm} \otimes \bar{L}Z^{\mp}] = [E \otimes F, Z^{\mp} \otimes \bar{L}Z^{\pm}] = 0$$

so that Θ commutes with \bar{R}_{α} , and the whole product $DR_{\mathbf{z}} \Theta \bar{R}_{\alpha}$ can be checked to fulfill (QT1)-(QT4), which can be checked directly using that U, H are both quasitriangular.

Thirdly, we need to verify (QT5) for all generators. But we see that

$$[\Delta(K), R_{\mathbf{z}} \bar{R}_{\alpha}] = [\Delta(E), R_{\mathbf{z}} \bar{R}_{\alpha}] = [\Delta(F), R_{\mathbf{z}} \bar{R}_{\alpha}] = 0$$

so for K, E, F (QT5) follows from its fulfillment in $u_q\mathfrak{sl}_2$. Also for nilpotent generators X_k , we have

$$[\Delta(K_a), \Theta] = [\Delta(X_k), \Theta] = 0,$$

so for K_a, X_k follows from $DR_{\mathbf{z}} \bar{R}_{\alpha}$ being a valid Nenciu type R-matrix. That is $DR_{\mathbf{z}} \bar{R}_{\alpha}$ satisfying (QT5) for K_a and X_k follows from Theorem 2.9. □

Lemma A.3. For the setting of Proposition A.2, the corresponding monodromy matrix is

$$M = D_{21}\Theta_{21}D\Theta(\bar{R}_\alpha)^2 := M_U(\bar{R}_\alpha)^2.$$

where $M_U := D_{21}\Theta_{21}D\Theta$.

Proof. Since Θ and \bar{R}_α commute we can rewrite it as

$$M = R_{\mathbf{z},21}D_{21}\Theta_{21}\bar{R}_{\alpha,21}R_{\mathbf{z}}D\bar{R}_\alpha\Theta$$

Now, we can check that $\bar{R}_{\alpha,21}R_{\mathbf{z}}D\bar{R}_\alpha = R_{\mathbf{z}}D(\bar{R}_\alpha)^2$. Moreover, $R_{\mathbf{z}}$ commutes with D, Θ , so can be brought to $R_{\mathbf{z},21}R_{\mathbf{z}} = \mathbf{1} \otimes \mathbf{1}$ and we find

$$= D_{21}\Theta_{21}D(\bar{R}_\alpha)^2\Theta.$$

We can commute $(\bar{R}_\alpha)^2$ past Θ and retrieve $M = M_U(\bar{R}_\alpha)^2$, as expected. \square

Proposition A.4. Let, using the setting of Proposition A.2, $U = u_q\mathfrak{sl}_2$ carry the ribbon structure $(D\Theta, Ku_U)$ and $H = H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$ carry the ribbon structure $(R_{\mathbf{z}}R_\alpha, g_H u_H)$, where u_U, u_H are Drinfeld elements of $D\Theta$ and $R_{\mathbf{z}}R_\alpha$ respectively, such that

$$R_\alpha = \exp\left(\sum_{l=0}^{t_2} \alpha_l(Z_l^+ \otimes L_l Z_l^- - Z_l^- \otimes L_l Z_l^+)\right),$$

and

$$v_H := \exp\left(-2\sum_{k=1}^{t_2} \alpha_k Z_k^+ Z_k^-\right)$$

with the corresponding pivotal element satisfying $g_H^2 = \mathbf{1}$ and the Drinfeld element

$$u_H := g_H \exp\left(-2\sum_{k=1}^{t_2} \alpha_k Z_k^+ Z_k^-\right).$$

Then $u_q\mathfrak{sl}_2 \times H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$ carries the ribbon structure $(DR_{\mathbf{z}}\Theta\bar{R}_\alpha, v_U u_H)$, with the ribbon element

$$v := v_U u_H =$$

$$\frac{1-i}{\sqrt{r'}} \sum_{a=0}^{r'-1} \sum_{b=0}^{r''-1} \frac{\{-1\}^a}{[a]!} q^{-\frac{(a+3)a}{2}+2b^2} E^a F^a K^{-a-2b-1} g_H \exp\left(-2\sum_{k=1}^{t_2} \alpha_k Z_k^+ Z_k^-\right),$$

for $\bar{L} = K^{r''}L$, with the pivotal element $g = K^{-1}$, and the Drinfeld element

$$u := u_U u_H = \frac{1-i}{\sqrt{r'}} \sum_{a=0}^{r'-1} \sum_{b=0}^{r''-1} \frac{\{-1\}^a}{[a]!} q^{-\frac{(a+3)a}{2}+2b^2} E^a F^a K^{-a-2b} g_H \exp\left(-2\sum_{k=1}^{t_2} \alpha_k Z_k^+ Z_k^-\right).$$

Proof. It is straightforward to find the Drinfeld element of $R = DR_{\mathbf{z}}\Theta\bar{R}_\alpha$ to be $u := u_U u_H$ since the two pieces coming from the respective algebras commute and the piece coming from \bar{R}_α is in fact just u_H as for R_α , because $(K^{r''})^2 = \mathbf{1}$. We need to find a pivotal element $g \in G(U \times H)$ that would make the ribbon element $v := g^{-1}u$ central. This is achieved by choosing $g = K$, as it verifies the criterion for K, E, F by assumption, and $\{K, X_k\} = 0$, which is the criterion necessary for $S^2(X_k) = KX_kK^{-1} = -X_k$. The ribbon axioms can be checked directly, using the relations in $U \times H$ and the fact that both U, H carry ribbon structures that we know explicitly. Let us denote them v_H and v_U . Since $S(v_H) = v_H$ and $S(g_H) = g_H$, we have $S(u_H) = u_H$, because $[g_H, u_H] = 0$. Then we have

$$S(v) = S(v_U u_H) = S(u_H)S(v_U) = u_H v_U = v_U u_H = v.$$

In a similar fashion (R2) is immediate. Finally, for (R3) we found the monodromy matrix in Lemma A.3 above

$$M = D_{21}\Theta_{21}D\Theta(\bar{R}_\alpha)^2 := M_U(\bar{R}_\alpha)^2,$$

so we have, using the commutations established so far

$$M\Delta(v) = M_U(\bar{R}_\alpha)^2\Delta(v_U)\Delta(u_H).$$

Now since $\Delta(g_H)^2\Delta(K)\Delta(K^{-1}) = \mathbf{1} \otimes \mathbf{1}$ we have

$$(\bar{R}_\alpha)^2\Delta(u_H) = (\bar{R}_\alpha)^2\Delta(K)\Delta(K^{-1})\Delta(g_H)^2\Delta(u_H).$$

Before we proceed, let us consider the subalgebra $\mathbb{C}\langle K, K_a, X_k \rangle \subset U \times H$, for $a = 1, \dots, s$ and $k = 1, \dots, t$. It is a Nenciu algebra. Using Theorems 2.9 and 3.21 it can be shown that $DR_{\mathbf{z}}\bar{R}_\alpha$ is an R-matrix with the monodromy $(\bar{R}_\alpha)^2$ for this subalgebra and $\bar{v}_H := K^{-1}g_H u_H$ is a compatible ribbon element with the corresponding pivotal element $g = K$. Hence we have,

$$(\bar{R}_\alpha)^2\Delta(\bar{v}_H) = \bar{v}_H \otimes \bar{v}_H.$$

Therefore, since $[\bar{R}_\alpha, K] = 0$,

$$(\bar{R}_\alpha)^2\Delta(K)\Delta(u_H) = \Delta(K)\Delta(g_H)(\bar{R}_\alpha)^2\Delta(\bar{v}_H) = \Delta(K)\Delta(g_H)(\bar{v}_H \otimes \bar{v}_H) = (u_H \otimes u_H).$$

Plugging it back to the main equation we find

$$M_U(\bar{R}_\alpha)^2\Delta(v_U)\Delta(u_H) = M_U\Delta(v_U)(\bar{R}_\alpha)^2\Delta(u_H) = (v_U \otimes v_U)(u_H \otimes u_H) = v \otimes v.$$

which is what we sought. \square

A.3 Proofs of Propositions 4.21 and 4.25

Proof of Proposition 4.21. What follows is the proof of Theorem 4.13 applied to the Hopf algebra of Example 4.20. We think it informative to spell it out in some details. We split the proof into several pieces.

1. We need to verify the properties of the integrals, see Preliminaries, except for the two-sidedness of the integral which is already not satisfied by the $u_q\mathfrak{sl}_2$ piece [BdR23].

$$\overline{(\text{Id} \otimes \lambda)\Delta(h) = \mathbf{1}\lambda(h):}$$

We know that only the element $E^{r'-1}F^{r'-1}K^{r'-1}X^+X^-$ is picked up by the integral, and it is vanishing for any other element of the monomial basis. We need to verify only those monomial basis elements where all nilpotent generators appear, so of the form $E^{r'-1}F^{r'-1}\mathbf{K}^{\mathbf{w}}X^+X^-$ for some $\mathbf{w} \in \mathbb{Z}_4^3$. Then the only term in the coproduct carrying the top piece in the right factor is

$$\Delta(E^{r'-1}F^{r'-1}\mathbf{K}^{\mathbf{w}}X^+X^-) = K^{1-r'}\mathbf{K}^{\mathbf{w}} \otimes E^{r'-1}F^{r'-1}\mathbf{K}^{\mathbf{w}}X^+X^- + \dots$$

If we now act with $\text{Id} \otimes \lambda$, we see that $\mathbf{K}^{\mathbf{w}} = K^{r'-1}$ is required for the top term to survive, all other vanish. But then we have

$$\Delta(E^{r'-1}F^{r'-1}K_1^{r'-1}X^+X^-) = \mathbf{1} \otimes E^{r'-1}F^{r'-1}K^{r'-1}X^+X^- + \dots$$

so the left factor is proportional to $\mathbf{1}$ as required. Thus, after acting by $\text{Id} \otimes \lambda$, we retrieve the axiom. $\overline{h\Lambda = \Lambda h = \epsilon(h)\Lambda:}$

Now consider Λ multiplying any other element of the monomial basis $h \in H$. If h contains nilpotents, we clearly get $h\Lambda = \Lambda h = \epsilon(h)\Lambda = 0$, because the cointegral of Proposition 4.2 is two-sided, and X^+X^- is central. Moreover, $E^{r'-1}F^{r'-1}X^+X^-$ commutes with all grouplike generators, so by the usual reasoning $K\Lambda = \Lambda K = \epsilon(K)\Lambda = \Lambda$ and $K_a\Lambda = \Lambda K_a = \epsilon(K_a)\Lambda = \Lambda$ for $a = 1, 2$. Thus, Λ is a two-sided cointegral for H .

$$\overline{\lambda(\Lambda)=1:}$$

Satisfied by construction.

2. It is easy to see that $R_{\mathbf{z}}$ commutes with D and Θ . From Remark 4.19 and Proposition 3.24 we know that it fulfills (QT1)-(QT4), as does the product $D\Theta$. Hence the entire R-matrix does. It is left to check (QT5). Firstly, for E and F , we see that $[E, R_{\mathbf{z}}] = [F, R_{\mathbf{z}}] = 0$, so for these two generators (QT5) follows from it being true for $u_q\mathfrak{sl}_2$. Then, for X^\pm , notice that $[\Delta(X^\pm), E \otimes F] = 0$ in $H \otimes H$. Thus, we determine (QT5) only for $R_{\mathbf{z}}D$. The two can be written as $R_{\bar{\mathbf{z}}}$ for

$$\bar{\mathbf{z}} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 2 & 3 \\ 0 & 1 & 0 \end{pmatrix}.$$

This is clearly a block matrix, so (QT5) can be checked for K and K_1, K_2 separately. For the latter this is the same as in Example 3.10, for the former, using Remark 4.3, we have

$$\begin{aligned} \Delta^{\text{cop}}(X^\pm)D &= (X^\pm \otimes \mathbf{1} + (K^{r''}K_1K_2)^{\pm 1} \otimes X^\pm) \sum_{v,w \in \mathbb{Z}_{r'}} (q^2)^{-wv} K^w \otimes K^v = \\ &\sum_{v,w \in \mathbb{Z}_{r'}} (q^2)^{-wv \mp wr''} K^w \otimes K^{v \mp r''} (X^\pm \otimes (K^{r''})^{\pm 1}) + \\ &+ \sum_{v,w \in \mathbb{Z}_{r'}} (q^2)^{-wv \mp vr''} K^{w \pm r''} \otimes K^v ((K_1K_2)^{\pm 1} \otimes X^\pm). \end{aligned}$$

The two sums can be restored to the original form by the substitutions $v \mapsto v \pm r''$ and $w \mapsto w \pm r''$, respectively. The remaining $(K_1K_2)^{\pm 1}$ factors are dealt with in the same way by $R_{\mathbf{z}}$.

3. Note that the modification to the ribbon element is given by K_2^2 , which recall from Remark 4.19 and Proposition 3.24 is the Drinfeld element u_H of $R_{\mathbf{z}}$. Moreover, u_H commutes with v_U , so the only thing left to check is whether K remains the pivotal element. Indeed, $\{K, X^\pm\} = 0$, which is consistent with the order of the antipode $S^4(X^\pm) = X^\pm$.

We can also check the ribbon axioms, as follows. Let v_U be the ribbon element of Proposition 4.2. For (R1), notice that $v = v_U K_2^2$. Immediately we have $\epsilon(v) = \epsilon(v_U)\epsilon(K_2^2) = 1$, since v_U is the ribbon element for $u_q\mathfrak{sl}_2$. Similarly, $S(v) = S(K_2^2)S(v_U) = K_1^2 v_U = v_U K_2^2$, since K_1^2 is of order 2 and commutes with v_U . The axiom (R2) is immediate.

Towards (R3), to verify $R_{21}R\Delta(v) = (v \otimes v)$, note first that the monodromy matrix is

$$R_{21}R = R_{\mathbf{z},21}D_{21}\Theta_{21}R_{\mathbf{z}}D\Theta = R_{\mathbf{z},21}R_{\mathbf{z}}D_{21}\Theta_{21}D\Theta = M_U,$$

where $M_U = D_{21}\Theta_{21}D\Theta$ is the monodromy matrix of $u_q\mathfrak{sl}_2$. This follows as $R_{\mathbf{z}}$ is on its own triangular and commutes with the other components. Now, we have

$$M\Delta(v) = M_U\Delta(v_U)\Delta(K_2^2) = (v_U \otimes v_U)(K_2^2 \otimes K_2^2) = v_U K_2^2 \otimes v_U K_2^2,$$

as required. □

Proof of Proposition 4.25. Most of the proof is the same as in the Proposition 4.21. We have to establish that the new R-matrix and ribbon element fulfil the axioms of a ribbon Hopf algebra. Firstly it is easy to check that

$$[Z^\pm \otimes LZ^\mp, E \otimes F] = [Z^\pm \otimes LZ^\mp, \Delta(E)] = [Z^\pm \otimes LZ^\mp, \Delta(F)] = 0.$$

Thus, the R_α commutes with Θ , while $R_{\mathbf{z}}D$ is its triangular piece of the R-matrix. So we can directly again execute the (QT1)-(QT4) axioms, as well as (QT5), since the appearance R_α does not interrupt their holding for E and F . In light of Theorem 2.9, we can note that the map is block on generators - are no cross-terms between E, F and Z^\pm .

To verify ribbon axioms call again v_U the ribbon element of Proposition 4.2, and let $u_H = K_4^2 \exp(-2\alpha Z^+ Z^-)$, so that $v = v_U u_H$. Then we obviously have $\epsilon(v) = \epsilon(v_U)\epsilon(u_H) = 1$, and we check $S(v_U u_H) = S(u_H v_U) = S(v_U)S(u_H) = v_U S(u_H)$ since both pieces commute. This is because, while u_H is not the ribbon element for the subalgebra $\langle K, K_1, K_2, K_3, X^\pm, Z^\pm \rangle$, but $\exp(-2\alpha Z^+ Z^-)$ is, so we compute

$$S(K_4^2 \exp(-2\alpha Z^+ Z^-)) = \exp(-2\alpha Z^+ Z^-) K_4^2 = u_H,$$

since $\exp(-2\alpha Z^+ Z^-)$ is central.

For the final axiom we need the monodromy matrix. Call M_U the monodromy matrix of $u_q \mathfrak{sl}_2$ and M_H of the subalgebra $\langle K, K_1, K_2, K_3, X^\pm, Z^\pm \rangle$, which can be easily shown to be $M_H = R_\alpha^2$. We can now write

$$M = R_{\mathbf{z},21} D_{21} \Theta_{21} R_{\alpha,21} R_{\mathbf{z}} D \Theta R_\alpha.$$

Since Θ and R_α commute we can rewrite this as

$$= R_{\mathbf{z},21} D_{21} \Theta_{21} R_{\alpha,21} R_{\mathbf{z}} D R_\alpha \Theta$$

Now, we can check that $R_{\alpha,21} R_{\mathbf{z}} D R_\alpha = R_{\mathbf{z}} D R_\alpha^2 = R_{\mathbf{z}} D M_H$. Moreover, $R_{\mathbf{z}}$ commutes with D, Θ , so can be brought to $R_{\mathbf{z},21} R_{\mathbf{z}} = \mathbf{1} \otimes \mathbf{1}$ and we find

$$= D_{21} \Theta_{21} D M_H \Theta.$$

Finally, since $M_H = R_\alpha^2$ we can commute it past Θ and retrieve $M = M_U M_H$. With this we now verify

$$M_U M_H \Delta(v_U u_H) = M_U M_H \Delta(u_H) \Delta(v_U) = M_U M_H \Delta(K_4^2) \Delta(\exp(-2\alpha Z^+ Z^-)) \Delta(v_U).$$

Since $\exp(-2\alpha Z^+ Z^-)$ is central, we easily rearrange

$$M_H \Delta(K_4^2) \Delta(\exp(-2\alpha Z^+ Z^-)) = \Delta(K_4^2) M_H \Delta(\exp(-2\alpha Z^+ Z^-)) = u_H \otimes u_H,$$

since the exponential part is ribbon in H with respect to M_H . We found so far

$$M_U M_H \Delta(v_U u_H) = M_U \Delta(v_U) M_H \Delta(u_H) = M_U \Delta(v_U) (u_H \otimes u_H),$$

but this rearranges to

$$M_U \Delta(v_U) (u_H \otimes u_H) = (v_U \otimes v_U) (u_H \otimes u_H) = (v_U u_H \otimes v_U u_H),$$

as required. \square

B Summary of examples

In this appendix we collect all the examples of both constructions, with their corresponding structures. Throughout we denote

- by Λ the two-sided cointegral
- by λ (respectively λ_L) the two-sided (respectively left) integral
- by R the R-matrix
- by u the Drinfeld element
- by g the pivotal element
- by v the ribbon element.

B.1 Nenciu type examples

Example B.1. Let N_1 be the Hopf algebra generated by K_a, X_k, Z_l^\pm for $a \in \{1, 2\}$, $j, k \in \{1, \dots, t_1\}$; $t_1 \in 4\mathbb{N}$ and $l, m \in \{1, \dots, t_2\}$; $t_2 \in \mathbb{N}$, subject to the following relations

$$K_a^4 = \mathbf{1}, \quad K_a X_j = i X_j K_a, \quad K_1 Z_l^\pm = \pm i Z_l^\pm K_1, \quad K_2 Z_l^\pm = Z_l^\pm K_2, \\ Z_l^\pm X_k = \mp i X_k Z_l^\pm, \quad \{X_j, X_k\} = \{Z_l^\pm, Z_m^\pm\} = \{Z_l^\pm, Z_m^\mp\} = 0,$$

where $i^2 = -1$. The Hopf structure is defined by

$$\epsilon(K_a) = 1, \quad \epsilon(X_k) = \epsilon(Z_l^\pm) = 0,$$

$$\Delta(K_a) = K_a \otimes K_a, \quad \Delta(X_k) = \mathbf{1} \otimes X_k + X_k \otimes K_1 K_2, \quad \Delta(Z_l^\pm) = \mathbf{1} \otimes Z_l^\pm + Z_l^\pm \otimes K_1^{\pm 2} K_2^{\pm 1}, \\ S(K_a) = K_a^{-1}, \quad S(X_k) = -X_k (K_1 K_2)^{-1}, \quad S(Z_l^\pm) = -Z_l^\pm K_1^{\mp 2} K_2^{\mp 1}.$$

The Nenciu data for this Hopf algebra is

- $\mathbf{m} = (4, 4)$, so $s = 2$
- $t = t_1 + 2t_2$
- \mathbf{d} and \mathbf{u} are $t \times 2$ matrices

$$\mathbf{d} = \begin{pmatrix} 1 & 1 \\ \vdots & \vdots \\ 1 & 1 \\ 1 & 0 \\ -1 & 0 \\ \vdots & \vdots \\ 1 & 0 \\ -1 & 0 \end{pmatrix}, \quad \mathbf{u} = \begin{pmatrix} 1 & 1 \\ \vdots & \vdots \\ 1 & 1 \\ 2 & 1 \\ -2 & -1 \\ \vdots & \vdots \\ 2 & 1 \\ -2 & -1 \end{pmatrix}$$

The case of minimal dimension of Example 3.10 occurs for $t_1 = 4$, $t_2 = 1$, and its dimension is $4 \times 4 \times 2^6 = 2^{10} = 1024$. It will be shown later this examples admits only triangular ribbon structures.

Proposition B.2. The algebra N_1 of Example 3.10 carries

$$\Lambda := \left(\sum_{a,b=0}^3 K_1^a K_2^b \right) \prod_{k=1}^{t_1} X_k \prod_{l=1}^{t_2} Z_l^+ \prod_{l=1}^{t_2} Z_l^-$$

$$\lambda(\mathbf{K}^{\mathbf{v}} \mathbf{X}^{\mathbf{r}}) := \begin{cases} 1 & \text{if } \mathbf{v} = (0, 0), \mathbf{r} = (1, 1, \dots, 1) \\ 0 & \text{otherwise.} \end{cases}$$

$$R_{\mathbf{z}} := \frac{1}{16} \sum_{\mathbf{v}, \mathbf{w} \in \mathbb{Z}_4^2} i^{-\mathbf{v}\mathbf{w}^T} \mathbf{K}^{\mathbf{w}} \otimes \mathbf{K}^{\mathbf{v}\mathbf{z}},$$

$$v := \mathbf{1},$$

where $\mathbf{z} = \begin{pmatrix} 2 & 3 \\ 1 & 0 \end{pmatrix}$ and $g = K_1^2$.

Example B.3. Let N_2 be the Hopf algebra generated by K_a , X_k^\pm and Z_l^\pm , for $a = 1, 2, 3$, $j, k = 1, \dots, t_1$; $t_1 \in \mathbb{N}$ and $l, m = 1, \dots, t_2$, $t_2 \in \mathbb{N}$, subject to the following relations

$$K_a^4 = \mathbf{1}, \quad K_a X_k^\pm = \pm i X_k^\pm K_a$$

$$K_1 Z_l^\pm = Z_l^\pm K_1, \quad K_2 Z_l^\pm = -Z_l^\pm K_2, \quad K_3 Z_l^\pm = \pm i Z_l^\pm K_3$$

$$\{X_j^\pm, X_k^\pm\} = \{X_j^\pm, X_k^\mp\} = \{Z_l^\pm, X_k^\pm\} = \{Z_l^\pm, X_k^\mp\} = \{Z_l^\pm, Z_m^\pm\} = \{Z_l^\pm, Z_m^\mp\} = 0.$$

Let also $L := K_3^2$ as a shorthand, note that $L^2 = \mathbf{1}$. The Hopf structure is defined by

$$\epsilon(K_a) = 1, \quad \epsilon(X_k^\pm) = \epsilon(Z_l^\pm) = 0,$$

$$\Delta(K_a) = K_a \otimes K_a, \quad \Delta(X_k^\pm) = \mathbf{1} \otimes X_k^\pm + X_k^\pm \otimes (K_1 K_2)^{\pm 1}, \quad \Delta(Z_l^\pm) = \mathbf{1} \otimes Z_l^\pm + Z_l^\pm \otimes L$$

$$S(K_a) = K_a^{-1} \quad S(X_k^\pm) = -X_k^\pm (K_1 K_2)^{\mp 1} \quad S(Z_l^\pm) = -Z_l^\pm L.$$

. The Nenciu data for this Hopf algebra is

- $\mathbf{m} = (4, 4, 4)$, so $s = 3$
- $t = 2t_1 + 2t_2$
- Then \mathbf{d} and \mathbf{u} are $t \times 3$ matrices

$$\mathbf{d} = \begin{pmatrix} 1 & 1 & 1 \\ -1 & -1 & -1 \\ \vdots & \vdots & \vdots \\ 1 & 1 & 1 \\ -1 & -1 & -1 \\ 0 & 2 & 1 \\ 0 & -2 & -1 \\ \vdots & \vdots & \vdots \\ 0 & 2 & 1 \\ 0 & -2 & -1 \end{pmatrix}, \quad \mathbf{u} = \begin{pmatrix} 1 & 1 & 0 \\ -1 & -1 & 0 \\ \vdots & \vdots & \vdots \\ 1 & 1 & 0 \\ -1 & -1 & 0 \\ 0 & 0 & 2 \\ 0 & 0 & -2 \\ \vdots & \vdots & \vdots \\ 0 & 0 & 2 \\ 0 & 0 & -2 \end{pmatrix}$$

The case of minimal dimension of Example 3.11 occurs for $t_1 = 1$, $t_2 = 1$, and its dimension is $4 \times 4 \times 4 \times 2^4 = 2^{10} = 1024$.

Proposition B.4. The algebra N_2 of Example 3.11 carries

$$\Lambda := \left(\sum_{a,b,c=0}^3 K_1^a K_2^b K_3^c \right) \prod_{k=1}^{t_1} X_k^+ \prod_{k=1}^{t_1} X_k^- \prod_{l=1}^{t_2} Z_l^+ \prod_{l=1}^{t_2} Z_l^-,$$

$$\lambda(\mathbf{K}^{\mathbf{v}} \mathbf{X}^{\mathbf{r}}) := \begin{cases} 1 & \text{if } \mathbf{v} = (0, 0, 0), \mathbf{r} = (1, 1, \dots, 1) \\ 0 & \text{otherwise.} \end{cases},$$

$$R_{\mathbf{z}} R_{\boldsymbol{\alpha}} := \frac{1}{64} \sum_{\mathbf{v}, \mathbf{w} \in \mathbb{Z}_4^3} i^{-\mathbf{v}\mathbf{w}^T} \mathbf{K}^{\mathbf{w}} \otimes \mathbf{K}^{\mathbf{v}\mathbf{z}} \exp \left(\sum_{l=1}^{t_2} \alpha_l (Z_l^+ \otimes LZ_l^- - Z_l^- \otimes LZ_l^+) \right),$$

$$v := \exp \left(-2 \sum_{l=0}^{t_2} \alpha_l Z_l^+ Z_l^- \right),$$

where $\mathbf{z} = \begin{pmatrix} 0 & 3 & 2 \\ 1 & 0 & 0 \\ 2 & 0 & 2 \end{pmatrix}$, $\boldsymbol{\alpha} = (\alpha_1, \dots, \alpha_{t_2}) \in \mathbb{C}^{2t_2}$, and $g = L = K_3^2$.

Example B.5. Let N_3 be the Hopf algebra generated by K_a, X_k^\pm, Y_k^\pm and Z_l^\pm , for $a = 1, 2, 3, j, k = 1, \dots, t_1$; $t_1 \in \mathbb{N}$ and $l, m = 1, \dots, t_2$; $t_2 \in \mathbb{N}$, subject to the following relations

$$K_a^4 = \mathbf{1}, \quad K_a X_k^\pm = \pm i X_k^\pm K_a,$$

$$K_1 Y_k^\pm = \pm i Y_k^\pm K_1, \quad K_2 Y_k^\pm = Y_k^\pm K_2, \quad K_3 Y_k^\pm = \pm i Y_l^\pm K_3$$

$$K_1 Z_l^\pm = Z_l^\pm K_1, \quad K_2 Z_l^\pm = -Z_l^\pm K_2, \quad K_3 Z_l^\pm = \pm i Z_l^\pm K_3$$

$$\{X_j^\pm, X_k^\pm\} = \{X_j^\pm, X_k^\mp\} = \{Y_j^\pm, Y_k^\pm\} = \{Y_j^\pm, Y_k^\mp\}, \quad X_j Y_k = i Y_k X_j$$

$$\{Z_l^\pm, X_k^\pm\} = \{Z_l^\pm, X_k^\mp\} = \{Z_l^\pm, Y_k^\pm\} = \{Z_l^\pm, Y_k^\mp\} = \{Z_l^\pm, Z_m^\pm\} = \{Z_l^\pm, Z_m^\mp\} = 0.$$

Let also $L := K_3^2$ as a shorthand. The Hopf structure is defined by

$$\epsilon(K_a) = 1, \quad \epsilon(X_k^\pm) = \epsilon(Y_k^\pm) = \epsilon(Z_l^\pm) = 0,$$

$$\Delta(K_a) = K_a \otimes K_a, \quad \Delta(X_k^\pm) = \mathbf{1} \otimes X_k^\pm + X_k^\pm \otimes (K_1 K_2)^{\pm 1},$$

$$\Delta(Y_k^\pm) = \mathbf{1} \otimes Y_k^\pm + Y_k^\pm \otimes (K_1 K_2^2)^{\pm 1} \quad \Delta(Z_l^\pm) = \mathbf{1} \otimes Z_l^\pm + Z_l^\pm \otimes L$$

$$S(K_a) = K_a^{-1} \quad S(X_k^\pm) = -X_k^\pm (K_1 K_2)^{\mp 1}, \quad S(Y_k^\pm) = -Y_k^\pm (K_1^2 K_2)^{\mp 1} \quad S(Z_l^\pm) = -Z_l^\pm L.$$

- $\mathbf{m} = (4, 4, 4)$, so $s = 3$
- $t = 4t_1 + 2t_2$
- Then \mathbf{d} and \mathbf{u} are $t \times 3$ matrices

$$\mathbf{d} = \begin{pmatrix} 1 & 1 & 1 \\ -1 & -1 & -1 \\ & \vdots & \\ 1 & 1 & 1 \\ -1 & -1 & -1 \\ 1 & 0 & 1 \\ -1 & 0 & -1 \\ & \vdots & \\ 1 & 0 & 1 \\ -1 & 0 & -1 \\ 0 & 2 & 1 \\ 0 & -2 & -1 \\ & \vdots & \\ 0 & 2 & 1 \\ 0 & -2 & -1 \end{pmatrix}, \quad \mathbf{u} = \begin{pmatrix} 1 & 1 & 0 \\ -1 & -1 & 0 \\ & \vdots & \\ 1 & 1 & 0 \\ -1 & -1 & 0 \\ 2 & 1 & 0 \\ -2 & -1 & 0 \\ & \vdots & \\ 2 & 1 & 0 \\ -2 & -1 & 0 \\ 0 & 0 & 2 \\ 0 & 0 & -2 \\ & \vdots & \\ 0 & 0 & 2 \\ 0 & 0 & -2 \end{pmatrix}$$

The case of minimal dimension of Example 3.12 occurs for $t_1 = 1, t_2 = 1$, and its dimension is $4 \times 4 \times 4 \times 2^6 = 2^{12} = 4096$.

Proposition B.6. The algebra N_3 of Example 3.12 carries

$$\Lambda := \left(\sum_{a,b,c=0}^3 K_1^a K_2^b K_3^c \right) \prod_{k=1}^{t_1} X_k^+ \prod_{k=1}^{t_1} X_k^- \prod_{k=1}^{t_1} Y_k^+ \prod_{k=1}^{t_1} Y_k^- \prod_{l=1}^{t_2} Z_l^+ \prod_{l=1}^{t_2} Z_l^-,$$

$$\lambda(\mathbf{K}^{\mathbf{v}} \mathbf{X}^{\mathbf{r}}) := \begin{cases} 1 & \text{if } \mathbf{v} = (0, 0, 0), \mathbf{r} = (1, 1, \dots, 1) \\ 0 & \text{otherwise.} \end{cases}$$

$$R_{\mathbf{z}} R_{\boldsymbol{\alpha}} := \frac{1}{64} \sum_{\mathbf{v}, \mathbf{w} \in \mathbb{Z}_4^3} i^{-\mathbf{v}\mathbf{w}^T} \mathbf{K}^{\mathbf{w}} \otimes \mathbf{K}^{\mathbf{v}\mathbf{z}} \exp \left(\sum_{l=1}^{t_2} \alpha_l (Z_l^+ \otimes LZ_l^- - Z_l^- \otimes LZ_l^+) \right),$$

$$v := \exp \left(-2 \sum_{k=1}^{t_2} \alpha_k Z_k^+ Z_k^- \right),$$

where $\mathbf{z} = \begin{pmatrix} 0 & 3 & 2 \\ 1 & 0 & 0 \\ 2 & 0 & 2 \end{pmatrix}$, $\boldsymbol{\alpha} = (\alpha_1, \dots, \alpha_{t_2}) \in \mathbb{C}^{2t_2}$ and $g = L = K_3^2$.

Example B.7. Let $N_4 := H(\mathbf{m}, t, \mathbf{d}, \mathbf{u})$ be a Nenciu type algebra generated by grouplike K_1, K_2 and nilpotent X^\pm generators, and the following relations

$$K_1^4 = K_2^4 = \mathbf{1}, \quad (X^\pm)^2 = 0,$$

$$KX^\pm K^{-1} = q^{r'} X^\pm = -X^\pm, \quad K_1 X^\pm K_2^{-1} = K_2 X^\pm K_2^{-1} = \pm i X^\pm.$$

Let $L := K^{r'/2} K_1 K_2$ for convenience. The Hopf structure is

$$\epsilon(K_1) = \epsilon(K_2) = 1, \quad \epsilon(X^\pm) = 0,$$

$$\Delta(K_1) = K_1 \otimes K_1, \quad \Delta(K_2) = K_2 \otimes K_2, \quad \Delta(X^\pm) = \mathbf{1} \otimes X^\pm + X^\pm \otimes L^{\pm 1}.$$

and

$$S(K_1) = K_1^{-1}, \quad S(K_2) = K_2^{-1}, \quad S(X^\pm) = -X^\pm L^{\mp 1}.$$

Proposition B.8. The Hopf algebra N_4 of Example 4.17 carries

$$\begin{aligned} \Lambda &:= \left(\sum_{a,b=0}^3 K_1^a K_2^b \right) X^+ X^- \\ \lambda(\mathbf{K}^{\mathbf{v}} \mathbf{X}^{\mathbf{r}}) &:= \begin{cases} 1 & \text{if } \mathbf{v} = (0, 0), \mathbf{r} = (1, 1) \\ 0 & \text{otherwise.} \end{cases} \\ R_{\mathbf{z}} &:= \frac{1}{16} \sum_{\mathbf{v}, \mathbf{w} \in \mathbb{Z}_4^2} i^{-\mathbf{v}\mathbf{w}^T} \mathbf{K}^{\mathbf{w}} \otimes \mathbf{K}^{\mathbf{v}\mathbf{z}}, \\ v &:= \mathbf{1}, \end{aligned}$$

where $\mathbf{z} = \begin{pmatrix} 2 & 3 \\ 1 & 0 \end{pmatrix}$ and $g = K_1^2$.

B.2 Non-factorizable extensions of $u_q \mathfrak{sl}_2$ examples

Example B.9. Let $u_q \mathfrak{sl}_2 \rtimes H$, where $H = \text{SF}_2$ be a Hopf algebra generated by K, E, F with relations and morphisms of Definition 4.1, as well as K_1, Z^\pm with the following relations

$$K_2^2 = \mathbf{1}, \quad (Z^\pm)^2 = 0,$$

$$KZ^\pm K^{-1} = q^{r'} Z^\pm = -Z^\pm, \quad K_1 Z^\pm K_2^{-1} = -Z^\pm.$$

$$[K_1, K] = [K_1, E] = [K_1, F] = 0$$

$$EZ^\pm = q^{r'} Z^\pm E = -Z^\pm E, \quad [Z^\pm, F] = 0.$$

Let $L := K^{r'/2} K_1$ for convenience. The Hopf structure is

$$\epsilon(K_1) = 1, \quad \epsilon(Z^\pm) = 0,$$

$$\Delta(K_1) = K_1 \otimes K_1, \quad \Delta(Z^\pm) = \mathbf{1} \otimes Z^\pm + Z^\pm \otimes L^{\pm 1}.$$

and

$$S(K_1) = K_1^{-1}, \quad S(Z^\pm) = -Z^\pm L^{\mp 1}.$$

The instance of smallest dimension with a non-trivial commutation relation $[E, F]$ occurs at $r = 8$. The dimension is then $4^3 \times 2^3 = 2^8 = 512$.

Proposition B.10. The algebra $u_q \mathfrak{sl}_2 \rtimes \text{SF}_2$ carries

$$\begin{aligned} \Lambda &:= \frac{\{1\}^{r'-1}}{\sqrt{r'} [r'-1]!} \sum_{a=0}^{r'-1} \sum_b^2 E^{r'-1} F^{r'-1} K^a L^b Z^+ Z^-, \\ \lambda_L(E^e F^f K^a L^b (Z^+)^g (Z^-)^h) &:= \\ &= \frac{\sqrt{r'} [r'-1]!}{\{1\}^{r'-1}} \delta_{a,r'-1} \delta_{b,0} \delta_{e,r'-1} \delta_{f,r'-1} \delta_{g,1} \delta_{h,1}, \\ R &:= R_{\mathbf{z}} D \Theta \bar{R}_{\boldsymbol{\alpha}} \\ \bar{R}_{\boldsymbol{\alpha}} &:= \exp(\alpha(Z^+ \otimes \bar{L} Z^- - Z^- \otimes \bar{L} Z^+)) \text{ for } \bar{L} = K^{r'/2} L, \\ v &:= \frac{1-i}{\sqrt{r'}} \sum_{a=0}^{r'-1} \sum_{b=0}^{r'-1} \frac{\{-1\}^a}{[a]!} q^{-\frac{(a+3)a}{2} + 2b^2} E^a F^a K^{-a-2b-1} L \exp(-2\alpha Z^+ Z^-), \end{aligned}$$

where D, Θ were defined in Proposition 4.2, $R_{\mathbf{z}}$ in Proposition 3.22, and $g = K$.

Example B.11. Let $u_q\mathfrak{sl}_2 \times N_4$, be the Hopf algebra generated by K, E, F with relations and morphisms of Definition 4.1, as well as K_1, K_2, X^\pm with the following relations

$$\begin{aligned} K_1^4 &= K_2^4 = \mathbf{1}, \quad (X^\pm)^2 = 0, \\ KX^\pm K^{-1} &= q^{r'} X^\pm = -X^\pm, \quad K_1 X^\pm K_2^{-1} = K_2 X^\pm K_3^{-1} = \pm i X^\pm, \\ [K_1, K_2] &= [K_1, K] = [K_1, E] = [K_1, F] = [K_2, K] = [K_2, E] = [K_2, F] = 0 \\ EX^\pm &= q^{r'} X^\pm E = -X^\pm E, \quad [X^\pm, F] = 0. \end{aligned}$$

Let $L := K^{r'/2} K_1 K_2$ for convenience. The Hopf structure is

$$\begin{aligned} \epsilon(K_1) &= \epsilon(K_2) = 1, \quad \epsilon(X^\pm) = 0, \\ \Delta(K_1) &= K_1 \otimes K_1, \quad \Delta(K_2) = K_2 \otimes K_2, \quad \Delta(X^\pm) = \mathbf{1} \otimes X^\pm + X^\pm \otimes L^{\pm 1}. \end{aligned}$$

and

$$S(K_1) = K_1^{-1}, \quad S(K_2) = K_2^{-1}, \quad S(X^\pm) = -X^\pm L^{\mp 1}.$$

The instance of smallest dimension with a non-trivial commutation relation $[E, F]$ occurs at $r = 8$. The dimension is then $4^3 \times 2^4 \times 2^2 = 2^{12} = 4096$.

Proposition B.12. The algebra $u_q\mathfrak{sl}_2 \times N_4$ carries

$$\begin{aligned} \Lambda &:= \frac{\{1\}^{r'-1}}{\sqrt{r'}[r'-1]!} \sum_{a=0}^{r'-1} \sum_{b,c=0}^4 E^{r'-1} F^{r'-1} K^a K_1^b K_2^c X^+ X^-, \\ \lambda_L(E^e F^f K^a K_1^b K_2^c (X^+)^g (X^-)^h) &:= \\ &= \frac{\sqrt{r'}[r'-1]!}{\{1\}^{r'-1}} \delta_{a,r'-1} \delta_{b,0} \delta_{c,0} \delta_{e,r'-1} \delta_{f,r'-1} \delta_{g,1} \delta_{h,1}, \\ R &:= R_{\mathbf{z}} D \Theta \\ R_{\mathbf{z}} &:= \frac{1}{16} \sum_{\mathbf{v}, \mathbf{w} \in \mathbb{Z}_4^2} i^{-\mathbf{v}\mathbf{w}^T} (K_1, K_2)^{\mathbf{w}} \otimes (K_1, K_2)^{\mathbf{v}\mathbf{z}}, \\ v &:= \frac{1-i}{\sqrt{r'}} \sum_{a=0}^{r'-1} \sum_{b=0}^{r''-1} \frac{\{-1\}^a}{[a]!} q^{-\frac{(a+3)a}{2} + 2b^2} E^a F^a K^{-a-2b-1} K_2^2, \end{aligned}$$

where D, Θ were defined in Proposition 4.2, $\mathbf{z} = \begin{pmatrix} 2 & 3 \\ 1 & 0 \end{pmatrix}$, and $g = K$.

Example B.13. Let $u_q\mathfrak{sl}_2 \times N_2$, be the Hopf algebra generated by K, E, F with relations and morphisms of Definition 4.1, as well as $K_1, K_2, K_3, X^\pm, Z^\pm$ with the following relations, for $a = 1, 2, 3$:

$$\begin{aligned} [K_a, K] &= [K_a, E] = [K_a, F] = 0, \quad KX^\pm K^{-1} = -X^\pm \quad K_a^4 = \mathbf{1}, \quad K_a X^\pm = \pm i X^\pm K_a \\ KZ^\pm K^{-1} &= -Z^\pm \quad K_1 Z^\pm K_2^{-1} = Z^\pm, \quad K_2 Z^\pm K_3^{-1} = -Z^\pm, \quad K_3 Z^\pm K_4^{-1} = \pm i Z^\pm \\ \{X^\pm, X^\pm\} &= \{X^\pm, X^\mp\} = \{Z^\pm, X^\pm\} = \{Z^\pm, X^\mp\} = \{Z^\pm, Z^\pm\} = \{Z^\pm, Z^\mp\} = 0 \\ EX^\pm &= -X^\pm E, \quad EZ^\pm = -Z^\pm E, \quad [X^\pm, F] = [Z^\pm, F] = 0. \end{aligned}$$

Let also $L := K^{r'/2} K_3^2$ as a shorthand, note that this time $L^2 = \mathbf{1}$. The Hopf structure is defined by

$$\begin{aligned} \epsilon(K_a) &= 1, \quad \epsilon(X^\pm) = \epsilon(Z^\pm) = 0, \\ \Delta(K_a) &= K_a \otimes K_a, \quad \Delta(X^\pm) = \mathbf{1} \otimes X^\pm + X^\pm \otimes (K^{r'/2} K_1 K_2)^{\pm 1}, \quad \Delta(Z^\pm) = \mathbf{1} \otimes Z^\pm + Z^\pm \otimes L \\ S(K_a) &= K_a^{-1} \quad S(X^\pm) = -X^\pm (K^{r'/2} K_1 K_2)^{\mp 1} \quad S(Z^\pm) = -Z^\pm L. \end{aligned}$$

The instance of smallest dimension occurs when q is a root of unity of order 8 and the dimension is $4^4 \times 2^4 \times 2^2 \times 2^2 = 2^{16} = 65536$.

Proposition B.14. The algebra $u_q\mathfrak{sl}_2 \times N_2$ carries

$$\begin{aligned} \Lambda &:= \frac{\{1\}^{r'-1}}{\sqrt{r'}[r'-1]!} \sum_{a=0}^{r'-1} \sum_{b,c,d=0}^4 E^{r'-1} F^{r'-1} K^a K_1^b K_2^c K_3^d X^+ X^- Z^+ Z^-, \\ \lambda_L(E^e F^f K^a K_1^b K_2^c K_3^d (X^+)^g (X^-)^h (Z^+)^i (Z^-)^j) &:= \\ &= \frac{\sqrt{r'}[r'-1]!}{\{1\}^{r'-1}} \delta_{a,r'-1} \delta_{b,0} \delta_{c,0} \delta_{d,0} \delta_{e,r'-1} \delta_{f,r'-1} \delta_{g,1} \delta_{h,1} \delta_{i,1} \delta_{j,1}, \\ R &:= R_{\mathbf{z}} D \Theta R_{\alpha}, \\ R_{\mathbf{z}} &:= \frac{1}{64} \sum_{\mathbf{v}, \mathbf{w} \in \mathbb{Z}_4^3} i^{-\mathbf{v}\mathbf{w}^T} (K_1, K_2, K_3)^{\mathbf{w}} \otimes (K_1, K_2, K_3)^{\mathbf{v}\mathbf{z}}, \\ R_{\alpha} &:= \exp(\alpha(Z^+ \otimes LZ^- - Z^- \otimes LZ^+)), \\ v &:= \frac{1-i}{\sqrt{r'}} \sum_{a=0}^{r'-1} \sum_{b=0}^{r''-1} \frac{\{-1\}^a}{[a]!} q^{-\frac{(a+3)a}{2} + 2b^2} K^{-a-2b-1} K_3^2 E^a F^a \exp(-2\alpha Z^+ Z^-), \end{aligned}$$

where D, Θ were defined in Proposition 4.2, $\mathbf{z} = \begin{pmatrix} 0 & 3 & 2 \\ 1 & 0 & 0 \\ 2 & 0 & 2 \end{pmatrix}$, and $g = K$.

References

- [AC65] James J Andrews and Morton L Curtis. Free groups and handlebodies. *Proceedings of the American Mathematical Society*, 16(2):192–195, 1965.
- [AEG02] Nicolas Andruskiewitsch, Pavel Etingof, and Shlomo Gelaki. Triangular hopf algebras with the chevalley property, 2002.
- [AS06] N. Andruskiewitsch and H. J. Schneider. On the classification of finite-dimensional pointed hopf algebras, 2006.
- [BBDRP23] Anna Beliakova, Ivelina Bobtcheva, Marco De Renzi, and Riccardo Piergallini. On algebraization in low-dimensional topology. *arXiv.org*, Dec 2023.
- [BDG00] M Beattie, S Dăscălescu, and L Grünenfelder. Constructing pointed hopf algebras by ore extensions. *Journal of Algebra*, 225(2):743–770, Mar 2000.
- [BDR22] Anna Beliakova and Marco De Renzi. Refined bobtcheva-messia invariants of 4-dimensional 2-handlebodies. *arXiv preprint arXiv:2205.11385*, 2022.
- [BdR23] Anna Beliakova and Marco de Renzi. Kerler–lyubashenko functors on 4-dimensional 2-handlebodies. *International mathematics research notices*, Feb 2023.
- [Bob23] Ivelina Bobtcheva. Algebraic characterisation of the category of cobordisms of 2-dimensional cw-complexes and the andrews-curtis conjecture. *arXiv.org*, Sep 2023.
- [BP06] Ivelina Bobtcheva and Riccardo Piergallini. A universal invariant of four-dimensional 2-handlebodies and three-manifolds, 2006.
- [CGHPM23] Francesco Costantino, Nathan Geer, Benjamin Haioun, and Bertrand Patureau-Mirand. Skein (3+1)-tqfts from non-semisimple ribbon categories, 2023.
- [CP95] Vyjayanthi Chari and Andrew N Pressley. *A Guide to Quantum Groups*. Cambridge University Press, Jul 1995.
- [CY93] Louis Crane and David N Yetter. A categorical construction of 4d tqfts. *arXiv preprint hep-th/9301062*, 1993.
- [EGNO16] Pavel Etingof, Shlomo Gelaki, Dmitri Nikshych, and Victor Ostrik. *Tensor Categories*. American Mathematical Soc., Aug 2016.
- [FGS24] Matthieu Faitg, Azat M Gainutdinov, and Christoph Schweigert. Davydov–yetter cohomology and relative homological algebra. *Selecta Mathematica*, 30(2), Feb 2024.
- [GR17] A.M. Gainutdinov and I. Runkel. Symplectic fermions and a quasi-hopf algebra structure on $\bar{U}_q sl(2)$. *Journal of Algebra*, 476:415–458, Apr 2017.
- [GS99] Robert E Gompf and András Stipsicz. *4-manifolds and Kirby Calculus*. American Mathematical Soc., 1999.
- [Kas12] Christian Kassel. *Quantum Groups*. Springer Science & Business Media, Dec 2012.
- [Ker03] Thomas Kerler. Homology tqft’s and the alexander–reidemeister invariant of 3-manifolds via hopf algebras and skein theory. *Canadian Journal of Mathematics*, 55(4):766–821, 2003.
- [KL01] Thomas Kerler and Volodymyr V Lyubashenko. *Non-semisimple topological quantum field theories for 3-manifolds with corners*, volume 1765. Springer Science & Business Media, 2001.
- [Maj00] Shahn Majid. *Foundations of quantum group theory*. Cambridge University Press, Cambridge ; New York, 2000.
- [Mol77] Richard K Molnar. Semi-direct products of hopf algebras. *Journal of Algebra*, 47(1):29–51, Jul 1977.
- [Nen04] Adriana Nenciu. Quasitriangular pointed hopf algebras constructed by ore extensions. *Algebras and Representation Theory*, 7(2):159–172, May 2004.
- [PO99] Florin Panaite and Freddy Van Oystaeyen. Quasitriangular structures for some pointed hopf algebras of dimension 2^n . *Communications in Algebra*, 27(10):4929–4942, Jan 1999.
- [Rad94] David E Radford. On kauffman’s knot invariants arising from finite-dimensional hopf algebras. *CRC Press eBooks*, page 205–266, 1994.
- [Rad12] David E Radford. *Hopf Algebras*. World Scientific, 2012.
- [Shi16] Kenichi Shimizu. Non-degeneracy conditions for braided finite tensor categories, 2016.
- [Tur94] Vladimir G Turaev. *Quantum invariants of knots and 3-manifolds*. de Gruyter, 1994.

INSTITUTE OF MATHEMATICS, UNIVERSITY OF ZURICH, WINTERTHURERSTRASSE 190, CH-8057 ZURICH, SWITZERLAND
E-mail address: `quentin.faes@math.uzh.ch`

INSTITUTE OF MATHEMATICS, UNIVERSITY OF ZURICH, WINTERTHURERSTRASSE 190, CH-8057 ZURICH, SWITZERLAND
E-mail address: `maksymilian.manko@math.uzh.ch`